

Quasinormal modes of a d -dimensional regular black hole featuring an integrable singularity

Zhongzhinan Dong^{1,*}, Dan Zhang^{1,†}, Guoyang Fu^{2,‡} and Jian-Pin Wu^{1,§}

¹ *Center for Gravitation and Cosmology,
College of Physical Science and Technology,
Yangzhou University, Yangzhou 225009, China*

² *Department of Physics and Astronomy,
Shanghai Jiao Tong University, Shanghai 200240, China*

Abstract

In this paper, we exhaustively investigate the quasinormal modes (QNMs) of a probe scalar field over a d -dimensional regular black hole (BH) characterized by the parameter A . The quasinormal frequencies (QNFs) exhibit different behaviors with respect to the parameter A for $d = 4$ and $d > 4$. Firstly, the trends of QNFs with respect to A exhibit completely opposite patterns for the case of $d = 4$ and $d > 4$. Secondly, in the 4-dimensional regular BH, a non-monotonic behavior with respect to A is observed in the imaginary part of the fundamental modes with vanishing angular quantum number. In contrast, this non-monotonic behavior only appears in the overtones when $d > 4$. Thirdly, an overtone outburst accompanied by an oscillatory pattern is observed only in the case of $d > 4$, but not in $d = 4$.

*dzzn_edu@163.com

†danzhanglnk@163.com

‡FuguoyangEDU@163.com

§jianpinwu@yzu.edu.cn

Contents

I. Introduction	2
II. Scalar field over the regular black hole	5
III. QUASINORMAL MODES	7
A. Fundamental modes	9
B. Overtones	12
IV. Conclusions and discussions	15
Acknowledgments	16
References	16

I. INTRODUCTION

Regular black holes (BHs) were initially proposed as a solution to avoid the core singularity that is inherent in ordinary BHs. The scalar curvature of these regular BHs remains finite everywhere, including at the core. In contrast to ordinary BHs, regular BHs have complete null and timelike geodesics in their spacetime. These BHs can be formed by incorporating quantum gravity effects [1–4], exotic matter like the nonlinear Maxwell field [5–7] that typically defy the standard energy conditions. For comprehensive reviews on regular BHs, refer to [8–10].

Regular BHs can be usually classified into two categories based on their asymptotic behavior near the center: those with a de-Sitter (dS) core and those with a Minkowski core. Regular BHs with a dS core feature an energy density that converges to a non-zero constant as one approaches the center of the BH, specifically, $\lim_{r \rightarrow 0} \rho(r) = \rho_0 \neq 0$. These consist of the widely recognized Bardeen BH [1], Hayward BH [2] and Frolov BH [3]. We would like to mention that in [11], the authors introduce a novel regular BH solution, which greatly extends the Hayward class. This solution incorporates mass-dependent regulators, resulting in significant, percent-level effects on observables for regular astrophysical BHs.

The BH with Minkowski core was first proposed in [12] by incorporating an exponentially

suppressed Newton potential. This solution has been further extended to encompass a category of regular BHs featuring various kinds of exponential potentials [4, 13–20]. In contrast to regular BHs with a dS core, those with a Minkowski core have an energy density that approaches zero at the center, specially, $\lim_{r \rightarrow 0} \rho(r) = 0$.

These regular BHs typically possess distinct outer and inner horizons, each exhibiting nonzero surface gravity. The presence of a nonzero inner horizon surface gravity typically leads to an exponential instability within the vicinity of the inner horizon. This phenomenon is known as the mass inflation instability [21, 22]. In order to circumvent the mass inflation instability, the authors in [23] propose a regular BH model where a surface gravity becomes zero at the inner horizon while preserving non-zero at the outer horizon. In contrast to scenarios with finite surface gravity at the inner horizon, this model does not exhibit an exponential growth associated with mass inflation instability [23].

Recently, an alternative approach to avoid the problem of mass inflation instability was proposed in [24]. The key idea is to incorporate quantum matter described by regular wave functions, resulting in a novel regular BH geometry with an integrable singularity¹ and no Cauchy horizon in spherically symmetric configurations. Following Ref. [24], we refer to this novel regular BH as the one with localized sources of matter (LSM). In contrast to those regular BHs that possess clearly defined outer and inner Cauchy horizons, this novel regular BH proposed in [24] allows radial geodesics to reach the core of the BH. Subsequently, this solution was extended to encompass higher-dimensional scenarios [26]. In this paper, we will study the characteristics of the quasinormal modes (QNMs) of a probe scalar field over this novel d-dimension regular BH.

During the ringdown phase, the BH emits gravitational waves (GWs) with characteristic discrete frequencies known as quasinormal frequencies (QNFs). The frequencies encode information regarding the decaying scales and damped oscillations of the BH [27]. Studying the properties of QNFs offers an opportunity to implement the strong field tests of general relativity (GR) [28–30]. Additionally, it allows for testing the nature of compact remnants formed after a coalescence [31] as well as potential imprint of quantum gravity [32–45]. Nevertheless, most regular BHs are commonly formed by integrating quantum gravity effects

¹ Integrable singularities are defined as the locations where the effective energy-momentum tensor and curvature invariants become infinite, but their “volume” integrals are still finite [25].

at the phenomenological level, which complicates the task of establishing consistently effective equations for gravitational perturbations. Fortunately, even when merely a probe matter field is taken into account over these regular BHs, their QNM spectra are likewise affected by the background spacetime. As a result, these QNM spectra may be utilized to phenomenologically simulate the GW form during the ringdown phase and also offer vital insights into the internal structure of the BH [30, 32–37, 39, 43, 44, 46–51].

At first, the majority of research efforts concentrated on the fundamental modes of QNM spectra. Nevertheless, the examination of GW data from LIGO/Virgo has lately shown compelling evidence in favor of the presence of overtone patterns [52, 53]. Compared to results obtained exclusively from the fundamental modes, incorporating overtones theoretically provide more exact information regarding the mass and spin of the remnant BH [54]. Moreover, several studies have emphasized the importance of overtones in accurately simulating the ringdown phase [54–57]. These results challenge the prevailing belief that the fundamental mode dominates the signal primarily and imply that the quasinormal ringing might begin earlier than previously anticipated.

Furthermore, a recent study has revealed a connection between the properties of the high overtones and the geometric structure surrounding the event horizon of a BH [58]. The research discovered that even a slight modification near the event horizon might have a substantial influence on the first few overtones [58], so offering an opportunity to explore the structure of the event horizon. In addition, in contrast to echoes, overtones display a significantly higher energy contribution, rendering them an essential domain for investigating diverse BHs [59–61]. These significant discoveries offer additional impetus for our ongoing exploration of the overtones of different BHs. Considering the aforementioned motivations, we will study the characteristics of both the fundamental modes and the high overtone modes of the scalar perturbation of the novel regular BHs [26]. In addition to studying the implications of the model’s characteristic parameters, we will also investigate the influence of extra dimensions.

This paper is structured as follows. In Section II, we provide a brief review of this regular BH with LSM, derive the expression for the scalar field equation over this d -dimensional BH, and then discuss the properties of the corresponding effective potentials. In Section III, we use the pseudo-spectral method (PSM) to obtain the QNFs and analyze their characteristics. Finally, the conclusions and discussions are presented in Section IV.

II. SCALAR FIELD OVER THE REGULAR BLACK HOLE

Following the framework presented in [24], the authors of [26] construct a d -dimensional analytical regular BH featuring an integrable singularity and no an internal horizon by incorporating the LSM. The model's exterior geometry can be written as [26]:

$$\begin{aligned} ds^2 &= -f(r)dt^2 + f(r)^{-1}dr^2 + r^2d\Omega_{d-2}, \\ f(r) &= 1 - \frac{2G_d m(r)}{r^{d-3}}, \end{aligned} \quad (1)$$

where $m(r)$ is the mass function with

$$m(r) = \frac{Mr^{d-3}}{A + r^{d-3}}. \quad (2)$$

Here, $d\Omega_{d-2}$ represents the transversal section of a $(d-2)$ -dimensional sphere, G_d denotes the d -dimensional Newton constant, A is a positive constant measured in units of ℓ_p^{d-3} , and M , quantified in units of ℓ_p^{-1} , corresponds to the mass parameter, where ℓ_p refers to Planck units. Then, the energy density can be explicitly calculated as [26]:

$$\rho = \frac{d-2}{8\pi} \frac{A(d-3)M}{r^2(A + r^{d-3})^2}. \quad (3)$$

From the above equations (2) and (3), it is evident that the mass function m and the energy density ρ exhibit behaviors of $m \sim r^{d-3}$ and $\rho \sim r^{-2}$ near the origin, respectively. The asymptotic behaviors of m and ρ guarantee that there is no internal horizon linked to an unstable core, and that the energy-momentum tensor and Ricci scalar are integrable at the origin. In addition, it can be shown that the radial timelike geodesic and the tidal forces of this model are finite [26]. Furthermore, the model's central integrable singularity is weak. This gravitationally weak singularity hence will not crush any object following a radial timelike world-line.

The Hawking temperature of this regular BH can be calculated as follows:

$$T = \frac{(d-3)M}{2\pi} \frac{r_h^{d-4}}{(A + r_h^{d-3})^2}. \quad (4)$$

Here, $r_h = (2M - A)^{1/(d-3)}$, with $A \leq 2M$, is the event horizon of the BH and is determined by taking the positive root of $f(r) = 0$. In this paper, without loss of generality, we set $M = 1/2$, implying $A \leq 1$. From Eq. (4), it is evident that when $d = 4$, the Hawking temperature approaches a constant value as A tends to 1. Nevertheless, for dimensions

greater than 4 ($d > 4$), the Hawking temperature decreases towards zero as A approaches 1. To visualize this picture, we present the Hawking temperature as a function of the parameter A in Fig.1.




FIG. 1: The Hawking temperature T as a function of the parameter A in different dimensions.

Next, we investigate how this regular BH responds to perturbations from a massless scalar field Φ . The scalar field's dynamics may be described using the Klein-Gordon (KG) equation:

$$\frac{1}{\sqrt{-g}}\partial_{\mu}(\sqrt{-g}g^{\mu\nu}\partial_{\nu}\Phi) = 0, \quad (5)$$

where $g_{\mu\nu}$ is the background metric. We will analyze the eigenvalue problem associated with the dynamics of the scalar field in the frequency domain. Given the hyperspherical symmetry of the spacetime under investigation, we will use hyperspherical harmonics to separate variables for the perturbation field Φ . Thus, the perturbation field Φ can be expressed as:

$$\Phi(t, r, \theta_1, \theta_2, \dots, \theta_n) = \int d\omega \sum_{l,m} r^{-(d-2)/2} \Psi_{lm}(r) Y_{lm}(\theta_1, \theta_2, \dots, \theta_n) e^{-i\omega t}, \quad (6)$$

where $Y_{lm}(\theta_1, \theta_2, \dots, \theta_n)$ is the hyperspherical harmonics, with l and m representing the angular and azimuthal quantum numbers, respectively. Then, the KG equation (5) can be

transformed into a Schrödinger-like form:

$$\frac{d^2\Psi}{dr_*^2} + (\omega^2 - V(r_*))\Psi(r_*) = 0. \quad (7)$$

Here, r_* is the tortoise coordinate, defined by the relation r by $dr_*/dr = 1/f(r)$. The effective potential V in the above equation is given by:

$$V(r) = f(r) \left(\frac{l(l+d-3)}{r^2} + \frac{(d-2)(d-4)}{4r^2} f(r) + \frac{d-2}{2r} \frac{df(r)}{dr} \right), \quad (8)$$

where the angular quantum number l takes the integer values of $0, 1, \dots$. For simplicity, we have removed the subscript l and m from $\Psi_{l,m}(t, r)$ and denoted it as Ψ in Eq.(7). In the effective potential V , the first and the second terms are the centrifugal potential, while the third term corresponds to the gravitational potential. In 4 dimensional spacetime, when $l = 0$, the centrifugal potential vanishes. While for $d > 4$, the centrifugal potential persists even when $l = 0$. Fig.2 and Fig.3 illustrate the effective potentials $V(r)$ for various values of the parameter A and different spacetime dimension d . The potentials are positive definite, ensuring the stability of the scalar field perturbations.




FIG. 2: The effective potential $V_{eff}(r)$ for $d=4$ with different parameter A

III. QUASINORMAL MODES

In this section, we will explore the characteristics of the QNMs associated with this regular BH, with a particular focus on the influence of the parameter A and the spacetime dimension d . The QNMs are determined by solving an eigenvalue problem that requires a




FIG. 3: The effective potential $V_{eff}(r)$ for $d=5$ with different parameter A

purely outgoing wave at spatial infinity ($r_* \rightarrow +\infty$) and a purely ingoing wave at the event horizon ($r_* \rightarrow -\infty$), specially,

$$\Psi \sim e^{+i\omega r_*}, \quad r_* \rightarrow +\infty, \quad (9)$$

$$\Psi \sim e^{-i\omega r_*}, \quad r_* \rightarrow -\infty. \quad (10)$$

The boundary conditions described above characterize the black hole's response to a transient perturbation after the source has stopped acting [27, 62, 63].

Numerous techniques have been developed to calculate QNMs, including the WKB method [64–69], the asymptotic iteration method (AIM) [70–72], the Horowitz-Hubeny method [73], the continued fraction method (CFM) [74], and the PSM [75, 76]. In this paper, we utilize the PSM to compute the QNM spectra, as it is a highly effective numerical tool [32–36, 75–82], particularly adept at identifying overtone modes [33–35].

The primary feature of the PSM is its discretization of differential equations, leading to the solution of corresponding generalized eigenvalue problems. In particular, continuous variables are substituted with a discrete set of collocation points, known as grid points, while functions are expanded using particular basis functions called cardinal functions. Typically, Chebyshev grids and Lagrange cardinal functions are used, defined as follows:

$$x_i = \cos\left(\frac{i}{N}\pi\right), \quad C_j(x) = \prod_{j=0, j \neq i}^N \frac{x - x_j}{x_i - x_j}, \quad i = 0, \dots, N. \quad (11)$$

When applying the PSM, it is advantageous to use the Eddington-Finkelstein (EF) coordinate, as this renders the wave equation (7) linear with respect to the frequency ω . To

achieve this, we perform the following coordinate transformation:

$$\begin{aligned} r &\rightarrow (1/u^{d-3} - A)^{1/(d-3)}, \\ v &= t + r_*. \end{aligned} \quad (12)$$

Under this transformation, the metric (1) becomes

$$ds_{EF}^2 = -f(u)dv^2 - 2dvdu(1 - Au^d)^{-(d-4)/(d-3)}u^{-2} + (u^{-d} - A)^{2/(d-3)}d\Omega_{d-2}, \quad (13)$$

where $f(u) = 1 - u^d$.

Now, based on the hyperspherical harmonics, the perturbation field Φ in the EF coordinate can be expressed as follows:

$$\Phi(v, u, \theta_1, \theta_2, \dots, \theta_n) = \int d\omega \sum_{l,m} u^{(d-4)/2} \psi_{lm}(u) Y_{lm}(\theta_1, \theta_2, \dots, \theta_n) e^{-i\omega v}. \quad (14)$$

Then, the wave equation (7) turns into the following form:

$$\begin{aligned} &u^2 f(u)(Au^{d-3} - 1)^2 \psi''(u) + (Au^{d-3} - 1)[2(d-3)Au^{d-2}f(u) \\ &+ u^2(Au^{d-3} - 1)f'(u) - 2i\omega(1 - Au^{d-3})^{1/(d-3)}] \psi'(u) + \left[-l(d+l-3) \right. \\ &- 2i\omega u^{-1}(1 - Au^{d-3})^{d-3}(1 + (d-4)Au^{d-3}/2) + (Au^{d-3} - 1)((d-2)(d-4)/4 \\ &\left. + (5 + 3(d-5))(d-4)Au^{d-3}/4)f(u) + u(d-4)(Au^{d-3} - 1)f'(u)/2 \right] \psi(u) = 0. \end{aligned} \quad (15)$$

Combining with the boundary conditions (9) and (10), we can solve Eq (15) using the PSM.

A. Fundamental modes

We firstly discuss the case of $l = 0$. Fig.4 and Fig.5 illustrate QNFs of fundamental modes as a function of the parameter A in 4-dimensional and 5-dimensional spacetimes, respectively. In 4-dimensional spacetime, the real part of QNFs, $\text{Re}(\omega)$, increases as A increases. Notably, as A approaches 1, $\text{Re}(\omega)$ experiences a more rapid rise. This suggests that the parameter A strengthens the oscillation. However, we observe that the absolute value of the imaginary part $|\text{Im}(\omega)|$ shows a non-monotonic change with respect to A . Specifically, as A increases, $|\text{Im}(\omega)|$ first increases, signifying a faster decay modes of this system, and then rapidly decreases when A approaches 1. In contrast, in the 5-dimensional spacetime, as A increases, $\text{Re}(\omega)$ increases almost linearly. Meanwhile, $|\text{Im}(\omega)|$ decreases with increasing A , which differs from the behavior observed in the 4-dimensional case.




FIG. 4: QNFs in 4-dimensional spacetime as a function of the parameter A for $l = 0$ and $n = 0$.




FIG. 5: QNFs in 5-dimensional spacetime as a function of the parameter A for $l = 0$ and $n = 0$.




FIG. 6: QNFs in 4-dimensional spacetime as a function of the parameter A for $l = 1$ and $n = 0$.




FIG. 7: QNFs in 5-dimensional spacetime as a function of the parameter A for $l = 1$ and $n = 0$.

When the angular quantum number l is turned on, it is observed that as A approaches 1, the real part $\text{Re}(\omega)$ of QNFs in 4-dimensional spacetime exhibits a more rapid increase compared to the $l = 0$ case (left plot in Fig.6). In the 5-dimensional spacetime, in sharp contrast to the $l = 0$ scenario, $\text{Re}\omega$ also rapidly increase as A approaches 1 when l is turned on (left plot in Fig.7). Additionally, the previously non-monotonic behavior of $\text{Im}\omega$ in 4-dimensional spacetime for $l = 0$ becomes almost linear increasing once l is turned on (right plot in Fig.6), while in 5-dimensional spacetime, the decrease becomes more rapid as A approaches 1 compared to the $l = 0$ case (right plot in Fig.7).

Typically, the fundamental modes are associated with variations in the peak of the effective potential, allowing us to understand the behaviors of QNFs in different scenarios. In 4-dimensional spacetime, the peak values and their positions change significantly as A increases, particularly as A approaches 1 for $l = 1$ compared to $l = 0$. However, this relative change between $l = 0$ and $l = 1$ in 5-dimensional spacetime is less pronounced. These changes in the peak of the effective potential are consistent with the variations in the QNFs of the fundamental modes.

To further clearly illustrate the effects of spacetime dimension d and the the angular quantum number l , we also show the the phase diagram of the real part $\text{Re}(\omega)$ and the absolute value of the imaginary part $|\text{Im}(\omega)|$ of the fundamental modes for various d and l in Fig.8. Green points represent the Schwarzschild case, i.e., $A = 0$, while the final points correspond to $A = 0.9$. The non-monotonic behavior is visible only in $\text{Im}\omega$ for $l = 0$ and $d = 4$. For $l > 0$ and $d > 4$, this non-monotonic behavior disappears, likely due to the pres-




FIG. 8: The phase diagram $\text{Re}(\omega) - |\text{Im}(\omega)|$ of the fundamental modes for various spacetime dimensions d and angular quantum numbers l . Green points represent the Schwarzschild case, i.e., $A = 0$, while the final points correspond to $A = 0.9$.

ence of a centrifugal potential that significantly alters the shape of the effective potential near its peak, suppressing the effect of the parameter A . Moreover, when both l and d are fixed, the real oscillation frequency of the fundamental mode increases monotonically with A . This increase is more pronounced in 4-dimensional spacetime compared to higher dimensions, suggesting that the influence of A on oscillatory behaviors reduces as d increases. Additionally, as A increases, the absolute value of the imaginary part $|\text{Im}(\omega)|$ rapidly decreases for dimensions beyond $d = 4$, indicating that the impact of A on the decay behavior of the QNFs becomes significant in higher dimensions.

B. Overtones

In this subsection, we explore the characteristics of the overtones. Fig.9 displays the QNFs of the overtones in 4-dimensional spacetime as a function of the parameter A for $l = 0$. It is evident that no overtone outburst is observed in this model with $d = 4$. Instead, the QNFs of the overtones exhibit similar behaviors to the fundamental modes with respect to A . What's slightly different is that as A approaches 1, $\text{Re}(\omega)$ increases rapidly while $|\text{Im}(\omega)|$ decreases with increasing overtone number n .

Next, we turn to study the case of $l = 1$ (see Fig.10). We find that the behaviors of




FIG. 9: QNFs of the overtones in 4-dimensional spacetime as a function of the parameter A for $l = 0$.




FIG. 10: QNFs of the overtones in 4-dimensional spacetime as a function of the parameter A for $l = 1$.

$\text{Re}(\omega)$ for the overtones is also similar to that of $l = 0$. However, compared to the $l = 0$ case, the non-monotonic behavior in $|\text{Im}(\omega)|$ vanishes for overtone numbers $n = 1, 2, 3, 4, 5$. This non-monotonic behavior reemerges for $n = 6, 7$. This observation suggests that the angular quantum number l suppresses the effects of quantum gravity or modified gravity. Similar observations have been reported in previous studies [34–36].

Now, we turn our attention to investigating the properties of overtones in 5-dimensional spacetime. As depicted in Fig.11, the QNFs of overtones in 5-dimensional spacetime are shown as a function of the parameter A for $l = 0$. In contrast to the case in 4-dimensional spacetime, an obvious overtone outburst is observed in 5-dimensional spacetime for higher




FIG. 11: QNFs of the overtones in 5-dimensional spacetime as a function of the parameter A for $l = 0$.




FIG. 12: QNFs of the overtones in 5-dimensional spacetime as a function of the parameter A for $l = 1$.

overtones ($n = 2, n = 3$). The quantum gravity effect is the cause behind this overtone outburst. This phenomenon has been widely noticed in modified gravity theories and effective quantum gravity models such as [30, 32–35, 46–51, 58]. After the initial outburst of overtones, an oscillatory behavior is detected. This particular pattern has been witnessed in RN-BH [83, 84] and other effective quantum gravity corrected BH [33–35]. However, when the angular quantum number l is activated, that is, $l > 0$, no overtone outburst is detected as shown in Fig.12. This further demonstrates that the angular quantum number l has a suppressive effect on the quantum gravity effect. Finally, for the convenience of readers, we also provide the QNF values for various spacetime dimensions d as well as for different values of A , l and n .

		$l = 1$		$l = 2$	
d	n	$A = 0$	$A = 0.2$	$A = 0$	$A = 0.2$
4	0	0.58587-0.19532i	0.67905-0.20406i	0.96729-0.19352i	1.12070-0.20253i
	1	0.52890-0.61251i	0.63092-0.63392i	0.92770-0.59121i	1.08747-0.61635i
	2	0.45908-1.08027i	0.56983-1.10604i	0.86109-1.01712i	1.03115-1.05316i
	3	0.40650-1.57660i	0.52200-1.60459i	0.78772-1.47619i	0.96766-1.51747i
5	0	1.01602-0.36233i	1.07415-0.355029i	1.51057-0.35754i	1.59582-0.34636i
	1	0.85638-1.15761i	0.94842-1.10679i	1.39272-1.10456i	1.50265-1.06501i
	2	0.66344-2.10918i	0.79353-1.98378i	1.19383-1.94574i	1.34691-1.85663i
	3	0.54135-3.13670i	0.69099-2.92551i	0.99443-2.89898i	1.18765-2.73406i
6	0	1.44651-0.50927i	1.48985-0.48545i	2.01153-0.50194i	2.07078-0.47964i
	1	1.14354-1.64146i	1.24912-1.54847i	1.78604-1.55879i	1.88947-1.48378i
	2	0.76082-3.11901i	0.93661-2.87161i	1.37829-2.81127i	1.56918-2.63945i
	3	0.58272-4.73201i	0.77050-4.31972i	1.00617-4.33625i	1.26325-3.99930i
7	0	1.88140-0.64108i	1.91560-0.60953i	2.49678-0.63188i	2.54151-0.60249i
	1	1.39266-2.06571i	1.52289-1.94852i	2.13727-1.96117i	2.24641-1.86623i
	2	0.76366-4.17660i	0.98524-3.79518i	1.40117-3.64147i	1.66855-3.39827i
	3	0.59931-6.39379i	0.79794-5.79251i	0.88946-5.89767i	1.20542-5.36718i
8	0	2.32073-0.76102i	2.34835-0.72396i	2.97469-0.75056i	3.00978-0.71618i
	1	1.60031-2.42111i	1.76649-2.29865i	2.45747-2.31307i	2.57798-2.20968i
	2	0.74822-5.32442i	0.98259-4.80134i	1.19049-4.55135i	1.58856-4.17849i
	3	0.63532-8.09335i	0.82107-7.32006i	0.79398-7.60744i	1.11959-6.87478i

TABLE I: The QNMs for the massless scalar field perturbation with different values of d , n , l and A by PSM.

IV. CONCLUSIONS AND DISCUSSIONS

In this paper, we systematically investigated the properties of the QNMs associated with a probe scalar field over a regular BH featuring an integrable singularity, which is characterized by the parameter A . Our primary focus is on how the QNFs vary with changes in the parameter A . We also explore the influence of the angular quantum number l and the spacetime dimension d on the QNFs.

In 4-dimensional spacetime, a non-monotonic behavior in the absolute value of the imaginary part of QNFs emerges with respect to the parameter A for vanishing angular quantum number l . This behavior can be ascribed to the effective quantum gravity effects, as discussed in references [32–36]. Notably, when the angular quantum number l is turned on, this

non-monotonic behavior ceases to exist, suggesting that the influence of the angular quantum number l supersedes the effects of effective quantum gravity [32–36]. Furthermore, our analysis reveals that this non-monotonic behavior reappears in higher overtones, indicating that the effective quantum gravity effects regain dominance in these higher overtones.

An additional significant finding of this study is the observation that an overtone outburst accompanied by an oscillatory pattern occurs exclusively when the dimensionality d exceeds 4. This phenomenon is notably absent in the scenario where $d = 4$. This distinction underscores the critical role of dimensionality in influencing the dynamic behavior of the system under investigation. The underlying reasons for this distinction warrant further investigation in future research.

Acknowledgments

This work is supported by National Key R&D Program of China (No. 2020YFC2201400), the Natural Science Foundation of China under Grants No. 12375055, 12347159 and 12447151.

-
- [1] J. Bardeen, *Non-singular general relativistic gravitational collapse*, in Proceedings of the 5th International Conference on Gravitation and the Theory of Relativity, p. 87, 1968.
 - [2] S. A. Hayward, *Formation and evaporation of regular black holes*, Phys. Rev. Lett. **96** (2006) 031103, [[gr-qc/0506126](#)].
 - [3] V. P. Frolov, *Notes on nonsingular models of black holes*, Phys. Rev. D **94** (2016), no. 10 104056, [[arXiv:1609.01758](#)].
 - [4] Y. Ling and M.-H. Wu, *Regular black holes with sub-Planckian curvature*, Class. Quant. Grav. **40** (2023), no. 7 075009, [[arXiv:2109.05974](#)].
 - [5] E. Ayon-Beato and A. Garcia, *Regular black hole in general relativity coupled to nonlinear electrodynamics*, Phys. Rev. Lett. **80** (1998) 5056–5059, [[gr-qc/9911046](#)].
 - [6] I. Dymnikova, *Regular electrically charged structures in nonlinear electrodynamics coupled to general relativity*, Class. Quant. Grav. **21** (2004) 4417–4429, [[gr-qc/0407072](#)].

- [7] M. Cataldo and A. Garcia, *Regular (2+1)-dimensional black holes within nonlinear electrodynamics*, Phys. Rev. D **61** (2000) 084003, [[hep-th/0004177](#)].
- [8] C. Bambi, Regular Black Holes. Springer Series in Astrophysics and Cosmology. Springer Singapore, 2023.
- [9] S. Vagnozzi et al., *Horizon-scale tests of gravity theories and fundamental physics from the Event Horizon Telescope image of Sagittarius A*, Class. Quant. Grav. **40** (2023), no. 16 165007, [[arXiv:2205.07787](#)].
- [10] C. Lan, H. Yang, Y. Guo, and Y.-G. Miao, *Regular Black Holes: A Short Topic Review*, Int. J. Theor. Phys. **62** (2023), no. 9 202, [[arXiv:2303.11696](#)].
- [11] J. Boos and C. D. Carone, *Note on black holes with kilometer-scale ultraviolet regulators*, [arXiv:2311.16319](#).
- [12] L. Xiang, Y. Ling, and Y. G. Shen, *Singularities and the Finale of Black Hole Evaporation*, Int. J. Mod. Phys. D **22** (2013) 1342016, [[arXiv:1305.3851](#)].
- [13] H. Culetu, *On a regular modified Schwarzschild spacetime*, [arXiv:1305.5964](#).
- [14] H. Culetu, *On a regular charged black hole with a nonlinear electric source*, Int. J. Theor. Phys. **54** (2015), no. 8 2855–2863, [[arXiv:1408.3334](#)].
- [15] M. E. Rodrigues, E. L. B. Junior, G. T. Marques, and V. T. Zanchin, *Regular black holes in $f(R)$ gravity coupled to nonlinear electrodynamics*, Phys. Rev. D **94** (2016), no. 2 024062, [[arXiv:1511.00569](#)]. [Addendum: Phys.Rev.D 94, 049904 (2016)].
- [16] A. Simpson and M. Visser, *Regular black holes with asymptotically Minkowski cores*, Universe **6** (2019), no. 1 8, [[arXiv:1911.01020](#)].
- [17] S. G. Ghosh, *A nonsingular rotating black hole*, Eur. Phys. J. C **75** (2015), no. 11 532, [[arXiv:1408.5668](#)].
- [18] S. G. Ghosh, D. V. Singh, and S. D. Maharaj, *Regular black holes in Einstein-Gauss-Bonnet gravity*, Phys. Rev. D **97** (2018), no. 10 104050.
- [19] X. Li, Y. Ling, Y.-G. Shen, C.-Z. Liu, H.-S. He, and L.-F. Xu, *Generalized uncertainty principles, effective Newton constant and the regular black hole*, Annals Phys. **396** (2018) 334–350, [[arXiv:1611.09016](#)].
- [20] M. Martinis and N. Perkovic, *Is exponential metric a natural space-time metric of Newtonian gravity?*, [arXiv:1009.6017](#).
- [21] E. Poisson and W. Israel, *Internal structure of black holes*, Phys. Rev. D **41** (1990)

1796–1809.

- [22] E. G. Brown, R. B. Mann, and L. Modesto, *Mass Inflation in the Loop Black Hole*, Phys. Rev. D **84** (2011) 104041, [[arXiv:1104.3126](#)].
- [23] R. Carballo-Rubio, F. Di Filippo, S. Liberati, C. Pacilio, and M. Visser, *Regular black holes without mass inflation instability*, JHEP **09** (2022) 118, [[arXiv:2205.13556](#)].
- [24] R. Casadio, A. Giusti, and J. Ovalle, *Quantum rotating black holes*, JHEP **05** (2023) 118, [[arXiv:2303.02713](#)].
- [25] V. N. Lukash and V. N. Strokov, *Space-Times with Integrable Singularity*, Int. J. Mod. Phys. A **28** (2013) 1350007, [[arXiv:1301.5544](#)].
- [26] M. Estrada, G. Alencar, T. M. Crispim, and C. R. Muniz, *New models of d-dimensional black holes without inner horizon and with an integrable singularity*, JCAP **06** (2024) 042, [[arXiv:2310.01734](#)].
- [27] E. Berti, V. Cardoso, and A. O. Starinets, *Quasinormal modes of black holes and black branes*, Class. Quant. Grav. **26** (2009) 163001, [[arXiv:0905.2975](#)].
- [28] **LIGO Scientific, VIRGO, KAGRA** Collaboration, R. Abbott et al., *Tests of General Relativity with GWTC-3*, [arXiv:2112.06861](#).
- [29] E. Berti et al., *Testing General Relativity with Present and Future Astrophysical Observations*, Class. Quant. Grav. **32** (2015) 243001, [[arXiv:1501.07274](#)].
- [30] E. Berti, K. Yagi, H. Yang, and N. Yunes, *Extreme Gravity Tests with Gravitational Waves from Compact Binary Coalescences: (II) Ringdown*, Gen. Rel. Grav. **50** (2018), no. 5 49, [[arXiv:1801.03587](#)].
- [31] V. Cardoso and P. Pani, *Testing the nature of dark compact objects: a status report*, Living Rev. Rel. **22** (2019), no. 1 4, [[arXiv:1904.05363](#)].
- [32] G. Fu, D. Zhang, P. Liu, X.-M. Kuang, Q. Pan, and J.-P. Wu, *Quasinormal modes and Hawking radiation of a charged Weyl black hole*, Phys. Rev. D **107** (2023), no. 4 044049, [[arXiv:2207.12927](#)].
- [33] G. Fu, D. Zhang, P. Liu, X.-M. Kuang, and J.-P. Wu, *Peculiar properties in quasinormal spectra from loop quantum gravity effect*, Phys. Rev. D **109** (2024), no. 2 026010, [[arXiv:2301.08421](#)].
- [34] H. Gong, S. Li, D. Zhang, G. Fu, and J.-P. Wu, *Quasinormal modes of quantum-corrected black holes*, [arXiv:2312.17639](#).

- [35] D. Zhang, H. Gong, G. Fu, J.-P. Wu, and Q. Pan, *Quasinormal modes of a regular black hole with sub-Planckian curvature*, Eur. Phys. J. C **84** (2024), no. 6 564, [[arXiv:2402.15085](#)].
- [36] Z. Song, H. Gong, H.-L. Li, G. Fu, L.-G. Zhu, and J.-P. Wu, *Quasinormal modes and ringdown waveform of the Frolov black hole*, [arXiv:2406.04787](#).
- [37] L.-G. Zhu, G. Fu, S. Li, D. Zhang, and J.-P. Wu, *Quasinormal modes of a charged loop quantum black hole*, [arXiv:2410.00543](#).
- [38] C. Tang, Y. Ling, Q.-Q. Jiang, and G.-P. Li, *Quasinormal modes of regular black holes with sub-Planckian curvature and Minkowskian core*, Eur. Phys. J. C **84** (2024), no. 12 1296, [[arXiv:2411.01764](#)].
- [39] R. A. Konoplya and O. S. Stashko, *Probing the Effective Quantum Gravity via Quasinormal Modes and Shadows of Black Holes*, [arXiv:2408.02578](#).
- [40] A. S. Koshelev, C. Li, and A. Tokareva, *Quasi-normal modes in non-perturbative quantum gravity*, [arXiv:2412.02678](#).
- [41] O. Stashko, *Quasinormal modes and gray-body factors of regular black holes in asymptotically safe gravity*, Phys. Rev. D **110** (2024), no. 8 084016, [[arXiv:2407.07892](#)].
- [42] A. Dubinsky, *Quasinormal modes of charged black holes in Asymptotically Safe Gravity*, Phys. Dark Univ. **46** (2024) 101657, [[arXiv:2405.08262](#)].
- [43] D. M. Gingrich, *Quasinormal modes of a nonsingular spherically symmetric black hole effective model with holonomy corrections*, [arXiv:2404.04447](#).
- [44] R. A. Konoplya and A. Zhidenko, *Infinite tower of higher-curvature corrections: Quasinormal modes and late-time behavior of D-dimensional regular black holes*, Phys. Rev. D **109** (2024), no. 10 104005, [[arXiv:2403.07848](#)].
- [45] A. F. Zinhailo,
Beyond Einstein: Quasinormal modes and Hawking radiation in higher curvature corrected and asymptotically safe gravity
 PhD thesis, Silesian U., Czech Republic, 2024.
- [46] E. Berti, V. Cardoso, and C. M. Will, *On gravitational-wave spectroscopy of massive black holes with the space interferometer LISA*, Phys. Rev. D **73** (2006) 064030, [[gr-qc/0512160](#)].
- [47] F. Moura and J. a. Rodrigues, *Eikonal quasinormal modes and shadow of string-corrected d-dimensional black holes*, Phys. Lett. B **819** (2021) 136407, [[arXiv:2103.09302](#)].
- [48] F. Moura and J. a. Rodrigues, *Asymptotic quasinormal modes of string-theoretical d-dimensional black holes*, JHEP **08** (2021) 078, [[arXiv:2105.02616](#)].

- [49] F. Moura and J. a. Rodrigues, *The isospectrality of asymptotic quasinormal modes of large Gauss-Bonnet d -dimensional black holes*, Nucl. Phys. B **993** (2023) 116255, [[arXiv:2206.11377](#)].
- [50] J. Lin, M. Bravo-Gaete, and X. Zhang, *Quasinormal modes, greybody factors, and thermodynamics of four dimensional AdS black holes in critical gravity*, Phys. Rev. D **109** (2024), no. 10 104039, [[arXiv:2401.02045](#)].
- [51] R. Ghosh, M. Rahman, and A. K. Mishra, *Regularized stable Kerr black hole: cosmic censorship, shadow and quasi-normal modes*, Eur. Phys. J. C **83** (2023), no. 1 91, [[arXiv:2209.12291](#)].
- [52] M. Isi, *Testing the no-hair theorem with LIGO and Virgo*, in 3rd World Summit on Exploring the Dark Side of the Universe, pp. 49–58, 2020.
- [53] **LIGO Scientific, Virgo** Collaboration, R. Abbott et al., *Tests of general relativity with binary black holes from the second LIGO-Virgo gravitational-wave transient catalog*, Phys. Rev. D **103** (2021), no. 12 122002, [[arXiv:2010.14529](#)].
- [54] M. Giesler, M. Isi, M. A. Scheel, and S. Teukolsky, *Black Hole Ringdown: The Importance of Overtones*, Phys. Rev. X **9** (2019), no. 4 041060, [[arXiv:1903.08284](#)].
- [55] N. Oshita, *Ease of excitation of black hole ringing: Quantifying the importance of overtones by the excitation factors*, Phys. Rev. D **104** (2021), no. 12 124032, [[arXiv:2109.09757](#)].
- [56] X. J. Forteza and P. Mourier, *High-overtone fits to numerical relativity ringdowns: Beyond the dismissed $n=8$ special tone*, Phys. Rev. D **104** (2021), no. 12 124072, [[arXiv:2107.11829](#)].
- [57] N. Oshita, *Thermal ringdown of a Kerr black hole: overtone excitation, Fermi-Dirac statistics and greybody factor*, JCAP **04** (2023) 013, [[arXiv:2208.02923](#)].
- [58] R. A. Konoplya and A. Zhidenko, *First few overtones probe the event horizon geometry*, [[arXiv:2209.00679](#)].
- [59] R. A. Konoplya, A. F. Zinhailo, J. Kunz, Z. Stuchlik, and A. Zhidenko, *Quasinormal ringing of regular black holes in asymptotically safe gravity: the importance of overtones*, JCAP **10** (2022) 091, [[arXiv:2206.14714](#)].
- [60] R. A. Konoplya, *Quasinormal modes in higher-derivative gravity: Testing the black hole parametrization and sensitivity of overtones*, Phys. Rev. D **107** (2023), no. 6 064039, [[arXiv:2210.14506](#)].

- [61] R. A. Konoplya, Z. Stuchlik, A. Zhidenko, and A. F. Zinhailo, *Quasinormal modes of renormalization group improved Dymnikova regular black holes*, Phys. Rev. D **107** (2023), no. 10 104050, [[arXiv:2303.01987](#)].
- [62] R. A. Konoplya and A. Zhidenko, *Quasinormal modes of black holes: From astrophysics to string theory*, Rev. Mod. Phys. **83** (2011) 793–836, [[arXiv:1102.4014](#)].
- [63] K. D. Kokkotas and B. G. Schmidt, *Quasinormal modes of stars and black holes*, Living Rev. Rel. **2** (1999) 2, [[gr-qc/9909058](#)].
- [64] V. Ferrari and B. Mashhoon, *New approach to the quasinormal modes of a black hole*, Phys. Rev. D **30** (1984) 295–304.
- [65] B. F. Schutz and C. M. Will, *BLACK HOLE NORMAL MODES: A SEMIANALYTIC APPROACH*, Astrophys. J. Lett. **291** (1985) L33–L36.
- [66] S. Iyer and C. M. Will, *Black Hole Normal Modes: A WKB Approach. 1. Foundations and Application of a Higher Order WKB Analysis of Potential Barrier Scattering*, Phys. Rev. D **35** (1987) 3621.
- [67] S. Iyer, *BLACK HOLE NORMAL MODES: A WKB APPROACH. 2. SCHWARZSCHILD BLACK HOLES*, Phys. Rev. D **35** (1987) 3632.
- [68] R. A. Konoplya, *Quasinormal behavior of the d -dimensional Schwarzschild black hole and higher order WKB approach*, Phys. Rev. D **68** (2003) 024018, [[gr-qc/0303052](#)].
- [69] J. Matyjasek and M. Opala, *Quasinormal modes of black holes. The improved semianalytic approach*, Phys. Rev. D **96** (2017), no. 2 024011, [[arXiv:1704.00361](#)].
- [70] H. Ciftci, R. L. Hall, and N. Saad, *Perturbation theory in a framework of iteration methods*, Phys. Lett. A **340** (2005) 388–396, [[math-ph/0504056](#)].
- [71] H. T. Cho, A. S. Cornell, J. Doukas, and W. Naylor, *Black hole quasinormal modes using the asymptotic iteration method*, Class. Quant. Grav. **27** (2010) 155004, [[arXiv:0912.2740](#)].
- [72] H. T. Cho, A. S. Cornell, J. Doukas, T. R. Huang, and W. Naylor, *A New Approach to Black Hole Quasinormal Modes: A Review of the Asymptotic Iteration Method*, Adv. Math. Phys. **2012** (2012) 281705, [[arXiv:1111.5024](#)].
- [73] G. T. Horowitz and V. E. Hubeny, *Quasinormal modes of AdS black holes and the approach to thermal equilibrium*, Phys. Rev. D **62** (2000) 024027, [[hep-th/9909056](#)].
- [74] E. W. Leaver, *An Analytic representation for the quasi normal modes of Kerr black holes*, Proc. Roy. Soc. Lond. A **402** (1985) 285–298.

- [75] J. P. Boyd, Chebyshev and Fourier spectral methods. Courier Corporation, 2001.
- [76] A. Jansen, *Overdamped modes in Schwarzschild-de Sitter and a Mathematica package for the numerical computation of quasinormal modes*, Eur. Phys. J. Plus **132** (2017), no. 12 546, [[arXiv:1709.09178](#)].
- [77] J.-P. Wu and P. Liu, *Quasi-normal modes of holographic system with Weyl correction and momentum dissipation*, Phys. Lett. B **780** (2018) 616–621, [[arXiv:1804.10897](#)].
- [78] G. Fu and J.-P. Wu, *EM Duality and Quasinormal Modes from Higher Derivatives with Homogeneous Disorder*, Adv. High Energy Phys. **2019** (2019) 5472310, [[arXiv:1812.11522](#)].
- [79] P. Liu, C. Niu, and C.-Y. Zhang, *Linear instability of charged massless scalar perturbation in regularized 4D charged Einstein-Gauss-Bonnet anti de-Sitter black holes*, Chin. Phys. C **45** (2021), no. 2 025111.
- [80] K. Destounis, R. P. Macedo, E. Berti, V. Cardoso, and J. L. Jaramillo, *Pseudospectrum of Reissner-Nordström black holes: Quasinormal mode instability and universality*, Phys. Rev. D **104** (2021), no. 8 084091, [[arXiv:2107.09673](#)].
- [81] J. L. Jaramillo, R. Panosso Macedo, and L. A. Sheikh, *Gravitational Wave Signatures of Black Hole Quasinormal Mode Instability*, Phys. Rev. Lett. **128** (2022), no. 21 211102, [[arXiv:2105.03451](#)].
- [82] W. Xiong, P. Liu, C.-Y. Zhang, and C. Niu, *Quasinormal modes of the Einstein-Maxwell-aether black hole*, Phys. Rev. D **106** (2022), no. 6 064057, [[arXiv:2112.12523](#)].
- [83] E. Berti and K. D. Kokkotas, *Asymptotic quasinormal modes of Reissner-Nordstrom and Kerr black holes*, Phys. Rev. D **68** (2003) 044027, [[hep-th/0303029](#)].
- [84] J. Jing and Q. Pan, *Quasinormal modes and second order thermodynamic phase transition for Reissner-Nordstrom black hole*, Phys. Lett. B **660** (2008) 13–18, [[arXiv:0802.0043](#)].