

THE SMALLEST NORMALIZED SIGNLESS ∞ -LAPLACIAN EIGENVALUE FOR NON-BIPARTITE CONNECTED GRAPHS

YI DAI

ABSTRACT. In this paper, we aim to study the smallest normalized signless ∞ -Laplacian eigenvalue μ_∞ , a generalisation of the smallest signless Laplacian eigenvalue. For a non-bipartite connected graph, we show that the invariant μ_∞ equals to the reciprocal of the minimal ∞ -norm of the generalized inverses of the weighted signless incidence matrix. An example is also given to illustrate the result.

1. INTRODUCTION

The graph invariants of Laplacian matrix and signless Laplacian matrix are well studied, see [2, 8, 12, 14, 15, 16, 17, 18]. As one of typical objects in the linear spectral theory, the eigenvalues of Laplacian matrix and signless Laplacian matrix are directly related to the connectedness and bipartiteness of graph.

Let $\Omega = (V, E)$ be a non-bipartite connected graph. Note that the signless Laplacian matrix Q of Ω is positive semi-definite. For $X = (x_i, v_i \in V) \in \mathbb{R}^V$, the quadratic form of Q is

$$X^T Q X = \sum_{i \sim j} (x_i + x_j)^2 = \|BX\|_2^2,$$

where $i \sim j$ denotes vertex v_i is adjacent to v_j . The smallest eigenvalue of Q is

$$q(\Omega) = \min_{X \in \mathbb{R}^V \setminus \{0\}} \frac{X^T Q X}{\|X\|_2^2} = \min_{X \in \mathbb{R}^V \setminus \{0\}} \frac{\|BX\|_2^2}{\|X\|_2^2} = \min_{\|X\|_2^2=1} \sum_{i \sim j} (x_i + x_j)^2.$$

In [9], Desai and Rao showed that the smallest eigenvalue $q(\Omega)$ is bounded above and below by functions of the measure of the bipartiteness of Ω .

The signless p -Laplacian is a nonlinear generalization of the quadratic form of the signless Laplacian matrix. For any $p \geq 1$, the smallest signless p -Laplacian eigenvalue

$$q_p(\Omega) = \min_{X \in \mathbb{R}^V \setminus \{0\}} \frac{\|BX\|_p^p}{\|X\|_p^p} = \min_{\|X\|_p^p=1} \sum_{i \sim j} (x_i + x_j)^p.$$

In [4], Borba and Schwerdtfeger proved that

$$q_\infty(\Omega) = \min_{\|X\|_\infty=1} \max_{i \sim j} |x_i + x_j| = \frac{2}{\max_{u \in V} l(u)},$$

Date: December 29, 2024.

2020 Mathematics Subject Classification. 90C47; 90C27; 15A60; 15A09; 05C50.

Key words and phrases. Min-max optimization, signless Laplacian eigenvalue, generalized inverse, norm.

where $l(u)$ is the length of the smallest odd cycle in graph Ω . In [6], Chang proved that the first nonzero 1-Laplacian eigenvalue equals to Cheeger's constant, which had only some upper and lower bounds in 2-Laplacian spectral theory. In [7], Chang, Shao, and Zhang introduced and developed the non-linear spectral theory of signless 1-Laplacian. They showed how the non-bipartiteness of a graph is related to the smallest normalized signless 1-Laplacian, which can be rewritten as follows:

$$\min_{\|X\|_1=1} \|\Delta_1 X\|_1 = \min_{\|X\|_1=1} \sum_{i \sim j} \left| \frac{x_i}{d_i} + \frac{x_j}{d_j} \right|.$$

In this paper, for a non-bipartite connected graph Ω , we study the smallest normalized signless ∞ -Laplacian eigenvalue

$$\mu_\infty(\Omega) = \min_{\|X\|_\infty=1} \|\Delta_\infty X\|_\infty = \min_{\|X\|_\infty=1} \max_{i \sim j} \left| \frac{x_i}{d_i} + \frac{x_j}{d_j} \right|. \quad (1.1)$$

We provide an equivalent algebraic characterization for this invariant. In Section 2, we recall some notations and definitions. In Section 3, we prove that μ_∞ equals to the reciprocal of the minimal ∞ -norm of generalized inverses of weighted signless incidence matrix W of graph, i.e.,

$$\min_{\|X\|_\infty=1} \|WX\|_\infty = \frac{1}{\min_{WGW=W} \|G\|_{\infty,\infty}}.$$

In Section 4, we apply the result to a class of non-bipartite bicyclic connected graph.

2. PRELIMINARY

In this paper, all vectors and matrices are real. We use \mathbb{R}^n to denote the set of all n dimensional column vectors over \mathbb{R} and $\mathbb{R}^{m \times n}$ to denote the set of all $m \times n$ matrices over \mathbb{R} .

Definition 2.1. [13] The infinity norm of a vector $x = (x_1, x_2, \dots, x_n)^T \in \mathbb{R}^n$ is $\|x\|_\infty = \max\{|x_i| : i = 1, 2, \dots, n\}$. The induced matrix norm for a real $m \times n$ matrix $A = (a_{ij})$ is

$$\|A\|_{\infty,\infty} = \max\{\|AX\|_\infty : \|X\|_\infty = 1\} = \max_{1 \leq i \leq m} \sum_{j=1}^n |a_{ij}|.$$

Definition 2.2. [5] Let Ω be a graph without loops or multiple edges, whose vertex set and edge set are $V = \{v_1, v_2, \dots, v_n\}$ and $E = \{e_1, e_2, \dots, e_m\}$. Let $B = (b_{ij})_{m \times n}$ be the edge-vertex incidence matrix of Ω , where

$$b_{ij} = \begin{cases} 1, & v_j \text{ is incident to } e_i, \\ 0, & \text{otherwise.} \end{cases}$$

Let $D = \text{diag}(d_1, d_2, \dots, d_n)$, where d_i is the degree of vertex v_i , $i = 1, 2, \dots, n$. Define $W = BD^{-1}$ as the weighted signless edge-vertex incidence matrix of Ω .

Remark 2.3. By equation (1.1), it is easy to check that the smallest normalized signless ∞ -Laplacian eigenvalue for graph Ω is

$$\mu_\infty(\Omega) = \min_{\|X\|_\infty=1} \|WX\|_\infty = \min_{\|X\|_\infty=1} \max_{i \sim j} \left| \frac{x_i}{d_i} + \frac{x_j}{d_j} \right|,$$

where $X = (x_i, v_i \in V) \in \mathbb{R}^V$ and $i \sim j$ denotes vertex v_i is adjacent to v_j .

Definition 2.4. [3] Let W be a real $m \times n$ matrix. Then a real $n \times m$ matrix G is called a generalized inverse of W if

$$WGW = W.$$

In particular, if W has full column rank, then a generalized inverse G of W should be left inverse of W , i.e.,

$$GW = I.$$

Note that there may be many generalized inverses for a matrix. The study of generalized inverses with certain minimal norms is important in matrix analysis, see [10, 11].

3. MAIN THEOREM AND ITS PROOF

Theorem 3.1. Suppose that Ω is a non-bipartite connected graph with n vertices and m edges where $m \geq n$. Let W be the weighted signless edge-vertex incidence matrix of Ω . Then

$$\mu_\infty(\Omega) = \min_{\|X\|_\infty=1} \|WX\|_\infty = \frac{1}{\min_{WGW=W} \|G\|_{\infty,\infty}}.$$

Proof. By the assumption on Ω , we know W has full column rank. For any generalized inverse $G \in \mathbb{R}^{n \times m}$ of W satisfying $GW = I$, and any $X \in \mathbb{R}^n \setminus \{\mathbf{0}\}$, we have

$$\|X\|_\infty = \|GWX\|_\infty \leq \|G\|_{\infty,\infty} \cdot \|WX\|_\infty.$$

So

$$\min\left\{\frac{\|WX\|_\infty}{\|X\|_\infty} : X \in \mathbb{R}^n \setminus \{\mathbf{0}\}\right\} \geq \frac{1}{\min_{GW=I} \|G\|_{\infty,\infty}}. \quad (3.1)$$

Now we turn to prove there exists a left inverse G of W such that

$$\min\left\{\frac{\|WX\|_\infty}{\|X\|_\infty} : X \in \mathbb{R}^n \setminus \{\mathbf{0}\}\right\} = \frac{1}{\|G\|_{\infty,\infty}}.$$

Let $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}^m : X \mapsto WX$. Then φ is injective. Therefore the mapping $\tilde{\varphi} : \mathbb{R}^n \rightarrow \text{Im}\varphi : X \mapsto WX$ is a linear isomorphism. Let $\tilde{\psi}$ be the unique inverse mapping of $\tilde{\varphi}$. Note that there exists some $\xi = (\xi_1, \xi_2, \dots, \xi_n)^T \in \mathbb{R}^n \setminus \{\mathbf{0}\}$ such that

$$c = \frac{\|\varphi(\xi)\|_\infty}{\|\xi\|_\infty} = \min\left\{\frac{\|\varphi(X)\|_\infty}{\|X\|_\infty} : X \in \mathbb{R}^n \setminus \{\mathbf{0}\}\right\}.$$

Take $\eta = \varphi(\xi)$, then $c = \frac{\|\eta\|_\infty}{\|\xi\|_\infty}$. Since $\tilde{\psi}(\eta) = \xi$, by definition of c , we can get

$$\max_{Y \in \text{Im}\varphi} \frac{\|\tilde{\psi}(Y)\|_\infty}{\|Y\|_\infty} = \frac{\|\tilde{\psi}(\eta)\|_\infty}{\|\eta\|_\infty} = \frac{\|\xi\|_\infty}{\|\varphi(\xi)\|_\infty} = \frac{1}{c},$$

i.e., $\|\tilde{\psi}\|_{\infty, \infty} = \frac{1}{c}$.

Write

$$\begin{aligned}\tilde{\psi} : \text{Im}\varphi &\rightarrow \mathbb{R}^n \\ \alpha &\mapsto (\tilde{\psi}_1(\alpha), \tilde{\psi}_2(\alpha), \dots, \tilde{\psi}_n(\alpha))^T.\end{aligned}$$

By Hahn-Banach Theorem (see [1], P. 34), there exists a norm-preserving linear extension $\psi : \mathbb{R}^m \rightarrow \mathbb{R}^n$ of $\tilde{\psi}$, where $i = 1, 2, \dots, n$. Take

$$\begin{aligned}\psi : \mathbb{R}^m &\rightarrow \mathbb{R}^n \\ \beta &\mapsto (\psi_1(\beta), \psi_2(\beta), \dots, \psi_n(\beta))^T.\end{aligned}$$

We can get

$$\begin{aligned}\|\psi\|_{\infty, \infty} &= \max_{Z \in \mathbb{R}^m \setminus \{\mathbf{0}\}} \frac{\|\psi(Z)\|_{\infty}}{\|Z\|_{\infty}} \\ &= \max_{i=1,2,\dots,n} \max_{Z \in \mathbb{R}^m \setminus \{\mathbf{0}\}} \frac{|\psi_i(Z)|}{\|Z\|_{\infty}} \\ &= \max_{i=1,2,\dots,n} \max_{Z \in \text{Im}\varphi \setminus \{\mathbf{0}\}} \frac{|\tilde{\psi}_i(Z)|}{\|Z\|_{\infty}} \\ &= \max_{i=1,2,\dots,n} \frac{|\tilde{\psi}_i(\eta)|}{\|\eta\|_{\infty}} \\ &= \|\tilde{\psi}\|_{\infty, \infty}.\end{aligned}$$

So ψ is a norm-preserving linear extension of $\tilde{\psi}$, and thus there exists some generalized inverse matrix G of W such that

$$\|G\|_{\infty, \infty} = \|\psi\|_{\infty, \infty} = \frac{1}{c} = \frac{1}{\min_{\|X\|_{\infty}=1} \|WX\|_{\infty}}. \quad (3.2)$$

Combining equations (3.1) and (3.2), we get

$$\mu_{\infty}(\Omega) = \min_{\|X\|_{\infty}=1} \|WX\|_{\infty} = \frac{1}{\min_{WG W=W} \|G\|_{\infty, \infty}}.$$

□

4. AN EXAMPLE

As an example, we introduce the definition for the class of non-bipartite bicyclic connected graphs $ET(n, a, b)$ in Definition 4.1. We compute the minimal ∞ -norm of generalized inverses of the weighted signless incidence matrices W for $ET(n, a, b)$ in Theorem 4.2. And we construct a vector $Y \in \mathbb{R}^n \setminus \{\mathbf{0}\}$ in Remark 4.3 such that

$$\mu_{\infty} = \frac{\|WY\|_{\infty}}{\|Y\|_{\infty}} = \min_{\|X\|_{\infty}=1} \|WX\|_{\infty}$$

to illustrate the result of Theorem 3.1.

Definition 4.1. Suppose that $n = a + b - 1$, where integers $a, b > 2$ and a is odd. Let $V = \{v_1, v_2, \dots, v_n\}$ and $E = \{e_1, e_2, \dots, e_{n+1}\}$, where

- (1) e_k is an edge between v_k and v_{k+1} , $k = 1, 2, \dots, a - 1$,
- (2) e_a is an edge between v_a and v_1 ,
- (3) e_k is an edge between v_{k-1} and v_k , $k = a + 1, a + 2, \dots, n$,
- (4) e_{n+1} is an edge between v_n and v_a .

We denote the bicyclic graph (V, E) by $ET(n, a, b)$.

For example, $ET(9, 3, 7)$ is as FIGURE 1.




FIGURE 1. $ET(9, 3, 7)$

Theorem 4.2. Let W be the weighted signless incidence matrix of $ET(n, a, b)$ and $G = (g_{ij})$ be a generalized inverse of W . Then we have

$$\min_{WGW=W} \|G\|_{\infty, \infty} = \begin{cases} a, & \text{if } b \text{ is odd and } a > 2b, \\ 2 \min\{a, b\}, & \text{if } b \text{ is odd and } a \leq 2b, \ b \leq 2a, \\ b, & \text{if } b \text{ is odd and } b > 2a, \\ \max\{2a, a + b\}, & \text{if } b \text{ is even.} \end{cases}$$

Proof. The weighted signless incidence matrix W of $ET(n, a, b)$ is

$$\begin{matrix} & 1 & & \dots & & a & & \dots & & n \\ 1 & \begin{pmatrix} \frac{1}{2} & \frac{1}{2} & 0 & \dots & 0 & 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & \frac{1}{2} & \frac{1}{2} & \dots & 0 & 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \dots & 0 & 0 & 0 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & \frac{1}{2} & \frac{1}{4} & 0 & 0 & \dots & 0 & 0 \\ a & \frac{1}{2} & 0 & 0 & \dots & 0 & \frac{1}{4} & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & \frac{1}{4} & \frac{1}{2} & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 & \frac{1}{2} & \frac{1}{2} & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & 0 & 0 & 0 & \dots & \frac{1}{2} & \frac{1}{2} \\ n+1 & 0 & 0 & 0 & \dots & 0 & \frac{1}{4} & 0 & 0 & \dots & 0 & \frac{1}{2} \end{pmatrix} & . \end{matrix}$$

Since W has full column rank, so its generalized inverse G satisfying $GW = I$.

Case 1: If b is odd, then let \hat{G} be

$$\begin{matrix} & 1 & & & \cdots & & a & & \cdots & & n+1 \\ 1 & \left(\begin{array}{cccccccccccccc} 1 & -1 & 1 & -1 & \cdots & 1 & -1 & 1 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 1 & 1 & -1 & 1 & \cdots & -1 & 1 & -1 & 0 & 0 & 0 & \cdots & 0 & 0 \\ -1 & 1 & 1 & -1 & \cdots & 1 & -1 & 1 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 1 & -1 & 1 & 1 & \cdots & -1 & 1 & -1 & 0 & 0 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ -1 & 1 & -1 & 1 & \cdots & 1 & -1 & 1 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 1 & -1 & 1 & -1 & \cdots & 1 & 1 & -1 & 0 & 0 & 0 & \cdots & 0 & 0 \\ a & -2 & 2 & -2 & 2 & \cdots & -2 & 2 & 2 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 1 & -1 & 1 & -1 & \cdots & 1 & -1 & -1 & 2 & 0 & 0 & \cdots & 0 & 0 \\ -1 & 1 & -1 & 1 & \cdots & -1 & 1 & 1 & -2 & 2 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & -1 & 1 & -1 & \cdots & 1 & -1 & -1 & 2 & -2 & 2 & \cdots & 0 & 0 \\ n & -1 & 1 & -1 & 1 & \cdots & -1 & 1 & 1 & -2 & 2 & -2 & \cdots & 2 & 0 \end{array} \right) \end{matrix}.$$

We may check that \hat{G} is a generalized inverse matrix of W . Any generalized inverse G of W can be expressed as

$$G = \hat{G} + \alpha\beta,$$

where $\alpha = (z_1, z_2, \dots, z_n)^T$ is an arbitrary vector in \mathbb{R}^n , and

$$\beta = \begin{pmatrix} 1 & & & \cdots & & a & & \cdots & & n+1 \\ 1 & -1 & 1 & -1 & \cdots & 1 & -1 & -1 & 1 & -1 & 1 & \cdots & -1 & 1 \end{pmatrix}.$$

As we know, if x_i is the median of a sequence of real numbers

$$m_{i1}, m_{i2}, \dots, m_{in},$$

then

$$\sum_{j=1}^n |m_{ij} - x_i| = \min \left\{ \sum_{j=1}^n |m_{ij} - y| : y \in \mathbb{R} \right\}.$$

Since

$$\min_{WGW=W} \|G\|_{\infty, \infty} = \min_{\alpha} \|G\|_{\infty, \infty} = \min_{\alpha} \max_{1 \leq i \leq n} \sum_{j=1}^{n+1} |g_{ij}| = \max_{1 \leq i \leq n} \min_{\alpha} \sum_{j=1}^{n+1} |g_{ij}|,$$

where α is an arbitrary vector in \mathbb{R}^n . Now we compute

$$\min_{\alpha} \sum_{j=1}^{n+1} |g_{ij}|$$

for each row.

(1) For $i = 1, 2, \dots, a-1$, let

$$s = \begin{cases} i, & \text{if } i \text{ is even,} \\ a-i, & \text{if } i \text{ is odd,} \end{cases}$$

and $t = a - s$. Then

$$\sum_{j=1}^{n+1} |g_{ij}| = |-1 - z_i| \cdot s + |1 - z_i| \cdot t + |z_i| \cdot b.$$

- (i) If $s > b + t$, we take $z_i = -1$, then $\min_{\alpha} \sum_{j=1}^{n+1} |g_{ij}| = 2t + b < s + t = a$.
 - (ii) If $t > b + s$, we take $z_i = 1$, then $\min_{\alpha} \sum_{j=1}^{n+1} |g_{ij}| = 2s + b < s + t = a$.
 - (iii) If $s \leq b + t$ and $t \leq b + s$, we take $z_i = 0$, then $\min_{\alpha} \sum_{j=1}^{n+1} |g_{ij}| = s + t = a$.
- (2) For $i = a$, we have

$$\sum_{j=1}^{n+1} |g_{ij}| = |2 - z_i| \cdot a + |z_i| \cdot b.$$

- (i) If $a > b$, we take $z_i = 2$, then $\min_{\alpha} \sum_{j=1}^{n+1} |g_{ij}| = 2b$.
 - (ii) If $a = b$, we take $z_i = 1$, then $\min_{\alpha} \sum_{j=1}^{n+1} |g_{ij}| = a + b$.
 - (iii) If $a < b$, we take $z_i = 0$, then $\min_{\alpha} \sum_{j=1}^{n+1} |g_{ij}| = 2a$.
- (3) For $i = a + 1, a + 3, \dots, a + b - 2$, let $s = i - a$ and $t = a + b - i$. Then

$$\sum_{j=1}^{n+1} |g_{ij}| = |-1 - z_i| \cdot a + |-2 - z_i| \cdot s + |z_i| \cdot t.$$

- (i) If $s > a + t$, we take $z_i = -2$, then $\min_{\alpha} \sum_{j=1}^{n+1} |g_{ij}| = a + 2t < s + t = b$.
 - (ii) If $t > a + s$, we take $z_i = 0$, then $\min_{\alpha} \sum_{j=1}^{n+1} |g_{ij}| = a + 2s < s + t = b$.
 - (iii) If $s \leq a + t$ and $t \leq a + s$, we take $z_i = -1$, then $\min_{\alpha} \sum_{j=1}^{n+1} |g_{ij}| = s + t = b$.
- (4) For $i = a + 2, a + 4, \dots, a + b - 1$, let $s = i - a$ and $t = a + b - i$. Then

$$\sum_{j=1}^{n+1} |g_{ij}| = |1 - z_i| \cdot a + |2 - z_i| \cdot s + |z_i| \cdot t.$$

- (i) If $s > a + t$, we take $z_i = 2$, then $\min_{\alpha} \sum_{j=1}^{n+1} |g_{ij}| = a + 2t < s + t = b$.
- (ii) If $t > a + s$, we take $z_i = 0$, then $\min_{\alpha} \sum_{j=1}^{n+1} |g_{ij}| = a + 2s < s + t = b$.
- (iii) If $s \leq a + t$ and $t \leq a + s$, we take $z_i = 1$, then $\min_{\alpha} \sum_{j=1}^{n+1} |g_{ij}| = s + t = b$.

So when b is odd,

$$\min_{WGW=W} \|G\|_{\infty, \infty} = \begin{cases} a, & \text{if } a > 2b, \\ 2 \min\{a, b\}, & \text{if } a \leq 2b \text{ and } b \leq 2a, \\ b, & \text{if } b > 2a. \end{cases}$$

The proof in Case 1 is complete.

Case 2: If b is even, then let \hat{G} be

$$\begin{matrix} & 1 & & \cdots & & a & & \cdots & & n+1 \\ 1 & \left(\begin{array}{cccccccccccccc} 1 & -1 & 1 & -1 & \cdots & 1 & -1 & 1 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 1 & 1 & -1 & 1 & \cdots & -1 & 1 & -1 & 0 & 0 & 0 & \cdots & 0 & 0 \\ -1 & 1 & 1 & -1 & \cdots & 1 & -1 & 1 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 1 & -1 & 1 & 1 & \cdots & -1 & 1 & -1 & 0 & 0 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ -1 & 1 & -1 & 1 & \cdots & 1 & -1 & 1 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 1 & -1 & 1 & -1 & \cdots & 1 & 1 & -1 & 0 & 0 & 0 & \cdots & 0 & 0 \\ a & -2 & 2 & -2 & 2 & \cdots & -2 & 2 & 2 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 1 & -1 & 1 & -1 & \cdots & 1 & -1 & -1 & 2 & 0 & 0 & \cdots & 0 & 0 \\ -1 & 1 & -1 & 1 & \cdots & -1 & 1 & 1 & -2 & 2 & 0 & \cdots & 0 & 0 \\ 1 & -1 & 1 & -1 & \cdots & 1 & -1 & -1 & 2 & -2 & 2 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ -1 & 1 & -1 & 1 & \cdots & -1 & 1 & 1 & -2 & 2 & -2 & \cdots & 0 & 0 \\ n & 1 & -1 & 1 & -1 & \cdots & 1 & -1 & -1 & 2 & -2 & 2 & \cdots & 2 & 0 \end{array} \right) \end{matrix}.$$

We may check that \hat{G} is a generalized inverse matrix of W . Any generalized inverse of W can be expressed as

$$G = \hat{G} + \alpha\beta,$$

where $\alpha = (z_1, z_2, \dots, z_n)^T$ is an arbitrary vector in \mathbb{R}^n , and

$$\beta = \begin{pmatrix} 1 & \cdots & a & \cdots & n+1 \\ 0 & 0 & \cdots & 0 & 0 & -1 & 1 & -1 & 1 & \cdots & -1 & 1 \end{pmatrix}$$

Similarly as in Case 1, now we can get

$$\min_{WGW=W} \|G\|_{\infty, \infty} = \min_{\alpha} \|G\|_{\infty, \infty} = \max\{2a, a+b\}.$$

The proof in Case 2 is complete. \square

Now we construct a vector $Y \in \mathbb{R}^n \setminus \{\mathbf{0}\}$ such that the smallest normalized signless ∞ -Laplacian eigenvalue

$$\mu_{\infty} = \frac{\|WY\|_{\infty}}{\|Y\|_{\infty}}$$

for the non-bipartite bicyclic connected graphs $ET(n, a, b)$.

Remark 4.3. Let W be the weighted signless incidence matrix of $ET(n, a, b)$. We construct the optimal vector $Y = (y_1, y_2, \dots, y_n)^T \in \mathbb{R}^n \setminus \{\mathbf{0}\}$ as follows,

such that

$$\mu_\infty = \frac{\|WY\|_\infty}{\|Y\|_\infty} = \frac{\max_{i \sim j} |\frac{y_i}{d_i} + \frac{y_j}{d_j}|}{\max_i |y_i|} = \min_{\|X\|_\infty=1} \|WX\|_\infty = \frac{1}{\min_{GW=I} \|G\|_{\infty,\infty}}.$$

If a and b are odd, without loss of generality, we only need consider two cases when $a \geq b$ as follows. If a, b are both odd, and $b \geq a$, we can easily get the vector Y by exchanging a and b in Case 1 and Case 2.

Case 1: If b is odd, and $a > 2b$. Suppose that $Y = (y_i) \in \mathbb{R}^n \setminus \{\mathbf{0}\}$ satisfying

$$y_i = \begin{cases} (-1)^i \cdot (a - 2 \cdot \lfloor \frac{a-1}{2} - i \rfloor), & \text{if } i = 1, 2, \dots, a-1, \\ 2, & \text{if } i = a, \\ (-1)^i \cdot (b - 2 \cdot \lfloor a + \frac{b+1}{2} - i \rfloor), & \text{if } i = a+1, a+2, \dots, a+b-1. \end{cases}$$

Then

$$\begin{aligned} \max_i |y_i| &= |y_{\frac{a-1}{2}}| = a, \\ \max_{i \sim j} |\frac{y_i}{d_i} + \frac{y_j}{d_j}| &= 1. \end{aligned}$$

So

$$\mu_\infty = \frac{\|WY\|_\infty}{\|Y\|_\infty} = \frac{\max_{i \sim j} |\frac{y_i}{d_i} + \frac{y_j}{d_j}|}{\max_i |y_i|} = \frac{1}{a}.$$

Case 2: If b is odd, and $b \leq a \leq 2b$. Suppose that $Y = (y_i) \in \mathbb{R}^n \setminus \{\mathbf{0}\}$ satisfying

$$y_i = \begin{cases} (-1)^i \cdot (b - 2i), & \text{if } i = 1, 2, \dots, \frac{b-1}{2}, \\ (-1)^{\frac{b-1}{2}}, & \text{if } i = \frac{b+1}{2}, \frac{b+1}{2} + 1, \dots, a - \frac{b+1}{2}, \\ (-1)^{a-i} \cdot (b - 2(a-i)), & \text{if } i = a - \frac{b-1}{2}, a - \frac{b-1}{2} + 1, \dots, a-1, \\ 2b, & \text{if } i = a, \\ (-1)^{i-a} \cdot [b - 2(i-a)], & \text{if } i = a+1, a+2, \dots, a + \frac{b-1}{2}, \\ (-1)^{a+b-i} \cdot [b - 2(a+b-i)], & \text{if } i = a + \frac{b+1}{2}, a + \frac{b+1}{2} + 1, \dots, a+b-1. \end{cases}$$

Then

$$\begin{aligned} \max_i |y_i| &= |y_a| = 2b, \\ \max_{i \sim j} |\frac{y_i}{d_i} + \frac{y_j}{d_j}| &= 1. \end{aligned}$$

So

$$\mu_\infty = \frac{\|WY\|_\infty}{\|Y\|_\infty} = \frac{\max_{i \sim j} |\frac{y_i}{d_i} + \frac{y_j}{d_j}|}{\max_i |y_i|} = \frac{1}{2b}.$$

Case 3: If a is odd and b is even, suppose that the vector $Y \in \mathbb{R}^n \setminus \{\mathbf{0}\}$ satisfying

$$y_i = \begin{cases} (-1)^i \cdot (a - 2i), & \text{if } i = 1, 2, \dots, \frac{a-1}{2}, \\ (-1)^{a-i} \cdot [a - 2 \cdot (a-i)], & \text{if } i = \frac{a+1}{2}, \frac{a+1}{2} + 1, \dots, a-1, \\ 2a, & \text{if } i = a, \\ (-1)^{i-a} \cdot [a + 2 \cdot (i-a)], & \text{if } i = a+1, a+2, \dots, a + \frac{b}{2}, \\ (-1)^{a+b-i} \cdot [a + 2 \cdot (a+b-i)], & \text{if } i = a + \frac{b}{2} + 1, a + \frac{b}{2} + 2, \dots, a+b-1. \end{cases}$$

When $a > b$, we have

$$\max_i |y_i| = |y_a| = 2a,$$

$$\max_{i \sim j} \left| \frac{y_i}{d_i} + \frac{y_j}{d_j} \right| = 1.$$

So

$$\mu_\infty = \frac{\|WY\|_\infty}{\|Y\|_\infty} = \frac{\max_{i \sim j} \left| \frac{y_i}{d_i} + \frac{y_j}{d_j} \right|}{\max_i |y_i|} = \frac{1}{2a}.$$

When $a < b$, we have

$$\max_i |y_i| = |y_{a+\frac{b}{2}}| = a + b,$$

$$\max_{i \sim j} \left| \frac{y_i}{d_i} + \frac{y_j}{d_j} \right| = 1.$$

So

$$\mu_\infty = \frac{\|WY\|_\infty}{\|Y\|_\infty} = \frac{\max_{i \sim j} \left| \frac{y_i}{d_i} + \frac{y_j}{d_j} \right|}{\max_i |y_i|} = \frac{1}{a + b}.$$

The computational results from these construction and Theorem 4.2 coincide. These verify the identity in Theorem 3.1 for $ET(n, a, b)$.

Conflict of interest: The authors declare there is no conflict of interest.

Funding: The research is supported by the Fundamental Research Funds for the Central Universities of China (Grant no. GK202304003).

REFERENCES

- [1] S. BANACH, *Theory of linear operations*, Translated from the French by F. Jellett. North-Holland Publishing Co., Amsterdam, 1987.
- [2] F. BELARDO, M. BRUNETTI, A. CIAMPELLA, *Signed bicyclic graphs minimizing the least Laplacian eigenvalue*, Linear Algebra Appl. 557 (2018) 201–233.
- [3] A. BEN-ISRAEL, T.N.E. GREVILLE, *Generalized Inverses: Theory and Applications*, Corrected reprint of the 1974 original. Robert E. Krieger Publishing Co., Inc., Huntington, New York, 1980.
- [4] E.M. BORBA, U. SCHWERDTFEGER, *Eigenvalue bounds for the signless p -Laplacian*, Electron. J. Combin. 25(2)(2018)1–18.
- [5] R.A. BRUALDI, H.J. RYSER, *Combinatorial matrix theory*, Cambridge University Press, Cambridge, 1991.
- [6] K.C. CHANG, *Spectrum of the 1-Laplacian and Cheeger's constant on graphs*, J. Graph Theory 81 (2016) 167–207.
- [7] K.C. CHANG, S.H. SHAO, D. ZHANG, *Spectrum of the Signless 1-Laplacian and the Dual Cheeger Constant on Graphs*, (2016) <https://arxiv.org/abs/1607.00489>.
- [8] J. CHEEGER, *A lower bound for the smallest eigenvalue of the Laplacian*, Problems in analysis, (1970), 195–199.
- [9] M. DESAI, V. RAO, *A characterization of the smallest eigenvalue of a graph*, J. Graph Theory 18 (1994), 181–194.
- [10] I. DOKMANIĆ, R. GRIBONVAL, *Beyond Moore-Penrose Part I: Generalized Inverses that Minimize Matrix Norms*, (2017) <https://arxiv.org/abs/1706.08349>.
- [11] I. DOKMANIĆ, R. GRIBONVAL, *Concentration of the Frobenius norm of generalized matrix inverses*, SIAM J. Matrix Anal. Appl. 40 (2019) 92–121.

- [12] S. FALLAT, Y.-Z. FAN, *Bipartiteness and the least eigenvalue of signless Laplacian of graphs*, Linear Algebra Appl. 436 (2012) 3254–3267.
- [13] G.H. GOLUB, C.F. VAN LOAN, *Matrix computations*, fourth ed., Johns Hopkins University Press, Baltimore, MD, 2013.
- [14] M. KHAN, R. FAROOQ, A.A. SIDDIQUI, *On the extremal energy of bicyclic digraphs*, J. Math. Inequal. 9 (2015), no. 3, 799–810.
- [15] A.D. MADEN, A.S. ÇEVİK, N. HABIBI, *New bounds for the spread of the signless Laplacian spectrum*, Math. Inequal. Appl., 17 (2014), no. 1, 283–294.
- [16] C. POIGNARD, T. PEREIRA, J.P. PADE, *Spectra of Laplacian matrices of weighted graphs: structural genericity properties*, SIAM J. Appl. Math., 78 (2018), no. 1, 372–394.
- [17] J.-M. ZHANG, T.-Z. HUANG, J.-M. GUO, *The smallest signless Laplacian spectral radius of graphs with a given clique number*, Linear Algebra Appl. 439 (2013) 2562–2576.
- [18] B. ZHOU, *Signless Laplacian spectral radius and Hamiltonicity*, Linear Algebra Appl. 432(2010) 566–570.

YI DAI, SCHOOL OF MATHEMATICS AND STATISTICS, SHAANXI NORMAL UNIVERSITY,
XI'AN 710062, CHINA.

Email address: daiyi@snnu.edu.cn