RELATIVE LANGLANDS DUALITY FOR $\mathfrak{osp}(2n+1|2n)$

ALEXANDER BRAVERMAN, MICHAEL FINKELBERG, DAVID KAZHDAN, AND ROMAN TRAVKIN

ABSTRACT. We establish an S-duality converse to the one of [BFT]; this is also a case of a twisted version of the relative Langlands duality of [BZSV]. Namely, we prove that the S-dual of $SO(2n + 1) \times Sp(2n) \circlearrowright \mathbb{C}^{2n+1}_+ \otimes \mathbb{C}^{2n}_-$ is the symplectic mirabolic space $Sp(2n) \times Sp(2n) \circlearrowright T^*Sp(2n) \times \mathbb{C}^{2n}_-$ (note that due to the anomaly, the dual of the second factor Sp(2n) is metaplectic dual, i.e. Sp(2n)). We also formulate the corresponding global conjecture, which describes explicitly the categorical theta-correspondence on the Langlands dual side.

CONTENTS

1. Introduction	2
1.1. Ring objects in the Satake category and relative Langlands duality	2
1.2. The non-cotangent case	2
1.3. The subject of [BFT]	2
1.4. The subject of this paper	3
1.5. The global conjecture	3
1.6. Acknowledgments	4
2. Setup	5
2.1. Weyl algebra	5
2.2. Main theorem	6
3. The proof	6
3.1. The two Hecke actions can be identified	6
3.2. The Hecke action generates \mathcal{W} -mod ^{SO(V')_0 \times Sp(V)_0}	8
3.3. A deequivariantized Ext-algebra	8
3.4. A pseudo-slice	10
3.5. Equivariant cohomology	11
References	14

1. INTRODUCTION

1.1. Ring objects in the Satake category and relative Langlands duality. Let G be a connected reductive group over \mathbb{C} . Let $\mathcal{K} = \mathbb{C}((t)) \supset \mathbb{O} = \mathbb{C}[t]$. The affine Grassmannian ind-scheme $\operatorname{Gr}_G = G_{\mathcal{K}}/G_0$ is the moduli space of G-bundles on the formal disc equipped with a trivialization on the punctured formal disc. One can consider the *derived Satake category* $D_{G_0}(\operatorname{Gr}_G)$.¹ This is a factorization monoidal category which is monoidally equivalent to $D^{G^{\vee}}(\operatorname{Sym}^{\bullet}(\mathfrak{g}^{\vee}[-2]))$: the derived category of dg-modules over $\operatorname{Sym}^{\bullet}(\mathfrak{g}^{\vee}[-2])$ endowed with a compatible action of G^{\vee} (the monoidal structure on this category is just given by tensor product over $\operatorname{Sym}^{\bullet}(\mathfrak{g}^{\vee}[-2])$); we shall denote the corresponding functor from $D_{G_0}(\operatorname{Gr}_G)$ to $D^{G^{\vee}}(\operatorname{Sym}^{\bullet}(\mathfrak{g}^{\vee}[-2])$ by Φ_G .

In [BFN] we have attached to any **N** as above a certain ring object $\mathcal{A}_{G,\mathbf{N}}$ in $D_{G_{\mathcal{O}}}(\mathrm{Gr}_{G})$. This construction was generalized in [BZSV] to the case when **N** is an arbitrary smooth affine variety with a *G*-action. Moreover, it is argued in [BFN], [BDFRT] and [BZSV] that the object $\mathcal{A}_{G,\mathbf{N}}$ should only depend on $\mathbf{M} = T^*\mathbf{N}$. One of the main conjectures of [BZSV] says that if **N** is a spherical *G*-variety then $H^*(\Phi_G(\mathcal{A}_{G,\mathbf{N}})$ should be the algebra of functions on certain (*relative Langlands dual* or *S*-dual) hyper-spherical Poisson *G*-variety \mathbf{M}^{\vee} , and this construction is expected to be involutive in some reasonable generality, cf. also [N2].

1.2. The non-cotangent case. The construction of $\mathcal{A}_{G,\mathbf{N}}$ should in principle make sense for any smooth affine symplectic *G*-variety \mathbf{M} . However, when \mathbf{M} is not of cotangent type some extra care is needed. This is discussed in detail in [BDFRT] when \mathbf{M} is a symplectic linear representation of *G*. In this case in *loc. cit.* the corresponding object $\mathcal{A}_{G,\mathbf{M}}$ was constructed, but in general it is not an object of $D_{G_0}(\mathbf{Gr}_G)$ but rather of some twisted version of it. The twisting is by a square root of some line bundle on \mathbf{Gr}_G which depends on \mathbf{M} ; in the case when the twisting is indeed non-trivial we say that an *anomaly* is present. The derived Satake equivalence can be extended to such twisted categories (but one has to change the notion of the Langlands dual group G^{\vee}) so even in the anomalous case all of the above constructions go through.

1.3. The subject of [BFT]. This note is a sequel to [BFT]. There we considered the symplectic group $\operatorname{Sp}(2n) \circlearrowright \mathbb{C}^{2n}_{-}$ and its Langlands dual group $\operatorname{SO}(2n+1) \circlearrowright \mathbb{C}^{2n+1}_{+}$. We also considered a symplectic vector space $\mathbf{M} = \mathbb{C}^{2n+1}_{+} \otimes \mathbb{C}^{2n}_{-}$ and the corresponding Weyl algebra \mathcal{W} of $\mathbf{M}_{\mathcal{K}}$. Here $\mathcal{K} = \mathbb{C}((t)) \supset \mathbb{C}[t] = 0$. We studied the category \mathcal{DW} -mod^{Sp(2n)_0,lc} of locally compact $\operatorname{Sp}(2n)_0$ -equivariant objects in

¹In fact we are going to work with a renormalized version of it, which by definition is equal to the ind-completition of the corresponding subcategory of bounded complexes with constructible cohomology.

the tensor product cagegory $D-\text{mod}_{1/2}(\text{Gr}_{\text{Sp}(2n)}) \otimes \mathcal{W}$ -mod (*D*-modules on the affine Grassmannian of Sp(2n) twisted by the square root of the determinant line bundle).

We established the following algebraic description of \mathcal{DW} -mod^{Sp(2n)₀,lc} conjectured by D. Gaiotto. We consider the (infinite-dimensional) graded algebra Sym[•]($\Pi \mathbf{M}[-1]$) (Π assigns to \mathbf{M} the odd parity) as a dg-superalgebra with trivial differential. We constructed an equivalence of categories

$$\mathcal{DW}\text{-}\mathrm{mod}^{\mathrm{Sp}(2n)_{\mathcal{O}},\mathrm{lc}} \xrightarrow{\sim} D_{\mathrm{perf}}^{\mathrm{SO}(2n+1)\times\mathrm{Sp}(2n)}(\mathrm{Sym}^{\bullet}(\Pi\mathbf{M}[-1])).$$

This is also a particular case of the general [BZSV, Conjecture 7.5.1]. In other words, it means that the S-dual of the symplectic variety $T^*\operatorname{Sp}(2n) \times \mathbb{C}_{-}^{2n}$ equipped with the hamiltonian action of $\operatorname{Sp}(2n) \times \operatorname{Sp}(2n)$, is $\mathbf{M} \oslash \operatorname{SO}(2n+1) \times \operatorname{Sp}(2n)$. (Note that due to the twisting by the square root of the determinant line bundle, the second factor $\operatorname{Sp}(2n)$ corresponds under the S-duality to its metaplectic Langlands dual $\operatorname{Sp}(2n)$.)

1.4. The subject of this paper. In the present note we confirm the converse claim: the S-dual of $\mathbf{M} \circlearrowleft \mathrm{SO}(2n+1) \times \mathrm{Sp}(2n)$ is $T^*\mathrm{Sp}(2n) \times \mathbb{C}^{2n} \circlearrowright \mathrm{Sp}(2n) \times \mathrm{Sp}(2n)$. That is, we construct an equivalence of categories

$$D_{\mathrm{perf}}^{\mathrm{Sp}(2n)\times\mathrm{Sp}(2n)}\left(\mathbb{C}[\mathrm{Sp}(2n)]\otimes\mathrm{Sym}^{\bullet}(\mathfrak{sp}(2n)[-2])\otimes\mathrm{Sym}^{\bullet}(\Pi(\mathbb{C}^{2n}_{-})[-1])\right)$$
$$\xrightarrow{\sim} \mathcal{W}\operatorname{-mod}^{\mathrm{SO}(2n+1)_{0}\times\mathrm{Sp}(2n)_{0},\mathrm{lc}}$$

1.5. The global conjecture. Another very important point of [BZSV] is that the local relative Langlands duality is expected to give rise to certain global geometric ("period") representation of automorphic *L*-functions. The precise formulation of [BZSV] is in the case when the spectral side is of cotangent type and when there is no anomaly. Hopefully, both assumptions can be overcome. We are not going to discuss how to do this in general, however, we are going to present a conjecture in the above case, which we would like to think of as a natural extension of the setting of [BZSV], Section 12] to our setting.

Let C be a smooth projective irreducible curve over \mathbb{C} .² For an algebraic group G we denote by Bun_G the moduli stack of G-bundles on C; in the case when $G = \operatorname{Sp}(2n)$ we shall also consider the twisted version of $\operatorname{Bun}_{\operatorname{Sp}(2n)}$ which we shall denote by $\operatorname{Bun}_{\operatorname{Sp}(2n)}^{\omega}$ — it classifies vector bundles M on C of rank 2n equipped with a non-degenerate skew-symmetric form $\Lambda^2(M) \to \omega_C$. We have a natural morphism ι : $\operatorname{Bun}_{\operatorname{Sp}(2n)}^{\omega} \times \operatorname{Bun}_{\operatorname{SO}(2n+1)} \to \operatorname{Bun}_{\operatorname{Sp}(2n(2n+1))}^{\omega}$. Let Θ denote the theta-sheaf of [L1] on $\operatorname{Bun}_{\operatorname{Sp}(2n(2n+1))}^{\omega}$. This is actually a sheaf twisted by the square root

²A similar discussion should make sense in the ℓ -adic setting when C is a curve over a finite field; however in that case we do not know how to formulate a precise conjecture.

of the natural determinant bundle on $\operatorname{Bun}_{\operatorname{Sp}(2n(2n+1))}^{\omega}$. Consider now $\iota^!\Theta$. This is a sheaf on $\operatorname{Bun}_{\operatorname{Sp}(2n)} \times \operatorname{Bun}_{\operatorname{SO}(2n+1)}$ twisted by the square root of the determinant line bundle along the first factor.

It is expected (cf. [GS]) that the twisted global geometric Langlands duality should assign to any sheaf \mathcal{F} on $\operatorname{Bun}_{\operatorname{Sp}(2n)}^{\omega} \times \operatorname{Bun}_{\operatorname{SO}(2n+1)}$ twisted by the square root of the determinant bundle along the first factor an ind-coherent sheaf $\mathbb{L}(\mathcal{F})$ (with nilpotent singular support) on $\operatorname{LocSys}_{\operatorname{Sp}(2n)}(C) \times \operatorname{LocSys}_{\operatorname{Sp}(2n)}(C)$ where for an algebraic group H over \mathbb{C} we denote by $\operatorname{LocSys}_{H}(C)$ the moduli stack of de Rham H-local systems on $C.^{3}$

To formulate our conjecture we need to introduce the following notation. Let \mathcal{E} be a symplectic local system on C. For simplicity let us assume that $H^0_{dR}(C, \mathcal{E}) = H^2_{dR}(C, \mathcal{E}) = 0$ where the subscript dR stands for de Rham cohomology. The space $H^1_{dR}(C, \mathcal{E})$ has a canonical symmetric non-degenerate bilinear form; it also has a canonical maximal isotropic subspace $L_{\mathcal{E}}$ which is equal to the image of $H^0(C, \mathcal{E} \otimes \Omega_C)$ in $H^1_{dR}(C, \mathcal{E})$. We set $S_{\mathcal{E}}$ to be the corresponding spinor representation of the Clifford algebra $\text{Cliff}(H^1_{dR}(C, \mathcal{E}))$ of $H^1_{dR}(C, \mathcal{E})$ (by definition, it is induced from the trivial representation of $\Lambda(L_{\mathcal{E}})$) which is naturally a subalgebra of $\text{Cliff}(H^1_{dR}(C, \mathcal{E}))$).

The following conjecture appears in a slightly weaker form in [L2, Conjecture 1.2.4] for n = 1:

- **Conjecture 1.5.1.** (1) There exists a sheaf Θ^{\vee} on $\operatorname{LocSys}_{\operatorname{Sp}(2n)}(C)$ such that $\mathbb{L}(i^{!}\Theta)$ is equal to $\Delta_{*}\Theta^{\vee}$ where $\Delta : \operatorname{LocSys}_{\operatorname{Sp}(2n)}(C) \to \operatorname{LocSys}_{\operatorname{Sp}(2n)}(C) \times \operatorname{LocSys}_{\operatorname{Sp}(2n)}(C)$ is the diagonal embedding.
 - (2) The fiber of Θ^{\vee} at \mathcal{E} as above is equal to $S_{\mathcal{E}}$.

Corollary 1.5.2. Let Ψ denote the natural functor from $D(\operatorname{Bun}_{\operatorname{SO}(2n+1)})$ to $D_{-1/2}(\operatorname{Bun}_{\operatorname{Sp}(2n)}^{\Omega})$ defined by the kernel $i^!\Theta$ (here $D_{-1/2}(\operatorname{Bun}_{\operatorname{Sp}(2n)}^{\Omega})$) stands for the corresponding twisted category of D-modules). Let Ψ^* denote the functor in the opposite direction given by $\mathbb{D}(i^!\Theta)$ (Verdier duality). Let also \mathcal{E} be as above and let $A_{\mathcal{E}}$ be a Hecke eigen-D-module on $\operatorname{Bun}_{\operatorname{SO}(2n+1)}$) with eigenvalue \mathcal{E} . Then $\Psi^* \circ \Psi(A_{\mathcal{E}}) \simeq A_{\mathcal{E}} \otimes \operatorname{Cliff}(H^1_{\mathrm{dR}}(C, \mathcal{E})).$

Note that $\text{Cliff}(H^1_{dR}(C, \mathcal{E}))$ can be thought of as a categorification of the value of the *L*-function of \mathcal{E} at 1/2, so the above corollary is a version of the correspondence between period sheaves and *L*-sheaves which is the main subject of [BZSV].

1.6. Acknowledgments. We are grateful to D. Gaiotto, N. Gurevich, S. Lysenko and H. Nakajima for the inspiring discussions. Our note follows in the footsteps of an unpublished work by T.-H. Chen and J. Wang, cf. [CW].

³In the untwisted case (in the de Rham setting) the global geometric Langlands conjecture has been recently proved, but the twisted case so far remains completely open.

The research of A.B. was partially supported by NSERC; in addition this was was started when A.B. was visiting the University of Geneva, and he thanks that institution for its hospitality. The research of M.F. was supported by the Israel Science Foundation (grant No. 994/24). The research of D.K. was partially supported by ERC grant 101142781.

2. Setup

2.1. Weyl algebra. Let $V = \mathbb{C}^{2n}$ be a symplectic vector space; the symplectic pairing is denoted by \langle, \rangle . Let $V' = \mathbb{C}^{2n+1}$ be a vector space equipped with a nondegenerate symmetric pairing (,). We set $\mathbf{M} := V' \otimes V$ and equip it with the tensor product symplectic form, also do be denoted by \langle, \rangle .

The symplectic form on \mathbf{M} extends to the same named \mathbb{C} -valued symplectic form on $\mathbf{M}_{\mathcal{K}}$: $\langle f, g \rangle = \operatorname{Res} \langle f, g \rangle_{\mathcal{K}} dt$. We denote by \mathcal{W} the completion of the Weyl algebra of $(\mathbf{M}_{\mathcal{K}}, \langle , \rangle)$ with respect to the left ideals generated by the compact subspaces of $\mathbf{M}_{\mathcal{K}}$. It has an irreducible representation $\mathbb{C}[\mathbf{M}_0]$.

We consider the dg-category \mathcal{W} -mod of discrete \mathcal{W} -modules. We recall this is a renormalization of the naive derived category \mathcal{W} -mod^{naive} of discrete \mathcal{W} modules, or more carefully its canonical dg-enhancement, defined as follows.

For each compact open subspace $\mathbf{K} \subset \mathbf{M}_{\mathcal{K}}$, consider the module $U_{\mathbf{K}}$ obtained as the quotient of \mathcal{W} by the left ideal generated by \mathbf{K} . Let us denote by \mathcal{E} the pretriangulated envelope of all such modules $U_{\mathbf{K}}$ within \mathcal{W} -mod^{naive}. By definition, \mathcal{W} -mod is the ind-completion of \mathcal{E} . It carries a unique *t*-structure for which the natural map

$$\mathcal{W}\text{-}\mathrm{mod} \to \mathcal{W}\text{-}\mathrm{mod}^{\mathrm{naive}}$$

is t-exact.

More concretely, we may identify \mathcal{W} with the ring of differential operators on a Lagrangian discrete lattice $\mathbf{L} \subset \mathbf{M}_{\mathcal{K}}$, e.g. $\mathbf{L} = t^{-1}\mathbf{M}_{\mathbb{C}[t^{-1}]}$. Then \mathcal{W} -mod is the inverse limit of D-mod(V) over finite dimensional subspaces $U \subset \mathbf{L}$ with respect to the functors $i'_{U \to U'}$. Equivalently, \mathcal{W} -mod is the colimit, in the sense of cocomplete dg-categories, of D-mod(U) with respect to the functors $i_{U \to U',*}$. The following lemma is a consequence of [R, §10], see [BDFRT, Lemma 2.4.1].

Lemma 2.1.1. There is a categorical action

 $D\operatorname{-mod}_{-1/2}(\operatorname{Sp}(\mathbf{M})_{\mathcal{K}}) \circlearrowright \mathcal{W}\operatorname{-mod}.$

In particular, upon taking spherical vectors, there is an action

 $\mathrm{D}\operatorname{-mod}_{-1/2}(\mathrm{Gr}_{\mathrm{Sp}(\mathbf{M})})^{\mathrm{Sp}(\mathbf{M})_{\mathbb{O}}} \circlearrowright (\mathcal{W}\operatorname{-mod})^{\mathrm{Sp}(\mathbf{M})_{\mathbb{O}}}.$

We have an embedding $\mathrm{SO}(V')_{\mathbb{O}} \times \mathrm{Sp}(V)_{\mathbb{O}} \hookrightarrow \mathrm{Sp}(\mathbf{M})_{\mathbb{O}}$, and we denote by \mathcal{W} -mod^{$\mathrm{SO}(V')_{\mathbb{O}} \times \mathrm{Sp}(V)_{\mathbb{O}}$} the category of $\mathrm{SO}(V')_{\mathbb{O}} \times \mathrm{Sp}(V)_{\mathbb{O}}$ -invariants in \mathcal{W} -mod. Furthermore, we denote by \mathcal{W} -mod^{$\mathrm{SO}(V')_{\mathbb{O}} \times \mathrm{Sp}(V)_{\mathbb{O}}$, the category of locally compact objects of \mathcal{W} -mod^{$\mathrm{SO}(V')_{\mathbb{O}} \times \mathrm{Sp}(V)_{\mathbb{O}}$}. (Recall that an equivariant object is called} locally compact if it becomes a compact object of W-mod after forgetting the equivariant structure.)

2.2. Main theorem. We consider a dg-superalgebra with trivial differential $\mathfrak{A}^{\bullet} := \mathbb{C}[\operatorname{Sp}(V)] \otimes \operatorname{Sym}^{\bullet}(\mathfrak{sp}(V)[-2]) \otimes \operatorname{Sym}^{\bullet}(\Pi(V)[-1])$ (here $\Pi(V)$ stands for the vector space V made odd, so that $\operatorname{Sym}^{\bullet}(\Pi(V)[-1])$ is infinite-dimensional). It is equipped with the following action of $\operatorname{Sp}(V) \times \operatorname{Sp}(V) : (g_1, g_2)(g, x, v) = (g_1gg_2^{-1}, \operatorname{Ad}_{g_1} x, g_1v)$. We also consider a dg-algebra with trivial differential $\mathfrak{G}^{\bullet} := \operatorname{Sym}^{\bullet}(\mathfrak{sp}(V)[-2]) \otimes \operatorname{Sym}^{\bullet}(\Pi(V)[-1])$ equipped with the following action of $\operatorname{Sp}(V) : g(x, v) = (\operatorname{Ad}_g x, gv)$. We have an obvious equivalence of categories $D_{\operatorname{perf}}^{\operatorname{Sp}(V) \times \operatorname{Sp}(V)}(\mathfrak{A}^{\bullet}) \cong D_{\operatorname{perf}}^{\operatorname{Sp}(V)}(\mathfrak{G}^{\bullet})$ (equivariant perfect dg-modules).

Our goal is the following

Theorem 2.2.1. There is an equivalence of triangulated categories $\Phi: D_{\text{perf}}^{\text{Sp}(V) \times \text{Sp}(V)}(\mathfrak{A}^{\bullet}) \xrightarrow{\sim} W\text{-mod}^{\text{SO}(V')_{0} \times \text{Sp}(V)_{0}, \text{lc}}$ commuting with the convolution action of the monoidal spherical Hecke category⁴

$$D_{\mathrm{perf}}^{\mathrm{Sp}(V)}(\mathrm{Sym}^{\bullet}(\mathfrak{sp}(V)[-2])) \otimes D_{\mathrm{perf}}^{\mathrm{Sp}(V)}(\mathrm{Sym}^{\bullet}(\mathfrak{sp}(V)[-2])) \\ \cong \mathrm{D}\operatorname{-mod}(\mathrm{Gr}_{\mathrm{SO}(V')})^{\mathrm{SO}(V')_{0}} \otimes \mathrm{D}\operatorname{-mod}_{-1/2}(\mathrm{Gr}_{\mathrm{Sp}(V)})^{\mathrm{Sp}(V)_{0}}$$

The proof occupies the rest of the paper.

3. The proof

3.1. The two Hecke actions can be identified. We identify \mathcal{W} with differential operators on $(V' \otimes V)_{\mathcal{K}}/(V' \otimes V)_{0}$, and consider a *D*-module E_{0} : the delta-function of the origin. Then $E_{0} \in \mathcal{W}$ -mod^{SO(V')_{0} \times Sp(V)_{0} is the unit object. We denote by ϕ_{Sp} (resp. ϕ_{SO}) the geometric Satake equivalence $\operatorname{Rep}(Sp(V)) \xrightarrow{\sim} D\operatorname{-mod}_{-1/2}(\operatorname{Gr}_{Sp(V)})^{Sp(V)_{0}, \mathfrak{V}}$ (resp. $\operatorname{Rep}(Sp(V)) \xrightarrow{\sim} D\operatorname{-mod}(\operatorname{Gr}_{SO(V')_{0}, \mathfrak{V}})$ (hearts of the natural *t*-structures).}

Lemma 3.1.1. Under the convolution actions of D-mod $(\operatorname{Gr}_{\operatorname{SO}(V')})^{\operatorname{SO}(V')_{0}, \heartsuit}$ and D-mod $_{-1/2}(\operatorname{Gr}_{\operatorname{Sp}(V)})^{\operatorname{Sp}(V)_{0}, \heartsuit}$ on \mathcal{W} -mod $^{\operatorname{SO}(V')_{0} \times \operatorname{Sp}(V)_{0}}$, for any representation U of $\operatorname{Sp}(V)$, the convolution actions $\phi_{\operatorname{SO}}(U) * E_{0}$ and $\phi_{\operatorname{Sp}}(U) * E_{0}$, coincide (compatibly with the tensor structures).

Proof. It suffices to

- a) identify the two convolution actions for the generator V of $\operatorname{Rep}(\operatorname{Sp}(V))$;
- b) under the above identification to identify the direct summands

⁴For the equivalence of first factors, see [BF], and for the equivalence of second factors, see [DLYZ].

RELATIVE LANGLANDS DUALITY FOR $\mathfrak{osp}(2n+1|2n)$

$$E_0 = \phi_{\mathrm{Sp}}(\mathbb{C}) * E_0 \subset \phi_{\mathrm{Sp}}(V) * \phi_{\mathrm{Sp}}(V) * E_0 \cong \phi_{\mathrm{Sp}}(V) * \phi_{\mathrm{SO}}(V) * E_0$$
$$\cong \phi_{\mathrm{SO}}(V) * \phi_{\mathrm{Sp}}(V) * E_0 \cong \phi_{\mathrm{SO}}(V) * \phi_{\mathrm{SO}}(V) * E_0 \supset \phi_{\mathrm{SO}}(\mathbb{C}) * E_0 = E_0,$$

corresponding to the embedding of the direct summand $\mathbb{C} \subset V \otimes V$ in $\operatorname{Rep}(\operatorname{Sp}(V))$.

a) We view \mathcal{W} -mod as D-modules on $(V' \otimes V)_{\mathfrak{K}}/(V' \otimes V)_{0}$, then both $\phi_{\mathrm{SO}}(V) * E_{0}$ and $\phi_{\mathrm{Sp}}(V) * E_{0}$ are supported on $t^{-1}(V' \otimes V)_{0}/(V' \otimes V)_{0}$. Here is a description of these D-modules. Recall the D-module $\exp(f) \in D$ -mod $(\mathrm{Gr}_{\mathrm{Sp}(\mathbf{M})}^{\omega} \times ((t^{-1}\mathbf{M}_{0})/\mathbf{M}_{0}))^{\mathrm{Sp}(\mathbf{M})_{0}}$ from [BFT, §2.13]. Here $\mathrm{Gr}_{\mathrm{Sp}(\mathbf{M})}^{\omega}$ is the minimal orbit in $\mathrm{Gr}_{\mathrm{Sp}(\mathbf{M})}$, but \mathbf{M} was denoted by V in *loc.cit*. The embedding $\mathrm{SO}(V') \times \mathrm{Sp}(V) \hookrightarrow \mathrm{Sp}(\mathbf{M})$ gives rise to embeddings $Q \hookrightarrow \mathrm{Gr}_{\mathrm{Sp}(\mathbf{M})}^{\omega}$ and $\mathrm{Gr}_{\mathrm{Sp}(V)}^{\min} \hookrightarrow \mathrm{Gr}_{\mathrm{Sp}(\mathbf{M})}^{\omega}$. Here $Q \subset \mathrm{Gr}_{\mathrm{SO}(V')}$ is the minuscule $\mathrm{SO}(V')_{0}$ -orbit, and $\mathrm{Gr}_{\mathrm{Sp}(V)}^{\min} \subset \mathrm{Gr}_{\mathrm{Sp}(V)}$ is the minimal $\mathrm{Sp}(V)_{0}$ -orbit. We denote by \mathcal{F}_{Q} (resp. $\mathcal{F}_{\mathrm{min}}$) the restriction of $\exp(f)$ to $Q \times (t^{-1}\mathbf{M}_{0})/\mathbf{M}_{0}$ (resp. to $\mathrm{Gr}_{\mathrm{Sp}(V)}^{\min} \times (t^{-1}\mathbf{M}_{0})/\mathbf{M}_{0}$). Then $\phi_{\mathrm{SO}}(V) * E_{0} = \mathrm{pr}_{2*}\mathcal{F}_{Q}$, and $\phi_{\mathrm{Sp}}(V) * E_{0} = \mathrm{pr}_{2*}\mathcal{F}_{\mathrm{min}}$, where pr_{2} stands for the projection to the $(t^{-1}\mathbf{M}_{0})/\mathbf{M}_{0}$ factor. (Note that the extension of $\mathcal{F}_{\mathrm{min}}$ to $\overline{\mathrm{Gr}_{\mathrm{Sp}(V)}^{\min} \times (t^{-1}\mathbf{M}_{0})/\mathbf{M}_{0}$ is clean.)

Since the image of Q is contained in the fixed point set of the loop rotation in $\operatorname{Gr}_{\operatorname{Sp}(\mathbf{M})}^{*}$, it follows that \mathcal{F}_{Q} is a regular D-module, and $\phi_{\operatorname{SO}}(V) * E_{0} = \operatorname{pr}_{2*}\mathcal{F}_{Q}$ is a regular D-module as well. Moreover, $\phi_{\operatorname{Sp}}(V) * E_{0} = \operatorname{pr}_{2*}\mathcal{F}_{\min}$ is also regular. Indeed, $\operatorname{pr}_{2*}\mathcal{F}_{\min} = \operatorname{pr}_{*}p_*\mathcal{F}_{\min}$, where

$$p: \operatorname{Gr}_{\operatorname{Sp}(V)}^{\min} \times (t^{-1}\mathbf{M}_{\mathcal{O}})/\mathbf{M}_{\mathcal{O}} \to \mathbb{P}((t^{-1}V_{\mathcal{O}})/V_{\mathcal{O}}) \times (t^{-1}\mathbf{M}_{\mathcal{O}})/\mathbf{M}_{\mathcal{O}}$$

is the projection (contraction by the loop rotation action) in the first factor, and pr: $\mathbb{P}((t^{-1}V_0)/V_0) \times (t^{-1}\mathbf{M}_0)/\mathbf{M}_0 \to (t^{-1}\mathbf{M}_0)/\mathbf{M}_0$ is the second projection. But $p_*\mathcal{F}_{\min}$ is already a regular *D*-module whose support along $(t^{-1}\mathbf{M}_0)/\mathbf{M}_0$ is contained in a quadratic cone f = 0 (notation of [BFT, §2.13]).

Now that we know that both $\phi_{SO}(V) * E_0$ and $\phi_{Sp}(V) * E_0$ are regular, we describe the corresponding constructible (perverse) sheaves.

Let $C \subset V' \otimes V \cong t^{-1}(V' \otimes V)_0/(V' \otimes V)_0$ be the cone formed by all the tensors of the form $v' \otimes v$, where $v' \in V'_0$ (the cone of isotropic vectors). Then Chas two small resolutions: $((V'_0 \setminus \{0\}) \times V))/\mathbb{C}^{\times}_{hyperb}$, and $(V'_0 \times (V \setminus \{0\}))/\mathbb{C}^{\times}_{hyperb}$ (note that the second "resolution" is only *rationally* smooth). The pushforwards of the constant sheaves coincide with $\phi_{SO}(V) * E_0$ and $\phi_{Sp}(V) * E_0$ respectively; and they both coincide with the Goresky-MacPherson sheaf of C. This completes the identification in a).

b) We have $\operatorname{Hom}(E_0, \phi_{\operatorname{Sp}}(V) * \phi_{\operatorname{Sp}}(V) * E_0) = \operatorname{Hom}(\phi_{\operatorname{Sp}}(V) * E_0, \phi_{\operatorname{Sp}}(V) * E_0) = \operatorname{Hom}(\operatorname{IC}_C, \operatorname{IC}_C) = \mathbb{C}$. So the direct summand E_0 in $\phi_{\operatorname{Sp}}(V) * \phi_{\operatorname{Sp}}(V) * E_0$ is uniquely determined. Similarly, the direct summand E_0 in $\phi_{\operatorname{SO}}(V) * \phi_{\operatorname{SO}}(V) * E_0$ is uniquely determined and must coincide with the former summand under the identification in b).

3.2. The Hecke action generates \mathcal{W} -mod^{SO(V')_0 \times Sp(V)_0}. We first classify irreducible $\operatorname{GL}(N)_0 \times \operatorname{GL}(N)_0$ -equivariant *D*-modules on $\operatorname{Mat}(N \times N, \mathcal{K})$. Here by *D*-modules we mean the ones with support in a lattice in $\operatorname{Mat}(N \times N, \mathcal{K})$ invariant under translations by a smaller lattice. Thus we need to classify the $\operatorname{GL}(N)_0 \times \operatorname{GL}(N)_0$ -orbits in $\operatorname{Mat}(N \times N, \mathcal{K})$ invariant under translations by a lattice. These orbits are exactly the ones in $\operatorname{GL}(N)_{\mathcal{K}}$, and they are indexed by the length N signatures $\boldsymbol{\lambda}$.

Note that \mathbb{O}_{μ} lies in $\overline{\mathbb{O}}_{\lambda}$ iff

 $\lambda_1 \ge \mu_1, \ \lambda_1 + \lambda_2 \ge \mu_1 + \mu_2, \ \dots, \ \lambda_1 + \dots + \lambda_N \ge \mu_1 + \dots + \mu_N.$

The last condition is different from the usual definition of dominating order, where we require $\lambda_1 + \cdots + \lambda_N = \mu_1 + \cdots + \mu_N$. Indeed, $\operatorname{Mat}(N \times N, \mathcal{K})$ is connected, while $\pi_0(\operatorname{GL}(N)_{\mathcal{K}}) = \mathbb{Z}$.

Similarly, one proves

Lemma 3.2.1. The $\operatorname{GL}(M)_0 \times \operatorname{GL}(N)_0$ -orbits in $\operatorname{Mat}(M \times N, \mathcal{K})$ invariant under translations by a sublattice are indexed by the set of length M signatures (we assume $M \leq N$). An orbit \mathbb{O}_{λ} has a representative $(t^{-\lambda}, 0)$ (the last N - M columns are all zero).

The Fourier transform gives an equivalence

 $D\operatorname{-mod}(\operatorname{Mat}(M \times N, \mathcal{K}))^{\operatorname{GL}(M)_{\mathcal{O}} \times \operatorname{GL}(N)_{\mathcal{O}}} \xrightarrow{\sim} D\operatorname{-mod}(\operatorname{Mat}(N \times M, \mathcal{K}))^{\operatorname{GL}(N)_{\mathcal{O}} \times \operatorname{GL}(M)_{\mathcal{O}}}.$

The irreducible objects in both sides are indexed by the (translation invariant) orbits of $\operatorname{GL}(N)_{0} \times \operatorname{GL}(M)_{0}$, i.e. by the length M signatures. Hence we obtain an involution FT of the set of length M signatures.

Lemma 3.2.2. $FT(\lambda_1 \ge \ldots \ge \lambda_M) = (-\lambda_M \ge \ldots \ge -\lambda_1)$

Proof. The Fourier transform is compatible with the action of $\operatorname{GL}(N)_{\mathcal{K}} \times \operatorname{GL}(M)_{\mathcal{K}}$, and hence is compatible with the Hecke action. But duality induces the Chevalley involution on the Hecke category.

Now we take M = 2n, N = 2n + 1, and realize $SO(V') \times Sp(V)$ as the (connected component of the) fixed point set of an appropriate involution on $GL(N) \times GL(M)$. The argument of [BFT, §2.6] establishes

Corollary 3.2.3. The irreducible objects of W-mod^{SO(V')_0 × Sp(V)_0, \heartsuit} are indexed by the cone of dominant weights of Sp(V). Hence W-mod^{SO(V')_0 × Sp(V)_0} is generated by a collection of objects $\phi_{SO}(U) * E_0$, $U \in \text{Rep}(Sp(V))$.

3.3. A deequivariantized Ext-algebra. Making use of the convolution action $\operatorname{Rep}(\operatorname{Sp}(V)) \circlearrowright \mathcal{W}\operatorname{-mod}^{\operatorname{SO}(V')_{0} \times \operatorname{Sp}(V)_{0}}$, we consider the deequivariantized category

 $\mathcal{W}\operatorname{-mod}_{\operatorname{deeq}}^{\operatorname{SO}(V')_{0} \times \operatorname{Sp}(V)_{0}}$. Recall the unit object E_{0} in $\mathcal{W}\operatorname{-mod}^{\operatorname{SO}(V')_{0} \times \operatorname{Sp}(V)_{0}}$ introduced in §3.1. We will keep the same notation E_{0} for the corresponding object in $\mathcal{W}\operatorname{-mod}_{\operatorname{deeq}}^{\operatorname{SO}(V')_{0} \times \operatorname{Sp}(V)_{0}}$.

The following lemma is proved the same way as [BFT, Lemma 2.7.1].

Lemma 3.3.1. The dg-algebra $\operatorname{RHom}_{W\operatorname{-mod}_{\operatorname{deeq}}^{\operatorname{SO}(V')_{\odot} \times \operatorname{Sp}(V)_{\odot}}}(E_0, E_0)$ is formal, i.e. it is quasiisomorphic to the graded algebra $\operatorname{Ext}_{W\operatorname{-mod}_{\operatorname{deeq}}}^{\bullet}(E_0, E_0)$ with trivial differential.

We denote the dg-algebra $\operatorname{Ext}_{W-\operatorname{mod}_{\operatorname{deeq}}^{\operatorname{SO}(V')_{\odot} \times \operatorname{Sp}(V)_{\odot}}}^{\bullet}(E_0, E_0)$ (with trivial differential) by \mathfrak{E}^{\bullet} . Since it is an Ext-algebra in the deequivariantized category between objects induced from the original category, it is automatically equipped with an action of $\operatorname{Sp}(V)$, and we can consider the corresponding triangulated category $D_{\operatorname{perf}}^{\operatorname{Sp}(V)}(\mathfrak{E}^{\bullet})$. The following lemma is proved the same way as [BFT, Lemma 2.7.2], making use of Corollary 3.2.3.

Lemma 3.3.2. There is a canonical equivalence $D_{\text{perf}}^{\text{Sp}(V)}(\mathfrak{E}^{\bullet}) \xrightarrow{\sim} \mathcal{W} \operatorname{-mod}^{\mathrm{SO}(V')_{\odot} \times \operatorname{Sp}(V)_{\odot}}$.

We choose a decomposition $V = L \oplus L^*$ of V into a direct sum of two transversal Lagrangian subspaces. It defines a Siegel Levi subgroup $\operatorname{GL}(L) \subset \operatorname{Sp}(V)$. We choose Kostant slices $\Sigma_{\mathfrak{gl}} \subset \mathfrak{gl}(L)$ and $\Sigma_{\mathfrak{sp}} \subset \mathfrak{sp}(V)$. Making use of the trace forms, we identify $H^{\bullet}_{\operatorname{GL}(L)_{\mathbb{O}}}(\operatorname{pt}) \cong \mathbb{C}[\Sigma_{\mathfrak{gl}}]$ and $H^{\bullet}_{\operatorname{Sp}(V)_{\mathbb{O}}}(\operatorname{pt}) \cong \mathbb{C}[\Sigma_{\mathfrak{sp}}] \cong H^{\bullet}_{\operatorname{SO}(V')_{\mathbb{O}}}(\operatorname{pt})$. Restricting the equivariance from $\operatorname{Sp}(V)_{\mathbb{O}}$ to $\operatorname{GL}(L)_{\mathbb{O}}$, we obtain the deequiv-

Restricting the equivariance from $\operatorname{Sp}(V)_{\mathbb{O}}$ to $\operatorname{GL}(L)_{\mathbb{O}}$, we obtain the deequivariantized Ext-algebra $\widetilde{\mathfrak{E}}^{\bullet} := \operatorname{Ext}^{\bullet}_{\mathcal{W}\operatorname{-mod}_{\operatorname{does}}^{\operatorname{SO}(V')_{\mathbb{O}} \times \operatorname{GL}(L)_{\mathbb{O}}}(E_0, E_0).$

The purity argument used in the proofs of Lemmas 3.3.1 and 3.3.2 also establishes the following

Lemma 3.3.3. (1) \mathfrak{E}^{\bullet} is a free module over $H^{\bullet}_{\mathrm{SO}(V')_{\mathbb{O}} \times \mathrm{Sp}(V)_{\mathbb{O}}}(\mathrm{pt}) \cong \mathbb{C}[\Sigma_{\mathfrak{sp}} \times \Sigma_{\mathfrak{sp}}].$ (2) $\widetilde{\mathfrak{E}}^{\bullet} \cong \mathbb{C}[\Sigma_{\mathfrak{sp}} \times \Sigma_{\mathfrak{gl}}] \otimes_{\mathbb{C}[\Sigma_{\mathfrak{sp}} \times \Sigma_{\mathfrak{sp}}]} \mathfrak{E}^{\bullet}.$

Now we can prove

Lemma 3.3.4. The algebras $\mathfrak{E}^{\bullet}, \widetilde{\mathfrak{E}}^{\bullet}$ are commutative.

Proof. Making use of the SO(V') × GL(L)-invariant decomposition $V' \otimes V = V' \otimes L \oplus V' \otimes L^*$, we identify the categories

$$\mathcal{W}$$
-mod^{SO(V')_{0} \times GL(L)_{0} \cong D-mod $((V' \otimes L)_{\mathcal{K}})^{SO(V')_{0} \times GL(L)_{0}}$.}

The action of the central $\mathbb{G}_m \subset \operatorname{GL}(L) \subset \operatorname{GL}(L)_0$ contracts $(V' \otimes L)_{\mathcal{K}}$ to the origin. Hence by the Localization Theorem we have an isomorphism $\widetilde{\mathfrak{E}}^{\bullet}_{\operatorname{loc}} \simeq \operatorname{Ext}^{\bullet}_{\operatorname{D-mod}(\operatorname{pt})^{\operatorname{SO}(V')_0 \times \operatorname{GL}(L)_0}_{\operatorname{deeq}}(\delta_0, \delta_0)_{\operatorname{loc}}$. Here for a

module M over $H^{\bullet}_{\mathrm{SO}(V')_{0} \times \mathrm{GL}(L)_{0}}(\mathrm{pt}) \cong \mathbb{C}[\Sigma_{\mathfrak{sp}} \times \Sigma_{\mathfrak{gl}}]$ we denote by M_{loc} the extension of scalars to the field of fractions. Obviously, $\mathrm{Ext}^{\bullet}_{\mathrm{D-mod}(\mathrm{pt})^{\mathrm{SO}(V')_{0} \times \mathrm{GL}(L)_{0}}(\delta_{0}, \delta_{0})_{\mathrm{loc}} \simeq \mathbb{C}[\mathrm{Sp}(V) \times \Sigma_{\mathfrak{sp}} \times \Sigma_{\mathfrak{gl}}]_{\mathrm{loc}}$ is commutative. By Lemma 3.3.3, $\widetilde{\mathfrak{E}}^{\bullet}$ embeds into $\widetilde{\mathfrak{E}}^{\bullet}_{\mathrm{loc}}$. Hence $\widetilde{\mathfrak{E}}^{\bullet}$ is commutative, and \mathfrak{E}^{\bullet} is commutative as well.

Recall that $\mathfrak{G}^{\bullet} := \operatorname{Sym}^{\bullet}(\mathfrak{sp}(V)[-2]) \otimes \operatorname{Sym}^{\bullet}(\Pi(V)[-1])$ (see §2.2). We construct a homomorphism $\psi : \mathfrak{G}^{\bullet} \to \mathfrak{E}^{\bullet}$ as follows. First, the derived Satake equivalence $D_{\operatorname{perf}}^{\operatorname{Sp}(V)}(\operatorname{Sym}^{\bullet}(\mathfrak{sp}(V)[-2])) \cong \operatorname{D-mod}(\operatorname{Gr}_{\operatorname{SO}(V')})^{\operatorname{SO}(V')_{0}}$ along with the convolution action $\operatorname{D-mod}(\operatorname{Gr}_{\operatorname{SO}(V')})^{\operatorname{SO}(V')_{0}} \circlearrowright \mathcal{W} \operatorname{-mod}^{\operatorname{SO}(V')_{0} \times \operatorname{Sp}(V)_{0}}$ gives rise to a homomorphism $\operatorname{Sym}^{\bullet}(\mathfrak{sp}(V)[-2]) \to \mathfrak{E}^{\bullet}$. Second, the computation of $\phi_{\operatorname{SO}}(V) * E_{0}$ in the proof of Lemma 3.1.1 produces a canonical nonzero element of cohomological degree 1 in $\iota_{0}^{!}(\phi_{\operatorname{SO}}(V) * E_{0})$. Here ι_{0} stands for the embedding of the origin into the colattice $(V' \otimes V)_{\mathcal{K}}/(V' \otimes V)_{0}$, and $E_{0} = \delta_{0}$ as a D-module over $(V' \otimes V)_{\mathcal{K}}/(V' \otimes V)_{0}$. Since $\operatorname{Ext}_{\mathcal{W}\operatorname{-mod}^{\operatorname{SO}(V')_{0} \times \operatorname{Sp}(V)_{0}}(E_{0}, \phi_{\operatorname{SO}}(V) * E_{0} \otimes V^{*})$ is a direct summand of \mathfrak{E}^{\bullet} (and we have $V^{*} \cong V$), we obtain the desired embedding $V[-1] \hookrightarrow \mathfrak{E}^{\bullet}$.

By Lemma 3.3.2, in order to establish the equivalence of Theorem 2.2.1, it remains to prove the following

Proposition 3.3.5. The homomorphism $\psi \colon \mathfrak{G}^{\bullet} \to \mathfrak{E}^{\bullet}$ is an isomorphism.

Its proof will be given in $\S3.5$ after some preparation from Invariant Theory.

3.4. A pseudo-slice. We choose two transversal Lagrangian subspaces $V = L \oplus L^*$ as in §3.3. We choose a regular \mathfrak{sl}_2 -triple (e, h, f) in $\mathfrak{sp}(V)$ such that $L = \operatorname{Ker}(f^n)$, and $L^* = \operatorname{Ker}(e^n)$. We choose a vector $v_0^* \in L^* \setminus \operatorname{Ker}(e^{n-1})$ such that $\langle v_0^*, f v_0^* \rangle = 1$. The Kostant slice $\Sigma_{\mathfrak{sp}} = e + \mathfrak{z}_{\mathfrak{sp}(V)}(f)$. We define $\Sigma := \Sigma_{\mathfrak{sp}} \times (v_0^* + L) \subset \mathfrak{sp}(V) \oplus V$.

According to [S, row 3 of Table 4a], $\mathbb{C}[\mathfrak{sp}(V) \oplus V]^{\operatorname{Sp}(V)}$ is a polynomial algebra with 2n generators $\operatorname{Tr}(x^2), \ldots, \operatorname{Tr} x^{2n}, \langle v, xv \rangle, \ldots, \langle v, x^{2n-1}v \rangle$. Note that the projection $\pi \colon \Sigma \to (\mathfrak{sp}(V) \oplus V)/\!\!/ \operatorname{Sp}(V) \cong (\mathfrak{sp}(V))/\!\!/ \operatorname{Sp}(V) \times (\mathfrak{sp}(V))/\!\!/ \operatorname{Sp}(V) \cong \Sigma_{\mathfrak{sp}} \times \Sigma_{\mathfrak{sp}} \simeq \mathbb{A}^{2n}, (x, v) \mapsto [x], [x - v^2], \text{ is a surjective (ramified) cover of degree } 2^n$. This cover is not Galois for n > 1. In fact, for any $n \ge 1$, the projection $\mathfrak{sp}(V) \oplus V \to (\mathfrak{sp}(V) \oplus V)/\!\!/ \operatorname{Sp}(V) \cong \Sigma_{\mathfrak{sp}} \times \Sigma_{\mathfrak{sp}}$ admits no Weierstraß section.

Note that the Lagrangian subspace L is invariant with respect to $x - v^2$ for $(x, v) \in \Sigma$. We consider the composed map $\Sigma \to \mathfrak{gl}(L) \to \mathfrak{gl}(L)/\!\!/ \mathrm{GL}(L) =: \Sigma_{\mathfrak{gl}},$ $(x, v) \mapsto \mathrm{Spec}(x - v^2)$. Together with the first projection $\Sigma \to \Sigma_{\mathfrak{sp}}, (x, v) \mapsto x$, we obtain a morphism $\varpi : \Sigma \to \Sigma_{\mathfrak{sp}} \times \Sigma_{\mathfrak{gl}}$.

Lemma 3.4.1. The map $\varpi \colon \Sigma \to \Sigma_{\mathfrak{sp}} \times \Sigma_{\mathfrak{gl}}$ is an isomorphism.

Proof. The map ϖ takes $(x, v_0^* + \ell) \in \Sigma$ to $(x, \operatorname{Spec}(\operatorname{pr}_L(x) + v_0^* \otimes \ell))$, where $\operatorname{pr}_L: \mathfrak{sp}(V) \to \mathfrak{gl}(L)$ is the projection to the L-block with respect to decomposition

 $V = L \oplus L^*. \text{ We set } v_i := f^{n+1-i}v_0^* \in \operatorname{Ker}(f^i) \smallsetminus \operatorname{Ker}(f^{i-1}), \text{ and set } v = v_0^* + \sum_{i=1}^n a_i v_i. \text{ Let } t^n + \sum_{i=1}^n c_i t^{n-i} \text{ be the characteristic polynomial of } \operatorname{pr}_L(x) + v_0^* \otimes \ell.$ Then $c_i = \varkappa_i a_i + \varphi_i$, where $\varkappa_i \neq 0$ is a constant, while φ_i is a function of a_j , $1 \leq j < i$, and coordinates on $\Sigma_{\mathfrak{sp}}$. \Box

Lemma 3.4.2. The saturation $\operatorname{Sp}(V) \cdot \Sigma \subset \mathfrak{sp}(V) \oplus V$ of Σ is a constructible subset of $\mathfrak{sp}(V) \oplus V$. It contains an open subset \mathfrak{U} whose complement \mathfrak{Z} has codimension 2.

Proof. The image of the morphism $\operatorname{Sp}(V) \times \Sigma \to \mathfrak{sp}(V) \times V \times \operatorname{LGr}(V)$ (Lagrangian Grassmannian), $(g, x, v) \mapsto (\operatorname{Ad}_g(x), gv, gL)$, contains a locally closed subvariety $X \subset \mathfrak{sp}(V) \times V \times \operatorname{LGr}(V)$ consisting of triples (x', v', L') satisfying the following conditions:

- a) $x' \in \mathfrak{sp}(V)$ is a regular element,
- b) $x' v'^2 \in \mathfrak{sp}(V)$ preserves L',
- c) $v' \pmod{L'}$ is a cyclic vector for $x' v'^2 \pmod{L'}$.

Furthermore, the natural morphism $\operatorname{Sp}(V) \times \Sigma \to \mathfrak{sp}(V) \oplus V$, $(g, x, v) \mapsto (\operatorname{Ad}_g(x) - (gv)^2, gv)$, factors as the composition $\operatorname{Sp}(V) \times \Sigma \to \mathfrak{sp}(V) \times V \times \operatorname{LGr}(V) \to \mathfrak{sp}(V) \oplus V$, where the second morphism takes (x', v', L') to $(x' - v'^2, v')$.

So we must find the desired open subspace \mathcal{U} in the image of X in $\mathfrak{sp}(V) \oplus V$. If $x' - v'^2$ has a zero eigenvalue (a codimension 1 condition), then for (x', v') in the image of X, the corresponding component v'_0 is a cyclic vector for the corresponding block $(x' - v'^2)_0$. The complement to the set of such pairs has codimension 1. All in all, the set of pairs (x', v') such that $x' - v'^2$ is not invertible, and v'_0 is not a cyclic vector for $(x' - v'^2)_0$ has codimension 2 in $\mathfrak{sp}(V) \oplus V$.

Now if $x' - v'^2$ is invertible, for each pair of opposite eigenvalues $(\lambda, -\lambda)$, either the corresponding component v'_{λ} is a cyclic vector for the corresponding block $(x' - v'^2)_{\lambda}$, or $v'_{-\lambda}$ is a cyclic vector for $(x' - v'^2)_{-\lambda}$. The complement to the set of such pairs again has codimension 2 in $\mathfrak{sp}(V) \oplus V$.

This completes the proof of the lemma.

We consider the fibre product (over the second copy of $\Sigma_{\mathfrak{sp}} = \mathfrak{sp}(V)/\!\!/ \operatorname{Sp}(V)$) $\mathfrak{M} := (\mathfrak{sp}(V) \oplus V) \times_{\Sigma_{\mathfrak{sp}}} \Sigma_{\mathfrak{gl}}$. The above maps $\Sigma \to \mathfrak{sp}(V) \oplus V$ and $\Sigma \to \Sigma_{\mathfrak{gl}}$ give rise to a closed embedding $\Sigma \hookrightarrow \mathfrak{M}$.

Corollary 3.4.3. (1) We have $\mathfrak{M}/\!\!/\mathrm{Sp}(V) = \Sigma_{\mathfrak{sp}} \times \Sigma_{\mathfrak{gl}}$, and $\Sigma \hookrightarrow \mathfrak{M}$ is a Weierstraß section of the $\mathrm{Sp}(V)$ -action.

(2) $\mathbb{C}[\mathfrak{sp}(V) \oplus V] = \mathbb{C}[\Sigma \times \operatorname{Sp}(V)] \times_{\mathbb{C}(\mathfrak{M})} \mathbb{C}(\mathfrak{sp}(V) \oplus V).$

3.5. Equivariant cohomology. Recall from the proof of Lemma 3.4.2 that Σ is equipped with a \mathbb{G}_m -action. It corresponds to the half-grading in the RHS of the isomorphism $\mathbb{C}[\Sigma] \cong H^{\bullet}_{\mathrm{SO}(V')_{0} \times \mathrm{GL}(L)_{0}}(\mathrm{pt})$. We have a functor χ : Rep(Sp(V)) \rightarrow Vect $^{\mathbb{G}_m}(\Sigma)$ (\mathbb{G}_m -equivariant vector bundles), $U \mapsto H^{\bullet}_{\mathrm{SO}(V')_{0} \times \mathrm{GL}(L)_{0}}((V' \otimes L)_{\mathcal{K}}, \phi_{\mathrm{SO}}(U) * E_{0}).$ Here we view $\phi_{\mathrm{SO}}(U) * E_{0}$ as an $\mathrm{SO}(V')_{0} \times \mathrm{GL}(L)_{0}$ -equivariant *D*-module on $(V' \otimes L)_{\mathcal{K}}.$

Lemma 3.5.1. χ is a tensor functor.

Proof. We equip \mathcal{W} -mod^{SO(V')_{\circ} \times GL(L)_{\circ} \cong D-mod $((V' \otimes L)_{\mathcal{K}})^{SO(V')_{\circ} \times GL(L)_{\circ}}$ with the fusion tensor structure. Then the functor}

$$\operatorname{Rep}(\operatorname{Sp}(V)) \to \operatorname{D-mod}((V' \otimes L)_{\mathcal{K}})^{\operatorname{SO}(V')_{\mathcal{O}} \times \operatorname{GL}(L)_{\mathcal{O}}}, \ U \mapsto \phi_{\operatorname{SO}}(U) * E_0$$

is a tensor functor. Its essential image lies in the full subcategory D-mod $((V' \otimes L)_{\mathcal{K}})_{ss}^{SO(V')_{0} \times GL(L)_{0}}$ of semisimple *D*-modules in the heart of the natural *t*-structure. The functor D-mod $((V' \otimes L)_{\mathcal{K}})_{ss}^{SO(V')_{0} \times GL(L)_{0}} \to \operatorname{Vect}^{\mathbb{G}_{m}}(\Sigma)$, $\mathcal{F} \mapsto H^{\bullet}_{SO(V')_{0} \times GL(L)_{0}}((V' \otimes L)_{\mathcal{K}}, \phi_{SO}(U) * E_{0})$ is also a tensor functor. Indeed, consider the dilation action of \mathbb{G}_{m} on $(V' \otimes L)_{\mathcal{K}}$. It coincides with the action of the central $\mathbb{G}_{m} \subset \operatorname{GL}(L) \subset \operatorname{SO}(V') \times \operatorname{GL}(L) \subset \operatorname{SO}(V')_{0} \times \operatorname{GL}(L)_{0}$. It contracts $(V' \otimes L)_{\mathcal{K}}$ to the origin, and \mathcal{F} is a monodromic *D*-module. Hence $H^{\bullet}_{\operatorname{SO}(V')_{0} \times \operatorname{GL}(L)_{0}}((V' \otimes L)_{\mathcal{K}}, \phi_{SO}(U) * E_{0}) \cong i_{0}^{*}\mathcal{F}$, where i_{0} stands for the embedding of the origin into $(V' \otimes L)_{\mathcal{K}}$. So the desired tensor property is the simplest instance of the commutation of nearby cycles and hyperbolic restriction [N1, Proposition 5.4.1(2)].

Finally, the tensor property of χ follows since it is a composition of two tensor functors.

The tensor functor χ defines a \mathbb{G}_m -equivariant $\operatorname{Sp}(V)$ -torsor \mathfrak{T} over Σ . The deequivariantized Ext-algebra \mathfrak{E}^{\bullet} by definition acts on $\mathbb{C}[\operatorname{Sp}(V)] * E_0$, and we obtain a morphism $\mathfrak{T} \to \operatorname{Spec} \mathfrak{E}$. In other words, we obtain a morphism $\alpha \colon \Sigma \to \operatorname{Spec} \mathfrak{E}/\operatorname{Sp}(V)$ (the stacky quotient).

The homomorphism $\psi \colon \mathfrak{G}^{\bullet} \to \mathfrak{E}^{\bullet}$ of Proposition 3.3.5 gives rise to the same named $\mathbb{G}_m \times \operatorname{Sp}(V)$ -equivariant morphism of spectra $\operatorname{Spec}(\mathfrak{E}) \to \operatorname{Spec}(\mathfrak{G}) = \mathfrak{sp}(V) \oplus V$. Consider the composition

$$\Sigma \xrightarrow{\alpha} \operatorname{Spec}(\mathfrak{E})/\operatorname{Sp}(V) \xrightarrow{\psi} (\mathfrak{sp}(V) \oplus V)/\operatorname{Sp}(V).$$

Lemma 3.5.2. (1) The composition $\psi \circ \alpha$ coincides (up to a unique isomorphism) with the tautological morphism $\Sigma \to (\mathfrak{sp}(V) \oplus V)/\mathrm{Sp}(V)$.

(2) The $\operatorname{Sp}(V)$ -torsor $\mathfrak{T} \to \Sigma$ is trivialized: $\mathfrak{T} \cong \operatorname{Sp}(V) \times \Sigma$.

Proof. (1) We know that the further composition $\Sigma \to \operatorname{Spec} \mathfrak{E}/\operatorname{Sp}(V) \to \operatorname{Spec} \mathfrak{G}/\operatorname{Sp}(V) \to (\mathfrak{sp}(V) \oplus V)/\!\!/\operatorname{Sp}(V)$ is the tautological map $\Sigma \to \Sigma_{\mathfrak{sp}} \times \Sigma_{\mathfrak{sp}}$ since our monoidal functor is compatible with the action of \mathfrak{E}^{\bullet} on E_0 . We also know that $\psi \circ \alpha \colon \Sigma \to (\mathfrak{sp}(V) \oplus V)/\operatorname{Sp}(V)$ is equivariant with respect to the \mathbb{G}_m -action on $\mathfrak{sp}(V) \oplus V$ given by $(2\rho^{\vee} - 2, 2\rho^{\vee} - 1)$ (that is $c(x, v) = (c^{-2}\operatorname{Ad}_{2\rho^{\vee}(c)} x, c^{-1}(2\rho^{\vee}(c))v)).$ So we have to check that there is a unique \mathbb{G}_m -equivariant morphism $\gamma: \Sigma \to (\mathfrak{sp}(V) \oplus V)/\mathrm{Sp}(V)$ whose composition with the map $(\mathfrak{sp}(V) \oplus V)/\mathrm{Sp}(V) \to (\mathfrak{sp}(V) \oplus V)//\mathrm{Sp}(V)$ is the tautological map $\Sigma \to \Sigma_{\mathfrak{sp}} \times \Sigma_{\mathfrak{sp}}$.

First we check that the base point $0 \in \Sigma$ must go to (the orbit of) the point $(e, v_0^*) \in \mathfrak{sp}(V) \oplus V$. Indeed, the first component of $\gamma(0)$ must be a regular nilpotent in $\mathfrak{sp}(V)$: otherwise the composition of γ with the first projection $(\mathfrak{sp}(V) \oplus V)/\mathrm{Sp}(V) \to \mathfrak{sp}(V)/\!/\mathrm{Sp}(V) = \Sigma_{\mathfrak{sp}}$ will not be smooth at $0 \in \Sigma$ (the differential will not be surjective). So we can assume that the first component of $\gamma(0)$ is $e \in \mathfrak{sp}(V)$. The second component of $\gamma(0)$ must be a \mathbb{G}_m -fixed point in V, so it is cv_0^* for a constant c, and we must prove that $c = \pm 1$. It is easy to see that otherwise the ramification locus of the map $\Sigma \to \Sigma_{\mathfrak{sp}} \times \Sigma_{\mathfrak{sp}}$ obtained as the composition of γ with the projection $(\mathfrak{sp}(V) \oplus V)/\mathrm{Sp}(V) \to (\mathfrak{sp}(V) \oplus V)//\mathrm{Sp}(V) = \Sigma_{\mathfrak{sp}} \times \Sigma_{\mathfrak{sp}}$, does not contain $\Sigma_{\mathfrak{sp}} \times \{0\}$. This contradicts our assumption since the tautological map $\Sigma \to \Sigma_{\mathfrak{sp}} \times \Sigma_{\mathfrak{sp}}$ is ramified over $\Sigma_{\mathfrak{sp}} \times \{0\}$.

Now consider the Borel subalgebra $\mathfrak{b} \subset \mathfrak{sp}(V)$ containing f. Let \mathfrak{u} be the nilpotent radical of \mathfrak{b} , and let $U \subset \operatorname{Sp}(V)$ be the corresponding unipotent subgroup. The attractor \mathcal{A} to the point (e, v_0^*) of the \mathbb{G}_m -action on $\mathfrak{sp}(V) \oplus V$ is $\{(e+\mathfrak{b}, v_0^* + L)\}$. It remains to note that given a \mathbb{G}_m -equivariant morphism $\gamma: \Sigma \to \mathcal{A}$ such that the composition of γ with the projection $\mathcal{A} \to (\mathfrak{sp}(V) \oplus V)//\operatorname{Sp}(V) = \Sigma_{\mathfrak{sp}} \times \Sigma_{\mathfrak{sp}}$ is the tautological map $\Sigma \to \Sigma_{\mathfrak{sp}} \times \Sigma_{\mathfrak{sp}}$, there is a morphism $\Upsilon: \Sigma \to U$ such that $\gamma = \operatorname{Ad}_{\Upsilon}(\sigma)$. Here $\sigma: \Sigma \hookrightarrow \mathfrak{sp}(V) \oplus V$ is the canonical embedding. Indeed, the map $U \times \Sigma \to \mathcal{A}$, $(u, x, v) \mapsto u \cdot \sigma(x, v)$ is an isomorphism. This completes the proof of (1).

(2) follows since the tautological morphism $\Sigma \to (\mathfrak{sp}(V) \oplus V)/\operatorname{Sp}(V)$ arises from the following map of the trivial $\operatorname{Sp}(V)$ -torsor $\operatorname{Sp}(V) \times \Sigma$ into $\mathfrak{sp}(V) \oplus V$, $(g, x, v) \mapsto g \cdot \sigma(x, v)$.

Now we can prove Proposition 3.3.5. Recall the algebras $\mathfrak{E} \subset \mathfrak{E}_{\text{loc}}$ introduced in §3.3. (We ignore the gradings from now on, so we omit the superscript •.) Similarly, we introduce an algebra $\mathfrak{G} := \mathbb{C}[\mathfrak{M}]$ (recall from §3.4 that $\mathfrak{M} := (\mathfrak{sp}(V) \oplus V) \times_{\Sigma_{\mathfrak{sp}}} \Sigma_{\mathfrak{gl}}$, and $\mathfrak{G} = \mathbb{C}[\mathfrak{sp}(V) \oplus V]$). Then the homomorphism ψ of Proposition 3.3.5 induces $\tilde{\psi} : \mathfrak{G} \to \mathfrak{E}$. From the proof of Lemma 3.3.4 we know that $\tilde{\psi}$ is injective, since both \mathfrak{G} and \mathfrak{E} embed into $\mathfrak{G}_{\text{loc}} \simeq \mathfrak{E}_{\text{loc}}$. It follows that ψ is an embedding $\mathfrak{G} \hookrightarrow \mathfrak{E} \subset \text{Frac}(\mathfrak{G}) = \mathbb{C}(\mathfrak{sp}(V) \oplus V)$. Hence we may view an element $f \in \mathfrak{E}$ as a rational function on $\mathfrak{sp}(V) \oplus V$. We know that the pullback of this rational function to $\operatorname{Sp}(V) \times \Sigma$ (under the natural morphism $\operatorname{Sp}(V) \times \Sigma \to \mathfrak{sp}(V) \oplus V$, $(g, x, v) \mapsto g \cdot \sigma(x, v)$) is regular by Lemma 3.5.2. By Corollary 3.4.3(2) we conclude that f is regular. Hence the embedding $\psi : \mathfrak{G} \hookrightarrow \mathfrak{E}$ is an equality.

This completes the proof of Proposition 3.3.5 along with Theorem 2.2.1.

References

- [BDFRT] A. Braverman, G. Dhillon, M. Finkelberg, S. Raskin, R. Travkin, Coulomb branches of noncotangent type (with appendices by Gurbir Dhillon and Theo Johnson-Freyd), arXiv:2201.09475.
- [BF] R. Bezrukavnikov, M. Finkelberg, *Equivariant Satake category and Kostant-Whittaker reduction*, Moscow Math. Journal **8** (2008), no. 1, 39–72.
- [BFN] A. Braverman, M. Finkelberg, H. Nakajima, *Ring objects in the equivariant derived Satake category arising from Coulomb branches* (with appendix by Gus Lonergan), Adv. Theor. Math. Phys. 23 (2019), no. 2, 253–344.
- [BFT] A. Braverman, M. Finkelberg, R. Travkin, Orthosymplectic Satake equivalence, II, arXiv:2207.03115.
- [BZSV] D. Ben-Zvi, Y. Sakellaridis and A. Venkatesh, *Relative Langlands duality*, arXiv:2409.04677.
- [CW] T.-H. Chen, J. Wang, Derived Satake equivalence for Godement-Jacquet monoids, https://www.jonathanpwang.com/notes/GJsatake_purdue_handout.pdf (2021).
- [DLYZ] G. Dhillon, Y.-W. Li, Z. Yun, X. Zhu, *Endoscopy for affine Hecke algebras*, in preparation.
- [GS] D. Gaitsgory and S. Lysenko, Parameters and duality for the metaplectic geometric Langlands theory, Selecta Math. (N.S.) 24 (2018), no.1, 227–301.
- [L1] S. Lysenko, Moduli of metaplectic bundles on curves and theta-sheaves, Ann. Sci.
 École Norm. Sup. (4) 39 (2006), no. 3, 415–466.
- [L2] S. Lysenko, *Geometric Waldspurger periods II*, Represent. Theory **24** (2020), 235–291.
- [N1] H. Nakajima, Lectures on perverse sheaves on instanton moduli spaces, in Geometry of Moduli Spaces and Representation Theory, IAS Park City Mathematics Series, 24, AMS, Providence, RI (2017), 381–436.
- [N2] H. Nakajima, S-dual of Hamiltonian G-spaces and relative Langlands duality, Abstracts of the 71st Geometry Symposium, arXiv:2409.06303.
- [R] S. Raskin, Homological methods in semi-infinite contexts, https://web.ma.utexas.edu/users/sraskin/topalg.pdf (2019).
- [S] G. Schwarz, Representations of simple Lie groups with regular rings of invariants, Inventiones Math. 49 (1978), 167–191.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO AND PERIMETER INSTITUTE OF THEORETICAL PHYSICS, WATERLOO, ONTARIO, CANADA, N2L 2Y5 Email address: braval@math.toronto.edu

EINSTEIN INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, ED-MOND J. SAFRA CAMPUS, GIV'AT RAM, JERUSALEM, 91904, ISRAEL; NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS; SKOLKOVO INSTITUTE OF SCIENCE AND TECHNOLOGY

Email address: fnklberg@gmail.com

EINSTEIN INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, ED-MOND J. SAFRA CAMPUS, GIV'AT RAM, JERUSALEM, 91904, ISRAEL Email address: kazhdan@math.huji.ac.il

SKOLKOVO INSTITUTE OF SCIENCE AND TECHNOLOGY, MOSCOW, RUSSIA *Email address:* roman.travkin2012@gmail.com

14