COLOR SYMMETRY AND FERROMAGNETISM IN POTTS SPIN GLASS

HONG-BIN CHEN

ABSTRACT. We consider the Potts spin glass with additional ferromagnetic interaction parametrized by t. It has long been observed that the Potts color symmetry breaking for the spin glass order parameter is closely related to the ferromagnetic phase transition. To clarify this, we identify a single critical value t_c , which marks the onset of both color symmetry breaking and the transition to ferromagnetism.

1. INTRODUCTION

1.1. Setting and notation. Throughout, for matrices or vectors a and b of the same dimension, we denote by $a \cdot b$ the entry-wise inner product and write $|a| = \sqrt{a \cdot a}$. Let **1** be the vector in \mathbb{R}^D with all entries equal to 1. We write $\mathbb{R}_+ = [0, \infty)$ and $\mathbb{R}_{++} = (0, \infty)$.

Fix an integer $D \ge 2$ and let μ be the uniform probability measure on the standard basis $\{e_1, \ldots, e_D\}$ of \mathbb{R}^D viewed as the state space of the *D*-Potts spins. We interpret $\{1, \ldots, D\}$ as the labels of *D* instinct colors. For each $N \in \mathbb{N}$, we sample column vectors $\sigma_{\bullet i} = (\sigma_{di})_{1 \le d \le D}$ independently from μ , for each $i \in \{1, \ldots, N\}$. We denote by $\sigma = (\sigma_{di})_{1 \le d \le D}$, $1 \le i \le N$ the spin configuration, which is viewed as a $D \times N$ matrix.

The \mathbb{R}^{D} -valued mean magnetization is defined as

(1.1)
$$m_N = N^{-1} \sum_{i=1}^N \sigma_{\bullet i}$$

Notice that for Potts spins, we have the following relations

(1.2)
$$N^{-1}\sigma\sigma^{\mathsf{T}} = \operatorname{diag}(m_N), \quad m_N = N^{-1} \Big(\# \{ i \in \{1, \dots, N\} : \sigma_{\bullet i} = e_d \} \Big)_{1 \leq d \leq D}.$$

Hence, the $\mathbb{R}^{D \times D}$ -valued self-overlap $N^{-1} \sigma \sigma^{\dagger}$ contains the same information of the mean magnetization m_N .

We consider the spin glass interaction

$$H_N(\sigma) = \frac{1}{\sqrt{N}} \sum_{i,j=1}^N g_{ij} \sigma_{\bullet i} \cdot \sigma_{\bullet j}.$$

where $(g_{ij})_{1 \leq i,j \leq N}$ is a collection of i.i.d. standard Gaussian random variables. Throughout, we fix $\beta \ge 0$ to be the inverse temperature. For $N \in \mathbb{N}$, $t \in \mathbb{R}_+$, and $x \in \mathbb{R}^D$, we consider the augmented Hamiltonian

(1.3)
$$H_{\beta,N}(t,x,\sigma) = \beta H_N(\sigma) + \left(t - \frac{\beta^2}{2}\right) N|m_N|^2 + Nx \cdot m_N$$

where on the right-hand side the second term accounts for the additional ferromagnetic interaction and the third term is an external magnetic field. At t = 0 and x = 0, we can

²⁰²⁰ Mathematics Subject Classification. 82B44, 82D30.

Key words and phrases. Potts spin glass, color symmetry, ferromagnetism, phase transition.

use (1.2) to write

(1.4)
$$H_{\beta,N}(0,0,\sigma) = \beta H_N(\sigma) - \frac{\beta^2}{2} N \left| \frac{\sigma \sigma^{\mathsf{T}}}{N} \right|^2$$

which is the Hamiltonian in the model with self-overlap correction considered in [16]. The self-overlap correction refers to $-\frac{\beta^2}{2}N \left|\frac{\sigma\sigma^{\intercal}}{N}\right|^2$ which is $-\frac{1}{2}$ times the variance of $H_N(\sigma)$ (viewed as a Gaussian random variable). The main result in [16] states that the Parisi formula for the model (1.4) is an infimum taken over paths respecting the color symmetry of the Potts spin (i.e. invariant under permutation of color labels $\{1, \ldots, D\}$). In the setting of vector spin glass which includes the Potts one as a special case, the model with self-overlap correction [14, 17] is easier to deal with than the standard ones, partly due to the reason that the Parisi formula is an infimum rather than a sup inf formula [43, 44] for the standard model without the self-overlap correction. This is the reason for us to view (1.4) as the base model and add ferromagnetic interactions on top of it as in (1.3).

We also mention that it is natural to view the term $\left(t - \frac{\beta^2}{2}\right) N |m_N|^2$ through (1.1) as shifting the mean of (g_{ij}) , which is a more common perspective used in, e.g., [27, 25, 26, 34, 30, 22, 8].

For $N \in \mathbb{N}$ and $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^D$, the free energy is defined as

(1.5)
$$F_{\beta,N}(t,x) = \frac{1}{N} \mathbb{E} \log \int \exp\left(H_{\beta,N}(t,x,\sigma)\right) \otimes_{i=1}^{N} \mathrm{d}\mu(\sigma_{\bullet i})$$

where \mathbb{E} averages over (g_{ij}) . The associated Gibbs measure is denoted as

(1.6)
$$\langle \cdot \rangle_{t,x} \propto \exp\left(H_{\beta,N}(t,x,\sigma)\right) \otimes_{i=1}^{N} \mathrm{d}\mu(\sigma_{\bullet i})$$

where we choose to omit β and N from the notation since β is fixed and N will be clear from the context.

1.2. Motivation, overview of results, and related works. It has long been observed in physics [27, 25, 26, 34, 30, 22, 8] that sometimes (i.e. for certain D and β) additional antiferromagnetic interaction (i.e. $t < \frac{\beta^2}{2}$) is needed to ensure that the spin glass order parameter respects the Potts color symmetry. This suggests a close link between ferromagnetism and color symmetry breaking. But to the best knowledge of the author, there has not been rigorous studies of it yet.

We describe our main results. For fixed D and β , at x = 0 (zero external field), we clarify the aforementioned link by finding a critical value $t_c > 0$ marking both the onset of color symmetry breaking (in terms of the spin glass order parameter) and the transition into ferromagnetism (in terms of the mean magnetization, also called the conventional order parameter in the context of spin glasses). Our results on phase transitions are summarized in the table below.

$t < t_{\rm c}$	$t > t_{ m c}$
Color symmetry	Color symmetry breaking
"Zero" magnetization	Spontaneous magnetization

Since the Potts spins do not average to zero, here "zero" means zero after proper recentering. We define t_c in (2.12) as the first time when $\lim_{N\to\infty} F_{\beta,N}(t,0)$ starts to be non-linear in t. Proposition 2.6 gives equivalent characterizations of t_c in terms of the differentiability of $\lim_{N\to\infty} F_{\beta,N}(t,\cdot)$ at 0 and in terms of the number of maximizers for the variational representation of the limit. The notion of color symmetry is given precisely in Definition 2.8. The color symmetry breaking result shown in the table is proved in Proposition 2.9. The ferromagnetic phase transition is proved in Proposition 2.10. It is reasonable to expect that ferromagnetism implies color symmetry breaking (since one color is favored over others in the ferromagnetic phase), but it is interesting to see that the two phase transitions happen simultaneously since they are associated with different kinds of order parameters. We attempt to heuristically explain this phenomenon in the following. The spin glass order parameter can be viewed as a increasing path that optimizes the Parisi formula (see (2.21)). The right endpoint of the path is "ideally" equal to the limit of self-overlap.¹ Due to (1.2), the endpoint is equivalent to the mean magnetization. We will verify that the optimal path is color-symmetric if and only if the endpoint is so. On the other hand, we will show that the mean magnetization is color-symmetric if and only if it is "zero". Hence, "zero" magnetization corresponds exactly to the color symmetry of the spin glass order parameter.

Next, we make two comments on the model:

- Notice that at x = 0, the law of the overlap (at each N) is always color-symmetric at any t, and thus so is its limit. When $t > t_c$, due to the color symmetry breaking, the optimal path of the Parisi formula is not color-symmetric. Hence, in this supercritical regime, the overlap does not converge to the optimal path. An analogous statement can be said about the self-overlap at $t > t_c$.
- Also, notice that, since $t_c > 0$, the color symmetry is always preserved at t = 0 for every β . In this case, the Hamiltonian is given by (1.4), where the self-overlap correction is present. This correction can be interpreted as subtracting from $\beta H_N(\sigma)$ the "correct" amount of ferromagnetic interaction. Furthermore, at t = 0, the Parisi formula (see (2.20)) is an infimum, as in the Sherrington-Kirkpatrick model, whereas for every t > 0, the Parisi formula takes the max-min form, as in (2.21).

We describe another source of motivation from [40]. When $t = \frac{\beta^2}{2}$ and x = 0, the model becomes standard. Hence, it is interesting to compare t_c with $\frac{\beta^2}{2}$. We display the dependence of t_c on D and β by writing $t_c = t_c(D, \beta)$. Recently, [40] showed that for sufficiently large D, there is a range of β such that $t_c(D, \beta) < \frac{\beta^2}{2}$ (color symmetry breaking in the standard Potts spin glass). In this work, however, we do not pursue questions along this line.

Lastly, we briefly mention other related works. When D = 2, the Potts spin glass is essentially the same as the Sherrington–Kirkpatrick model [48, 45, 46, 32, 49, 42] (but we do not see the phenomenon discussed here, since the self-overlap of Ising spins is constantly one). The Potts spin glass was introduced in [27] and the associated Parisi formula was first established in [43]. Recently, [7] showed that the spin-glass order parameter is color symmetric in the constrained model, which inspired [16] to show a similar result for the model with self-overlap correction. The argument in the two works was further generalized in [33]. The studies of the non-disordered Curie–Weiss–Potts model include [28, 21, 24, 35]. The Potts spin glass model in consideration here has a conventional order parameter, the mean magnetization, in addition to the spin glass order parameter. Studies of the interplay of the two kinds of parameters include [36, 20, 9, 3, 15]. Spin glass models with self-overlap correction have been considered in [14, 17], inspired by the Hamilton–Jacobi equation approach to spin glass. Also, at the core of the analysis is the Hopf–Lax formula for the solution of such an equation. Considerations along this line include [38, 41, 37, 39, 23, 18, 31, 5, 4, 1, 6, 29, 12, 19, 11].

¹The right endpoint is in general below (or equal to) the limit of self-overlap as the overlap is bounded by the self-overlap. For instance, in the replica symmetric regime of the Sherrington–Kirkpatrick model, the overlap is zero while the self-overlap is always one. But, we can always approximate optimal true path by a near-optimal path with the right endpoint equal to the limit of self-overlap.

HONG-BIN CHEN

2. Proofs

We prove the main results as described in the introduction. We start with some preparation.

2.1. Properties of the initial condition. We view $F_{\beta,N}$ (see (1.5)) as a function on $\mathbb{R}_+ \times \mathbb{R}^D$. We denote the limit of initial condition as

(2.1)
$$\mathscr{P}_{\beta}(x) = \lim_{N \to \infty} F_{\beta,N}(0,x), \quad \forall x \in \mathbb{R}^{D}.$$

Recall the discussion below (1.4) that the case t = 0 corresponds to the model with self-overlap correction. The existence of the limit in (2.1) is ensured by [14, Theorem 1.1]. One should think of $\mathscr{P}_{\beta}(x)$ as the Parisi formula associated with the model $H_{\beta,N}(0, x, \sigma)$ (see (1.3)) or the model $H_{\beta,N}(0,0,\sigma)$ with single spin distribution $e^{x\cdot\tau}\mu(d\tau)$ (instead of $\mu(d\tau)$). Just as the Parisi formula for the Sherrington–Kirkpatrick model, due to the presence of the self-overlap correction $-\frac{\beta^2}{2}N|m_N|^2$, $\mathscr{P}_{\beta}(x)$ is an infimum over increasing paths. This will be later described in more detail in Section 2.4.

Remark 2.1 (Matching notation in [17]). Since we will use results from [17], we clarify that

$$F_{\beta,N}(0,x) = F_N(0,\operatorname{diag}(x)), \quad \forall x \in \mathbb{R}^D,$$

in the Potts setting, where free energy on the right-hand side is the one considered in [17]. The relation holds due to (1.2). As a result of (2.1) and [17, (1.3) and (1.10)], we have

$$\mathscr{P}_{\beta}(x) = \mathscr{P}(\operatorname{diag}(x)), \quad \forall x \in \mathbb{R}^{D}$$

where \mathscr{P} is defined in [17, (1.10)]. There, the domain for F_N and \mathscr{P} is the set of $D \times D$ real symmetric matrices. We also need one more important identity. Since the self-overlap $N^{-1}\sigma\sigma^{\dagger}$ is diagonal (see (1.2)), we can infer from [17, Theorem 1.1 (1)] that the symmetric matrix-valued $\nabla \mathscr{P}$ is always diagonal. Using this and the previous display, we have

(2.2)
$$\operatorname{diag}\left(\nabla \mathscr{P}_{\beta}(x)\right) = \nabla \mathscr{P}(\operatorname{diag}(x)), \quad \forall x \in \mathbb{R}^{D}$$

Here, the everywhere differentiability of \mathscr{P} (and thus \mathscr{P}_{β}) is ensured by [17, Proposition 2.3].

Lemma 2.2 (Properties of the initial condition). The function $\mathscr{P}_{\beta} : \mathbb{R}^{D} \to \mathbb{R}$ is differentiable everywhere, Lipschitz, convex, and semi-concave. Moreover, $\mathbf{1} \cdot \nabla \mathscr{P}_{\beta}(x) = 1$ for every $x \in \mathbb{R}^{D}$.

Here, a function is semi-concave if, after subtracting some quadratic function, it becomes concave.

Proof. We use Remark 2.1 and results from [17]. We know from [17, Lemma 2.2] that the first and second order derivatives of $\mathscr{P}_{\beta}(\pi, \cdot)$ are bounded everywhere on \mathbb{R}^{D} and uniformly in π . It is well known (e.g. see [10, Proposition 1.5]) that the infimum of functions with uniform bounds on concavity is semi-concave. Hence, \mathscr{P}_{β} is semi-concave. Other properties in the first sentence of the statement can be found in [17, Lemma 2.1 and Proposition 2.3]. To show the second part, we use Remark 2.1 to see

(2.3)
$$\mathbf{1} \cdot \nabla \mathscr{P}_{\beta}(x) = \operatorname{diag}(\mathbf{1}) \cdot \nabla \mathscr{P}(\operatorname{diag}(x)), \quad \forall x \in \mathbb{R}^{D}.$$

Here, on the right-hand side, $\nabla \mathscr{P}$ is the derivative with respect to the entry-wise inner product of $D \times D$ matrices (Frobenius inner product). So, $\nabla \mathscr{P}$ is $\mathbb{R}^{D \times D}$ -valued. The dot product on the right-hand side is this inner product. Recall that the differentiability of

 \mathscr{P} is ensured by [17, Proposition 2.3]. Applying [17, Theorem 1.1 (1)] to the right-hand side of (2.3), we have

$$\mathbf{1} \cdot \nabla \mathscr{P}_{\beta}(x) = \lim_{N \to \infty} \operatorname{diag}(\mathbf{1}) \cdot \mathbb{E} \left\langle N^{-1} \sigma \sigma^{\mathsf{T}} \right\rangle$$

where the Gibbs measure $\langle \cdot \rangle$ is equal to the one in (1.6) at t = 0 and this x. From (1.2), we can deduce that $\operatorname{diag}(\mathbf{1}) \cdot (N^{-1}\sigma\sigma^{\intercal}) = 1$. Inserting this into the above display gives the second part of this lemma.

2.2. Limit of free energy. Notice that the convex conjugate of $|\cdot|^2$ is $\frac{1}{4}|\cdot|^2$. The following formula in (2.4) is known as the Hopf–Lax formula.

Proposition 2.3 (Hopf–Lax formula for the limit). As $N \to \infty$, $F_{\beta,N}$ converges pointwise on $\mathbb{R}_+ \times \mathbb{R}^D$ to a Lipschitz and convex function f_β given by

(2.4)
$$f_{\beta}(t,x) = \sup_{y \in \mathbb{R}^D} \left\{ \mathscr{P}_{\beta}(x+y) - \frac{|y|^2}{4t} \right\}, \quad \forall (t,x) \in \mathbb{R}_+ \times \mathbb{R}^D.$$

Moreover, for every $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^D$, maximizers of the right-hand side in (2.4) exist.

Proof. This proposition can be deduced following the proof of [14, Proposition 5.2]. The difference is that, there, the domain for the second variable is the space of real symmetric matrices, while the domain here is \mathbb{R}^D . The simplification here is allowed by the fact that the self-overlap $N^{-1}\sigma\sigma^{\dagger}$ is diagonal as in (1.2). The convexity and Lipschitzness of f_{β} are due to the fact that $F_{N,\beta}$ is convex and Lipschitz uniformly in N, which can be deduced as in [14, Lemma 5.1]. Since \mathscr{P}_{β} is Lipschitz (see Lemma 2.2) and thus grows at most linearly, the existence of maximizers follows.

We need the following lemma to characterize the differentiability of $f_{\beta}(t, \cdot)$.

Lemma 2.4 (Envelop theorem). Let $x \in \mathbb{R}^D$. Then, $f_\beta(t, \cdot)$ is differentiable at x if and only if the set

$$\{\nabla \mathscr{P}_{\beta}(x+y) \mid y \text{ maximizes the right-hand side of } (2.4)\}$$

is a singleton. In this case, $\nabla f_{\beta}(t, x) = \nabla \mathscr{P}_{\beta}(x+y)$ for every maximizer y.

Proof. This is a result of the envelope theorem, for which we refer to the version stated as [23, Theorem 2.21]. The linear growth of \mathscr{P}_{β} as implied by Lemma 2.2 ensures the conditions in that theorem are satisfied.

Next, we introduce some notations relevant to the color symmetry of Potts spins. Let $\operatorname{Sym}(D)$ be the group consisting of permutations on $\{1, \ldots, D\}$ (labels of colors). For every $\mathbf{s} \in \operatorname{Sym}(D)$ and $x = (x_1, \ldots, x_D) \in \mathbb{R}^D$, we write

(2.5)
$$x^{\mathbf{s}} = (x_{\mathbf{s}(1)}, \dots, x_{\mathbf{s}(D)}).$$

In this notation, any $\mathbf{s} \in \text{Sym}(D)$ acts on the spin configuration σ through

(2.6)
$$\sigma^{\mathbf{s}} = (\sigma^{\mathbf{s}}_{\bullet i})_{1 \leqslant i \leqslant N}$$

where we apply the same permutation \mathbf{s} to every single spin (column vector).

Since the uniform distribution of Potts spins is invariant under the action as in (2.6), a simple observation from (1.5), (2.1), and (2.4) is that

(2.7)
$$F_{\beta,N}(t,x) = F_{\beta,N}(t,x^{\mathbf{s}}), \qquad \mathscr{P}_{\beta}(x) = \mathscr{P}_{\beta}(x^{\mathbf{s}}), \qquad f_{\beta}(t,x) = f_{\beta}(t,x^{\mathbf{s}})$$

for every $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^D$ and $\mathbf{s} \in \text{Sym}(D)$. As a result, we also have

(2.8)
$$x \in \mathbb{R}\mathbf{1} \implies \nabla \mathscr{P}_{\beta}(x) = D^{-1}\mathbf{1}.$$

Here and henceforth, $\mathbb{R}\mathbf{1} = \{r\mathbf{1} : r \in \mathbb{R}\}$. Indeed, from (2.7), we get $\nabla \mathscr{P}_{\beta}(x^{\mathbf{s}}) = (\nabla \mathscr{P}_{\beta}(x))^{\mathbf{s}}$ for every $x \in \mathbb{R}^{D}$. Then, if $x \in \mathbb{R}\mathbf{1}$, we have $x^{\mathbf{s}} = x$ and thus $\nabla \mathscr{P}_{\beta}(x) = (\nabla \mathscr{P}_{\beta}(x))^{\mathbf{s}}$ for every $\mathbf{s} \in \operatorname{Sym}(D)$. This along with $\mathbf{1} \cdot \nabla \mathscr{P}(x) = 1$ in Lemma 2.2 verifies (2.8).

We are mostly interested in the zero external field case. From Proposition 2.3, we have, for every $t \in \mathbb{R}_+$,

(2.9)
$$\lim_{N \to \infty} F_{\beta,N}(t,0) = f_{\beta}(t,0) = \sup_{y \in \mathbb{R}^D} \left\{ \mathscr{P}_{\beta}(y) - \frac{|y|^2}{4t} \right\}$$

The only subset of \mathbb{R}^D that is invariant under the action of Sym(D) as in (2.5) is $\mathbb{R}\mathbf{1}$. We are particularly interested in the case where the supremum in (2.9) is achieved on $\mathbb{R}\mathbf{1}$.

Lemma 2.5. For every $t \in \mathbb{R}_+$, we have

(2.10)
$$f_{\beta}(t,0) \ge \sup_{y \in \mathbb{R}^1} \left\{ \mathscr{P}_{\beta}(y) - \frac{|y|^2}{4t} \right\} = \mathscr{P}_{\beta}(0) + \frac{t}{D}$$

where the supremum has a unique maximizer $y = \frac{2t}{D}\mathbf{1}$.

Proof. The inequality follows from (2.4). Since $\mathbf{1} \cdot m_N = 1$ (see (1.2)), using the definition of $F_{\beta,N}$ in (1.5) and (2.1), we have

(2.11)
$$F_{\beta,N}(t, x + r\mathbf{1}) = F_{\beta,N}(t, x) + r, \quad \forall x \in \mathbb{R}^D, r \in \mathbb{R};$$
$$\mathscr{P}_{\beta}(x + r\mathbf{1}) = \mathscr{P}_{\beta}(x) + r, \quad \forall x \in \mathbb{R}^D, r \in \mathbb{R}.$$

Thus, the equality in (2.10) follows from (2.11) at x = 0 and simple computation.

2.3. Critical point. We define

(2.12)
$$t_{\rm c} = \inf\left\{t > 0: f_{\beta}(t,0) > \mathscr{P}_{\beta}(0) + \frac{t}{D}\right\}$$

This is the critical value of t that marks the onset of color symmetry breaking and the transition.

Proposition 2.6 (Properties of t_c). The following holds:

- (1) We have $0 < t_c < \infty$.
- (2) For every $t < t_c$, we have that $f_{\beta}(t, \cdot)$ is differentiable at 0 (with $\nabla f_{\beta}(t, 0) = D^{-1}\mathbf{1}$) and that the Hopf-Lax formula of $f_{\beta}(t, 0)$ as in (2.4) has a unique maximizer $\frac{2t}{D}\mathbf{1}$.
- (3) For every $t > t_c$, we have that $f_{\beta}(t,0) > \mathscr{P}_{\beta}(0) + \frac{t}{D}$; that $f_{\beta}(t,\cdot)$ is not differentiable at 0; and that the Hopf-Lax formula of $f_{\beta}(t,0)$ has more than one maximizer, none of which is equal to $\frac{2t}{D}\mathbf{1}$.

From this proposition, we can obtain different equivalent definitions of t_c : as the supremum of t such that $f_{\beta}(t,0) = \mathscr{P}(0) + \frac{t}{D}$; or that $f_{\beta}(t,\cdot)$ is differentiable at 0; or that the Hopf-Lax formula has a unique maximizer.

In the remainder of this subsection, we prove this proposition part by part.

Proof of $t_c > 0$ in Proposition 2.6 (1). Since \mathscr{P}_{β} is semi-concave (see Lemma 2.2), we have that $y \mapsto \mathscr{P}_{\beta}(y) - \frac{|y|^2}{4t}$ is strictly concave for sufficiently small t. Hence, the maximizer in (2.4) must be unique, which we denote by y_0 . Due to the symmetry of \mathscr{P}_{β} as in (2.7), we can see that $y_0^{\mathbf{s}}$ (defined as in (2.5)) is also a maximizer for any $\mathbf{s} \in \text{Sym}(D)$. Hence, the uniqueness of maximizers implies $y_0 \in \mathbb{R}\mathbf{1}$. Then, Lemma 2.5 gives $f_{\beta}(t,0) = \mathscr{P}_{\beta}(0) + \frac{t}{D}$ for all sufficiently small t > 0, which ensures $t_c > 0$.

Proof Proposition 2.6 (2). Let $t < t_c$. By definition, we have that $f_{\beta}(t,0) = \mathscr{P}_{\beta}(0) + \frac{t}{D}$. We set $y_0 = \frac{2t}{D}\mathbf{1}$ and we know from Lemma 2.5 that y_0 is a maximizer of the Hopf–Lax formula (2.4). Due to (2.8), we have $\nabla \mathscr{P}_{\beta}(y_0) = D^{-1}\mathbf{1}$. Now, we argue by contradiction and assume that $f_{\beta}(t, \cdot)$ is not differentiable at 0. By the envelope theorem as stated in Lemma 2.4, there must be a maximizer y of (2.4) such that

(2.13)
$$\nabla \mathscr{P}_{\beta}(y) \neq D^{-1}\mathbf{1}.$$

In particular, this along with (2.8) implies $y \notin \mathbb{R}\mathbf{1}$. Since y is a maximizer and $t < t_c$, we have

(2.14)
$$\mathscr{P}_{\beta}(y) - \frac{|y|^2}{4t} = \mathscr{P}_{\beta}(0) + \frac{t}{D}.$$

In the following we define functions $g, g_0 : \mathbb{R}_{++} \to \mathbb{R}$ by

$$g(s) = \mathscr{P}_{\beta}(y) - \frac{|y|^2}{4s}, \qquad g_0(s) = \mathscr{P}_{\beta}(0) + \frac{s}{D}, \qquad \forall s \in \mathbb{R}_{++}.$$

We then analyze the behavior of these two functions as s varies.

Since y is a maximizer of (2.4), we get

(2.15)
$$\nabla \mathscr{P}_{\beta}(y) = \frac{y}{2t}$$

As $\mathbf{1} \cdot \nabla \mathscr{P}_{\beta}(y) = 1$ due to Lemma 2.2, we have $\sum_{i=1}^{D} \frac{y_i}{2t} = 1$, which along with Jensen's inequality implies

(2.16)
$$\frac{|y|^2}{4t^2} \ge \frac{1}{D}$$

where the equality is achieved if and only if $\frac{y}{2t} = \frac{1}{D}\mathbf{1}$. This equality does not hold in this case due to (2.13) and (2.15). Hence, the inequality in (2.16) is strict, which implies $g'(t) > g'_0(t)$. Meanwhile, (2.14) gives $g(t) = g_0(t)$. Therefore, for all sufficiently small $\varepsilon > 0$, we have $g(t + \varepsilon) > g_0(t + \varepsilon)$, which along with the Hopf–Lax formula (2.4) yields

$$f_{\beta}(t+\varepsilon,0) > \mathscr{P}_{\beta}(0) + \frac{t+\varepsilon}{D}.$$

We can choose ε to satisfy $t + \varepsilon < t_c$, which then reaches a contradiction with the definition of t_c . Hence, $f_{\beta}(t, \cdot)$ must be differentiable at 0 by the argument of contradiction.

Let y be any maximizer of the Hopf–Lax formula. The envelope theorem stated as in Lemma 2.4 gives $\nabla f_{\beta}(t,0) = \nabla \mathscr{P}_{\beta}(y)$. By (2.7), we also have $\nabla f_{\beta}(t,0) = (\nabla f_{\beta}(t,0))^{\mathbf{s}}$ for every $\mathbf{s} \in \text{Sym}(D)$, which along with $\mathbf{1} \cdot \nabla f_{\beta}(t,0) = \mathbf{1} \cdot \nabla \mathscr{P}_{\beta}(y) = 1$ (due to Lemma 2.2) implies $\nabla f_{\beta}(t,0) = D^{-1}\mathbf{1}$. Hence, this along with (2.15) yields $\frac{y}{2t} = \nabla f_{\beta}(t,0) = D^{-1}\mathbf{1}$, which completes proof.

Proof Proposition 2.6 (3). Notice that $s \mapsto f_{\beta}(s,0)$ is convex (see (2.4)), $f_{\beta}(0,0) = \mathscr{P}_{\beta}(0)$, and $f_{\beta}(s,0) \geq \mathscr{P}_{\beta}(0) + \frac{s}{D}$ for every s. Hence, if $f_{\beta}(t,0) = \mathscr{P}_{\beta}(0) + \frac{t}{D}$ at some $t > t_{\rm c}$, then we have $f_{\beta}(s,0) = \mathscr{P}_{\beta}(0) + \frac{s}{D}$ for every $s \in [t_{\rm c},t]$, which contradicts the definition of $t_{\rm c}$. Therefore, we have $f_{\beta}(t,0) > \mathscr{P}_{\beta}(0) + \frac{t}{D}$ for every $t > t_{\rm c}$.

For the second statement, suppose that $f_{\beta}(t, \cdot)$ is differentiable at 0 at some $t > t_c$. Let y be any maximizer of the Hopf–Lax formula (2.4). Then, from the last paragraph in the proof of Part (2), we have $y = 2tD^{-1}\mathbf{1}$. This implies that the Hopf–Lax formula (2.4) of $f_{\beta}(t, \cdot)$ maximizes over $\mathbb{R}\mathbf{1}$ which by Lemma 2.5 gives $f_{\beta}(t, 0) = \mathscr{P}_{\beta}(0) + \frac{t}{D}$. This contradicts the result in the previous paragraph. Hence, $f_{\beta}(t, \cdot)$ cannot be differentiable at 0 for any $t > t_c$.

Lastly, the Hopf–Lax formula must have more than one maximizer because otherwise, we can use the envelope theorem as stated in Lemma 2.4 to deduce that $f_{\beta}(t, \cdot)$ is differentiable at 0. Also, none of the maximizers can be $\frac{2t}{D}\mathbf{1}$, because otherwise we can use Lemma 2.5 to deduce $f_{\beta}(t,0) = \mathscr{P}_{\beta}(0) + \frac{t}{D}$, which contradicts the first part of the statement.

To show that $t_c < \infty$, we need some preparation. Denote by $F_N^{CW}(t, x)$ be the free energy in (1.5) with β set to be zero. This is the free energy in the Curie–Weiss– Potts model. We denote by $\psi = F_1^{CW}(0, \cdot)$ the initial condition. In this case, we have $\psi = F_N^{CW}(0, \cdot)$ for every $N \in \mathbb{N}$ and the explicit expression

(2.17)
$$\psi(x) = \log\left(\sum_{d=1}^{D} e^{x_d}\right) - \log D, \quad \forall x \in \mathbb{R}^D.$$

Proposition 2.3 yields that, for every $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^D$,

(2.18)
$$\lim_{N \in \infty} F_N^{CW}(t, x) = \sup_{y \in \mathbb{R}^D} \left\{ \psi(x+y) - \frac{|y|^2}{4t} \right\}.$$

The difference between the free energy of the Potts spin glass and that of Curie–Weiss–Potts is also bounded.

Lemma 2.7. We have

$$F_N^{\text{CW}}(t,x) - \frac{\beta^2}{2} \leqslant F_{\beta,N}(t,x) \leqslant F_N^{\text{CW}}(t,x)$$

uniformly in $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^D$ and $N \in \mathbb{N}$.

Proof. For $s \in [0, 1]$, we define $\phi(s) = F_{\beta\sqrt{s},N}(t, x)$. Then, we can compute the derivative $\phi'(s)$ of $\phi(s)$. By using the Gaussian integration by parts and the boundedness of Potts spins, we can show $-\frac{\beta^2}{2} \leq \phi'(s) \leq 0$. Since $\phi(1) = F_{\beta,N}(t, x)$ and $\phi(0) = F_N^{\text{CW}}(t, x)$, we can get the desired result.

Proof of $t_c < \infty$ in Proposition 2.6 (1). For any t > 0, set $y(t) = (2t, 0, \dots, 0) \in \mathbb{R}^D$. Using (2.17), we can compute

(2.19)
$$\psi(y(t)) - \frac{|y(t)|^2}{4t} = \log(e^{2t} + D - 1) - \log D - t \ge t - \log D$$

Then, we have

$$f_{\beta}(t,0) = \lim_{N} F_{\beta,N}(t,0) \overset{\text{L.2.7}}{\geqslant} \lim_{N} F_{N}^{\text{CW}}(t,0) - \frac{\beta^{2}}{2} \overset{(2.18),(2.19)}{\geqslant} t - \log D - \frac{\beta^{2}}{2}$$

which implies that, for all t sufficiently large,

$$f_{\beta}(t,0) > \mathscr{P}_{\beta}(0) + \frac{t}{D}$$

and thus $t_c < \infty$ by its definition in (2.12).

2.4. Color symmetry breaking. We explain our definition of color symmetry in terms of the spin glass order parameter and show that t_c in (2.12) marks the color symmetry breaking.

We start with describing the Parisi formula. Recall that \mathscr{P}_{β} (as in (2.1)) is the limit of free energy in the model with self-overlap correction. The existence of the limit is ensured by [14, Theorem 1.1], which states that, for each $x \in \mathbb{R}^D$, $\mathscr{P}_{\beta}(x)$ is given by the Parisi formula

(2.20)
$$\mathscr{P}_{\beta}(x) = \inf_{\pi \in \Pi} \mathscr{P}_{\beta}(\pi, x).$$

We explain the formula on the right-hand side. Denote by \mathbf{S}^{D}_{+} the set of $D \times D$ real symmetric positive semi-definite matrices. Here, Π is the collection of all left-continuous paths $\pi : [0, 1) \to \mathbf{S}^{D}_{+}$ that are increasing in the sense that $\pi(s') - \pi(s) \in \mathbf{S}^{D}_{+}$ whenever $s' \geq s$. The Parisi functional $\mathscr{P}_{\beta}(\pi, x)$ is defined explicitly as in [14, (1.5)] (see also [17, (1.7)]).

Since any path $\pi \in \Pi$ is monotone, π has right limits and we define the right endpoint of π as $\pi(1) = \lim_{s \uparrow 1} \pi(s)$. For each $z \in \mathbb{R}^D$, we consider the collection $\Pi(z) = \{\pi \in \Pi : \pi(1) = \operatorname{diag}(z)\}$ of paths whose endpoints are equal to $\operatorname{diag}(z)$. Using (2.2) in Remark 2.1, we can restate [17, Theorem 1.1 (3)] as that, for every $x \in \mathbb{R}^D$,

$$\mathscr{P}_{\beta}(x) = \inf_{\pi \in \Pi(\nabla \mathscr{P}_{\beta}(x))} \mathscr{P}_{\beta}(\pi, x)$$

This refines the formula in the previous display by telling us that the Parisi formula optimizes over paths with a fixed endpoint specified by $\nabla \mathscr{P}_{\beta}(x)$. Using this, we can rewrite the Hopf–Lax formula in (2.4) as, for $t \in \mathbb{R}_+$,

(2.21)
$$f_{\beta}(t,0) = \sup_{y \in \mathbb{R}^{D}} \inf_{\pi \in \Pi\left(\nabla \mathscr{P}_{\beta}(y)\right)} \left\{ \mathscr{P}_{\beta}(\pi,y) - \frac{|y|^{2}}{4t} \right\}.$$

This will be the formula to be work this.

We set \mathcal{Z} to be the collection of increasing left-continuous paths $\zeta : [0,1] \to [0,1]$ such that $\zeta(1) = \lim_{s \uparrow 1} \zeta(s) = 1$. We also define

(2.22)
$$\Psi(s) = \frac{s}{D} \mathbf{Id} + \frac{1-s}{D^2} \mathbf{11}^{\mathsf{T}} \quad \forall s \in [0,1].$$

These will be needed in the following discussion.

We explain the physical meaning of the parameters in (2.21). Let y and π be optimal. Heuristically, the asymptotic distribution of the overlap $N^{-1}\sigma\sigma'^{\mathsf{T}}$ under $\mathbb{E}\langle \cdot \rangle_{t,0}$ (where σ and σ' are independent samples from $\langle \cdot \rangle_{t,0}$) should correspond to the law of $\pi(U)$ where U is the uniform random variable over [0, 1]. The limit of the self-overlap $N^{-1}\sigma\sigma^{\mathsf{T}}$ should correspond to the endpoint $\pi(1)$ which is equal to diag $(\nabla \mathscr{P}_{\beta}(y))$. We heuristically view this correspondence as an equality, which is not correct (see the footnote in Section 1.2). The color symmetry should mean that the optimal π is invariant under the action induced by permuting labels of the spin (see (2.6)), more precisely,

$$(\pi_{dd'}(s))_{1 \leqslant d, d' \leqslant D} = \left(\pi_{\mathbf{s}(d)\mathbf{s}(d')}(s)\right)_{1 \leqslant d, d' \leqslant D}, \quad \forall s \in [0, 1), \quad \forall \mathbf{s} \in \operatorname{Sym}(D).$$

On the other hand, since the entries of $N^{-1}\sigma\sigma'^{\mathsf{T}}$ always sum to 1, we also expect $\sum_{d,d'=1}^{D} \pi_{dd'} = 1$. Hence, assuming that π takes the form in the above display, we can deduce that π must be of the form $\pi = \Psi \circ \zeta$ for some $\zeta \in \mathcal{Z}$. In particular, the endpoint satisfies $\pi(1) = D^{-1}\mathbf{Id}$ and thus we necessarily have $\nabla \mathscr{P}_{\beta}(y) = D^{-1}\mathbf{1}$. This paragraph is not rigorous, but it motivates the following definition.

Definition 2.8. At $t \in \mathbb{R}_+$, we say that the **color symmetry is preserved** provided that there is a maximizer $y \in \mathbb{R}^D$ of the supremum in (2.21) such that $\nabla \mathscr{P}_{\beta}(y) = D^{-1}\mathbf{1}$ and the remaining infimum in (2.21) optimizes over symmetric paths, namely,

$$f_{\beta}(t,0) = \inf_{\zeta \in \mathcal{Z}} \mathscr{P}_{\beta}(\Psi \circ \zeta, y) - \frac{|y|^2}{4t}$$

where Ψ is given in (2.22) and \mathcal{Z} is defined above (2.22). Otherwise, we say that the color symmetry is broken.

Proposition 2.9 (Color symmetry). *The following holds:*

(1) If $t \in [0, t_c]$, then the color symmetry is preserved and we have

(2.23)
$$f_{\beta}(t,0) = \inf_{\zeta \in \mathcal{Z}} \mathscr{P}_{\beta}(\Psi \circ \zeta, 0) + \frac{t}{D}$$

(2) If $t > t_c$, then the color symmetry is broken and we have

$$f_{\beta}(t,0) > \inf_{\zeta \in \mathcal{Z}} \mathscr{P}_{\beta}(\Psi \circ \zeta, 0) + \frac{t}{D}$$

We remark that [16, Theorem 1.2] gives a rewriting of the infimum in (2.23) which admits a unique minimizer. This result is based on an extension of the arguments in [2] (also see [13]).

Proof. To show the first part, let $t \in [0, t_c]$. By the definition of t_c in (2.12) and continuity (needed for $t = t_c$), we have $f_{\beta}(t, 0) = \mathscr{P}_{\beta}(0) + \frac{t}{D}$. Lemma 2.5 implies that $y = \frac{2t}{D}\mathbf{1}$ is a maximizer of the Hopf–Lax formula in (2.4) and thus also of the supremum in (2.21). By (2.8), we have $\nabla \mathscr{P}_{\beta}(y) = D^{-1}\mathbf{1}$. By [16, Theorem 1.1], we have

(2.24)
$$\mathscr{P}_{\beta}(0) = \inf_{\zeta \in \mathcal{Z}} \mathscr{P}_{\beta}(\Psi \circ \zeta, 0)$$

and thus

$$f_{\beta}(t,0) = \inf_{\zeta \in \mathcal{Z}} \mathscr{P}_{\beta}(\Psi \circ \zeta, 0) + \frac{t}{D} \stackrel{(2.11)}{=} \inf_{\zeta \in \mathcal{Z}} \mathscr{P}_{\beta}(\Psi \circ \zeta, y) - \frac{|y|^2}{4t}$$

which proves the first part.

Now, let $t > t_c$. Suppose that the color symmetry is preserved. Then, there is a maximizer y of the Hopf–Lax formula (2.4) such that $\nabla \mathscr{P}_{\beta}(y) = D^{-1}\mathbf{1}$. The maximality implies that $\nabla \mathscr{P}_{\beta}(y) = \frac{y}{2t}$ and thus $y = \frac{2t}{D}\mathbf{1}$, which contradicts Proposition 2.6 (3). Hence, the color symmetry is broken. Proposition 2.6 (3) also gives $f_{\beta}(t,0) > \mathscr{P}_{\beta}(0) + \frac{t}{D}$ which together with (2.24) gives the desired inequality.

2.5. Ferromagnetism. We describe the phase transition at $t_{\rm c}$ in terms of the change of magnetization.

Proposition 2.10 (Magnetization). The following holds.

(1) ("Zero" magnetization in the sub-critical regime) Let $t \in [0, t_c)$ and let $(x_N)_{N \in \mathbb{N}}$ be any sequence converging to 0. Then,

$$\lim_{N \to \infty} \mathbb{E} \left\langle \left| m_N - D^{-1} \mathbf{1} \right| \right\rangle_{t, x_N} = 0.$$

(2) (Spontaneous magnetization in the super-critical regime) Let $t > t_c$. Then, there is a strictly increasing sequence $(N_n)_{n \in \mathbb{N}}$ of positive integers and $(x_n)_{n \in \mathbb{N}}$ converging to 0 such that $\lim_{n\to\infty} \mathbb{E} \langle m_{N_n} \rangle_{t,x_n}$ exists but is not $D^{-1}\mathbf{1}$.

Proof of Proposition 2.10 (1). By computing the first-order derivative of $F_{\beta,N}(t,\cdot)$ using the expression in (1.5), we have

(2.25)
$$\nabla F_{\beta,N}(t,x) = \mathbb{E} \langle m_N \rangle_{t,x}, \quad \forall x \in \mathbb{R}^D.$$

In particular, this implies that $F_{\beta,N}(t,\cdot)$ is Lipschitz uniformly in N. We can use this to upgrade the pointwise convergence in Proposition 2.3 to that $F_{\beta,N}(t,\cdot)$ converges to $f_{\beta}(t,\cdot)$ uniformly on every compact set. By computing the second-order derivatives of $F_{\beta,N}(t,\cdot)$ (similar to [17, (2.4)]), we can verify that $F_{\beta,N}(t,\cdot)$ is convex. Then, for any $y \in \mathbb{R}^D$ and r > 0, the convexity gives

$$\frac{F_{\beta,N}(t,x_N) - F_{\beta,N}(t,x_N - ry)}{r} \leqslant y \cdot \nabla F_{\beta,N}(t,x_N) \leqslant \frac{F_{\beta,N}(t,x_N + ry) - F_{\beta,N}(t,x_N)}{r}.$$

First sending $N \to \infty$ and then $r \to 0$, we can use the local uniform convergence of $F_{\beta,N}$ and the differentiability of $f_{\beta}(t, \cdot)$ at 0 to get $\lim_{N\to\infty} y \cdot \nabla F_{\beta,N}(t, x_N) = y \cdot \nabla f_{\beta}(t, 0)$. Using this along with (1.5) and $\nabla f_{\beta}(t, 0) = D^{-1}\mathbf{1}$ (Proposition 2.6 (2)), we get

$$\lim_{N \to \infty} \mathbb{E} \left\langle m_N \right\rangle_{t, x_N} = D^{-1} \mathbf{1}.$$

The proof will be complete if we can show

$$\lim_{N \to \infty} \mathbb{E} \left\langle \left| m_N - \mathbb{E} \left\langle m_N \right\rangle_{t, x_N} \right| \right\rangle_{t, x_N} = 0$$

This is a standard consequence of the concentration of free energy, the convexity of $F_{\beta,N}(t,\cdot)$ (and its un-averaged version), and the differentiability of $f_{\beta}(t,\cdot)$ at 0 (due to $t < t_c$). For instance, one can use the same argument for [17, Proposition 2.4] verbatim by substituting m_N , x_N , $F_{\beta,N}(t,\cdot)$, $f_{\beta}(t,\cdot)$ for $N^{-1}\sigma\sigma^{\intercal}$, x, F_N , \mathscr{P} therein. The detail is tedious and thus omitted here.

Proof of Proposition 2.10 (2). Recall that $f_{\beta}(t, \cdot)$ is convex (see Proposition 2.3). If for every sequence $(x_n)_{n \in \mathbb{N}}$ of differentiable points of $f_{\beta}(t, \cdot)$ converging to 0 we have $\lim_{n\to\infty} \nabla f_{\beta}(t, x_n) = D^{-1}\mathbf{1}$, then we can use the characterization of differentials of convex function as stated in [47, Theorem 25.6] to deduce that $f_{\beta}(t, \cdot)$ is differentiable at 0. Since we know that $f_{\beta}(t, \cdot)$ is not differentiable at 0 due to Proposition 2.6 (3), there must be $(x_n)_{n\in\mathbb{N}}$ such that $(\nabla f_{\beta}(t, x_n))_{n\in\mathbb{N}}$ does not converge to $D^{-1}\mathbf{1}$. Since $f_{\beta}(t, \cdot)$ is Lipschitz (see Proposition 2.3), the derivatives of $f_{\beta}(t, \cdot)$ are bounded, which allows us to pass to a subsequence (still denoted as $(x_n)_{n\in\mathbb{N}}$ for convenience) along which $\nabla f_{\beta}(t, x_n)$ converges to some $a \in \mathbb{R}^D$ other than $D^{-1}\mathbf{1}$. As argued in the previous proof, we have $\lim_{N\to\infty} \nabla F_{\beta,N}(t, x_n) = \nabla f_{\beta}(t, x_n)$ for each n. Then, we can find a strictly increasing sequence $(N_n)_{n\in\mathbb{N}}$ such that $\nabla F_{\beta,N_n}(t, x_n)$ approximates $\nabla f_{\beta}(t, x_n)$ and thus $\lim_{n\to\infty} \nabla F_{\beta,N_n}(t, x_n) = a$. This along with (2.25) gives the result.

Acknowledgments. The author thanks Victor Issa and Jean-Christophe Mourrat for helpful discussions.

Funding. The author is funded by the Simons Foundation.

Data availability. No datasets were generated during this work.

Conflict of interests. The author has no conflicts of interest to declare.

Competing interests. The author has no competing interests to declare.

References

- E. Agliari, A. Barra, R. Burioni, and A. Di Biasio. Notes on the p-spin glass studied via Hamilton-Jacobi and smooth-cavity techniques. J. Math. Phys., 53(6):063304, 29, 2012.
- [2] A. Auffinger and W.-K. Chen. The Parisi formula has a unique minimizer. Comm. Math. Phys., 335(3):1429–1444, 2015.
- [3] C. L. Baldwin and B. Swingle. Revisiting the replica trick: Competition between spin glass and conventional order. *Journal of Statistical Physics*, 190(7):125, Jul 2023.
- [4] A. Barra, G. Del Ferraro, and D. Tantari. Mean field spin glasses treated with PDE techniques. Eur. Phys. J. B, 86(7):Art. 332, 10, 2013.
- [5] A. Barra, A. Di Biasio, and F. Guerra. Replica symmetry breaking in mean-field spin glasses through the Hamilton–Jacobi technique. *Journal of Statistical Mechanics: Theory and Experiment*, 2010(09):P09006, 2010.
- [6] A. Barra, G. Genovese, and F. Guerra. Equilibrium statistical mechanics of bipartite spin systems. J. Phys. A, 44(24):245002, 22, 2011.
- [7] E. Bates and Y. Sohn. Parisi formula for balanced Potts spin glass. arXiv preprint arXiv:2310.06745, 2023.
- [8] F. Caltagirone, G. Parisi, and T. Rizzo. Dynamical critical exponents for the mean-field Potts glass. *Physical Review E—Statistical, Nonlinear, and Soft Matter Physics*, 85(5):051504, 2012.

HONG-BIN CHEN

- [9] F. Camilli, P. Contucci, and E. Mingione. An inference problem in a mismatched setting: a spin-glass model with mattis interaction. *SciPost Physics*, 12(4):125, 2022.
- [10] P. Cannarsa and C. Sinestrari. Semiconcave functions, Hamilton-Jacobi equations, and optimal control, volume 58. Springer Science & Business Media, 2004.
- [11] H. Chen, J.-C. Mourrat, and J. Xia. Statistical inference of finite-rank tensors. Ann. H. Lebesgue, 5:1161–1189, 2022.
- [12] H.-B. Chen. Hamilton-Jacobi equations for nonsymmetric matrix inference. Ann. Appl. Probab., 32(4):2540-2567, 2022.
- [13] H.-B. Chen. Parisi PDE and convexity for vector spins. Preprint, arXiv:2311.10446, 2023.
- [14] H.-B. Chen. Self-overlap correction simplifies the Parisi formula for vector spins. *Electronic Journal of Probability*, 28:1–20, 2023.
- [15] H.-B. Chen. Free energy in spin glass models with conventional order. Journal of Statistical Physics, 191(4):49, 2024.
- [16] H.-B. Chen. On Parisi measures of Potts spin glasses with correction. *Electronic Communications in Probability*, 29(none):1 13, 2024.
- [17] H.-B. Chen. On the self-overlap in vector spin glasses. Journal of Mathematical Physics, 65(3), 2024.
- [18] H.-B. Chen and J.-C. Mourrat. On the free energy of vector spin glasses with non-convex interactions. arXiv preprint arXiv:2311.08980, 2023.
- [19] H.-B. Chen and J. Xia. Hamilton-Jacobi equations for inference of matrix tensor products. Ann. Inst. Henri Poincaré Probab. Stat., 58(2):755–793, 2022.
- [20] W.-K. Chen. On the mixed even-spin Sherrington-Kirkpatrick model with ferromagnetic interaction. Annales de l'IHP Probabilités et statistiques, 50(1):63-83, 2014.
- [21] M. Costeniuc, R. S. Ellis, and H. Touchette. Complete analysis of phase transitions and ensemble equivalence for the Curie–Weiss–Potts model. *Journal of Mathematical Physics*, 46(6), 2005.
- [22] E. De Santis, G. Parisi, and F. Ritort. On the static and dynamical transition in the mean-field potts glass. Journal of Physics A: Mathematical and General, 28(11):3025, 1995.
- [23] T. Dominguez and J.-C. Mourrat. Statistical mechanics of mean-field disordered systems: a Hamilton-Jacobi approach. Preprint, arXiv:2311.08976, 2023.
- [24] P. Eichelsbacher and B. Martschink. On rates of convergence in the curie-weiss-potts model with an external field. Annales de l'IHP Probabilités et statistiques, 51(1):252-282, 2015.
- [25] D. Elderfield and D. Sherrington. Novel non-ergodicity in the Potts spin glass. Journal of Physics C: Solid State Physics, 16(32):L1169, 1983.
- [26] D. Elderfield and D. Sherrington. Spin glass, ferromagnetic and mixed phases in the disordered Potts model. Journal of Physics C: Solid State Physics, 16(27):L971, 1983.
- [27] D. Elderfield and D. Sherrington. The curious case of the Potts spin glass. Journal of Physics C: Solid State Physics, 16(15):L497, 1983.
- [28] R. S. Ellis and K. Wang. Limit theorems for maximum likelihood estimators in the Curie-Weiss-Potts model. Stochastic processes and their applications, 40(2):251–288, 1992.
- [29] G. Genovese and A. Barra. A mechanical approach to mean field spin models. Journal of Mathematical Physics, 50(5):053303, 2009.
- [30] D. J. Gross, I. Kanter, and H. Sompolinsky. Mean-field theory of the Potts glass. *Physical review letters*, 55(3):304, 1985.
- [31] F. Guerra. Sum rules for the free energy in the mean field spin glass model. *Fields Institute Communications*, 30(11), 2001.
- [32] F. Guerra. Broken replica symmetry bounds in the mean field spin glass model. Comm. Math. Phys., 233(1):1–12, 2003.
- [33] V. Issa. Existence and uniqueness of permutation-invariant optimizers for parisi formula. arXiv preprint arXiv:2407.13846, 2024.
- [34] E. Lage and A. Erzan. Mixed phase for the infinite-range Potts spin glass. Journal of Physics C: Solid State Physics, 16(24):L873, 1983.
- [35] J. Lee. Energy landscape and metastability of Curie–Weiss–Potts model. Journal of Statistical Physics, 187(1):2, 2022.
- [36] P. Mottishaw. First-order spin glass transitions: an exact solution. *Europhysics Letters*, 1(8):409, 1986.
- [37] J.-C. Mourrat. Nonconvex interactions in mean-field spin glasses. Probab. Math. Phys., 2(2):281–339, 2021.
- [38] J.-C. Mourrat. The Parisi formula is a Hamilton-Jacobi equation in Wasserstein space. Canad. J. Math., 74(3):607–629, 2022.
- [39] J.-C. Mourrat. Free energy upper bound for mean-field vector spin glasses. Ann. Inst. Henri Poincaré Probab. Stat., 59(3):1143–1182, 2023.

- [40] J.-C. Mourrat. Color symmetry breaking in the Potts spin glass. arXiv preprint arXiv:2409.10437, 2024.
- [41] J.-C. Mourrat and D. Panchenko. Extending the Parisi formula along a Hamilton-Jacobi equation. Electronic Journal of Probability, 25:Paper No. 23, 17, 2020.
- [42] D. Panchenko. The Sherrington-Kirkpatrick model. Springer Monographs in Mathematics. Springer, New York, 2013.
- [43] D. Panchenko. Free energy in the Potts spin glass. Ann. Probab., 46(2):829–864, 2018.
- [44] D. Panchenko. Free energy in the mixed p-spin models with vector spins. Ann. Probab., 46(2):865–896, 2018.
- [45] G. Parisi. Infinite number of order parameters for spin-glasses. Phys. Rev. Lett., 43(23):1754, 1979.
- [46] G. Parisi. A sequence of approximated solutions to the SK model for spin glasses. J. Phys. A, 13(4):L115–L121, 1980.
- [47] R. T. Rockafellar. Convex Analysis, volume 36. Princeton university press, 1970.
- [48] D. Sherrington and S. Kirkpatrick. Solvable model of a spin-glass. Physical Review Letters, 35(26):1792, 1975.
- [49] M. Talagrand. The Parisi formula. Ann. of Math. (2), 163(1):221-263, 2006.

(Hong-Bin Chen) INSTITUT DES HAUTES ÉTUDES SCIENTIFIQUES, BURES-SUR-YVETTE, FRANCE *Email address*: hbchen@ihes.fr