# arXiv:2412.20664v1 [gr-qc] 30 Dec 2024

# The Advantage of Early Detection and Localization from Eccentricity-Induced Higher Harmonic Modes in Second-Generation Ground-Based Detector Networks

Tao Yang<sup>(b),1,\*</sup> Rong-Gen Cai<sup>(b),2,3,4,†</sup> Zhoujian Cao<sup>(b),5,4,‡</sup> and Hyung Mok Lee<sup>(b)6,§</sup>

<sup>1</sup>School of Physics and Technology, Wuhan University, Wuhan 430072, China

<sup>2</sup>CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,

Chinese Academy of Sciences, Beijing 100190, China

<sup>3</sup>School of Physical Science and Technology, Ningbo University, Ningbo, 315211, China

<sup>4</sup>School of Fundamental Physics and Mathematical Sciences, Hangzhou Institute for Advanced Study (HIAS),

University of Chinese Academy of Sciences, Hangzhou 310024, China

<sup>5</sup>Department of Astronomy, Beijing Normal University, Beijing 100875, China

<sup>6</sup>Center for the Gravitational-Wave Universe, Astronomy Research Center,

Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Korea

(Dated: December 31, 2024)

Early detection and localization of gravitational waves (GWs) are crucial for identifying and capturing their electromagnetic (EM) counterparts, playing a significant role in multi-messenger astronomy. For second-generation (2G) detectors such as LIGO, Virgo, and KAGRA, the typical duration of GW signals ranges from  $\mathcal{O}(0.1)$  seconds to several tens of seconds due to their optimal sensitivities only at higher frequencies. This limited duration is insufficient to provide adequate early warning and localization for potential GWs and their associated EM counterparts. In this paper, we investigate whether eccentricity-induced higher harmonic modes, which enter the detector band much earlier than the dominant mode, can aid early detection and localization of GW sources using the 2G ground-based detector network. We select two typical events, a GW170817-like BNS and a GW150914-like BBH, as illustrative examples. Our focus is on presenting the evolution of signal-to-noise ratios (SNRs) and localization accuracy as functions of time-to-merger, examining the time gained for achieving certain SNRs and localization accuracy with the help of higher modes in eccentric cases compared to the circular case. For a GW170817-like BNS, we find that the eccentric case with  $e_0 = 0.4$  at 10 Hz can achieve an SNR of 4 and the threshold SNR of 8 approximately 12 minutes and 5 minutes before merger, representing 4.5- and 1.5-minute improvements in timeto-merger compared to the circular case, respectively. Additionally, with  $e_0 = 0.4$ , it can achieve a localization of  $1000 (100) \text{ deg}^2$  at 5 (1) minutes before the merger, reflecting improvements of 2 minutes (15 seconds) compared to the circular case. For a typical GW150914-like BBH, due to the much shorter signal duration, the time-to-merger gained from higher modes for achieving the same SNR and localization is limited to  $\mathcal{O}(0.1) - \mathcal{O}(1)$  seconds. Our results demonstrate the usefulness of eccentricity-induced higher harmonic modes in improving early warning and localization of GW and EM counterparts, particularly for BNS systems.

# I. INTRODUCTION

Early detection and precise localization of gravitational waves (GWs) are pivotal in advancing multimessenger GW astronomy and enhancing their applications in cosmology, astrophysics, and fundamental physics [1–10]. In particular, low-latency or even negative-latency GW alerts, as well as preparations to search for electromagnetic (EM) counterparts, can significantly benefit from earlier pre-merger detection and localization of potential sources. Achieving pre-merger detection and localization relies heavily on the in-band duration of the signal and the configuration of the detector network. For the LIGO-Virgo-KAGRA (LVK) detectors operating at their current sensitivity levels, the inband time is quite limited, typically ranging from  $\mathcal{O}(0.1)$  to several tens of seconds, assuming an initial frequency starting at 30 Hz. This duration is insufficient to accommodate the slew times of many electromagnetic telescopes, which range from 30 to 60 seconds [11]. The localization of sources in the third Gravitational-Wave Transient Catalog (GWTC-3) of LVK spans from several tens to several thousand square degrees [12]. Therefore, achieving pre-merger detection and localization with the current LVK network remains very challenging.

The enhancement of early warning and source localization through additional features of gravitational waves has been explored from various perspectives, including the multipole higher modes arising from asymmetric mass ratios [11, 13, 14] and the higher harmonic modes induced by eccentricity [15–18]. Our previous studies [19–21] shows that the eccentricity of longinspiraling compact binaries can enhance distance estimation and source localization accuracy by several orders of magnitude when observed with space-based decihertz observatories. Intriguingly, our more recent study [22] demonstrates that by utilizing eccentricityinduced higher modes, the third-generation (3G) detec-

<sup>\*</sup> yangtao@whu.edu.cn

<sup>&</sup>lt;sup>†</sup> cairg@itp.ac.cn

<sup>&</sup>lt;sup>‡</sup> zjcao@bnu.edu.cn

<sup>&</sup>lt;sup>§</sup> hmlee@snu.ac.kr

tor network ET+2CE can detect and localize the typical sources even before the dominant mode enters the band, enabling the earliest possible warning and maximizing preparation time for observing EM counterparts. These advantages stem from two key factors: on the one hand, higher modes, with their higher frequencies compared to the dominant mode, can enter the detector band earlier; on the other hand, eccentricity can break the degeneracy among GW parameters, including angular parameters. In [22], we focused on the period before the dominant  $\ell = 2$  mode enters the band. Our findings revealed that the second-generation (2G) network, comprising advanced LIGO (Hanford and Livingston), advanced Virgo, KAGRA, and India-LIGO (HLVKI), was unable to achieve enough signal-to-noise ratios (SNRs)  $\rho \geq 8$  from the pure higher modes before the dominant mode enters the band. However, since the eccentricityinduced higher modes enter the band much earlier than the dominant mode, we believe a certain SNR can be accumulated before the dominant mode is in-band, which subsequently enhances the SNR and improves the localization of sources further after the dominant mode enters the band. This effect is particularly beneficial for the early warning and localization with 2G detectors, which have a very limited in-band signal duration.

In this paper, we aim to supplement our previous studies on the early detection and localization of eccentric GWs using 2G detector networks. Our focus lies on the evolution of SNR and localization accuracy as functions of time-to-merger, as well as the time gained for pre-merger detection and localization enabled by eccentricity-induced higher harmonic modes. By simulating typical compact binaries, we quantify the extent to which eccentric cases can achieve earlier accumulation of a given SNR and localization accuracy compared to non-eccentric cases.

This paper is organized as follows. In Section II, we introduce our methodology, including the waveform models and simulation setup. In Section III, we present our main results, comparing the SNR and localization as a function of time to merger between eccentric and non-eccentric cases. Finally, in Section IV, we provide our conclusions along with potential caveats and a discussion of our findings.

### **II. METHODOLOGY**

Orbital eccentricity is a crucial feature for distinguishing between different binary black hole (BBH) formation scenarios [23–26]. Over the last decade, various studies have explored eccentric gravitational waves (GWs) from multiple perspectives, including the template [27–36], detection [37–46], and parameter estimation [20, 47–52]. In this paper, as we need the frequency decomposition of higher harmonic modes and focus on the pre-merger SNR and localization, we follow our previous research [22] and adopt the non-spinning, inspiral-only, frequencydomain EccentricFD waveform approximant provided by the LALSUITE software package [53]. The waveform can be expressed as the sum of harmonics [28]:

$$\tilde{h}(f) = \sum_{\ell=1}^{10} \tilde{h}_{\ell}(f) , \qquad (1)$$

where  $\tilde{h}_{\ell}(f)$  represents the  $\ell$ -th harmonic, which has a frequency  $\ell$  times the orbital frequency F. The detailed definitions and expressions in  $\tilde{h}_{\ell}(f)$  can be found in [27, 28]. The eccentric waveforms are then generated using PYCBC [54]. When  $e_0 = 0$ , only the  $\ell = 2$  mode is present in the waveform described by Eq. 1, reducing it to the TaylorF2 waveform model. In contrast, for eccentric cases, additional subdominant higher modes also contribute to the waveform.

Since the higher modes (e.g.,  $\ell = 10$ ) can remain inband (assuming an initial frequency of 10 Hz) for approximately 20 hours before merger, we account for the Earth's rotational effect throughout this paper. Consequently, the time- (and hence frequency-) dependent beam pattern functions  $F_{+,\times}$  of the detector for different higher modes must be calculated separately. We modify LALSUITE to extract each harmonic's polarization  $h_{\ell+,\times}(f)$  individually. The positions and orientations of the arms for the 2G detector network HLVKI are detailed in Table I of [18]. The time-dependent  $F_{+,\times}(t)$  are constructed using a geocentric coordinate system. For the eccentric cases, we follow [27] to numerically solve the phase evolution of the eccentric orbits and obtain  $t(f_{ref})$ , with the reference frequency corresponding to that of the dominant  $\ell = 2$  mode. We then derive the frequencydependent  $F_{+,\times}(f_{\rm ref})$  based on the reference frequency and for each  $\ell$ -th mode, the corresponding  $F_{+,\times}(f)$  is given by  $F_{+\times}(2f/\ell)$ .

We select a GW170817-like binary neutron star (BNS) and a GW150914-like BBH for simulating typical compact binaries <sup>1</sup>. The component masses  $(m_1, m_2)$ , redshift (z), and luminosity distance  $(d_L)$  are set to match the median values of the actual events<sup>2</sup>. To account for the varying sky positions and orientations of the systems, we generate 1000 random sets of angular parameters  $(P_{\text{ang}})$ , including the inclination angle  $(\iota)$ , sky location  $(\theta, \phi)$ , polarization angle  $(\psi)$ , and longitude of the ascending nodes axis  $(\beta)$ , sampled from a uniform and isotropic distribution. Given the validated use of this eccentric waveform for initial eccentricities:  $e_0 = 0$ , 0.05, 0.1, 0.2, and 0.4 at the initial frequency of the 2G detectors,  $f_0 = 10$  Hz. In total, we simulate

<sup>&</sup>lt;sup>1</sup> In this paper, we do not present results for a typical neutron star-black hole (NSBH) binary, as it does not provide additional conclusions. For such unequal-mass systems, the multipolar higher modes become more prominent, and we leave the exploration of these systems for future research.

<sup>&</sup>lt;sup>2</sup> https://gwosc.org/eventapi/html/GWTC/

10,000 events. We present the averaged results across the 1000 samples of  $P_{\rm ang}$  for each typical binary at a given eccentricity.

Since we need to analyze 10,000 events at various specific times to calculate the corresponding SNR and perform parameter estimation for localization, it is impractical to use stochastic sampling algorithms such as the Markov Chain Monte Carlo (MCMC) method. We adopt the approach of [19] and utilize the Fisher matrix technique for gravitational waves [55] to estimate the uncertainties and covariances of the waveform parameters. The Fisher matrix is defined as

$$\Gamma_{ij} = (\partial_i h, \partial_j h), \qquad (2)$$

where  $\partial_i h = \partial h / \partial P_i$  and  $P_i$  represents a parameter in the waveform. The network Fisher matrix is computed as the sum of  $\Gamma_{ij}$  from each detector. The inner product is defined as

$$(a,b) = 4 \int_{f_{\min}}^{f_{\max}} \frac{\tilde{a}^*(f)\tilde{b}(f) + \tilde{b}^*(f)\tilde{a}(f)}{2S_n(f)} df, \qquad (3)$$

where  $f_{\min}$  and  $f_{\max}$  denote the low- and max-frequency cutoffs. We set  $f_{\min} = 10$  Hz as the initial frequency of the 2G detectors, while  $f_{\max}$  is determined based on the time to merger at which the calculation is performed. Note that each mode in Eq. (1) must be truncated at its respective  $f_{\max}$ . The noise power spectral densities  $S_n(f)$  are the designed sensitivities for advanced LIGO, Virgo, and KAGRA. Then the SNR is

$$\rho = (h, h) \,. \tag{4}$$

The network SNR is calculated as the quadrature sum of the SNRs from each detector. The sky localization error is given by [56]

$$\Delta \Omega = 2\pi |\sin(\theta)| \sqrt{C_{\theta\theta} C_{\phi\phi} - C_{\theta\phi}^2}, \qquad (5)$$

with the covariance matrix of the parameters as  $C_{ij} = (\Gamma^{-1})_{ij}$ . We need to note the limitations of the Fisher matrix in the parameter estimation of GWs, especially for low SNRs [57]. We incorporate Gaussian priors  $\Gamma_{ii}^p = 1/(\delta P_i)^2$  into the Fisher information matrix, where  $\delta P_i$  represents the maximum allowable variation in the parameter [47, 55, 58, 59]. The Fisher matrix approach, with the inclusion of Gaussian priors, has been shown to be consistent with the more computationally intensive MCMC method [58, 59].

We select the reference frequencies  $f_2 = f(\ell = 2)$  to correspond to the time-to-merger  $t_c - t$ . The calculations of SNR and localization are performed at several tens of  $f_2$  points with adaptive step sizes to ensure a sufficiently smooth evolution. Our focus is on quantifying how much earlier the eccentric cases can, on average, accumulate a specific SNR and localization compared to the noneccentric cases.

### III. RESULTS

### A. SNR versus time-to-merger

We begin calculating the accumulated SNR once the higher mode  $\ell = 10$  enters the detector band at 10 Hz. This allows us to evaluate how much SNR is accumulated before the dominant mode enters the band and to observe the enhancement of the accumulated SNR contributed by higher modes after the dominant mode becomes in-band. The results for the evolution of average SNR as a function of time-to-merger are shown in Fig. 1. The left panel shows the results for GW170817-like BNS cases with various eccentricities, while the right panel illustrates the results for GW150914-like BBH cases. The stars indicate the time when the dominant  $\ell = 2$  mode enters the band at 10 Hz. As larger eccentricities shorten the inspiral time, the time-to-merger when the dominant mode becomes in-band is smaller for cases with higher eccentricities.

To quantify the time-to-merger gained by eccentricityinduced higher modes compared to non-eccentric cases, we set two reference SNR values:  $\rho = 4$ , corresponding to the lowest SNR in the LVK GW catalogs, and  $\rho = 8$ , the conventional threshold for confirming a GW event, as indicated by the two horizontal lines. For the non-eccentric case of GW170817-like BNS, the SNR of 4 is achieved at a time-to-merger of approximately 7.5 minutes, while for the eccentric case with  $e_0 = 0.4$ , the time-to-merger improves to 12 minutes, representing a 4.5-minute improvement. For the threshold SNR of 8, the time-to-merger increases from 3.5 minutes to 5 minutes, vielding a 1.5minute improvement. However, for smaller eccentricities, the improvement is less significant. In the typical GW150914-like BBH case, where the inspiral time is relatively short, the eccentric case with  $e_0 = 0.4$  achieves an SNR of 4 approximately 1.5 seconds earlier than the non-eccentric case. For the threshold SNR of 8, the improvement is only 0.5 seconds.

We observe that the eccentric cases begin to accumulate SNR much earlier than the non-eccentric cases, although we only present the SNR results for the period starting 20 minutes (for BNS) and 10 seconds (for BBH) before the merger. Additionally, with larger eccentricities, the SNR accumulates more rapidly due to the increased prominence of higher modes. An exception is observed for the cases with  $e_0 = 0.2$  and  $e_0 = 0.4$ , where the accumulated SNR for  $e_0 = 0.4$  is smaller than that for  $e_0 = 0.2$  at earlier times (for BBH cases). This can be attributed to two competing effects: while higher eccentricity excites stronger higher modes, the shortened inspiral time may reduce the SNR. In some cases,  $e_0 = 0.2$ represents the optimal scenario. A similar feature was also identified in our previous research.

We also observe that the SNR for all cases converges in the final stage before merger. This indicates that the dominant  $\ell = 2$  mode accumulates SNR very rapidly in the later stages, contributing the majority of the total SNR compared to the higher modes. A further explanation is as follows: the strength of the subdominant modes is only  $\mathcal{O}(10\%)$  of the dominant mode. Thus, for both eccentric and non-eccentric cases, the primary contribution to the SNR comes from the  $\ell = 2$  mode. Additionally, since the initial eccentricity is set at 10 Hz, the remaining eccentricity decreases significantly by the time the system evolves to frequencies where the noise power spectral density is optimal (recall the inverse relation between eccentricity and orbital frequency). As a result, the higher modes mainly contribute at earlier times when the noise is relatively large. Consequently, the SNR for eccentric cases does not show substantial improvement compared to the non-eccentric cases near the merger.

### B. Localization versus time-to-merger

Our previous research concluded that localization based solely on higher modes before the dominant  $\ell = 2$ mode enters the detector at 10 Hz is guite insufficient  $(> 10^3 \text{ deg}^2)$ . Therefore, we calculate the evolution of localization only after the  $\ell = 2$  mode has entered the band. The results are presented in Fig. 2, where we plot the evolution of localization only when its accuracy  $\Delta \Omega < 10^4 \, \text{deg}^2$ . For a typical BNS, achieving a localization of  $1000 \, \text{deg}^2$  requires 3 minutes before merger for the non-eccentric case, while the  $e_0 = 0.4$  eccentric case achieves the same localization at 5 minutes of time-tomerger, representing a 2-minute improvement. For a localization of  $\Delta \Omega = 100 \, \text{deg}^2$ , the eccentric case reaches this level 1 minute before merger, compared to about 45 seconds before merger for the non-eccentric case. In the typical BBH case, a localization of  $1000 \text{ deg}^2$  can be achieved 2 seconds before the merger for  $e_0 = 0.4$ , compared to 1.2 seconds before the merger for  $e_0 = 0$ . Due to the short inspiral time, the improvement in localization to  $100 \,\mathrm{deg}^2$  from eccentricity-induced higher modes is negligible, around  $\mathcal{O}(0.1)$  seconds.

We observe that a higher eccentricity results in better localization at early times, while all cases converge at late times just before the merger. This behavior is similar to what was explained for the SNR results. The final localization near merger time is on the order of  $\mathcal{O}(1) \deg^2$ for all BNS and BBH cases. For BNS cases, eccentricity does not yield an improvement in localization. However, for BBH cases, there is a slight improvement, as seen in the right panel of Fig. 2 (note that  $f_{\text{max}}$  is truncated at the innermost stable orbit, resulting in a flat evolution beyond that point). This conclusion aligns with the findings of [18], which examined the improvement in source localization due to eccentricity when the entire inspiral period is taken into account (it means around the merger time) for the 2G ground-based detector network.

### IV. CONCLUSION AND DISCUSSION

In this paper, we investigate the improvement in early detection and localization of compact binaries enabled by eccentricity-induced higher harmonic modes using the 2G detector network HLVKI. We conclude that in eccentric cases, the higher harmonic modes, which enter the detector band much earlier than the dominant mode, can enhance the SNR and localization at earlier times compared to non-eccentric cases.

For the evolution of accumulated SNR versus timeto-merger, we find that for a GW170817-like BNS, the eccentric case with  $e_0 = 0.4$  at 10 Hz can achieve  $\rho = 4$ and the threshold  $\rho = 8$  approximately 12 minutes and 5 minutes before merger, representing 4.5- and 1.5-minute improvements in time-to-merger compared to the circular case, respectively. In the case of a GW150914-like BBH, the improvement is only 1.5 and 0.5 seconds for achieving the same SNR. For the evolution of localization accuracy versus time-to-merger, an eccentric GW170817-like BNS with  $e_0 = 0.4$  at 10 Hz can achieve  $\Delta \Omega = 1000 (100) \text{ deg}^2$ 5(1) minutes before the merger, representing improvements of 2 minutes (15 seconds) compared to the circular case. In contrast, due to the very short period of inspiral, for a typical GW150914-like BBH, the improvement to circular case is less than 1 seconds for achieving the same localization. Our results demonstrate the advantages of early detection and localization enabled by eccentricityinduced higher modes for the 2G detector network.

The search for electromagnetic (EM) counterparts relies heavily on the early alert of possible GW events [60, 61], particularly for BNS systems, which are the most promising sources for accompanying various EM signals. For 2G detectors, the duration of detectable GW signals is still limited compared to 3G and space-based detectors. However, eccentricity-induced higher modes can provide earlier alerts for GW candidates, both in terms of detection and localization. This allows more preparation time for EM telescopes and maximizes the likelihood of capturing potential EM counterparts.

In the simulation of the typical GW events, We fixed their distances to match the median values of the true events reported by LVK. Our focus is on examining the newly proposed idea, and these typical events with their true distances suffice as examples for demonstration. However, our methodology can easily be extended to other distances, as there is a rough relationship in our calculation of SNR and localization:  $\rho \sim 1/d_L$  and  $\Delta\Omega \sim 1/\rho^2 \sim d_L^2$ .

The rate of eccentric binaries has been a topic of active research [62–65]. These estimates crucially depend on the unknown astrophysics of these classes of compact binaries. For BNS, [62] estimated that the rate of the eccentric case could be rare. However, we note that a single detection of such event would be highly significant. Given the extreme rarity of EM counterparts in current GW detections, an early warning of a potential GW + EM signal could be of great interest to both the



FIG. 1: The average SNR versus time-to-merger for the typical BNS (left) and BBH (right) event with 2G detector networks HLVKI. The stars indicate the time when the dominant  $\ell = 2$  mode enters the band at 10 Hz. The two dashed lines indicate the SNR of 4 and 8, respectively. We only present the SNR results for the period starting 20 minutes (for BNS) and 10 seconds (for BBH) before the merger.



FIG. 2: The average localization versus time-to-merger for the typical BNS (left) and BBH (right) event with 2G detector networks HLVKI. The two dashed lines indicate the localization of 1000 and 100 deg<sup>2</sup>, respectively. We only present the localization results for the period starting 15 minutes (for BNS) and 5 seconds (for BBH) before the merger.

GW community and EM follow-up teams.

# ACKNOWLEDGMENTS

This work is supported by "the Fundamental Research Funds for the Central Universities" under the reference No. 2042024FG0009. R.G.C is supported by the National Key Research and Development Program of China Grant No. 2020YFC2201502 and 2021YFA0718304 and by National Natural Science Foundation of China Grants No. 11821505, No. 11991052, No. 11947302, No. 12235019. Z.C is supported in part by the National Key Research and Development Program of China Grant No. 2021YFC2203001, in part by "the Fundamental Research Funds for the Central Universities". H.M.L is supported by the National Research Foundation of Korea 2021M3F7A1082056.

[1] B. P. Abbott *et al.* (LIGO Scientific, Virgo, Fermi GBM, INTEGRAL, IceCube, AstroSat Cadmium Zinc Telluride Imager Team, IPN, Insight-Hxmt, ANTARES,

Swift, AGILE Team, 1M2H Team, Dark Energy Camera GW-EM, DES, DLT40, GRAWITA, Fermi-LAT, ATCA, ASKAP, Las Cumbres Observatory Group, OzGrav, DWF (Deeper Wider Faster Program), AST3, CAAS-TRO, VINROUGE, MASTER, J-GEM, GROWTH, JAGWAR, CaltechNRAO, TTU-NRAO, NuSTAR, Pan-STARRS, MAXI Team, TZAC Consortium, KU, Nordic Optical Telescope, ePESSTO, GROND, Texas Tech University, SALT Group, TOROS, BOOTES, MWA, CALET, IKI-GW Follow-up, H.E.S.S., LOFAR, LWA, HAWC, Pierre Auger, ALMA, Euro VLBI Team, Pi of Sky, Chandra Team at McGill University, DFN, AT-LAS Telescopes, High Time Resolution Universe Survey, RIMAS, RATIR, SKA South Africa/MeerKAT), Multi-messenger Observations of a Binary Neutron Star Merger, Astrophys. J. Lett. 848, L12 (2017), arXiv:1710.05833 [astro-ph.HE].

- [2] B. P. Abbott *et al.* (LIGO Scientific, Virgo, Fermi-GBM, INTEGRAL), Gravitational Waves and Gammarays from a Binary Neutron Star Merger: GW170817 and GRB 170817A, Astrophys. J. Lett. 848, L13 (2017), arXiv:1710.05834 [astro-ph.HE].
- [3] B. P. Abbott *et al.* (LIGO Scientific, Virgo, 1M2H, Dark Energy Camera GW-E, DES, DLT40, Las Cumbres Observatory, VINROUGE, MASTER), A gravitationalwave standard siren measurement of the Hubble constant, Nature 551, 85 (2017), arXiv:1710.05835 [astroph.CO].
- [4] P. Creminelli and F. Vernizzi, Dark Energy after GW170817 and GRB170817A, Phys. Rev. Lett. 119, 251302 (2017), arXiv:1710.05877 [astro-ph.CO].
- [5] J. M. Ezquiaga and M. Zumalacárregui, Dark Energy After GW170817: Dead Ends and the Road Ahead, Phys. Rev. Lett. **119**, 251304 (2017), arXiv:1710.05901 [astroph.CO].
- [6] T. Baker, E. Bellini, P. G. Ferreira, M. Lagos, J. Noller, and I. Sawicki, Strong constraints on cosmological gravity from GW170817 and GRB 170817A, Phys. Rev. Lett. 119, 251301 (2017), arXiv:1710.06394 [astro-ph.CO].
- [7] K. P. Mooley, A. T. Deller, O. Gottlieb, E. Nakar, G. Hallinan, S. Bourke, D. A. Frail, A. Horesh, A. Corsi, and K. Hotokezaka, Superluminal motion of a relativistic jet in the neutron-star merger GW170817, Nature 561, 355 (2018), arXiv:1806.09693 [astro-ph.HE].
- [8] K. Hotokezaka, E. Nakar, O. Gottlieb, S. Nissanke, K. Masuda, G. Hallinan, K. P. Mooley, and A. T. Deller, A Hubble constant measurement from superluminal motion of the jet in GW170817, Nature Astron. 3, 940 (2019), arXiv:1806.10596 [astro-ph.CO].
- [9] T. Dietrich, M. W. Coughlin, P. T. H. Pang, M. Bulla, J. Heinzel, L. Issa, I. Tews, and S. Antier, Multimessenger constraints on the neutron-star equation of state and the Hubble constant, Science **370**, 1450 (2020), arXiv:2002.11355 [astro-ph.HE].
- [10] B. P. Abbott *et al.* (LIGO Scientific, Virgo), Lowlatency Gravitational-wave Alerts for Multimessenger Astronomy during the Second Advanced LIGO and Virgo Observing Run, Astrophys. J. 875, 161 (2019), arXiv:1901.03310 [astro-ph.HE].
- [11] S. J. Kapadia, M. K. Singh, M. A. Shaikh, D. Chatterjee, and P. Ajith, Of Harbingers and Higher Modes: Improved gravitational-wave early-warning of compact binary mergers, Astrophys. J. Lett. 898, L39 (2020), arXiv:2005.08830 [astro-ph.HE].

- [12] R. Abbott *et al.* (KAGRA, VIRGO, LIGO Scientific), GWTC-3: Compact Binary Coalescences Observed by LIGO and Virgo during the Second Part of the Third Observing Run, Phys. Rev. X 13, 041039 (2023), arXiv:2111.03606 [gr-qc].
- [13] M. K. Singh, S. J. Kapadia, M. A. Shaikh, D. Chatterjee, and P. Ajith, Improved early warning of compact binary mergers using higher modes of gravitational radiation: A population study, Mon. Not. Roy. Astron. Soc. 502, 1612 (2021), arXiv:2010.12407 [astro-ph.HE].
- [14] T. Islam, S. E. Field, C.-J. Haster, and R. Smith, High precision source characterization of intermediate massratio black hole coalescences with gravitational waves: The importance of higher order multipoles, Phys. Rev. D 104, 084068 (2021), arXiv:2105.04422 [gr-qc].
- [15] B. Mikoczi, B. Kocsis, P. Forgacs, and M. Vasuth, Parameter estimation for inspiraling eccentric compact binaries including pericenter precession, Phys. Rev. D 86, 104027 (2012), arXiv:1206.5786 [gr-qc].
- [16] K. Kyutoku and N. Seto, Pre-merger localization of eccentric compact binary coalescences with secondgeneration gravitational-wave detector networks, Mon. Not. Roy. Astron. Soc. 441, 1934 (2014), arXiv:1312.2953 [astro-ph.HE].
- [17] S. Ma, Z. Cao, C.-Y. Lin, H.-P. Pan, and H.-J. Yo, Gravitational wave source localization for eccentric binary coalesce with a ground-based detector network, Phys. Rev. D 96, 084046 (2017), arXiv:1710.02965 [gr-qc].
- [18] H.-P. Pan, C.-Y. Lin, Z. Cao, and H.-J. Yo, Accuracy of source localization for eccentric inspiraling binary mergers using a ground-based detector network, Phys. Rev. D 100, 124003 (2019), arXiv:1912.04455 [gr-qc].
- [19] T. Yang, R.-G. Cai, Z. Cao, and H. M. Lee, Eccentricity of Long Inspiraling Compact Binaries Sheds Light on Dark Sirens, Phys. Rev. Lett. **129**, 191102 (2022), arXiv:2202.08608 [gr-qc].
- [20] T. Yang, R.-G. Cai, Z. Cao, and H. M. Lee, Parameter estimation of eccentric gravitational waves with a decihertz observatory and its cosmological implications, Phys. Rev. D 107, 043539 (2023), arXiv:2212.11131 [gr-qc].
- [21] T. Yang, R.-G. Cai, and H. M. Lee, Space-borne atom interferometric gravitational wave detections. Part III. Eccentricity on dark sirens, JCAP 10, 061, arXiv:2208.10998 [gr-qc].
- [22] T. Yang, R.-G. Cai, Z. Cao, and H. M. Lee, Eccentricity enables the earliest warning and localization of gravitational waves with ground-based detectors, Phys. Rev. D 109, 104041 (2024), arXiv:2310.08160 [gr-qc].
- [23] A. Nishizawa, E. Berti, A. Klein, and A. Sesana, eLISA eccentricity measurements as tracers of binary black hole formation, Phys. Rev. D 94, 064020 (2016), arXiv:1605.01341 [gr-qc].
- [24] A. Nishizawa, A. Sesana, E. Berti, and A. Klein, Constraining stellar binary black hole formation scenarios with eLISA eccentricity measurements, Mon. Not. Roy. Astron. Soc. 465, 4375 (2017), arXiv:1606.09295 [astroph.HE].
- [25] K. Breivik, C. L. Rodriguez, S. L. Larson, V. Kalogera, and F. A. Rasio, Distinguishing Between Formation Channels for Binary Black Holes with LISA, Astrophys. J. Lett. 830, L18 (2016), arXiv:1606.09558 [astroph.GA].
- [26] M. Zevin, I. M. Romero-Shaw, K. Kremer, E. Thrane, and P. D. Lasky, Implications of Eccentric Observations

on Binary Black Hole Formation Channels, Astrophys. J. Lett. **921**, L43 (2021), arXiv:2106.09042 [astro-ph.HE].

- [27] N. Yunes, K. G. Arun, E. Berti, and C. M. Will, Post-Circular Expansion of Eccentric Binary Inspirals: Fourier-Domain Waveforms in the Stationary Phase Approximation, Phys. Rev. D 80, 084001 (2009), [Erratum: Phys.Rev.D 89, 109901 (2014)], arXiv:0906.0313 [gr-qc].
- [28] E. A. Huerta, P. Kumar, S. T. McWilliams, R. O'Shaughnessy, and N. Yunes, Accurate and efficient waveforms for compact binaries on eccentric orbits, Phys. Rev. D 90, 084016 (2014), arXiv:1408.3406 [gr-qc].
- [29] S. Tanay, M. Haney, and A. Gopakumar, Frequency and time domain inspiral templates for comparable mass compact binaries in eccentric orbits, Phys. Rev. D 93, 064031 (2016), arXiv:1602.03081 [gr-qc].
- [30] E. A. Huerta *et al.*, Complete waveform model for compact binaries on eccentric orbits, Phys. Rev. D 95, 024038 (2017), arXiv:1609.05933 [gr-qc].
- [31] Z. Cao and W.-B. Han, Waveform model for an eccentric binary black hole based on the effective-one-bodynumerical-relativity formalism, Phys. Rev. D 96, 044028 (2017), arXiv:1708.00166 [gr-qc].
- [32] E. A. Huerta *et al.*, Eccentric, nonspinning, inspiral, Gaussian-process merger approximant for the detection and characterization of eccentric binary black hole mergers, Phys. Rev. D 97, 024031 (2018), arXiv:1711.06276 [gr-qc].
- [33] B. Moore and N. Yunes, A 3PN Fourier Domain Waveform for Non-Spinning Binaries with Moderate Eccentricity, Class. Quant. Grav. 36, 185003 (2019), arXiv:1903.05203 [gr-qc].
- [34] A. Ramos-Buades, A. Buonanno, M. Khalil, and S. Ossokine, Effective-one-body multipolar waveforms for eccentric binary black holes with nonprecessing spins, Phys. Rev. D 105, 044035 (2022), arXiv:2112.06952 [gr-qc].
- [35] X. Liu, Z. Cao, and Z.-H. Zhu, Effective-one-body numerical-relativity waveform model for eccentric spinprecessing binary black hole coalescence, Class. Quant. Grav. 41, 195019 (2024), arXiv:2310.04552 [gr-qc].
- [36] J. N. Arredondo, A. Klein, and N. Yunes, Efficient gravitational-wave model for fully-precessing and moderately eccentric, compact binary inspirals, Phys. Rev. D 110, 044044 (2024), arXiv:2402.06804 [gr-qc].
- [37] M. E. Lower, E. Thrane, P. D. Lasky, and R. Smith, Measuring eccentricity in binary black hole inspirals with gravitational waves, Phys. Rev. D 98, 083028 (2018), arXiv:1806.05350 [astro-ph.HE].
- [38] B. P. Abbott *et al.* (LIGO Scientific, Virgo), Search for Eccentric Binary Black Hole Mergers with Advanced LIGO and Advanced Virgo during their First and Second Observing Runs, Astrophys. J. 883, 149 (2019), arXiv:1907.09384 [astro-ph.HE].
- [39] I. M. Romero-Shaw, P. D. Lasky, and E. Thrane, Searching for Eccentricity: Signatures of Dynamical Formation in the First Gravitational-Wave Transient Catalogue of LIGO and Virgo, Mon. Not. Roy. Astron. Soc. 490, 5210 (2019), arXiv:1909.05466 [astro-ph.HE].
- [40] A. H. Nitz, A. Lenon, and D. A. Brown, Search for Eccentric Binary Neutron Star Mergers in the first and second observing runs of Advanced LIGO, Astrophys. J. 890, 1 (2019), arXiv:1912.05464 [astro-ph.HE].
- [41] S. Wu, Z. Cao, and Z.-H. Zhu, Measuring the eccentricity of binary black holes in GWTC-1 by using the inspiral-

only waveform, Mon. Not. Roy. Astron. Soc. **495**, 466 (2020), arXiv:2002.05528 [astro-ph.IM].

- [42] I. M. Romero-Shaw, P. D. Lasky, E. Thrane, and J. C. Bustillo, GW190521: orbital eccentricity and signatures of dynamical formation in a binary black hole merger signal, Astrophys. J. Lett. **903**, L5 (2020), arXiv:2009.04771 [astro-ph.HE].
- [43] V. Gayathri, J. Healy, J. Lange, B. O'Brien, M. Szczepanczyk, I. Bartos, M. Campanelli, S. Klimenko, C. O. Lousto, and R. O'Shaughnessy, Eccentricity estimate for black hole mergers with numerical relativity simulations, Nature Astron. 6, 344 (2022), arXiv:2009.05461 [astroph.HE].
- [44] H. L. Iglesias *et al.*, Eccentricity Estimation for Five Binary Black Hole Mergers with Higher-order Gravitational-wave Modes, Astrophys. J. **972**, 65 (2024), arXiv:2208.01766 [gr-qc].
- [45] A. G. Abac *et al.* (LIGO Scientific, KAGRA, VIRGO), Search for Eccentric Black Hole Coalescences during the Third Observing Run of LIGO and Virgo, Astrophys. J. 973, 132 (2024), arXiv:2308.03822 [astro-ph.HE].
- [46] N. Gupte *et al.*, Evidence for eccentricity in the population of binary black holes observed by LIGO-Virgo-KAGRA, (2024), arXiv:2404.14286 [gr-qc].
- [47] M. Favata, Systematic parameter errors in inspiraling neutron star binaries, Phys. Rev. Lett. **112**, 101101 (2014), arXiv:1310.8288 [gr-qc].
- [48] B. Sun, Z. Cao, Y. Wang, and H.-C. Yeh, Parameter estimation of eccentric inspiraling compact binaries using an enhanced post circular model for ground-based detectors, Phys. Rev. D 92, 044034 (2015).
- [49] L. Gondán, B. Kocsis, P. Raffai, and Z. Frei, Accuracy of Estimating Highly Eccentric Binary Black Hole Parameters with Gravitational-Wave Detections, Astrophys. J. 855, 34 (2018), arXiv:1705.10781 [astro-ph.HE].
- [50] P. Saini, M. Favata, and K. G. Arun, Systematic bias on parametrized tests of general relativity due to neglect of orbital eccentricity, Phys. Rev. D 106, 084031 (2022), arXiv:2203.04634 [gr-qc].
- [51] P. Narayan, N. K. Johnson-McDaniel, and A. Gupta, Effect of ignoring eccentricity in testing general relativity with gravitational waves, Phys. Rev. D 108, 064003 (2023), arXiv:2306.04068 [gr-qc].
- [52] H. Gil Choi, T. Yang, and H. M. Lee, Importance of eccentricities in parameter estimation of compact binary inspirals with decihertz gravitational-wave detectors, Phys. Rev. D 110, 024025 (2024), arXiv:2210.09541 [gr-qc].
- [53] LIGO Scientific Collaboration, Virgo Collaboration, and KAGRA Collaboration, LVK Algorithm Library - LAL-Suite, Free software (GPL) (2018).
- [54] A. Nitz, I. Harry, D. Brown, C. M. Biwer, J. Willis, T. D. Canton, C. Capano, T. Dent, L. Pekowsky, A. R. Williamson, S. De, M. Cabero, B. Machenschalk, D. Macleod, P. Kumar, F. Pannarale, S. Reyes, G. S. C. Davies, dfinstad, S. Kumar, M. Tápai, L. Singer, S. Khan, S. Fairhurst, A. Nielsen, S. Singh, T. Massinger, K. Chandra, shasvath, and veronica villa, gwastro/pycbc: v2.0.5 release of pycbc (2022).
- [55] C. Cutler and E. E. Flanagan, Gravitational waves from merging compact binaries: How accurately can one extract the binary's parameters from the inspiral wave form?, Phys. Rev. D 49, 2658 (1994), arXiv:grqc/9402014.
- [56] C. Cutler, Angular resolution of the LISA gravitational

wave detector, Phys. Rev. D 57, 7089 (1998), arXiv:gr-qc/9703068.

- [57] M. Vallisneri, Use and abuse of the Fisher information matrix in the assessment of gravitationalwave parameter-estimation prospects, Phys. Rev. D 77, 042001 (2008), arXiv:gr-qc/0703086.
- [58] M. Favata, C. Kim, K. G. Arun, J. Kim, and H. W. Lee, Constraining the orbital eccentricity of inspiralling compact binary systems with Advanced LIGO, Phys. Rev. D 105, 023003 (2022), arXiv:2108.05861 [gr-qc].
- [59] H.-S. Cho, Systematic bias due to eccentricity in parameter estimation for merging binary neutron stars, Phys. Rev. D 105, 124022 (2022), arXiv:2205.12531 [gr-qc].
- [60] Q. Chu, E. J. Howell, A. Rowlinson, H. Gao, B. Zhang, S. J. Tingay, M. Boër, and L. Wen, Capturing the electromagnetic counterparts of binary neutron star mergers through low latency gravitational wave triggers, Mon. Not. Roy. Astron. Soc. 459, 121 (2016), arXiv:1509.06876 [astro-ph.HE].
- [61] B. Banerjee *et al.*, Pre-merger alert to detect prompt emission in very-high-energy gamma-rays from binary

neutron star mergers: Einstein Telescope and Cherenkov Telescope Array synergy, Astron. Astrophys. **678**, A126 (2023), arXiv:2212.14007 [astro-ph.HE].

- [62] C. S. Ye, W.-f. Fong, K. Kremer, C. L. Rodriguez, S. Chatterjee, G. Fragione, and F. A. Rasio, On the Rate of Neutron Star Binary Mergers from Globular Clusters, Astrophys. J. Lett. 888, L10 (2020), arXiv:1910.10740 [astro-ph.HE].
- [63] J. Samsing, M. MacLeod, and E. Ramirez-Ruiz, The Formation of Eccentric Compact Binary Inspirals and the Role of Gravitational Wave Emission in Binary-Single Stellar Encounters, Astrophys. J. 784, 71 (2014), arXiv:1308.2964 [astro-ph.HE].
- [64] J. Samsing, Eccentric Black Hole Mergers Forming in Globular Clusters, Phys. Rev. D 97, 103014 (2018), arXiv:1711.07452 [astro-ph.HE].
- [65] J. Samsing and E. Ramirez-Ruiz, On the Assembly Rate of Highly Eccentric Binary Black Hole Mergers, Astrophys. J. Lett. 840, L14 (2017), arXiv:1703.09703 [astroph.HE].