

The quark-lepton portal beyond lepto-quarks

Linda M Carpenter and Katherine Schwind and Taylor Murphy

December 31, 2024

Abstract

We explore models where single new exotic states interact with the Standard Model through an asymmetric Standard Model portal with couplings to at least one quark and one lepton. We write down all effective operators up to dimension six where such interactions couple the SM to spin 0 and spin 1/2 particles. We identify the exotic states accessible through the portal and find the interactions couple the SM to new particles with exotic combinations of baryon and lepton number, and particles with unusual SM charge, including states in higher dimensional representations of both SU(3) and SU(2), and states of higher electric charge. We discuss the phenomenology of these interactions including novel particle decays, and we classify some of the collider production modes for exotic states at LHC, LHeC and the muon collider.

1 Introduction

In this work we explore an unusual portal to new physics that allows us to access new particles with highly unusual standard model gauge and global charges. There have been scattered efforts to make general studies of particles of exotic charge[1][2][3][4][5][6][7][8][9][10]. This work attempt to take a more systematized approach by specifying all possible new states which can be accessed through a specific Standard Model portal[11][12].In this work, we are considering the lepton-quark portal. Here we catalog all new exotic particles of spin 0 and 1/2 which can be produced in interactions that contain at least one SM quark and one SM lepton. This portal will allow us to access new states with very unusual combinations of SU(3) and SU(2) charges, and unusual combinations of baryon and lepton number. This portal will be especially relevant to new physics at lepton-quark colliders like LHeC[13], but also produced many new interesting stand out discovery channels at LHC and a possible future muon collider[14].

The typical exotic particle that is accessible through quark-lepton couplings is the lepto-quark. This is a scalar or vector color triplet object that carries unit lepton number and baryon number 1/3. Leptoquarks may be singly produced at e-p colliders and pair produced in quark and gluon fusion processes at hadron colliders. As an interesting fact, there are still some light phenomenological

windows for leptoquarks in the few hundred GeV mass range. Standard leptoquarks, which couple to the standard model with renormalizable couplings, come in a limited variety of SU(2) representations -triplet, doublet, and singlet- and limited hypercharges[15],[16] [17].


In this work we will explore a catalog of exotic states with a quark-lepton bilinear interactions that occur at dimension 5 and 6. We consider operators that contain a quark lepton and single light exotic state (LEX). These operators may also contain one or more additional SM states. We are interesting operators which will contribute to single production modes of the LEX particles, since these production modes will allow us to access the highest mass ranges of new particles. This new portal will contain new dimension 5 and 6 interactions of the standard leptoquarks with SM. However, we will also catalog large classes of new and interesting exotics-beyond standard lepto quarks- can be accessed through the extended lepton quark portal. By studying these interactions, we find collider production modes for new light exotic states with highly unusual SM quantum numbers. To study these interactions we we invoke effective field theory. We will consider effective interactions of dimension greater than 4 that involve SM and exotic states, the Light Exotics Effective Field Theory (LEXEFT)[12]. Interactions have the form

$$L \sup \frac{1}{\Lambda^n} \Psi_{i..} \Psi_n \phi_\alpha \phi_\beta \dots \quad (1)$$

Here the fields Ψ are exotic states, the fields ϕ are Standard Model states, and Λ is the scale of the effective cut-off. In our case we will study all such operators that could involve a single LEX state, where the SM states involve at least one quark and at least one lepton.

$$L \sup \frac{1}{\Lambda^n} \Psi_{LEX}(Q_L, u_r, d_r)(L_l, l_r) \phi \quad (2)$$

were ϕ are additional SM fields. We can visualize the resultant interactions in the diagram below where X is LEX state, we are looking for interactions that couple a LEX single particle to one quark and one lepton, either alone or in conjunction with other SM particles.



These operators will tell us all processes in which we can produce a single LEX state in the quark-lepton portal up to the desired dimension in effective field theory. In this case we will list operators up to dimension 6. This portal allows us to explore extended exotics with lepto-quark features. Many of these

exotic states will have unusual combinations of higher lepton and baryon number. They can have a variety of highly unusual combinations of color and weak charge. In addition, this allows us to explore interesting collider production scenarios at different type of collider.

In particular, this allows the associated production of an exotic state with one or more SM particles from direct quark-lepton collision at LHEC in processes that look like $ql \rightarrow \psi_{LEX} + SM$. We expect from lepto-quark analysis that at least some of these states have mass windows unconstrained from LHC production that are in the few hundred GeV range[18][19]. In addition, we see that some of these operators will allow interesting 2-2 and 2-3 LHC processes involving the single production of exotic sates in association with Standard Model particles. Finally we can identify some processes by which LEX states may be singly produced at the muon collider or other lepton colliders leading to interesting hadronic signatures.

This paper is organized as follows, in section 2 we enumerate effective operators that contain spin 0 LEX states that can be accessed through the lepton-quark portal. We explain the quantum numbers of the LEX states found in the portal and detail some of the possible collider production and decay modes for the scalars. In section 3 we enumerate effective operators that contain fermionic LEX states. We describe the possible baryon and lepton number assignments of the various states as well as their color weak and hypercharge assignments. We then describe some of the possible particle decays and collider production modes. Section 5 mentions lepton-quark portal operators for vector LEX states. Section 6 concludes.

2 Spin 0 LEX states

We will now enumerate the operators of the extended lepton-quark portal up to effective dimension 6. Note that we will not include in this catalog the dimension 4 operators which lead renormalizable couplings of the standard scalar lepto-quarks. These operators have been laid out in a variety of existing phenomenological works. We will however, enumerate operators for effective interactions higher than dimension 4, these will represent new couplings in the lepton-quark portal. Some of these operators lead to new types of interactions for standard lepto-quarks. We will also see that we are able to access new types exotic particles which have standard model charges very different than renormalizable lepto-quark models. We will find several new fields in higher dimensional representation of the SM gauge groups. We note that all of the spin-zero particles that appear in the lepton-quark portal carry lepton number -1 and baryon number 1/3. In classifying the operators we have adopted notation similar to the operator classifications of SMEFT [20] to indicate operator Lorentz structure.

Below we begin by enumerating the spin 0 LEX states which can couple to a quark-lepton bilinear. In the tables below we list the effective dimension of the

operator, the operator and the $SU(3) \times SU(2) \times Y$ charges of the LEX state. In the below operators we denote the LEX state as ϕ . We begin with the CP-even scalar state. Below roman indices indicate fundamental SU(2) indices.

2.1 Operators Containing Higgs Couplings and Exotic Scalar Quarks

We first discuss new operators of the simplest forms that contain couplings between the quark-lepton bilinear, the new exotic state, and one or two Higgs fields. We will sub categorize these operators by the left and right handed fields Q_L, q_R, L_L, ℓ_R that appear in the operators. In these operators the exotic scalar ϕ must be an SU(3) fundamental. These particles then will be exotic types of scalar quark. Notice that in this class of spin 0 operators, the LEX states carry unit lepton number and a baryon number 1/3. We have written out SU(2) indices explicitly, and these indices are given by lower case roman letters.

2.1.1 Operators of the Form $QLH\phi$

We begin by listing operators with couple the LEX state to a single SM Higgs particle and the quark-lepton bilinear. These operators are effective dimension 5. We will begin by listing operators that contain both a left handed quark and left handed lepton in the fermion bilinear.

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 5	$\overline{Q_{L_i}} L_j \phi^{ijk} H_k$	(3,4,1/3)
dim 5	$\overline{Q_L^i} L_j \phi^j H_i, \overline{Q_L^i} L_i \phi^j H_j, \overline{Q_L^i} L_j \phi_i H^j$	(3,2,1/3)

Dimension 5 Higgs operators containing the QL bilinear

In the operators above we have fields that are SM SU(2) doublets. We find the SU(2) charges of the LEX state using the tensor product rule $\frac{1}{2} \otimes \frac{1}{2} \otimes \frac{1}{2} = \frac{3}{2} \oplus \frac{1}{2}$ which yields SU(2) representation of 4 and 2 for the LEX scalar. Recall that the SU(2) quadruplet is a symmetric object with 3 fundamental indices. For a doublet scalar there are various independent SU(2) contractions, which are represented in the table.

2.1.2 Operators of the Form $Q\ell H\phi$ and $qLH\phi$

We now list operators that contain one left handed fermion and one right handed fermion in the bilinear. We thus have operators that contain two SM SU(2) doublets. The allowed SU(2) representations of the LEX state follow the SU(2) tensor product rule $1/2 \otimes 1/2 = 0 \oplus 1$; that is they are triplets or singlets. The operators are given in the table below

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 5	$\overline{Q}_L \ell \phi_i^j H_j$	(3,3,4/3)
dim5	$\overline{Q}_L \ell \phi H_i$	(3,1,4/3)
dim5	$\overline{u}_R L_i \phi^{ij} H_j$	(3,3,4/3)
dim5	$\overline{u}_R L_i \phi H^i$	(3,1,4/3)
dim5	$\overline{d}_R L_i \phi^{ij} H_j$	(3,3,-2/3)
dim5	$\overline{d}_R L_i \phi H^i$	(3,1,-2/3)

Dimension 5 Higgs operators containing qL or Q ℓ bilinear

2.1.3 Operators of the Form $q\ell H\phi$

We finally list operators which contain both a right handed quark and right handed lepton in the fermion bilinear. In these operators, the Higgs is the only SM field that is an SU(2) doublet. This means the LEX state ϕ must also be an SU(2) doublet. We list operators in the table below.

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim5	$\overline{u}_R \ell \phi^i H_i$	(3,2,7/3)
dim5	$\overline{u}_R \ell \phi^i H_i^\dagger$	(3,2,13/3)
dim5	$\overline{d}_R \ell \phi^i H_i$	(3,3,1/3)
dim5	$\overline{d}_R \ell \phi^i H_i^\dagger$	(3,1,7/3)

Dimension 5 Higgs operators containing the q ℓ bilinear

2.2 Operators with two Higgs fields

Next we explore operators with a quark lepton bilinear couples to a LEX state and two Higgs fields. These operators will be of dimension 6.

Noting SU(2) contractions, we can see when listing di-Higgs operators that any terms containing $H_j \epsilon^{ij} H_i$ are zero.

2.2.1 Operators of the Form $QLHH\phi$

We first begin by listing the operators in which the quark and lepton in the fermion bi-linear are SU(2) doublets. Again we have written out SU(2) fundamental indices explicitly with roman letters. The LEX state is denoted as ϕ . Operators and LEX charge assignments are given in the table below.

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 6	$\overline{Q}_{L_i} L_j \phi^{ijkl} H_k H_l$	(3,5,-2/3)
dim 6	$\overline{Q}_{L_i} L_j \phi_i^j H^{k\dagger} H_k$	(3,3,4/3)
dim6	$\overline{Q}_{L_i} L_j \phi^{jk} H_i H_k,$	(3,3,-2/3)
dim6	$\overline{Q}_{L_i} L_j \phi H_i H^j$	(3,1,-2/3)
dim6	$\overline{Q}_{L_i} L_i \phi H_j H^{j\dagger}$	(3,1,4/3)

Dimension 6 di-Higgs operators containing the QL bilinear

The LEX states must be SU(3) fundamentals. There are four SM SU(2) doublets that appear in the operators. We construct the available SU(2) charges of LEX states with the SU(2) tensor product rule $\frac{1}{2} \otimes \frac{1}{2} \otimes \frac{1}{2} \otimes \frac{1}{2} = 2 \oplus 1 \oplus 0$. in spinor notation. This gives us SU(2) quintuplet, triplet or singlet representations for the LEX state. The SU(2) quintuplet is a symmetric object with 4 fundamental indices.

2.2.2 Operators of the Form $qLHH\phi$ and $Q\ell HH\phi$

We next enumerate the operators that contain one left handed and one right handed fermion in the bilinear. The fermions couple to two SM Higgs fields and a scalar LEX state. Again, the LEX state must be an SU(3) fundamental. To find the allowed SU(2) representations of the LEX state we note that there are three SU(2) doublets in these operators. The SU(2) tensor product rule tells us the available LEX representations follow $1/2 \otimes 1/2 \otimes 1/2 = 3/2 \oplus 1/2$. This gives us SU(2) quadruplet and doublet representations for ϕ . We write the operators in the table below with SU(2) indices are given by roman letters

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 6	$\overline{Q}_{L_i} \ell \phi^{ijk} H_j H_k$	(3,4,1/3)
dim 5	$\overline{Q}_{L_i} \ell \phi_i H^{k\dagger} H_k$	(3,2,7/3)
dim5	$\overline{Q}_{L_i} \ell \phi^i H_j H_j$	(3,2,1/3)
dim5	$\overline{u}_R L_i \phi^{ijk} H_j H_k$	(3,4,1/3)
dim5	$\overline{u}_R L_i \phi^i H_j H^{j\dagger}$	(3,2,7/3)
dim5	$\overline{u}_R L_i \phi^i H_j H^j$	(3,2,1/3)
dim5	$\overline{d}_R L_i \phi^{ijk} H_j H_k$	(3,4,-5/3)
dim5	$\overline{d}_R L_i \phi^i H_j H^{j\dagger}$	(3,2,1/3)
dim5	$\overline{d}_R L_i \phi^i H_j H^j$	(3,2,-5/3)

Dimension 6 di-Higgs operators containing the qL or Q ℓ bilinears

2.2.3 Operators of the Form and $q\ell HH\phi$

Finally we enumerate the di-Higgs operators that couple to a fermion bilinear in which both fermions are right handed. In this case, there are two SU(2) Higgs fundamentals in the operator. The SU(2) charges of the LEX state are given by

the tensor product rule $1/2 \otimes 1/2 = 0 \oplus 1$; the LEX states are SU(2) triplets or singlets. We list the operators in the table below.


dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 6	$\overline{u}_R \ell \phi^{ij} H_i H_j$	(3,3,4/3)
dim6	$\overline{u}_R \ell \phi H_i H^{i\dagger}$	(3,1,-10/3)
dim6	$\overline{d}_R \ell \phi^{ij} H_i H_j$	(3,3,-2/3)
dim6	$\overline{d}_R \ell \phi H_i H^{i\dagger}$	(3,1,4/3)

Dimension 6 di-Higgs operators containing the $q\ell$ bilinears

2.3 Some Collider Phenomenology With Higgs Couplings


Glancing at the Higgs operator entries in the lepto-quark portal we can extract a few interesting LEX states. Most interesting, we see states which are SU(3) fundamentals that have higher SU(2) charge. There are several new LEX states that are quadruplets and quintuplets of SU(2). These states also carry unit lepton number and baryon number 1/3. The SU(2) charge is a significant departure from the standard leptoquarks which are in the singlet, doublet and triplet representations of SU(2). The new multiplets contain multiple charged components. For example, the largest SU(2) charge we find is the quintuplet LEX state in the (3,5,-2/3) representation. This field appears in the dimension 6 operator $\overline{Q}_{L_i} L_j \phi^{ijkl} H_k H_l$. This field is an extended scalar lepto-quark with charge components (5/3,2/3,-1/3,-4/3,-7/3). Similarly the (3,4,1/3) field which appears in the dimension 5 operator $\frac{1}{\Lambda} \overline{Q}_{L_i} L_j \phi^{ijk} H_k$ has charged components (5/3,2/3,-1/3,-4/3). The exotic quadruplet that appears in the dimension 6 operator $\frac{1}{\Lambda^2} \overline{d}_R L_i \phi^{ijk} H_j H_k$ with quantum numbers (3,4,-5/3) has charge components (2/3,-1/3,-4/3,-7/3). These multiplets allow for interesting cascade decays between components if the masses can be modestly split using Higgs couplings. Another small point is that in the above operator list, we also see a different patters of hypercharge assignments in the triplet, doublet and singlet SU(2) representations of our exotic leptoquarks than in the standard dimension 4 leptoquarks models. This tells us the components of the LEX states will have different charge patters than in standard leptoquark models.

We can extract some interesting collider phenomenology from the above lepton-quark portal operators which contain LEX states and one or two Higgs fields. In the preceding operators, the Higgses can either appear in the operator as a propagating fields, or can be replaced by vevs. From the Higgs operators we can extract two types of interaction vertices that will play an interesting role in 2 to 2 collider production processes. If all Higgses are replaces by their vevs, these operators contain a three point interaction between a lepton-quark and LEX scalar. Instead, there may be a four point interactions between a lepton, quark, Higgs and LEX scalar. This four point interaction may follow from the one Higgs operators, or from the two-Higgs operators with one vev insertion. We represent the interactions below




Three and four point interactions of the Higgs operators in the lepton-quark portal

These interaction contribute to several interesting production modes in hadron and e-p and muon colliders. First, any of the leptoquarks can be pair produced at LHC through gluon fusion, including all of the exotic leptoquarks charge components. In addition, electroweak pair production can occur through quark fusion at LHC and through lepton fusion at the muon collider. However, we can see from our operator list there are LEX single production processes. These single production processes may be the dominant production modes if the LEX states are several TeV in mass. As we can see directly from 3 and 4 point interactions above, the exotic scalar lepto-quarks in the LEX catalog can be produced in an e-p collider such as LHeC in association with a Higgs field directly from lepton-quark fusion in the process $lq \rightarrow \phi + h$. There is also a very interesting LHC process in which an exotic lepto-quark could be produced in association with a standard model lepton from quark gluon fusion



LHC production of lepton and LEX state from quark-gluon fusion

In the process $qg \rightarrow l + \phi$. We also briefly mention a 2-3 process in which LEX leptoquark states can be singly produced at a future muon collider. Through a process where an incoming lepton fuses with an off shell gauge boson to produce a leptoquark, quark and forward lepton. The process $\bar{l}l \rightarrow l + q + \phi$



Single production of LEX state and quark in association with a hard lepton at a muon collider

These production processes will be especially interesting in the production of the LEX states with higher SU(2) charge, the quintuplets and quadruplets which contain multiply-charged states. We note that the quintuplet and quadruplet components with highest charged will not themselves be singly produced in the processes above, but can be accessed through SU(2) cascades if they are the lightest states. The multiply charged states may decay through other effective operators. The details of single production cascades will depend on the mass splittings in the specific model but there is potential for quite unusual collider signatures. An especially interesting collider scenario involves very small mass splittings which can result in longer lived particles.

2.4 Operators Containing Field Strength tensors: exotic SU(3) charged states

We now discuss a different class of effective operators in the quark-lepton portal which contain Gauge Field strength tensors. These operators will couple the lepton-quark bilinear to an exotic scalar state and one or more SM gauge bosons. These are dimension 6 operators of the general form $\bar{f}_1 \sigma^{\mu\nu} f_2 F_{\mu\nu} \phi$. Where $F_{\mu\nu}$ can be any of the three SM field strength tensors. In this sector of the lepton-quark portal, the SM can couple to LEX states that have even more exotic SM representations than the Higgs-type operators. We will see that operators containing the gluon field strength tensor allow us to access LEX states that are 15-plets, and sextets of SU(3). Operators containing the SU(2) field strength tensor will also allow couplings to LEX states in higher dimensional representation of SU(2). Once again the LEX states in this portal will have unit lepton number -1 and baryon number 1/3-the same baryon and lepton number assignments as standard lepto-quarks.

2.4.1 Operators of the form $QLG\phi$

Operators coupling to an SU(3) field strength tensor $G_{\mu\nu}$ are listed in the table below. Color charge assignments are quite interesting as operators contain one quark field and one gluon. Thus color representation of the LEX states is thus given by the the SU(3) tensor product rule [21]

$$\mathbf{8} \otimes \mathbf{3} = \mathbf{15} \oplus \mathbf{6} \oplus \mathbf{3}$$

The 15-plet of SU(3) is the highest possible SU(3) charge assignment. The 15 in question is an object with two symmetrized lower indices and one upper index [22]. This LEX state accessible through the portal has baryon number 1/3 and lepton number -1, we can refer to it as a scalar decapenta-leptoquark . The sextet of SU(3) is an object with two symmetrized lower indices. The scalar color sextet LEX state have lepton number -1 and baryon number 1/3, so we can refer to as a scalar lepto-diquark. The SU(3) triplet will have quantum numbers assigned like the regular lepto-quarks from renormalizable models; the $QLG\phi$ operators thus provide additional SM couplings to normal lepto-quark models.

Operators with both left handed quark and left handed lepton field are listed in the table below. The possible SU(2) charge assignments for LEX states are triplet and singlet, given by the SU(2) tensor product rule in spinor notation $\mathbf{1}/2 \otimes \mathbf{1}/2 = \mathbf{0} \oplus \mathbf{1}$., the LEX states maybe SU(2) triplets or singlets. We thus have 6 operators involving left handed SM fermions.

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 6	$\overline{Q}_{L_i} \sigma^{\mu\nu} L_j \phi^{ij} G_{\mu\nu}$	(15,3,4/3)
dim6	$\overline{Q}_L^i \sigma^{\mu\nu} L_i \phi G_{\mu\nu}$	(15,1,4/3)
dim 6	$\overline{Q}_{L_i} \sigma^{\mu\nu} L_j \phi^{ij} G_{\mu\nu}$	(6,3,4/3)
dim6	$\overline{Q}_L^i \sigma^{\mu\nu} L_i \phi G_{\mu\nu}$	(6,1,4/3)
dim6	$\overline{Q}_{L_i} \sigma^{\mu\nu} L_j \phi^{ij} G_{\mu\nu}$	(3,3,4/3)
dim5	$\overline{Q}_L^i \sigma^{\mu\nu} L_i \phi G_{\mu\nu}$	(3,1,4/3)

Dimension 6 operators of the lepton-quark portal with gluon field strength tensor and QL bilinear

Operators containing one left handed and one right handed SM fermion in the are simpler. We maintain the same SU(3) charge contractions. Now, however, our only choice for SU(2) charge of the LEX state is a doublet.

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim6	$\overline{Q}_L^i \sigma^{\mu\nu} \ell \phi_i G_{\mu\nu}$	(15,2,7/3)
dim6	$\overline{u}_R \sigma^{\mu\nu} L_i \phi^i G_{\mu\nu}$	(6,2,7/3)
dim6	$\overline{d}_R \sigma^{\mu\nu} L_i \phi^i G_{\mu\nu}$	(3,1,1/3)

Dimension 6 operators of the lepton-quark portal with gluon field strength tensor and qL or Q ℓ bilinear

and of course operators which contain two right handed fermions in the bilinear only allow LEX states that are SU(2) singlets

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim6	$\overline{u_R}\sigma^{\mu\nu}\ell\phi G_{\mu\nu}$	(15,1,10/3)
dim6	$\overline{d_R}\sigma^{\mu\nu}\ell\phi G_{\mu\nu}$	(15,1,4/3)
dim6	$\overline{u_R}\sigma^{\mu\nu}\ell\phi G_{\mu\nu}$	(6,1,10/3)
dim6	$\overline{d_R}\sigma^{\mu\nu}\ell\phi G_{\mu\nu}$	(6,1,4/3)
dim6	$\overline{u_R}\sigma^{\mu\nu}\ell\phi G_{\mu\nu}$	(3,1,10/3)
dim6	$\overline{d_R}\sigma^{\mu\nu}\ell\phi G_{\mu\nu}$	(3,1,4/3)

Dimension 6 operators of the lepton-quark portal with gluon field strength tensor and ql bilinear

2.4.2 Operators of the form $QLW\phi$

We now consider operators which contain the SU(2) field strength tensor $W_{\mu\nu}$ -an adjoint of SU(2). In these operators, the only possible SU(3) charge allowed for the LEX state is a 3 of SU(3). The LEX states will have standard lepto-quark baryon and lepton number. The SU(2) charges, however can be quite complex depending on the type of fermions in the bilinear.

If two left handed fermions in the fermion bi-linear the allowed SU(2) charges of the LEX state follow the SU(2) tensor product rule in spinor notation

$$\mathbf{1} \otimes \mathbf{1}/2 \otimes \mathbf{1}/2 = \mathbf{2} \oplus \mathbf{1} \oplus \mathbf{0}$$

This means allowed SU(2) representations of the LEX state are the quintuplet, the triplet, or the singlet. Recall that the SU(2) quintuplet is a 4 index symmetric object. We give operators with two left handed fermions in the table below

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 6	$\overline{Q_{L_i}}\sigma^{\mu\nu}L_jW_{\mu\nu kl}\phi^{ijkl}$	(3,5,4/3)
dim 6	$\overline{Q_L^i}\sigma^{\mu\nu}L_jW_{\mu\nu l}^j\phi_i^l$	(3,3,4/3)
dim6	$\overline{Q_L^i}\sigma^{\mu\nu}L_jW_{i\mu\nu}^j\phi$	(3,1,4/3)

Dimension 6 operators of the lepton-quark portal with SU(2) field strength tensor and QL bilinear

If one of the fields in the fermion bilinear is right handed and the other is left handed then we have one SU(2) fundamental and one adjoint in the operator. The SU(2) representation of the LEX state is given by the tensor product rule

$$\mathbf{1} \otimes \mathbf{1}/2 = \mathbf{3}/2 \oplus \mathbf{1}/2$$

in spinor notation. The available SU(2) representations for the LEX state are the quadruplet and the doublet. The 4 of SU(2) is a three index symmetric object. We write the allowed operators in the table below.

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 6	$\overline{Q_L^i} \sigma^{\mu\nu} \ell W_{jk\mu\nu} \phi^{ijk}$	(3,4,7/3)
dim 6	$\overline{Q_L^i} \sigma^{\mu\nu} \ell W_{\mu\nu}^{ij} \phi_j$	(3,2,7/3)
dim 6	$\overline{u_R} \sigma^{\mu\nu} L_i W_{jk\mu\nu} \phi^{ijk}$	(3,4,7/3)
dim 6	$\overline{u_R} \sigma^{\mu\nu} L_i W_{j\mu\nu}^i \phi^j$	(3,2,7/3)
dim 6	$\overline{d_R} \sigma^{\mu\nu} L_i W_{jk\mu\nu} \phi^{ijk}$	(3,4,1/3)
dim 6	$\overline{d_R} \sigma^{\mu\nu} L_i W_{j\mu\nu}^i \phi^j$	(3,2,1/3)

Dimension 6 operators of the lepton-quark portal with SU(2) field strength tensor and qL or QL bilinear

Finally, if the the fermion bilinear in the operator contains only right handed fields then the LEX state must be an SU(2) triplet to contact indices with the field strength tensor. We give these operators in the table below

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 6	$\overline{u_R} \sigma^{\mu\nu} \ell W_{\mu\nu}^{ij} \phi_{ij}$	(3,3,10/3)
dim 6	$\overline{d_R} \sigma^{\mu\nu} \ell W_{\mu\nu}^{ij} \phi_{ij}$	(3,3,4/3)

Dimension 6 operators of the lepton-quark portal with gluon field strength tensor and ql bilinear

This charge assignment for the LEX state is one of the standard leptoquarks.

2.4.3 Operators of the form $QLB\phi$

Finally we may insert a U(1) field strength tensor into the operator. In this case the charge assignments of the LEX states ensure the field will appear as one of standard leptoquarks. The LEX state must be an SU(3) fundamental. It will also have standard lepto-quark lepton and baryon number. If the operator contains two left handed SM fermions the allowed SU(2) charges are given by the SU(2) tensor product $\mathbf{1/2} \otimes \mathbf{1/2} = \mathbf{1} \oplus \mathbf{0}$; that is an SU(2) triplet or singlet. If only one fermion of the bi-linear is left handed, the LEX state must be an SU(2) doublet. If both fermions are right handed the LEX state is an SU(2) singlet. We write the allowed operators in the table below.

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 6	$\overline{Q_L^i} \sigma^{\mu\nu} L_j \phi_i^j B_{\mu\nu}$	(3,3 4/3)
dim 6	$\overline{Q_L^i} \sigma^{\mu\nu} L_i \phi B_{\mu\nu}$	(3,1 4/3)
dim 6	$\overline{Q_L^i} \sigma^{\mu\nu} \ell \phi_i B_{\mu\nu}$	(3,3 7/3)
dim 6	$\overline{u_R} \sigma^{\mu\nu} L_i \phi^i B_{\mu\nu}$	(3,3 7/3)
dim 6	$\overline{d_R} \sigma^{\mu\nu} L_i \phi^i B_{\mu\nu}$	(3,3 1/3)
dim 6	$\overline{u_R} \sigma^{\mu\nu} \ell \phi B_{\mu\nu}$	(3,3 10/3)
dim 6	$\overline{d_R} \sigma^{\mu\nu} \ell \phi B_{\mu\nu}$	(3,1,4/3)


Dimension 6 operators of the lepton-quark portal with U(1) field strength tensor

These interactions represent a new effective coupling between a standard scalar lepto-quark with a lepton-quark and photon.

2.5 Some Collider Phenomenology with Field Strength Tensor operators.

The catalog of dimension 6 operators involving field strength tensors reveals some very interesting types of exotic states. We note that the states of higher SU(3) charge such as the sextet and 15 will be come with a large Clebsch-Gordon coefficient to contract SU(3) charges. This will provide a large enhancement for production processes involving those states. We also find exotic scalar sextets that carry lepton and baryon number. The phenomenology of scalar sextets has usually focuses on renormalizable operators pertaining to SU(2) singlet states [23][24][7] though there have been recent efforts to study the phenomenology of effective operators and sextets of higher SU(2) charge [8][25] [26][27][10][1]. The operators that involve the SU(2) field strength tensor allow us to access LEX states in high representation of SU(2). For this operator type, the highest SU(2) charge accessible is a quintuplet in representation (3,5,4/3) resulting from the dimension 6 operator $\bar{Q}_{Li}\sigma^{\mu\nu}L_jW_{\mu\nu kl}\phi^{ijkl}$. This state is different from the quintuplet that was accessible through the di-Higgs operator. This state has charge components (8/3,5/3,2/3,-1/3,-4/3). Various states in this catalog have quantum numbers which depart from the dimension 4 lepto-quark models. However, standard lepto-quarks also appear in the operator catalog and this portal provides additional beyond tree-level interactions.

We can see there are several new types of four point four-particle vertices induces by the operators. These vertices contain a SM lepton and quark, a LEX scalar, and a SM gauge boson which can be gluon, W,Z or photon. We visualize the new interactions in the diagrams below




Four point vertices from field strength tensor operators in the lepto-quark portal

These lead to a host of new collider signatures. We first note that the operators lead to interesting 3 body decays of the exotic lepto-quarks. Including the resonance decay of a lepto-quark into two jets and a lepton $\phi \rightarrow l + q + g$ through the $qLG\phi$ operators and $\phi \rightarrow l + q + \gamma$ decay through the $qLW\phi$ and

$qlB\phi$ operators. There are also three body decays to a lepton quark and heavy gauge boson $\phi \rightarrow l + q + W/Z$. These three body decays will also be open to the standard lepto-quarks. These decays may become important in the region of standard-leptoquark parameter space where leptoquark couplings to SM fermions are suppressed-yet they can still be pair produced through gluon fusion or electroweak processes.


We also find new single LEX state production mechanisms at various colliders. For example, the above diagrams make clear a standard or exotic lepto-quark, lepto-diquark, or, decapenta-leptoquark could be produced at an e-p collider in association with a gluon jet. The process is $lq \rightarrow \phi + g$. This could be an interesting production mode for the states of unusual color-charge, especially the 15. Further, we see that various LEX states in the SU(3) triplet representation can be singly produced at e-p colliders in association with an electro-weak gauge boson. Most interesting would be the photon associated production channel through the process $lq \rightarrow \phi + \gamma$.

There is an interesting LHC production mode that follows from the four point vertex of the $qlG\phi$ operator. This allows the associated production of a single LEX state with an associated lepton from quark gluon fusion. This process is pictured in the diagram below



LHC production of LEX scalar and one SM lepton from quark gluon fusion

and the process is $qg \rightarrow \phi+l$. There are also a few interesting 2-3 LHC production processes that produce a single LEX state through the operators. One is a $ql\phi$ final state produced from gluon fusion, the other is a ϕVl final state produced from quark fusion. The diagrams for these processes are shown below



LHC production of a)LEX with a quark and lepton from gluon fusion b) LEX state with lepton and SM gauge boson from quark gluon fusion

In the first process there will be another contributing diagram from the 5 point vertex involving the non-abelian gluon field strength tensor. This process is $gg \rightarrow q+l+\phi$. The hard lepton in the process might make this an interesting discovery signature for some LEX models. In a second production process, a quark and gluon can fuse into an off-shell quark which produces a lepton, LEX state, and electroweak boson in the process $qg \rightarrow q^* \rightarrow l+\phi+W/Z/\gamma$. A possible $l\gamma\phi$ final state contains both a hard lepton and photon and presents a very interesting new discovery channel for new physics in lepto-quark and exotic lepto-quark models.

2.6 Operators of Type $FFD\phi$

We can also create alternate types of operators in the lepton-quark portal that involve derivatives. We will now discuss dimension 5 operators of the form

$$\bar{f}_1\gamma^\mu L f_2 D\phi.$$

In the Lorentz structure of these operators, a vector fermion bi-linear is contracted with one covariant derivative and a scalar field. In principle, the derivative may appear in front of any of the three other fields in the operator. For each type of fermion bi-linear $\bar{f}_1 f_2$, we can identify the operator Lorentz structures.

$$\bar{f}_2\gamma^\mu \overleftarrow{D}_\mu f_2\phi, \bar{f}_1\gamma^\mu f_2 D_\mu\phi \text{ and } \bar{f}_1\gamma^\mu \overrightarrow{D}_\mu f_2\phi.$$

We will show that these operators of this type are redundant to the portal. We will show how we can reduce the number of these operators with integration by parts and other QFT identities in a procedure that is familiar from SMEFT[20][28][29].

First we note that the operators which contain d-slash are proportional to fermion equations of motion. We can insert the tree level equations of motion for the SM fermions into the operators above. Using the example of the $\bar{Q}L$ bilinear, for instance, we have

$$\begin{aligned} \bar{Q}\gamma^\mu \overrightarrow{D}_\mu L\phi &= \bar{Q}_L y_\ell H\ell\phi \\ &\text{and} \\ \bar{Q}_L\gamma^\mu \overleftarrow{D}_\mu L\phi &= \bar{d}_R y_d H L\phi + \bar{u}_R y_u H^\dagger L\phi \end{aligned}$$

We can see that the resulting operators after equations of motion are inserted are of the type $Q\ell H\phi$ and $qLH\phi$ which we have already cataloged. This means that the operators appearing with d-slash are redundant within our portal. They have already been accounted for by the Higgs-type operators and can be removed. We also have an integration by parts relation,

$$f_1\gamma^\mu\overleftarrow{D}_\mu f_2\phi = f_1\gamma^\mu\overrightarrow{D}_\mu f_2\phi + \overline{f}_1\gamma^\mu L f_2 D_\mu\phi$$

This condition allows us to remove the final operator proportional to the derivative of ϕ . This leaves us with no new non-redundant operators under the class $FFD\phi$.

2.7 Operators of Type $FFD\phi H$

At mass dimension 6, we can add a Higgs scalar to the operators in the class above. These operators involve one covariant derivative. These represent interactions between the lepton-quark bilinear, an exotic scalar, a Higgs particle and one or more SM gauge bosons. If a Higgs vev is inserted, this operator couples lepton-quark bilinear, an exotic scalar and one or more SM gauge bosons. The derivative in the operator may act on any of the 4 fields in the operator. The allowed operators are of types

$$\overline{f}_2\gamma^\mu\overleftarrow{D}_\mu f_2\phi H, f_1\gamma^\mu\overrightarrow{D}_\mu f_2\phi H, \overline{f}_1\gamma^\mu f_2 D_\mu\phi H, \overline{f}_1\gamma^\mu f_2\phi D_\mu H$$

making four operators. Following the discussion in the preceding subsection we can remove the two operators with $\gamma^\mu D_\mu$ from the portal as they are equivalent to existing operators. Inserting the fermion equations of motion we have

$$\begin{aligned} \overline{f}_1\gamma^\mu\overleftarrow{D}_\mu f_2\phi H &= y_{f_1} H \overline{f}_{1R/L} f_2\phi H \\ &\text{and} \\ \overline{f}_1\gamma^\mu\overrightarrow{D}_\mu f_{2L/R}\phi H &= y_{f_2} \overline{f}_1 f_{2R/L}\phi H \end{aligned}$$

We see that these operators are equivalent in form to operators of the form $ff\phi HH$ and thus redundant.

There is one integration by parts relation between the remaining operators

$$D_\mu\overline{f}_1\gamma^\mu L f_2\phi H = \overline{f}_1\gamma^\mu D_\mu f_2 L\phi H + \overline{f}_1\gamma^\mu f_2 D_\mu\phi H + \overline{f}_1\gamma^\mu f_2\phi D_\mu H.$$

We are thus left with a single non-redundant operator which we can choose to be $\overline{f}_1\gamma^\mu f_2 D_\mu\phi H$. Then possible SM charges of the LEX states are the same as the charges for operators of the form $ff\phi H$, but we have an additional covariant derivative in the operator. Operators are given in the table below

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 6	$\overline{Q_{L_i}}\gamma^\mu L_j D_\mu \phi^{ijk} H_k$	(3,4,1/3)
dim 6	$\overline{Q_{L_i}} L_j \phi^j H_i, \overline{Q_{L_i}} L_i \phi^j H_j, \overline{Q_{L_i}} L_j \phi_i H^j$	(3,2,1/3)
dim 6	$\overline{Q_{L_i}} \ell D_\mu \phi_i^j H_j$	(3,3,4/3)
dim6	$\overline{Q_{L_i}} \ell D_\mu \phi H_i$	(3,1,4/3)
dim6	$\overline{u_{R_i}} L_i D_\mu \phi^{ij} H_j$	(3,3,4/3)
dim6	$\overline{u_{R_i}} L_i D_\mu \phi H^i$	(3,1,4/3)
dim6	$\overline{d_{R_i}} L_i D_\mu \phi^{ij} H_j$	(3,3,-2/3)
dim6	$\overline{d_{R_i}} L_i D_\mu \phi H^i$	(3,1,-2/3)
dim6	$\overline{u_{R_i}} \ell D_\mu \phi^i H_i$	(3,2,7/3)
dim6	$\overline{u_{R_i}} \ell D_\mu \phi^i H_i^\dagger$	(3,2,13/3)
dim6	$\overline{d_{R_i}} \ell D_\mu \phi^i H_j$	(3,3,1/3)
dim6	$\overline{d_{R_i}} \ell D_\mu \phi^i H_i^\dagger$	(3,1,7/3)

Dimension 6 operators of the type $FFD\phi H$

2.8 Operators of Type $ffDD\phi$

The final type of operator involving scalar LEX states we will consider involves a s LEX field coupling to a quark-lepton bilinear with two covariant derivatives. The LEX states accessible through this operator have the same SM charge assignments as those of the dimension 4 lepto-quark couplings of the form $\overline{f_1} f_2 \phi$. These operators will allow a LEX state to couple to a quark-fermion bi-linear and at least one SM gauge boson. We can show that all of the operators of this type are redundant. This argument follows closely calculations done in reference [11]. Operators of this type have three possible Lorentz structures for the derivatives

$$\gamma^\mu D_\mu \gamma^\nu D_\nu, D_\mu D_\nu \sigma_{\mu\nu}, D^2$$

We will first explain the redundancy in the simplest operators which contain $D_\mu D^\mu$, arguments for the redundancy of the rest of the operators will follow. Our operators contain three fields and 3 derivatives. There are thus 6 possible types of operators. However these operators are related by three integration by parts relations. We have 6 operators and three conditions, we are thus free to eliminate three operators. We choose to keep operators of the type

$$D^2 \overline{f_1} f_2 \phi, \overline{f_1} D^2 f_2 \phi, \overline{f_1} f_2 D^2 \phi.$$

We can now use field theory relation and equations of motion to show these operators are redundant. First, we can see that the $\overline{f_1} f_2 D^2 \phi$ operators is proportional to the ϕ equation of motion.

$$D^2 \phi = m\phi + \partial V / \partial \phi.$$

We see from substituting the equation of motion back into the operator that we can reduce the operator $\bar{f}_1 f_2 m \phi$ + terms with multiple ϕ fields from the ϕ potential. The single ϕ term in the operator is proportional to standard dimension 4 lepto-quark operators which are already known in the literature. Terms with multiple ϕ fields are not considered in this portal. We can thus remove the operator $\bar{f}_1 f_2 D^2 \phi$. We now turn to the other two operators.

The other terms have two derivatives acting on a fermion field, for example $\bar{f}_1 D^2 f_2 \phi$. Here we can use the known QFT relation

$$D^2 f = \gamma^\mu D_\mu \gamma^\mu D_\mu f - i \sigma^{\mu\nu} F_{\mu\nu}.$$

Which is worked out in references [29][11]. The term proportional to $F_{\mu\nu}$ gives operators of the type $\bar{f}_1 \sigma^{\mu\nu} f_2 F^{\mu\nu} \phi$ which are detailed in section 2.4 These are thus already counted in the portal and are redundant. The first term containing D-slash is proportional to the fermion equations of motion.

$$D^\mu \gamma_\mu f = y H f_{L,R}$$

Inserting this equation of motion into the operator we get

$$\bar{f}_1 D^\mu \gamma_\mu y H f_{2L/R} \phi = \bar{f}_1 \gamma_\mu y D^\mu H f_{2L/R} \phi + \bar{f}_1 y H D^\mu \gamma_\mu f_{2R/L} \phi$$

Where the first term has the same form as the operators $\bar{f}_1 f_2 D H \phi$ discussed in the previous subsection. Once another fermion equation of motion is inserted, the second term has the form $\bar{f}_1 y^2 H H f_2 \phi$ which is one of the double Higgs form discussed in section 2.2 We can see that the operators

$$D^2 \bar{f}_1 f_2 \phi \text{ and } \bar{f}_1 D^2 f_2 \phi$$

are redundant.

Next, one can check that operators with two slashed derivatives acting on fermions are also redundant. These operators can be reduced directly using the fermion equations of motion to operators of the form $f f D H \phi$ and $f f H H \phi$ as just discussed above. These operators are also redundant.

Finally, we consider operators that contain the $D_\nu D_\mu \sigma^{\mu\nu}$ structure. We can see that, like the $D^\mu D_\mu$ operators we will have six possibilities for derivatives acting on the fields in the operators. Once again we will have three integration by parts relations which allows us to reduce to only three operators.

$$\bar{f}_1 \sigma^{\mu\nu} f_2 D_\mu D_\nu \phi, \bar{f}_1 \sigma^{\mu\nu} D_\mu D_\nu f_2 \phi, D_\mu D_\nu \bar{f}_1 \sigma^{\mu\nu} f_2 \phi$$

We now use a field theory relation to reduce these operators, expressing $D_\mu D_\nu$ as the sum of a commutator and anti-commutator we have

$$D_\mu D_\nu = \frac{g}{2} F_{\mu\nu} + \frac{1}{2} \{D_\mu, D_\nu\}$$

When inserted into the three operators above, the piece proportional to the commutator of derivatives will vanish when contracted with the antisymmetric $\sigma^{\mu\nu}$. The first term reduced the operators to the form $\bar{f}_1 \sigma^{\mu\nu} f_2 F_{\mu\nu} \phi$ which have already been cataloged in the text. We have thus shown that these two derivative operators are redundant to the lepton-quark portal.

3 Pseudoscalar LEX states

We note that pseudoscalar, CP odd spin 0 particles, can couple through the lepton-quark portal with operators that differ from the scalar operators only slightly. Operators of the type $FF\phi H$ and $FF\phi HH$ differ through extra insertion of γ^5 in the fermion bilinear. As an example, we give the table for $QL\phi H$ operators with pseudoscalars

dimension	Operators with scalar LEX fields (CP odd)	(SU(3),SU(2),Y)
dim 5	$\overline{Q_{L_i} \gamma^5 L_j} \phi^{ijk} H_k$	(3,4,1/3)
dim 5	$\overline{Q_{L_i} \gamma^5 L_j} \phi^j H_i, \overline{Q_{L_i} \gamma^5 L_i} \phi^j H_j, \overline{Q_{L_i} \gamma^5 L_j} \phi_i H^j$	(3,2,1/3)

Dimension 5 Higgs type pseudoscalar operators with QL bilinear

where here the pseudoscalar LEX state is represented as $\tilde{\phi}$. This formula holds for the rest of the $qL\phi H$, $Q\ell\phi H$, $q\ell\phi H$, $qL\phi HH$, $Q\ell\phi HH$, $q\ell\phi HH$ operators. For brevity we will not reproduce all of the Higgs type operator tables.

For operators containing gauge field strength tensors we have two options for constructing the operators that involve pseudoscalar LEX states. The operators are similar to those in subsections 2.7-2.9 but they can contain either an additional γ^5 insertion in the fermion bilinear, or they can contain a dual field strength tensor in place of the regular field strength tensor. For example, $QL\phi G$ operators with a pseudoscalar ϕ coupling to the pseudoscalar current are

dimension	Operators with scalar LEX fields (CP odd)	(SU(3),SU(2),Y)
dim 6	$\overline{Q_{L_i} \sigma^{\mu\nu} \gamma^5 L_j} \phi^{ij} G_{\mu\nu}$	(15,3,4/3)
dim6	$\overline{Q_{L_i} \sigma^{\mu\nu} \gamma^5 L_i} \phi G_{\mu\nu}$	(15,1,4/3)
dim 6	$\overline{Q_{L_i} \sigma^{\mu\nu} \gamma^5 L_j} \phi^{ij} G_{\mu\nu}$	(6,3,4/3)
dim6	$\overline{Q_{L_i} \sigma^{\mu\nu} \gamma^5 L_i} \phi G_{\mu\nu}$	(6,1,4/3)
dim6	$\overline{Q_{L_i} \sigma^{\mu\nu} \gamma^5 L_j} \phi^{ij} G_{\mu\nu}$	(3,3,4/3)
dim5	$\overline{Q_{L_i} \sigma^{\mu\nu} \gamma^5 L_i} \phi G_{\mu\nu}$	(3,14,3)

Dimension 6 pseudoscalar operators with gluon field strength tensor and QL bilinear

In stead operators with pseudoscalar ϕ that couple to the dual field strength may be written

dimension	Operators with scalar LEX fields (CP)	(SU(3),SU(2),Y)
dim 6	$\overline{Q_{L_i}} \sigma^{\mu\nu} L_j \phi^{ij} \tilde{G}_{\mu\nu}$	(15,3,4/3)
dim6	$\overline{Q_L}^i \sigma^{\mu\nu} L_i \phi \tilde{G}_{\mu\nu}$	(15,1,4/3)
dim 6	$\overline{Q_{L_i}} \sigma^{\mu\nu} L_j \phi^{ij} \tilde{G}_{\mu\nu}$	(6,3,4/3)
dim6	$\overline{Q_L}^i \sigma^{\mu\nu} L_i \phi \tilde{G}_{\mu\nu}$	(6,1,4/3)
dim6	$\overline{Q_{L_i}} \sigma^{\mu\nu} L_j \phi^{ij} \tilde{G}_{\mu\nu}$	(3,3,4/3)
dim5	$\overline{Q_L}^i \sigma^{\mu\nu} L_i \phi \tilde{G}_{\mu\nu}$	(3,1,4/3)

Dimension 6 pseudoscalar operators with gluon field strength tensor and QL bilinear

Through a similar procedure we can generate all of the operators of the type $f_1 \sigma^{\mu\nu} f_2 F_{\mu\nu} \tilde{\phi}$. The pseudoscalar operators may be of interest as collider production for pseudo-scalar states is generally high [30].

4 Fermionic LEX States

We will now enumerate operators in the lepton-quark portal where the SM couples to a fermionic spin 1/2 LEX state. These are dimension 6 operators in which a lepton-quark bilinear couples to a LEX fermion and one other standard model fermion. In These operators we denote the spin 1/2 LEX state as ψ . If we denote the Standard Model fermions as $f_{1,2,3}$ then the LEX operators have the form $f_1 \Gamma f_2 \psi \Gamma f_3$.

The operators may contain the scalar, pseudoscalar, vector, axial vector, or tensor currents which we have written as $\Gamma \supset 1, \gamma^5, \gamma^\mu \gamma^\mu \gamma^5, \sigma^{\mu\nu}$. This sector of the lepton-quark portal is interesting because the lepton and baryon numbers of the LEX state vary depending on SM fermions appear in the operator. As we will see we will be able to access LEX states with baryon number 2/3 and lepton number -1 as well as states with lepton number -2 and baryon number 1. We will also be able to access exotic fermionic with color charge and lepton number but no baryon number.

4.1 Operators with two quarks

We first enumerate the fermionic dimension 6 operators which contain a spin 1/2 LEX state ψ , two SM quarks and one SM lepton. The operators may either contain a quark and anti-quark or two (anti)quarks. In the first case the color assignments of the LEX state are given by the SU(3) tensor product rules $\bar{3} \otimes 3 = 1 \oplus 8$ and in the second case by $\bar{3} \otimes 3 = \bar{3} \oplus 6$. Therefore the allowed SU(3) representations are singlet, triplet, sextet and octet. Since there is one lepton in the operators, the LEX states will have lepton number -1. If there are is a quark anti-quark pair (color singlet or octet) the baryon number will be 0. If there is are two (anti)quarks (color triplet or sextet) the baryon number will be 2/3. The available SU(2) representations depend on the combination of left and right handed SM fermions in the operator.

4.1.1 operators of the form $QL\psi Q$

If all of the SM fermions in the operators are left handed then the available SU(2) representations of the LEX state are $1/2 \otimes 1/2 \otimes 1/2 = 3/2 \oplus 1/2$ or quadruplet and doublet. In the tables below the lower case roman letters represent SU(2) fundamental indices. We have written out the SU(3) contractions using upper case roman letters for fundamental SU(3) indices.

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 6	$\overline{Q}_{Li}{}^A \Gamma L_j \overline{\psi}_A^{ijkB} \Gamma Q_{kB}$	(8,4,1)
dim6	$\overline{Q}_{Li}{}^A \Gamma L_j \overline{Q}_k{}^B \Gamma \psi_{AB}^{ijk}$	(6,4,5/3)
dim 6	$\overline{Q}_L{}^{iA} \Gamma L_i \overline{\psi}_A^{jB} \Gamma Q_{jB}$	(8,2,1)
dim6	$\overline{Q}_L{}^{iA} \Gamma L_j \overline{\psi}_{iA}{}^j \Gamma Q_B^j$	(8,2,1)
dim6	$\overline{Q}_L{}^{iA} \Gamma L_j \overline{\psi}_A^{Bj} \Gamma Q_{iB}$	(8,2,1)
dim6	$\overline{Q}_L{}^{iA} \Gamma L_i \overline{Q}^j{}^B \Gamma \psi_{jAB}$	(6,2,5/3)
dim6	$\overline{Q}_L{}^{iA} \Gamma L_j \overline{Q}^j{}^B \Gamma \psi_{iAB}$	(6,2,5/3)
dim6	$\overline{Q}_L{}^{iA} \Gamma L^j \overline{Q}_i{}^B \Gamma \psi_{jAB}$	(6,2,5/3)
dim 6	$\overline{Q}_{Li}{}^A \Gamma L_j \overline{Q}_k{}^B \Gamma \psi^{ijkC} f_{ABC}$	(3,4,1)
dim6	$\overline{Q}_L{}^{iA} \Gamma L_i \overline{Q}^j{}^B \Gamma \psi_j^C f_{ABC}$	(3,2,1)
dim6	$\overline{Q}_L{}^{iA} \Gamma L_j \overline{Q}^j{}^B \Gamma \psi_i^C f_{ABC}$	(3,2,1)
dim6	$\overline{Q}_L{}^{iA} \Gamma L_j \overline{Q}_i{}^B \Gamma \psi^{Cj} f_{ABC}$	(3,2,1)
dim6	$\overline{Q}_{Li}{}^A \Gamma L_j \overline{\psi}^{ijk} \Gamma Q_{kA}$	(1,4,1)
dim6	$\overline{Q}_L{}^{iA} \Gamma L_i \overline{\psi}^j \Gamma Q_{jA}$	(1,2,1)
dim6	$\overline{Q}_L{}^{iA} \Gamma L_j \overline{\psi}_i \Gamma Q_A^j$	(1,2,1)
dim6	$\overline{Q}_L{}^{iA} \Gamma L_j \overline{\psi}^j \Gamma Q_{iA}$	(1,2,1)

Dimision 6 fermionic lepton-quark portal operators Containing QQL

4.1.2 operators of the form $Ql\psi Q$, and $qL\psi Q$

In the next set of operators two of the SM fermions in the operator are left handed. The SU(3) representations of the LEX state ψ are still 1,3,6 and 8 however the allowed SU(2) representations are given by $1/2 \otimes 1/2 = 0 \oplus 1$ or triplet and singlet. Once again in the table of operators below SU(2) indices are given by lower case roman letters and SU(3) fundamental indices are given by upper case roman letters

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 6	$\overline{Q}_L^{Ai} \Gamma \ell \overline{\psi}_{iA}^j \Gamma Q_{jB}$	(8,3,2)
dim 6	$\overline{u}_R^A \Gamma L^i \overline{\psi}_{iA}^j \Gamma Q_{jB}$	(8,3,2)
dim 6	$\overline{d}_R^A \Gamma L^i \overline{\psi}_{iA}^j \Gamma Q_{jB}$	(8,3,0)
dim 6	$\overline{Q}_L^{Ai} \Gamma L_j \overline{\psi}_{iA}^j \Gamma u_{RB}$	(8,3,0)
dim 6	$\overline{Q}_L^{Ai} \Gamma L_j \overline{\psi}_{iA}^j \Gamma d_{RB}$	(8,3,2)
dim6	$\overline{Q}_L^{Ai} \Gamma \ell \overline{\psi}_A^B \Gamma Q_{iB}$	(8,1,2)
dim 6	$\overline{u}_R^A \Gamma L^i \overline{\psi}_A^B \Gamma Q_{iB}$	(8,1,2)
dim 6	$\overline{d}_R^A \Gamma L^i \overline{\psi}_A^B \Gamma Q_{iB}$	(8,1,0)
dim 6	$\overline{Q}_L^{Ai} \Gamma L_i \overline{\psi}_A^B \Gamma u_{RB}$	(8,1,0)
dim 6	$\overline{Q}_L^{Ai} \Gamma L_i \overline{\psi}_A^B \Gamma d_{RB}$	(8,1,2)
dim6	$\overline{Q}_L^{iA} \Gamma \ell \overline{Q}^j \Gamma \psi_{ijAB}$	(6,3,8/3)
dim6	$\overline{u}_R^A \Gamma L^i \overline{Q}_j^B \Gamma \psi_{iAB}^j$	(6,3,8/3)
dim6	$\overline{d}_R^A \Gamma L^i \overline{Q}_j^B \Gamma \psi_{iAB}^j$	(6,3,2/3)
dim6	$\overline{Q}_L^{iA} \Gamma \ell \overline{Q}_j^B \Gamma \psi_{AB}$	(6,1,8/3)
dim6	$\overline{u}_R^A \Gamma L_i \overline{Q}^j \Gamma \psi_{AB}$	(6,1,8/3)
dim6	$\overline{d}_R^A \Gamma L_i \overline{Q}^j \Gamma \psi_{AB}$	(6,1,2/3)
dim 6	$\overline{Q}_L^{iA} \Gamma \ell \overline{Q}^{kB} \Gamma \psi_{ij}^C f_{ABC}$	(3,3,8/3)
dim6	$\overline{u}_R^A \Gamma L_i \overline{Q}^j \Gamma \psi_j^i f_{ABC}$	(3,3,8/3)
dim6	$\overline{d}_R^A \Gamma L_i \overline{Q}^j \Gamma \psi_j^i f_{ABC}$	(3,3,2/3)
dim6	$\overline{Q}_L^{iA} \Gamma \ell \overline{Q}_i^B \Gamma \psi^C f_{ABC}$	(3,1,8/3)
dim6	$\overline{u}_R^A \Gamma L_i \overline{Q}^i \Gamma \psi^C f_{ABC}$	(3,1,8/3)
dim6	$\overline{d}_R^A \Gamma L_i \overline{Q}^i \Gamma \psi^C f_{ABC}$	(3,1,2/3)
dim6	$\overline{Q}_L^{iA} \Gamma \ell \overline{\psi}_i^j \Gamma Q_{jA}$	(1,3,2)
dim6	$\overline{u}_R^A \Gamma L_i \overline{\psi}^{ij} \Gamma Q_{jA}$	(1,3,0)
dim6	$\overline{d}_R^A \Gamma L_i \overline{\psi}^{ij} \Gamma Q_{jA}$	(1,3,0)
dim6	$\overline{Q}_L^{iA} \Gamma L_j \overline{\psi}_i^j \Gamma u_{RA}$	(1,3,0)
dim6	$\overline{Q}_L^{iA} \Gamma L_j \overline{\psi}_i^j \Gamma d_{RA}$	(1,3,2)
dim6	$\overline{Q}_L^{iA} \Gamma \ell \overline{\psi} \Gamma Q_{iA}$	(1,1,2)
dim6	$\overline{u}_R^A \Gamma L^i \overline{\psi} \Gamma Q_{iA}$	(1,1,2)
dim6	$\overline{d}_R^A \Gamma L^i \overline{\psi} \Gamma Q_{iA}$	(1,1,0)
dim6	$\overline{Q}_L^{iA} \Gamma L_i \overline{\psi} \Gamma u_{RA}$	(1,1,0)
dim6	$\overline{Q}_L^{iA} \Gamma L_i \overline{\psi} \Gamma d_{RA}$	(1,1,2)

Dimension 6 fermionic operators containing QqL and QQ ℓ

4.1.3 operators of the form $q\ell\psi Q$, and $qL\psi q$

In this set of operators the LEX state ψ must be an SU(2) doublet

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 6	$\overline{u}_R^A \Gamma L^i \overline{\psi}_{iA}^B \Gamma u_{RB}$	(8,2,1)
dim 6	$\overline{u}_R^A \Gamma L^i \overline{\psi}_{iA}^B \Gamma d_{RB}$	(8,2,3)
dim6	$\overline{d}_R^A \Gamma L^i \overline{\psi}_{iA}^B \Gamma u_{RB}$	(8,2,-1)
dim6	$\overline{d}_R^A \Gamma L^i \overline{\psi}_{iA}^B \Gamma d_{RB}$	(8,2,1)
dim 6	$\overline{Q}_L^{Ai} \Gamma \ell \overline{\psi}_{iA}^B \Gamma u_{RB}$	(8,2,1)
dim 6	$\overline{Q}_L^{Ai} \Gamma \ell \overline{\psi}_{iA}^B \Gamma d_{RB}$	(8,2,3)
dim 6	$\overline{u}_R^A \Gamma \ell \overline{\psi}_A^{iB} \Gamma Q_{iB}$	(8,2,3)
dim 6	$\overline{d}_R^A \Gamma \ell \overline{\psi}_A^{iB} \Gamma Q_{iB}$	(8,2,1)
dim 6	$\overline{Q}_L^{Ai} \Gamma L_i \overline{\psi}_A^B \Gamma u_{RB}$	(8,1,0)
dim6	$\overline{u}_R^A \Gamma L_i \overline{u}_R^B \Gamma \psi_{AB}^i$	(6,2,11/3)
dim6	$\overline{u}_R^A \Gamma L_i \overline{d}_R^B \Gamma \psi_{AB}^i$	(6,2,5/3)
dim6	$\overline{d}_R^A \Gamma L_i \overline{d}_R^B \Gamma \psi_{AB}^i$	(6,2,-1/3)
dim6	$\overline{Q}_L^{iA} \Gamma \ell \overline{u}_R^B \Gamma \psi_{iAB}$	(6,2,11/3)
dim6	$\overline{Q}_L^{iA} \Gamma \ell \overline{d}_R^B \Gamma \psi_{iAB}$	(6,2,5/3)
dim6	$\overline{u}_R^A \Gamma \ell \overline{Q}^{iB} \Gamma \psi_{iAB}$	(6,2,11/3)
dim6	$\overline{d}_R^A \Gamma \ell \overline{Q}^{iB} \Gamma \psi_{iAB}$	(6,2,5/3)
dim6	$\overline{u}_R^A \Gamma L_i \overline{u}_R^B \Gamma \psi^{iC} f_{ABC}$	(3,2,11/3)
dim6	$\overline{u}_R^A \Gamma L_i \overline{d}_R^B \Gamma \psi^{iC} f_{ABC}$	(3,2,5/3)
dim6	$\overline{d}_R^A \Gamma L_i \overline{u}_R^B \Gamma \psi^{iC} f_{ABC}$	(3,2,5/3)
dim6	$\overline{d}_R^A \Gamma L_i \overline{d}_R^B \Gamma \psi^{iC} f_{ABC}$	(3,2,-1/3)
dim6	$\overline{u}_R^A \Gamma \ell \overline{Q}^{iB} \Gamma \psi_i^C f_{ABC}$	(3,2,11/3)
dim6	$\overline{d}_R^A \Gamma \ell \overline{Q}^{iB} \Gamma \psi_i^C f_{ABC}$	(3,2,5/3)
dim6	$\overline{Q}_L^{iA} \Gamma \ell \overline{u}_R^B \Gamma \psi_i^C f_{ABC}$	(3,2,11/3)
dim6	$\overline{Q}_L^{iA} \Gamma \ell \overline{d}_R^B \Gamma \psi_i^C f_{ABC}$	(3,2,5/3)
dim6	$\overline{u}_R^A L^i \overline{\psi}_i^j \Gamma u_{RjA}$	(1,2,1)
dim6	$\overline{u}_R^A L^i \overline{\psi}_i^j \Gamma d_{RjA}$	(1,2,3)
dim6	$\overline{d}_R^A L^i \overline{\psi}_i^j \Gamma u_{RjA}$	(1,2,-1)
dim6	$\overline{d}_R^A L^i \overline{\psi}_i^j \Gamma d_{RjA}$	(1,2,1)
dim6	$\overline{Q}_L^{iA} \Gamma \ell \overline{\psi}_i \Gamma u_{RA}$	(1,2,1)
dim6	$\overline{Q}_L^{iA} \Gamma \ell \overline{\psi}_i \Gamma d_{RA}$	(1,2,3)
dim6	$\overline{u}_R^A \Gamma \ell \overline{\psi}^i \Gamma Q_{iA}$	(1,2,3)
dim6	$\overline{d}_R^A \Gamma L^i \overline{\psi} \Gamma Q_{iA}$	(1,2,0)

Dimension 6 fermionic operators containing $Qq\ell$ and qqL

4.1.4 operators of the form $ql\psi q$

Finally we consider operators with two quarks where all of the SM fermions are right handed. In this case the LEX state must be an SU(2) singlet.

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 6	$\overline{u}_R^A \Gamma \ell \overline{\psi}_A^B \Gamma u_{RB}$	(8,1,2)
dim 6	$\overline{u}_R^A \Gamma \ell \overline{\psi}_A^B \Gamma d_{RB}$	(8,1,4)
dim6	$\overline{d}_R^A \Gamma \ell \overline{\psi}_A^B \Gamma u_{RB}$	(8,1,0)
dim6	$\overline{d}_R^A \Gamma \ell \overline{\psi}_A^B \Gamma d_{RB}$	(8,1,2)
dim6	$\overline{u}_R^A \Gamma \ell u_R^B \Gamma \psi_{AB}$	(6,1,14/3)
dim6	$\overline{d}_R^A \Gamma \ell d_R^B \Gamma \psi_{AB}$	(6,1,2/3)
dim6	$\overline{u}_R^A \Gamma \ell d_R^B \Gamma \psi_{AB}, \overline{d}_R^A \Gamma \ell u_R^B \Gamma \psi_{AB}$	(6,1,8/3)
dim6	$\overline{u}_R^A \Gamma \ell \overline{u}_R^B \Gamma \psi^C f_{ABC}$	(3,1,14/3)
dim6	$\overline{u}_R^A \Gamma \ell \overline{d}_R^B \Gamma \psi^C f_{ABC}, \overline{d}_R^A \Gamma \ell \overline{u}_R^B \Gamma \psi^C f_{ABC},$	(3,1,8/3)
dim6	$\overline{d}_R^A \Gamma \ell \overline{d}_R^B \Gamma \psi^C f_{ABC}$	(3,1,2/3)
dim6	$\overline{u}_R^A \ell \overline{\psi} \Gamma u_{RjA}$	(1,1,2)
dim6	$\overline{u}_R^A \ell \overline{\psi} \Gamma d_{RjA}$	(1,1,4)
dim6	$\overline{d}_R^A \ell \overline{\psi} \Gamma u_{RjA}$	(1,1,0)
dim6	$\overline{d}_R^A \ell \overline{\psi} \Gamma d_{RjA}$	(1,1,2)

Dimension 6 fermionic operators containing $qq\ell$

4.2 Operators with two leptons

We now enumerate the dimension 6 operators in the lepton-quark portal that contain two SM leptons. These operators are dimension 6 operators which contain a quark-lepton bilinear and another bi-linear consisting of the spin 1/2 LEX state ψ and a left or right handed SM lepton. In this case the color assignments of the LEX state must be a fundamental or antifundamental of SU(3). Since there is a single (anti)quark in the operators, we see that the LEX state must have baryon number 1/3. The operators further may contain either a lepton-anti lepton pair or they may contain two leptons. In the first case the LEX states will have lepton number 0. In the second case they will have lepton number -2. The SU(2) assignments of the LEX states will depend on the the left or right handedness of the SM fermions as we shall discuss.

4.2.1 operators of the form $QL\psi L$

We will first discuss the operators in that contain two leptons and in which all three SM fermions are left handed. There are 2 possible SU(2) charge assignments of the LEX state following the SU(2) tensor product rule

$$\mathbf{1}/2 \otimes \mathbf{1}/2 \otimes \mathbf{1}/2 = \mathbf{3}/2 \oplus \mathbf{1}/2.$$

The possible SU(2) representation of ψ the quadruplet and the doublet. Operators are given in the table below.

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 6	$\overline{Q_{Li}{}^A} \Gamma L_j \psi_A^{ijk} \Gamma L_k$	(3,4,7/3)
dim 6	$\overline{Q_L}{}^{iA} \Gamma L_i \overline{\psi}_A^j \Gamma L_j, \overline{Q_L}{}^{iA} \Gamma L_j \overline{\psi}_A^j \Gamma L_i$	(3,2,7/3)
dim 6	$\overline{Q_L}{}^{iA} \Gamma L_j \overline{\psi}_{Ai} \Gamma L^j$	(3,2,7/3)
dim 6	$\overline{Q_{Li}{}^A} \Gamma L_j \overline{L}_k \Gamma \psi_A^{ijk}$	(3,4,1/3)
dim 6	$\overline{Q_L}{}^{iA} \Gamma L_i \overline{L}^j \Gamma \psi_{jA}, \overline{Q_L}{}^{iA} \Gamma L_j \overline{L}^j \Gamma \psi_{iA}$	(3,2,1/3)
dim 6	$\overline{Q_L}{}^{iA} \Gamma L_j \overline{L}_i \Gamma \psi_A^j$	(3,2,1/3)

Dimension 6 fermionic operators containing QLL

4.2.2 operators of the form $Q\ell\psi L$ and $qL\psi L$

We will now discuss the four fermion operators of the lepton-quark portal which have two left handed SM fermions. In this case the available SU(2) charges of the LEX states follow the SU(2) tensor product rule $1/2 \otimes 1/2 = 0 \oplus 1$, that is the LEX states are SU(2) singlets or triplets. Again the LEX states in these operators must also be fundamental of SU(3).

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 6	$\overline{Q_L}{}^{iA} \Gamma \ell \overline{\psi}_{iA} \Gamma L_j, \overline{Q_L}{}^{iA} \Gamma L_j \overline{\psi}_i^j \Gamma \ell$	(3,3,10/3)
dim6	$\overline{Q_L}{}^{iA} \Gamma \ell \overline{L}_j \Gamma \psi_j^i$	(3,3,4/3)
dim6	$\overline{Q_L}{}^{iA} \Gamma L_j \overline{\ell} \Gamma \psi_j^i$	(3,3,-2/3)
dim 6	$\overline{u_R}{}^A \Gamma L_i \overline{\psi}_{Aj} \Gamma L^j$	(3,3,10/3)
dim 6	$\overline{d_R}{}^A \Gamma L_i \overline{\psi}_{Aj} \Gamma L^j$	(3,3,4/3)
dim 6	$\overline{u_R}{}^A \Gamma L_i \overline{L}^j \Gamma \psi_{jA}^i$	(3,3,4/3)
dim 6	$\overline{d_R}{}^A \Gamma L_i \overline{L}^j \Gamma \psi_{jA}^i$	(3,3,-2/3)
dim 6	$\overline{Q_L}{}^{iA} \Gamma \ell \overline{\psi}_A \Gamma L_i, \overline{Q_L}{}^{iA} \Gamma L_i \overline{\psi} \Gamma \ell$	(3,1,10/3)
dim6	$\overline{Q_L}{}^{iA} \Gamma \ell \overline{L}_i \Gamma \psi$	(3,1,4/3)
dim6	$\overline{Q_L}{}^{iA} \Gamma L_i \overline{\ell} \Gamma \psi$	(3,1,-2/3)
dim 6	$\overline{u_R}{}^A \Gamma L_i \overline{\psi}_A \Gamma L^i$	(3,1,10/3)
dim 6	$\overline{d_R}{}^A \Gamma L_i \overline{\psi}_A \Gamma L^i$	(3,1,4/3)
dim 6	$\overline{u_R}{}^A \Gamma L_i \overline{L}^j \Gamma \psi_A$	(3,1,4/3)
dim 6	$\overline{d_R}{}^A \Gamma L_i \overline{L}^j \Gamma \psi_A$	(3,1,-2/3)

Dimension 6 fermionic operators that contain QL ℓ and qLL

4.2.3 operators of the form $q\ell\psi L$ and $Q\ell\psi\ell$

We will now consider operators where only one of the SM fermions in the four fermion interaction is a SM weak doublet. In this case the LEX field must be an SU(2) doublet.

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 6	$\overline{Q}_L^{iA}\Gamma\ell\overline{\psi}_{iA}\Gamma\ell$	(3,2,13/3)
dim6	$\overline{Q}_L^{iA}\Gamma\ell\overline{\ell}\Gamma\psi_{iA}$	(3,2,1/3)
dim 6	$\overline{u}_R^A\Gamma\ell\overline{\psi}_A^i\Gamma L_i$	(3,2,13/3)
dim 6	$\overline{u}_R^A\Gamma\overline{\ell}^i\Gamma\psi_{iA}$	(3,2,7/3)
dim 6	$\overline{d}_R^A\Gamma\ell\overline{\psi}_A^i\Gamma L_i$	(3,3,7/3)
dim 6	$\overline{d}_R^A\Gamma\overline{\ell}^i\Gamma\psi_{iA}$	(3,3,1/3)

Dimension 6 fermionic operators that contain $Q\ell\ell$ or $q\ell L$

4.2.4 operators of the form $q\ell\psi\ell$

We now enumerate the 4 fermion operators of the lepton-quark portal where the LEX fermion is an SU(2) singlet and all SM fermions in the operators are SU(2) singlets.

dimension	Operators with scalar LEX fields (CP even)	(SU(3),SU(2),Y)
dim 6	$\overline{u}_R^A\Gamma\ell\overline{\psi}_A\Gamma\ell$	(3,1,16/3)
dim6	$\overline{u}_R^A\Gamma\ell\overline{\ell}\Gamma\psi_A$	(3,1,4/3)
dim 6	$\overline{d}_R^A\Gamma\ell\overline{\psi}_A\Gamma\ell$	(3,1,10/3)
dim 6	$\overline{d}_R^A\Gamma\ell\overline{\ell}\Gamma\psi_A$	(3,3,-2/3)

Dimension 6 fermionic operators than contain $q\ell\ell$

4.3 Some Collider Phenomenology of the Fermionic Operators


The fermionic operators list contains a large number of interesting states that can be accessed through the quark-lepton portal. Some of these objects are extended types of lepto-quarks with higher baryon and lepton number. Some of them are other types of exotic states as we will see. Among the fields in the two quark operator list are fermionic objects that couple to a lepton and a $\bar{q}q$ pair. These objects have zero baryon number but they do carry unit lepton number. The fermions in the 8 and 1 representations fall into this category. These are not lepto-quarks, as they carry no baryon number, but are an interesting type of exotic states. The fermionic color octet with lepton number 1 and baryon number 0 is a type of exotic lepto-gluon [5] which may carry highly nontrivial SU(2) charge. The a fermionic color singlets with lepton number 1 and zero baryon number are exotic leptons which can couple to the SM through the

lepton quark portal. Conversely, the objects that appear in operators with a quarks and one lepton carry baryon number $2/3$ and unit lepton number. These are the fermionic LEX states in the 6 and 3 representation of $SU(3)$. These are exotic states with another unusual baryon and lepton number assignment.

The objects that couple to twos lepton and one quark are always in the fundamental representation of $SU(3)$. Some of these objects that couple to $\bar{l}l$ pairs and one quark, are just themselves, exotic quarks with zero lepton number and baryon number $1/3$. The lepto-quark portal operators allow these exotics to be accessed through new dimension 6 operators. The fermionic LEX states that are $SU(3)$ fermions that couple to $\ell\ell$ pairs and a single quark are an interesting type of object with baryon number $1/3$ but lepton number 2. This is an interesting bilepto-quark object.

We can see that some of the LEX states have very interesting SM charge assignments. For example, there fermionic color octet and color sextet $SU(2)$ quadruplets. There are also objects with large hyper charges. Many operator contain a coupling to two leptons(as opposed to a lepton anti-lepton pair) and one quark. For example the $\bar{u}_R^A \Gamma \ell \bar{\psi}_A \Gamma \ell$ operator contains a LEX state with charges $(3,1,16/3)$, a color triplet fermionic object with electric charge $8/3$.

As we have notes, the fermionic operators contain various four particle couplings that fall into two classes. The first class are couplings between two quarks, a lepton, and a LEX state. The second type are four particle interactions between to leptons, a quark and a LEX state. We visualize these interactions in the diagrams below.




Four point vertices of the fermionic lepton-quark portal operators

We first note that the above four point operators allow interesting dimension 6 decay processes for the various LEX states. For example, color sextet or triplet LEX states may decay via a three body process to a di-quark plus lepton resonance $\psi \rightarrow qql$ involving two quarks or two antiquarks. Color octet or singlet states may decay to a quark-antiquark and lepton $\psi \rightarrow \bar{q}ql$. Even more striking, exotic color triplets may decay to two leptons and a quark. LEX states with lepton number zero may decay to a lepton anti-lepton pair $\psi \rightarrow \bar{l}lq$. LEX states with lepton number 2 may decay to a same sign di-lepton plus quark resonance $\psi \rightarrow llq$.


At e-p colliders like LHeC, we see that these dimension 6 operators lead the the direct production of LEX extended lepto-quarks. in association with a

quark or lepton through the processes $ql \rightarrow q + \psi$ and $q \rightarrow l + \psi$. If the exotic states decay back through the dimension 6 operators, there could be extremely interesting multilepton processes. For example, if we consider the $llq\psi$ operators we may have a production and decay process $\bar{q}l \rightarrow \bar{l}\psi \rightarrow \bar{l}(llq)$ at LHeC. There are several 2-2 and 2-3 LHC processes that could singly produce an exotic state. The $qql\psi$ type operators can lead to the single production of a LEX state in association with a hard lepton from quark fusion through the process $qq \rightarrow l + \psi$ as in the diagram below



LHC production of lepton plus fermionic LEX state from quark fusion

There are also interesting 2-3 production processes at LHC that might follow from the new operators such as a quark gluon fusion process the diagram below



LHC production of fermionic LEX state inassociation with two leptons from quark gluon fusion

Here a quark and gluon fuse into an off-shell quark which produces a final state with two leptons and a LEX state. The process is $gq \rightarrow q \rightarrow ll\psi$. If the ψ state decays through the same operator, there will be a final state with four final state leptons. Though this process may have a smaller cross section as it is a 2-3 process and results from a dimension 6 operator, it may still stand out as a new physics channel due to the possibility of a multi-lepton(and perhaps same sign lepton) final state.

We finally note that the unusual $llq\psi$ operators may have substantial production cross section at future muon colliders. If there is an allowed lepton fusion channel, a single LEX fermion may be produced in association with a quark from a process $\bar{l}l \rightarrow q + \psi$. This is interesting because of it is a hadronic signature, and involves the production of quite unusual exotics states.

5 Vector LEX states

We briefly mention that spin 1 vector LEX states are also accessible in the lepton-quark portal. The Higgs type, di-Higgs type, and field strength tensor type operators involving the vector LEX state are very similar to the scalar operators. We can refer to the vector LEX state as V^μ . Then, for example, the dimension 5 Higgs type operators has structure $\overline{f_1}\gamma^\mu f_2 V_\mu H$. This structure is very similar to the scalar LEX states, but with a vector current substituted for the scalar current. The charge assignments of the vector LEX states will be the same as the scalar LEX states in Higgs type operators. Similarly the dimension 6 di-Higgs operators have the structure $\overline{f_1}\gamma^\mu f_2 V_\mu HH$ again with the same charge assignments as the scalar LEX states in analogous operators.

LEX vectors can also couple to SM field strength tensors through dimension six operators of the form $\overline{f_1}\gamma^\mu f_2 F_{\mu\nu} V^\nu$. This structure is very similar to the scalar operators with a vector current substituted for the fermion tensor current. Again the LEX charge assignments are the same as the scalar LEX field strength operators. Thus the appropriate substitution can be done in the scalar tables to get the quantum numbers and couplings of the vector states.

6 Conclusions

We have produced an operator catalog for the lepton-quark portal. We have written down operators for spin 0 and spin 1/2 states up to dimension 6 in which a single light exotic state couples to the standard model through operators involving at least one quark and at least one lepton. We have found that the quark-lepton portal allows us to singly produce interesting exotic fields. These are states may be in unusual larger dimensional representations of both the SM color, and weak gauge groups. For example, we show that we can access 15plets of SU(3), exotic color sextets, a variety of unusual color adjoints, and states in quintuplet and quadruplets representation of SU(2). We have show that these states can have simultaneous interesting lepton and baryon number, including states with baryon number 2/3 and lepton number 1, states with lepton number 2 and baryon number 1/3, color charged objects with baryon number 0 and unit lepton number, new types of lepto quarks with baryon number 1/3 and lepton number 1, and exotic leptons with unusual hypercharges. We have shown relation between operators in the portal so we can present the operator catalog without redundancy of operators. We have reviewed some of the most promising collider production processes at LHC, at e-p colliders like LHeC and even a few muon collider production processes.

There are many possible new directions that could evolve from this work. It would be interesting to complete collider studies for some of the more stand out signatures of LEX states explored in this paper. Most interesting would be HL-LHC collider studies of operators that yield multi-lepton and hard photon final states. It would be interesting to peruse lepton collider studies for some of the exotic lepton-quark model portals as they would provide an interesting

class of models with hadronic signatures at lepton colliders. It would also be interesting to peruse studies of the the exotic 15-plets of $SU(3)$ as these states are in a quite unusual $SU(3)$ representation and would be produced with large Clebsch-Gordon coefficients which raise the production cross section.

It would be worth while to peruse the operator catalog and identify some of the dimension 5 and 6 operators that could provide alternate production and decay channels for the standard leptoquarks which usually appear in dimension 4 interactions. There may be interesting regions of their parameter space where the effective operators give alternate discovery or constraint channels. The lepton-quark portal tells us that dozens of LEX states are accessible through quark-lepton interactions. As these LEX states have varied SM quantum numbers, it would also be useful to study which phenomenological discriminants could identify and discriminate between the models, if an excess was found.

It would also be interesting to peruse some model building with theories of exotic states in higher dimensional representations of $SU(2)$, the quadruplets and quintuplets. It would be enlightening to study the Higgs coupling operators which could split the masses of the charged fields within these multiplets, and it would be worthwhile to study some of the decay chains of the charged states, especially if they contain longer lived particles. These models would also yield interesting constrains from Higgs sector measurements as Higgs couplings to colored scalars are constrained by loop level enhancements to Higgs di-boson couplings.[31][32]. Finally another extension of this work would be to build UV completions for some of the operators in this catalog. , especially the states in higher dimensional representations of the SM gauge groups.

References

- [1] L. M. Carpenter, T. Murphy, and T. M. P. Tait, *Phenomenological cornucopia of $SU(3)$ exotica*, *Phys. Rev. D* **105** (2022), no. 3 035014, [[arXiv:2110.11359](#)].
- [2] L. M. Carpenter, M. J. Smylie, J. M. C. Ramirez, C. McDowell, and D. Whiteson, *New physics in triboson event topologies*, *Phys. Rev. D* **105** (2022), no. 7 075027, [[arXiv:2112.00137](#)].
- [3] L. M. Carpenter, T. Murphy, and M. J. Smylie, *Changing patterns in electroweak precision fits with new color-charged states: Oblique corrections and the W -boson mass*, *Phys. Rev. D* **106** (2022), no. 5 055005, [[arXiv:2204.08546](#)].
- [4] V. Miralles and A. Pich, *LHC bounds on colored scalars*, *Phys. Rev. D* **100** (2019), no. 11 115042, [[arXiv:1910.07947](#)].
- [5] A. Çelikel, M. Kantar, and S. Sultansoy, *A search for sextet quarks and leptogluons at the LHC*, *Phys. Lett. B* **443** (1998), no. 1 359–364.

- [6] K. Blum, A. Efrati, C. Frugiuele, and Y. Nir, *Exotic colored scalars at the LHC*, *JHEP* **02** (2017) 104, [[arXiv:1610.06582](#)].
- [7] T. Han, I. Lewis, and Z. Liu, *Colored Resonant Signals at the LHC: Largest Rate and Simplest Topology*, *JHEP* **12** (2010) 085, [[arXiv:1010.4309](#)].
- [8] T. Han, I. M. Lewis, H. Liu, Z. Liu, and X. Wang, *A guide to diagnosing colored resonances at hadron colliders*, *JHEP* **08** (2023) 173, [[arXiv:2306.00079](#)].
- [9] S. Biondini, T. A. Chowdhury, and S. Saad, *Non-perturbative effects for dark sectors with QCD portals*, *JHEP* **08** (2023) 002, [[arXiv:2306.09428](#)].
- [10] L. M. Carpenter and T. Murphy, *Dark matter with exotic mediators: The diquark portal*, [arXiv:2312.17607](#).
- [11] L. M. Carpenter and K. Schwind, *Unusual portals to new exotics, the W-gluon portal*, [arXiv:2410.08195](#).
- [12] L. M. Carpenter, T. Murphy, and M. J. Smylie, *LEX-EFT: the Light Exotics Effective Field Theory*, *JHEP* **08** (2023) 050, [[arXiv:2302.01344](#)].
- [13] **LHeC, FCC-he Study Group** Collaboration, P. Agostini et al., *The Large Hadron–Electron Collider at the HL-LHC*, *J. Phys. G* **48** (2021), no. 11 110501, [[arXiv:2007.14491](#)].
- [14] M. Narain et al., *The Future of US Particle Physics - The Snowmass 2021 Energy Frontier Report*, [arXiv:2211.11084](#).
- [15] W. Buchmüller, R. Ratz, and D. Wyler, *Leptoquarks in lepton-quark collisions*, *Phys. Lett. B* **191** (1987), no. 4 442–448.
- [16] H. Murayama and T. Yanagida, *Viable $su(5)$ gut with light leptoquark bosons*, *Mod. Phys. Lett. A* **07** (1992), no. 02 147–151.
- [17] **Particle Data Group** Collaboration, S. Navas et al., *Review of particle physics*, *Phys. Rev. D* **110** (2024), no. 3 030001.
- [18] **D0** Collaboration, V. M. Abazov et al., *Search for pair production of first-generation leptoquarks in p anti- p collisions at $s^{1/2} = 1.96$ -TeV*, *Phys. Lett. B* **681** (2009) 224–232, [[arXiv:0907.1048](#)].
- [19] **CDF** Collaboration, D. Acosta et al., *Search for first-generation scalar leptoquarks in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV*, *Phys. Rev. D* **72** (2005) 051107, [[hep-ex/0506074](#)].
- [20] B. Grzadkowski, M. Iskrzynski, M. Misiak, and J. Rosiek, *Dimension-six terms in the Standard Model Lagrangian*, *J. High Energy Phys.* **10** (Oct, 2010) 085.

- [21] R. Slansky, *Group Theory for Unified Model Building*, *Phys. Rept.* **79** (1981) 1–128.
- [22] S. Coleman, *Fun with $SU(3)$* , in *Seminar on high-energy physics and elementary particles*, pp. 331–352, 1965.
- [23] E. C. F. S. Fortes, K. S. Babu, and R. N. Mohapatra, *Flavor Physics Constraints on TeV Scale Color Sextet Scalars*, in *6th International Workshop on Charm Physics*, 11, 2013. [arXiv:1311.4101](#).
- [24] C.-R. Chen, W. Klemm, V. Rentala, and K. Wang, *Color Sextet Scalars at the CERN Large Hadron Collider*, *Phys. Rev. D* **79** (2009) 054002, [[arXiv:0811.2105](#)].
- [25] L. M. Carpenter, K. Schwind, and T. Murphy, *Leptonic signatures of color-sextet scalars. II. Exploiting unique large- ET_{miss} signals at the LHC*, *Phys. Rev. D* **109** (2024), no. 7 075010, [[arXiv:2312.09273](#)].
- [26] L. M. Carpenter, T. Murphy, and K. Schwind, *Leptonic signatures of color-sextet scalars*, *Phys. Rev. D* **106** (2022), no. 11 115006, [[arXiv:2209.04456](#)].
- [27] L. M. Carpenter, T. Murphy, and T. M. P. Tait, *Distinctive signals of frustrated dark matter*, *JHEP* **09** (2022) 175, [[arXiv:2205.06824](#)].
- [28] C. W. Murphy, *Dimension-8 operators in the Standard Model Effective Field Theory*, *JHEP* **10** (2020) 174, [[arXiv:2005.00059](#)].
- [29] L. Lehman, *Extending the Standard Model Effective Field Theory with the Complete Set of Dimension-7 Operators*, *Phys. Rev. D* **90** (2014), no. 12 125023, [[arXiv:1410.4193](#)].
- [30] B. Field, *Distinguishing scalar from pseudoscalar Higgs production at the CERN LHC*, *Phys. Rev. D* **66** (2002) 114007, [[hep-ph/0208262](#)].
- [31] J. Cao, P. Wan, J. M. Yang, and J. Zhu, *The SM extension with color-octet scalars: diphoton enhancement and global fit of LHC Higgs data*, *JHEP* **08** (2013) 009, [[arXiv:1303.2426](#)].
- [32] R. Boughezal and F. Petriello, *Precise predictions for Higgs production in models with color-octet scalars*, *Nucl. Phys. B Proc. Suppl.* **205-206** (2010) 289–294, [[arXiv:1006.5773](#)].