

# REALIZING MODULAR DATA FROM CENTERS OF NEAR-GROUP CATEGORIES

ZHIQIANG YU<sup>1</sup> AND QING ZHANG<sup>2</sup>

ABSTRACT. In this paper, we show the existence of a near-group category of type  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} + 16$  and compute the modular data of its Drinfeld center. We prove that a modular data of rank 10 can be obtained through condensation of the Drinfeld center of the near-group category  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} + 16$ . Moreover, we compute the modular data for the Drinfeld center of a near-group category  $\mathbb{Z}/8\mathbb{Z} + 8$  and show that the non-pointed factor of its condensation has the same modular data as the quantum group category  $\mathcal{C}(\mathfrak{g}_2, 4)$ .

## 1. INTRODUCTION

Modular categories over  $\mathbb{C}$  are non-degenerate ribbon fusion categories over  $\mathbb{C}$ . These categories originally appeared in conformal field theory [13] and were formally defined in [27] to study (2+1)-dimensional topological quantum field theories. Algebraically, modular categories arise as representation categories of quantum groups, von Neumann algebras, vertex operator algebras, and local conformal nets.

One of the central problems in the study of modular categories is their classification. As shown in [4], there are finitely many classes of modular tensor categories of a given rank (i.e., the number of isomorphism classes of simple objects) up to tensor equivalence. This finiteness property motivates the rank classification of modular categories. The classification of low-rank modular categories has been achieved in [22, 3, 15, 16], at least up to their modular data.

A key invariant used in the classification of modular categories is their modular data, which is a pair of  $S$  and  $T$  matrices. Given any modular category  $\mathcal{C}$ , the modular data defines a projective representation of the modular group  $\mathrm{SL}(2, \mathbb{Z})$ . The kernel of this projective representation is a congruence subgroup of level  $N$  thus factors through  $\mathrm{SL}(2, \mathbb{Z}/N\mathbb{Z})$ , where  $N$  is the order of the  $T$  matrix [17]. Furthermore, this projective representation can be lifted to linear representations, which also have congruence kernels [7].

In [15], a method for classifying modular categories was introduced by constructing modular data directly from the irreducible representations of  $\mathrm{SL}(2, \mathbb{Z}/n\mathbb{Z})$  for certain integers  $n$ . This technique was successfully applied in [15] to classify rank 6 modular categories up to their modular data. The classification was further extended in [16], where all modular data up to rank 11 are explicitly constructed.

One of the main motivations of this paper is to find a modular category that realize a rank 10 modular data constructed in [16]. It was conjectured in [16] that this modular data could be realized through the condensation of the Drinfeld center of a near-group category of type  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} + 16$ , provided such a category exists.

To achieve this, we first prove the existence of such near-group category using results from [11, 10]. The next step is deriving the modular data for the center of  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} + 16$ , a category of rank 304. One of the main challenges is to solve a large system of non-linear

equations in [11]. Through this process, we obtain the explicit modular data of the center. Within the center of this near-group category, there exists a Tannakian fusion subcategory  $\mathcal{E} = \text{Rep}(\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z})$ . We show that the condensation processed by  $\mathcal{E}$  yields a modular category of rank 10. By further analyzing the congruence representations of  $\text{SL}(2, \mathbb{Z})$ , we identify the resulting rank 10 modular data with that from [16]. Explicitly, we have the following result:

**Theorem.** *Let  $\mathcal{C}$  be a near-group category of type  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} + 16$  associated with the symmetric bi-character  $\langle (g_1, g_2), (h_1, h_2) \rangle = (\zeta_4)^{g_1 h_1 - g_2 h_2}$ ,  $g_1, g_2, h_1, h_2 \in \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ . Then  $\mathcal{Z}(\mathcal{C})$  contains a Tannakian fusion subcategory  $\mathcal{E} = \text{Rep}(G)$ , where  $G = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ . Moreover, the condensation  $\mathcal{Z}(\mathcal{C})_G^0$  has the same modular data as in Equation (12).*

Several low-rank modular categories can be constructed using similar methods, such as by factorization and/or condensation of the centers of near-group categories. See [16, 23] for examples of verified cases and conjectures. However, it remains unclear whether some of these categories can be realized through quantum group categories, vertex operator algebras or their condensations.

To verify a conjecture from [23, Section 4], we compute the center of a near-group category of type  $\mathbb{Z}/8\mathbb{Z} + 8$ . The existence of this fusion category was established in [10]. By computing and analyzing the modular data of the center, we obtain the following result:

**Theorem.** *Let  $\mathcal{C}$  be a near-group category of type  $\mathbb{Z}/8\mathbb{Z} + 8$ . Then  $\mathcal{Z}(\mathcal{C})$  contains a Tannakian subcategory  $\mathcal{E} = \text{Rep}(\mathbb{Z}/2\mathbb{Z})$  such that there is a braided tensor equivalence  $\mathcal{Z}(\mathcal{C})_{\mathbb{Z}/2\mathbb{Z}}^0 \cong \mathcal{C}(\mathbb{Z}/4\mathbb{Z}, q) \boxtimes \mathcal{D}$ , where  $\mathcal{C}(\mathbb{Z}/4\mathbb{Z}, q)$  is a pointed modular category and  $\mathcal{D}$  has the same modular data as  $\mathcal{C}(\mathfrak{g}_2, 4)$ .*

The paper is organized as follows. In Section 2, we review some basic notions of modular categories. In Section 3, we first show the existence of a near-group category  $\mathcal{C}$  of type  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} + 16$  (Proposition 3.1.1). We then compute the modular data of  $\mathcal{Z}(\mathcal{C})$ , and prove that the condensation by Tannakian fusion category  $\text{Rep}(\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z})$  matches the modular data in [16] (Theorem 3.2.6). Finally, we compute the Drinfeld center of a near-group category of type  $\mathbb{Z}/8\mathbb{Z} + 8$  and obtain the modular data for the non-pointed factor from the condensation of the center (Theorem 3.3.3).

Throughout this paper, we use the notation  $\mathbb{T} = \{x \in \mathbb{C}; |x| = 1\}$ ,  $\zeta_n = \exp(2\pi i/n)$  and  $\chi_n^m = m + \sqrt{n}$ .

## 2. PRELIMINARY

In this section, we recall some basic notions on braided fusion categories. For further details, readers are referred to [1, 9, 11, 10].

**2.1. Modular data.** Let  $\mathcal{C}$  be a braided fusion category with a braiding  $c$ . We use  $\mathcal{O}(\mathcal{C})$  to denote the set of isomorphism classes of simple objects in  $\mathcal{C}$ . Let  $\mathcal{D}$  be a braided fusion subcategory of  $\mathcal{C}$ , the Müger centralizer  $\mathcal{D}'_{\mathcal{C}}$  of  $\mathcal{D}$  in  $\mathcal{C}$  is the fusion subcategory generated by objects  $X$  of  $\mathcal{C}$  such that  $c_{Y,X}c_{X,Y} = \text{id}_{X \otimes Y}$  for all  $Y$  of  $\mathcal{D}$  [14]. The Müger center of  $\mathcal{C}$  is the fusion subcategory  $\mathcal{C}' = \mathcal{C}'_{\mathcal{C}}$ . A braided fusion category  $\mathcal{C}$  is *symmetric* if  $\mathcal{C}' = \mathcal{C}$ .

A ribbon fusion category  $\mathcal{C}$  is called *modular*, if its Müger center is equivalent to  $\text{Vec}$ , the category of finite-dimensional vector space over  $\mathbb{C}$ . Equivalently,  $\mathcal{C}$  is modular if the  $S$ -matrix is non-degenerate [14], where  $S = \text{Tr}(c_{Y,X^*}c_{X^*,Y})$ ,  $X, Y \in \mathcal{O}(\mathcal{C})$ . Let  $T = (\delta_{X,Y}\theta_X)$ , where  $\theta_X$

is the scalar associated with the ribbon twist of  $X \in \mathcal{O}(\mathcal{C})$ . The pair of  $(S, T)$  is referred as the *modular data*. Let  $\mathfrak{s}, \mathfrak{t} \in \mathrm{SL}(2, \mathbb{Z})$  be the generators given by  $\mathfrak{s} := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$  and  $\mathfrak{t} := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ . For a modular category, the assignment  $\mathfrak{s} \mapsto S, \mathfrak{t} \mapsto T$  gives a projective representation of  $\mathrm{SL}(2, \mathbb{Z})$ . Let  $N$  be the order of the  $T$  matrix. The kernel of this projective representation is a congruence subgroup of  $\mathrm{SL}(2, \mathbb{Z})$  at level  $N$ . Consequently, the image factors through the finite group  $\mathrm{SL}(2, \mathbb{Z}/N\mathbb{Z})$  [17]. This projective representations can be lifted to linear representations, which also have congruence kernels [7]. The congruence kernel results have been used in proving key results such as rank finiteness of modular categories [4], and in the classifications of modular categories [3, 15, 16, 18, 21].

Let  $\rho$  be an arbitrary irreducible finite-dimensional congruence representation of  $\mathrm{SL}(2, \mathbb{Z})$  of level  $n$ , where  $n$  is a positive integer. By the Chinese Remainder Theorem,  $\rho$  factors through the finite groups

$$\mathrm{SL}(2, \mathbb{Z}/n\mathbb{Z}) \cong \mathrm{SL}(2, \mathbb{Z}/p_1^{n_1}\mathbb{Z}) \times \cdots \times \mathrm{SL}(2, \mathbb{Z}/p_r^{n_r}\mathbb{Z}),$$

where  $n = p_1^{n_1} \cdots p_r^{n_r}$ , and  $p_j$  are distinct primes,  $1 \leq j \leq r$ . The finite-dimensional irreducible representations of  $\mathrm{SL}(2, \mathbb{Z}/p^m\mathbb{Z})$  are classified in [19, 20].

Let  $\mathcal{C}$  be a ribbon fusion category with  $S$ - and  $T$ - matrix. For any simple objects  $X$  and  $Y$ , we have the balancing equation [1]

$$\theta_X \theta_Y S_{X,Y} = \sum_{Z \in \mathcal{O}(\mathcal{C})} N_{X^*,Y}^Z \theta_Z d_Z,$$

where  $N_{X^*,Y}^Z = \dim \mathrm{Hom}(X^* \otimes Y, Z)$  is the fusion coefficient and  $d_Z$  is the quantum dimension of the simple object  $Z$ . If  $\mathcal{C}$  is modular, the fusion coefficients can be expressed in terms of the  $S$ -matrix via the Verlinde formula (see, for example, [1])

$$N_{X^*,Y}^Z = \frac{1}{\dim(\mathcal{C})} \sum_{W \in \mathcal{O}(\mathcal{C})} \frac{S_{X,W} S_{Y,W} S_{Z^*,W}}{S_{\mathbf{1},W}},$$

where  $\mathbf{1}$  denotes the tensor unit of  $\mathcal{C}$ .

Let  $\mathcal{C}$  be modular with modular data  $(S, T)$ . Denote by  $\mathbb{Q}(S)$  the smallest field containing all entries of the  $S$ -matrix. It is known that  $\mathbb{Q}(S)$  is an abelian Galois extension over  $\mathbb{Q}$  [6, 5]. Let  $\mathbb{Q}_N := \mathbb{Q}(\zeta_N)$ , where  $\zeta_N$  is a primitive  $N$ th root of unity. It is proved in [17] that  $\mathbb{Q}(S) \subset \mathbb{Q}(T) = \mathbb{Q}_N$ , where  $N$  is the order of the  $T$  matrix. Let  $\mathrm{Gal}(\mathcal{C}) = \mathrm{Gal}(\mathbb{Q}(S)/\mathbb{Q})$ . The assignments  $\chi_Y : X \mapsto \frac{S_{X,Y}}{S_{\mathbf{1},Y}}$  define linear characters of the fusion ring  $\mathcal{K}_0(\mathcal{C})$ . Since  $S$  is non-degenerate,  $\{\chi_Y\}_{Y \in \mathcal{O}(\mathcal{C})}$  forms all characters of  $\mathcal{K}_0(\mathcal{C})$ . For any  $\sigma \in \mathrm{Gal}(\mathcal{C})$ ,  $\sigma(\chi_Y)$  is also a linear character of  $\mathcal{K}_0(\mathcal{C})$ . Thus, there is a unique  $\hat{\sigma} \in \mathrm{Sym}(\mathcal{O}(\mathcal{C}))$  such that

$$\sigma \left( \frac{S_{X,Y}}{S_{\mathbf{1},Y}} \right) = \frac{S_{X,\hat{\sigma}(Y)}}{S_{\mathbf{1},\hat{\sigma}(Y)}}$$

for all  $X, Y \in \mathcal{O}(\mathcal{C})$ . For more details on the Galois action on modular data, see [4, 7] and the references therein.

A fusion category  $\mathcal{C}$  is called *pointed* if all of its simple objects are invertible. In this case, the set of isomorphism classes of simple objects  $\mathcal{O}(\mathcal{C})$  forms a finite group  $G$ . Given a braided pointed fusion category  $\mathcal{C}$  with  $\mathcal{O}(\mathcal{C}) = G$ , the braiding  $c$  naturally determines a quadratic form  $q$  on  $G$ , given by  $q(g) = c_{g,g}$ . The pair  $(G, q)$  is called a *pre-metric group*.

In fact, there is a bijective correspondence between pre-metric groups and ribbon pointed fusion categories [8]. We denote by  $\mathcal{C}(G, q)$  the ribbon pointed fusion category associated with the pre-metric group  $(G, q)$ . A pre-metric group  $(G, q)$  is called *metric* if the quadratic form  $q$  is non-degenerate, that is, the bi-character  $\langle \cdot, \cdot \rangle$  determined by  $q$  is non-degenerate, where  $\langle g, h \rangle = \frac{q(g+h)}{q(g)q(h)}$ ,  $g, h \in G$ . In this case, the corresponding fusion category is modular.

**2.2. Near-group categories.** Let  $G$  be a finite abelian group of order  $n$ , and let  $m$  be a non-negative integer. A *near-group category* of type  $G + m$  [24] is a fusion category with simple objects labeled by elements  $g \in G$  with an additional simple object  $\rho$  such that the fusion rules are generated by the multiplication in  $G$ , with  $g\rho = \rho g = \rho$  for all  $g \in G$ , and  $\rho^{\otimes 2} = m\rho \oplus_{g \in G} g$ . As shown in [10, Theorem 2], the only possible values of  $m$  are  $n$ ,  $n - 1$  or  $m = kn$  for some nonnegative integer  $k$ . When  $m = 0$ , the corresponding near-group categories are called the Tambara-Yamagami categories, which are completely classified in [25]. When  $m = n$ , the near-group categories  $G + n$  are characterized by the following:

**Theorem 2.2.1.** [11, Theorem 5.3], [10, Corollary 5] *Let  $G$  be a finite abelian group of order  $n$ ,  $\langle \cdot, \cdot \rangle$  a non-degenerate symmetric bicharacter on  $G$  and define  $d := \frac{n + \sqrt{n^2 + 4n}}{2}$ . Let  $c \in \mathbb{T}$ ,  $a : G \rightarrow \mathbb{T}$ ,  $b : G \rightarrow \mathbb{C}$  be such that*

$$a(0) = 1, \quad a(x) = a(-x), \quad a(x+y)\langle x, y \rangle = a(x)a(y), \quad \sum_{a \in G} a(x) = \sqrt{nc}^{-3}, \quad (1)$$

$$b(0) = -\frac{1}{d}, \quad \sum_y \overline{\langle x, y \rangle} b(y) = \sqrt{nc} \overline{b(x)}, \quad a(x)b(-x) = \overline{b(x)}, \quad (2)$$

$$\sum_x b(x+y)\overline{b(x)} = \delta_{y,0} - \frac{1}{d}, \quad \sum_x b(x+y)b(x+z)\overline{b(x)} = \overline{\langle y, z \rangle} b(y)b(z) - \frac{c}{d\sqrt{n}}. \quad (3)$$

Then the data  $\langle \cdot, \cdot \rangle, c, a, b$  determine a near-group category of type  $G + n$ . Two such categories  $\mathcal{C}_1$  and  $\mathcal{C}_2$  determined by  $\langle \cdot, \cdot \rangle_1, c_1, a_1, b_1$  and  $\langle \cdot, \cdot \rangle_2, c_2, a_2, b_2$  are equivalent as fusion categories if and only if there is  $\psi \in \text{Aut}(G)$  such that  $\langle x, y \rangle_1 = \langle \psi x, \psi y \rangle_2, a_1(x) = a_2(\psi x), b_1(x) = b_2(\psi x)$  and  $c_1 = c_2$ .

The modular data for the Drinfeld center of a near-group category of type  $G + n$ , for  $|G| = n$ , is given as follows [11]. First, we need to find all functions  $\xi : G \rightarrow \mathbb{T}$  and values  $\tau \in G, \omega \in \mathbb{T}$  which satisfy

$$\sum_g \xi(g) = \sqrt{n}\omega^2 a(\tau) c^3 - nd^{-1} \quad (4)$$

$$\bar{c} \sum_k b(g+k)\xi(k) = \omega^2 c^3 a(\tau) \overline{\xi(g+\tau)} - \sqrt{nd}^{-1} \quad (5)$$

$$\xi(\tau-g) = \omega c^4 a(g) a(\tau-g) \overline{\xi(g)} \quad (6)$$

$$\sum_k \xi(k) b(k-g) b(k-h) = c^{-2} b(g+h-\tau) \xi(g) \xi(h) \overline{a(g-h)} - c^2 d^{-1} \quad (7)$$

There are  $n(n+3)/2$  triples  $(\xi_i, \tau_i, \omega_i)$  that satisfy the above equations. The corresponding center has rank  $n(n+3)$  with the following 4 subsets of simple objects:

- (1)  $\mathfrak{a}_g, g \in G, \dim(\mathfrak{a}_g) = 1;$
- (2)  $\mathfrak{b}_h, h \in G, \dim(\mathfrak{b}_h) = d + 1;$

- (3)  $\mathbf{c}_{l,k} = \mathbf{c}_{k,l}$ ,  $l, k \in G$ ,  $l \neq k$ ,  $\dim(\mathbf{c}_{k,l}) = d + 2$   
(4)  $\mathfrak{d}_j$ , where  $j$  corresponds to a triple  $(\xi_j, \tau_j, \omega_j)$ ,  $\dim(\mathfrak{d}_j) = d$ .

The  $T$  and  $S$  matrices are given by the following block form

$$T = \text{diag}(\langle g, g \rangle, \langle h, h \rangle, \langle k, l \rangle, \omega_j) \quad (8)$$

$$S = \begin{pmatrix} \langle g, g' \rangle^{-2} & (d+1)\langle g, h' \rangle^{-2} & (d+2)\overline{\langle g, k' + l' \rangle} & d\langle g, \tau_{j'} \rangle \\ (d+1)\langle h, g' \rangle^{-2} & \langle h, h' \rangle^{-2} & (d+2)\overline{\langle h, k' + l' \rangle} & -d\langle h, \tau_{j'} \rangle \\ (d+2)\overline{\langle k + l, g' \rangle} & (d+2)\overline{\langle k + l, h' \rangle} & S_{(k,l),(k',l')} & \mathbf{0} \\ d\langle \tau_j, g' \rangle & -d\langle \tau_j, h' \rangle & \mathbf{0} & S_{j,j'} \end{pmatrix}, \quad (9)$$

where

$$S_{j,j'} = \omega_j \omega_{j'} \sum_{g \in G} \langle \tau_j + \tau_{j'} + g, g \rangle \\ + d\omega_j \omega_{j'} c^6 a(\tau_j) a(\tau_{j'}) n^{-1} \sum_{g, h \in G} \overline{\xi_j(g) \xi_{j'}(h) \langle \tau_j - \tau_{j'} + h - g, h - g \rangle}, \quad (10)$$

and

$$S_{(k,l),(k',l')} = (d+2) \left( \overline{\langle k, k' \rangle \langle l, l' \rangle} + \overline{\langle k, l' \rangle \langle l, k' \rangle} \right). \quad (11)$$

### 3. REALIZATION OF MODULAR DATA

In [16], the following modular data of rank 10 labeled as  $10_{0,1435}^{20,676}$  was constructed:

$$S = \begin{pmatrix} 1 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & 4\chi_5^2 & 4\chi_5^2 & \chi_{80}^9 \\ \chi_{20}^5 & 3\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & 0 & 0 & \chi_{20}^5 \\ \chi_{20}^5 & -\chi_{20}^5 & 3\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & 0 & 0 & \chi_{20}^5 \\ \chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & s\chi_{20}^5 & \bar{s}\chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & 0 & 0 & \chi_{20}^5 \\ \chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & \bar{s}\chi_{20}^5 & s\chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & 0 & 0 & \chi_{20}^5 \\ \chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \bar{s}\chi_{20}^5 & s\chi_{20}^5 & 0 & 0 & \chi_{20}^5 \\ \chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & s\chi_{20}^5 & \bar{s}\chi_{20}^5 & 0 & 0 & \chi_{20}^5 \\ 4\chi_5^2 & 0 & 0 & 0 & 0 & 0 & 0 & -2\chi_5^3 & \chi_{180}^{14} & -4\chi_5^2 \\ 4\chi_5^2 & 0 & 0 & 0 & 0 & 0 & 0 & \chi_{180}^{14} & -2\chi_5^3 & -4\chi_5^2 \\ \chi_{80}^9 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & -4\chi_5^2 & -4\chi_5^2 & 1 \end{pmatrix}, \quad (12)$$

$$T = \text{diag} \left( 1, 1, 1, i, i, -i, -i, e^{\frac{4i\pi}{5}}, e^{\frac{3i\pi}{5}}, 1 \right),$$

where  $\chi_n^m = m + \sqrt{n}$  and  $s = -1 + 2i$ . It was conjectured in [16] that they can be realized from the condensation reductions of the Drinfeld center of the near-group category of type  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} + 16$ , with the symmetric bi-character  $\langle (g_1, g_2), (h_1, h_2) \rangle = (\zeta_4)^{g_1 h_1 - g_2 h_2}$ ,  $(g_1, g_2), (h_1, h_2) \in \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ .

In this section, we prove this conjecture in Theorem 3.2.6. Additionally, we obtain the modular data for the condensation of the Drinfeld center of a near-group category of type  $\mathbb{Z}/8\mathbb{Z} + 8$  in Theorem 3.3.3. This modular data is a Galois conjugate to that of the modular category  $\mathcal{C}(\mathfrak{g}_2, 4)$  (see Corollary 3.3.4), confirming a conjecture from [23, Table 8].

#### 3.1. Near-group categories $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} + 16$ .

3.1.1. *Existence of  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} + 16$ .* Let  $\langle \cdot, \cdot \rangle$  be the non-degenerate symmetric bi-character on  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$  defined by:

$$\begin{aligned} \langle \cdot, \cdot \rangle : \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} &\rightarrow \mathbb{T} \\ \langle (g_1, g_2), (h_1, h_2) \rangle &= (\zeta_4)^{g_1 h_1 - g_2 h_2} \end{aligned} \quad (13)$$

One solution associated with  $\langle \cdot, \cdot \rangle$  by solving the equations in Theorem 2.2.1 is

$$\begin{aligned} a(g_1, g_2) &= \zeta_8^{3(g_1^2 - g_2^2)}, \quad (g_1, g_2) \in \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}, \quad c = 1, \quad b(0, 0) = \frac{1}{4}(2 - \sqrt{5}), \\ b(1, 2) &= \overline{b(2, 1)}, \quad b(2, 0) = \overline{b(0, 2)}, \quad b(1, 3) = \frac{1}{8} \left( 1 - \sqrt{5} + i\sqrt{2(-1 + \sqrt{5})} \right), \\ b(1, 0) &= \overline{b(0, 1)}, \quad b(1, 1) = b(2, 2) = \frac{1}{4}, \quad b(0, 2) = -\frac{i}{4}, \\ b(0, 1) &= \frac{1}{32} \left( (2 - 4i) + (1 - 3i)\sqrt{2} - 2\sqrt{5} + (1 + i)\sqrt{10} \right. \\ &\quad \left. - 2(-1)^{1/8} 2^{1/4} (2 + \sqrt{2}) \sqrt{(-1 + \sqrt{2})(-1 + \sqrt{5})} \right), \\ b(2, 1) &= \frac{1}{32} \left( (2 - 4i) - (1 - 3i)\sqrt{2} - 2\sqrt{5} - 2(-1)^{1/4} \sqrt{5} \right. \\ &\quad \left. - 2i\sqrt{-1 + \sqrt{5}} + 2^{3/4} \sqrt{(-4 + 3\sqrt{2})(-1 + \sqrt{5})} \right). \end{aligned} \quad (14)$$

The rest values of  $b$  on  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$  can be computed via the formula  $b(-x) = \overline{a(x)b(x)}$  from Equation (2). Up to equivalence of fusion categories, the above solution is the unique one corresponds to the bicharacter in Equation (13):

**Proposition 3.1.1.** *Let  $\langle \cdot, \cdot \rangle$  be the non-degenerate symmetric bicharacter on  $\mathbb{Z}/4 \times \mathbb{Z}/4$  defined in Equation (13). Then, up to equivalence of fusion categories, there exists a unique near-group category of type  $\mathbb{Z}/4 \times \mathbb{Z}/4 + 16$  corresponding to this bicharacter.*

*Proof.* Solving Equation (1) in Theorem 2.2.1 with respect to the bicharacter in Equation (13), we found 4 solutions for the quadratic forms:

$$a_1(g_1, g_2) = \zeta_8^{3(g_1^2 - g_2^2)}, \quad a_2(g_1, g_2) = \zeta_8^{3g_1^2 + g_2^2}, \quad a_3(g_1, g_2) = \zeta_8^{-g_1^2 + g_2^2}, \quad a_4(g_1, g_2) = \zeta_8^{-g_1^2 - 3g_2^2}.$$

Only the quadratic form  $a_1$  leads to further solutions for all the equations in Theorem 2.2.1. Furthermore, the equation  $\sum_{a \in G} a(x) = \sqrt{n}c^{-3}$  in Equation (1) implies  $c^{-3} = 1$ . This yields four possible solutions for  $b$ , all corresponding to  $c = 1$ . These solutions are listed in Table 1. By Equations (2) and (3), we have  $b(0, 0) = \frac{1}{4}(2 - \sqrt{5})$ ,  $b(-x) = \overline{a(x)b(x)}$  and  $b(x) = \frac{1}{\sqrt{n}} \exp(ij(x))$  for some  $j(x) \in [-\pi, \pi]$  and  $x \in \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ . Thus it suffices to present values in the first row of Table 1 to fully determine  $b$ . In particular,  $J_1$  in Table 1 corresponds to the solution in Equation (14). Next, consider the automorphisms on

$\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$  defined by  $\psi_1(g_1, g_2) = \begin{cases} (g_1, g_2) & \text{if } g_1 = 0 \text{ or } 2 \\ (-g_1, g_2) & \text{if } g_1 = 1 \text{ or } 3 \end{cases}$ ,  $\psi_2(g_1, g_2) = (-g_1, -g_2)$ , and

$\psi_3(g_1, g_2) = \begin{cases} (g_1, g_2) & \text{if } g_2 = 0 \text{ or } 2 \\ (g_1, -g_2) & \text{if } g_2 = 1 \text{ or } 3 \end{cases}$ , where  $(g_1, g_2) \in \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ . It is straightforward

to verify  $\psi_1, \psi_2$  and  $\psi_3$  establish equivalences from  $J_1$  to  $J_2, J_3$  and  $J_4$  as described in Theorem 2.2.1, respectively. Thus the result follows.

id	$j(0, 1)$	$j(0, 2)$	$j(1, 0)$	$j(1, 1)$	$j(1, 2)$	$j(1, 3)$	$j(2, 0)$	$j(2, 1)$	$j(2, 2)$
$J_1$	1.99103	1.57080	-1.99103	0	-2.44331	-2.23704	-1.57080	2.44331	0
$J_2$	1.99103	1.57080	-0.36516	2.23704	-3.05447	0	-1.57080	2.44331	0
$J_3$	0.36516	1.57080	-0.36516	0	-3.05447	2.23704	-1.57080	3.05447	0
$J_4$	0.36516	1.57080	-1.99103	-2.23704	-2.44331	0	-1.57080	3.05447	0

TABLE 1. Four solutions for  $b$  in Proposition 3.1.1

□

**Remark 3.1.2.** *We do not know if there are any near-group categories of type  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} + 16$  other than the one in Proposition 3.1.1 up to Galois conjugation and equivalence of fusion categories. For the remainder of this paper, we will use  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} + 16$  to refer specifically to the category in Proposition 3.1.1.*

3.1.2. *Modular data of the center.* By solving the Equations (4)–(7), we found 152 triples of solutions for  $(\xi_i, \tau_i, \omega_i)$ . Apply Equations (8) – (11), we obtain the modular data of the center of  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} + 16$ . In particular, the center of the near-group category  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} + 16$  is a modular category of rank 304 with the following simple objects:

- 16 invertible objects  $X_g, g \in \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ ;
- 16  $\chi_{80}^9$ -dimensional simple objects  $Y_h, h \in \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ ;
- 120  $\chi_{80}^{10}$ -dimensional objects  $Z_{k,l}, k, l \in \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ , and  $k \neq l$ ;
- 152  $\chi_{80}^8$ -dimensional objects  $W_{\omega_i, \tau_i, \xi_i}$ , where  $(\omega_i, \tau_i, \xi_i)$  are solutions in Table 6.

We provide detailed modular data in the Mathematica notebook `MDZ4Z4.nb` in the arXiv source. The pointed subcategory is essential for the next section, and the relevant data is presented below. Using Equations (8) and (9), the  $S$  and  $T$  matrices for the pointed subcategory of the center are given by:

$$\begin{aligned} S_{\text{pt}} &= \left( \zeta_4^{g_1 h_1 - g_2 h_2} \right), \text{ where } (g_1, g_2), (h_1, h_2) \in \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}, \\ T_{\text{pt}} &= \text{diag} \left( \zeta_4^{g_1^2 - g_2^2} \right), \text{ where } (g_1, g_2) \in \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}. \end{aligned} \tag{15}$$

3.2. **Condensation of the Drinfeld center of  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} + 16$ .** Let  $\mathcal{C}$  be a modular category. Assume that  $\mathcal{C}$  contains a Tannakian fusion subcategory  $\text{Rep}(G)$ , then the de-equivariantization of  $\mathcal{C}$  by  $\text{Rep}(G)$ , denoted by  $\mathcal{C}_G$ , is a  $G$ -crossed braided fusion category with a faithful  $G$ -grading. The trivial component  $\mathcal{C}_G^0$  is also a modular category, which is called the condensation of  $\mathcal{C}$  by  $\text{Rep}(G)$  (see [8, 2, 12] for further details). Moreover,  $\mathcal{C}_G^0$  is the category of local modules  $\mathcal{C}_A^0$  over the étale algebra  $A = \text{Fun}(G) \in \text{Rep}(G)$  in  $\mathcal{C}$ . It is also braided tensor equivalent to the de-equivariantization of centralizer of  $\text{Rep}(G)$  by  $\text{Rep}(G)$ . The dimension of  $\mathcal{C}_G^0$  is given by  $\dim(\mathcal{C}_G^0) = \frac{\dim(\mathcal{C})}{|G|^2}$  [8].

Let  $\mathcal{C}$  be the near-group category of type  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} + 16$ , then the Drinfeld center of  $\mathcal{C}$ , denoted by  $\mathcal{Z}(\mathcal{C})$ , is modular. By analyzing the twists of the pointed subcategory of  $\mathcal{Z}(\mathcal{C})$  in Equation (15), we find that  $\mathcal{Z}(\mathcal{C})$  contains a Tannakian fusion subcategory  $\mathcal{E} = \text{Rep}(\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z})$ . Furthermore, we have the dimensions  $\dim(\mathcal{E}'_{\mathcal{Z}(\mathcal{C})}) = 640(9 + 4\sqrt{5})$  and  $\dim(\mathcal{Z}(\mathcal{C})_G^0) = 80(2 + \sqrt{5})^2$ , where  $G = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ .

Denote  $Y_i$  and  $Z_j, W_k$  be the simple objects of  $\mathcal{Z}(\mathcal{C})$  of dimensions  $\chi_{80}^9, \chi_{80}^{10}, \chi_{80}^8$ , respectively. For simplicity of the indices, we take the ordering of indices as following:  $Y_i$ 's are indexed by elements in  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$  with order  $(0, 0), \dots, (0, 3), (1, 0), \dots, (3, 3)$ ,  $Z_j$ 's are indexed by 120 pairs of elements  $\{(g_1, g_2), (h_1, h_2)\}$  in  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$  with order  $\{(0, 0), (0, 1)\}, \dots, \{(0, 1), (0, 2)\}, \dots, \{(3, 2), (3, 3)\}$  and  $W_k$  are indexed with the ordering in Table 6 in Appendix A. Then  $\text{rank}(\mathcal{E}'_{\mathcal{Z}(\mathcal{C})}) = 44$ , and

$$\mathcal{O}(\mathcal{E}'_{\mathcal{Z}(\mathcal{C})}) = \mathcal{O}(\mathcal{E}) \cup \left\{ \begin{array}{l} Y_1, Y_3, Y_6, Y_8, Y_9, Y_{11}, Y_{14}, Y_{16}, \\ Z_{10}, Z_{17}, Z_{23}, Z_{35}, Z_{50}, Z_{56}, Z_{62}, Z_{75}, Z_{83}, Z_{90}, Z_{101}, Z_{116}, \\ W_1, W_2, W_3, W_4, W_5, W_6, W_7, W_8, \\ W_{95}, W_{96}, W_{97}, W_{98}, W_{99}, W_{100}, W_{101}, W_{102} \end{array} \right\}. \quad (16)$$

Next we determine the set of simple objects of the condensation  $\mathcal{D} := \mathcal{Z}(\mathcal{C})_G^0$  of  $\mathcal{Z}(\mathcal{C})$  by  $\mathcal{E}$ . Let  $A := \text{Fun}(\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z})$ , we list the action of  $A$  on simple objects of  $\mathcal{E}'_{\mathcal{Z}(\mathcal{C})}$  in Table 2. Hence we obtain the simple objects of  $\mathcal{D}$ :

$$\{F(\mathbf{1}), F(Y_1), (Z_{10})_1, (Z_{10})_2, (Z_{17})_1, (Z_{17})_2, (Z_{23})_1, (Z_{23})_2, F(W_1), F(W_5)\}.$$

Next, we determine the modular data of  $\mathcal{D}$ . Since  $\theta_{Y_1} = \theta_{Z_{10}} = 1$ ,  $\theta_{Z_{17}} = \zeta_4$ ,  $\theta_{Z_{23}} = \zeta_4^3$ ,  $\theta_{W_1} = \zeta_5^2$  and  $\theta_{W_5} = \zeta_5^3$ , it follows from [12] that the  $T$ -matrix of  $\mathcal{D}$  is given by:

$$\text{diag}(1, 1, 1, 1, \zeta_4, \zeta_4, \zeta_4^3, \zeta_4^3, \zeta_5^2, \zeta_5^3).$$

The remaining task is to determine the  $S$ -matrix of  $\mathcal{D}$ .

Dim	$A \otimes -$	
$\chi_{80}^9$	$A \otimes Y_i = Y_1 \oplus Y_3 \oplus Y_6 \oplus Y_8 \oplus Y_9 \oplus Y_{11} \oplus Y_{14} \oplus Y_{16}$	$i = 1, 3, 6, 8, 9, 11, 14, 16$
$\chi_{80}^{10}$	$A \otimes Z_j = 2(Z_{10} \oplus Z_{35} \oplus Z_{75} \oplus Z_{90})$	$j = 10, 35, 75, 90$
$\chi_{80}^{10}$	$A \otimes Z_j = 2(Z_{17} \oplus Z_{56} \oplus Z_{101} \oplus Z_{116})$	$j = 17, 56, 101, 116$
$\chi_{80}^{10}$	$A \otimes Z_j = 2(Z_{23} \oplus Z_{50} \oplus Z_{62} \oplus Z_{83})$	$j = 23, 50, 62, 83$
$\chi_{80}^8$	$A \otimes W_k = \bigoplus_{k=1}^4 (W_k \oplus W_{k+94})$	$1 \leq k \leq 4, 95 \leq k \leq 98$
$\chi_{80}^8$	$A \otimes W_k = \bigoplus_{k=5}^8 (W_k \oplus W_{k+94})$	$5 \leq k \leq 8, 99 \leq k \leq 102$

TABLE 2. Action of étale algebra  $A$

**Lemma 3.2.1.**  $S_{F(Y_1), F(Y_1)} = 1$ ,  $S_{F(Y_1), F(W_j)} = -\chi_{80}^8$ ,  $S_{F(W_1), F(W_1)} = S_{F(W_5), F(W_5)} = -\chi_{20}^6$ ,  $S_{F(W_1), F(W_5)} = \chi_{180}^{14}$ .

*Proof.* By [12, Lemma 4.2], we have  $S_{F(V_1), F(V_2)} = \frac{1}{64} S_{A \otimes V_1, A \otimes V_2}$ , for any objects  $V_1, V_2 \in \{W_1, W_5, Y_1\}$ . Notice that the étale algebra  $A$  acts freely on these simple objects, hence  $S_{F(V_1), F(V_2)} = \frac{1}{64} \sum_{g, h \in \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}} S_{g \otimes V_1, h \otimes V_2} = S_{V_1, V_2}$ . Thus the result follows.  $\square$

**Lemma 3.2.2.** For all  $V \in \mathcal{O}(\mathcal{D})$ ,  $S_{F(Y_1), V} = \chi_{80}^9 \sigma(\dim(V))$ , where  $\sigma(\sqrt{5}) = -\sqrt{5}$ . Additionally,  $S_{F(W_1), (Z_k)_j} = S_{F(W_5), (Z_k)_j} = 0$ .

*Proof.* Let  $\sigma \in \text{Gal}(\mathbb{Q}(T)/\mathbb{Q})$  be the Galois automorphism such that  $\sigma(\sqrt{5}) = -\sqrt{5}$ . Then  $\sigma(\dim(\mathcal{D})) = \frac{\dim(\mathcal{D})}{\dim(\hat{\sigma}(F(\mathbf{1})))^2}$ . This implies  $\dim(\hat{\sigma}(F(\mathbf{1}))) = \chi_{80}^9$ , so  $\hat{\sigma}(F(\mathbf{1})) = F(Y_1)$  and consequently,  $S_{F(Y_1), V} = \chi_{80}^9 \sigma(\dim(V))$ . In particular,  $S_{(Z_k)_j, F(Y_1)} = \chi_{20}^5$ .



As  $\theta_{W_1} = \zeta_5^2$  and  $\theta_{W_5} = \zeta_5^3$ ,  $F(W_1)$  and  $F(W_5)$  are in the same orbit under the action of  $\sigma$ . Since  $S_{F(W_i), F(W_j)}$  are real algebraic integers,  $F(W_1)$  and  $F(W_5)$  are self-dual. Therefore  $S_{(Z_k)_j, F(W_l)}$  are also real algebraic integers, where  $l = 1, 5$ . Since  $\dim(\mathcal{D}) = 2(\chi_{80}^8)^2 + (\chi_{180}^{14})^2 + (\chi_{20}^6)^2$ , we conclude that  $S_{(Z_k)_j, F(W_l)} = 0$  for  $l = 1, 5$ .  $\square$

Thus the  $S$ -matrix of  $\mathcal{D}$  is of the form

$$\begin{pmatrix} 1 & \chi_{80}^9 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{80}^8 & \chi_{80}^8 \\ \chi_{80}^9 & 1 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & -\chi_{80}^8 & -\chi_{80}^8 \\ \chi_{20}^5 & \chi_{20}^5 & & & & & & & & 0 & 0 \\ \chi_{20}^5 & \chi_{20}^5 & & & & & & & & 0 & 0 \\ \chi_{20}^5 & \chi_{20}^5 & & & * & * & * & & & 0 & 0 \\ \chi_{20}^5 & \chi_{20}^5 & & & & & & & & 0 & 0 \\ \chi_{20}^5 & \chi_{20}^5 & & & & & & & & 0 & 0 \\ \chi_{20}^5 & \chi_{20}^5 & & & & & & & & 0 & 0 \\ \chi_{20}^5 & \chi_{20}^5 & & & & & & & & 0 & 0 \\ \chi_{80}^8 & -\chi_{80}^8 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -\chi_{20}^6 & \chi_{180}^{14} \\ \chi_{80}^8 & -\chi_{80}^8 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \chi_{180}^{14} & -\chi_{20}^6 \end{pmatrix} \quad (17)$$

To determine the remaining entries of the  $S$ -matrix of  $\mathcal{D}$  in (17), we apply the method introduced in the reconstruction program in [15, 16]. Specifically, we begin by identifying all possible representation types of the associated modular congruence representation. This involves decomposing the representation into a direct sum of irreducible components. Next, we reconstruct the modular data of  $\mathcal{D}$  from the representation of  $\mathrm{SL}(2, \mathbb{Z})$ , leveraging properties of modular categories, such as Galois symmetry, the Verlinde formula, and other structural constraints.

Let  $\rho_{\mathcal{D}}$  be the  $\mathrm{SL}(2, \mathbb{Z})$  modular congruence representation of  $\mathcal{D}$ , then it follows from the spectrum of  $\rho_{\mathcal{D}}(\mathfrak{t})$  that  $\rho_{\mathcal{D}}$  contains a unique direct summand  $\rho_1$  of dimension 3 of

level 5, where  $\rho_1(\mathfrak{s}) = \frac{-1}{\sqrt{5}} \begin{pmatrix} 1 & \sqrt{2} & \sqrt{2} \\ \sqrt{2} & 2 \cos \frac{2\pi}{5} & 2 \cos \frac{6\pi}{5} \\ \sqrt{2} & 2 \cos \frac{6\pi}{5} & 2 \cos \frac{2\pi}{5} \end{pmatrix}$  and  $\rho_1(\mathfrak{t}) = \mathrm{diag}(1, \zeta_5^2, \zeta_5^3)$ . Assume

$\rho_{\mathcal{D}} \cong \rho_1 \oplus \rho'$ , then  $\rho'$  is a direct sum of irreducible representations with level 4, whose  $\mathfrak{t}$ -spectrum belongs to  $\{1, \zeta_4, \zeta_4^3\}$ . Notice that these irreducible direct summands of level 4 are  $\rho_2, \rho_3, \rho_4, \rho_5$  [19, 20, 21], where

$$\rho_2(\mathfrak{s}) = \frac{1}{2} \begin{pmatrix} -1 & 1 & \sqrt{2} \\ 1 & -1 & \sqrt{2} \\ \sqrt{2} & -\sqrt{2} & 0 \end{pmatrix}, \rho_2(\mathfrak{t}) = \mathrm{diag}(\zeta_4, \zeta_4^3, 1),$$

$$\rho_3(\mathfrak{s}) = \frac{1}{2} \begin{pmatrix} \zeta_4 & \sqrt{3}\zeta_4 \\ \sqrt{3}\zeta_4 & -\zeta_4 \end{pmatrix}, \rho_3(\mathfrak{t}) = \mathrm{diag}(\zeta_4, \zeta_4^3),$$

$$\rho_4(\mathfrak{s}) = \zeta_4^3, \rho_4(\mathfrak{t}) = \zeta_4, \rho_5(\mathfrak{s}) = \zeta_4, \rho_5(\mathfrak{t}) = \zeta_4^3.$$

**Lemma 3.2.3.** *Let  $\rho_{\mathcal{D}} = \rho_1 \oplus \rho'$  be the  $\mathrm{SL}(2, \mathbb{Z})$  representation of  $\mathcal{D}$ . The  $\rho_{\mathcal{D}}$  is equivalent to one of the following*

$$\rho_1 \oplus 2\rho_2 \oplus \rho_0, \rho_1 \oplus \rho_2 \oplus \rho_3 \oplus 2\rho_0, \rho_1 \oplus \rho_2 \oplus \rho_4 \oplus \rho_5 \oplus 2\rho_0.$$

*Proof.* By comparing the  $\mathfrak{t}$ -spectrum of  $\rho'(\mathfrak{t})$  with that of  $\rho_{\mathcal{D}}(\mathfrak{t})$ , a direct computation shows  $\rho'$  is equivalent to one of the following

$$2\rho_2 \oplus \rho_0, \rho_2 \oplus \rho_3 \oplus 2\rho_0, \rho_2 \oplus \rho_4 \oplus \rho_5 \oplus 2\rho_0, 2\rho_3 \oplus 3\rho_0, \rho_3 \oplus \rho_4 \oplus \rho_5 \oplus 3\rho_0, 2\rho_4 \oplus 2\rho_5 \oplus 3\rho_0.$$

If  $\rho_{\mathcal{D}} \cong (\rho_1 \oplus 3\rho_0) \oplus \rho''$  for  $\rho'' = 2\rho_3, \rho_3 \oplus \rho_4 \oplus \rho_5$ , or  $2\rho_4 \oplus 2\rho_5$ , then  $\rho_{\mathcal{D}}$  is a direct sum of two sub-representations with disjoint  $\mathfrak{t}$ -spectrums. However, it is not possible by [3, Lemma 3.18]. This completes the proof of the lemma.  $\square$

**Proposition 3.2.4.**  $\rho_{\mathcal{D}} \not\cong \rho_1 \oplus \rho_2 \oplus \rho_3 \oplus 2\rho_0$ .

*Proof.* Assume  $\rho_{\mathcal{D}} \cong \tilde{\rho} := \rho_1 \oplus \rho_2 \oplus \rho_3 \oplus 2\rho_0$ . Notice that

$$\tilde{\rho}(\mathfrak{s}) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -\frac{1}{\sqrt{5}} & 0 & 0 & 0 & 0 & 0 & -\frac{\sqrt{2}}{\sqrt{5}} & -\frac{\sqrt{2}}{\sqrt{5}} & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{\sqrt{2}} & -\frac{1}{2} & 0 & \frac{1}{2} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{\zeta_4}{2} & 0 & \frac{\sqrt{3}\zeta_4}{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{\sqrt{2}} & \frac{1}{2} & 0 & -\frac{1}{2} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{\sqrt{3}\zeta_4}{2} & 0 & -\frac{\zeta_4}{2} & 0 & 0 & 0 \\ 0 & 0 & -\frac{\sqrt{2}}{\sqrt{5}} & 0 & 0 & 0 & 0 & 0 & -\frac{2\cos\frac{2\pi}{5}}{\sqrt{5}} & -\frac{2\cos\frac{6\pi}{5}}{\sqrt{5}} & 0 \\ 0 & 0 & -\frac{\sqrt{2}}{\sqrt{5}} & 0 & 0 & 0 & 0 & 0 & -\frac{2\cos\frac{6\pi}{5}}{\sqrt{5}} & -\frac{2\cos\frac{2\pi}{5}}{\sqrt{5}} & 0 \end{pmatrix}.$$

It follows from [15, Theorem 3.4] that there is a real orthogonal matrix  $U = (a_{ij})$  such

that  $\rho_{\mathcal{D}}(\mathfrak{s}) = U\tilde{\rho}(\mathfrak{s})U^t$  and  $U\rho_{\mathcal{D}}(\mathfrak{t}) = \rho_{\mathcal{D}}(\mathfrak{t})U$ . Then  $U = \begin{pmatrix} A_1 & & & \\ & A_2 & & \\ & & A_3 & \\ & & & A_4 \end{pmatrix}$  and  $A_4 =$

$$\begin{pmatrix} \mu_1 & \\ & \mu_2 \end{pmatrix}, \mu_1, \mu_2 \in \{\pm 1\}.$$

Hence, we obtain  $\rho_{\mathcal{D}}(\mathfrak{s}) = \begin{pmatrix} B_1 & B_2 \\ B_2^t & B_3 \end{pmatrix}$ , where

$$B_1 = \begin{pmatrix} 1 - \frac{a_{13}^2+1}{\sqrt{5}} - a_{14}^2 & -a_{13}a_{23}\frac{\sqrt{5}+1}{\sqrt{5}} - a_{14}a_{24} & -a_{13}a_{33}\frac{\sqrt{5}+1}{\sqrt{5}} - a_{14}a_{34} & -a_{13}a_{43}\frac{\sqrt{5}+1}{\sqrt{5}} - a_{14}a_{44} \\ -a_{13}a_{23}\frac{\sqrt{5}+1}{\sqrt{5}} - a_{14}a_{24} & 1 - \frac{a_{23}^2+1}{\sqrt{5}} - a_{24}^2 & -a_{23}a_{33}\frac{\sqrt{5}+1}{\sqrt{5}} - a_{24}a_{34} & -a_{23}a_{43}\frac{\sqrt{5}+1}{\sqrt{5}} - a_{24}a_{44} \\ -a_{13}a_{33}\frac{\sqrt{5}+1}{\sqrt{5}} - a_{14}a_{34} & -a_{23}a_{33}\frac{\sqrt{5}+1}{\sqrt{5}} - a_{24}a_{34} & 1 - \frac{a_{33}^2+1}{\sqrt{5}} - a_{34}^2 & -a_{33}a_{43}\frac{\sqrt{5}+1}{\sqrt{5}} - a_{34}a_{44} \\ -a_{13}a_{43}\frac{\sqrt{5}+1}{\sqrt{5}} - a_{14}a_{44} & -a_{23}a_{43}\frac{\sqrt{5}+1}{\sqrt{5}} - a_{24}a_{44} & -a_{33}a_{43}\frac{\sqrt{5}+1}{\sqrt{5}} - a_{34}a_{44} & 1 - \frac{a_{43}^2+1}{\sqrt{5}} - a_{44}^2 \end{pmatrix}$$

$$B_2 = \begin{pmatrix} \frac{a_{14}a_{55}}{\sqrt{2}} & \frac{a_{14}a_{65}}{\sqrt{2}} & \frac{a_{14}a_{77}}{\sqrt{2}} & \frac{a_{14}a_{87}}{\sqrt{2}} & -\frac{\sqrt{2}a_{13}\mu_1}{\sqrt{5}} & -\frac{\sqrt{2}a_{13}\mu_2}{\sqrt{5}} \\ \frac{a_{24}a_{55}}{\sqrt{2}} & \frac{a_{24}a_{65}}{\sqrt{2}} & \frac{a_{24}a_{77}}{\sqrt{2}} & \frac{a_{24}a_{87}}{\sqrt{2}} & -\frac{\sqrt{2}a_{23}\mu_1}{\sqrt{5}} & -\frac{\sqrt{2}a_{23}\mu_2}{\sqrt{5}} \\ \frac{a_{34}a_{55}}{\sqrt{2}} & \frac{a_{34}a_{65}}{\sqrt{2}} & \frac{a_{34}a_{77}}{\sqrt{2}} & \frac{a_{34}a_{87}}{\sqrt{2}} & -\frac{\sqrt{2}a_{33}\mu_1}{\sqrt{5}} & -\frac{\sqrt{2}a_{33}\mu_2}{\sqrt{5}} \\ \frac{a_{44}a_{55}}{\sqrt{2}} & \frac{a_{44}a_{65}}{\sqrt{2}} & \frac{a_{44}a_{77}}{\sqrt{2}} & \frac{a_{44}a_{87}}{\sqrt{2}} & -\frac{\sqrt{2}a_{43}\mu_1}{\sqrt{5}} & -\frac{\sqrt{2}a_{43}\mu_2}{\sqrt{5}} \end{pmatrix}$$

$$B_3 = \frac{1}{2} \begin{pmatrix} B_3^{(1,2,3,4)} & \mathbf{0}_{4 \times 2} \\ \mathbf{0}_{2 \times 4} & B_3^{(5,6)} \end{pmatrix}, \text{ with } B_3^{(5,6)} = \begin{pmatrix} \frac{-4 \cos \frac{2\pi}{5}}{\sqrt{5}} & \frac{-4\mu_1\mu_2 \cos \frac{6\pi}{5}}{\sqrt{5}} \\ \frac{-4\mu_1\mu_2 \cos \frac{6\pi}{5}}{\sqrt{5}} & \frac{-4 \cos \frac{2\pi}{5}}{\sqrt{5}} \end{pmatrix}, \text{ and } B_3^{(1,2,3,4)} =$$

$$\begin{pmatrix} -a_{55}^2 + \zeta_4 a_{56}^2 & -a_{55}a_{65} + \zeta_4 a_{56}a_{66} & a_{55}a_{77} + \sqrt{3}\zeta_4 a_{56}a_{78} & a_{55}a_{65} + \sqrt{3}\zeta_4 a_{56}a_{88} \\ -a_{55}a_{65} + \zeta_4 a_{56}a_{66} & -a_{65}^2 + \zeta_4 a_{66}^2 & a_{65}a_{77} + \sqrt{3}\zeta_4 a_{66}a_{78} & a_{65}a_{87} + \sqrt{3}\zeta_4 a_{66}a_{88} \\ a_{55}a_{77} + \sqrt{3}\zeta_4 a_{56}a_{78} & -a_{65}a_{77} + \zeta_4 a_{66}a_{78} & -(a_{77}^2 + \zeta_4 a_{78}^2) & -(a_{77}a_{87} + \sqrt{3}\zeta_4 a_{78}a_{88}) \\ a_{55}a_{87} + \zeta_4 a_{56}a_{88} & a_{65}a_{87} + \sqrt{3}\zeta_4 a_{66}a_{88} & -(a_{77}a_{87} + \sqrt{3}\zeta_4 a_{78}a_{88}) & -(a_{87}^2 + \zeta_4 a_{88}^2) \end{pmatrix}.$$

Without loss of generality, assume the first row and the second row are determined by the unit object and its Galois conjugate. Hence we have  $a_{13}\mu_j = a_{23}\mu_j = -\frac{1}{\sqrt{2}}$ ,  $a_{33} = a_{43} = 0$ ,  $a_{13}a_{23} = -\frac{1}{2}$ ,  $a_{14} = a_{24}$ ,  $a_{14}^2 = \frac{1}{4}$ . Let  $a_{13} = \frac{1}{\sqrt{2}}$  (the other case is similar),  $\mu_1 = \mu_2 = -1$ ,  $a_{14} = \frac{1}{2}$ . From coefficients in the first row and second row of the  $S$ -matrix, we have  $a_{14}a_{55} = a_{14}a_{65} = a_{14}a_{77} = a_{14}a_{87} = \frac{1}{2\sqrt{2}}$ , which implies  $a_{55} = a_{65} = a_{77} = a_{87} = \frac{1}{\sqrt{2}}$ . The orthogonality of matrices  $A_2$  and  $A_3$  implies  $a_{56}a_{66} = -\frac{1}{2} = a_{78}a_{88}$ . Then the  $S$ -matrix  $S = \sqrt{\dim(\mathcal{D})} \rho_{\mathcal{D}}(\mathfrak{s}) =$

$$\begin{pmatrix} 1 & \chi_{80}^9 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{80}^8 & \chi_{80}^8 \\ \chi_{80}^9 & 1 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & -\chi_{80}^8 & -\chi_{80}^8 \\ \chi_{20}^5 & \chi_{20}^5 & 3\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & 0 & 0 \\ \chi_{20}^5 & \chi_{20}^5 & -\chi_{20}^5 & 3\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & 0 & 0 \\ \chi_{20}^5 & \chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & c_1\chi_{20}^5 & \overline{c_1}\chi_{20}^5 & c_2\chi_{20}^5 & \overline{c_2}\chi_{20}^5 & 0 & 0 \\ \chi_{20}^5 & \chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & \overline{c_1}\chi_{20}^5 & c_1\chi_{20}^5 & \overline{c_2}\chi_{20}^5 & c_2\chi_{20}^5 & 0 & 0 \\ \chi_{20}^5 & \chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & c_2\chi_{20}^5 & \overline{c_2}\chi_{20}^5 & c_1\chi_{20}^5 & \overline{c_1}\chi_{20}^5 & 0 & 0 \\ \chi_{20}^5 & \chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & \overline{c_2}\chi_{20}^5 & c_2\chi_{20}^5 & \overline{c_1}\chi_{20}^5 & c_1\chi_{20}^5 & 0 & 0 \\ \chi_{80}^8 & -\chi_{80}^8 & 0 & 0 & 0 & 0 & 0 & 0 & -\chi_{20}^6 & \chi_{180}^{14} \\ \chi_{80}^8 & -\chi_{80}^8 & 0 & 0 & 0 & 0 & 0 & 0 & \chi_{180}^{14} & -\chi_{20}^6 \end{pmatrix},$$

where  $c_1 := (-1 + \zeta_4)$  and  $c_2 = (-1 + \sqrt{3}\zeta_4)$ . However, the fourth and fifth rows violate the orthogonality of  $S$ -matrix. Therefore,  $\rho_{\mathcal{D}} \not\cong \rho_1 \oplus \rho_2 \oplus \rho_3 \oplus 2\rho_0$ .  $\square$

**Proposition 3.2.5.**  $\rho_{\mathcal{D}} \not\cong \rho_1 \oplus 2\rho_2 \oplus \rho_0$ .

*Proof.* This proposition can be proved by using the same argument as Proposition 3.2.4, we omit the details here.  $\square$

**Theorem 3.2.6.**  $\rho_{\mathcal{D}} \cong \rho_1 \oplus \rho_2 \oplus \rho_4 \oplus \rho_5 \oplus 2\rho_0$ . The  $S$ -matrix of  $\mathcal{D}$  is

$$\begin{pmatrix} 1 & \chi_{80}^9 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{80}^8 & \chi_{80}^8 \\ \chi_{80}^9 & 1 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & -\chi_{80}^8 & -\chi_{80}^8 \\ \chi_{20}^5 & \chi_{20}^5 & 3\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & 0 & 0 \\ \chi_{20}^5 & \chi_{20}^5 & -\chi_{20}^5 & 3\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & 0 & 0 \\ \chi_{20}^5 & \chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & s\chi_{20}^5 & \overline{s}\chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & 0 & 0 \\ \chi_{20}^5 & \chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & \overline{s}\chi_{20}^5 & s\chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & 0 & 0 \\ \chi_{20}^5 & \chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & s\chi_{20}^5 & \overline{s}\chi_{20}^5 & 0 & 0 \\ \chi_{20}^5 & \chi_{20}^5 & -\chi_{20}^5 & -\chi_{20}^5 & \chi_{20}^5 & \chi_{20}^5 & \overline{s}\chi_{20}^5 & s\chi_{20}^5 & 0 & 0 \\ \chi_{80}^8 & -\chi_{80}^8 & 0 & 0 & 0 & 0 & 0 & 0 & -\chi_{20}^6 & \chi_{180}^{14} \\ \chi_{80}^8 & -\chi_{80}^8 & 0 & 0 & 0 & 0 & 0 & 0 & \chi_{180}^{14} & -\chi_{20}^6 \end{pmatrix},$$

where  $s := (-1 + 2\zeta_4)$ .

*Proof.* By Lemma 3.2.3, Proposition 3.2.4 and Proposition 3.2.5, we know  $\rho_{\mathcal{D}} \cong \rho_1 \oplus \rho_2 \oplus \rho_4 \oplus \rho_5 \oplus 2\rho_0$ . Let  $\tilde{\rho} := \rho_1 \oplus \rho_2 \oplus \rho_4 \oplus \rho_5 \oplus 2\rho_0$ , then

$$\tilde{\rho}(\mathfrak{s}) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -\frac{1}{\sqrt{5}} & 0 & 0 & 0 & 0 & 0 & -\frac{\sqrt{2}}{\sqrt{5}} & -\frac{\sqrt{2}}{\sqrt{5}} \\ 0 & 0 & 0 & 0 & \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{\sqrt{2}} & -\frac{1}{2} & 0 & \frac{1}{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -\zeta_4 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{\sqrt{2}} & \frac{1}{2} & 0 & -\frac{1}{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \zeta_4 & 0 & 0 \\ 0 & 0 & -\frac{\sqrt{2}}{\sqrt{5}} & 0 & 0 & 0 & 0 & 0 & -\frac{2 \cos \frac{2\pi}{5}}{\sqrt{5}} & -\frac{2 \cos \frac{6\pi}{5}}{\sqrt{5}} \\ 0 & 0 & -\frac{\sqrt{2}}{\sqrt{5}} & 0 & 0 & 0 & 0 & 0 & -\frac{2 \cos \frac{6\pi}{5}}{\sqrt{5}} & -\frac{2 \cos \frac{2\pi}{5}}{\sqrt{5}} \end{pmatrix}.$$

Again, we obtain  $\rho_{\mathcal{D}}(\mathfrak{s}) = U\tilde{\rho}(\mathfrak{s})U^t = \begin{pmatrix} B_1 & B_2 \\ B_2^t & \widetilde{B}_3 \end{pmatrix}$ , where

$$\widetilde{B}_3 = \frac{1}{2} \begin{pmatrix} \widetilde{B}_3^{(1,2,3,4)} & \mathbf{0}_{4 \times 2} \\ \mathbf{0}_{2 \times 4} & B_3^{(5,6)} \end{pmatrix}, \text{ with } \widetilde{B}_3^{(1,2,3,4)} = \begin{pmatrix} -a_{55}^2 + 2\zeta_4 a_{56}^2 & -a_{55}a_{65} + 2\zeta_4 a_{56}a_{66} & 0 & 0 \\ -a_{55}a_{65} + 2\zeta_4 a_{56}a_{66} & -a_{65}^2 + 2\zeta_4 a_{66}^2 & 0 & 0 \\ 0 & 0 & -(a_{77}^2 + 2\zeta_4 a_{78}^2) & -(a_{77}a_{87} + 2\zeta_4 a_{78}a_{88}) \\ 0 & 0 & -(a_{77}a_{87} + 2\zeta_4 a_{78}a_{88}) & -(a_{87}^2 + 2\zeta_4 a_{88}^2) \end{pmatrix}.$$

Applying the same method as in Proposition 3.2.4, we obtain the same relations for the orthogonal matrix  $U$ . Consequently, the desired  $S$ -matrix follows immediately as  $S = \sqrt{\dim(\mathcal{D})} \rho_{\mathcal{D}}(\mathfrak{s})$ .  $\square$

**3.3. Near-group category  $\mathbb{Z}/8\mathbb{Z} + 8$ .** There are 8 inequivalent near-group categories of type  $\mathbb{Z}/8\mathbb{Z} + 8$  [10]. The data for  $\mathbb{Z}/8\mathbb{Z} + 8$  is listed in Table 3. The non-degenerate symmetric bi-character for a cyclic group  $\mathbb{Z}/n\mathbb{Z}$  is of the form  $\langle x, y \rangle = \exp(2\pi i mxy/n)$ , where  $m \in \mathbb{Z}$  and  $\gcd(m, n) = 1$ . When  $n$  is even, the function  $a$  is in the form of  $a(x) = \varepsilon^x \exp(-\pi i mx^2/n)$ , where  $\varepsilon = \pm 1$ . By taking  $m \in \mathbb{Z}/2n\mathbb{Z}$  and coprime to  $n$ , we can eliminate the potential factor of  $-1$ . This  $m$  is recorded in the second column of Table 1. Recall that  $b(0) = -\frac{1}{d}$ ,  $b(-x) = \overline{a(x)b(x)}$  and  $b(x) = \frac{1}{\sqrt{n}} \exp(ij(x))$ , where  $j(x) \in [-\pi, \pi]$ . It suffices to list the values of  $j(1), \dots, j(4)$  to determine  $b(x)$ .

**3.3.1. The center of  $\mathbb{Z}/8\mathbb{Z} + 8$ .** We focus on the near-group category labeled by  $J_8^1$  in Table 3. The solutions for the triples  $(\omega_i, \tau_i, \xi_i)$  for Equations (4) – (7) are listed in Table 7 in Appendix A. Additional details regarding these solutions and the modular data of the center are provided in the Mathematica notebook `MDZ8.nb`, available in the arXiv source.

The center of  $\mathbb{Z}/8\mathbb{Z} + 8$  is a rank 88 modular category with the following simple objects:

- eight invertible objects  $X_g$ ,  $g \in \mathbb{Z}/8\mathbb{Z}$ ;
- eight  $\chi_{24}^5$ -dimensional simple objects  $Y_h$ ,  $h \in \mathbb{Z}/8\mathbb{Z}$ ;

id	$m$	$c$	$j(1), j(2), j(3), j(4)$
$\frac{J_8^1}{J_8^1}$	-1	$\zeta_{24}^{-1}$	0.872276, -2.70426, 2.9768, 3.14159
$\frac{J_8^1}{J_8^1}$	1	$\zeta_{24}$	-0.872276, 2.70426, -2.9768, 3.14159
$\frac{J_8^2}{J_8^2}$	-1	$\zeta_{24}^{-1}$	-1.26498, 1.13347, -0.227903, 3.14159
$\frac{J_8^2}{J_8^2}$	1	$\zeta_{24}$	1.26498, -1.13347, 0.227903, 3.14159
$\frac{J_8^3}{J_8^3}$	-3	$\zeta_{24}^{17}$	-2.46405, 3.07557, 0.491887, 0
$\frac{J_8^3}{J_8^3}$	3	$\zeta_{24}^{-17}$	2.46405, -3.07557, -0.491887, 0
$\frac{J_8^4}{J_8^4}$	-3	$\zeta_{24}^{17}$	1.28595, -1.50478, 1.47161, 0
$\frac{J_8^4}{J_8^4}$	3	$\zeta_{24}^{-17}$	-1.28595, 1.50478, -1.47161, 0

TABLE 3. Near-group categories of type  $\mathbb{Z}/8\mathbb{Z} + 8$ .

- twenty-eight  $2\chi_6^3$ -dimensional objects  $Z_{k,l}$ ,  $k, l \in \mathbb{Z}/8\mathbb{Z}$ , and  $k < l$ ;
- forty-four  $\chi_{24}^4$ -dimensional objects  $W_{\omega_i, \tau_i, \xi_i}$ , where  $(\omega_i, \tau_i, \xi_i)$  are solutions in Table 7.

The  $T$ -matrix is given as the following:

$$T_{X_g} = e^{-\frac{1}{4}\pi ig^2}, T_{Y_h} = e^{-\frac{1}{4}\pi ih^2}, T_{Z_{k,l}} = e^{-\frac{1}{4}\pi ikl}, g, h, k, l \in \mathbb{Z}/8\mathbb{Z}, \text{ and}$$

$$T_{W_{\omega, \tau, \xi}} = \text{diag} \left( e^{-\frac{i\pi}{2}}, e^{\frac{5i\pi}{6}}, e^{\frac{5i\pi}{6}}, e^{-\frac{2i\pi}{3}}, e^{-\frac{2i\pi}{3}}, e^{\frac{13i\pi}{16}}, e^{-\frac{3i\pi}{16}}, e^{\frac{7i\pi}{48}}, e^{\frac{7i\pi}{48}}, e^{-\frac{41i\pi}{48}}, e^{-\frac{41i\pi}{48}}, e^{-\frac{3i\pi}{4}}, e^{\frac{7i\pi}{12}}, e^{\frac{7i\pi}{12}}, e^{-\frac{11i\pi}{12}}, e^{-\frac{11i\pi}{12}}, e^{\frac{5i\pi}{16}}, e^{-\frac{11i\pi}{16}}, e^{\frac{31i\pi}{48}}, e^{\frac{31i\pi}{48}}, e^{-\frac{17i\pi}{48}}, e^{-\frac{17i\pi}{48}}, e^{\frac{i\pi}{2}}, e^{\frac{i\pi}{3}}, e^{\frac{i\pi}{3}}, e^{-\frac{i\pi}{6}}, e^{-\frac{i\pi}{6}}, e^{\frac{5i\pi}{16}}, e^{-\frac{11i\pi}{16}}, e^{\frac{31i\pi}{48}}, e^{\frac{31i\pi}{48}}, e^{-\frac{17i\pi}{48}}, e^{-\frac{17i\pi}{48}}, e^{-\frac{3i\pi}{4}}, e^{\frac{7i\pi}{12}}, e^{\frac{7i\pi}{12}}, e^{-\frac{11i\pi}{12}}, e^{-\frac{11i\pi}{12}}, e^{\frac{13i\pi}{16}}, e^{-\frac{3i\pi}{16}}, e^{\frac{7i\pi}{48}}, e^{\frac{7i\pi}{48}}, e^{-\frac{41i\pi}{48}}, e^{-\frac{41i\pi}{48}} \right).$$

And the  $S$ -matrix is

$$S = \begin{pmatrix} s_{X,X} & \chi_{24}^5 s_{X,X} & 2\chi_6^3 s_{X,Z} & \chi_{24}^4 s_{X,W} \\ \chi_{24}^5 s_{X,X} & s_{X,X} & 2\chi_6^3 s_{X,Z} & -\chi_{24}^4 s_{X,W} \\ 2\chi_6^3 s_{X,Z}^T & 2\chi_6^3 s_{X,Z}^T & 2\chi_6^3 s_{Z,Z} & \mathbf{0} \\ \chi_{24}^4 s_{X,W}^T & -\chi_{24}^4 s_{X,W}^T & \mathbf{0} & s_{W,W} \end{pmatrix},$$

where  $\lambda = 16\sqrt{3}\chi_{24}^5$ ,  $s_{X_g, X_h} = e^{\frac{1}{2}\pi igh}$ ,  $s_{X_g, Z_{k,l}} = e^{\frac{1}{4}\pi ig(k+l)}$ ,  $s_{Z_{k,l}, Z_{k',l'}}$  =  $e^{\frac{1}{4}\pi ikl' + \frac{1}{4}\pi ik'l} + e^{\frac{1}{4}\pi ikk' + \frac{1}{4}\pi ill'}$ , where  $g, h, k, l, k', l' \in \mathbb{Z}/8\mathbb{Z}$ ,  $k < l$  and  $k' < l'$ ; and  $s_{X,W}$  and  $s_{W,W}$  can be computed from equations in [10] using the data from Table 7.

As  $\theta_{X_4} = 1$  for the invertible object  $X_4$ , we know  $b = X_4$  is a boson. Therefore we can apply the boson condensation to  $\mathcal{Z}(\mathcal{C})$  to obtain  $\mathcal{Z}(\mathcal{C})_{\mathbb{Z}/2\mathbb{Z}}^0$ . By the Verlinde formula, the fusion rules for tensoring with the boson  $b$  can be computed. The fusion rules for tensoring with  $b$  and the objects centralized by  $b$  are summarized in Table 4.

By examining the table, we find that  $\mathcal{Z}(\mathcal{C})_{\mathbb{Z}/2\mathbb{Z}}^0$  has rank 36. The dimensions and twists of the simple objects are summarized in Table 5.

Since  $\mathcal{Z}(\mathcal{C})_{\mathbb{Z}/2\mathbb{Z}}^0$  contains a pointed modular subcategory  $\mathcal{C}(\mathbb{Z}_4, q)$ , there exists a braided tensor equivalence  $\mathcal{Z}(\mathcal{C})_{\mathbb{Z}/2\mathbb{Z}}^0 \cong \mathcal{C}(\mathbb{Z}_4, q) \boxtimes \mathcal{D}$  by [8, Theorem 3.13]. Hence, it suffices to determine the  $S$ -matrix of  $\mathcal{D}$ . From Table 5, the set of simple objects of  $\mathcal{D}$  is:

$$\mathcal{O}(\mathcal{D}) = \{F(\mathbf{1}), F(Y_0), F(Z_{1,7}), (Z_{2,6})_1, (Z_{2,6})_2, F(W_2), F(W_4), (W_1)_1, (W_1)_2\}.$$

**Lemma 3.3.1.**  $S_{(Z_{2,6})_i, F(Z_{1,7})} = -2\chi_6^3$ ,  $S_{(W_1)_i, F(Z_{1,7})} = 0$ ,  $i = 1, 2$ .

*Proof.* The  $S$ -coefficients between objects  $F(Y_0), F(Z_{1,7}), F(W_2), F(W_4)$  are the same of that in the Drinfeld center by [23, Theorem 2.2] or [12, Lemma 4.2]. Notice that  $\frac{\dim(\mathcal{D})}{\dim(F(Z_{1,7}))^2}$  is an

Dim	$b \otimes -$			in $\langle b \rangle'$ ?
1	$bX_g = X_{g+4}$	$g \in \mathbb{Z}/8\mathbb{Z}$		Yes
$\chi_{24}^5$	$bY_h = Y_{h+4}$	$h \in \mathbb{Z}/8\mathbb{Z}$		Yes
$2\chi_6^3$	$bZ_{(k,l)} = Z_{(k',l')}$	$l' = \begin{cases} k+4, & k < 4, l \geq 4 \\ l+4, & k < 4, l < 4k' \\ l-4, & k \geq 4 \end{cases}$	$k+l$ even	Yes
			$k+l$ odd	No
$\chi_{24}^4$	$bW_i = W_i$	$i = 1, 12, 23, 34$		Yes
	$bW_i = W_{i+1}$	$i = 2, 4, 13, 15, 24, 26, 35, 37$		Yes
		$i = 6, 17, 20, 28, 31, 39$		No
	$bW_i = W_{i+2}$	$i = 8, 9, 41, 42$		No
$bW_i = W_{i+3}$	$i = 19, 30$		No	

TABLE 4. Action of boson  $b$

dim	Objects	Twists	Number Count
1	$F(1), F(X_1), F(X_2), F(X_3)$	$1, e^{-\frac{\pi i}{4}}, -1, e^{-\frac{\pi i}{4}}$	4
$\chi_{24}^5$	$F(Y_0), F(Y_1), F(Y_2), F(Y_3)$	$1, e^{-\frac{\pi i}{4}}, -1, e^{-\frac{\pi i}{4}}$	4
$2\chi_6^3$	$F(Z_{0,2}), F(Z_{0,6}), F(Z_{1,3}), F(Z_{1,7})$	$1, 1, e^{-\frac{3\pi i}{4}}, e^{\frac{\pi i}{4}}$	4
$\chi_6^3$	$(Z_{0,4})_1, (Z_{0,4})_2, (Z_{1,5})_1, (Z_{1,5})_2$ $(Z_{2,6})_1, (Z_{2,6})_2, (Z_{3,7})_1, (Z_{3,7})_2$	$1, 1, e^{\frac{3\pi i}{4}}, e^{\frac{3\pi i}{4}}, -1, -1, e^{\frac{3\pi i}{4}}, e^{\frac{3\pi i}{4}}$	8
$\chi_{24}^4$	$F(W_2), F(W_4), F(W_{13}), F(W_{15}),$ $F(W_{24}), F(W_{26}), F(W_{35}), F(W_{37})$	$e^{\frac{5\pi i}{6}}, e^{-\frac{2\pi i}{3}}, e^{\frac{7\pi i}{12}}, e^{-\frac{11\pi i}{12}}$ $e^{\frac{\pi i}{3}}, e^{-\frac{\pi i}{6}}, e^{\frac{7\pi i}{12}}, e^{-\frac{11\pi i}{12}}$	8
$\chi_6^2$	$(W_1)_1, (W_1)_2, (W_{12})_1, (W_{12})_2$ $(W_{23})_1, (W_{23})_2, (W_{34})_1, (W_{34})_2$	$-i, -i, e^{-\frac{3\pi i}{4}}, e^{-\frac{3\pi i}{4}}$ $i, i, e^{\frac{\pi i}{3}}, e^{\frac{\pi i}{3}}$	8

TABLE 5. Twists of simple objects

integer and  $F(Z_{1,7})$  is the unique simple object with dimension  $2\chi_6^3$ . Therefore,  $\hat{\sigma}(F(Z_{1,7})) = F(Z_{1,7})$  for all  $\sigma \in \text{Gal}(\mathbb{Q}(T)/\mathbb{Q})$ , which implies that  $\frac{S_{V,F(Z_{1,7})}}{2\chi_6^3}$  is an integer for all  $V \in \mathcal{O}(\mathcal{D})$ . By [23, Theorem 2.2], we also have  $S_{(Z_{2,6})_1, F(Z_{1,7})} + S_{(Z_{2,6})_2, F(Z_{1,7})} = -4\chi_6^3$  and  $S_{(W_1)_1, F(Z_{1,7})} + S_{(W_1)_2, F(Z_{1,7})} = 0$ . Since  $\dim(\mathcal{D}) = \sum_{V \in \mathcal{O}(\mathcal{D})} S_{V,F(Z_{1,7})}^2$ , it follows that  $S_{(Z_{2,6})_i, F(Z_{1,7})} = -2\chi_6^3$  and  $S_{(W_1)_i, F(Z_{1,7})} = 0$  for  $i = 1, 2$ .  $\square$

**Lemma 3.3.2.**  $S_{(Z_{2,6})_i, F(W_j)} = 0$ ,  $S_{(W_1)_i, F(W_j)} = \chi_{24}^4$  for  $i = 1, 2$  and  $j = 2, 4$ .

*Proof.* By [23, Theorem 2.2], we have  $S_{(Z_{2,6})_1, F(W_j)} + S_{(Z_{2,6})_2, F(W_j)} = 0$  and  $S_{(W_1)_1, F(W_j)} + S_{(W_1)_2, F(W_j)} = 0$ . Let  $\sigma \in \text{Gal}(\mathbb{Q}(T)/\mathbb{Q})$  such that  $\sigma(\sqrt{6}) = -\sqrt{6}$ , then  $\sigma\left(\frac{S_{F(W_2), F(W_2)}}{\dim(F(W_2))}\right) = -1 = \frac{S_{F(W_2), \hat{\sigma}(F(W_2))}}{\dim(\hat{\sigma}(F(W_2)))}$ . It follows that  $S_{F(W_2), \hat{\sigma}(F(W_2))} = -\chi_{24}^4$  and  $\dim(\hat{\sigma}(F(W_2))) = \chi_{24}^4$ . Thus,  $\hat{\sigma}(F(W_2)) = F(W_2)$  and  $\hat{\sigma}(F(W_4)) = F(W_4)$ . In particular,  $\frac{S_{V,F(W_2)}}{\dim(F(W_2))}$  is an integer. Together with the relation  $\dim(\mathcal{D}) = \sum_{V \in \mathcal{O}(\mathcal{D})} S_{V,F(W_2)}^2$ , we have  $S_{(Z_{2,6})_i, F(W_2)} = 0$ ,  $S_{(W_1)_i, F(W_2)} = \chi_{24}^4$  for  $i = 1, 2$ . The same argument also works for object  $F(W_4)$ .  $\square$

**Theorem 3.3.3.** *The S-matrix and T-matrix of modular category  $\mathcal{D}$  are*

$$S = \begin{pmatrix} 1 & \chi_{24}^5 & 2\chi_6^3 & \chi_6^3 & \chi_6^3 & 2\chi_6^2 & 2\chi_6^2 & \chi_6^2 & \chi_6^2 \\ \chi_{24}^5 & 1 & 2\chi_6^3 & \chi_6^3 & \chi_6^3 & -2\chi_6^2 & -2\chi_6^2 & -\chi_6^2 & -\chi_6^2 \\ 2\chi_6^3 & 2\chi_6^3 & 0 & -2\chi_6^3 & -2\chi_6^3 & 0 & 0 & 0 & 0 \\ \chi_6^3 & \chi_6^3 & -2\chi_6^3 & u_1 & u_2 & 0 & 0 & u_3 & -u_3 \\ \chi_6^3 & \chi_6^3 & -2\chi_6^3 & u_2 & u_1 & 0 & 0 & -u_3 & u_3 \\ 2\chi_6^2 & -2\chi_6^2 & 0 & 0 & 0 & -2\chi_6^2 & 2\chi_6^2 & 2\chi_6^2 & 2\chi_6^2 \\ 2\chi_6^2 & -\chi_{24}^4 & 0 & 0 & 0 & 2\chi_6^2 & 2\chi_6^2 & -2\chi_6^2 & -2\chi_6^2 \\ \chi_6^2 & -\chi_6^2 & 0 & \sqrt{2}\chi_6^3 & -\sqrt{2}\chi_6^3 & 2\chi_6^2 & -2\chi_6^2 & u_5 & u_6 \\ \chi_6^2 & -\chi_6^2 & 0 & -\sqrt{2}\chi_6^3 & \sqrt{2}\chi_6^3 & 2\chi_6^2 & -2\chi_6^2 & u_6 & u_5 \end{pmatrix},$$

$$T = \text{diag}(1, 1, e^{\frac{\pi i}{4}}, -1, -1, e^{\frac{5\pi i}{6}}, e^{-\frac{2\pi i}{3}}, -e^{\frac{2\pi i}{4}}, -e^{\frac{2\pi i}{4}}),$$

where  $u_1 = \chi_2^1 \chi_6^3$ , and  $u_2 = (2 - \chi_2^1) \chi_6^3$ ,  $u_5 = \chi_3^1 \chi_6^2$ ,  $u_6 = (2 - \chi_3^1) \chi_6^2$ ,  $u_3 = \sqrt{2} \chi_6^3$ .

*Proof.* Let

$$\begin{aligned} u_1 &:= S_{(Z_{2,6})_1, (Z_{2,6})_1}, \quad u_2 := S_{(Z_{2,6})_1, (Z_{2,6})_2}, \quad u_3 := S_{(Z_{2,6})_1, (W_1)_1}, \quad u_4 := S_{(Z_{2,6})_2, (W_1)_1}, \\ u_5 &:= S_{(W_1)_1, (W_1)_1}, \quad u_6 := S_{(W_1)_1, (W_1)_2}, \quad u_7 := S_{(W_1)_2, (W_1)_2}. \end{aligned}$$

Applying orthogonality conditions, we obtain the following:

- From  $S_{F(Z_{1,7}), -}$  and  $S_{(Z_{2,6})_1, -}$ :  $u_1 + u_2 = 2\chi_6^3$ .
- From  $S_{(Z_{2,6})_1, -}$  and  $S_{(Z_{2,6})_2, -}$ :  $u_1 = S_{(Z_{2,6})_2, (Z_{2,6})_2}$ .
- By  $S_{(Z_{2,6})_1, -}$  and  $S_{F(W_2), -}$ :  $S_{(Z_{2,6})_1, (W_1)_2} = -u_3$ .
- Similarly, by  $S_{(Z_{2,6})_2, -}$  and  $S_{F(W_2), -}$  we have:  $S_{(Z_{2,6})_2, (W_1)_2} = -u_4$ .
- From  $S_{F(Z_{1,7}), -}$  and  $S_{(W_1)_1, -}$ :  $u_3 + u_4 = 0$ .
- From  $S_{(W_1)_1, -}$  and  $S_{F(W_2), -}$ , we get  $u_5 + u_6 = 2\chi_6^2$ .
- By  $S_{(W_1)_2, -}$  and  $S_{F(W_2), -}$ :  $u_7 + u_6 = 2\chi_6^2$ , implying  $u_5 = u_7$ .
- Finally, orthogonality between  $S_{(W_1)_1, -}$  and  $S_{(Z_{2,6})_1, -}$  gives  $u_3(u_1 - u_2 + u_5 - u_6) = 0$ .

Note that  $u_3$  can't be zero. Otherwise,  $u_1 u_2 = -3(\chi_6^3)^2$  and  $u_5 u_6 = -10(5 + 2\sqrt{6})$ , then  $N_{F(Y_6), (W_1)_2}^{(W_1)_2} = -\frac{1}{2}$ , which gives a contradiction. Therefore,  $(u_1 - u_2 + u_5 - u_6) = 0$ . We claim that  $(W_1)_1, (W_1)_2$  must be in the same orbit under the action of  $\text{Gal}(\mathcal{C})$ . Suppose not. Then for any simple object  $X$  and all  $\sigma \in \text{Gal}(\mathbb{Q}(T)/\mathbb{Q})$ ,  $\sigma\left(\frac{S_{X, (W_1)_1}}{\chi_6^2}\right) = \frac{S_{X, (W_1)_1}}{\chi_6^2}$ , implying  $S_{X, (W_1)_1} = a_X \chi_6^2$  for some integer  $a_X$ . Let  $S_{(Z_{2,6})_1, (W_1)_1} = a \chi_6^2$  and  $S_{(W_1)_1, (W_1)_1} = b \chi_6^2$ . This leads to the equation  $2a^2 + b^2 + (2 - b)^2 = 14$ , which has no integer solution. Notice that  $\sigma\left(\frac{S_{(Z_{2,6})_1, (W_1)_1}}{\chi_6^2}\right) = \frac{S_{(Z_{2,6})_1, (W_1)_2}}{\chi_6^2} = -\frac{S_{(Z_{2,6})_1, (W_1)_1}}{\chi_6^2}$  and  $\sigma\left(\frac{u_5}{\chi_6^2}\right) = \frac{u_6}{\chi_6^2}$ . Using a similar argument, we conclude that  $\frac{u_3}{\chi_6^3}$ ,  $\frac{u_1}{\chi_6^3}$  and  $\frac{u_2}{\chi_6^3}$  cannot be integers. Thus  $(Z_{2,6})_1$  and  $(Z_{2,6})_2$  are in the same Galois orbit, and for some  $\tau \in \text{Gal}(\mathbb{Q}(T)/\mathbb{Q})$ ,  $\tau\left(\frac{u_1}{\chi_6^3}\right) = \frac{u_2}{\chi_6^3}$ .

Applying the Verlinde formula, we obtain

$$N_{(W_1)_1, (W_1)_1}^{F(W_4)} = \frac{1}{\dim(\mathcal{D})} \sum_{V \in \mathcal{O}(\mathcal{D})} \frac{S_{(W_1)_1, V}^2 S_{F(W_4), V}}{\dim(V)} = \frac{2}{3} - \frac{u_5^2 + u_6^2}{12(\chi_6^2)^2}.$$

If  $u_5$  and  $u_6$  are real algebraic integers, notice that  $u_5, u_6 \neq 0$  while the fusion coefficient must be non-negative, we deduce  $u_5^2 + u_6^2 = 8(\chi_6^2)^2$ . Together with  $u_5 + u_6 = \chi_{24}^4$ , we have  $u_5, u_6 = \chi_6^2 \pm \sqrt{3}\chi_6^2$ . Thus we find that  $u_3 = \pm\sqrt{2}\chi_6^3$ , as desired.

If  $u_5^2 + u_6^2$  is negative, i.e.,  $\overline{u_5} = u_6$ , then  $((W_1)_1)^* = (W_1)_2$ , and  $u_5^2 + u_6^2 = -4(3k+1)(\chi_6^2)^2$  for some integer  $k \geq 0$ . Additionally,  $\dim(\mathcal{D}) = 10(\chi_6^2)^2 - 2u_3^2 + 2|u_5|^2$ , which implies  $|u_5|^2 - u_3^2 = 7(\chi_6^2)^2$ . By the balancing equation, we have  $|u_5| \leq \dim((W_1)_1)^2$  and  $|u_6| \leq \dim((W_1)_1)\dim((W_1)_2)$ , so

$$2(\chi_6^2)^4 \geq 2|u_5|^2 \geq |u_5^2 + u_6^2| = 4(3k+1)(\chi_6^2)^2,$$

implying  $k \leq 2$ . We now show this case cannot occur. In fact, notice that  $u_5, u_6$  are roots of equation  $x^2 - 2\chi_6^2x + \frac{1}{2}((2\chi_6^2)^2 - (u_5^2 + u_6^2)) = 0$ . Then

$$N_{(W_1)_1, (W_1)_1}^{(W_1)_1} = \frac{1}{\dim(\mathcal{D})} \sum_{V \in \mathcal{O}(\mathcal{D})} \frac{S_{(W_1)_1, V}^2 S_{(W_1)_2, V}}{\dim(V)} = \frac{k+1}{2},$$

thus  $k = 1$ . If  $k = 1$ , then

$$N_{(W_1)_1, (W_1)_1}^{F(W_2)} = \frac{1}{\dim(\mathcal{D})} \sum_{V \in \mathcal{O}(\mathcal{D})} \frac{S_{(W_1)_1, V} S_{F(W_2), V}}{\dim(V)} = -1,$$

it is a contradiction. Hence  $u_5, u_6, u_3$  must be real algebraic integers. Similarly, by determining the fusion coefficient  $N_{(Z_{2,6})_1, (Z_{2,6})_1}^{F(Z_{1,7})}$ , we obtain  $u_1^2 + u_2^2 = 8(\chi_6^3)^2$  and  $u_1 u_2 = -(\chi_6^3)^2$ , yielding  $u_1, u_2 = \chi_6^3 \pm \sqrt{2}\chi_6^3$ . Thus, we derive the desired modular data.  $\square$

The following modular data of  $\mathcal{C}(\mathfrak{g}_2, 4)$  can be produced using Sagemath [26]

$$S = \begin{pmatrix} 1 & \chi_6^2 & \chi_6^3 & 2\chi_6^2 & 2\chi_6^3 & \chi_6^2 & \chi_{24}^5 & 2\chi_6^2 & \chi_6^3 \\ \chi_6^2 & \chi_3^1 \chi_6^2 & \sqrt{2}\chi_6^3 & 2\chi_6^2 & 0 & (2 - \chi_3^1)\chi_6^2 & -\chi_6^2 & -2\chi_6^2 & -\sqrt{2}\chi_6^3 \\ \chi_6^3 & \sqrt{2}\chi_6^3 & (2 - \chi_2^1)\chi_6^3 & 0 & -2\chi_6^3 & -\sqrt{2}\chi_6^3 & \chi_6^3 & 0 & \chi_2^1 \chi_6^3 \\ 2\chi_6^2 & 2\chi_6^2 & 0 & -2\chi_6^2 & 0 & 2\chi_6^2 & -2\chi_6^2 & 2\chi_6^2 & 0 \\ 2\chi_6^3 & 0 & -2\chi_6^3 & 0 & 0 & 0 & 2\chi_6^3 & 0 & -2\chi_6^3 \\ \chi_6^2 & (2 - \chi_3^1)\chi_6^2 & -\sqrt{2}\chi_6^3 & 2\chi_6^2 & 0 & \chi_3^1 \chi_6^2 & -\chi_6^2 & -2\chi_6^2 & \sqrt{2}\chi_6^3 \\ \chi_{24}^5 & -\chi_6^2 & \chi_6^3 & -2\chi_6^2 & 2\chi_6^3 & -\chi_6^2 & 1 & -2\chi_6^2 & \chi_6^3 \\ 2\chi_6^2 & -2\chi_6^2 & 0 & 2\chi_6^2 & 0 & -2\chi_6^2 & -2\chi_6^2 & 2\chi_6^2 & 0 \\ \chi_6^3 & -\sqrt{2}\chi_6^3 & \chi_2^1 \chi_6^3 & 0 & -2\chi_6^3 & \sqrt{2}\chi_6^3 & \chi_6^3 & 0 & (2 - \chi_2^1)\chi_6^3 \end{pmatrix}$$

$$T = \text{diag}(1, \zeta_4, -1, \zeta_{12}^7, -\zeta_8^3, \zeta_4, 1, \zeta_3, -1).$$

**Corollary 3.3.4.**  $\mathcal{D}$  has the same modular data as  $\mathcal{C}(\mathfrak{g}_2, 4)$ .

*Proof.* We replace  $\zeta_8$  and  $\zeta_3$  by their inverses in modular category  $\mathcal{C}(\mathfrak{g}_2, 4)$  (i.e., replace  $\zeta_{24}$  by its inverse), then we get the desired ribbon twists.  $\square$

#### ACKNOWLEDGMENTS

The authors thank Yilong Wang for helpful conversations. The second author also thanks Eric Rowell, Siu-Hung Ng, Pinhas Grossman and Andrew Schopieray for their discussions during the SLMath Quantum Symmetry Reunion in 2024. The first author is supported by the National Natural Science Foundation of China (no.12101541).



APPENDIX A. SOLUTIONS FOR  $(\omega_j, \tau_j, \xi_j)$

In this section, we list the solutions of  $(\omega_j, \tau_j, \xi_j)$  from Equations (4) – (7) for the near-group categories in Section 3.1 and Section 3.3.

The following table gives the solutions  $(\omega_j, \tau_j, \xi_j)$  for the near-group category  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} + 16$ , where  $j = 1, \dots, 152$ . Each  $\omega_j$  is of the form  $\zeta_{80}^k$  for some integer  $k$ , with the corresponding  $k$  values provided in the relevant column. Since  $\xi_j(x) = \exp(i\theta_{j,x})$ , the table records the values  $\theta_{j,x}$  in the column for  $\xi$ , ordered by  $x = (0, 0), \dots, (0, 3), (1, 0), \dots, (3, 3) \in \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ .

#	$\omega$	$\tau$	$\xi$
1	32	(0, 0)	1.25664, 0.13318, -1.88496, -2.33229, -2.55613, -0.706082, 1.11454, 0.35208, -1.88496, -1.57477, -1.88496, -0.62434, -2.78457, 2.16119, -0.172061, -3.06383
2	32	(0, 0)	-1.88496, -2.33229, 1.25664, 0.13318, 1.11454, 0.35208, -2.55613, -0.706082, -1.88496, -0.62434, -1.88496, -1.57477, -0.172061, -3.06383, -2.78457, 2.16119
3	32	(0, 0)	-1.88496, -0.62434, -1.88496, -1.57477, -0.172061 - 3.06383, -2.78457, 2.16119, -1.88496, -2.33229, 1.25664, 0.13318, 1.11454, 0.35208, -2.55613, -0.706082
4	32	(0, 0)	-1.88496, -1.57477, -1.88496, -0.62434, -2.78457, 2.16119, -0.172061, -3.06383, 1.25664, 0.13318, -1.88496, -2.33229, -2.55613, -0.706082, 1.11454, 0.35208
5	48	(0, 0)	1.88496, -1.11454, 1.88496, 0.172061, 2.33229, -0.35208, 0.62434, 3.06383, -1.25664, 2.55613, 1.88496, 2.78457, -0.13318, 0.706082, 1.57477, -2.16119
6	48	(0, 0)	1.88496, -1.11454, 1.88496, 0.172061, 2.33229, -0.35208, 0.62434, 3.06383, -1.25664, 2.55613, 1.88496, 2.78457, -0.13318, 0.706082, 1.57477, -2.16119
7	48	(0, 0)	1.88496, 0.172061, 1.88496, -1.11454, 0.62434, 3.06383, 2.33229, -0.35208, 1.88496, 2.78457, -1.25664, 2.55613, 1.57477, -2.16119, -0.13318, 0.706082
8	48	(0, 0)	1.88496, 2.78457, -1.25664, 2.55613, 1.57477, -2.16119, -0.13318, 0.706082, 1.88496, 0.172061, 1.88496, -1.11454, 0.62434, 3.06383, 2.33229, -0.35208
9	7	(0, 1)	-0.883402, -0.923013, 3.0903, -1.75512, -3.07932, -2.22175, -1.62208, 1.30524, 0.974834, -2.78125, -0.625598, 1.96077, -1.15546, -0.297893, -1.54086, 1.38646
10	7	(0, 1)	0.974834, -2.78125, -0.625598, 1.96077, -1.15546, -0.297893, -1.54086, 1.38646, -0.883402, -0.923013, 3.0903, -1.75512, -3.07932, -2.22175, -1.62208, 1.30524
11	15	(0, 1)	-2.88492, 1.70682, -0.169554, 2.13305, -0.937357, -1.81154, 0.572197, -0.179498, -2.88492, 1.70682, -0.169554, 2.13305, -0.937357, -1.81154, 0.572197, -0.179498
12	23	(0, 1)	0.521724, -1.0715, -0.173026, 2.76484, 2.2799, 2.94948, -1.23134, -1.42571, -1.38096, 0.83118, 1.16544, 1.42637, 1.21313, 1.88271, 2.44672, 2.25236
13	23	(0, 1)	-1.38096, 0.83118, 1.16544, 1.42637, 1.21313, 1.88271, 2.44672, 2.25236, 0.521724, -1.0715, -0.173026, 2.76484, 2.2799, 2.94948, -1.23134, -1.42571
14	47	(0, 1)	-0.625598, 1.96077, 0.974834, -2.78125, -1.54086, 1.38646, -1.15546, -0.297893, 3.0903, -1.75512, -0.883402, -0.923013, -1.62208, 1.30524, -3.07932, -2.22175
15	47	(0, 1)	3.0903, -1.75512, -0.883402, -0.923013, -1.62208, 1.30524, -3.07932, -2.22175, -0.625598, 1.96077, 0.974834, -2.78125, -1.54086, 1.38646, -1.15546, -0.297893
16	55	(0, 1)	-0.169554, 2.13305, -2.88492, 1.70682, 0.572197, -0.179498, -0.937357, -1.81154, -0.169554, 2.13305, -2.88492, 1.70682, 0.572197, -0.179498, -0.937357, -1.81154
17	63	(0, 1)	1.16544, 1.42637, -1.38096, 0.83118, 2.44672, 2.25236, 1.21313, 1.88271, -0.173026, 2.76484, 0.521724, -1.0715, -1.23134, -1.42571, 2.2799, 2.94948
18	63	(0, 1)	-0.173026, 2.76484, 0.521724, -1.0715, -1.23134, -1.42571, 2.2799, 2.94948, 1.16544, 1.42637, -1.38096, 0.83118, 2.44672, 2.25236, 1.21313, 1.88271

*Continued on next page*

#	$\omega$	$\tau$	$\xi$
19	12	(0, 2)	0.13318, -1.88496, -2.33229, 1.25664, -0.706082, 1.11454, 0.35208, -2.55613, -1.57477, -1.88496, -0.62434, -1.88496, 2.16119, -0.172061, -3.06383, -2.78457
20	12	(0, 2)	-1.57477, -1.88496, -0.62434, -1.88496, 2.16119, -0.172061, -3.06383, -2.78457, 0.13318, -1.88496, -2.33229, 1.25664, -0.706082, 1.11454, 0.35208, -2.55613
21	12	(0, 2)	-0.62434, -1.88496, -1.57477, -1.88496, -3.06383, -2.78457, 2.16119, -0.172061, -2.33229, 1.25664, 0.13318, -1.88496, 0.35208, -2.55613, -0.706082, 1.11454
22	12	(0, 2)	-2.33229, 1.25664, 0.13318, -1.88496, 0.35208, -2.55613, -0.706082, 1.11454, -0.62434, -1.88496, -1.57477, -1.88496, -3.06383, -2.78457, 2.16119, -0.172061
23	28	(0, 2)	2.55613, 1.88496, 2.78457, -1.25664, 0.706082, 1.57477, -2.16119, -0.13318, -1.11454, 1.88496, 0.172061, 1.88496, -0.35208, 0.62434, 3.06383, 2.33229
24	28	(0, 2)	0.172061, 1.88496, -1.11454, 1.88496, 3.06383, 2.33229, -0.35208, 0.62434, 2.78457, -1.25664, 2.55613, 1.88496, -2.16119, -0.13318, 0.706082, 1.57477
25	28	(0, 2)	2.78457, -1.25664, 2.55613, 1.88496, -2.16119, -0.13318, 0.706082, 1.57477, 0.172061, 1.88496, -1.11454, 1.88496, 3.06383, 2.33229, -0.35208, 0.62434
26	28	(0, 2)	-1.11454, 1.88496, 0.172061, 1.88496, -0.35208, 0.62434, 3.06383, 2.33229, 2.55613, 1.88496, 2.78457, -1.25664, 0.706082, 1.57477, -2.16119, -0.13318
27	7	(0, 3)	-2.78125, -0.625598, 1.96077, 0.974834, -0.297893, -1.54086, 1.38646, -1.15546, -0.923013, 3.0903, -1.75512, -0.883402, -2.22175, -1.62208, 1.30524, -3.07932
28	7	(0, 3)	-0.923013, 3.0903, -1.75512, -0.883402, -2.22175, -1.62208, 1.30524, -3.07932, -2.78125, -0.625598, 1.96077, 0.974834, -0.297893, -1.54086, 1.38646, -1.15546
29	15	(0, 3)	1.70682, -0.169554, 2.13305, -2.88492, -1.81154, 0.572197, -0.179498, -0.937357, 1.70682, -0.169554, 2.13305, -2.88492, -1.81154, 0.572197, -0.179498, -0.937357
30	23	(0, 3)	0.83118, 1.16544, 1.42637, -1.38096, 1.88271, 2.44672, 2.25236, 1.21313, -1.0715, -0.173026, 2.76484, 0.521724, 2.94948, -1.23134, -1.42571, 2.2799
31	23	(0, 3)	-1.0715, -0.173026, 2.76484, 0.521724, 2.94948, -1.23134, -1.42571, 2.2799, 0.83118, 1.16544, 1.42637, -1.38096, 1.88271, 2.44672, 2.25236, 1.21313
32	47	(0, 3)	1.96077, 0.974834, -2.78125, -0.625598, 1.38646, -1.15546, -0.297893, -1.54086, -1.75512, -0.883402, -0.923013, 3.0903, 1.30524, -3.07932, -2.22175, -1.62208
33	47	(0, 3)	-1.75512, -0.883402, -0.923013, 3.0903, 1.30524, -3.07932, -2.22175, -1.62208, 1.96077, 0.974834, -2.78125, -0.625598, 1.38646, -1.15546, -0.297893, -1.54086
34	55	(0, 3)	2.13305, -2.88492, 1.70682, -0.169554, -0.179498, -0.937357, -1.81154, 0.572197, 2.13305, -2.88492, 1.70682, -0.169554, -0.179498, -0.937357, -1.81154, 0.572197
35	63	(0, 3)	2.76484, 0.521724, -1.0715, -0.173026, -1.42571, 2.2799, 2.94948, -1.23134, 1.42637, -1.38096, 0.83118, 1.16544, 2.25236, 1.21313, 1.88271, 2.44672
36	63	(0, 3)	1.42637, -1.38096, 0.83118, 1.16544, 2.25236, 1.21313, 1.88271, 2.44672, 2.76484, 0.521724, -1.0715, -0.173026, -1.42571, 2.2799, 2.94948, -1.23134
37	17	(1, 0)	-1.16544, -2.44672, 0.173026, 1.23134, -1.42637, -2.25236, -2.76484, 1.42571, 1.38096, -1.21313, -0.521724, -2.2799, -0.83118, -1.88271, 1.0715, -2.94948
38	17	(1, 0)	0.173026, 1.23134, -1.16544, -2.44672, -2.76484, 1.42571, -1.42637, -2.25236, -0.521724, -2.2799, 1.38096, -1.21313, 1.0715, -2.94948, -0.83118, -1.88271
39	25	(1, 0)	0.169554, -0.572197, 0.169554, -0.572197, -2.13305, 0.179498, -2.13305, 0.179498, 2.88492, 0.937357, 2.88492, 0.937357, -1.70682, 1.81154, -1.70682, 1.81154
40	33	(1, 0)	0.625598, 1.54086, -3.0903, 1.62208, -1.96077, -1.38646, 1.75512, -1.30524, -0.974834, 1.15546, 0.883402, 3.07932, 2.78125, 0.297893, 0.923013, 2.22175
41	33	(1, 0)	-3.0903, 1.62208, 0.625598, 1.54086, 1.75512, -1.30524, -1.96077, -1.38646, 0.883402, 3.07932, -0.974834, 1.15546, 0.923013, 2.22175, 2.78125, 0.297893

Continued on next page

#	$\omega$	$\tau$	$\xi$
42	57	(1, 0)	-0.521724, -2.2799, 1.38096, -1.21313, 1.0715, -2.94948, -0.83118, -1.88271, 0.173026, 1.23134, -1.16544, -2.44672, -2.76484, 1.42571, -1.42637, -2.25236
43	57	(1, 0)	1.38096, -1.21313, -0.521724, -2.2799, -0.83118, -1.88271, 1.0715, -2.94948, -1.16544, -2.44672, 0.173026, 1.23134, -1.42637, -2.25236, -2.76484, 1.42571
44	65	(1, 0)	2.88492, 0.937357, 2.88492, 0.937357, -1.70682, 1.81154, -1.70682, 1.81154, 0.169554, -0.572197, 0.169554, -0.572197, -2.13305, 0.179498, -2.13305, 0.179498
45	73	(1, 0)	0.883402, 3.07932, -0.974834, 1.15546, 0.923013, 2.22175, 2.78125, 0.297893, -3.0903, 1.62208, 0.625598, 1.54086, 1.75512, -1.30524, -1.96077, -1.38646
46	73	(1, 0)	-0.974834, 1.15546, 0.883402, 3.07932, 2.78125, 0.297893, 0.923013, 2.22175, 0.625598, 1.54086, -3.0903, 1.62208, -1.96077, -1.38646, 1.75512, -1.30524
47	12	(1, 1)	2.14417, 0.33666, 0.50165, 2.34429, 0.605818, -1.20169, 1.73978, -2.70077, 3.05334, 1.2107, -2.63533, 2.03946, 2.87337, 1.03073, -1.09698, -2.70538
48	12	(1, 1)	-2.63533, 2.03946, 3.05334, 1.2107, -1.09698, -2.70538, 2.87337, 1.03073, 0.50165, 2.34429, 2.14417, 0.33666, 1.73978, -2.70077, 0.605818, -1.20169
49	20	(1, 1)	-1.23768, -1.23768, 1.23768, -2.80847, 2.80847, 2.80847, 1.23768, -2.80847, 1.23768, -2.80847, -1.23768, -1.23768, 1.23768, -2.80847, 2.80847, 2.80847
50	28	(1, 1)	-3.05334, -2.87337, -2.14417, -0.605818, -1.2107, -1.03073, -0.33666, 1.20169, 2.63533, 1.09698, -0.50165, -1.73978, -2.03946, 2.70538, -2.34429, 2.70077
51	28	(1, 1)	-0.50165, -1.73978, 2.63533, 1.09698, -2.34429, 2.70077, -2.03946, 2.70538, -2.14417, -0.605818, -3.05334, -2.87337, -0.33666, 1.20169, -1.2107, -1.03073
52	52	(1, 1)	3.05334, 1.2107, -2.63533, 2.03946, 2.87337, 1.03073, -1.09698, -2.70538, 2.14417, 0.33666, 0.50165, 2.34429, 0.605818, -1.20169, 1.73978, -2.70077
53	52	(1, 1)	0.50165, 2.34429, 2.14417, 0.33666, 1.73978, -2.70077, 0.605818, -1.20169, -2.63533, 2.03946, 3.05334, 1.2107, -1.09698, -2.70538, 2.87337, 1.03073
54	60	(1, 1)	1.23768, -2.80847, -1.23768, -1.23768, 1.23768, -2.80847, 2.80847, 2.80847, -1.23768, -1.23768, 1.23768, -2.80847, 2.80847, 2.80847, 1.23768, -2.80847
55	68	(1, 1)	-2.14417, -0.605818, -3.05334, -2.87337, -0.33666, 1.20169, -1.2107, -1.03073, -0.50165, -1.73978, 2.63533, 1.09698, -2.34429, 2.70077, -2.03946, 2.70538
56	68	(1, 1)	2.63533, 1.09698, -0.50165, -1.73978, -2.03946, 2.70538, -2.34429, 2.70077, -3.05334, -2.87337, -2.14417, -0.605818, -1.2107, -1.03073, -0.33666, 1.20169
57	5	(1, 2)	-0.572197, 0.169554, -0.572197, 0.169554, 0.179498, -2.13305, 0.179498, -2.13305, 0.937357, 2.88492, 0.937357, 2.88492, 1.81154, -1.70682, 1.81154, -1.70682
58	13	(1, 2)	1.54086, -3.0903, 1.62208, 0.625598, -1.38646, 1.75512, -1.30524, -1.96077, 1.15546, 0.883402, 3.07932, -0.974834, 0.297893, 0.923013, 2.22175, 2.78125
59	13	(1, 2)	1.62208, 0.625598, 1.54086, -3.0903, -1.30524, -1.96077, -1.38646, 1.75512, 3.07932, -0.974834, 1.15546, 0.883402, 2.22175, 2.78125, 0.297893, 0.923013
60	37	(1, 2)	-2.2799, 1.38096, -1.21313, -0.521724, -2.94948, -0.83118, -1.88271, 1.0715, 1.23134, -1.16544, -2.44672, 0.173026, 1.42571, -1.42637, -2.25236, -2.76484
61	37	(1, 2)	-1.21313, -0.521724, -2.2799, 1.38096, -1.88271, 1.0715, -2.94948, -0.83118, -2.44672, 0.173026, 1.23134, -1.16544, -2.25236, -2.76484, 1.42571, -1.42637
62	45	(1, 2)	0.937357, 2.88492, 0.937357, 2.88492, 1.81154, -1.70682, 1.81154, -1.70682, -0.572197, 0.169554, -0.572197, 0.169554, 0.179498, -2.13305, 0.179498, -2.13305
63	43	(1, 2)	1.15546, 0.883402, 3.07932, -0.974834, 0.297893, 0.923013, 2.22175, 2.78125, 1.54086, -3.0903, 1.62208, 0.625598, -1.38646, 1.75512, -1.30524, -1.96077
64	53	(1, 2)	3.07932, -0.974834, 1.15546, 0.883402, 2.22175, 2.78125, 0.297893, 0.923013, 1.62208, 0.625598, 1.54086, -3.0903, -1.30524, -1.96077, -1.38646, 1.75512

Continued on next page

#	$\omega$	$\tau$	$\xi$
65	77	(1, 2)	1.23134, -1.16544, -2.44672, 0.173026, 1.42571, -1.42637, -2.25236, -2.76484, -2.2799, 1.38096, -1.21313, -0.521724, -2.94948, -0.83118, -1.88271, 1.0715
66	77	(1, 2)	-2.44672, 0.173026, 1.23134, -1.16544, -2.25236, -2.76484, 1.42571, -1.42637, -1.21313, -0.521724, -2.2799, 1.38096, -1.88271, 1.0715, -2.94948, -0.83118
67	12	(1, 3)	0.33666, 0.50165, 2.34429, 2.14417, -1.20169, 1.73978, -2.70077, 0.605818, 1.2107, -2.63533, 2.03946, 3.05334, 1.03073, -1.09698, -2.70538, 2.87337
68	12	(1, 3)	2.03946, 3.05334, 1.2107, -2.63533, -2.70538, 2.87337, 1.03073, -1.09698, 2.34429, 2.14417, 0.33666, 0.50165, -2.70077, 0.605818, -1.20169, 1.73978
69	20	(1, 3)	-1.23768, 1.23768, -2.80847, -1.23768, 2.80847, 1.23768, -2.80847, 2.80847, -2.80847, -1.23768, -1.23768, 1.23768, -2.80847, 2.80847, 2.80847, 1.23768
70	28	(1, 3)	-2.87337, -2.14417, -0.605818, -3.05334, -1.03073, -0.33666, 1.20169, -1.2107, 1.09698, -0.50165, -1.73978, 2.63533, 2.70538, -2.34429, 2.70077, -2.03946
71	28	(1, 3)	-1.73978, 2.63533, 1.09698, -0.50165, 2.70077, -2.03946, 2.70538, -2.34429, -0.605818, -3.05334, -2.87337, -2.14417, 1.20169, -1.2107, -1.03073, -0.33666
72	52	(1, 3)	2.34429, 2.14417, 0.33666, 0.50165, -2.70077, 0.605818, -1.20169, 1.73978, 2.03946, 3.05334, 1.2107, -2.63533, -2.70538, 2.87337, 1.03073, -1.09698
73	52	(1, 3)	1.2107, -2.63533, 2.03946, 3.05334, 1.03073, -1.09698, -2.70538, 2.87337, 0.33666, 0.50165, 2.34429, 2.14417, -1.20169, 1.73978, -2.70077, 0.605818
74	60	(1, 3)	-2.80847, -1.23768, -1.23768, 1.23768, -2.80847, 2.80847, 2.80847, 1.23768, -1.23768, 1.23768, -2.80847, -1.23768, 2.80847, 1.23768, -2.80847, 2.80847
75	68	(1, 3)	-0.605818, -3.05334, -2.87337, -2.14417, 1.20169, -1.2107, -1.03073, -0.33666, -1.73978, 2.63533, 1.09698, -0.50165, 2.70077, -2.03946, 2.70538, -2.34429
76	68	(1, 3)	1.09698, -0.50165, -1.73978, 2.63533, 2.70538, -2.34429, 2.70077, -2.03946, -2.87337, -2.14417, -0.605818, -3.05334, -1.03073, -0.33666, 1.20169, -1.2107
77	52	(2, 0)	-2.55613, -0.706082, 1.11454, 0.35208, -1.88496, -1.57477, -1.88496, -0.62434, -2.78457, 2.16119, -0.172061, -3.06383, 1.25664, 0.13318, -1.88496, -2.33229
78	52	(2, 0)	-0.172061, -3.06383, -2.78457, 2.16119, -1.88496, -2.33229, 1.25664, 0.13318, 1.11454, 0.35208, -2.55613, -0.706082, -1.88496, -0.62434, -1.88496, -1.57477
79	52	(2, 0)	-2.78457, 2.16119, -0.172061, -3.06383, 1.25664, 0.13318, -1.88496, -2.33229, -2.55613, -0.706082, 1.11454, 0.35208, -1.88496, -1.57477, -1.88496, -0.62434
80	52	(2, 0)	1.11454, 0.35208, -2.55613, -0.706082, -1.88496, -0.62434, -1.88496, -1.57477, -0.172061, -3.06383, -2.78457, 2.16119, -1.88496, -2.33229, 1.25664, 0.13318
81	68	(2, 0)	-0.13318, 0.706082, 1.57477, -2.16119, 1.88496, -1.11454, 1.88496, 0.172061, 2.33229, -0.35208, 0.62434, 3.06383, -1.25664, 2.55613, 1.88496, 2.78457
82	68	(2, 0)	1.57477, -2.16119, -0.13318, 0.706082, 1.88496, 0.172061, 1.88496, -1.11454, 0.62434, 3.06383, 2.33229, -0.35208, 1.88496, 2.78457, -1.25664, 2.55613
83	68	(2, 0)	0.62434, 3.06383, 2.33229, -0.35208, 1.88496, 2.78457, -1.25664, 2.55613, 1.57477, -2.16119, -0.13318, 0.706082, 1.88496, 0.172061, 1.88496, -1.11454
84	68	(2, 0)	2.33229, -0.35208, 0.62434, 3.06383, -1.25664, 2.55613, 1.88496, 2.78457, -0.13318, 0.706082, 1.57477, -2.16119, 1.88496, -1.11454, 1.88496, 0.172061
85	3	(2, 1)	-1.23134, -1.42571, 2.2799, 2.94948, 1.16544, 1.42637, -1.38096, 0.83118, 2.44672, 2.25236, 1.21313, 1.88271, -0.173026, 2.76484, 0.521724, -1.0715
86	3	(2, 1)	2.44672, 2.25236, 1.21313, 1.88271, -0.173026, 2.76484, 0.521724, -1.0715, -1.23134, -1.42571, 2.2799, 2.94948, 1.16544, 1.42637, -1.38096, 0.83118
87	27	(2, 1)	-1.15546, -0.297893, -1.54086, 1.38646, -0.883402, -0.923013, 3.0903, -1.75512, -3.07932, -2.22175, -1.62208, 1.30524, 0.974834, -2.78125, -0.625598, 1.96077

Continued on next page

#	$\omega$	$\tau$	$\xi$
88	27	(2, 1)	-3.07932, -2.22175, -1.62208, 1.30524, 0.974834, -2.78125, -0.625598, 1.96077, -1.15546, -0.297893, -1.54086, 1.38646, -0.883402, -0.923013, 3.0903, -1.75512
89	35	(2, 1)	-0.937357, -1.81154, 0.572197, -0.179498, -2.88492, 1.70682, -0.169554, 2.13305, -0.937357, -1.81154, 0.572197, -0.179498, -2.88492, 1.70682, -0.169554, 2.13305
90	43	(2, 1)	2.2799, 2.94948, -1.23134, -1.42571, -1.38096, 0.83118, 1.16544, 1.42637, 1.21313, 1.88271, 2.44672, 2.25236, 0.521724, -1.0715, -0.173026, 2.76484
91	43	(2, 1)	1.21313, 1.88271, 2.44672, 2.25236, 0.521724, -1.0715, -0.173026, 2.76484, 2.2799, 2.94948, -1.23134, -1.42571, -1.38096, 0.83118, 1.16544, 1.42637
92	67	(2, 1)	-1.54086, 1.38646, -1.15546, -0.297893, 3.0903, -1.75512, -0.883402, -0.923013, -1.62208, 1.30524, -3.07932, -2.22175, -0.625598, 1.96077, 0.974834, -2.78125
93	67	(2, 1)	-1.62208, 1.30524, -3.07932, -2.22175, -0.625598, 1.96077, 0.974834, -2.78125, -1.54086, 1.38646, -1.15546, -0.297893, 3.0903, -1.75512, -0.883402, -0.923013
94	75	(2, 1)	0.572197, -0.179498, -0.937357, -1.81154, -0.169554, 2.13305, -2.88492, 1.70682, 0.572197, -0.179498, -0.937357, -1.81154, -0.169554, 2.13305, -2.88492, 1.70682
95	32	(2, 2)	-0.706082, 1.11454, 0.35208, -2.55613, -1.57477, -1.88496, -0.62434, -1.88496, 2.16119, -0.172061, -3.06383, -2.78457, 0.13318, -1.88496, -2.33229, 1.25664
96	32	(2, 2)	-3.06383, -2.78457, 2.16119, -0.172061, -2.33229, 1.25664, 0.13318, -1.88496, 0.35208, -2.55613, -0.706082, 1.11454, -0.62434, -1.88496, -1.57477, -1.88496
97	32	(2, 2)	2.16119, -0.172061, -3.06383, -2.78457, 0.13318, -1.88496, -2.33229, 1.25664, -0.706082, 1.11454, 0.35208, -2.55613, -1.57477, -1.88496, -0.62434, -1.88496
98	32	(2, 2)	0.35208, -2.55613, -0.706082, 1.11454, -0.62434, -1.88496, -1.57477, -1.88496, -3.06383, -2.78457, 2.16119, -0.172061, -2.33229, 1.25664, 0.13318, -1.88496
99	48	(2, 2)	0.706082, 1.57477, -2.16119, -0.13318, -1.11454, 1.88496, 0.172061, 1.88496, -0.35208, 0.62434, 3.06383, 2.33229, 2.55613, 1.88496, 2.78457, -1.25664
100	48	(2, 2)	-2.16119, -0.13318, 0.706082, 1.57477, 0.172061, 1.88496, -1.11454, 1.88496, 3.06383, 2.33229, -0.35208, 0.62434, 2.78457, -1.25664, 2.55613, 1.88496
101	48	(2, 2)	3.06383, 2.33229, -0.35208, 0.62434, 2.78457, -1.25664, 2.55613, 1.88496, -2.16119, -0.13318, 0.706082, 1.57477, 0.172061, 1.88496, -1.11454, 1.88496
102	48	(2, 2)	-0.35208, 0.62434, 3.06383, 2.33229, 2.55613, 1.88496, 2.78457, -1.25664, 0.706082, 1.57477, -2.16119, -0.13318, -1.11454, 1.88496, 0.172061, 1.88496
103	3	(2, 3)	2.25236, 1.21313, 1.88271, 2.44672, 2.76484, 0.521724, -1.0715, -0.173026, -1.42571, 2.2799, 2.94948, -1.23134, 1.42637, -1.38096, 0.83118, 1.16544
104	3	(2, 3)	-1.42571, 2.2799, 2.94948, -1.23134, 1.42637, -1.38096, 0.83118, 1.16544, 2.25236, 1.21313, 1.88271, 2.44672, 2.76484, 0.521724, -1.0715, -0.173026
105	27	(2, 3)	-0.297893, -1.54086, 1.38646, -1.15546, -0.923013, 3.0903, -1.75512, -0.883402, -2.22175, -1.62208, 1.30524, -3.07932, -2.78125, -0.625598, 1.96077, 0.974834
106	27	(2, 3)	-2.22175, -1.62208, 1.30524, -3.07932, -2.78125, -0.625598, 1.96077, 0.974834, -0.297893, -1.54086, 1.38646, -1.15546, -0.923013, 3.0903, -1.75512, -0.883402
107	35	(2, 3)	-1.81154, 0.572197, -0.179498, -0.937357, 1.70682, -0.169554, 2.13305, -2.88492, -1.81154, 0.572197, -0.179498, -0.937357, 1.70682, -0.169554, 2.13305, -2.88492
108	43	(2, 3)	1.88271, 2.44672, 2.25236, 1.21313, -1.0715, -0.173026, 2.76484, 0.521724, 2.94948, -1.23134, -1.42571, 2.2799, 0.83118, 1.16544, 1.42637, -1.38096
109	43	(2, 3)	2.94948, -1.23134, -1.42571, 2.2799, 0.83118, 1.16544, 1.42637, -1.38096, 1.88271, 2.44672, 2.25236, 1.21313, -1.0715, -0.173026, 2.76484, 0.521724
110	67	(2, 3)	1.30524, -3.07932, -2.22175, -1.62208, 1.96077, 0.974834, -2.78125, -0.625598, 1.38646, -1.15546, -0.297893, -1.54086, -1.75512, -0.883402, -0.923013, 3.0903

Continued on next page

#	$\omega$	$\tau$	$\xi$
111	67	(2, 3)	1.38646, -1.15546, -0.297893, -1.54086, -1.75512, -0.883402, -0.923013, 3.0903, 1.30524, -3.07932, -2.22175, -1.62208, 1.96077, 0.974834, -2.78125, -0.625598
112	75	(2, 3)	-0.179498, -0.937357, -1.81154, 0.572197, 2.13305, -2.88492, 1.70682, -0.169554, -0.179498, -0.937357, -1.81154, 0.572197, 2.13305, -2.88492, 1.70682, -0.169554
113	17	(3, 0)	-2.76484, 1.42571, -1.42637, -2.25236, -0.521724, -2.2799, 1.38096, -1.21313, 1.0715, -2.94948, -0.83118, -1.88271, 0.173026, 1.23134, -1.16544, -2.44672
114	17	(3, 0)	-1.42637, -2.25236, -2.76484, 1.42571, 1.38096, -1.21313, -0.521724, -2.2799, -0.83118, -1.88271, 1.0715, -2.94948, -1.16544, -2.44672, 0.173026, 1.23134
115	25	(3, 0)	-2.13305, 0.179498, -2.13305, 0.179498, 2.88492, 0.937357, 2.88492, 0.937357, -1.70682, 1.81154, -1.70682, 1.81154, 0.169554, -0.572197, 0.169554, -0.572197
116	33	(3, 0)	-1.96077, -1.38646, 1.75512, -1.30524, -0.974834, 1.15546, 0.883402, 3.07932, 2.78125, 0.297893, 0.923013, 2.22175, 0.625598, 1.54086, -3.0903, 1.62208
117	33	(3, 0)	1.75512, -1.30524, -1.96077, -1.38646, 0.883402, 3.07932, -0.974834, 1.15546, 0.923013, 2.22175, 2.78125, 0.297893, -3.0903, 1.62208, 0.625598, 1.54086
118	57	(3, 0)	-0.83118, -1.88271, 1.0715, -2.94948, -1.16544, -2.44672, 0.173026, 1.23134, -1.42637, -2.25236, -2.76484, 1.42571, 1.38096, -1.21313, -0.521724, -2.2799
119	57	(3, 0)	1.0715, -2.94948, -0.83118, -1.88271, 0.173026, 1.23134, -1.16544, -2.44672, -2.76484, 1.42571, -1.42637, -2.25236, -0.521724, -2.2799, 1.38096, -1.21313
120	65	(3, 0)	-1.70682, 1.81154, -1.70682, 1.81154, 0.169554, -0.572197, 0.169554, -0.572197, -2.13305, 0.179498, -2.13305, 0.179498, 2.88492, 0.937357, 2.88492, 0.937357
121	73	(3, 0)	2.78125, 0.297893, 0.923013, 2.22175, 0.625598, 1.54086, -3.0903, 1.62208, -1.96077, -1.38646, 1.75512, -1.30524, -0.974834, 1.15546, 0.883402, 3.07932
122	73	(3, 0)	0.923013, 2.22175, 2.78125, 0.297893, -3.0903, 1.62208, 0.625598, 1.54086, 1.75512, -1.30524, -1.96077, -1.38646, 0.883402, 3.07932, -0.974834, 1.15546
123	12	(3, 1)	0.605818, -1.20169, 1.73978, -2.70077, 3.05334, 1.2107, -2.63533, 2.03946, 2.87337, 1.03073, -1.09698, -2.70538, 2.14417, 0.33666, 0.50165, 2.34429
124	12	(3, 1)	-1.09698, -2.70538, 2.87337, 1.03073, 0.50165, 2.34429, 2.14417, 0.33666, 1.73978, -2.70077, 0.605818, -1.20169, -2.63533, 2.03946, 3.05334, 1.2107
125	20	(3, 1)	2.80847, 2.80847, 1.23768, -2.80847, 1.23768, -2.80847, -1.23768, -1.23768, 1.23768, -2.80847, 2.80847, 2.80847, -1.23768, -1.23768, 1.23768, -2.80847
126	28	(3, 1)	-2.34429, 2.70077, -2.03946, 2.70538, -2.14417, -0.605818, -3.05334, -2.87337, -0.33666, 1.20169, -1.2107, -1.03073, -0.50165, -1.73978, 2.63533, 1.09698
127	28	(3, 1)	-1.2107, -1.03073, -0.33666, 1.20169, 2.63533, 1.09698, -0.50165, -1.73978, -2.03946, 2.70538, -2.34429, 2.70077, -3.05334, -2.87337, -2.14417, -0.605818
128	52	(3, 1)	2.87337, 1.03073, -1.09698, -2.70538, 2.14417, 0.33666, 0.50165, 2.34429, 0.605818, -1.20169, 1.73978, -2.70077, 3.05334, 1.2107, -2.63533, 2.03946
129	52	(3, 1)	1.73978, -2.70077, 0.605818, -1.20169, -2.63533, 2.03946, 3.05334, 1.2107, -1.09698, -2.70538, 2.87337, 1.03073, 0.50165, 2.34429, 2.14417, 0.33666
130	60	(3, 1)	1.23768, -2.80847, 2.80847, 2.80847, -1.23768, -1.23768, 1.23768, -2.80847, 2.80847, 2.80847, 1.23768, -2.80847, 1.23768, -2.80847, -1.23768, -1.23768
131	68	(3, 1)	-0.33666, 1.20169, -1.2107, -1.03073, -0.50165, -1.73978, 2.63533, 1.09698, -2.34429, 2.70077, -2.03946, 2.70538, -2.14417, -0.605818, -3.05334, -2.87337
132	68	(3, 1)	-2.03946, 2.70538, -2.34429, 2.70077, -3.05334, -2.87337, -2.14417, -0.605818, -1.2107, -1.03073, -0.33666, 1.20169, 2.63533, 1.09698, -0.50165, -1.73978
133	5	(3, 2)	0.179498, -2.13305, 0.179498, -2.13305, 0.937357, 2.88492, 0.937357, 2.88492, 1.81154, -1.70682, 1.81154, -1.70682, -0.572197, 0.169554, -0.572197, 0.169554

*Continued on next page*

#	$\omega$	$\tau$	$\xi$
134	13	(3, 2)	-1.30524, -1.96077, -1.38646, 1.75512, 3.07932, -0.974834, 1.15546, 0.883402, 2.22175, 2.78125, 0.297893, 0.923013, 1.62208, 0.625598, 1.54086, -3.0903
135	13	(3, 2)	-1.38646, 1.75512, -1.30524, -1.96077, 1.15546, 0.883402, 3.07932, -0.974834, 0.297893, 0.923013, 2.22175, 2.78125, 1.54086, -3.0903, 1.62208, 0.625598
136	37	(3, 2)	-1.88271, 1.0715, -2.94948, -0.83118, -2.44672, 0.173026, 1.23134, -1.16544, -2.25236, -2.76484, 1.42571, -1.42637, -1.21313, -0.521724, -2.2799, 1.38096
137	37	(3, 2)	-2.94948, -0.83118, -1.88271, 1.0715, 1.23134, -1.16544, -2.44672, 0.173026, 1.42571, -1.42637, -2.25236, -2.76484, -2.2799, 1.38096, -1.21313, -0.521724
138	45	(3, 2)	1.81154, -1.70682, 1.81154, -1.70682, -0.572197, 0.169554, -0.572197, 0.169554, 0.179498, -2.13305, 0.179498, -2.13305, 0.937357, 2.88492, 0.937357, 2.88492
139	53	(3, 2)	0.297893, 0.923013, 2.22175, 2.78125, 1.54086, -3.0903, 1.62208, 0.625598, -1.38646, 1.75512, -1.30524, -1.96077, 1.15546, 0.883402, 3.07932, -0.974834
140	53	(3, 2)	2.22175, 2.78125, 0.297893, 0.923013, 1.62208, 0.625598, 1.54086, -3.0903, -1.30524, -1.96077, -1.38646, 1.75512, 3.07932, -0.974834, 1.15546, 0.883402
141	77	(3, 2)	-2.25236, -2.76484, 1.42571, -1.42637, -1.21313, -0.521724, -2.2799, 1.38096, -1.88271, 1.0715, -2.94948, -0.83118, -2.44672, 0.173026, 1.23134, -1.16544
142	77	(3, 2)	1.42571, -1.42637, -2.25236, -2.76484, -2.2799, 1.38096, -1.21313, -0.521724, -2.94948, -0.83118, -1.88271, 1.0715, 1.23134, -1.16544, -2.44672, 0.173026
143	12	(3, 3)	-2.70538, 2.87337, 1.03073, -1.09698, 2.34429, 2.14417, 0.33666, 0.50165, -2.70077, 0.605818, -1.20169, 1.73978, 2.03946, 3.05334, 1.2107, -2.63533
144	12	(3, 3)	-1.20169, 1.73978, -2.70077, 0.605818, 1.2107, -2.63533, 2.03946, 3.05334, 1.03073, -1.09698, -2.70538, 2.87337, 0.33666, 0.50165, 2.34429, 2.14417
145	20	(3, 3)	2.80847, 1.23768, -2.80847, 2.80847, -2.80847, -1.23768, -1.23768, 1.23768, -2.80847, 2.80847, 2.80847, 1.23768, -1.23768, 1.23768, -2.80847, -1.23768
146	28	(3, 3)	-1.03073, -0.33666, 1.20169, -1.2107, 1.09698, -0.50165, -1.73978, 2.63533, 2.70538, -2.34429, 2.70077, -2.03946, -2.87337, -2.14417, -0.605818, -3.05334
147	28	(3, 3)	2.70077, -2.03946, 2.70538, -2.34429, -0.605818, -3.05334, -2.87337, -2.14417, 1.20169, -1.2107, -1.03073, -0.33666, -1.73978, 2.63533, 1.09698, -0.50165
148	52	(3, 3)	1.03073, -1.09698, -2.70538, 2.87337, 0.33666, 0.50165, 2.34429, 2.14417, -1.20169, 1.73978, -2.70077, 0.605818, 1.2107, -2.63533, 2.03946, 3.05334
149	52	(3, 3)	-2.70077, 0.605818, -1.20169, 1.73978, 2.03946, 3.05334, 1.2107, -2.63533, -2.70538, 2.87337, 1.03073, -1.09698, 2.34429, 2.14417, 0.33666, 0.50165
150	60	(3, 3)	-2.80847, 2.80847, 2.80847, 1.23768, -1.23768, 1.23768, -2.80847, -1.23768, 2.80847, 1.23768, -2.80847, 2.80847, -2.80847, -1.23768, -1.23768, 1.23768
151	68	(3, 3)	2.70538, -2.34429, 2.70077, -2.03946, -2.87337, -2.14417, -0.605818, -3.05334, -1.03073, -0.33666, 1.20169, -1.2107, 1.09698, -0.50165, -1.73978, 2.63533
152	68	(3, 3)	1.20169, -1.2107, -1.03073, -0.33666, -1.73978, 2.63533, 1.09698, -0.50165, 2.70077, -2.03946, 2.70538, -2.34429, -0.605818, -3.05334, -2.87337, -2.14417

Table 6:  $(\omega_j, \tau_j, \xi_j)$  for  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} + 16$

The table below gives the solutions  $(\omega_j, \tau_j, \xi_j)$  in Table 3 for the near-group category  $J_8^1$  of type  $\mathbb{Z}/8\mathbb{Z} + 8$ , where  $j = 1, \dots, 44$ . The  $\omega_i$  equals to  $\zeta_{48}^k$  for some  $k \in \mathbb{Z}$ , which are recorded in the corresponding column.

#	$\omega$	$\tau$	$\xi$
1	24	0	1.8326, 0.840155, -2.87979, -2.67275, 1.8326, 0.840155, -2.87979, -2.67275
2	40	0	-2.35619, -2.03783, -2.19092, 2.32649, -2.35619, 0.0297022, 0.620122, -1.88916
3	40	0	-2.35619, 0.0297022, 0.620122, -1.88916, -2.35619, -2.03783, -2.19092, 2.32649
4	16	0	-1.5708, 1.76553, -1.06745, 0.914113, 1.5708, 3.01288, 1.06745, 2.16146
5	16	0	1.5708, 3.01288, 1.06745, 2.16146, -1.5708, 1.76553, -1.06745, 0.914113
6	39	1	2.86523, -0.967181, -2.38552, -2.68973, 0.113755, -1.3573, 3.01698, -0.428819
7	39	1	0.113755, -1.3573, 3.01698, -0.428819, 2.86523, -0.967181, -2.38552, -2.68973
8	7	1	-0.960866, 0.764517, 0.941698, 2.72147, -2.6727, -0.665243, 1.79457, 0.432749
9	7	1	0.7022, -0.89855, -0.94287, -2.38689, 0.955559, 1.98968, 0.619742, 2.31732
10	7	1	-2.6727, -0.665243, 1.79457, 0.432749, -0.960866, 0.764517, 0.941698, 2.72147
11	7	1	0.955559, 1.98968, 0.619742, 2.31732, 0.7022, -0.89855, -0.94287, -2.38689
12	12	2	-2.67275, 1.8326, 0.840155, -2.87979, -2.67275, 1.8326, 0.840155, -2.87979
13	28	2	2.32649, -2.35619, 0.0297022, 0.620122, -1.88916, -2.35619, -2.03783, -2.19092
14	28	2	-1.88916, -2.35619, -2.03783, -2.19092, 2.32649, -2.35619, 0.0297022, 0.620122
15	4	2	2.16146, -1.5708, 1.76553, -1.06745, 0.914113, 1.5708, 3.01288, 1.06745
16	4	2	0.914113, 1.5708, 3.01288, 1.06745, 2.16146, -1.5708, 1.76553, -1.06745
17	15	3	-0.428819, 2.86523, -0.967181, -2.38552, -2.68973, 0.113755, -1.3573, 3.01698
18	15	3	-2.68973, 0.113755, -1.3573, 3.01698, -0.428819, 2.86523, -0.967181, -2.38552
19	31	3	2.72147, -2.6727, -0.665243, 1.79457, 0.432749, -0.960866, 0.764517, 0.941698
20	31	3	-2.38689, 0.955559, 1.98968, 0.619742, 2.31732, 0.7022, -0.89855, -0.94287
21	31	3	2.31732, 0.7022, -0.89855, -0.94287, -2.38689, 0.955559, 1.98968, 0.619742
22	31	3	0.432749, -0.960866, 0.764517, 0.941698, 2.72147, -2.6727, -0.665243, 1.79457
23	24	4	-2.87979, -2.67275, 1.8326, 0.840155, -2.87979, -2.67275, 1.8326, 0.840155
24	16	4	1.06745, 2.16146, -1.5708, 1.76553, -1.06745, 0.914113, 1.5708, 3.01288
25	16	4	-1.06745, 0.914113, 1.5708, 3.01288, 1.06745, 2.16146, -1.5708, 1.76553
26	40	4	-2.19092, 2.32649, -2.35619, 0.0297022, 0.620122, -1.88916, -2.35619, -2.03783
27	40	4	0.620122, -1.88916, -2.35619, -2.03783, -2.19092, 2.32649, -2.35619, 0.0297022
28	15	5	-2.38552, -2.68973, 0.113755, -1.3573, 3.01698, -0.428819, 2.86523, -0.967181
29	15	5	3.01698, -0.428819, 2.86523, -0.967181, -2.38552, -2.68973, 0.113755, -1.3573
30	31	5	1.79457, 0.432749, -0.960866, 0.764517, 0.941698, 2.72147, -2.6727, -0.665243
31	31	5	0.619742, 2.31732, 0.7022, -0.89855, -0.94287, -2.38689, 0.955559, 1.98968
32	31	5	-0.94287, -2.38689, 0.955559, 1.98968, 0.619742, 2.31732, 0.7022, -0.89855
33	31	5	0.941698, 2.72147, -2.6727, -0.665243, 1.79457, 0.432749, -0.960866, 0.764517
34	12	6	0.840155, -2.87979, -2.67275, 1.8326, 0.840155, -2.87979, -2.67275, 1.8326
35	28	6	-2.03783, -2.19092, 2.32649, -2.35619, 0.0297022, 0.620122, -1.88916, -2.35619
36	28	6	0.0297022, 0.620122, -1.88916, -2.35619, -2.03783, -2.19092, 2.32649, -2.35619
37	4	6	3.01288, 1.06745, 2.16146, -1.5708, 1.76553, -1.06745, 0.914113, 1.5708
38	4	6	1.76553, -1.06745, 0.914113, 1.5708, 3.01288, 1.06745, 2.16146, -1.5708
39	39	7	-0.967181, -2.38552, -2.68973, 0.113755, -1.3573, 3.01698, -0.428819, 2.86523
40	39	7	-1.3573, 3.01698, -0.428819, 2.86523, -0.967181, -2.38552, -2.68973, 0.113755
41	7	7	-0.89855, -0.94287, -2.38689, 0.955559, 1.98968, 0.619742, 2.31732, 0.7022
42	7	7	0.764517, 0.941698, 2.72147, -2.6727, -0.665243, 1.79457, 0.432749, -0.960866
43	7	7	1.98968, 0.619742, 2.31732, 0.7022, -0.89855, -0.94287, -2.38689, 0.955559
44	7	7	-0.665243, 1.79457, 0.432749, -0.960866, 0.764517, 0.941698, 2.72147, -2.6727

Table 7:  $(\omega_j, \tau_j, \xi_j)$  for  $J_g^1$  in Table 3.



## REFERENCES

- [1] B. Bakalov and A. Kirillov, Jr. *Lectures on tensor categories and modular functors*, volume 21 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2001.
- [2] A. Bruguières. Catégories prémodulaires, modularization et invariants des variétés de dimension. *Math. Ann.*, 316(2):215–236, 2000.
- [3] P. Bruillard, S.-H. Ng, E. C. Rowell, and Z. Wang. On classification of modular categories by rank,. *Int. Math. Res. Not.*, 2016(24):7546–7588, 2016.
- [4] P. Bruillard, S.-H. Ng, E. C. Rowell, and Z. Wang. Rank-finiteness for modular categories. *J. Amer. Math. Soc.*, 29(3):857–881, 2016.
- [5] A. Coste and T. Gannon. Remarks on Galois symmetry in rational conformal field theories. *Phys. Lett. B*, 323(3-4):316–321, 1994.
- [6] J. de Boer and J. Goeree. Markov traces and  $II_1$  factors in conformal field theory. *Comm. Math. Phys.*, 139(2):267–304, 1991.
- [7] C. Dong, X. Lin, and S.-H. Ng. Congruence property in conformal field theory. *Algebra Number Theory*, 9(9):2121–2166, 2015.
- [8] V. Drinfeld, S. Gelaki, D. Nikshych, and V. Ostrik. On braided fusion category I. *Sel. Math. New Ser.*, 16:1–119, 2010.
- [9] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik. *Tensor categories*. Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2015.
- [10] D. E. Evans and T. Gannon. Near-group fusion categories and their doubles. *Adv. Math.*, 255:586–640, 2014.
- [11] M. Izumi. The structure of sectors associated with Longo–Rehren inclusions II: examples. *Rev. Math. Phys.*, 13(05):603–674, 2001.
- [12] A. Kirillov, Jr and V. Ostrik. On a q-analogue of the McKay correspondence and the ade classification of  $\mathfrak{sl}_2$  conformal field theories. *Adv. Math.*, 171(2):183–227, 2002.
- [13] G. Moore and N. Seiberg. Classical and quantum conformal field theory. *Comm. Math. Phys.*, 123(2):177–254, 1989.
- [14] M. Müger. On the structure of modular categories. *Proc. London Math. Soc.*, 87(2):291–308, 2003.
- [15] S.-H. Ng, E. C. Rowell, Z. Wang, and X. Wen. Reconstruction of modular data from  $SL_2(\mathbb{Z})$  representations. *Comm. Math. Phys.*, 402:2465–2545, 2023.
- [16] S.-H. Ng, E. C. Rowell, and X.-G. Wen. Classification of modular data up to rank 11. *arXiv preprint arXiv:2308.09670*, 2023.
- [17] S.-H. Ng and P. Schauenburg. Congruence subgroups and generalized Frobenius-Schur indicators. *Comm. Math. Phys.*, 300(1):1–46, 2010.
- [18] S.-H. Ng, Y. Wang, and Q. Zhang. Modular categories with transitive galois actions. *Comm. Math. Phys.*, 390(3):1271–1310, 2022.
- [19] A. Nobs. Die irreduziblen Darstellungen der Gruppen  $SL_2(\mathbb{Z}_p)$ , insbesondere  $SL_2(\mathbb{Z}_2)$ . I. *Comment. Math. Helv.*, 51(4):465–489, 1976.
- [20] A. Nobs and J. Wolfart. Die irreduziblen darstellungen der gruppen  $SL_2(\mathbb{Z}_p)$ , insbesondere  $SL_2(\mathbb{Z}_p)$ . II. *Comment. Math. Helv.*, 51(4):491–526, 1976.

- [21] J. Plavnik, A. Schopieray, Z. Yu, and Q. Zhang. Modular tensor categories, subcategories, and Galois orbits. *Transform. Groups*, 29:1623–1648, 2024.
- [22] E. Rowell, R. Stong, and Z. Wang. On classification of modular tensor categories. *Comm. Math. Phys.*, 292(2):343–389, 2009.
- [23] E. C. Rowell, H. Solomon, and Q. Zhang. On near-group centers and super-modular categories. *arXiv:2305.09108v1*, 2023.
- [24] J. Siehler. Near-group categories. *Algebr. Geom. Topol.*, 3:719–775, 2003.
- [25] D. Tambara and S. Yamagami. Tensor categories with fusion rules of self-duality for finite abelian groups. *J. Algebra*, 209(2):692–707, 1998.
- [26] The Sage Developers. *SageMath: Fusion Rings*. The Sage Development Team, 2024. <https://www.sagemath.org>.
- [27] V. G. Turaev. Modular categories and 3-manifold invariants. *Int. J. Mod. Phys. B*, 6(11n12):1807–1824, 1992.

*Email address:* zhiqyumath@yzu.edu.cn

*Email address:* qingzhang@ucsb.edu

(1) SCHOOL OF MATHEMATICAL SCIENCE, YANGZHOU UNIVERSITY, YANGZHOU, 225002, CHINA

(2) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SANTA BARBARA, CA 93106, USA