

# Higher rank Nichols algebras of diagonal type with finite arithmetic root systems in positive characteristic

Chen Yuan, Chen Qian, Jing Wang <sup>\*†</sup>

January 7, 2025

## Abstract

The classification of finite dimensional Nichols algebras of diagonal type plays an important role in the classification of Hopf algebras by the lifting method of N. Andruskiewitsch and H.-J. Schneider over fields of characteristic zero. In this paper, we obtain the classification theorem of all finite-dimensional rank 5, rank 6 and rank 7 Nichols algebras of diagonal type over fields of positive characteristic. Weyl groupoids and finite arithmetic root systems are crucial tools to the classification theorem.

**Key Words:** Nichols algebra, Cartan graph, Weyl groupoid, arithmetic root system

## 1 Introduction

The theory of Nichols algebras is motivated by the Hopf algebra theory and plays an important role in Hopf algebra, especially in quantum group structures [1, 3, 8, 33]. Besides, it has interesting applications to other research fields such as Kac-Moody Lie superalgebras [2] and conformal field theory [34–36].

In 1978, W. Nichols [31] first introduced bialgebras of type one, which was later named Nichols algebra in honor of him. In the early 1990s, researchers began to study the relationship between Nichols algebras and representation theory, laying the foundation for classification work. In the 2000s, the classification problem of Nichols algebras became a research hot spot. Nichols algebras appeared naturally in the classification of pointed Hopf algebras in the following way. Given a Hopf algebra  $H$ , consider its coradical filtration

$$H_0 \subset H_1 \subset \dots$$

such that  $H_0$  is a Hopf subalgebra of  $H$  and the associated graded coalgebra

$$\mathrm{gr}H = \bigoplus_i H_i/H_{i-1}.$$

---

<sup>\*</sup>Corresponding author: wang\_jing619@163.com

<sup>†</sup>The author is partially supported by National Natural Science Foundation of China(No.12471038).

Then  $\text{gr}H$  is a graded Hopf algebra, since the coradical  $H_0$  of  $H$  is a Hopf subalgebra. In addition, consider a projection  $\pi : \text{gr}H \rightarrow H_0$ ; let  $R$  be the algebra of coinvariants of  $\pi$ . Then, by a result of Radford and Majid,  $R$  is a braided Hopf algebra and  $\text{gr}H$  is the bosonization (or biproduct) of  $R$  and  $H_0$ :  $\text{gr}H \simeq R\#H_0$ . The principle of the "Lifting method" introduced in [6, 8] is first to study  $R$ , then to transfer the information to  $\text{gr}H$  via bosonization, and finally to lift to  $H$ . The braided Hopf algebra  $R$  is generated by the vector space  $V$  of  $H_0$ -coinvariants of  $H_1/H_0$ , namely, Nichols algebra [6]  $\mathcal{B}(V)$  is generated by  $V$ . Nichols algebras can be described in many different but alternative ways; for example, see [29, 30, 32, 40, 41].

The main theoretical tools for the classification of Nichols algebra  $\mathcal{B}(V)$  are the root system and the Weyl groupoid of  $\mathcal{B}(V)$ . I. Heckenberger [18] defined the root system and Weyl groupoid of  $\mathcal{B}(V)$  of diagonal type which based on the ideas of V. Kharchenko on Poincaré–Birkhoff–Witt basis of  $\mathcal{B}(V)$  of diagonal type [28, Theorem 2]. Later, M. Cuntz, I. Heckenberger and H. Yamane developed the combinatorial theory of these two structures [13, 27]. Then the theory of root systems and Weyl groupoids was carried out in more general Nichols algebras [4, 23, 24]. Further, all finite Weyl groupoids were classified in [14, 15].

The crucial step to classify pointed Hopf algebras is to determine that all Nichols algebras  $\mathcal{B}(V)$  are finite-dimensional. N. Andruskiewitsch stated the following question.

**Question 1.1. (N. Andruskiewitsch [1])** *Given a braiding matrix  $(q_{ij})_{1 \leq i, j \leq \theta}$  whose entries are roots of 1, when  $\mathcal{B}(V)$  is finite-dimensional, where  $V$  is a vector space with basis  $x_1, \dots, x_\theta$  and braiding  $c(x_i \otimes x_j) = q_{ij}(x_j \otimes x_i)$ ? If so, compute  $\dim_{\mathbb{k}} \mathcal{B}(V)$ , and give a "nice" presentation by generators and relations.*

Several authors obtained the classification results for infinite and finite dimensional Nichols algebras of Cartan type, see [7, 18, 32]. I. Heckenberger classified all finite dimensional Nichols algebras of diagonal type, see [16, 17, 19, 21]. The generators and relations of such Nichols algebras were also given [10, 11]. With the classification results, N. Andruskiewitsch and H.-J. Schneider [9] obtained a classification theorem about finite-dimensional pointed Hopf algebras under some technical assumptions. Based on such successful applications, the analysis to the structure of Nichols algebras over arbitrary fields is crucial and has potential applications. Over fields of positive characteristic, finite-dimensional rank 2, rank 3, and rank 4 Nichols algebras of diagonal type were classified in [26, 38, 39]. Those classification results in positive characteristic yield some important applications in pointed Hopf algebras and Nichols algebras [5, 22]. In this paper, we classify rank 5, rank 6 and rank 7 Nichols algebras of diagonal type with finite root systems over fields of positive characteristic. Ultimately, we can answer the first half of the Question 1.1.

The structure of this paper is as follows. In Section 2, we mainly recall some important notations and general results. In Section 3, we introduce the concept of good neighborhoods in order to explicitly characterize finite connected indecomposable Cartan graphs. And we formulate the main properties in Theorem 3.4. Finally, based on Theorem 3.4, we obtain the classification of rank 5, rank 6 and rank 7 Nichols algebras of diagonal type with finite root systems over the positive characteristic fields in Section 4. Then we present the corresponding generalized Dynkin diagrams in Table 1 and the exchange graphs of  $\mathcal{C}(M)$  in Table 2.

In this paper,  $\mathbb{k}$  is a field of characteristic  $p > 0$ . Assume  $\mathbb{k}^* = \mathbb{k} \setminus \{0\}$ ,  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ ,  $\theta \in \mathbb{N}_0$ , and  $I = \{1, \dots, \theta\}$ .

## 2 Preliminaries

In this section, we briefly recall notations, conventions, and related results, following [12, 26, 27, 38, 39]. For further details, we refer to the book [25].

### 2.1 Yetter–Drinfel’d modules and braided vector space

In this section, we recall Yetter–Drinfel’d modules, braided vector spaces, and their relations. We start by recalling the main objects.

**Definition 2.1.** *Let  $V$  be a  $\theta$ -dimensional vector space over  $\mathbb{k}$ . The pair  $(V, c)$  is called a braided vector space, if  $c \in \text{Aut}(V \otimes V)$  is a solution of the braid equation, that is*

$$(c \otimes \text{id})(\text{id} \otimes c)(c \otimes \text{id}) = (\text{id} \otimes c)(c \otimes \text{id})(\text{id} \otimes c).$$

*A braided vector space  $(V, c)$  is of diagonal type if  $V$  admits a basis  $\{x_i | i \in I\}$  such that for all  $i, j \in I$  one has*

$$c(x_i \otimes x_j) = q_{ij} x_j \otimes x_i \quad \text{for some } q_{ij} \in \mathbb{k}^*.$$

**Definition 2.2.** *The matrix  $(q_{ij})_{i, j \in I}$  is termed of the braiding matrix of  $V$ . The braiding matrix  $(q_{ij})_{i, j \in I}$  is indecomposable if for any  $i \neq j$  there exists a sequence  $i_1 = i, i_2, \dots, i_t = j$  of elements of  $I$  such that  $q_{i_s i_{s+1}} q_{i_{s+1} i_s} \neq 1$ , where  $1 \leq s \leq t - 1$ .*

In this paper, we mainly concern the braided vector spaces of diagonal type with an indecomposable braiding matrix.

**Definition 2.3.** *Let  $H$  be a Hopf algebra. A Yetter–Drinfel’d module  $V$  over  $H$  is a left  $H$ -module with left action  $\lambda_L : H \otimes V \rightarrow V$  and a left  $H$ -comodule with left coaction  $\delta_L : V \rightarrow H \otimes V$  satisfying the compatibility condition*

$$\delta_L(h.v) = h_{(1)} v_{(-1)} \kappa(h_{(3)}) \otimes h_{(2)}.v_{(0)}, \quad h \in H, v \in V.$$

*We say that  $V$  is of diagonal type if  $H = \mathbb{k}G$  and  $V$  is a direct sum of one-dimensional Yetter–Drinfel’d modules over the group algebra  $\mathbb{k}G$ , where  $G$  is abelian.*

We denote by  ${}^H_H\mathcal{YD}$  the category of Yetter–Drinfel’d modules over  $H$ , where morphisms preserve both the action and the coaction of  $H$ . The category  ${}^H_H\mathcal{YD}$  is braided with braiding

$$c_{V, W}(v \otimes w) = v_{(-1)}.w \otimes v_{(0)} \tag{1}$$

for all  $V, W \in {}^H_H\mathcal{YD}$ ,  $v \in V$ , and  $w \in W$ . Actually, the category  ${}^H_H\mathcal{YD}$  is a braided monoidal category, where the monoidal structure is given by the tensor product over  $\mathbb{k}$ . Then any Yetter–Drinfel’d module  $V \in {}^H_H\mathcal{YD}$  over  $H$  admits a braiding  $c_{V, V}$  and hence  $(V, c_{V, V})$  is a braided vector space. Conversely, any braided vector space can be realized as a Yetter–Drinfel’d module over some Hopf algebras if and only if the braiding is rigid [37, Section 2]. Notice that Yetter–Drinfel’d module structures on  $V$  with different Hopf algebras can give the same braiding and not all braidings of  $V$  are induced by the above Equation (1).

Let  $H = \mathbb{k}G$ . We write  ${}^G_G\mathcal{YD}$  for the category of Yetter–Drinfel’d modules over  $\mathbb{k}G$  and say that  $V \in {}^G_G\mathcal{YD}$  is a Yetter–Drinfel’d module over  $G$ . Notice that if  $V$  is of diagonal

type, then  $(V, c_{V,V})$  is a braided vector space of diagonal type. Any braided vector space of diagonal type is also a Yetter–Drinfel’d module of diagonal type. Indeed, assume that  $(V, c)$  is a braided vector space of diagonal type with an indecomposable braiding matrix  $(q_{ij})_{i,j \in I}$  of a basis  $\{x_i | i \in I\}$ . Let  $G_0$  be a free abelian group generated by elements  $\{g_i | i \in I\}$ . Define the left coaction and left action by

$$\delta_L(x_i) = g_i \otimes x_i \in G_0 \otimes V, \quad g_i \cdot x_j = q_{ij} x_j \in V.$$

Then  $V = \bigoplus_{i \in I} \mathbb{k}x_i$  and each  $\mathbb{k}x_i$  is one-dimensional Yetter–Drinfel’d modules over  $G_0$ . Hence  $V$  is a Yetter–Drinfel’d module of diagonal type over  $G_0$ .

## 2.2 Nichols Algebras of Diagonal Type

Let  $(V, c)$  be a  $\theta$ -dimensional braided vector space of diagonal type. In this section, we recall a definition of the Nichols algebra  $\mathcal{B}(V)$  generated by  $(V, c)$ . In order to do that, we start by introducing one more notion in the category  ${}^H_H\mathcal{YD}$ .

**Definition 2.4.** *Let  $H$  be a Hopf algebra. A braided Hopf algebra in  ${}^H_H\mathcal{YD}$  is a 6-tuple  $\mathcal{B} = (\mathcal{B}, \mu, 1, \delta, \epsilon, \kappa_B)$ , where  $(\mathcal{B}, \mu, 1)$  is an algebra in  ${}^H_H\mathcal{YD}$  and a coalgebra  $(\mathcal{B}, \delta, \epsilon)$  in  ${}^H_H\mathcal{YD}$ , and  $\kappa_B : \mathcal{B} \rightarrow \mathcal{B}$  is a morphism in  ${}^H_H\mathcal{YD}$  such that  $\delta, \epsilon$  and  $\kappa_B$  satisfy  $\kappa_B(b^{(1)})b^{(2)} = b^{(1)}\kappa_B(b^{(2)}) = \epsilon(b)1$ , where we define  $\delta(b) = b^{(1)} \otimes b^{(2)}$  as the coproduct of  $\mathcal{B}$  to avoid the confusion.*

We define Nichols algebra in the following way. And the equivalent descriptions of the definition of Nichols algebra can be seen in [29, 30, 32, 40, 41].

**Definition 2.5.** *The Nichols algebra generated by  $V \in {}^H_H\mathcal{YD}$  is defined as the quotient*

$$\mathcal{B}(V) = T(V)/\mathcal{I}(V) = (\bigoplus_{n=0}^{\infty} V^{\otimes n})/\mathcal{I}(V)$$

where  $\mathcal{I}(V)$  is the unique maximal coideal among all coideals of  $T(V)$  which are contained in  $\bigoplus_{n \geq 2} V^{\otimes n}$ . Nichols algebra  $\mathcal{B}(V)$  is said to be of diagonal type if  $V$  is a Yetter–Drinfel’d module of diagonal type. The dimension of  $V$  is the rank of Nichols algebra  $\mathcal{B}(V)$ .

Note that if  $B \in {}^H_H\mathcal{YD}$  and  $B$  is an algebra in  ${}^H_H\mathcal{YD}$  then  $B \otimes B$  is an algebra in  ${}^H_H\mathcal{YD}$  with the product given by

$$(a \otimes b)(c \otimes d) = a(b_{(-1)} \cdot c) \otimes b_{(0)}d, \tag{2}$$

for all  $a, b, c, d \in B$ , where  $\cdot$  denotes the left action of  $H$  on  $B$ .

The tensor algebra  $T(V)$  admits a natural structure of a Yetter–Drinfel’d module and an algebra structure in  ${}^H_H\mathcal{YD}$ . It is an  $\mathbb{N}_0$ -graded braided Hopf algebra in  ${}^H_H\mathcal{YD}$  with coproduct  $\delta(v)$  and counit  $\epsilon$  given by  $\delta(v) = 1 \otimes v + v \otimes 1$  and  $\epsilon(v) = 0$  for all  $v \in V$  such that  $\delta$  and  $\epsilon$  are the algebra morphisms. The antipode of  $T(V)$  exists; see [8, Section 2.1]. Notice that the product defined by Equation (2) on  $T(V)$  is the unique algebra structure such that  $\delta(v) = 1 \otimes v + v \otimes 1 \in T(V) \otimes T(V)$  for all  $v \in V$ . And the coproduct can be extended from

$V$  to  $T(V)$ . For example, for all  $v, w \in V$  one gets (we write the elements of  $T(V)$  without the tensor product sign for brevity)

$$\begin{aligned}\delta(vw) &= \delta(v)\delta(w) \\ &= (1 \otimes v + v \otimes 1)(1 \otimes w + w \otimes 1) \\ &= 1 \otimes vw + v_{(-1)}.w \otimes v_{(0)} + v \otimes w + vw \otimes 1.\end{aligned}$$

Let  $(I_i)_{i \in I}$  be the family of all ideals of  $T(V)$  contained in  $\bigoplus_{n \geq 2} V^{\otimes n}$ , i.e.

$$\delta(I_i) \subset I_i \otimes T(V) + T(V) \otimes I_i.$$

Then the coideal  $\mathcal{I}(V) := \sum_{i \in I} I_i$  is the largest element in  $(I_i)_{i \in I}$ . Hence  $\mathcal{B}(V)$  is a braided Hopf algebra in  ${}^H_H\mathcal{YD}$ . By [3, Proposition 3.2.12], Nichols algebra  $\mathcal{B}(V)$  is the unique  $\mathbb{N}_0$ -graded braided Hopf algebra generated by  $V$  in  ${}^H_H\mathcal{YD}$  with homogenous components  $\mathcal{B}(V)(0) = \mathbb{k}$ ,  $\mathcal{B}(V)(1) = V$ , and  $P(\mathcal{B}(V)) = V$ , where  $P(\mathcal{B}(V))$  is the space of primitive elements of  $\mathcal{B}(V)$ .

## 2.3 Cartan graphs, root systems, and Weyl groupoids

In this section, we mainly recall semi-Cartan graphs, root systems and Weyl groupoids. We start by introducing the notion of a generalized Cartan matrix.

**Definition 2.6.** *A generalized Cartan matrix is a matrix  $A = (a_{ij})_{i,j \in I}$  with integer entries such that*

- (1)  $a_{ii} = 2$  and  $a_{jk} \leq 0$  for any  $i, j, k \in I$  with  $j \neq k$ ,
- (2) if  $i, j \in I$  and  $a_{ij} = 0$ , then  $a_{ji} = 0$ .

We say that a generalized Cartan matrix  $A \in \mathbb{Z}^{I \times I}$  is *decomposable* if there exists a nonempty proper subset  $I_1 \subset I$  such that  $a_{ij} = 0$  for any  $i \in I_1$  and  $j \in I \setminus I_1$ . And  $A$  is called *indecomposable* if it is not decomposable.

**Definition 2.7.** *Let  $\mathcal{X}$  be a non-empty set and  $A^X = (a_{ij}^X)_{i,j \in I}$  be a generalized Cartan matrix for all  $X \in \mathcal{X}$ . For any  $i \in I$  let  $r_i : \mathcal{X} \rightarrow \mathcal{X}$ ,  $X \mapsto r(i, X)$ , where  $r : I \times \mathcal{X} \rightarrow \mathcal{X}$  is a map. The quadruple*

$$\mathcal{C} = \mathcal{C}(I, \mathcal{X}, r, (A^X)_{X \in \mathcal{X}})$$

*is called a semi-Cartan graph if  $r_i^2 = \text{id}_{\mathcal{X}}$  for all  $i \in I$ , and  $a_{ij}^X = a_{ij}^{r_i(X)}$  for all  $X \in \mathcal{X}$  and  $i, j \in I$ . We say that a semi-Cartan graph  $\mathcal{C}$  is indecomposable if  $A^X$  is indecomposable for all  $X \in \mathcal{X}$ .*

The elements of the set  $\{r_i(X), i \in I\}$  are called the *neighbors* of  $X$  for all  $X \in \mathcal{X}$ . The cardinality of  $I$  is called the *rank* of  $\mathcal{C}$  and the elements of  $\mathcal{X}$  are called the *points* of  $\mathcal{C}$ .

**Definition 2.8.** *The exchange graph of  $\mathcal{C}$  is a labeled non-oriented graph with vertices set  $\mathcal{X}$  and edges set  $I$ , where two vertices  $X, Y$  are connected by an edge  $i$  if and only if  $X \neq Y$  and  $r_i(X) = Y$  (and  $r_i(Y) = X$ ).*

Note that the exchange graph of  $\mathcal{C}$  may have multiple edges with different labels. For simplicity, we display only one edge with several labels instead of several edges.

**Definition 2.9.** Let  $\mathcal{C}$  be a semi-Cartan graph.  $\mathcal{C}$  is said to be connected if there is no proper non-empty subset  $\mathcal{Y} \subseteq \mathcal{X}$  such that  $r_i(Y) \in \mathcal{Y}$  for all  $i \in I$ ,  $Y \in \mathcal{Y}$ .

For the remaining part of this section, let  $\mathcal{C} = \mathcal{C}(I, \mathcal{X}, r, (A^X)_{X \in \mathcal{X}})$  be a connected semi-Cartan graph. Let  $(\alpha_i)_{i \in I}$  be the standard basis of  $\mathbb{Z}^I$ . For any  $X \in \mathcal{X}$  and any  $i \in I$  let

$$s_i^X \in \text{Aut}(\mathbb{Z}^I), \quad s_i^X \alpha_j = \alpha_j - a_{ij}^X \alpha_i, \quad (3)$$

for all  $j \in I$ . Let  $\mathcal{D}(\mathcal{X}, I)$  be the category such that  $\text{Ob}\mathcal{D}(\mathcal{X}, I) = \mathcal{X}$  and morphisms  $\text{Hom}(X, Y) = \{(Y, f, X) | f \in \text{End}(\mathbb{Z}^I)\}$  for  $X, Y \in \mathcal{X}$ , where composition of morphism is defined by  $(Z, g, Y) \circ (Y, f, X) = (Z, gf, X)$  for all  $X, Y, Z \in \mathcal{X}$ ,  $f, g \in \text{End}(\mathbb{Z}^I)$ . Let  $\mathcal{W}(\mathcal{C})$  be the smallest subcategory of  $\mathcal{D}(\mathcal{X}, I)$  which contains all morphisms  $(r_i(X), s_i^X, X)$  with  $i \in I$ ,  $X \in \mathcal{X}$ . For the sake of simplicity, we write  $s_i^X$  instead of  $(r_i(X), s_i^X, X)$  if no confusion is possible. Notice that  $\mathcal{W}(\mathcal{C})$  is a groupoid since all generators are invertible.

**Definition 2.10.** For all  $X \in \mathcal{X}$ , the set

$$\Delta^{X \text{ re}} = \{\omega \alpha_i \in \mathbb{Z}^I | \omega \in \text{Hom}(\mathcal{W}(\mathcal{C}), X)\} \quad (4)$$

is called the set of real roots of  $\mathcal{C}$  at  $X$ . The real roots  $\alpha_i$ ,  $i \in I$ , are called simple. The elements of  $\Delta_+^{X \text{ re}} = \Delta^{X \text{ re}} \cap \mathbb{N}_0^I$  and  $\Delta_-^{X \text{ re}} = \Delta^{X \text{ re}} \cap -\mathbb{N}_0^I$  are called positive roots and negative roots, respectively. We say that  $\mathcal{C}$  is finite if the set  $\Delta^{X \text{ re}}$  is finite for all  $X \in \mathcal{X}$ .

**Definition 2.11.** The semi-Cartan graph  $\mathcal{C}$  is called a Cartan graph if the following hold :

- (1) For all  $X \in \mathcal{X}$ , the set  $\Delta^{X \text{ re}}$  consists of positive and negative roots.
- (2) If  $m_{ij}^X := |\Delta^{X \text{ re}} \cap (\mathbb{N}_0 \alpha_i + \mathbb{N}_0 \alpha_j)|$  is finite, then  $(r_i r_j)^{m_{ij}^X}(X) = X$ , where  $i, j \in I$  and  $X \in \mathcal{X}$ .

In this case,  $\mathcal{W}(\mathcal{C})$  is called the Weyl groupoid of  $\mathcal{C}$ .

**Definition 2.12.**  $\mathcal{R} = \mathcal{R}(\mathcal{C}, (\Delta^X)_{X \in \mathcal{X}})$  is a root system of type  $\mathcal{C}$  if for all  $X \in \mathcal{X}$ , the sets  $\Delta^X$  are the subsets of  $\mathbb{Z}^I$  such that

- (1)  $\Delta^X = (\Delta^X \cap \mathbb{N}_0^I) \cup -(\Delta^X \cap \mathbb{N}_0^I)$ ,
- (2)  $\Delta^X \cap \mathbb{Z} \alpha_i = \{\alpha_i, -\alpha_i\}$  for all  $i \in I$ ,
- (3)  $s_i^X(\Delta^X) = \Delta^{r_i(X)}$  for all  $i \in I$ ,
- (4)  $(r_i r_j)^{m_{ij}^X}(X) = X$  for all  $i \neq j \in I$  where  $m_{ij}^X = |\Delta^X \cap (\mathbb{N}_0 \alpha_i + \mathbb{N}_0 \alpha_j)|$  is finite.

We say that  $\mathcal{R}$  is finite if  $\Delta^X$  is finite for all  $X \in \mathcal{X}$ .

We call  $\mathcal{W}(\mathcal{R}) := \mathcal{W}(\mathcal{C})$  is the groupoid of  $\mathcal{R}$ . As in [13, definition 4.3] we say that  $\mathcal{R}$  is *reducible* if there exist non-empty disjoint subsets of  $I', I'' \subset I$  such that  $I = I' \cup I''$  and  $a_{ij} = 0$  for all  $i \in I', j \in I''$  and

$$\Delta^X = \left( \Delta^X \cap \sum_{i \in I'} \mathbb{Z}\alpha_i \right) \cup \left( \Delta^X \cap \sum_{j \in I''} \mathbb{Z}\alpha_j \right) \quad \text{for all } X \in \mathcal{X}.$$

In this case, we write  $\mathcal{R} = \mathcal{R}|_{I_1} \oplus \mathcal{R}|_{I_2}$ . Conversely, we say that  $\mathcal{R}$  is irreducible if it is not reducible.

Let  $\mathcal{R} = \mathcal{R}(\mathcal{C}, (\Delta^X)_{X \in \mathcal{X}})$  be a root system of type  $\mathcal{C}$ . Next we recall some properties of  $\mathcal{R}$  from [13] and [14].

**Lemma 2.13.** *Let  $X \in \mathcal{X}$ ,  $k \in \mathbb{Z}$ , and  $i, j \in I$  such that  $i \neq j$ . Then  $\alpha_j + k\alpha_i \in \Delta^{X \text{ re}}$  if and only if  $0 \leq k \leq -a_{ij}^X$ .*

Notice that the finiteness of  $\mathcal{R}$  does not mean that  $\mathcal{W}(\mathcal{R})$  is also finite, since the set  $\mathcal{X}$  might be infinite.

**Lemma 2.14.** *Let  $\mathcal{C}$  be a connected semi-Cartan graph and  $\mathcal{R} = \mathcal{R}(\mathcal{C}, (\Delta^X)_{X \in \mathcal{X}})$  be a root system of type  $\mathcal{C}$ . Then the following are equivalent.*

- (1)  $\mathcal{R}$  is finite.
- (2)  $\Delta^X$  is finite for some  $X \in \mathcal{X}$ .
- (3)  $\mathcal{C}$  is finite.
- (4)  $\mathcal{W}(\mathcal{R})$  is finite.

When  $\mathcal{C}$  is a connected semi-Cartan graph, we can get the following.

**Proposition 2.15.** *Let  $\mathcal{R} = \mathcal{R}(\mathcal{C}, (\Delta^X)_{X \in \mathcal{X}})$  be a root system of type  $\mathcal{C}$ , then*

- (1) *there exists  $X \in \mathcal{X}$  such that  $\Delta^X$  is indecomposable if and only if the semi-Cartan graph  $\mathcal{C}$  is indecomposable.*
- (2) *the semi-Cartan graph  $\mathcal{C}$  is indecomposable if and only if the root system  $\mathcal{R}$  is irreducible if  $\mathcal{R}$  is finite.*

The following proposition implies that if  $\mathcal{R}$  is a finite root system of type  $\mathcal{C}$  then all roots are real and  $\mathcal{R}$  is uniquely determined by  $\mathcal{C}$ .

**Proposition 2.16.** *Let  $\mathcal{R} = \mathcal{R}(\mathcal{C}, (\Delta^X)_{X \in \mathcal{X}})$  be a root system of type  $\mathcal{C}$ . Let  $X \in \mathcal{X}$ ,  $m \in \mathbb{N}_0$ , and  $i_1, \dots, i_m \in I$  such that*

$$\omega = \text{id}_X s_{i_1} s_{i_2} \cdots s_{i_m} \in \text{Hom}(\mathcal{W}(\mathcal{C}), X)$$

and  $\ell(\omega) = m$ . Then the elements

$$\beta_n = \text{id}_X s_{i_1} s_{i_2} \cdots s_{i_{n-1}}(\alpha_{i_n}) \in \Delta^X \cap \mathbb{N}_0^I,$$

are pairwise different, where  $n \in \{1, 2, \dots, m\}$  (and  $\beta_1 = \alpha_{i_1}$ ). Here,

$$\ell(\omega) = \min\{m \in \mathbb{N}_0 \mid \omega = \text{id}_X s_{i_1} s_{i_2} \cdots s_{i_m}, i_1, i_2, \dots, i_m \in I\}$$

is the length of  $\omega \in \text{Hom}(\mathcal{W}(\mathcal{C}), X)$ . In particular, if  $\mathcal{R}$  is finite and  $\omega \in \text{Hom}(\mathcal{W}(\mathcal{C}))$  is the longest element, then

$$\{\beta_n \mid 1 \leq n \leq \ell(\omega) = |\Delta^X|/2\} = \Delta^X \cap \mathbb{N}_0^I.$$

**Remark 2.17.** If  $\mathcal{C}$  is a finite Cartan graph then  $\mathcal{R}$  is finite and hence

$$\mathcal{R}^{re} = \mathcal{R}^{re}(\mathcal{C}, (\Delta^{X^{re}})_{X \in \mathcal{X}})$$

is the unique root system of type  $\mathcal{C}$  by Proposition 2.16, that is,  $\mathcal{R}$  is uniquely determined by  $\mathcal{C}$ .

## 2.4 The semi-Cartan Graph of Nichols Algebras of Diagonal Type

In this section, we attach a semi-Cartan graph to a tuple of finite-dimensional Yetter–Drinfel’d modules under some finiteness conditions. Assume that  $G$  is an abelian group and  $\mathbb{k}G$  is the group algebra over  $G$ . Let  ${}^G_G\mathcal{YD}$  be the category of Yetter–Drinfel’d modules over  $\mathbb{k}G$ . Let  $\mathcal{F}_\theta^G$  be the set of  $\theta$ -tuples of finite-dimensional irreducible objects in  ${}^G_G\mathcal{YD}$  and  $\mathcal{X}_\theta^G$  be the set of  $\theta$ -tuples of isomorphism classes of finite-dimensional irreducible objects in  ${}^G_G\mathcal{YD}$ . For any  $(M_1, \dots, M_\theta) \in \mathcal{F}_\theta^G$ , write  $[M] := ([M_1], \dots, [M_\theta]) \in \mathcal{X}_\theta^G$  the corresponding isomorphism class of  $(M_1, \dots, M_\theta)$ .

We fix that  $M = (\mathbb{k}x_1, \mathbb{k}x_2, \dots, \mathbb{k}x_\theta) \in \mathcal{F}_\theta^G$  is a tuple of one-dimensional Yetter–Drinfel’d over  $G$  and  $[M] \in \mathcal{X}_\theta^G$ . Let  $V_M = \bigoplus_{i \in I} \mathbb{k}x_i \in {}^G_G\mathcal{YD}$  be a Yetter–Drinfel’d module of diagonal type over  $G$ , where  $\{x_i \mid i \in I\}$  is a basis of  $V$ . Then there exist a matrix  $(q_{ij})_{i,j \in I}$  and elements  $\{g_i \mid i \in I\} \subset G$  such that  $\delta(x_i) = g_i \otimes x_i$  and  $g_i \cdot x_j = q_{ij} x_j$  for all  $i, j \in I$ . We say that the matrix  $(q_{ij})_{i,j \in I}$  is the braiding matrix of  $M$ . Recall that the matrix does not depend on the basis  $\{x_i \mid i \in I\}$  up to permutation of  $I$ . We say that  $\mathcal{B}(V_M) = \mathcal{B}(\bigoplus_{i=1}^n \mathbb{k}x_i)$  is the Nichols algebra of the tuple  $M$ , denoted by  $\mathcal{B}(M)$ .

Recall that the adjoint representation  $\text{ad}$  of a Nichols algebra  $\mathcal{B}(V)$  from [6] is the linear map  $\text{ad}_c : V \rightarrow \text{End}(\mathcal{B}(V))$

$$\text{ad}_c x(y) = \mu(\text{id} - c)(x \otimes y) = xy - (x_{(-1)} \cdot y)x_{(0)}$$

for all  $x \in V$ ,  $y \in \mathcal{B}(V)$ , where  $\mu$  is the multiplication map of  $\mathcal{B}(V)$  and  $c$  is defined by Equation (1). In particular, the braided commutator  $\text{ad}_c$  of  $\mathcal{B}(M)$  takes the form

$$\text{ad}_c x_i(y) = x_i y - (g_i \cdot y)x_i \text{ for all } i \in I, y \in \mathcal{B}(M).$$

In order to construct a semi-Cartan graph to  $M$ , we recall some finiteness conditions from [8] and [23].

**Definition 2.18.** Let  $i \in I$ .  $M$  is  $i$ -finite if for any  $j \in I \setminus \{i\}$ ,  $(\text{ad}_c x_i)^m(x_j) = 0$  for some  $m \in \mathbb{N}$ .



**Lemma 2.19.** For any  $i, j \in I$  with  $i \neq j$ , the following are equivalent.

- (1)  $(m+1)_{q_{ii}}(q_{ii}^m q_{ij} q_{ji} - 1) = 0$  and  $(k+1)_{q_{ii}}(q_{ii}^k q_{ij} q_{ji} - 1) \neq 0$  for all  $0 \leq k < m$ .
- (2)  $(ad_c x_i)^{m+1}(x_j) = 0$  and  $(ad_c x_i)^m(x_j) \neq 0$  in  $\mathcal{B}(V)$ .

Here  $(n)_q := 1 + q + \cdots + q^{n-1}$ , and  $n = 0$  if and only if  $q^n = 1$  for  $q \neq 1$  or  $p|n$  for  $q = 1$ . Notice that  $(1)_q \neq 0$  for any  $q \in \mathbb{N}$ .

By Lemma 2.19, we get the following lemma.

**Lemma 2.20.** Let  $i \in I$ . Then  $M = (\mathbb{k}x_j)_{j \in I}$  is  $i$ -finite if and only if for any  $j \in I \setminus \{i\}$  there is a non-negative integer  $m$  satisfying  $(m+1)_{q_{ii}}(q_{ii}^m q_{ij} q_{ji} - 1) = 0$ .

Let  $i \in I$ . Assume that  $M$  is  $i$ -finite. Let  $(a_{ij}^M)_{j \in I} \in \mathbb{Z}^I$  and  $R_i(M) = (R_i(M)_j)_{j \in I}$ , where

$$a_{ij}^M = \begin{cases} 2 & \text{if } j = i, \\ -\max\{m \in \mathbb{N}_0 \mid (ad_c x_i)^m(x_j) \neq 0\} & \text{if } j \neq i. \end{cases}$$

$$R_i(M)_i = \mathbb{k}y_i, \quad R_i(M)_j = \mathbb{k}(ad_c x_i)^{-a_{ij}^M}(x_j), \quad (5)$$

where  $y_i \in (\mathbb{k}x_i)^* \setminus \{0\}$ . If  $M$  is not  $i$ -finite, then let  $R_i(M) = M$ . Then  $R_i(M)$  is a  $\theta$ -tuple of one-dimensional Yetter–Drinfel’d modules over  $G$ .

Let

$$\mathcal{F}_\theta^G(M) = \{R_{i_1} \cdots R_{i_n}(M) \in \mathcal{F}_\theta^G \mid n \in \mathbb{N}_0, i_1, \dots, i_n \in I\}$$

and

$$\mathcal{X}_\theta^G(M) = \{[R_{i_1} \cdots R_{i_n}(M)] \in \mathcal{X}_\theta^G \mid n \in \mathbb{N}_0, i_1, \dots, i_n \in I\}.$$

**Definition 2.21.** We say that  $M$  admits all reflections if  $N$  is  $i$ -finite for all  $N \in \mathcal{F}_\theta^G(M)$ .

Notice that the reflections depend only on the braiding matrix  $(q_{ij})_{i,j \in I}$ . We recall the notion of generalized Dynkin diagram for a braided vector spaces of diagonal type [20].

**Definition 2.22.** Let  $V$  be a  $\theta$ -dimensional braided vector space of diagonal type with the braiding matrix  $(q_{ij})_{i,j \in I}$ . The generalized Dynkin diagram of  $V$  is a non-directed graph  $\mathcal{D}$  with the following properties:

- (1) there is a bijective map  $\phi$  from  $I$  to the vertices of  $\mathcal{D}$ ,
- (2) for all  $i \in I$  the vertex  $\phi(i)$  is labeled by  $q_{ii}$ ,
- (3) for all  $i, j \in I$  with  $i \neq j$ , the number  $n_{ij}$  of edges between  $\phi(i)$  and  $\phi(j)$  is either 0 or 1. If  $q_{ij} q_{ji} = 1$  then  $n_{ij} = 0$ , otherwise  $n_{ij} = 1$  and the edge is labeled by  $q_{ij} q_{ji}$ .

We say that the generalized Dynkin diagram of  $M$  is the generalized Dynkin diagram of braided vector space  $\bigoplus_{i \in I} M_i$ . The generalized Dynkin diagram of  $M$  is connected if the braiding matrix of  $M$  is indecomposable.

We can get the labels of the generalized Dynkin diagram of  $R_i(M) = (R_i(M)_j)_{j \in I}$  by the following lemma.

**Lemma 2.23.** *Let  $i \in I$ . Assume that  $M$  is  $i$ -finite and let  $a_{ij} := a_{ij}^M$  for all  $j \in I$ . Let  $(q'_{jk})_{j,k \in I}$  be the braiding matrix of  $R_i(M)$  with respect to  $(y_j)_{j \in I}$ . Then*

$$q'_{jj} = \begin{cases} q_{ii} & \text{if } j = i, \\ q_{jj} & \text{if } j \neq i, q_{ij}q_{ji} = q_{ii}^{a_{ij}}, \\ q_{ii}q_{jj}(q_{ij}q_{ji})^{-a_{ij}} & \text{if } j \neq i, q_{ii} \in G'_{1-a_{ij}}, \\ q_{jj}(q_{ij}q_{ji})^{-a_{ij}} & \text{if } j \neq i, q_{ii} = 1, \end{cases}$$

$$q'_{ij}q'_{ji} = \begin{cases} q_{ij}q_{ji} & \text{if } j \neq i, q_{ij}q_{ji} = q_{ii}^{a_{ij}}, \\ q_{ii}^2(q_{ij}q_{ji})^{-1} & \text{if } j \neq i, q_{ii} \in G'_{1-a_{ij}}, \\ (q_{ij}q_{ji})^{-1} & \text{if } j \neq i, q_{ii} = 1, \end{cases}$$

and

$$q'_{jk}q'_{kj} = \begin{cases} q_{jk}q_{kj} & \text{if } q_{ir}q_{ri} = q_{ii}^{a_{ir}}, r \in \{j, k\}, \\ q_{jk}q_{kj}(q_{ik}q_{ki}q_{ii}^{-1})^{-a_{ij}} & \text{if } q_{ij}q_{ji} = q_{ii}^{a_{ij}}, q_{ii} \in G'_{1-a_{ik}}, \\ q_{jk}q_{kj}(q_{ij}q_{ji})^{-a_{ik}}(q_{ik}q_{ki})^{-a_{ij}} & \text{if } q_{ii} = 1, \\ q_{jk}q_{kj}q_{ii}^2(q_{ij}q_{ji}q_{ik}q_{ki})^{-a_{ij}} & \text{if } q_{ii} \in G'_{1-a_{ik}}, q_{ii} \in G'_{1-a_{ij}}, \end{cases}$$

for  $j \neq k$ ,  $|j - i| = 1$ ,  $|k - i| = 1$ . For  $|j - i| > 1$ ,  $|k - i| > 1$ ,  $q'_{jk}q'_{kj} = q_{jk}q_{kj}$ . Here,  $G'_n$  denotes the set of primitive  $n$ -th roots of unity in  $\mathbb{k}$ , that is  $G'_n = \{q \in \mathbb{k}^* \mid q^n = 1, q^k \neq 1 \text{ for all } 1 \leq k < n\}$  for  $n \in \mathbb{N}$ .

If  $M$  admits all reflections, then we are able to construct a semi-Cartan graph  $\mathcal{C}(M)$  of  $M$  by [38, Proposition 1.5].

**Theorem 2.24.** *Assume that  $M$  admits all reflections. For all  $X \in \mathcal{X}_\theta^G(M)$  let*

$$[X]_\theta = \{Y \in \mathcal{X}_\theta^G(M) \mid Y \text{ and } X \text{ have the same generalized Dynkin diagram}\}.$$

Let  $\mathcal{Y}_\theta(M) = \{[X]_\theta \mid X \in \mathcal{X}_\theta^G(M)\}$  and  $A^{[X]_\theta} = A^X$  for all  $X \in \mathcal{X}_\theta^G(M)$ . Let  $t : I \times \mathcal{Y}_\theta(M) \rightarrow \mathcal{Y}_\theta(M)$ ,  $(i, [X]_\theta) \mapsto [R_i(X)]_\theta$ . Then the tuple

$$\mathcal{C}(M) = \{I, \mathcal{Y}_\theta(M), t, (A^Y)_{Y \in \mathcal{Y}_\theta(M)}\}$$

is a connected semi-Cartan graph. We say that  $\mathcal{C}(M)$  is the semi-Cartan graph attached to  $M$ .

Furthermore, we can attach a groupoid  $\mathcal{W}(M) := \mathcal{W}(\mathcal{C}(M))$  to  $M$  if  $M$  admits all reflections.

Notice that Nichols algebra  $\mathcal{B}(M)$  is  $\mathbb{N}_0^\theta$ -graded with  $\deg M_i = \alpha_i$  for all  $i \in I$ . Following the terminology in [23], we say that the Nichols algebra  $\mathcal{B}(M)$  is *decomposable* if there exists a totally ordered index set  $(L, \leq)$  and a sequence  $(W_l)_{l \in L}$  of finite-dimensional irreducible  $\mathbb{N}_0^\theta$ -graded objects in  ${}^G\mathcal{YD}$  such that

$$\mathcal{B}(M) \simeq \bigotimes_{l \in L} \mathcal{B}(W_l). \quad (6)$$

For each decomposition (6), we define the set of *positive roots*  $\Delta_+^{[M]} \subset \mathbb{Z}^I$  and the set of *roots*  $\Delta^{[M]} \subset \mathbb{Z}^I$  of  $[M]$  by

$$\Delta_+^{[M]} = \{\deg(W_l) \mid l \in L\}, \quad \Delta^{[M]} = \Delta_+^{[M]} \cup -\Delta_+^{[M]}.$$

By [23, Theorem 4.5] we obtain that the set of roots  $\Delta^{[M]}$  of  $[M]$  does not depend on the choice of the decomposition.

**Remark 2.25.** *If  $\dim M_i = 1$  for all  $i \in I$ , then the Nichols algebra  $\mathcal{B}(M)$  is decomposable based on the theorem of V. Kharchenko [28, Theorem 2]. The set of roots of Nichols algebra  $\mathcal{B}(M)$  can be always defined and it is denoted by  $\Delta^{[M]}$ . If the set of roots  $\Delta^{[M]}$  is finite then we can check that  $M$  admits all reflections by [23, Corollary 6.12].*

If  $M$  admits all reflections and  $\Delta^{[M]}$  is finite, then we can define a finite root system  $\mathcal{R}(M)(\mathcal{C}(M), (\Delta^{[N]})_{N \in \mathcal{F}_\theta^G(M)})$  of type  $\mathcal{C}(M)$ .

**Theorem 2.26.** *Assume that  $M$  admits all reflections. Then the following are equivalent.*

- (1)  $\Delta^{[M]}$  is finite.
- (2)  $\mathcal{C}(M)$  is a finite Cartan graph.
- (3)  $\mathcal{W}(M)$  is finite.
- (4)  $\mathcal{R}(M) := \mathcal{R}(M)(\mathcal{C}(M), (\Delta^{[N]})_{N \in \mathcal{F}_\theta^G(M)})$  is finite.

In all cases,  $\mathcal{R}(M)$  is the unique root system of type  $\mathcal{C}(M)$ .

*Proof.* Since  $M$  is a  $\theta$ -tuple of one-dimensional Yetter–Drinfel’d modules, we obtain that the Nichols algebra  $\mathcal{B}(M)$  generated by  $V_M$  is decomposable and hence  $\Delta^{[M]}$  is defined. Then  $\mathcal{R}(M) = \mathcal{R}(M)(\mathcal{C}(M), (\Delta^{[N]})_{N \in \mathcal{F}_\theta^G(M)})$  is the root system of type  $\mathcal{C}(M)$  by [23, Theorem 6.11]. Hence the claim is true by Lemma 2.14.  $\square$

### 3 Good Neighborhoods

Let  $I = \{1, 2, \dots, \theta\}$  and  $M := (\mathbb{k}x_1, \dots, \mathbb{k}x_\theta)$  be a tuple of one-dimensional Yetter–Drinfel’d modules over  $\mathbb{k}G$ . Assume that  $M$  admits all reflections and  $\mathcal{C}(M) = \mathcal{C}(I, \mathcal{X}, r, (A^X)_{X \in \mathcal{X}})$  be the attached indecomposable semi-Cartan graph to  $M$ . In general, the points of a finite Cartan graph  $\mathcal{C}$  could have many different neighborhoods. We analyze the positive real roots  $\Delta^{[M]}$  by Gap Software. Depending on the output, we construct the following ”good neighborhoods” in order to cover all the finite connected indecomposable Cartan graphs in such way that at least one point of  $\mathcal{C}$  has one of the good neighborhoods.

$$\text{Assume that } A_5 = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix}, \quad A_6 = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}$$

$$\text{and } A_7 = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}.$$

**Definition 3.1.** We say that  $X$  has a **good  $A_5$  neighborhood** if there exists a permutation of  $I$  and integers  $a, b, c, d, e \in \mathbb{N}$  such that

$$A^X = A^{r_1(X)} = A^{r_2(X)} = A_5, \quad A^{r_3(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & -\mathbf{a} & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & -\mathbf{b} & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix},$$

$$A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -\mathbf{c} \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -\mathbf{d} & -1 & 2 \end{pmatrix}, \quad A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & -\mathbf{e} & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix},$$

$a_{15}^{r_2 r_3 r_4(X)} = 0$ ,  $a_{45}^{r_3 r_2 r_1(X)} \neq -3$ , and satisfies one of the following.

$$(A_{5a}) \quad (a, b, c, d, e) = (1, 1, 0, 0, 1), \quad [a_{41}^{r_2 r_3(X)}, a_{45}^{r_2 r_3(X)}] = [0, -1], \quad [a_{32}^{r_4 r_5(X)}, a_{34}^{r_2 r_1(X)}] \neq [-2, -2];$$

$$(A_{5b}) \quad (a, b, c, d, e) = (0, 0, 1, 1, 2), \quad a_{25}^{r_3 r_4(X)} \in \{0, -1\}, \quad [a_{45}^{r_3 r_2 r_1(X)}, a_{45}^{r_3 r_2(X)}] \neq [-2, -2];$$

$$(A_{5c}) \quad (a, b, c, d, e) = (0, 0, 1, 1, 1), \quad [a_{52}^{r_3 r_4(X)}, a_{32}^{r_5 r_4(X)}] = [-1, -1].$$

**Definition 3.2.** We say that  $X$  has a **good  $A_6$  neighborhood** if there exists a permutation of  $I$  and an integer  $a \in \mathbb{N}$  such that

$$A^X = A^{r_1(X)} = A^{r_2(X)} = A^{r_3(X)} = A^{r_6(X)} = A_6,$$

$$A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 & 0 \\ 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & -1 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}, \quad A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & -\mathbf{a} \\ 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & -\mathbf{a} & -1 & 2 \end{pmatrix},$$

$a_{36}^{r_5 r_4(X)} = 0$ , and  $a_{25}^{r_3 r_4(X)} = 0$ , where  $a$  satisfies one of the following.

$$(A_{6a}) \quad a = 0, \quad a_{32}^{r_5 r_4(X)} = -1, \quad a_{43}^{r_5 r_6(X)} = -1, \quad [a_{45}^{r_3 r_2 r_1(X)}, a_{45}^{r_3 r_2(X)}] \neq [-2, -2];$$

$$(A_{6b}) \quad a = 1, \quad a_{45}^{r_3 r_2(X)} = -1, \quad a_{45}^{r_3 r_2 r_1(X)} = -1.$$

**Definition 3.3.** We say that  $X$  has a **good  $A_7$  neighborhood** if there exists a permutation of  $I$  such that

$$A^X = A^{r_1(X)} = A^{r_2(X)} = A^{r_4(X)} = A^{r_5(X)} = A^{r_6(X)} = A^{r_7(X)} = A_7,$$

$$A^{r_3(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}, [a_{25}^{r_4 r_3(X)}, a_{21}^{r_4 r_3(X)}] = [0, -1],$$

$$a_{45}^{r_2 r_3(x)} = -1, a_{34}^{r_2 r_1(X)} = -1, a_{32}^{r_4 r_5(X)} = -1, a_{32}^{r_4 r_5 r_6(x)} = -1, \text{ and } a_{32}^{r_4 r_5 r_6 r_7(X)} = -1.$$

**Theorem 3.4.** *Let  $I = \{1, 2, \dots, \theta\}$  and  $M := (\mathbb{k}x_1, \dots, \mathbb{k}x_\theta)$  be a tuple of one-dimensional Yetter–Drinfel’d modules over  $\mathbb{k}G$ . Assume that  $\mathcal{C}(M) = \mathcal{C}(I, \mathcal{X}, r, (A^X)_{X \in \mathcal{X}})$  is the attached indecomposable semi-Cartan graph to  $M$ . Assume there exists a point  $X \in \mathcal{X}$  satisfying that the set  $\Delta_+^{X \text{ re}}$  appears in Appendix, then up to equivalence, there exists a point  $Y \in \mathcal{X}$  such that  $Y$  has a good  $A_\theta$  neighborhood, where  $\theta = 5, 6, 7$ .*

*Proof.* Assume that  $\Delta_+^{X \text{ re}}$  appears in Appendix. Then there are precisely 6 possible sets of real roots for the rank 5 case, 3 sets of real roots for the rank 6 case (Except  $E_6$  type), and 1 set of real roots for the rank 7 case (Except  $E_7$  type). We analyze each set of the real roots in the list. Since the reflection  $s_i^X$  maps  $\Delta_+^{X \text{ re}} \setminus \{\alpha_i\}$  bijectively to  $\Delta_+^{r_i(X) \text{ re}} \setminus \{\alpha_i\}$  for any  $i \in I$ , the Cartan matrices of all neighbors of  $X$  can be obtained from  $\Delta_+^{X \text{ re}}$  by Lemma 2.13. If there exists a point  $Y$  of the Cartan graph  $\mathcal{C}(M)$  which has a good  $A_\theta$  neighborhood, then the claim is true. Otherwise repeat the previous step to the neighbors of  $X$ . Since  $\mathcal{X}$  is finite, this algorithm terminates.  $\square$

## 4 The classification theorem

In this section, we give the classification of all the finite Cartan graphs  $\mathcal{C}(M)$  with finite set of roots in Appendix. In addition, we present the corresponding generalized Dynkin diagrams in Table 1 and the exchange graphs of  $\mathcal{C}(M)$  in Table 2.

To avoid confusion, we label the vertices of the Dynkin diagrams from left to right and top to bottom by  $1, 2, \dots, \theta$  ( $\theta = 5, 6, 7$ ) (for example: label  $D^7$  as in Fig. 1).




Figure 1: Dynkin diagram of type  $D^7$

For a Dynkin diagram  $\mathcal{D}$  and  $i, j, k, l, m, n, t \in I$ , write  $\tau_{ijklmnt} \mathcal{D}^\theta$  for the graph  $\mathcal{D}$  where the vertices of  $\mathcal{D}$  change to  $i, j, k, l, m, n, t$  respectively.

## 4.1 Classification theorem for rank 5

**Lemma 4.1.** *If  $\mathcal{D}_{\chi,E}$  is of the form*

$$\begin{array}{ccccccccc} q_{11} & p & q_{22} & r & q_{33} & s & q_{44} & t & q_{55} \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}$$

and has a good  $A_5$  neighborhood, then there exists  $i \in \{2, 3, 4\}$  satisfies  $q_{ii} = -1$ .

*Proof.* By the definition of the good  $A_5$  neighborhood, the claim is clear.  $\square$

**Lemma 4.2.** *If  $\mathcal{D}_{\chi,E}$  is of the form*

$$\begin{array}{ccccccccc} q_{11} & p & q_{22} & r & q_{33} & s & q_{44} & t & q_{55} \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}$$

and has a good  $A_5$  neighborhood, then the following are impossible to appear.


- (a<sub>1</sub>)  $\begin{array}{ccccccccc} q^{-1} & q & q^{-1} & q & -1 & -1 & -1 & -1 & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$
- (a<sub>2</sub>)  $\begin{array}{ccccccccc} -1 & -1 & -1 & -1 & -1 & q & q^{-1} & q & q^{-1} \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$
- (a<sub>3</sub>)  $\begin{array}{ccccccccc} -1 & -1 & -1 & -1 & -1 & q & q^{-1} & q & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$
- (a<sub>4</sub>)  $\begin{array}{ccccccccc} -1 & q & q^{-1} & q & -1 & -1 & -1 & -1 & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$
- (b<sub>1</sub>)  $\begin{array}{ccccccccc} -1 & q^{-2} & q^2 & q^{-2} & q^2 & q^{-2} & -1 & q & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \notin G'_3, p \neq 3$
- (b<sub>2</sub>)  $\begin{array}{ccccccccc} -1 & q^2 & -1 & q^{-2} & q^2 & q^{-2} & -1 & q & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \notin G'_3, p \neq 3$
- (b<sub>3</sub>)  $\begin{array}{ccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & q & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$
- (c<sub>1</sub>)  $\begin{array}{ccccccccc} q^{-1} & q & -1 & q^{-1} & q & q^{-1} & -1 & -1 & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$
- (c<sub>2</sub>)  $\begin{array}{ccccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & -1 & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$
- (c<sub>3</sub>)  $\begin{array}{ccccccccc} -1 & q & -1 & q^{-1} & q & q^{-1} & -1 & -1 & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$
- (c<sub>4</sub>)  $\begin{array}{ccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & q^{-1} & q \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}, \quad q \neq -1, p \neq 2$

*Proof.* Consider the following cases.

- (a<sub>1</sub>) If  $X$  is of the type  $\begin{array}{ccccccccc} q_{11} & q & q^{-1} & q & -1 & -1 & -1 & -1 & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}$  then it satisfies the case ( $A_{5a}$ ) in the definition of the good  $A_5$  neighborhood. Then the transformation

$$\begin{array}{ccccccccc} q_{11} & q & q^{-1} & q & -1 & -1 & -1 & -1 & -1 \\ \circ & & \circ & & \bullet & & \circ & & \circ \end{array} \Rightarrow \begin{array}{ccccccc} & & & & -1 & & \\ & & & & \circ & & \\ & & q^{-1} & & -1 & & \\ & & / & & -1 & & \\ q_{11} & q & -1 & & -q & & -1 & -1 & -1 \\ \circ & & \circ & & \circ & & \circ & & \circ \end{array}$$

implies  $q_{11} = q^{-1}$  or  $q_{11} = -1$ . By  $a_{41}^{r_2 r_3(X)} = 0$ , we get  $q \in G'_4$  and hence the further transformation




where

$$q'_{11} = \begin{cases} -1 & \text{if } q_{11} = q^{-1} \\ q & \text{if } q_{11} = -1 \end{cases}$$


implies  $a_{45}^{r_2 r_3(X)} = -2$ , which contradicts to  $a_{45}^{r_2 r_3(X)} = -1$ . Hence we also get the situation in  $(a_4)$ .

(a<sub>2</sub>) If  $X$  is of the type  $\overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q_{55}}{\circ}$  then the transformation




implies  $q_{55} = q^{-1}$  or  $q_{55} = -1$ . By  $a_{41}^{r_2 r_3(X)} = 0$ , we get  $q = 1$ , which gives a contradiction. Hence the case in  $(a_3)$  is also impossible to appear.

(b<sub>1</sub>) If  $X$  is of the type  $\overset{-1}{\circ} \overset{q^{-2}}{\circ} \overset{q^2}{\circ} \overset{q^{-2}}{\circ} \overset{q^2}{\circ} \overset{q^{-2}}{\circ} \overset{-1}{\circ} \overset{q}{\circ} \overset{-1}{\circ}$  then it satisfies the case  $(A_{5b})$  in the definition of the good  $A_5$  neighborhood. Then the transformation




implies  $a_{25}^{r_3 r_4(X)} = -1$  by  $q \notin G'_3$ . By  $a_{15}^{r_2 r_3 r_4(X)} = 0$ , we get  $q \in G'_5$ . Then the following transformation




gives  $a_{45}^{r_3 r_2 r_1(X)} = -3$ , which contradicts to  $a_{45}^{r_3 r_2 r_1(X)} \neq -3$  in Definition 3.1.

(b<sub>2</sub>) If  $X$  is of the type  $\overset{-1}{\circ} \overset{q^2}{\circ} \overset{-1}{\circ} \overset{q^{-2}}{\circ} \overset{q^2}{\circ} \overset{q^{-2}}{\circ} \overset{-1}{\circ} \overset{q}{\circ} \overset{-1}{\circ}$  then it satisfies the case (A<sub>5b</sub>) in Definition 3.1. Then we get




and hence  $q = 1$  by  $a_{15}^{r_2 r_3 r_4(X)} = 0$ , which gives a contradiction.

(b<sub>3</sub>) If  $X$  is of the type  $\overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{q}{\circ} \overset{-1}{\circ}$  it satisfies the case (A<sub>5b</sub>) in Definition 3.1. We obtain




and hence  $q \in G'_4$  by  $a_{53}^{r_4(X)} = -1$ . Then the further transformation is




and hence  $a_{15}^{r_2 r_3 r_4(X)} = -1$ , which contradicts to  $a_{15}^{r_2 r_3 r_4(X)} = 0$  in Definition 3.1.

(c<sub>1</sub>) If  $X$  is of the type  $\overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{-1}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ}$  then it satisfies the case (A<sub>5c</sub>) in Definition 3.1. On the one hand, we get



and hence  $q \in G'_3$  by  $a_{52}^{r_3 r_4(X)} = -1$ . On the other, the transformation



implies that  $a_{32}^{r_5 r_4(X)} = -2$  by  $q \in G'_3$ , which contradicts to  $a_{32}^{r_5 r_4(X)} = -1$ . The similar reflections and conditions happen in cases (c<sub>2</sub>) and (c<sub>3</sub>). Hence the following two cases (c<sub>2</sub>) and (c<sub>3</sub>) can be discussed by the same argument as in the proof of (c<sub>1</sub>).



(c<sub>4</sub>) If  $X$  is of the type  $\overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ}$  then we get

$$\begin{array}{c} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{q^{-1}}{\bullet} \overset{q}{\circ} \\ \Rightarrow \begin{array}{c} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \begin{array}{l} \overset{-1}{\circ} \\ \overset{q}{\circ} \\ \overset{-q^{-1}}{\bullet} \\ \overset{-1}{\circ} \end{array} \end{array} \\ \Rightarrow \tau_{12354} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-q^{-1}-q}{\circ} \overset{-1}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \end{array}$$

which implies  $q = 1$  by  $a_{32}^{r_5 r_4(X)} = -1$ , which is a contradiction. □

**Theorem 4.3.** *All the possible Dynkin diagrams of  $V$  such that  $\mathcal{B}(V)$  has a finite root system listed in Appendix A appear in rows 11 to 15 of Table 1.*

*Proof.* Let  $X = [M]_5^s$ ,  $I = \{1, 2, 3, 4, 5\}$ , and  $A^X := (a_{ij})_{i,j \in I}$  be the Cartan matrix of  $X$ . Let  $(q_{i,j})_{i,j \in I}$  be the braiding matrix of  $X$  and  $(q_{i,j}^{r_i(X)})_{i,j \in I}$  be the braiding matrix of  $r_i(X)$ . Let  $a := -a_{24}^{r_3(X)}$ ,  $b := -a_{42}^{r_3(X)}$ ,  $c := -a_{35}^{r_4(X)}$ ,  $d := -a_{53}^{r_4(X)}$ , and  $e := -a_{43}^{r_5(X)}$ . To simply the labels, we write  $q'_{ij} := q_{ij}q_{ji}$  for  $1 \leq i, j \leq 5$ . By Theorem 3.4,  $A^X$  has a good  $A_5$  neighborhood. By  $A^X = A_5$ , we get  $(2)_{q_{ii}}(q_{ii}q'_{i,i+1} - 1) = (2)_{q_{jj}}(q_{jj}q'_{j-1,j} - 1) = 0$ , for all  $i \in \{1, 2, 3, 4\}$  and  $j \in \{2, 3, 4, 5\}$ . Hence we distinguish the following cases:  $a_1, a_2, \dots, a_{20}$ .

Case  $a_1$ . Consider that  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 3, 4\}$  and  $j \in \{2, 3, 4, 5\}$ . By Lemma 4.1, this case is impossible to appear.

Case  $a_2$ . Consider that  $q_{11} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 3, 4\}$  and  $j \in \{2, 3, 4, 5\}$ . By Lemma 4.1, this case is impossible to appear. Further consider the symmetric case of this case that  $q_{55} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 3, 4\}$  and  $j \in \{2, 3, 4\}$ . By Lemma 4.1, this case is impossible to appear.

Case  $a_3$ . Consider that  $q_{22} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 3, 4\}$  and  $j \in \{3, 4, 5\}$ . Let  $q := q_{11}$  and  $r := q_{33}$ . Since  $X$  has a good  $A_5$  neighborhood, we get  $qr = 1$  by  $A^{r_2(X)} = A_5$ . Rewrite  $X$  as

$$\overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ}$$

which implies  $(a, b, c, d, e) = (0, 0, 0, 0, 1)$ . By Definition 3.1 this case is impossible to appear. Further consider the symmetric case of this case that  $q_{44} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 3\}$  and  $j \in \{2, 3, 5\}$ . Let  $q := q_{11}$  and  $r := q_{55}$ . Hence  $(a, b, e) = (0, 0, 1)$ . Since  $X$  has a good  $A_5$  neighborhood, we have  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . Then  $qr \neq 1$ . On the one hand, the reflections of  $X$

$$X : \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{-1}{\bullet} \overset{r^{-1}}{\circ} \overset{r}{\circ} \Rightarrow r_4(X) : \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q^{-1}-1}{\circ} \begin{array}{l} \overset{q}{\circ} \\ \overset{r}{\circ} \\ \overset{(qr)^{-1}}{\bullet} \\ \overset{-1}{\circ} \end{array}$$



implies  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . Then the reflections of  $X$

$$X : \begin{array}{c} \circ \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \bullet \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} \circ \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array}$$

$$\Rightarrow r_5 r_4(X) : \tau_{12354} \begin{array}{c} \circ \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} -q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} -q \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array}$$

imply  $q^2 = -1$  by  $a_{32}^{r_5 r_4(X)} = -1$ . Thus  $q \in G'_4$ . In addition, the reflection of  $X$

$$r_4(X) : \begin{array}{c} \circ \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \bullet \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \Rightarrow r_3 r_4(X) : \tau_{12534} \begin{array}{c} \circ \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} -q^{-2} \\ | \\ \circ \end{array} \begin{array}{c} -q \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array}$$

implies  $q^3 = 1$  and  $p \neq 3$  by  $a_{52}^{r_3 r_4(X)} = -1$ . Thus we get  $q \in G'_3$ , which contradicts to  $q \in G'_4$ .

Subcase  $a_{5b}$ . We discuss the subcase  $(a, b, c, d, e) = (0, 0, 1, 1, 2)$ . That is  $A^{r_3(X)} = A_5$ ,

$$A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & -0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix} \text{ and } A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & -2 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix}. \text{ Then } qr \neq 1.$$

Then the reflection of  $X$

$$X : \begin{array}{c} \circ \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \bullet \end{array} \begin{array}{c} r^{-1} \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} \circ \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} r \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array}$$

implies  $r = -1$  or  $qr^2 = 1$  by  $a_{53}^{r_4(X)} = -1$ . If  $r = -1$ , the reflection of  $X$  is

$$X : \begin{array}{c} \circ \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \bullet \end{array} \Rightarrow r_5(X) : \begin{array}{c} \circ \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} q \\ | \\ \circ \end{array} \begin{array}{c} q^{-1} \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array}$$

which contradicts to  $e = 2$ . If  $qr^2 = 1$ , we have either  $a_{25}^{r_3 r_4(X)} = 0$  or  $a_{25}^{r_3 r_4(X)} = -1$  by Definition 3.1 ( $A_{5b}$ ). When  $a_{25}^{r_3 r_4(X)} = 0$ , we get  $r^3 = 1$ . Then  $q = r \in G'_3$  and  $p \neq 3$ . Hence  $\mathcal{D} = \mathcal{D}_{12,2}^5$ . When  $a_{25}^{r_3 r_4(X)} = -1$ , the reflection of  $X$

$$r_4(X) : \begin{array}{c} \circ \\ | \\ \circ \end{array} \begin{array}{c} r^{-2} \\ | \\ \circ \end{array} \begin{array}{c} r^2 \\ | \\ \circ \end{array} \begin{array}{c} r^{-2} \\ | \\ \circ \end{array} \begin{array}{c} r^2 \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \bullet \end{array} \begin{array}{c} r^{-2} \\ | \\ \circ \end{array} \begin{array}{c} r \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \begin{array}{c} r^{-1} \\ | \\ \circ \end{array} \Rightarrow r_3 r_4(X) : \tau_{12534} \begin{array}{c} \circ \\ | \\ \circ \end{array} \begin{array}{c} r^{-2} \\ | \\ \circ \end{array} \begin{array}{c} r^2 \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \begin{array}{c} r^{-2} \\ | \\ \circ \end{array} \begin{array}{c} r^3 \\ | \\ \circ \end{array} \begin{array}{c} r^{-1} \\ | \\ \circ \end{array} \begin{array}{c} -1 \\ | \\ \circ \end{array} \begin{array}{c} r^2 \\ | \\ \circ \end{array} \begin{array}{c} r^{-2} \\ | \\ \circ \end{array}$$

implies  $r \in G'_5$ . Hence  $\mathcal{D} = \mathcal{D}_{15,6}^5$ ,  $p \neq 5$ .

Case  $a_6$ . Consider that  $q_{11} = q_{33} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 4\}$  and  $j \in \{2, 4, 5\}$ . Let  $q := q_{22}$  and  $r := q_{44}$ . Hence  $(c, d, e) = (0, 0, 1)$ . Since  $X$  has a good  $A_5$  neighborhood, we have  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . Then  $qr \neq 1$ . The reflection of  $X$

$$X : \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{r^{-1}} \begin{array}{c} r \\ \circ \end{array} \xrightarrow{r^{-1}} \begin{array}{c} r \\ \circ \end{array} \Rightarrow r_3(X) : \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{(qr)^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{r^{-1}} \begin{array}{c} r \\ \circ \end{array} \xrightarrow{r^{-1}} \begin{array}{c} r \\ \circ \end{array}$$

implies  $q^2r = 1$  by  $a_{41}^{r_2r_3(X)} = 0$ . And  $a_{45}^{r_2r_3(X)} = -1$  gives  $qr^2 = 1$  or  $qr = -1$ . Summing up all proceeding conditions, we get  $q = r \in G'_3$  and  $p \neq 3$ . Hence  $\mathcal{D} = \mathcal{D}_{12,13}^5$ . Further consider the symmetric case of this case that  $q_{33} = q_{55} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 4\}$  and  $j \in \{2, 4\}$ . Let  $q := q_{11}$  and  $r := q_{44}$ . Hence  $(c, d) = (0, 0)$ . Since  $X$  has a good  $A_5$  neighborhood, we have  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . Then  $qr \neq 1$ . Similar to the proof above, we get  $q^2r = 1$  by  $a_{41}^{r_2r_3(X)} = 0$ . And  $a_{45}^{r_2r_3(X)} = -1$  gives  $qr^2 = 1$  or  $qr = -1$ . Summing up all proceeding conditions, we get  $q = r \in G'_3$  and  $p \neq 3$ . Hence  $\mathcal{D} = \tau_{54321}\mathcal{D}_{12,13}^5$ .

Case  $a_7$ . Consider that  $q_{11} = q_{44} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 3\}$  and  $j \in \{2, 3, 5\}$ . Let  $q := q_{22}$  and  $r := q_{55}$ . Hence  $(a, b, e) = (0, 0, 1)$ . Since  $X$  has a good  $A_5$  neighborhood, we have  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . Then  $qr \neq 1$ . On the one hand, the reflections of  $X$

$$X : \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{r^{-1}} \begin{array}{c} r \\ \circ \end{array} \xrightarrow{r^{-1}} \begin{array}{c} r \\ \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{(qr)^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{r} \begin{array}{c} -1 \\ \circ \end{array}$$

$$\Rightarrow r_3r_4(X) : \tau_{12534} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{(qr)^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array}$$
(7)

imply  $qr = -1$  or  $q^3r^2 = 1$  by  $a_{52}^{r_3r_4(X)} = -1$ . On the other hand, the reflections of  $X$

$$X : \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{r^{-1}} \begin{array}{c} r \\ \circ \end{array} \xrightarrow{r^{-1}} \begin{array}{c} r \\ \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{(qr)^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{r} \begin{array}{c} -1 \\ \bullet \end{array}$$

$$\Rightarrow r_5r_4(X) : \tau_{12354} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} (qr)^{-1} \\ \circ \end{array} \xrightarrow{qr} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{r^{-1}} \begin{array}{c} r \\ \circ \end{array} \xrightarrow{r^{-1}} \begin{array}{c} r \\ \circ \end{array}$$

imply  $qr = -1$  or  $q^2r = 1$  by  $a_{32}^{r_5r_4(X)} = -1$ . Summing up all proceeding conditions, we get  $qr = -1$  and  $p \neq 2$ . Then the reflection (7) of  $X$  implies  $q^2 = -1$  by  $a_{15}^{r_2r_3r_4(X)} = 0$ . Hence  $q \in G'_4$ . Then the reflections of  $X$

$$X : \begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{-q^{-q^{-1}}} \begin{array}{c} -1 \\ \circ \end{array} \Rightarrow r_1(X) : \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{q} \begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q \\ \circ \end{array} \xrightarrow{q^{-1}} \begin{array}{c} -1 \\ \circ \end{array} \xrightarrow{-q^{-q^{-1}}} \begin{array}{c} -1 \\ \circ \end{array}$$

$$\Rightarrow r_2 r_1(X) : \begin{array}{c} q \quad q^{-1} \quad -1 \quad q \quad -1 \quad q^{-1} \quad -1 \quad -q^{-q^{-1}} \\ \circ \quad \circ \quad \bullet \quad \circ \quad \circ \end{array} \Rightarrow r_3 r_2 r_1(X) : \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad q \quad q^{-1} \quad -q^{-q^{-1}} \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}$$

imply  $a_{45}^{r_3 r_2 r_1(X)} = -3$ , which contradicts to  $a_{45}^{r_3 r_2 r_1(X)} \neq -3$ . Further consider the symmetric case of this case that  $q_{22} = q_{55} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 3, 4\}$  and  $j \in \{3, 4\}$ . Let  $q := q_{11}$  and  $r := q_{33}$ . Hence  $(a, b, c, d) = (0, 0, 0, 0)$ , which gives a contradiction by Definition 3.1.

Case  $a_8$ . Consider that  $q_{11} = q_{55} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 3, 4\}$  and  $j \in \{2, 3, 4\}$ . By Lemma 4.1, this case is impossible to appear.

Case  $a_9$ . Consider that  $q_{22} = q_{33} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 4\}$  and  $j \in \{4, 5\}$ . Let  $q := q_{11}$ ,  $r^{-1} := q'_{23}$  and  $s := q_{44}$ . Hence  $(c, d, e) = (0, 0, 1)$ . Since  $X$  has a good  $A_5$  neighborhood, we have  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . Then  $rs \neq 1$ . In addition, we have  $qr = 1$  by  $A^{r_2(X)} = A_5$ . Then the reflection of  $X$

$$X : \begin{array}{c} r^{-1} \quad r \quad -1 \quad r^{-1} \quad -1 \quad s^{-1} \quad s \quad s^{-1} \quad s \\ \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \Rightarrow r_2(X) : \begin{array}{c} -1 \quad r^{-1} \quad -1 \quad r \quad r^{-1} \quad s^{-1} \quad s \quad s^{-1} \quad s \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.2 ( $a_2$ ), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{33} = q_{44} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2\}$  and  $j \in \{2, 5\}$ . Let  $q := q_{11}$ ,  $r^{-1} := q'_{34}$  and  $s := q_{55}$ . Hence  $e = 1$ . Since  $X$  has a good  $A_5$  neighborhood, we have either  $(a, b, c, d) = (0, 0, 1, 1)$  or  $(a, b, c, d) = (1, 1, 0, 0)$ . Next we discuss in two subcases.

Subcase  $a_{9a}$ . We discuss the subcase  $(a, b, c, d) = (0, 0, 1, 1)$ . That is  $A^{r_3(X)} = A_5$  and

$$A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix}. \text{ Then } qr = 1 \text{ and } rs \neq 1. \text{ The reflection of } X$$

$$X : \begin{array}{c} r^{-1} \quad r \quad r^{-1} \quad r \quad -1 \quad r^{-1} \quad -1 \quad s^{-1} \quad s \\ \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \Rightarrow r_3(X) : \begin{array}{c} r^{-1} \quad r \quad -1 \quad r^{-1} \quad -1 \quad r \quad r^{-1} \quad s^{-1} \quad s \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.2 ( $c_4$ ), this subcase is impossible to appear.

Subcase  $a_{9b}$ . We discuss the subcase  $(a, b, c, d) = (1, 1, 0, 0)$ . That is

$$A^{r_3(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & -1 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & -1 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix} \text{ and } A^{r_4(X)} = A_5. \text{ Then } qr \neq 1 \text{ and } rs = 1. \text{ The}$$

reflection of  $X$

$$X : \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad -1 \quad r \quad r^{-1} \\ \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad r^{-1} \quad r \quad -1 \quad r^{-1} \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.2 ( $a_1$ ), this subcase is impossible to appear.

Case  $a_{10}$ . Consider that  $q_{22} = q_{44} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 3\}$  and  $j \in \{3, 5\}$ . Let  $q := q_{11}$ ,  $r := q_{33}$  and  $s := q_{55}$ . Hence  $(a, b, e) = (0, 0, 1)$ . Since  $X$  has a good  $A_5$  neighborhood, we have  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . Then  $rs \neq 1$ . In addition, we have  $qr = 1$  by  $A^{r_2(X)} = A_5$ . On the one hand, the reflections of  $X$

$$\begin{aligned}
X : \begin{array}{c} r^{-1} \quad r \quad -1 \quad r^{-1} \quad r \quad r^{-1} \quad -1 \quad s^{-1} \quad s \\ \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \end{array} &\Rightarrow r_4(X) : \begin{array}{c} r^{-1} \quad r \quad -1 \quad r^{-1} \quad -1 \quad r \\ \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \end{array} \begin{array}{c} \circ^{-1} \\ r \\ \circ^{-1} \\ s \\ \circ^{-1} \\ (rs)^{-1} \end{array} \\
&\Rightarrow r_3 r_4(X) : \begin{array}{c} (rs)^{-1} \\ r^{-2} s^{-1} \quad r s \\ r^{-1} \quad r \quad r^{-1} \quad r \quad -1 \quad r^{-1} \quad r \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}
\end{aligned}$$

imply  $rs = -1$  or  $r^3 s^2 = 1$  by  $a_{52}^{r_3 r_4(X)} = -1$ . On the other hand, the reflections of  $X$

$$\begin{aligned}
X : \begin{array}{c} r^{-1} \quad r \quad -1 \quad r^{-1} \quad r \quad r^{-1} \quad -1 \quad s^{-1} \quad s \\ \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \end{array} &\Rightarrow r_4(X) : \begin{array}{c} r^{-1} \quad r \quad -1 \quad r^{-1} \quad -1 \quad r \\ \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \end{array} \begin{array}{c} \circ^{-1} \\ r \\ \circ^{-1} \\ s \\ \bullet^{-1} \\ (rs)^{-1} \end{array} \\
&\Rightarrow r_5 r_4(X) : \begin{array}{c} r^{-1} \quad r \quad -1 \quad r^{-1} (rs)^{-1} r s \quad -1 \quad s^{-1} \quad s \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}
\end{aligned}$$

imply  $rs = -1$  or  $r^2 s = 1$  by  $a_{32}^{r_5 r_4(X)} = -1$ . Summing up all proceeding conditions, we get  $rs = -1$  and  $p \neq 2$ . Then  $r \in G'_4$  by  $a_{15}^{r_2 r_3 r_4(X)} = 0$ . Similar to the proof in Case  $a_7$ , we have  $a_{45}^{r_3 r_2 r_1(X)} = -3$ , which contradicts to  $a_{45}^{r_3 r_2 r_1(X)} \neq -3$ .

Case  $a_{11}$ . Consider that  $q_{11} = q_{22} = q_{33} = -1$ ,  $q_{44}q'_{45} - 1 = 0$  and  $q_{jj}q'_{j-1,j} - 1 = 0$  for  $j \in \{4, 5\}$ . Let  $q^{-1} := q'_{12}$ ,  $r^{-1} := q'_{23}$  and  $s := q_{44}$ . Hence  $(c, d, e) = (0, 0, 1)$ . Since  $X$  has a good  $A_5$  neighborhood, we have  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . Then  $rs \neq 1$ . In addition, we get  $qr = 1$  by  $A^{r_2(X)} = A_5$ . Then the reflection of  $X$

$$X : \begin{array}{c} -1 \quad r \quad -1 \quad r^{-1} \quad -1 \quad s^{-1} \quad s \quad s^{-1} \quad s \\ \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \Rightarrow r_2(X) : \begin{array}{c} r \quad r^{-1} \quad -1 \quad r \quad r^{-1} \quad s^{-1} \quad s \quad s^{-1} \quad s \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.2 ( $a_2$ ), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{33} = q_{44} = q_{55} = -1$ ,  $q_{22}q'_{12} - 1 = 0$  and  $q_{ii}q'_{i,i+1} - 1 = 0$  for  $i \in \{1, 2\}$ . Let  $q := q_{11}$ ,  $r^{-1} := q'_{34}$  and  $s^{-1} := q'_{45}$ . Since  $X$  has a good  $A_5$  neighborhood, we have either  $(a, b, c, d) = (1, 1, 0, 0)$  or  $(a, b, c, d) = (0, 0, 1, 1)$ . Next we discuss in two subcases.

Subcase  $a_{11a}$ . We discuss the subcase  $(a, b, c, d) = (1, 1, 0, 0)$ . That is

$$A^{r_3(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & -1 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & -1 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix} \text{ and } A^{r_4(X)} = A_5. \text{ Then } qr \neq 1 \text{ and } rs = 1. \text{ The}$$

reflection of  $X$

$$X : \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad -1 \quad r \quad -1 \\ \circ \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} \quad r^{-1} \quad r \quad -1 \quad r^{-1} \quad r \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.2 ( $a_1$ ), this subcase is impossible to appear.

Subcase  $a_{11b}$ . We discuss the subcase  $(a, b, c, d) = (0, 0, 1, 1)$ . That is  $A^{r_3(X)} = A_5$  and  $A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix}$ . Then  $qr = 1$  and  $rs \neq 1$ . The reflection of  $X$

$$X : \begin{array}{ccccccccc} r^{-1} & r & r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccccc} r^{-1} & r & -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.2 ( $b_3$ ), this subcase is impossible to appear.

Case  $a_{12}$ . Consider that  $q_{11} = q_{22} = q_{44} = -1$ ,  $q_{33}q'_{34} - 1 = 0$  and  $q_{jj}q'_{j-1,j} - 1 = 0$  for  $j \in \{3, 5\}$ . Let  $q^{-1} := q'_{12}$ ,  $r := q_{33}$  and  $s := q_{55}$ . Hence  $(a, b, e) = (0, 0, 1)$ . Since  $X$  has a good  $A_5$  neighborhood, we have  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . Then  $rs \neq 1$ . In addition, we get  $qr = 1$  by  $A^{r_2(X)} = A_5$ . On the one hand, the reflections of  $X$

$$X : \begin{array}{ccccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{ccccccc} -1 & r & -1 & r^{-1} & -1 & r & -1 \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ \\ & & & & & & \circ \\ & & & & & & | \\ & & & & & & s \\ & & & & & & \circ \\ & & & & & & -1 \\ & & & & & & (rs)^{-1} \end{array}$$

$$\Rightarrow r_3r_4(X) : \tau_{12534} \begin{array}{ccccccc} & & & & & & (rs)^{-1} \\ & & & & & & \circ \\ & & & & & & / \\ & & & & & & r^{-2}s^{-1} \\ & & & & & & \circ \\ & & & & & & | \\ & & & & & & rs \\ & & & & & & \circ \\ & & & & & & | \\ & & & & & & -1 \\ & & & & & & \circ \\ & & & & & & | \\ & & & & & & r \\ & & & & & & \circ \\ & & & & & & | \\ & & & & & & -1 \\ & & & & & & \circ \\ & & & & & & | \\ & & & & & & r \\ & & & & & & \circ \end{array}$$

imply  $rs = -1$  or  $r^3s^2 = 1$  by  $a_{52}^{r_3r_4(X)} = -1$ . On the other hand, the reflections of  $X$

$$X : \begin{array}{ccccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{ccccccc} -1 & r & -1 & r^{-1} & -1 & r & -1 \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ \\ & & & & & & \circ \\ & & & & & & | \\ & & & & & & s \\ & & & & & & \circ \\ & & & & & & -1 \\ & & & & & & (rs)^{-1} \end{array}$$

$$\Rightarrow r_5r_4(X) : \tau_{12354} \begin{array}{ccccccc} -1 & r & -1 & r^{-1} & (rs)^{-1} & rs & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ \\ & & & & & & \circ \\ & & & & & & | \\ & & & & & & s^{-1} \\ & & & & & & \circ \\ & & & & & & | \\ & & & & & & s \\ & & & & & & \circ \end{array}$$

imply  $rs = -1$  or  $r^2s = 1$  by  $a_{32}^{r_5r_4(X)} = -1$ . Summing up all proceeding conditions, we get  $rs = -1$  and  $p \neq 2$ . By  $a_{15}^{r_2r_3r_4(X)} = 0$ , we get  $rs = 1$ , which contradicts to  $rs = -1$ . Further consider the symmetric case of this case that  $q_{22} = q_{44} = q_{55} = -1$ ,  $q_{33}q'_{23} - 1 = 0$  and  $q_{ii}q'_{i,i+1} - 1 = 0$  for  $i \in \{1, 3\}$ . Let  $q := q_{11}$ ,  $r := q_{33}$  and  $s^{-1} := q'_{45}$ . Hence  $(a, b) = (0, 0)$ . Since  $X$  has a good  $A_5$  neighborhood, we have either  $(a, b, c, d, e) = (0, 0, 1, 1, 2)$  or  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . In addition, we get  $qr = 1$  by  $A^{r_2(X)} = A_5$ . Next we discuss in two subcases.

Subcase  $a_{12a}$ . We discuss the subcase  $(a, b, c, d, e) = (0, 0, 1, 1, 2)$ . That is  $A^{r_3(X)} = A_5$ ,

$$A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix} \text{ and } A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & -2 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix}. \text{ Then } rs \neq 1.$$

On the one hand, the reflection of  $X$

$$X : \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\bullet} \overset{s^{-1}}{\circ} \overset{-1}{\circ} \Rightarrow r_4(X) : \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \begin{array}{l} \overset{r}{\circ} \\ \diagdown \quad \diagup \\ \overset{(rs)^{-1}}{\circ} \\ \diagup \quad \diagdown \\ \overset{s}{\circ} \\ \overset{s^{-1}}{\circ} \end{array} \quad (8)$$

implies  $s = -1$  or  $rs^2 = 1$  by  $a_{53}^{r_4(X)} = -1$ . On the other hand, the reflection of  $X$

$$X : \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{s^{-1}}{\circ} \overset{-1}{\bullet} \Rightarrow r_5(X) : \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{-1}{\circ}$$

implies  $s \neq -1$  by  $a_{43}^{r_5(X)} = -2$ . Hence  $rs^2 = 1$ . By Definition 3.1 ( $A_{5b}$ ) we have either  $a_{25}^{r_3r_4(X)} = 0$  or  $a_{25}^{r_3r_4(X)} = -1$ . When  $a_{25}^{r_3r_4(X)} = 0$ , the reflection (8) of  $X$  implies  $r^2s = 1$ . Hence  $r = s \in G'_3$  and  $p \neq 3$ . Then the reflections of  $X$

$$\begin{aligned} X : \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{-1}{\bullet} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} &\Rightarrow r_2(X) : \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{r}{\circ} \overset{-1}{\bullet} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \\ &\Rightarrow r_3r_2(X) : \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \end{aligned}$$

imply  $[a_{45}^{r_3r_2r_1(X)}, a_{45}^{r_3r_2(X)}] = [-2, -2]$ , which contradicts to  $[a_{45}^{r_3r_2r_1(X)}, a_{45}^{r_3r_2(X)}] \neq [-2, -2]$ . When  $a_{25}^{r_3r_4(X)} = -1$ , the reflection of  $X$

$$r_4(X) : \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \begin{array}{l} \overset{r}{\circ} \\ \diagdown \quad \diagup \\ \overset{(rs)^{-1}}{\circ} \\ \diagup \quad \diagdown \\ \overset{s}{\circ} \\ \overset{s^{-1}}{\circ} \end{array} \Rightarrow r_3r_4(X) : \tau_{12534} \begin{array}{l} \overset{-r^{-1}s^{-2}}{\circ} \\ \overset{r^{-2}s^{-1}}{\circ} \quad \overset{rs}{\circ} \\ \overset{r}{\circ} \quad \overset{r}{\circ} \quad \overset{r}{\circ} \\ \overset{-1}{\circ} \quad \overset{r^{-1}}{\circ} \quad \overset{r}{\circ} \end{array}$$

implies  $r = -1$  or  $r^3s = 1$ . If  $r = -1$ , we have  $s = -1$  and  $p \neq 2$  by  $a_{15}^{r_2r_3r_4(X)} = 0$ , which gives a contradiction. If  $r^3s = 1$ , we have  $r \in G'_5$  by  $rs^2 = 1$ . Similar to the proof in Case  $a_7$ , we have  $a_{45}^{r_3r_2r_1(X)} = -3$ , which contradicts to  $a_{45}^{r_3r_2r_1(X)} \neq -3$ .

Subcase  $a_{12b}$ . We discuss the subcase  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . That is  $A^{r_3(X)} = A_5$ ,

$$A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix} \text{ and } A^{r_5(X)} = A_5. \text{ Then } rs \neq 1. \text{ The reflection of } X$$

$$X : \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{s^{-1}}{\circ} \overset{-1}{\bullet} \Rightarrow r_5(X) : \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{-1}{\circ}$$





implies  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . By Lemma 4.2 ( $c_2$ ) this subcase is impossible to appear.

Case  $a_{14}$ . Consider that  $q_{11} = q_{33} = q_{44} = -1$ ,  $q_{22}q'_{23} - 1 = 0$  and  $q_{jj}q'_{j-1,j} - 1 = 0$  for  $j \in \{2, 5\}$ . Let  $q := q_{22}$ ,  $r^{-1} := q'_{34}$  and  $s := q_{55}$ . Hence  $e = 1$ . Since  $X$  has a good  $A_5$  neighborhood, we have either  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$  or  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . Next we discuss in two subcases.

Subcase  $a_{14a}$ . We discuss the subcase  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . That is

$$A^{r_3(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & -1 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & -1 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix}, A^{r_4(X)} = A_5 \text{ and } A^{r_5(X)} = A_5. \text{ Then } qr \neq 1 \text{ and } rs = 1.$$

The reflection of  $X$

$$X : \begin{array}{ccccccccc} -1 & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & r & r^{-1} \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{ccccccccc} -1 & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & r^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . Thus this subcase is impossible to appear by Lemma 4.2 ( $a_4$ ).

Subcase  $a_{14b}$ . We discuss the subcase  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . That is  $A^{r_3(X)} = A_5$ ,

$$A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix} \text{ and } A^{r_5(X)} = A_5. \text{ Then } qr = 1 \text{ and } rs \neq 1. \text{ The}$$

reflection of  $X$

$$X : \begin{array}{ccccccccc} -1 & r & r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccccc} -1 & r & -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . Thus this subcase is impossible to appear by Lemma 4.2 ( $c_4$ ).

Further consider the symmetric case of this case that  $q_{22} = q_{33} = q_{55} = -1$ ,  $q_{44}q'_{34} - 1 = 0$  and  $q_{ii}q'_{i,i+1} - 1 = 0$  for  $i \in \{1, 4\}$ . Let  $q := q_{11}$ ,  $r^{-1} := q'_{23}$  and  $s := q_{44}$ . Hence  $(c, d) = (0, 0)$ . Since  $X$  has a good  $A_5$  neighborhood, we have  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . Then  $rs \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{ccccccccc} q & q^{-1} & -1 & r^{-1} & -1 & s^{-1} & s & s^{-1} & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{ccccccccc} -1 & q & -1 & r & r^{-1} & s^{-1} & s & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$  and  $r = -1$  by  $a_{34}^{r_2(X)} = -1$ . Then  $q = r = -1$  and  $p \neq 2$ . Thus this subcase is impossible to appear by Lemma 4.2 ( $a_3$ ).

Case  $a_{15}$ . Consider that  $q_{11} = q_{33} = q_{55} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 4\}$  and  $j \in \{2, 4\}$ . Let  $q := q_{22}$  and  $r := q_{44}$ . Hence  $(c, d) = (0, 0)$ . Since  $X$  has a good  $A_5$  neighborhood, we have  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . Then  $qr \neq 1$ . The reflection of  $X$

$$X : \begin{array}{ccccccccc} -1 & q^{-1} & q & q^{-1} & -1 & r^{-1} & r & r^{-1} & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccccc} -1 & q^{-1} & -1 & -1 & -1 & r^{-1} & -1 & r^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $q^2r = 1$  by  $a_{41}^{r_2r_3(X)} = 0$ . And  $a_{45}^{r_2r_3(X)} = -1$  gives  $qr = -1$  or  $qr^2 = 1$ . Summing up all proceeding conditions, we get  $q = r \in G'_3$  and  $p \neq 3$ . Hence the reflections of  $X$

$$\begin{aligned} X : \begin{array}{ccccccccc} -1 & q^{-1} & q & q^{-1} & -1 & r^{-1} & r & r^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ \end{array} & \Rightarrow & r_5(X) : \begin{array}{ccccccccc} -1 & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & r & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ \end{array} \\ & & \Rightarrow & r_4r_5(X) : \begin{array}{ccccccccc} -1 & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & r^{-1} & r \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \end{aligned}$$

imply  $a_{32}^{r_4r_5(X)} = -2$ . And the reflections of  $X$

$$\begin{aligned} X : \begin{array}{ccccccccc} -1 & q^{-1} & q & q^{-1} & -1 & r^{-1} & r & r^{-1} & -1 \\ \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} & \Rightarrow & r_1(X) : \begin{array}{ccccccccc} -1 & q & -1 & q^{-1} & -1 & r^{-1} & r & r^{-1} & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \\ & & \Rightarrow & r_2r_1(X) : \begin{array}{ccccccccc} q & q^{-1} & -1 & q & q^{-1} & r^{-1} & r & r^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \end{aligned}$$

imply  $a_{34}^{r_2r_1(X)} = -2$ , which contradicts to  $[a_{32}^{r_4r_5(X)}, a_{34}^{r_2r_1(X)}] \neq [-2, -2]$ .

Case  $a_{16}$ . Consider that  $q_{22} = q_{33} = q_{44} = -1$ ,  $q_{11}q'_{12} - 1 = 0$  and  $q_{55}q'_{45} - 1 = 0$ . Let  $q := q_{11}$ ,  $r^{-1} := q'_{23}$ ,  $s^{-1} := q'_{34}$  and  $t := q_{55}$ . Hence  $e = 1$ . Since  $X$  has a good  $A_5$  neighborhood, we have either  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$  or  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . In addition, we have  $qr = 1$  by  $A^{r_2(X)} = A_5$ . Next we discuss in two subcases.

Subcase  $a_{16a}$ . We discuss the subcase  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . That is

$$A^{r_3(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & -1 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & -1 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix}, A^{r_4(X)} = A_5 \text{ and } A^{r_5(X)} = A_5. \text{ Then } rs \neq 1 \text{ and } st = 1.$$

The reflection of  $X$

$$X : \begin{array}{ccccccccc} r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & -1 & s & s^{-1} \\ \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{ccccccccc} -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & -1 & s & s^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccccc} -1 & -1 & -1 & -1 & -1 & s^{-1} & -1 & s & s^{-1} \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{ccccccccc} -1 & -1 & -1 & -1 & s^{-1} & s & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ , which contradicts to  $rs \neq 1$ .

Subcase  $a_{16b}$ . We discuss the subcase  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . That is  $A^{r_3(X)} = A_5$ ,

$$A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix} \text{ and } A^{r_5(X)} = A_5. \text{ Then } rs = 1 \text{ and } st \neq 1. \text{ The}$$

reflection of  $X$

$$X : \begin{array}{ccccccccc} s & s^{-1} & -1 & s & -1 & s^{-1} & -1 & t^{-1} & t \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccccc} s & s^{-1} & s & s^{-1} & -1 & s & s^{-1} & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.2 ( $c_4$ ), this subcase is impossible to appear.

Case  $a_{17}$ . Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = -1$  and  $q_{55}q'_{45} - 1 = 0$ . Let  $q^{-1} := q'_{12}$ ,  $r^{-1} := q'_{23}$ ,  $s^{-1} := q'_{34}$  and  $t := q_{55}$ . Hence  $e = 1$ . Since  $X$  has a good  $A_5$  neighborhood, we have either  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$  or  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . In addition, we have  $qr = 1$  by  $A^{r_2(X)} = A_5$ . Next we discuss in two subcases.

Subcase  $a_{17a}$ . We discuss the subcase  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . That is

$$A^{r_3(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & -1 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & -1 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix}, A^{r_4(X)} = A_5 \text{ and } A^{r_5(X)} = A_5. \text{ Then } rs \neq 1 \text{ and } st = 1.$$

The reflection of  $X$

$$X : \begin{array}{ccccccccc} -1 & r & -1 & r^{-1} & -1 & s^{-1} & -1 & s & s^{-1} \\ \circ & \text{---} & \bullet & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{ccccccccc} r & r^{-1} & -1 & r & r^{-1} & s^{-1} & -1 & s & s^{-1} \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccccc} -1 & -1 & -1 & -1 & -1 & s^{-1} & -1 & s & s^{-1} \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \bullet & \text{---} & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{ccccccccc} -1 & -1 & -1 & -1 & s^{-1} & s & -1 & s^{-1} & -1 \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ , which contradicts to  $rs \neq 1$ .

Subcase  $a_{17b}$ . We discuss the subcase  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . That is  $A^{r_3(X)} = A_5$ ,

$$A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix} \text{ and } A^{r_5(X)} = A_5. \text{ Then } rs = 1 \text{ and } st \neq 1. \text{ Then the}$$

reflection of  $X$

$$X : \begin{array}{ccccccccc} -1 & s^{-1} & -1 & s & -1 & s^{-1} & -1 & t^{-1} & t \\ \circ & \text{---} & \circ & \text{---} & \bullet & \text{---} & \circ & \text{---} & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccccc} -1 & s^{-1} & s & s^{-1} & -1 & s & s^{-1} & t^{-1} & t \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.2 ( $c_4$ ), this subcase is impossible to appear.

Further consider the symmetric case of this case that  $q_{22} = q_{33} = q_{44} = q_{55} = -1$  and  $q_{11}q'_{12} - 1 = 0$ . Let  $q := q_{11}$ ,  $r^{-1} := q'_{23}$ ,  $s^{-1} := q'_{34}$  and  $t^{-1} := q'_{45}$ . Since  $X$  has a good  $A_5$  neighborhood, we have either  $(a, b, c, d) = (1, 1, 0, 0)$  or  $(a, b, c, d) = (0, 0, 1, 1)$ . In addition, we have  $qr = 1$  by  $A^{r_2(X)} = A_5$ . Next we discuss in two subcases.

Subcase  $a_{17c}$ . We discuss the subcase  $(a, b, c, d) = (1, 1, 0, 0)$ . That is

$$A^{r_3(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & -1 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & -1 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix} \text{ and } A^{r_4(X)} = A_5. \text{ Then } rs \neq 1 \text{ and } st = 1. \text{ The}$$

reflection of  $X$

$$X : \begin{array}{ccccccccc} r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & -1 & s & -1 \\ \circ & \text{---} & \bullet & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{ccccccccc} -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & -1 & s & -1 \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccccc} -1 & -1 & -1 & -1 & -1 & s^{-1} & -1 & s & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{ccccccccc} -1 & -1 & -1 & -1 & s^{-1} & s & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ , which contradicts to  $rs \neq 1$ .

Subcase  $a_{17d}$ . We discuss the subcase  $(a, b, c, d) = (0, 0, 1, 1)$ . That is  $A^{r_3(X)} = A_5$  and

$$A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix}. \text{ Then } rs = 1 \text{ and } st \neq 1. \text{ The reflection of } X$$

$$X : \begin{array}{ccccccccc} s & s^{-1} & -1 & s & -1 & s^{-1} & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccccc} s & s^{-1} & s & s^{-1} & -1 & s & s^{-1} & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.2 ( $b_3$ ), this subcase is impossible to appear.

Case  $a_{18}$ . Consider that  $q_{11} = q_{22} = q_{33} = q_{55} = -1$ ,  $q_{44}q'_{45} - 1 = 0$  and  $q_{44}q'_{34} - 1 = 0$ . Let  $q^{-1} := q'_{12}$ ,  $r^{-1} := q'_{23}$  and  $s := q_{44}$ . Hence  $(c, d) = (0, 0)$ . Since  $X$  has a good  $A_5$  neighborhood, we have  $(a, b, c, d, e) = (1, 1, 0, 0, 1)$ . Then  $rs \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{ccccccccc} -1 & q^{-1} & -1 & r^{-1} & -1 & s^{-1} & s & s^{-1} & -1 \\ \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{ccccccccc} q^{-1} & q & -1 & r & r^{-1} & s^{-1} & s & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$  and  $r = -1$  by  $a_{34}^{r_2(X)} = -1$ . Then  $q = r = -1$  and  $p \neq 2$ . Thus this case is impossible to appear by Lemma 4.2 ( $a_3$ ). Further consider the symmetric case of this case that  $q_{11} = q_{33} = q_{44} = q_{55} = -1$ ,  $q_{22}q'_{23} - 1 = 0$  and  $q_{22}q'_{12} - 1 = 0$ . Let  $q := q_{22}$ ,  $r^{-1} := q'_{34}$  and  $s^{-1} := q'_{45}$ . Since  $X$  has a good  $A_5$  neighborhood, we have either  $(a, b, c, d) = (1, 1, 0, 0)$  or  $(a, b, c, d) = (0, 0, 1, 1)$ . Next we discuss in two subcases.

Subcase  $a_{18a}$ . We discuss the subcase  $(a, b, c, d) = (1, 1, 0, 0)$ . That is

$$A^{r_3(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & -1 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & -1 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix} \text{ and } A^{r_4(X)} = A_5. \text{ Then } qr \neq 1 \text{ and } rs = 1. \text{ The}$$

reflection of  $X$

$$X : \begin{array}{ccccccccc} -1 & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & r & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \bullet & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{ccccccccc} -1 & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & r^{-1} & r \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.2 ( $a_4$ ), this subcase is impossible to appear.

Subcase  $a_{18b}$ . We discuss the subcase  $(a, b, c, d) = (0, 0, 1, 1)$ . That is  $A^{r_3(X)} = A_5$  and

$$A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix}. \text{ Then } qr = 1 \text{ and } rs \neq 1. \text{ The reflection of } X$$

$$X : \begin{array}{ccccccccc} -1 & r & r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccccc} -1 & r & -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$



Subcase  $a_{19b}$ . We discuss the subcase  $(a, b, c, d, e) = (0, 0, 1, 1, 1)$ . That is  $A^{r_3(X)} = A_5$ ,

$$A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix} \text{ and } A^{r_5(X)} = A_5. \text{ Then } rs \neq 1. \text{ The reflection of } X$$

$$X : \begin{array}{ccccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{ccccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & s^{-1} & s & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . By Lemma 4.2 ( $c_3$ ), this subcase is impossible to appear.

Case  $a_{20}$ . Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = q_{55} = -1$ . Let  $q^{-1} := q'_{12}$ ,  $r^{-1} := q'_{23}$ ,  $s^{-1} := q'_{34}$  and  $t^{-1} := q'_{45}$ . Since  $X$  has a good  $A_5$  neighborhood, we have either  $(a, b, c, d) = (1, 1, 0, 0)$  or  $(a, b, c, d) = (0, 0, 1, 1)$ . In addition, we have  $qr = 1$  by  $A^{r_2(X)} = A_5$ . Next we discuss in two subcases.

Subcase  $a_{20a}$ . We discuss the subcase  $(a, b, c, d) = (1, 1, 0, 0)$ . That is

$$A^{r_3(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & -1 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & -1 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix} \text{ and } A^{r_4(X)} = A_5. \text{ Then } rs \neq 1 \text{ and } st = 1. \text{ The reflection of } X$$

$$X : \begin{array}{ccccccccc} -1 & r & -1 & r^{-1} & -1 & s^{-1} & -1 & s & -1 \\ \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{ccccccccc} r & r^{-1} & -1 & r & r^{-1} & s^{-1} & -1 & s & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccccc} -1 & -1 & -1 & -1 & -1 & s^{-1} & -1 & s & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{ccccccccc} -1 & -1 & -1 & -1 & s^{-1} & s & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ , which contradicts to  $rs \neq 1$ .

Subcase  $a_{20b}$ . We discuss the subcase  $(a, b, c, d) = (0, 0, 1, 1)$ . That is  $A^{r_3(X)} = A_5$  and

$$A^{r_4(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix}. \text{ Then } rs = 1 \text{ and } st \neq 1. \text{ The reflection of } X$$


$$X : \begin{array}{ccccccccc} -1 & s^{-1} & -1 & s & -1 & s^{-1} & -1 & t^{-1} & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccccc} -1 & s^{-1} & s & s^{-1} & -1 & s & s^{-1} & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.2 ( $b_3$ ), this subcase is impossible to appear.  $\square$






(a<sub>2</sub>) Assume that  $X$  is of type  $\overset{q^{-1}}{\circ} - \overset{q}{\circ} - \overset{-1}{\circ} - \overset{q^{-1}}{\circ} - \overset{q}{\circ} - \overset{q^{-1}}{\circ} - \overset{-1}{\circ} - \overset{-1}{\circ} - \overset{-1}{\circ} - \overset{-1}{\circ}$  where  $q \neq -1$ . We have



Then we get  $q = 1$  by  $a_{36}^{r_5 r_4} = 0$ , which contradicts to  $q \neq 1$ . The similar reflections and conditions happen in cases (a<sub>3</sub>), (a<sub>4</sub>) and (a<sub>5</sub>). Hence the three cases (a<sub>3</sub>), (a<sub>4</sub>) and (a<sub>5</sub>) can be discussed by the same argument as in the proof of (a<sub>2</sub>).

(b<sub>1</sub>) If  $X$  is of type  $\overset{-1}{\circ} - \overset{-1}{\circ} - \overset{-1}{\circ} - \overset{-1}{\circ} - \overset{-1}{\circ} - \overset{-1}{\circ} - \overset{-1}{\circ} - \overset{q^{-1}}{\circ} - \overset{-1}{\circ} - \overset{-1}{\circ} - \overset{-1}{\circ}$  where  $q \in G'_4$ , then we get





We obtain  $q = 1$  by  $a_{25}^{r_3 r_4(X)} = 0$ , which contradicts to  $q \neq 1$ . The similar reflections and conditions happen in cases (b<sub>2</sub>), (b<sub>3</sub>), (b<sub>4</sub>), (b<sub>5</sub>) and (b<sub>6</sub>). Hence the cases (b<sub>2</sub>)–(b<sub>6</sub>) can be discussed by the same argument as in the proof of (b<sub>1</sub>).

□

**Theorem 4.6.** *All the possible Dynkin diagrams of  $V$  such that  $\mathcal{B}(V)$  has a finite root system listed in Appendix B appear in rows 17 to 19 of Table 1.*

*Proof.* Let  $X = [M]_6^s$ ,  $I = \{1, 2, 3, 4, 5, 6\}$  and  $A^X := (a_{ij})_{i,j \in I}$  be the Cartan matrix of  $X$ . Let  $(q_{i,j})_{i,j \in I}$  be the braiding matrix of  $X$  and  $(q_{i,j}^{r_i(X)})_{i,j \in I}$  be the braiding matrix of  $r_i(X)$ . Let  $a := -a_{46}^{r_5(X)}$ . To simplify the labels, we write  $q'_{ij} := q_{ij}q_{ji}$  for  $1 \leq i, j \leq 6$ . By Theorem 3.4,  $A^X$  has a good  $A_6$  neighborhood. By  $A^X = A_6$ , we get  $(2)_{q_{ii}}(q_{ii}q'_{i,i+1} - 1) = (2)_{q_{jj}}(q_{jj}q'_{j-1,j} - 1) = 0$ , for all  $i \in \{1, 2, 3, 4, 5\}$  and  $j \in \{2, 3, 4, 5, 6\}$ . From Lemma 4.4, we know  $q_{44} = -1$  for all cases. Depending on the number of vertices that satisfy  $q_{ii} = -1$ , we divide the cases into the following categories: Step 1 to Step 6.

Step 1. Assume that there is one vertex labeled by  $-1$ . Since Lemma 4.4 implies  $q_{44} = -1$ , there is only one case:  $q_{44} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 3, 5\}$  and  $j \in \{2, 3, 5, 6\}$ . Let  $q := q_{11}$  and  $r := q_{55}$ . Hence  $a = 0$ . Since  $X$  has a good  $A_6$  neighborhood, we have  $qr \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . The reflections of  $X$



imply  $rq = -1$  or  $q^2r = 1$  by  $a_{32}^{r_5r_4(X)} = -1$ . And  $a_{36}^{r_5r_4(X)} = 0$  gives  $qr^2 = 1$ . Summing up all proceeding conditions, we get  $q = r \in G'_3$ . Hence  $\mathcal{D} = \mathcal{D}_{17,1}^6$  and  $p \neq 3$ .

Step 2. Assume that there are two vertices labeled by  $-1$ . By  $q_{44} = -1$ , we distinguish the following five steps.

Step 2.1. Consider that  $q_{11} = q_{44} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 3, 5\}$  and  $j \in \{2, 3, 5, 6\}$ . Let  $q := q_{22}$  and  $r := q_{55}$ . Hence  $a = 0$ . Since  $X$  has a good  $A_6$  neighborhood, we have  $qr \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . The reflection of  $X$

$$X : \begin{array}{cccccccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & r^{-1} & r & r^{-1} & r \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccccccc} & & & & & & & & & & & -1 \\ & & & & & & & & & & & \circ \\ & & & & & & & & & & & / \quad \backslash \\ & & & & & & & & & & & q \quad r \\ & & & & & & & & & & & \circ \\ & & & & & & & & & & & \backslash \quad / \\ & & & & & & & & & & & (qr)^{-1} \\ & & & & & & & & & & & \circ \\ & & & & & & & & & & & -1 \\ & & & & & & & & & & & \circ \\ & & & & & & & & & & & \circ \end{array}$$

implies  $r = q^2$  by  $a_{25}^{r_3r_4(X)} = 0$  and  $q = r^2$  by  $a_{36}^{r_5r_4(X)} = 0$ . Then  $r = q \in G'_3$ . Hence  $\mathcal{D} = \mathcal{D}_{18,19}^6$  and  $p \neq 3$ . Further consider the symmetric case of this case that  $q_{33} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 4, 5\}$  and  $j \in \{2, 4, 5\}$ . By Lemma 4.4, this case is impossible to appear.

Step 2.2. Consider that  $q_{22} = q_{44} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 3, 5\}$  and  $j \in \{3, 5, 6\}$ . Let  $q := q_{11}$ ,  $r := q_{33}$  and  $s := q_{55}$ . Hence  $a = 0$ . Since  $X$  has a good  $A_6$  neighborhood, we get  $qr = 1$  and  $rs \neq 1$ . The reflection of  $X$

$$X : \begin{array}{cccccccccccc} q & q^{-1} & -1 & q & q^{-1} & q & -1 & s^{-1} & s & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccccccc} & & & & & & & & & & & -1 \\ & & & & & & & & & & & \circ \\ & & & & & & & & & & & / \quad \backslash \\ & & & & & & & & & & & q^{-1} \quad s \\ & & & & & & & & & & & \circ \\ & & & & & & & & & & & \backslash \quad / \\ & & & & & & & & & & & qs^{-1} \\ & & & & & & & & & & & \circ \\ & & & & & & & & & & & -1 \\ & & & & & & & & & & & \circ \\ & & & & & & & & & & & s^{-1} \\ & & & & & & & & & & & \circ \\ & & & & & & & & & & & s \end{array}$$

implies  $s = q^2$  by  $a_{25}^{r_3r_4(X)} = 0$  and  $q = s^2$  by  $a_{36}^{r_5r_4(X)} = 0$ . Hence  $q \in G'_3$  and  $p \neq 3$ . Then the reflections of  $X$

$$X : \begin{array}{cccccccccccc} q & q^{-1} & -1 & q & q^{-1} & q & -1 & s^{-1} & s & s^{-1} & s \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccccccc} -1 & q & -1 & q^{-1} & -1 & q & -1 & s^{-1} & s & s^{-1} & s \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \\ \Rightarrow r_3r_2(X) : \begin{array}{cccccccccccc} -1 & q & q^{-1} & q & -1 & q^{-1} & q & s^{-1} & s & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

imply  $[a_{45}^{r_3r_2r_1(X)}, a_{45}^{r_3r_2(X)}] = [-2, -2]$ , which contradicts to  $[a_{45}^{r_3r_2r_1(X)}, a_{45}^{r_3r_2(X)}] \neq [-2, -2]$ . Further by Lemma 4.4 the symmetric case is impossible to appear.

Step 2.3. Consider that  $q_{33} = q_{44} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 5\}$  and  $j \in \{2, 5, 6\}$ . Let  $q := q_{11}$ ,  $r^{-1} := q'_{34}$ ,  $s := q_{55}$ . Hence  $a = 0$ . Since  $X$  has a good  $A_6$  neighborhood, we have  $qr = 1$  and  $rs \neq 1$ . Then the reflection of  $X$

$$X : \begin{array}{cccccccccccc} r^{-1} & r & r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & s & s^{-1} & s \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{cccccccccccc} r^{-1} & r & -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & s & s^{-1} & s \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.5 (b<sub>2</sub>), this case is impossible to appear.

Step 2.4. Consider that  $q_{44} = q_{55} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 3\}$  and  $j \in \{2, 3, 6\}$ . Let  $q := q_{11}$ ,  $r^{-1} := q'_{45}$  and  $s := q_{66}$ . Since  $X$  has a good

$A_6$  neighborhood, we have  $qr \neq 1$ . Then we get  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . The reflection of  $X$

$$X : \begin{array}{c} q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & -1 & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} & & & & & & & & -1 \\ & & & & & & & & \circ \\ & & & & & & & & / \quad \backslash \\ q & & & & & & & & -1 \\ & & & & & & & & \circ \\ & & & & & & & & / \quad \backslash \\ & & & & & & & & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $q^2 = -1$  by  $a_{25}^{r_3 r_4(X)} = 0$  and  $qs = -1$  by  $a_{36}^{r_5 r_4(X)} = 0$ . Hence  $q = s \in G'_4$  and  $p \neq 2$ . Rewrite  $X$  as

$$\begin{array}{c} q & -q & q & -q & q & -q & -1 & -1 & -1 & -q & q \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

and we have  $\mathcal{D} = \mathcal{D}_{19,1}^6$  and  $p \neq 2$ . Further by Lemma 4.4 the symmetric case is impossible to appear.

Step 2.5. Consider that  $q_{44} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 3, 5\}$  and  $j \in \{2, 3, 5\}$ . Let  $q := q_{11}$  and  $r := q_{55}$ . Hence  $a = 0$ . Since  $X$  has a good  $A_6$  neighborhood, we get  $qr \neq 1$ . The reflection of  $X$

$$X : \begin{array}{c} q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & r^{-1} & r & r^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} & & & & & & & & -1 \\ & & & & & & & & \circ \\ & & & & & & & & / \quad \backslash \\ q & & & & & & & & r \\ & & & & & & & & \circ \\ & & & & & & & & / \quad \backslash \\ & & & & & & & & -1 & r^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = q^{-2}$  by  $a_{25}^{r_3 r_4(X)} = 0$  and  $q = r^{-2}$  by  $a_{36}^{r_5 r_4(X)} = 0$ . Hence  $r = q \in G'_3$  and  $p \neq 3$ . Then the reflections of  $X$

$$X : \begin{array}{c} q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & q^{-1} & q & q^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_6(X) : \begin{array}{c} q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & q^{-1} & -1 & q & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \\ \Rightarrow r_5 r_6(X) : \begin{array}{c} q & q^{-1} & q & q^{-1} & q & q^{-1} & q^{-1} & q & -1 & q^{-1} & q \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

imply  $q = -1$  and  $p \neq 2$  by  $a_{43}^{r_5 r_6(X)} = -1$ , which contradicts to  $q \in G'_3$ . Further by Lemma 4.4 the symmetric case is impossible to appear.

Step 3. Assume that there are three vertices labeled by  $-1$ . By  $q_{44} = -1$ , we distinguish the following eight steps.

Step 3.1. Consider that  $q_{11} = q_{22} = q_{44} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{3, 5\}$  and  $j \in \{3, 5, 6\}$ . Let  $q^{-1} := q'_{12}$ ,  $r := q_{33}$  and  $s := q_{55}$ . Hence  $a = 0$ . Since  $X$  has a good  $A_6$  neighborhood, we get  $rs \neq 1$  by  $a_{35}^{r_4(X)} = -1$  and  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$ . The reflection of  $X$

$$X : \begin{array}{c} -1 & r & -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & s & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{c} & & & & & & & & -1 \\ & & & & & & & & \circ \\ & & & & & & & & / \quad \backslash \\ r & & & & & & & & s \\ & & & & & & & & \circ \\ & & & & & & & & / \quad \backslash \\ & & & & & & & & (rs)^{-1} & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = r^{-2}$  by  $a_{25}^{r_3 r_4(X)} = 0$  and  $r = s^{-2}$  by  $a_{36}^{r_5 r_4(X)} = 0$ . Hence  $r = s \in G'_3$  and  $p \neq 3$ . Rewrite  $X$  as

$$\begin{array}{c} -1 & r & -1 & r^{-1} & r & r^{-1} & -1 & r^{-1} & r & r^{-1} & r \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

and we have  $\mathcal{D} = \mathcal{D}_{18,16}^6$ . Further consider the symmetric case of this case that  $q_{33} = q_{55} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 4\}$  and  $j \in \{2, 4\}$ . By Lemma 4.4, this case is impossible to appear.

Step 3.2. Consider that  $q_{11} = q_{33} = q_{44} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 5\}$  and  $j \in \{2, 5, 6\}$ . Let  $q := q_{22}, r^{-1} := q'_{34}$  and  $s := q_{55}$ . Hence  $a = 0$ . Since  $X$  has a good  $A_6$  neighborhood, we get  $rs \neq 1$  and  $qr = 1$ . The reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & r & r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & s & s^{-1} & s \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{cccccccc} -1 & r & -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & s & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.5 (b<sub>2</sub>), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{33} = q_{44} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 5\}$  and  $j \in \{2, 5\}$ . Let  $q := q_{11}, r^{-1} := q'_{34}$  and  $s := q_{55}$ . Hence  $a = 0$ . Since  $X$  has a good  $A_6$  neighborhood, then  $rs \neq 1$  and  $qr = 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} q & q^{-1} & q & q^{-1} & -1 & q & -1 & s^{-1} & s & s^{-1} & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{cccccccc} q & q^{-1} & -1 & q & -1 & q^{-1} & q & s^{-1} & s & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $q = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.5 (b<sub>4</sub>), this case is impossible to appear.

Step 3.3. Consider that  $q_{11} = q_{44} = q_{55} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 3\}$  and  $j \in \{2, 3, 6\}$ . Let  $q := q_{22}, r^{-1} := q'_{45}$  and  $s := q_{66}$ . Since  $X$  has a good  $A_6$  neighborhood, we have either  $a = 0$  or  $a = 1$ . In addition, we have  $qr \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . Next we discuss in two steps.

Step 3.3.1. We discuss the subcase  $a = 0$ . That is


$$A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}. \text{ Then } rs = 1 \text{ and the reflection of } X$$

$$X : \begin{array}{cccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & r & r^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{cccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & r^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . By Lemma 4.5 (a<sub>2</sub>), this subcase is impossible to appear.

Step 3.3.2. We discuss the subcase  $a = 1$ . That is

$$A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & -1 & 2 \end{pmatrix}. \text{ Then } rs \neq 1. \text{ The reflection of } X$$

$$X : \begin{array}{cccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{cccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & r^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$


implies  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . Then the reflection of  $X$

$$\begin{aligned}
X : & \begin{array}{cccccccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & -1 & -1 & s^{-1} & s \\ \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_1(X) : \begin{array}{cccccccccccc} -1 & q & -1 & q^{-1} & q & q^{-1} & -1 & -1 & -1 & s^{-1} & s \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \\
\Rightarrow r_2 r_1(X) : & \begin{array}{cccccccccccc} q & q^{-1} & -1 & q & -1 & q^{-1} & -1 & -1 & -1 & s^{-1} & s \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \\
\Rightarrow r_3 r_2 r_1(X) : & \begin{array}{cccccccccccc} q & q^{-1} & q & q^{-1} & -1 & q & q^{-1} & -1 & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}
\end{aligned}$$

imply  $q = -1$  and  $p \neq 2$  by  $a_{45}^{r_3 r_2 r_1(X)} = -1$ , which contradicts to  $qr \neq 1$ .

Further by Lemma 4.4 the symmetric case is impossible to appear.

Step 3.4. Consider that  $q_{11} = q_{44} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 3, 5\}$  and  $j \in \{2, 3, 5\}$ . Let  $q := q_{22}$  and  $r := q_{55}$ . Hence  $a = 0$ . Since  $X$  has a good  $A_6$  neighborhood, we get  $qr \neq 1$ . The reflections of  $X$

$$\begin{aligned}
X : & \begin{array}{cccccccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & r^{-1} & r & r^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ \end{array} \Rightarrow r_6(X) : \begin{array}{cccccccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & r & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ \end{array} \\
\Rightarrow r_5 r_6(X) : & \begin{array}{cccccccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & r^{-1} & r \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}
\end{aligned}$$

imply  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5 r_6(X)} = -1$ . Thus this case is impossible to appear by Lemma 4.5 (a<sub>2</sub>). Further by Lemma 4.4 the symmetric case is impossible to appear.


Step 3.5. Consider that  $q_{22} = q_{33} = q_{44} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 5\}$  and  $j \in \{5, 6\}$ . Let  $q := q_{11}$ ,  $r^{-1} := q'_{23}$ ,  $s^{-1} := q'_{34}$  and  $t := q_{55}$ . Hence  $a = 0$ . Since  $X$  has a good  $A_6$  neighborhood, we get  $qr = 1$ ,  $rs = 1$  and  $st \neq 1$ . Then the reflection of  $X$

$$X : \begin{array}{cccccccccccc} s & s^{-1} & -1 & s & -1 & s^{-1} & -1 & t^{-1} & t & t^{-1} & t \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{cccccccccccc} s & s^{-1} & s & s^{-1} & -1 & s & s^{-1} & t^{-1} & t & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.5 (b<sub>2</sub>), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{33} = q_{44} = q_{55} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2\}$  and  $j \in \{2, 6\}$ . Let  $q := q_{11}$ ,  $r^{-1} := q'_{34}$ ,  $s^{-1} := q'_{45}$  and  $t := q_{66}$ . Since  $X$  has a good  $A_6$  neighborhood, then  $qr = 1$  and  $rs \neq 1$ . The reflection of  $X$

$$X : \begin{array}{cccccccccccc} r^{-1} & r & r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & -1 & t^{-1} & t \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{cccccccccccc} r^{-1} & r & -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & -1 & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . Hence  $s \neq -1$  and the reflection of  $X$

$$X : \begin{array}{cccccccccccc} r^{-1} & r & r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & -1 & t^{-1} & t \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccccccc} r^{-1} & r & r^{-1} & r & r^{-1} & r & r^{-1} & s^{-1} & -1 & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$


implies  $st = 1$  by  $a_{56}^{r_4(X)} = -1$ . By Lemma 4.5 (b<sub>6</sub>), this case is impossible to appear.

Step 3.6. Consider that  $q_{22} = q_{44} = q_{55} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 3\}$  and  $j \in \{3, 6\}$ . Let  $q := q_{11}$ ,  $r := q_{33}$ ,  $s^{-1} := q'_{45}$  and  $t := q_{66}$ . Since  $X$  has a good  $A_6$  neighborhood, we have either  $a = 0$  or  $a = 1$ . In addition, we have  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$  and  $rs \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . Next we discuss in two steps.

Step 3.6.1. We discuss the subcase  $a = 0$ . That is


$$A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}. \text{ Then } st = 1 \text{ and the reflection of } X$$

$$X : \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{s^{-1}}{\circ} \overset{-1}{\bullet} \overset{s}{\circ} \overset{s^{-1}}{\circ} \Rightarrow r_5(X) : \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{-1}{\circ} \overset{s^{-1}}{\circ} \overset{-1}{\circ}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . By Lemma 4.5 ( $a_3$ ), this subcase is impossible to appear.

Step 3.6.2. We discuss the subcase  $a = 1$ . That is

$$A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & -1 & 2 \end{pmatrix}. \text{ Then } st \neq 1. \text{ The reflection of } X$$

$$X : \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{s^{-1}}{\circ} \overset{-1}{\bullet} \overset{t^{-1}}{\circ} \overset{t}{\circ} \Rightarrow r_5(X) : \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{(st)^{-1}}{\circ} \overset{-1}{\circ} \overset{-1}{\circ}$$


implies  $s = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . Then the reflection of  $X$

$$X : \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{-1}{\bullet} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{t^{-1}}{\circ} \overset{t}{\circ} \Rightarrow r_2(X) : \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{r}{\circ} \overset{-1}{\bullet} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{t^{-1}}{\circ} \overset{t}{\circ}$$

$$\Rightarrow r_3 r_2(X) : \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{-1}{\circ} \overset{t^{-1}}{\circ} \overset{t}{\circ}$$

imply  $r = -1$  and  $p \neq 2$  by  $a_{45}^{r_3 r_2(X)} = -1$ , which contradicts to  $rs \neq 1$ .

Further by Lemma 4.4 the symmetric case is impossible to appear.

Step 3.7. Consider that  $q_{22} = q_{44} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 3, 5\}$  and  $j \in \{3, 5\}$ . Let  $q := q_{11}$ ,  $r := q_{33}$  and  $s := q_{55}$ . Hence  $a = 0$ . Since  $X$  has a good  $A_6$  neighborhood, we get  $qr = 1$  and  $rs \neq 1$ . Then the reflections of  $X$

$$X : \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ} \overset{-1}{\bullet} \Rightarrow r_6(X) : \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{s^{-1}}{\circ} \overset{-1}{\circ} \overset{s}{\circ} \overset{-1}{\circ}$$

$$\Rightarrow r_5 r_6(X) : \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{-1}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ}$$

imply  $s = -1$  and  $p \neq 2$  by  $a_{43}^{r_5 r_6(X)} = -1$ . By Lemma 4.5 ( $a_3$ ), this case is impossible to appear. Further by Lemma 4.4 the symmetric case is impossible to appear.

Step 3.8. Consider that  $q_{44} = q_{55} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 3\}$  and  $j \in \{2, 3\}$ . Let  $q := q_{11}$ ,  $r^{-1} := q'_{45}$ ,  $s^{-1} := q'_{56}$ . Since  $X$  has a good  $A_6$  neighborhood, we have either  $a = 0$  or  $a = 1$ . In addition, we have  $qr \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . Next we discuss in two steps.

Step 3.8.1. We discuss the subcase  $a = 0$ . That is

$$A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}. \text{ Then } rs = 1 \text{ and the reflection of } X$$

$$X : \begin{array}{ccccccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{ccccccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & s & s^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . Hence  $r = s = -1$ . By Lemma 4.5 ( $a_1$ ), this subcase is impossible to appear.

Step 3.8.2. We discuss the subcase  $a = 1$ . That is

$$A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & -1 & 2 \end{pmatrix}. \text{ Then } rs \neq 1 \text{ and the reflection of } X$$

$$X : \begin{array}{ccccccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{ccccccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & s & s^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & -1 & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ \end{array} \Rightarrow r_6(X) : \begin{array}{ccccccccccc} q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & -1 & s^{-1} & s & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{54}^{r_6(X)} = -1$ , which contradicts to  $rs \neq 1$ .

Further by Lemma 4.4 the symmetric case is impossible to appear.

Step 4. Assume that there are four vertices labeled by  $-1$ . By  $q_{44} = -1$ , we distinguish the following eight steps.

Step 4.1. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = -1$ ,  $q_{55}q'_{56} - 1 = 0$  and  $q_{jj}q'_{j-1,j} - 1 = 0$  for  $j \in \{5, 6\}$ . Let  $q^{-1} := q'_{12}$ ,  $r^{-1} := q'_{23}$ ,  $s^{-1} := q'_{34}$  and  $t := q_{55}$ . Hence  $a = 0$ . Since  $X$  has a good  $A_6$  neighborhood, we get  $st \neq 1$  by  $a_{35}^{r_4(X)} = -1$ ,  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$  and  $rs = 1$  by  $a_{24}^{r_3(X)} = 0$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccccccc} -1 & s^{-1} & -1 & s & -1 & s^{-1} & -1 & t^{-1} & t & t^{-1} & t \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccccccc} -1 & s^{-1} & s & s^{-1} & -1 & s & s^{-1} & t^{-1} & t & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.5 (b<sub>2</sub>), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{33} = q_{44} = q_{55} = q_{66} = -1$ ,  $q_{22}q'_{21} - 1 = 0$  and  $q_{ii}q'_{i,i+1} - 1 = 0$  for  $i \in \{1, 2\}$ . Let  $q := q_{11}$ ,  $r^{-1} := q'_{34}$ ,  $s^{-1} := q'_{45}$  and  $t^{-1} := q'_{56}$ . Since  $X$  has a good  $A_6$  neighborhood, then  $qr = 1$  and  $rs \neq 1$ . The reflection of  $X$

$$X : \begin{array}{cccccccc} r^{-1} & r & r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{cccccccc} r^{-1} & r & -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . Then the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & s^{-1} & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ \end{array} \Rightarrow r_6(X) : \begin{array}{cccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & s^{-1} & t^{-1} & t & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $st = 1$  or  $t = -1$  by  $a_{54}^{r_6(X)} = -1$ . If  $st = 1$ , this case is impossible to appear by Lemma 4.5 (b<sub>5</sub>). If  $t = -1$ , then  $s \neq -1$  and the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & s^{-1} & -1 & -1 & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & s^{-1} & -1 & -1 & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s^2 = -1$  by  $a_{53}^{r_4(X)} = -1$ . Hence  $s \in G'_4$ . By Lemma 4.5 (b<sub>1</sub>), this case is impossible to appear.

Step 4.2. Consider that  $q_{11} = q_{22} = q_{44} = q_{55} = -1$ ,  $q_{33}q'_{34} - 1 = 0$  and  $q_{jj}q'_{j-1,j} - 1 = 0$  for  $j \in \{3, 6\}$ . Let  $q^{-1} := q'_{12}$ ,  $r := q_{33}$ ,  $s^{-1} := q'_{45}$  and  $t := q_{66}$ . Since  $X$  has a good  $A_6$  neighborhood, we have either  $a = 0$  or  $a = 1$ . In addition, we have  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$  and  $rs \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . Next we discuss in two steps.

Step 4.2.1. We discuss the subcase  $a = 0$ . That is

$$A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}. \text{ Then } st = 1. \text{ The reflection of } X$$

$$X : \begin{array}{cccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & -1 & s & s^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{cccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & s^{-1} & s & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . By Lemma 4.5 (a<sub>4</sub>), this subcase is impossible to appear.

Step 4.2.2. We discuss the subcase  $a = 1$ . That is

$$A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & -1 & 2 \end{pmatrix}. \text{ Then } st \neq 1. \text{ Similar to the proof in Step 3.6.2,}$$

this subcase is impossible to appear by  $a_{43}^{r_5(X)} = -1$  and  $a_{45}^{r_3r_2(X)} = -1$ .



Further consider the symmetric case of this case that  $q_{22} = q_{33} = q_{55} = q_{66} = -1$ ,  $q_{44}q'_{43} - 1 = 0$  and  $q_{ii}q'_{i,i+1} - 1 = 0$  for  $i \in \{1, 4\}$ . By Lemma 4.4, this case is impossible to appear.

Step 4.3. Consider that  $q_{11} = q_{22} = q_{44} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{3, 5\}$  and  $j \in \{3, 5\}$ . Let  $q^{-1} := q'_{12}$ ,  $r := q_{33}$  and  $s := q_{55}$ . Hence  $a = 0$ . Since  $X$  has a good  $A_6$  neighborhood, we get  $qr = 1$  and  $rs \neq 1$ . The reflections of  $X$

$$\begin{aligned}
X : \begin{array}{cccccccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & s & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \bullet \end{array} & \Rightarrow & r_6(X) : \begin{array}{cccccccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & -1 & s & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ \end{array} \\
& \Rightarrow & r_5 r_6(X) : \begin{array}{cccccccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & s^{-1} & s & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}
\end{aligned}$$

imply  $s = -1$  and  $p \neq 2$  by  $a_{43}^{r_5 r_6(X)} = -1$ . By Lemma 4.5 ( $a_4$ ), this case is impossible to appear. Further by Lemma 4.4 the symmetric case is impossible to appear.

Step 4.4. Consider that  $q_{11} = q_{33} = q_{44} = q_{55} = -1$ ,  $q_{22}q'_{23} - 1 = 0$  and  $q_{jj}q'_{j-1,j} - 1 = 0$  for  $j \in \{2, 6\}$ . Let  $q := q_{22}$ ,  $r^{-1} := q'_{34}$ ,  $s^{-1} := q'_{45}$  and  $t := q_{66}$ . Since  $X$  has a good  $A_6$  neighborhood, we get  $qr = 1$  and  $rs \neq 1$ . The reflection of  $X$

$$\begin{aligned}
X : \begin{array}{cccccccccccc} -1 & r & r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & -1 & t^{-1} & t \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} & \Rightarrow & r_3(X) : \begin{array}{cccccccccccc} -1 & r & -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & -1 & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}
\end{aligned}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . Hence  $s \neq -1$  and the reflection of  $X$

$$\begin{aligned}
X : \begin{array}{cccccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & s^{-1} & -1 & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} & \Rightarrow & r_4(X) : \begin{array}{cccccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & s^{-1} & -1 & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}
\end{aligned}$$

implies  $st = 1$  by  $a_{56}^{r_4(X)} = -1$ . Then this case is impossible to appear by Lemma 4.5 ( $b_6$ ). Further consider the symmetric case of this case that  $q_{22} = q_{33} = q_{44} = q_{66} = -1$ ,  $q_{55}q'_{45} - 1 = 0$  and  $q_{ii}q'_{i,i+1} - 1 = 0$  for  $i \in \{1, 5\}$ . Let  $q := q_{11}$ ,  $r^{-1} := q'_{23}$ ,  $s^{-1} := q'_{34}$  and  $t := q_{55}$ . Hence  $a = 0$ . Since  $X$  has a good  $A_6$  neighborhood, then  $qr = 1$ ,  $rs = 1$  and  $st \neq 1$ . The reflections of  $X$

$$\begin{aligned}
X : \begin{array}{cccccccccccc} s & s^{-1} & -1 & s & -1 & s^{-1} & -1 & t^{-1} & t & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \bullet \end{array} & \Rightarrow & r_6(X) : \begin{array}{cccccccccccc} s & s^{-1} & -1 & s & -1 & s^{-1} & -1 & t^{-1} & -1 & t & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ \end{array} \\
& \Rightarrow & r_5 r_6(X) : \begin{array}{cccccccccccc} s & s^{-1} & -1 & s & -1 & s^{-1} & t^{-1} & t & -1 & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}
\end{aligned}$$

imply  $t = -1$  and  $p \neq 2$  by  $a_{43}^{r_5 r_6(X)} = -1$ . By Lemma 4.5 ( $a_5$ ), this case is impossible to appear.

Step 4.5. Consider that  $q_{11} = q_{33} = q_{44} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 5\}$  and  $j \in \{2, 5\}$ . Let  $q := q_{22}$ ,  $r^{-1} := q'_{34}$  and  $s := q_{55}$ . Hence  $a = 0$ . Since  $X$  has a good  $A_6$  neighborhood, we get  $rs \neq 1$  and  $qr = 1$ . Then the reflection of  $X$

$$\begin{aligned}
X : \begin{array}{cccccccccccc} -1 & r & r^{-1} & r & -1 & r^{-1} & -1 & s^{-1} & s & s^{-1} & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} & \Rightarrow & r_3(X) : \begin{array}{cccccccccccc} -1 & r & -1 & r^{-1} & -1 & r & r^{-1} & s^{-1} & s & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}
\end{aligned}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.5 ( $b_4$ ), this case is impossible to appear.

Step 4.6. Consider that  $q_{11} = q_{44} = q_{55} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 3\}$  and  $j \in \{2, 3\}$ . Let  $q := q_{22}$ ,  $r^{-1} := q'_{45}$  and  $s^{-1} := q'_{56}$ . Since  $X$  has a good  $A_6$  neighborhood, we have either  $a = 0$  or  $a = 1$ . In addition, we have  $qr \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . Next we discuss in two steps.

Step 4.6.1. We discuss the subcase  $a = 0$ . That is

$$A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}. \text{ Then } rs = 1. \text{ The reflection of } X$$

$$X : \begin{array}{cccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & r & -1 \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{cccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & r^{-1} & r \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . By Lemma 4.5 ( $a_2$ ), this subcase is impossible to appear.

Step 4.6.2. We discuss the subcase  $a = 1$ . That is

$$A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & -1 & 2 \end{pmatrix}. \text{ Then } rs \neq 1. \text{ The reflection of } X$$

$$X : \begin{array}{cccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & r^{-1} & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{cccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & r^{-1} & r & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & -1 & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_6(X) : \begin{array}{cccccccc} -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & -1 & s^{-1} & s & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{54}^{r_6(X)} = -1$ , which contradicts to  $rs \neq 1$ .

Further by Lemma 4.4 the symmetric case is impossible to appear.

Step 4.7. Consider that  $q_{22} = q_{33} = q_{44} = q_{55} = -1$ ,  $q_{11}q'_{12} - 1 = 0$  and  $q_{66}q'_{56} - 1 = 0$ . Let  $q := q_{11}$ ,  $r^{-1} := q'_{23}$ ,  $s^{-1} := q'_{34}$ ,  $t^{-1} := q'_{45}$  and  $u := q_{66}$ . Since  $X$  has a good  $A_6$  neighborhood, we have either  $a = 0$  or  $a = 1$ . In addition, we have  $rs = 1$ ,  $qr = 1$  and  $st \neq 1$ . Then  $qt \neq 1$ . Next we can discuss in two steps.

Step 4.7.1. We discuss the subcase  $a = 0$ . That is

$$A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}. \text{ Then } tu = 1. \text{ The reflection of } X$$

$$X : \begin{array}{ccccccccc} q & q^{-1} & -1 & q & -1 & q^{-1} & -1 & t^{-1} & -1 & t & t^{-1} \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{ccccccccc} q & q^{-1} & -1 & q & -1 & q^{-1} & t^{-1} & t & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . Thus this subcase is impossible to appear by Lemma 4.5 ( $a_5$ ).

Step 4.7.2. We discuss the subcase  $a = 1$ . That is

$$A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & -1 & 2 \end{pmatrix}. \text{ Then } tu \neq 1. \text{ The reflection of } X$$

$$X : \begin{array}{ccccccccc} q & q^{-1} & -1 & q & -1 & q^{-1} & -1 & t^{-1} & -1 & u^{-1} & u \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{ccccccccc} q & q^{-1} & -1 & q & -1 & q^{-1} & t^{-1} & t & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccccc} q & q^{-1} & -1 & q & -1 & q^{-1} & -1 & -1 & u^{-1} & u \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccccc} q & q^{-1} & q & q^{-1} & -1 & q & q^{-1} & -1 & -1 & u^{-1} & u \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $q = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ , which contradicts to  $qt \neq 1$ .

Step 4.8. Consider that  $q_{22} = q_{44} = q_{55} = q_{66} = -1$ ,  $q_{33}q'_{32} - 1 = 0$  and  $q_{ii}q'_{i,i+1} - 1 = 0$  for  $i \in \{1, 3\}$ . Let  $q := q_{11}$ ,  $r := q_{33}$ ,  $s^{-1} := q'_{45}$  and  $t^{-1} := q'_{56}$ . Since  $X$  has a good  $A_6$  neighborhood, we have either  $a = 0$  or  $a = 1$ . In addition, we have  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$  and  $rs \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . The further discussion is similar to Step 4.6. Thus this case is impossible to appear. Further by Lemma 4.4 the symmetric case is impossible to appear.

Step 5. Assume that there are five vertices labeled by  $-1$ . By  $q_{44} = -1$ , we distinguish the following three steps.

Step 5.1. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = q_{55} = -1$  and  $q_{66}q'_{56} - 1 = 0$ . Let  $q^{-1} := q'_{12}$ ,  $r^{-1} := q'_{23}$ ,  $s^{-1} := q'_{34}$ ,  $t^{-1} := q'_{45}$  and  $u := q_{66}$ . Since  $X$  has a good  $A_6$  neighborhood, we get  $rs = 1$  by  $a_{24}^{r_3(X)} = 0$ ,  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$  and  $st \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . Then the reflection of  $X$

$$X : \begin{array}{ccccccccc} -1 & s^{-1} & -1 & s & -1 & s^{-1} & -1 & t^{-1} & -1 & u^{-1} & u \\ \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{ccccccccc} -1 & s^{-1} & s & s^{-1} & -1 & s & s^{-1} & t^{-1} & -1 & u^{-1} & u \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . Hence  $t \neq -1$  and the reflection of  $X$

$$X : \begin{array}{ccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & t^{-1} & -1 & u^{-1} & u \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{ccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & t^{-1} & -1 & u^{-1} & u \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $tu = 1$  by  $a_{56}^{r_4(X)} = -1$ . By Lemma 4.5 (b<sub>6</sub>), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{22} = q_{33} = q_{44} = q_{55} = q_{66} = -1$  and  $q_{11}q'_{12} - 1 = 0$ . Let  $q := q_{11}$ ,  $r^{-1} := q'_{23}$ ,  $s^{-1} := q'_{34}$ ,  $t^{-1} := q'_{45}$  and  $u^{-1} := q'_{56}$ . Since  $X$  has a good  $A_6$  neighborhood, then  $rs = 1$ ,  $qr = 1$  and  $st \neq 1$ . Similar to the proof above, we get  $q = -1$ ,  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$  and  $tu = 1$  by  $a_{56}^{r_4(X)} = -1$ . Rewrite  $X$  as

$$\begin{array}{cccccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & t^{-1} & -1 & t & -1 \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array}$$

By Lemma 4.5 (b<sub>5</sub>), this case is also impossible to appear.

Step 5.2. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = q_{66} = -1$  and  $q_{55}q'_{56} - 1 = q_{55}q'_{45} - 1 = 0$ . Let  $q^{-1} := q'_{12}$ ,  $r^{-1} := q'_{23}$ ,  $s^{-1} := q'_{34}$  and  $t := q_{55}$ . Hence  $a = 0$ . Since  $X$  has a good  $A_6$  neighborhood, we get  $qr = 1$ ,  $rs = 1$  and  $st \neq 1$ . The reflection of  $X$

$$X : \begin{array}{cccccccccccc} -1 & q^{-1} & -1 & q & -1 & q^{-1} & -1 & t^{-1} & t & t^{-1} & -1 \\ \circ & \text{---} & \circ & \text{---} & \bullet & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{cccccccccccc} -1 & q^{-1} & q & q^{-1} & -1 & q & q^{-1} & t^{-1} & t & t^{-1} & -1 \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array}$$

implies  $q = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . By Lemma 4.5 (b<sub>4</sub>), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{11} = q_{33} = q_{44} = q_{55} = q_{66} = -1$ , and  $q_{22}q'_{23} - 1 = q_{22}q'_{12} - 1 = 0$ . Let  $q := q_{22}$ ,  $r^{-1} := q'_{34}$ ,  $s^{-1} := q'_{45}$  and  $t^{-1} := q'_{56}$ . Since  $X$  has a good  $A_6$  neighborhood, then  $qr = 1$  and  $rs \neq 1$ . Similar to the proof in Step 5.1, we get  $q = -1$ ,  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$  and  $st = 1$  by  $a_{56}^{r_4(X)} = -1$ . Rewrite  $X$  as

$$\begin{array}{cccccccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & s^{-1} & -1 & s & -1 \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array}$$

and this case is impossible to appear by Lemma 4.5 (b<sub>5</sub>).

Step 5.3. Consider that  $q_{11} = q_{22} = q_{44} = q_{55} = q_{66} = -1$ , and  $q_{33}q'_{34} - 1 = q_{33}q'_{23} - 1 = 0$ . Let  $q^{-1} := q'_{12}$ ,  $r := q_{33}$ ,  $s^{-1} := q'_{45}$  and  $t^{-1} := q'_{56}$ . Since  $X$  has a good  $A_6$  neighborhood, we have either  $a = 0$  or  $a = 1$ . In addition, we have  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$  and  $rs \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . Next we discuss in two steps.

Step 5.3.1. We discuss the subcase  $a = 0$ . That is

$$A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}. \text{ Then } st = 1. \text{ The reflection of } X$$

$$X : \begin{array}{cccccccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & -1 & s & -1 \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \bullet & \text{---} & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{cccccccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & s^{-1} & s & -1 & s^{-1} & s \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . By Lemma 4.5 (a<sub>4</sub>), this subcase is impossible to appear.

Step 5.3.2. We discuss the subcase  $a = 1$ . That is

$$A^{r_5(X)} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & -1 & 2 \end{pmatrix}. \text{ Then } st \neq 1. \text{ The reflection of } X$$

$$X : \begin{array}{cccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{cccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & s^{-1} & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{43}^{r_5(X)} = -1$ . Then the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & -1 & -1 & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_6(X) : \begin{array}{cccccccc} -1 & r & -1 & r^{-1} & r & r^{-1} & -1 & -1 & t^{-1} & t & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{54}^{r_6(X)} = -1$ , which contradicts to  $st \neq 1$ .

Further by Lemma 4.4 the symmetric case is impossible to appear.

Step 6. Assume that all vertices are labeled by  $-1$ . Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = q_{55} = q_{66} = -1$ . Let  $q^{-1} := q'_{12}$ ,  $r^{-1} := q'_{23}$ ,  $s^{-1} := q'_{34}$ ,  $t^{-1} := q'_{45}$  and  $u^{-1} := q'_{56}$ . Since  $X$  has a good  $A_6$  neighborhood, we get  $qr = 1$  by  $a_{13}^{r_2(X)} = 0$ ,  $rs = 1$  by  $a_{24}^{r_3(X)} = 0$  and  $st \neq 1$  by  $a_{35}^{r_4(X)} = -1$ . The reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & s^{-1} & -1 & t^{-1} & -1 & u^{-1} & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_3(X) : \begin{array}{cccccccc} -1 & s^{-1} & s & s^{-1} & -1 & s & s^{-1} & t^{-1} & -1 & u^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{45}^{r_3(X)} = -1$ . Hence  $t \neq -1$  and the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & t^{-1} & -1 & u^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & -1 & -1 & -1 & -1 & -1 & -1 & t^{-1} & -1 & u^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $tu = 1$  by  $a_{56}^{r_4(X)} = -1$ . By Lemma 4.5 ( $b_5$ ), this case is impossible to appear.  $\square$

### 4.3 Classification theorem for rank 7

**Lemma 4.7.** *If  $\mathcal{D}_{\chi,E}$  is of the form*

$$\begin{array}{cccccccccccc} q_{11} & r & q_{22} & s & q_{33} & t & q_{44} & u & q_{55} & v & q_{66} & w & q_{77} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

and has a good  $A_7$  neighborhood, then  $q_{33} = -1$ .

*Proof.* By the definition of the good  $A_7$  neighborhood, the claim is clear.  $\square$

**Lemma 4.8.** *If  $\mathcal{D}_{\chi,E}$  is of the form*

$$\begin{array}{cccccccccccc} q_{11} & r & q_{22} & s & q_{33} & t & q_{44} & u & q_{55} & v & q_{66} & w & q_{77} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

and has a good  $A_7$  neighborhood, then the following are impossible to appear.

- (a<sub>1</sub>)  $\begin{array}{c} -1 \ q^{-1} \ -1 \ q \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}$  ,  $q \neq -1, p \neq 2$
- (a<sub>2</sub>)  $\begin{array}{c} q \ q^{-1} \ -1 \ q \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}$  ,  $q \neq -1, p \neq 2$
- (b<sub>1</sub>)  $\begin{array}{c} -1 \ q^{-1} \ q \ q^{-1} \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}$  ,  $q \neq -1, p \neq 2$
- (b<sub>2</sub>)  $\begin{array}{c} q \ q^{-1} \ q \ q^{-1} \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}$  ,  $q \neq -1, p \neq 2$
- (c<sub>1</sub>)  $\begin{array}{c} -1 \ -1 \ -1 \ -1 \ -1 \ q^{-1} \ q \ q^{-1} \ q \ q^{-1} \ q \ q^{-1} \ q \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}$  ,  $q \neq -1, p \neq 2$
- (c<sub>2</sub>)  $\begin{array}{c} -1 \ -1 \ -1 \ -1 \ -1 \ q^{-1} \ q \ q^{-1} \ q \ q^{-1} \ q \ q^{-1} \ -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}$  ,  $q \neq -1, p \neq 2$
- (c<sub>3</sub>)  $\begin{array}{c} -1 \ -1 \ -1 \ -1 \ -1 \ q^{-1} \ q \ q^{-1} \ q \ q^{-1} \ -1 \ q \ q^{-1} \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}$  ,  $q \neq -1, p \neq 2$
- (c<sub>4</sub>)  $\begin{array}{c} -1 \ -1 \ -1 \ -1 \ -1 \ q^{-1} \ q \ q^{-1} \ q \ q^{-1} \ -1 \ q \ -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}$  ,  $q \neq -1, p \neq 2$
- (c<sub>5</sub>)  $\begin{array}{c} -1 \ -1 \ -1 \ -1 \ -1 \ q^{-1} \ q \ q^{-1} \ -1 \ q \ -1 \ q^{-1} \ -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}$  ,  $q \neq -1, p \neq 2$
- (c<sub>6</sub>)  $\begin{array}{c} -1 \ -1 \ -1 \ -1 \ -1 \ q^{-1} \ q \ q^{-1} \ -1 \ q \ -1 \ q^{-1} \ q \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}$  ,  $q \neq -1, p \neq 2$
- (c<sub>7</sub>)  $\begin{array}{c} -1 \ -1 \ -1 \ -1 \ -1 \ q^{-1} \ q \ q^{-1} \ -1 \ q \ q^{-1} \ q \ -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}$  ,  $q \neq -1, p \neq 2$
- (c<sub>8</sub>)  $\begin{array}{c} -1 \ -1 \ -1 \ -1 \ -1 \ q^{-1} \ q \ q^{-1} \ -1 \ q \ q^{-1} \ q \ q^{-1} \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}$  ,  $q \neq -1, p \neq 2$

*Proof.* Consider the following cases.

- (a<sub>1</sub>) Assume that  $X$  is of the type  $\begin{array}{c} q_{11} \ q^{-1} \ -1 \ q \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}$  where  $q_{11} = -1$  or  $q_{11} = q$ . Since  $X$  has a good  $A_7$  neighborhood, the transformation

$$\begin{array}{c} q_{11} \ q^{-1} \ -1 \ q \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \\ \circ \text{---} \bullet \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array} \Rightarrow \begin{array}{c} q'_{11} \ q \ -1 \ q^{-1} \ q \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}$$

shows  $q = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ , where



$$q'_{11} = \begin{cases} q^{-1} & \text{if } q_{11} = -1 \\ -1 & \text{if } q_{11} = q \end{cases}$$

which contradicts to  $q \neq -1$ . Hence the case in (a<sub>2</sub>) is also impossible to appear.

- (b<sub>1</sub>) Assume that  $X$  is of the type  $\begin{array}{c} q^{11} \ q^{-1} \ q \ q^{-1} \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \end{array}$  where  $q_{11} = -1$  or  $q_{11} = q$ . We can discuss in two subcases  $q^2 = -1$  and  $q^2 \neq -1$ .



(c<sub>12</sub>) If  $q^{-2} \neq -1$ , then the transformations



show  $q = 1$  by  $a_{21}^{r_4 r_3(x)} = -1$ , which gives a contradiction.



The reflections and conditions of the cases (c<sub>2</sub>), (c<sub>3</sub>), (c<sub>4</sub>), (c<sub>5</sub>), (c<sub>6</sub>), (c<sub>7</sub>) and (c<sub>8</sub>) happen similarly to the case (c<sub>1</sub>). Hence the cases (c<sub>2</sub>) – (c<sub>8</sub>) can be proved by the same argument as in (c<sub>1</sub>).

□

**Theorem 4.9.** *All the possible Dynkin diagrams of  $V$  such that  $\mathcal{B}(V)$  has a finite root system listed in Appendix C appear in row 21 of Table 1.*

*Proof.* Let  $X = [M]_7^s$ ,  $I = \{1, 2, 3, 4, 5, 6, 7\}$ , and  $A^X := (a_{ij})_{i,j \in I}$  be the Cartan matrix of  $X$ . Let  $(q_{i,j})_{i,j \in I}$  be the braiding matrix of  $X$  and  $(q_{i,j}^{r_i(X)})_{i,j \in I}$  be the braiding matrix of  $r_i(X)$ . To simplify the labels, we write  $q'_{ij} := q_{ij}q_{ji}$  for  $1 \leq i, j \leq 7$ . By Theorem 3.4,  $A^X$  has a good  $A_7$  neighborhood. By  $A^X = A_7$ , we get  $(2)_{q_{ii}}(q_{ii}q'_{i,i+1} - 1) = (2)_{q_{jj}}(q_{jj}q'_{j-1,j} - 1) = 0$ , for all  $i \in \{1, 2, 3, 4, 5, 6\}$  and  $j \in \{2, 3, 4, 5, 6, 7\}$ . In addition, we obtain  $q_{33} = -1$  for all cases by Lemma 4.7. Depending on the number of vertices labeled by  $-1$ , we divide the cases into the following categories: Step 1 to Step 7.

Step 1. Assume that there is one vertex labeled by  $-1$ . That is  $q_{33} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 4, 5, 6\}$  and  $j \in \{2, 4, 5, 6, 7\}$ . Let  $r := q_{11}$  and  $q := q_{44}$ . Since  $X$  has a good  $A_7$  neighborhood, we get  $rq \neq 1$  by  $a_{24}^{r_3(X)} = -1$ . The reflections of  $X$



imply  $rq^2 = 1$  or  $rq = -1$  by  $a_{45}^{r_2 r_3(X)} = -1$ . We discuss in two steps.



Step 1.1. If  $rq^2 = 1$  then the reflection of  $X$

$$r_3(X) : \begin{array}{c} \circ \\ \swarrow q^{-2} \\ \circ \\ \swarrow q^{-2} \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \bullet \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \end{array} \Rightarrow r_4r_3(X) : \begin{array}{c} \circ \\ \swarrow q^{-2} \\ \circ \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \end{array}$$

implies that  $q = -1$ ,  $p \neq 2$  or  $q \in G'_3$ ,  $p \neq 3$  by  $a_{21}^{r_4r_3(X)} = -1$ . Since  $q = -1$  yields  $r = 1$ , it gives a contradiction. Rewrite  $X$  as

$$\begin{array}{c} q \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \end{array}$$

where  $q \in G'_3$ ,  $p \neq 3$ . Hence  $\mathcal{D} = \tau_{7654321}\mathcal{D}_{21,1}^7$  and  $p \neq 3$ .

Step 1.2. Assume that  $rq = -1$ . Then the reflection of  $X$

$$r_3(X) : \begin{array}{c} \circ \\ \swarrow -q^{-1} \\ \circ \\ \swarrow -q \\ \circ \\ \swarrow -1 \\ \circ \\ \swarrow -1 \\ \circ \\ \swarrow q \\ \bullet \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \end{array} \Rightarrow r_4r_3(X) : \begin{array}{c} \circ \\ \swarrow -q^{-1} \\ \circ \\ \swarrow -q \\ \circ \\ \swarrow -1 \\ \circ \\ \swarrow -1 \\ \circ \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \end{array}$$

implies  $q = -1$  and  $p \neq 2$  by  $a_{25}^{r_4r_3(X)} = 0$ . Rewrite  $X$  as

$$\begin{array}{c} r \\ \swarrow r^{-1} \\ \circ \\ \swarrow r \\ \circ \\ \swarrow r^{-1} \\ \circ \\ \swarrow -1 \\ \circ \\ \swarrow -1 \\ \circ \\ \swarrow -1 \\ \circ \\ \swarrow -1 \\ \circ \\ \swarrow -1 \\ \circ \\ \swarrow -1 \\ \circ \\ \swarrow -1 \\ \circ \end{array}$$

and this subcase is impossible to appear by Lemma 4.8 ( $b_2$ ).

Step 2. Assume that there are two vertices labeled by  $-1$ . By  $q_{33} = -1$ , we distinguish the following six steps.

Step 2.1. Consider that  $q_{11} = q_{33} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 4, 5, 6\}$  and  $j \in \{2, 4, 5, 6, 7\}$ . Let  $r := q_{22}$  and  $q := q_{44}$ . Since  $X$  has a good  $A_7$  neighborhood, we get  $rq \neq 1$  by  $a_{24}^{r_3(X)} = -1$ . The reflections of  $X$

$$X : \begin{array}{c} -1 \\ \swarrow r^{-1} \\ \bullet \\ \swarrow r \\ \circ \\ \swarrow r^{-1} \\ \circ \\ \swarrow -1 \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \end{array} \Rightarrow r_1(X) : \begin{array}{c} -1 \\ \swarrow r \\ \bullet \\ \swarrow r^{-1} \\ \circ \\ \swarrow -1 \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \end{array}$$

$$\Rightarrow r_2r_1(X) : \begin{array}{c} r \\ \swarrow r^{-1} \\ \circ \\ \swarrow r \\ \circ \\ \swarrow r^{-1} \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \\ \swarrow q^{-1} \\ \circ \\ \swarrow q \\ \circ \end{array}$$

imply that  $r = -1$  and  $p \neq 2$  by  $a_{34}^{r_2r_1(X)} = -1$ . By Lemma 4.8 ( $c_1$ ), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{55} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 3, 4, 6\}$  and  $j \in \{2, 3, 4, 6\}$ . By Lemma 4.7, this case is impossible to appear.

Step 2.2. Consider that  $q_{22} = q_{33} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 4, 5, 6\}$  and  $j \in \{4, 5, 6, 7\}$ . Let  $r := q_{11}$ ,  $q := q'_{23}$  and  $s := q_{44}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $r^{-1}q = 1$  and  $qs^{-1} \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{c} q \\ \swarrow q^{-1} \\ \bullet \\ \swarrow -1 \\ \circ \\ \swarrow s^{-1} \\ \circ \\ \swarrow s \\ \circ \\ \swarrow s^{-1} \\ \circ \\ \swarrow s \\ \circ \\ \swarrow s^{-1} \\ \circ \\ \swarrow s \\ \circ \end{array} \Rightarrow r_2(X) : \begin{array}{c} -1 \\ \swarrow q \\ \bullet \\ \swarrow q^{-1} \\ \circ \\ \swarrow s^{-1} \\ \circ \\ \swarrow s \\ \circ \\ \swarrow s^{-1} \\ \circ \\ \swarrow s \\ \circ \\ \swarrow s^{-1} \\ \circ \\ \swarrow s \\ \circ \end{array}$$

implies  $q = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.8 ( $c_1$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 2.3. Consider that  $q_{33} = q_{44} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 5, 6\}$  and  $j \in \{2, 5, 6, 7\}$ . Let  $r := q_{11}$ ,  $q := q'_{34}$  and  $s := q_{55}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $r^{-1}q \neq 1$  and  $qs^{-1} = 1$ . Then the reflection of  $X$

$$X : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & -1 & q & -1 & q^{-1} & q & q^{-1} & q \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & q & q^{-1} & -1 & q & -1 & q^{-1} & q \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $q = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 ( $b_2$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 2.4. Consider that  $q_{33} = q_{55} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 4, 6\}$  and  $j \in \{2, 4, 6, 7\}$ . Let  $r := q_{11}$ ,  $q := q_{44}$  and  $s := q_{66}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rq \neq 1$  and  $qs = 1$ . Then the reflections of  $X$

$$X : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & -1 & q^{-1} & q & q^{-1} & -1 & q & q^{-1} & q \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & -1 & q^{-1} & -1 & q & -1 & q^{-1} & -1 & q & q^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

$$\Rightarrow r_4r_5(X) : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & q^{-1} & q & -1 & q^{-1} & q & q^{-1} & -1 & q & q^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

imply  $q = -1$  and  $p \neq 2$  by  $a_{32}^{r_4r_5(X)} = -1$ . By Lemma 4.8 ( $b_2$ ), this case is impossible to appear.

Step 2.5. Consider that  $q_{33} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 4, 5\}$  and  $j \in \{2, 4, 5, 7\}$ . Let  $r := q_{11}$ ,  $q := q_{44}$  and  $s := q_{77}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rq \neq 1$  and  $qs = 1$ . The reflections of  $X$

$$X : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & q & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_6(X) : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & -1 & q^{-1} & q & q^{-1} & -1 & q & -1 & q^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ \end{array}$$

$$\Rightarrow r_5r_6(X) : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & -1 & q^{-1} & -1 & q & -1 & q^{-1} & q & q^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ \end{array}$$

$$\Rightarrow r_4r_5r_6(X) : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & q^{-1} & q & -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

imply  $q = -1$  and  $p \neq 2$  by  $a_{32}^{r_4r_5r_6(X)} = -1$ . By Lemma 4.8 ( $b_2$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 2.6. Consider that  $q_{33} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 4, 5, 6\}$  and  $j \in \{2, 4, 5, 6\}$ . Let  $r := q_{11}$  and  $q := q_{44}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rq \neq 1$  by  $a_{24}^{r_3(X)} = -1$ . Similar to the proof in Step 2.5, we have  $q = -1$  and  $p \neq 2$  by  $a_{32}^{r_4r_5r_6r_7(X)} = -1$ . Rewrite  $X$  as

$$\begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ \end{array}$$

and this case is impossible to appear by Lemma 4.8 ( $b_2$ ). Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 3. Assume that there are three vertices labeled by  $-1$ . By  $q_{33} = -1$ , we distinguish the following thirteen steps.

Step 3.1. Consider that  $q_{11} = q_{22} = q_{33} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{4, 5, 6\}$  and  $j \in \{4, 5, 6, 7\}$ . Let  $s := q'_{12}$ ,  $r := q'_{23}$  and  $q := q_{44}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rq^{-1} \neq 1$  by  $a_{24}^{r_3(X)} = -1$  and  $sr = 1$  by  $a_{13}^{r_2(X)} = 0$ . The reflection of  $X$

$$X : \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{-1}{\bullet} \overset{r}{\circ} \overset{-1}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \Rightarrow r_2(X) : \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ} \overset{q^{-1}}{\circ} \overset{q}{\circ}$$

implies  $r = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.8 ( $c_1$ ), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{55} = q_{66} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 3, 4\}$  and  $j \in \{2, 3, 4\}$ . By Lemma 4.7, this case is impossible to appear.

Step 3.2. Consider that  $q_{11} = q_{33} = q_{44} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 5, 6\}$  and  $j \in \{2, 5, 6, 7\}$ . Let  $r := q_{22}$ ,  $s := q'_{34}$ , and  $q := q_{55}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $sq^{-1} = 1$  and  $r^{-1}s \neq 1$ . Hence the reflection of  $X$

$$X : \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{s}{\circ} \overset{-1}{\bullet} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \Rightarrow r_4(X) : \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ} \overset{-1}{\circ} \overset{s}{\circ} \overset{-1}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 ( $b_1$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 3.3. Consider that  $q_{11} = q_{33} = q_{55} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 4, 6\}$  and  $j \in \{2, 4, 6, 7\}$ . Let  $r := q_{22}$ ,  $s := q_{44}$ , and  $q := q_{66}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $sq = 1$  and  $rs \neq 1$ . Hence the reflections of  $X$

$$X : \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\bullet} \overset{-1}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ} \Rightarrow r_5(X) : \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{s^{-1}}{\circ} \overset{-1}{\bullet} \overset{s}{\circ} \overset{-1}{\circ} \overset{s^{-1}}{\circ} \overset{-1}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ}$$

$$\Rightarrow r_4 r_5(X) : \overset{-1}{\circ} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{-1}{\circ} \overset{s^{-1}}{\circ} \overset{-1}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ}$$

imply  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4 r_5(X)} = -1$ . By Lemma 4.8 ( $b_1$ ), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{33} = q_{55} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 4, 6\}$  and  $j \in \{2, 4, 6\}$ . Let  $r := q_{11}$ ,  $s := q_{44}$  and  $q := q_{66}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rs \neq 1$  and  $sq = 1$ . Similar to the proof above, we obtain  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4 r_5(X)} = -1$ . By Lemma 4.8 ( $b_2$ ), this case is impossible to appear.

Step 3.4. Consider that  $q_{11} = q_{33} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 4, 5\}$  and  $j \in \{2, 4, 5, 7\}$ . Let  $r := q_{22}$ ,  $s := q_{44}$ , and  $q := q_{77}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $sq = 1$  and  $rs \neq 1$ . Hence the reflections of  $X$

$$X : \overset{-1}{\bullet} \overset{r^{-1}}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ} \overset{-1}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ} \Rightarrow r_1(X) : \overset{-1}{\circ} \overset{r}{\circ} \overset{-1}{\bullet} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ} \overset{-1}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ}$$

$$\Rightarrow r_2 r_1(X) : \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{-1}{\circ} \overset{r}{\circ} \overset{r^{-1}}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ} \overset{-1}{\circ} \overset{s}{\circ} \overset{s^{-1}}{\circ}$$

imply  $r = -1$  and  $p \neq 2$  by  $a_{34}^{r_2 r_1(X)} = -1$ . By Lemma 4.8 ( $c_3$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 3.5. Consider that  $q_{11} = q_{33} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 4, 5, 6\}$  and  $j \in \{2, 4, 5, 6\}$ . Let  $r := q_{22}$  and  $q := q_{44}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rq \neq 1$ . Similar to the proof in Step 3.4, we get  $r = -1$  and  $p \neq 2$  by  $a_{34}^{r_2 r_1(X)} = -1$ . Rewrite  $X$  as

$$\begin{array}{cccccccccccc} -1 & -1 & -1 & -1 & q^{-1} & q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array}$$

and this case is impossible to appear by Lemma 4.8 ( $c_2$ ). Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 3.6. Consider that  $q_{22} = q_{33} = q_{44} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 5, 6\}$  and  $j \in \{5, 6, 7\}$ . Let  $q := q_{11}$ ,  $s := q'_{23}$ ,  $t := q'_{34}$  and  $r := q_{55}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $r^{-1}s = 1$ ,  $tq^{-1} = 1$  and  $st \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccccccc} s & s^{-1} & -1 & s & -1 & t & -1 & t^{-1} & t & t^{-1} & t & t^{-1} & t \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \bullet & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccccccc} s & s^{-1} & -1 & s & t & t^{-1} & -1 & t & -1 & t^{-1} & t & t^{-1} & t \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 ( $a_2$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 3.7. Consider that  $q_{22} = q_{33} = q_{55} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 4, 6\}$  and  $j \in \{4, 6, 7\}$ . Let  $r := q_{11}$ ,  $s := q'_{23}$ ,  $q := q_{44}$  and  $t := q_{66}$ . Since  $X$  has a good  $A_7$  neighborhood, then  $r^{-1}s = 1$ ,  $qt = 1$  and  $sq^{-1} \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccccccc} s & s^{-1} & -1 & s & -1 & q^{-1} & q & q^{-1} & -1 & q & q^{-1} & q & q^{-1} \\ \circ & \text{---} & \bullet & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccccccc} -1 & s & -1 & s^{-1} & s & q^{-1} & q & q^{-1} & -1 & q & q^{-1} & q & q^{-1} \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.8 ( $c_8$ ), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{33} = q_{55} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 4\}$  and  $j \in \{2, 4, 7\}$ . Let  $r := q_{11}$ ,  $s := q_{44}$ ,  $q := q'_{56}$  and  $t := q_{77}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $s^{-1}q = 1$ ,  $qt^{-1} = 1$  and  $rs \neq 1$ . Hence the reflections of  $X$

$$X : \begin{array}{cccccccccccc} r & r^{-1} & r & r^{-1} & -1 & s^{-1} & s & s^{-1} & -1 & s & -1 & s^{-1} & s \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \bullet & \text{---} & \circ & \text{---} & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{cccccccccccc} r & r^{-1} & r & r^{-1} & -1 & s^{-1} & -1 & s & -1 & s^{-1} & s & s^{-1} & s \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \bullet & \text{---} & \circ & \text{---} & \circ \end{array}$$

$$\Rightarrow r_4 r_5(X) : \begin{array}{cccccccccccc} r & r^{-1} & r & r^{-1} & s^{-1} & s & -1 & s^{-1} & s & s^{-1} & s & s^{-1} & s \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array}$$

imply  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4 r_5(X)} = -1$ . By Lemma 4.8 ( $b_2$ ), this case is impossible to appear.

Step 3.8. Consider that  $q_{22} = q_{33} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 4, 5\}$  and  $j \in \{4, 5, 7\}$ . Let  $r := q_{11}$ ,  $s := q'_{23}$ ,  $q := q_{44}$  and  $t := q_{77}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $r^{-1}s = 1$ ,  $qt = 1$  and  $sq^{-1} \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccccccc} s & s^{-1} & -1 & s & -1 & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & q & q^{-1} \\ \circ & \text{---} & \bullet & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccccccc} -1 & s & -1 & s^{-1} & s & q^{-1} & q & q^{-1} & q & q^{-1} & -1 & q & q^{-1} \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.8 ( $c_3$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 3.9. Consider that  $q_{22} = q_{33} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 4, 5, 6\}$  and  $j \in \{4, 5, 6\}$ . Let  $r := q_{11}$ ,  $s := q'_{23}$  and  $q := q_{44}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $r^{-1}s = 1$  and  $sq^{-1} \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} s & s^{-1} & -1 & s & -1 & q^{-1} & q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccc} -1 & s & -1 & s^{-1} & s & q^{-1} & q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.8 ( $c_2$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 3.10. Consider that  $q_{33} = q_{44} = q_{55} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 6\}$  and  $j \in \{2, 6, 7\}$ . Let  $r := q_{11}$ ,  $s := q'_{34}$ ,  $q := q'_{45}$  and  $t := q_{66}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $sq = 1$ ,  $qt^{-1} = 1$  and  $r^{-1}s \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & -1 & s & -1 & s^{-1} & -1 & s & s^{-1} & s & s^{-1} \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & s & s^{-1} & -1 & s & s^{-1} & s & s^{-1} & s & s^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 ( $b_2$ ), this case is impossible to appear.

Step 3.11. Consider that  $q_{33} = q_{44} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 5\}$  and  $j \in \{2, 5, 7\}$ . Let  $r := q_{11}$ ,  $s := q'_{34}$ ,  $q := q_{55}$  and  $t := q_{77}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $sq^{-1} = 1$ ,  $qt = 1$  and  $r^{-1}s \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & -1 & s & -1 & s^{-1} & -1 & s & s^{-1} \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & s & s^{-1} & -1 & s & -1 & s^{-1} & -1 & s & s^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 ( $b_2$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 3.12. Consider that  $q_{33} = q_{44} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 5, 6\}$  and  $j \in \{2, 5, 6\}$ . Let  $r := q_{11}$ ,  $s := q'_{34}$  and  $q := q_{55}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $sq^{-1} = 1$  and  $r^{-1}s \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & -1 & s & -1 & s^{-1} & s & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & s & s^{-1} & -1 & s & -1 & s^{-1} & s & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 ( $b_2$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 3.13. Consider that  $q_{33} = q_{66} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 4, 5\}$  and  $j \in \{2, 4, 5\}$ . Let  $q := q_{11}$ ,  $r := q_{44}$  and  $s := q'_{67}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $qr \neq 1$  and  $r^{-1}s = 1$ . Similar to the proof in Step 2.5, we obtain  $r = -1$  and  $p \neq 2$  by  $a_{32}^{r_4 r_5 r_6(X)} = -1$ . Rewrite  $X$  as

$$\begin{array}{cccccccccccc} q & q^{-1} & q & q^{-1} & -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

and this case is impossible to appear by Lemma 4.8 ( $b_2$ ). Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 4. Assume that there are four vertices labeled by  $-1$ . By  $q_{33} = -1$ , we distinguish the following sixteen steps.

Step 4.1. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{5, 6\}$  and  $j \in \{5, 6, 7\}$ . Let  $r := q'_{12}$ ,  $s := q'_{23}$ ,  $t := q'_{34}$  and  $q := q_{55}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rs = 1$  by  $a_{13}^{r_2(X)} = 0$ ,  $tq^{-1} = 1$  by  $a_{35}^{r_4(X)} = 0$  and  $st \neq 1$  by  $a_{24}^{r_3(X)} = -1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & t & -1 & t^{-1} & t & t^{-1} & t & t^{-1} & t \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & t & t^{-1} & -1 & t & -1 & t^{-1} & t & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 ( $a_1$ ), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{44} = q_{55} = q_{66} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 3\}$  and  $j \in \{2, 3\}$ . By Lemma 4.7, this case is impossible to appear.

Step 4.2. Consider that  $q_{11} = q_{22} = q_{33} = q_{55} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{4, 6\}$  and  $j \in \{4, 6, 7\}$ . Let  $r := q'_{12}$ ,  $s := q'_{23}$ ,  $t := q_{44}$  and  $q := q_{66}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rs = 1$ ,  $tq = 1$  and  $st^{-1} \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & t^{-1} & t & t^{-1} & -1 & t & t^{-1} & t & t^{-1} \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccc} s^{-1} & s & -1 & s^{-1} & s & t^{-1} & t & t^{-1} & -1 & t & t^{-1} & t & t^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.8 ( $c_8$ ), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{33} = q_{55} = q_{66} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 4\}$  and  $j \in \{2, 4\}$ . Let  $r := q_{11}$ ,  $s := q_{44}$ ,  $t := q'_{56}$  and  $q := q'_{67}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $s^{-1}t = 1$ ,  $tq = 1$  and  $rs \neq 1$ . Similar to the proof in Step 2.4, we obtain  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4 r_5(X)} = -1$ . Rewrite  $X$  as

$$\begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

and this case is impossible to appear by Lemma 4.8 ( $b_2$ ).

Step 4.3. Consider that  $q_{11} = q_{22} = q_{33} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{4, 5\}$  and  $j \in \{4, 5, 7\}$ . Let  $r := q'_{12}$ ,  $s := q'_{23}$ ,  $t := q_{44}$  and  $q := q_{77}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rs = 1$ ,  $tq = 1$  and  $st^{-1} \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & t^{-1} & t & t^{-1} & t & t^{-1} & -1 & t & t^{-1} \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccc} s^{-1} & s & -1 & s^{-1} & s & t^{-1} & t & t^{-1} & t & t^{-1} & -1 & t & t^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.8 ( $c_3$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 4.4. Consider that  $q_{11} = q_{22} = q_{33} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{4, 5, 6\}$  and  $j \in \{4, 5, 6\}$ . Let  $r := q'_{12}$ ,  $s := q'_{23}$  and  $q := q_{44}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rs = 1$  and  $sq^{-1} \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & q^{-1} & q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccc} s^{-1} & s & -1 & s^{-1} & s & q^{-1} & q & q^{-1} & q & q^{-1} & q & q^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.8 ( $c_2$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 4.5. Consider that  $q_{11} = q_{33} = q_{44} = q_{55} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 6\}$  and  $j \in \{2, 6, 7\}$ . Let  $r := q_{22}$ ,  $s := q'_{34}$ ,  $t := q'_{45}$  and  $q = q_{66}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $st = 1$ ,  $tq^{-1} = 1$  and  $r^{-1}s \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & -1 & s & -1 & s^{-1} & -1 & s & s^{-1} & s & s^{-1} \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & s & s^{-1} & -1 & s & s^{-1} & s & s^{-1} & s & s^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 (b<sub>1</sub>), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{33} = q_{44} = q_{55} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 6\}$  and  $j \in \{2, 6\}$ . Let  $r := q_{11}$ ,  $s := q'_{34}$ ,  $t := q'_{45}$  and  $q := q_{66}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $st = 1$ ,  $tq^{-1} = 1$  and  $r^{-1}s \neq 1$ . Similar to the proof above, we have  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 (b<sub>2</sub>), this case is impossible to appear.

Step 4.6. Consider that  $q_{11} = q_{33} = q_{44} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 5\}$  and  $j \in \{2, 5, 7\}$ . Let  $r := q_{22}$ ,  $s := q'_{34}$ ,  $t := q_{55}$  and  $q = q_{77}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $st^{-1} = 1$ ,  $tq = 1$  and  $r^{-1}s \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & -1 & s & -1 & s^{-1} & s & s^{-1} & -1 & s & s^{-1} \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & s & s^{-1} & -1 & s & -1 & s^{-1} & -1 & s & s^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 (b<sub>1</sub>), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 4.7. Consider that  $q_{11} = q_{33} = q_{44} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 5, 6\}$  and  $j \in \{2, 5, 6\}$ . Let  $r := q_{22}$ ,  $s := q'_{34}$  and  $q := q_{55}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $sq^{-1} = 1$  and  $r^{-1}s \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & -1 & s & -1 & s^{-1} & s & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & s & s^{-1} & -1 & s & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 (b<sub>1</sub>), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 4.8. Consider that  $q_{11} = q_{33} = q_{55} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 4\}$  and  $j \in \{2, 4, 7\}$ . Let  $r := q_{22}$ ,  $s := q_{44}$ ,  $t := q'_{56}$  and  $q = q_{77}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $s^{-1}t = 1$ ,  $tq^{-1} = 1$  and  $rs \neq 1$ . Hence the reflections of  $X$

$$X : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & s & s^{-1} & -1 & s & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & -1 & s & -1 & s^{-1} & s & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

$$\Rightarrow r_4 r_5(X) : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & s^{-1} & s & -1 & s^{-1} & s & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

imply  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4 r_5(X)} = -1$ . By Lemma 4.8 (b<sub>1</sub>), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{22} = q_{33} = q_{55} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 4, 6\}$  and  $j \in \{4, 6\}$ . Let  $r := q_{11}$ ,  $s := q'_{23}$ ,  $t := q_{44}$  and  $q := q_{66}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $r^{-1}s = 1$ ,  $tq = 1$  and  $st^{-1} \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} s & s^{-1} & -1 & s & -1 & t^{-1} & t & t^{-1} & -1 & t & t^{-1} & t & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccc} -1 & s & -1 & s^{-1} & s & t^{-1} & t & t^{-1} & -1 & t & t^{-1} & t & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.8 (c<sub>7</sub>), this case is impossible to appear.

Step 4.9. Consider that  $q_{11} = q_{33} = q_{55} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 4, 6\}$  and  $j \in \{2, 4, 6\}$ . Let  $r := q_{22}$ ,  $s := q_{44}$  and  $q := q_{66}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $sq = 1$  and  $rs \neq 1$ . Hence the reflections of  $X$

$$\begin{aligned}
X : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & s & s^{-1} \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ \end{array} \Rightarrow r_5(X) : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & -1 & s \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ \end{array} \\
\Rightarrow r_4 r_5(X) : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & s^{-1} & s & -1 & s^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}
\end{aligned}$$

imply  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4 r_5(X)} = -1$ . By Lemma 4.8 (b<sub>1</sub>), this case is impossible to appear.

Step 4.10. Consider that  $q_{11} = q_{33} = q_{66} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 4, 5\}$  and  $j \in \{2, 4, 5\}$ . Let  $r := q_{22}$ ,  $s := q_{44}$  and  $q = q'_{67}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $s^{-1}q = 1$  and  $rs \neq 1$ . Hence the reflections of  $X$

$$\begin{aligned}
X : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & s & s^{-1} \\ \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_1(X) : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & s & s^{-1} \\ \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \\
\Rightarrow r_2 r_1(X) : \begin{array}{cccccccc} r & r^{-1} & -1 & r & r^{-1} & s^{-1} & s & s^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}
\end{aligned}$$

imply  $r = -1$  and  $p \neq 2$  by  $a_{34}^{r_2 r_1(X)} = -1$ . By Lemma 4.8 (c<sub>4</sub>), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 4.11. Consider that  $q_{22} = q_{33} = q_{44} = q_{55} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 6\}$  and  $j \in \{6, 7\}$ . Let  $r := q_{11}$ ,  $s := q'_{23}$ ,  $t := q'_{34}$ ,  $u := q'_{45}$  and  $q := q_{66}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $r^{-1}s = 1$ ,  $tu = 1$ ,  $uq^{-1} = 1$  and  $st \neq 1$ . Hence the reflection of  $X$

$$\begin{aligned}
X : \begin{array}{cccccccc} s & s^{-1} & -1 & s & -1 & t & -1 & t^{-1} \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} s & s^{-1} & -1 & s & t & t^{-1} & -1 & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}
\end{aligned}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 (a<sub>2</sub>), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{33} = q_{44} = q_{55} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2\}$  and  $j \in \{2, 7\}$ . Let  $r := q_{11}$ ,  $s := q'_{34}$ ,  $t := q'_{45}$ ,  $u := q'_{56}$  and  $q := q_{77}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $st = 1$ ,  $tu = 1$ ,  $uq^{-1} = 1$  and  $r^{-1}s \neq 1$ . Hence the reflection of  $X$

$$\begin{aligned}
X : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & -1 & s & -1 & s^{-1} \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & s & s^{-1} & -1 & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}
\end{aligned}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 (b<sub>2</sub>), this case is impossible to appear.

Step 4.12. Consider that  $q_{22} = q_{33} = q_{44} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 5\}$  and  $j \in \{5, 7\}$ . Let  $r := q_{11}$ ,  $s := q'_{23}$ ,  $t := q'_{34}$ ,  $u := q_{55}$  and  $q := q_{77}$ . Since



$X$  has a good  $A_7$  neighborhood, we have  $r^{-1}s = 1$ ,  $tu^{-1} = 1$ ,  $uq = 1$  and  $st \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} s & s^{-1} & -1 & s & -1 & t & -1 & t^{-1} & t & t^{-1} & -1 & t & t^{-1} \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} s & s^{-1} & -1 & s & t & t^{-1} & -1 & t & -1 & t^{-1} & -1 & t & t^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 ( $a_2$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 4.13. Consider that  $q_{22} = q_{33} = q_{44} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 6\}$  and  $j \in \{2, 6\}$ . Let  $r := q_{11}$ ,  $s := q'_{23}$ ,  $t := q'_{34}$  and  $q := q_{55}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $r^{-1}s = 1$ ,  $tq^{-1} = 1$  and  $st \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} s & s^{-1} & -1 & s & -1 & t & -1 & t^{-1} & t & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} s & s^{-1} & -1 & s & t & t^{-1} & -1 & t & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 ( $a_2$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 4.14. Consider that  $q_{22} = q_{33} = q_{55} = q_{66} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 4\}$  and  $j \in \{4, 7\}$ . Let  $r := q_{11}$ ,  $s := q'_{23}$ ,  $t := q_{44}$  and  $q := q_{77}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $r^{-1}s = 1$ ,  $t^{-1}u = 1$ ,  $uq^{-1} = 1$  and  $st^{-1} \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} s & s^{-1} & -1 & s & -1 & t^{-1} & t & t^{-1} & -1 & t & -1 & t^{-1} & t \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccc} -1 & s & -1 & s^{-1} & s & t^{-1} & t & t^{-1} & -1 & t & -1 & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.8 ( $c_6$ ), this case is impossible to appear.

Step 4.15. Consider that  $q_{22} = q_{33} = q_{66} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 4, 5\}$  and  $j \in \{4, 5\}$ . Let  $r := q_{11}$ ,  $s := q'_{23}$ ,  $t := q_{44}$  and  $q := q'_{67}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $r^{-1}s = 1$ ,  $t^{-1}q = 1$  and  $st^{-1} \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} s & s^{-1} & -1 & s & -1 & t^{-1} & t & t^{-1} & t & t^{-1} & -1 & t & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccc} -1 & s & -1 & s^{-1} & s & t^{-1} & t & t^{-1} & t & t^{-1} & -1 & t & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.8 ( $c_4$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 4.16. Consider that  $q_{33} = q_{44} = q_{66} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{1, 2, 5\}$  and  $j \in \{2, 5\}$ . Let  $r := q_{11}$ ,  $s := q'_{34}$ ,  $t := q_{55}$  and  $q := q'_{67}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $st^{-1} = 1$ ,  $t^{-1}q = 1$  and  $r^{-1}s \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & -1 & s & -1 & s^{-1} & s & s^{-1} & -1 & s & -1 \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} r & r^{-1} & r & r^{-1} & s & s^{-1} & -1 & s & -1 & s^{-1} & -1 & s & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 ( $b_2$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 5. Assume that there are five vertices labeled by  $-1$ . By  $q_{33} = -1$ , we distinguish the following eleven steps.

Step 5.1. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = q_{55} = -1$ ,  $q_{66}q'_{67} - 1 = 0$  and  $q_{jj}q'_{j-1,j} - 1 = 0$  for  $j \in \{6, 7\}$ . Let  $r := q'_{12}$ ,  $s := q'_{23}$ ,  $t := q'_{34}$ ,  $u := q'_{45}$  and  $q := q_{66}$ . Since  $X$  has a good

$A_7$  neighborhood, we have  $rs = 1$  by  $a_{13}^{r_2(X)} = 0$ ,  $tu = 1$  by  $a_{35}^{r_4(X)} = 0$  and  $uq^{-1} = 1$  by  $a_{46}^{r_5(X)} = 0$ . In addition, we obtain  $st \neq 1$  by  $a_{24}^{r_3(X)} = -1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & t & -1 & t^{-1} & -1 & t & t^{-1} & t & t^{-1} \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & t & t^{-1} & -1 & t & t^{-1} & t & t^{-1} & t & t^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 ( $a_1$ ), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{33} = q_{44} = q_{55} = q_{66} = q_{77} = -1$ ,  $q_{22}q'_{12} - 1 = 0$  and  $q_{ii}q'_{i,i+1} - 1 = 0$  for  $i \in \{2, 4, 5\}$ . Let  $r := q_{11}$ ,  $s := q'_{34}$ ,  $t := q'_{45}$ ,  $u := q'_{56}$  and  $q := q'_{67}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $st = 1$ ,  $tu = 1$ ,  $uq = 1$  and  $r^{-1}s \neq 1$ . Similar to the proof above, we have  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 ( $b_2$ ), this case is impossible to appear.

Step 5.2. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = q_{66} = -1$ ,  $q_{55}q'_{56} - 1 = 0$  and  $q_{jj}q'_{j-1,j} - 1 = 0$  for  $j \in \{5, 7\}$ . Let  $r := q'_{12}$ ,  $s := q'_{23}$ ,  $t := q'_{34}$ ,  $u := q_{55}$  and  $q := q_{77}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rs = 1$ ,  $tu^{-1} = 1$ ,  $uq = 1$  and  $st \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & t & -1 & t^{-1} & t & t^{-1} & -1 & t & t^{-1} \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & t & t^{-1} & -1 & t & -1 & t^{-1} & -1 & t & t^{-1} \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 ( $a_1$ ), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{22} = q_{44} = q_{55} = q_{66} = q_{77} = -1$ ,  $q_{33}q'_{23} - 1 = 0$  and  $q_{ii}q'_{i,i+1} - 1 = 0$  for  $i \in \{1, 3\}$ . By Lemma 4.7, this case is impossible to appear.

Step 5.3. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{5, 6\}$  and  $j \in \{5, 6\}$ . Let  $r := q'_{12}$ ,  $s := q'_{23}$ ,  $t := q'_{34}$  and  $q := q_{55}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rs = 1$ ,  $tq^{-1} = 1$  and  $st \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & t & -1 & t^{-1} & t & t^{-1} & t & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & t & t^{-1} & -1 & t & -1 & t^{-1} & t & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 ( $a_1$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 5.4. Consider that  $q_{11} = q_{22} = q_{33} = q_{55} = q_{66} = -1$ ,  $q_{44}q'_{45} - 1 = 0$  and  $q_{jj}q'_{j-1,j} - 1 = 0$  for  $j \in \{4, 7\}$ . Let  $r := q'_{12}$ ,  $s := q'_{23}$ ,  $t := q_{44}$ ,  $u := q'_{56}$  and  $q := q_{77}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rs = 1$ ,  $t^{-1}u = 1$ ,  $uq^{-1} = 1$  and  $st^{-1} \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & t^{-1} & t & t^{-1} & -1 & t & -1 & t^{-1} & t \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccc} s^{-1} & s & -1 & s^{-1} & s & t^{-1} & t & t^{-1} & -1 & t & -1 & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.8 ( $c_6$ ), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{22} = q_{33} = q_{55} = q_{66} = q_{77} = -1$ ,  $q_{44}q'_{34} - 1 = 0$  and  $q_{ii}q'_{i,i+1} - 1 = 0$  for  $i \in \{1, 4\}$ . Let  $r := q_{11}$ ,  $s := q'_{23}$ ,  $t := q_{44}$ ,  $u := q'_{56}$  and  $q := q'_{67}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $r^{-1}s = 1$ ,  $t^{-1}u = 1$ ,  $uq = 1$  and  $st^{-1} \neq 1$ . Similar to the proof above, we have  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.8 ( $c_5$ ), this case is impossible to appear.

Step 5.5. Consider that  $q_{11} = q_{22} = q_{33} = q_{55} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{4, 6\}$  and  $j \in \{4, 6\}$ . Let  $r := q'_{12}$ ,  $s := q'_{23}$ ,  $t := q_{44}$  and  $q := q_{66}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rs = 1$ ,  $tq = 1$  and  $st^{-1} \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & t^{-1} & t & t^{-1} & -1 & t & t^{-1} & t & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccc} s^{-1} & s & -1 & s^{-1} & s & t^{-1} & t & t^{-1} & -1 & t & t^{-1} & t & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.8 ( $c_7$ ), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{11} = q_{33} = q_{55} = q_{66} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 4\}$  and  $j \in \{2, 4\}$ . Let  $r := q_{22}$ ,  $s := q_{44}$ ,  $t := q'_{56}$  and  $q := q'_{67}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $s^{-1}t = 1$ ,  $tq = 1$  and  $rs \neq 1$ . Hence the reflections of  $X$

$$\begin{aligned} X : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & s & s^{-1} & -1 & s & -1 & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} & \Rightarrow & r_5(X) : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & -1 & s^{-1} & -1 & s & -1 & s^{-1} & s & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \\ & \Rightarrow & r_4r_5(X) : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & s^{-1} & s & -1 & s^{-1} & s & s^{-1} & s & s^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \end{aligned}$$

imply  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4r_5(X)} = -1$ . By Lemma 4.8 ( $b_1$ ), this case is impossible to appear.

Step 5.6. Consider that  $q_{11} = q_{22} = q_{33} = q_{66} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{4, 5\}$  and  $j \in \{4, 5\}$ . Let  $r := q'_{12}$ ,  $s := q'_{23}$ ,  $t := q_{44}$  and  $q := q'_{67}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rs = 1$ ,  $t^{-1}q = 1$  and  $st^{-1} \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & t^{-1} & t & t^{-1} & t & t^{-1} & -1 & t & -1 \\ \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccc} s^{-1} & s & -1 & s^{-1} & s & t^{-1} & t & t^{-1} & t & t^{-1} & -1 & t & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.8 ( $c_4$ ), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 5.7. Consider that  $q_{11} = q_{33} = q_{44} = q_{55} = q_{66} = -1$ ,  $q_{22}q'_{23} - 1 = 0$  and  $q_{jj}q'_{j-1,j} - 1 = 0$  for  $j \in \{2, 7\}$ . Let  $r := q_{22}$ ,  $s := q'_{34}$ ,  $t := q'_{45}$ ,  $u := q'_{56}$  and  $q := q_{77}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $st = 1$ ,  $tu = 1$ ,  $uq^{-1} = 1$  and  $r^{-1}s \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & -1 & s & -1 & s^{-1} & -1 & s & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & s & s^{-1} & -1 & s & s^{-1} & s & -1 & s^{-1} & s \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 ( $b_1$ ), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{22} = q_{33} = q_{44} = q_{55} = q_{77} = -1$ ,  $q_{66}q'_{56} - 1 = 0$  and  $q_{ii}q'_{i,i+1} - 1 = 0$  for  $i \in \{1, 6\}$ . Let  $r := q_{11}$ ,  $s := q'_{23}$ ,  $t := q'_{34}$ ,  $u := q'_{45}$  and  $q := q_{66}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $r^{-1}s = 1$ ,  $tu = 1$ ,  $uq^{-1} = 1$  and  $st \neq 1$ . Similar to the proof above, we have  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 ( $a_2$ ), this case is impossible to appear.

Step 5.8. Consider that  $q_{11} = q_{33} = q_{44} = q_{55} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 6\}$  and  $j \in \{2, 6\}$ . Let  $r := q_{22}$ ,  $s := q'_{34}$ ,  $t := q'_{45}$  and  $q := q_{66}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $st = 1$ ,  $tq^{-1} = 1$  and  $r^{-1}s \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & -1 & s & -1 & s^{-1} & -1 & s & s^{-1} & s & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & s & s^{-1} & -1 & s & s^{-1} & s & s^{-1} & s & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 ( $b_1$ ), this case is impossible to appear.

Step 5.9. Consider that  $q_{11} = q_{33} = q_{44} = q_{66} = q_{77} = -1$  and  $q_{ii}q'_{i,i+1} - 1 = q_{jj}q'_{j-1,j} - 1 = 0$  for all  $i \in \{2, 5\}$  and  $j \in \{2, 5\}$ . Let  $r := q_{22}$ ,  $s := q'_{34}$ ,  $t := q_{55}$  and  $q := q'_{67}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $st^{-1} = 1$ ,  $t^{-1}q = 1$  and  $r^{-1}s \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & -1 & s & -1 & s^{-1} & s & s^{-1} & -1 & s & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & r^{-1} & r & r^{-1} & s & s^{-1} & -1 & s & -1 & s^{-1} & -1 & s & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  by  $a_{32}^{r_4(X)} = -1$ ,  $p \neq 2$ . By Lemma 4.8 (b<sub>1</sub>), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 5.10. Consider that  $q_{22} = q_{33} = q_{44} = q_{55} = q_{66} = -1$  and  $q_{11}q'_{12} - 1 = q_{77}q'_{67} - 1 = 0$ . Let  $r := q_{11}$ ,  $s := q'_{23}$ ,  $t := q'_{34}$ ,  $u := q'_{45}$ ,  $v := q'_{56}$  and  $q := q_{77}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $r^{-1}s = 1$ ,  $tu = 1$ ,  $uv = 1$ ,  $vq^{-1} = 1$  and  $st \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} s & s^{-1} & -1 & s & -1 & t & -1 & t^{-1} & -1 & t & -1 & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} s & s^{-1} & -1 & s & t & t^{-1} & -1 & t & t^{-1} & t & -1 & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 (a<sub>2</sub>), this case is impossible to appear.

Step 5.11. Consider that  $q_{22} = q_{33} = q_{44} = q_{66} = q_{77} = -1$ ,  $q_{55}q'_{45} - 1 = 0$  and  $q_{ii}q'_{i,i+1} - 1 = 0$  for  $i \in \{1, 5\}$ . Let  $r := q_{11}$ ,  $s := q'_{23}$ ,  $t := q'_{34}$ ,  $u := q_{55}$  and  $q := q'_{67}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $r^{-1}s = 1$ ,  $tu^{-1} = 1$ ,  $u^{-1}q = 1$  and  $st \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} s & s^{-1} & -1 & s & -1 & t & -1 & t^{-1} & t & t^{-1} & -1 & t & -1 \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} s & s^{-1} & -1 & s & t & t^{-1} & -1 & t & -1 & t^{-1} & -1 & t & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 (a<sub>2</sub>), this case is impossible to appear. Further by Lemma 4.7 the symmetric case is impossible to appear.

Step 6. Assume that there are six vertices labeled by  $-1$ . By  $q_{33} = -1$ , we distinguish the following four steps.

Step 6.1. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = q_{55} = q_{66} = -1$  and  $q_{77}q'_{67} - 1 = 0$ . Let  $r := q'_{12}$ ,  $s := q'_{23}$ ,  $t := q'_{34}$ ,  $u := q'_{45}$ ,  $v := q'_{56}$  and  $q := q_{77}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rs = 1$ ,  $tu = 1$ ,  $uv = 1$ ,  $vq^{-1} = 1$  and  $st \neq 1$  by Definition 3.3. Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & t & -1 & t^{-1} & -1 & t & -1 & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & t & t^{-1} & -1 & t & t^{-1} & t & -1 & t^{-1} & t \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 (a<sub>1</sub>), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{22} = q_{33} = q_{44} = q_{55} = q_{66} = q_{77} = -1$  and  $q_{11}q'_{12} - 1 = 0$ . Let  $r := q_{11}$ ,  $s := q'_{23}$ ,  $t := q'_{34}$ ,  $u := q'_{45}$ ,  $v := q'_{56}$  and  $q := q'_{67}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $r^{-1}s = 1$ ,  $tu = 1$ ,  $uv = 1$ ,  $vq = 1$  and  $st \neq 1$ . Similar to the proof above, we have  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 (a<sub>2</sub>), this case is impossible to appear.

Step 6.2. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = q_{55} = q_{77} = -1$  and  $q_{66}q'_{56} - 1 = q_{66}q'_{67} - 1 = 0$ . Let  $r := q'_{12}$ ,  $s := q'_{23}$ ,  $t := q'_{34}$ ,  $u := q'_{45}$  and  $q := q_{66}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rs = 1$ ,  $tu = 1$ ,  $uq^{-1} = 1$  and  $st \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & t & -1 & t^{-1} & -1 & t & t^{-1} & t & -1 \\ \circ & \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & t & t^{-1} & -1 & t & t^{-1} & t & t^{-1} & t & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 (a<sub>1</sub>), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{11} = q_{33} = q_{44} = q_{55} = q_{66} = q_{77} = -1$  and  $q_{22}q'_{12} - 1 = q_{22}q'_{23} - 1 = 0$ . Let  $r := q_{22}$ ,  $s := q'_{34}$ ,  $t := q'_{45}$ ,  $u := q'_{56}$  and

$q := q'_{67}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $st = 1$ ,  $tu = 1$ ,  $uq = 1$  and  $r^{-1}s \neq 1$ . Similar to the proof above, we have  $s = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 (b<sub>1</sub>), this case is impossible to appear.

Step 6.3. Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = q_{66} = q_{77} = -1$  and  $q_{55}q'_{45} - 1 = q_{55}q'_{56} - 1 = 0$ . Let  $r := q'_{12}$ ,  $s := q'_{23}$ ,  $t := q'_{34}$ ,  $u := q_{55}$  and  $q := q'_{67}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rs = 1$ ,  $tu^{-1} = 1$ ,  $u^{-1}q = 1$  and  $st \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & t & -1 & t^{-1} & t & t^{-1} & -1 & t & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & t & t^{-1} & -1 & t & -1 & t^{-1} & -1 & t & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  by  $a_{32}^{r_4(X)} = -1$ ,  $p \neq 2$ . By Lemma 4.8 (a<sub>1</sub>), this case is impossible to appear. Further consider the symmetric case of this case that  $q_{11} = q_{22} = q_{44} = q_{55} = q_{66} = q_{77} = -1$  and  $q_{33}q'_{23} - 1 = q_{33}q'_{34} - 1 = 0$ . By Lemma 4.7, this case is impossible to appear.

Step 6.4. Consider that  $q_{11} = q_{22} = q_{33} = q_{55} = q_{66} = q_{77} = -1$  and  $q_{44}q'_{34} - 1 = q_{44}q'_{45} - 1 = 0$ . Let  $r := q'_{12}$ ,  $s := q'_{23}$ ,  $t := q_{44}$ ,  $u := q'_{56}$  and  $q := q'_{67}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rs = 1$ ,  $t^{-1}u = 1$ ,  $uq = 1$  and  $st^{-1} \neq 1$ . Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & t^{-1} & t & t^{-1} & -1 & t & -1 & t^{-1} & -1 \\ \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_2(X) : \begin{array}{cccccccc} s^{-1} & s & -1 & s^{-1} & s & t^{-1} & t & t^{-1} & -1 & t & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $s = -1$  and  $p \neq 2$  by  $a_{34}^{r_2(X)} = -1$ . By Lemma 4.8 (c<sub>5</sub>), this case is impossible to appear.

Step 7. Assume that all vertices are labeled by  $-1$ . Consider that  $q_{11} = q_{22} = q_{33} = q_{44} = q_{55} = q_{66} = q_{77} = -1$ . Let  $r := q'_{12}$ ,  $s := q'_{23}$ ,  $t := q'_{34}$ ,  $u := q'_{45}$ ,  $v := q'_{56}$  and  $q := q'_{67}$ . Since  $X$  has a good  $A_7$  neighborhood, we have  $rs = 1$ ,  $tu = 1$ ,  $uv = 1$ ,  $vq = 1$  and  $st \neq 1$  by Definition 3.3. Hence the reflection of  $X$

$$X : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & -1 & t & -1 & t^{-1} & -1 & t & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \bullet & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \Rightarrow r_4(X) : \begin{array}{cccccccc} -1 & s^{-1} & -1 & s & t & t^{-1} & -1 & t & t^{-1} & t & -1 & t^{-1} & -1 \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array}$$

implies  $t = -1$  and  $p \neq 2$  by  $a_{32}^{r_4(X)} = -1$ . By Lemma 4.8 (a<sub>1</sub>), this case is impossible to appear.  $\square$

**Remark 4.10.** We present the exchange graphs of  $\mathcal{C}(M)$  in Table 2.

## 5 Generalized Dynkin diagrams and exchanges graphs

In this section, we give a complete list of all the Nichols algebras of diagonal type with finite root systems listed in Appendix 6.

### 5.1 Generalized Dynkin diagrams

| row | Dynkin diagrams            | fixed parameters | char $\mathbb{k}$ |
|-----|----------------------------|------------------|-------------------|
| 11  |                            | $\zeta \in R_3$  | $p \neq 3$        |
| 12  |                            | $\zeta \in R_3$  | $p \neq 3$        |
|     | row continues on next page |                  |                   |

| row | Dynkin diagrams            | fixed parameters | char $\mathbb{k}$ |
|-----|----------------------------|------------------|-------------------|
|     |                            |                  |                   |
| 13  |                            | $\zeta \in R_3$  | $p \neq 3$        |
|     | row continues on next page |                  |                   |

| row | Dynkin diagrams                   | fixed parameters | char $\mathbb{k}$ |
|-----|-----------------------------------|------------------|-------------------|
|     |                                   |                  |                   |
| 14  | <p>row continues on next page</p> | $\zeta \in R_4$  | $p \neq 2$        |



| row | Dynkin diagrams            | fixed parameters | char $\mathbb{k}$ |
|-----|----------------------------|------------------|-------------------|
|     |                            |                  |                   |
| 15  |                            | $\zeta \in R_5$  | $p \neq 5$        |
| 17  |                            | $\zeta \in R_3$  | $p \neq 3$        |
|     | row continues on next page |                  |                   |

| row | Dynkin diagrams | fixed parameters | char $\mathbb{k}$ |
|-----|-----------------|------------------|-------------------|
|     |                 |                  |                   |
| 18  |                 | $\zeta \in R_3$  | $p \neq 3$        |

| row | Dynkin diagrams            | fixed parameters | char $\mathbb{k}$ |
|-----|----------------------------|------------------|-------------------|
|     |                            |                  |                   |
|     |                            |                  |                   |
|     |                            |                  |                   |
|     |                            |                  |                   |
|     |                            |                  |                   |
|     |                            |                  |                   |
|     |                            |                  |                   |
|     |                            |                  |                   |
|     |                            |                  |                   |
|     |                            |                  |                   |
| 19  |                            | $\zeta \in R_4$  | $p \neq 2$        |
|     |                            |                  |                   |
|     | row continues on next page |                  |                   |

| row | Dynkin diagrams | fixed parameters | char $\mathbb{k}$ |
|-----|-----------------|------------------|-------------------|
|     |                 |                  |                   |
| 21  |                 | $\zeta \in R_3$  | $p \neq 3$        |

Table 1: Dynkin diagrams in all positive characteristic  $p > 0$

## 5.2 Exchange graphs

|    | exchange graph   | row | char $\mathbb{k}$ |
|----|--|-----|-------------------|
| 11 | $  \begin{array}{c}  \mathcal{D}_{11,1}^5 \xrightarrow{3} \mathcal{D}_{11,2}^5 \xrightarrow{2} \tau_{54321} \mathcal{D}_{11,3}^5 \xrightarrow{1} \tau_{54321} \mathcal{D}_{11,4}^5 \\    \\  \tau_{12435} \mathcal{D}_{11,3}^5 \xrightarrow{5} \tau_{12435} \mathcal{D}_{11,4}^5  \end{array}  $   | 11  | $p \neq 3$        |
| 12 | $  \begin{array}{c}  \tau_{13245} \mathcal{D}_{12,14}^5 \xrightarrow{2} \mathcal{D}_{12,11}^5 \xrightarrow{3} \mathcal{D}_{12,13}^5 \\  \swarrow 4 \quad   \quad   \\  \mathcal{D}_{12,9}^5 \xrightarrow{3} \mathcal{D}_{12,12}^5 \xrightarrow{2} \mathcal{D}_{12,15}^5 \\    \\  \tau_{12435} \mathcal{D}_{12,7}^5 \xrightarrow{1} \tau_{12435} \mathcal{D}_{12,6}^5 \xrightarrow{2} \tau_{12435} \mathcal{D}_{12,5}^5 \\    \quad   \quad   \\  \tau_{12435} \mathcal{D}_{12,10}^5 \xrightarrow{1} \tau_{12435} \mathcal{D}_{12,8}^5 \xrightarrow{2} \tau_{12435} \mathcal{D}_{12,4}^5 \\    \\  \tau_{12435} \mathcal{D}_{12,3}^5 \xrightarrow{3} \tau_{12435} \mathcal{D}_{12,2}^5 \xrightarrow{5} \tau_{12435} \mathcal{D}_{12,1}^5  \end{array}  $   | 12  | $p \neq 3$        |
| 13 | $  \begin{array}{c}  \mathcal{D}_{13,21}^5 \xrightarrow{1} \mathcal{D}_{13,20}^5 \xrightarrow{2} \mathcal{D}_{13,19}^5 \xrightarrow{3} \mathcal{D}_{13,18}^5 \\    \\  \tau_{12435} \mathcal{D}_{13,16}^5 \xrightarrow{1} \tau_{12435} \mathcal{D}_{13,15}^5 \xrightarrow{2} \tau_{12435} \mathcal{D}_{13,17}^5 \\    \quad   \quad   \\  \tau_{12435} \mathcal{D}_{13,13}^5 \xrightarrow{1} \tau_{12435} \mathcal{D}_{13,10}^5 \xrightarrow{2} \tau_{12435} \mathcal{D}_{13,14}^5 \\    \quad   \\  \tau_{12435} \mathcal{D}_{13,12}^5 \xrightarrow{2} \tau_{12435} \mathcal{D}_{13,9}^5 \xrightarrow{1} \tau_{12435} \mathcal{D}_{13,6}^5 \\    \quad   \quad   \\  \tau_{12435} \mathcal{D}_{13,11}^5 \xrightarrow{4} \tau_{12435} \mathcal{D}_{13,8}^5 \xrightarrow{2} \tau_{12435} \mathcal{D}_{13,5}^5 \xrightarrow{1} \tau_{12435} \mathcal{D}_{13,3}^5 \\    \quad   \quad   \\  \tau_{12435} \mathcal{D}_{13,7}^5 \xrightarrow{4} \tau_{12435} \mathcal{D}_{13,4}^5 \xrightarrow{2} \tau_{12435} \mathcal{D}_{13,2}^5 \xrightarrow{1} \tau_{12435} \mathcal{D}_{13,1}^5  \end{array}  $ | 13  | $p \neq 3$        |
| 14 | $  \begin{array}{c}  \tau_{12354} \mathcal{D}_{14,6}^5 \xrightarrow{1} \tau_{12354} \mathcal{D}_{14,5}^5 \\    \\  \mathcal{D}_{14,4}^5 \xrightarrow{4} \tau_{12354} \mathcal{D}_{14,3}^5 \xrightarrow{5} \tau_{12354} \mathcal{D}_{14,1}^5 \xrightarrow{3} \tau_{12354} \mathcal{D}_{14,2}^5 \\    \\  \tau_{12534} \mathcal{D}_{14,2}^5 \xrightarrow{5} \tau_{12534} \mathcal{D}_{14,1}^5 \xrightarrow{3} \tau_{12534} \mathcal{D}_{14,3}^5 \\    \\  \tau_{12534} \mathcal{D}_{14,5}^6 \xrightarrow{1} \tau_{12534} \mathcal{D}_{14,6}^6 \quad \tau_{12543} \mathcal{D}_{14,4}^5  \end{array}  $  | 14  | $p \neq 2$        |
| 15 | $  \begin{array}{c}  \mathcal{D}_{15,1}^5 \xrightarrow{3} \mathcal{D}_{15,2}^5 \xrightarrow{2} \mathcal{D}_{15,4}^5 \xrightarrow{1} \mathcal{D}_{15,5}^5 \\    \\  \tau_{12435} \mathcal{D}_{15,3}^5 \xrightarrow{5} \tau_{12453} \mathcal{D}_{15,6}^5 \xrightarrow{3} \tau_{12453} \mathcal{D}_{15,7}^5  \end{array}  $   | 15  | $p \neq 5$        |





## Acknowledgment

This work is supported by National Natural Science Foundation of China (No.12471038).

The authors thank N. Andruskiewitsch and H. Yamane for the invaluable suggestions to the references. Furthermore, the author J. Wang would like to thank her PhD advisor Prof. I. Heckenberger for the fruitful discussions during her PhD study in Germany, which have been crucial to the entire project.



## References

- [1] N. Andruskiewitsch. About finite dimensional Hopf algebras. *Contemp. Math.*, 294:1–57, 01 2002.
- [2] N. Andruskiewitsch. On finite-dimensional Hopf algebras. *Proc. Int. Cong. Math.*, 2:117–141, 2014.
- [3] N. Andruskiewitsch and M. Graña. Braided Hopf algebras over non abelian finite groups. *Bol. Acad. Nac. Cienc. (Córdoba)*, 63:45–78, 1998.
- [4] N. Andruskiewitsch, I. Heckenberger, and H.-J. Schneider. The Nichols algebra of a semisimple Yetter-Drinfeld module. *Amer. J. Math.*, 132(6):1493–1547, 2010.
- [5] N. Andruskiewitsch, I. Heckenberger, and L. Vendramin. Pointed hopf algebras of odd dimension and nichols algebras over solvable groups. *arXiv preprint 2411.02304*, 2024.
- [6] N. Andruskiewitsch and H.-J. Schneider. Lifting of quantum linear spaces and pointed Hopf algebras of order  $p^3$ . *J. Algebra*, 209:658–691, 1998.
- [7] N. Andruskiewitsch and H.-J. Schneider. Finite Quantum Groups and Cartan Matrices. *Adv. Math.*, 154(1):1–45, 2000.
- [8] N. Andruskiewitsch and H.-J. Schneider. Pointed Hopf algebras. *Math. Sci. Res. Inst. Publ.*, 43, 10 2001.
- [9] N. Andruskiewitsch and H.-J. Schneider. On the classification of finite-dimensional pointed Hopf algebras. *Ann. Math.*, 171(1):375–417, 2010.
- [10] I. Angiono. On Nichols algebras of diagonal type. *J. Reine Angew. Math.*, 2013(683):189–251, 2013.
- [11] I. Angiono. A presentation by generators and relations of Nichols algebras of diagonal type and convex orders on root systems. *J. Europ. Math. Soc.*, 17(10):2643–2671, 2015.
- [12] M. Cuntz and I. Heckenberger. Weyl groupoids of rank two and continued fractions. *Algebra Number Theory*, 3(3):317–340, 2009.
- [13] M. Cuntz and I. Heckenberger. Weyl groupoids with at most three objects. *J. Pure Appl. Algebra*, 213(6):1112–1128, 2009.
- [14] M. Cuntz and I. Heckenberger. Finite Weyl groupoids of rank three. *Trans. Amer. Math. Soc.*, 364(3):1369–1393, 2012.
- [15] M. Cuntz and I. Heckenberger. Finite Weyl groupoids. *Reine Angew. Math.*, 2015(702):77–108, 2015.
- [16] I. Heckenberger. Classification of arithmetic root systems of rank 3. *arXiv preprint math/0509145*, 2005.

- [17] I. Heckenberger. Weyl equivalence for rank 2 Nichols algebras of diagonal type. *Ann. Uni. Ferrara*, 51(1):281–289, 2005.
- [18] I. Heckenberger. The Weyl groupoid of a Nichols algebra of diagonal type. *Invent. Math.*, 164(1):175–188, 2006.
- [19] I. Heckenberger. Examples of finite-dimensional rank 2 Nichols algebras of diagonal type. *Compos. Math.*, 143(1):165–190, 2007.
- [20] I. Heckenberger. Rank 2 Nichols algebras with finite arithmetic root system. *Algebr. Represent. Theory*, 11(2):115–132, 2008.
- [21] I. Heckenberger. Classification of arithmetic root systems. *Adv. Math.*, 220(1):59–124, 2009.
- [22] I. Heckenberger, E. Meir, and L. Vendramin. Finite-dimensional nichols algebras of simple yetter–drinfeld modules (over groups) of prime dimension. *Advances in Mathematics*, 444:109637, 2024.
- [23] I. Heckenberger and H.-J. Schneider. Root systems and Weyl groupoids for Nichols algebras. *Proc. Lond. Math. Soc.*, 101(3):623–654, 2010.
- [24] I. Heckenberger and H.-J. Schneider. Right coideal subalgebras of Nichols algebras and the duflo order on the Weyl groupoid. *Israel J. Math.*, 197(1):139–187, 2013.
- [25] I. Heckenberger and H.-J. Schneider. *Hopf algebras and root systems*, volume 247. American Mathematical Soc., 2020.
- [26] I. Heckenberger and J. Wang. Rank 2 Nichols algebras of diagonal type over fields of positive characteristic. *SIGMA Symmetry. Integr. Geom.*, 11:011, 2015.
- [27] I. Heckenberger and H. Yamane. A generalization of Coxeter groups, root systems, and Matsumoto’s theorem. *Math. Z.*, 259(2):255–276, 2008.
- [28] V. Kharchenko. A quantum analog of the Poincare-Birkhoff-Witt theorem. *Algebra Logic*, 38(4):259–276, 1999.
- [29] G. Lusztig. Noncommutative Differentials and Yang-Mills on Permutation Groups  $s_n$ . *Lect. Notes Pure Appl. Math.*, 239, 2001.
- [30] G. Lusztig. *Introduction to quantum groups*. Springer Science & Business Media, 2010.
- [31] W.D. Nichols. Bialgebras of type one. *Commun. Alg.*, 6(15):1521–1552, 1978.
- [32] M. Rosso. Quantum groups and quantum shuffles. *Invent. Math.*, 133(2):399–416, 1998.
- [33] P. Schauenburg. A characterization of the Borel-like subalgebras of quantum enveloping algebras. *Commun. Alg.*, 24(9):2811–2823, 1996.
- [34] A.M Semikhatov. Virasoro central charges for Nichols algebras. In *Conformal Field Theories and Tensor Categories*, pages 67–92. Springer, 2014.

- [35] A.M. Semikhatov and I.Yu. Tipunin. The Nichols algebra of screenings. *Commun. Contemp. Math.*, 14(04):1250029, 2012.
- [36] A.M. Semikhatov and I.Yu. Tipunin. Logarithmic  $sl_2$  CFT models from Nichols algebras: I. *J. Phys.*, 46(49):494011, 2013.
- [37] M. Takeuchi. Survey of braided hopf algebras. *Contemp. Math.*, 267:301–324, 2000.
- [38] J. Wang. Rank three Nichols algebras of diagonal type over arbitrary fields. *Israel J. Math.*, 218(1):1–26, 2017.
- [39] J. Wang. Rank 4 finite-dimensional Nichols algebras of diagonal type in positive characteristic. *J. Algebra*, 559:547–579, 2020.
- [40] S.L. Woronowicz. Compact matrix pseudogroups. *Comm. Math. Phys.*, 111(4):613–665, 1987.
- [41] S.L. Woronowicz. Differential calculus on compact matrix pseudogroups (quantum groups). *Comm. Math. Phys.*, 122(1):125–170, 1989.