Tensor structure on the module category of the triplet superalgebra $\mathcal{SW}(m)$

Hiromu Nakano

Abstract

We discuss the tensor structure on the category of modules of the $N = 1$ triplet vertex operator superalgebra $\mathcal{SW}(m)$ introduced by Adamović and Milas. Based on the theory of vertex tensor supercategories, we determine the structure of fusion products between the simple and projective $\mathcal{SW}(m)$ -modules and show that the tensor supercategory on $\mathcal{SW}(m)$ -mod is rigid. Technically, explicit solutions of a fourth-order Fuchsian differential equation are important to show the rigidity of $\mathcal{SW}(m)$ -modules. We construct solutions of this Fuchsian differential equation using the theory of the Dotsenko-Fateev integrals developed by Sussman.

1 Introduction

In recent years, comprehensive studies of logarithmic vertex operator algebras have developed, where the adjective "logarithmic" comes from the property that L_0 the generator of scale transformations is no longer diagonalizable. In the logarithmic conformal field theories, this non-diagonalizability leads to interesting examples of physical systems such as polymers, spin chains, percolations and sand-pile models [\[CR,](#page-66-0) [GK2,](#page-67-0) [Gu,](#page-68-0) [JPR,](#page-69-0) [MR,](#page-70-0) [Ni,](#page-70-1) [PR,](#page-70-2) [Ri,](#page-70-3) [RS\]](#page-70-4). In the representation theories of logarithmic vertex operator algebras, indecomposable modules appear such that L_0 acts non-semisimply, and this non-semisimplicity made it challenging to formulate the category theories associated with vertex operator algebras.

Among the logarithmic vertex operator algebras, the triplet W-algebra \mathcal{W}_p [\[FGST1,](#page-67-1) [FGST2,](#page-67-2) [FGST3,](#page-67-3) [Ka\]](#page-69-1) is particularly famous and is known for satisfying a cetain finiteness called C_2 -cofinite condition. By the rigorous studies [\[AM1,](#page-65-0) [MY,](#page-70-5) [NT,](#page-70-6) [TW1\]](#page-71-0), the structure of the category of \mathcal{W}_n -modules is completely determined. Furthermore, by [\[CLR,](#page-66-1) [GN\]](#page-68-1), it is shown that the category of W_p -modules is equivalent to a quasi-Hopf modification of the category of finite-dimensional representations for the restricted quantum group $\overline{U}_q(sl_2)$ at $q = e^{\frac{\pi i}{p}}$.

As a natural super analogue of the triplet algebra \mathcal{W}_p , a $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded (logarithmic) vertex operaor superalgebra, usually denoted by $\mathcal{SW}(m)$, was introduced by Adamović and Milas in the paper [\[AM3\]](#page-65-1). This vertex operaor superalgebra $\mathcal{SW}(m)$ is called $N = 1$ triplet vertex operaor superalgebra, and contains conformal and superconformal vectors at the central charge

$$
c_{1,2m+1} = \frac{15}{2} - 3(2m+1+\frac{1}{2m+1}), \quad m \in \mathbb{Z}_{\geq 1}.
$$

Adamović and Milas proved the C_2 -cofiniteness of $\mathcal{SW}(m)$, classified all simple $\mathcal{SW}(m)$ -modules and conjectured the equivalence between the module category of $\mathcal{SW}(m)$ and the category of finite dimensional modules over the small quantum group $U_q^{small}(sl_2)$ at $q = e^{\frac{2\pi i}{2m+1}}$. Furthermore they showed that the characters of the simple $\mathcal{SW}(m)$ -modules can be expressed in terms of the characters of the simple \mathcal{W}_p -modules.

For a C_2 -cofinite, $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded vertex operator superalgebra V, let Vmod be the category of grading-restricted generalized modules. It is shown by Creutzig, Genra, Nakatsuka and Sato [\[CGNS\]](#page-66-2) that the category V -mod admits the structure of the vertex tensor supercategory developed by Huang-Lepowsky-Zhang [\[HLZ1\]](#page-68-2)-[\[HLZ8\]](#page-69-2) and Creutzig-Kanade-McRae [\[CKM\]](#page-66-3). Recently, as a deeper result, Creutzig, McRae, Orosz Hunziker and Yang [\[CMOY\]](#page-66-4) have shown that the category of C_1 -cofinite grading-restricted generalized modules for a vertex operator superalgebra has the structure of the above vertex tensor supercategory when the module category satisfy some appropriate conditions. By these important results, in particular, $\mathcal{SW}(m)$ -mod admits the structure of the vertex tensor supercategory.

When studying the structure of $\mathcal{SW}(m)$ -mod, in addition to general theories of vertex operator superalgebras, the rigidity of modules are also important. In the case of \mathcal{W}_p [\[CMY1,](#page-66-5) [MY,](#page-70-5) [TW1\]](#page-71-0), a second-order Fuchsian differential equation, called BPZ differential equation [\[BPZ\]](#page-65-2), becomes important for examining the rigidity of modules. In the case of $\mathcal{SW}(m)$, the investigation of the rigidity of simple modules leads to a fourth-order Fuchsian differential equation. This Fuchsian differential equation is derived from the four point function with the minimal conformal weight vectors of $N = 1$ super Virasoro simple modules $L(c_{1,2m+1}, h_{2,2})$ inserted, where $h_{2,2}$ is the minimal conformal weight defined by

$$
h_{2,2} = \frac{3}{8}(2m+1) - \frac{3}{4} + \frac{3}{8(2m+1)}.
$$

In this paper, we construct the fundamental system of solutions of this fourthorder Fuchsian differential equation using the theory of Dotsenko-Fateev integrals [\[DF1,](#page-66-6) [DF2\]](#page-66-7) developed by Sussman [\[Su1,](#page-70-7) [Su2\]](#page-70-8), and determine some monodromy properties of the solutions. Those monodromy data are important to show the rigidity of a simple $\mathcal{SW}(m)$ -module whose minimal conformal weight is $h_{2,2}$ (Theorem [5.10\)](#page-52-0).

The following are the main results and goals of this paper:

- We determine the structure of the projective covers of the simple $\mathcal{SW}(m)$ modules (Propositions [5.17](#page-59-0)[-5.18\)](#page-60-0).
- We show that the tensor supercategory on $\mathcal{SW}(m)$ -mod is rigid.
- We determine the structure of fusion products between the simple and projective $\mathcal{SW}(m)$ -modules (Subsection [5.4\)](#page-62-0).

We show the above results in Subsections [5.3-](#page-54-0)[5.4,](#page-62-0) using properties of vertex tensor supercategories and the Dotsenko-Fateev integrals. These results are partially based on our thesis [\[Nak\]](#page-70-9).

Recently, in [\[CMOY\]](#page-66-4), Creutzig, McRae, Orosz Hunziker and Yang obtain important results for the rigidity and the structure of the C_1 -cofinite module category of the $N = 1$ super Virasoro vertex operator superalgebra. They show the rigidity and self-duality of the simple module $L(c_{1,2m+1}, h_{2,2})$ using a certain embedding technique for Virasoro vertex operator algebras. Just as the structure of W_p -mod can be determined from the Virasoro tensor category at central charge $13 - 6p - 6p^{-1}$ $(p \in \mathbb{Z}_{\geq 2})$ [\[MY,](#page-70-5) Section 7], we expect that their results will rederive the rigidity and structure of $\mathcal{SW}(m)$ mod and further develop the theory of the super singlet algebra $\mathcal{SM}(m)$ introduced in [\[AM3\]](#page-65-1).

This paper is organized as follows.

In Section [2,](#page-3-0) we review some facts of the representation theory of $\frac{1}{2}$ $\frac{1}{2} \mathbb{Z}_{\geq 0}$ graded vertex operator superalgebras and $N = 1$ Neveu-Schwarz algebra in accordance with [\[BMRW,](#page-65-3) [CKM,](#page-66-3) [IK1,](#page-69-3) [IK2,](#page-69-4) [KW\]](#page-69-5).

In Section [3,](#page-15-0) we review some important properties for $\mathcal{SW}(m)$ -mod, such as the structure of the simple $\mathcal{SW}(m)$ -modules and the Zhu-algebra $A(SW(m))$ determined by Admović and Milas in [\[AM3,](#page-65-1) [AM4\]](#page-65-4).

In Section [4,](#page-20-0) we introduce a $\mathcal{SW}(m)$ correlation function and examine some monodromy properties of this correlation function. In Subsection [4.1,](#page-20-1) we show that this correlation function satisfies a complicated fourth-order Fuchsian differential equation. In Subsection [4.2,](#page-23-0) we introduce the Dotsenko-Fateev integrals [\[DF1,](#page-66-6) [DF2\]](#page-66-7) and regularization results of these integrals recently established by Sussman [\[Su2\]](#page-70-8). In Subsection [4.3,](#page-32-0) we introduce some

formulas for the Dotsenko-Fateev integrals given by Forrester [\[Fo1,](#page-67-4) [Fo2\]](#page-67-5) and some results for meromorphic continuations of the Dotsenko-Fateev integrals given by Sussman [\[Su1\]](#page-70-7). We also introduce some transformation formulas between Dotsenko-Fateev integrals of special types. In Subsection [4.4,](#page-37-0) we construct the fundamental system of solutions of the Fuchsian differential equation given in Subsection [4.1,](#page-20-1) using the results in Subsection [4.3](#page-32-0) and a certain deformation technique by Tsuchiya and Wood [\[TW2\]](#page-71-1). As a result, we get some monodromy properties of this Fuchsian differential equation.

In Section [5,](#page-43-0) we examine the structure of fusion products and the rigidity of $\mathcal{SW}(m)$ -modules. In Subsection [5.2,](#page-46-0) using the results in Section [4](#page-20-0) and certain vertex tensor categorical techniques introduced by Creutzig-McRae-Yang [\[CMY1,](#page-66-5) [CMY2\]](#page-66-8) and Tsuchiya-Wood [\[TW1\]](#page-71-0), we show that a simple $\mathcal{SW}(m)$ -module X_2 is rigid and self-dual. By using self-duality of the simple module X_2 , we show that all simple and projective modules can be obtained by repeatedly multiplying X_2 . As a result, we can determine the structure of all projective modules and show that the tensor supercategory on $\mathcal{SW}(m)$ -mod is rigid. In Subsection [5.4,](#page-62-0) we introduce a non-semisimple fusion ring $P(\mathcal{SW}(m))$ and determine the ring structure.

2 Basic definitions and notation

The $N = 1$ Neveu-Schwarz algebra is the Lie superalgebra

$$
\mathfrak{ns} = \bigoplus_{n \in \mathbb{Z}} \mathbb{C}L_n \oplus \bigoplus_{r \in \frac{1}{2} + \mathbb{Z}} \mathbb{C}G_r \oplus \bigoplus \mathbb{C}C
$$

with the relations $(k, l \in \mathbb{Z}, r, s \in \mathbb{Z} + \frac{1}{2})$ $(\frac{1}{2})$:

$$
[L_k, L_l] = (k-l)L_{k+l} + \delta_{k+l,0} \frac{k^3 - k}{12}C,
$$

\n
$$
[L_k, G_r] = (\frac{1}{2}k - r)G_{k+r},
$$

\n
$$
\{G_r, G_s\} = 2L_{r+s} + \frac{1}{3}(r^2 - \frac{1}{4})\delta_{r+s,0}C,
$$

\n
$$
[L_k, C] = 0, [G_r, C] = 0,
$$

where $\{ , \}$ is the anti-commutator. We identify C with a scalar multiple of the identity, $C = c \cdot id$, when acting on modules and refer to the number $c \in \mathbb{C}$ as the central charge. In this section, we briefly review basic facts of representation theories of the Neveu-Schwarz algebra.

2.1 Vertex operator superalgebras

In this subsection we briefly review the definitions of $N = 1$ vertex operator superalgebras and the notion such as vertex algebra modules and intertwining operators used in later section. See [\[CKL,](#page-66-9) [CKM,](#page-66-3) [CMOY,](#page-66-4) [FLM,](#page-67-6) [KW\]](#page-69-5) for details.

Let us recall the definition of $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded vertex operator superalgebras.

Definition 2.1. A four pairs $(V, |0\rangle, T, Y)$ is called a $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded vertex operator superalgebra where

1. V is a $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded C-vector space

$$
V = \bigoplus_{n \in \frac{1}{2}\mathbb{Z}_{\geq 0}} V[n].
$$

For $\overline{0}$, $\overline{1} \in \mathbb{Z}/2\mathbb{Z}$, set

$$
V^{\bar{0}} := \bigoplus_{n \in \mathbb{Z}_{\geq 0}} V[n], \qquad V^{\bar{1}} := \bigoplus_{n \in \mathbb{Z}_{\geq 0} + \frac{1}{2}} V[n].
$$

- 2. $|0\rangle \in V[0]$ is called the vacuum vector.
- 3. $T \in V[2]$ is called the conformal vector.
- 4. Y is a C-linear map

$$
Y: V \to \text{End}_{\mathbb{C}}(V)[[z, z^{-1}]].
$$

These data are subject to the following axioms:

- $\dim_{\mathbb{C}} V[0] = 1$ and $0 < \dim_{\mathbb{C}} V[n] < \infty$ for any $n \in \frac{1}{2}$ $rac{1}{2}\mathbb{Z}_{\geq 0}$.
- For each $v \in V^{\bar{i}}[h]$ there exists a field

$$
Y(v, z) = \sum_{n \in \mathbb{Z} + \frac{i}{2}} v_n z^{-n-h}.
$$

• $Y(|0\rangle, z) = id_V$ and

$$
Y(v, z)|0\rangle - v \in V[[z]]z
$$

for all $v \in V$.

• The modes of the field $Y(T, z) = T(z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2}$ define the commutation relations of the Virasoro algebra with fixed central charge $c = c_V$:

$$
[L_k, L_l] = (k-l)L_{k+l} + \delta_{k+l,0} \frac{k^3 - k}{12} c_V.
$$

• The zero mode L_0 of $T(z)$ acts semisimply on V and

$$
V[h] = \{ v \in V \mid L_0 v = hv \}.
$$

• For all $v \in V$

$$
Y(L_{-1}v, z) = \frac{\mathrm{d}}{\mathrm{d}z}Y(v, z).
$$

• For $v_1 \in V^{\bar{i}}$ and $v_2 \in V^{\bar{j}}$, the following super Jacobi identity holds $z_0^{-1}\delta\left(\frac{z_1-z_2}{z_1-z_2}\right)$ z_0 $Y(v_1, z_1)Y(v_2, z_2) - (-1)^{ij}z_0^{-1}\delta\left(\frac{z_2 - z_1}{z_2}\right)$ $-z_0$ $Y(v_2, z_2)Y(v_1, z_1)$ $= z_2^{-1} \delta \left(\frac{z_1 - z_0}{z_1 - z_0} \right)$ z_2 $Y(Y(v_1, z_0)v_2, z_2).$

In the above defintion, we call $V^{\bar{0}}$ the *even part* of V and $V^{\bar{1}}$ the *odd part* of V. For any $v \in V^{\bar{i}} (i = 1, 0)$, we call v parity-homogeneous vector in V, and we denote by $|v| = i$ the parity of v.

The $N = 1$ Neveu-Schwarz vertex operator superalgebras are special cases of the $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded vertex operator superalgebras, which are subject to an additional axiom:

There exists $G \in V[\frac{3}{2}]$ $\frac{3}{2}$ (super conformal vector) such that the modes of fields

$$
Y(T, z) = T(z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2}, \qquad Y(G, z) = G(z) = \sum_{r \in \mathbb{Z} + \frac{1}{2}} G_r z^{-r - \frac{3}{2}},
$$

define the commutation relations of the Neveu-Schwarz algebra with fixed central charge $c = c_V$:

$$
[L_k, L_l] = (k - l)L_{k+l} + \delta_{k+l,0} \frac{k^3 - k}{12} c_V,
$$

\n
$$
[L_k, G_r] = (\frac{1}{2}k - r)G_{m+r},
$$

\n
$$
\{G_r, G_s\} = 2L_{r+s} + \frac{1}{3}(r^2 - \frac{1}{4})\delta_{r+s,0}c_V.
$$
\n(2.1)

Let us recall the definition of modules of $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded vertex operator superalgebras.

Definition 2.2. Given a $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded vertex operator superalgebra $(V, |0\rangle, T, G, Y)$, a grading restricted generalised V-module is a pair (M, Y_M) of a vector space M and a linear map Y_M from V to $\text{End}M[[z,z^{-1}]]$ satisfying the following conditions

1. $Y_M(|0\rangle, z) = \text{Id}_M$ and the modes of

$$
Y_M(T, z) = \sum_{n \in \mathbb{Z}} L_n^M z^{-n-2}
$$

satisfy the commutation relations of the Virasoro algebra with the central charge c_V .

2. For all $v \in V$,

$$
Y_M(L_{-1}v, z) = \frac{\mathrm{d}}{\mathrm{d}z} Y_M(v, z).
$$

3. For $v_1 \in V^{\bar{i}}$ and $v_2 \in V^{\bar{j}}$, the following super Jacobi identity holds

$$
z_0^{-1}\delta\left(\frac{z_1 - z_2}{z_0}\right)Y_M(v_1, z_1)Y_M(v_2, z_2)
$$

$$
-(-1)^{ij}z_0^{-1}\delta\left(\frac{z_2 - z_1}{z_0}\right)Y_M(v_2, z_2)Y_M(v_1, z_1)
$$

\n
$$
= z_2^{-1}\delta\left(\frac{z_1 - z_0}{z_2}\right)Y_M(Y(v_1, z_0)v_2, z_2),
$$

where $\delta(z)$ is the formal delta function $\delta(z) = \sum_{n \in \mathbb{Z}} z^n$.

4. M is a C-graded superspace

$$
M = \bigoplus_{\bar{i} \in \mathbb{Z}/2\mathbb{Z}} M^{\bar{i}} = \bigoplus_{h \in H(M)} M[h]
$$

such that

- For some finite subset $H_0(M)$ of \mathbb{C} , $H(M) = H_0(M) + \frac{1}{2}\mathbb{Z}_{\geq 0}$.
- For $h \in H(M)$, $M[h] = \{ \psi \in M : \exists n \ge 0 \text{ s.t. } (L_0 h)^n \psi = 0 \}.$
- $0 < \dim_{\mathbb{C}} M[h] < \infty$.
- For all $v \in V$, $v_h M[h'] \subset M[h+h']$.
- For $\bar{i} = \bar{0}, \bar{1}, M^{\bar{i}} = \bigoplus_{h \in H(M)} M^{\bar{i}}[h]$, where $M^{\bar{i}}[h] = M^{\bar{i}} \cap M[h]$.
- For $v \in V^{\bar{i}}$ and $\psi \in M^{\bar{j}}$ $(\bar{i}, \bar{j} \in \mathbb{Z}/2\mathbb{Z})$, $Y_M(v; z)\psi \in M^{\bar{i}+\bar{j}}[[z, z^{-1}]]$.

In the above defintion, we call $M^{\bar{0}}$ the *even part* of M and $M^{\bar{1}}$ the *odd* part of M. For any $\psi \in M^{\bar{i}} (i = 1, 0)$, we call ψ parity-homogeneous vector in M, and we denote by $|\psi| = i$ the parity of ψ .

Given an $N = 1$ Neveu-Schwarz vertex operator superalgebra V, a grading restricted generalised V-module M is a special case of Definition [2.2,](#page-6-0) which is subject to an additional axiom: The modes of

$$
Y_M(T, z) = \sum_{n \in \mathbb{Z}} L_n^M z^{-n-2} \qquad Y_M(G, z) = \sum_{r \in \mathbb{Z} + \frac{1}{2}} G_r^M z^{-r - \frac{3}{2}}
$$

satisfy the commutation relations of the Neveu-Schwarz algebra with the central charge c_V .

Analogous to non-super cases, contragredient modules and intertwining operators can be defined as follows.

Definition 2.3. Let V be a $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded vertex operator superalgebra and M be a a grading restricted generalised V -module. Let

$$
M^* = \bigoplus_{h \in H(M)} M^*[h]
$$

be the graded dual space of M, where $M^*[h] = \text{Hom}_{\mathbb{C}}(M[h], \mathbb{C})$ with parity decomposition

$$
(M^*)^{\overline{i}} = \bigoplus_{h \in H(M)} (M^*[h])^{\overline{i}}, \qquad (M^*[h])^{\overline{i}} = \text{Hom}_{\mathbb{C}}(M^{\overline{i}}[h], \mathbb{C}).
$$

Let \langle , \rangle be the natural dual pairing between M^* and M. Then we define the V-module structure Y_{M^*} as follows

$$
\langle Y_{M^*}(v,z)\psi^*,\psi\rangle := (-1)^{ij}\langle \psi^*, Y_M(e^{zL_1}(-z^{-2})^{L_0}v,z^{-1})\psi\rangle, \qquad (2.2)
$$

where $\psi^* \in (M^*)^{\overline{i}}, \psi \in M$ and $v \in V^{\overline{j}},$ for $\overline{i}, \overline{j} \in \mathbb{Z}/2\mathbb{Z}$.

Definition 2.4. Let V be a a $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded vertex operator superalgebra and M_1 , M_2 and M_3 a triple of V-module. Denote by $M_3\{z\}$ [logz] the space of formal power series in z and $\log z$ with coefficient in M_3 , where the exponents of z can be arbitrary complex numbers and with only finitely many $\log z$ terms. A parity-homogeneous intertwining operator $\mathcal{Y}(\cdot, z)$ of type $\begin{pmatrix} M_3 \\ M_1 \end{pmatrix}$ M_1 M_2 $\overline{ }$ is a parity-homogeneous linear map

$$
\mathcal{Y}: M_1 \to \text{End}(M_2, M_3) \{z\} [\log z],
$$

$$
\psi_1 \mapsto \mathcal{Y}(\psi_1, z) = \sum_{t \in \mathbb{C}} \sum_{s \ge 0} (\psi_1)_{t,s}^{\mathcal{Y}} z^{-t-1} (\log z)^s
$$

satisfying the following conditions for parity-homogeneous vectors $\psi_1 \in M_1$, $\psi_2 \in M_2$ and $v \in V$:

- 1. $\mathcal{Y}(L_{-1}\psi_1, z) = \frac{d}{dz}\mathcal{Y}(\psi_1, z)$.
- 2. $(\psi_1)_{t,s}^{\mathcal{Y}} \psi_2 = 0$ for Re(t) sufficiently large.
- 3. The following super Jacobi identity holds

$$
(-1)^{|v||\mathcal{Y}|}z_0^{-1}\delta\left(\frac{z_1-z_2}{z_0}\right)Y_{M_3}(v,z_1)\mathcal{Y}(\psi_1,z_2)
$$

$$
-(-1)^{|v||\psi_1|}z_0^{-1}\delta\left(\frac{z_2-z_1}{-z_0}\right)\mathcal{Y}(\psi_1,z_2)Y_{M_2}(v,z_1)
$$

$$
=z_2^{-1}\delta\left(\frac{z_1-z_0}{z_2}\right)\mathcal{Y}(Y_{M_1}(v,z_0)\psi_1,z_2).
$$
 (2.3)

A general intertwining operator of type $\begin{pmatrix} M_3 \\ M_1 \end{pmatrix}$ M_1 M_2 $\big)$ is a sum of parity-homogeneous ones.

Given a $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded vertex operator superalgebra V and V-modules M_1, M_2, M_3 , we denote by $I_V \begin{pmatrix} M_3 \\ M_2 \end{pmatrix}$ M_2 M_1) the vector superspace of the intertwining operators of type $\begin{pmatrix} \tilde{M}_3 \\ M_{3} \end{pmatrix}$ M_2 M_1 . This vector superspace has the parity decomposition

$$
I_V\begin{pmatrix}M_3\\ M_2\;\;M_1\end{pmatrix}=I_V^0\begin{pmatrix}M_3\\ M_2\;\;M_1\end{pmatrix}\oplus I_V^1\begin{pmatrix}M_3\\ M_2\;\;M_1\end{pmatrix},
$$

where I_V^0 and I_V^1 are the parity-homogeneous subspaces of I_V whose parity are even and odd, respectively.

2.2 Free field realization of the Neveu-Scwarz algebra

In this subsection, we review the free field realization of the Neveu-Scwarz algebra in accordance with the papers [\[BMRW,](#page-65-3) [IK1,](#page-69-3) [IK2\]](#page-69-4).

First, let us introduce the bosonic Fock modules and the bosonic vertex operators. The Heisenberg algebra is the Lie algebra

$$
\mathfrak{h}=\bigoplus_{n\in\mathbb{Z}}\mathbb{C} a_n\oplus\mathbb{C}\mathbf{1},
$$

with commutation relations:

$$
[a_k, a_l] = k\delta_{k+l,0} \mathbf{1}, \qquad [a_k, \mathbf{1}] = 0 \quad (k, l \in \mathbb{Z}).
$$

The Heisenberg algebra $\mathfrak h$ has the triangular decomposition

$$
\mathfrak{h}^{\pm} = \bigoplus_{n>0} \mathbb{C}a_{\pm}, \qquad \qquad \mathfrak{h}^0 = \mathbb{C}a_0 \oplus \mathbb{C}1.
$$

For $\beta \in \mathbb{C}$, let $\mathbb{C}|\beta; B\rangle$ be the one dimensional representation of $\mathfrak{h}^{\ge} = \mathfrak{h}^+ \oplus \mathfrak{h}^0$, which satisfies

$$
a_0|\beta; B\rangle = \beta|\beta; B\rangle, \quad 1|\beta; B\rangle = |\beta; B\rangle, \quad \mathfrak{h}^+|\beta; B\rangle = 0.
$$

Definition 2.5. The bosonic Fock module is defined by induced representation

$$
F_{\beta}^{\mathrm{B}} = \mathrm{Ind}_{\mathfrak{h}^{\geq}}^{\mathfrak{h}} \mathbb{C} | \beta; \mathrm{B} \rangle.
$$

Let $a(z) = \sum_{n \in \mathbb{Z}} a_n z^{-n-1}$ and we define the following bosonic energymomentum tensor

$$
T^{(\text{B};\alpha)}(z) = \frac{1}{2}(:a(z)^{2}: +\alpha \partial a(z)) = \sum_{n \in \mathbb{Z}} L_{n}^{(\text{B};\alpha)} z^{-n-2}
$$

where : : is the normal ordered product. The modes $\{L_n^{(B;\alpha)}\}_{n\in\mathbb{Z}}$ generate the Virasoso algebra with the central charge fixed to $1-3\alpha^2$. By the energymomentum tensor $T^{(\text{B};\alpha)}(z)$, each bosonic Fock module $F^{\text{B}}_{\beta}(\beta \in \mathbb{C})$ becomes a Virasoro module with

$$
L_0^{(\mathbf{B};\alpha)}|\beta;\mathbf{B}\rangle = h_{\beta}|\beta;\mathbf{B}\rangle, \qquad C|\beta;\mathbf{B}\rangle = (1 - 3\alpha^2)|\beta;\mathbf{B}\rangle \tag{2.4}
$$

where $h_{\beta} = \frac{1}{2}$ $\frac{1}{2}\beta(\beta-\alpha)$. Set $T^{(\mathbf{B};\alpha)} := \frac{1}{2}(a_{-1}^2 + \alpha a_{-2})|0;\mathbf{B}\rangle$. The Fock module F_0^{B} carries the structure of a vertex operator algebra, with

$$
Y(|0; B\rangle, z) = id
$$
, $Y(a_{-1}|0; B\rangle, z) = a(z)$, $Y(T^{(B;\alpha)}, z) = T^{(B;\alpha)}(z)$.

We denote by $\mathcal{F}_{\alpha}^{\text{B}}$ this vertex operator algebra.

We extend the Heisenberg algebra $\mathfrak h$ by a generator $\hat a$ satisfying the relations

$$
[a_n, \hat{a}] = \delta_{n,0}, \qquad [\hat{a}, 1] = 0.
$$

For $\beta \in \mathbb{C}$, we define the following vertex operator

$$
V_{\beta}(z) = e^{\beta \hat{a}} z^{\beta a_0} \prod_{n \ge 1} \left[\exp \left(\beta \frac{a_{-n}}{n} z^n \right) \exp \left(-\beta \frac{a_n}{n} z^{-n} \right) \right].
$$

The composition of k vertex operators is given by (cf. [\[BMRW\]](#page-65-3))

$$
V_{\beta_1}(z_1)\cdots V_{\beta_k}(z_k) = e^{\sum_{i=1}^k \beta_i \hat{a}} \prod_{1 \le i \le j \le k} (z_i - z_j)^{\beta_i \beta_j} \prod_{i=1}^k z_i^{\beta_i a_0}
$$

$$
\cdot \prod_{n \ge 1} \left[\exp\left(\frac{a_{-n}}{n} \sum_{i=1}^k \beta_i z_i^n \right) \exp\left(-\frac{a_n}{n} \sum_{i=1}^k \beta_i z_i^{-n}\right) \right].
$$
\n(2.5)

By identifying $e^{\beta \hat{a}}|\gamma; B\rangle = |\beta + \gamma; B\rangle$, $V_{\beta}(z)$ becomes a linear map

$$
V_{\beta}(z): F_{\gamma} \to F_{\beta+\gamma}[[z, z^{-1}]] z^{\beta\gamma}
$$

such that

$$
\langle \beta + \gamma; B | V_{\beta}(z) | \gamma; B \rangle = z^{\beta \gamma}, \qquad (2.6)
$$

where $\langle \beta; \mathbf{B} |$ is the bra vector of $(F^{\mathbf{B}}_{\beta})^*$.

Since all F^{B}_{β} ($\beta \in \mathbb{C}$) are simple $\mathcal{F}^{\text{B}}_{\alpha}$ -modules, we have $\dim_{\mathbb{C}}I_{\mathcal{F}^{\text{B}}_{\alpha}}$ $\begin{cases} F^{\text{B}}_{\beta+\beta'} \\ F^{\text{B}}_{\beta} \hspace{0.1cm} F^{\text{B}}_{\beta'} \end{cases}$ $=$ 1, and $\dim_{\mathbb{C}}I_{\mathcal{F}_{\alpha}^{\mathbf{B}}}$ $\begin{pmatrix} F^{\rm B}_\gamma\\ F^{\rm B}_\beta\;\;F^{\rm B}_{\beta'} \end{pmatrix}$ $= 0$ for $\gamma \neq \beta + \beta'$. For any non-zero intertwining operator $I_{\beta,\beta'}^{\text{B}} \in I_{\mathcal{F}_{\alpha}^{\text{B}}}$ $\begin{pmatrix} F^{\text{B}}_{\beta+\beta'} \\ F^{\text{B}}_{\beta} & F^{\text{B}}_{\beta'} \end{pmatrix}$), $I_{\beta,\beta'}(|\beta\rangle, z)$ is equal to $V_{\beta}(z)$, up to scalar multiples.

Next, let us introduce the Neveu-Schwarz fermionic Fock module and some notation related to it. The Neveu-Schwarz fermion algebra f is the Lie superalgebra

$$
\mathfrak{f}=\bigoplus_{r\in\mathbb{Z}+\frac{1}{2}}\mathbb{C}b_r\oplus\mathbb{C}\mathbf{1}
$$

with anti-commutation relations ${b_r, b_s} = \delta_{r+s,0}, {b_r, 1} = 0$. The Neveu-Schwarz fermion algebra f has the triangular decomposition

$$
\mathfrak{f}^{\pm}=\bigoplus_{r>0}\mathbb{C} b_r,\quad \mathfrak{f}^0=\mathbb{C}\mathbf{1}.
$$

Let $\mathbb{C}|\text{NS}\rangle$ be the one dimensional representation of $\mathfrak{f}^{\ge} = \mathfrak{f}^+ \oplus \mathfrak{f}^0$ defined by

$$
1|NS\rangle = |NS\rangle, \quad f^+|NS\rangle = 0.
$$

Definition 2.6. The Neveu-Schwarz fermionic Fock module F^{NS} is defined by

$$
F^{\text{NS}} = \text{Ind}_{\mathfrak{f}^{\geq}}^{\mathfrak{f}} \mathbb{C}|\text{NS}\rangle.
$$

Let $b(z) = \sum_{n \in \mathbb{Z}+\frac{1}{2}} b_n z^{-n-\frac{1}{2}}$. Then this field satisfies the operator product expansion

$$
b(z)b(w) = \frac{1}{z - w} + \dotsb, \tag{2.7}
$$

where " \cdots " denote the holomorphic parts about $z = w$. We define the following energy-momentum tensor

$$
T^{(\mathfrak{f})}(z) = \frac{1}{2} : \partial b(z)b(z) := \sum_{n \in \mathbb{Z}} L_n^{(\mathfrak{f})} z^{-n-2}.
$$

The modes $\{L_n^{(f)}\}_{n\in\mathbb{Z}}$ generate the Virasoso algebra with the central charge fixed to $\frac{1}{2}$. By the energy-momentum tensor $T^{(\mathfrak{f})}(z)$, the Neveu-Schwarz fermionic Fock module F^{NS} becomes a Virasoro module with

$$
L_0^{(\dagger)}|NS\rangle = 0
$$
, $C|NS\rangle = \frac{1}{2}|NS\rangle$.

We introduce an even field and an odd field

$$
T^{(\alpha)}(z) = T^{(\text{B};\alpha)}(z) \otimes \mathbf{1} + \mathbf{1} \otimes T^{(\text{f})}(z),
$$

\n
$$
G^{(\alpha)}(z) = a(z) \otimes b(z) + \alpha \mathbf{1} \otimes \partial b(z).
$$
\n(2.8)

We see that $T^{(\alpha)}(z)$ and $G^{(\alpha)}(z)$ satisfy the operator product expansions

$$
T^{(\alpha)}(z)T^{(\alpha)}(w) = \frac{c_{\alpha}/2}{(z-w)^4} + \frac{2T^{(\alpha)}(w)}{(z-w)^3} + \frac{\partial T^{(\alpha)}(w)}{z-w} + \cdots,
$$

\n
$$
T^{(\alpha)}(z)G^{(\alpha)}(w) = \frac{\frac{3}{2}G^{(\alpha)}(w)}{(z-w)^2} + \frac{\partial G^{(\alpha)}(w)}{z-w} + \cdots,
$$

\n
$$
G^{(\alpha)}(z)G^{(\alpha)}(w) = \frac{2c_{\alpha}/3}{(z-w)^3} + \frac{2T^{(\alpha)}(w)}{z-w} + \cdots,
$$
\n(2.9)

where $c_{\alpha} := \frac{3}{2} - 3\alpha^2$. For the Fourier mode expansions of fields

$$
T^{(\alpha)}(z) := \sum_{n \in \mathbb{Z}} L_n^{(\alpha)} z^{-n-2}, \quad G^{(\alpha)}(w) := \sum_{r \in \mathbb{Z} + \frac{1}{2}} G_r^{(\alpha)} z^{-r - \frac{3}{2}}, \quad (2.10)
$$

the modes $\{L_n^{(\alpha)}\}$ and $\{G_r^{(\alpha)}\}$ define the following commutation and anticommutation relations

$$
[L_k^{(\alpha)}, L_l^{(\alpha)}] = (k-l)L_{k+l}^{(\alpha)} + \delta_{k+l,0} \frac{k^3 - k}{12} c_{\alpha},
$$

$$
[L_k^{(\alpha)}, G_r^{(\alpha)}] = (\frac{1}{2}k - r)G_{k+r}^{(\alpha)},
$$

$$
\{G_r^{(\alpha)}, G_s^{(\alpha)}\} = 2L_{r+s}^{(\alpha)} + \frac{1}{3}(r^2 - \frac{1}{4})\delta_{r+s,0}c_{\alpha}.
$$

Thus the modes of the fields $T^{(\alpha)}(z)$ and $G^{(\alpha)}(z)$ generate the Neveu-Schwarz algebra with the central charge fixed to $c_{\alpha} = \frac{3}{2} - 3\alpha^2$.

Definition 2.7. For $\beta \in \mathbb{C}$, we set

$$
F_{\beta} := F_{\beta}^{\mathcal{B}} \otimes F^{\mathcal{NS}}
$$

and call this tensor product Fock module simply.

We set

$$
|\beta\rangle = |\beta; B\rangle \otimes |NS\rangle.
$$

In the following, we omit the tensor product in $a_n \otimes \mathbf{1}$ and $\mathbf{1} \otimes b_r$ and simply denote them as a_n and b_r . We also use the following shorthand notation

$$
V_{\beta}(z) = Y(|\beta\rangle, z) = Y(|\beta; \mathbf{B}\rangle, z) \otimes Y(|\mathbf{NS}\rangle, z).
$$

We define the following two vectors in F_0

$$
T^{(\alpha)} = \frac{1}{2}(a_{-1}^2 + \alpha a_{-2} + b_{-\frac{1}{2}}b_{-\frac{3}{2}})|0\rangle, \qquad G^{(\alpha)} = (a_{-1}b_{-\frac{1}{2}} + \alpha b_{-\frac{3}{2}})|0\rangle.
$$

The Fock module F_0 carries the structure of an $N = 1$ Neveu-Schwarz vertex operator superalgebra, with

$$
Y(|0\rangle, z) = id
$$
, $Y(a_{-1}|0\rangle, z) = a(z)$, $Y(b_{-\frac{1}{2}}|0\rangle, z) = b(z)$,
\n $Y(G^{(\alpha)}, z) = G^{(\alpha)}(z)$, $Y(T^{(\alpha)}, z) = T^{(\alpha)}(z)$.

We denote by \mathcal{F}_{α} this vertex operator superalgebra. We use the shorthand notation $T(z) = T^{(\alpha)}(z)$, $G(z) = G^{(\alpha)}(z)$, $L_n^{(\alpha)}$ and $G_r^{(\alpha)}$, unless otherwise stated.

Before introducing the structure of Fock modules, let us review the construction of the intertwining operators between Fock modules. Given $\alpha, \beta, \beta' \in$ $\mathbb C$ and $\gamma \in \mathbb C$ satisfying $\gamma \neq \beta + \beta'$, we have

$$
\dim I_{\mathcal{F}_{\alpha}}^{0}\left(\begin{matrix}F_{\beta+\beta'}\\F_{\beta}&F_{\beta'}\end{matrix}\right)=1, \quad \dim I_{\mathcal{F}_{\alpha}}^{1}\left(\begin{matrix}F_{\beta+\beta'}\\F_{\beta}&F_{\beta'}\end{matrix}\right)=1, \quad \dim I_{\mathcal{F}_{\alpha}}\left(\begin{matrix}F_{\gamma}\\F_{\beta}&F_{\beta'}\end{matrix}\right)=0.
$$

For $I_{\beta,\beta'}^0 \in I_{\mathcal{F}_{\alpha}}^0$ $\int F_{\beta+\beta'}$ F_β $F_{\beta'}$ \setminus $\setminus \{0\}$ and $I_{\beta,\beta'}^1 \in I_{\mathcal{F}_{\alpha}}^1$ $\int F_{\beta+\beta'}$ F_{β} $F_{\beta'}$ $\overline{}$ $\setminus \{0\}$, we have $I^0_{\beta,\beta'}(|\beta\rangle,z) = V_{\beta}(z)$ and $I^1_{\beta,\beta'}(|\beta\rangle,z) = b(z)V_{\beta}(z)$ up to scalar multiples. Thus the non-trivial \mathcal{F}_{α} -intertwining operators of type $\begin{pmatrix} F_{\beta+\beta'} \\ F_{\beta} & F_{\beta} \end{pmatrix}$ F_β $F_{\beta'}$ $\big)$ can be obtained by the tensor product of the bosonic intertwining operators of type $\begin{cases} F^{\textrm{B}}_{\beta+\beta'} \\ F^{\textrm{B}}_{\beta} \hspace{0.1cm} F^{\textrm{B}}_{\beta'} \end{cases}$ and the Neveu-Schwarz fermion vertex operator.

2.3 Structure of Fock modules

Let $m \in \mathbb{Z}_{\geq 1}$. In this subsection we review the structure of Fock modules whose central charges are

$$
c = c_{1,2m+1} = \frac{15}{2} - 3(2m + 1 + \frac{1}{2m + 1})
$$

in accordance with the papers [\[BMRW,](#page-65-3) [IK1,](#page-69-3) [IK2\]](#page-69-4).

We set

$$
\alpha_{+} = \sqrt{2m+1}, \qquad \alpha_{-} = -\sqrt{\frac{1}{2m+1}}, \qquad \alpha_{0} = \alpha_{+} + \alpha_{-}.
$$
\n(2.11)

Note that $c_{\alpha_0} = \frac{3}{2} - 3\alpha_0^2 = c_{1,2m+1}$. For $r, s, n \in \mathbb{Z}$, we set

$$
\beta_{r,s;n} := \frac{1-r}{2}\alpha_+ + \frac{1-s}{2}\alpha_- + \frac{n}{2}\alpha_+, \qquad \beta_{r,s} = \beta_{r,s;0}, \qquad (2.12)
$$

and we use the shorthand notation $F_{r,s;n} = F_{\beta_{r,s;n}}$ and $F_{r,s} = F_{\beta_{r,s}}$. For $r, s, n \in \mathbb{Z}$, we introduce the notation

$$
h_{r,s} := h_{\beta_{r,s}} = \frac{1}{8}(r^2 - 1)(2m + 1) - \frac{1}{4}(rs - 1) + \frac{1}{8}(s^2 - 1)\frac{1}{2m + 1},
$$

$$
h_{r,s;n} := h_{\beta_{r,s;n}} = h_{r-n,s} = h_{r,s+(2m+1)n}
$$

and denote by $L(h)$ the simple **ns**-module whose minimal conformal weight and central charge are h and $c_{1,2m+1}$.

Before describing the structure of Fock modules, let us introduce the notion of socle series.

Definition 2.8. Let V be a vertex operator superalgebra or the **ns** algebra, and let M be a finite length V-module. We denote by $Soc(M)$ the socle of M, that is $Soc(M)$ is the maximal semisimple submodule of M. Then we have the sequence of the submodules

$$
Soc_1(M) \subsetneq Soc_2(M) \subsetneq \cdots \subsetneq Soc_n(M) = M
$$

such that $Soc_1(M) = Soc(M)$ and $Soc_{i+1}(M)/Soc_i(M) = Soc(M/Soc_i(M)).$ We call such a sequence of the submodules of M the socle series of M.

Proposition 2.9 ([\[IK2\]](#page-69-4)). For $(r, s) \in \mathbb{Z}^2$ such that $r - s \in 2\mathbb{Z}$, the Fock modules $F_{r,s} \in \mathcal{F}_{\alpha_0}$ -mod have the following socle series as the **ns**-modules:

1. For each $F_{1,s;n}$ $(1 \leq s < 2m+1, n \in \mathbb{Z}, s-n \in 2\mathbb{Z}+1)$, we have

$$
Soc(F_{1,s;n}) = \bigoplus_{k \ge 0} L(h_{1,2m+1-s;|n|+2k+1}),
$$

$$
F_{1,s;n}/Soc(F_{1,s;n}) = \bigoplus_{k \ge a} L(h_{1,s;|n|+2k}),
$$

where $a = 0$ if $n \ge 0$, $a = 1$ if $n < 0$.

2. For each $F_{1,2m+1;2n}(n \in \mathbb{Z})$, we have

$$
Soc(F_{1,2m+1;2n}) = F_{1,2m+1;2n} = \bigoplus_{k \ge 0} L(h_{1,2m+1;|2n|+2k}).
$$

We introduce the following two fields

$$
Q_{+}(z) := b(z)V_{\alpha_{+}}(z), \qquad Q_{-}(z) := b(z)V_{\alpha_{-}}(z). \qquad (2.13)
$$

These fields are the so-called screening currents, which satisfy the operator product expansions

$$
T(z)Q_{\pm}(w) = \partial_w \frac{Q_{\pm}(w)}{z - w} + \cdots, \quad G(z)Q_{\pm}(w) = \frac{1}{\alpha_{\pm}} \partial_w \frac{V_{\alpha_{\pm}}(w)}{z - w} + \cdots.
$$
\n(2.14)

By [\(2.14\)](#page-14-0), the operators

$$
Q_{+} := \oint_{z=0} Q_{+}(z)dz : F_{1,2k+1} \to F_{-1,2k+1} \quad (k \in \mathbb{Z}),
$$

$$
Q_{-} := \oint_{z=0} Q_{-}(z)dz : F_{2k+1,1} \to F_{2k+1,-1} \quad (k \in \mathbb{Z})
$$

become commutative with the **ns**-action of \mathcal{F}_{α_0} -Mod. These zero-modes of $Q_{\pm}(z)$ are called screening operators.

We define fields

$$
Q_{-}^{[s]}(z): F_{s+2k,s} \to F_{s+2k,-s}[[z,z^{-1}]] \quad (s \geq 2, \ k \in \mathbb{Z}),
$$

as follows

$$
Q_{-}^{[s]}(z) = \int_{\Gamma_s} Q_{-}(z) Q_{-}(zx_1) Q_{-}(zx_2) \cdots Q_{-}(zx_{s-1}) z^{s-1} dx_1 \cdots dx_{s-1},
$$

where Γ_s are certain regularized cycles constructed from the simplexes

$$
\Delta_{s-1} = \{ (x_1, \ldots, x_{s-1}) \in \mathbb{R}^{s-1} \mid 1 > x_1 > \cdots > x_{s-1} > 0 \}
$$

(see [\[TK\]](#page-71-2) for the detatiled construction of the cycles Γ_s). Then by the results in [\[IK2,](#page-69-4) [TK\]](#page-71-2), the zero-modes

$$
Q_-^{[s]} := \oint_{z=0} Q_-^{[s]}(z) dz : F_{s+2k,s} \to F_{s+2k,-s} \quad (s \ge 2, \ k \in \mathbb{Z})
$$

are non trivial and commutative with the **ns**-action of \mathcal{F}_{α_0} -Mod. These fields $Q_-^{[s]}(z)$ are called screening currents and the zero-modes $Q_-^{[s]}$ are called screening operators.

We set $Q_{-}^{[1]} := Q_{-}$. The structure of the kernels of the screening operators $Q_{-}^{[s]}$ is given by the following proposition.

Proposition 2.10 ([\[IK2\]](#page-69-4)). For any $1 \leq s \leq 2m$ and $n \in \mathbb{Z}$ such that $s - n$ is odd, let

$$
K_{s;n} = \ker Q_{-}^{[s]} : F_{1,s;n} \to F_{1,-s;n}.
$$

Then we have $K_{s;n} = \text{Soc}(F_{1,s;n}).$

3 The abelian category $\mathcal{SW}(m)$ -mod

In this section, we introduce the $N = 1$ triplet vertex operator superalgebra $\mathcal{SW}(m)$ and review some important results for the abelian category of untwisted $\mathcal{SW}(m)$ -modules given in [\[AM3\]](#page-65-1) (for the twisted sector, see [\[AM2\]](#page-65-5)).

3.1 The triplet vertex operator superalgebra $\mathcal{SW}(m)$

Let $m \in \mathbb{Z}_{\geq 1}$. Let $L = \mathbb{Z}\alpha_+ = \mathbb{Z}\sqrt{2m+1}$ be an integral lattice.

Definition 3.1. The lattice vertex operator superalgebra V_L is the quadruple

$$
\left(\bigoplus_{\beta\in L}F_{\beta}, |0\rangle, T, G, Y\right)
$$

where the fields corresponding to $|0\rangle$, $a_{-1}|0\rangle$, $b_{-\frac{1}{2}}|0\rangle$, T and G are those of \mathcal{F}_{α_0} and $Y(|\beta\rangle, z) = V_{\beta}(z)$ $(\beta \in L).$

For each $i \in \mathbb{Z}$, we introduce the following symbol

$$
\gamma_i = \frac{i}{2m+1}\alpha_+ = -i\alpha_-.
$$

It is a known fact that simple \mathcal{V}_L -modules are given by

$$
\mathcal{V}_{L+\gamma_i} := \bigoplus_{n \in \mathbb{Z}} F_{\beta_{1,1;2n} + \gamma_i} = \bigoplus_{n \in \mathbb{Z}} F_{1,1+2i;2n} \quad (i = 0, \dots, 2m). \tag{3.1}
$$

We define $2m + 1$ vector spaces X_s $(1 \leq s \leq 2m + 1)$ as follows:

1. For each $i \in \{0, \ldots, m-1\}$, we define

$$
X_{2i+1} := \ker \left. Q_{-}^{[2i+1]} \right|_{\mathcal{V}_{L+\gamma_i}}, \quad X_{2(m-i)} := \ker \left. Q_{-}^{[2(m-i)]} \right|_{\mathcal{V}_{L+\gamma_{2m-i}}}.
$$
 (3.2)

2. For $s = 2m + 1$, we define $X_{2m+1} := \mathcal{V}_{L+\gamma_m}$.

By Propositions [2.9-](#page-13-0)[2.10,](#page-15-1) X_s satisfy the following decomposition as nsmodules

$$
X_{2i+1} \simeq \bigoplus_{n \in \mathbb{Z}_{\geq 0}} (2n+1)L(h_{1,2i+1;-2n}) \quad (i = 0, \dots, m),
$$

$$
X_{2(m-j)} \simeq \bigoplus_{n \in \mathbb{Z}_{\geq 1}} (2n)L(h_{1,2(m-j);-2n+1}) \quad (j = 0, \dots, m-1).
$$
 (3.3)

Proposition 3.2 ([\[AM3\]](#page-65-1)). Let $\mathcal{SW}(m) = X_1$. Then $\mathcal{SW}(m)$ has the structure of an $N = 1$ Neveu-Schwarz vertex operator superalgebra.

This vertex operator superalgebra is called $N = 1$ triplet vertex operator superalgebra or $N = 1$ triplet superalgebra.

3.2 Simple $\mathcal{SW}(m)$ -modules

Proposition 3.3 ([\[AM3\]](#page-65-1)).

- 1. The vector spaces X_{2i+1} $(0 \leq i \leq m)$ and $X_{2(m-i)}$ $(0 \leq j \leq m-1)$ become simple $\mathcal{SW}(m)$ -modules.
- 2. For each $0 \leq i \leq m-1$, the simple \mathcal{V}_L -modules $\mathcal{V}_{L+\gamma_i}$ and $\mathcal{V}_{L+\gamma_{2m-i}}$ become $\mathcal{SW}(m)$ -modules by restriction, and satisfy the following exact sequences:

$$
0 \to X_{2i+1} \to \mathcal{V}_{L+\gamma_i} \to X_{2(m-i)} \to 0,
$$

$$
0 \to X_{2(m-i)} \to \mathcal{V}_{L+\gamma_{2m-i}} \to X_{2i+1} \to 0.
$$

Remark 3.4. For the simple $\mathcal{SW}(m)$ -modules, the notation of [\[AM3\]](#page-65-1) and ours correspond as follows:

$$
S\Lambda(i+1) = X_{2i+1} \ (0 \le i \le m), \quad S\Pi(m-i) = X_{2(m-i)} \ (0 \le i \le m-1).
$$

We define the following three elements in \mathcal{V}_L

$$
W^- := |\beta_{1,1;-2}\rangle, \quad W^0 := Q_+ W^-, \quad W^+ := Q_+^2 W^-.
$$

These elements have the same L_0 -weight $h_{3,1} = 2m + \frac{1}{2}$ $\frac{1}{2}$. We define the following three elements

$$
\widehat{W}^- := b_{-\frac{1}{2}} |\beta_{1,1;-2}\rangle, \quad \widehat{W}^0 := Q_+ \widehat{W}^-, \quad \widehat{W}^+ := Q_+^2 \widehat{W}^-.
$$
 (3.4)

These elements have the same L_0 -weight $2m + 1$.

The following three theorems are very important in examining the detailed structure of the module category of $\mathcal{SW}(m)$.

Theorem 3.5 ([\[AM3\]](#page-65-1)). The $N = 1$ triplet vertex operator superalgebra $\mathcal{SW}(m)$ is generated by $Y(W^{\pm}, z), Y(W^0, z), G(z)$. Furthermore $\mathcal{SW}(m)$ is strongly generated by

$$
G(z)
$$
, $T(z)$, $Y(W^{\pm}, z)$, $Y(W^0, z)$, $Y(\widehat{W}^{\pm}, z)$, $Y(\widehat{W}^0, z)$.

Theorem 3.6 ([\[AM3\]](#page-65-1)). The $N = 1$ triplet vertex operator superalgebra $\mathcal{SW}(m)$ is C_2 -cofinite.

Theorem 3.7 ([\[AM3\]](#page-65-1)). All simple $\mathcal{SW}(m)$ -modules are completed by $2m+1$ simple $\mathcal{SW}(m)$ -modules in the set $\{X_s \mid 1 \leq s \leq 2m+1\}$.

The following proposition is straightforward from [\(3.2\)](#page-16-0), Proposition [2.10](#page-15-1) and the definition of Q_+ .

Proposition 3.8.

1. For $0 \leq i \leq m$, $n \geq 0$ and $-n \leq k \leq n$, we define $w_k^{(n)}$ $\binom{n}{k}(X_{2i+1}) := Q^{n+k}_+ \ket{\beta_{1,2i+1;-2n}}.$

Then the set $\{w_k^{(n)}\}$ $\{f_k^{(n)}(X_{2i+1})\}_{k=-n}^n$ gives a basis of the minimal conformal weight spaces of $(2n + 1)L(h_{1,2i+1;-2n}) \subset X_{2i+1}.$

2. For $0 \leq i \leq m-1$, $n \geq 0$ and $-n \leq k \leq n+1$, we define

$$
v_{\frac{2k-1}{2}}^{(n)}(X_{2(m-i)}) := Q_{+}^{n+k} | \beta_{1,2(m-i);-2n-1} \rangle.
$$

Then the set ${v_{\frac{2k-1}{2}}^{(n)}(X_{2(m-i)})\}_{k=-}^{n+1}$ $\sum_{k=-n}^{n+1}$ gives a basis of the minimal conformal weight spaces of $(2n + 2)L(h_{1,2(m-i);-2n-1})$ ⊂ $X_{2(m-i)}$.

In the following, we use the shorthand notation $w_k^{(n)} = w_k^{(n)}$ $k^{(n)}(X_{2i+1})$ and $v_{\frac{2k-1}{2}}^{(n)}=v_{\frac{2k-1}{2}}^{(n)}(X_{2(m-i)})$. The transitive $\mathcal{SW}(m)$ -actions on the simple nsmodules of (3.3) are given by the following proposition.

Proposition 3.9 ([\[AM3\]](#page-65-1)). Let $0 \leq i \leq m$ and $0 \leq j \leq m-1$. Then the fields $Y(W^{\pm}, z)$, $Y(W^0, z)$, $Y(W^{\pm}, z)$ and $Y(W^0, z)$ act on the vectors $w_k^{(n)} \in X_{2i+1}$ and $v_{\frac{2k-1}{2}}^{(n)} \in X_{2(m-j)}$ as follows:

1. The vectors $w_k^{(n)}$ $\binom{n}{k}$ $(n \geq 0, -n \leq k \leq n)$ satisfy

$$
W^{\pm}[-h]w_0^{(0)} = 0, \quad h < h_{1,2i+1;-2} - h_{1,2i+1} = \frac{1}{2} - i + 2m,\tag{3.5}
$$

and

$$
\begin{aligned}\n\widehat{W}^{\pm}[0]w_k^{(n)} &\in \mathbb{C}^{\times}w_{k\pm 1}^{(n)} + \sum_{l=0}^{n-1} U(\mathfrak{ns})w_{k\pm 1}^{(l)}, \\
w_{k\pm 1}^{(n+1)} &\in \mathbb{C}^{\times}W^{\pm}[h_{1,2i+1;-2n} - h_{1,2i+1;-2n-2}]w_k^{(n)} + \sum_{l=0}^{n} U(\mathfrak{ns})w_{k\pm 1}^{(l)}, \\
w_k^{(n+1)} &\in \mathbb{C}^{\times}W^0[h_{1,2i+1;-2n} - h_{1,2i+1;-2n-2}]w_k^{(n)} + \sum_{l=0}^{n} U(\mathfrak{ns})w_k^{(l)},\n\end{aligned}
$$

where $w_{n+1}^{(n)} = w_{-n-1}^{(n)} = w_k^{(-1)} = 0$, $W^{\bullet}[s] = W_s^{\bullet}$, $\widehat{W}^{\bullet}[t] = \widehat{W}_t^{\bullet}$ (s, $t \in \frac{1}{\sqrt[n]{\pi}}$ and $U(\mathfrak{ns})$ is the universal enveloping algebra of \mathfrak{ns} $\frac{1}{2}\mathbb{Z}$) and $U(\mathfrak{ns})$ is the universal enveloping algebra of $\mathfrak{ns}.$

2. The vectors $v_{\frac{2k+1}{2}}^{(n)}$ $(n \geq 0, -n-1 \leq k \leq n)$ satisfy

$$
W^\delta[-h]v^{(0)}_{\frac{\pm 1}{2}} \begin{cases} = 0 & \delta = \pm \\ \in U(\mathfrak{ns})v^{(0)}_{\frac{\pm 1}{2}} & \delta = 0 \\ \in U(\mathfrak{ns})v^{(0)}_{\frac{\mp 1}{2}} & \delta = \mp \end{cases} \quad h < h_{1,2(m-j);-3} - h_{1,2(m-j);-1}
$$

and

$$
\widehat{W}^{\pm}[0]v_{\frac{2k+1}{2}}^{(n)} \in \mathbb{C}^{\times}v_{\frac{2k+1\pm 2}{2}}^{(n)} + \sum_{l=0}^{n-1}U(\mathfrak{ns})v_{\frac{2k+1}{2}\pm 1}^{(l)},
$$
\n
$$
v_{\frac{2k+1}{2}\pm 1}^{(n+1)} \in \mathbb{C}^{\times}W^{\pm}[h_{1,2(m-j);-2n-1}-h_{1,2(m-j);-2n-3}]v_{\frac{2k+1}{2}}^{(n)} + \sum_{l=0}^{n}U(\mathfrak{ns})v_{\frac{2k+1}{2}\pm 1}^{(l)},
$$
\n(3.6)

$$
v^{(n+1)}_{\frac{2k+1}{2}}\in \mathbb{C}^{\times}W^0[h_{1,2(m-j);-2n-1}-h_{1,2(m-j);-2n-3}]v^{(n)}_{\frac{2k+1}{2}}+\sum_{l=0}^nU(\mathfrak{ns})v^{(l)}_{\frac{2k+1}{2}},
$$

where $v^{(n)}_{\frac{-2n-3}{2}} = v^{(n)}_{\frac{2n+3}{2}} = v^{(-1)}_{\frac{2k+1}{2}} = 0.$

Remark 3.10. The above proposition can be shown in a similar way as for the triplet \mathcal{W}_p case [\[AM1\]](#page-65-0), using the free field realizations and the screening operators (see also [\[FGST4,](#page-67-7) [TW2\]](#page-71-1)).

Let $A(\mathcal{SW}(m))$ be the Zhu-algebra of $\mathcal{SW}(m)$ (for the definition of Zhu-algebras, see [\[KW\]](#page-69-5), [\[Zh\]](#page-71-3)). For the structure of the Zhu-algebra $A(\mathcal{SW}(m))$, the following theorem holds.

Theorem 3.11 ([\[AM4\]](#page-65-4)). The Zhu-algebra $A(SW(m))$ decomposes as a sum of ideals

$$
A(\mathcal{SW}(m)) = \bigoplus_{i=2m+1}^{3m} \mathbb{M}_{h_{1,2i+1}} \oplus \bigoplus_{i=0}^{m-1} \mathbb{I}_{h_{1,2i+1}} \oplus \mathbb{C}_{h_{1,2m+1}},
$$

where $\mathbb{M}_{h_1,2i+1} \simeq M_2(\mathbb{C})$, $\dim(\mathbb{I}_{h_1,2i+1}) = 2$ and $\dim(\mathbb{C}_{h_1,2m+1}) = 1$.

Remark 3.12. Each $\mathbb{M}_{h_{1,2i+1}}$ $(i = 2m+1, \ldots, 3m)$ corresponds to the minimal conformal weight space of the simple module $X_{2(3m+1-i)}$, $\mathbb{C}_{h_{1,2m+1}}$ to the minimal conformal weight space of X_{2m+1} and each $\mathbb{I}_{h_{1,2i+1}}$ $(i = 0, \ldots, m-1)$ to the minimal conformal weight space of the projective cover of X_{2i+1} .

3.3 The block decomposition of $\mathcal{SW}(m)$ -mod

Let $\mathcal{SW}(m)$ -mod be the abelian category of grading restricted generalised $\mathcal{SW}(m)$ -modules (for the definition of grading restricted generalised modules, see Definition [2.2\)](#page-6-0). Since $\mathcal{SW}(m)$ is C_2 -cofinite, all objects of $\mathcal{SW}(m)$ -mod have finite length [\[Hu\]](#page-68-3). Note that $\mathcal{SW}(m)$ -mod is closed under contragredient.

We denote $\text{Ext}^n_{\mathcal{SW}(m)}(\bullet,\bullet)$ by the *n*-th Ext groups in the abelian category $\mathcal{SW}(m)$ -mod. The following proposition can be proved in a similar way as [\[AM1,](#page-65-0) Theorem 4.4] by using Theorem [3.7](#page-17-0) and the results for the semisimple category of \mathfrak{ns} in $|IK2|$, so we omit the proof.

Proposition 3.13. For all $i \neq j$, we have

$$
Ext^1_{\mathcal{SW}(m)}(X_{2i+1}, X_{2j+1}) = Ext^1_{\mathcal{SW}(m)}(X_{2(m-i)}, X_{2(m-j)}) = 0,
$$

$$
Ext^1_{\mathcal{SW}(m)}(X_{2i+1}, X_{2(m-j)}) = 0.
$$

For each $0 \leq i \leq m-1$ we denote by \mathcal{B}_{i+1} the full abelian subcategory of $\mathcal{SW}(m)$ -mod such that

 $M \in \mathcal{B}_{i+1} \Leftrightarrow$ every composition factors of M are given by $X_{2i+1}, X_{2(m-i)}$.

We denote by \mathcal{B}_{m+1} the full abelian subcategory of $\mathcal{SW}(m)$ -mod such that

 $M \in \mathcal{B}_{m+1} \Leftrightarrow$ every composition factors of M are given by X_{2m+1} .

By Proposition [3.13,](#page-19-0) we have the following proposition.

Proposition 3.14. The abelian category $\mathcal{SW}(m)$ -mod has the following block decomposition

$$
\mathcal{SW}(m)\text{-}\mathrm{mod}=\bigoplus_{i=0}^m\mathcal{B}_{i+1}.
$$

4 Correlation functions

In this section, using a certain free field realization technique for vertex operators by [\[DF1,](#page-66-6) [DF2,](#page-66-7) [Fe,](#page-67-8) [FS\]](#page-67-9) and some properties of the Dotsenko-Fateev integrals [\[DF1,](#page-66-6) [DF2\]](#page-66-7) given by [\[Su1,](#page-70-7) [Su2\]](#page-70-8), we will construct the fundamental system of solutions of a fourth-order differential equation, and determine the connection matrix between the solutions arround $z = 0$ and $z = 1$. The results of this section will be important to show the self-duality of the simple module X_2 (see Subsection [5.2\)](#page-46-0).

4.1 A fourth-order Fuchsian differential equation

Let \mathcal{Y}_1 and \mathcal{Y}_2 be even $\mathcal{SW}(m)$ -intertwining operators of type $\begin{pmatrix} X_2 \\ X_2 \end{pmatrix}$ X_2 M) and $\left(M \right)$ M X_2 X_2 , respectively, for some $\mathcal{SW}(m)$ -module M. Fix any minimal conformal weight vector $v_{X_2} \in X_2[h_{2,2}]$ and let $v_{X_2}^*$ be a minimal conformal weight vector of $X_2^*(\simeq X_2)$ such that $\langle v_{X_2}^*, v_{X_2} \rangle \neq 0$. We define two correlation functions

$$
R^{0}(z_{1}, z_{2}) = \langle v_{X_{2}}^{*}, \mathcal{Y}_{1}(v_{X_{2}}, z_{1})\mathcal{Y}_{2}(v_{X_{2}}, z_{2})v_{X_{2}}\rangle, R^{1}(z_{1}, z_{2}) = \langle v_{X_{2}}^{*}, \mathcal{Y}_{1}(G_{-\frac{1}{2}}v_{X_{2}}, z_{1})\mathcal{Y}_{2}(G_{-\frac{1}{2}}v_{X_{2}}, z_{2})v_{X_{2}}\rangle,
$$
(4.1)

where $|z_1| > |z_2| > |z_1 - z_2| > 0$. Using [\(2.1\)](#page-5-0) and [\(2.3\)](#page-8-0), we have (cf. [\[CMOY,](#page-66-4) Subsection 2.4])

$$
[G_{-n-\frac{1}{2}}, \mathcal{Y}_i(v_{X_2}, z)] = z^{-n} \mathcal{Y}_i(G_{-\frac{1}{2}} v_{X_2}, z),
$$

\n
$$
[G_{-n-\frac{1}{2}}, \mathcal{Y}_i(G_{-\frac{1}{2}} v_{X_2}, z)] = (z^{-n} \partial_z - 2nz^{-n-1} h_{2,2}) \mathcal{Y}_i(v_{X_2}, z),
$$

\n
$$
[L_{-n}, \mathcal{Y}_i(v_{X_2}, z)] = (z^{-n+1} \partial_z + (1-n)z^{-n} h_{2,2}) \mathcal{Y}_i(v_{X_2}, z),
$$

\n
$$
[L_{-n}, \mathcal{Y}_i(G_{-\frac{1}{2}} v_{X_2}, z)] = (z^{-n+1} \partial_z + (1-n)z^{-n} (h_{2,2} + \frac{1}{2})) \mathcal{Y}_i(G_{-\frac{1}{2}} v_{X_2}, z).
$$

\n(4.2)

We set

$$
S_{2,2} := \frac{4t}{t^2 - 1} G_{-\frac{1}{2}}^4 + \frac{t+1}{t-1} G_{-\frac{1}{2}} G_{-\frac{3}{2}} + \frac{t-1}{t+1} G_{-\frac{3}{2}} G_{-\frac{1}{2}} \quad (t = -2m - 1). \tag{4.3}
$$

This element of $U(\mathfrak{ns})$ gives an \mathfrak{ns} -singular vector of the Verma \mathfrak{ns} -module whose lowest weight and central charge are given by $h_{2,2}$ and $c_{1,2m+1}$ (cf. [\[BS,](#page-65-6) [CMOY\]](#page-66-4)). Since $U(\mathfrak{ns})v_{X_2} \simeq L(h_{2,2}), v_{X_2}$ satisfies the relation $S_{2,2}v_{X_2} =$ 0. Then, similar to the arguments in [\[HM\]](#page-69-6) (see also [\[Ba,](#page-65-7) [Be\]](#page-65-8)), using [\(4.2\)](#page-20-2), we can show that $R^0(z_1, z_2)$ and $R^1(z_1, z_2)$ satisfy the following differential equations

$$
\begin{split}\n&\left\{\frac{m^2 + 2h_{2,2}(m+1)^2}{m(m+1)} \left(\frac{1}{z_1^2} + \frac{1}{z_2^2}\right) - \frac{2m+1}{m(m+1)} (\partial_{z_1} + \partial_{z_2})^2 - \frac{2m^2 + 2m + 1}{m(m+1)} \left(\frac{\partial_{z_1}}{z_1} + \frac{\partial_{z_2}}{z_2}\right) \right\} R^1 \\
&+ \left\{\frac{2m+1}{m(m+1)} \left(\frac{1}{z_2} - \frac{1}{z_1}\right) \partial_{z_1} \partial_{z_2} - \frac{2(2m+1)h_{2,2}}{m(m+1)} \left(\frac{\partial_{z_1}}{z_2^2} - \frac{\partial_{z_2}}{z_1^2}\right) \right\} R^0 = 0, \\
&\left\{\frac{2mh_{2,2}}{m+1} \left(\frac{1}{z_1^2} + \frac{1}{z_2^2}\right) - \frac{2m+1}{m(m+1)} (\partial_{z_1} + \partial_{z_2})^2 - \frac{2m^2 + 2m + 1}{m(m+1)} \left(\frac{\partial_{z_1}}{z_1} + \frac{\partial_{z_2}}{z_2}\right) \right\} R^0 + \frac{2m+1}{m(m+1)} \left(\frac{1}{z_1} - \frac{1}{z_2}\right) R^1 = 0.\n\end{split} \tag{4.4}
$$

We define two functions $\overline{R}^0(z) = R^0(1, z)$ and $\overline{R}^1(z) = R^1(1, z)$. From the L₀-conjugation formula for intertwining operators, we have $R^i(z_1, z_2) =$ $z_1^{-2h_{2,2}-i}\overline{R}^i(z_2/z_1)$, $(i=0,1)$. Then, from [\(4.4\)](#page-21-0), we can show that $\overline{R}^0(z)$ and $\overline{R}^1(z)$ satisfy

$$
\begin{aligned} \left\{ \frac{d^2}{dz^2} + \frac{(4m^2 + 2m + 1)z - 2m^2 - 2m - 1}{(2m + 1)z(z - 1)} \frac{d}{dz} - \frac{3m^4}{(2m + 1)^2 z^2 (z - 1)^2} \right\} \overline{R}^0(z) \\ &+ \frac{\overline{R}^1(z)}{z(1 - z)} = 0, \end{aligned}
$$

and

$$
\begin{aligned}\n&\left\{\frac{d^2}{dz^2} + \frac{(4m^2 + 6m + 3)z - 2m^2 - 2m - 1}{(2m + 1)z(z - 1)}\frac{d}{dz} + \frac{(2m + 1)^2z^2 - m^2(3m^2 + 8m + 4)}{(2m + 1)^2z^2(z - 1)^2}\right\}\overline{R}^1(z) \\
&+ \left\{\frac{1}{1 - z}\frac{d^2}{dz^2} - \frac{(6m^2 + 2m + 1)z - 2m - 1}{(2m + 1)z(z - 1)^2}\frac{d}{dz} - \frac{9m^4}{(2m + 1)^2z^2(z - 1)^2}\right\}\overline{R}^0(z) = 0.\n\end{aligned}
$$

From these equations, we can see that $\overline{R}^0(z)$ satisfies the following Fuchsian differential equation

$$
\left(\frac{d^4}{dz^4} + \frac{p_3(z)}{z(z-1)}\frac{d^3}{dz^3} + \frac{p_2(z)}{z^2(z-1)^2}\frac{d^2}{dz^2} + \frac{p_1(z)}{z^3(z-1)^3}\frac{d}{dz} + \frac{p_0(z)}{z^4(z-1)^4}\right)\Phi(z) = 0,
$$
\n(4.5)

where

$$
p_0(z) = \frac{3m^4}{(2m+1)^4} \left((16m^3 - 8m^2 - 16m - 4)z^2 + (-16m^3 + 8m^2 + 16m + 4)z + (3m^4 + 12m^3 + 2m^2 - 4m - 1) \right),
$$

\n
$$
p_1(z) = \frac{2}{(2m+1)^3} \left((16m^5 + 48m^4 + 56m^3 + 34m^2 + 12m + 2)z^3 + (-24m^5 - 72m^4 - 84m^3 - 51m^2 - 18m - 3)z^2 + (-12m^6 - 8m^5 + 12m^3 + 13m^2 + 6m + 1)z + (6m^6 + 8m^5 + 12m^4 + 8m^3 + 2m^2) \right),
$$

\n
$$
p_2(z) = \frac{2}{(2m+1)^2} \left((8m^4 + 32m^3 + 44m^2 + 28m + 7)z^2 + (-8m^4 - 32m^3 - 44m^2 - 28m - 7)z + (-m^4 + 2m^3 + 5m^2 + 4m + 1) \right),
$$

\n
$$
p_3(z) = \frac{4(m+1)^2(2z-1)}{(2m+1)}.
$$

The Riemann scheme of the Fuchsian differential equation [\(4.5\)](#page-22-0) is given by

$$
\begin{bmatrix}\n0 & 1 & \infty \\
\rho_{1,1} & \rho_{1,1} & 0 \\
\rho_{0,1} & \rho_{0,1} & \frac{1}{2m+1} \\
\rho_{1,0} & \rho_{1,0} & \frac{4m^2}{2m+1} \\
\rho_{0,0} & \rho_{0,0} & 2m+1\n\end{bmatrix}.
$$
\n(4.6)

where

$$
\rho_{1,1} = h_{3,3} - 2h_{2,2} = \frac{m^2}{2m+1}, \qquad \rho_{0,1} = h_{3,1} + \frac{1}{2} - 2h_{2,2} = \frac{m^2 + 4m + 1}{2m+1},
$$

$$
\rho_{1,0} = h_{1,3} + \frac{1}{2} - 2h_{2,2} = \frac{1 - 3m^2}{2m+1}, \quad \rho_{0,0} = h_{1,1} - 2h_{2,2} = -\frac{3m^2}{2m+1}
$$
(4.7)

(see [\[Ha\]](#page-68-4) for basic facts of the Fuchsian differential equations). Note that $\rho_{1,1} - \rho_{1,0}, \ \rho_{0,1} - \rho_{0,0} \in \mathbb{Z}_{\geq 1}$. In Subsection [4.4,](#page-37-0) we show that [\(4.5\)](#page-22-0) has no logarithmic solutions at $z = 0, 1$.

Similarly, we can see that the correlation function

$$
\langle v_{X_2}^*, \mathcal{Y}^1(\mathcal{Y}^2(v_{X_2}, 1-z)v_{X_2}, z)v_{X_2} \rangle \tag{4.8}
$$

satisfies the Fuchsian differential equation (4.5) , where \mathcal{Y}^1 and \mathcal{Y}^2 are even $\mathcal{SW}(m)$ -intertwining operators of type $\begin{pmatrix} X_2 \\ N & X \end{pmatrix}$ N X_2) and $\left(\frac{N}{\mathbf{r}_{\alpha}}\right)$ X_2 X_2 , respectively, for some $\mathcal{SW}(m)$ -module N.

Remark 4.1. In [\[CMOY,](#page-66-4) Subsection 4.3], unlike in our cases $((4.1), (4.8)),$ $((4.1), (4.8)),$ $((4.1), (4.8)),$ $((4.1), (4.8)),$ $((4.1), (4.8)),$ a slightly different correlation function is examined using a certain embedding technique for Virasoro vertex operator algebras.

4.2 Regularization of the Dotsenko-Fateev integrals

For each $l, m, n \in \mathbb{N}$ not all zero, let

$$
\Box_{\bm{x}}^{l,m,n} = [-\infty, 0]_{x_1,\dots,x_l}^l \times [0, 1]_{x_{l+1},\dots,x_{l+m}}^m \times [1, \infty]_{x_{l+m+1},\dots,x_N}^n \subseteq \overline{\mathbb{R}}^N, \quad (4.9)
$$

where $N = l + m + n$. We define

$$
J_{l,m,n}(\boldsymbol{a},\boldsymbol{b},\boldsymbol{\gamma}) = \int_{\square_{\boldsymbol{x}}^{l,m,n}} \prod_{i=1}^N x_i^{a_i} (x_i - 1)^{b_i} \prod_{1 \le j < k \le N} (x_k - x_j + i0)^{-2\gamma_{j,k}} \mathrm{d} x_1 \dots \mathrm{d} x_N,
$$
\n(4.10)

where $\boldsymbol{a} = \{a_i\}_{i=1}^N$, $\boldsymbol{b} = \{b_i\}_{i=1}^N$ and $\boldsymbol{\gamma} = \{\gamma_{j,k} = \gamma_{k,j}\}_{1 \leq j < k \leq N}$. These integrals are called *Dotsenko-Fateev integrals* [\[DF1,](#page-66-6) [DF2\]](#page-66-7). Note that for $N = 1$, $J_{l,m,n}(\boldsymbol{a},\boldsymbol{b},\boldsymbol{\gamma})$ is just Euler Beta integral. The Dotsenko-Fateev integrals were introduced in [\[DF1,](#page-66-6) [DF2,](#page-66-7) [Fe,](#page-67-8) [FS\]](#page-67-9) for the approachs to the construction of the BPZ-minimal models [\[BPZ\]](#page-65-2). In these papers [\[DF1,](#page-66-6) [DF2,](#page-66-7) [Fe,](#page-67-8) [FS\]](#page-67-9), the parameters $\gamma_{i,k}$ are fixed as $\gamma_{i,k} = 1$, a condition that follows naturally from the free-field realization of correlation functions (see also Subsection [4.4\)](#page-37-0). However, this integer condition makes the problem for the regularization of integrals very difficult. In fact, we see that the construction of cycles in [\[TK\]](#page-71-2) is not applicable under condition $\gamma_{i,k} = 1$. Thus, the regularization of the Dotsenko-Fateev integrals has been an open problem until a recent result by [\[Su2\]](#page-70-8).

Let us introduce some notation in accordance with [\[Su2,](#page-70-8) Section 1]. For $0 \leq r < 1 < R \leq \infty$, we define

$$
\mathcal{M}_{1,0;N} := \{ \boldsymbol{x} \in \mathbb{C}^N \mid (x_i \neq x_j, i \neq j) \land (x_i \neq 1,0) \},
$$

$$
\mathcal{M}_{1,0;N}(r,R) := \{ (u,v) \in \mathcal{M}_{1,0;N} \mid r < |x_i| < R \}.
$$

The manifold $\mathcal{M}_{1,0:N}(r, R)$ is the moduli space of two pairwise distinct elements of the punctured annulus $\{z \in \mathbb{C} \mid r < |x_i| < R, x_i \neq 1\}$. Let $\mathcal{M}_{1,0;N}(r, R)$ be the universal cover of $\mathcal{M}_{1,0;N}(r, R)$. Then the integrand

$$
\mathcal{V}_N(\boldsymbol{a}, \boldsymbol{b}, \boldsymbol{\gamma}; \boldsymbol{x}) := \prod_{i=1}^N x_i^{a_i} (x_i - 1)^{b_i} \prod_{1 \leq j < k \leq N} (x_k - x_j + i0)^{-2\gamma_{j,k}} \tag{4.11}
$$

of $J_{l,m,n}(\boldsymbol{a},\boldsymbol{b},\boldsymbol{\gamma})$ is a *single-valued* analytic function on the monodromy cover

$$
\widehat{\mathcal{M}}_{1,0;N}(r,R):=\widehat{\mathcal{M}}_{1,0;N}(r,R)/[\pi_1(\mathcal{M}_{1,0;N}(r,R)),\pi_1(\mathcal{M}_{1,0;N}(r,R))].
$$

Remark 4.2. The above setting is simplified by considering the case of $N = 1$ and $(r, R) = (0, \infty)$. Let g and h be the generators of $\pi_1(\mathcal{M}_{1,0;1}) =$ $\pi_1(\mathbb{C} \setminus \{1,0\})$ corresponding to the one counterclockwise circuit around 0, 1, respectively. Then the element

$$
g^{-1}h^{-1}gh \in [\pi_1(\mathbb{C} \setminus \{1,0\}), \pi_1(\mathbb{C} \setminus \{1,0\})]
$$

defines the Pochhammer contour arround 1 and 0 (Figure [4.1\)](#page-24-0). Circling along the Pochhammer contour $g^{-1}h^{-1}gh$, we see that the monodromy from $x^a(x-1)^b$ becomes trivial. Thus the integrand $x^a(x-1)^b$ is well-defined on the quotient space $\widetilde{\mathcal{M}}_{1,0;1}/[\pi_1(\mathbb{C} \setminus \{1,0\}), \pi_1(\mathbb{C} \setminus \{1,0\})].$

Figure 4.1: The Pochhammer contour arround 1 and 0.

According to [\[Su2\]](#page-70-8), the problem of regularizing the Dotsenko–Fateev integral $J_{l,m,n}(\boldsymbol{a},\boldsymbol{b},\boldsymbol{\gamma})$ can be formulated as finding multi-contours

$$
\Gamma_{l,m,n} \in H_N(\mathcal{\tilde{M}}_{1,0;N}(r,R);\mathbb{Z})
$$

satisfying

$$
\int_{\Gamma_{l,m,n}} \mathcal{V}_N(\boldsymbol{a},\boldsymbol{b},\boldsymbol{\gamma};\boldsymbol{x}) dx_1 \wedge \cdots \wedge dx_N \propto J_{l,m,n}(\boldsymbol{a},\boldsymbol{b},\boldsymbol{\gamma})
$$
(4.12)

where " \propto " denotes proportionality up to constants and trigonometric functions of $\boldsymbol{a}, \boldsymbol{b}, \gamma$. In [\[Su2\]](#page-70-8), these cycles $\Gamma_{l,m,n}$ are constructed explicitly, and each proportional constant in [\(4.12\)](#page-25-0) is determined. For each $S \subseteq \{1, \ldots, N\}$, let

$$
a_{\mathcal{S}} = \sum_{j \in \mathcal{S}} a_j - 2 \sum_{\substack{1 \le j < k \le N \\ j, k \in \mathcal{S}}} \gamma_{j,k}, \qquad b_{\mathcal{S}} = \sum_{j \in \mathcal{S}} b_j - 2 \sum_{\substack{1 \le j < k \le N \\ j, k \in \mathcal{S}}} \gamma_{j,k},
$$
\n
$$
\zeta_{\mathcal{S}} = -\sum_{j \in \mathcal{S}} (a_j + b_j) + 2 \sum_{\substack{1 \le j < k \le N \\ j \in \mathcal{S} \text{ or } k \in \mathcal{S}}} \gamma_{j,k}.
$$

Theorem 4.3 ([\[Su2\]](#page-70-8)). There exists a cycle $\Gamma_{l,m,n} \in H_N(\widehat{\mathcal{M}}_{1,0;N}(r,R); \mathbb{Z})$ such that

$$
\int_{\Gamma_{l,m,n}} \mathcal{V}_N(\boldsymbol{a},\boldsymbol{b},\boldsymbol{\gamma};\boldsymbol{x}) dx_1 \wedge \cdots \wedge dx_N = c_{l,m,n}(\boldsymbol{a},\boldsymbol{b},\boldsymbol{\gamma}) J_{l,m,n}(\boldsymbol{a},\boldsymbol{b},\boldsymbol{\gamma}) \quad (4.13)
$$

for all $a, b \in \mathbb{C}^N$ and $\gamma \in \mathbb{C}^{\frac{N(N-1)}{2}}$ for which $J_{l,m,n}(a, b, \gamma)$ is defined, where by denoting $\mathfrak{e}(x) = 1 - \exp(2\pi i x), c_{l,m,n}(\mathbf{a}, \mathbf{b}, \gamma)$ is given by

$$
c_{l,m,n} = \prod_{\varnothing \subsetneq S \subseteq \{1,\ldots,l+m\}} \mathfrak{e}(a_{\mathcal{S}}) \prod_{\varnothing \subsetneq S \subseteq \{l+1,\ldots,N\}} \mathfrak{e}(b_{\mathcal{S}}) \prod_{\varnothing \subsetneq S \subseteq \{1,\ldots,l\} \cup \{l+m+1,\ldots,N\}} \mathfrak{e}(\zeta_{\mathcal{S}}).
$$
\n(4.14)

Remark 4.4. 1. Let $\mathcal{H}_{l,m,n}$ be the following collection of hyperplanes

$$
\mathcal{H}_{l,m,n} = \Big(\bigcup_{\mathcal{S} \subseteq \{1,\dots,N\}} \{a_{\mathcal{S}} \in \mathbb{Z}\}\Big) \cup \Big(\bigcup_{\mathcal{S}} \{b_{\mathcal{S}} \in \mathbb{Z}\}\Big) \cup \Big(\bigcup_{\mathcal{S}} \{\zeta_{\mathcal{S}} \in \mathbb{Z}\}\Big).
$$

Then from the above theorem, $J_{l,m,n}$ can be extended to an analytic function

$$
J_{l,m,n}: \left(\mathbb{C}^N\times \mathbb{C}^N\times \mathbb{C}^{\frac{N(N-1)}{2}}\right)\setminus \mathcal{H}_{l,m,n}\to \mathbb{C}.
$$

In particular, since $\mathcal{H}_{l,m,n}$ does not contain any affine hyperplanes $\{\gamma_{i,k} =$ c} (c $\in \mathbb{C}$), the variables $\gamma_{j,k}$ of $J_{l,m,n}(\boldsymbol{a},\boldsymbol{b},\boldsymbol{\gamma})$ are apparent singularities at $\gamma_{i,k} = 1$.

2. Each cycle $\Gamma_{l,m,n} \in H_N(\widehat{\mathcal{M}}_{1,0:N}(r,R);\mathbb{Z})$ is defined by a lifting of a cycle in $\mathcal{M}_{1,0;N}(r, R)$. These cycles and the lifting are constructed explicitly in [\[Su2,](#page-70-8) Sections 4 and 5].

In this paper we are mainly interested in the properties of the Dotsenko–Fateev integrals in two dimensional case. Then in the following, we introduce our notation for $N = 2$. Fix $z_1, z_2 \in \mathbb{R}_{>0}$ satisfying $z_2 > z_1 - z_2 > 0$. Given $\mathbf{a} = \{a_1, a_2\}, \mathbf{b} = \{b_1, b_2\}, \mathbf{c} = \{c_1, c_2\} \in \mathbb{C}^2$ and $\gamma \in \mathbb{C}$, we define a new multivalued function

$$
\mathcal{U}(\mathbf{a}, \mathbf{b}, \mathbf{c}, \gamma; u, v; z_1, z_2) \n:= u^{a_1}(u - z_2)^{b_1}(u - z_1)^{c_1}v^{a_2}(v - z_2)^{b_2}(v - z_1)^{c_2}(u - v)^{-2\gamma}
$$
\n(4.15)

on $\mathcal{N}_{z_1,z_2,0} = \{(u,v) \in \mathbb{C}^2 \mid (u \neq v) \wedge (u \neq z_1,z_2,0) \wedge (v \neq z_1,z_2,0)\}.$ For $i, j = 1, 0$, let $\Box_{i,j}^{\pm}$ and $\overline{\Box}_{i,j}^{\pm}$ be the open subsets of $\mathcal{N}_{z_1, z_2, 0}$ and $\mathcal{M}_{1,0;2}$ defined by

$$
\Box_{1,1}^{+} := \{z_1 < u, z_1 < v\}, \qquad \Box_{0,1}^{+} := \{0 < u < z_2, z_1 < v\},
$$
\n
$$
\Box_{1,0}^{+} := \{z_1 < u, 0 < v < z_2\}, \qquad \Box_{0,0}^{+} := \{0 < u < z_2, 0 < v < z_2\},
$$
\n
$$
\overline{\Box}_{1,1}^{+} := \{1 < u, 1 < v\}, \qquad \overline{\Box}_{0,1}^{+} := \{0 < u < 1, 1 < v\},
$$
\n
$$
\overline{\Box}_{1,0}^{+} := \{1 < u, 0 < v < 1\}, \qquad \overline{\Box}_{0,0}^{+} := \{0 < u < 1, 0 < v < 1\},
$$

and

$$
\Box_{1,1}^{-} := \{u < 0, v < 0\}, \qquad \Box_{0,1}^{-} := \{z_2 < u < z_1, v < 0\},
$$
\n
$$
\Box_{1,0}^{-} := \{u < 0, z_2 < v < z_1\}, \qquad \Box_{0,0}^{-} := \{z_2 < u < z_1, z_2 < v < z_1\},
$$
\n
$$
\overline{\Box}_{1,1}^{-} := \{u < 0, v < 0\}, \qquad \overline{\Box}_{0,1}^{-} := \{0 < u < 1, v < 0\},
$$
\n
$$
\overline{\Box}_{1,0}^{-} := \{u < 0, 0 < v < 1\}, \qquad \overline{\Box}_{0,0}^{-} := \overline{\Box}_{0,0}^{+}
$$

(see Figure [4.2\)](#page-27-0). Note that by using the notation [\(4.9\)](#page-23-2), we have

$$
\overline{\Box}_{1,1}^{+} = \Box_{u,v}^{0,0,2}, \qquad \overline{\Box}_{0,1}^{+} = \Box_{u,v}^{0,1,1}, \qquad \overline{\Box}_{0,0}^{\pm} = \Box_{u,v}^{0,2,0}, \qquad \overline{\Box}_{1,0}^{+} = \Box_{v,u}^{0,1,1},
$$

$$
\overline{\Box}_{1,0}^{-} = \Box_{u,v}^{1,1,0}, \qquad \overline{\Box}_{1,1}^{-} = \Box_{u,v}^{2,0,0}, \qquad \overline{\Box}_{0,1}^{-} = \Box_{v,u}^{1,1,0}.
$$

We set

$$
P_{x,y,z}(u,v) := \mathbb{C}[(u-x)^{\pm 1}, (v-x)^{\pm 1}, (u-y)^{\pm 1}, (v-y)^{\pm 1}, (u-z)^{\pm 1}, (v-z)^{\pm 1}],
$$

\n
$$
P_{x,y}(u,v) := P_{x,y,y}(u,v),
$$

\n
$$
P_x(u,v) := P_{x,x}(u,v),
$$

Figure 4.2: The open subsets $\Box_{i,j}^{\pm}$ and $\overline{\Box}_{i,j}^{\pm}$.

for $x, y, z \in \mathbb{C}$. We define

 $l_{i,j}^{\pm,s}, l_{i,j}^{\pm,t}, r_{i,j}^{\pm,s}, r_{i,j}^{\pm,t} \in \{-\infty, 0, z_1, z_2, \infty\}, \qquad \bar{l}_{i,j}^{\pm,s}, \bar{l}_{i,j}^{\pm,t}, \bar{r}_{i,j}^{\pm,s}, \bar{r}_{i,j}^{\pm,t} \in \{-\infty, 0, 1, \infty\}$ as follows

$$
\Box_{i,j}^{\pm} = \{ l_{i,j}^{\pm,s} < u < l_{i,j}^{\pm,t}, r_{i,j}^{\pm,s} < v < r_{i,j}^{\pm,t} \},
$$

$$
\overline{\Box}_{i,j}^{\pm} = \{ \overline{l}_{i,j}^{\pm,s} < u < \overline{l}_{i,j}^{\pm,t}, \overline{r}_{i,j}^{\pm,s} < v < \overline{r}_{i,j}^{\pm,t} \}.
$$

Then we introduce the following two types of integrals.

Definition 4.5. 1. For $i, j \in \{1, 0\}$ and $E \in P_{z_1, z_2, 0}$, we define

$$
\mathcal{I}_{i,j}^{\pm}[E](\boldsymbol{a},\boldsymbol{b},\boldsymbol{c},\gamma;z_1,z_2) \n:= \int_{\Box_{i,j}^{\pm}} \mathcal{U}(\boldsymbol{a},\boldsymbol{b},\boldsymbol{c},\gamma;u,v;z_1,z_2) E(u,v) du dv \n:= \int_{l_{i,j}^{\pm,s}}^{l_{i,j}^{\pm,t}} \int_{r_{i,j}^{\pm,s}}^{r_{i,j}^{\pm,t}} u^{a_1}(u-z_2)^{b_1}(u-z_1)^{c_1} v^{a_2}(v-z_2)^{b_2}(v-z_1)^{c_2} \n\times (u-v+i0)^{-2\gamma} E(u,v) du dv.
$$

2. For $i, j \in \{1, 0\}$ and $F \in P_{1,0}$, we define

$$
\mathcal{J}_{i,j}^{\pm}[F](\boldsymbol{a},\boldsymbol{b},\gamma)
$$

\n
$$
:= \int_{\overline{\square}^{\pm}_{i,j}} \mathcal{V}_2(\boldsymbol{a},\boldsymbol{b},\gamma;u,v) F(u,v) du dv
$$

\n
$$
:= \int_{\overline{\square}^{\pm}_{i,j}}^{\overline{\square}^{\pm,t}} \int_{\overline{r}_{i,j}^{\pm,s}}^{\overline{r}_{i,j}^{\pm,t}} u^{a_1}(u-1)^{b_1} v^{a_2}(v-1)^{b_2}(u-v+i0)^{-2\gamma} F(u,v) du dv,
$$

where we set $\mathcal{J}_{0,0}^+[F](\boldsymbol{a},\boldsymbol{b},\gamma)=\mathcal{J}_{0,0}^-[F](\boldsymbol{a},\boldsymbol{b},\gamma).$

We refer to these integrals collectively as the Dotsenko-Fateev integrals. Applying Theorem [4.3](#page-25-1) to our notation for $N = 2$, we have the following theorem.

Theorem 4.6 ([\[Su2\]](#page-70-8)). Let $i, j \in \{1, 0\}$. Then for each (r, R) such that $0 \leq r < 1 < R \leq \infty$, there exists a cycle $\Gamma^{\pm}_{i,j} \in H_2(\widehat{\mathcal{M}}_{1,0;2}(r,R);\mathbb{Z})$ such that

$$
\int_{\Gamma_{i,j}^{\pm}} \mathcal{V}_2(\boldsymbol{a}, \boldsymbol{b}, \gamma; u, v) \, \mathrm{d}u \wedge \mathrm{d}v = c_{i,j}^{\pm}(\boldsymbol{a}, \boldsymbol{b}, \gamma) \mathcal{J}_{i,j}^{\pm}[1](\boldsymbol{a}, \boldsymbol{b}, \gamma) \tag{4.16}
$$

for all $(\bm{a},\bm{b},\gamma) \in \mathbb{C}^5_{\bm{a},\bm{b},\gamma}$ for which $\mathcal{J}_{i,j}^{\pm}[1](\bm{a},\bm{b},\gamma)$ is defined, where we set $\Gamma^{+}_{0,0}=\Gamma^{-}_{0,0}$ and $c_{i,j}^{\pm}(\boldsymbol{a},\boldsymbol{b},\gamma)$ are given by

$$
c_{1,1}^{+}(\boldsymbol{a}, \boldsymbol{b}, \gamma) = \mathbf{e}(b_{1})\mathbf{e}(b_{2})\mathbf{e}(b_{1} + b_{2} - 2\gamma)\mathbf{e}(a_{1} + b_{1} - 2\gamma)\mathbf{e}(a_{2} + b_{2} - 2\gamma)
$$

\n
$$
\times \mathbf{e}(a_{1} + a_{2} + b_{1} + b_{2} - 2\gamma),
$$

\n
$$
c_{1,1}^{-}(\boldsymbol{a}, \boldsymbol{b}, \gamma) = c_{1,1}^{+}(\boldsymbol{b}, \boldsymbol{a}, \gamma),
$$

\n
$$
c_{0,0}^{+}(\boldsymbol{a}, \boldsymbol{b}, \gamma) = \mathbf{e}(a_{1})\mathbf{e}(a_{2})\mathbf{e}(b_{1})\mathbf{e}(b_{2})\mathbf{e}(a_{1} + a_{2} - 2\gamma)\mathbf{e}(b_{1} + b_{2} - 2\gamma),
$$

\n
$$
c_{0,1}^{+}(\boldsymbol{a}, \boldsymbol{b}, \gamma) = \mathbf{e}(a_{1})\mathbf{e}(b_{1})\mathbf{e}(b_{2})\mathbf{e}(b_{1} + b_{2} - 2\gamma)\mathbf{e}(a_{2} + b_{2} - 2\gamma),
$$

\n
$$
c_{1,0}^{+}(\boldsymbol{a}, \boldsymbol{b}, \gamma) = \mathbf{e}(a_{2})\mathbf{e}(b_{1})\mathbf{e}(b_{2})\mathbf{e}(b_{1} + b_{2} - 2\gamma)\mathbf{e}(a_{1} + b_{1} - 2\gamma),
$$

\n
$$
c_{0,1}^{-}(\boldsymbol{a}, \boldsymbol{b}, \gamma) = c_{0,1}^{+}(\boldsymbol{b}, \boldsymbol{a}, \gamma),
$$

\n
$$
c_{1,0}^{-}(\boldsymbol{a}, \boldsymbol{b}, \gamma) = c_{1,0}^{+}(\boldsymbol{b}, \boldsymbol{a}, \gamma).
$$
\n(4.17)

By [\(4.16\)](#page-28-0), each $\mathcal{J}_{i,j}^{\pm}[1](\boldsymbol{a},\boldsymbol{b},\gamma)$ admits an analytic continuation for the variables (a, b, γ) . Since the left-hand side of [\(4.16\)](#page-28-0) is holomorphic with respect to $(\bm{a},\bm{b},\gamma),$ from the explicit form of $c_{i,j}^{\pm}(\bm{a},\bm{b},\gamma),$ we have the following theorem.

Theorem 4.7 ([\[Su1,](#page-70-7) [Su2\]](#page-70-8)). Let $i, j = 1, 0$. Then $\mathcal{J}_{i,j}^{\pm}[1](a, b, \gamma)$ admits an analytic extension to $\mathbb{C}^5_{a,b,\gamma} \setminus \mathcal{H}_{a,b,\gamma}$, where

$$
\mathcal{H}_{a,b,\gamma} = \{a_1 \in \mathbb{Z}\} \cup \{a_2 \in \mathbb{Z}\} \cup \{a_1 + a_2 - 2\gamma \in \mathbb{Z}\} \cup \{b_1 \in \mathbb{Z}\} \cup \{b_2 \in \mathbb{Z}\} \cup \{b_1 + b_2 - 2\gamma \in \mathbb{Z}\} \cup \{a_1 + b_1 \in \mathbb{Z}\} \cup \{a_2 + b_2 \in \mathbb{Z}\} \cup \{a_1 + a_2 + b_1 + b_2 - 2\gamma \in \mathbb{Z}\}.
$$
\n(4.18)

Let z and w be real numbers satisfying $z > w \geq 0$. Then we set

$$
\mathcal{M}_{z,w;N} := \{ \boldsymbol{x} \in \mathbb{C}^N \mid (x_i \neq x_j, i \neq j) \land (x_i \neq z, w) \}.
$$

For $0 \le r \le z - w \le R \le \infty$ we define

$$
\mathcal{M}_{z,w;N}(r,R):=\{\boldsymbol{x}\in\mathbb{C}^N\mid r<|x_i-w|
$$

Let $\mathcal{M}_{z,w;N}(r, R)$ be the universal cover of $\mathcal{M}_{z,w;N}(r, R)$. Then

$$
\mathcal{V}_N(\bm{a},\bm{b},\bm{\gamma};(z-w)^{-1}(x_1-w),\ldots,(z-w)^{-1}(x_N-w))
$$

is a single-valued analytic function on the monodromy cover

$$
\widehat{\mathcal{M}}_{z,w;N}(r,R):=\widetilde{\mathcal{M}}_{z,w;N}(r,R)/[\pi_1(\mathcal{M}_{z,w;N}(r,R)),\pi_1(\mathcal{M}_{z,w;N}(r,R))].
$$

Note that there exists a diffeomorphism

$$
t_{z,w;N;r,R}: \mathcal{M}_{z,w;N}(r,R) \to \mathcal{M}_{1,0;N}((z-w)^{-1}r,(z-w)^{-1}R)
$$

defined by

$$
t_{z,w;N;r,R}((x_1,\ldots,x_N))=((z-w)^{-1}(x_1-w),\ldots,(z-w)^{-1}(x_N-w)).
$$

This diffeomorphism leads to a natural isomorphism

 $s_{z,w;N;r,R}: H_N(\widehat{\mathcal{M}}_{1,0;N}((z-w)^{-1}r,(z-w)^{-1}R); \mathbb{Z}) \xrightarrow{\simeq} H_N(\widehat{\mathcal{M}}_{z,w;N}(r,R); \mathbb{Z}).$

Then, by Theorem [4.3,](#page-25-1) we have

$$
\int_{s_{z,w;N;r,R}(\Gamma_{l,m,n})} \mathcal{V}_N(\boldsymbol{a},\boldsymbol{b},\boldsymbol{\gamma};(z-w)^{-1}(x_1-w),\ldots,(z-w)^{-1}(x_N-w))\mathrm{d}\boldsymbol{x} \n= (z-w)^{N+\sum_{i=1}^N (a_i+b_i)-2\sum_{j\n(4.19)
$$

where $dx = dx_1 \wedge \cdots \wedge dx_N$. Using these settings, let us define the regularization of the integrals $\mathcal{I}_{i,j}^{\pm}[E](a,b,c,\gamma;z_1,z_2)$. Since it is sufficient to consider the case where $E \in P_{z_1,z_2,0}$ is a monomial, we can set $E = 1$. First we consider the case $i = j$. Note that for each $(z, w) \in \{(z_1, z_2), (z_1, 0), (z_2, 0)\}$, by choosing (r, R) appropriately, the integrand $\mathcal{U}(\boldsymbol{a}, \boldsymbol{b}, \boldsymbol{c}, \gamma; u, v; z_1, z_2)$ is a singlevalued analytic function on $\widehat{\mathcal{M}}_{z,w:2}(r,R)$. Then for each (i, \pm) , by choosing (r, R, z, w) appropriately, we can pair the two form $\mathcal{U}(\boldsymbol{a}, \boldsymbol{b}, \boldsymbol{c}, \gamma; u, v; z_1, z_2)du \wedge$ dv with the cycles $s_{z,w;2;r,R}(\Gamma_{i,i}^{\pm})$. Let $\delta > 0$ be a small real number satisfying $z_2 > z_1 - z_2 + \delta$ and let

$$
\begin{aligned}\n[\Box^+_{1,1}] &:= (c^+_{1,1}(\boldsymbol{a} + \boldsymbol{b}, \boldsymbol{c}, \gamma))^{-1} s_{z_1,0;2;r,R}(\Gamma^+_{1,1}), & (r, R) &= (z_1 - \delta, \infty), \\
[\Box^+_{0,0}] &:= (c^+_{0,0}(\boldsymbol{a}, \boldsymbol{b}, \gamma))^{-1} s_{z_2,0;2;r,R}(\Gamma^+_{0,0}), & (r, R) &= (0, z_2 + \delta), \\
[\Box^-_{1,1}] &:= (c^-_{1,1}(\boldsymbol{a}, \boldsymbol{c} + \boldsymbol{b}, \gamma))^{-1} s_{z_1,0;2;r,R}(\Gamma^-_{1,1}), & (r, R) &= (z_1 - \delta, \infty), \\
[\Box^-_{0,0}] &:= (c^-_{0,0}(\boldsymbol{b}, \boldsymbol{c}, \gamma))^{-1} s_{z_1, z_2;2;r,R}(\Gamma^-_{0,0}), & (r, R) &= (0, z_1 - z_2 + \delta).\n\end{aligned}
$$

Similar to the argument in [\[Su2,](#page-70-8) Section 6], we can show that each pairing

$$
\int_{\left[\Box_{i,i}^{\kappa}\right]} \mathcal{U}(\boldsymbol{a},\boldsymbol{b},\boldsymbol{c},\gamma;u,v;z_1,z_2) \mathrm{d}u \wedge \mathrm{d}v, \qquad \kappa = \pm, \ i = 1,0 \tag{4.20}
$$

is well-defined and defines the regularization of $\mathcal{I}_{i,i}^{\kappa}[1](a, b, c, \gamma; z_1, z_2)$. This fact can also be seen from the following argument. By the Taylor expansion of the integrand $\mathcal{U}(\mathbf{a}, \mathbf{b}, \mathbf{c}, \gamma; u, v; z_1, z_2)$ and by [\(4.19\)](#page-29-0), we see that the above pairing [\(4.20\)](#page-30-0) admits the following expansion:

1. For $\kappa = +$, we have

$$
\int_{\left[\Box_{1,1}^{+}\right]} U(\mathbf{a}, \mathbf{b}, \mathbf{c}, \gamma; u, v; z_1, z_2) du \wedge dv
$$
\n
$$
= z_1^{\sum_{i=1,2}^{\infty} (a_i + b_i + c_i) - 2\gamma} \int_{\left[\Box_{1,1}^{+}\right]} V_2(\mathbf{a} + \mathbf{b}, \mathbf{c}, \gamma; z_1^{-1}u, z_1^{-1}v) \left(\sum_{k\geq 0} z_2^k F_{1;k}^{+}\right) du \wedge dv
$$
\n
$$
= z_1^{\sum_{i=1,2}^{\infty} (a_i + b_i + c_i) - 2\gamma + 2} \sum_{k\geq 0} \left(\frac{z_2}{z_1}\right)^k \mathcal{J}_{1,1}^{+}[F_{1,k}^{+}](\mathbf{a} + \mathbf{b}, \mathbf{c}, \gamma),
$$

$$
\int_{\left[\Box_{0,0}^{+}\right]} U(\mathbf{a}, \mathbf{b}, \mathbf{c}, \gamma; u, v; z_1, z_2) du \wedge dv
$$
\n
$$
= z_2^{\sum_{i=1,2}^{(a_i+b_i)-2\gamma} z_1^{c_1+c_2} \int_{\left[\Box_{0,0}^{+}\right]} \mathcal{V}_2(\mathbf{a}, \mathbf{b}, \gamma; z_2^{-1}u, z_2^{-1}v) \left(\sum_{k\geq 0} z_1^{-k} F_{0,k}^{+}\right) du \wedge dv
$$
\n
$$
= z_2^{\sum_{i=1,2}^{(a_i+b_i)-2\gamma+2} z_1^{c_1+c_2} \sum_{k\geq 0} \left(\frac{z_2}{z_1}\right)^k \mathcal{J}_{0,0}^{+}[F_{0,k}^{+}](\mathbf{a}, \mathbf{b}, \gamma),
$$
\n(4.21)

where

$$
F_{1;k}^+(u,v) = (k!)^{-1} \partial_z^k (1 - zu^{-1})^{b_1} (1 - zv^{-1})^{b_2} \Big|_{z=0},
$$

$$
F_{0;k}^+(u,v) = (k!)^{-1} \partial_z^k (1 - zu)^{c_1} (1 - zv)^{c_2} \Big|_{z=0},
$$

up to phases, and we use notation $\mathbf{x} + \mathbf{y} = \{x_1 + y_1, x_2 + y_2\}$ for $\boldsymbol{x} = \{x_1, x_2\}, \ \boldsymbol{y} = \{y_1, y_2\}.$

2. For $\kappa = -$, we have

$$
\int_{\left[\Box_{1,1}^{-}\right]} \mathcal{U}(\mathbf{a}, \mathbf{b}, \mathbf{c}, \gamma; u, v; z_1, z_2) du \wedge dv
$$
\n
$$
= z_1^{\sum_{i=1,2}^{\infty} (a_i + b_i + c_i) - 2\gamma} \int_{\left[\Box_{1,1}^{-}\right]} \mathcal{V}_2(\mathbf{a}, \mathbf{c} + \mathbf{b}, \gamma; z_1^{-1} u, z_1^{-1} v)
$$
\n
$$
\cdot \left(\sum_{k \geq 0} (z_1 - z_2)^k F_{1,k}^-(u + 1 - z_1, v + 1 - z_1) \right) du \wedge dv
$$
\n
$$
= z_1^{\sum_{i=1,2}^{\infty} (a_i + b_i + c_i) - 2\gamma + 2} \sum_{k \geq 0} \left(\frac{z_1 - z_2}{z_1} \right)^k \mathcal{J}_{1,1}^-[F_{1,k}^-](\mathbf{a}, \mathbf{c} + \mathbf{b}, \gamma),
$$

$$
(z_1 - z_2)^{-\sum_{i=1,2}(c_i + b_i) + 2\gamma} \int_{\left[\Box_{0,0}^{-}\right]} \mathcal{U}(\mathbf{a}, \mathbf{b}, \mathbf{c}, \gamma; u, v; z_1, z_2) du \wedge dv
$$

\n
$$
= z_1^{a_1 + a_2} \int_{\left[\Box_{0,0}^{-}\right]} \mathcal{V}_2(\mathbf{b}, \mathbf{c}, \gamma; (z_1 - z_2)^{-1}(u - z_2), (z_1 - z_2)^{-1}(v - z_2))
$$

\n
$$
\cdot \left(\sum_{k \ge 0} z_1^{-k} F_{0,k}^{-}(u + 1 - z_1, v + 1 - z_1) \right) du \wedge dv
$$

\n
$$
= (z_1 - z_2)^2 z_1^{a_1 + a_2} \sum_{k \ge 0} \left(\frac{z_1 - z_2}{z_1} \right)^k \mathcal{J}_{0,0}^{-}[F_{0,k}^{-}](\mathbf{b}, \mathbf{c}, \gamma),
$$

\n(4.22)

where

$$
F_{1;k}^-(u,v) = (k!)^{-1} \partial_z^k (1 - z(1 - u)^{-1})^{b_1} (1 - z(1 - v)^{-1})^{b_2} \Big|_{z=0},
$$

$$
F_{0;k}^-(u,v) = (k!)^{-1} \partial_z^k (1 - z(1 - u))^{a_1} (1 - z(1 - v))^{a_2} \Big|_{z=0}.
$$

Since $\mathcal{I}_{i,i}^{\kappa}[1](a, b, c, \gamma; z_1, z_2)$ has the same expansion for an appropriate range of (a, b, c, γ) , the pairing [\(4.20\)](#page-30-0) gives the regularization.

Next, let us define the regularization of $\mathcal{I}_{i,j}^{\pm}[1](a, b, c, \gamma; z_1, z_2)$ for the case $i \neq j$. Note that in this case, the factor $(u - v)^{-2\gamma}$ of the integrand $\mathcal{U}(\boldsymbol{a}, \boldsymbol{b}, \boldsymbol{c}, \gamma; u, v; z_1, z_2)$ is holomorphic on an appropriate neighborhood of $\Box_{i,j}^{\pm}$. Then we define the products of one dimensional cycles

$$
[\Box_{0,1}^+] := (d_{0,1}^+(\boldsymbol{a}, \boldsymbol{b}, \boldsymbol{c}, \gamma))^{-1} s_{z_2,0;1;0,z_2+\delta}(\Gamma_{0,1,0}) \times s_{z_1,0;1;z_1-\delta,\infty}(\Gamma_{0,0,1}),
$$

\n
$$
[\Box_{1,0}^+] := (d_{1,0}^+(\boldsymbol{a}, \boldsymbol{b}, \boldsymbol{c}, \gamma))^{-1} s_{z_1,0;1;z_1-\delta,\infty}(\Gamma_{0,0,1}) \times s_{z_2,0;1;0,z_2+\delta}(\Gamma_{0,1,0}),
$$

\n
$$
[\Box_{0,1}^-] := (d_{0,1}^-(\boldsymbol{a}, \boldsymbol{b}, \boldsymbol{c}, \gamma))^{-1} s_{z_1,z_2;1;0,z_1-z_2+\delta}(\Gamma_{0,1,0}) \times s_{z_1,0;1;z_1-\delta,\infty}(\Gamma_{1,0,0}),
$$

\n
$$
[\Box_{1,0}^-] := (d_{1,0}^-(\boldsymbol{a}, \boldsymbol{b}, \boldsymbol{c}, \gamma))^{-1} s_{z_1,0;1;z_1-\delta,\infty}(\Gamma_{1,0,0}) \times s_{z_1,z_2;1;0,z_1-z_2+\delta}(\Gamma_{0,1,0}),
$$

where

$$
d_{0,1}^+(a, b, c, \gamma) = \mathfrak{e}(a_1)\mathfrak{e}(b_1)\mathfrak{e}(c_2)\mathfrak{e}(a_2 + b_2 + c_2 - 2\gamma),
$$

\n
$$
d_{1,0}^+(a, b, c, \gamma) = \mathfrak{e}(a_2)\mathfrak{e}(b_2)\mathfrak{e}(c_1)\mathfrak{e}(a_1 + b_1 + c_1 - 2\gamma),
$$

\n
$$
d_{0,1}^-(a, b, c, \gamma) = \mathfrak{e}(b_1)\mathfrak{e}(c_1)\mathfrak{e}(a_2)\mathfrak{e}(a_2 + b_2 + c_2 - 2\gamma),
$$

\n
$$
d_{1,0}^-(a, b, c, \gamma) = \mathfrak{e}(b_2)\mathfrak{e}(c_2)\mathfrak{e}(a_1)\mathfrak{e}(a_1 + b_1 + c_1 - 2\gamma).
$$
\n(4.23)

Similar to the arguments for the case $i = j$, we can show

$$
\int_{\left[\Box_{i,j}^{\pm}\right]} U(\boldsymbol{a},\boldsymbol{b},\boldsymbol{c},\gamma;u,v;z_1,z_2) du \wedge dv = \mathcal{I}_{i,j}^{\pm}[1](\boldsymbol{a},\boldsymbol{b},\boldsymbol{c},\gamma;z_1,z_2), \quad i \neq j,
$$

for all $a, b \in \mathbb{C}^2$ and $\gamma \in \mathbb{C}$ for which $\mathcal{I}_{i,j}^{\pm}[1](a, b, c, \gamma; z_1, z_2)$ is defined. Let $\mathcal{H}_{a,b,c,\gamma}$ be a collection of hyperplanes

$$
\mathcal{H}_{a,b,c,\gamma} := \mathcal{H}_{a,b,\gamma} \cup \mathcal{H}_{b,c,\gamma} \cup \mathcal{H}_{a,c,\gamma} \cup \mathcal{H}_{a+b,c,\gamma} \cup \mathcal{H}_{b+c,a,\gamma} \cup \mathcal{H}_{a+c,b,\gamma}, \quad (4.24)
$$

where we use the notation (4.18) . From the explicit forms of the trigonometric functions [\(4.17\)](#page-28-2), [\(4.23\)](#page-32-1), we obtain the following proposition.

Proposition 4.8. Let $i, j \in \{1, 0\}$ and let $z_1, z_2 \in \mathbb{R}_{>0}$ be real numbers satisfying $z_2 > z_1 - z_2 > 0$. Then for $E(u, v) \in P_{z_1, z_2, 0}, \mathcal{I}_{i,j}^{\pm}[E](a, b, c, \gamma; z_1, z_2)$ admits an analytic extension to $\mathbb{C}^7_{a,b,c,\gamma} \setminus \mathcal{H}_{a,b,c,\gamma}$.

4.3 Properties of the Dotsenko-Fateev integrals

From this subsection, we mainly use the following notation

$$
U(a, \rho; u, v; z_1, z_2) := \mathcal{U}(\{a, a'\}, \{a, a'\}, \{a, a'\}, 1; u, v; z_1, z_2),
$$

\n
$$
I_{i,j}^{\pm}[E](a, \rho; z_1, z_2) := \mathcal{I}_{i,j}^{\pm}[E](\{a, a'\}, \{a, a'\}, \{a, a'\}, 1; z_1, z_2),
$$

\n
$$
J_{i,j}^{\pm}[F](a, b, \rho) := \mathcal{J}_{i,j}^{\pm}[F](\{a, a'\}, \{b, b'\}, 1),
$$
\n(4.25)

where

$$
a' = -a\rho', \quad b' = -b\rho', \quad \rho' = 1/\rho \tag{4.26}
$$

(see (4.15)) for the notation U). The condition (4.26) appears naturally in a certain free-field realization of correlation functions (see [\[DF1,](#page-66-6) [DF2\]](#page-66-7) and Subsection [4.4\)](#page-37-0).

Definition 4.9 ([\[Su1\]](#page-70-7)). Fix $x \in \mathbb{C}$ and $\rho \in \mathbb{C} \setminus \{0,1\}$. We define a \mathbb{C} subalgebra $P_x^{DF,\rho}(u, v)$ of $P_x(u, v) = \mathbb{C}[(u-x)^{\pm 1}, (v-x)^{\pm 1}]$ as follows

$$
P_x^{DF,\rho}(u,v) := \left\{ F(u,v) \in P_x(u,v) \middle| (1 - \frac{1}{\rho}) \left(\frac{\partial}{\partial u} F \right) \bigg|_{u=v} = \frac{\partial}{\partial v} \left(F \big|_{u=v} \right) \right\}.
$$

Let us call the Laurent polynomials in Definition [4.9](#page-32-3) "DF-symmetric poly-nomials" according to [\[Su1\]](#page-70-7). For example, we have

$$
(u-x)^n + (-\rho^{-1})(v-x)^n \in P_x^{\text{DF},\rho}(u,v) \quad (n \in \mathbb{Z}).
$$

Note that the relation of $P_x^{\text{DF},\rho}(u, v)$ is equivalent to

$$
(1 - \rho) \left. \left(\frac{\partial}{\partial v} F \right) \right|_{v = u} = \frac{\partial}{\partial u} \left. \left(F \right|_{v = u} \right).
$$

Given a DF-symmetric polynomial $F \in P_x^{\text{DF},\rho}$ $(x = 1,0)$, $J_{0,0}^+[F](a, b, \rho)$ satisfies the following very important property.

Theorem 4.10 ([\[Su1\]](#page-70-7)). Fix $x \in \{1, 0\}$, $\rho \in \mathbb{C} \setminus \{0, 1\}$ and $F(u, v) \in P_x^{\text{DF}, \rho}$. Then, for $J^+_{0,0}[F](a, b, \rho)$, there exists an entire function $\tilde{J}[F] : \mathbb{C}^2 \to \mathbb{C}$ such that

$$
J_{0,0}^+[F](a,b,\rho) = \frac{\sin(\pi(a+b))\sin(\pi(a'+b'))}{\sin(\pi a)\sin(\pi b)\sin(\pi a')\sin(\pi b')} \tilde{J}[F](a,b),
$$

where $a' = -\rho^{-1}a$ and $b' = -\rho^{-1}b$ (see the notation [\(4.26\)](#page-32-2)).

The following transformation formulas hold among $J_{i,j}^{\pm}[F](a, b, \rho)$ $(i, j \in$ $\{1,0\}$).

Theorem 4.11 ([\[DF2,](#page-66-7) [Fo2,](#page-67-5) [Su1\]](#page-70-7)). Fix $x \in \{1,0\}$, $\rho \in \mathbb{C} \setminus \{0,1\}$ and $F(u, v) \in P_x^{\text{DF}, \rho}$. Let $J_{i,j}^{\pm} = J_{i,j}^{\pm}[F](a, b, \rho)$ $(i, j \in \{1, 0\})$ and $s(x) = \sin(\pi x)$. Then we have

$$
J_{1,0}^{+} = \frac{s(a)}{s(a+b)} J_{0,0}^{+},
$$

\n
$$
J_{1,0}^{-} = \frac{s(b)}{s(a+b)} J_{0,0}^{+},
$$

\n
$$
J_{0,1}^{+} = \frac{s(a')}{s(a'+b')} J_{0,0}^{+},
$$

\n
$$
J_{0,1}^{-} = \frac{s(b')}{s(a'+b')} J_{0,0}^{+},
$$

\n
$$
J_{1,1}^{+} = \frac{s(a)s(a')}{s(a+b)s(a'+b')} J_{0,0}^{+},
$$

\n
$$
J_{1,1}^{-} = \frac{s(b)s(b')}{s(a+b)s(a'+b')} J_{0,0}^{+},
$$

up to phase factors.

The following formula was first given explicitly in [\[Fo1\]](#page-67-4).

Theorem 4.12 ([\[DF2,](#page-66-7) [Fo1,](#page-67-4) [Fo2\]](#page-67-5)). We have

$$
J_{0,0}^+[1](a,b,\rho)=(\rho')^2\frac{\sin\pi(a+b)}{\sin(\pi a)}\frac{\Gamma(\rho'-1)}{\Gamma(\rho')}\frac{\Gamma(1+b)\Gamma(1-a-b)}{\Gamma(-a)}\frac{\Gamma(a')\Gamma(b')}{\Gamma(1+a'+b')},
$$

up to a phase factor.

These three theorems will be important in constructing the solutions of the Fuchsian differential equation [\(4.5\)](#page-22-0).

In the following, we introduce transformation and expansion formulas for $I_{i,j}^{\pm}[1](a,\rho;1,z)$. We use the shorthand notation $I_{i,j}^{\pm}(a,\rho;z)=I_{i,j}^{\pm}[1](a,\rho;1,z)$ for $i, j = 1, 0$, and set

$$
\mathcal{H}_{a,\rho}^{\rm DF}:=\mathcal{H}_{\{a,-\rho^{-1}a\},\{a,-\rho^{-1}a\},\{a,-\rho^{-1}a\},1}
$$

(see (4.18) and (4.24)).

Lemma 4.13. Fix $(a, \rho) \in \mathbb{C}_{a,\rho}^2 \setminus \mathcal{H}_{a,\rho}^{\mathrm{DF}}$. We define an involutory matrix

$$
\mathcal{C} = \begin{pmatrix} (cc')^{-1} & -(cc')^{-1} & -(cc')^{-1} & (cc')^{-1} \\ (-c + c^{-1})(c')^{-1} & -(cc')^{-1} & (c - c^{-1})(c')^{-1} & (cc')^{-1} \\ -c^{-1}(c' - (c')^{-1}) & c^{-1}(c' - (c')^{-1}) & -(cc')^{-1} & (cc')^{-1} \\ (c - c^{-1})(c' - (c')^{-1}) & c^{-1}(c' - (c')^{-1}) & (c - c^{-1})(c')^{-1} & (cc')^{-1} \end{pmatrix},
$$

where $c = 2\cos(\pi a)$ and $c' = 2\cos(\pi a')$. Then we have

$$
\begin{pmatrix} I_{1,1}^+(a, \rho; z) \\ I_{0,1}^+(a, \rho; z) \\ I_{1,0}^+(a, \rho; z) \\ I_{0,0}^+(a, \rho; z) \end{pmatrix} = \mathcal{C} \begin{pmatrix} I_{1,1}^-(a, \rho; z) \\ I_{0,1}^-(a, \rho; z) \\ I_{1,0}^-(a, \rho; z) \\ I_{0,0}^-(a, \rho; z) \end{pmatrix}.
$$

Proof. We prove this lemma by similar methods in [\[DF1\]](#page-66-6), [\[Ha,](#page-68-4) Chapter 9] and [\[Su1,](#page-70-7) Proposition 3.10]. Let us show the identity

$$
I_{1,1}^{+}(a, \rho; z) = (cc')^{-1} \big(I_{1,1}^{-}(a, \rho; z) - I_{0,1}^{-}(a, \rho; z) - I_{1,0}^{-}(a, \rho; z) + I_{0,0}^{-}(a, \rho; z) \big).
$$
\n(4.27)

The other identities can be proved in the same way. Let us derive [\(4.27\)](#page-34-0) using $\mathcal{I}_{i,j}^{\pm}[1](\{a,a'\},\{a,a'\},\{a,a'\},\gamma;1,z)$. We use the shorthand notation

$$
\mathcal{I}_{i,j}^{\pm} = \mathcal{I}_{i,j}^{\pm}[1](\{a,a'\},\{a,a'\},\{a,a'\},\gamma;1,z),
$$

$$
\mathcal{U}(u,v) = \mathcal{U}(\{a,a'\},\{a,a'\},\{a,a'\},\gamma;u,v;1,z).
$$

Suppose that a, ρ and γ satisfy $(a, \rho, \gamma) \in \mathbb{C}^3_{a,\rho,\gamma} \setminus \mathcal{H}_{\{a,\rho'^a\},\{a,\rho'^a\},\{a,\rho'^a\},\gamma}$ and

$$
-\frac{1}{2} < \text{Re}(a) \ll 0, \quad -\frac{1}{2} < \text{Re}(a') \ll 0, \quad 0 < \text{Re}(\gamma) \ll 1. \tag{4.28}
$$

Let C^+ be a counterclockwise simple closed contour in the upper half plane ${u \in \mathbb{C} \mid \text{Im}(u) > 0}$, and let C^- be a clockwise simple closed contour in the lower half plane $\{u \in \mathbb{C} \mid \text{Im}(u) < 0\}$. For $\mathcal{U}(u, v)$ with $v \in \mathbb{R}_{>1}$ fixed, we take the branch cut along the real axis $\{u \in \mathbb{R} \mid u \leq v\}$. Then by noting the condition [\(4.28\)](#page-34-1), from the Cauchy's theorem, we obtain

$$
0 = e^{2i\pi\gamma} \oint_{C^+} \left(\int_1^\infty \mathcal{U}(u + i0, v) dv \right) du
$$

= $e^{2i\pi\gamma} \mathcal{I}_{1,1}^+ + e^{i\pi a} \int_z^1 \int_1^\infty \mathcal{U} dv du + e^{2i\pi a} \int_0^z \int_1^\infty \mathcal{U} dv du + e^{3i\pi a} \int_{-\infty}^0 \int_1^\infty \mathcal{U} dv du$

and

$$
0 = e^{-2i\pi\gamma} \oint_{C^-} \left(\int_1^{\infty} \mathcal{U}(u - i0, v) dv \right) du
$$

= $e^{-6i\pi\gamma} \mathcal{I}_{1,1}^+ + e^{-i\pi a} \int_z^1 \int_1^{\infty} \mathcal{U} dv du + e^{-2i\pi a} \int_0^z \int_1^{\infty} \mathcal{U} dv du + e^{-3i\pi a} \int_{-\infty}^0 \int_1^{\infty} \mathcal{U} dv du.$

From these identities, we have

$$
e^{-2i\pi\gamma}s(2a-4\gamma)\mathcal{I}_{1,1}^{+} = s(a)\int_{1}^{z} \int_{1}^{\infty} \mathcal{U} \mathrm{d}v \mathrm{d}u - s(a)\int_{0}^{-\infty} \int_{1}^{\infty} \mathcal{U} \mathrm{d}v \mathrm{d}u, \tag{4.29}
$$

where $s(x) = \sin(\pi x)$. Similarly, we can show the identities

$$
s(2(a' - \gamma)) \int_1^z \int_1^\infty \mathcal{U} \mathrm{d}v \mathrm{d}u = e^{-2i\pi\gamma} s(a' - 4\gamma) \mathcal{I}_{0,0}^- - s(a') \mathcal{I}_{0,1}^-
$$

$$
s(2a') \int_0^{-\infty} \int_1^\infty \mathcal{U} \mathrm{d}v \mathrm{d}u = s(a') \mathcal{I}_{1,0}^- - e^{-2i\pi\gamma} s(a' - 2\gamma) \mathcal{I}_{1,1}^-.
$$

Thus, from these identities and [\(4.29\)](#page-35-0), we obtain the desired identity [\(4.27\)](#page-34-0) by analytically continuing $\mathcal{I}_{i,j}^{\pm}$ to $\gamma = 1$. \Box

From the expansions [\(4.21\)](#page-30-1)-[\(4.22\)](#page-31-0), we obtain the following lemma.

Lemma 4.14. Fix $(a, \rho) \in \mathbb{C}^2_{a,\rho} \setminus \mathcal{H}_{a,\rho}^{\text{DF}}$. Then $I^{\pm}_{i,j}(a, \rho; z)$ admits the following expansion.

1. The functions $z^{2(i-1)(a+a')} I_{i,i}^+(a, \rho; z)$ and $(1-z)^{2(i-1)(a+a')} I_{i,i}^-(a, \rho; z)$ $(i =$ 1, 0) admit analytic continuations to the complex domains ${|z| < 1}$ and ${|z-1| < 1}$, respectively, and on these domains, satisfy convergent series

$$
z^{2(i-1)(a+a')} I_{i,i}^+(a,\rho;z) = \sum_{k\geq 0} J_{i,i}^+[F_{i,k}^+]((i+1)a,a,\rho)z^k,
$$

$$
(1-z)^{2(i-1)(a+a')} I_{i,i}^-(a,\rho;z) = \sum_{k\geq 0} J_{i,i}^-[F_{i,k}^-](a,(i+1)a,\rho)(1-z)^k,
$$
where $F_{1:}^+$ $\hat{u}_{1;k}^{+}(u,v)$ is defined by

$$
F_{1;k}^{+}(u,v) = (k!)^{-1} \partial_z^k ((1 - zu^{-1})^a (1 - zv^{-1})^{a'}) \Big|_{z=0},
$$

and

$$
F_{0;k}^+(u,v) = F_{1;k}^+(u^{-1}, v^{-1}), \qquad F_{1;k}^-(u,v) = F_{1;k}^+(u-1, v-1),
$$

$$
F_{0;k}^-(u,v) = F_{1;k}^+((1-u)^{-1}, (1-v)^{-1}).
$$

up to phases.

2. For $i \neq j$, $z^{2(i-1)a+2(j-1)a'} I_{i,j}^+(a, \rho; z)$ and $(1-z)^{2(i-1)a+2(j-1)a'} I_{i,j}^-(a, \rho; z)$ admit analytic continuations to the complex domains $\{|z| < 1\}$ and ${|z - 1| < 1}$, respectively, and on these domains, satisfy convergent series

$$
\sum_{k\geq 0} \mathcal{J}_{i,j}^+[F^+_{\{i,j\};k}] (\{(i+1)a-2i,(j+1)a'-2j\},\{a,a'\},0) z^{1+k},
$$

$$
\sum_{k\geq 0} \mathcal{J}_{i,j}^- [F^-_{\{i,j\};k}] (\{a,a'\},\{(i+1)a-2i,(j+1)a'-2j\},0) (1-z)^{1+k},
$$

respectively, where by setting

$$
G_k(x_1, x_2) = (k!)^{-1} \partial_z^k ((1 - zx_1)^a (1 - zx_2)^{a'} (1 - zx_1x_2)^{-2}) \Big|_{z=0},
$$

 $F^\pm_{\ell i}$ $\mathcal{E}^{\pm}_{\{i,j\};k}$ are defined by

$$
F_{\{0,1\};k}^+(u,v) = G_k(u,v^{-1}), \qquad F_{\{1,0\};k}^+(u,v) = G_k(u^{-1},v),
$$

$$
F_{\{0,1\};k}^-(u,v) = G_k(1-u,(1-v)^{-1}), \quad F_{\{1,0\};k}^-(u,v) = G_k((1-u)^{-1},1-v)
$$

up to phases.

- Remark 4.15. 1. We can see that $\partial_z^n((1-zu)^a(1-zv)^{a'})\big|_{z=0} \in P_0^{\text{DF},\rho}$ $\frac{D\mathbf{r}, \rho}{0}$. Then we have $F_{1:}^+$ $T_{1;k}^+, F_{0;k}^+ \in \mathcal{P}_0^{\mathcal{DF},\rho}$ and $F_{1;k}^ T_{1;k}^-, F_{0;k}^- \in \mathcal{P}_1^{\mathcal{DF},\rho}$ $1^{\text{DF }, \rho}$.
	- 2. From Definition [4.5](#page-27-0) and the formulas of the beta integrals, we have

$$
\mathcal{J}_{0,1}^{\pm}[1](\boldsymbol{a},\boldsymbol{b},0)=\frac{\Gamma(a_1+1)\Gamma(b_1+1)\Gamma(-a_2-b_2-1)\Gamma(b_2+1)^{\frac{1\pm1}{2}}\Gamma(a_2+1)^{\frac{1\pm1}{2}}}{\Gamma(a_1+b_1+2)\Gamma(-a_2)^{\frac{1\pm1}{2}}\Gamma(-b_2)^{\frac{1\pm1}{2}}},\\ \mathcal{J}_{1,0}^{\pm}[1](\boldsymbol{a},\boldsymbol{b},0)=\frac{\Gamma(a_2+1)\Gamma(b_2+1)\Gamma(-a_1-b_1-1)\Gamma(b_1+1)^{\frac{1\pm1}{2}}\Gamma(a_1+1)^{\frac{1\pm1}{2}}}{\Gamma(a_2+b_2+2)\Gamma(-a_1)^{\frac{1\pm1}{2}}\Gamma(-b_1)^{\frac{1\pm1}{2}}},
$$

up to phase factors (see [\[Su2,](#page-70-0) Appendix A]).

4.4 Construction of solutions

We want to use the Dotsenko-Fateev integrals $I_{i,j}^{\pm}[1](a,\rho;1,z)$ $(i,j=1,0)$ with parameters $a = \alpha_{-}\beta_{2,2}, \ \rho = \alpha_{-}^2$ (for the definition of $I_{i,j}^{\pm}[1](a,\rho;1,z)$, see [\(4.25\)](#page-32-0)). However, in this case, we have $a' = \alpha_{+}\beta_{2,2} = -m \in \mathbb{Z}$, and then the integral cannot be defined. So let us add a small complex parameter to the variables (a, ρ) to make the integrals well-defined. Before that, let us define some notation. We set

$$
\mathbb{D}(x) := \{ x \in \mathbb{C} \mid |x| < 1 \}, \quad \mathbb{D}^{\times}(x) := \mathbb{D}(x) \setminus \{ x = 0 \}.
$$

We also use the shorthand notation $\mathbb{D} = \mathbb{D}(x)$ and $\mathbb{D}^{\times} = \mathbb{D}^{\times}(x)$. For any non-empty open set $U \in \mathbb{C}^n$, we define

$$
\mathcal{O}(U) := \{ f \in F(U) \mid f \text{ is holomorphic on } U \},
$$

where $F(U)$ is the set of complex functions on U.

Remark 4.16. In the above definition, we say that $f(z_1, \ldots, z_n) \in F(U)$ is holomorphic on $U \in \mathbb{C}^n$, when $f(z_1, \ldots, z_n)$ is holomorphic for each variable z_i $((z_1, \ldots, z_n) \in U)$.

Following [\[TW2\]](#page-71-0), we introduce ϵ -deformations of α_{\pm} , α_0 and $\beta_{r,s}$ ($r, s \in \mathbb{Z}$) as follows (see $(2.11)-(2.12)$ $(2.11)-(2.12)$)

$$
\widetilde{\alpha}_{+}(\epsilon) = \alpha_{+} + \theta \epsilon, \qquad \widetilde{\alpha}_{-}(\epsilon) = -\frac{1}{\widetilde{\alpha}_{+}(\epsilon)},
$$

$$
\widetilde{\alpha}_{0}(\epsilon) = \widetilde{\alpha}_{+}(\epsilon) + \widetilde{\alpha}_{-}(\epsilon), \quad \widetilde{\beta}_{r,s} = \frac{(1-r)}{2}\widetilde{\alpha}_{+} + \frac{(1-s)}{2}\widetilde{\alpha}_{-},
$$

where we fix a sufficiently small $\theta \in \mathbb{R}_{>0}$ to satisfy the following condition

$$
(a,\rho) = (\widetilde{\alpha}_{-}(\epsilon)\widetilde{\beta}_{2,2}(\epsilon), \widetilde{\alpha}_{-}(\epsilon)^2) \notin \mathcal{H}_{a,\rho}^{\mathrm{DF}} \qquad (\epsilon \in \mathbb{D}^{\times}(\epsilon)). \tag{4.30}
$$

Note that $\tilde{\alpha}_{\pm}, \tilde{\alpha}_0, \beta_{r,s} \in \mathcal{O}(\mathbb{D}), \ \tilde{\alpha}_{\pm}(0) = \alpha_{\pm}, \ \tilde{\alpha}_0(0) = \alpha_0$ and $\beta_{r,s}(0) =$ $\beta_{r,s}$. Consider the integrals $I_{i,j}^{\pm}[1](\widetilde{\alpha}_{-}(\epsilon)\widetilde{\beta}_{2,2}(\epsilon), \widetilde{\alpha}_{-}(\epsilon)^2; 1, z)$ for $i, j = 1, 0$. Since [\(4.30\)](#page-37-0), from Proposition [4.8,](#page-32-1) we see that these integrals are welldefined for $\epsilon \in \mathbb{D}^{\times}$. We choose branches of $I_{i,j}^{\pm}[1](\widetilde{\alpha}_{-}(\epsilon)\widetilde{\beta}_{2,2}(\epsilon), \widetilde{\alpha}_{-}(\epsilon)^2; 1, z)$ at $z=\frac{1}{2}$ $\frac{1}{2}$ as $\arg z = \arg(1-z) = 0$. Then by the analytic continuations, $I_{i,j}^{\pm}[1](\widetilde{\alpha}_{-}(\epsilon)\widetilde{\beta}_{2,2}(\epsilon),\widetilde{\alpha}_{-}(\epsilon)^{2};1,z)$ are single-valued functions on the simplyconnected domain $\mathbb{D}(z) \cap \mathbb{D}(1-z) = \{|z| < 1\} \cap \{|z-1| < 1\}.$

Proposition 4.17. 1. For $i, j = 1, 0$, we have

$$
\epsilon I_{i,j}^{\pm}[1](\widetilde{\alpha}_{-}(\epsilon)\widetilde{\beta}_{2,2}(\epsilon),\widetilde{\alpha}_{-}(\epsilon)^{2};1,z)\in\mathcal{O}((\mathbb{D}(z)\cap\mathbb{D}(1-z))\times\mathbb{D}(\epsilon)).
$$

2. Let $C_0 \subset \mathbb{D}^{\times}(\epsilon)$ be a simple closed curve arround $\epsilon = 0$. Then for $z \in \mathbb{D}(z) \cap \mathbb{D}(1-z)$ and $i, j = 1, 0$, we have

$$
\int_{C_0} I_{i,j}^{\pm}[1](\widetilde{\alpha}_{-}(\epsilon)\widetilde{\beta}_{2,2}(\epsilon),\widetilde{\alpha}_{-}(\epsilon)^2;1,z)d\epsilon \neq 0,
$$

and asymptotic behaviors

$$
\int_{C_0} I_{i,j}^{\kappa}[1](\widetilde{\alpha}_{-}(\epsilon)\widetilde{\beta}_{2,2}(\epsilon), \widetilde{\alpha}_{-}(\epsilon)^2; 1, z) d\epsilon \sim\n\begin{cases}\nC_{i,j}^+ z^{\lambda_{i,j}} & (\kappa = +, z \to 0), \\
C_{i,j}^-(1-z)^{\lambda_{i,j}} & (\kappa = -, z \to 1),\n\end{cases}
$$
\nwhere $\lambda_{i,j} = 2(1-i)\alpha_{-}\beta_{2,2} + 2(1-j)\alpha_{+}\beta_{2,2} + (1-i)j + (1-j)i$ and $C_{i,j}^{\pm}$ are some non-zero constants.

Proof. We only prove the case $i = j$. The cases $i \neq j$ can be proved in the same way. We use the shorthand notation

$$
J_{i,i}^+[F_{i,k}^+](\epsilon) = J_{i,i}^+[F_{i,k}^+]((i+1)a,a,\rho), \quad J_{i,i}^-[F_{i,k}^-](\epsilon) = J_{i,i}^-[F_{i,k}^-](a,(i+1)a,\rho),
$$

$$
I_{i,j}^{\pm}(z,\epsilon) = I_{i,j}^{\pm}[1](a,\rho;1,z),
$$

as $a = \tilde{\alpha}_{-}(\epsilon)\tilde{\beta}_{2,2}(\epsilon)$, $a' = \tilde{\alpha}_{+}(\epsilon)\tilde{\beta}_{2,2}(\epsilon)$ and $\rho = \tilde{\alpha}_{-}(\epsilon)^2$, where $F_{1;i}^{+}$ $E_{1;k}^+$, $F_{0;k}^+$ \in ${\rm P}_0^{{\rm DF},\widetilde\alpha_-(\epsilon)^2}$ $_{0}^{\text{DF},\tilde{\alpha}_{-}(\epsilon)^{2}}$ and $F_{1,i}^{-}$ $F_{1;k}^-$, $F_{0;k}^ \in$ $\mathrm{P}_1^{\mathrm{DF},\widetilde{\alpha}_{-}(\epsilon)^2}$ $1^{\text{Dr}, \alpha = (\epsilon)}$ are defined in Lemma [4.14](#page-35-0) (see also Remark [4.15\)](#page-36-0).

By Proposition [4.8](#page-32-1) and by Lemma [4.14,](#page-35-0) we see that

$$
I_{i,i}^{\pm}(z,\epsilon) \in \mathcal{O}((\mathbb{D}(z) \cap \mathbb{D}(1-z)) \times \mathbb{D}^{\times}(\epsilon)) \quad (i=1,0). \tag{4.32}
$$

By Theorems [4.7,](#page-28-0) [4.10](#page-33-0) and [4.11,](#page-33-1) we see that

$$
\epsilon J_{i,i}^+[F_{i,k}^+](\epsilon), \ \epsilon J_{i,i}^-[F_{i,k}^-](\epsilon) \in \mathcal{O}(\mathbb{D}(\epsilon)) \quad (k \ge 0, \ i = 1, 0). \tag{4.33}
$$

Furthermore, by Theorems [4.11](#page-33-1)[-4.12,](#page-33-2) we can see that

$$
\text{Res}_{\epsilon=0} J_{i,j}^{\pm}[1](\epsilon) \neq 0 \quad (i, j = 1, 0). \tag{4.34}
$$

Then by Lemma [4.14](#page-35-0) and by [\(4.32\)](#page-38-0)-[\(4.33\)](#page-38-1), we see that $\epsilon I_{i,i}^{\pm}(z,\epsilon)$ $(i = 1,0)$ are holomorphic for $\epsilon \in \mathbb{D}(\epsilon)$.

Let $C_0 \subset \mathbb{D}^{\times}(\epsilon)$ be a simple closed curve arround $\epsilon = 0$. By Lemma [4.14](#page-35-0) and by [\(4.33\)](#page-38-1)-[\(4.34\)](#page-38-2), we see that $\int_{C_0} I_{i,i}^{\pm}(z,\epsilon) d\epsilon \neq 0$ (i = 1,0) and $\int_{C_0} I_{i,i}^{\pm}(z,\epsilon) d\epsilon$ satisfy the asymptotic behavior [\(4.31\)](#page-38-3). \Box We introduce ϵ -deformations of the screening currents [\(2.13\)](#page-14-0) as follows

$$
Q_{\pm}^{(\epsilon)}(z) = b(z)V_{\widetilde{\alpha}_{\pm}(\epsilon)}(z).
$$

We set $T^{(\epsilon)}(z) = T^{(\tilde{\alpha}_0(\epsilon))}(z)$, $G^{(\epsilon)}(z) = G^{(\tilde{\alpha}_0(\epsilon))}(z)$, $L_n^{(\epsilon)} = L_n^{(\tilde{\alpha}_0(\epsilon))}$ and $G_r^{(\epsilon)} =$ $G_r^{(\tilde{\alpha}_0(\epsilon))}$ (see [\(2.8\)](#page-11-0)-[\(2.10\)](#page-11-1) for the right-hand notation). From the definition of these operators, we have the operator product expansions

$$
T^{(\epsilon)}(z)Q_{\pm}^{(\epsilon)}(w) = \partial_w \frac{Q_{\pm}^{(\epsilon)}(w)}{z-w} + \cdots, \quad G^{(\epsilon)}(z)Q_{\pm}^{(\epsilon)}(w) = \frac{1}{\widetilde{\alpha}_{\pm}(\epsilon)} \partial_w \frac{V_{\widetilde{\alpha}_{\pm}(\epsilon)}(w)}{z-w} + \cdots \tag{4.35}
$$

Then the operators $Q_{\pm}^{(\epsilon)}(z)$ define the screening currents for $U^{(\epsilon)}(\mathfrak{ns})$, where $U^{(\epsilon)}(\mathfrak{ns})$ is the universal enveloping algebra of \mathfrak{ns} with the central charge fixed to $c_{\widetilde{\alpha}_0(\epsilon)}$.

For β, β' , let $Y_{\beta,\beta'}^{(\epsilon)}$ be a non-zero even free field $\mathcal{F}_{\alpha_0(\epsilon)}$ -intertwining operator of type $\begin{pmatrix} F_{\beta+\beta'} \\ F_{\beta+\beta'} \end{pmatrix}$ F_β $F_{\beta'}$). Note that by the restriction, $Y_{\beta,\beta'}^{(\epsilon)}$ gives an even **ns**-intertwining operator of type $\begin{pmatrix} U^{(\epsilon)}(\mathfrak{n}\mathfrak{s})|\beta+\beta' \end{pmatrix}$ $U^{(\epsilon)}(\mathfrak{ns}) \ket{\beta} \ \, U^{(\epsilon)}(\mathfrak{ns}) \ket{\beta'}$. From the last argu-ment in Subsection [2.2,](#page-8-0) we can fix the normalization of $Y_{\beta,\beta'}^{(\epsilon)}$ as $Y_{\beta,\beta'}^{(\epsilon)}(|\beta\rangle, z)$ $V_\beta(z)$. Then, based on the free-field realization techniques in [\[DF1,](#page-66-0) [DF2,](#page-66-1) [Fe\]](#page-67-0). we define

$$
R_{i,j}^{0,\pm}(z_1, z_2, \epsilon)
$$
\n
$$
= \int_{\left[\Box_{i,j}^{\pm}\right]} \langle \widetilde{\beta}_{2,2}(\epsilon)|Q_{-}^{(\epsilon)}(u)Q_{+}^{(\epsilon)}(v)V_{\widetilde{\beta}_{2,2}(\epsilon)}(z_1)V_{\widetilde{\beta}_{2,2}(\epsilon)}(z_2)|\widetilde{\beta}_{2,2}(\epsilon)\rangle \, du dv
$$
\n
$$
= z_1^{\widetilde{\beta}_{2,2}(\epsilon)\widetilde{\beta}_{2,2}(\epsilon)} z_2^{\widetilde{\beta}_{2,2}(\epsilon)\widetilde{\beta}_{2,2}(\epsilon)}(z_1 - z_2)^{\widetilde{\beta}_{2,2}(\epsilon)\widetilde{\beta}_{2,2}(\epsilon)} I_{i,j}^{\pm}[1](\widetilde{\alpha}_{-}(\epsilon)\widetilde{\beta}_{2,2}(\epsilon), \widetilde{\alpha}_{-}(\epsilon)^2; z_1, z_2),
$$

$$
R_{i,j}^{1,\pm}(z_1, z_2, \epsilon)
$$
\n
$$
= \int_{\left[\Box_{i,j}^{\pm}\right]} \langle \widetilde{\beta}_{2,2}(\epsilon)|Q_{-}^{(\epsilon)}(u)Q_{+}^{(\epsilon)}(v)[G_{-\frac{1}{2}}^{(\epsilon)}, V_{\widetilde{\beta}_{2,2}(\epsilon)}(z_1)][G_{-\frac{1}{2}}^{(\epsilon)}, V_{\widetilde{\beta}_{2,2}(\epsilon)}(z_2)]|\widetilde{\beta}_{2,2}(\epsilon)\rangle \text{d}u \text{d}v
$$
\n
$$
= \widetilde{\beta}_{2,2}(\epsilon)\widetilde{\beta}_{2,2}(\epsilon)z_1^{\widetilde{\beta}_{2,2}(\epsilon)}\widetilde{\beta}_{2,2}(\epsilon)z_2^{\widetilde{\beta}_{2,2}(\epsilon)}z_2^{\widetilde{\beta}_{2,2}(\epsilon)}(z_1 - z_2)^{\widetilde{\beta}_{2,2}(\epsilon)}\widetilde{\beta}_{2,2}(\epsilon) - 1
$$
\n
$$
\times I_{i,j}^{\pm}[E](\widetilde{\alpha}_{-}(\epsilon)\widetilde{\beta}_{2,2}(\epsilon), \widetilde{\alpha}_{-}(\epsilon)^{2}; z_1, z_2), \tag{4.36}
$$

for $i, j = 1, 0$ and $\epsilon \in \mathbb{D}^{\times}$, where $E(u, v) = ((u - z_1)(u - z_2)(v - z_1)(v - z_2))$ (z_2))⁻¹ and we use the operator product expansions [\(2.5\)](#page-10-0) and [\(2.7\)](#page-11-2). By Proposition [4.8,](#page-32-1) these functions are well-defined for $\epsilon \in \mathbb{D}^{\times}$. Then, from the following lemma, we see that each $R_{i,j}^{0,\pm}(z_1, z_2, \epsilon)$ realizes a four point correlation function with respect to $U^{(\epsilon)}(\mathfrak{ns}).$

Lemma 4.18. Let $i, j \in \{1, 0\}$, $\epsilon \in \mathbb{D}^{\times}$ and $e_1, e_2 \in F_{\tilde{\beta}_{2,2}(\epsilon)}$. Then for any $A \in U^{(\epsilon)}(\mathfrak{ns}),\$ the function

$$
\int_{\left[\Box_{i,j}^{\pm}\right]} \langle \widetilde{\beta}_{2,2}(\epsilon)|[A,Q_{-}^{(\epsilon)}(u)Q_{+}^{(\epsilon)}(v)]Y^{(\epsilon)}(e_1,z_1)Y^{(\epsilon)}(e_2,z_2)|\widetilde{\beta}_{2,2}(\epsilon)\rangle \mathrm{d}u\mathrm{d}v
$$

is identically zero, where we omit the subscripts of the intertwining operators $Y_{\beta,\beta'}^{(\epsilon)}$.

Proof. By the operator expansions (2.9) and (4.35) , we see that the function becomes a total derivative form

$$
z_1^{\tilde{\beta}_{2,2}(\epsilon)\tilde{\beta}_{2,2}(\epsilon)} z_2^{\tilde{\beta}_{2,2}(\epsilon)\tilde{\beta}_{2,2}(\epsilon)}(z_1 - z_2)^{\tilde{\beta}_{2,2}(\epsilon)\tilde{\beta}_{2,2}(\epsilon)}
$$

$$
\times \left[f \int_{\left[\Box_{i,j}^{\pm}\right]} d_{u,v} \left(U(\tilde{\alpha}_{-}(\epsilon)\tilde{\beta}_{2,2}(\epsilon), \tilde{\alpha}_{-}(\epsilon)^2; u, v; z_1, z_2) \left(E \mathrm{d}u + F \mathrm{d}v \right) \right) + g \int_{\left[\Box_{i,j}^{\pm}\right]} d_{u,v} \left(U^{\frac{1}{2}}(\tilde{\alpha}_{-}(\epsilon)\tilde{\beta}_{2,2}(\epsilon), \tilde{\alpha}_{-}(\epsilon)^2; u, v; z_1, z_2) \left(E' \mathrm{d}u + F' \mathrm{d}v \right) \right)
$$

for some $f, g \in \mathbb{C}[z_1^{\pm 1}, z_2^{\pm 1}, (z_1 - z_2)^{-1}]$ and Laurent polynomials E, F, E', F' in $\mathbb{C}[u^{\pm 1}, v^{\pm 1}, (u-z_1)^{-1}, (v-z_1)^{-1}, (u-z_2)^{-1}, (v-z_2)^{-1}, (u-v)^{-1}]$, where $d_{u,v}$ is the total derivative with respect to u, v and we use the notation

$$
U^{\frac{1}{2}}(a,\rho;u,v;z_1,z_2) = \mathcal{U}(\{a,a'\},\{a,a'\},\{a,a'\},2^{-1};u,v;z_1,z_2)
$$

as $a = \tilde{\alpha}_{-}(\epsilon)\beta_{2,2}(\epsilon)$ and $\rho = \tilde{\alpha}_{-}(\epsilon)^2$. Thus, by Theorem [4.6](#page-28-1) and by the Stokes theorem, this function must vanish. \Box

We define $\overline{R}_{i,j}^{0,\pm}(z,\epsilon)=R_{i,j}^{0,\pm}(1,z,\epsilon),$ $\overline{R}_{i,j}^{1,\pm}(z,\epsilon)=R_{i,j}^{1,\pm}(1,z,\epsilon).$ From Lemma [4.18](#page-40-0) and the $L_0^{(\epsilon)}$ $_{0}^{(e)}$ -conjugation formula for intertwining operators, we have

$$
R_{i,j}^{0,\pm}(z_1, z_2, \epsilon) = z_1^{-2h_{\tilde{\beta}_{2,2}(\epsilon)}} \overline{R}_{i,j}^{0,\pm}(z_2/z_1, \epsilon),
$$

\n
$$
R_{i,j}^{1,\pm}(z_1, z_2, \epsilon) = z_1^{-2h_{\tilde{\beta}_{2,2}(\epsilon)} - 1} \overline{R}_{i,j}^{1,\pm}(z_2/z_1, \epsilon),
$$
\n(4.37)

where $h_{\tilde{\beta}_{2,2}(\epsilon)} = 2^{-1} \tilde{\beta}_{2,2}(\epsilon) (\tilde{\beta}_{2,2}(\epsilon) - \tilde{\alpha}_0(\epsilon))$ (see [\(2.4\)](#page-9-0)). Then by Proposition [4.17](#page-38-2) and by the definition [\(4.36\)](#page-39-1), we see that $\int_{C_0} \overline{R}_{i,j}^{0,\pm}(z,\epsilon) d\epsilon$ is nonzero for any simple closed curve $C_0 \subset \mathbb{D}^{\times}(\epsilon)$ arround $\epsilon = 0$. Then we define

$$
\Psi_{i,j}^{\pm}(z) := \int_{C_0} \overline{R}_{i,j}^{0,\pm}(z,\epsilon) d\epsilon
$$
\n(4.38)

for $i, j = 1, 0$. Note that these functions are invariant under the homotopy deformation of C_0 in $\mathbb{D}^{\times}(\epsilon)$.

Proposition 4.19. The functions $\Psi_{i,j}^{\pm}(z)$ $(i, j = 1, 0)$ satisfy the Fuchsian differential equation (4.5) .

Proof. From the structure of the Fock module $F_{2,2}$ (see Proposition [2.9\)](#page-13-2), we see that the lowest weight vector $|\beta_{2,2}(0)\rangle = |\beta_{2,2}\rangle \in F_{2,2}$ satisfies the relation $S_{2,2}^0 \ket{\beta_{2,2}} = 0$, where $S_{2,2}^0$ is defined by [\(4.3\)](#page-21-0) with the central charge fixed to $c_{1,2m+1}$. We define an ϵ -deformation of $S_{2,2}^0$ as follows

$$
S_{2,2}(\epsilon) = s_1(\epsilon)(L_{-1}^{(\epsilon)})^2 + s_2(\epsilon)L_{-2}^{(\epsilon)} - s_1(\epsilon)G_{-\frac{3}{2}}^{(\epsilon)}G_{-\frac{1}{2}}^{(\epsilon)} \in U^{(\epsilon)}(\mathfrak{ns}),
$$

where

$$
s_1(\epsilon) = \frac{4\widetilde{\alpha}_+(\epsilon)^2}{1 - \widetilde{\alpha}_+(\epsilon)^4}, \qquad s_2(\epsilon) = -\frac{2 - 2\widetilde{\alpha}_+(\epsilon)^2}{\widetilde{\alpha}_+(\epsilon)^2 + 1}.
$$

From the definition, we see that

$$
S_{2,2}(0) = S_{2,2}^0, \quad s_1(\epsilon), s_2(\epsilon) \in \mathcal{O}(\mathbb{D}(\epsilon)) \setminus \epsilon \mathcal{O}(\mathbb{D}(\epsilon)), \tag{4.39}
$$

where $\epsilon \mathcal{O}(\mathbb{D}(\epsilon))$ is the subring of $\mathcal{O}(\mathbb{D}(\epsilon))$ defined by

$$
\left\{ f(\epsilon) \in \mathcal{O}(\mathbb{D}(\epsilon)) \mid \frac{f(\epsilon)}{\epsilon} \in \mathcal{O}(\mathbb{D}(\epsilon)) \right\}.
$$

By a straightforward calculation, we see that the lowest weight vector $|\tilde{\beta}_{2,2}(\epsilon)\rangle$ satisfies the relation $S_{2,2}(\epsilon)|\beta_{2,2}(\epsilon)\rangle = 0$ (cf. [\[BS,](#page-65-0) [IK2\]](#page-69-0)). Then, by Lemma [4.18,](#page-40-0) similar to [\(4.4\)](#page-21-1), we can show that $R^{0,\pm}_{i,j}(z_1, z_2, \epsilon)$ and $R^{1,\pm}_{i,j}(z_1, z_2, \epsilon)$ satisfy the following differential equations

$$
\begin{split}\n&\left\{ \left(-2s_{1}(\epsilon)h_{\tilde{\beta}_{2,2}(\epsilon)} + s_{2}(\epsilon)h_{\tilde{\beta}_{2,2}(\epsilon)} + \frac{s_{2}(\epsilon)}{2} \right) \left(\frac{1}{z_{1}^{2}} + \frac{1}{z_{2}^{2}} \right) + s_{1}(\epsilon)(\partial_{z_{1}} + \partial_{z_{2}})^{2} \right. \\
&\left. + (s_{1}(\epsilon) - s_{2}(\epsilon)) \left(\frac{\partial_{z_{1}}}{z_{1}} + \frac{\partial_{z_{2}}}{z_{2}} \right) \right\} R_{i,j}^{1,\pm}(z_{1},z_{2},\epsilon) \\
&+ \left\{ -s_{1}(\epsilon) \left(\frac{1}{z_{2}} - \frac{1}{z_{1}} \right) \partial_{z_{1}} \partial_{z_{2}} + 2s_{1}(\epsilon) h_{\tilde{\beta}_{2,2}(\epsilon)} \left(\frac{\partial_{z_{1}}}{z_{2}^{2}} - \frac{\partial_{z_{2}}}{z_{1}^{2}} \right) \right\} R_{i,j}^{0,\pm}(z_{1},z_{2},\epsilon) = 0, \\
&\left\{ s_{2}(\epsilon)h_{\tilde{\beta}_{2,2}(\epsilon)} \left(\frac{1}{z_{1}^{2}} + \frac{1}{z_{2}^{2}} \right) + s_{1}(\epsilon)(\partial_{z_{1}} + \partial_{z_{2}})^{2} \right\} R_{i,j}^{0,\pm}(z_{1},z_{2},\epsilon) \\
&+ (s_{1}(\epsilon) - s_{2}(\epsilon)) \left(\frac{\partial_{z_{1}}}{z_{1}} + \frac{\partial_{z_{2}}}{z_{2}} \right) R_{i,j}^{0,\pm}(z_{1},z_{2},\epsilon) - s_{1}(\epsilon) \left(\frac{1}{z_{1}} - \frac{1}{z_{2}} \right) R_{i,j}^{1,\pm}(z_{1},z_{2},\epsilon) = 0. \tag{4.40}\n\end{split}
$$

Thus by [\(4.37\)](#page-40-1) and [\(4.40\)](#page-41-0), $\overline{R}_{i,j}^{0,\pm}(z,\epsilon)$ satisfy a fourth-order differential equation. Let $\mathcal{D}_z^{(\epsilon)}$ be the fourth-order differential operator defined by

$$
\mathcal{D}_z^{(\epsilon)} \overline{R}_{i,j}^{0,\pm}(z,\epsilon) = 0, \quad \mathcal{D}_z^{(\epsilon)} = \partial_z^4 + \text{(lower order terms for } \partial_z^k),
$$

and let \mathcal{D}_z be the fourth-order differential operator of [\(4.5\)](#page-22-0). Then, by [\(4.39\)](#page-41-1), we can see that

$$
\mathcal{D}_z^{(\epsilon)} - \mathcal{D}_z \in \epsilon \mathcal{O}(\mathbb{D}(\epsilon))[z, z^{-1}, (z-1)^{-1}]\langle \partial_z \rangle, \tag{4.41}
$$

where the right hand side is the ring of linear differential operators with coefficients in the ring $\epsilon \mathcal{O}(\mathbb{D}(\epsilon))[z, z^{-1}, (z-1)^{-1}] := \epsilon \mathcal{O}(\mathbb{D}(\epsilon)) \otimes \mathbb{C}[z, z^{-1}, (z-1)^{-1}]$ 1)[−]¹]. Thus by Proposition [4.17,](#page-38-2) we have

$$
0 = \int_{C_0} \mathcal{D}_z^{(\epsilon)} \overline{R}_{i,j}^{0,\pm}(z,\epsilon) d\epsilon
$$

=
$$
\int_{C_0} (\mathcal{D}_z^{(\epsilon)} - \mathcal{D}_z) \overline{R}_{i,j}^{0,\pm}(z,\epsilon) d\epsilon + \int_{C_0} \mathcal{D}_z \overline{R}_{i,j}^{0,\pm}(z,\epsilon) d\epsilon
$$

=
$$
\mathcal{D}_z \int_{C_0} \overline{R}_{i,j}^{0,\pm}(z,\epsilon) d\epsilon.
$$

Therefore, $\Psi_{i,j}^{\pm}$ satisfy the Fuchsian differential equation [\(4.5\)](#page-22-0).

 \Box

By noting the Riemann scheme [\(4.6\)](#page-22-1) of [\(4.5\)](#page-22-0), from Propositions [4.17](#page-38-2) and [4.19,](#page-41-2) we obtain the following proposition.

Proposition 4.20. The sets $\{\Psi_{i,j}^{+}(z) | i,j = 1,0\}$ and $\{\Psi_{i,j}^{-}(z) | i,j = 1,1\}$ 1, 0} are fumdamental systems of solutions of (4.5) at $z = 0$ and $z = 1$, respectively. Let $\rho_{i,j}^+$ and $\rho_{i,j}^-$ be the characteristic exponents of $\Psi_{i,j}^+(z)$, $\Psi_{i,j}^-(z)$ at $z = 0$ and $z = \tilde{1}$, respectively. Then we have $\rho_{i,j}^{\pm} = \rho_{i,j}$ for all $i, j = \tilde{1}, 0$, where $\rho_{i,j}$ are defined by [\(4.7\)](#page-23-0).

Remark 4.21. The important parts of the above construction of solutions are Lemma [4.18,](#page-40-0) Proposition [4.17](#page-38-2) and [\(4.41\)](#page-42-0). The explicit forms of \mathcal{D}_z and $\mathcal{D}_z^{(\epsilon)}$ are not important. Hence, we believe that our construction is applicable to the correlation functions of other logarithmic minimal models.

From Lemma [4.13,](#page-34-0) we obtain the following proposition.

Proposition 4.22. We have the the following connection formulas:

$$
\begin{pmatrix} \Psi^+_{1,1}(z) \\ \Psi^+_{0,1}(z) \\ \Psi^+_{1,0}(z) \\ \Psi^+_{0,0}(z) \end{pmatrix} = \frac{(-1)^m}{2} \begin{pmatrix} c^{-1} & -c^{-1} & -c^{-1} & c^{-1} \\ -c + c^{-1} & -c^{-1} & c - c^{-1} & c^{-1} \\ -3c^{-1} & 3c^{-1} & -c^{-1} & c^{-1} \\ 3(c - c^{-1}) & 3c^{-1} & c - c^{-1} & c^{-1} \end{pmatrix} \begin{pmatrix} \Psi^-_{1,1}(z) \\ \Psi^-_{0,1}(z) \\ \Psi^-_{1,0}(z) \\ \Psi^-_{0,0}(z) \end{pmatrix},
$$

where $c = 2\cos(\pi \alpha - \beta_{2,2}).$

Remark 4.23. From Proposition [4.22,](#page-42-1) we obtain

$$
\begin{pmatrix} \Psi_{1,1}^{+}(z) + \Psi_{1,0}^{+}(z) \\ \Psi_{0,1}^{+}(z) + \Psi_{0,0}^{+}(z) \end{pmatrix} = (-1)^{m} \begin{pmatrix} -c^{-1} & c^{-1} \\ c - c^{-1} & c^{-1} \end{pmatrix} \begin{pmatrix} \Psi_{1,1}^{-}(z) + \Psi_{1,0}^{-}(z) \\ \Psi_{0,1}^{-}(z) + \Psi_{0,0}^{-}(z) \end{pmatrix} . \tag{4.42}
$$

Then, from [\(4.42\)](#page-43-0), it seems that the monodoromy of the Fuchsian differential equation [\(4.5\)](#page-22-0) is reducible and $\{\Psi_{1,1}^{\pm}(z) + \Psi_{1,0}^{\pm}(z), \Psi_{0,1}^{\pm}(z) + \Psi_{0,0}^{\pm}(z)\}\$ gives a two dimensional subspace of the monodromy representation of [\(4.5\)](#page-22-0).

5 Tensor structure on $\mathcal{SW}(m)$ -mod

Since the super triplet W-algebra $\mathcal{SW}(m)$ is C_2 -cofinite, Proposition 2.1 in [\[CGNS\]](#page-66-2) (see also [\[CMOY,](#page-66-3) Theorem 2.25] and [\[Hu,](#page-68-0) Theorem 4.13]) show that $\mathcal{SW}(m)$ has braided tensor supercategory structure developed in the papers [\[CKM\]](#page-66-4) and [\[HLZ1\]](#page-68-1)-[\[HLZ8\]](#page-69-1). We denote by $(\mathcal{SW}(m)$ -mod, $\boxtimes)$ the tensor supercategory on $\mathcal{SW}(m)$ -mod, where the unit object is given by X_1 and the symbol ⊠ denotes the tensor product. It is known that the tensor product ⊠ of $(\mathcal{SW}(m)$ -mod, \boxtimes) is right exact [\[CGNS,](#page-66-2) Proposition 2.1]. In this section, we study the fusion structure of $\mathcal{SW}(m)$ and determine the structure of the projective covers of all simple $\mathcal{SW}(m)$ -modules. See [\[CRR,](#page-66-5) [CMOY\]](#page-66-3) for the detailed structure of fusion rules and the tensor category for the $N = 1$ super Virasoro minimal models.

5.1 Tensor product \boxtimes and $P(w)$ -intertwining operators

In this subsection, we review the definition of the tensor product \boxtimes and $P(w)$ intertwining operators in accordance with [\[CKM,](#page-66-4) [HLZ3,](#page-68-2) [KR\]](#page-69-2) and derive some identities known as the Nahm-Gaberdiel-Kausch fusion algorithm(cf. [\[CRR,](#page-66-5) [GK1,](#page-67-1) [Nah\]](#page-70-1)).

Definition 5.1. Let V be a $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded vertex operator superalgebra and let C be a category of grading-restricted generalized V-modules. A tensor product (or fusion product) of M_1 and M_2 in C is a pair $(M_1 \boxtimes M_2, \mathcal{Y}_{\boxtimes})$, with $M_1 \boxtimes M_2$ and \mathcal{Y}_{\boxtimes} an intertwining operator of type $M_1 \boxtimes M_2 \choose M_1 \ M_2$ M_1 M_2 $\big)$, which satisfies the following universal property: For any $M_3 \in \mathcal{C}$ and intertwining operator $\mathcal Y$ of $\delta type\left(\frac{\tilde{M}_{3}}{M_{1}}\right)$ M_1 M_2 $\Big),$ there is a unique V-module homomorphism $f: M_1 \boxtimes M_2 \rightarrow M_3$ such that $\mathcal{Y} = f \circ \mathcal{Y}_{\boxtimes}$.

In the paper [\[HLZ3\]](#page-68-2), the notion of $P(w)$ -intertwining operators and the $P(w)$ -tensor product are introduced. The definitions are as follows.

Definition 5.2. Fix $w \in \mathbb{C}^{\times}$. Let V be a $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded vertex operator superalgebra and let C be a category of grading-restricted generalized V-modules. Given M_1 , M_2 and M_3 in C, a parity-homogeneous $P(w)$ -intertwining operator I of type $\binom{M_3}{M_1\ldots M_k}$ M_1 M_2 $\Big)$ is a parity-homogeneous bilinear map $I : M_1{\otimes} M_2 \rightarrow \overline{M}_3$ that satisfies the following properties:

- 1. For any $\psi_1 \in M_1$ and $\psi_2 \in M_2$, $\pi_h(I[\psi_1 \otimes \psi_2]) = 0$ for all $h \ll 0$, where π_h denotes the projection onto the generalised eigenspace $M_3[h]$ of L_0 -eigenvalue h.
- 2. For any $\psi_1 \in M_1$, $\psi_2 \in M_2$, $\psi_3^* \in M_3^*$ and $v \in V$, the three point functions

$$
\langle \psi_3^*, Y_3(v, z)I[\psi_1 \otimes \psi_2] \rangle, \qquad \langle \psi_3^*, I[Y_1(v, z - w)\psi_1 \otimes \psi_2] \rangle, \langle \psi_3^*, I[\psi_1 \otimes Y_2(v, z)\psi_2] \rangle
$$

are absolutely convergent in the regions $|z| > |w| > 0$, $|w| > |z-w| > 0$, $|w| > |z| > 0$, respectively, where Y_i is the action of V-module on M_i .

3. Given any $f(t) \in R_{P(w)} := \mathbb{C}[t, t^{-1}, (t-w)^{-1}]$ and parity-homogeneous vectors $v \in V$, $\psi_1 \in M_1$, $\psi_2 \in M_2$, $\psi_3^* \in M_3^*$, we have the following identity

$$
(-1)^{|v||I|} \oint_{0,w} f(z) \langle \psi_3^*, Y_3(v, z) I [\psi_1 \otimes \psi_2] \rangle \frac{dz}{2\pi i}
$$

=
$$
\oint_w f(z) \langle \psi_3^*, I[Y_1(v, z - w)\psi_1 \otimes \psi_2] \rangle \frac{dz}{2\pi i}
$$

+
$$
(-1)^{|v||\psi_1|} \oint_0 f(z) \langle \psi_3^*, I[\psi_1 \otimes Y_2(v, z)\psi_2] \rangle \frac{dz}{2\pi i}.
$$
 (5.1)

A general $P(w)$ -intertwining operator of type $\begin{pmatrix} M_3 \\ M_1 \end{pmatrix}$ M_1 M_2 is a sum of parityhomogeneous ones.

Definition 5.3. Let V be a $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded vertex operator superalgebra and let C be a category of grading-restricted generalized V-modules. A $P(w)$ -tensor product of M_1 and M_2 in C is a pair $(M_1 \boxtimes_{P(w)} M_2, \boxtimes_{P(w)})$, with $M_1 \boxtimes_{P(w)} M_2$ and $\boxtimes_{P(w)}$ a $P(w)$ -intertwining operator of type $\binom{M_1 \boxtimes_{P(w)} M_2}{M_1 \cdots M_2}$ M_1 M_2 , which satisfies the following universal property: For any $M_3 \in \mathcal{C}$ and $P(w)$ -intertwining operator I of type $\begin{pmatrix} M_3 \\ M_1 \end{pmatrix}$ M_1 M_2), there is a unique V-module homomorphism η : $M_1 \boxtimes_{P(w)} M_2 \rightarrow M_3$ such that

$$
\overline{\eta} \circ \boxtimes_{P(w)} [\psi_1 \otimes \psi_2] = I[\psi_1 \otimes \psi_2]
$$

for all $\psi_1 \in M_1$ and $\psi_2 \in M_2$, where $\overline{\eta}$ denotes the extension of η to a map between the completions of $M_1 \boxtimes_{P(w)} M_2$ and M_3 .

It is known that the definition $P(w)$ -tensor product $\boxtimes_{P(w)}$ does not depend on the choice of $w \in \mathbb{C}^{\times}$. Precisely, the following proposition holds [\[CKM,](#page-66-4) Corollary 3.36] (see also [\[HLZ3,](#page-68-2) Remark 4.22]).

Proposition 5.4 ([\[CKM\]](#page-66-4)). Let V be a $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded vertex operator superalgebra and let M_1, M_2 be V-modules. Suppose that for some $w_0 \in \mathbb{C}^\times$, $P(w_0)$ tensor product $M_1 \boxtimes_{P(w_0)} M_2$ exists. Then, for any $w, w' \in \mathbb{C}^\times$, $M_1 \boxtimes_{P(w)} M_2$ and $M_1 \boxtimes_{P(w')} M_2$ are isomorphic.

It is known that certain specializations for intertwining operators yield $P(w)$ -intertwining operators and by these specializations give linear isomorphisms from the the spaces of intertwining operators to the spaces of $P(w)$ intertwining operators of the same types [\[CKM,](#page-66-4) Proposition 3.15](see also [\[HLZ3\]](#page-68-2)). Thus, from Proposition [5.4,](#page-45-0) $\mathcal{SW}(m)$ has a braided tensor supercategory structure with $\mathbb{Z} = \mathbb{Z}_{P(1)}$.

In the following, we will introduce some useful formulas derived from the $P(w)$ -compatibility conditions. We define a translation map

$$
T_1: \mathbb{C}(t) \to \mathbb{C}(t)
$$
, by $f(t) \mapsto f(t+1)$,

and a expansion map

$$
\iota_+ : \mathbb{C}(t) \hookrightarrow \mathbb{C}((t))
$$

that expands a given rational function in t as a power series around $t = 0$. Given a $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded vertex operator superalgebra V and a homogeneous vector $v \in V[h]$, we use notation $v_{n-h+1} = vt^n = v \otimes t^n \in V \otimes \mathbb{C}[t, t^{-1}]$ for $n \in \mathbb{Z}$. Given V-modules M_1 , M_2 , M_3 and a parity-homogeneous $P(1)$ -intertwining operator I of type $\begin{pmatrix} M_3 \\ M_1 & M_2 \end{pmatrix}$ M_1 M_2), as detailed in $[KR]$, by using the action (2.2) and the identities [\(5.1\)](#page-44-0), we can define the action of $V \otimes \mathbb{C}[t, t^{-1}, (t-1)^{-1}]$ or $V \otimes \mathbb{C}((t))$ on M_3^* as

$$
(-1)^{|v||I|}\langle vf(t)\psi_3^*, I[\psi_1 \otimes \psi_1] \rangle = (-1)^{|v||I|}\langle vt_+(f(t))\psi_3^*, I[\psi_1 \otimes \psi_1] \rangle
$$

$$
= \langle \psi_3^*, I[\iota_+ \circ T_1(v^{\text{opp}}f(t^{-1}))\psi_1 \otimes \psi_2] \rangle
$$

$$
+ (-1)^{|v||\psi_1|}\langle \psi_3^*, I[\psi_1 \otimes \iota_+(v^{\text{opp}}f(t^{-1}))\psi_2] \rangle
$$

where $\psi_3^* \in M_3^*, \psi_i \in M_i$, and

$$
v^{\text{opp}} := e^{t^{-1}L_1}(-t^2)^{L_0}vt^{-2}.
$$

Lemma 5.5. Let V be a $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded vertex operator superalgebra and let $v \in V$ be a non-zero Virasoro primary vector with the conformal weight h:

$$
L_0 v = hv, \qquad L_n v = 0 \quad (n \ge 1).
$$

Given V-modules M_1 , M_2 , M_3 and a parity-homogeneous $P(1)$ -intertwining *operator I* of type $\begin{pmatrix} M_3 \\ M_1 & M_2 \end{pmatrix}$ M_1 M_2 $\big)$, we have the following identities:

$$
\begin{split}\n& (-1)^{|v||I|} \langle v_n \psi_3^*, I[\psi_1 \otimes \psi_2] \rangle \\
&= \sum_{i=0}^{\infty} {h - n - 1 \choose i} \langle \psi_3^*, I[(v_{i-h+1}\psi_1) \otimes \psi_2] \rangle + (-1)^{|v||\psi_1|} \langle \psi_3^*, I[\psi_1 \otimes (v_{-n}\psi_2)] \rangle, \\
& (-1)^{|v||I|} \sum_{i=0}^{\infty} {n + h - 1 \choose i} (-1)^i \langle v_{i-n} \psi_3^*, I[\psi_1 \otimes \psi_2] \rangle \\
&= \langle \psi_3^*, I[(v_n\psi_1) \otimes \psi_2] \rangle + (-1)^{|v||\psi_1|} \sum_{i=0}^{\infty} {n + h - 1 \choose i} (-1)^i \langle \psi_3^*, I[\psi_1 \otimes (v_{n-i}\psi_2)] \rangle, \\
& (-1)^{|v||I|} \sum_{i=0}^{\infty} {n + h - 2 \choose i} (-1)^i \langle v_{i-n} \psi_3^*, I[\psi_1 \otimes \psi_2] \rangle \\
&= \langle \psi_3^*, I[((v_{n-1} + v_n)\psi_1) \otimes \psi_2]) \\
&+ (-1)^{|v||\psi_1|} \sum_{i=0}^{\infty} {n + h - 2 \choose i} (-1)^{n-i+h-2} \langle \psi_3^*, I[\psi_1 \otimes (v_{i-h+2}\psi_2)] \rangle, \n\end{split}
$$

where $\psi_3^* \in M_3^*$, $\psi_i \in M_i$ and $n \in \mathbb{Z}$.

 $i=0$

Proof. For the first identity, let $f(t) = t^{n+h-1}$ in [\(5.2\)](#page-46-0), for the second, let $f(t) = t^{2h-2}(t^{-1} - 1)^{n+h-1}$ in [\(5.2\)](#page-46-0), and for the third identity, let $f(t) =$ $t^{2h-3}(t^{-1}-1)^{n+h-2}$ in [\(5.2\)](#page-46-0).

From the next subsection, we use the shorthand notation

i

$$
\langle \psi_3^*, \psi_1 \otimes \psi_2 \rangle = \langle \psi_3^*, I[\psi_1 \otimes \psi_2] \rangle \tag{5.2}
$$

for $P(1)$ -intertwining operators I.

5.2 Self duality of the simple module X_2

Note that all simple V_L -modules can be written as forms

$$
\mathcal{V}_{L+\beta} := \bigoplus_{n \in \mathbb{Z}} F_{\beta + n\alpha_+}, \quad \beta \in \{ \beta_{r,s,n} \mid r, s, n \in \mathbb{Z} \}
$$

(for the definition of the simple V_L -modules, see [\(3.1\)](#page-16-0)). Given simple V_L modules $\mathcal{V}_{L+\beta}$, $\mathcal{V}_{L+\beta'}$ and $\mathcal{V}_{L+\beta''}$, it can be proved easily that there are no \mathcal{V}_L -intertwining operators of type $\begin{pmatrix} \mathcal{V}_{L+\beta'} \\ \mathcal{V}_{L+\beta'} \end{pmatrix}$ $V_{L+\beta'} V_{L+\beta}$) unless $\beta'' \equiv \beta' + \beta \mod L$, and $\dim_{\mathbb{C}} I_{\mathcal{V}_L}^0$ \int , $V_{L+\beta'+\beta}$ $V_{L+\beta'} V_{L+\beta}$ = 1. Let $\mathcal Y$ be the even $\mathcal V_L$ -intertwining operator of type $\left(\begin{array}{cc} \mathcal{V}_{L+\beta'+\beta'} & \mathcal{V}_{L+\beta'+\beta'} \end{array}\right)$ $V_{L+\beta'} V_{L+\beta}$). Then, by restricting the action of \mathcal{V}_L to $\mathcal{SW}(m)$, \mathcal{Y} defines an $\mathcal{SW}(m)$ -intertwining operator of type $\begin{pmatrix} \mathcal{V}_{L+\beta'+\beta'} \\ \mathcal{V}_{L+\beta'} & \mathcal{V}_{L+\beta'} \end{pmatrix}$ $V_{L+\beta'} V_{L+\beta}$. We denote by $\mathcal{Y}_{\beta',\beta}$ this even $\mathcal{SW}(m)$ -intertwining operator.

Lemma 5.6. For $2 \leq s \leq 2m$, we have

$$
I_{\mathcal{SW}(m)}\begin{pmatrix} X_{s-1} \\ X_2 \ X_s \end{pmatrix} \neq 0, \qquad I_{\mathcal{SW}(m)}\begin{pmatrix} X_{s+1} \\ X_2 \ X_s \end{pmatrix} \neq 0.
$$

Proof. We only prove

$$
I_{\mathcal{SW}(m)}\left(\frac{X_{2i-1}}{X_2\ X_{2i}}\right)\neq 0.
$$

The other cases can be shown in similar ways.

Let us consider the $\mathcal{SW}(m)$ -intertwining operator $\mathcal{Y} = \mathcal{Y}_{\beta_1,\beta_2}$, where $\beta_1 =$ $\beta_{2,2}$ and $\beta_2 = \beta_{1,2(m-i)+1,2}$. By [\(2.6\)](#page-10-1), we have

$$
\langle \beta_{1,2(m-i+1);1} | \mathcal{Y}(|\beta_{2,2}\rangle, z) | \beta_{1,2(m-i)+1;2} \rangle \neq 0. \tag{5.3}
$$

From Proposition [3.3,](#page-16-1) we have

$$
X_2 \simeq \mathcal{SW}(m)|\beta_{2,2}\rangle, \qquad \qquad \mathcal{V}_{L+\beta_2} \simeq \mathcal{SW}(m)|\beta_{1,2(m-i)+1;2}\rangle, \n\mathcal{V}_{L+\beta_2+\beta_1} \simeq \mathcal{SW}(m)|\beta_{1,2(m-i+1);1}\rangle,
$$

and the exact sequence

$$
0 \to X_{2(m-i+1)} \to \mathcal{V}_{L+\beta_2+\beta_1} \to X_{2i-1} \to 0.
$$

Thus, by (5.3) , we have

$$
I_{\mathcal{SW}(m)} \begin{pmatrix} X_{2i-1} \\ X_2 \ \mathcal{V}_{L+\beta_2} \end{pmatrix} \neq 0. \tag{5.4}
$$

From Proposition [3.3,](#page-16-1) we have the exact sequence

$$
0 \to X_{2(m-i)+1} \to \mathcal{V}_{L+\beta_2} \to X_{2i} \to 0.
$$

Then we have the following exact sequence

$$
X_2 \boxtimes X_{2(m-i)+1} \to X_2 \boxtimes \mathcal{V}_{L+\beta_2} \to X_2 \boxtimes X_{2i} \to 0.
$$

From this exact sequence, we have the following exact sequence

$$
0 \to \text{Hom}_{\mathcal{SW}(m)}(X_2 \boxtimes X_{2i}, X_{2i-1}) \to \text{Hom}_{\mathcal{SW}(m)}(X_2 \boxtimes \mathcal{V}_{L+\beta_2}, X_{2i-1})
$$

$$
\to \text{Hom}_{\mathcal{SW}(m)}(X_2 \boxtimes X_{2(m-i)+1}, X_{2i-1}). \tag{5.5}
$$

By Lemma [5.8,](#page-48-0) we have $\text{Hom}_{\mathcal{SW}(m)}(X_2 \boxtimes X_{2(m-i)+1}, X_{2i-1}) = 0$. Therefore by (5.4) and (5.5) , we obatin $\text{Hom}_{\mathcal{SW}(m)}(X_2 \boxtimes X_{2i}, X_{2i-1}) \neq 0$. \Box

From this subsection, we will use the following notation.

Definition 5.7. For any $M \in \mathcal{SW}(m)$ -mod, we define the following vector space

$$
A_0(M) = \left\{ \psi \in M \setminus \{0\} \mid v_n \psi = 0 \text{ for } v \in \mathcal{SW}(m) \text{ and } n \in \frac{1}{2} \mathbb{Z}_{>0} \right\}.
$$

Lemma 5.8. For $i = 1, ..., m$, the vector space $A_0((X_2 \boxtimes X_{2i})^*)$ is at most two dimensional. L_0 acts semisimply on $A_0((X_2 \boxtimes X_{2i})^*)$ and any L_0 eigenvalue of this space is contained in $\{h_{1,2i-1}, h_{1,2i+1}\}$, where $h_{1,2i-1}$ and $h_{1,2i+1}$ are the minimal conformal weights of X_{2i-1} and X_{2i+1} , respectively.

Similar results are obtained for the $N = 1$ super Virasoro algebra by using Zhu bimodules [\[CMOY,](#page-66-3) Subsection 4.1].

Proof. By Lemma [5.6,](#page-47-2) we see that the tensor product $X_2 \boxtimes X_{2i}$ is non-zero. Let ψ^* be an arbitrary non zero L_0 homogeneous vector of $A_0((X_2 \boxtimes X_{2i})^*)$. Let ϕ_1 and ϕ_2 be arbitrary non zero L_0 homogeneous vectors X_2 and X_{2i} such that $\langle \psi^*, \phi_1 \otimes \phi_2 \rangle \neq 0$, where we use the shorthand notation [\(5.2\)](#page-46-0) as $I = \boxtimes_{P(1)}$. For $1 \leq j \leq m$, let $\{v_j^+\}$ j^+ , v_j^- } be a basis of the minimal conformal weight space of X_{2j} such that

$$
\widehat{W}_0^{\pm} v_j^{\pm} = 0, \qquad \qquad \widehat{W}_0^{\pm} v_j^{\mp} \in \mathbb{C}^{\times} v_j^{\pm},
$$

where \widehat{W}^{\pm}_0 are the zero-mode of the fields $Y(\widehat{W}^{\pm}, z)$. For $n \geq 1$, let

$$
v_{\frac{2k-1}{2}}^{(n)} \t (k=-n,\ldots,n+1)
$$

be the minimal conformal weight vectors of the subspace $(2n+2)L(h_{2n+2,2i}) \subset$ X_{2i} defined in Proposition [3.8.](#page-17-0) First let us show

$$
\langle \psi^*, U(\mathfrak{ns}) v_1^{\pm} \otimes v_{\frac{2k-1}{2}}^{(n)} \rangle = 0 \quad (n \ge 1, \ k \in \mathbb{Z}). \tag{5.6}
$$

Note that v_1^{\pm} satisfy the following relations (cf.[\[BS,](#page-65-0) [CMOY\]](#page-66-3))

$$
\left\{\frac{4t}{t^2-1}G_{-\frac{1}{2}}^4 + \frac{t+1}{t-1}G_{-\frac{1}{2}}G_{-\frac{3}{2}} + \frac{t-1}{t+1}G_{-\frac{3}{2}}G_{-\frac{1}{2}}\right\}v_1^{\pm} = 0 \quad (t = -2m - 1). \tag{5.7}
$$

By using Lemma [5.5](#page-46-1) and the relation [\(5.7\)](#page-49-0), we see that, depending on whether ψ^* is even or odd, the values $\langle \psi^*, U(\mathfrak{ns})v_1^{\pm} \otimes \otimes v_{\frac{2k-1}{2}}^{(n)} \rangle$ is determined by the numbers

$$
\langle \psi^*, v_1^{\delta} \otimes v_{\frac{2l-1}{2}}^{(2n)} \rangle, \qquad \langle \psi^*, L_{-1} v_1^{\delta} \otimes v_{\frac{2l-1}{2}}^{(2n)} \rangle
$$

or

$$
\langle \psi^*, G_{-\frac{1}{2}} v_1^{\delta} \otimes v_{\frac{2l-1}{2}}^{(2n+1)} \rangle, \qquad \langle \psi^*, G_{-\frac{1}{2}}^3 v_1^{\delta} \otimes v_{\frac{2l-1}{2}}^{(2n+1)} \rangle,
$$

for $\delta = \pm$ and some finite n and l (cf. [\[KR,](#page-69-2) Section 7]). By using Lemma [5.5](#page-46-1) and (5.7) , we have

$$
\begin{aligned}\n&\left(\langle L_0 \psi^*, v_1^{\delta} \otimes v_{\frac{2k-1}{2}}^{(n)} \rangle \right) \\
&\left(\langle L_0 \psi^*, (L_{-1} v_1^{\delta}) \otimes v_{\frac{2k-1}{2}}^{(n)} \rangle \right) \\
&= \left(\begin{matrix} h_{2,2} + h_{2n+2,2i} & 1 \\ \frac{2m^2}{2m+1} h_{2n+2,2i} & h_{2,2} + h_{2n+2,2i} + 1 - \frac{(2m+1)^2+1}{2(2m+1)} \end{matrix} \right) \left(\begin{matrix} \langle \psi^*, v_1^{\delta} \otimes v_{\frac{2k-1}{2}}^{(n)} \rangle \\ \langle \psi^*, (L_{-1} v_1^{\delta}) \otimes v_{\frac{2k-1}{2}}^{(n)} \rangle \end{matrix} \right).\n\end{aligned}
$$

We see that the eigenvalues of this matrix do not correspond to the minimal conformal weights of all simple $\mathcal{SW}(m)$ -modules. Thus we have

$$
\langle \psi^*, v_1^{\delta} \otimes v_{\frac{2k-1}{2}}^{(n)} \rangle = 0, \qquad \langle \psi^*, (L_{-1}v_1^{\delta}) \otimes v_{\frac{2k-1}{2}}^{(n)} \rangle = 0
$$

for any $n \in \mathbb{Z}_{\geq 1}$ and $k \in \mathbb{Z}$. Similary we can show that

$$
\langle \psi^*, G_{-\frac{1}{2}} v_1^{\delta} \otimes v_{\frac{2k-1}{2}}^{(n)} \rangle = 0, \qquad \langle \psi^*, G_{-\frac{1}{2}}^3 v_1^{\delta} \otimes v_{\frac{2k-1}{2}}^{(n)} \rangle = 0
$$

for any $n \in \mathbb{Z}_{\geq 1}$ and $k \in \mathbb{Z}$. Therefore we obtain [\(5.6\)](#page-48-2).

By Proposition [3.9,](#page-18-0) Lemma [5.5,](#page-46-1) [\(5.7\)](#page-49-0) and [\(5.6\)](#page-48-2), we see that, depending on whether ψ^* is even or odd, $\langle \psi^*, \phi_1 \otimes \phi_2 \rangle$ is determined by the numbers

$$
\langle \psi^*, G_{-\frac{1}{2}}^l v_1^{\delta} \otimes v_i^{\delta'} \rangle \quad (l = 0, 2, \ \delta = \pm, \ \delta' = \pm)
$$
 (5.8)

or

$$
\langle \psi^*, G^l_{-\frac{1}{2}} v_1^{\delta} \otimes v_i^{\delta'} \rangle
$$
 $(l = 1, 3, \delta = \pm, \delta' = \pm).$ (5.9)

Let us assume that the parity of ψ^* is odd. Then $\langle \psi^*, \phi_1 \otimes \phi_2 \rangle$ is determined by the numbers [\(5.9\)](#page-49-1). By using Lemma [5.5](#page-46-1) and the relation [\(5.7\)](#page-49-0), we have

$$
\begin{pmatrix}\n\langle L_0 \psi^*, G_{-\frac{1}{2}} v_1^\delta \otimes v_i^{\delta'} \rangle \\
\langle L_0 \psi^*, G_{-\frac{1}{2}}^3 v_1^\delta \otimes v_i^{\delta'} \rangle\n\end{pmatrix} = M_1 \begin{pmatrix}\n\langle \psi^*, G_{-\frac{1}{2}} v_1^\delta \otimes v_i^{\delta'} \rangle \\
\langle \psi^*, G_{-\frac{1}{2}}^3 v_1^\delta \otimes v_i^{\delta'} \rangle\n\end{pmatrix},
$$

where

$$
M_1 = \begin{pmatrix} h_{2,2} + h_{2,2i} + \frac{1}{2} & 1\\ \frac{m^2}{2m+1} + \frac{2m^2}{2m+1}h_{2,2i} & h_{2,2} + h_{2,2i} + \frac{3}{2} - \frac{(2m+1)^2 + 1}{2(2m+1)} \end{pmatrix}
$$

We see that this matrix M_1 is diagonalizable and the eigenvalues are given by $h_{1,2i+1}$ and $h_{3,2i-1}$. Note that the eigenvalue $h_{3,2i-1}$ does not correspond to any minimal conformal weight of the simple $\mathcal{SW}(m)$ -modules. Thus L_0 acts semisimply on ψ^* and the L₀-weight of ψ^* is $h_{1,2i+1}$ which is the minimal conformal weight of X_{2i+1} .

Next let us assume that the parity of ψ^* is even. Then $\langle \psi^*, \phi_1 \otimes \phi_2 \rangle$ is determined by the numbers [\(5.8\)](#page-49-2). By using Lemma [5.5](#page-46-1) and [\(5.7\)](#page-49-0), we have

$$
\begin{pmatrix}\n\langle L_0 \psi^*, v_1^{\delta} \otimes v_i^{\delta'} \rangle \\
\langle L_0 \psi^*, (L_{-1} v_1^{\delta}) \otimes v_i^{\delta'} \rangle\n\end{pmatrix} = M_2 \begin{pmatrix}\n\langle \psi^*, v_1^{\delta} \otimes v_i^{\delta'} \rangle \\
\langle \psi^*, (L_{-1} v_1^{\delta}) \otimes v_i^{\delta'} \rangle\n\end{pmatrix},
$$

where

$$
M_2 = \begin{pmatrix} h_{2,2} + h_{2,2i} & 1\\ \frac{2m^2}{2m+1}h_{2,2i} & h_{2,2} + h_{2,2i} + 1 - \frac{(2m+1)^2 + 1}{2(2m+1)} \end{pmatrix}.
$$

We see that this matrix M_2 is diagonalizable and eigenvalues are given by $h_{1,2i-1}$ and $h_{3,2i+1}$. Note that the eigenvalue $h_{3,2i+1}$ does not correspond to any minimal conformal weight of the simple $\mathcal{SW}(m)$ -modules. Thus L_0 acts semisimply on ψ^* and the L₀-weight of ψ^* is $h_{1,2i-1}$ which is the minimal conformal weight of X_{2i-1} .

Hence the L₀-weight of ψ^* is given by $h_{1,2i-1}$ or $h_{1,2i+1}$. Note that from Proposition [3.9,](#page-18-0) \hat{W}^{\pm}_{0} act trivially on the minimal conformal weight spaces of X_{2i-1} and X_{2i+1} . Then we have $\widehat{W}^{\pm}_{0} \psi^* = 0$. Thus, by Lemma [5.5,](#page-46-1) we see that $\langle \psi^*, \phi_1 \otimes \phi_2 \rangle$ is determined by the numbers

$$
\langle \psi^*, v_1^+ \otimes v_i^- \rangle, \qquad \langle \psi^*, L_{-1} v_1^+ \otimes v_i^- \rangle
$$

in the case of $L_0 \psi^* = h_{1,2i-1} \psi^*$, and

$$
\langle \psi^*, G_{-\frac{1}{2}}v^+_1 \otimes v^-_i \rangle, \qquad \qquad \langle \psi^*, G^3_{-\frac{1}{2}}v^+_1 \otimes v^-_i \rangle
$$

in the case of $L_0 \psi^* = h_{1,2i+1} \psi^*$. Let $\binom{\mu_1}{\mu_2}$ $\binom{\mu_1}{\mu_2}$ be an eigenvector of tM_1 with the eigenvalue $h_{3,2i-1}$ and $\binom{\nu_1}{\nu_2}$ $\begin{bmatrix} \nu_1 \\ \nu_2 \end{bmatrix}$ be an eigenvector of tM_2 with the eigenvalue $h_{3,2i+1}$. Assume $L_0 \psi^* = h_{1,2i+1} \psi^*$. Then we have

$$
h_{1,2i+1}(\mu_1, \mu_2) \begin{pmatrix} \langle \psi^*, G_{-\frac{1}{2}} v_1^+ \otimes v_i^- \rangle \\ \langle \psi^*, G_{-\frac{1}{2}}^3 v_1^+ \otimes v_i^- \rangle \end{pmatrix} = (\mu_1, \mu_2) \begin{pmatrix} \langle L_0 \psi^*, G_{-\frac{1}{2}} v_1^+ \otimes v_i^- \rangle \\ \langle L_0 \psi^*, G_{-\frac{1}{2}}^3 v_1^+ \otimes v_i^- \rangle \end{pmatrix}
$$

= $(\mu_1, \mu_2) M_1 \begin{pmatrix} \langle \psi^*, G_{-\frac{1}{2}} v_1^+ \otimes v_i^- \rangle \\ \langle \psi^*, G_{-\frac{1}{2}}^3 v_1^+ \otimes v_i^- \rangle \end{pmatrix}$
= $h_{3,2i-1}(\mu_1, \mu_2) \begin{pmatrix} \langle \psi^*, G_{-\frac{1}{2}} v_1^+ \otimes v_i^- \rangle \\ \langle \psi^*, G_{-\frac{1}{2}}^3 v_1^+ \otimes v_i^- \rangle \end{pmatrix}.$

Thus we obtain

$$
\mu_1 \langle \psi^*, G_{-\frac{1}{2}} v_1^+ \otimes v_i^- \rangle + \mu_2 \langle \psi^*, G_{-\frac{1}{2}}^3 v_1^+ \otimes v_i^- \rangle = 0.
$$

Similarly, assuming $L_0 \psi^* = h_{1,2i-1} \psi^*$, we obtain

$$
\nu_1 \langle \psi^*, v_1^+ \otimes v_i^- \rangle + \nu_2 \langle \psi^*, L_{-1} v_1^+ \otimes v_i^- \rangle = 0.
$$

Therefore the vector space $A_0((X_2 \boxtimes X_{2i})^*)$ is at most two dimensional. \Box

Lemma 5.9. We have

$$
X_2 \boxtimes X_2 = X_1 \oplus \Gamma(X_3), \tag{5.10}
$$

where $\Gamma(X_3)$ is a lowest weight module whose top composition factor is X_3 .

Proof. From Lemmas [5.6](#page-47-2) and [5.8,](#page-48-0) we have

$$
X_2 \boxtimes X_2 = \Gamma(X_1) \oplus \Gamma(X_3),
$$

where $\Gamma(X_{2i+1})$ $(i = 0, 1)$ are lowest weight modules whose top composition factors are given by X_{2i+1} . Let us show $\Gamma(X_1) = X_1$. Assume $\Gamma(X_1) \ncong X_1$. Then by Proposition [3.13,](#page-19-0) $\Gamma(X_1)$ must have a composition factor X_{2m} . In particular, we have $\psi^* \in A_0((X_2 \boxtimes X_2)^*)$ such that

$$
L_0 \psi^* = h_{1,1} \psi^* = 0, \qquad G_{-\frac{1}{2}} \psi^* \neq 0.
$$

Note that $G_{-\frac{1}{2}}\psi^* \in A_0((X_2 \boxtimes X_2)^*)$. Then, from the proof of Lemma [5.8,](#page-48-0) we see that the L₀-weight of $G_{-\frac{1}{2}}\psi^*$ must be $h_{1,3}$. But, since $L_0G_{-\frac{1}{2}}\psi^* =$ $h_{2,2m}G_{-\frac{1}{2}}\psi^*$, we have a contradiction. \Box

Recall that in Subsection [4.4,](#page-37-1) we construct the solutions $\{\Psi_{i,j}^{+}(z) \mid i,j =$ 1, 0} and $\{\Psi_{i,j}(z) \mid i,j = 1, 0\}$ of the Fuchsian differential equation [\(4.5\)](#page-22-0) (see [\(4.38\)](#page-40-2) and Proposition [4.20\)](#page-42-2). By the connection formulas in Proposition [4.22,](#page-42-1) we can show the following theorem (cf. [\[CMOY,](#page-66-3) Theorem 4.7]).

Theorem 5.10. X_2 is rigid and self-dual.

Proof. We show the rigidity of X_2 following the methods in [\[CMY1,](#page-66-6) [CMY2,](#page-66-7) [MY,](#page-70-2) [TW1\]](#page-71-1). By Lemma [5.9,](#page-51-0) we have parity-homogeneous homomorphisms

$$
i_1: X_1 \to X_2 \boxtimes X_2,
$$

\n
$$
i_3: \Gamma(X_3) \to X_2 \boxtimes X_2,
$$

\n
$$
p_1: X_2 \boxtimes X_2 \to X_1,
$$

\n
$$
p_3: X_2 \boxtimes X_2 \to \Gamma(X_3)
$$

such that

$$
p_1 \circ i_1 = id_{X_1}, \qquad p_3 \circ i_3 = id_{\Gamma(X_3)}, \n i_1 \circ p_1 + i_3 \circ p_3 = id_{X_2 \boxtimes X_2},
$$
\n(5.11)

where $\Gamma(X_3)$ is the lowest weight module defined by [\(5.10\)](#page-51-1). The maps i_1 and p_1 are candidates of the coevaluation and evaluation, respectively.

We define two homomorphisms $f, g: X_2 \to X_2$ as the compositions

$$
f: X_2 \xrightarrow{r^{-1}} X_2 \boxtimes X_1 \xrightarrow{\mathrm{id} \boxtimes i_1} X_2 \boxtimes (X_2 \boxtimes X_2) \xrightarrow{A} (X_2 \boxtimes X_2) \boxtimes X_2 \xrightarrow{p_1 \boxtimes \mathrm{id}} X_1 \boxtimes X_2 \xrightarrow{l} X_2,
$$

$$
g: X_2 \xrightarrow{l^{-1}} X_1 \boxtimes X_2 \xrightarrow{i_1 \boxtimes \mathrm{id}} (X_2 \boxtimes X_2) \boxtimes X_2 \xrightarrow{A^{-1}} X_2 \boxtimes (X_2 \boxtimes X_2) \xrightarrow{\mathrm{id} \boxtimes p_1} X_2 \boxtimes X_1 \xrightarrow{r} X_2,
$$

where A is the associativity isomorphism, and l, r are the left and right unit isomorphisms. The left and right unit isomorphisms l, r are characterized by

$$
\begin{aligned} \bar{l}_{X_2}(u_{X_1}\boxtimes u_{X_2})&=Y_{X_2}(u_{X_1},1)u_{X_2},\\ \bar{r}_{X_2}(u_{X_2}\boxtimes u_{X_1})&=(-1)^{|u_{X_1}||u_{X_2}|}e^{L_{-1}}Y_{X_2}(u_{X_1},-1)u_{X_2}, \end{aligned}
$$

for parity-homogeneous $u_{X_1} \in X_1$ and $u_{X_2} \in X_2$. By Lemma 4.2.1 and Corollary 4.2.2 of [\[CMY2\]](#page-66-7), it is enough to show that one of f and g is nonzero in order to show that X_2 is rigid and self-dual. Let us show $f \neq 0$.

Let $\mathcal{Y}_{2\boxtimes 2}$ and $\mathcal{Y}_{2\boxtimes (2\boxtimes 2)}$ be non-zero even intertwining operators of type

$$
\left(\begin{array}{c} X_2 \boxtimes X_2 \\ X_2 \end{array}\right), \qquad \left(\begin{array}{c} X_2 \boxtimes (X_2 \boxtimes X_2) \\ X_2 \end{array}\right),
$$

respectively. We introduce even intertwining operators

$$
\mathcal{Y}_{21}^2 = l_{X_2} \circ (p_1 \boxtimes id_{X_2}) \circ \mathcal{A}_{X_2, X_2, X_2} \circ \mathcal{Y}_{2\boxtimes (2\boxtimes 2)} \circ (id_{X_2} \otimes i_1), \n\mathcal{Y}_{23}^2 = l_{X_2} \circ (p_1 \boxtimes id_{X_2}) \circ \mathcal{A}_{X_2, X_2, X_2} \circ \mathcal{Y}_{2\boxtimes (2\boxtimes 2)} \circ (id_{X_2} \otimes i_3).
$$

The first intertwining operator corresponds to f . Fix a minimal conformal weight vector $v \in X_2[h_{2,2}]$ and let v^* be a minimal conformal weight vector of $X_2^*(\simeq X_2)$ such that $\langle v^*, v \rangle \neq 0$. For some $x \in \mathbb{R}$ such that $1 > x > 1-x > 0$, we set

$$
\phi_1(x) = \langle v^*, \mathcal{Y}_{21}^2(v, 1)(p_1 \circ \mathcal{Y}_{2\boxtimes 2})(v, x)v \rangle, \n\phi_3(x) = \langle v^*, \mathcal{Y}_{23}^2(v, 1)(p_3 \circ \mathcal{Y}_{2\boxtimes 2})(v, x)v \rangle.
$$

To prove $f \neq 0$, it is enough to show that $\phi_1(x) \neq 0$. Note that ϕ_1 and ϕ_3 satisfy the Fuchsian differential equation [\(4.5\)](#page-22-0), and admit series

$$
\phi_1(x) \in \mathbb{C}x^{h_{1,1}-2h_{2,2}}\big(1+x\mathbb{C}[[x]]\big), \quad \phi_3(x) \in \mathbb{C}x^{h_{1,3}+\frac{1}{2}-2h_{2,2}}\big(1+x\mathbb{C}[[x]]\big).
$$

Then, by noting the characteristic exponents of [\(4.5\)](#page-22-0), we have

$$
\phi_1(x) \in \mathbb{C}\Psi_{0,1}^+(x) + \mathbb{C}\Psi_{0,0}^+(x), \qquad \phi_3(x) \in \mathbb{C}\Psi_{1,1}^+(x) + \mathbb{C}\Psi_{1,0}^+(x). \tag{5.12}
$$

By (5.11), we have

$$
\phi_1(x) + \phi_3(x)
$$
\n
$$
= \langle v^*, \overline{l_{X_2} \circ (p_1 \boxtimes id_{X_2}) \circ A_{X_2, X_2, X_2}} (\mathcal{Y}_{2\boxtimes(2\boxtimes 2)}(v, 1) \mathcal{Y}_{2\boxtimes 2}(v, x)v) \rangle
$$
\n
$$
= \langle v^*, \overline{l_{X_2} \circ (p_1 \boxtimes id_{X_2})} (\mathcal{Y}_{(2\boxtimes 2)\boxtimes 2}(\mathcal{Y}_{2\boxtimes 2}(v, 1 - x)v, x)v) \rangle
$$
\n
$$
= \langle v^*, \overline{l_{X_2}} (\mathcal{Y}_{1\boxtimes 2}((p_1 \circ \mathcal{Y}_{2\boxtimes 2})(v, 1 - x)v, x)v) \rangle,
$$
\n
$$
= \langle v^*, Y_{X_2}((p_1 \circ \mathcal{Y}_{2\boxtimes 2})(v, 1 - x)v, x) \rangle,
$$
\n(5.13)

where $\mathcal{Y}_{1\boxtimes 2}$ is a non-zero even intertwining operator of type $\begin{pmatrix} X_2 \\ X_1 \end{pmatrix}$ X_1 X_2 . Note that $p_1 \circ \mathcal{Y}_{2\boxtimes 2}$ is a non-zero even intertwining operator of type $\begin{pmatrix} X_1 \\ X_2 \end{pmatrix}$ X_2 X_2 . Then we have

$$
\langle v^*, Y_{X_2}((p_1 \circ \mathcal{Y}_{2\boxtimes 2})(v, 1-x)v, x) \rangle v \rangle \in \mathbb{C}^\times \Psi_{0,0}^-(x) + \mathbb{C} \Psi_{0,1}^-(x). \tag{5.14}
$$

Assume that $\phi_1(x) = 0$. Then, from [\(5.13\)](#page-53-0) and [\(5.14\)](#page-53-1), we have $\phi_3(x) \neq 0$ and

$$
\phi_3(x) \in \mathbb{C}^\times \Psi^-_{0,0}(x) + \mathbb{C} \Psi^-_{0,1}(x).
$$

Then by [\(5.12\)](#page-53-2), we see that there exist $(k, l) \in \mathbb{C}^2 \setminus \{(0, 0)\}\$ such that

$$
k\Psi_{1,1}^+(x) + l\Psi_{1,0}^+(x) \in \mathbb{C}^\times \Psi_{0,0}^-(x) + \mathbb{C} \Psi_{0,1}^-(x). \tag{5.15}
$$

On the other hand, from Proposition [4.22,](#page-42-1) we have

$$
\Psi_{1,1}^{+}(x) = \frac{(-1)^{m}}{4\cos(\pi\alpha_{-}\beta_{2,2})} (\Psi_{1,1}^{-}(x) - \Psi_{0,1}^{-}(x) - \Psi_{1,0}^{-}(x) + \Psi_{0,0}^{-}(x)),
$$
\n
$$
\Psi_{1,0}^{+}(x) = \frac{(-1)^{m}}{4\cos(\pi\alpha_{-}\beta_{2,2})} (-3\Psi_{1,1}^{-}(x) + 3\Psi_{0,1}^{-}(x) - \Psi_{1,0}^{-}(x) + \Psi_{0,0}^{-}(x)).
$$
\n(5.16)

By [\(5.16\)](#page-53-3), we see that [\(5.15\)](#page-53-4) contradicts the linear independence of ${\Psi}_{i,j}^-(x)$. Therefore, we obtain $\phi_1(x) \neq 0$.

5.3 Non-semisimple fusion rules

Lemma 5.11. For $i = 1, ..., m-1$, the vector space $A_0((X_2 \boxtimes X_{2i+1})^*)$ is at most four dimensional. L₀ acts semisimply on $A_0((X_2 \boxtimes X_{2i+1})^*)$ and any L_0 eigenvalue of this space is contained in $\{h_{2,2i}, h_{2,2i+2}\}$, where $h_{2,2i}$ and $h_{2,2i+2}$ are the minimal conformal weights of X_{2i} and X_{2i+2} , respectively.

Proof. By Lemma [5.6,](#page-47-2) the tensor product $X_2 \boxtimes X_{2i+1}$ is non zero. Let ψ^* be an arbitrary non zero L_0 homogeneous vector of $A_0((X_2 \boxtimes X_{2i+1})^*)$. Let ϕ_1 and ϕ_2 be arbitrary non zero L_0 homogeneous vectors of X_2 and X_{2i+1} such that $\langle \psi^*, \phi_1 \otimes \phi_2 \rangle \neq 0$, where we use the shorthand notation [\(5.2\)](#page-46-0) as $I = \boxtimes_{P(1)}$. Let $\{v^+, v^-\}$ be a basis of the minimal conformal weight space of X_2 such that $\widehat{W}^{\pm}_0 v^{\pm} = 0$ and $\widehat{W}^{\pm}_0 v^{\mp} \in \mathbb{C}^{\times} v^{\pm}$. For $n \geq 1$, let $w_k^{(n)}$ $k_k^{(n)}(k=-n,\ldots,n)$ be the minimal conformal weight vectors of the subspace $(2n+1)L(h_{2n+1,2i+1}) \subset X_{2i+1}$ defined in Proposition [3.8.](#page-17-0) Similar to the arguments in Lemma [5.8,](#page-48-0) we have

$$
\langle \psi^*, U(\mathfrak{ns})v^{\pm} \otimes w_k^{(n)} \rangle = 0 \quad (n \ge 2, -n \le k \le n). \tag{5.17}
$$

Let $u (= w_0^{(0)}$ $\binom{0}{0}$ be the minimal conformal weight vector of X_{2i+1} . Then, by Lemma [5.5](#page-46-1) and by the relations [\(5.7\)](#page-49-0) and [\(5.17\)](#page-54-0), we see that $\langle \psi^*, \phi_1 \otimes \phi_2 \rangle$ is determined by the numbers

$$
\langle \psi^*, G^l_{-\frac{1}{2}} v^{\pm} \otimes u \rangle, \qquad \langle \psi^*, G^l_{-\frac{1}{2}} v^{\pm} \otimes w_k^{(1)} \rangle,
$$

for $l = 0, 1, 2, 3$ and $k = -1, 0, 1$. From Proposition [3.9,](#page-18-0) we have

$$
\begin{aligned}\n\widehat{W}_{-h}^{\pm}v^{\delta} &\in U(\mathfrak{ns})v^{\delta} + U(\mathfrak{ns})v^{-\delta} \\
W_{-h}^{\pm}v^{\delta} &\in U(\mathfrak{ns})v^{\delta} + U(\mathfrak{ns})v^{-\delta}\n\end{aligned}\n\quad h < h_{3,1} = \frac{1}{2} + 2m, \ \delta = \pm,\n\tag{5.18}
$$

and

$$
w_{\pm 1}^{(1)} \in \mathbb{C}^{\times} W^{\pm}[h_{1,2i+1} - h_{3,2i+1}]u + U(\mathfrak{ns})u,
$$

\n
$$
w_0^{(1)} \in \mathbb{C}^{\times} W^0[h_{1,2i+1} - h_{3,2i+1}]u + U(\mathfrak{ns})u.
$$
\n(5.19)

From the definition [\(3.4\)](#page-17-1), we have

$$
[G_{-\frac{1}{2}}, Y(W^a, z)] \in \mathbb{C}^{\times} Y(\widehat{W}^a, z), \quad [G_{-\frac{1}{2}}, Y(\widehat{W}^a, z)] \in \mathbb{C}^{\times} Y(L_{-1}W^a, z)
$$

for $a = \pm$, 0. Thus, by using [\(5.18\)](#page-54-1)-[\(5.19\)](#page-54-2) and the identities in Lemma [5.5,](#page-46-1) we see that each $\langle \psi^*, G^s_{-\frac{1}{2}}. v^{\pm} \otimes w_k^{(1)} \rangle$ $\langle k^{\perp} \rangle$ $(s = 0, 1, 2, 3)$ is determined by the numbers

$$
\langle \psi^*, G^l_{-\frac{1}{2}} v^{\pm} \otimes u \rangle
$$
 $(l = 0, 1, 2, 3).$

Therefore $\langle \psi^*, \phi_1 \otimes \phi_2 \rangle$ is determined by the numbers

$$
\langle \psi^*, G^l_{-\frac{1}{2}} v^\delta \otimes u \rangle \quad (l = 0, 2, \ \delta = \pm)
$$
 (5.20)

or

$$
\langle \psi^*, G^l_{-\frac{1}{2}} v^\delta \otimes u \rangle \quad (l = 1, 3, \ \delta = \pm). \tag{5.21}
$$

(a) Assume that $\langle \psi^*, \phi_1 \otimes \phi_2 \rangle$ is determined by the numbers [\(5.21\)](#page-55-0). Then, by using Lemma 5.5 and (5.7) , we have

$$
\begin{pmatrix} \langle L_0 \psi^*, G_{-\frac{1}{2}} v^\delta \otimes u \rangle \\ \langle L_0 \psi^*, G_{-\frac{1}{2}}^3 v_1^\delta \otimes u \rangle \end{pmatrix} = N_1 \begin{pmatrix} \langle \psi^*, G_{-\frac{1}{2}} v^\delta \otimes u \rangle \\ \langle \psi^*, G_{-\frac{1}{2}}^3 v^\delta \otimes u \rangle \end{pmatrix},
$$

where

$$
N_1 = \begin{pmatrix} h_{2,2} + h_{1,2i+1} + \frac{1}{2} & 1\\ \frac{m^2}{2m+1} + \frac{2m^2}{2m+1}h_{1,2i+1} & h_{2,2} + h_{1,2i+1} + \frac{3}{2} - \frac{(2m+1)^2 + 1}{2(2m+1)} \end{pmatrix}.
$$

We see that this matrix N_1 is diagonalizable and the eigenvalues are given by $h_{2,2i}$ and $h_{1,2(m-i)-1}$ which are the minimal conformal weights of X_{2i} and $X_{2(m-i)-1}$ respectively.

(b) Let us assume that $\langle \psi^*, \phi_1 \otimes \phi_2 \rangle$ is determined by the numbers [\(5.20\)](#page-55-1). By using Lemma 5.5 and (5.7) , we have

$$
\begin{pmatrix} \langle L_0 \psi^*, v^{\delta} \otimes u \rangle \\ \langle L_0 \psi^*, (L_{-1} v^{\delta}) \otimes u \rangle \end{pmatrix} = N_2 \begin{pmatrix} \langle \psi^*, v^{\delta} \otimes u \rangle \\ \langle \psi^*, (L_{-1} v^{\delta}) \otimes u \rangle \end{pmatrix},
$$

where

$$
N_2 = \begin{pmatrix} h_{2,2} + h_{1,2i+1} & 1 \\ \frac{2m^2}{2m+1}h_{1,2i+1} & h_{2,2} + h_{1,2i+1} + 1 - \frac{(2m+1)^2 + 1}{2(2m+1)} \end{pmatrix}.
$$

We see that this matrix N_2 is diagonalizable and eigenvalues are given by $h_{2,2i+2}$ and $h_{1,2(m-i)+1}$ which are the minimal conformal weights of X_{2i+2} and $X_{2(m-i)+1}$ respectively.

(c) Let us assume that the L_0 -weight of ψ^* is $h_{1,2(m-i)+1}$ or $h_{1,2(m-i)-1}$. Then, similar to the arguments in Lemma [5.8,](#page-48-0) we see that

$$
\langle \psi^*, U(\mathfrak{ns})v^{\pm} \otimes w_k^{(1)} \rangle = 0 \quad (k = 1, 0, -1). \tag{5.22}
$$

From the structure of the simple modules $X_{2(m-i)+1}$ and $X_{2(m-i)-1}$, we have

$$
\widehat{W}_0^{\pm} \psi^* = 0. \tag{5.23}
$$

By [\(3.5\)](#page-18-1) in Proposition [3.9\)](#page-18-0), we have

$$
\widehat{W}_{-h}^{\pm}u = 0 \quad (h < \frac{1}{2} - i + 2m). \tag{5.24}
$$

Then, by using the second formula in Lemma [5.5](#page-46-1) as $v = \widehat{W}^{\pm}$ and $n = 0$, and by using the relations (3.6) , (5.17) , (5.22) , (5.23) and (5.24) , we have $\langle \psi^*, v^{\pm} \otimes u \rangle = 0$. Thus we have a contradiction.

Hence, from (a), (b) and (c), the L_0 -weight of ψ^* is $h_{2,2i}$ or $h_{2,2i+2}$. Note that by Proposition [3.9,](#page-18-0)

$$
W_{-h}^{\pm}v^{\pm} = 0, \qquad \qquad \widehat{W}_{-h}^{\pm}v^{\pm} = 0 \qquad (5.25)
$$

for $h < h_{3,1}$. Then, noting the argument just before (a) and using Lemma [5.5](#page-46-1) and [\(5.24\)](#page-56-0)-[\(5.25\)](#page-56-1), we see that

$$
\langle \widehat{W}^{\pm}_0 \psi^*, v^{\pm} \otimes U(\mathfrak{ns})u \rangle = 0.
$$

Thus, noting that the minimal conformal weight spaces of X_{2i} and X_{2i+2} are two dimensional, we see that $\langle \psi^*, \phi_1 \otimes \phi_2 \rangle$ is determined by the numbers

$$
\langle \widehat{W}_0^{\pm} \psi^* + \psi^*, v^{\mp} \otimes u \rangle, \qquad \langle \widehat{W}_0^{\pm} \psi^* + \psi^*, L_{-1} v^{\mp} \otimes u \rangle
$$

or

$$
\langle \widehat{W}^{\pm}_0 \psi^* + \psi^*, G_{-\frac{1}{2}} \psi^{\mp} \otimes u \rangle, \qquad \qquad \langle \widehat{W}^{\pm}_0 \psi^* + \psi^*, G_{-\frac{1}{2}}^3 \psi^{\mp} \otimes u \rangle.
$$

Let $\binom{\kappa_1}{\kappa_2}$ $\binom{\kappa_1}{\kappa_2}$ be an eigenvector of ^tN₁ with the eigenvalue $h_{1,2(m-i)-1}$ and $\binom{\lambda_1}{\lambda_2}$ $\begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix}$ be an eigenvector of tN_2 with the eigenvalue $h_{1,2(m-i)+1}$. Similar to the argument in the proof of Lemma [5.8,](#page-48-0) we have

$$
\kappa_1 \langle \widehat{W}_0^{\pm} \psi^* + \psi^*, G_{-\frac{1}{2}} v^{\mp} \otimes u \rangle + \kappa_2 \langle \widehat{W}_0^{\pm} \psi^* + \psi^*, G_{-\frac{1}{2}}^3 v^{\mp} \otimes u \rangle = 0,
$$

$$
\lambda_1 \langle \widehat{W}_0^{\pm} \psi^* + \psi^*, v^{\mp} \otimes u \rangle + \lambda_2 \langle \widehat{W}_0^{\pm} \psi^* + \psi^*, L_{-1} v^{\mp} \otimes u \rangle = 0.
$$

Therefore $A_0((X_2 \boxtimes X_{2i+1})^*)$ is at most four dimensional.

Let us recall some properties of rigid and dual objects in tensor categories. For the following proposition, see, for example, [\[ESNO,](#page-66-8) [JS,](#page-69-3) [KL\]](#page-69-4).

 \Box

Proposition 5.12. Let (C, \otimes) be a tensor category, then we have:

1. Let V be a rigid object in \mathcal{C} . Then there is a natural adjunction isomorphism

$$
\operatorname{Hom}_{\mathcal{C}}(U \otimes V, W) \simeq \operatorname{Hom}_{\mathcal{C}}(U, W \otimes V^{\vee}),
$$

where U and W are any objects in \mathcal{C} , and V^{\vee} is the dual object of V.

2. Let V_1 and V_2 be rigid objects in C. Then $V_1 \otimes V_2$ is also rigid and

$$
(V_1 \otimes V_2)^{\vee} = V_2^{\vee} \otimes V_1^{\vee}.
$$

3. Let V be a rigid object in $\mathcal C$ and let P be a projective object in $\mathcal C$. Then $V^{\vee} \otimes P$ is projective.

4. Assume that

- ^C has enough projective and injective objects.
- All projective objects are injective and all injective objects are projective.
- All projective objects are rigid.

Then if

$$
0 \to V_1 \to V_2 \to V_3 \to 0
$$

is an exact sequence in C such that two of V_1 , V_2 , V_3 are rigid, then the third object is also rigid.

Proposition 5.13. For $s = 2, \ldots, 2m$, we have

$$
X_2 \boxtimes X_s = X_{s-1} \oplus X_{s+1}.
$$

Proof. First, let us show

$$
X_2 \boxtimes X_{2i} = X_{2i-1} \oplus X_{2i+1} \quad (i = 1, ..., m). \tag{5.26}
$$

By Lemma [5.6,](#page-47-2) we have

$$
I_{\mathcal{SW}(m)} \begin{pmatrix} X_{2i-1} \\ X_2 \ X_{2i} \end{pmatrix} \neq 0, \qquad I_{\mathcal{SW}(m)} \begin{pmatrix} X_{2i+1} \\ X_2 \ X_{2i} \end{pmatrix} \neq 0. \tag{5.27}
$$

By Lemma [5.8,](#page-48-0) Proposition [5.12](#page-56-2) and the self-duality of X_2 , we see that

Hom_{SW(m)}
$$
(X_{2(m-i+1)} \oplus X_{2(m-i)}, X_2 \boxtimes X_{2i}) = 0.
$$

Thus, by (5.27) , we obtain (5.26) .

Next, let us show

$$
X_2 \boxtimes X_{2i+1} = X_{2i} \oplus X_{2i+2} \quad (i = 0, \dots, m-1). \tag{5.28}
$$

By Lemma [5.6,](#page-47-2) we have

$$
I_{\mathcal{SW}(m)}\left(X_2\ X_{2i+1}\right) \neq 0, \qquad I_{\mathcal{SW}(m)}\left(X_{2i+2}\ X_{2i+1}\right) \neq 0. \tag{5.29}
$$

By Lemma [5.11,](#page-54-3) Proposition [5.12](#page-56-2) and the self-duality of X_2 , we see that

 $\text{Hom}_{\mathcal{SW}(m)}(X_{2(m-i)-1} \oplus X_{2(m-i)+1}, X_2 \boxtimes X_{2i+1}) = 0.$

Thus, by (5.29) , we obtain (5.28) .

Since $\mathcal{SW}(m)$ is C_2 -cofinite, every simple module has a projective cover [\[Hu\]](#page-68-0). In the following, we introduce an indecomposable $\mathcal{SW}(m)$ -module and some lemmas to determine the structure of these projective covers. Let us introduce an operator $\Delta_{Q_{-}}(-,z)$ - : $\mathcal{SW}(m) \times \mathcal{V}_L \rightarrow \overline{\mathcal{V}_{L+\gamma_{2m}}}[[z,z^{-1}]]$ as follows

$$
\Delta_{Q_{-}}(A, z) = \sum_{n \ge 1} \frac{(-1)^{n+1}}{nz^n} \oint_{z'=z} (z'-z)^n Q_{-}(z') Y(A, z) \mathrm{d} z' \quad (A \in \mathcal{SW}(m)).
$$

This operator $\Delta_{Q_{-}}$ is called logarithmic deformation [\[FFHST\]](#page-67-2) or Li's oper-ator [\[Li\]](#page-70-3). Let $W = \mathcal{V}_L \oplus \mathcal{V}_{L+\gamma_{2m}} (\in \mathcal{B}_1)$ and let Y_W be the ordinary $\mathcal{SW}(m)$ action on W. We define an operator $J(-, z) - : \mathcal{SW}(m) \times W \to \overline{W}[[z, z^{-1}]]$ as follows

$$
J(A, z) = \begin{cases} Y_W(A, z) + \Delta_{Q_{-}}(A, z) & \text{on } \mathcal{V}_L, \\ Y_W(A, z) & \text{on } \mathcal{V}_{L + \gamma_{2m}}, \end{cases}
$$

where $A \in \mathcal{SW}(m)$. By the results in [\[FFHST,](#page-67-2) [Li\]](#page-70-3), the operator J defines an $\mathcal{SW}(m)$ -action on W. We set $\mathcal{P} = (W, J)$. For the conformal vector T, the action $J(T, z)$ on the subspace $\mathcal{V}_L \subset \mathcal{P}$ is given by

$$
J(T, z) = T(z) + \frac{Q_{-}(z)}{z}.
$$

Thus, by Propositions [2.10,](#page-15-0) P is indecomposable and has L_0 -nilpotent rank two. By the definition, P has X_{2m} as a submodule and a quotient, and the total composition factors of P are given by $X_1 \oplus X_1 \oplus X_{2m} \oplus X_{2m}$.

Lemma 5.14. \mathcal{P}/X_{2m} or \mathcal{P}^*/X_{2m} is indecomposable.

Proof. Let X_{2m} be the minimal conformal weight space of X_{2m} . By the results in [\[KW,](#page-69-5) [Zh\]](#page-71-2), X_{2m} has the structure of an $A(\mathcal{SW}(m))$ -module. Then, from Theorem [3.7,](#page-17-2) we see that

$$
\operatorname{Ind}_{A(\mathcal{SW}(m))}^{\mathcal{SW}(m)} \overline{X}_{2m} \simeq X_{2m}.\tag{5.30}
$$

If neither \mathcal{P}/X_{2m} nor \mathcal{P}^*/X_{2m} is indecomposable, then, as the quotient of \mathcal{P} or \mathcal{P}^* , we have a non trivial extension in

$$
\text{Ext}^1_{\mathcal{SW}(m)}(X_{2m}, X_{2m}).
$$

But from [\(5.30\)](#page-58-0), this contradicts Theorem [3.11.](#page-19-1)

 \Box

 \Box

The following lemma can be proved in the same way as Lemmas [5.8](#page-48-0) and [5.11.](#page-54-3)

Lemma 5.15. The vector space $A_0((X_2 \boxtimes X_{2m+1})^*)$ is at most six dimensional. Any L_0 eigenvalue of $A_0((X_2 \boxtimes X_{2m+1})^*)$ is contained in $\{h_{1,1}, h_{2,2m}\} =$ ${0, \frac{1}{2}}$ $\frac{1}{2}$ }, where $h_{1,1}$ and $h_{2,2m}$ are the minimal conformal weights of X_1 and X_{2m} , respectively.

Lemma 5.16. For any simple $\mathcal{SW}(m)$ -module X, we have

$$
\operatorname{Hom}_{\mathcal{SW}(m)}(X_2 \boxtimes X_{2m+1}, X) = \begin{cases} \mathbb{C} & X = X_{2m}, \\ 0 & \text{otherwise} \end{cases}
$$

$$
\operatorname{Hom}_{\mathcal{SW}(m)}(X, X_2 \boxtimes X_{2m+1}) = \begin{cases} \mathbb{C} & X = X_{2m}, \\ 0 & \text{otherwise} \end{cases}
$$

Proof. By Propositions [5.12-](#page-56-2)[5.13](#page-57-4) and the self-duality of X_2 , we obtain the above eaqualities. \Box

Let us use the following notation:

- 1. For $1 \leq i \leq m$, let P_{2i} be the projective cover of $X_{2(m-i)+1}$.
- 2. For $0 \leq j \leq m-1$, let P_{2j+1} be the projective cover of $X_{2(m-j)}$.

By Proposition [3.14,](#page-20-0) we have $P_{2i} \in \mathcal{B}_{m-i+1}$ and $P_{2i+1} \in \mathcal{B}_{j+1}$.

Proposition 5.17. 1. We have

$$
X_2 \boxtimes X_{2m+1} = P_1.
$$

2. P_1 has the socle series $Soc_1(P_1) \subsetneq Soc_2(P_1) \subsetneq Soc_3(P_1) = P_1$ such that

$$
Soc1(P1) \simeq X2m, Soc2(P1)/Soc1(P1) \simeq X1 \oplus X1,Soc3(P1)/Soc2(P1) \simeq X2m.
$$
\n(5.31)

Proof. Note that from Proposition [3.14,](#page-20-0) X_{2m+1} is projective. Then, by the self-duality of X_2 , $X_2 \boxtimes X_{2m+1}$ must be projective. Thus, by Lemma [5.16,](#page-59-0) we obtain

$$
X_2 \boxtimes X_{2m+1} = P_1.
$$

By Lemma [5.14](#page-58-1) and the projectivity of P_1 , we see that P_1 has $X_1 \oplus X_1$ as composition factors. Thus, by Lemmas $5.15-5.16$, we see that P_1 satisfies the socle series [\(5.31\)](#page-59-2). \Box **Proposition 5.18.** 1. For each $1 \leq s \leq 2m$, the tensor product $X_2 \boxtimes P_s$ is given by:

• For $s = 1$, we have

$$
X_2 \boxtimes P_1 = X_{2m+1} \oplus X_{2m+1} \oplus P_2.
$$

• For $2 \leq s \leq 2m-1$, we have

$$
X_2 \boxtimes P_s = P_{s-1} \oplus P_{s+1}.
$$

• For $s = 2m$, we have

$$
X_2 \boxtimes P_{2m} = X_{2m+1} \oplus X_{2m+1} \oplus P_{2m-1}.
$$

- 2. The socle series of the projective covers of the simple modules are given by:
	- For $1 \leq i \leq m$, we have $\text{Soc}_1(P_{2i}) \subsetneq \text{Soc}_2(P_{2i}) \subsetneq \text{Soc}_3(P_{2i}) = P_{2i}$ such that

 $\text{Soc}_1(P_{2i}) \simeq X_{2(m-i)+1}, \ \text{Soc}_2(P_{2i})/\text{Soc}_1(P_{2i}) \simeq X_{2i} \oplus X_{2i},$ $\text{Soc}_3(P_{2i})/\text{Soc}_2(P_{2i}) \simeq X_{2(m-i)+1}.$

• For $0 \leq i \leq m-1$, we have $\text{Soc}_1(P_{2i+1}) \subsetneq \text{Soc}_2(P_{2i+1}) \subsetneq \text{Soc}_3(P_{2i+1}) =$ P_{2i+1} such that

 $\text{Soc}_1(P_{2i+1}) \simeq X_{2(m-i)}, \text{ Soc}_2(P_{2i+1})/\text{Soc}_1(P_{2i+1}) \simeq X_{2i+1} \oplus X_{2i+1},$ $\text{Soc}_3(P_{2i+1})/\text{Soc}_2(P_{2i+1}) \simeq X_{2(m-i)}$.

Proof. We only prove that $X_2 \boxtimes P_1 = X_{2m+1} \oplus X_{2m+1} \oplus P_2$ and

$$
Soc1(P2) \simeq X2m-1, Soc2(P2)/Soc1(P2) \simeq X2 \oplus X2,P2/Soc2(P2) \simeq X2m-1.
$$
\n(5.32)

The other cases can be proved in the same way.

From Propositions [5.12,](#page-56-2) [5.13,](#page-57-4) [5.17](#page-59-3) and the self-duality of X_2 , we can see that for any simple $\mathcal{SW}(m)$ -module X

$$
\text{Hom}_{\mathcal{SW}(m)}(X_2 \boxtimes P_1, X) = \begin{cases} \mathbb{C}^2 & X = X_{2m+1}, \\ \mathbb{C} & X = X_{2m-1}, \\ 0 & \text{otherwise} \end{cases} \tag{5.33}
$$
\n
$$
\text{Hom}_{\mathcal{SW}(m)}(X, X_2 \boxtimes P_1) = \begin{cases} \mathbb{C}^2 & X = X_{2m+1} \\ \mathbb{C} & X = X_{2m-1}, \\ 0 & \text{otherwise} \end{cases} \tag{5.33}
$$

Since X_2 is rigid and P_1 is projective, $X_2 \boxtimes P_1$ is also projective. Thus, from [\(5.33\)](#page-60-0), we obtain $X_2 \boxtimes P_1 = X_{2m+1} \oplus X_{2m+1} \oplus P_2$. Therefore, by the rigidity of X_2 and Proposition [5.13,](#page-57-4) we see that P_2 satisfies the socle series [\(5.32\)](#page-60-1). \Box

In [\[AM3\]](#page-65-1), the equivalence between $\mathcal{SW}(m)$ -mod and $U_q^{small}(sl_2)$ -mod is conjectured, where $U_q^{small}(sl_2)$ -mod is the abelian category of finite dimensional modules over the small quantum group $U_q^{small}(sl_2)$ at $q = e^{\frac{2\pi i}{2m+1}}$. At the level of the arbelian category, we see that this conjecture is true. That is, the following corollary holds.

Corollary 5.19. Two categories $\mathcal{SW}(m)$ -mod and $U_q^{small}(sl_2)$ -mod are equivalent as abelian categories.

Proof. Similar to the arguments in [\[NT,](#page-70-4) Section 6], using the structure of the projective $\mathcal{SW}(m)$ -modules given in Propositions [5.17-](#page-59-3)[5.18](#page-60-2) and the projective $U_q^{small}(sl_2)$ -modules classified by [Kü, [Sut,](#page-71-3) [Xi\]](#page-71-4), we can prove the equivalence of the two abelian categories. We omit the details. \Box

Remark 5.20. Let $q = e^{\frac{2\pi i}{2m+1}}$. The small quantum group $U_q^{small}(sl_2)$ is an associative \mathbb{C} -algebra which is generated by E, F, K, K^{-1} satisfying the following fundamental relations

$$
KK^{-1} = K^{-1}K = 1, \ KEK^{-1} = q^2E, \ KFK^{-1} = q^{-2}F,
$$

\n
$$
EF - FE = \frac{K - K^{-1}}{q - q^{-1}},
$$

\n
$$
E^{2m+1} = F^{2m+1} = 0, \ K^{2m+1} = 1.
$$

See $[K\ddot{u}, Sut, Xi]$ $[K\ddot{u}, Sut, Xi]$ $[K\ddot{u}, Sut, Xi]$ for details.

Finally, let us show that the tensor supercategory $(\mathcal{SW}(m)\text{-mod}, \mathbb{Z})$ is rigid. Note that the tensor supercategory $(\mathcal{SW}(m) \text{-mod}, \boxtimes)$ is weakly rigid. In fact, we have the following proposition.

Proposition 5.21 ([\[ALSW,](#page-65-2) [HLZ2,](#page-68-3) [Xu\]](#page-71-5)). Let C be a vertex tensor (super)category with the unit 1. Given $U, V \in \mathcal{C}$, assume $V^* \in \mathcal{C}$, where V^* is the contragredient of V . Then, we have a natural isomorphism

$$
\operatorname{Hom}_{\mathcal{C}}(U,V) \simeq \operatorname{Hom}_{\mathcal{C}}(U \boxtimes V^*, \mathbf{1}).
$$

Theorem 5.22. The braided tensor supercategory $(\mathcal{SW}(m)$ -mod, $\boxtimes)$ is rigid. For any $M \in \mathcal{SW}(m)$ -mod, we have $M^{\vee} = M^*$, where M^{\vee} is the dual of M. Proof. From Propositions [5.13](#page-57-4) and [5.18,](#page-60-2) all simple and projective modules are rigid and self-dual. By the structure of the projective modules, we see that all indecomposable modules $M \in \mathcal{SW}(m)$ -mod except the simple modules and the projective modules satisfy exact sequence

$$
0 \to L \to M \to N \to 0
$$

such that L and N are direct sum of simple modules. Then, from Proposition [5.12,](#page-56-2) we see that M is rigid. Since

$$
\operatorname{Hom}_{\mathcal{SW}(m)}(M \boxtimes M^*, X_1) = \operatorname{Hom}_{\mathcal{SW}(m)}(M, M) \simeq \mathbb{C}
$$

from Proposition [5.21,](#page-61-0) we have $M^{\vee} = M^*$.

5.4 Fusion rings

In this subsection, following the argument in [\[TW1,](#page-71-1) Subsection 5.3], we introduce two fusion rings $P(SW(m))$ and $K(SW(m))$, and determine their structure.

Let \mathbb{I}_m be the set consisting of all simple modules $X_s(1 \leq s \leq 2m+1)$ and all projective modules $P_s(1 \leq s \leq 2m)$. We introduce the free abelian group $P(\mathcal{SW}(m))$ of rank $4m+1$ generated by the elements of \mathbb{I}_m :

$$
P(\mathcal{SW}(m)) = \bigoplus_{s=1}^{2m+1} \mathbb{Z}[X_s]_P \oplus \bigoplus_{s=1}^{2m} \mathbb{Z}[P_s]_P.
$$

Then, from the results presented in the previous subsection, we can define the structure of a commutative ring on $P(SW(m))$ such that the product as a ring is given by

$$
[M_1]_P \cdot [M_2]_P = [M_1 \boxtimes M_2]_P
$$

for $M_1, M_2 \in \mathbb{I}_m$, where we extend the symbol $[\bullet]_P$ as follows

$$
\left[\bigoplus_{i\geq 1}^n N_i\right]_P = \bigoplus_{i\geq 1}^n [N_i]_P
$$

for any $N_i \in \mathbb{I}_m$ and any $n \in \mathbb{Z}_{\geq 1}$.

The operator

$$
X = X_2 \boxtimes -
$$

 \Box

define a Z-linear endomorphism of $P(SW(m))$. Thus $P(SW(m))$ is a module over $\mathbb{Z}[X]$. We define the following $\mathbb{Z}[X]$ -module map

$$
\psi : \mathbb{Z}[X] \to P(\mathcal{SW}(m)),
$$

$$
f(X) \mapsto f(X) \cdot [X_1]_P.
$$

Before examining the action of $\mathbb{Z}[X]$ on $P(\mathcal{SW}(m))$, we introduce the following Chebyshev polynomials.

Definition 5.23. We define Chebyshev polynomials $U_n(A)$, $n = 0, 1, \dots \in$ $\mathbb{Z}[A]$ recursively

$$
U_0(A) = 1,
$$
 $U_1(A) = A,$ $U_{n+1}(A) = AU_n(A) - U_{n-1}(A).$

Remark 5.24. The coefficient of the leading term of any Chebyshev polynomial $U_n(A)$ is 1. Thus we have

$$
\mathbb{Z}[A] = \bigoplus_{n=0}^{\infty} \mathbb{Z}U_n(A).
$$

From the results of previous subsection, we obtain the following lemma.

Lemma 5.25. 1. For $s = 1, ..., 2m + 1$, we have

$$
[X_s]_P = U_{s-1}(X)[X_1]_P.
$$

2. For $s = 1, \ldots, 2m$, we have

$$
[P_s]_P = (U_{2m+s}(X) + U_{2m-s}(X))[X_1]_P.
$$

3. We have the relation

$$
U_{4m+1}(X)[X_1]_P = 2U_{2m}(X)[X_1]_P.
$$

From this lemma, we obtain the following theorem.

Theorem 5.26. The $\mathbb{Z}[X]$ -module map ψ is surjective and the kernel of ψ is given by the ideal ker $\psi = \langle U_{4m+1}(X) - 2U_{2m}(X)\rangle.$

Proof. By Lemma [5.25,](#page-63-0) we see that ψ is surjective. We define the following ideal of $\mathbb{Z}[X]$

$$
I = \langle U_{4m+1}(X) - 2U_{2m}(X) \rangle.
$$

By the third statement in Lemma [5.25,](#page-63-0) we see that I is contained in ker ψ . It is easy to see that the dimension of $\mathbb{Z}[X]/I$ is $4m+1$. Therefore we obtain $\ker \psi = I.$

 \Box

Next, let us state the results for the Grothendieck fusion ring of $\mathcal{SW}(m)$. We introduce the rank $2m + 1$ Grothendieck group

$$
K(\mathcal{SW}(m)) = \bigoplus_{s=1}^{2m+1} \mathbb{Z}[X_s]_K.
$$

From the results presented in the previous subsection, we see that $K(SW(m))$ has the structure of a commutative ring whose unit object is $[X_1]_K$. The operator $X = X_2 \boxtimes -$ define a Z-linear endomorphism of $K(\mathcal{SW}(m))$. Thus $K(SW(m))$ is a module over $\mathbb{Z}[X]$. Then we can define the following $\mathbb{Z}[X]$ module map

$$
\phi: \mathbb{Z}[X] \to K(\mathcal{SW}(m)),
$$

$$
f(X) \mapsto f(X) \cdot [X_1]_K.
$$

Similar to the arguments in the case of $P(SW(m))$, we obtain the following proposition.

Proposition 5.27. The $\mathbb{Z}[X]$ -module map ϕ is surjective and the kernel of ϕ is given by the ideal ker $\phi = \langle U_{2m+1}(X) - U_{2m-1}(X) - 2 \rangle$.

Remark 5.28. In [\[TW1,](#page-71-1) Subsection 5.3], a non-semisimple fusion ring $P(\mathcal{W}_p)$ of the triplet algebra \mathcal{W}_p is introduced. As in the case of $P(\mathcal{SW}(m))$, $P(\mathcal{W}_p)$ is defined by adding a ring structure determined from the tensor product to the free abelian group generated from all simple and projective W_p -modules. As shown in [\[TW1\]](#page-71-1), $P(\mathcal{W}_p)$ is isomorphic to the quotient ring

$$
\frac{\mathbb{Z}[X,Y]}{\langle Y^2-1, U_{2p-1}(X)-2YU_{p-1}(X)\rangle}
$$

.

The Y variable corresponds to the simple current of \mathcal{W}_p and the X variable to a simple \mathcal{W}_p -module which has a weight two Virasoro null vector (according to the notation in [\[FGST3,](#page-67-3) [NT,](#page-70-4) [TW1\]](#page-71-1), Y to \mathcal{X}_1^- and X to \mathcal{X}_2^+). For this ring, setting $p = 2m + 1$ and $Y = 1$, we have a quotient ring isomorphic to $P(SW(m))$ (see Theorem [5.27\)](#page-64-0). Similarly, we can obtain the Grothendieck ring $K(SW(m))$ as a quotient of the Grothendieck ring $K(W_{2m+1})$ determined by $|TW1|$. In $|AMS|$, it is shown that the characters of the simple $\mathcal{SW}(m)$ -modules can be expressed in terms of the characters of the simple \mathcal{W}_{2m+1} -modules. From these results, $\mathcal{SW}(m)$ -mod and \mathcal{W}_{2m+1} -mod seem to be closely related at the level of tensor categories.

Acknowledgement

We would like to thank Koji Hasegawa for reading the manuscript and for valuable comments. We would also like to thank Yuto Moriwaki and Shigenori Nakatsuka for stimulating discussions.

References

- [AM1] D. Adamović and A. Milas, "On the triplet vertex algebra $W(p)$," Advances in Mathematics 217 (2008) 2664-2699.
- [AM2] D. Adamović and A. Milas, "The $N=1$ triplet vertex operator superalgebras: twisted sector," SIGMA. Symmetry, Integrability and Geometry: Methods and Applications, 4 (2008) 087.
- [AM3] D. Adamović and A. Milas, "The $N = 1$ triplet vertex operator superalgebras," Communications in Mathematical Physics, 288 (2009) 225- 270.
- [AM4] D. Adamović and A. Milas, "The structure of Zhu's algebras for certain W-algebras," Advances in Mathematics 227 (2011) 2425-2456.
- [ALSW] R. Allen, S. Lentner, C. Schweigert and S. Wood, "Duality structures for module categories of vertex operator algebras and the Feigin Fuchs boson," (2021), [arXiv:2107.05718.](http://arxiv.org/abs/2107.05718)
- [Ba] K. D. Barron, "N=1 Neveu-Schwarz vertex operator superalgebras over Grassmann algebras and with odd formal variables," (1999), [arXiv:math/9910007\[math.QA\].](https://arxiv.org/abs/math/9910007)
- $[Be]$ V. A. Belavin, "On the N= 1 super Liouville four-point functions," Nuclear Physics B, 798 (2008) 423-442.
- [BPZ] A. A. Belavin, A. M. Polyakov and A. B. Zamolodchikov, "Infinite conformal symmetry in two-dimensional quantum field theory," Nuclear Physics B, 241 (1984) 333-380.
- [BS] L. Benoit and Y. Saint-Aubin, "Fusion and the Neveu-Schwarz singular vectors," International Journal of Modern Physics A 9 (1994) 547.
- [BMRW] O. Blondeau-Fournier, P. Mathieu, D. Ridout and S. Wood, "Superconformal minimal models and admissible Jack polynomials," Advances in Mathematics 314 (2017) 71-123.
- [CRR] M. Canagasabey, J. Rasmussen and D. Ridout, "Fusion rules for the $N = 1$ superconformal logarithmic minimal models I: The Neveu-Schwarz sector," Journal of Physics A: Mathematical and Theoretical, 48 (2015) 415402.
- [CGNS] T. Creutzig, N. Genra, S. Nakatsuka and R. Sato, "Correspondences of categories for subregular W-algebras and principal W-superalgebras," Communications in Mathematical Physics, 393 (2022) 1-60.
- [CKL] T. Creutzig, S. Kanade and A. R. Linshaw, "Simple current extensions beyond semi-simplicity," *Commun. Contemp. Math.* 22 (2020), no. 1, 1950001, 49 pp.
- [CKM] T. Creutzig, S. Kanade and R. McRae, "Tensor categories for vertex operator superalgebra extensions," (2017), [arXiv:1705.05017\[math.QA\].](https://arxiv.org/abs/1705.05017)
- [CLR] T. Creutzig, S. Lentner and M. Rupert, "An algebraic theory for logarithmic Kazhdan-Lusztig correspondences," (2023), [arXiv:2306.11492\[math.QA\].](https://arxiv.org/abs/2306.11492)
- [CMY1] T. Creutzig, R. McRae and J. Yang, "On ribbon categories for singlet vertex algebras," Communications in Mathematical Physics, 387 (2021) 865-925.
- [CMY2] T. Creutzig, R. McRae and J. Yang, "Tensor structure on the Kazhdan–Lusztig category for affine $\mathfrak{gl}(1|1)$," International Mathematics Research Notices, 16 (2022) 12462-12515.
- [CMOY] T. Creutzig, R. McRae, F. Orosz Hunziker and J. Yang, " $N = 1$ " super Virasoro tensor categories," (2024), [arXiv:2412.18127\[math.QA\].](https://arxiv.org/abs/2412.18127)
- [CR] T. Creutzig and D. Ridout, "Logarithmic conformal field theory: Beyond an introduction," J. Phys. A46 (2013) 494006.
- [DF1] V. S. Dotsenko and V. A. Fateev, "Conformal algebra and multipoint correlation functions in 2D statistical models," Nucl. Phys. B240 (1984) 312-348.
- [DF2] V. S. Dotsenko and V. A. Fateev, "Four-point correlation functions and the operator algebra in 2D conformal invariant theories with central charge $c \le 1$," *Nucl. Phys.* **B251** (1985) 691-734.
- [ESNO] P. Etingof, G. Shlomo, D. Nikshych and V. Ostrik, Tensor Categories. Number volume 205 in Mathematical Surveys and Monographs. American Mathematical Society 2015.
- [FGST1] B. L. Feigin, A. M. Gainutdinov, A. M. Semikhatov and I. Y. Tipunin, "The Kazhudan-Lusztig correspondence for the representation category of the triplet W-algebra in logarithmic conformal field theories," Theoret. and Math. Phys. 148 (2006) 1210-1235.
- [FGST2] B. L. Feigin, A. M. Gainutdinov, A. M. Semikhatov and I. Y. Tipunin, "Modular group representations and fusion in logarithmic conformal field theories and in the quantum group center," Comm. Math. Phys. 265 (2006) 47–93.
- [FGST3] B. L. Feigin, A. M. Gainutdinov, A. M. Semikhatov and I. Y. Tipunin, "Kazhdan-Lusztig correspondence for the representation category of the triplet W-algebra in logarithmic CFT," Theor.Math.Phys. 148 (2006) 1210-1235; Teor.Mat.Fiz. 148 (2006) 398-427.
- [FGST4] B. L. Feigin, A. M. Gainutdinov, A. M. Semikhatov and I. Y. Tipunin, "Logarithmic extensions of minimal models: characters and modular transformation," Nucl. Phys. B757 (2006) 303-343.
- [Fe] G. Felder, "BRST approach to minimal models," Nuclear Phys. B317 (1989) 215-236.
- [FS] G. Felder and R. Silvotti, "Conformal blocks of minimal models on a Riemann surface," Commun. Math. Phys. 144 (1992) 17-40.
- [FFHST] J. Fjelstad, J. Fuchs, S. Hwang, A. M. Semikhatov and I. Y. Tipunin, "Logarithmic conformal field theories via logarithmic deformation," Nucl. Phys. B633 (2002) 379-413.
- [Fo1] P. J. Forrester, "Integration formulas and exact calculations in the Calogero–Sutherland model," Modern Physics Letters B, 9 (1995) 359- 371.
- [Fo2] P. J. Forrester, Log-gases and random matrices (LMS-34). Princeton University Press 2010.
- [FLM] I. Frenkel, J. Lepowsky and A. Meurman, Vertex operator algebras and the Monster. Academic press 1989.
- [GK1] M. R. Gaberdiel and H. G. Kausch, "Indecomposable fusion products," Nucl. Phys. B477 (1996) 293-318.
- [GK2] M. R. Gaberdiel and H. G. Kausch, "A Local Logarithmic Conformal Field Theory," Nucl. Phys. B538 (1999) 631-658.
- [GN] T. Gannon and C. Negron, "Quantum SL(2) and logarithmic vertex operator algebras at $(p, 1)$ -central charge," (2021) , [arXiv:2104.12821\[math.QA\].](https://arxiv.org/abs/2104.12821)
- [Gu] V. Gurarie, "Logarithmic Operators in Conformal Field Theory," Nucl. Phys. **B410** (1993) 535-549.
- [Ha] Y. Haraoka, Linear Differential Equations in the Complex Domain. Lecture Notes in Mathematics Vol. 2271, Springer, New York 2020.
- [Hu] Y. Z. Huang, "Cofiniteness conditions, projective covers and the logarithmic tensor product theory," J. Pure Appl. Algebra, 213 (2009) 458- 475.
- [HLZ1] Y. Z. Huang, J. Lepowsky and L. Zhang, "Logarithmic tensor category theory for generalized modules for a conformal vertex algebra, I: Introduction and strongly graded algebras and their generalized modules," Proceedings of a Workshop Held at Beijing International Center for Mathematical Research. Springer Berlin Heidelberg, (2014) 169-248.
- [HLZ2] Y. Z. Huang, J. Lepowsky and L. Zhang, "Logarithmic tensor category theory for generalized modules for a conformal vertex algebra, II: Logarithmic formal calculus and properties of logarithmic intertwining operators," (2010), [arXiv:1012.4196\[math.QA\].](https://arxiv.org/abs/1012.4196)
- [HLZ3] Y. Z. Huang, J. Lepowsky and L. Zhang, "Logarithmic tensor category theory for generalized modules for a conformal vertex algebra, III: Intertwining maps and tensor product bifunctors," (2010), [arXiv:1012.4197\[math.QA\].](https://arxiv.org/abs/1012.4197)
- [HLZ4] Y. Z. Huang, J. Lepowsky and L. Zhang, "Logarithmic tensor category theory for generalized modules for a conformal vertex algebra, IV: Constructions of tensor product bifunctors and the compatibility conditions," (2010), [arXiv:1012.4198\[math.QA\].](https://arxiv.org/abs/1012.4198)
- [HLZ5] Y. Z. Huang, J. Lepowsky and L. Zhang, "Logarithmic tensor category theory for generalized modules for a conformal vertex algebra, V: Convergence condition for intertwining maps and the corresponding compatibility condition," (2010), [arXiv:1012.4199\[math.QA\].](https://arxiv.org/abs/1012.4199)
- [HLZ6] Y. Z. Huang, J. Lepowsky and L. Zhang, "Logarithmic tensor category theory for generalized modules for a conformal vertex algebra, VI: Expansion condition, associativity of logarithmic intertwining operators, and the associativity isomorphisms," (2010), [arXiv:1012.4202\[math.QA\].](https://arxiv.org/abs/1012.4202)
- [HLZ7] Y. Z. Huang, J. Lepowsky and L. Zhang, "Logarithmic tensor category theory for generalized modules for a conformal vertex algebra, VII: Convergence and extension properties and applications to expansion for intertwining maps," (2011), [arXiv:1110.1929\[math.QA\].](https://arxiv.org/abs/1110.1929)
- [HLZ8] Y. Z. Huang, J. Lepowsky and L. Zhang, "Logarithmic tensor category theory for generalized modules for a conformal vertex algebra, VIII: Braided tensor category structure on categories of generalized modules for a conformal vertex algebra," (2011), [arXiv:1110.1931\[math.QA\].](https://arxiv.org/abs/1110.1931)
- [HM] Y. Z. Huang and A. Milas, "Intertwining operator superalgebras and vertex tensor categories for superconformal algebras, I," Communications in Contemporary Mathematics, 4 (2002) 327-355.
- [IK1] K. Iohara and Y. Koga, "Representation theory of Neveu-Schwarz and Ramond algebras I: Verma modules," Adv. Math., 178 (2003) 1-65.
- [IK2] K. Iohara and Y. Koga, "Representation theory of Neveu-Schwarz and Ramond Algebra II: Fock modules," Ann. Inst. Fourier, Grenoble, 53, 6 (2003) 1755-1818.
- [JPR] M. Jeng, G. Piroux and P. Ruelle, "Height variables in the Abelian sandpile model: scaling fields and correlations," J. Stat. Mech, **0610** (2006) P105.
- [JS] A. Joyal and R. Street, "Braided tensor categories," Adv. Math. 102 (1993) 20-78.
- [KW] V. Kac and W. Wang, "Vertex Operator Superalgebras and Their," Mathematical aspects of conformal and topological field theories and quantum groups, 175, 161 (1994).
- [KR] S. Kanade and D. Ridout, "NGK and HLZ: Fusion for physicists and mathematicians," Affine, Vertex and W-algebras (2019) 135-181.
- [Ka] H. G. Kausch, "Extended conformal algebras generated by multiplet of primary fields," Phys. Lett. B, 259 (1991) 448-455
- [KL] D. Kazhdan and G. Lusztig, "Tensor structures arising from affine Lie algebras. IV," Journal of the American Mathematical Society, 7 (1994) 383-453.
- [Kü] J. Külshammer, "Representation type and Auslander-Reiten theory of Frobenius-Lusztig kernels," Ph.D. thesis, Christian-Albrechts-Universität zu Kiel (2012) .
- [Li] H. Li, "The physics superselection principle in vertex operator algebra theory," Journal of Algebra 196 (1997), no. 2, 436-457.
- [MR] P. Mathieu and D. Ridout, "From Percolation to Logarithmic Conformal Field Theory," Phys. Lett. B657 (2007) 120-129.
- [MY] R. McRae and J. Yang, "Structure of Virasoro tensor categories at central charge $13 - 6p - 6p^{-1}$ for integers $p > 1$," (2021), [arXiv:2011.02170\[math.QA\].](https://arxiv.org/abs/2011.02170)
- [NT] K. Nagatomo and A. Tsuchiya, "The Triplet Vertex Operator Algebra $W(p)$ and Restricted Quantum Group at Root of Unity," *Exploring new* structures and natural constructions in mathematical physics. Vol. 61. Mathematical Society of Japan, (2011) 1-50.
- [Nah] W. Nahm, "Quasirational fusion products," Int. J. Mod. Phys. B8 (1994) 3693-3702.
- [Nak] H. Nakano, "The category of modules of the triplet W-algebras associated to the Virasoro minimal models," the Doctor Thesis, Mathematical Institute, Tohoku University (2023).
- [Ni] A. Nigro, "Integrals of Motion for Critical Dense Polymers and Symplectic Fermions," J. Stat. Mech. 0910 (2009) P10007.
- [PR] P. A. Pearce and J. Rasmussen, "Solvable critical dense polymers," J. Stat. Mech. 0702 (2007) P015.
- [RS] N. Read and H. Saleur, "Associative-algebraic approach to logarithmic conformal field theories," Nucl. Phys. B777 (2007) 316-351.
- [Ri] D. Ridout, "On the Percolation BCFT and the Crossing Probability of Watts," Nucl. Phys, B810 (2009) 503-526.
- [RW] D. Ridout and S. Wood, "Modular transformations and Verlinde formulae for logarithmic (p_+, p_-) -models," *Nucl. Phys.* **B880** (2014) 175-202.
- [Su1] E. Sussman, "The singularities of Selberg–and Dotsenko–Fateev-like integrals," (2023), [arXiv:2301.03750\[math.QA\].](https://arxiv.org/abs/2301.03750)
- [Su2] E. Sussman, "The regularization of Dotsenko–Fateev integrals," Letters in Mathematical Physics, 113 (2023) 1-29.
- [Sut] R. Suter, "Modules over $\mathfrak{U}_{q}(\mathfrak{sl}_2)$," *Communications in Mathematical* Physics, 163 (1994) 359-393.
- [TK] A. Tsuchiya and Y. Kanie, "Fock space representations of the Virasoro algebra - Intertwining operators," Publ. RIMS, Kyoto Univ. 22 (1986) 259-327.
- [TW1] A. Tsuchiya and S. Wood, "The tensor structure on the representation category of the \mathcal{W}_p triplet algebra," J. Phys. A 46 (2013) 445203.
- [TW2] A. Tsuchiya and S. Wood, "On the extended W-algebra of type sl_2 at positive rational level," International Mathematics Research Notices 2015.14 (2015) 5357-5435.
- [Xi] J. Xiao, "Restricted representations of $U(sl(2))$ -quantizations," Algebra Colloquium, 1 (1994) 56-66.
- [Xu] X. P. Xu, "Intertwining operators for twisted modules of a colored vertex operator superalgebra," Journal of Algebra, 175 (1995) 241-273.
- [Zh] Y. Zhu, "Modular invariance of characters of of vertex operator algebras," J. Amer. Math. Soc. 9 (1996) 237-302.

H. Nakano, Advanced Mathematical Institute, Osaka Metropolitan University, Osaka 558-8585, Japan E-mail address : hiromutakati@gmail.com