Quantum tunneling and its absence in deep wells and strong magnetic fields

Charles L. Fefferman a,1 , Jacob Shapiro a,1 , and Michael I. Weinstein b,1

^aDepartment of Mathematics, Princeton University; Princeton, New Jersey USA; ^bDepartment of Applied Physics and Applied Mathematics, and Department of Mathematics, Columbia University; New York, NY USA

This manuscript was compiled on January 3, 2025

We present new results on quantum tunneling between deep potential wells, in the presence of a strong constant magnetic field. We construct a family of double well potentials containing examples for which the low-energy eigenvalue splitting vanishes, and hence quantum tunneling is eliminated. Further, by deforming within this family, the magnetic ground state can be made to transition from symmetric to anti-symmetric. However, for typical double wells in a certain regime, tunneling is not suppressed, and we provide a lower bound for the eigenvalue splitting.

Quantum tunneling | hopping probability | magnetic fields | eigenvalue degeneracy

T unneling is a phenomenon of central importance in quantum systems. An important paradigm is a quantum particle, such as an electron, in a double well potential. The ground state tunneling rate (tunneling probability) between weakly coupled wells is related to the magnitude of the energy splitting of the two lowest eigenvalues of the double well Hamiltonian: in a purely classical description the lowest two states of the system are degenerate. Physicists and chemists have long known (1), and mathematicians have proven (2, 3), that in the regime of deep potential wells (one way to realize the weakly-coupled regime), the energy splitting in non-magnetic quantum systems (a) is always non-zero and exponentially small in the depth of the wells, and (b) is asymptotically related to the hopping coefficient, an overlap integral of ground state atomic orbitals.

In this paper we consider two dimensional double well quantum systems in a constant perpendicular magnetic field. In (4) we studied a Schrödinger double well model in which each single potential well is radially symmetric; see also (5, 6). The constant magnetic field strength of order $\sim \lambda$ and the potential well depths $\sim \lambda^2$ are taken to be large. We found (see Theorem 1) that (a) the eigenvalue splitting (and hence tunneling rate) is controlled by the hopping coefficient and (b) is always non-zero and exponentially small in λ .

In this article we announce new results on the general case, where the single well is no longer constrained to be radially symmetric. We exhibit a family of double well potentials containing cases for which the low-energy eigenvalue splitting vanishes and quantum tunneling is eliminated. Further, by deforming within this family, the magnetic ground state can be made to transition from symmetric to anti-symmetric. For <u>typical</u> double wells in a certain regime, we prove that tunneling is not suppressed, and we provide a lower bound for the eigenvalue splitting.

Mathematical setup. We consider a quantum mechanical particle in two space dimensions, with a constant magnetic field perpendicular to the plane. We shall study two related Hamil-

tonians: the one well Hamiltonian h_{λ} and the two well Hamiltonian H_{λ} .

The magnetic one well Hamiltonian, h_{λ} . Let $x = (x_1, x_2)^{\top}$ and $x^{\perp} = (x_2, -x_1)^{\top}$. In the symmetric gauge for the magnetic vector potential, the one well magnetic Hamiltonian is given by

$$h_{\lambda} = \left(P + \frac{1}{2}\lambda bX^{\perp}\right)^{2} + \lambda^{2}v(X).$$

Here, Xf(x) = xf(x) and $P = -i\nabla$ denote the position operator and the momentum operator, respectively. We take the one well $v : \mathbb{R}^2 \to (-\infty, 0]$ to be smooth, nonzero and compactly supported within $B_a(0)$, the disc of radius a > 0in \mathbb{R}^2 . The parameter $\lambda > 0$, which sets the scale of both the depth of the potential and the magnetic field strength, is taken to be sufficiently large. The parameter b > 0 is the relative (order 1) strength of the magnetic field. For fixed b, the effect of the magnetic field and that of a potential well with a non-degenerate minimum are of the same order of magnitude as $\lambda \to \infty$.

Let $e_{\lambda} \sim -\lambda^2$ be the ground state eigenvalue of h_{λ} . Throughout this paper, we assume that (i) e_{λ} is simple, and that (ii) e_{λ} remains bounded away from all other energies in the spectrum of h_{λ} as $\lambda \to \infty$. Assumption (ii) is valid if, for example, v has an isolated non-degenerate minimum or if v is very flat near its minimum, as for the square well.

We denote the corresponding $L^2(\mathbb{R}^2)$ -normalized eigenfunction by $\varphi_{\lambda}(x)$; to simplify notation we will often write merely $\varphi(x)$. We emphasize that, in contrast to the non-magnetic

Significance Statement

We establish surprising phenomena in magnetic systems, in strong contrast to the long-understood phenomena in nonmagnetic systems. We construct quantum double well systems such that, in a strong magnetic field, the tunneling effect between wells is completely eliminated. Moreover, through appropriate deformation of the double well, the ground state can be made to transition from symmetric to anti-symmetric. This raises the question whether such phenomena could be observable in suitable experimental systems. Further, our results suggest an approach to achieving flat bands, which are of great interest in the study of strongly-correlated quantum systems. Some of our results rely on basic estimates on analytic functions, and could prove useful for other problems in mathematical physics.

All authors contributed equally to the research and writing of this article. No conflicts of interest.

²To whom correspondence should be addressed. E-mail: author.twoemail.com

case, it may not be possible to choose the ground state to be real-valued.

The magnetic double well Hamiltonian, H_{λ} . In some of the discussion below, we shall allow for $v(x) = v(x; \lambda)$, i.e., the one well potential depends on the large parameter λ . We replace the one well potential $v(x; \lambda)$ by two copies of v, one displaced by $d \in \mathbb{R}^2$, the other by -d. The corresponding double well Hamiltonian is:

$$H_{\lambda} = \left(P + \frac{1}{2}\lambda bX^{\perp}\right)^2 + v^L(X;\lambda) + v^R(X;\lambda), \qquad [1a]$$

In the most basic case, we take

$$v^{L}(X;\lambda) = \lambda^{2}v(X+d) \quad v^{R}(X;\lambda) = \lambda^{2}v(X-d).$$
 [1b]

Below, we shall also allow for more general $\lambda-$ dependence of v^L and v^R .

We center the two wells on the x_1 -axis and impose that their supports do not overlap by taking:

$$d = \begin{bmatrix} d_1 \\ 0 \end{bmatrix}, \quad d_1 > a$$

If $\lambda \gg 1$, corresponding to the regime of deep wells and strong magnetic field, then under the constraints given above on v, H_{λ} has two eigenvalues in a small neighborhood of the single well ground state energy, e_{λ} : $E_{0,\lambda} \leq E_{1,\lambda}$. The two-dimensional eigenspace corresponding to this pair of eigenvalues is approximately spanned by the (nearly orthonormal) symmetric and antisymmetric linear combinations of the "atomic ground state orbitals", $\varphi^L(x)$ and $\varphi^R(x)$ arising separately from $v^L(x;\lambda)$ and $v^R(x;\lambda)$. The states $\varphi^L(x)$ and $\varphi^R(x)$ are obtained via magnetic translation of $\varphi(x)$ (7):

$$\varphi^L(x) = e^{i\frac{\lambda}{2}bd_1x_2}\varphi(x+d)$$
 and $\varphi^R(x) = e^{-i\frac{\lambda}{2}bd_1x_2}\varphi(x-d).$

We introduce the magnetic eigenvalue splitting:

$$\Delta_{\lambda} := E_{1,\lambda} - E_{0,\lambda} \tag{2}$$

and the magnetic hopping coefficient, an overlap integral of the single well ground state, $\varphi_{\lambda}(x)$, magnetically translated to x = -d and x = +d; (7):

$$\rho_{\lambda} \equiv \int \overline{\varphi^{L}(x;\lambda)} v^{R}(x;\lambda) \varphi^{R}(x;\lambda) dx.$$
 [3a]

In the most basic case [1b], we have

$$\rho_{\lambda} = \int_{x \in \mathbb{R}^2} \overline{\varphi(x+d)} \ \lambda^2 v(x-d) \ \varphi(x-d) \ e^{-i\lambda b d_1 x_2} \ \mathrm{d}x \,,$$
[3b]

Note that ρ_{λ} is in general complex-valued. However, ρ_{λ} is real if $v^{L}(x; \lambda) + v^{R}(x; \lambda)$ is inversion symmetric (even).

For large λ , a formal analysis suggests that the emergent quantity, ρ_{λ} , controls the eigenvalue splitting Δ_{λ} , and hence tunneling between two wells. In non-magnetic systems (b = 0) this has been proved. But as we shall see below, the magnetic case is far more subtle.

The hopping coefficient also emerges in the strong binding regime for an infinite crystal Hamiltonian, H_{crystal} , whose potential is a superposition of atomic wells, centered on a discrete set $\mathbb{G} \subset \mathbb{R}^2$, in the sense that

$$\left[J\left(H_{\text{crystal}} - e_{\lambda}\mathbb{1}\right)J^{*}\right]_{nm} \approx \rho_{\lambda} \ e^{i\frac{\rho_{\lambda}}{2}n \wedge m} \delta_{\|m-n\|, d_{0}}, \quad n, m \in \mathbb{C}.$$
[4]

. .

Here, d_0 , the minimal distance within \mathbb{G} , is strictly positive and the minimal next-nearest-neighbor distance is uniformly bounded away from d_0 . The operator J relates functions on the continuum with vectors whose amplitudes are indexed by \mathbb{G} ; see (8).

Previous work. Upper bounds for the eigenvalue splitting and the magnitude of the hopping coefficient are straightforward. Concerning previous work on lower bounds:

- Non-magnetic case (b = 0): Under the condition that vhas a single non-degenerate minimum, it was proved in (2, 3) that Δ_{λ} satisfies an exponential lower bound; the relationship between Δ_{λ} and ρ_{λ} , and an asymptotic expansion of ρ_{λ} was given in (3). In (9), a lower bound on ρ_{λ} was proved with no assumption on the minimum of v, but assuming v is of compact support.
- Perturbative magnetic case $(b \sim \frac{1}{\lambda})$: Lower bounds on $|\rho_{\lambda}|$ have been obtained under the assumption of a weak magnetic field, where $b \sim \frac{1}{\lambda}$ in (10). This regime is perturbative, relative to the non-magnetic case, since the magnetic field strength is of order 1 and the potential wells are deep, of order λ^2 .

The first result providing lower bounds for deep wells and strong magnetic fields was proved in (4):

Theorem 1: Magnetic hopping and magnetic splitting for radially symmetric atomic potentials. Assume that v is smooth and radial, such that φ_{λ} is also radial. Consider the hopping, ρ_{λ} and splitting Δ_{λ} associated with the magnetic double well Hamiltonian [1a-1b]. There exists $d_{\star} > 0$, c > 0 and $\lambda_{\star} > 0$ such that for all $d_1 > d_{\star}$:

$$-\rho_{\lambda} \ge e^{-c\lambda \|d\|^2}$$
 for $\lambda \ge \lambda_{\star}$, and
 $\lim_{\lambda \to \infty} \quad \frac{\Delta_{\lambda}}{2|\rho_{\lambda}|} \to 1.$

The articles (5, 6) sharpen Theorem 1 by providing leading order asymptotics of ρ_{λ} for $\lambda \to \infty$.

The radial symmetry assumption on v in Theorem 1 is very restrictive; we used it to rewrite the expression for ρ_{λ} as a real integral with non-oscillatory phase, to which we applied Laplace asymptotics for λ large.

The Non-Radial Case. For double well systems, where the single well is not radially symmetric, the situation is surprisingly different.

Theorem 2: (Magnetic double wells with zero tunneling) Fix a radial one well potential $v_0(x)$ satisfying the hypotheses of Theorem 1. Then there is a λ -dependent family \mathcal{F}_{λ} of (nonradial) one well potentials, such that the following hold for all large enough λ :

1. Potentials in \mathcal{F}_{λ} are exponentially close to $\lambda^2 v_0$: Let $\overline{v(\cdot;\lambda)} \in \mathcal{F}_{\lambda}$. Then $|\lambda^2 v_0(x) - v(x;\lambda)| < \exp(-C\lambda)$ for all $x \in \mathbb{R}^2$. Moreover, $v(x;\lambda) \neq \lambda^2 v_0(x)$ only for x in a set of area less than $\exp(-C\lambda)$. Here, C > 0 is determined by v_0 .




Fig. 1. Inversion-symmetric potential of the type used in the proof of Theorem 2. Large circles are the supports, B_0^L and B_0^R , of $v_0^L(x)$, centered at $x_0^L = (-d_1, 0)$, and $v_0^R(x)$, centered at $x_0^R = (+d_1, 0)$. For $\alpha = L, R$, we add four potentials ("sophons"), which are inversion symmetrically placed relative to x_0^α . The sophons are supported on the small red discs, B_{μ}^α , which are centered at x_{μ}^α ; $\mu = 1, 2, 3, 4$, the vertical components of which determine the phase θ in [8]; the amplitudes, support radii and centers of the sophon potentials are specially tuned λ – dependent functions.

2. Inversion symmetry: The two well potential

$$v^{L}(x;\lambda) + v^{R}(x;\lambda) = v(x+d;\lambda) + v(x-d;\lambda)$$

is inversion-symmetric for all $v \in \mathcal{F}_{\lambda}$.

- 3. Absence of quantum tunneling for H_{λ} : The family \mathcal{F}_{λ} contains a potential for which the eigenvalue splitting vanishes, i.e., $\Delta_{\lambda} = 0$. Hence, no tunneling occurs for such double wells. Moreover, \mathcal{F}_{λ} also contains a potential for which the hopping coefficient $\rho_{\lambda} = 0$.
- 4. <u>Odd ground state</u>: By deforming within the family \mathcal{F}_{λ} , the two well ground state eigenfunction of H_{λ} , ψ , can be made to transition between symmetric ($\psi(x) = \psi(-x)$) and antisymmetric ($\psi(x) = -\psi(-x)$).

Figure 1 is a schematic showing the support of a potential in \mathcal{F}_{λ} . Such a potential is a perturbation of the double well potential of Theorem 1, which is comprised of two radially symmetric single wells. A discussion of the proof, with the aid of Figure 1, is given below.

We do not have an example of a non-inversion symmetric potential for which $\Delta_{\lambda} = 0$. However, we have the following result.

Theorem 3: There exist non-inversion symmetric magnetic double well potentials for which $\rho_{\lambda} = 0$.

The potentials $v(\cdot; \lambda)$ realizing zero tunneling or zero hopping coefficient in Theorems 2 and 3 do not arise by simply scaling a λ -independent potential. We conjecture that a potential v_0 , nonradial and independent of λ , can be chosen so that the double well Hamiltonian arising from $\lambda^2 v_0$ gives rise to zero eigenvalue splitting for an infinite sequence of λ values tending to infinity, and similarly for the hopping coefficient.

We next claim that, in a precise sense, the counterexamples of Theorems 2 and 3 are exceptional in a regime where b (the magnetic strength parameter in [1a]) is small but fixed as $\lambda \to \infty$. And further, in that regime, for typical large λ , there is a quantitative lower bound on both $|\rho_{\lambda}|$ and Δ_{λ} . Recall that a set $S \subseteq (0, \infty)$ has zero density if $\mu(S \cap [0, M])/M \to 0$ as $M \to \infty$, where μ denotes Lebesgue measure.

Theorem 4 : Fix $\gamma \in (0, 1)$. Suppose the one well potential v(x) is C^3 and has a unique minimum at the origin, with its Hessian there obeying $v''(0) = \mathbb{1}_{2\times 2}$. Consider the magnetic double well operator [1a-1b]. Suppose $|b| < c_{\gamma}$ for some small constant c_{γ} . Then for all λ outside a set $\mathcal{Z}(\gamma, v) \subseteq (0, \infty)$ of density zero, we have

$$|\rho_{\lambda}| > \exp(-\lambda^{\frac{1}{\gamma}})$$
 and $\Delta_{\lambda} > \exp(-\lambda^{\frac{1}{\gamma}})$. [5]

Note that here, although b is assumed small, it remains independent of λ , which may be taken arbitrarily large. Recall that our two well potential is $\lambda^2(v(x+d) + v(x-d))$ and the magnetic field strength is $b\lambda$. Finally, we expect the condition $v''(0) = \mathbb{1}_{2\times 2}$ can be relaxed.

Flat band materials. The ideas of Theorems 2 and 3 can be applied to the design of periodic structures, which have nearly flat band spectrum in a neighborhood of the one well ground state energy. We explain this in a section below.

Ideas underlying the proofs. Let us perturb a radial one-well potential v_0 by adding a small correction δv . Then the hopping coefficient ρ will change by an amount $\delta \rho$. Due to the strong magnetic field ($\lambda \gg 1$), $\delta \rho$ is given by an oscillatory integral. However, if the support of δv has sufficiently small diameter (depending on λ), then the expected cancellation due to the rapidly oscillating exponential in the relevant integrand will not occur. Moreover, due to the rapid (Gaussian) decay of the ground state for v_0 , the effect of δv on $\delta \rho$ is enhanced if the support of δv is favorably placed. For instance, in Figure 1, two of the small discs on the left are closer than the large disc on the left to everything on the right. Consequently, although δv is small compared to v_0 , it may happen that $\delta \rho$ is large compared to ρ . We find that we can arrange $\delta \rho$ to have greater, smaller, or the same order of magnitude as ρ , as we please. This intuition plays a role in the proofs of Theorems 2 and 3. Here, we focus on the proof of Theorem 2.

Sketch of proof of Theorem 2

1

We refer to Figure 1. We denote the large disc (a "planet") on the left by B_0^L , centered at $x_0^L = (-d_1, 0)$ and the four small discs on the left by B_{μ}^L with $\mu = 1, \ldots, 4$; ("sophons" (11)); similarly for the right with $L \to R$ and B_0^R centered at $x_0^R = (d_1, 0)$. The centers of these discs are denoted as x_{μ}^{α} for $\alpha = L, R$; $\mu = 0, \ldots, 4$. The centers, x_0^L and x_0^R are independent of λ . The centers of the small discs, x_{μ}^{α} for $\alpha = L, R$; $\mu = 1, \ldots, 4$, and the radii of $B_{\mu}^{\alpha}, \mu = 1, 2, 3, 4$ and $\alpha = L, R$ are chosen to depend on λ . The radii of these discs are chosen to shrink exponentially fast as $\lambda \to \infty$.

Our "two-well" potential is $v^L(x;\lambda) + v^R(x;\lambda)$, where

$$v^{\alpha}(x;\lambda) = \lambda^2 v_0^{\alpha}(x) + \sum_{\mu=1}^{4} v_{\mu}^{\alpha}(x;\lambda), \quad \alpha = L, R.$$

Here, (a) $v_0^{\alpha}(x)$ is independent of λ , and supported on B_0^{α} (the planet), and (b) for $\mu = 1, \ldots, 4$, $v_{\mu}^{\alpha}(\cdot; \lambda)$ is supported on B_{μ}^{α} (a sophon), and $\|v_{\mu}^{\alpha}\|_{\infty} \sim \tau$ with $\tau = \tau(\lambda)$ a carefully chosen constant which is exponentially small in λ .

For $\alpha = L, R$, we write φ^{α} to denote the ground state of the one-well magnetic Hamiltonian $h^{\alpha} = \left(P + \frac{1}{2}\lambda b X^{\perp}\right)^2 + v^{\alpha}(X;\lambda)$. We study the magnetic hopping coefficient ρ_{λ} , defined in [3a]. Due to inversion symmetry, ρ_{λ} is real. Since v^R is supported on the five disjoint discs B^R_{ν} , we have

$$\rho = \sum_{\nu=0}^4 \int_{B^R_\nu} \overline{\varphi^L} v^R \varphi^R$$

Carefully justified perturbation theory lets us write

$$\varphi^L = \varphi_0^L + \sum_{\mu=1}^4 \varphi_\mu^L$$

modulo a negligible error. Here, φ_0^L denotes the magnetic ground state arising from $\lambda^2 v_0^L(x)$ while, for $\mu = 1, 2, 3, 4$, the function φ_{μ}^L denotes the correction to φ_0^L arising from the added potential $v_{\mu}^L(x; \lambda)$ in first order perturbation theory. Hence, modulo a negligible error, we find

$$\rho_{\lambda} = \sum_{\mu=0}^{4} \left[\lambda^2 \int_{B_0^R} \overline{\varphi_{\mu}^L} \, v_0^R \, \varphi^R + \sum_{\nu=1}^{4} \int_{B_{\nu}^R} \overline{\varphi_{\mu}^L} \, v_{\nu}^R \, \varphi^R \right] \qquad [6]$$
$$=: \sum_{\mu,\nu=0}^{4} \mathrm{T}(\mu,\nu) \, .$$

Each term, $T(\mu, \nu)$, may be thought of as a contribution to the tunneling coming from an interaction between planets and sophons on the left, L, and right, R. In particular, T(0,0)corresponds to an interaction between planets; it is the hopping coefficient for the unperturbed radial one well potential, $\lambda^2 v_0(x)$, to which Theorem 1 applies. The terms $T(\mu, \nu)$, with $\mu + \nu \geq 1$, are corrections due to the introduction of sophons.

Now eigenstates φ of the one well Hamiltonian satisfy $\left((P + \frac{1}{2}\lambda bX^{\perp})^2 - e_{\lambda}\right)\varphi = 0$ outside of the support of v, and hence have Gaussian decay. Using this, we find that as $\lambda \to \infty$

$$|T(0,0)| \sim \exp\left(-\frac{\lambda}{4} ||x_0^L - x_0^R||^2 (1+o(1))\right)$$
 [7a]

$$|\mathrm{T}(\mu,\nu)| \sim \tau \exp\left(-\frac{\lambda}{4} ||x_{\mu}^{L} - x_{\nu}^{R}||^{2} (1+o(1))\right),$$
 [7b]

where
$$\mu = 0$$
 and $\nu \in \{1, 2, 3, 4\}$
or $\nu = 0$ and $\mu \in \{1, 2, 3, 4\}$
 $|T(\mu, \nu)| \sim \tau^2 \exp\left(-\frac{\lambda}{4} \|x_{\mu}^L - x_{\nu}^R\|^2 (1 + o(1))\right),$ [7c]

where $\mu, \nu \in \{1, 2, 3, 4\}.$

Here we have used the fact that the diameter of the supports of $v_{\mu}^{L}(x)$ and $v_{\mu}^{R}(x)$ shrinks to zero for $\mu = 1, 2, 3, 4$ at an exponential rate as $\lambda \to \infty$. Due to the effects discussed above in the section: *Ideas underlying the proofs*, we may choose $\tau =$ $\tau(\lambda)$ to be exponentially small in λ , and such that $T(0, \mu)$ and $T(\mu, 0)$ ($\mu = 1, 2, 3, 4$), which are formally of order τ , dominate T(0, 0), the unperturbed radial hopping coefficient. Further, one can derive from the expression for ρ_{λ} in [3b] that the vertical displacements of the sophons (see Figure 1) introduce complex phases into $T(\mu, \nu)$ for $(\mu, \nu) \neq (0, 0)$. The relative phase, θ , of the dominant terms depends sensitively on the dimensions of the rectangle formed by $x_{1}^{R}, \ldots, x_{4}^{R}$. Accordingly, [7] implies that

$$\rho_{\lambda} = \tau \, \exp\left(-\frac{\lambda}{4}D^2 \left(1 + o(1)\right)\right) \left[\cos(\theta) + o(1)\right], \quad \text{for } \lambda \gg 1,$$
[8]

where $D := \min_{1 \le \mu \le 4} ||x_{\mu}^{L} - x_{0}^{R}||$. Here, the cosine appears in [8] due to inversion symmetry, which dictates that ρ is real. From [8] we see that by slightly varying the placement of the sophons we can arrange for ρ_{λ} to be either positive or negative.

Let us now consider the spectral gap Δ_{λ} . Since the twowell potential $v^L + v^R$ is even, the magnetic Hamiltonian $H_{\lambda} = \left(P + \frac{1}{2}\lambda bX^{\perp}\right)^2 + v^L + v^R$ commutes with the parity operator $f(x) \mapsto (\mathcal{P}f)(x) = f(-x)$, and hence preserves the subspaces L^2_{even} , L^2_{odd} consisting of even wavefunctions $\psi(x) =$ $\psi(-x)$ and odd wave functions $\psi(x) = -\psi(-x)$, respectively. Let $E_{\text{even},\lambda}$ and $E_{\text{odd},\lambda}$ denote the ground state eigenvalues of H_{λ} restricted to L^2_{even} and L^2_{odd} , respectively. Then, for the lowest two eigenvalues of H_{λ} , we have $\{E_{0,\lambda}, E_{1,\lambda}\} = \{E_{\text{even},\lambda}, E_{\text{odd},\lambda}\}$. Hence, by [2], the spectral gap of H is then

$$\Delta_{\lambda} = \left| E_{\text{even},\lambda} - E_{\text{odd},\lambda} \right|.$$

Moreover, modulo a small error, $E_{\text{even},\lambda} - E_{\text{odd},\lambda} = 2\rho_{\lambda}$. By deforming our potential (by moving the discs B^{α}_{μ} , $\mu = 1, 2, 3, 4$, $\alpha = R, L$) we can, by [8] and the ensuing discussion, continuously vary $E_{\text{even},\lambda} - E_{\text{odd},\lambda}$ from a positive value to a negative value. Consequently,

- At some point during the deformation, we have $E_{\text{even},\lambda} = E_{\text{odd},\lambda}$, and hence $\Delta_{\lambda} = 0$.
- We may take $E_{\text{even},\lambda} E_{\text{odd},\lambda}$ to be positive, i.e., the ground state of the magnetic double well may be odd rather than even.

The class \mathcal{F}_{λ} in Theorem 2 consists of all one-well potentials arising as above, for all possible placements of the four "sophons", B_{μ}^{L} ($\mu = 1, 2, 3, 4$) in a rectangle about the "planet" B_{0}^{L} , as in Figure 1. This completes our rough sketch of the proof of Theorem 2.

Sketch of proof of Theorem 4:. We study the ground state φ_{λ} of the one well Hamiltonian h_{λ} , and the orthogonal projection Π_{λ} from $L^2(\mathbb{R}^2)$ to the two lowest eigenstates of the two well Hamiltonian H_{λ} . We show that the L^2 -valued function $\lambda \mapsto \varphi_{\lambda}$ and the operator-valued function $\lambda \mapsto \Pi_{\lambda}$ can be continued to analytic functions on the region

$$\Gamma_{\gamma} := \{ \Re \lambda > C_1, \, |\arg(\lambda)| < \gamma \, \pi/2 \}, \qquad [9]$$

where $C_1 < \infty$ is a large fixed constant and $0 < \gamma < 1$, provided $|b| < c_{\gamma}$, a sufficiently small and positive constant independent of λ . From this we conclude that $\lambda \mapsto \rho_{\lambda}$ and $\lambda \mapsto (\Delta_{\lambda})^2$ also continue to analytic functions on Γ_{γ} . We first prove Theorem 4 assuming the analyticity of $\lambda \mapsto \rho_{\lambda}$ and $\lambda \mapsto (\Delta_{\lambda})^2$. Then, in a section below we outline the proof of analyticity.

The strategy for deducing Theorem 4 from analyticity is to combine (i) upper bounds on the above analytic functions in Γ_{γ} with, (ii) a lower bound on these functions at a single point in Γ_{γ} and then apply the complex analysis theorem below to prove an averaged lower bound on on $|\rho_{\lambda}|$ and Δ_{λ} . This average lower bound implies that the lower bounds in [5] hold except for a set of density zero.

The upper bounds (i) follow from our analysis of $\lambda \mapsto \varphi_{\lambda}$ and $\lambda \mapsto \Pi_{\lambda}$ for complex $\lambda \in \Gamma_{\gamma}$, promised above. To explain our derivation of the lower bound (ii), we now write $\Delta_{\lambda} \equiv \Delta(\lambda, b)$ to denote the spectral gap arising from the Hamiltonian H_{λ} for a given λ and b. First set b = 0. Then, $\Delta(\lambda, 0)$ is the spectral gap for the non-magnetic Hamiltonian. By classical results for the non-magnetic Hamiltonian (see, for example, (2, 3, 9)), for fixed real $\lambda_0 > C_1$ (see [9]), we have $\Delta(\lambda_0, 0) > \delta_1$ for some positive constant δ_1 . Consequently, for |b| small enough - say, $|b| < b_0(\lambda_0) -$ we have $\Delta(\lambda_0, b) > \frac{1}{2}\delta_1$. This is the required lower bound (ii) for $|\Delta(\lambda, b)|^2$ for a single value of λ and small b.

To deduce Theorem 4 from the above bounds (i) and (ii), we apply the following result to $z \in \Lambda_{\gamma} \mapsto \rho_z$ and $z \in \Lambda_{\gamma} \mapsto \Delta(z, b)^2$, for $|b| < b_0(\lambda_0)$: A theorem in complex analysis Let Γ_{γ} be the region defined in [9], for $\gamma \in (0, 1]$. Assume that $F : \Gamma_{\gamma} \to \mathbb{C}$ is analytic and such that there exists $\beta > 0$ for which

$$|F(\lambda_0)| \ge e^{-\beta}, \qquad [10]$$

for some real $\lambda_0 > C_1$. Further assume that there exists a non-vanishing analytic function $U: \Gamma_{\gamma} \to \mathbb{C}$ such that for all $z \in \Gamma_{\gamma}$,

$$|F(z)| \le |U(z)|. \tag{11}$$

There exists a constant C (depending on β , γ , λ_0 and U) such that for all R sufficiently large,

$$\frac{1}{R} \int_{t=R}^{2R} -\log\left(|F(t)|\right) \mathrm{d}\, t \le CR^{1/\gamma} +$$

$$+ \frac{1}{R} \int_{t=R}^{2R} -\log\left(|U(t)|\right) \mathrm{d}\, t \,.$$
[13]

The above theorem is an application of the standard Blaschke factorization of analytic functions on the disc, see Theorem 15.21 of (12).

The proof of analytic continuation: Making the above analytic continuations is not straightforward, since for typical $\mu \in \mathbb{C}$ the resolvents $(h_{\lambda} - \mu \mathbb{1})^{-1}$ and $(H_{\lambda} - \mu \mathbb{1})^{-1}$ do <u>not</u> admit analytic continuations as functions of λ taking values in the space of bounded operators on L^2 . However, when applied to functions supported on $B_a(0)$, these resolvents do admit analytic continuation to $\lambda \in \Gamma_{\gamma}$. Once this is proved, it is then not hard to control the above ground state φ_{λ} and projection Π_{λ} as functions of complex $\lambda \in \Gamma_{\gamma}$, where $0 < \gamma < 1$.

To establish the analyticity of $\lambda \mapsto (h_{\lambda} - \mu \mathbb{1})^{-1} \chi$, where $\operatorname{supp}(\chi) \subset B_a(0)$, we fix $\Lambda \gg 1$. For $\lambda \in \Gamma_{\gamma}$, with $\frac{1}{2}\Lambda \leq |\lambda| \leq 2\Lambda$, we localize χ into the regions

$$U_{\text{inner}} = \left\{ |x| \le \Lambda^{-\frac{1}{2} + \varepsilon} \right\} \text{ and } U_k = \left\{ 2^{-(k+1)} \le |x| \le 2^{-(k-1)} \right\},$$

for $|\Lambda|^{-\frac{1}{2}+\varepsilon} \leq 2^{-k} \leq a$. In the region U_{inner} , h_{λ} is closely approximated by a magnetic harmonic oscillator Hamiltonian. Consequently, for $\operatorname{supp}(\chi) \subset U_{\text{inner}}$, we can control $(h_{\lambda} - \mu \mathbb{1})^{-1}\chi$ in terms of the resolvent of the magnetic harmonic oscillator (MHO):

$$H_{\lambda}^{\mathrm{MHO}} = (-\mathrm{i}\nabla + \frac{1}{2}\lambda bX^{\perp})^2 + \frac{1}{2}\lambda^2\omega_0^2X^2.$$

The resolvent kernel of H_{λ}^{MHO} can be expressed via the Laplace transform of the (magnetic) Mehler (heat) kernel exp $\left(-tH_{\lambda}^{\text{MHO}}\right)(x,y)$. This kernel has poles for $\lambda t \in i\pi\mathbb{Z}$ (13, Prop. 2.1), hence we cannot hope to obtain an analytic extension beyond the open right half plane.

To control $(h_{\lambda} - \mu \mathbb{1})^{-1}$ localized in U_k , we cover U_k by finitely many discs $B(x_*, \delta)$, with δ = small constant $\times 2^{-k}$. On each $B(x_*, \delta)$, we rescale via the change of variables $x = x_* + \delta y$. Then, $B(x_*, \delta)$ goes over to the unit disc B(0, 1), and $h_{\lambda} - \mu \mathbb{1}$ becomes a constant multiple of a second order differential operator:

$$L = -\Delta_y + \vec{\gamma}(y) \cdot \nabla_y + \tilde{\gamma}(y),$$




Fig. 2. Single-well potential for flat band crystal.

whose symbol $\sigma(L)(y,\eta) = |\eta|^2 + i\vec{\gamma}(y) \cdot \eta + \tilde{\gamma}(y)$ satisfies, for all $y \in B(0,1), \eta \in \mathbb{R}^2$ and all α, β , the following estimates:

$$\begin{split} \left| \partial_y^{\alpha} \partial_{\eta}^{\beta} \left[\sigma(L)(y,\eta) \right] \right| &\leq C_{\alpha\beta} \left[\Lambda^2 \delta^4 + |\eta|^2 \right]^{1 - \frac{|\beta|}{2}} \\ \left| \sigma(L)(y,\eta) \right| &\geq c \left[\Lambda^2 \delta^4 + |\eta|^2 \right]. \end{split}$$

Second order differential operators satisfying the above conditions behave like $-\Delta + (\Lambda^2 \delta^4) \mathbf{1}$ and their resolvents may be controlled accordingly; in particular, their resolvents may be analytically continued. To make this precise, it is convenient to use pseudodifferential operator calculus (see, e.g. (14)). This concludes our discussion of the case of functions supported in U_k .

Combining our results on U_{inner} and U_k , we establish the desired analytic continuation of $\lambda \mapsto (h_\lambda - \mu \mathbb{1})^{-1}\chi$ for $\operatorname{supp}(\chi) \subset B_a(0), \ \lambda \in \Gamma_{\gamma}$, and $\frac{1}{2}\Lambda \leq |\lambda| \leq 2\Lambda$. Finally, combining our results for $\Lambda = 2^n$ as n varies, we control $\lambda \mapsto (h_\lambda - \mu \mathbb{1})^{-1}\chi$ for $\lambda \in \Gamma_{\gamma}$, where $\operatorname{supp}(\chi) \subset B_a(0)$. Our outline of the proof of analyticity of $(h_\lambda - \mu \mathbb{1})^{-1}\chi$ is now complete. Our analysis of the double-well resolvent $(H_\lambda - \mu \mathbb{1})^{-1}\chi$ is similar.

On the construction of periodic crystals with flat bands. Consider a single planet surrounded by eight sophons as in Figure 2. Now form a periodic arrangement by placing central planets at the vertices of $D'\mathbb{Z}^2$, where D' > 0 is fixed and taken sufficiently large. By carefully adjusting the position of the sophons, we can again make the hopping coefficient vanish between any two nearest neighbors of the lattice $D'\mathbb{Z}^2$. Thanks to [4] this leads to a crystal structure with nearly flat band spectrum centered near e_{λ} . Further, as one varies within the family \mathcal{F}_{λ} , the structure can be tuned, between one that supports a flat band and one that does not. This may lead to a novel tunable platform for exploring flat band physics.

Open questions and conjectures.

- 1. Is ρ_{λ} analytic in a half-plane $\Re \lambda \geq \lambda_0 > 0$? Do the bounds of Theorem 4 hold with $\gamma = 1$ for typical large real λ ?
- 2. Are there double well potentials v, independent of λ , such that the Hamiltonian $(P + \frac{1}{2}\lambda bX^{\perp})^2 + \lambda^2 v(X)$ gives rise to an infinite sequence of λ 's, tending to infinity, for which the magnetic eigenvalue splitting vanishes, and another infinite sequence of λ 's for which the hopping coefficient vanishes?
- 3. Can the transition from a symmetric to antisymmetric ground state be observed experimentally? Or the flat band crystal, by the mechanism described?

4. Are there analogous phenomena in three dimensions? If the magnetic field is perpendicular to the displacement between the wells, then we suspect that producing counterexamples in the 3D case is actually easier than in the 2D case. However, if the magnetic field is not perpendicular to the displacement, new questions may arise.

Acknowledgements. MIW was supported in part by NSF grant DMS-1908657, DMS-1937254 and Simons Foundation Math + X Investigator Award # 376319 (MIW). Part of this research was carried out during the 2023-24 academic year, when MIW was a Visiting Member in the School of Mathematics - Institute of Advanced Study, Princeton, supported by the Charles Simonyi Endowment, and a Visiting Fellow in the Department of Mathematics at Princeton University. The authors wish to thank Antonio Cordoba and David Huse for stimulating discussions. We thank P.A. Deift and J. Lu for their careful reading of the paper, and insightful comments and questions.

- LD Landau, E Lifshitz, <u>Quantum Mechanics Non-Relativistic Theory, Third Edition: Volume 3</u> (Course of Theoretical Physics) (Vol. 3). (Butterworth-Heinemann), (1981).
- B Simon, Semiclassical analysis of low lying eigenvalues, ii. tunneling. <u>Annals Math.</u> 120, 89–118 (1984).
- B Helffer, J Sjostrand, Multiple wells in the semi-classical limit i. <u>Commun. Partial. Differ. Equations</u> 9, 337–408 (1984).
- C Fefferman, J Shapiro, MI Weinstein, Lower bound on quantum tunneling for strong magnetic fields. SIAM J. on Math. Analysis 54, 1105–1130 (2022).
- B Helffer, A Kachmar, Quantum tunneling in deep potential wells and strong magnetic field revisited (2023).
- 6. L Morin, Tunneling effect between radial electric wells in a homogeneous magnetic field (2023).
- 7. J Zak, Magnetic translation group. Phys. Rev. 134, A1602–A1606 (1964).
- J Shapiro, MI Weinstein, Tight-binding reduction and topological equivalence in strong magnetic fields. <u>Adv. Math.</u> 403, 108343 (2022).
- CL Fefferman, JP Lee-Thorp, MI Weinstein, Honeycomb Schrödinger operators in the strong binding regime. <u>Commun. on Pure Appl. Math.</u> **71**, 1178–1270 (2018).
 B Helffer, J Sjöstrand, Effet tunnel pour l'équation de schrödinger avec champ magnétique.
- Annali della Scuola Norm. Super. di Pisa Cl. di Scienze **4e série, 14**, 625–657 (1987).
- C Liu, K Liu, <u>The Three-Body Problem</u>, The Three-Body Problem Series. (Tor Publishing Group), (2014).
- 12. W Rudin, Real and complex analysis. (McGraw-Hill, New York), (1987).
- H Matsumoto, Semiclassical asymptotics of eigenvalues for schrödinger operators with magnetic fields. <u>J. Funct. Analysis</u> **129**, 168 – 190 (1995).
- L H
 örmander, <u>Pseudo-Differential Operators</u>. (Springer Berlin Heidelberg, Berlin, Heidelberg), pp. 63–179 (2007).