

Vector-like quark doublets, weak-basis invariants and CP violation

F. Albergaria^{a,*}, J. F. Bastos^{a,†}, B. Belfatto^{b,‡}, G. C. Branco^{a,§},
J. T. Penedo^{c,¶}, A. Rodríguez-Sánchez^{d,||}, J. I. Silva-Marcos^{a,**}

^a *CFTP, Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa,
Avenida Rovisco Pais 1, 1049-001 Lisboa, Portugal*

^b *Institut für Theoretische Teilchenphysik, Karlsruhe Institute of Technology,
Engesserstraße 7, D-76128 Karlsruhe, Germany*

^c *INFN, Sezione di Roma Tre, Via della Vasca Navale 84, 00146, Rome, Italy*

^d *Departament de Física Teòrica, IFIC, Universitat de València - CSIC*

Abstract

We study Standard Model extensions with isodoublet vector-like quarks with standard charges. Their presence induces right-handed charged and neutral currents. We identify minimal sets of independent parameters characterizing these extensions, describe useful weak bases, and provide parameterizations for all quark mixing. We analyze the intricacies of CP violation in such scenarios, finding a complete set of CP-odd invariants for the single doublet case. Crucially, we uncover a connection between weak-basis invariants and effective rephasing invariants involving only standard quarks. These results allow us to explore the phenomenology of doublet vector-like quarks through a rephasing-invariant analysis, with an emphasis on CP violation, including the potential role of these fields in explaining the Cabibbo angle anomalies.

*francisco.albergaria@tecnico.ulisboa.pt

†jose.bastos@tecnico.ulisboa.pt

‡benedetta.belfatto@kit.edu

§gbranco@tecnico.ulisboa.pt

¶jpenedo@roma3.infn.it

||anrosanz@ific.uv.es

**juca@cftp.tecnico.ulisboa.pt

Contents

1	Introduction	5
2	Setup	7
2.1	Gauge and Higgs interactions	9
3	Weak bases and physical parameters	12
3.1	Convenient weak bases	13
3.1.1	A minimal weak basis	13
3.1.2	A “stepladder” weak basis ($N = 1$)	15
3.2	Alternate ways of counting physical parameters	17
3.2.1	Counting via a spurion analysis	17
3.2.2	Counting in the mass basis ($N = 1$)	18
3.3	Rephasing invariants	18
3.3.1	The general case ($N \geq 1$)	19
3.3.2	A single doublet ($N = 1$)	20
4	Rotating to the mass basis	22
4.1	Approximate masses and mixing ($N = 1$)	22
4.1.1	The RH sector	24
4.1.2	The LH sector	26
4.1.3	Effective rephasing invariants	28
4.2	Parameterizing the mixing exactly	30
4.2.1	A single doublet ($N = 1$)	30
4.2.2	More than one doublet ($N > 1$)	32
5	Weak-basis invariants	36
5.1	Constructing WBIs	36
5.2	Reconstructing minimal parameterizations from WBIs	39
5.2.1	Reconstructing the stepladder WB	40
5.2.2	Reconstructing the minimal WB	43
5.2.3	A numerical example	46
6	CP violation	48
6.1	CP-odd weak-basis invariants	48
6.1.1	One or more vector-like doublets	48
6.1.2	Other CP-odd WBIs with more than one vector-like doublet	51
6.2	Effective CP-odd WBIs	53
6.3	Conditions for CP invariance	56
6.4	Motivated scenarios	60
6.4.1	With $N = 1$ doublet	60
6.4.2	With $N = 2$ doublets	61
6.4.3	The extreme chiral limit with $N = 1$ doublet	64

6.4.4	The extreme chiral limit with $N = 2$ doublets	65
7	Integrating out the VLQs	66
7.1	SMEFT description	66
7.2	Electroweak symmetry breaking	67
7.3	Low-energy EFT	69
8	Phenomenological effects of mixing with VLQ doublets	71
8.1	Weak currents	71
8.2	Phenomenology of effective rephasing invariants	72
8.3	Bilinears and the Cabibbo sector	74
8.3.1	Cabibbo angle anomalies	74
8.3.2	The neutral kaon system	76
8.3.3	Electric dipole moments	85
8.3.4	Low-energy EW flavour-conserving observables and Z decay	86
8.4	Trilinears and flavour changing neutral currents	87
8.4.1	Kaon mixing	87
8.4.2	$K_L \rightarrow \pi^0 \nu \nu$, $K_{S,L} \rightarrow \mu \mu$ and other rare kaon decays	90
8.4.3	CP violation in $K \rightarrow \pi \pi$ by trilinears	93
8.5	CAA-motivated scenarios	95
8.5.1	One VLQ doublet ($N = 1$)	95
8.5.2	Two VLQ doublets ($N = 2$)	97
9	Summary and conclusions	102
A	Notation	104
B	Effective CKM angles vs. Lagrangian parameters	106
C	Conditions for CP Invariance	108
C.1	Case-by-case analysis	110

List of symbols

BAU	baryon asymmetry of the Universe
BSM	beyond-the-Standard-Model
CAA	Cabibbo angle anomaly
CKM	Cabibbo–Kobayashi–Maskawa
CPI	CP invariance
CPV	CP violation
ECL	extreme chiral limit
EFT	effective field theory
EW	electroweak
EWSB	electroweak symmetry breaking
FCNC	flavour-changing neutral current
GIM	Glashow–Iliopoulos–Maiani
LH	left-handed
NP	New Physics
RH	right-handed
SM	Standard Model
SMEFT	Standard Model effective field theory
UV	ultraviolet
VEV	vacuum expectation value
VLQ	vector-like quark
WB	weak basis
WBI	weak-basis invariant
WBT	weak-basis transformation

1 Introduction

Vector-like quarks (VLQs) are one of the simplest and best-motivated extensions of the Standard Model (SM), which may play an important role in solving some of the open questions in particle physics. They appear in models of grand unification [1–5], extra dimensions [6], and emerge in models addressing the electroweak hierarchy problem in which the light Higgs arises as a pseudo-Goldstone boson of a global symmetry [7–11]. They play a fundamental part in models with inter-family symmetries which explain the origin of fermion mass hierarchies and mixings [12–18], and provide a natural realization of the minimal flavour violation scenario [19–21]. They can also be relevant to axion models [22, 23] as well as axion-less models of the Nelson-Barr type solving the strong CP problem [24–28].

These extensions also present a very rich phenomenology that can affect many important flavour observables, including neutral meson mixing parameters, meson decay amplitudes, and electroweak precision measurements (see e.g. [29–45]). Moreover, since they introduce extra — and in general complex — couplings in the Yukawa Lagrangian, new sources of CP violation (CPV) arise which may be crucial in explaining the baryon asymmetry of the Universe (BAU), given that CPV within the SM is too suppressed to generate a large enough BAU, consistent with observations.

Most of the analyses of VLQs have been done in the framework of VLQs which are isosinglets of weak isospin $SU(2)_L$ (see [46] for a recent review). However, VLQs were also studied in higher dimensional representations of $SU(2)_L$ [32, 47, 48]. If the scalar sector of a model only includes $SU(2)_L$ doublets, VLQs can only appear in seven types of $SU(2)_L$ multiplets: singlets with $Y = -1/3$ or $Y = 2/3$; doublets with $Y = -5/6$, $Y = 1/6$ or $Y = 7/6$; or triplets with $Y = -1/3$ or $Y = 2/3$. The next-to-minimal realization of VLQs corresponds to doublets of $SU(2)_L$.

In this paper, we are specifically interested in studying models of VLQ isodoublets with non-exotic charges, i.e. VLQs with $Y = 1/6$, similarly to the quark doublets present in the SM. These appear to be capable of addressing several current anomalies. Most notably, it was shown that VLQ doublets mixing with the light generations of SM quarks are favoured candidates in explaining the Cabibbo angle anomalies (CAAs) [45, 49–51]. Indeed, models of VLQ doublets i) induce right-handed currents, which can potentially accommodate all CAA tensions, and ii) are in agreement with electroweak precision observables, e.g. the W -boson mass. Note that, unlike in the case of singlet extensions, there are no significant deviations from Cabibbo–Kobayashi–Maskawa (CKM) unitarity at leading order in these models.

We study these new-physics effects and sources of CP violation using a weak-basis invariant (WBI) description. The usefulness of WBI quantities lies in the fact that they remain unmodified under weak-basis changes. Thus, WBI quantities can be used to identify the physical information contained in any arbitrary parameterization of the Yukawa matrices. This allows for an unambiguous connection to physical observables such as quark masses and mixings [52–60]. The existence of complex WBIs signals the presence of physical phases and CP violation. The imaginary parts of such invariants —

the so-called CP-odd WBIs — are one of the focuses of this work.

In the SM, the only independent CP-odd WBI is of mass dimension $M = 12$ [61] and is proportional to CKM quartets $V_{\alpha i} V_{\beta j} V_{\alpha j}^* V_{\beta i}^*$, resulting in a large suppression of CP violation effects. In the extensively-studied isosinglet VLQ case, the potential for enhancement is well-known, given the presence of a larger number of independent CP-odd WBIs, including some of dimension $M = 8$ [56, 58, 60]. However, as we emphasize in this work, the doublet representation can allow for lower-dimension WBIs, starting from $M = 6$ invariants, which suggests an even greater enhancement.¹ This is directly related to the fact that in the doublet scenario new kinds of rephasing invariants arise from the interplay between the two types of charged currents. In particular, bilinears of the form $V_{\alpha i}^R V_{\alpha i}^{L*}$ are now allowed, where $V_{L(R)}$ represents the mixing matrix for the left(right)-handed charged currents. In this way, the number of insertions of the mixing elements needed to construct CP-odd WBIs can be brought to a minimum, potentially allowing for strong CPV effects.

This paper is divided into the following sections, each of which is as self-contained as possible. In section 2, we set up the framework and notation (see also Appendix A) and display the gauge and Higgs interactions of the model. In section 3, we discuss various convenient weak bases, count the physical parameters and present the most relevant rephasing invariants. In section 4, we present approximate expressions for the quark masses and the mixings as a function of the Lagrangian parameters, in the realistic limit in which the VLQs mass scale is at least a few times larger than the electroweak scale. In this section, we also derive an exact parameterization for the quark mixing matrices for any number N of VLQ doublets. In section 5, we explain how to construct weak-basis invariants (WBIs) and how they are related to physical parameters. We also identify a set of WBIs totally characterizing the single VLQ doublet scenario ($N = 1$), and demonstrate how such a characterization can be generalized to an arbitrary number of doublets ($N > 1$). In section 6, we delve into the topic of CP violation, presenting CP-odd WBIs, their connection to effective rephasing invariants, and conditions for CP conservation with one VLQ doublet. In section 7, we present an effective field theory (EFT) description of models with $Y = 1/6$ VLQ doublets. In section 8, we analyze some of the phenomenology of these models using a WBI description, particularly in relation to the CAAs and CPV effects. Finally, we summarize our results in section 9.

¹It has been pointed out that models with isodoublet VLQs are uniquely capable of generating significant CP violation in the coupling of third-generation quarks to the Higgs boson, whereas for all other VLQ realizations CP violation is suppressed in these couplings [62].

2 Setup

We extend the SM by adding N vector-like quark (VLQ) isodoublets $Q_{L,R} = (U, D)_{L,R}$ in the same representation of $SU(3)_c \times SU(2)_L \times U(1)_Y$ as standard left-handed (LH) quarks, i.e. with the SM quantum numbers $(\mathbf{3}, \mathbf{2})_{1/6}$. Thus, U and D have electric charges $+2/3$ and $-1/3$, respectively. We define

$$Q_{L\alpha}^0 = \begin{pmatrix} U_{L\alpha}^0 \\ D_{L\alpha}^0 \end{pmatrix}, \quad Q_{R\alpha}^0 = \begin{pmatrix} U_{R\alpha}^0 \\ D_{R\alpha}^0 \end{pmatrix} \quad (\alpha = 1, \dots, N). \quad (2.1)$$

The ‘0’ superscript indicates that fields are in a flavour basis, as opposed to the mass basis. SM quark doublets and singlets are denoted by $q_{Li}^0 = (u_{Li}^0, d_{Li}^0)^T$, and u_{Ri}^0 and d_{Ri}^0 , respectively, using Latin family indices ($i = 1, 2, 3$).

Along with the standard Yukawa terms, the relevant Lagrangian includes new couplings and mass terms. Prior to electroweak symmetry breaking (EWSB), it reads

$$\begin{aligned} -\mathcal{L} = & (Y_u)_{ij} \bar{q}_{Li}^0 \tilde{\Phi} u_{Rj}^0 + (Y_d)_{ij} \bar{q}_{Li}^0 \Phi d_{Rj}^0 + \text{h.c.} \\ & + (Z_u)_{\alpha j} \bar{Q}_{L\alpha}^0 \tilde{\Phi} u_{Rj}^0 + (Z_d)_{\alpha j} \bar{Q}_{L\alpha}^0 \Phi d_{Rj}^0 + \text{h.c.} \\ & + (\bar{M})_{i\beta} \bar{q}_{Li}^0 Q_{R\beta}^0 + (M_Q)_{\alpha\beta} \bar{Q}_{L\alpha}^0 Q_{R\beta}^0 + \text{h.c.}, \end{aligned} \quad (2.2)$$

where the first line contains the standard terms, and $i, j = 1, 2, 3$ and $\alpha, \beta = 1, \dots, N$. Here, Φ denotes the Higgs doublet, while $\tilde{\Phi} \equiv \epsilon\Phi^*$. The last line in \mathcal{L} contains bare mass terms, controlled by \bar{M} and M_Q . It may also prove useful to collect all LH $SU(2)_L$ doublet quark fields in a single $(3 + N)$ -dimensional flavour vector

$$\mathcal{Q}_L^0 = \begin{pmatrix} q_L^0 \\ Q_L^0 \end{pmatrix}, \quad (2.3)$$

which allows us to rewrite the Lagrangian of eq. (2.2) in a compact form:

$$-\mathcal{L} = \bar{\mathcal{Q}}_L^0 \tilde{\Phi} \mathcal{Y}_u u_R^0 + \bar{\mathcal{Q}}_L^0 \Phi \mathcal{Y}_d d_R^0 + \bar{\mathcal{Q}}_L^0 M Q_R^0 + \text{h.c.}, \quad (2.4)$$

where the $(3 + N) \times 3$ matrices \mathcal{Y}_u and \mathcal{Y}_d contain all Yukawa couplings to the Higgs doublet and the $(3 + N) \times N$ matrix M contains all bare mass terms. Given that the $3 + N$ species of LH doublets have identical quantum numbers, one can consider a basis in field space where \bar{M} vanishes and M_Q is real and diagonal ($M_Q \rightarrow D_Q$; see also section 3.1.1). From section 3 onwards, we will take a Lagrangian with $\bar{M} = 0$ and $M_Q = D_Q$ as our starting point, without loss of generality, cf. eqs. (3.5) and (3.6).

Following EWSB, one has $\Phi \rightarrow (0, v + h/\sqrt{2})^T$, with the vacuum expectation value (VEV) $v \simeq 174$ GeV, and obtains the tree-level mass terms

$$\begin{aligned} -\mathcal{L}_m = & \begin{pmatrix} \bar{u}_L^0 & \bar{U}_L^0 \end{pmatrix} \begin{pmatrix} v Y_u & \bar{M} \\ v Z_u & M_Q \end{pmatrix} \begin{pmatrix} u_R^0 \\ U_R^0 \end{pmatrix} + \begin{pmatrix} \bar{d}_L^0 & \bar{D}_L^0 \end{pmatrix} \begin{pmatrix} v Y_d & \bar{M} \\ v Z_d & M_Q \end{pmatrix} \begin{pmatrix} d_R^0 \\ D_R^0 \end{pmatrix} + \text{h.c.} \\ \equiv & \bar{\mathcal{U}}_L^0 \mathcal{M}_u \mathcal{U}_R^0 + \bar{\mathcal{D}}_L^0 \mathcal{M}_d \mathcal{D}_R^0 + \text{h.c.}, \end{aligned} \quad (2.5)$$

where the \mathcal{M}_q are $(3 + N)$ -dimensional square matrices, with

$$\mathcal{M}_q = \begin{pmatrix} & \vdots & \\ m_q & & M \end{pmatrix}, \quad m_q \equiv v \mathcal{Y}_q = v \begin{pmatrix} Y_q \\ Z_q \end{pmatrix}, \quad M \equiv \begin{pmatrix} \overline{M} \\ M_Q \end{pmatrix}, \quad (2.6)$$

and $q = u, d$. We have also defined the vectors

$$\mathcal{U}_{L,R}^0 = \begin{pmatrix} u_{L,R}^0 \\ U_{L,R}^0 \end{pmatrix}, \quad \mathcal{D}_{L,R}^0 = \begin{pmatrix} d_{L,R}^0 \\ D_{L,R}^0 \end{pmatrix}, \quad (2.7)$$

in flavour space. The mass matrices \mathcal{M}_u and \mathcal{M}_d can be diagonalized via bi-unitary transformations (a singular value decomposition). To this end, one defines the mass basis fields via

$$\mathcal{U}_{L,R}^0 = \mathcal{V}_{L,R}^u \mathcal{U}_{L,R} = \mathcal{V}_{L,R}^u \begin{pmatrix} u_{L,R} \\ U_{L,R} \end{pmatrix}, \quad \mathcal{D}_{L,R}^0 = \mathcal{V}_{L,R}^d \mathcal{D}_{L,R} = \mathcal{V}_{L,R}^d \begin{pmatrix} d_{L,R} \\ D_{L,R} \end{pmatrix}, \quad (2.8)$$

where the four \mathcal{V} matrices are $(3 + N)$ -dimensional unitary matrices. One then has

$$-\mathcal{L}_m = \overline{\mathcal{U}}_L \mathcal{D}_u \mathcal{U}_R + \overline{\mathcal{D}}_L \mathcal{D}_d \mathcal{D}_R + \text{h.c.} \quad (2.9)$$

and

$$\mathcal{V}_L^{q\dagger} \mathcal{M}_q \mathcal{V}_R^q = \mathcal{D}_q \equiv \text{diag}(m_1^q, m_2^q, m_3^q, M_1^q, \dots, M_N^q) \quad (q = u, d), \quad (2.10)$$

where the m_i^q and M_α^q are the (non-negative) physical quark masses in each sector, henceforth assumed positive for simplicity. In the case of a single VLQ doublet ($N = 1$), the M_1^q are simply denoted $M_{T'}$ and $M_{B'}$, for $q = u, d$ respectively. As an alternative, one can denote light (standard) quark masses by m_α ($\alpha = u, c, t$) and m_i ($i = d, s, b$), where up- and down-type quarks are labelled by Greek and Latin indices respectively.

Note that eq. (2.10) describes the singular value decompositions of the \mathcal{M}_q . As with any such decomposition, one can find the unitary matrices \mathcal{V}_L^q and \mathcal{V}_R^q via the usual (diagonalization) relations

$$\mathcal{V}_L^{q\dagger} \mathcal{M}_q \mathcal{M}_q^\dagger \mathcal{V}_L^q = \mathcal{V}_R^{q\dagger} \mathcal{M}_q^\dagger \mathcal{M}_q \mathcal{V}_R^q = \mathcal{D}_q^2. \quad (2.11)$$

Before proceeding, it also proves convenient to decompose each \mathcal{V} matrix into its upper and lower blocks, according to

$$\mathcal{V}_{L,R}^q = \begin{pmatrix} A_{L,R}^q \\ \dots\dots\dots \\ B_{L,R}^q \end{pmatrix}, \quad (2.12)$$

where, independently of q , the A^q are $3 \times (3 + N)$ matrices while the B^q are $N \times (3 + N)$ matrices. Equation (2.10) directly implies the relations

$$m_q = \mathcal{V}_L^q \mathcal{D}_q A_R^{q\dagger}, \quad M = \mathcal{V}_L^q \mathcal{D}_q B_R^{q\dagger}. \quad (2.13)$$

Finally, for each chirality, the matrices A^q and B^q obey the relations

$$A^q A^{q\dagger} = \mathbb{1}_3, \quad B^q B^{q\dagger} = \mathbb{1}_N, \quad A^q B^{q\dagger} = 0, \quad A^{q\dagger} A^q + B^{q\dagger} B^q = \mathbb{1}_{3+N}, \quad (2.14)$$

which follow from the unitarity of \mathcal{V}^q .

2.1 Gauge and Higgs interactions

The introduction of doublet VLQs will also modify the form of the electroweak (EW) gauge interactions. The interactions of quarks with the W boson are encoded in the charged-current Lagrangian, which reads

$$\begin{aligned} \mathcal{L}_W &= -\frac{g}{\sqrt{2}} W_\mu^+ \left[\overline{U}_L^0 \gamma^\mu \mathcal{D}_L^0 + \overline{U}_R^0 \gamma^\mu D_R^0 \right] + \text{h.c.} \\ &= -\frac{g}{\sqrt{2}} W_\mu^+ \left[\overline{U}_L \gamma^\mu V_L \mathcal{D}_L + \overline{U}_R \gamma^\mu V_R \mathcal{D}_R \right] + \text{h.c.}, \end{aligned} \quad (2.15)$$

where we have defined the $(3+N)$ -dimensional unitary matrix V_L , and the $(3+N)$ -dimensional right-handed (RH) current matrix V_R , with

$$V_L = \mathcal{V}_L^{u\dagger} \mathcal{V}_L^d, \quad V_R = B_R^{u\dagger} B_R^d = \mathcal{V}_R^{u\dagger} \text{diag}(0, 0, 0, 1, \dots, 1) \mathcal{V}_R^d. \quad (2.16)$$

Important differences arise with respect to an extension of the SM with VLQ isosinglets (see e.g. [46]). Namely, there are two physical mixing matrices, one for each chirality, with V_R originating from the $W_\mu^+ \overline{U}_R^0 \gamma^\mu D_R^0$ term after performing the rotation to the mass eigenbasis, see eq. (2.15). The charged current couplings are determined by the elements of these matrices. In the simplest case of $N = 1$, these are 4×4 square matrices, with

$$V_R = \mathcal{V}_R^{u\dagger} \text{diag}(0, 0, 0, 1) \mathcal{V}_R^d, \quad (\text{for } N = 1). \quad (2.17)$$

While the enlarged LH quark mixing matrix V_L is necessarily unitary, taking the form

$$V_L = \left(\begin{array}{c|ccc} & \begin{matrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{matrix} & \cdots \\ \hline & \vdots & \vdots & \ddots \end{array} \right), \quad (2.18)$$

the new RH mixing matrix V_R cannot be unitary, since $\det V_R = 0$. Here, the 3×3 upper-left block of V_L can be identified as the CKM quark mixing matrix V_{CKM} , which is no longer unitary in general. Nevertheless, at leading order in VLQ corrections and unlike in SM extensions with isosinglet VLQs, the matrix V_{CKM} is still unitary to a good approximation (see also sections 4.1.2 and 7.2). Finally, note that factorizable phases in V_L are not necessarily unphysical as in the SM, where one can exploit the freedom

of identically rephasing LH and RH quark fields while keeping the (diagonalized) mass terms invariant. Indeed, the mixing of RH quarks here means that any such rephasing will modify V_R . Hence, in any physical basis, at least one of V_L or V_R will contain factorizable phases that cannot be eliminated.

Additionally, the mixing of SM quarks with the VLQs induces couplings to the Higgs and Z bosons which are flavour-non-diagonal in the mass basis and originate flavour-changing phenomena. The weak neutral current Lagrangian describing the interactions of quarks with the Z boson is

$$\begin{aligned}\mathcal{L}_Z &= -\frac{g}{2c_W} Z_\mu \left[\overline{U}_L^0 \gamma^\mu U_L^0 - \overline{\mathcal{D}}_L^0 \gamma^\mu \mathcal{D}_L^0 + \overline{U}_R^0 \gamma^\mu U_R^0 - \overline{D}_R^0 \gamma^\mu D_R^0 - 2s_W^2 J_{\text{em}}^\mu \right] \\ &= -\frac{g}{2c_W} Z_\mu \left[\overline{U}_L \gamma^\mu U_L - \overline{\mathcal{D}}_L \gamma^\mu \mathcal{D}_L + \overline{U}_R \gamma^\mu F_u U_R - \overline{\mathcal{D}}_R \gamma^\mu F_d \mathcal{D}_R - 2s_W^2 J_{\text{em}}^\mu \right],\end{aligned}\tag{2.19}$$

with $c_W = \cos \theta_W$ and $s_W = \sin \theta_W$, while the electromagnetic current reads

$$J_{\text{em}}^\mu = +\frac{2}{3} \overline{U} \gamma^\mu U - \frac{1}{3} \overline{\mathcal{D}} \gamma^\mu \mathcal{D},\tag{2.20}$$

with $U = U_L + U_R$ and $\mathcal{D} = \mathcal{D}_L + \mathcal{D}_R$, and preserves its diagonal form in moving from the flavour to the mass basis. In eq. (2.19) we have introduced the $(3 + N)$ -dimensional Hermitian matrices

$$\begin{aligned}F_u &= B_R^{u\dagger} B_R^u = V_R V_R^\dagger = \mathcal{V}_R^{u\dagger} \text{diag}(0, 0, 0, 1, \dots, 1) \mathcal{V}_R^u, \\ F_d &= B_R^{d\dagger} B_R^d = V_R^\dagger V_R = \mathcal{V}_R^{d\dagger} \text{diag}(0, 0, 0, 1, \dots, 1) \mathcal{V}_R^d.\end{aligned}\tag{2.21}$$

Note that LH couplings remain diagonal at tree level, as in the SM, while flavour-changing neutral currents (FCNCs) appear only in the RH sector, controlled by these matrices F_q . This is in contrast with the case of VLQ singlets, for which only LH Z -mediated FCNCs are present. In the simplest case of $N = 1$, the matrices F_q determining FCNCs couplings are 4×4 square matrices, with

$$F_q = \mathcal{V}_R^{q\dagger} \text{diag}(0, 0, 0, 1) \mathcal{V}_R^q, \quad (\text{for } N = 1).\tag{2.22}$$

The relations between the F_q and V_R in eq. (2.21) follow from eq. (2.14). They directly connect the existence of such tree-level FCNCs to the non-unitarity of the RH charged-current mixing matrix.

As for the interactions with the Higgs boson h , the mass matrices are not proportional to the Yukawa matrices and flavour non-diagonal couplings are also generated. The relevant Lagrangian reads

$$\begin{aligned}\mathcal{L}_h &= -\frac{h}{\sqrt{2}} \left[\overline{U}_L^0 \begin{pmatrix} Y_u & 0 \\ Z_u & 0 \end{pmatrix} U_R^0 + \overline{\mathcal{D}}_L^0 \begin{pmatrix} Y_d & 0 \\ Z_d & 0 \end{pmatrix} \mathcal{D}_R^0 \right] + \text{h.c.} \\ &= -\frac{h}{\sqrt{2}} \left[\overline{U}_L \frac{\mathcal{D}_u}{v} (\mathbf{1} - F_u) U_R + \overline{\mathcal{D}}_L \frac{\mathcal{D}_d}{v} (\mathbf{1} - F_d) \mathcal{D}_R \right] + \text{h.c.},\end{aligned}\tag{2.23}$$

where we have used the fact that

$$A_R^{q\dagger} A_R^q = \mathbb{1} - F_q = \mathcal{V}_R^{q\dagger} \text{diag}(1, 1, 1, 0, \dots, 0) \mathcal{V}_R^q. \quad (2.24)$$

Hence, Higgs-mediated FCNCs are controlled by the same quantities as Z -mediated FCNCs, namely F_u and F_d . Given the presence of the matrices \mathcal{D}_q in eq. (2.23), one sees that, as in the case of singlet VLQs, Higgs-mediated transitions between the lighter quarks are suppressed by ratios m_i^q/v with respect to Z -mediated transitions. On the other hand, there emerge non-negligible couplings between right-handed light states and left-handed heavy states, while heavy-light transitions of the opposite chirality are generically suppressed by small SM Yukawas.²

It is worth noting that the last N columns of the mass matrices are shared between \mathcal{M}_u and \mathcal{M}_d , i.e. the $(3 + N) \times N$ bare mass matrix M is common to both sectors. Keeping in mind that

$$B_R^{q\dagger} = \mathcal{V}_R^{q\dagger} \begin{pmatrix} 0_{3 \times N} \\ \mathbb{1}_{N \times N} \end{pmatrix}, \quad (2.25)$$

and recalling eq. (2.13), which implies

$$\mathcal{V}_L^u \mathcal{D}_u B_R^{u\dagger} = \mathcal{V}_L^d \mathcal{D}_d B_R^{d\dagger}, \quad (2.26)$$

one finds the relations

$$\mathcal{D}_u V_R = V_L \mathcal{D}_d F_d, \quad \mathcal{D}_d V_R^\dagger = V_L^\dagger \mathcal{D}_u F_u, \quad (2.27)$$

and

$$\mathcal{D}_u F_u \mathcal{D}_u V_L = V_L \mathcal{D}_d F_d \mathcal{D}_d \quad (2.28)$$

between LH and RH couplings in charged and neutral currents.

The discussion above applies to a general number N of SM-like VLQs. In the simplest case of $N = 1$, M_Q is a number, \overline{M} , Z_u and Z_d are 3-dimensional vectors, and the matrices $\mathcal{D}_u = \text{diag}(m_u, m_c, m_t, M_{T'})$, $\mathcal{D}_d = \text{diag}(m_d, m_s, m_b, M_{B'})$, V_L (unitary), V_R (not unitary), F_u and F_d (Hermitian) are 4×4 square matrices. The counting of independent physical parameters contained in these matrices, for both this case and the most general one, is carried out in the next section.

²In the isosinglet VLQ case, the products $F_q \mathcal{D}_q$ appear in \mathcal{L}_h instead of $\mathcal{D}_q F_q$, so that the reverse happens, i.e. the non-negligible transitions are those between LH light states and RH heavy states.

3 Weak bases and physical parameters

Not all of the $18 + 9N + N^2$ complex Yukawa couplings and bare mass terms in the Lagrangian of eq. (2.4) represent independent physical parameters. Indeed, owing to the invariance of the S-matrix under field redefinitions and the underlying symmetries of QFT, many choices are equivalent. Each choice corresponds to a so-called weak basis (WB) in flavour space. These are connected by weak-basis transformations (WBTs) which keep the kinetic terms invariant, i.e. do not modify the form of the EW gauge interactions of section 2.1. Any two sets of Yukawa and mass matrices related by a WBT represent the same physical system.

In the class of models under consideration, it is straightforward to identify the following transformations as the most general WBTs, describing the redundancy in the determination of the number of independent parameters:

$$\begin{aligned} U_L^0 &\rightarrow \mathcal{W}_L U_L^0, & \mathcal{D}_L^0 &\rightarrow \mathcal{W}_L \mathcal{D}_L^0, \\ u_R^0 &\rightarrow W_R^u u_R^0, & d_R^0 &\rightarrow W_R^d d_R^0, \\ U_R^0 &\rightarrow W_R U_R^0, & D_R^0 &\rightarrow W_R D_R^0, \end{aligned} \quad (3.1)$$

where the matrices W_R^q ($q = u, d$) and W_R are respectively 3×3 and $N \times N$ unitary matrices. Note that the $3 + N$ species of LH doublets have identical quantum numbers, and can thus be connected via a larger $(3 + N) \times (3 + N)$ unitary matrix, \mathcal{W}_L . In the case $N = 1$, $W_R = e^{i\varphi_R}$ is simply a phase.

Under a WBT, as given in eq. (3.1), the Yukawa and mass matrices are not left invariant, but instead transform as

$$\mathcal{Y}_q \rightarrow \mathcal{W}_L^\dagger \mathcal{Y}_q W_R^q, \quad M \rightarrow \mathcal{W}_L^\dagger M W_R, \quad (3.2)$$

where $v \mathcal{Y}_q$ is the $(3 + N) \times 3$ submatrix of \mathcal{M}_q arising from Yukawa couplings to the Higgs doublet and M is the $(3 + N) \times N$ matrix corresponding to the last N columns of the mass matrices \mathcal{M}_q , see eq. (2.6). It follows that the Hermitian combinations

$$h_q \equiv m_q m_q^\dagger = v^2 \mathcal{Y}_q \mathcal{Y}_q^\dagger, \quad H \equiv M M^\dagger \quad (3.3)$$

transform as

$$h_q \rightarrow \mathcal{W}_L^\dagger h_q \mathcal{W}_L, \quad H \rightarrow \mathcal{W}_L^\dagger H \mathcal{W}_L, \quad (3.4)$$

where $m_q = v \mathcal{Y}_q$, c.f. eq. (2.6). Since any product involving these matrices transforms in the same way, they are useful to build WBIs, i.e. physical quantities which do not depend on the WB choice, via traces or determinants.³ WBIs allow one to extract the actual physical content of a set of Yukawa and mass matrices (see sections 5 and 6).

³Notice that also the determinant of a matrix can always be rewritten as a simple polynomial of traces of powers of that matrix and thus traces alone can be used to form complete bases of WBIs.

3.1 Convenient weak bases

3.1.1 A minimal weak basis

The freedom present in the transformations of eq. (3.1) allows one to shape the mass matrices \mathcal{M}_q ($q = u, d$) without affecting their physical content, via eq. (3.2). For instance, one can always move to a WB where the mixed bare mass terms vanish, $\overline{M} = 0$, by an appropriate choice of \mathcal{W}_L , using the fact that the $3 + N$ species of LH doublets have identical quantum numbers. Further WBTs can be used to make the M_Q block become diagonal and non-negative, $M_Q \rightarrow D_Q$. Thus, here and in what follows, without loss of generality, we take as a starting point the pre-EWSB Lagrangian

$$\begin{aligned}
-\mathcal{L} = & (Y_u)_{ij} \overline{q_{Li}^0} \tilde{\Phi} u_{Rj}^0 + (Y_d)_{ij} \overline{q_{Li}^0} \Phi d_{Rj}^0 + \text{h.c.} \\
& + (Z_u)_{\alpha j} \overline{Q_{L\alpha}^0} \tilde{\Phi} u_{Rj}^0 + (Z_d)_{\alpha j} \overline{Q_{L\alpha}^0} \Phi d_{Rj}^0 + \text{h.c.} \\
& + (D_Q)_{\alpha\beta} \overline{Q_{L\alpha}^0} Q_{R\beta}^0 + \text{h.c.},
\end{aligned} \tag{3.5}$$

and the corresponding mass matrices

$$\mathcal{M}_u = \begin{pmatrix} v Y_u & 0 \\ v Z_u & D_Q \end{pmatrix}, \quad \mathcal{M}_d = \begin{pmatrix} v Y_d & 0 \\ v Z_d & D_Q \end{pmatrix}, \tag{3.6}$$

following EWSB. At this stage, one has $18 + 7N$ complex parameters, which are still not independent and physical.

The 3×3 block containing Y_q can be made diagonal (non-negative) via bidiagonalization in one of the two sectors, say the up sector. Explicitly, one may write the singular value decompositions

$$Y_u = \hat{U}_{uL} \hat{Y}_u \hat{U}_{uR}^\dagger, \quad Y_d = \hat{U}_{dL} \hat{Y}_d \hat{U}_{dR}^\dagger, \tag{3.7}$$

where the \hat{Y}_q are diagonal and non-negative,

$$\hat{Y}_u \equiv \text{diag}(\hat{y}_u, \hat{y}_c, \hat{y}_t), \quad \hat{Y}_d \equiv \text{diag}(\hat{y}_d, \hat{y}_s, \hat{y}_b), \tag{3.8}$$

while \hat{U}_{qL} and \hat{U}_{qR} are 3×3 unitary matrices. One is free to choose the order of the non-zero elements in any of the diagonal matrices. Then, \mathcal{W}_L and W_R^u can be used to cancel \hat{U}_{uL} and \hat{U}_{uR} in the matrix Y_u , leading to $Y_u \rightarrow \hat{Y}_u$, while $Y_d \rightarrow Y'_d = \hat{U}_{uL}^\dagger Y_d$.

One is also free to use the matrix W_R^d to remove the rightmost unitary rotation from the (general) matrix $Y'_d = \hat{U}_{uL}^\dagger \hat{U}_{dL} \hat{Y}_d \hat{U}_{dR}^\dagger$, which becomes $Y'_d \rightarrow \hat{V} \hat{Y}_d$, where we have defined

$$\hat{V} \equiv \hat{U}_{uL}^\dagger \hat{U}_{dL}. \tag{3.9}$$

After these WBTs, the matrices Z_u and Z_d now read

$$Z_u \rightarrow z_u = Z_u \hat{U}_{uR}, \quad Z_d \rightarrow z_d = Z_d \hat{U}_{dR}. \tag{3.10}$$

	\hat{Y}_u	\hat{z}_u	\hat{V}	\hat{Y}_d	z_d	D_Q	Total
Moduli	3	$3N$	3	3	$3N$	N	$9 + 7N$
Phases	0	$2N$	1	0	$3N$	0	$1 + 5N$
Total	3	$5N$	4	3	$6N$	N	$10 + 12N$

Table 1: Physical parameter count in the minimal WB of eqs. (3.11) and (3.12), in the presence of N isodoublet VLQs.

As in the SM, 5 phases can be removed from \hat{V} by rephasing standard quark fields.⁴ Such rephasings can be included from the start in the definition of the matrices \hat{U}_{qL} , \hat{U}_{qR} . Then, \hat{V} can be parameterized as the standard CKM, with three mixing angles and a phase. Thanks to the possibility of removing these phases, the angles can be chosen to lie in the first quadrant, while the range of the (sole) physical phase is a priori unrestricted. With these choices, \hat{V} becomes the SM V_{CKM} in the limit of VLQ doublet decoupling ($z_u, z_d \rightarrow 0$, $D_Q \rightarrow \infty \mathbf{1}$).

Note that z_u and z_d have remained general complex, up to this point. One can finally use the $N \times N$ lower-right block of \mathcal{W}_L , which we will denote P , to remove N of the $6N$ phases contained in z_u and z_d .⁵ As a concrete possibility, we consider removing one phase in each row of z_u , making its first column real.⁶ As a result of the above sequence of transformations, the Yukawa and mass matrices now take the form

$$\mathcal{Y}_u = \begin{pmatrix} \hat{Y}_u \\ \hat{z}_u \end{pmatrix}, \quad \mathcal{Y}_d = \begin{pmatrix} \hat{V} \hat{Y}_d \\ z_d \end{pmatrix}, \quad M = \begin{pmatrix} 0 \\ D_Q \end{pmatrix} \quad (3.11)$$

(*minimal WB*), which after EWSB correspond to the mass matrices

$$\mathcal{M}_u = \begin{pmatrix} v \hat{Y}_u & 0 \\ v \hat{z}_u & D_Q \end{pmatrix}, \quad \mathcal{M}_d = \begin{pmatrix} v \hat{V} \hat{Y}_d & 0 \\ v z_d & D_Q \end{pmatrix}. \quad (3.12)$$

Here, $\hat{z}_u = P z_u$ has N phases less than the general complex z_u .

Having exhausted the WB freedom, the minimal WB of eqs. (3.11) and (3.12) allows one to count the number of physical parameters. One thus finds a total of $10 + 12N$ parameters, with the corresponding breakdown being shown in Table 1. Out of this total, $9 + 7N$ parameters correspond to physical moduli, while $1 + 5N$ correspond to physical phases. Anticipating the correspondence of the moduli to physical masses and mixing angles, the parameter count in the scenarios of interest is summarized in Table 2. Independent ways of counting physical parameters are presented in section 3.2.

⁴The sixth phase is not removable due to the unbroken baryon number. In this procedure, the diagonal matrix \hat{Y}_u is unchanged by an appropriate rephasing with W_R^u . The columns of both z_d and z_u are rephased.

⁵The diagonal matrix D_Q is unchanged by an appropriate rephasing with W_R affecting both sectors.

⁶One could just as well have removed N phases from the first column of $z_d \rightarrow \hat{z}_d$, as considered in sections 5.2.2 and 7.1.

	SM	New ($N = 1$)	Total ($N = 1$)	New ($N \geq 1$)	Total ($N \geq 1$)
Quark masses	6	2	8	$2N$	$6 + 2N$
Mixing angles	3	5	8	$5N$	$3 + 5N$
Total moduli	9	7	16	$7N$	$9 + 7N$
Phases	1	5	6	$5N$	$1 + 5N$
Total angles + phases	4	10	14	$10N$	$4 + 10N$
Total (parameters)	10	12	22	$12N$	$10 + 12N$

Table 2: Summary of the number of physical parameters in SM extensions with one ($N = 1$) and several ($N > 1$) VLQ doublets with hypercharge $1/6$.

3.1.2 A “stepladder” weak basis ($N = 1$)

In the case $N = 1$, one can move to a special “stepladder” WB which allows for a sequential and straightforward determination of the Yukawa couplings from a set of WBIs.

We take the minimal WB of eq. (3.12) as a starting point. Then, one can move all 5 physical phases in \hat{z}_u and z_d to \hat{V} by rephasing standard quarks, undoing the aforementioned rephasing and restoring \hat{V} in this context to a general unitary matrix $\hat{V} \rightarrow \tilde{V}$ (not CKM-like), while z_u and z_d are now real 3-dimensional row vectors. The mass matrices will have the structure

$$\mathcal{M}_u \sim \begin{pmatrix} \mathbb{R} & & & \\ & \mathbb{R} & & \\ & & \mathbb{R} & \\ \mathbb{R} & \mathbb{R} & \mathbb{R} & \mathbb{R} \end{pmatrix}, \quad \mathcal{M}_d \sim \begin{pmatrix} \tilde{V} & & & \\ & 1 & & \end{pmatrix} \begin{pmatrix} \mathbb{R} & & & \\ & \mathbb{R} & & \\ & & \mathbb{R} & \\ \mathbb{R} & \mathbb{R} & \mathbb{R} & \mathbb{R} \end{pmatrix}. \quad (3.13)$$

One can use the matrices W_R^q to perform an orthogonal rotation in each sector, mixing the first two columns of \mathcal{M}_u and of (the rightmost matrix of) \mathcal{M}_d in order to cancel the $(4, 1)$ element in each of these matrices. Then, via a similar orthogonal rotation of the second and third columns in each sector, one may cancel the $(4, 2)$ entries, leading to a WB where, generically,

$$\mathcal{M}_u \sim \begin{pmatrix} \mathbb{R} & \mathbb{R} & \mathbb{R} & \\ \mathbb{R} & \mathbb{R} & \mathbb{R} & \\ & \mathbb{R} & \mathbb{R} & \\ & & \mathbb{R} & \mathbb{R} \end{pmatrix}, \quad \mathcal{M}_d \sim \begin{pmatrix} \tilde{V} & & & \\ & 1 & & \end{pmatrix} \begin{pmatrix} \mathbb{R} & \mathbb{R} & \mathbb{R} & \\ \mathbb{R} & \mathbb{R} & \mathbb{R} & \\ & \mathbb{R} & \mathbb{R} & \\ & & \mathbb{R} & \mathbb{R} \end{pmatrix}. \quad (3.14)$$

Now, the upper-left block of \mathcal{W}_L can be used as an orthogonal rotation between the first two lines, enforcing a zero in the $(1, 3)$ position of \mathcal{M}_u . A zero in the same position in the rightmost matrix of \mathcal{M}_d can be obtained by absorbing orthogonal rotations (including the one from \mathcal{W}_L) in the general \tilde{V} . Similarly, one can enforce zeros in the $(2, 3)$ positions

by mixing the second and third rows and reach a WB where

$$\mathcal{M}_u \sim \begin{pmatrix} \mathbb{R} & \mathbb{R} & & \\ \mathbb{R} & \mathbb{R} & & \\ \mathbb{R} & \mathbb{R} & \mathbb{R} & \\ & & \mathbb{R} & \mathbb{R} \end{pmatrix}, \quad \mathcal{M}_d \sim \begin{pmatrix} \tilde{V} & & & \\ & 1 & & \end{pmatrix} \begin{pmatrix} \mathbb{R} & \mathbb{R} & & \\ \mathbb{R} & \mathbb{R} & & \\ \mathbb{R} & \mathbb{R} & \mathbb{R} & \\ & & \mathbb{R} & \mathbb{R} \end{pmatrix}. \quad (3.15)$$

Finally, one may use RH rotations between the first two columns to cancel the (3,1) elements and LH rotations between the first two rows to cancel the (1,2) elements, arriving at the desired *stepladder WB*,

$$\mathcal{M}_u \sim \begin{pmatrix} \mathbb{R} & & & \\ \mathbb{R} & \mathbb{R} & & \\ & \mathbb{R} & \mathbb{R} & \\ & & \mathbb{R} & \mathbb{R} \end{pmatrix}, \quad \mathcal{M}_d \sim \begin{pmatrix} \tilde{V} & & & \\ & 1 & & \end{pmatrix} \begin{pmatrix} \mathbb{R} & & & \\ \mathbb{R} & \mathbb{R} & & \\ & \mathbb{R} & \mathbb{R} & \\ & & \mathbb{R} & \mathbb{R} \end{pmatrix}. \quad (3.16)$$

Note that all intermediate basis are valid WB choices. Details on the extraction of couplings in the WB of eq. (3.16) from a set of invariants is postponed to section 5.2.1. In what follows, we will consider the following explicit notation for the entries of the mass matrices in the stepladder WB:

$$\mathcal{M}_u = M_Q \begin{pmatrix} r_{u5} & 0 & 0 & 0 \\ r_{u4} & r_{u3} & 0 & 0 \\ 0 & r_{u2} & r_{u1} & 0 \\ 0 & 0 & r_{u0} & 1 \end{pmatrix}, \quad \mathcal{M}_d = M_Q \begin{pmatrix} \tilde{V} & & & \\ & 1 & & \end{pmatrix} \begin{pmatrix} r_{d5} & 0 & 0 & 0 \\ r_{d4} & r_{d3} & 0 & 0 \\ 0 & r_{d2} & r_{d1} & 0 \\ 0 & 0 & r_{d0} & 1 \end{pmatrix}, \quad (3.17)$$

where r_{qi} ($i = 0, \dots, 5$) are dimensionless (note the overall M_Q factors). Given these definitions, one expects $r_{qi} \lesssim v/M_Q$.

We emphasize that, in the $N = 1$ case, one can always switch between this WB and that of eq. (3.12) via WBTs, while keeping the number of parameters at a minimum. In this sense, both WBs can be said to be minimal. This remains true even when some of the parameters characterizing these WBs vanish. In fact, one can prove that, for a given sector, when any number of off-diagonal real parameters vanishes in the stepladder WB, the same number of real couplings between SM quarks and the VLQ doublets must vanish in the minimal WB. Namely, it is straightforward to show that:

- If $r_{q4} = 0$, then when transforming into the minimal WB one finds that the Yukawa coupling vector \mathbf{z}_q defined in section 3.1.1 has one vanishing entry.
- If $r_{q2} = 0$, then r_{q4} can be eliminated using WBTs and when switching into the minimal WB one finds that \mathbf{z}_q has two vanishing entries.
- If $r_{q0} = 0$, then both r_{q4} and r_{q2} can be eliminated using WBTs, resulting in $\mathbf{z}_q = 0$ when considering the minimal WB. In that case the VLQ is decoupled from that sector.
- $r_{q5,3,1} \neq 0$ correspond to non-zero Yukawa couplings \hat{y}_α or \hat{y}_i .

This link between both WBs will be relevant later on to establish a connection between the results of section 5.2.1 and those of section 6.

3.2 Alternate ways of counting physical parameters

3.2.1 Counting via a spurion analysis

To independently check the (physical) parameter count of Table 1, one may perform a spurion analysis [63] (see also [46, 64]), which does not require fixing a WB. One starts by identifying the general flavour symmetry available in the absence of mass and Yukawa terms — in direct correspondence with the WB freedom of eq. (3.1), as mentioned above. Mass and Yukawa terms break the symmetry of the kinetic terms to the conserved baryon number symmetry. Schematically, one has

$$U(3+N)_{\mathcal{Q}_L} \times U(3)_{u_R} \times U(3)_{d_R} \times U(N)_{\mathcal{Q}_R} \rightarrow U(1)_B. \quad (3.18)$$

The number of physical parameters $N_{\text{phys}} = N_{\text{phys}}^{\text{moduli}} + N_{\text{phys}}^{\text{phases}}$ can be obtained from the differences

$$\begin{aligned} N_{\text{phys}}^{\text{moduli}} &= N_{\text{general}}^{\text{moduli}} - N_{\text{broken}}^{\text{moduli}}, \\ N_{\text{phys}}^{\text{phases}} &= N_{\text{general}}^{\text{phases}} - N_{\text{broken}}^{\text{phases}}, \end{aligned} \quad (3.19)$$

where N_{general} refers to the number of parameters (even if unphysical) describing the \mathcal{M}_q matrices in a generic WB, i.e.

$$\begin{aligned} N_{\text{general}}^{\text{moduli}} &= N_{\text{general}}^{\text{phases}} = \overbrace{2(9+3N)}^{Y_q, Z_q} + \overbrace{3N}^{\overline{M}} + \overbrace{N^2}^{M_Q} = 18 + 9N + N^2 \\ \Rightarrow N_{\text{general}} &= N_{\text{general}}^{\text{moduli}} + N_{\text{general}}^{\text{phases}} = 36 + 18N + 2N^2. \end{aligned} \quad (3.20)$$

As for N_{broken} , each $U(k)$ factor contributes with $\frac{1}{2}k(k-1)$ moduli and $\frac{1}{2}k(k+1)$ phases. The chain of eq. (3.18) then implies

$$\begin{cases} N_{\text{broken}}^{\text{moduli}} = \frac{1}{2}(3+N)(2+N) + 3 + 3 + \frac{1}{2}N(N-1) - 0 = 9 + 2N + N^2 \\ N_{\text{broken}}^{\text{phases}} = \frac{1}{2}(3+N)(4+N) + 6 + 6 + \frac{1}{2}N(N+1) - 1 = 17 + 4N + N^2 \end{cases} \quad (3.21)$$

$$\Rightarrow N_{\text{broken}} = N_{\text{broken}}^{\text{moduli}} + N_{\text{broken}}^{\text{phases}} = 26 + 6N + 2N^2.$$

As a result, we obtain

$$\begin{cases} N_{\text{phys}}^{\text{moduli}} = 9 + 7N \\ N_{\text{phys}}^{\text{phases}} = 1 + 5N \end{cases} \Rightarrow N_{\text{phys}} = 10 + 12N, \quad (3.22)$$

in agreement with Table 1.

The results above apply to a general number N of SM-like VLQs. In the simplest case of $N = 1$, one counts 12 extra parameters in the quark sector (7 moduli and 5 phases), for a total of 22 physical parameters (16 moduli and 6 phases).

3.2.2 Counting in the mass basis ($N = 1$)

Consider the mass basis and the simplest case of $N = 1$. We have 8 physical masses. The mixing matrix in the LH sector, V_L , is a unitary 4×4 matrix which can be parameterized using 6 angles and 10 phases. As regards the RH sector, the matrices V_R , F_u and F_d are built from B_R^u and B_R^d , i.e. from the last rows of the unitary matrices \mathcal{V}_R^u and \mathcal{V}_R^d , which contain a total of 6 independent mixing angles (3 in each B_R^q) and 7 independent phases, as only differences of the 8 phases (4 in each B_R^q) appear.

However, not all of the $8 + 6 + 6 = 20$ moduli and $10 + 7 = 17$ phases identified above are physical. Recall that the extra species is vector-like. In the mass basis, this information is encoded in the relations of eq. (2.26), which in this case read

$$\mathcal{D}_u \mathcal{V}_R^{u\dagger} \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} = V_L \mathcal{D}_d \mathcal{V}_R^{d\dagger} \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}. \quad (3.23)$$

This equations represent 4 real and 4 imaginary conditions. Then, we have $8 + 6 + 6 - 4 = 16$ moduli describing the model. As regards the phases, note that, beyond the 4 imaginary constraints set by eq. (3.23), 7 phases can be eliminated (in total, between the RH and LH sectors) by rephasing the quark fields. Namely, one is free to rephase all 8 quark fields u, c, t, T' , and d, s, b, B' in the mass basis with common phases for both chiralities (leaving the mass terms real), cf. eq. (3.25). Due to the conserved baryon number, this procedure can remove at most 7 unphysical phases, as indicated above. Thus, we are left with $10 + 7 - 4 - 7 = 6$ physical phases describing the model.

As an example, out of the 10 phases contained in V_L , one may choose to eliminate 5 phases related to the LH mixing of the three light quark species by rephasing the corresponding fields. Then, 4 of the remaining 5 phases are determined by the relations in eq. (3.23). This means that only one physical phase is contained in V_L . By rephasing the T' and B' fields, 2 out of 7 phases can be eliminated from the RH currents, leaving 5 phases as free parameters in the RH sector. This results in a total of 6 physical phases, as anticipated.

This discussion can be extended to the case of more VLQ doublets. Counting in the context of a specific parameterization, for $N \geq 1$, is carried out in section 4.2.

3.3 Rephasing invariants

Observables can only depend on phase-convention-independent quantities. In the SM, one can construct rephasing-invariant quantities in terms of the mixing using as building blocks the moduli $|V_{\alpha i}|^2$ of CKM entries or quartets of the form $Q_{\alpha i \beta j} = V_{\alpha i} V_{\beta j} V_{\alpha j}^* V_{\beta i}^*$, involving four different quarks. When $\alpha \neq \beta$ and $i \neq j$, these are the simplest generically-complex rephasing invariants. The non-vanishing of the imaginary parts of the quartets is then directly connected to the existence of CP violation in the SM. Namely, given the unitarity of V_{CKM} , all nine quantities $\text{Im } Q_{\alpha i \beta j}$ ($\alpha \neq \beta, i \neq j$) are equal to the invariant

J [52, 65, 66] up to a sign,

$$J \sum_{\gamma, k} \epsilon_{\alpha\beta\gamma} \epsilon_{ijk} = \text{Im}(V_{\alpha i} V_{\beta j} V_{\alpha j}^* V_{\beta i}^*), \quad (3.24)$$

and all CP-violating effects in the SM are proportional to J .

When introducing VLQ doublets, apart from the mixing of LH quarks in the charged currents, we will also have mixing of RH quarks in both charged and neutral currents. Hence, one can expect new types of rephasing-invariant quantities to be relevant and potentially new sources of CP violation to arise. In this subsection, we briefly present some of the simplest examples of such quantities in the context of these extensions.

We turn our attention to the gauge interactions in eqs. (2.15) and (2.19), from which we can conclude that under general rephasings of the quark mass eigenstates $U_\alpha \rightarrow U_\alpha e^{i\varphi_\alpha}$ and $D_i \rightarrow D_i e^{i\varphi_i}$, the mixing matrices transform as

$$\begin{aligned} V_{\alpha i}^L &\rightarrow e^{-i(\varphi_\alpha - \varphi_i)} V_{\alpha i}^L, \\ V_{\alpha i}^R &\rightarrow e^{-i(\varphi_\alpha - \varphi_i)} V_{\alpha i}^R, \\ F_{\alpha\beta}^u &\rightarrow e^{-i(\varphi_\alpha - \varphi_\beta)} F_{\alpha\beta}^u, \\ F_{ij}^d &\rightarrow e^{-i(\varphi_i - \varphi_j)} F_{ij}^d, \end{aligned} \quad (3.25)$$

where chirality and quark-type subscripts were changed to superscripts to avoid index cluttering. Now, taking these transformations into account, we provide some examples of rephasing invariants in the general case of $N \geq 1$ doublets, and then focus on the particular case of a single doublet ($N = 1$), for which the number of independent rephasing invariants is lower.

3.3.1 The general case ($N \geq 1$)

Since V_L and V_R transform in the same manner, any bilinear of the form

$$\mathcal{B}_{\alpha i}^{\chi\chi'} \equiv V_{\alpha i}^\chi V_{\alpha i}^{\chi'*} = \left(\mathcal{B}_{\alpha i}^{\chi'\chi}\right)^* \quad (3.26)$$

is a rephasing-invariant quantity. Here $\chi, \chi' = L, R$ are chirality indices. For $\chi = \chi'$, these quantities reduce to the squared moduli $|V_{\alpha i}^L|^2$ or $|V_{\alpha i}^R|^2$. When $\chi \neq \chi'$, we obtain bilinears $\mathcal{B}_{\alpha i}^{LR} = V_{\alpha i}^L V_{\alpha i}^{R*}$ which are complex in general and have no analogue in the SM (or even in extensions with VLQ singlets). These are potentially related to new sources of CP violation. In particular, this means one may expect CP-violating effects in charged currents which involve only two quarks. The next simplest rephasing invariants one can build are complex trilinears such as

$$\begin{aligned} \mathcal{T}_{i,\alpha\beta}^{\chi\chi'} &\equiv V_{\alpha i}^{\chi*} V_{\beta i}^{\chi'} F_{\alpha\beta}^u = \left(\mathcal{T}_{i,\beta\alpha}^{\chi'\chi}\right)^*, \\ \mathcal{T}_{\alpha,ij}^{\chi\chi'} &\equiv V_{\alpha i}^\chi V_{\alpha j}^{\chi'*} F_{ij}^d = \left(\mathcal{T}_{\alpha,ji}^{\chi'\chi}\right)^*. \end{aligned} \quad (3.27)$$

These invariants, which are associated with FCNCs, involve three quarks. Other meaningful rephasing invariant quantities that have analogues in the SM are the quartets. Here, in general we have

$$\mathcal{Q}_{\alpha i \beta j}^{\chi \chi' \xi \xi'} \equiv V_{\alpha i}^{\chi} V_{\beta j}^{\chi'} V_{\alpha j}^{\xi*} V_{\beta i}^{\xi'*} = \left(\mathcal{Q}_{\beta i \alpha j}^{\xi' \xi \chi' \chi} \right)^* = \left(\mathcal{Q}_{\alpha j \beta i}^{\xi \xi' \chi \chi'} \right)^* = \mathcal{Q}_{\beta j \alpha i}^{\chi' \chi \xi' \xi}, \quad (3.28)$$

where $\xi, \xi' = L, R$ are also chirality indices.

Note that trilinears can be written as sums of quartets. For instance,

$$\begin{aligned} \sum_i \mathcal{Q}_{\alpha i \beta j}^{R \chi \chi' R} &= F_{\alpha \beta}^u V_{\beta j}^{\chi} V_{\alpha j}^{\chi'*} = \mathcal{T}_{j, \alpha \beta}^{\chi' \chi}, \\ \sum_{\alpha} \mathcal{Q}_{\alpha i \beta j}^{R \chi R \chi'} &= F_{j i}^d V_{\beta j}^{\chi} V_{\beta i}^{\chi'*} = \mathcal{T}_{\beta, j i}^{\chi \chi'}. \end{aligned} \quad (3.29)$$

Additionally, one has

$$\sum_{i, j} \mathcal{Q}_{\alpha i \beta j}^{R R R R} = |F_{\alpha \beta}^u|^2 \quad \text{and} \quad \sum_{\alpha, \beta} \mathcal{Q}_{\alpha i \beta j}^{R R R R} = |F_{i j}^d|^2. \quad (3.30)$$

In these equations, the sums go over all $3 + N$ quarks of a certain type (including the new states). Replacing $R \rightarrow L$ in these relations and making use of the unitarity of V_L , we can relate sums of quartets to the bilinears of eq. (3.26), namely

$$\begin{aligned} \sum_i \mathcal{Q}_{\alpha i \beta j}^{L \chi \chi' L} &= \delta_{\alpha \beta} V_{\beta j}^{\chi} V_{\alpha j}^{\chi'*} = \delta_{\alpha \beta} \mathcal{B}_{\alpha j}^{\chi \chi'}, \\ \sum_{\alpha} \mathcal{Q}_{\alpha i \beta j}^{L \chi L \chi'} &= \delta_{j i} V_{\beta j}^{\chi} V_{\beta i}^{\chi'*} = \delta_{i j} \mathcal{B}_{\beta i}^{\chi \chi'}. \end{aligned} \quad (3.31)$$

We further obtain the constraints

$$\sum_{i, j} \mathcal{Q}_{\alpha i \beta j}^{L L L L} = \delta_{\alpha \beta} \quad \text{and} \quad \sum_{\alpha, \beta} \mathcal{Q}_{\alpha i \beta j}^{L L L L} = \delta_{i j}, \quad (3.32)$$

which follow directly from the above result and also hold in the absence of VLQs.

In principle one can continue building other distinct rephasing invariants by appropriately assembling arbitrarily large products of the entries of V_L , V_R and F_q . However, here we choose to leave this discussion at the quartet level, as, at least in the case of one VLQ doublet, after this point rephasing-invariant quantities are expected to be less relevant from a phenomenological standpoint.

3.3.2 A single doublet ($N = 1$)

The case of a single doublet is special because the number of distinct invariants is considerably reduced when compared to cases with $N > 1$. This is connected to the fact that, in this specific case, we have

$$V_{\alpha i}^R V_{\beta j}^R = V_{\beta i}^R V_{\alpha j}^R \quad \Rightarrow \quad \begin{cases} V_{\beta i}^R F_{\alpha \beta}^u = V_{\beta i}^R \sum_j V_{\alpha j}^R V_{\beta j}^{R*} = V_{\alpha i}^R \sum_j |V_{\beta j}^R|^2, \\ V_{\alpha i}^R F_{i j}^d = V_{\alpha i}^R \sum_{\beta} V_{\beta i}^{R*} V_{\beta j}^R = V_{\alpha j}^R \sum_{\beta} |V_{\beta i}^R|^2, \end{cases} \quad (3.33)$$

as a consequence of having $V_{\alpha i}^R = (B_R^u)_\alpha (B_R^d)_i$, where, for $N = 1$, the B_R^q are vectors in flavour space. It then follows that the only independent trilinears are those of the form

$$\mathcal{T}_{i,\alpha\beta}^{LL} = (\mathcal{T}_{i,\beta\alpha}^{LL})^* = V_{\alpha i}^{L*} V_{\beta i}^L F_{\alpha\beta}^u \quad \text{and} \quad \mathcal{T}_{\alpha,ij}^{LL} = (\mathcal{T}_{\alpha,ji}^{LL})^* = V_{\alpha i}^L V_{\alpha j}^{L*} F_{ij}^d, \quad (3.34)$$

while the only generically-complex independent quartets are those of the type

$$\mathcal{Q}_{\alpha i \beta j}^{LLL\chi} = \mathcal{Q}_{\beta j \alpha i}^{LL\chi L} = \left(\mathcal{Q}_{\alpha j \beta i}^{L\chi LL} \right)^* = \left(\mathcal{Q}_{\beta i \alpha j}^{\chi LLL} \right)^* = V_{\alpha i}^L V_{\beta j}^L V_{\alpha j}^{L*} V_{\beta i}^{\chi*}, \quad (3.35)$$

with $\chi = L, R$. Quartets like $\mathcal{Q}_{\alpha i \beta j}^{RRLL}$, for instance, can be written as the product of two bilinears $\mathcal{B}_{\alpha j}^{RL} \mathcal{B}_{\beta i}^{RL}$ and do not contain new information on complex phases.

Note that no complex trilinears depend on V_R and that no complex quartets depend solely on V_R . This can be traced to the fact that when $N = 1$ all phases in V_R can be brought into V_L via rephasings, while the reciprocal is not true. In fact, under the rephasing of the quark mass eigenstates $\mathcal{U}_\alpha \rightarrow \mathcal{U}_\alpha e^{i\varphi_\alpha}$ and $\mathcal{D}_i \rightarrow \mathcal{D}_i e^{i\varphi_i}$ one has

$$(B_R^u)_\alpha \rightarrow (B_R^u)_\alpha e^{i\varphi_\alpha} \quad \text{and} \quad (B_R^d)_i \rightarrow (B_R^d)_i e^{i\varphi_i}, \quad (3.36)$$

where we again emphasize that the B_R^q are vectors in flavour space. Hence, with an appropriate choice of φ_α and φ_i , all B_R^q can be made real, and consequently also $V_R = B_R^{u\dagger} B_R^d$, $F_u = B_R^{u\dagger} B_R^u$ and $F_d = B_R^{d\dagger} B_R^d$ will be real. One may thus eliminate all phases from the RH currents in the $N = 1$ case, which then emerge in the LH currents.

4 Rotating to the mass basis

As previously seen, the relevant parts of the Lagrangian in the mass basis read:

$$\begin{aligned}
\mathcal{L}_m &= -\overline{U}_L \mathcal{D}_u U_R - \overline{D}_L \mathcal{D}_d \mathcal{D}_R + \text{h.c.}, \\
\mathcal{L}_W &= -\frac{g}{\sqrt{2}} W_\mu^+ [\overline{U}_L \gamma^\mu V_L \mathcal{D}_L + \overline{U}_R \gamma^\mu V_R \mathcal{D}_R] + \text{h.c.}, \\
\mathcal{L}_Z &= -\frac{g}{2c_W} Z_\mu [\overline{U}_L \gamma^\mu U_L - \overline{D}_L \gamma^\mu \mathcal{D}_L + \overline{U}_R \gamma^\mu F_u U_R - \overline{D}_R \gamma^\mu F_d \mathcal{D}_R - 2s_W^2 J_{\text{em}}^\mu], \\
\mathcal{L}_h &= -\frac{h}{\sqrt{2}} \left[\overline{U}_L \frac{\mathcal{D}_u}{v} (\mathbb{1} - F_u) U_R + \overline{D}_L \frac{\mathcal{D}_d}{v} (\mathbb{1} - F_d) \mathcal{D}_R \right] + \text{h.c.},
\end{aligned} \tag{4.1}$$

after EWSB. In what follows, we write the physical quark masses, the mixing matrices V_L and V_R , and the FCNC matrices F_u and F_d in terms of flavour-basis Yukawa couplings for $N = 1$ (section 4.1). This allows us to identify an alternative set of rephasing-invariant quantities. We then present exact parameterizations for the mixing matrices V_L and V_R in these theories, for any $N \geq 1$ (section 4.2).

4.1 Approximate masses and mixing ($N = 1$)

We consider for simplicity the case $N = 1$.⁷ Let us start from the general Lagrangian of eq. (3.5), where mixed mass terms of the type $\overline{M} \overline{q}_L Q_R$ have been rotated away, so that the mass matrices of up-type and down-type quarks read

$$\mathcal{M}_u = \begin{pmatrix} v Y_u & 0 \\ v Z_u & M_Q \end{pmatrix}, \quad \mathcal{M}_d = \begin{pmatrix} v Y_d & 0 \\ v Z_d & M_Q \end{pmatrix}. \tag{4.2}$$

These mass matrices can be diagonalized via bi-unitary transformations, as in eq. (2.10):

$$\begin{aligned}
\mathcal{V}_L^{u\dagger} \mathcal{M}_u \mathcal{V}_R^u &= \mathcal{D}_u = \text{diag}(y_u v, y_c v, y_t v, M_{T'}), \\
\mathcal{V}_L^{d\dagger} \mathcal{M}_d \mathcal{V}_R^d &= \mathcal{D}_d = \text{diag}(y_d v, y_s v, y_b v, M_{B'}),
\end{aligned} \tag{4.3}$$

and the rotation matrices \mathcal{V} can be obtained from the Hermitian products $\mathcal{M}_q \mathcal{M}_q^\dagger$ and $\mathcal{M}_q^\dagger \mathcal{M}_q$, see eq. (2.11). The initial states are related to the mass eigenstates via

$$\begin{pmatrix} u_1 \\ u_2 \\ u_3 \\ T \end{pmatrix}_{L,R} = \mathcal{V}_{L,R}^u \begin{pmatrix} u \\ c \\ t \\ T' \end{pmatrix}_{L,R}, \quad \begin{pmatrix} d_1 \\ d_2 \\ d_3 \\ B \end{pmatrix}_{L,R} = \mathcal{V}_{L,R}^d \begin{pmatrix} d \\ s \\ b \\ B' \end{pmatrix}_{L,R}, \tag{4.4}$$

where we have expanded and simplified the notation, cf. eq. (2.8). Finally, it is useful to keep in mind the singular value decompositions of Y_u and Y_d introduced in eqs. (3.7) and (3.8). Namely, recall that

$$\hat{U}_{uL}^\dagger Y_u \hat{U}_{uR} = \hat{Y}_u = \text{diag}(\hat{y}_u, \hat{y}_c, \hat{y}_t), \quad \hat{U}_{dL}^\dagger Y_d \hat{U}_{dR} = \hat{Y}_d = \text{diag}(\hat{y}_d, \hat{y}_s, \hat{y}_b), \tag{4.5}$$

⁷This analysis can be extended to $N > 1$. We make use of such a formalism for the cases with $N = 2$ in sections 6 and 8.

where the diagonal elements are not yet the physical masses in units of v (only approximately so, see later eq. (4.10)). We also define

$$\mathbf{z}_u \equiv Z_u \hat{U}_{uR} = (z_u, z_c, z_t), \quad \mathbf{z}_d \equiv Z_d \hat{U}_{dR} = (z_d, z_s, z_b). \quad (4.6)$$

Then, one can directly decompose the mass matrices as

$$\mathcal{M}_q = \begin{pmatrix} \hat{U}_{qL} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} v \hat{Y}_q & 0 \\ v \mathbf{z}_q & M_Q \end{pmatrix} \begin{pmatrix} \hat{U}_{qR}^\dagger & 0 \\ 0 & 1 \end{pmatrix} \quad (q = u, d), \quad (4.7)$$

and carry out their diagonalization. For both chiralities $\chi = L, R$, one finds that the rotation matrices \mathcal{V} can be schematically parameterized as

$$\mathcal{V}_\chi^q = \begin{pmatrix} \hat{U}_{q\chi} & \\ & 1 \end{pmatrix} \begin{pmatrix} \delta U_{q\chi} & \\ & 1 \end{pmatrix} \begin{pmatrix} R_4^{q\chi} \\ & \end{pmatrix}, \quad (4.8)$$

in each sector, where each R_4 matrix is a product of (1, 4), (2, 4) and (3, 4) unitary rotations.

The present experimental bounds on VLQ masses are $M_{T'}, M_{B'} > 1\text{--}3$ TeV [67, 68], depending on the decay modes. For perturbative couplings, the bare mass M_Q (here a number) is expected to source the VLQ scale at the TeV or beyond, and thus the ratio v/M_Q is a small parameter. Taking this into account, masses and mixings in both the RH and LH sectors can be directly related to Lagrangian parameters through power series in v/M_Q (see also Ref. [49]). Note that, beyond agreeing with the EFT expansion of section 7 when computing tree-level amplitudes at relatively low energies, this v/M_Q expansion can be used at energies above the VLQ threshold, unlike the EFT one.

Before discussing the rotations bidiagonalizing the \mathcal{M}_q , we first obtain expressions for the eigenvalues of $\mathcal{M}_q \mathcal{M}_q^\dagger$, i.e. for the squares of physical quark masses. One finds

$$\begin{aligned} M_{T'}^2 &= M_Q^2 \left[1 + |\mathbf{z}_u|^2 \frac{v^2}{M_Q^2} + \mathcal{O}\left(\frac{v^4}{M_Q^4}\right) \right], \\ M_{B'}^2 &= M_Q^2 \left[1 + |\mathbf{z}_d|^2 \frac{v^2}{M_Q^2} + \mathcal{O}\left(\frac{v^4}{M_Q^4}\right) \right], \end{aligned} \quad (4.9)$$

for the masses of the new states, implying $M_{T'} \simeq M_{B'}$, and

$$\begin{aligned} m_\alpha^2 &= v^2 y_\alpha^2 = v^2 \hat{y}_\alpha^2 \left[1 - |z_\alpha|^2 \frac{v^2}{M_Q^2} + \mathcal{O}\left(\frac{v^4}{M_Q^4}\right) \right] \quad (\alpha = u, c, t), \\ m_i^2 &= v^2 y_i^2 = v^2 \hat{y}_i^2 \left[1 - |z_i|^2 \frac{v^2}{M_Q^2} + \mathcal{O}\left(\frac{v^4}{M_Q^4}\right) \right] \quad (i = d, s, b), \end{aligned} \quad (4.10)$$

for the masses of standard quarks, where up- and down-type quarks are labelled by Greek and Latin indices respectively. Note that, to first order in v/M_Q , the physical Yukawa couplings defined above as $y_{\alpha,i} = m_{\alpha,i}/v$ coincide with the elements $\hat{y}_{\alpha,i}$ of the diagonal matrices \hat{Y}_q and with the couplings of standard quarks to the Higgs boson.

4.1.1 The RH sector

An explicit parameterization for the down-sector RH rotation reads

$$\mathcal{V}_R^d \equiv \begin{pmatrix} \mathcal{V}_{1d}^R & \mathcal{V}_{1s}^R & \mathcal{V}_{1b}^R & \mathcal{V}_{1B'}^R \\ \mathcal{V}_{2d}^R & \mathcal{V}_{2s}^R & \mathcal{V}_{2b}^R & \mathcal{V}_{2B'}^R \\ \mathcal{V}_{3d}^R & \mathcal{V}_{3s}^R & \mathcal{V}_{3b}^R & \mathcal{V}_{3B'}^R \\ \mathcal{V}_{Bd}^R & \mathcal{V}_{Bs}^R & \mathcal{V}_{Bb}^R & \mathcal{V}_{BB'}^R \end{pmatrix} = \begin{pmatrix} \hat{U}_{dR} & & & \\ & & & \\ & & & \\ & & & 1 \end{pmatrix} \begin{pmatrix} \delta U_{dR} & & & \\ & & & \\ & & & \\ & & & 1 \end{pmatrix} \times \underbrace{\begin{pmatrix} c_{R1}^d & & -\tilde{s}_{R1}^{d*} & \\ & 1 & & \\ & & 1 & \\ \tilde{s}_{R1}^d & & & c_{R1}^d \end{pmatrix} \begin{pmatrix} 1 & & & \\ & c_{R2}^d & & -\tilde{s}_{R2}^{d*} \\ & & 1 & \\ & \tilde{s}_{R2}^d & & c_{R2}^d \end{pmatrix} \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & c_{R3}^d & -\tilde{s}_{R3}^{d*} \\ & & \tilde{s}_{R3}^d & c_{R3}^d \end{pmatrix}}_{R_4^{dR}}, \quad (4.11)$$

resulting in

$$R_4^{dR} = \begin{pmatrix} c_{R1}^d & -\tilde{s}_{R2}^d \tilde{s}_{R1}^{d*} & -c_{R2}^d \tilde{s}_{R3}^d \tilde{s}_{R1}^{d*} & -\tilde{s}_{R1}^{d*} c_{R2}^d c_{R3}^d \\ 0 & c_{R2}^d & -\tilde{s}_{R3}^d \tilde{s}_{R2}^{d*} & -\tilde{s}_{R2}^{d*} c_{R3}^d \\ 0 & 0 & c_{R3}^d & -\tilde{s}_{R3}^{d*} \\ \tilde{s}_{R1}^d & \tilde{s}_{R2}^d c_{R1}^d & \tilde{s}_{R3}^d c_{R1}^d c_{R2}^d & c_{R1}^d c_{R2}^d c_{R3}^d \end{pmatrix}, \quad (4.12)$$

and a similar one is valid for the up-type quarks. Here,

$$c_{Ri}^q \equiv \cos \theta_{Ri4}^q, \quad \tilde{s}_{Ri}^q = s_{Ri}^q e^{i\delta_{Ri}^q} \equiv \sin \theta_{Ri4}^q e^{i\delta_{Ri}^q} \quad (4.13)$$

are real cosines and complex sines of angles in the (1, 4), (2, 4), and (3, 4) family planes parameterizing the mixing of the first three families with the VLQs. We assume the analogous parameterization with real cosines and complex sines also for the other unitary matrices δU_{qR} , \hat{U}_{qR} .

As it is clear from eqs. (2.17) and (2.22), the RH charged currents and flavour changing couplings exclusively depend on the last rows of the matrices \mathcal{V}_R^u , \mathcal{V}_R^d , that is, on the mixings of SM quarks with the vector-like doublet. For couplings $\lesssim \mathcal{O}(1)$, these mixings are determined by the small parameter v/M_Q . In fact, up to terms of order v^2/M_Q^2 , one can write \mathcal{V}_R^d as

$$\mathcal{V}_R^d = \begin{pmatrix} \hat{U}_{dR} & & & \\ & & & \\ & & & \\ & & & 1 \end{pmatrix} \begin{pmatrix} \mathbb{1}_{3 \times 3} & z_d^\dagger \frac{v}{M_Q} \\ -z_d \frac{v}{M_Q} & 1 \end{pmatrix} + \mathcal{O}\left(\frac{v^2}{M_Q^2}\right), \quad (4.14)$$

where we have used the fact that $\delta U_{dR} \simeq \mathbb{1}$ at this order. A similar expression holds for the rotation \mathcal{V}_R^u of up-type quarks. Thus, the mixing elements of interest, corresponding to the last row of \mathcal{V}_R^d , are given by

$$\mathcal{V}_{Bd}^R = \tilde{s}_{R1}^d \simeq -z_d \frac{v}{M_Q}, \quad \mathcal{V}_{Bs}^R \simeq \tilde{s}_{R2}^d \simeq -z_s \frac{v}{M_Q}, \quad \mathcal{V}_{Bb}^R \simeq \tilde{s}_{R3}^d \simeq -z_b \frac{v}{M_Q}, \quad (4.15)$$

in first approximation. In particular, notice that the phases of these mixing elements exactly coincide with the phases of the complex sines. As is clear from eq. (4.15), these coincide with those of the extra Yukawa couplings z_d ,

$$\arg(\mathcal{V}_{Bi}^R) = \arg \tilde{s}_{Ri}^d (\equiv \delta_{Ri}^d) = \arg z_i, \quad (4.16)$$

at all orders (this result is also independent on the parameterization choice of the order of the product in R_4^{dR}). In other words, the higher order corrections to eq. (4.15) share the same phase as the leading term. The matrix $\delta U_{dR} \simeq \mathbb{1}$ contains tiny corrections given by small complex sines

$$\tilde{s}_{i<j} \simeq z_i z_j^* \frac{\hat{y}_i^2}{\hat{y}_j^2 - \hat{y}_i^2} \frac{v^2}{M_Q^2} \simeq z_i z_j^* \frac{y_i^2}{y_j^2} \frac{v^2}{M_Q^2} \quad (i, j = 1, 2, 3), \quad (4.17)$$

at leading order, where we have taken into account eq. (4.10) and the hierarchy between SM quark masses.⁸ A completely analogous discussion applies to up-type quarks.

Focusing on the last rows of \mathcal{V}_R^u and \mathcal{V}_R^d , one can find explicit expressions for the mixings up to order v^3/M_Q^3 . In the up sector, for instance, we have

$$\begin{aligned} \mathcal{V}_{T\alpha}^R &\simeq -z_\alpha \frac{v}{M_Q} + z_\alpha \left(\frac{1}{2} |z_\alpha|^2 - y_\alpha^2 + \sum_{\beta < \alpha} |z_\beta|^2 \right) \frac{v^3}{M_Q^3} + \mathcal{O} \left(\frac{v^5}{M_Q^5} \right), \\ \mathcal{V}_{T'T'}^R &= 1 - \frac{1}{2} |z_u|^2 \frac{v^2}{M_Q^2} + \mathcal{O} \left(\frac{v^4}{M_Q^4} \right), \end{aligned} \quad (4.18)$$

with $\alpha = u, c, t$ and where we have once again taken into account the hierarchies between standard quark masses. Similar expressions hold for down-type quarks. Given the above, one can construct an approximation for the non-unitary mixing matrix in the RH charged current. At leading order, it reads

$$\begin{aligned} V_R &= \mathcal{V}_R^{u\dagger} \text{diag}(0, 0, 0, 1) \mathcal{V}_R^d = \begin{pmatrix} V_{ud}^R & V_{us}^R & V_{ub}^R & V_{uB'}^R \\ V_{cd}^R & V_{cs}^R & V_{cb}^R & V_{cB'}^R \\ V_{td}^R & V_{ts}^R & V_{tb}^R & V_{tB'}^R \\ V_{T'd}^R & V_{T's}^R & V_{T'b}^R & V_{T'B'}^R \end{pmatrix} \\ &= \begin{pmatrix} z_u^* z_d \frac{v^2}{M_Q^2} & z_u^* z_s \frac{v^2}{M_Q^2} & z_u^* z_b \frac{v^2}{M_Q^2} & -z_u^* \frac{v}{M_Q} \\ z_c^* z_d \frac{v^2}{M_Q^2} & z_c^* z_s \frac{v^2}{M_Q^2} & z_c^* z_b \frac{v^2}{M_Q^2} & -z_c^* \frac{v}{M_Q} \\ z_t^* z_d \frac{v^2}{M_Q^2} & z_t^* z_s \frac{v^2}{M_Q^2} & z_t^* z_b \frac{v^2}{M_Q^2} & -z_t^* \frac{v}{M_Q} \\ -z_d \frac{v}{M_Q} & -z_s \frac{v}{M_Q} & -z_b \frac{v}{M_Q} & 1 - \frac{1}{2} (|z_u|^2 + |z_d|^2) \frac{v^2}{M_Q^2} \end{pmatrix} + \mathcal{O} \left(\frac{v^3}{M_Q^3} \right). \end{aligned} \quad (4.19)$$

⁸With a slight abuse of notation, it should be clear that $z_i = (z_d)_i$ when $i = 1, 2, 3$.

Similarly, one can construct the matrices of RH FCNC couplings. In the up sector, $F_u = \mathcal{V}_R^{u\dagger} \text{diag}(0, 0, 0, 1) \mathcal{V}_R^u$ explicitly reads

$$F_u = \begin{pmatrix} |z_u|^2 \frac{v^2}{M_Q^2} & z_u^* z_c \frac{v^2}{M_Q^2} & z_u^* z_t \frac{v^2}{M_Q^2} & -z_u^* \frac{v}{M_Q} \\ z_c^* z_u \frac{v^2}{M_Q^2} & |z_c|^2 \frac{v^2}{M_Q^2} & z_c^* z_t \frac{v^2}{M_Q^2} & -z_c^* \frac{v}{M_Q} \\ z_t^* z_u \frac{v^2}{M_Q^2} & z_t^* z_c \frac{v^2}{M_Q^2} & |z_t|^2 \frac{v^2}{M_Q^2} & -z_t^* \frac{v}{M_Q} \\ -z_u \frac{v}{M_Q} & -z_c \frac{v}{M_Q} & -z_t \frac{v}{M_Q} & 1 - |z_u|^2 \frac{v^2}{M_Q^2} \end{pmatrix} + \mathcal{O}\left(\frac{v^3}{M_Q^3}\right), \quad (4.20)$$

while the down sector $F_d = \mathcal{V}_R^{d\dagger} \text{diag}(0, 0, 0, 1) \mathcal{V}_R^d$ has an analogous structure.

4.1.2 The LH sector

Regarding the LH sector, recall that, since the four quark species make up $SU(2)_L$ doublets, the mixing matrix $V_L = \mathcal{V}_L^{u\dagger} \mathcal{V}_L^d$ is a unitary matrix. The extra elements in the \mathcal{V}_L^q describe the mixing of SM quarks with the VLQ doublets. These mixings can be parameterized analogously to what was done in eq. (4.12), with the substitution $R \rightarrow L$. The corresponding mixing angles are suppressed by the ratio v^2/M_Q^2 , and are associated to complex sines which at leading order read $\tilde{s}_{Li} \simeq -y_i z_i v^2/M_Q^2$ in the down sector ($i \rightarrow \alpha$ in the up sector).

Focusing on the last rows of \mathcal{V}_L^u and \mathcal{V}_L^d , one can once again find explicit expressions for the mixing matrix elements. In the up sector, for instance, we have

$$\begin{aligned} \mathcal{V}_{T\alpha}^L &\simeq -z_\alpha y_\alpha \frac{v^2}{M_Q^2} - z_\alpha y_\alpha \left(y_\alpha^2 - \frac{1}{2}|z_\alpha|^2 - \sum_{\beta < \alpha} |z_\beta|^2 \right) \frac{v^4}{M_Q^4} + \mathcal{O}\left(\frac{v^6}{M_Q^6}\right), \\ \mathcal{V}_{TT'}^L &= 1 - \frac{1}{2} \sum_{\beta} |z_\beta|^2 y_\beta^2 \frac{v^4}{M_Q^4} + \mathcal{O}\left(\frac{v^6}{M_Q^6}\right), \end{aligned} \quad (4.21)$$

with $\alpha = u, c, t$ and where we took into account the hierarchy between SM quark masses in writing terms of order v^4/M_Q^4 in the first line. At this order, the last row and last columns of the matrix R_4^{uL} in the parameterization of \mathcal{V}_L^u are related by $(R_4^{uL})_{4i} \simeq -(R_4^{uL})_{i4}^*$ to a good approximation, as one can anticipate from eq. (4.12). Using the fact that δU_{uL} deviates from $\mathbf{1}$ by small rotations with complex sines given by $\tilde{s}_{i < j} \simeq z_i z_j^* \frac{y_i}{y_j} \frac{v^2}{M_Q^2}$ ($i, j = 1, 2, 3$) at leading order, one finds

$$\mathcal{V}_{kT'}^L \simeq \sum_{\alpha=u,c,t} (\hat{U}_{uL})_{k\alpha} \left[z_\alpha^* y_\alpha \frac{v^2}{M_Q^2} + z_\alpha^* y_\alpha \left(y_\alpha^2 + \frac{1}{2}|z_\alpha|^2 - |z_u|^2 \right) \frac{v^4}{M_Q^4} \right] + \mathcal{O}\left(\frac{v^6}{M_Q^6}\right), \quad (4.22)$$

where $k = 1, 2, 3$. Similar expressions hold for down-type quarks. Keeping only terms

up to order v^2/M_Q^2 , the mixing matrix in the LH charged currents reads

$$\begin{aligned}
V_L &= \begin{pmatrix} c_{L1}^u & 0 & 0 & \tilde{s}_{L1}^{u*} \\ -\tilde{s}_{L2}^{u*}\tilde{s}_{L1}^u & c_{L2}^u & 0 & \tilde{s}_{L2}^{u*}c_{L1}^u \\ -c_{L2}^u\tilde{s}_{L3}^*\tilde{s}_{L1}^u & -\tilde{s}_{L3}^{u*}\tilde{s}_{L2}^u & c_{L3}^u & \tilde{s}_{L3}^{u*}c_{L1}^u c_{L2}^u \\ -\tilde{s}_{L1}^u c_{L2}^u c_{L3}^u & -\tilde{s}_{L2}^u c_{L3}^u & -\tilde{s}_{L3}^u & c_{L1}^u c_{L2}^u c_{L3}^u \end{pmatrix} \begin{pmatrix} \hat{V}_L & \\ & 1 \end{pmatrix} \begin{pmatrix} c_{L1}^d & -\tilde{s}_{L2}^d \tilde{s}_{L1}^{d*} & -c_{L2}^d \tilde{s}_{L3}^d \tilde{s}_{L1}^{d*} & -\tilde{s}_{L1}^{d*} c_{L2}^d c_{L3}^d \\ 0 & c_{L2}^d & -\tilde{s}_{L3}^d \tilde{s}_{L2}^{d*} & -\tilde{s}_{L2}^{d*} c_{L3}^d \\ 0 & 0 & c_{L3}^d & -\tilde{s}_{L3}^{d*} \\ \tilde{s}_{L1}^d & \tilde{s}_{L2}^d c_{L1}^d & \tilde{s}_{L3}^d c_{L1}^d c_{L2}^d & c_{L1}^d c_{L2}^d c_{L3}^d \end{pmatrix} \\
&\simeq \begin{pmatrix} \mathbb{1}_{3 \times 3} & -\hat{Y}_u z_u^\dagger \frac{v^2}{M_Q^2} \\ z_u \hat{Y}_u \frac{v^2}{M_Q^2} & 1 \end{pmatrix} \begin{pmatrix} \hat{V}_L & \\ & 1 \end{pmatrix} \begin{pmatrix} \mathbb{1}_{3 \times 3} & \hat{Y}_d z_d^\dagger \frac{v^2}{M_Q^2} \\ -z_d \hat{Y}_d \frac{v^2}{M_Q^2} & 1 \end{pmatrix} + \mathcal{O}\left(\frac{v^4}{M_Q^4}\right), \tag{4.23}
\end{aligned}$$

where $\hat{V}_L \equiv \delta U_{uL}^\dagger \hat{U}_{uL}^\dagger \hat{U}_{dL} \delta U_{dL}$. Here, c_{Li}^q and \tilde{s}_{Li}^q are the cosines and complex sines which make up the R_4^{qL} matrices, defined as in the RH sector. For the last row and column of V_L , one has

$$\begin{aligned}
V_{T'i}^L &= \left(\sum_\alpha z_\alpha y_\alpha \hat{V}_{\alpha i}^L - z_i y_i \right) \frac{v^2}{M_Q^2} + \mathcal{O}\left(\frac{v^4}{M_Q^4}\right), \\
V_{\alpha B'}^L &= \left(\sum_i \hat{V}_{\alpha i}^L z_i^* y_i - z_\alpha^* y_\alpha \right) \frac{v^2}{M_Q^2} + \mathcal{O}\left(\frac{v^4}{M_Q^4}\right), \tag{4.24}
\end{aligned}$$

where $i = d, s, b$ and $\alpha = u, c, t$. At this order, $V_{T'B'}^L \simeq 1$.

The matrix \hat{V}_L introduced in eq. (4.23) can be approximately obtained from the rotations defined in eq. (4.5), namely $\hat{V}_L \simeq \hat{U}_{uL}^\dagger \hat{U}_{dL}$, up to small extra rotations:

$$\hat{V}_{\alpha i}^L \simeq \left[\hat{U}_{uL}^\dagger \hat{U}_{dL} \right]_{\alpha i} + \left(\sum_{j \neq i} z_i z_j^* \frac{y_i y_j}{y_j^2 - y_i^2} \left[\hat{U}_{uL}^\dagger \hat{U}_{dL} \right]_{\alpha j} + \sum_{\beta \neq \alpha} z_\alpha^* z_\beta \frac{y_\alpha y_\beta}{y_\beta^2 - y_\alpha^2} \left[\hat{U}_{uL}^\dagger \hat{U}_{dL} \right]_{\beta i} \right) \frac{v^2}{M_Q^2}, \tag{4.25}$$

with $\alpha, \beta = u, c, t$ and $i, j = d, s, b$. Indeed, aside from the v^2/M_Q^2 suppression, one can see that these corrections are always additionally suppressed by ratios of quark masses of different generations and FCNC constraints (note that, for instance, $F_{ij}^d \simeq z_i^* z_j v^2/M_Q^2$). Recall that $\hat{U}_{uL}^\dagger \hat{U}_{dL}$ is actually the 3×3 CKM matrix in the limit of a decoupled VLQ doublet. Moreover, \hat{V}_L is a unitary 3×3 matrix by definition — the corrections in eq. (4.25) are induced by the unitary transformations δU_{uL} and δU_{dL} between the three SM quarks. One can also see that, to a very good approximation, \hat{V}_L corresponds to the 3×3 upper-left submatrix of V_L , i.e.

$$V_{\alpha i}^L = \hat{V}_{\alpha i}^L + \mathcal{O}\left(y^2 z^2 \frac{v^4}{M_Q^4}\right), \tag{4.26}$$

for $\alpha = u, c, t$ and $i = d, s, b$. In summary, we have schematically

$$V_L = \begin{pmatrix} V_{ud}^L & V_{us}^L & V_{ub}^L & V_{uB'}^L \\ V_{cd}^L & V_{cs}^L & V_{cb}^L & V_{cB'}^L \\ V_{td}^L & V_{ts}^L & V_{tb}^L & V_{tB'}^L \\ V_{T'd}^L & V_{T's}^L & V_{T'b}^L & V_{T'B'}^L \end{pmatrix} \simeq \begin{pmatrix} \hat{U}_{uL}^\dagger \hat{U}_{dL} \simeq V_{\text{CKM}} & \mathcal{O}\left(yz \frac{v^2}{M_Q^2}\right) \\ \hline \mathcal{O}\left(yz \frac{v^2}{M_Q^2}\right) & 1 \end{pmatrix} + \mathcal{O}\left(\frac{v^4}{M_Q^4}, \frac{m}{m'} \frac{v^2}{M_Q^2}\right), \quad (4.27)$$

at leading order, where m/m' indicates mass ratios of different-generation quarks. The explicit form of the $\mathcal{O}(yzv^2/M_Q^2)$ off-diagonal blocks can be found in eq. (4.24).

4.1.3 Effective rephasing invariants

Let us note that the SM quark fields can be rephased in order to absorb 5 phases in \hat{V}_L , e.g. one row and one column of \hat{V}_L can always be made real. Hence, \hat{V}_L can be parameterized by three angles and one phase, as in the usual CKM parameterization. When rephasing the LH quark fields, the corresponding RH fields must go through the same phase transformation in order to keep the diagonal Yukawa terms real. The effect of these transformations (e.g. $u_{L,R} \rightarrow u_{L,R} e^{i\delta_u}$) can be absorbed in the R_4 matrices, and corresponds to changing the phases of the complex sines, i.e. of the Yukawa couplings z_α and z_i (for instance as $s_{R,L1}^u \rightarrow s_{R,L1}^u e^{i\delta_u}$ and $z_u \rightarrow z_u e^{i\delta_u}$), where we have used the fact that relations analogous to eq. (4.16) hold for the LH sector. One can then readily identify the following rephasing-invariant quantities:

$$z_\alpha z_i^* \hat{V}_{\alpha i}^L, \quad z_\alpha z_\beta^* \hat{V}_{\alpha i}^L \hat{V}_{\beta i}^{L*}, \quad z_i^* z_j \hat{V}_{\alpha i}^L \hat{V}_{\alpha j}^{L*}, \quad z_\alpha z_j^* \hat{V}_{\beta i}^{L*} \hat{V}_{\alpha i}^L \hat{V}_{\beta j}^L, \quad (4.28)$$

with $\alpha = u, c, t$ and $i = d, s, b$. These quantities are related to the bilinears, trilinears and quartets described in section 3.3. In fact, let us define⁹

$$\begin{aligned} \hat{B}_{\alpha i} &\equiv \frac{v^2}{M_Q^2} z_\alpha^* z_i \hat{V}_{\alpha i}^{L*} \equiv \hat{V}_{\alpha i}^R \hat{V}_{\alpha i}^{L*}, \\ \hat{T}_{i, \alpha\beta} &\equiv \frac{v^2}{M_Q^2} z_\alpha^* z_\beta \hat{V}_{\alpha i}^{L*} \hat{V}_{\beta i}^L \equiv \hat{F}_{\alpha\beta}^u \hat{V}_{\alpha i}^{L*} \hat{V}_{\beta i}^L, \\ \hat{T}_{\alpha, ij} &\equiv \frac{v^2}{M_Q^2} z_i^* z_j \hat{V}_{\alpha i}^L \hat{V}_{\alpha j}^{L*} \equiv \hat{F}_{ij}^d \hat{V}_{\alpha i}^L \hat{V}_{\alpha j}^{L*}, \\ \hat{Q}_{\alpha i\beta j} &\equiv \frac{v^2}{M_Q^2} z_\alpha^* z_i \hat{V}_{\beta j}^L \hat{V}_{\alpha j}^{L*} \hat{V}_{\beta i}^{L*} \equiv \hat{V}_{\alpha i}^R \hat{V}_{\beta j}^L \hat{V}_{\alpha j}^{L*} \hat{V}_{\beta i}^{L*}. \end{aligned} \quad (4.29)$$

⁹ The quantities $\hat{V}_{\alpha i}^R \equiv \frac{v^2}{M_Q^2} z_\alpha^* z_i$, $\hat{F}_{\alpha\beta}^u \equiv \frac{v^2}{M_Q^2} z_\alpha^* z_\beta$, and $\hat{F}_{ij}^d \equiv \frac{v^2}{M_Q^2} z_i^* z_j$, defined here for $N = 1$, are useful shorthands used in the following sections (and generalized in sections 7 and 8 for $N > 1$). To leading order in v^2/M_Q^2 , they approximate the 3×3 upper-left submatrices of V_R , F_u , and F_d , respectively.

While these do not exactly coincide with the previously-defined unhatted versions of section 3.3 — consider eqs. (4.18), (4.19) and (4.26) — they differ from them by rephasing-invariant quantities. Moreover, they are approximately equal at leading order, i.e.

$$\begin{aligned}
\hat{\mathcal{B}}_{\alpha i} &\simeq V_{\alpha i}^R V_{\alpha i}^{L*} = \mathcal{B}_{\alpha i}^{RL}, \\
\hat{\mathcal{T}}_{i,\alpha\beta} &\simeq F_{\alpha\beta}^u V_{\alpha i}^{L*} V_{\beta i}^L = \mathcal{T}_{\alpha\beta,i}^{LL}, \\
\hat{\mathcal{T}}_{\alpha,ij} &\simeq F_{ij}^d V_{\alpha i}^L V_{\alpha j}^{L*} = \mathcal{T}_{ij,\alpha}^{LL}, \\
\hat{\mathcal{Q}}_{\alpha i\beta j} &\simeq V_{\alpha i}^R V_{\beta j}^L V_{\alpha j}^{L*} V_{\beta i}^{L*} = \mathcal{Q}_{\alpha i\beta j}^{RLLL},
\end{aligned} \tag{4.30}$$

meaning that for the purposes of phenomenology (see section 8) one may focus on the hatted rephasing invariants. We also define the non-calligraphic hatted invariants

$$\hat{\mathcal{Q}}_{\alpha i\beta j} \equiv \hat{V}_{\alpha i}^L \hat{V}_{\beta j}^L \hat{V}_{\alpha j}^{L*} \hat{V}_{\beta i}^{L*} \simeq V_{\alpha i}^L V_{\beta j}^L V_{\alpha j}^{L*} V_{\beta i}^{L*} = \mathcal{Q}_{\alpha i\beta j}^{LLLL}, \tag{4.31}$$

which are the practical generalization of the SM quartets ($\hat{V}_{\alpha i}^L = V_{\alpha i}^L$ in the SM limit).

Let us also notice that in presence of only one doublet, if $z_\alpha \neq 0$ and $z_i \neq 0$ the trilinears and quartets can be written as

$$\begin{aligned}
\hat{\mathcal{T}}_{i,\alpha\beta} &= \frac{\hat{\mathcal{B}}_{\alpha i} \hat{\mathcal{B}}_{\beta i}^*}{\frac{v^2}{M_Q^2} |z_i|^2} \simeq \frac{\hat{\mathcal{B}}_{\alpha i} \hat{\mathcal{B}}_{\beta i}^*}{F_{ii}^d}, \\
\hat{\mathcal{T}}_{\alpha,ij} &= \frac{\hat{\mathcal{B}}_{\alpha i}^* \hat{\mathcal{B}}_{\alpha j}}{\frac{v^2}{M_Q^2} |z_\alpha|^2} \simeq \frac{\hat{\mathcal{B}}_{\alpha i}^* \hat{\mathcal{B}}_{\alpha j}}{F_{\alpha\alpha}^u}, \\
\hat{\mathcal{Q}}_{\alpha i\beta j} &= \frac{\hat{\mathcal{T}}_{\beta,ij}^* \hat{\mathcal{B}}_{\alpha j}}{\frac{v^2}{M_Q^2} |z_j|^2} = \frac{\hat{\mathcal{T}}_{j,\alpha\beta} \hat{\mathcal{B}}_{\beta i}}{\frac{v^2}{M_Q^2} |z_\beta|^2} = \frac{\hat{\mathcal{B}}_{\alpha j} \hat{\mathcal{B}}_{\beta j}^* \hat{\mathcal{B}}_{\beta i}}{\frac{v^4}{M_Q^4} |z_j|^2 |z_\beta|^2} \simeq \frac{\hat{\mathcal{B}}_{\alpha j} \hat{\mathcal{B}}_{\beta j}^* \hat{\mathcal{B}}_{\beta i}}{F_{jj}^d F_{\beta\beta}^u}, \\
\hat{\mathcal{Q}}_{\alpha i\beta j} &= \frac{\hat{\mathcal{Q}}_{\alpha i\beta j} \hat{\mathcal{B}}_{\alpha i}^*}{\frac{v^4}{M_Q^4} |z_\alpha|^2 |z_i|^2} = \frac{\hat{\mathcal{B}}_{\alpha j} \hat{\mathcal{B}}_{\beta j}^* \hat{\mathcal{B}}_{\beta i} \hat{\mathcal{B}}_{\alpha i}^*}{\frac{v^8}{M_Q^8} |z_j|^2 |z_\beta|^2 |z_\alpha|^2 |z_i|^2} \simeq \frac{\hat{\mathcal{B}}_{\alpha j} \hat{\mathcal{B}}_{\beta j}^* \hat{\mathcal{B}}_{\beta i} \hat{\mathcal{B}}_{\alpha i}^*}{|F_{ij}^d|^2 |F_{\alpha\beta}^u|^2}.
\end{aligned} \tag{4.32}$$

Thus, the physical phases may be identified as the six independent phases of the bilinears. These relations do not hold in scenarios with vanishing couplings, e.g. when the vector-like doublet is coupling only to one sector (up or down). In these cases, the phases of the trilinears and quartets are independent CP-violating phases¹⁰ (see section 6 for further details). This correspondence is strictly related to the type of WBIs needed to describe CP violation.

We also have the rephasing-invariant elements

$$\hat{F}_{\alpha\alpha}^u = \frac{v^2}{M_Q^2} |z_\alpha|^2 \simeq F_{\alpha\alpha}^u, \quad \hat{F}_{ii}^d = \frac{v^2}{M_Q^2} |z_i|^2 \simeq F_{ii}^d, \tag{4.33}$$

together with the other CP-even quantities which are explicitly moduli ($|\hat{V}_{\alpha i}^L|$, $|\hat{V}_{\alpha i}^R|$, etc.).

¹⁰It is interesting to note that the SM limit can be regarded as such a singular case, where only the quartet phase is relevant for CP violation.

4.2 Parameterizing the mixing exactly

In this section, we derive exact parameterizations for the mixings in these extensions, namely for the matrices V_L , V_R and consequently for the F_q , for an arbitrary number of VLQ doublets N with hypercharge $1/6$. These are given explicitly in terms of (physical) angles and phases, in a way that is reminiscent of the PDG parameterization for the standard CKM matrix [67,69] or the Botella-Chau parameterization [70] for the mixing in SM extensions with a fourth generation. While this is done for completeness and may be useful for comprehensive scans of the parameter space of the model(s), an expansion like the one considered in the previous section should be enough for most phenomenological applications.

Throughout this section we will make use of the following definitions for the $(3 + N) \times (3 + N)$ orthogonal rotation matrices \mathcal{O}_{ij} ,

$$(\mathcal{O}_{ij})_{\alpha\beta} \equiv \begin{cases} \sin \theta_{ij}, & \text{for } \alpha = i, \beta = j, \\ -\sin \theta_{ij}, & \text{for } \alpha = j, \beta = i, \\ \cos \theta_{ij}, & \text{for } \alpha = \beta = i \text{ or } j, \\ \delta_{\alpha\beta}, & \text{otherwise,} \end{cases} \quad (4.34)$$

with $\theta_{ij} \in [0, \pi/2]$, and for the $(3 + N) \times (3 + N)$ diagonal phase matrices \mathcal{K}_{ij} ,

$$(\mathcal{K}_{ij})_{\alpha\beta} \equiv \begin{cases} e^{i\varphi_{ij}}, & \text{for } \alpha = \beta = i, \\ \delta_{\alpha\beta}, & \text{otherwise,} \end{cases} \quad (4.35)$$

with $\varphi_{ij} \in [0, 2\pi]$, implying e.g. that $\mathcal{K}_{31} = \text{diag}(1, 1, e^{i\varphi_{31}}, 1)$ and that $\mathcal{K}_{24}\mathcal{K}_{14} = \text{diag}(e^{i\varphi_{14}}, e^{i\varphi_{24}}, 1, 1)$, for $N = 1$.

4.2.1 A single doublet ($N = 1$)

For the case of the SM extended with a single doublet VLQ we can parameterize the unitary mixing matrix of the charged LH currents as a general 4×4 unitary matrix

$$V_L = \mathcal{K} \mathcal{O}_{34} \mathcal{O}_{24} \mathcal{O}_{14} \mathcal{K}_{24} \mathcal{K}_{14} \mathcal{O}_{23} \mathcal{K}_{31} \mathcal{O}_{13} \mathcal{O}_{12} \mathcal{K}', \quad (4.36)$$

where we have used the 4×4 matrices \mathcal{O}_{ij} and \mathcal{K}_{ij} defined above, as well as the diagonal phase matrices given by

$$\begin{aligned} \mathcal{K} &= \text{diag}(e^{i\varphi_1}, e^{i\varphi_2}, e^{i\varphi_3}, e^{i\varphi_4}), \\ \mathcal{K}' &= \text{diag}(1, e^{i\varphi'_2}, e^{i\varphi'_3}, e^{i\varphi'_4}). \end{aligned} \quad (4.37)$$

Obtaining a parameterization for V_R is less immediate due to its non-unitarity. Nonetheless, it is helpful to recall that V_R is given by the product

$$V_R = \mathcal{V}_R^{u\dagger} \text{diag}(0, 0, 0, 1) \mathcal{V}_R^d, \quad (4.38)$$

where the \mathcal{V}_R^q are 4×4 unitary matrices. Then, using again the general parameterization for a unitary matrix in eq. (4.36) to express both \mathcal{V}_R^u and \mathcal{V}_R^d ,¹¹ one can write

$$V_R = \mathcal{K}_u^* \mathcal{O}_{34}^{uT} \mathcal{O}_{24}^{uT} \mathcal{O}_{14}^{uT} \text{diag}(0, 0, 0, 1) \mathcal{O}_{14}^d \mathcal{O}_{24}^d \mathcal{O}_{34}^d \mathcal{K}_d, \quad (4.39)$$

so that the matrices controlling the FCNCs are given by

$$F_q = \mathcal{K}_q^* \mathcal{O}_{34}^{qT} \mathcal{O}_{24}^{qT} \mathcal{O}_{14}^{qT} \text{diag}(0, 0, 0, 1) \mathcal{O}_{14}^q \mathcal{O}_{24}^q \mathcal{O}_{34}^q \mathcal{K}_q, \quad (4.40)$$

for $q = u, d$. Here, the \mathcal{O}_{ij}^q matrices are 4×4 orthogonal matrices as defined in eq. (4.34) and the \mathcal{K}_u and \mathcal{K}_d matrices are diagonal phase matrices similar to \mathcal{K} in eq. (4.37).

Now, by rephasing the LH fields as $\mathcal{U}_L \rightarrow \mathcal{K} \mathcal{U}_L$ and $\mathcal{D}_L \rightarrow \mathcal{K}'^* \mathcal{D}_L$, followed by an identical transformation of the RH fields, i.e. $\mathcal{U}_R \rightarrow \mathcal{K} \mathcal{U}_R$ and $\mathcal{D}_R \rightarrow \mathcal{K}'^* \mathcal{D}_R$ which maintains the mass terms in eq. (2.9) invariant, one can simply write

$$V_L = \mathcal{O}_{34} \mathcal{O}_{24} \mathcal{O}_{14} \mathcal{K}_{24} \mathcal{K}_{14} \mathcal{O}_{23} \mathcal{K}_{31} \mathcal{O}_{13} \mathcal{O}_{12}, \quad (4.41)$$

which has a form identical to the Botella-Chau parameterization of the quark mixing matrix for the case of the SM extended with one VLQ isosinglet. On the other hand, V_R maintains the form in eq. (4.39), of course with \mathcal{K}_u and \mathcal{K}_d being redefined to absorb the effects of the rephasing

$$\mathcal{K}_u \mathcal{K} \rightarrow \mathcal{K}_u, \quad \mathcal{K}_d \mathcal{K}'^* \rightarrow \mathcal{K}_d, \quad (4.42)$$

so that all factorizable phases are brought to the RH charged currents (absorbed in V_R).

Note that, right now, we have parameterizations for V_L and V_R that appear to depend on a number of parameters that largely exceeds the 14 mixing parameters we counted in section 3, see Table 2. This issue is resolved when enforcing the constraint of eq. (2.26), which allowed us to derive eqs. (2.27) and (2.28). For instance by writing

$$B_R^q = (0 \ 0 \ 0 \ 1) \mathcal{V}_R^q = (0 \ 0 \ 0 \ 1) \mathcal{O}_{14}^q \mathcal{O}_{24}^q \mathcal{O}_{34}^q \mathcal{K}_q \quad (4.43)$$

and using eq. (3.23) which directly follows from eq. (2.26) for $N = 1$, one finds

$$\mathcal{D}_u \mathcal{K}_u^* \mathcal{O}_{34}^{uT} \mathcal{O}_{24}^{uT} \mathcal{O}_{14}^{uT} \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} = V_L \mathcal{D}_d \mathcal{K}_d^* \mathcal{O}_{34}^{dT} \mathcal{O}_{24}^{dT} \mathcal{O}_{14}^{dT} \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}, \quad (4.44)$$

that demonstrates the existence of relations between the (interdependent) parameters.

One can choose to keep the 6 angles and 3 phases of V_L as given in eq. (4.41) as independent parameters — different choices are possible, c.f. section 3.2.2. In this case, one can plug eq. (4.40) into the leftmost part of eq. (2.27) to obtain

$$V_R = \mathcal{D}_u^{-1} V_L \mathcal{D}_d \mathcal{K}_d^* \mathcal{O}_{34}^{dT} \mathcal{O}_{24}^{dT} \mathcal{O}_{14}^{dT} \text{diag}(0, 0, 0, 1) \mathcal{O}_{14}^d \mathcal{O}_{24}^d \mathcal{O}_{34}^d \mathcal{K}_d, \quad (4.45)$$

¹¹Without loss of generality, we reverse the order of the product in eq. (4.36) to parameterize the \mathcal{V}_R^q .

after multiplying by \mathcal{D}_u^{-1} from the left.¹² Here, one can parameterize the phase matrix \mathcal{K}_d as

$$\mathcal{K}_d = \text{diag} \left(e^{i\varphi_1^d}, e^{i\varphi_2^d}, e^{i\varphi_3^d}, 1 \right), \quad (4.46)$$

with the fourth phase being factorized out as a global phase, which cancels with that from \mathcal{K}_d^* . Note that any mention of the \mathcal{O}_{i4}^u ($i = 1, 2, 3$) and \mathcal{K}_u matrices has disappeared in this parameterization of V_R , and consequently from $F_u = V_R V_R^\dagger$ and $F_d = V_R^\dagger V_R$. Finally, using the relation $B_R^u B_R^{u\dagger} = 1$, see eq. (2.14), and recalling that $B_R^{u\dagger} = \mathcal{D}_u^{-1} V_L \mathcal{D}_d B_R^{d\dagger}$, one finds an extra constraint on B_R^d and V_L . Taking into account eq. (4.43), this constraint can be expressed as

$$\left[\mathcal{O}_{14}^d \mathcal{O}_{24}^d \mathcal{O}_{34}^d \mathcal{K}_d \mathcal{D}_d V_L^\dagger \mathcal{D}_u^{-2} V_L \mathcal{D}_d \mathcal{K}_d^* \mathcal{O}_{34}^{dT} \mathcal{O}_{24}^{dT} \mathcal{O}_{14}^{dT} \right]_{44} = 1. \quad (4.47)$$

Since the right-hand side of this equation is trivially real, this relation will correspond to one single condition that has to be met by the mixing parameters. This allows us to eliminate, for instance, the dependence on the mixing angle in \mathcal{O}_{14}^d . In summary, the parameters of the theory can be chosen as:

- the 8 quark masses,
- the 6 mixing angles ($\theta_{12}, \theta_{13}, \theta_{23}, \theta_{14}, \theta_{24}$ and θ_{34}) and 3 non-factorizable phases (δ_{13}, δ_{14} and δ_{24}) in V_L ,
- the 3 independent phases in \mathcal{K}_d and the 2 independent mixing angles present in the product $\mathcal{O}_{14}^d \mathcal{O}_{24}^d \mathcal{O}_{34}^d$.

Hence all the mixing is parameterized by 8 mixing angles and 6 phases, for a total of 14 physical mixing parameters, as anticipated in Table 2.

4.2.2 More than one doublet ($N > 1$)

In a more general extension of the SM with N doublets, one can parameterize the $(3+N) \times (3+N)$ unitary mixing matrix of the charged LH currents in a way analogous to that of eq. (4.36) for the $N = 1$ case (see also e.g. [29]). We have

$$V_L = \mathcal{U}_{3+N} \dots \mathcal{U}_5 \mathcal{U}_4 \mathcal{U}_3, \quad (4.48)$$

where the upper-left 3×3 block of the unitary matrix $\mathcal{U}_3 \equiv \mathcal{O}_{23} \mathcal{K}_{31} \mathcal{O}_{13} \mathcal{O}_{12}$ corresponds to the SM CKM matrix in the limit of decoupled VLQ doublets. The remaining \mathcal{U}_i ($i > 3$) matrices are also unitary and are given by

$$\mathcal{U}_i \equiv \left(\prod_{j=1}^{i-1} \mathcal{O}_{i-j,i} \right) \left(\prod_{k=2}^{i-1} \mathcal{K}_{i-k,i} \right) \quad (i = 4, \dots, 3+N), \quad (4.49)$$

¹²We do not consider cases where lighter masses vanish, which would have to be treated differently.

where the matrices \mathcal{O}_{ij} and \mathcal{K}_{ij} have been defined in eqs. (4.34) and (4.35).

For $N > 1$, eq. (2.27) still holds, implying V_R can be obtained from V_L and F_d via

$$V_R = \mathcal{D}_u^{-1} V_L \mathcal{D}_d F_d, \quad (4.50)$$

where

$$F_d = B_R^{d\dagger} B_R^d = \mathcal{V}_R^{d\dagger} \text{diag}(0, 0, 0, 1, \dots, 1) \mathcal{V}_R^d. \quad (4.51)$$

The matrix $B_R^{d\dagger}$ can be written as¹³

$$B_R^{d\dagger} = \mathcal{V}_R^{d\dagger} \begin{pmatrix} 0_{3 \times N} \\ \mathbb{1}_{N \times N} \end{pmatrix} = \begin{pmatrix} c & c & \cdots & c & c \\ c & c & \cdots & c & c \\ c & c & \cdots & c & c \\ r & c & \cdots & c & c \\ 0 & r & \cdots & c & c \\ 0 & 0 & \ddots & \vdots & \vdots \\ \vdots & \vdots & \ddots & r & c \\ 0 & 0 & \cdots & 0 & r \end{pmatrix} \mathcal{U}_{N \times N} \equiv \overline{B_R^d}^\dagger \mathcal{U}_{N \times N}, \quad (4.52)$$

with $\mathcal{U}_{N \times N}$ being an $N \times N$ unitary matrix and $\overline{B_R^d}^\dagger$ a $(3+N) \times N$ rectangular matrix. Here, each r represents a real entry while each c represents a complex entry. Note that, just like for $B_R^{d\dagger}$, the columns of $\overline{B_R^d}^\dagger$ are orthonormal. This is because

$$\overline{B_R^d} \overline{B_R^d}^\dagger = \mathcal{U}_{N \times N} B_R^d B_R^{d\dagger} \mathcal{U}_{N \times N}^\dagger = \mathbb{1}_{N \times N}. \quad (4.53)$$

At the same time, we also have

$$F_d = \overline{B_R^d}^\dagger B_R^d. \quad (4.54)$$

One can now decompose $\overline{B_R^d}^\dagger$ as

$$\overline{B_R^d}^\dagger = \mathcal{K}_{34}^{d*} \mathcal{K}_{24}^{d*} \mathcal{K}_{14}^{d*} \mathcal{O}_{34}^{dT} \mathcal{O}_{24}^{dT} \mathcal{O}_{14}^{dT} \begin{pmatrix} 0 & c & \cdots & c & c \\ 0 & c & \cdots & c & c \\ 0 & c & \cdots & c & c \\ r_{41} & c_{42} & \cdots & c_{4,N-1} & c_{4N} \\ 0 & r & \cdots & c & c \\ 0 & 0 & \ddots & \vdots & \vdots \\ \vdots & \vdots & \ddots & r & c \\ 0 & 0 & \cdots & 0 & r \end{pmatrix}, \quad (4.55)$$

¹³Note that any square matrix can be written as the product of an upper-triangular matrix and a unitary matrix (RQ decomposition). By decomposing the last N rows of $B_R^{d\dagger}$ in this way, one can extract the unitary $\mathcal{U}_{N \times N}$ to the right while keeping the 3 first rows generically complex. The diagonal elements can be made real by making an appropriate rephasing of each column, which is absorbed by the unitary matrix.

where \mathcal{O}_{ij}^d and \mathcal{K}_{ij}^d are also of the forms described in eqs. (4.34) and (4.35), respectively. From the orthonormality condition in eq. (4.53), one must have $r_{41} = 1$ and $c_{4i} = 0$, so that

$$\overline{B}_R^{d\dagger} = \mathcal{K}_{34}^{d*} \mathcal{K}_{24}^{d*} \mathcal{K}_{14}^{d*} \mathcal{O}_{34}^{dT} \mathcal{O}_{24}^{dT} \mathcal{O}_{14}^{dT} \begin{pmatrix} 0 & c & \cdots & c & c \\ 0 & c & \cdots & c & c \\ 0 & c & \cdots & c & c \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & r & \cdots & c & c \\ 0 & 0 & \ddots & \vdots & \vdots \\ \vdots & \vdots & \ddots & r & c \\ 0 & 0 & \cdots & 0 & r \end{pmatrix}. \quad (4.56)$$

Now, by repeating this process $N - 1$ more times, once for each remaining column of $\overline{B}_R^{d\dagger}$, one can arrive at the form

$$\overline{B}_R^{d\dagger} = \left(\prod_{i=4}^{N+3} \mathcal{K}_{3i}^{d*} \mathcal{K}_{2i}^{d*} \mathcal{K}_{1i}^{d*} \mathcal{O}_{3i}^{dT} \mathcal{O}_{2i}^{dT} \mathcal{O}_{1i}^{dT} \right) \begin{pmatrix} 0_{3 \times N} \\ \mathbf{1}_{N \times N} \end{pmatrix} \equiv \overline{\mathcal{V}}_R^{d\dagger} \begin{pmatrix} 0_{3 \times N} \\ \mathbf{1}_{N \times N} \end{pmatrix}. \quad (4.57)$$

Note that each time we apply this procedure, only three phases and three orthogonal rotations can be factorised from each column $j = 1, \dots, N$. After that, eq. (4.53) implies that $r_{3+j,j} = 1$ and $c_{3+j,k} = 0$ for all $k \neq j$. In the end, since one has

$$F_d = \overline{B}_R^{d\dagger} \overline{B}_R^d = \overline{\mathcal{V}}_R^{d\dagger} \text{diag}(0, 0, 0, 1, \dots, 1) \overline{\mathcal{V}}_R^d, \quad (4.58)$$

by plugging this expression for F_d in eq. (4.50), we obtain a parameterization for V_R in terms of a reduced number of parameters, which only grows linearly with N .

Finally, using the relation $B_R^u B_R^{u\dagger} = \mathbf{1}_{N \times N}$ and recalling that $B_R^{u\dagger} = \mathcal{D}_u^{-1} V_L \mathcal{D}_d B_R^{d\dagger}$ and $B_R^d = \mathcal{U}_{N \times N}^\dagger \overline{B}_R^d$, we have

$$\overline{B}_R^d \mathcal{D}_d V_L^\dagger \mathcal{D}_u^{-2} V_L \mathcal{D}_d \overline{B}_R^{d\dagger} = \mathbf{1}_{N \times N}, \quad (4.59)$$

or alternatively, given eq. (4.57),

$$(0_{N \times 3} \quad \mathbf{1}_{N \times N}) \mathcal{H} \begin{pmatrix} 0_{3 \times N} \\ \mathbf{1}_{N \times N} \end{pmatrix} = \mathbf{1}_{N \times N}, \quad (4.60)$$

where

$$\mathcal{H} \equiv \overline{\mathcal{V}}_R^{d\dagger} \mathcal{D}_d V_L \mathcal{D}_u^{-2} V_L^\dagger \mathcal{D}_d \overline{\mathcal{V}}_R^d, \quad (4.61)$$

which results in N^2 independent constraints for the mixing parameters. Of these, N can be expressed as

$$\mathcal{H}_{ii} = 1, \quad (4.62)$$

with $i = 4, \dots, 3 + N$, while the remaining $N(N - 1)$ constraints can be written as

$$\mathcal{H}_{ij} = 0, \quad (4.63)$$

with $i < j$. Note that since \mathcal{H} is Hermitian, eq. (4.62) corresponds to one constraint for each value of i , while eq. (4.63) corresponds to two distinct constraints for each pair (i, j) — one on a modulus and one on a phase.

A quick counting of parameters shows us that this method results in a minimal parameterization of the mixing in a general 1/6-hypercharge VLQ doublet model. For instance, to parameterize V_L we have:

- $(N + 3)(N + 2)/2$ mixing angles,
- $(N + 2)(N + 1)/2$ physical phases,

as can be inferred from eqs. (4.48) and (4.49), whereas to parameterize V_R we are additionally using:

- $3N$ mixing angles,
- $3N$ physical phases,

as can be seen from eqs. (4.50), (4.54) and (4.57). Moreover, the naïve quadratic dependence of the number of mixing parameters on the number of VLQ doublets is removed by taking into account the constraints in eqs. (4.62) and (4.63). In summary, our parameterization depends on

$$\frac{(N + 3)(N + 2)}{2} + \frac{(N + 2)(N + 1)}{2} + 6N - N^2 = 4 + 10N \quad (4.64)$$

independent mixing parameters ($3+5N$ angles and $1+5N$ phases), in agreement with Table 2. Note that when one accounts for the $6 + 2N$ quark masses of theory, one finds the total of $10 + 12N$ physical parameters we obtained before.

5 Weak-basis invariants

As mentioned in section 3, weak-basis-invariant quantities can be constructed in the quark sector from Hermitian combinations of the quark mass matrices. In this section we show how these quantities can be built in an explicit and systematic way and how they can be related to the physical quark masses and mixings. We then demonstrate how one can determine all parameters describing $N \geq 1$ VLQ doublet extensions in terms of weak-basis invariants (WBIs).

5.1 Constructing WBIs

To construct WBI quantities, it is useful to define the Hermitian squared mass matrices

$$\mathcal{H}_q \equiv \mathcal{M}_q \mathcal{M}_q^\dagger, \quad \mathcal{H}_{qR} \equiv \mathcal{M}_q^\dagger \mathcal{M}_q, \quad (5.1)$$

with $\mathcal{M}_q = (m_q \mid M)$, see eq. (2.6). Recall the Hermitian combinations defined in section 3, namely $h_q \equiv m_q m_q^\dagger$ and $H \equiv MM^\dagger$, which transform under weak-basis transformations (WBTs) as

$$h_q \rightarrow \mathcal{W}_L^\dagger h_q \mathcal{W}_L, \quad H \rightarrow \mathcal{W}_L^\dagger H \mathcal{W}_L. \quad (5.2)$$

Noting that $\mathcal{H}_q = h_q + H$, one sees that also \mathcal{H}_q transforms as

$$\mathcal{H}_q \rightarrow \mathcal{W}_L^\dagger \mathcal{H}_q \mathcal{W}_L \quad (5.3)$$

under WBTs. Since they transform in the same way under WBTs, combinations of the three matrices \mathcal{H}_q , h_q , H and of their powers can be used to construct WBI quantities. This can be done, for instance, by taking traces or determinants, since e.g. $\text{Tr}(\mathcal{W}_L^\dagger \mathcal{H}_q \mathcal{W}_L) = \text{Tr} \mathcal{H}_q$ and $\det(\mathcal{W}_L^\dagger \mathcal{H}_q \mathcal{W}_L) = \det \mathcal{H}_q$. Given that the three building blocks are related, not all combinations necessarily lead to different WBIs. All independent WBIs can then be built from traces of products of matrices involving only two of the three matrices \mathcal{H}_q , h_q , and H .

Let us note that, in the case of a single VLQ doublet ($N = 1$), some simplification is possible. Namely, since M is a column vector, $M^\dagger M$ is a number and $H = MM^\dagger$ is a rank-1 matrix. Then, $M^\dagger M = \text{Tr}(MM^\dagger)$, resulting in

$$H^2 = (\text{Tr} H) H, \quad \text{for } N = 1, \quad (5.4)$$

and thus reducing the actual degrees of freedom when building independent WBIs. Indeed, eq. (5.4) directly implies

$$\text{Tr}(AH^r) = (\text{Tr} H)^{r-1} \text{Tr}(AH), \quad \text{for } N = 1. \quad (5.5)$$

Moreover, since $M^\dagger AM$ is a number, equalling its own trace, one also finds that

$$\text{Tr}(HAHB) = \text{Tr}(HA) \text{Tr}(HB), \quad \text{for } N = 1. \quad (5.6)$$

The matrix \mathcal{H}_{qR} defined above can also be relevant when building WBIs. One has

$$\mathcal{H}_{qR} = \mathcal{M}_q^\dagger \mathcal{M}_q = \begin{pmatrix} m_q^\dagger m_q & m_q^\dagger M \\ M^\dagger m_q & M^\dagger M \end{pmatrix}, \quad (5.7)$$

which under a general WBT transforms as

$$\mathcal{H}_{qR} \rightarrow \begin{pmatrix} W_R^{q\dagger} & \\ & W_R^\dagger \end{pmatrix} \mathcal{H}_{qR} \begin{pmatrix} W_R^q & \\ & W_R \end{pmatrix} = \begin{pmatrix} W_R^{q\dagger} m_q^\dagger m_q W_R^q & W_R^{q\dagger} m_q^\dagger M W_R \\ W_R^\dagger M^\dagger m_q W_R^q & W_R^\dagger M^\dagger M W_R \end{pmatrix}. \quad (5.8)$$

Inspection of the matrix blocks motivates defining the 3×3 Hermitian matrices

$$\begin{aligned} h_{qR} &\equiv m_q^\dagger m_q && \rightarrow W_R^{q\dagger} h_{qR} W_R^q, \\ \bar{h}_{qR} &\equiv m_q^\dagger M M^\dagger m_q = m_q^\dagger H m_q && \rightarrow W_R^{q\dagger} \bar{h}_{qR} W_R^q, \end{aligned} \quad (5.9)$$

and the $N \times N$ Hermitian matrices

$$\begin{aligned} H_R &\equiv M^\dagger M && \rightarrow W_R^\dagger H_R W_R, \\ \bar{H}_{qR} &\equiv M^\dagger m_q m_q^\dagger M = M^\dagger h_q M && \rightarrow W_R^\dagger \bar{H}_{qR} W_R. \end{aligned} \quad (5.10)$$

These are also viable building blocks of WBIs, directly connected to the transformations of RH quark fields, which are physical in this beyond-the-Standard-Model (BSM) scenario. Within each pair, matrices share the same dimensions and transformation properties, so that their powers can be combined within traces to form two sets of independent WBIs. Note, however, that WBIs built from these four RH Hermitian matrices are not independent from the ones built from the LH Hermitian matrices, \mathcal{H}_q , h_q , and H , due to the cyclicity of the traces.¹⁴

Additionally, note that powers of \mathcal{H}_{qR} transform as ($k \geq 1$)

$$\mathcal{H}_{qR}^k \rightarrow \begin{pmatrix} W_R^{q\dagger} [m_q^\dagger \mathcal{H}_q^{k-1} m_q] W_R^q & W_R^{q\dagger} [m_q^\dagger \mathcal{H}_q^{k-1} M] W_R \\ W_R^\dagger [M^\dagger \mathcal{H}_q^{k-1} m_q] W_R^q & W_R^\dagger [M^\dagger \mathcal{H}_q^{k-1} M] W_R \end{pmatrix}, \quad (5.11)$$

so that one can generalize the blocks in eqs. (5.9) and (5.10) to

$$\begin{aligned} h_{qR}^{(k)} &\equiv m_q^\dagger \mathcal{H}_q^{k-1} m_q && \rightarrow W_R^{q\dagger} h_{qR}^{(k)} W_R^q, \\ \bar{h}_{qR}^{(k,l)} &\equiv m_q^\dagger \mathcal{H}_q^{k-1} H \mathcal{H}_q^{l-1} m_q && \rightarrow W_R^{q\dagger} \bar{h}_{qR}^{(k,l)} W_R^q, \end{aligned} \quad (5.12)$$

and

$$\begin{aligned} H_{qR}^{(k)} &\equiv M^\dagger \mathcal{H}_q^{k-1} M && \rightarrow W_R^\dagger H_{qR}^{(k)} W_R, \\ \bar{H}_{qR}^{(k,l)} &\equiv M^\dagger \mathcal{H}_q^{k-1} h_q \mathcal{H}_q^{l-1} M && \rightarrow W_R^\dagger \bar{H}_{qR}^{(k,l)} W_R. \end{aligned} \quad (5.13)$$

¹⁴Working with WBIs written in terms of the RH Hermitian matrices may nevertheless prove to be useful in certain situations, see section 5.2.1.

Note that only for $k = 1$ does one have $H_{dR}^{(k)} = H_{uR}^{(k)} = H_R$ independent of the sector q .

It is useful to relate these building blocks to the quark masses and the unitary rotations connecting the flavour and mass bases. Using eqs. (2.11) and (2.13), one has

$$\begin{aligned}\mathcal{H}_q &= \mathcal{V}_L^q \mathcal{D}_q^2 \mathcal{V}_L^{q\dagger}, & \mathcal{H}_{qR} &= \mathcal{V}_R^q \mathcal{D}_q^2 \mathcal{V}_R^{q\dagger}, \\ h_q &= \mathcal{V}_L^q \mathcal{D}_q A_R^{q\dagger} A_R^q \mathcal{D}_q \mathcal{V}_L^{q\dagger}, & h_{qR}^{(k)} &= A_R^q \mathcal{D}_q^{2k} A_R^{q\dagger}.\end{aligned}\quad (5.14)$$

The remaining Hermitian matrices depend on M , which can be written in two different ways, using matrices from either one sector or the other, recall eqs. (2.13) and (2.26). Therefore, the remaining matrices can be connected to the masses and unitary rotations in more than one way. For instance,

$$H = \mathcal{V}_L^u \mathcal{D}_u V_R \mathcal{D}_d \mathcal{V}_L^{d\dagger} = \mathcal{V}_L^q \mathcal{D}_q F_q \mathcal{D}_q \mathcal{V}_L^{q\dagger} \quad (q = u, d). \quad (5.15)$$

Additionally, using $\mathcal{H}_q = h_q + H$, we have $h_q = \mathcal{V}_L^q \mathcal{D}_q^2 \mathcal{V}_L^{q\dagger} - H$, which could also be obtained from eq. (5.14) via eq. (2.24). Finally, let us remark that eq. (2.13) allows us to write explicitly

$$H_{uR}^{(k)} = B_R^d \mathcal{D}_d V_L^\dagger \mathcal{D}_u^{2(k-1)} V_L \mathcal{D}_d B_R^d = B_R^u \mathcal{D}_u^{2k} B_R^{u\dagger}, \quad (5.16)$$

and thus re-express the conditions in eqs. (4.47) and (4.59) — which stem from the unitarity relation $B_R^u B_R^{u\dagger} = \mathbb{1}_{N \times N}$ — in the form

$$H_{uR}^{(0)} \equiv M^\dagger \mathcal{H}_u^{-1} M = \mathbb{1}_{N \times N}. \quad (5.17)$$

Now, we are in a position to construct WBIs and relate them to the quark masses and mixings. The masses of all q -sector quarks can be obtained from a set of $3 + N$ WBIs constructed solely from \mathcal{H}_q . These are given by

$$\chi_\sigma(\mathcal{H}_q) = \frac{1}{\sigma!} \left[\frac{\partial^\sigma}{\partial x^\sigma} \det(\mathcal{H}_q + x \mathbb{1}_{3+N}) \right]_{x=0}, \quad (5.18)$$

where $0 \leq \sigma \leq 2 + N$. They are closely related to the coefficients of a characteristic polynomial, with e.g. $\chi_0(\mathcal{H}_q) = \det \mathcal{H}_q = (m_1^q \dots M_N^q)^2$ and $\chi_{2+N}(\mathcal{H}_q) = \text{Tr} \mathcal{H}_q = (m_1^q)^2 + \dots + (M_N^q)^2$.

WBIs that are written in terms of the mixings involve matrices from both quark sectors. We have, for instance, the following set of CP-even WBIs

$$\text{Tr}(\mathcal{H}_u^r \mathcal{H}_d^s) = \text{Tr} \left(V_L^\dagger \mathcal{D}_u^{2r} V_L \mathcal{D}_d^{2s} \right) = m_\alpha^{2r} m_i^{2s} |V_{\alpha i}^L|^2, \quad (5.19)$$

which relate to the mixing of LH quarks. Here, the implied sums in α and i go over all $3 + N$ quark masses in the up and down sectors, respectively. Similarly, after straightforward manipulations, one finds

$$\text{Tr} \left(H_{uR}^{(r)} H_{dR}^{(s)} \right) = \text{Tr} \left(V_R^\dagger \mathcal{D}_u^{2r} V_R \mathcal{D}_d^{2s} \right) = m_\alpha^{2r} m_i^{2s} |V_{\alpha i}^R|^2, \quad (5.20)$$

which are related to the mixing of RH quarks. It is also worth pointing out that, for $N = 1$, both $H_{qR}^{(k)}$ and $\bar{H}_{qR}^{(k,l)}$ are scalars, which are WBIs by themselves. In that case, and in terms of physical observables, one can write

$$H_{qR}^{(k)} = \text{Tr} \left(\mathcal{D}_q^{2k} F_q \right) \Rightarrow \begin{cases} H_{dR}^{(k)} = \sum_{i=d,s,b} m_i^{2k} F_{ii}^d + M_{B'}^{2k} F_{44}^d, \\ H_{uR}^{(k)} = \sum_{\alpha=u,c,t} m_\alpha^{2k} F_{ii}^u + M_{T'}^{2k} F_{44}^u, \end{cases} \quad (5.21)$$

and

$$\bar{H}_{qR}^{(k,l)} = \text{Tr} \left(F_q \mathcal{D}_q^{2k} (\mathbb{1} - F_q) \mathcal{D}_q^{2l} \right) \Rightarrow \begin{cases} \bar{H}_{dR}^{(k,l)} = H_{dR}^{(k+l)} - \sum_{i,j}^4 m_i^{2k} m_j^{2l} |F_{ij}^d|^2, \\ \bar{H}_{uR}^{(k,l)} = H_{uR}^{(k+l)} - \sum_{\alpha,\beta}^4 m_\alpha^{2k} m_\beta^{2l} |F_{\alpha\beta}^u|^2. \end{cases} \quad (5.22)$$

As for CP-odd WBIs, i.e. WBIs with non-vanishing imaginary parts, one has to look into combinations of at least three distinct Hermitian matrices and their powers. Some of these are of the form

$$\begin{aligned} \text{Tr} ([\mathcal{H}_u^r, \mathcal{H}_d^s] H) &= 2i \text{Im} \text{Tr} (\mathcal{H}_u^r \mathcal{H}_d^s H) \\ &= 2i \text{Im} \text{Tr} \left(\mathcal{D}_u^{2r+1} V_L \mathcal{D}_d^{2s+1} V_R^\dagger \right) \\ &= 2i m_\alpha^{2r+1} m_i^{2s+1} \text{Im} \mathcal{B}_{\alpha i}^{LR}, \end{aligned} \quad (5.23)$$

where we have recalled the definition of the bilinears $\mathcal{B}_{\alpha i}^{LR} = V_{\alpha i}^L V_{\alpha i}^{R*}$ from eq. (3.26). As before, the implied sums in α and i go over all $3 + N$ quark masses in the up and down sectors, respectively. Note that, while in the SM the CP-odd WBI with the lowest mass dimension has dimension $M = 12$, here, for the case $r = s = 1$, one finds a CP-odd WBI of dimension $M = 6$ which is non-zero in general. In that specific case we have

$$\text{Tr} ([\mathcal{H}_u, \mathcal{H}_d] H) = \text{Tr} ([h_u, h_d] H) = 2i m_\alpha^3 m_i^3 \text{Im} \mathcal{B}_{\alpha i}^{LR}. \quad (5.24)$$

In principle, this could imply the presence of new and important sources of CP violation in the quark sector, as we will show in sections 6 and 8. In section 6, we also identify other types of CP-odd WBIs which can be relevant in enhancing CP violation.

5.2 Reconstructing minimal parameterizations from WBIs

In the last section, we showed how one might construct WBIs and relate them to quark masses and mixings. Here, we are instead interested in the possibility of reconstructing the $(3 + N)$ -dimensional quark mass matrices from a set of WBIs. To do this in a comprehensive way, it is important to fix a suitable weak basis. In fact, this is a crucial ingredient; for certain WB choices, the mass-matrix parameters present in the relations may be entangled in a convoluted way, that obscures or even obstructs a viable solution.

In section 5.2.1, we will show how to characterize all 22 parameters of the stepladder WB, in the $N = 1$ case, in terms of a relatively small number of WBIs. In section 5.2.2, we present an alternative procedure which considers the minimal WB. Although slightly less direct, the latter can be applied to the case of $N \geq 1$ doublets. A numerical example is presented in section 5.2.3.

5.2.1 Reconstructing the stepladder WB

Recall the stepladder weak basis introduced in section 3.1, eq. (3.17), for the case $N = 1$:

$$\mathcal{M}_u = \begin{pmatrix} \tilde{m}_u & 0 \\ M_Q r_u & M_Q \end{pmatrix}, \quad \mathcal{M}_d = \begin{pmatrix} \tilde{V} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \tilde{m}_d & 0 \\ M_Q r_d & M_Q \end{pmatrix}. \quad (5.25)$$

Here \tilde{V} is a general 3×3 unitary matrix having three real angles and six complex phases, one of which is an internal phase, while the other five are factorizable phases. The real vectors $r_{d,u}$ and real matrices $\tilde{m}_{d,u}$ are, in both sectors, of the special form

$$r_q = (0 \quad 0 \quad r_{q0}), \quad \tilde{m}_q = M_Q \begin{pmatrix} r_{q5} & 0 & 0 \\ r_{q4} & r_{q3} & 0 \\ 0 & r_{q2} & r_{q1} \end{pmatrix}, \quad (5.26)$$

while M_Q is a real number. In fact, we may choose M_Q and all the (dimensionless) r_{qi} to be non-negative, as any physical minus sign can be incorporated into \tilde{V} via suitable rephasings of the quark fields. In this way, the matrices

$$\begin{pmatrix} \tilde{m}_q & 0 \\ M_Q r_q & M_Q \end{pmatrix} = M_Q \begin{pmatrix} r_{q5} & 0 & 0 & 0 \\ r_{q4} & r_{q3} & 0 & 0 \\ 0 & r_{q2} & r_{q1} & 0 \\ 0 & 0 & r_{q0} & 1 \end{pmatrix} \quad (5.27)$$

have a “stepladder” shape. As we shall see, this distinctive structure will be crucial to establish direct relations between the mass-matrix parameters and WBI quantities. For starters, in terms of the former, the RH Hermitian squared mass matrices $\mathcal{H}_{qR} \equiv \mathcal{M}_q^\dagger \mathcal{M}_q$ are real even for $q = u$ (they do not depend on \tilde{V}) and read

$$\mathcal{H}_{qR} = \begin{pmatrix} \tilde{m}_q^T \tilde{m}_q + M_Q^2 r_q^T r_q & M_Q^2 r_q^T \\ M_Q^2 r_q & M_Q^2 \end{pmatrix}. \quad (5.28)$$

Before proceeding, recall that this is also a “minimal” WB. Indeed, by counting the parameters that characterize this WB — the 12 real parameters r_{qi} , the mass scale M_Q , and the 9 parameters that determine \tilde{V} — one finds a total of 22 parameters, which coincides with the number of physical parameters of the model (cf. Table 2 and the discussion at the end of section 3.1.2).

A “step-by-step” solution

Now, as it is clear from the discussion in section 5.1, using \mathcal{H}_{qR} one may construct the following CP-even WBIs,

$$I_k^{q'} \equiv \text{Tr} \left(\mathcal{K}_0 \mathcal{H}_{qR}^{k+2} \right), \quad \mathcal{K}_0 \equiv \text{diag}(0, 0, 0, 1), \quad (5.29)$$

where $k \in \{-1, 0, 1, 2, \dots\}$. Note that, in the notation of last section and in this $N = 1$ VLQ doublet case we have

$$I_k^{q'} = \text{Tr} \left(\mathcal{K}_0 \mathcal{H}_{qR}^{k+2} \right) = H_{qR}^{(k+2)}, \quad (5.30)$$

which we have computed in terms of quark masses and RH mixings in eq. (5.21).

Note that these invariants differ in nature from those in the SM. We recall that in the SM, with the 3×3 RH Hermitian squared mass matrix $h = m^\dagger m$, we can only obtain information about the masses. There are no Z -mediated currents involving the RH fields. The limited information regarding physical parameters that is contained in h can be accessed by computing, e.g. $\text{Tr } h$, $\text{Tr}(h^2)$, and $\text{Tr}(h^3)$ or $\det h$. The same applies to the RH Hermitian squared mass matrix of extensions of the SM with VLQ isosinglets, as in that case, there are also no Z -mediated currents involving the RH fields. Clearly this is not the case for extensions of the SM with doublet VLQs, where we must deal with extra New Physics (NP) present in the mixing of RH quarks. Thus, in the considered model with one doublet VLQ, one expects more information to be contained in WBIs built from RH Hermitian matrices. Indeed, and in contrast with the SM, there are RH weak-basis rotations which are necessarily shared between the up and down sectors, allowing for the construction of the new types of WBIs considered here.

The real parameters r_{qi} and M_Q , contained in eq. (5.27), can be determined in terms of the $I_k^{q'}$. We find, for both sectors ($q = u, d$),

$$I_{-1}^{q'} = M_Q^2, \quad (5.31)$$

and then, making use of the rescaled invariants

$$I_k^q \equiv \frac{1}{M_Q^{2(k+2)}} I_k^{q'}, \quad k = 0, 1, 2, \dots, \quad (5.32)$$

we obtain

$$\begin{aligned} I_0 &= 1 + r_0^2, \\ I_1 &= (1 + r_0^2)^2 + r_0^2 r_1^2, \\ I_2 &= \left[(1 + r_0^2)^3 + 2r_0^2 r_1^2 + r_0^2 r_1^4 + 2r_0^4 r_1^2 \right] + r_0^2 r_1^2 r_2^2, \\ I_3 &= P_3(r_0^2, r_1^2, r_2^2) + r_0^2 r_1^2 r_2^2 r_3^2, \\ I_4 &= P_4(r_0^2, r_1^2, r_2^2, r_3^2) + r_0^2 r_1^2 r_2^2 r_3^2 r_4^2, \end{aligned} \quad (5.33)$$

where P_k are polynomial functions of r_0^2, \dots, r_{k-2}^2 and r_{k-1}^2 . Here, and for now, we omit the superscript q when expressions apply to both sectors. It should, however, be stressed that one generically needs to compute two invariants of each kind, one for each sector.

Given the structure of the stepladder WB, the relations of eq. (5.33) also follow a ‘‘stepladder pattern’’. By this we mean the following: the invariant I_1 is the sum of a polynomial in r_0 , which can be determined from I_0 , and a single term depending on the next parameter r_1 , to wit, the term $r_0^2 r_1^2$. This pattern repeats itself: each invariant I_k can be written as the sum of a polynomial $P_k(r_0^2, \dots, r_{k-1}^2)$ and a term proportional to r_k^2 . This allows one to iteratively extract each of the r_i . Finally to obtain r_5 , one may compute I_5 or, alternatively, evaluate

$$\det \mathcal{H}_R = M_Q^8 r_1^2 r_3^2 r_5^2, \quad (5.34)$$

for each sector. Thus, in both sectors, one can directly extract M_Q and all r_k from a small set of WBIs.

However, it should be noted that this method appears to fail when some $r_k = 0$. In fact, if a given $r_k = 0$, then none of these invariants depends on any of the $r_{l>k}$. For example, if $r_2 = 0$ then I_3 and I_4 (as well as all $I_{k>4}$) are independent of r_3 and r_4 . Still, in those cases¹⁵ one can always determine all non-vanishing r_i parameters, as there is a stepladder WB where the off-diagonal $r_{l>k}$ also vanish (see the discussion at the end of section 3.1.2). In this stepladder WB, the mass matrix is block-diagonal and only the r_i of the upper-left (diagonal) block remain undetermined. They can be obtained with the help of the determinant of \mathcal{H}_R or traces of its powers. For instance, if $r_2 = 0$, then $r_4 = 0$ in this WB, and one would extract $r_{3,5}$ by computing $\text{Tr } \mathcal{H}_R = M_Q^2(1 + r_0^2 + r_1^2 + r_3^2 + r_5^2)$ and $\det \mathcal{H}_R$ given in eq. (5.34).

To complete our characterization, we need to express all parameters of the unitary matrix \tilde{V} in terms of WBIs. Here we will assume that all $r_i \neq 0$. From the form of \mathcal{M}_d in eq. (5.25), it is clear that the WBIs that may “capture” the entries of \tilde{V} are those which relate to WBTs of LH fields. Since \tilde{V} is in general complex (in fact, it contains all physical phases) one must also consider CP-odd WBIs.

There is an analogous iterative way to fully determine \tilde{V} . We start by defining

$$J_{rs} \equiv \frac{1}{M_Q^{2(1+r+s)}} \text{Tr} (\mathcal{H}_u^r \mathcal{H}_d^s H) . \quad (5.35)$$

Then,

$$J_{11} = \frac{1}{M_Q^6} \text{Tr} (\mathcal{H}_u \mathcal{H}_d H) = (1 + r_{u0}^2) (1 + r_{d0}^2) + r_{u0} r_{u1} r_{d0} r_{d1} \tilde{V}_{33} , \quad (5.36)$$

which is linear in \tilde{V}_{33} , and depends on the already-determined real parameters r_{qi} and M_Q . Thus, by computing J_{11} , which is complex in general, both $|\tilde{V}_{33}|$ and $\arg \tilde{V}_{33}$ can be determined without discrete ambiguities. Next, we evaluate

$$J_{12} = \frac{1}{M_Q^8} \text{Tr} (\mathcal{H}_u \mathcal{H}_d^2 H) = \mathcal{P}_{12} + r_{u0} r_{u1} r_{d0} r_{d1} r_{d2} r_{d3} \tilde{V}_{32} , \quad (5.37)$$

where \mathcal{P}_{12} is again a simple polynomial which depends on already-known parameters, including \tilde{V}_{33} . Explicitly, we have

$$\begin{aligned} \mathcal{P}_{12} = & (1 + r_{d0}^2) \left[(1 + r_{d0}^2) (1 + r_{u0}^2) + r_{d0} r_{d1} r_{u0} r_{u1} \tilde{V}_{33} \right] \\ & + r_{d0} r_{d1} \left[r_{d0} r_{d1} (1 + r_{u0}^2) + r_{u0} r_{u1} (r_{d1}^2 + r_{d2}^2) \tilde{V}_{33} \right] . \end{aligned} \quad (5.38)$$

Notice that J_{12} is linear in \tilde{V}_{32} . Thus, like \tilde{V}_{33} , this entry can be fully determined without ambiguity. Further computing

$$J_{13} = \frac{1}{M_Q^{10}} \text{Tr} (\mathcal{H}_u \mathcal{H}_d^3 H) = \mathcal{P}_{13} + r_{u0} r_{u1} r_{d0} r_{d1} r_{d2} r_{d3} r_{d4} r_{d5} \tilde{V}_{31} \quad (5.39)$$

¹⁵We disregard the cases where r_1 , r_3 or r_5 vanish, as those amount to vanishing masses, cf. eq. (5.34).

allows one to find \tilde{V}_{31} . Once again, \mathcal{P}_{13} obeys the stepladder pattern: it is a linear combination of the previously-determined entries of \tilde{V} , whose coefficients are polynomials in the r_{qi} . The new invariant J_{13} is again linear in \tilde{V}_{31} , which is found directly.

To obtain all other \tilde{V}_{ij} , one needs to repeat this scheme, carefully choosing which invariants to compute. By computing, in the following order, the invariants

$$\begin{aligned}
J_{21} &= \frac{1}{M_Q^8} \text{Tr}(\mathcal{H}_u^2 \mathcal{H}_d H) = \mathcal{P}_{21} + r_{u0} r_{u1} r_{u2} r_{u3} r_{d0} r_{d1} \tilde{V}_{23}, \\
J_{31} &= \frac{1}{M_Q^{10}} \text{Tr}(\mathcal{H}_u^3 \mathcal{H}_d H) = \mathcal{P}_{31} + r_{u0} r_{u1} r_{u2} r_{u3} r_{u4} r_{u5} r_{d0} r_{d1} \tilde{V}_{13}, \\
J_{22} &= \frac{1}{M_Q^{10}} \text{Tr}(\mathcal{H}_u^2 \mathcal{H}_d^2 H) = \mathcal{P}_{22} + r_{u0} r_{u1} r_{u2} r_{u3} r_{d0} r_{d1} r_{d2} r_{d3} \tilde{V}_{22}, \\
J_{32} &= \frac{1}{M_Q^{12}} \text{Tr}(\mathcal{H}_u^3 \mathcal{H}_d^2 H) = \mathcal{P}_{32} + r_{u0} r_{u1} r_{u2} r_{u3} r_{u4} r_{u5} r_{d0} r_{d1} r_{d2} r_{d3} \tilde{V}_{12}, \\
J_{23} &= \frac{1}{M_Q^{12}} \text{Tr}(\mathcal{H}_u^2 \mathcal{H}_d^3 H) = \mathcal{P}_{23} + r_{u0} r_{u1} r_{u2} r_{u3} r_{u4} r_{u5} r_{d0} r_{d1} r_{d2} r_{d3} \tilde{V}_{21}, \\
J_{33} &= \frac{1}{M_Q^{14}} \text{Tr}(\mathcal{H}_u^3 \mathcal{H}_d^3 H) = \mathcal{P}_{33} + r_{u0} r_{u1} r_{u2} r_{u3} r_{u4} r_{u5} r_{d0} r_{d1} r_{d2} r_{d3} r_{d4} r_{d5} \tilde{V}_{11},
\end{aligned} \tag{5.40}$$

one fully determines all the remaining \tilde{V}_{ij} in succession, where again the \mathcal{P}_{ij} follow the stepladder pattern, being linear combinations of the \tilde{V}_{ij} obtained in previous steps. As a corollary, one sees that the vanishing of all nine J_{ij} is a sufficient condition for CP invariance in the case of non-vanishing r_{qi} , with $i = 0, 1, \dots, 5$. In section 6.3 we study the conditions for CP conservation in the various cases where r_{q0} , r_{q2} or r_{q4} vanish.

A direct connection is thus established between WBIs and WB parameters. In summary, we considered 13 CP-even WBIs, namely I'_{-1} and the 6 invariants $I_{0,\dots,5}^q$ for each sector ($q = u, d$), where the I_5^q can be exchanged for $\det \mathcal{H}_{qR}$; as well as the 9 WBIs J_{rs} , with $r, s = 1, 2, 3$. This set of 22 WBIs can be described collectively by

$$\mathcal{I}_{rs} \equiv \frac{1}{M_Q^{2(1+r+s)}} \text{Tr}(\mathcal{H}_u^r \mathcal{H}_d^s H), \tag{5.41}$$

for $(rs) = \{(00), (01), \dots, (06), (10), \dots, (60), (11), (12), (13), (21), (22), (23), (31), (32), (33)\}$, with $\mathcal{I}_{0s} = I_{s-1}^d$ and $\mathcal{I}_{r0} = I_{r-1}^u$. In general, the 31 real constraints described above are required to determine the relevant 22 parameters. In most cases, the unitarity of \tilde{V} can reduce the number of necessary constraints.

5.2.2 Reconstructing the minimal WB

In a less direct manner, one may yet connect the minimal weak basis of eq. (3.12) to WBIs which are straightforward to compute, given the numerical values of the Yukawa matrices in any weak basis, for any value of N . Thus, the corresponding connection gives a general procedure to map any parameterization of the Yukawa and VLQ (bare) mass matrices to

the minimal one studied above, also used to integrate out VLQs (see section 7). Finding this mapping is specially relevant in a top-down approach, where we may start from a different parameterization of the same model (e.g. from a specific flavour basis). In order to do so, we use the projection technique described below, which is valid up to singular directions, such as degenerate eigenvalues.

We define a projector P as an object that under WBTs transforms as one of the relevant building blocks and that satisfies $(P_i)_{ab} = \delta_{ai}\delta_{bi}$ in the considered WB. In our case, we are interested in the minimal WB and require that $P \rightarrow \mathcal{W}_L^\dagger P \mathcal{W}_L$ under WBT, just like the building blocks of eq. (3.4). In the minimal WB, one has

$$h_u = v^2 \begin{pmatrix} \hat{Y}_u^2 & \hat{Y}_u \hat{z}_u^\dagger \\ \hat{z}_u \hat{Y}_u & \hat{z}_u \hat{z}_u^\dagger \end{pmatrix}, \quad h_d = v^2 \begin{pmatrix} \hat{V} \hat{Y}_d^2 \hat{V}^\dagger & \hat{V} \hat{Y}_d \hat{z}_d^\dagger \\ \hat{z}_d \hat{Y}_d \hat{V}^\dagger & \hat{z}_d \hat{z}_d^\dagger \end{pmatrix}, \quad H = \begin{pmatrix} 0 & 0 \\ 0 & D_Q^2 \end{pmatrix}. \quad (5.42)$$

We assume, without loss of generality, that $\hat{Y}_{qi} > \hat{Y}_{qj} > 0$ and $D_{Qi} > D_{Qj} > 0$, for $i > j$, with $i, j = 1, 2, 3$ and $q = u, d$. Here and from now on, we use a single index when referring to matrix elements of the diagonal matrices \hat{Y}_q and D_Q . We also take the first column of \hat{z}_d to be real and positive, $\hat{z}_d \rightarrow \hat{z}_d$. For the unitary matrix \hat{V} , we choose a (standard) parameterization in terms of three mixing angles $\theta_{13} \in [0, \pi/2]$, $\theta_{12} \in [0, \pi/2]$, $\theta_{23} \in [0, \pi/2]$ and a phase $\delta_{13} \in [-\pi, \pi]$.

For the WB under consideration, one defines

$$P_{i \geq 4} \equiv \sum_{m=1}^N c_m^{(i)} H^m. \quad (5.43)$$

The corresponding projector condition, $(P_i)_{ab} = \delta_{ai}\delta_{bi}$ in a basis where H takes the form in eq. (5.42), fixes the $c_m^{(i)}$ coefficients as functions of the (WBI) non-zero eigenvalues of H , the D_{Qa}^2 . These coefficients may be re-expressed in terms of $\{\text{Tr } H, \text{Tr } H^2, \dots, \text{Tr } H^N\}$, i.e.

$$c_m^{(i)} \equiv c_m^{(i)}(\text{Tr } H, \text{Tr } H^2, \dots, \text{Tr } H^N) \quad (5.44)$$

are WBIs themselves and can be computed in any basis. Given that $P_{i \geq 4}$ is a polynomial in H , it transforms as desired, and any traces of combinations of the matrices h_u, h_d, H, P_i and their powers is also a WBI.

We may define now

$$P_{123} \equiv \mathbb{1}_N - \sum_{i=1}^N P_{i+3}. \quad (5.45)$$

In the minimal weak basis,

$$h_u^P \equiv P_{123} h_u P_{123} = v^2 \begin{pmatrix} \hat{Y}_u^2 & 0 \\ 0 & 0 \end{pmatrix}. \quad (5.46)$$

Defining

$$P_{i < 4} \equiv \sum_{m=1}^3 a_m^{(i)} \left(\text{Tr } h_u^P, \text{Tr } (h_u^P)^2, \text{Tr } (h_u^P)^3 \right) (h_u^P)^m, \quad (5.47)$$

with $a_m^{(i)}$ such that these are projectors in the minimal weak basis, $(P_i)_{ab} = \delta_{ai}\delta_{bi}$, we now have all the P_i , with $i = 1, \dots, N + 3$, as (dimensionless) polynomials in h_u and H . Hence, $P_i \rightarrow \mathcal{W}_L^\dagger P_i \mathcal{W}_L$ under WBTs and they, together with P_{123} , can be used to build WBIs.

Now that we have all the projectors, we can trivially express all the moduli of some matrix $C = h_u, h_d, H, h_u^2, h_u h_d, \dots$, in this minimal basis as a function of WBIs, via

$$|C_{ij}|^2 = \text{Tr}(P_i C P_j C). \quad (5.48)$$

Taking this into account, the mapping of the Yukawa and bare mass matrices in any basis to the minimal one can be achieved by following these steps:

1. Find the $N + 3$ projectors P_i as explained above. This step involves computing the non-zero eigenvalues of H and h_u^P , giving us the values of D_Q and \hat{Y}_u .
2. \hat{Y}_d^2 are the non-zero eigenvalues of $P_{123} h_d P_{123}$. These weak-basis invariants can be trivially obtained, for example, by solving

$$\text{Tr} \left[(P_{123} h_d P_{123})^k \right] = \sum_{i=1}^3 \left(v \hat{Y}_{di} \right)^{2k}, \quad (5.49)$$

where $k = 1, 2, 3$.

3. The moduli of \hat{V} can be reconstructed from

$$\begin{aligned} \text{Tr}(P_i h_d) &= \sum_j |\hat{V}_{ij}|^2 \left(v \hat{Y}_{dj} \right)^2, \\ \text{Tr}(P_i h_d P_{123} h_d) &= \sum_j |\hat{V}_{ij}|^2 \left(v \hat{Y}_{dj} \right)^4, \end{aligned} \quad (5.50)$$

where $i = 1, 2$ only. Taking into account the unitarity of \hat{V} , this gives θ_{12} , θ_{13} , θ_{23} , and $|\delta_{13}|$ via $\cos \delta_{13}$. The sign of δ_{13} is the same as the one of $\text{Im}[\text{Tr}(P_1 h_d P_2 h_d P_3 h_d)]$, since

$$\begin{aligned} \text{Im}[\text{Tr}(P_1 h_d P_2 h_d P_3 h_d)] &= v^6 \left((\hat{Y}_{d3})^2 - (\hat{Y}_{d2})^2 \right) \left((\hat{Y}_{d3})^2 - (\hat{Y}_{d1})^2 \right) \left((\hat{Y}_{d2})^2 - (\hat{Y}_{d1})^2 \right) \\ &\quad \times \frac{1}{8} \sin 2\theta_{12} \sin 2\theta_{13} \sin 2\theta_{23} \cos \theta_{13} \sin \delta_{13}. \end{aligned} \quad (5.51)$$

4. For $|z_u|_{\alpha i}$ one simply has

$$|z_u|_{\alpha i} = \frac{\sqrt{\text{Tr}(P_\alpha h_u P_i h_u)}}{v^2 \hat{Y}_{ui}}, \quad (5.52)$$

while $|\hat{z}_d|_{\alpha i}$ can be obtained from

$$\text{Tr} \left[P_\alpha h_d (P_{123} h_d)^k \right] = \sum_i \left(v \hat{Y}_{di} \right)^{2k} v^2 |\hat{z}_d|_{\alpha i}^2, \quad (5.53)$$

where $k = 1, 2, 3$.

5. The most involved step is finding the remaining phases. For each row α , we first take

$$\text{Tr} [P_\alpha (h_d P_{123} h_d) P_i h_d] = \sum_{k,j}^3 v^6 (\hat{z}_d)_{\alpha j} (\hat{z}_d)_{\alpha k}^* (\hat{Y}_{dj})^3 \hat{Y}_{dk} \hat{V}_{ij}^* \hat{V}_{ik}, \quad (5.54)$$

with $i = 1, 2$. Since we have $\arg[(\hat{z}_d)_{\alpha 1}] = 0$, at this stage we can use those equations to fix the only remaining unknown parameters there, i.e., $\arg[(\hat{z}_d)_{\alpha 2}]$ and $\arg[(\hat{z}_d)_{\alpha 3}]$. This completes the determination of all parameters in \mathcal{Y}_d and thus h_d in this weak basis, including the phases in h_d . The remaining $3N$ phases in z_u can be simply obtained from

$$\arg[(z_u)_{\alpha i}] = \arg[(h_u)_{\alpha i}] = \arg[\text{Tr}(P_\alpha h_u P_i h_d)] + \arg[(h_d)_{\alpha i}], \quad (5.55)$$

since $\text{Tr}(P_\alpha h_u P_i h_d) = (h_u)_{\alpha i} (h_d)_{i\alpha}$.

5.2.3 A numerical example

For illustration, let us give a numerical example with one doublet. Starting from an arbitrarily convoluted set of values for the original Yukawa and mass matrices in eq. (2.2),¹⁶

$$\begin{aligned} \mathcal{Y}_u^0 &\simeq \begin{pmatrix} 0.200 - 0.242i & -0.244 - 0.228i & -0.157 + 0.101i \\ 0.209 + 0.217i & 0.244 - 0.207i & -0.099 - 0.143i \\ 0.242 - 0.244i & -0.240 - 0.244i & 0.089 - 0.180i \\ -0.244 - 0.244i & -0.243 + 0.244i & -0.180 - 0.088i \end{pmatrix}, \\ \mathcal{Y}_d^0 &\simeq \begin{pmatrix} 0.084 - 0.080i & 0.085 - 0.083i & 0.101 - 0.080i \\ 0.080 + 0.080i & 0.083 + 0.081i & 0.080 + 0.095i \\ -0.082 + 0.080i & -0.080 + 0.080i & -0.080 + 0.087i \\ 0.080 + 0.082i & 0.080 + 0.080i & 0.088 + 0.080i \end{pmatrix}, \\ M^0 &\simeq \begin{pmatrix} 4.21 - 4.22i \\ 4.21 + 3.95i \\ -3.69 + 4.24i \\ 4.25 + 3.69i \end{pmatrix} v, \end{aligned} \quad (5.56)$$

the algorithm described in section 5.2.1 returns the equivalent stepladder form

$$v \mathcal{Y}_u = \begin{pmatrix} \tilde{m}_u \\ \tilde{r}_u \end{pmatrix}, \quad v \mathcal{Y}_d = \begin{pmatrix} \tilde{V} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \tilde{m}_d \\ \tilde{r}_d \end{pmatrix}, \quad M = \begin{pmatrix} 0 \\ M_Q \end{pmatrix}, \quad (5.57)$$

¹⁶As it will become apparent below, it corresponds to a physically meaningful solution. In this specific basis, all Yukawa couplings are of the same order.

with $M_Q \simeq 11.49 v$ ($\simeq 2$ TeV) and

$$\begin{aligned}
\begin{pmatrix} \tilde{m}_u \\ \tilde{r}_u \end{pmatrix} &\simeq \begin{pmatrix} 0.00014 & 0 & 0 \\ 0.0034 & 9.0 \times 10^{-4} & 0 \\ 0 & 0.92 & 0.18 \\ 0 & 0 & 0.37 \end{pmatrix} v, \\
\begin{pmatrix} \tilde{m}_d \\ \tilde{r}_d \end{pmatrix} &\simeq \begin{pmatrix} 1.4 \times 10^{-5} & 0 & 0 \\ 3.7 \times 10^{-7} & 0.0035 & 0 \\ 0 & 0.015 & 0.0013 \\ 0 & 0 & 0.41 \end{pmatrix} v, \\
\tilde{V} &\simeq \begin{pmatrix} 0.98 + 5.6 \times 10^{-6} i & 0.18 + 0.00040 i & -0.043 + 0.0018 i \\ 0.18 + 0.00014 i & -0.95 + 0.0081 i & 0.24 + 0.036 i \\ -0.0081 + 0.0033 i & 0.15 + 0.19 i & 0.48 + 0.84 i \end{pmatrix}.
\end{aligned} \tag{5.58}$$

Here, the row vectors $\tilde{r}_q = (0 \ 0 \ M_Q r_{q0})$ have mass dimension one.

Instead, the algorithm of section 5.2.2 applied to any of the two equivalent Yukawa sets,¹⁷ leads to the equivalent minimal form

$$\mathcal{Y}_u = \begin{pmatrix} \hat{Y}_u \\ \mathbf{z}_u \end{pmatrix}, \quad \mathcal{Y}_d = \begin{pmatrix} \hat{V}(\vec{\theta}_{\hat{V}}) \hat{Y}_d \\ \hat{\mathbf{z}}_d \end{pmatrix}, \quad M = \begin{pmatrix} 0 \\ D_Q \end{pmatrix}, \tag{5.59}$$

with again $D_Q = M_Q \simeq 11.49 v$, and

$$\begin{aligned}
\hat{Y}_u &= \text{diag}(6.91 \times 10^{-6}, 0.00344, 0.940), \\
\hat{Y}_d &= \text{diag}(1.45 \times 10^{-5}, 0.000292, 0.0158), \\
\mathbf{z}_u &= (0.365, 0.0180, -0.0702), \\
\hat{\mathbf{z}}_d &= (0.000501, -0.404, 0.0334 e^{2\pi i/3}), \\
\vec{\theta}_{\hat{V}} &\equiv (\theta_{12}^{\hat{V}}, \theta_{13}^{\hat{V}}, \theta_{23}^{\hat{V}}, \delta_{13}^{\hat{V}}) = (0.227, 0.00369, 0.0418, 0.36\pi).
\end{aligned} \tag{5.60}$$

This particular benchmark satisfies the constraints discussed in section 8, while providing a solution to one of the so-called Cabibbo angle anomalies (in this case CAA2, with $z_u, z_s \neq 0$; see also section 8.5.1).

A **Mathematica** implementation of the stepladder algorithm for $N = 1$ and of the minimal WB algorithm for $N = 1, 2, 3$ can be found as supplementary material, mapping any set of (non-singular) $\mathcal{Y}_u, \mathcal{Y}_d$ and M to these minimal bases.

¹⁷We have also checked that applying the stepladder algorithm to the equivalent minimal weak basis form also returns back the same stepladder result.

6 CP violation

In this section, we examine the CP-violating properties of a model containing vector-like quark doublets in a weak-basis-independent way. CPV must be connected to the non-vanishing imaginary parts of weak-basis-invariant quantities, e.g. imaginary parts of traces of mass matrix combinations [52, 53, 55, 56, 58]. We will show that there is *a crucial connection between weak-basis invariants (WBIs) and effective rephasing invariants*, which provides a compelling tool to analyze CP-violating phenomena (see also section 8).

We start by illustrating the construction of CP-odd WBIs in section 6.1. In section 6.2, we will show in the scenario with one VLQ doublet ($N = 1$) how the information carried by each WBI becomes evident using the parameters defined in section 4.1. These parameters are directly connected to observables. We show that the WBIs can be expressed in terms of the masses and mixings of the three families of SM quarks (providing an “effective” description of the WBIs). In particular, WBIs can be given in terms of the rephasing invariants introduced in section 4.1.3. In section 6.3, we derive a complete set of conditions for CP conservation in the case of $N = 1$, under the assumption of non-vanishing and non-degenerate quark masses.

6.1 CP-odd weak-basis invariants

In the SM, the presence of CPV is connected to the non-vanishing of one weak-basis invariant [52, 65, 66]:

$$\begin{aligned} \text{Tr}[\mathcal{H}_u, \mathcal{H}_d]^3 &= 3 \det[\mathcal{H}_u, \mathcal{H}_d] \\ &= -6i (m_t^2 - m_c^2)(m_t^2 - m_u^2)(m_c^2 - m_u^2)(m_b^2 - m_s^2)(m_b^2 - m_d^2)(m_s^2 - m_d^2) J, \end{aligned} \quad (6.1)$$

where $\mathcal{H}_q \equiv \mathcal{M}_q \mathcal{M}_q^\dagger$ as defined in section 5.1, while $J = \text{Im}(V_{ud}V_{cs}V_{us}^*V_{cd}^*)$ is the rephasing invariant of eq. (3.24). The vanishing of the trace in eq. (6.1) is a necessary and sufficient condition for CP conservation in the SM. This trace is the non-real WBI with lowest mass dimension which can appear in the context of the SM, as J is the simplest complex rephasing invariant that can be built. In a scenario involving VLQs, the extra couplings can act as new sources of CPV, since new physical phases emerge. Also, different kinds of complex WBIs can be built with lower mass matrix powers with respect to the SM, which can be traced to the presence of the new rephasing-invariant quantities, involving two or more quarks (recall section 3.3). In the following, we analyze the structure of different invariants of increasing mass dimension.

6.1.1 One or more vector-like doublets

The ingredients to build WBIs were provided in section 5.1. To construct CP-odd WBIs, one has to look into combinations of at least three distinct Hermitian WBI “building

blocks”, i.e. \mathcal{H}_u , \mathcal{H}_d and H . Then, the simplest WBIs are of the form (see also eq. (5.23)):

$$\begin{aligned} \text{Im Tr} \left[\mathcal{H}_u^n \mathcal{H}_d^k H \right] &= \text{Im Tr} \left[\left(\mathcal{M}_u \mathcal{M}_u^\dagger \right)^n \left(\mathcal{M}_d \mathcal{M}_d^\dagger \right)^k M M^\dagger \right] = \\ &= \text{Im Tr} \left[\mathcal{D}_u^{2n+1} V_L \mathcal{D}_d^{2k+1} V_R^\dagger \right] = \sum_{\alpha, i=1}^{3+N} m_\alpha^{2n+1} m_i^{2k+1} \text{Im} \left(V_{\alpha i}^L V_{\alpha i}^{R*} \right), \end{aligned} \quad (6.2)$$

where, as before, the Greek (Latin) indices refer to the up (down) sector and $H \equiv M M^\dagger$ as defined in section 3. Ordering them by mass dimension, the first CP-odd weak-basis invariant appears at mass dimension $M = 6$, using three Hermitian blocks:

$$\text{Im Tr} \left[\mathcal{H}_u \mathcal{H}_d H \right] = \text{Im Tr} \left[\mathcal{D}_u^3 V_L \mathcal{D}_d^3 V_R^\dagger \right] = \sum_{\alpha, i=1}^{3+N} m_\alpha^3 m_i^3 \text{Im} \left(V_{\alpha i}^L V_{\alpha i}^{R*} \right). \quad (6.3)$$

We also specified the expression of the invariant in the mass basis. In the last term, the sum runs over all species, the three standard model families and the vector-like doublets. Note that this lowest mass dimension CP-odd WBI, brings with it the rephasing invariant quantities $\mathcal{B}_{\alpha i}^{LR} = V_{\alpha i}^L V_{\alpha i}^{R*}$, which suggests the emergence of new sources of CPV arising from the mixing with vector-like doublets. In fact, the presence of these bilinears points to the fact that these extensions can induce CP-violating effects involving only two quarks. For instance, contributions to electric dipole moments and direct CP violation in kaon decays ϵ' emerge at tree level from the imaginary part of $V_{ud}^L V_{ud}^{R*}$ and $V_{us}^L V_{us}^{R*}$ (see section 8.3). Let us note that in the mass basis, using the relation in eq. (2.26), the WBI in eq. (6.3) can be written also in other ways involving more insertions of mixing matrices:

$$\begin{aligned} \text{Tr} \left[\mathcal{H}_u \mathcal{H}_d H \right] &= \text{Tr} \left[\mathcal{D}_u^3 V_L \mathcal{D}_d^3 V_R^\dagger \right] = \sum_{\alpha, i=1}^{3+N} m_\alpha^3 m_i^3 V_{\alpha i}^L V_{\alpha i}^{R*} \\ &= \text{Tr} \left[\mathcal{D}_u^2 V_L \mathcal{D}_d^3 F_d \mathcal{D}_d V_L^\dagger \right] = \sum_{\alpha, i=1}^{3+N} m_\alpha^2 m_i^3 m_j V_{\alpha i}^L V_{\alpha j}^{L*} F_{ij}^d \\ &= \text{Tr} \left[\mathcal{D}_u^3 V_L \mathcal{D}_d^2 V_L^\dagger \mathcal{D}_u F_u \right] = \sum_{\alpha, i=1}^{3+N} m_\alpha^3 m_i^2 m_\beta V_{\alpha i}^L V_{\beta i}^{L*} F_{\alpha\beta}^{u*} \\ &= \text{Tr} \left[\mathcal{D}_u^2 V_L \mathcal{D}_d^2 V_L^\dagger \mathcal{D}_u V_R \mathcal{D}_d V_L^\dagger \right] = \sum_{\alpha, i=1}^{3+N} m_\alpha^2 m_i^2 m_\beta m_j V_{\alpha i}^L V_{\beta i}^{L*} V_{L\alpha j}^{L*} V_{\beta j}^R, \end{aligned} \quad (6.4)$$

that is, in terms of the other rephasing invariants, trilinears and quartets (see section 3.3). Let us remark that the sum runs over all quarks, SM and vector-like, involving the whole $(3+N) \times (3+N)$ mixing matrices V_L , V_R , $F_{u,d}$. In the next section, it will be shown in the case of one vector-like doublet (and in an example with two vector-like doublets) that there exists a direct connection between the form of the weak basis invariants

and the effective rephasing invariants studied in section 4.1.3, which involve only the standard model quarks. In particular, it will be shown that the invariants built with three Hermitian matrices (of the type of eq. (6.2)) only involve bilinears $\hat{\mathcal{B}}_{\alpha i}$, where $\alpha = u, c, t$ and $i = d, s, b$.

We may, of course, obtain higher order WBIs, in general of mass dimension $M = 2(n + k + 1)$, by choosing different values of n and/or m . This means that, apart from the single 3-block WBI for $M = 6$, we can obtain two WBI for $M = 8$, three for $M = 10$, four for $M = 12$ and so on. In each one of those cases we could also make use of eq. (2.26) to write different but equivalent expressions.

Other distinct structures emerge when we consider WBIs with four Hermitian blocks of the form

$$\begin{aligned} \text{Im Tr} \left[\mathcal{H}_u^m \mathcal{H}_d^\ell \mathcal{H}_u^n H \right] &= \sum_{\alpha, i=1}^{3+N} m_\alpha^{2m+1} m_i^{2\ell} m_\beta^{2n+1} \text{Im} \left(V_{\alpha i}^L V_{\beta i}^{L*} F_{\beta \alpha}^u \right), \\ \text{Im Tr} \left[\mathcal{H}_d^m \mathcal{H}_u^\ell \mathcal{H}_d^n H \right] &= \sum_{\alpha, i=1}^{3+N} m_i^{2m+1} m_\alpha^{2\ell} m_j^{2n+1} \text{Im} \left(V_{\alpha i}^{L*} V_{\alpha j}^L F_{ji}^d \right), \end{aligned} \quad (6.5)$$

where $n \neq m$ in order to get a CP-odd invariant. The invariants of these type with lowest mass dimension are the two $M = 10$ WBIs obtained for $m = \ell = 1$ and $n = 2$. All these invariants contain now the imaginary part of the rephasing invariants $V_{\alpha i}^L V_{\alpha j}^{L*} F_{ij}^d$ and $V_{\alpha i}^L V_{\beta i}^{L*} F_{\beta \alpha}^u$ (trilinears) in their expression with lowest number of mixing matrices insertions. The trilinears involve three quarks and are related to CP-violation in flavour changing neutral currents (see section 8.4). Similarly to the 3-block case, these WBIs can be expressed with more insertions mixing matrices, i.e.

$$\begin{aligned} \text{Im Tr} \left[\mathcal{H}_u^m \mathcal{H}_d^\ell \mathcal{H}_u^n H \right] &= \sum_{\alpha, i=1}^{3+N} m_\alpha^{2m+1} m_i^{2\ell} m_\beta^{2n+1} \text{Im} \left(V_{\alpha i}^L V_{\beta i}^{L*} F_{\beta \alpha}^u \right) \\ &= \sum_{\alpha, i=1}^{3+N} m_\alpha^{2m+1} m_i^{2\ell} m_\beta^{2n} m_j \text{Im} \left(V_{\alpha i}^L V_{\beta i}^{L*} V_{\beta j}^L V_{\alpha j}^{R*} \right) \\ &= \sum_{\alpha, i=1}^{3+N} m_\alpha^{2m} m_i^{2\ell} m_\beta^{2n+1} m_j \text{Im} \left(V_{\alpha i}^L V_{\beta i}^{L*} V_{\alpha j}^{L*} V_{\beta j}^{R*} \right) \\ &= \sum_{\alpha, i=1}^{3+N} m_\alpha^{2m} m_i^{2\ell} m_\beta^{2n} m_j m_k \text{Im} \left(V_{\alpha i}^L V_{\beta i}^{L*} V_{\beta j}^L V_{\alpha k}^{L*} F_{jk}^d \right), \end{aligned} \quad (6.6)$$

where the largest rephasing invariants appear, quartets and quintuplets $V_{\alpha i}^L V_{\beta i}^{L*} V_{\beta j}^L V_{\alpha k}^{L*} F_{jk}^d$, and similarly interchanging up and down quarks (by considering WBIs with one power of H , quintuplets appear in WBIs with at least four blocks). The content of the weak basis invariants will become clearer in the next section in terms of the effective rephasing invariants described in section 4.1.3. We will show that in the case of one vector-like doublet the invariants built with four Hermitian blocks (of the

type of eq. (6.5)) involve bilinears $\hat{\mathcal{B}}_{\alpha i}$ and trilinears $\hat{\mathcal{T}}_{i,\alpha\beta}$ (or $\hat{\mathcal{T}}_{\alpha,ij}$), where the indices only refer to standard model quarks $\alpha = u, c, t$, $i = d, s, b$.

A sixth invariant of mass dimension 10 emerge in the form with five Hermitian blocks:

$$\text{Im Tr} \left[\mathcal{H}_u^m \mathcal{H}_d^\ell \mathcal{H}_u^n \mathcal{H}_d^k H \right] = \sum_{\alpha,i=1}^{3+N} m_\alpha^{2m+1} m_i^{2\ell} m_\beta^{2n} m_j^{2k+1} \text{Im} \left(V_{\alpha i}^L V_{\beta i}^{L*} V_{\beta j}^L V_{\alpha j}^{R*} \right), \quad (6.7)$$

where $n = m = \ell = k = 1$ at mass dimension 10. This invariant contains in its simplest expression the rephasing invariant quartet $V_{\alpha i}^L V_{\beta i}^{L*} V_{\beta j}^L V_{\alpha j}^{R*}$ involving four quarks. By doing the same exercise of expressing the matrix H in different ways in the mass basis with more insertions of mixing matrices, as in eq. (6.6), one can find again larger rephasing invariants. In particular, we can notice the new rephasing invariant sextet $V_{\alpha i}^L V_{\beta i}^{L*} V_{\beta j}^L V_{\delta j}^{L*} V_{\alpha k}^{L*} V_{\delta k}^R$. We will show that in the case of one vector-like doublet the invariants built with five Hermitian blocks are given by combinations of bilinears $\hat{\mathcal{B}}_{\alpha i}$, trilinears $\hat{\mathcal{T}}_{i,\alpha\beta}$, $\hat{\mathcal{T}}_{\alpha,ij}$ and the quartets $\hat{\mathcal{Q}}_{\alpha i\beta j}$ defined in section 4.1.3, involving only standard model quarks $\alpha = u, c, t$, $i = d, s, b$.

At $M = 12$, apart from the WBIs of eqs. (6.2), (6.5) and (6.7) containing the H block, we have the SM-like WBI built with four Hermitian blocks which are not vanishing in the SM limit (i.e. without the Hermitian matrix H),

$$\text{Im Tr} \left[\mathcal{H}_u^2 \mathcal{H}_d^2 \mathcal{H}_u \mathcal{H}_d \right] = \sum_{\alpha,i=1}^{3+N} m_\alpha^4 m_i^4 m_\beta^2 m_j^2 \text{Im} \left(V_{\alpha i}^L V_{\beta i}^{L*} V_{\beta j}^L V_{\alpha j}^{L*} \right), \quad (6.8)$$

which contains the quartet of mixing elements of the left unitary matrices.

From the results found in this section we can already expect that the CP-violating content embedded in the invariants depends on both the mass dimension and the number of Hermitian blocks from which it is built. For instance, the WBI of mass dimension 6 is built using three Hermitian matrices. The WBIs of mass dimension 8, which are also built with three blocks, bring different powers of the quark masses but share the same structure of the invariant of mass dimension 6, whereas the invariants of mass dimension 10 built from four or five blocks also bring new types of rephasing invariants. We will uncover and analyze this connection in detail in section 6.2. We will use $I(M, k)$ to identify each CP-odd weak-basis invariant, by the mass dimension M and by the number of blocks k (e.g. the 3-block WBI with $M = 6$ is dubbed $I(6, 3)$).

6.1.2 Other CP-odd WBIs with more than one vector-like doublet

By proceeding to higher masses, weak basis invariants with the same number of blocks will manifest the same expressions of rephasing invariants and mass eigenvalues, differing only by the powers of the mass eigenvalues. New structures and new rephasing invariants will appear when the weak basis invariant can be built by a larger number of Hermitian blocks. For instance, at mass dimension 14, the weak basis invariants built by six

Hermitian blocks appear, of the type:

$$\begin{aligned} \text{Im Tr} \left[\mathcal{H}_u^m \mathcal{H}_d^\ell \mathcal{H}_u^n \mathcal{H}_d^p \mathcal{H}_u^r H \right] &= \sum_{\alpha, i=1}^{3+N} m_\alpha^{2m+1} m_i^{2\ell} m_\beta^{2n} m_j^{2p} m_\delta^{2r+1} \text{Im} \left(V_{\alpha i}^L V_{\beta i}^{L*} V_{\beta j}^L V_{\delta j}^{L*} F_{\delta \alpha}^u \right), \\ \text{Im Tr} \left[\mathcal{H}_d^m \mathcal{H}_u^\ell \mathcal{H}_d^n \mathcal{H}_u^p \mathcal{H}_d^r H \right] &= \sum_{\alpha, i=1}^{3+N} m_i^{2m+1} m_\alpha^{2\ell} m_j^{2n} m_\beta^{2p} m_k^{2r+1} \text{Im} \left(V_{\alpha i}^{L*} V_{\alpha j}^L V_{\beta j}^{L*} V_{\beta k}^L F_{ki}^d \right), \end{aligned} \quad (6.9)$$

with $m \neq r$. Similarly other invariants can be constructed at higher mass dimension. We can also mention that by adding the pair of blocks $\mathcal{H}_u^m \mathcal{H}_d^n$ to the SM-like weak basis invariant of eq. (6.8) one finds the rephasing invariants built with the left-handed mixing matrix V_L . For instance, at mass dimension 16

$$\text{Im Tr} \left[\mathcal{H}_u^2 \mathcal{H}_d^2 \mathcal{H}_u \mathcal{H}_d \mathcal{H}_u \mathcal{H}_d \right] = \sum_{\alpha, i=1}^{3+N} m_\alpha^4 m_i^4 m_\beta^2 m_j^2 m_\delta^2 m_k^2 \text{Im} \left(V_{\alpha i}^L V_{\beta i}^{L*} V_{\beta j}^L V_{\delta j}^{L*} V_{\delta k}^L V_{\alpha k}^{L*} \right). \quad (6.10)$$

In the presence of more than one vector-like quark doublet, at mass dimension 8 one also encounters other types of WBIs, such as:

$$\begin{aligned} \text{Im Tr} \left[\mathcal{H}_u \mathcal{H}_d H H \right] &= \text{Im Tr} \left[\mathcal{D}_u^3 V_L \mathcal{D}_d^3 V_R^\dagger \mathcal{D}_u^2 F_u \right] = \sum_{\alpha, \beta, i=1}^{3+N} m_\alpha^3 m_i^3 m_\beta^2 \text{Im} \left(V_{\alpha i}^L V_{\beta i}^{R*} F_{\beta \alpha}^u \right) \\ &= \text{Im Tr} \left[\mathcal{D}_u^3 V_L \mathcal{D}_d^3 F_d \mathcal{D}_d^2 V_R^\dagger \right] = \sum_{\alpha, i, j=1}^{3+N} m_\alpha^3 m_i^3 m_j^2 \text{Im} \left(V_{\alpha i}^L V_{\alpha j}^{R*} F_{ij}^d \right), \end{aligned} \quad (6.11)$$

where other trilinear rephasing invariants appear. Starting from mass dimension 10, one can also construct the weak basis invariants using the product $\mathcal{H}_u H \mathcal{H}_d$, which brings in the rephasing invariants the entries of V_R instead of V_L :

$$\begin{aligned} \text{Im Tr} \left[\mathcal{H}_u H \mathcal{H}_d H H \right] &= \text{Im Tr} \left[\mathcal{D}_u^4 V_R \mathcal{D}_d^4 V_R^\dagger \mathcal{D}_u^2 F_u \right] = \sum_{\alpha, \beta, i=1}^{3+N} m_\alpha^4 m_i^4 m_\beta^2 \text{Im} \left(V_{\alpha i}^R V_{\beta i}^{R*} F_{\beta \alpha}^u \right) \\ &= \text{Im Tr} \left[\mathcal{D}_u^4 V_R \mathcal{D}_d^4 F_d \mathcal{D}_d^2 V_R^\dagger \right] = \sum_{\alpha, i, j=1}^{3+N} m_\alpha^4 m_i^4 m_j^2 \text{Im} \left(V_{\alpha i}^R V_{\alpha j}^{R*} F_{ij}^d \right). \end{aligned} \quad (6.12)$$

Larger rephasing invariants containing the mixings of the right handed sector emerge by

building weak basis invariants of higher mass dimension with the Hermitian matrix H :

$$\begin{aligned}
& \text{Im Tr} \left[\mathcal{H}_u \mathcal{H}_d H H H \right] \\
&= \text{Im Tr} \left[\mathcal{D}_u^3 V_R \mathcal{D}_d^3 V_R^\dagger \mathcal{D}_u^2 F_u \mathcal{D}_u^2 F_u \right] = \sum_{\alpha, \beta, \delta, i=1}^{3+N} m_\alpha^3 m_i^3 m_\beta^2 m_\delta^2 \text{Im} \left(V_{\alpha i}^L V_{\beta i}^{R*} F_{\beta \delta}^u F_{\delta \alpha}^u \right) \\
&= \text{Im Tr} \left[\mathcal{D}_u^3 V_L \mathcal{D}_d^3 F_d \mathcal{D}_d^2 F_d \mathcal{D}_d^2 V_R^\dagger \right] = \sum_{\alpha, i, j, k=1}^{3+N} m_\alpha^3 m_i^3 m_j^2 m_k^2 \text{Im} \left(V_{\alpha i}^L F_{ij}^d F_{jk}^d V_{\alpha k}^{R*} \right) \\
&= \text{Im Tr} \left[\mathcal{D}_u^3 V_L \mathcal{D}_d^3 F_d \mathcal{D}_d^2 V_R^\dagger \mathcal{D}_u^2 F_u \right] = \sum_{\alpha, \beta, i, j=1}^{3+N} m_\alpha^3 m_i^3 m_j^2 m_\beta^2 \text{Im} \left(V_{\alpha i}^L F_{ij}^d V_{\beta j}^{R*} F_{\beta \alpha}^u \right) \\
&= \text{Im Tr} \left[\mathcal{D}_u^3 V_L \mathcal{D}_d^3 V_R^\dagger \mathcal{D}_u^2 V_R \mathcal{D}_d^2 V_R^\dagger \right] = \sum_{\alpha, \beta, i, j=1}^{3+N} m_\alpha^3 m_i^3 m_j^2 m_\beta^2 \text{Im} \left(V_{\alpha i}^L V_{\beta i}^{R*} V_{\beta j}^R V_{\alpha j}^{R*} \right).
\end{aligned} \tag{6.13}$$

Note that the different expressions in this equation all appear with the same minimal number of insertion of mixing matrices.

Let us remark that in case of only one vector-like doublet the rephasing invariants (and weak-basis invariants) containing only matrix elements related to the right handed sector as in eq. (6.12) are real, whereas weak basis invariants built with more than one H block can be written in terms of weak basis invariants with lower mass dimension containing only one power of H as shown in eqs. (5.4) to (5.6). The same situation happens in presence of more vector-like doublets if the row vectors $z_{n\alpha(i)}$ are orthogonal between each other.

6.2 Effective CP-odd WBIs

In this section we provide an “effective” description of CP violating weak-basis invariants. Namely, we express the invariants in terms of the couplings involved in SM quarks interactions, which are connected to the 3×3 submatrices of V_L , V_R , F_d and F_u describing the coupling of SM quarks to W , Z and Higgs bosons. We show that in this way the information carried by weak-basis invariants can be more easily understood and connected to observables. Moreover it is possible to find a set of CP-odd weak-basis invariants which can give a complete description of the CP-violating effects in presence of one vector-like doublet mixing with the SM quarks.

This analysis is possible by taking into account that the ratio v/M_Q is a small parameter. As described in section 4.1, masses and mixings can be expressed in terms of Lagrangian parameters, more precisely the parameters in eq. (4.6), by an expansion in v/M_Q . Then, these relations can be used to write expressions for the weak-basis invariants, which are simply related to both the minimal set of Lagrangian parameters and mass basis quantities, and ultimately, with observables. In particular, for CP-odd weak-basis invariants this expansion also allows to clarify the content of weak-basis invariants in terms of rephasing invariant quantities involving only the three SM quarks.

Let us consider first CP-odd invariants of the form:

$$I(\mathbf{M}, 3) \equiv \text{Im Tr}[\mathcal{H}_u^n \mathcal{H}_d^m H], \quad (6.14)$$

where we used $I(\mathbf{M}, 3)$ to denote the family of invariants with the mass dimension $\mathbf{M} = 2(n+m+1)$ and built from 3 Hermitian blocks. Using the expressions of mass eigenvalues and mixing elements found in section 4.1 we find:

$$\frac{1}{M_Q^{2(m+n+1)}} \text{Im}(\text{Tr}[\mathcal{H}_u^n \mathcal{H}_d^m H]) = \frac{v^4}{M_Q^4} \sum_{i,\alpha=1}^3 \text{Im} \left(\hat{V}_{\alpha i}^L z_\alpha z_i^* \right) y_\alpha y_i \left(1 + k_{\alpha i}^{(m,n)} \right), \quad (6.15)$$

where $k_{\alpha i}^{(m,n)}$ stand for real corrections of order v^2/M_Q^2 or higher. CP violation is carried by the imaginary part of the rephasing invariant quantities

$$\frac{v^2}{M_Q^2} \hat{V}_{\alpha i}^L z_\alpha z_i^* = \hat{V}_{\alpha i}^L \hat{V}_{\alpha i}^{R*} = \hat{\mathcal{B}}_{\alpha i}^* \simeq V_{\alpha i}^L V_{\alpha i}^{R*}. \quad (6.16)$$

In the general case, all couplings z_α, z_i are different from zero, and thus all physical phases would appear in this class of invariants (see section 4.1.3). Even if the new physics only brings a real contribution, in the sense that only one phase survives in the model (e.g. all z_α, z_i are real in this parameterization) the CP-violating effect of the remaining phase would appear in these invariants. If the condition $\text{Im}(\hat{V}_{\alpha i}^L z_\alpha z_i^*) = 0$ held for each α, i (or for six independent combinations) then there would be no CP violation. These conditions change when some couplings are vanishing. In this case, not all physical phases emerge in bilinears and rephasing invariant trilinears and quartets become essential in the description of the CP-violating effects. In fact, in a situation in which one or more $z_{\alpha/i}$ are zero, the rephasing invariant trilinears and quartets $\hat{\mathcal{T}}_{i,\alpha\beta}, \hat{\mathcal{T}}_{\alpha,ij}, \hat{\mathcal{Q}}_{\alpha i\beta j}$ (see eq. (4.29)) cannot be written in terms of bilinears $\hat{\mathcal{B}}_{\alpha i}$ as shown in eq. (4.32) and they become non-redundant. As a consequence of this observation, we can already anticipate that in these scenarios the vanishing of the imaginary part of invariants containing only the bilinears will not suffice as conditions for CP conservation.

For completeness, let us notice that similar expansions hold for the real part of the same invariants. In first approximation, one has

$$\begin{aligned} \frac{1}{M_Q^{2(n+m+1)}} \text{Tr} \left[\mathcal{H}_u^n \mathcal{H}_d^m H \right] &\simeq 1 + \frac{v^2}{M_Q^2} \sum_{\alpha i} (n|z_\alpha|^2 + m|z_i|^2) \\ &= 1 + n \sum_{\alpha} \hat{F}_{\alpha\alpha}^u + m \sum_i \hat{F}_{ii}^d. \end{aligned} \quad (6.17)$$

If only two couplings are switched on (i.e. only $z_u, z_s \neq 0$), this means that, at first order

$$\begin{aligned} \frac{1}{M_Q^6} \text{Tr} \left[\mathcal{H}_u \mathcal{H}_d H \right] &\simeq 1 + \hat{F}_{uu}^u + \hat{F}_{ss}^d, \\ \frac{1}{M_Q^6} \text{Im} \left(\text{Tr} \left[\mathcal{H}_u \mathcal{H}_d H \right] \right) &\simeq y_u y_s \text{Im}(\hat{V}_{us}^L \hat{V}_{us}^{R*}). \end{aligned} \quad (6.18)$$

The next type of CP-odd invariants we can consider are of the form:

$$\begin{aligned} \text{I}(\mathbf{M}, 4)_d &\equiv \text{Im Tr}[\mathcal{H}_d^m \mathcal{H}_u^\ell \mathcal{H}_d^n H], \\ \text{I}(\mathbf{M}, 4)_u &\equiv \text{Im Tr}[\mathcal{H}_u^m \mathcal{H}_d^\ell \mathcal{H}_u^n H], \end{aligned} \quad (6.19)$$

with $\mathbf{M} = 2(n + m + \ell + 1)$. Taking $n > m$, we have

$$\begin{aligned} &\frac{1}{M_Q^{2m+2n+2\ell+2}} \text{Im}(\text{Tr}[\mathcal{H}_d^m \mathcal{H}_u^\ell \mathcal{H}_d^n H]) \\ &= \sum_{i,j,\alpha,\beta=1}^3 \left[\frac{v^{4+2m}}{M_Q^{4+2m}} \text{Im} \left(\hat{V}_{\alpha i}^L z_\alpha z_i^* \right) y_\alpha y_i^{2m+1} \left(1 + k_{i\alpha}^{(n,m,\ell)} \right) + \right. \\ &\quad + \frac{v^{4+2m+2\ell}}{M_Q^{4+2m+2\ell}} \text{Im} \left(\hat{V}_{\alpha i}^L \hat{V}_{\alpha j}^{L*} z_i^* z_j \right) y_i^{2m+1} y_j y_\alpha^{2\ell} \left(1 + k_{\alpha i j}^{(n,m,\ell)} \right) \\ &\quad \left. + \frac{v^{8+2m}}{M_Q^{8+2m}} \text{Im} \left(\hat{V}_{\alpha i}^L z_i^* z_\alpha \hat{V}_{\beta j}^{L*} z_j z_\beta^* \right) y_i^{2m+1} y_j y_\alpha y_\beta \left(1 + k_{i j \alpha \beta}^{(n,m,\ell)} \right) \right], \end{aligned} \quad (6.20)$$

and similarly for the second invariant with the exchange of down and up sectors. These invariants contain the rephasing invariant quantities involving three quarks associated with flavour changing neutral currents

$$\begin{aligned} \frac{v^2}{M_Q^2} z_i^* z_j \hat{V}_{\alpha i}^L \hat{V}_{\alpha j}^{L*} &= \hat{F}_{ij}^d \hat{V}_{\alpha i}^L \hat{V}_{\alpha j}^{L*} = \hat{\mathcal{T}}_{\alpha,ij} \simeq V_{\alpha i}^L V_{\alpha j}^{L*} F_{ij}^d, \\ \frac{v^2}{M_Q^2} z_\alpha z_\beta^* \hat{V}_{\alpha i}^L \hat{V}_{\beta i}^{L*} &= \hat{F}_{\alpha\beta}^{u*} \hat{V}_{\alpha i}^L \hat{V}_{\beta i}^{L*} = \hat{\mathcal{T}}_{i,\alpha\beta}^* \simeq V_{\alpha i}^L V_{\beta i}^{L*} F_{\beta\alpha}^u. \end{aligned} \quad (6.21)$$

The last term of eq. (6.20) involves the quantities of the type $\hat{V}_{\alpha i}^L \hat{V}_{\beta j}^{L*} z_i^* z_j z_\alpha z_\beta^*$ which can be expressed in terms of the rephasing invariants $\hat{V}_{\alpha i}^L z_\alpha z_i^*$.

Last, we can consider the 5-block weak-basis invariants

$$\text{I}(\mathbf{M}, 5) \equiv \text{Im}(\text{Tr}[\mathcal{H}_u^n \mathcal{H}_d^k \mathcal{H}_u^\ell \mathcal{H}_d^m H]), \quad (6.22)$$

with $\mathbf{M} = 2(n + k + m + \ell + 1)$. Besides the leading order terms proportional to $\hat{V}_{\alpha i}^L z_\alpha z_i^*$ and those associated with flavour changing neutral currents $\hat{V}_{\alpha i}^L \hat{V}_{\beta i}^{L*} z_\alpha z_\beta^*$, $\hat{V}_{\alpha i}^{L*} \hat{V}_{\alpha j}^L z_i z_j^*$, these invariants contain the rephasing invariant quantities involving four quarks

$$\frac{v^2}{M_Q^2} z_\beta z_j^* \hat{V}_{\alpha i}^{L*} \hat{V}_{\beta i}^L \hat{V}_{\alpha j}^L = \hat{V}_{\beta j}^{R*} \hat{V}_{\alpha i}^{L*} \hat{V}_{\beta i}^L \hat{V}_{\alpha j}^L = \hat{\mathcal{Q}}_{\beta j \alpha i}^* \simeq V_{\beta j}^{R*} V_{\alpha i}^{L*} V_{\beta i}^L V_{\alpha j}^L. \quad (6.23)$$

For instance, for the WBI of mass dimension 10 we can write:

$$\begin{aligned}
& \frac{1}{M_Q^{10}} \text{Im}(\text{Tr}[\mathcal{H}_u \mathcal{H}_d \mathcal{H}_u \mathcal{H}_d H]) \\
&= \frac{v^4}{M_Q^4} \sum_{i,\alpha=1}^3 \text{Im} \left[\hat{V}_{\alpha i}^L z_\alpha z_i^* \right] y_\alpha y_i \left(1 + k_{\alpha i}^{(10a)} \right) + \\
&+ \frac{v^8}{M_Q^8} \sum_{i,j,\alpha,\beta=1}^3 \text{Im} \left[\hat{V}_{\alpha j}^L \hat{V}_{\beta i}^L \hat{V}_{\beta j}^{L*} z_\alpha z_i^* \right] y_\beta^2 y_\alpha y_j^2 y_i \left(1 + k_{\alpha \beta i j}^{(10b)} \right) + \mathcal{O} \left(\frac{v^{12}}{M_Q^{12}} \right).
\end{aligned} \tag{6.24}$$

Finally we can consider the SM-like weak basis invariant shown in eq. (6.8) which is built with four Hermitian building blocks without the insertion of $H = MM^\dagger$ in eq. (6.8). This invariant contains, besides bilinears, trilinears and quartets, the SM-like quartet

$$\hat{Q}_{\alpha i \beta j} = \hat{V}_{\alpha i}^L \hat{V}_{\beta j}^L \hat{V}_{\alpha j}^{L*} \hat{V}_{\beta i}^{L*}, \tag{6.25}$$

recall eq. (4.31). Note that if we consider the rephasing invariant extracted solely from the LH mixing, $J = \text{Im}(V_{ud}^L V_{cs}^L V_{us}^{L*} V_{cd}^{L*})$ (see eq. (3.24)), then we have

$$|\text{Im} \hat{Q}_{\alpha i \beta j}| \simeq |\text{Im} V_{\alpha i}^L V_{\beta i}^{L*} V_{\beta j}^L V_{\alpha j}^{L*}| \simeq |J|. \tag{6.26}$$

We can conclude from the result of this section that there is a unique connection between weak basis invariants and rephasing invariants. Weak basis invariants built with three Hermitian blocks only contain bilinears, while trilinears appear in 4-block invariants. The non-standard quartets appear in 5-block invariants. The SM quartet explicitly emerges in the SM-like WBI of eq. (6.8).

6.3 Conditions for CP invariance

We can now study the weak-basis independent conditions for CP invariance (CPI) in presence of one vector-like doublet. If the imaginary part of any weak-basis invariant is different from zero, then at least one physical phase is present in the model, inducing CP-violation. Then, the conditions for CPI can be obtained by imposing that CP-odd weak-basis invariants vanish, i.e.

$$\text{I}(\mathbf{M}, k) = 0. \tag{6.27}$$

More precisely, we want to find a set of invariants which may completely describe the CP properties of the model. Then, the vanishing of these invariants would also imply CP conservation. We assume non-vanishing and non-degeneracy of quark masses. The results obtained in section 6.2 provide a simple guide in the identification of the set of invariants relevant in different scenarios, depending on the mixing of the VLQ doublet with the standard quarks. In the following, we illustrate the choice of weak basis invariants as based on these results. A more rigorous proof is given in Appendix C. We summarize the results in Table 3. In the first two columns of this table the number of couplings between SM-quarks and the extra VLQ doublet which are switched on are

$\#z_i$	$\#z_\alpha$	$\#\delta$	$\#I(M \geq 6, 3)$	$\#I(M \geq 8, 4)_d$	$\#I(M \geq 8, 4)_u$	$I(10, 5)$	$I_{(12,4)}^{\text{SM-like}}$
3	3	6	9				
3(2)	2(3)	5	7				
2	2	4	4				
3	1	4	3	1			
2	1	3	2	1			
3	0	3	= 0	6	= 0	= 0	
2	0	2	= 0	2	= 0	= 0	
1	3	4	3		1		
1	2	3	2		1		
0	3	3	= 0	= 0	6	= 0	
0	2	2	= 0	= 0	3	= 0	
1	1	2	1			1	
1	0	1	= 0	= 0	= 0	= 0	1
0	1	1	= 0	= 0	= 0	= 0	1

Table 3: CP-odd weak-basis invariants needed for CP conservation in different scenarios, in the presence of one vector-like doublet, assuming non-vanishing quark Yukawas with a SM-like hierarchy (see the text for more details).

indicated. In the third column we write the number of physical phases in each case. For each scenario, we indicate the type and number of weak-basis invariants which can provide a complete characterization of the CP properties of the model and whose vanishing would correspond to CP conservation. The entries with zeroes indicate that the imaginary part of the corresponding invariant vanishes in that scenario. Note that if the VLQ doublet is decoupled, or exhibits non-vanishing coupling only with one generation in only one sector, then the sole non-zero CP-odd invariant is the usual SM-like invariant for $3 + N$ generations, with mass dimension $M = 12$ and built from 4 blocks (see eq. (6.8)). The blank space indicates that the CP-odd WBIs is non-zero, but the condition is redundant. In fact, the set of invariants indicated in the table is not necessarily minimal, as we use the most convenient set of WBIs in each scenario. However, other non-zero invariants can be used in order to reduce the total number of WBIs needed for the necessary and sufficient conditions for CP conservation.

All $z_i \neq 0$, $z_\alpha \neq 0$. Let us start by the most generic case in which all couplings are allowed to be different from zero. In this scenario all the elements of the matri-

ces V_R , $F_{u,d}$ are non-vanishing and right-handed charged and neutral currents, both flavour-changing and flavour-conserving, are generated between SM quarks. In this case, the WBIs with three blocks in eq. (6.15) are linear combinations of the 9 bilinears $\text{Im}(\hat{V}_{\alpha i}^L z_\alpha z_i^*)$ (see eq. (6.15)). The 9 phases $\arg(\hat{V}_{\alpha i}^L z_\alpha z_i^*)$ are not independent, but in this scenario all the 6 physical phases emerge in these combinations. Moreover, the vanishing of all the $\text{Im}(\hat{V}_{\alpha i}^L z_\alpha z_i^*)$ for each α, i implies the vanishing of all physical phases, that is, CP invariance. In fact, as we have found in section 5.2.1 using the “stepladder” weak basis, the sufficient and necessary conditions for CPI in this scenario correspond to the vanishing of the imaginary part of 9 WBIs built from three Hermitian blocks.

One $z_{i/\alpha} = 0$. In this scenario one row or one column of V_R is zero, as well as one row and one column of F_u or F_d . There are 5 physical phases. Since there are 6 bilinears $\text{Im}(\hat{V}_{\alpha i}^L z_\alpha z_i^*)$ the 5 independent physical phases can be chosen among the 6 phases of the bilinears. Hence, the situation is similar to the previous case, with 3-block WBIs containing the information on CP-violation. It can be shown using the stepladder WB that the vanishing of the imaginary part of 7 WBIs with three blocks are the necessary and sufficient conditions for CP conservation in this scenario (see Appendix C).

One $z_i = 0$ and one $z_\alpha = 0$. In this scenario one row and one column of V_R are vanishing, as well as one row and one column of F_u and F_d . There are 4 physical phases and 4 combinations $\text{Im}(\hat{V}_{\alpha i}^L z_\alpha z_i^*)$. Also in this case, the conditions $\text{Im}(\hat{V}_{\alpha i}^L z_\alpha z_i^*) = 0$ for each α, i imply that all physical phases are zero. This can be seen e.g. by choosing the elements $\hat{V}_{\beta j}^L$ of a row $\beta \neq \alpha$ and a row $j \neq i$ to be real (one column and one row of \hat{V}_L can always be made real, together with one of the couplings $z_{i/\alpha}$) and also z_β real and assigning three phases to the other three non-vanishing z -couplings and the remaining one to the 2×2 submatrix of \hat{V}_L . Then these 4 phases are directly related to the phases of the bilinears in a trivial relation. Then, also in this scenario, 3-block WBIs can characterize all the CP properties. It can be shown that the vanishing of 4 WBIs of the type $\text{I}(n, 3) = 0$ are indeed necessary and sufficient conditions for CPI (see Appendix C).

Two $z_{\alpha(i)} = 0$ in the same sector. In this scenario 2 rows or 2 columns of V_R are zero, as well as 2 rows and 2 columns of F_u or F_d , implying no flavour changing couplings at tree level between the SM quarks in one sector. There are 4 physical phases. However, there are only 3 non-vanishing bilinears $\text{Im}(\hat{V}_{\alpha i}^L z_\alpha z_i^*)$. Then, the 3-block WBIs can be written as a linear combination of these 3 bilinears but one physical phase is not captured by these invariants (neither by the quartet $z_\beta z_j^* \hat{V}_{\alpha i}^{L*} \hat{V}_{\beta i}^L \hat{V}_{\alpha j}^L$). The remaining phase emerges in the trilinears of the type $z_i^* z_j \hat{V}_{\alpha i}^L \hat{V}_{\alpha j}^{L*}$ (for 2 couplings in the up-sector $z_\alpha = 0$) or $z_\alpha z_\beta^* \hat{V}_{\alpha i}^L \hat{V}_{\beta i}^{L*}$ (for 2 couplings in the down-sector $z_i = 0$). These trilinears appear in weak-basis invariants with at least 4 Hermitian blocks. In Appendix C we show that 3 conditions of the type $\text{I}(n, 3) = 0$ and one of the type $\text{I}(n, 4) = 0$ are necessary and sufficient for CPI.

Two $z_{\alpha(i)} = 0$ in one sector and one $z_{i(\alpha)} = 0$ in the other sector. This scenario is analogous to the previous one. There are 3 physical phases and there are 2 non-vanishing bilinears $\text{Im}(\hat{V}_{\alpha i}^L z_\alpha z_i^*)$. The condition $I(n, 3) = 0$ for two or more different 3-block invariants would imply the vanishing of the imaginary parts of the 2 bilinears $\text{Im}(\hat{V}_{\alpha i}^L z_\alpha z_i^*) = 0$. The remaining phase emerges in the trilinears of the type $z_i^* z_j \hat{V}_{\alpha i}^L \hat{V}_{\alpha j}^{L*}$ (or $z_\alpha z_\beta^* \hat{V}_{\alpha i}^L \hat{V}_{\beta i}^{L*}$). For instance, by further setting $z_b = 0$ in the previous scenario we have, for the 4-block WBI,

$$\text{Im}(\text{Tr}[\mathcal{H}_d \mathcal{H}_u \mathcal{H}_d^2 H]) \propto \text{Im}\left(\hat{V}_{td}^L \hat{V}_{ts}^{L*}\right) z_d z_s y_d y_s (y_t^2 - y_c^2) (y_d^2 - y_s^2). \quad (6.28)$$

The additional condition $I(n, 4) = 0$ would imply CP conservation.

Three $z_{\alpha(i)} = 0$ in one sector. In this scenario, the 3×3 submatrix of V_R involving SM quarks is vanishing, only its last column or last row survives, as well as one of the two matrices F_u or F_d depending on which sector is decoupled. All bilinears disappear, so that every 3-block WBI vanishes. Still, there are 3 physical phases and these phases appear in the trilinear terms of the form $\hat{V}_{\alpha i}^L \hat{V}_{\alpha j}^{L*} z_i^* z_j$ or $\hat{V}_{\alpha i}^L \hat{V}_{\beta i}^{L*} z_\alpha z_\beta^*$. Assuming for instance the case $z_\alpha = 0$, $\alpha = u, c, t$ (the down-sector case being analogous), for the 4-block invariants we have ($n > m$):

$$\text{Im}(\text{Tr}[\mathcal{H}_d^m \mathcal{H}_u^\ell \mathcal{H}_d^n H]) \propto \sum_{i,j,\alpha=1}^3 \text{Im}\left(\hat{V}_{\alpha i}^L \hat{V}_{\alpha j}^{L*} z_i^* z_j\right) y_i^{2m+1} y_j y_\alpha^{2\ell} \left(1 + k_{\alpha ij}^{(n,m,\ell)}\right). \quad (6.29)$$

It can be shown that the vanishing of 6 different invariants of this type would provide the necessary and sufficient conditions for CP conservation in this scenario (see Appendix C).

Three $z_{\alpha(i)} = 0$ in one sector and one $z_{i(\alpha)} = 0$ in the other sector. Let us take for illustration the case $z_\alpha = 0$, $\alpha = u, c, t$ and $z_b = 0$. Then, for instance

$$\begin{aligned} & \frac{1}{M_Q^{10}} \text{Im}(\text{Tr}[\mathcal{H}_d \mathcal{H}_u \mathcal{H}_d^2 H]) \\ & \simeq \frac{v^8}{M_Q^8} y_d y_s (y_d^2 - y_s^2) \left((y_u^2 - y_t^2) \text{Im}\left(\hat{V}_{ud}^L \hat{V}_{us}^{L*} z_d^* z_s\right) + (y_c^2 - y_t^2) \text{Im}\left(\hat{V}_{cd}^L \hat{V}_{cs}^{L*} z_d^* z_s\right) \right). \end{aligned} \quad (6.30)$$

The vanishing of physical phases would be signalled by the disappearance of invariants with at least 4 blocks. In particular, the vanishing of two 4-block invariants (of the relevant type) imply CP conservation in this scenario (see Appendix C).

Two $z_{\alpha/i} = 0$ in both sectors. In this scenario, only one coupling in each sector survives. There are no flavour-changing neutral currents at tree level between SM quarks and only one non-vanishing entry in the 3×3 submatrix of V_R involving the SM quarks.

There are two physical phases. One phase appears in the surviving bilinear, and consequently in the 3-block invariants. Let us assume as an illustration that only $z_u \neq 0$, $z_d \neq 0$. Then we have

$$\frac{1}{M_Q^6} \text{Im}(\text{Tr}[\mathcal{H}_u \mathcal{H}_d H]) \propto \frac{v^4}{M_Q^4} \text{Im} \left(\hat{V}_{ud}^L z_u z_d^* \right) y_u y_d. \quad (6.31)$$

However, the second phase cannot manifest in neither 3-block nor 4-block invariants. The last phase only emerge in quartets, or in terms of WBIs, in the $M = 10$ invariant made up of 5 blocks:

$$\begin{aligned} \frac{1}{M_Q^{10}} \text{Im}(\text{Tr}[\mathcal{H}_u \mathcal{H}_d \mathcal{H}_u \mathcal{H}_d H]) &\simeq \frac{v^4}{M_Q^4} \text{Im} \left(\hat{V}_{ud}^L z_u z_d^* \right) y_u y_d \\ &+ \frac{v^8}{M_Q^8} \text{Im} \left(\hat{V}_{us}^L \hat{V}_{cd}^L \hat{V}_{cs}^{L*} z_u z_d^* \right) y_u y_d (y_c^2 - y_t^2) (y_s^2 - y_b^2). \end{aligned} \quad (6.32)$$

When the invariant of mass dimension 6 is zero, the invariant of mass dimension 10 is proportional to the imaginary part of the quartet. The vanishing of the 2 invariants implies CP conservation (in the realistic case of non-degeneracy). Again, we provide an equivalent proof in Appendix C.

6.4 Motivated scenarios

We now analyze four specific scenarios. The first two scenarios are relevant from the phenomenological point of view and they will be motivated in section 8. The last two address the extreme chiral limit, i.e. the limit of extremely high energies, where SM quark masses become negligible and CP violation can emerge only from NP sources.

6.4.1 With $N = 1$ doublet

In some scenarios, the first type of invariants built with three blocks may not be enough to describe the CP violation. As an illustration, let us consider the case in which there are no flavour changing neutral currents at tree level in one sector (up or down) while in the other sector one vector-like species couples only to two families. Let us consider a scenario which can be relevant for the Cabibbo angle anomaly related to kaon decays, with $z_u, z_s \neq 0$, in which the mass matrices can be written in the form

$$\mathcal{M}_u = \begin{pmatrix} \hat{y}_u v & 0 & 0 & 0 \\ 0 & \hat{y}_u v & 0 & 0 \\ 0 & 0 & \hat{y}_t v & 0 \\ z_u v & 0 & 0 & M_Q \end{pmatrix}, \quad \mathcal{M}_d = \begin{pmatrix} & & 0 & \\ & Y_d v & 0 & \\ & & 0 & \\ 0 & z_s v & z_b v & M_Q \end{pmatrix} \quad (6.33)$$

(we could consider in an analogous way $z_b = 0$, $z_t \neq 0$). In this scenario there still remain three physical phases. We have in this case

$$\begin{aligned}
& \frac{1}{M_Q^{2m+2n+2}} \text{Im}(\text{Tr}[\mathcal{H}_u^n \mathcal{H}_d^m H]) \\
&= \frac{v^4}{M_Q^4} \text{Im} \left(z_s^* z_u \hat{V}_{us}^L \right) y_u y_s (1 + k_{mn}) + \frac{v^4}{M_Q^4} \text{Im} \left(z_b^* z_u \hat{V}_{ub}^L \right) y_u y_s (1 + k'_{mn}) \quad (6.34) \\
&\simeq \frac{m_u m_s}{M_Q^2} \text{Im} (V_{us}^{R*} V_{us}^L) + \frac{m_u m_b}{M_Q^2} \text{Im} (V_{ub}^{R*} V_{ub}^L) .
\end{aligned}$$

Let us assume that in this parameterization the rephasing invariant quantities $V_{us}^L \hat{V}_{us}^{R*}$ and $V_{ub}^L \hat{V}_{ub}^{R*}$ are real. For instance, we can write \hat{V}_L in the Kobayashi-Maskawa parameterization and consider that V_{us}^L , V_{ub}^L , z_u , z_s and z_b are all real parameters. Then, all the invariants in the above category would vanish. However, there still is one physical phase which cannot be caught in these expressions. The physical phase of the model would show in the other categories of invariants, in particular the invariant of mass dimension 10 in eq. (6.7). The same effect manifests also in the scenario in which one doublet couples only to one generation, e.g. $z_b = 0$ in the previous example. In this case there are no flavour changing neutral currents and all the CP-odd invariants with two and three building blocks give null result. The one physical phase would show in the invariant of mass dimension 10 and the non-zero imaginary part would result from the rephasing invariants involving four quarks,

$$\frac{v^2}{M_Q^2} \text{Im} \left(\hat{V}_{\alpha i}^{L*} \hat{V}_{ui}^L \hat{V}_{\alpha s}^L z_u z_s^* \right) \simeq \text{Im} (V_{\alpha i}^{L*} V_{ui}^L V_{\alpha s}^L V_{us}^{R*}) . \quad (6.35)$$

We can also imagine a situation in which the extra doublet only couples to one sector, up or down, that is, the 3×3 submatrix of V_R involving SM quarks vanishes (only the last column or the last row survives) as well as one of the two matrices F_u (or F_d) depending on which sector is decoupled. Then, the invariants of the type in eq. (6.15) would give null result. The three physical CP phases would be contained in the rephasing invariant quantities associated with FCNCs,

$$\frac{v^2}{M_Q^2} \text{Im} \left(\hat{V}_{\alpha s}^L \hat{V}_{\alpha b}^{L*} z_s^* z_b \right) \simeq \text{Im} \left(\hat{V}_{\alpha s}^L \hat{V}_{\alpha b}^{L*} F_{sb}^d \right) . \quad (6.36)$$

In particular, these non vanishing quantities would indicate a contribution to CP violation, for instance, in neutral meson mixing in interference with the SM.

6.4.2 With $N = 2$ doublets

Let us consider a scenario in which two vector-like doublets of quarks couple only with the light generations. More specifically, let us imagine a specific pattern of couplings, with the doublets coupling only to the first generation in the up-sector, while coupling either with the first or the second in the down-sector. This texture will be phenomenologically

motivated in section 8 in the context of Cabibbo angle anomalies. The mass matrices can be written as:

$$\mathcal{M}_u = \begin{pmatrix} & \hat{U}_{uL} \hat{Y}_u v & 0 & 0 \\ z_{1u} v & 0 & 0 & M_Q \\ z_{2u} v & 0 & 0 & 0 & aM_Q \end{pmatrix}, \quad (6.37)$$

$$\mathcal{M}_d = \begin{pmatrix} & \hat{U}_{dL} \hat{Y}_d v & 0 & 0 \\ z_{1d} v & 0 & 0 & M_Q \\ 0 & z_{2s} v & 0 & 0 & aM_Q \end{pmatrix},$$

where \hat{U}_{qL} are the unitary matrices diagonalizing Y_q from the left, $\hat{U}_{qL}^\dagger Y_q Y_q^\dagger \hat{U}_{qL} = \hat{Y}_q$, and we are in the weak basis where the rotations \hat{U}_{uR} , \hat{U}_{dR} have already been applied (as in the rest of this section).

The mixing matrix of the right-handed charged currents reads, at order v^2/M_Q^2 :

$$V_R \simeq \begin{pmatrix} z_{1u}^* z_{1d} \frac{v^2}{M_Q^2} & z_{2u}^* z_{2s} \frac{v^2}{a^2 M_Q^2} & 0 & -z_{1u}^* \frac{v}{M_Q} & -z_{2u}^* \frac{v}{aM_Q} \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ -c_{45} z_{1d} \frac{v}{M_Q} & -s_{45}^* z_{2s} \frac{v}{aM_Q} & 0 & c_{45} & s_{45}^* \\ z_{1d} \left(s_{45} + c_{45} z_{1u}^* z_{2u} \frac{v}{aM_Q} \right) \frac{v}{M_Q} & -c_{45} z_{2s} \frac{v}{aM_Q} & 0 & -s_{45} - c_{45} z_{1u}^* z_{2u} \frac{v^2}{aM_Q^2} & c_{45} \end{pmatrix}, \quad (6.38)$$

where c_{45} is the real cosine and s_{45} is the complex sine of the angle,

$$s_{45} = \sin \theta_{45} \frac{z_{1u}^* z_{2u}}{|z_{1u} z_{2u}|}, \quad \frac{1}{2} \tan(2\theta_{45}) = \frac{a |z_{1u} z_{2u}| \frac{v^2}{M_Q^2}}{1 - a^2 + (|z_{1u}|^2 - |z_{2u}|^2) \frac{v^2}{M_Q^2}}. \quad (6.39)$$

For simplicity, in the lower-right 2×2 submatrix in eq. (6.38) we have omitted relative corrections of order $O(v^2/M_Q^2)$ i.e. $s_{45}(1 + O(v^2/M_Q^2)) \simeq s_{45}$ and $c_{45}(1 + O(v^2/M_Q^2)) \simeq c_{45}$.

As for the neutral currents, we have

$$F_u \simeq \begin{pmatrix} \left(|z_{1u}|^2 + \frac{|z_{2u}|^2}{a^2} \right) \frac{v^2}{M_Q^2} & 0 & 0 & - \left(z_{1u}^* c_{45} + \frac{z_{2u}^*}{a} s_{45} \right) \frac{v}{M_Q} & \left(z_{1u}^* s_{45} - \frac{z_{2u}^*}{a} c_{45} \right) \frac{v}{M_Q} \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ - \left(z_{1u} c_{45} + \frac{z_{2u}}{a} s_{45}^* \right) \frac{v}{M_Q} & 0 & 0 & 1 & O\left(\frac{v^2}{M_Q^2} \right) \\ \left(z_{1u} s_{45} - \frac{z_{2u}}{a} c_{45} \right) \frac{v}{M_Q} & 0 & 0 & O\left(\frac{v^2}{M_Q^2} \right) & 1 \end{pmatrix} \quad (6.40)$$

and

$$F_d \simeq \begin{pmatrix} |z_{1d}|^2 \frac{v^2}{M_Q^2} & 0 & 0 & -z_{1d}^* \frac{v}{M_Q} & 0 \\ 0 & |z_{2s}|^2 \frac{v^2}{a^2 M_Q^2} & 0 & 0 & -z_{2s}^* \frac{v}{a M_Q} \\ 0 & 0 & 0 & 0 & 0 \\ -z_{1d} \frac{v}{M_Q} & 0 & 0 & 1 - |z_{1d}|^2 \frac{v^2}{M_Q^2} & 0 \\ 0 & -z_{2s} \frac{v}{a M_Q} & 0 & 0 & 1 - |z_{2s}|^2 \frac{v^2}{a^2 M_Q^2} \end{pmatrix}. \quad (6.41)$$

In the limit where $a \neq 1$ and

$$\frac{|a^2 - 1|}{a} \gg \frac{v^2}{M_Q^2} |z_{1u} z_{2u}|, \quad |a^2 - 1| \gg \frac{v^2}{M_Q^2} \left| |z_{1u}|^2 - |z_{2u}|^2 \right|, \quad (6.42)$$

so that θ_{45} is a small angle and we have

$$s_{45} \simeq -z_{1u} z_{2u}^* \frac{a}{a^2 - 1} \frac{v^2}{M_Q^2}, \quad c_{45} \simeq 1, \quad (6.43)$$

then the right handed currents can be written as

$$V_R \simeq \begin{pmatrix} z_{1u}^* z_{1d} \frac{v^2}{M_Q^2} & z_{2u}^* z_{2s} \frac{v^2}{a^2 M_Q^2} & 0 & -z_{1u}^* \frac{v}{M_Q} & -z_{2u}^* \frac{v}{a M_Q} \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ -z_{1d} \frac{v}{M_Q} & \frac{z_{2s} z_{1u} z_{2u}^* v^3}{M_Q^3 (a^2 - 1)} & 0 & 1 - \frac{1}{2} \frac{(|z_{1d}|^2 + |z_{1u}|^2) v^2}{M_Q^2} & -\frac{z_{1u} z_{2u}^* a v^2}{M_Q^2 (a^2 - 1)} \\ -\frac{z_{1d} z_{1u}^* z_{2u} v^3}{M_Q^3 a (a^2 - 1)} & -z_{2s} \frac{v}{a M_Q} & 0 & \frac{z_{1u}^* z_{2u} v^2}{M_Q^2 a (a^2 - 1)} & 1 - \frac{1}{2} \frac{(|z_{2s}|^2 + |z_{2u}|^2) v^2}{a^2 M_Q^2} \end{pmatrix}, \quad (6.44)$$

at order v^2/M_Q^2 . The CAAs can be explained in the presence of the couplings

$$\begin{aligned} V_{ud}^R &\simeq z_{1d}^* z_{1u} \frac{v^2}{M_Q^2} \simeq -0.78(27) \times 10^{-3}, \\ V_{us}^R &\simeq z_{2s}^* z_{2u} \frac{v^2}{a^2 M_Q^2} \simeq -1.26(38) \times 10^{-3}, \end{aligned} \quad (6.45)$$

while flavour changing neutral currents are suppressed since only diagonal elements are present in the 3×3 submatrix of F_u and F_d (see Refs. [45, 49]).

In what concerns the left rotations, the unitary matrices can be written as

$$\begin{aligned}
V_L \simeq & \begin{pmatrix} 1 & 0 & 0 & z_{1u}^* \hat{y}_u \frac{v^2}{M_Q^2} & z_{2u}^* \hat{y}_u \frac{v^2}{a^2 M_Q^2} \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ -z_{1u} \hat{y}_u \frac{v^2}{M_Q^2} & 0 & 0 & 1 & z_{1u} z_{2u}^* \frac{v^2}{M_Q^2 (a^2 - 1)} \\ -z_{2u} \hat{y}_u \frac{v^2}{a^2 M_Q^2} & 0 & 0 & -z_{1u}^* z_{2u} \frac{v^2}{M_Q^2 (a^2 - 1)} & 1 \end{pmatrix} \\
& \times \begin{pmatrix} & 0 & 0 & & \\ \hat{V}_L & 0 & 0 & & \\ & 0 & 0 & & \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & z_{1d}^* \hat{y}_d \frac{v^2}{M_Q^2} & 0 \\ 0 & 1 & 0 & 0 & z_{2s}^* \hat{y}_s \frac{v^2}{a^2 M_Q^2} \\ 0 & 0 & 1 & 0 & 0 \\ -z_{1d} \hat{y}_d \frac{v^2}{M_Q^2} & 0 & 0 & 1 & 0 \\ 0 & -z_{2s} \hat{y}_s \frac{v^2}{a^2 M_Q^2} & 0 & 0 & 1 \end{pmatrix}, \tag{6.46}
\end{aligned}$$

with $\hat{V}_L \simeq \hat{U}_{uL}^\dagger \hat{U}_{dL}$, in the limit of eqs. (6.42) and (6.43).

As for the $M = 6$ CP-odd weak-basis invariant we have

$$\begin{aligned}
\frac{1}{M_Q^6} \text{Im Tr}[\mathcal{H}_u \mathcal{H}_d H] &= \frac{v^4}{M_Q^4} \text{Im} \left(z_{1d}^* z_{1u} \hat{V}_{ud}^L \right) \hat{y}_u \hat{y}_d + \frac{v^4}{M_Q^4} \text{Im} \left(z_{2s}^* z_{2u} \hat{V}_{us}^L \right) a^2 \hat{y}_u \hat{y}_s \\
&\simeq \frac{m_u m_d}{M_Q^2} \text{Im} (V_{ud}^{R*} V_{ud}^L) + \frac{m_u m_s}{M_Q^2} a^4 \text{Im} (V_{us}^{R*} V_{us}^L), \tag{6.47}
\end{aligned}$$

which is valid even when $a = 1$. One should also note that, the vanishing of this category of CP-odd weak-basis invariants would not imply absence of CP-violation. The missing physical phase is contained in other rephasing invariant quantities as $\hat{V}_{\alpha i}^{L*} \hat{V}_{\beta i}^L \hat{V}_{\alpha j}^L z_\beta z_j^*$, which is contained in the CP-odd invariant of mass dimension 10 eq. (6.7).

6.4.3 The extreme chiral limit with $N = 1$ doublet

Let us consider the limit in which only the third generation has non-zero mass. This so-called extreme chiral limit (ECL) is relevant when considering scenarios with extremely high energies, where to good approximation the masses of the first two generations of quarks can be taken as zero. Notably, in VLQ models CPV is possible in this limit, in contrast to what happens in the SM. For instance, the VLQ singlet scenario has been studied in the ECL [56, 58], and it was found that one independent CP-odd WBI still survives.

In the ECL, for the doublet case, the mass matrices can take the form:

$$\mathcal{M}_u = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \hat{y}_t v & 0 \\ 0 & z_c v & z_t v & M_Q \end{pmatrix}, \quad \mathcal{M}_d = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & y_{23} v & 0 \\ 0 & 0 & y_{33} v & 0 \\ 0 & z_s v & z_b v & M_Q \end{pmatrix}. \tag{6.48}$$

Note that all phases except one can be absorbed by rephasing of quarks fields, so that there is a single physical phase which cannot be eliminated, and we can consider all parameters to be real except e.g. z_b . Similarly to what happens in the VLQ singlet scenario, this single surviving phase induces CPV which means all CP properties of the model are determined by a single CP-odd WBI built from 3 Hermitian blocks. In this case, that invariant can take the simple form

$$\frac{1}{M_Q^6} \text{Im Tr} [\mathcal{H}_u \mathcal{H}_d H] = \frac{v^4}{M_Q^4} \text{Im} (\hat{y}_t y_{33} z_t z_b^*) \simeq \frac{m_t m_b}{M_Q^2} \text{Im} (V_{tb}^{R*} V_{tb}^L), \quad (6.49)$$

where we make use of eqs. (4.10), (4.18) and (4.23), as well as the relation $y_{33} = \hat{y}_b \hat{V}_{tb}^L \simeq \hat{y}_b V_{tb}^L$, with $\hat{y}_b^2 = y_{23}^2 + y_{33}^2$, which arises from the diagonalization of the down sector mass matrix.

6.4.4 The extreme chiral limit with $N = 2$ doublets

We now briefly present an even more extreme limit of high energies, where the top quark is the only standard quark that has non-zero mass. In that case, the mass matrices can take the minimal form:

$$\mathcal{M}_u = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \hat{y}_t v & 0 & 0 \\ 0 & z_{1c} v & z_{1t} v & M_Q & 0 \\ z_{2u} v & z_{2c} v & z_{2t} v & 0 & a M_Q \end{pmatrix}, \quad \mathcal{M}_d = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & z_{1b} v & M_Q & 0 \\ 0 & z_{2s} v & z_{2b} v & 0 & a M_Q \end{pmatrix}, \quad (6.50)$$

and all phases except two can be eliminated via rephasings of the quark fields. Hence, we can write

$$\frac{1}{M_Q^6} \text{Im Tr} [\mathcal{H}_u \mathcal{H}_d H] = \frac{v^4}{M_Q^4} (a^2 - 1) \text{Im} [z_{1b} z_{2b}^* (z_{1c}^* z_{2c} + z_{1t}^* z_{2t})], \quad (6.51)$$

so that in general there is still CPV, even when $\hat{y}_t = 0$ (i.e. the limit of vanishing top mass). Note that, although this invariant vanishes for $a = 1$, in that case we still have $\text{Im Tr} [\mathcal{H}_u \mathcal{H}_d \mathcal{H}_u^2 H] \neq 0$, meaning that CPV is still possible as far as $\hat{y}_t \neq 0$.

7 Integrating out the VLQs

7.1 SMEFT description

Below the VLQ mass thresholds, physical observables can be obtained from an effective field theory (EFT) that does not include the VLQ fields as explicit degrees of freedom. The footprints of these fields, as well as those from any other heavy BSM particle, are fully captured by c -numbers, known as Wilson coefficients, which are suppressed by powers of the heavy masses. Under general assumptions, this description is provided by the Standard Model effective field theory (SMEFT), which extends the SM Lagrangian to incorporate operators of dimension greater than four ($D > 4$):

$$\mathcal{L}_{\text{SMEFT}} = \mathcal{L}_{\text{SM}} + \sum_{i, D > 4} C_{i,D} \mathcal{O}_{i,D}, \quad (7.1)$$

where $\mathcal{O}_{i,D}$ are operators of mass dimension D constructed from SM fields and $C_{i,D}$ are the corresponding Wilson coefficients. Their contributions to the amplitudes are suppressed by powers of the heavy masses, $1/M_{\text{BSM}}^{D-4}$, and thus, by truncating at $D = 6$, one can obtain the leading $\mathcal{O}(1/M_{\text{BSM}}^2)$ corrections to the amplitudes. The resulting amplitudes in terms of the Wilson coefficients $C_{i,D}$ are identical for any heavy extension of the SM, and one can consistently compute them at any loop order. Provided that these amplitudes and that the corresponding observables are known at a given order in terms of the Wilson coefficients, one needs only to match them to the parameters of the ultraviolet (UV) model at the required level of approximation. Apart from providing a clean comparison of the phenomenological consequences of different UV models, this can simplify the low-energy phenomenology (see next section).

The tree-level SMEFT Wilson coefficients for the different UV completions of the SM, including VLQ doublets, can be found for example in Ref. [71].¹⁸ Let us thus start from the Lagrangian in the form of eq. (3.5),¹⁹

$$\begin{aligned} -\mathcal{L} = & (Y_u)_{ij} \bar{q}_{Li}^0 \tilde{\Phi} u_{Rj}^0 + (Y_d)_{ij} \bar{q}_{Li}^0 \Phi d_{Rj}^0 + \text{h.c.} \\ & + (Z_u)_{\alpha j} \bar{Q}_{L\alpha}^0 \tilde{\Phi} u_{Rj}^0 + (Z_d)_{\alpha j} \bar{Q}_{L\alpha}^0 \Phi d_{Rj}^0 + \text{h.c.} \\ & + (D_Q)_{\alpha\beta} \bar{Q}_{L\alpha}^0 Q_{R\beta}^0 + \text{h.c.} \end{aligned} \quad (7.2)$$

As shown in section 3.1.1, a minimal parameterization is obtained taking Y_u diagonal and Y_d as the product of a CKM-like unitary matrix times a diagonal matrix, plus setting to zero N phases from Z_d (e.g. in its first column), thus avoiding redundancies that would

¹⁸While for the EFT applications of this work tree-level results suffice, it is worth mentioning that some partial results regarding the one-loop matching corrections can be found in the literature, for example in Refs. [50, 72]. With the fast development of automated tools [73], such as `MatchMakerEFT` [74] or `Matchete` [75], finding the SMEFT expressions in the same (Warsaw) basis and implementing them consistently all the way to the observable level will soon become straightforward.

¹⁹One could write it in the more general form of eq. (2.2) by simply undoing the set of unitary transformations that leads to this form.

translate into unphysical flat directions in the different fits. Finding the mapping of any weak basis to this minimal one was done in section 5.2.2, by making use of weak-basis invariants. One can thus immediately obtain the EFT parameters in terms of any set of values for the Yukawa matrices of the original UV VLQ completion.

After integrating them out, the low-energy tree-level effect of N VLQ doublets, up to $D = 6$, is fully encoded in

$$\begin{aligned} \mathcal{L}_{\text{SMEFT}} \supset & (C_{\Phi u})_{ij} (\Phi^\dagger i \overleftrightarrow{D}_\mu \Phi) \overline{u_{Ri}^0} \gamma^\mu u_{Rj}^0 + (C_{\Phi d})_{ij} (\Phi^\dagger i \overleftrightarrow{D}_\mu \Phi) \overline{d_{Ri}^0} \gamma^\mu d_{Rj}^0 \\ & + [(C_{\Phi ud})_{ij} (\tilde{\Phi}^\dagger i D_\mu \Phi) \overline{u_{Ri}^0} \gamma^\mu d_{Rj}^0 + \text{h.c.}] \\ & + [(C_{u\Phi})_{ij} (\Phi^\dagger \Phi) \overline{q_{Li}^0} \tilde{\Phi} u_{Rj}^0 + (C_{d\Phi})_{ij} (\Phi^\dagger \Phi) \overline{q_{Li}^0} \tilde{\Phi} d_{Rj}^0 + \text{h.c.}], \end{aligned} \quad (7.3)$$

with

$$\begin{aligned} C_{\Phi u} &= -\frac{1}{2v^2} Z_u^\dagger \frac{v^2}{D_Q^2} Z_u, \quad C_{\Phi d} = \frac{1}{2v^2} Z_d^\dagger \frac{v^2}{D_Q^2} Z_d, \\ C_{\Phi ud} &= Z_u^\dagger \frac{1}{D_Q^2} Z_d, \\ C_{u\Phi} &= \frac{1}{2v^2} Y_u Z_u^\dagger \frac{v^2}{D_Q^2} Z_u, \quad C_{d\Phi} = \frac{1}{2v^2} Y_d Z_d^\dagger \frac{v^2}{D_Q^2} Z_d. \end{aligned} \quad (7.4)$$

Notice how, regardless of the number of doublets one has, the full low-energy effect at tree level and of $\mathcal{O}(1/M_Q^2)$, where $M_Q = M_{\text{BSM}}$, is encoded into three 3×3 matrices, two of which are Hermitian, dramatically restricting the number of linear combinations of physical parameters entering this regime. Additionally, if the number of VLQ doublets is small, one finds simple relations between the parameters in the charged-current sector and the ones in the neutral-current one.

7.2 Electroweak symmetry breaking

After EWSB, the pure-VEV terms in eq. (7.3) modify both the effective quark gauge currents and the mass matrices,

$$\mathcal{L} \supset \left[-\frac{g}{\sqrt{2}} W_\mu^+ J_W^\mu + \text{h.c.} \right] - \left[\sum_{q=u,d} \overline{q_L^0} M_{\text{eff}}^q q_R^0 + \text{h.c.} \right] - \frac{g}{2c_W} Z_\mu J_Z^\mu. \quad (7.5)$$

The SM Higgs-fermion interactions are also modified,

$$\mathcal{L} \supset -\frac{h}{\sqrt{2}} \sum_{q=u,d} \overline{q_L^0} Y_{\text{eff}}^q q_R^0. \quad (7.6)$$

One has²⁰

$$M_{\text{eff}}^q = v Y_q \left[\mathbb{1} - \frac{1}{2} \hat{F}_q \right], \quad (7.7)$$

$$Y_{\text{eff}}^q = Y_q \left[\mathbb{1} - \frac{3}{2} \hat{F}_q \right] = \frac{M_{\text{eff}}^q}{v} \left[\mathbb{1} - \hat{F}_q \right], \quad (7.8)$$

and

$$\begin{aligned} J_W^\mu &= \bar{\nu}_L \gamma^\mu e_L + \bar{u}_L^0 \gamma^\mu d_L^0 + \bar{u}_R^0 \gamma^\mu \hat{V}_R d_R^0, \\ J_Z^\mu &= J_{Z,\text{SM}}^\mu + \bar{u}_R^0 \gamma^\mu \hat{F}_u u_R^0 - \bar{d}_R^0 \gamma^\mu \hat{F}_d d_R^0, \end{aligned} \quad (7.9)$$

where

$$J_{Z,\text{SM}}^\mu \supset \bar{u}_L^0 \gamma^\mu u_L^0 - \bar{d}_L^0 \gamma^\mu d_L^0 - 2s_W^2 \left(\frac{2}{3} \bar{u}^0 \gamma^\mu u^0 - \frac{1}{3} \bar{d}^0 \gamma^\mu d^0 \right), \quad (7.10)$$

with $u^0 = u_L^0 + u_R^0$ and $d^0 = d_L^0 + d_R^0$. We are omitting the well-known leptonic part of $J_{Z,\text{SM}}^\mu$, which is not used in this work.

Without loss of generality, from now on we can work in the weak basis in which $Y_u = \hat{Y}_u$ is diagonal and $Y_d = \hat{V} \hat{Y}_d$, i.e., a CKM-like unitary matrix (free from external phases) times a diagonal matrix, so that $Z_q \rightarrow z_q$ (corresponding to the trivial generalization of eq. (4.6) for N doublets). In eqs. (7.7) to (7.9) we have used

$$\hat{F}_u \equiv z_u^\dagger \frac{v^2}{D_Q^2} z_u, \quad \hat{F}_d \equiv z_d^\dagger \frac{v^2}{D_Q^2} z_d, \quad \hat{V}_R \equiv z_u^\dagger \frac{v^2}{D_Q^2} z_d, \quad (7.11)$$

generalizing the definitions introduced in eq. (4.29) and footnote 9 for the $N = 1$ case.

The resulting mass matrices are not proportional to Y_q and thus do not inherit the same form. Nevertheless we can perform the infinitesimal unitary field redefinitions

$$\begin{aligned} u_L^0 &\rightarrow (\mathbb{1} + i \delta L_q) u_L^0, & d_L^0 &\rightarrow (\mathbb{1} + i \delta L_q) d_L^0, \\ u_R^0 &\rightarrow (\mathbb{1} + i \delta R_u) u_R^0, & d_R^0 &\rightarrow (\mathbb{1} + i \delta R_d) d_R^0, \end{aligned} \quad (7.12)$$

where the infinitesimal matrices are $\mathcal{O}(1/M_Q^2)$. Unitarity requires them to be Hermitian. Since $D \geq 8$ corrections of $\mathcal{O}(1/M_Q^4)$ are neglected, any effect of this redefinition in the terms above that already contain an $\mathcal{O}(1/M_Q^2)$ insertion can be neglected. The only effect of the field redefinition at the working order is then in eqs. (7.7) and (7.8),

$$M_{\text{eff}}^u = v (\mathbb{1} - i \delta L_q) \hat{Y}_u \left[\mathbb{1} - \frac{1}{2} \hat{F}_u \right] (\mathbb{1} + i \delta R_u), \quad (7.13)$$

$$M_{\text{eff}}^d = v (\mathbb{1} - i \delta L_q) \hat{V} \hat{Y}_d \left[\mathbb{1} - \frac{1}{2} \hat{F}_d \right] (\mathbb{1} + i \delta R_d). \quad (7.14)$$

²⁰Note that in this work we are defining the mass (and Yukawa) terms using a left-right convention, i.e. $\mathcal{L} \sim -\bar{\psi}_L \mathcal{M} \psi_R$, while in SMEFT works (see e.g. [76]) they are typically defined using a right-left convention, $\mathcal{L} \sim -\bar{\psi}_R \mathcal{M} \psi_L$. Notice also the different VEV normalization.




Figure 1: Graphical representation of the VLQ diagrams generating the new effective gauge interactions.

Taking appropriate unitary matrices, one can always find a basis where $M_{\text{eff}}^u = D^u \equiv \text{diag}(m_u, m_c, m_t)$ and $M_{\text{eff}}^d = \hat{V}_L D^d \equiv \hat{V}_L \text{diag}(m_d, m_s, m_b)$, up to (but not including) $\mathcal{O}(1/M_Q^4)$ effects. Notice that $\hat{V}_L \neq \hat{V}$ in general. Then, in the mass basis and at this order, the effective Lagrangian is described by

$$M_{\text{eff}}^q = D^q, \quad Y_{\text{eff}}^q = \frac{D^q}{v} \left[\mathbb{1} - \hat{F}_q \right], \quad (7.15)$$

and

$$\begin{aligned} J_W^\mu &= \bar{\nu}_L \gamma^\mu e_L + \bar{u}_L \gamma^\mu \hat{V}_L d_L + \bar{u}_R \gamma^\mu \hat{V}_R d_R, \\ J_Z^\mu &= J_{Z,\text{SM}}^\mu + \bar{u}_R \gamma^\mu \hat{F}_u u_R - \bar{d}_R \gamma^\mu \hat{F}_d d_R, \end{aligned} \quad (7.16)$$

consistently with the full-theory result, see e.g. eqs. (4.19) and (4.20). The matrix \hat{V}_L can be parameterized as the usual CKM matrix and thus, at order $\mathcal{O}(1/M_Q^2)$, the tree-level LH sector is left unmodified, cf. eq. (4.26).

A diagrammatic interpretation of the origin of the new vertices is depicted in Figure 1. For completeness, we relate \hat{V}_L and $D^{u,d}$ to the original Lagrangian parameters in Appendix B.

7.3 Low-energy EFT

Below the EW scale, the top quark and the electroweak bosons, as well as the Higgs, cannot be directly produced. Physical observables, both within and beyond the SM,

can, once again, be obtained from an EFT free from the top quark, W and Z bosons or any other heavy BSM particle. In this case, the Wilson coefficients, now suppressed by $\mathcal{O}(1/M_Q^2, 1/M_{W,Z}^2, 1/m_t^2, 1/m_h^2)$ factors, also contain pure-SM contributions, whose corresponding coefficients can be explicitly computed in terms of the heavy SM masses,

$$\mathcal{L}_{\text{LEFT}} = \mathcal{L}_{\text{QCD+QED}} + \sum_{i,D>4} (C_{i,D}^{\text{SM}} + C_{i,D}^{\text{BSM}}) \mathcal{O}_{i,D}. \quad (7.17)$$

At tree level, the main operators induced both in the SM and in our VLQ scenario are the four-fermion operators generated by the interactions of two-fermion currents mediated by EW bosons.²¹ In that approximation, one simply recovers a generalized version of the Fermi theory,

$$\mathcal{L}_{\text{LEFT}} \supset \mathcal{L}_{\text{QCD+QED}} - \frac{1}{v^2} J_W^\mu J_{W,\mu}^\dagger - \frac{1}{4v^2} J_Z^\mu J_{Z,\mu}. \quad (7.18)$$

Any process mediated in the SM by EW currents is suppressed by $1/v^2$ and thus, even at very low energies, VLQ interactions are suppressed with respect to the SM only by $\mathcal{O}(v^2/M_Q^2)$. In practice, we do not need to limit ourselves to the approximation of eq. (7.18) for the SM computation, but can use the results of SM works at all known orders. Instead, eq. (7.18) is often sufficient to compute BSM effects, since the neglected contributions are suppressed by both loop factors and powers of v^2/M_Q^2 .²²

²¹Higgs exchanges are also tree-level, but they are suppressed by $1/m_h^2$ times either light Yukawas or extra BSM factors.

²²One should however keep in mind that, when quark currents are involved, it is important to use renormalization group equations to, at least, re-sum large logarithms associated to gluonic corrections $\sum_n \alpha_s^n(\mu) (\ln M_W^2/\mu^2)^n$, since at the typical scale μ of the corresponding hadronic processes they can easily become $\mathcal{O}(1)$. The corresponding running equations can be found e.g. in Ref. [76].

8 Phenomenological effects of mixing with VLQ doublets

The phenomenological effects of the presence of vector-like quarks have been the object of study of several works (see [29–45]). However, it is compelling to conduct a comprehensive phenomenological analysis of scenarios with VLQ doublets adopting a new perspective based on invariants.

Invariants under WBTs have been a main object of study in this work. Any observable, as any minimal set of parameters in a specific weak basis, can ultimately be written in terms of weak basis invariants. The possibility of such description in terms of invariants is intriguing, as it would avoid any potential misinterpretations due to the choice of weak basis (minimal or not). This feature appears particularly attractive when describing CP-violating phenomena, since the results cannot be the consequence of any parameterization choices as well as of any unphysical complex phase. Moreover, the different structures of invariants emerging in presence of VLQ doublets compared to the SM, as studied in detail in sections 5 and 6, provide hints on the processes which one can expect to be most affected, particularly when involving CP-violation.

In this section, we show how different observables can be written in terms of the invariant quantities analyzed in the previous sections, and how one can gain clearer insight into which observables are modified, as well as how CP-even and CP-odd probes relate. This exercise also highlights the distinct imprints of VLQ doublets compared to other models that generate similar invariant structures.

We start by summarising the low-energy effects of VLQ doublets in section 8.1. In section 8.2, we identify effective rephasing invariants as the most relevant objects for low-energy phenomenology, which provide a glimpse on which kind of observables are going to be more sensitive to VLQs prior to any computation. In sections 8.3 and 8.4, we study the most relevant phenomena in terms of these rephasing invariants, in particular bilinears and trilinears respectively. In this way, we can provide unambiguous constraints on the model and a clearer connection between different observables.

8.1 Weak currents

Let us first summarize the effects of the presence of vector-like quarks doublets mixing with SM quarks (see also sections 4.1 and 7). Weak charged and neutral currents are modified as

$$\begin{aligned} \mathcal{L} \supset & -\frac{g}{\sqrt{2}} W_\mu^+ \left(\bar{\nu}_L \gamma^\mu e_L + \bar{u}_L \gamma^\mu \hat{V}_L d_L + \bar{u}_R \gamma^\mu \hat{V}_R d_R \right) + \text{h.c.} \\ & -\frac{g}{2c_W} Z_\mu \left(J_\mu^{Z,\text{SM}} + \bar{u}_R \gamma^\mu \hat{F}_u u_R - \bar{d}_R \gamma^\mu \hat{F}_d d_R \right). \end{aligned} \quad (8.1)$$

The whole effect of vector-like quarks is captured by the matrices \hat{V}_L , \hat{V}_R , \hat{F}_u , \hat{F}_d in good approximation (see eqs. (4.19), (4.20) and (7.11)). In particular, it is evident from the description in sections 4.1 and 7 that the set of Lagrangian parameters z_α , z_i is directly connected to couplings of charged and neutral currents, and hence to observables. For

N VLQ doublets one has

$$\begin{aligned} \hat{V}_{\alpha i}^R &\equiv \sum_n z_{n\alpha}^* z_{ni} \frac{v^2}{[D_Q]_{nn}^2}, \\ \hat{F}_{\alpha\beta}^u &= \sum_n z_{n\alpha}^* z_{n\beta} \frac{v^2}{[D_Q]_{nn}^2}, \quad \hat{F}_{ij}^d = \sum_n z_{ni}^* z_{nj} \frac{v^2}{[D_Q]_{nn}^2}, \end{aligned} \quad (8.2)$$

where the mass elements $[D_Q]_{nn}$ are defined in eq. (3.5), the couplings $z_{ni/\alpha}$ are defined analogously to eqs. (4.6) and (7.11) and the index n runs from 4 to $3 + N$, while $\alpha, \beta = u, c, t$ and $i, j = d, s, b$. Then:

- FCNCs are generated at tree level, in contrast to the SM, where they only appear at loop level and are further suppressed by the Glashow–Iliopoulos–Maiani (GIM) mechanism [77–79]. The mixing of SM quarks with the vector-like quarks is thus strongly constrained.
- The new couplings introduce potential CP-violating RH quark currents at low energies, only suppressed by v^2/M_Q^2 .

Let us emphasize that the RH *charged* currents are intrinsically linked to the RH *neutral* currents, as both depend on the same Yukawa parameters. This connection is particularly relevant when comparing the phenomenological consequences of VLQ doublets with other models that also generate effective RH currents in the light-quark sectors. The relationship becomes even stronger with a lower number of doublets. In fact, for the case of a single doublet, knowing all non-zero couplings associated with FCNCs would fully determine the new Yukawa parameters, including the allowed CP-violation.

8.2 Phenomenology of effective rephasing invariants

Establishing which WBIs will be more or less relevant in low-energy observables is not trivial, due to the strong hierarchy of the SM parameters and the possible presence of hierarchy in VLQ couplings. Nevertheless, in section 6 we have seen how in a scenario with one vector-like doublet (and some phenomenologically motivated scenario with more doublets) starting from the UV description and expanding the mixing matrices in powers of v/M_Q , WBIs are naturally related to rephasing invariants involving only the three SM quarks (see section 4.1.3). This dependence provides a direct connection to low-energy observables, which can always be written in terms of rephasing invariants.

In a scenario with one or more vector-like doublets, we can combine the restrictions that rephasing invariants provide on the form that any observable may take with the power counting given by the effective description at the amplitude level. In this way, we can find some physical insight at the observable level prior to any computation, especially regarding CP-violating probes.²³ Let us thus recall the different possible rephasing invariants appearing in the effective description.

²³Let us note how this exercise can be generalized to other models generating effective RH currents. This generalization goes beyond the reach of this work.

The mixing matrix \hat{V}_L appears at dimension $D = 4$ in the effective Lagrangian. In principle one may think that up to 6 phases can appear in \hat{V}_L . However, only one (physical) of them can be obtained from rephasing invariants with vertex insertions appearing only at $D = 4$, that is only with \hat{V}_L insertions. In fact, the lowest rephasing invariant which can be obtained from the elements of \hat{V}_L is the SM-like one, corresponding to the imaginary part of the left-handed rephasing invariant quartet:

$$Q_{\alpha i \beta j} = \hat{V}_{\alpha i}^L \hat{V}_{\beta j}^L \hat{V}_{\alpha j}^{L*} \hat{V}_{\beta i}^{L*}. \quad (8.3)$$

In the limit $M_Q \rightarrow \infty$ (M_Q the lowest mass scale of vector-like quark doublets), the imaginary part $\text{Im}(Q_{\alpha i \beta j})$ reduces to the J invariant of eq. (3.24), and thus to the one physical CP phase of the SM with the same 4–vertex/4–quark SM suppression.

At $\mathcal{O}(1/M_Q^2)$ (but with different suppression of standard model mixing) one finds the new CP-odd structures given by the imaginary part of the following rephasing invariants:

$$\begin{aligned} - \text{bilinears:} \quad & \hat{\mathcal{B}}_{\alpha i} = \hat{V}_{\alpha i}^{L*} \hat{V}_{\alpha i}^R, \\ - \text{trilinears:} \quad & \hat{\mathcal{T}}_{i, \alpha \beta} = \hat{V}_{\alpha i}^{L*} \hat{V}_{\beta i}^L \hat{F}_{\alpha \beta}^u, \quad \hat{\mathcal{T}}_{\alpha, ij} = \hat{V}_{\alpha i}^L \hat{V}_{\alpha j}^{L*} \hat{F}_{ij}^d, \\ - \text{quartets:} \quad & \hat{\mathcal{Q}}_{\alpha i \beta j} = \hat{V}_{\beta j}^L \hat{V}_{\alpha j}^{L*} \hat{V}_{\beta i}^{L*} \hat{V}_{\alpha i}^R, \end{aligned} \quad (8.4)$$

corresponding to the ones defined in eq. (4.29) for one VLQ doublet. The following information can be gathered from the presence of the invariants in eq. (8.4):

- The imaginary part of the bilinears $\text{Im} \hat{\mathcal{B}}_{\alpha i}$ is directly linked to potential CP violation in the charged-current sector, produced by a phase shift of the BSM right-handed sector with respect to the corresponding SM (left-handed) one.
- In the presence of VLQs, $\hat{F}_{u(d)}$ induces tree-level FCNCs with two $\hat{V}_{\alpha i}^L$ insertions. A phase difference between the SM amplitude (which is always induced at loop level) and the new contributions can induce CP violation in the flavour-changing up or down sector driven respectively by $\text{Im} \hat{\mathcal{T}}_{i, \alpha \beta}$ and $\text{Im} \hat{\mathcal{T}}_{\alpha, ij}$.
- The last type of invariant comes with the same insertions as in the SM, but with a right-handed vertex. In general, there will be stronger contributions to the same process generated by the phases of bilinears and trilinears.

Let us mention that one cannot get CP-odd rephasing invariants at order $\mathcal{O}(1/M_Q^4)$ through the interference of two tree-level $\mathcal{O}(1/M_Q^2)$ amplitudes in the right-handed sector alone. Namely, CP-odd rephasing invariants cannot be made by two right-handed vertices (with elements of the mixing matrices \hat{F}_u , \hat{F}_d , \hat{V}_R) without CKM vertex insertions. Moreover, in presence of only one vector-like doublet, one cannot obtain new CP-odd rephasing invariants with more than one insertion of a right-handed vertex.

In the following, we analyze how the new CP-violating invariant types directly relate to physical processes, focusing on some key flavour observables involving light quarks.

8.3 Bilinears and the Cabibbo sector

The bilinears $\hat{\mathcal{B}}_{\alpha i} = \hat{V}_{\alpha i}^{L*} \hat{V}_{\alpha i}^R$ could potentially induce the largest VLQ imprints, both in their CP-conserving $\text{Re } \hat{\mathcal{B}}_{\alpha i}$ and CP-violating $\text{Im } \hat{\mathcal{B}}_{\alpha i}$ part, since they can appear in the charged current sectors at $\mathcal{O}(1/M_Q^2)$ at tree level. The presence of these right-handed couplings acquires highly motivated phenomenological importance given the presence of the Cabibbo angle anomalies, that is the tensions between three different determinations of the Cabibbo angle θ_C . In fact, it was shown that the vector-like quark charged as a doublet of $SU(2)_L$ appears as the favoured candidate which can explain the Cabibbo angle anomalies [45, 49–51].

8.3.1 Cabibbo angle anomalies

Three types of independent determinations of the Cabibbo angle θ_C are extracted with high precision and show tension between each other. In particular, the determination of $|V_{ud}| = \cos \theta_C$ is obtained from β decays. Recent calculations of short-distance radiative corrections (see e.g. [80, 81]) in β -decays led to an improved determination of $|V_{ud}|$. The most precise determination is obtained from super-allowed $0^+ - 0^+$ nuclear β decays, which are pure Fermi transitions [82] (despite in this last survey the uncertainty is increased by a factor of 2.6 due to new contributions in nuclear structure corrections δ_{NS} [80, 83], which now dominate the uncertainty). One can determine $|V_{ud}|$ also from free neutron β -decay, using the average of neutron lifetime from eight bottle experiments [84–91] (with rescaled uncertainty) and the axial-vector coupling from the latest experiments measuring parity-violating β -asymmetry parameter A from polarized neutrons [92–94]. Present lattice computations [95, 96] and analysis of electromagnetic radiative corrections [97, 98] provide a precise determination of $|V_{us}| = \sin \theta_C$ from semi-leptonic $K_{\ell 3}$ kaon decays $K \rightarrow \pi \ell \nu$ (K_{Le3} , K_{Se3} , K_{e3}^\pm , $K_{\mu 3}^\pm$, $K_{L\mu 3}$, $K_{S\mu 3}$) [99]. The ratio $|V_{us}/V_{ud}| = \tan \theta_C$ can be independently determined from the ratio of the kaon and pion leptonic decay rates $K_{\mu 2}$ and $\pi_{\mu 2}$, i.e. $K \rightarrow \mu \nu(\gamma)$ and $\pi \rightarrow \mu \nu(\gamma)$ [100] using lattice QCD calculations [95, 96] and including electroweak radiative corrections [101–103]. We get:²⁴

$$|V_{ud}|_\beta = 0.97372(26) , \quad |V_{us}|_{K\ell 3} = 0.22308(55) , \quad \left| \frac{V_{us}}{V_{ud}} \right|_{K\mu 2/\pi\mu 2} = 0.23126(48) . \quad (8.5)$$

After using these determinations, there is approximately $\sim 3\sigma$ deficit in the CKM first row unitarity (dubbed CAA1 in [45]) when using the value of $|V_{ud}|$ from β -decays with the value of $|V_{us}|$ from kaon decays. Additionally, there is about $\sim 3\sigma$ tension between the determination of $|V_{us}|$ from $K_{\ell 3}$ decays and the one obtained from leptonic $K_{\mu 2}$ and $\pi_{\mu 2}$ decay rates (CAA2 in Ref. [45]).

Various models have been suggested as possible solutions [42–44, 49, 50, 104–119]. A remarkable solution for the anomalies is given by the vector-like quark charged as a

²⁴We follow Ref. [45] updating the result $f_K/f_\pi = 1.1934(19)$ from ref. [96]. The obtained determinations are also in good agreement with the fit made in Ref. [51], with a slightly different choice of inputs.

doublet of $SU(2)_L$ as found in Refs. [45, 49, 50], which can potentially explain all the tensions through right-handed charged currents [45, 49–51, 104].

In fact, one must take into account that the value of V_{ud} which is obtained from β decays depends uniquely on the vector part of the weak interaction $G_V = G_F|V_{ud}|$. Also semi-leptonic kaon decays $K_{\ell 3}$ determine the weak vector coupling, while leptonic decays $K_{\mu 2}$ and $\pi_{\mu 2}$ depend on the axial-vector current. In the SM there is no difference between these determinations since they are all given by left-handed charged currents and they determine the same mixing angle θ_C . In a model with right-handed currents instead, vector and axial-vector couplings are not equal and the three determinations correspond to different couplings. In the scenario with vector-like doublets right-handed currents are generated. Then, the tensions in the different determinations can be explained by the mixing of vector-like doublets with light standard model quarks. From eq. (4.1) we have

$$\mathcal{L}_{cc} \supset -\frac{g}{\sqrt{2}} W_\mu^+ \left[\bar{u}_L \gamma^\mu \hat{V}_{ud}^L d_L + \bar{u}_L \gamma^\mu \hat{V}_{us}^L s_L + \bar{u}_R \gamma^\mu \hat{V}_{ud}^R d_R + \bar{u}_R \gamma^\mu \hat{V}_{us}^R s_R \right] + \text{h.c.} \quad (8.6)$$

Thus, in this scenario we have:

$$\begin{aligned} |V_{ud}|_\beta &= \left| \hat{V}_{ud}^L + \hat{V}_{ud}^R \right| \simeq |\hat{V}_{ud}^L| + \frac{\text{Re} \left(\hat{V}_{ud}^L \hat{V}_{ud}^{R*} \right)}{|\hat{V}_{ud}^L|}, \\ |V_{us}|_{K\ell 3} &= \left| \hat{V}_{us}^L + \hat{V}_{us}^R \right| \simeq |\hat{V}_{us}^L| + \frac{\text{Re} \left(\hat{V}_{us}^L \hat{V}_{us}^{R*} \right)}{|\hat{V}_{us}^L|}, \\ \left| \frac{V_{us}}{V_{ud}} \right|_{\frac{K\mu 2}{\pi\mu 2}} &= \left| \frac{\hat{V}_{us}^L - \hat{V}_{us}^R}{\hat{V}_{ud}^L - \hat{V}_{ud}^R} \right| \simeq \frac{|\hat{V}_{us}^L|}{|\hat{V}_{ud}^L|} \left(1 - \frac{\text{Re} \left(\hat{V}_{us}^L \hat{V}_{us}^{R*} \right)}{|\hat{V}_{us}^L|^2} + \frac{\text{Re} \left(\hat{V}_{ud}^L \hat{V}_{ud}^{R*} \right)}{|\hat{V}_{ud}^L|^2} \right). \end{aligned} \quad (8.7)$$

Then, using the determinations in eq. (8.5), one obtains the rephasing invariants:

$$\begin{aligned} |\hat{V}_{ud}^L| &= 0.97450 \quad (8), & \frac{\text{Re} \hat{\mathcal{B}}_{ud}}{|\hat{V}_{ud}^L|} &= \frac{\text{Re} \left(\hat{V}_{ud}^{L*} \hat{V}_{ud}^R \right)}{|\hat{V}_{ud}^L|} = -0.79(27) \times 10^{-3}, \\ |\hat{V}_{us}^L| &= 0.22431 \quad (35), & \frac{\text{Re} \hat{\mathcal{B}}_{us}}{|\hat{V}_{us}^L|} &= \frac{\text{Re} \left(\hat{V}_{us}^{L*} \hat{V}_{us}^R \right)}{|\hat{V}_{us}^L|} = -1.24(37) \times 10^{-3}. \end{aligned} \quad (8.8)$$

Therefore, the Cabibbo angle anomalies indicate towards the presence of right handed couplings, with a consistent part of the mixing in the right-handed sector aligned in phase to the left-handed sector, except for the opposite sign (π phase shift). The diagram of the processes is sketched in Figure 2.

In principle, a single vector-like doublet mixing with the up, the down and the strange SM quarks could explain all inconsistencies. However, as shown in Ref. [49], the required couplings are excluded by limits on flavour-changing neutral currents. Nevertheless, the presence of more than one doublet can be at the origin of the anomalies without




Figure 2: Diagrams contributing to a semileptonic process in the Cabibbo sector, leading to $\Gamma_H^{\text{Cab.}} \sim |\hat{V}_{1i}^L(\pm)_H \hat{V}_{1i}^R|^2 \simeq |\hat{V}_{1i}^L|^2 (\pm)_H 2 \text{Re} \hat{V}_{1i}^L \hat{V}_{1i}^{R*}$. Here, H refers to the specific hadronic process, typically mediated either by a vector current (plus sign) or by an axial-vector one (minus sign).

contradicting experimental constraints. In particular, a vector-like doublet mixing with the up and the down SM quarks ($z_u \neq 0$ and $z_d \neq 0$) can resolve the tension between V_{ud} and V_{us} (CAA1), while a vector-like quark doublet coupling predominantly to the up and strange quarks $z_u \neq 0$ and $z_s \neq 0$ can be the cause of the tension between the $K_{\ell 3}$ and $K_{\mu 2}/\pi_{\mu 2}$ determinations of V_{us} (CAA2) [45, 49].

8.3.2 The neutral kaon system

The success of the SM in high-precision measurements leads to very stringent bounds on NP models. This is particularly true for observables that are highly suppressed in the SM, such as CP-violating effects or FCNC processes. One of the most relevant testing fields for the SM is provided by kaon physics, e.g. rare kaon decays, kaon mixing both in the CP-conserving (mass splitting Δm_K) and CP-violating effects (ϵ), direct CP-violation, but also the decay $K \rightarrow \pi\pi$ into pions with isospin $I = 2$. The SM prediction for the latter does not require loops or FCNCs, but the decay into pions with isospin $I = 2$ is still suppressed, making it an additional interesting probe for BSM physics.

In particular, when facing the Cabibbo angle anomalies, the same new couplings in eq. (8.6) which can be responsible for the anomalies and contribute to leptonic and semileptonic kaon decays, also impact kaon hadronic decays, other leptonic and semileptonic decay channels, and kaon mixing. Strong constraints from kaon mixing and flavour-changing kaon decays prohibit large couplings with both the down and strange quarks of the same vector-like doublet. Then, as already mentioned, the anomalies cannot be simultaneously explained by one doublet [49].

These FCNC processes are determined by trilinears, and we will discuss some of these constraints in section 8.4. We begin by addressing constraints from charged currents and flavour-conserving processes, which are controlled by bilinears. In the context of this work we are interested in a rephasing invariant description of physical processes. In the following we will focus on some processes in which a rephasing invariant description can provide useful and unambiguous information on the constraints on new physics couplings.

Quantities related to observables should be invariant under the rephasing of the states

($|K^0\rangle, |\bar{K}^0\rangle, |K_{S,L}\rangle$, final states $|f\rangle$, etc.), rephasing of the quark fields, and independent on phase choices in the CP transformations. We are also interested in a description which is independent on the choice of parameterization of the mixing matrices. Nevertheless, assuming CPT is a good symmetry, it is convenient, for simplicity, to fix the relative phase of $|K_S\rangle$ and $|K_L\rangle$ in such a way that the coefficients of $|K^0\rangle$ have the same phase and are therefore equal. This choice also implies that the coefficients of $|\bar{K}^0\rangle$ have equal phases (see e.g. Ref. [61]). This is the only phase convention that we assume. Then, the Hamiltonian eigenstates can be written as

$$|K_L\rangle = p_K |K^0\rangle + q_K |\bar{K}^0\rangle, \quad |K_S\rangle = p_K |K^0\rangle - q_K |\bar{K}^0\rangle, \quad (8.9)$$

with $|p_K|^2 + |q_K|^2 = 1$ and $\delta = |p_K|^2 - |q_K|^2$.

$K \rightarrow \pi\pi$. As noted in Ref. [51], significant bounds on right-handed currents can come from the theoretical and experimental result on the decay $K \rightarrow \pi\pi$. The decays to two pions $\pi^+\pi^-$ and $\pi^0\pi^0$ are dominant for K_S . Because of the small CP violation in the kaon system, also K_L can decay to two pions (CP conservation would prevent the decay into two pions for the CP-odd state). Additionally, the kaons predominantly decay to the state of two pions with isospin $I = 0$, while the decay to two pions with isospin $I = 2$ is suppressed.²⁵ Therefore, the decay of kaons to two pions is usually described by three useful parameters [61]

$$\begin{aligned} \omega &= \frac{\langle 2 | H_{\Delta S=1} | K_S \rangle}{\langle 0 | H_{\Delta S=1} | K_S \rangle}, & \epsilon &= \frac{\langle 0 | H_{\Delta S=1} | K_L \rangle}{\langle 0 | H_{\Delta S=1} | K_S \rangle}, \\ \epsilon' &= \frac{\langle 2 | H_{\Delta S=1} | K_L \rangle \langle 0 | H_{\Delta S=1} | K_S \rangle - \langle 0 | H_{\Delta S=1} | K_L \rangle \langle 2 | H_{\Delta S=1} | K_S \rangle}{\sqrt{2} \langle 0 | H_{\Delta S=1} | K_S \rangle^2}, \end{aligned} \quad (8.10)$$

where $\langle 0 |$ and $\langle 2 |$ indicate the states $\langle 2\pi, I = 0 |$ and $\langle 2\pi, I = 2 |$ respectively. With the choice of relative phase between K_L and K_S already mentioned in eq. (8.9), these three parameters are rephasing invariants. ω is the parameter associated to $\Delta I = 1/2$ rule violation, ϵ indicates the mixing CP-violating part, and ϵ' direct CP-violation. Bilinear rephasing invariants $\hat{B}_{\alpha i}$ affect ω and ϵ' while ϵ is affected by trilinears $\hat{T}_{\alpha, ij}$. We will discuss the contribution to ϵ in the next section.

In presence of vector-like weak doublets of quarks mixing with SM quarks at the weak scale we have from eq. (7.18) the effective Hamiltonian

$$H_{\Delta S=1}^{\text{NP}} = \frac{4G_F}{\sqrt{2}} \left[\hat{V}_{us}^{L*} \hat{V}_{ud}^R (\bar{s}_L \gamma^\mu u_L) (\bar{u}_R \gamma_\mu d_R) + \hat{V}_{us}^{R*} \hat{V}_{ud}^L (\bar{s}_R \gamma^\mu u_R) (\bar{u}_L \gamma_\mu d_L) \right] + \text{h.c.} \quad (8.11)$$

²⁵Within the SM, the leading contribution to the $\Delta I = 3/2$ amplitude arises from the low-energy realization of the $(27_L, 1_R)$ -plet part of the product of two quark currents transforming as $(8_L, 1_R) \times (8_L, 1_R)$ under the $SU(3)_L \times SU(3)_R$ symmetry of the approximate massless QCD Lagrangian. The observed suppression of the corresponding non-perturbative coupling, leading to the $\Delta I = 1/2$ rule, is not present for VLQ doublets, whose corresponding $(8_L, 8_R)$ contributions come with an additional chiral enhancement. See for example Ref. [120] for more detailed discussion.

Then, the amplitudes of the decays $K \rightarrow \pi\pi$ receive additional new physics contribution:

$$A_I e^{i\delta_I} = A_I^{\text{SM}} e^{i\delta_I} + A_I^{\text{NP}} e^{i\delta_I}, \quad (8.12)$$

where

$$\langle I | H_{\Delta S=1} | K^0 \rangle = A_I e^{i\delta_I} \sqrt{2}, \quad \langle I | H_{\Delta S=1} | \bar{K}^0 \rangle = \bar{A}_I e^{i\delta_I} \sqrt{2}, \quad (8.13)$$

for $I = 0$ and $I = 2$ and the phases δ_I are the final-state-interaction phase shifts of the two pion states.²⁶ In the SM the main contribution to the amplitudes A_I is at tree level:

$$\sqrt{2} A_I^{\text{SM}} e^{i\delta_I} \simeq \frac{4G_F}{\sqrt{2}} \hat{V}_{us}^{L*} \hat{V}_{ud}^L \langle I | (\bar{s}_L \gamma^\mu u_L) (\bar{u}_L \gamma_\mu d_L) | K^0 \rangle = \frac{G_F}{\sqrt{2}} \hat{V}_{us}^{L*} \hat{V}_{ud}^L \langle I | Q_2 | K^0 \rangle. \quad (8.15)$$

The presence of vector-like doublets generate the additional contribution

$$\sqrt{2} A_I^{\text{NP}} e^{i\delta_I} = \frac{G_F}{\sqrt{2}} \left(\hat{V}_{us}^{L*} \hat{V}_{ud}^R - \hat{V}_{us}^{R*} \hat{V}_{ud}^L \right) (c_1 \langle I | Q_1^{LR} | K^0 \rangle + c_2 \langle I | Q_2^{LR} | K^0 \rangle), \quad (8.16)$$

where $c_1(3 \text{ GeV}) = 0.9$ and $c_2(3 \text{ GeV}) = 0.4$ are the factors resulting from the QCD renormalization [125] and the operators are defined as [124] (we rename $1 \leftrightarrow 2$):

$$\begin{aligned} Q_1^{LR} &= (\bar{s} \gamma_\mu (1 - \gamma_5) u) (\bar{u} \gamma_\mu (1 + \gamma_5) d), \\ Q_2^{LR} &= (\bar{s}_\alpha \gamma_\mu (1 - \gamma_5) u_\beta) (\bar{u}_\beta \gamma_\mu (1 + \gamma_5) d_\alpha). \end{aligned} \quad (8.17)$$

The hadronic matrix elements of left-right operators are chirally enhanced with respect to the tree-level SM ones.²⁷ Moreover, the corresponding amplitudes of the decay in the state with isospin $I = 2$ do not undergo a suppression relative to the $I = 0$ final state. Therefore, the new contributions to the amplitudes may be significant.

In order to obtain both the amplitudes A_0 and A_2 , it is useful to start with the computation of A_2 first. In the decay of K to two pions with isospin $I = 2$ a relation

²⁶The factor $\sqrt{2}$ comes from the convention on the normalization of the amplitudes. Depending on whether the charged pions are considered as identical particles or not, different normalizations are chosen for the isospin decomposition of the two pions states (see e.g. Ref. [61]). If π^+ and π^- are considered as distinguishable, the branching ratios are written as

$$\begin{aligned} \Gamma(K_S \rightarrow \pi^+ \pi^-) &= \frac{1}{16\pi m_K} \sqrt{1 - \frac{4m_\pi^2}{m_K^2}} \left| \sqrt{\frac{1}{3}} \langle 2 | H_{\Delta S=1} | K_S \rangle + \sqrt{\frac{2}{3}} \langle 0 | H_{\Delta S=1} | K_S \rangle \right|^2, \\ \Gamma(K_S \rightarrow \pi^0 \pi^0) &= \frac{1}{32\pi m_K} \sqrt{1 - \frac{4m_\pi^2}{m_K^2}} \left| \frac{2}{\sqrt{3}} \langle 2 | H_{\Delta S=1} | K_S \rangle - \sqrt{\frac{2}{3}} \langle 0 | H_{\Delta S=1} | K_S \rangle \right|^2. \end{aligned} \quad (8.14)$$

This normalization is used in Refs. [121–123] as well as in Refs. [124, 125] and gives $|A_0|_{\text{exp}} = 3.3201(18) \times 10^{-7} \text{ GeV}$ [121], while in Ref. [61] π^+ and π^- are considered as identical particles and the amplitudes A_I are $\sqrt{2}$ times larger giving $|A_0|_{\text{exp}} = 4.6953(25) \times 10^{-7} \text{ GeV}$. However, in Refs. [121, 122, 125] the amplitudes A_I are defined as $\sqrt{2} A_I e^{i\delta_I} = \langle I | H_{\Delta S=1} | K^0 \rangle$ so that the matrix elements are the same as in Ref. [61], while in Refs. [123, 124] $A_I e^{i\delta_I} = \langle I | H_{\Delta S=1} | K^0 \rangle$ so that the matrix elements are $\sqrt{2}$ times smaller than in Refs. [61, 121, 122]. We will use in the following the convention $\sqrt{2} A_I e^{i\delta_I} = \langle I | H_{\Delta S=1} | K^0 \rangle$ with $|A_0|_{\text{exp}} = 3.3201(18) \times 10^{-7} \text{ GeV}$.

²⁷The matrix elements of left-right operators are $\mathcal{O}(p^0)$ in the chiral counting, i.e. they do not vanish in the limit of zero quark/meson masses.

exists between the matrix elements of the operators $Q_{1,2}^{LR}$ and the matrix elements of the electroweak penguin operators $Q_{7,8}$. In the isospin limit [124, 126],

$$\langle 2|Q_1^{RL}|K^0\rangle = \frac{2}{3}\langle 2|Q_7|K^0\rangle, \quad \langle 2|Q_2^{RL}|K^0\rangle = \frac{2}{3}\langle 2|Q_8|K^0\rangle. \quad (8.18)$$

The final-state strong-interaction phase in eq. (8.16) arises from the hadronic matrix element. Then, we can use the lattice results in Ref. [121] to get the hadronic matrix elements at the scale $\mu = 3$ GeV:

$$\frac{\text{Re}\langle 2|Q_1^{RL}|K^0\rangle}{\cos\delta_2} = 0.238(13) \text{ GeV}^3, \quad \frac{\text{Re}\langle 2|Q_2^{RL}|K^0\rangle}{\cos\delta_2} = 1.078(61) \text{ GeV}^3, \quad (8.19)$$

in the suitable phase convention for $|K^0\rangle$. However, let us remark that any rephasing of the initial or final states is irrelevant whenever considering ratios of hadronic matrix elements with same initial and final states.²⁸ In the following analysis we will consider rephasing invariant ratios. Hence, we can write:

$$\begin{aligned} \sqrt{2} A_2^{\text{NP}} &\simeq \frac{G_F}{\sqrt{2}} \left(\hat{V}_{us}^{L*} \hat{V}_{ud}^R - \hat{V}_{us}^{R*} \hat{V}_{ud}^L \right) \times 0.645(27) \text{ GeV}^3 \\ &\simeq \sqrt{2} \left(\hat{V}_{us}^{L*} \hat{V}_{ud}^R - \hat{V}_{us}^{R*} \hat{V}_{ud}^L \right) \times 3.76(16) \times 10^{-6} \text{ GeV}. \end{aligned} \quad (8.22)$$

As regards the amplitude of the decay to the $I = 0$ state, we can use the relation $A_0^{\text{NP}} = -2\sqrt{2}A_2^{\text{NP}}$ in Ref. [125], valid in the chiral limit. Substituting in eq. (8.16), one finds

$$\begin{aligned} \sqrt{2} A_0^{\text{NP}} &\simeq \frac{G_F}{\sqrt{2}} \left(\hat{V}_{us}^{L*} \hat{V}_{ud}^R - \hat{V}_{us}^{R*} \hat{V}_{ud}^L \right) \times 1.825(77) \text{ GeV}^3 \\ &\simeq \sqrt{2} \left(\hat{V}_{us}^{L*} \hat{V}_{ud}^R - \hat{V}_{us}^{R*} \hat{V}_{ud}^L \right) \times 1.064(45) \times 10^{-5} \text{ GeV}. \end{aligned} \quad (8.23)$$

These new contributions should be compared with the SM expectation and experimental values.

²⁸The SM contribution to the amplitudes (also including the loop level contribution) as well as the new physics one present a structure (see eqs. (8.15) and (8.16)):

$$\sqrt{2} \lambda_u^* A_I e^{i\delta_I} = \frac{G_F}{\sqrt{2}} \lambda_u^* \sum_i \lambda^{(i)} \langle 0|Q_i|K^0\rangle, \quad (8.20)$$

where $\lambda^{(i)}$ include couplings and eventual Inami-Lim functions and we also explicitly multiply by $\lambda_u^* = \hat{V}_{us}^L \hat{V}_{ud}^{L*}$ since in the expression of observables the amplitudes always emerge in the combination $\lambda_u^* A_I$ (see e.g. eqs. (8.33) and (8.49)). The final-state strong-interaction phase in eq. (8.16) arises from the hadronic matrix element. Thus, one gets

$$\begin{aligned} \sqrt{2} \text{Re}(\lambda_u^* A_I) &= \frac{G_F}{\sqrt{2}} \sum_i \text{Re}(\lambda_u^* \lambda^{(i)}) \frac{\text{Re}\langle 0|Q_i|K^0\rangle}{\cos\delta_I}, \\ \sqrt{2} \text{Im}(\lambda_u^* A_I) &= \frac{G_F}{\sqrt{2}} \sum_i \text{Im}(\lambda_u^* \lambda^{(i)}) \frac{\text{Re}\langle 0|Q_i|K^0\rangle}{\cos\delta_I}. \end{aligned} \quad (8.21)$$

$K \rightarrow \pi\pi, I = 0$. The experimental value of $|A_0|$ can be obtained from the decay rates in eq. (8.14), footnote 26 [121]:

$$|A_0|_{\text{exp}} = 3.3201(18) \times 10^{-7} \text{ GeV}. \quad (8.24)$$

In the SM the main contribution to the amplitude A_0 is at tree level:

$$\sqrt{2} A_0^{\text{SM}} e^{i\delta_0} \simeq \frac{4G_F}{\sqrt{2}} \hat{V}_{us}^{L*} \hat{V}_{ud}^L \langle 0 | (\bar{s}_L \gamma^\mu u_L) (\bar{u}_L \gamma_\mu d_L) | K^0 \rangle = \frac{G_F}{\sqrt{2}} \lambda_u \langle 0 | Q_2 | K^0 \rangle, \quad (8.25)$$

with $\text{Re} \langle 0 | Q_2 | K^0 \rangle / \cos \delta_0 = 0.147(15) \text{ GeV}^3$ [123], which accounts for $\sim 97\%$ of the SM prediction or A_0 . We defined also $\lambda_u = \hat{V}_{us}^{L*} \hat{V}_{ud}^L$ for convenience. Therefore, the dominant contribution to the amplitude is proportional to $\hat{V}_{us}^{L*} \hat{V}_{ud}^L$ and in very good approximation

$$\lambda_u^* A_0^{\text{SM}} \simeq \text{Re} \left[\lambda_u^* A_0^{\text{SM}} \right]. \quad (8.26)$$

The SM expectation was computed in Ref. [123]:

$$\frac{\text{Re} \left[\lambda_u^* A_0^{\text{SM}} \right]}{|\lambda_u|} = 2.99(67) \times 10^{-7} \text{ GeV}. \quad (8.27)$$

Then, we can compare the SM expectation with the experimental determination:

$$\frac{|A_0|_{\text{exp}} - |A_0^{\text{SM}}|}{|A_0|_{\text{exp}}} = 1 - \frac{|\text{Re} [\lambda_u^* A_0^{\text{SM}}]|}{|\lambda_u| |A_0|_{\text{exp}}} = 0.10 \pm 0.20. \quad (8.28)$$

When including extra quark doublets, we get an additional contribution to the amplitudes:²⁹

$$|A_0^{\text{SM}} + A_0^{\text{NP}}| \simeq |A_0^{\text{SM}}| \left(1 + \frac{\text{Re} (A_0^{\text{SM}*} A_0^{\text{NP}})}{|A_0^{\text{SM}}|^2} \right) \simeq |A_0^{\text{SM}}| \left(1 + \frac{\text{Re} (\lambda_u^* A_0^{\text{NP}})}{\text{Re} (\lambda_u^* A_0^{\text{SM}})} \right). \quad (8.29)$$

Then, it is useful to compare the new rephasing invariant contribution given by the extra quark doublet with the SM contribution:

$$\frac{\text{Re} \left[\lambda_u^* A_0^{\text{NP}} \right]}{\text{Re} \left[\lambda_u^* A_0^{\text{SM}} \right]} \simeq \frac{-2\sqrt{2} \text{Re} \left[\left(|\hat{V}_{us}^L|^2 \hat{V}_{ud}^{L*} \hat{V}_{ud}^R - |\hat{V}_{ud}^L|^2 \hat{V}_{us}^L \hat{V}_{us}^{R*} \right) \right] \text{Re} \left[(0.9 \langle 0 | Q_1^{LR} | K^0 \rangle + 0.4 \langle 0 | Q_2^{LR} | K^0 \rangle) \right]}{|\hat{V}_{us}^L|^2 |\hat{V}_{ud}^L|^2 \text{Re} \left[\langle 0 | Q_2 | K^0 \rangle \right]}. \quad (8.30)$$

²⁹Since the dominant contributions to the SM amplitudes are proportional to λ_u , we have $|\text{Im} (\lambda_u^* A_I^{\text{SM}})| \ll |\text{Re} (\lambda_u^* A_I^{\text{SM}})|$ and we can write:

$$\frac{\text{Re} (A_I^{\text{SM}*} A_I^{\text{NP}})}{|A_I^{\text{SM}}|^2} = \frac{\text{Re} (\lambda_u^* A_I^{\text{NP}}) \text{Re} (\lambda_u A_I^{\text{SM}*}) - \text{Im} (\lambda_u^* A_I^{\text{NP}}) \text{Im} (\lambda_u A_I^{\text{SM}*})}{\text{Re} (\lambda_u^* A_I^{\text{SM}})^2 + \text{Im} (\lambda_u^* A_I^{\text{SM}})^2} \simeq \frac{\text{Re} (\lambda_u^* A_I^{\text{NP}})}{\text{Re} (\lambda_u^* A_I^{\text{SM}})}.$$

Let us underline that this ratio emerges in a rephasing-invariant form. Using the values needed to explain the Cabibbo angle anomalies and the SM estimate in Ref. [123] in their central value we get

$$\frac{\text{Re} \left[\left(\hat{V}_{us}^L \hat{V}_{ud}^{L*} \right) A_0^{\text{NP}} \right]}{\text{Re} \left[\left(\hat{V}_{us}^L \hat{V}_{ud}^{L*} \right) A_0^{\text{SM}} \right]} \simeq -0.037, \quad (8.31)$$

i.e. the extra contribution would be ~ 27 times smaller than the SM one (and, by comparing with eq. (8.28), ~ 30 times smaller than the experimental determination) and also 6 times smaller than the 1σ uncertainty on the lattice SM prediction. Therefore, we can neglect this effect.

$\Delta = \frac{1}{2}$ **isospin rule.** The $\Delta = 1/2$ rule violation and the CP-violating effects are suppressed in the SM context. Then, they may provide stringent constraints on the mixings with the extra vector-like doublets. Regarding the CP conserving contribution of the isospin-suppressed part of the decay $K \rightarrow \pi\pi$, it is conveniently described by the parameter ω [61]:

$$\omega = \frac{\langle 2 | H_{\Delta S=1} | K_S \rangle}{\langle 0 | H_{\Delta S=1} | K_S \rangle} \simeq e^{i(\delta_2 - \delta_0)} \text{Re} \frac{A_2}{A_0} = e^{i(\delta_2 - \delta_0)} \frac{\text{Re}(A_2 A_0^*)}{|A_0|^2}, \quad (8.32)$$

where the approximation follows from taking into account the small value of ϵ . The main contribution to the CP-conserving part of the decay of kaon into two pions with isospin $I = 2$ in the SM is again given by the operator Q_2 , which accounts for the $\sim 93\%$ of the amplitude [121]. Using the fact that the dominant contribution to the amplitudes is proportional to $\hat{V}_{us}^{L*} \hat{V}_{ud}^L$ so that $|\text{Re}(A_I \hat{V}_{us}^L \hat{V}_{ud}^{L*})| \gg |\text{Im}(A_I \hat{V}_{us}^L \hat{V}_{ud}^{L*})|$ we can write [61]

$$|\omega| \simeq \frac{\text{Re}(A_2 A_0^*)}{|A_0|^2} \simeq \frac{\text{Re}(A_2 \lambda_u^*)}{\text{Re}(A_0 \lambda_u^*)}. \quad (8.33)$$

The SM lattice prediction for A_2 gives $\text{Re}(\lambda_u^* A_2)/|\lambda_u| = 1.50(15) \times 10^{-8}$ GeV [122]. The experimental value can be determined from the decay rate of $K^+ \rightarrow \pi^+ \pi^0$ as reported in Ref. [121]:

$$|A_2|_{\text{exp}} = 1.479(4) \times 10^{-8} \text{ GeV}, \quad (8.34)$$

while from K_S decays $|A_2|_{\text{exp}, K_S} = 1.570(53) \times 10^{-8}$. The ω parameter represents the CP-conserving part of the ratio A_2/A_0 , while the parameter ϵ' represents the CP-violating part (direct CP violation). Taking into account that the CP violation is very suppressed, the experimental value of the amplitudes gives [121]:

$$|\omega|_{\text{exp}} = 0.04454(12). \quad (8.35)$$

We can compare the experimental result with the SM expectation:

$$\frac{|A_2|_{\text{exp}} - |A_2^{\text{SM}}|}{|A_2|_{\text{exp}}} = 1 - \frac{|\text{Re}[\lambda_u^* A_2^{\text{SM}}]|}{|\lambda_u| |A_2|_{\text{exp}}} = -0.014 \pm 0.098, \quad (8.36)$$

and

$$\frac{|\omega|_{\text{exp}} - |\omega|_{\text{SM}}}{|\omega|_{\text{exp}}} = 1 - \frac{1}{|\omega|_{\text{exp}}} \frac{\text{Re} \left(A_2 \hat{V}_{us}^L \hat{V}_{ud}^{L*} \right)}{\text{Re} \left(A_0 \hat{V}_{us}^L \hat{V}_{ud}^{L*} \right)} = -0.13 \pm 0.28. \quad (8.37)$$

In a scenario with VLQ doublets, the amplitudes receive additional contributions. Following the same steps as in eq. (8.29) we can write

$$|A_2^{\text{SM}} + A_2^{\text{NP}}| \simeq |A_2^{\text{SM}}| \left(1 + \frac{\text{Re} \left(\lambda_u^* A_2^{\text{NP}} \right)}{\text{Re} \left(\lambda_u^* A_2^{\text{SM}} \right)} \right), \quad (8.38)$$

where we have

$$\text{Re} \left[\lambda_u^* A_2^{\text{NP}} \right] = \text{Re} \left[|\hat{V}_{us}^L|^2 \hat{V}_{ud}^{L*} \hat{V}_{ud}^R - |\hat{V}_{ud}^L|^2 \hat{V}_{us}^L \hat{V}_{us}^{R*} \right] \times 3.76(16) \times 10^{-6} \text{ GeV}. \quad (8.39)$$

As regards the parameter ω , from eq. (8.33) we have:

$$\omega_{\text{SM}} + \omega_{\text{NP}} = e^{i(\delta_2 - \delta_0)} \frac{\text{Re} \left[\lambda_u^* (A_2^{\text{SM}} + A_2^{\text{NP}}) \right]}{\text{Re} \left[\lambda_u^* A_0^{\text{SM}} \right]} = \omega_{\text{SM}} \frac{\text{Re} \left[\lambda_u^* (A_2^{\text{SM}} + A_2^{\text{NP}}) \right]}{\text{Re} \left[\lambda_u^* A_2^{\text{SM}} \right]}. \quad (8.40)$$

Using the values needed to explain the Cabibbo angle anomalies in eq. (8.8) we get:

$$\frac{\text{Re} \left[\lambda_u^* A_2^{\text{NP}} \right]}{\text{Re} \left[\lambda_u^* A_2^{\text{SM}} \right]} = \frac{\omega_{\text{NP}}}{\omega_{\text{SM}}} = 0.26 \pm 0.10. \quad (8.41)$$

Let us emphasize again that this ratio arises as rephasing invariant.

Then, we can evaluate the impact of the new contribution against the experimental determinations. At 2σ we get

$$-0.29 \lesssim \frac{|A_2^{\text{exp}}| - |A_2^{\text{SM}} + A_2^{\text{NP}}|}{|A_2^{\text{exp}}|} \lesssim 0.26 \quad (8.42)$$

and from eqs. (8.38) and (8.39)

$$-1.1 \times 10^{-3} \lesssim |\hat{V}_{us}^L| \frac{\text{Re} \left(\hat{V}_{ud}^{L*} \hat{V}_{ud}^R \right)}{|\hat{V}_{ud}^L|} - |\hat{V}_{ud}^L| \frac{\text{Re} \left(\hat{V}_{us}^L \hat{V}_{us}^{R*} \right)}{|\hat{V}_{us}^L|} \lesssim 1.0 \times 10^{-3} \quad (8.43)$$

By inserting the values needed for Cabibbo angle anomalies

$$|\hat{V}_{us}^L| \frac{\text{Re} \left(\hat{V}_{ud}^{L*} \hat{V}_{ud}^R \right)}{|\hat{V}_{ud}^L|} - |\hat{V}_{ud}^L| \frac{\text{Re} \left(\hat{V}_{us}^L \hat{V}_{us}^{R*} \right)}{|\hat{V}_{us}^L|} = 1.05(37) \times 10^{-3} \quad (8.44)$$

we receive

$$\frac{|A_2^{\text{exp}}| - |A_2^{\text{SM}} + A_2^{\text{NP}}|}{|A_2^{\text{exp}}|} \simeq 1 - \frac{|A_2^{\text{SM}}|}{|A_2^{\text{exp}}|} \frac{\text{Re} \left[\lambda_u^* (A_2^{\text{SM}} + A_2^{\text{NP}}) \right]}{\text{Re} \left[\lambda_u^* A_2^{\text{SM}} \right]} = -0.28 \pm 0.14, \quad (8.45)$$

$$1 - \frac{|\omega_{\text{SM}} + \omega_{\text{NP}}|}{|\omega|_{\text{exp}}} = -0.42 \pm 0.35,$$

i.e., by comparison with eqs. (8.34) and (8.35), the CAA values agree with the experimental results within 2σ .

Direct CP violation. $\epsilon' \neq 0$ represents direct CP violation, i.e. CP violation in the decay amplitudes:

$$\epsilon' = \frac{\langle 2|H_{\Delta S=1}|K_L \rangle \langle 0|H_{\Delta S=1}|K_S \rangle - \langle 0|H_{\Delta S=1}|K_L \rangle \langle 2|H_{\Delta S=1}|K_S \rangle}{\sqrt{2} \langle 0|H_{\Delta S=1}|K_S \rangle^2} \quad (8.46)$$

$$\simeq \frac{i}{\sqrt{2}} e^{i(\delta_2 - \delta_0)} \text{Im} \frac{A_2}{A_0},$$

where the approximation makes use of the small value of ϵ [61]. The ratio ϵ'/ϵ can be experimentally determined from the relation

$$\left| \frac{\eta_{00}}{\eta_{+-}} \right| \simeq 1 - 6 \text{Re} \left(\frac{\epsilon'}{\epsilon} \right), \quad (8.47)$$

where the parameters η are defined as $|\eta_f| = \sqrt{(\Gamma_L/\Gamma_S) \text{Br}(K_L \rightarrow f)/\text{Br}(K_S \rightarrow f)}$. Since ϵ and ϵ' exhibit the same phase, we also have

$$\left(\frac{\epsilon'}{\epsilon} \right)_{\text{SM}} \simeq \text{Re} \left(\frac{\epsilon'}{\epsilon} \right)_{\text{SM}}. \quad (8.48)$$

Then, using the fact that the dominant contribution to the amplitudes is proportional to $\hat{V}_{us}^{L*} \hat{V}_{ud}^L$ and one expects $|\text{Re}(A_I \lambda_u^*)| \gg |\text{Im}(A_I \lambda_u^*)|$ we can write [61]

$$\left(\frac{\epsilon'}{\epsilon} \right)_{\text{SM}} \simeq \frac{\text{Im}(A_2 A_0^*)}{\sqrt{2} |\epsilon| |A_0|^2} \simeq \frac{\text{Im}(A_2 \lambda_u^*) \text{Re}(A_0 \lambda_u^*) - \text{Re}(A_2 \lambda_u^*) \text{Im}(A_0 \lambda_u^*)}{\sqrt{2} |\epsilon| [\text{Re}(A_0 \lambda_u^*)]^2} \quad (8.49)$$

$$\simeq \frac{\text{Im}(A_2 \lambda_u^*) - |\omega| \text{Im}(A_0 \lambda_u^*)}{\sqrt{2} |\epsilon A_0 \lambda_u^*|}.$$

Experimentally it is found that [127]

$$\text{Re} \left(\frac{\epsilon'}{\epsilon} \right)_{\text{exp}} = 1.66(23) \times 10^{-3}. \quad (8.50)$$

The lattice SM prediction gives [123]

$$\left(\frac{\epsilon'}{\epsilon} \right)_{\text{SM}} \simeq \text{Re} \left(\frac{\epsilon'}{\epsilon} \right)_{\text{SM}} = 2.17(84) \times 10^{-3}, \quad (8.51)$$

in agreement with the estimate from chiral perturbation theory [128], especially if one considers isospin breaking corrections [129].

From eq. (8.49) we get

$$\frac{\epsilon'}{\epsilon} = \left(\frac{\epsilon'}{\epsilon}\right)_{\text{SM}} + \frac{\text{Im}\left(A_2^{\text{NP}} \hat{V}_{us}^L \hat{V}_{ud}^{L*}\right) (1 + 2\sqrt{2}|\omega|_{\text{exp}})}{\sqrt{2} |\epsilon A_0^{\text{exp}} \hat{V}_{us}^L \hat{V}_{ud}^{L*}|}. \quad (8.52)$$

The comparison of the new contribution with the SM one gives:

$$\begin{aligned} \frac{(\epsilon'/\epsilon)_{\text{NP}}}{(\epsilon'/\epsilon)_{\text{SM}}} &\simeq \frac{(1 + 2\sqrt{2}|\omega|_{\text{exp}}) \text{Im}\left(A_2^{\text{NP}} \hat{V}_{us}^L \hat{V}_{ud}^{L*}\right)}{\text{Im}\left(A_2^{\text{SM}} \hat{V}_{us}^L \hat{V}_{ud}^{L*}\right) - |\omega|_{\text{exp}} \text{Im}\left(A_0^{\text{SM}} \hat{V}_{us}^L \hat{V}_{ud}^{L*}\right)} \simeq \\ &\simeq \frac{\text{Im}\left[\hat{V}_{us}^L \hat{V}_{ud}^{L*} \left(\hat{V}_{us}^{L*} \hat{V}_{ud}^R - \hat{V}_{us}^{R*} \hat{V}_{ud}^L\right)\right] \times 1.9(6) \times 10^6}{|\hat{V}_{us}^L \hat{V}_{ud}^{L*}|}. \end{aligned} \quad (8.53)$$

After inserting the values needed for the Cabibbo angle anomalies (see eq. (8.8)), this ratio indicates that in order to get a contribution less than the SM the CP violating part of the bilinears $\hat{\mathcal{B}}_{us}$, $\hat{\mathcal{B}}_{ud}$ should be a factor $\sim 10^3$ smaller than the CP conserving part.

By confronting the contribution of the model including vector-like quark doublets with the experimental result we can get a 95% confidence level limit on the CP violating part of the new physics contribution:

$$-5.7 \times 10^{-7} \lesssim \frac{|\hat{V}_{us}^L|}{|\hat{V}_{ud}^L|} \frac{\text{Im}\left[\hat{V}_{ud}^{L*} \hat{V}_{ud}^R\right]}{|\hat{V}_{ud}^L|} + \frac{\text{Im}\left[\hat{V}_{us}^{L*} \hat{V}_{us}^R\right]}{|\hat{V}_{us}^L|} \lesssim 3.1 \times 10^{-7}. \quad (8.54)$$

This constraint implies that the imaginary part of that combination of bilinears should be at least 3 orders of magnitude smaller than the CP conserving part requested by the Cabibbo angle anomalies:

$$\frac{\text{Im}\left[|\hat{V}_{us}^L|^2 \hat{V}_{ud}^{L*} \hat{V}_{ud}^R - |\hat{V}_{ud}^L|^2 \hat{V}_{us}^L \hat{V}_{us}^{R*}\right]}{\text{Re}\left[|\hat{V}_{us}^L|^2 \hat{V}_{ud}^{L*} \hat{V}_{ud}^R - |\hat{V}_{ud}^L|^2 \hat{V}_{us}^L \hat{V}_{us}^{R*}\right]} = \frac{\text{Im}\left[|\hat{V}_{us}^L|^2 \hat{\mathcal{B}}_{ud} + |\hat{V}_{ud}^L|^2 \hat{\mathcal{B}}_{us}\right]}{\text{Re}\left[|\hat{V}_{us}^L|^2 \hat{\mathcal{B}}_{ud} - |\hat{V}_{ud}^L|^2 \hat{\mathcal{B}}_{us}\right]} \lesssim 10^{-3}. \quad (8.55)$$

It is evident from this description the requirement on new physics which follows the fact that the observed direct CP violation is small, that is, the phases of the bilinears $\hat{\mathcal{B}}_{us}^*/|\hat{V}_{us}^L|$ and $\sim 0.23 \hat{\mathcal{B}}_{ud}/|\hat{V}_{ud}^L|$ should tend to align:

$$\arg\left(\frac{0.23 \hat{V}_{ud}^{L*} \hat{V}_{ud}^R}{|\hat{V}_{ud}^L|} - \frac{\hat{V}_{us}^L \hat{V}_{us}^{R*}}{|\hat{V}_{us}^L|}\right) \sim 10^{-3}. \quad (8.56)$$

The diagram of the process is sketched in Figure 4.

Quantity	Experimental value
ω_{exp}	0.04454(12) [121]
$ A_0 _{\text{exp}}$	$3.3201(18) \times 10^{-7}$ GeV [121]
$ A_2 _{\text{exp}}$	$1.479(4) \times 10^{-8}$ GeV [121]
$ A_2 _{\text{exp}, K_S}$	$1.570(53) \times 10^{-8}$ GeV [121]
ϵ_{exp}	$2.228(11) \times 10^{-3}$ [127]
$\text{Re}(\epsilon'/\epsilon)_{\text{exp}}$	$1.66(23) \times 10^{-3}$ [127]

Table 4: Values of the quantities used in the text, determined from experimental data.

8.3.3 Electric dipole moments

A combination of the imaginary parts of the bilinears $\hat{\mathcal{B}}_{ud}$, $\hat{\mathcal{B}}_{us}$ is constrained by the electric dipole moments. The experimental constraints for the neutron and proton electric dipole moments d_n and d_p at 90% CL respectively are

$$\begin{aligned} d_n &< 1.8 \times 10^{-13} \text{ e fm} \quad [130], \\ d_p &< 2.1 \times 10^{-12} \text{ e fm} \quad [131]. \end{aligned} \tag{8.57}$$

The contribution of right-handed couplings to electric dipole moments was computed in Ref. [132] (neglecting subleading contributions from couplings other than to light generations (u, d, s)):

$$d_n \simeq \text{Im} \left[(1.4 \pm 0.7) \hat{V}_{ud}^{L*} \hat{V}_{ud}^R + (2.7 \pm 1.3) \hat{V}_{us}^{L*} \hat{V}_{us}^R \right] \times 10^{-7} \text{ e fm}, \tag{8.58}$$

$$d_p \simeq \text{Im} \left[-(2.7 \pm 1.3) \hat{V}_{ud}^{L*} \hat{V}_{ud}^R - (3.6 \pm 1.5) \hat{V}_{us}^{L*} \hat{V}_{us}^R \right] \times 10^{-7} \text{ e fm}. \tag{8.59}$$

We can use the bounds on the neutron and proton electric dipole moment to obtain a limit on the combination of bilinears:

$$\begin{aligned} \left| (2.3 \pm 1.1) \frac{\text{Im} \left[\hat{V}_{ud}^{L*} \hat{V}_{ud}^R \right]}{|\hat{V}_{ud}^L|} + (1.0 \pm 0.5) \frac{\text{Im} \left[\hat{V}_{us}^{L*} \hat{V}_{us}^R \right]}{|\hat{V}_{us}^L|} \right| &\lesssim 3 \times 10^{-6}, \\ \left| (3.3 \pm 1.6) \frac{\text{Im} \left[\hat{V}_{ud}^{L*} \hat{V}_{ud}^R \right]}{|\hat{V}_{ud}^L|} + (1.0 \pm 0.4) \frac{\text{Im} \left[\hat{V}_{us}^{L*} \hat{V}_{us}^R \right]}{|\hat{V}_{us}^L|} \right| &\lesssim 6 \times 10^{-6}, \end{aligned} \tag{8.60}$$

which can be compared to the constraint in eq. (8.54). This condition requires for the imaginary part of the combinations of bilinears to be at least about 2 orders of magnitude smaller than the CP conserving part indicated by the Cabibbo angle anomalies.

Quantity	SM value
$\text{Re } A_0$	$2.99(67) \times 10^{-7} \text{ GeV}$ [123]
$\text{Im } A_0$	$-6.98(1.57) \times 10^{-11} \text{ GeV}$ [123]
$\text{Re } A_2$	$1.50(15) \times 10^{-8} \text{ GeV}$ [122]
$\text{Im } A_2$	$-8.34(1.03) \times 10^{-13} \text{ GeV}$ [123]
$\text{Re}(\epsilon'/\epsilon)_{\text{SM}}$	$2.17(84) \times 10^{-3}$ [123]

Table 5: Prediction of the SM values. The values of the amplitudes are given in a particular phase convention, in which the tree-level contribution is real. The amplitudes are given in the renormalization indicated in the text.

8.3.4 Low-energy EW flavour-conserving observables and Z decay

Cabibbo angle anomalies imply a large mixing of the vector-like species with the light quarks. In this context, Z -boson physics provide constraints on the magnitude of the flavour conserving couplings. In fact, mixing with the heavy doublet changes the prediction of the Z decay rate into hadrons (and consequently the total decay rate) as

$$\begin{aligned} \Gamma(Z \rightarrow \text{had}) - \Gamma(Z \rightarrow \text{had})_{\text{SM}} &= \Gamma(Z) - \Gamma(Z)_{\text{SM}} \\ &\simeq \frac{G_F M_Z^3}{\sqrt{2}\pi} \left[-\frac{2}{3} \sin^2 \theta_W (\hat{F}_{uu} + \hat{F}_{cc}) - \frac{1}{3} \sin^2 \theta_W (\hat{F}_{dd} + \hat{F}_{ss} + \hat{F}_{bb}) \right], \end{aligned} \quad (8.61)$$

where

$$\hat{F}_{\alpha\alpha} = \frac{v^2}{M_Q^2} |z_\alpha|^2 \simeq F_{\alpha\alpha}^u, \quad \hat{F}_{ii} = \frac{v^2}{M_Q^2} |z_i|^2 \simeq F_{ii}^d \quad (8.62)$$

are rephasing invariant moduli. Equation (8.61) implies that the predicted decay rate is lower than the SM expectation.

First, we consider the total decay rate of the Z boson as well as the partial decay rate into hadrons. The experimental measurements yield [127]

$$\begin{aligned} \Gamma(Z)_{\text{exp}} &= 2.4955 \pm 0.0023 \text{ GeV}, \\ \Gamma(Z \rightarrow \text{hadr})_{\text{exp}} &= 1.7432 \pm 0.0019 \text{ GeV}, \end{aligned} \quad (8.63)$$

while the corresponding SM predictions are $\Gamma(Z)_{\text{SM}} = 2.4940 \pm 0.0009 \text{ GeV}$ and $\Gamma(Z \rightarrow \text{hadr})_{\text{SM}} = 1.74088 \pm 0.00086 \text{ GeV}$ [127]. At 2σ CL we obtain the limit

$$\hat{F}_{uu} + \hat{F}_{cc} + \frac{1}{2} (\hat{F}_{dd} + \hat{F}_{ss} + \hat{F}_{bb}) \lesssim 5.7 \times 10^{-3}. \quad (8.64)$$

Another set of flavour-conserving constraints originates from parity violating effects at low-energy electron-hadron processes with Z -boson exchange, as well as other low energy electroweak observables as the oblique parameters (see Refs. [49, 50] for the effects of vector-like doublets; in the subsequent analysis we use $g_{AV,\text{exp}}^{ep} = -0.0356 \pm 0.0023$ for

the weak charge of the proton and we update the constraint $55g_{AV}^{ep} + 78g_{AV}^{en} = 36.25 \pm 0.21$ for atomic parity violation is in Cesium $Q_W^{55,78}(Cs)_{\text{exp}}$ [127]).

The importance of the vector-like quarks doublet in explaining the Cabibbo angle anomalies emerges particularly when considering these observables [45, 49–51]. In fact, in general, explanations for the CKM unitarity can be in tension with electroweak low energy observables, e.g. the value of m_W . For example, it was suggested in Ref. [42] that a solution to CAA1 which modifies the Fermi constant G_F with respect to the muon decay constant G_μ also implies a deficit of m_W . Conversely, in Ref. [133], it is shown that the models that predict a positive shift in the W mass may also predict a huge violation of CKM unitarity, much larger than the one indicated by the current anomaly. The vector-like quark charged as a doublet of $SU(2)_L$ emerges as a favoured candidate for explaining the Cabibbo angle anomalies since the mixing with the light generations of SM quarks can resolve the CAA at tree-level, while the contribution to m_W is at one-loop level and it can be relevant only in presence of a large coupling to the top quark.

8.4 Trilinears and flavour changing neutral currents

In the presence of VLQs, flavour changing neutral currents are induced at tree level as well as at loop level. These flavour-changing effects interfere with the SM, both in CP-conserving and CP-violating processes. In this section, we illustrate the effects on relevant kaon decays and on kaon mixing, in particular concerning mechanisms of CP violation. In fact, we have treated CP violation in decay amplitudes (direct CP violation) in section 8.3.2. In what follows, we will illustrate the rephasing-invariant effects of the vector-like doublet in CP violation in neutral kaon mixing (indirect CP violation) and CP violation due to the phase mismatch between mixing parameters and decay amplitudes (‘interference CP violation’).

8.4.1 Kaon mixing

In the SM the short-distance contribution to the transition $K^0(d\bar{s}) \leftrightarrow \bar{K}^0(\bar{d}s)$ arises from weak box diagrams. The effective Lagrangian describing this contribution is given by:

$$\mathcal{L}_{\Delta S=2}^{\text{SM}} = -\frac{G_F^2 m_W^2}{4\pi^2} (\lambda_c^2 S_0(x_c) + \lambda_t^2 S_0(x_t) + 2\lambda_c \lambda_t S_0(x_c, x_t)) (\bar{s}_L \gamma^\mu d_L)^2 + \text{h.c.}, \quad (8.65)$$

where $\lambda_a = V_{as}^* V_{ad}$, $x_a = m_a^2/m_W^2$ and $S_0(x_i)$ are the Inami-Lim functions [134]. The weak short-distance contribution to the mass splitting $\Delta m_K = m_{K_L} - m_{K_S}$ and the CP-violating effects are described by the off-diagonal term M_{12} of the mass matrix of neutral kaons $M_{12} = -\langle K^0 | \mathcal{L}_{\Delta S=2} | \bar{K}^0 \rangle / (2m_{K^0})$, which in the SM is [61]

$$M_{12}^{\text{SM}} = -\frac{G_F^2 m_W^2 f_K^2 m_K B_K}{12\pi^2} (\lambda_c^{*2} S(x_c) + \lambda_t^{*2} S(x_t) + 2\lambda_c^* \lambda_t^* S(x_c, x_t)), \quad (8.66)$$

where f_K is the kaon decay constant, which can be estimated in lattice QCD to be $f_K = 155.7(0.7)$ MeV [95], $m_{K^0} = 497.611 \pm 0.013$ MeV is the neutral kaon




Figure 3: Upper limits obtained from kaon decays and neutral kaon mixing on the flavour changing coupling $|\hat{\mathcal{T}}_{u,ds}|/|\hat{V}_{ud}^L \hat{V}_{us}^{L*}| = |\hat{F}_{ds}| = |z_s^* z_d| v^2 / M_Q^2 = |\hat{V}_{ud}^R \hat{V}_{us}^{R*}| / \hat{F}_{uu}$ as a function of the phase of the rephasing invariant $\hat{\mathcal{T}}_{u,ds} = \hat{V}_{ud}^L \hat{V}_{us}^{L*} \hat{F}_{ds}$ (see text and Refs. [45, 49] for details). We also indicated the bound from ϵ'/ϵ (dashed brown line) considering only the contribution of $\hat{\mathcal{T}}_{u,ds}$ which however would require z_u vanishingly small, which is not the case addressing the Cabibbo angle anomalies (when $z_u \neq 0$ the CP violation in ϵ' can receive contribution by the bilinears and this bound disappears). The black lines indicate the 1σ values of the mixings $|\hat{V}_{ud}^R \hat{V}_{us}^{R*}|$ needed to explain the Cabibbo angle anomalies, assuming $z_u = 0.8$ with $M_Q = 2$ TeV, or equivalently $|z_u|v/M_Q = 0.07$ (the maximum allowed value by the Z decay constraint).

mass, the factor \hat{B}_K is the correction to the vacuum insertion approximation, which is calculated lattice QCD $B_K = 0.7533(91)$ [96] and $S(x_i)$ are the Inami-Lim functions [134] (corrected by short-distance QCD effects [135]). The modulus of the mixing mass M_{12} describe short-distance contributions in the mass splitting. The mass splitting is given by $\Delta m_K \simeq 2|M_{12}| + \Delta m_{K,LD}$ [135], where $\Delta m_{K,LD}$ is the long-distance contribution which is difficult to evaluate [136, 137], however the short distance contribution gives the dominant contribution to the experimental determination $\Delta m_{K,exp} = (3.484 \pm 0.006) \times 10^{-15}$ GeV [127].

From eq. (8.10) the CP violating parameter ϵ can be written as [61]:

$$\epsilon \simeq \frac{e^{i\pi/4}}{\sqrt{2}} \frac{2 \operatorname{Im}(M_{12}^* \Gamma_{12})}{4|M_{12}|^2 + |\Gamma_{12}|^2} \simeq -e^{i\pi/4} \frac{\operatorname{Im}(M_{12} A_0 \bar{A}_0^*)}{\sqrt{2} (\Delta m_K) |A_0 \bar{A}_0|} \quad (8.67)$$

(recall that $\Delta m_K \simeq 2|M_{12}| \simeq -\frac{1}{2}\Delta\Gamma \simeq |\Gamma_{12}| \simeq \frac{1}{2}\Gamma_S$). The long-distance contribution to M_{12} should have a phase $\arg \lambda_u^{*2} + \xi_s - \xi_d - \xi_K$ which does not modify ϵ [61], so that the relevant contributions are the short-distance ones (box diagrams). The crucial approximation in the last formula is considering that the decay channel $|\pi\pi, I=0\rangle$ is the dominant channel $\Gamma_{12} = \sum_f A_f^* \bar{A}_f \simeq A_0^* \bar{A}_0$ (where the sum is over all decay modes f). In this approximation the parameter ϵ represents the mixing CP violation in the neutral kaons system and the phase $\arg \epsilon$ is given by the superweak phase $\phi_\epsilon = 43.52 \pm 0.04^\circ$.

In the SM, at tree level

$$\frac{A_0}{\bar{A}_0} = \frac{V_{us}^* V_{ud}}{V_{us} V_{ud}^*} e^{i(\xi_K + \xi_d - \xi_s)}. \quad (8.68)$$

The phases ξ_K, ξ_d, ξ_s are arbitrary phases in the definition of the CP transformations of $|K^0\rangle$ and the quark fields d and s respectively which are cancelled by the same phases arising from the hadronic matrix element $\langle K^0 | (\bar{d}_L \gamma^\mu s_L) (\bar{d}_L \gamma_\mu s_L) | \bar{K}^0 \rangle \propto -e^{-i(\xi_K + \xi_d - \xi_s)} f_K^2 m_K / 3$ in M_{12} , and thus for brevity we will drop them in the formula for the parameter ϵ which is independent of it. The relation in eq. (8.68) holds when A_0 and \bar{A}_0 are dominated by the tree level diagram. Loop contributions are also proportional to $V_{ts}^* V_{td}$. However, because of the small experimental value of ϵ'/ϵ , loop contributions to the phases of A_0 and \bar{A}_0 must be very small. Then, from eqs. (8.67) and (8.68) we have

$$\epsilon \simeq -e^{i\pi/4} \frac{\text{Im} [M_{12} \lambda_u^2]}{\sqrt{2} \Delta m_K |\lambda_u|^2}. \quad (8.69)$$

From experimental data one obtains [67]

$$|\epsilon|_{\text{exp}} = (2.228 \pm 0.011) \times 10^{-3}. \quad (8.70)$$

In the standard parameterization of the CKM matrix, λ_u is real and it is usually omitted. However, we are interested in a rephasing-invariant analysis. Then, in the SM we have:

$$\begin{aligned} \epsilon \simeq & -\frac{e^{i\pi/4}}{\sqrt{2} \Delta m_K} \frac{\text{Im} [M_{12}^{\text{SM}} \lambda_u^2]}{|\lambda_u|^2} = \frac{e^{i\pi/4} f_K^2 m_K B_K G_F^2 m_W^2}{\sqrt{2} \Delta m_K 12\pi^2} \\ & \times \frac{1}{|\lambda_u|^2} \text{Im} \left[(Q_{csud})^2 S(x_c) + (Q_{tsud})^2 S_0(x_t) + 2 Q_{tsud} Q_{csud} S(x_c, x_t) \right]. \end{aligned} \quad (8.71)$$

It is clear that the CP-violating effects in the SM are given by the imaginary parts of quartets appearing in the weak-basis invariant of mass dimension 12 in eq. (6.1). This is directly related to the fact that CP-violation emerges at loop level, in particular, in the case of kaon mixing, in box diagrams with four quarks involved. The last line can be explicitly written in terms of the J invariant of eq. (3.24):

$$\frac{\text{Im} [M_{12}^{\text{SM}} \lambda_u^2]}{|\lambda_u|^2} \propto \frac{2J}{|\lambda_u|^2} \left(\text{Re} (Q_{csud}) S(x_c) - \text{Re} (Q_{tsud}) S_0(x_t) + \text{Re} (Q_{tsud} - Q_{csud}) S(x_c, x_t) \right). \quad (8.72)$$

In the presence of VLQs, a new contribution to the mixing mass term is induced at tree level as well as at loop level. The loop level effects can be sizeable since they increase with higher masses of the vector-like species. Moreover, in presence of the vector-like doublets, also kinetic enhanced left-right contributions must be taken into

account. Then, $M_{12} = M_{12}^{\text{SM}} + M_{12}^{\text{NP}}$ with

$$M_{12}^{\text{NP}} \simeq -\frac{1}{3}m_{K^0}f_K^2 0.43 \left\{ \frac{G_F}{\sqrt{2}} F_{ds}^2 + \frac{G_F^2}{4\pi^2} \left[\frac{1}{2}M_Q^2 F_{ds}^2 - 3.1 \frac{m_{K^0}^2}{(m_d + m_s)^2} F_{ds} V_{td}^{L*} V_{ts}^L m_W^2 f(x_Q, x_t) \right] \right\}, \quad (8.73)$$

where $f(x_Q, x_t) \simeq x_t \ln(x_Q) / 4$ and we used the results calculated in Refs. [138–140]. We showed in section 8.3.2 that, in the context of Cabibbo angle anomalies, the new physics contribution to the amplitudes A_0 , \bar{A}_0 is negligible (as expected). Then, we can use eqs. (8.67) and (8.68) and write the new contribution to ϵ :

$$\frac{\text{Im} \left[M_{12}^{\text{NP}} (\lambda_u)^2 \right]}{|\lambda_u|^2} = \frac{1}{3}m_{K^0}f_K^2 0.43 \text{Im} \left\{ \frac{G_F}{\sqrt{2}} \frac{(\hat{\mathcal{T}}_{ds,u})^2}{|\lambda_u|^2} + \frac{G_F^2}{4\pi^2} \left[\frac{1}{2}M_Q^2 \frac{(\hat{\mathcal{T}}_{ds,u})^2}{|\lambda_u|^2} - 3.1 \frac{m_{K^0}^2}{(m_d + m_s)^2} \frac{\hat{\mathcal{T}}_{ds,u} Q_{tsud}}{|\lambda_u|^2} m_W^2 f(x_Q, x_t) \right] \right\}, \quad (8.74)$$

with $\lambda_u = \hat{V}_{us}^{L*} \hat{V}_{ud}^L$. Let us remark how the new contribution enters in ϵ only with rephasing invariants. In particular, this flavour-changing and CP-violating is determined by the trilinear $\hat{\mathcal{T}}_{ds,u}$. Constraints can be estimated as $|M_{12}^{\text{NP}}| < |M_{12}^{\text{SM}}| \Delta_K$, $|\text{Im} M_{12}^{\text{NP}}| < |\text{Im} M_{12}^{\text{SM}}| \Delta_{\epsilon_K}$. Then, the bound on the new physics will be a function of the modulus $|\hat{\mathcal{T}}_{ds,u}|$ and phase $\arg(\hat{\mathcal{T}}_{ds,u})$ of the rephasing invariant. Setting $\Delta_K = 1$ and using the results in Ref. [141] at 95% CL (which approximately corresponds to $\Delta_{\epsilon_K} = 0.3$) we obtain, for $M_Q \simeq 2$ TeV,

$$|\hat{F}_{ds}| < 6 \times 10^{-7} - 2 \times 10^{-4}, \quad (8.75)$$

depending on the relative phase of the couplings. We show the result in Figure 3. The limit becomes stronger with increasing mass M_Q because of the loop level contribution (second term in eq. (8.74)).

Notice that after considering flavour changing processes together with the constraint from Z decay in the one doublet case we would have

$$|\hat{V}_{ud}^R| = \frac{\hat{F}_{uu} |F_{sd}|}{|\hat{V}_{us}^R|} \lesssim \frac{5 \times 10^{-3} \times 5 \times 10^{-6}}{10^{-3}} \simeq 2.5 \times 10^{-5} \ll 10^{-3} \quad (8.76)$$

after assuming the value of \hat{V}_{us}^R needed for the Cabibbo angle anomaly, that is, the other tension would not be explained with one doublet.

8.4.2 $K_L \rightarrow \pi^0 \nu \nu$, $K_{S,L} \rightarrow \mu \mu$ and other rare kaon decays

The rephasing invariant $\hat{\mathcal{T}}_{ds,u}$ also generates contributions to flavour changing rare kaon decays, as $K \rightarrow \pi \nu \nu$, $K \rightarrow \mu \mu$, both CP-conserving and CP-violating. As regards CP violation, while ϵ' represents CP violation in the decay amplitudes (direct, $|\bar{A}/A| \neq 1$)

and ϵ indicates CP violation in the mixing (indirect, $\arg(M_{12}^* \Gamma_{12}) \neq 0$), CP violation in these decays belongs to the phase mismatch between the mixing parameters and the decay amplitudes.

The decay $K_L \rightarrow \pi^0 \nu \nu$ is theoretically very clean. It can be shown that if lepton flavour is conserved (ν and $\bar{\nu}$ are each others antiparticle and they are generated by the $\bar{\nu}_{\ell L} \gamma^\mu \nu_{\ell L}$ operator) the final state is CP even [142–145]. Then the decay $K_L \rightarrow \pi^0 \nu \nu$ is CP-violating. It is short-distance dominated and the effective Lagrangian in the SM reads [135]

$$\mathcal{L}(K \rightarrow \pi \nu \nu)_{\text{SM}} = -\frac{g^4}{16\pi^2 m_W^2} \left[V_{cs}^{L*} V_{cd}^L X(x_c) + V_{ts}^{L*} V_{td}^L X(x_t) \right] (\bar{s}_L \gamma^\mu d_L) \sum_{e,\mu,\tau} (\bar{\nu}_{\ell L} \gamma^\mu \nu_{\ell L}), \quad (8.77)$$

where $x_{c,t} = m_{c,t}^2/M_W^2$ and $X(x_a)$ are the Inami-Lim functions [134]. The mixing with the vector-like quark doublets induces the new contribution

$$\mathcal{L}(K \rightarrow \pi \nu \nu)_{\text{NP}} = \frac{4G_F}{\sqrt{2}} \frac{1}{2} F_{ds} (\bar{s}_R \gamma^\mu d_R) \sum_{e,\mu,\tau} (\bar{\nu}_{\ell L} \gamma_\mu \nu_{\ell L}). \quad (8.78)$$

The decay amplitude for the decay $K_L \rightarrow \pi^0 \nu \nu$ is given by

$$A(K_L \rightarrow \pi^0 \nu \nu) = p_K A(K^0 \rightarrow \pi^0 \nu \nu) + q_K A(\bar{K}^0 \rightarrow \pi^0 \nu \nu). \quad (8.79)$$

The relevant CP-violating parameter in this decay is [61]

$$\begin{aligned} \lambda_{\pi \nu \bar{\nu}} &= \frac{q_K A(\bar{K}^0 \rightarrow \pi^0 \nu \nu)}{p_K A(K^0 \rightarrow \pi^0 \nu \nu)} \\ &\simeq -\frac{\hat{V}_{us}^{L*} \hat{V}_{ud}^L}{\hat{V}_{us}^L \hat{V}_{ud}^{L*}} \frac{\frac{\alpha}{2\pi \sin^2 \theta_W} \left(\hat{V}_{cs}^L \hat{V}_{cd}^{L*} X(x_c) + \hat{V}_{ts}^L \hat{V}_{td}^{L*} X(x_t) \right) - \frac{1}{2} \hat{F}_{ds}^*}{\frac{\alpha}{2\pi \sin^2 \theta_W} \left(\hat{V}_{cs}^{L*} \hat{V}_{cd}^L X(x_c) + \hat{V}_{ts}^{L*} \hat{V}_{td}^L X(x_t) \right) - \frac{1}{2} \hat{F}_{ds}}, \end{aligned} \quad (8.80)$$

and the decay amplitude can be written as

$$A(K_L \rightarrow \pi^0 \nu \nu) = p_K (1 + \lambda_{\pi \nu \bar{\nu}}) A(\bar{K}^0 \rightarrow \pi^0 \nu \nu), \quad (8.81)$$

where $\lambda_{\pi \nu \bar{\nu}} \neq -1$ implies CP violation. Since strong final state interaction phases are absent and diagrams with intermediate up quarks are suppressed by the GIM mechanism, there is no direct CP violation ($|\bar{A}/A| = 1$). Indirect CP violation is very small ($|q/p|$ can be considered as 1). The main reason why $|\lambda_{\pi \nu \bar{\nu}}| \neq 1$ in the SM as well as in the new physics scenario is because the phases of q/p and \bar{A}/A do not match, that is, interference between mixing and decay [142].

Then, the total branching ratio results in

$$\frac{\text{Br}(K_L \rightarrow \pi^0 \nu \bar{\nu})}{\text{Br}(K_L \rightarrow \pi^0 \nu \bar{\nu})_{\text{SM}}} \simeq \left| 1 - \frac{\frac{1}{2} \text{Im} \left[\hat{V}_{us}^L \hat{V}_{ud}^{L*} \hat{F}_{ds} \right]}{\text{Im} \left[\hat{V}_{us}^L \hat{V}_{ud}^{L*} \frac{\alpha}{2\pi \sin^2 \theta_W} \left(\hat{V}_{cs}^{L*} \hat{V}_{cd}^L X(x_c) + \hat{V}_{ts}^{L*} \hat{V}_{td}^L X(x_t) \right) \right]} \right|^2. \quad (8.82)$$

The experimental limit on this decay at 90% CL is [127]

$$\text{Br}(K_L \rightarrow \pi^0 \nu \bar{\nu})_{\text{exp}} < 3.0 \times 10^{-9}, \quad (8.83)$$

which is two orders of magnitude larger than the SM expectation [146]

$$\text{Br}(K_L \rightarrow \pi^0 \nu \bar{\nu})_{\text{SM}} = (3.00 \pm 0.30) \times 10^{-11}, \quad (8.84)$$

so that the experimental limit can be applied on the NP contribution. Then, we obtain the rephasing invariant bound:

$$\frac{\frac{1}{2} \left| \text{Im} \left[\left(\hat{V}_{us}^L \hat{V}_{ud}^{L*} \right) \hat{F}_{ds} \right] \right|}{\frac{\alpha}{2\pi \sin^2 \theta_W} \left| \text{Im} \left[\left(\hat{V}_{us}^L \hat{V}_{ud}^{L*} \right) \left(\hat{V}_{cs}^{L*} \hat{V}_{cd}^L X(x_c) + \hat{V}_{ts}^{L*} \hat{V}_{td}^L X(x_t) \right) \right] \right|} < 9.3, \quad (8.85)$$

that is, the new contribution can be ~ 10 times larger than the SM contribution.

A similar analysis leads to rephasing invariant constraints on the decays $K_L \rightarrow \mu\mu$ and $K_S \rightarrow \mu\mu$, which involve the real part and the imaginary part of the invariant $\hat{V}_{us}^L \hat{V}_{ud}^{L*} \hat{F}_{ds}$ respectively.

The same effective Lagrangian in eqs. (8.77) and (8.78) leads to the decay $K^+ \rightarrow \pi^+ \nu \bar{\nu}$. Ref. [127] estimates the SM contribution as $\text{Br}(K^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{SM}} = (0.81 \pm 0.04) \times 10^{-10}$ (it also states that parametric uncertainty in the CKM angles can result in shifts of the central value differing from this one by up to 10%). The experimental limit is $\text{Br}(K^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{exp}} = 1.14_{-0.33}^{+0.40} \times 10^{-10}$ at 2σ [127]. One obtains

$$0.98 \lesssim \left| -\frac{\frac{1}{2} \left(\hat{V}_{us}^L \hat{V}_{ud}^{L*} \right) \hat{F}_{ds}}{\frac{\alpha}{2\pi \sin^2 \theta_W} \left(\hat{V}_{us}^L \hat{V}_{ud}^{L*} \right) \left(\hat{V}_{cs}^{L*} \hat{V}_{cd}^L X(x_c) + \hat{V}_{ts}^{L*} \hat{V}_{td}^L X(x_t) \right)} + 1 \right| \lesssim 1.44, \quad (8.86)$$

Notice that the SM expectation is at the edge of the 2σ region determined by the experimental value. Then, a new contribution with modulus and phase

$$|\hat{F}_{ds}| \lesssim 3 \times 10^6, \quad 0 \leq \arg(\hat{\mathcal{T}}_{ds,u}) \lesssim 1.3, \quad 4.3 \leq \arg(\hat{\mathcal{T}}_{ds,u}) \lesssim 2\pi \quad (8.87)$$

would align the SM prediction to the experimental result and it would always be allowed. The result is shown in Figure 3.

Let us note that CP-violating effects in the RH sector are determined by the rephasing-invariant trilinears (see eq. (4.29)), which appear in the weak-basis invariants of mass dimension 10. This is related to the fact that CP violation in neutral meson mixing due to the mixing with the vector-like doublet only involve two quarks of the same sector (down for K and B , up for D mesons). As a consequence, the strength of CP violation can be expected to be stronger than in the SM.




Figure 4: Simplified sketch of possible Feynman diagrams producing CP interference in $K \rightarrow \pi\pi$, probed in ϵ' . On the left, the leading SM contribution, proportional to $\hat{V}_{ud}^L \hat{V}_{us}^{L*}$. In the middle the contribution due to the mixing with the vector-like doublet in charged currents, proportional to $\hat{V}_{ud}^L \hat{V}_{ud}^{R*}$ (see top panel of Figure 1 and the operator in eq. (8.11)). On the right, the new contribution from neutral currents to \hat{F}_{ds}^* (see bottom panel of Figure 1 and eq. (8.88)).

8.4.3 CP violation in $K \rightarrow \pi\pi$ by trilinears

We observe how the new CP-odd rephasing invariant associated to RH FCNCs appears and possible deviations from the SM could be hinting for non-zero values of it.

Additional strong bounds to the same effective 'triplet' invariant $\hat{\mathcal{T}}_{u,ds}$ can be obtained from ϵ' . The mechanism is sketched in Figure 4. The corresponding CP-violating Z -mediated RH $s \rightarrow d$ transition, generates $\Delta S = 1$ four-fermion operators below the EW scale again potentially inducing a very large CP violating $K \rightarrow (\pi\pi)_{I=2}$ amplitude. Indeed, using eq. (7.18) at the EW scale and singling out the chirality-enhanced $8_L \times 8_R$ operators under $SU(3)_L \times SU(3)_R$ transformations ($q_{L(R)}^l \rightarrow L(R)q_{L(R)}^l$, $q^l \equiv (u, d, s)^T$), one finds

$$\begin{aligned}
\mathcal{L} &\supset \frac{4G_F}{\sqrt{2}} \left[\bar{u}_L \gamma^\mu u_L \left(\frac{1}{2} - \frac{2}{3} \sin^2 \theta_W \right) - (\bar{d}_L \gamma^\mu d_L + \bar{s}_L \gamma^\mu s_L) \left(\frac{1}{2} - \frac{1}{3} \sin^2 \theta_W \right) \right] (\bar{s}_R \hat{F}_{sd} d_R) \\
&= \frac{4G_F}{\sqrt{2}} \sum_{q=u,d,s} \left(-\frac{1}{3} \bar{q}_L^q \gamma^\mu q_L^q + \cos^2 \theta_W \bar{q}_L^q Q_q \gamma^\mu q_L^q \right) (\bar{s}_R \gamma^\mu d_R) \\
&\supset \frac{4G_F}{\sqrt{2}} \cos^2 \theta_W \hat{F}_{sd} \left(\bar{q}_L^l Q \gamma^\mu q_L^l \right) (\bar{s}_R \gamma^\mu d_R) .
\end{aligned} \tag{8.88}$$

At leading order in the chiral counting, i.e. in the limit of exact $SU(3)_V$ symmetry, the $K \rightarrow (\pi\pi)_{I=2}$ amplitudes generated by $\bar{q}_L^l Q \gamma^\mu q_L^l \bar{s}_R \gamma^\mu d_R$ and $\bar{u}_L \gamma^\mu d_L \bar{s}_R \gamma^\mu u_R$ entering in eq. (8.11) are identical.³⁰ Thus the sensitivity of $\frac{c_W^2}{v^2} \hat{F}_{ds}$ to A_2 is approximately the same as $-\frac{1}{v^2} (\hat{V}_{us}^R \hat{V}_{ud}^{L*})$, that is, in this scenario the sensitivity to $c_W^2 \text{Im} \hat{\mathcal{T}}_{u,ds}$ is the same as the sensitivity to $-|\hat{V}_{ud}^L|^2 \text{Im} \hat{\mathcal{B}}_{us}$ above. Thus, using eqs. (8.53) and (8.54) we can

³⁰One way of seeing this is using eqs. (25) and (29) of Ref. [120]. The operator $\bar{q}_L^l Q q_L^l \bar{s}_R \gamma^\mu d_R$ induces $\text{Tr}[\frac{\lambda_{6-i7}}{2} U Q U^\dagger]$ with the same non-perturbative strength as $\bar{u}_L \gamma^\mu d_L \bar{s}_R \gamma^\mu u_R$ induces $\text{Tr}[\frac{\lambda_{4-i5}}{2} U \frac{\lambda_{1+i2}}{2} U^\dagger]$. Using that the explicit U convention is the one of eq. (4) of Ref. [147], one easily obtains the $K \rightarrow \pi\pi$ amplitudes.

Conventional name	Rephasing-invariant form	Bound on VLQ invariant
V_{ud}^β	$ \hat{V}_{ud}^L + \hat{V}_{ud}^R $	
$V_{us}^{K\ell 3}$	$ \hat{V}_{us}^L + \hat{V}_{us}^R $	$-\frac{\text{Re } \hat{\mathcal{B}}_{ud}}{ \hat{V}_{ud}^L } \simeq 10^{-3}, -\frac{\text{Re } \hat{\mathcal{B}}_{us}}{ \hat{V}_{us}^L } \simeq 10^{-3}$
$\left(\frac{V_{us}}{V_{ud}}\right)_{\frac{K\mu 2}{\pi\mu 2}}$	$\frac{ \hat{V}_{us}^L - \hat{V}_{us}^R }{ \hat{V}_{ud}^L - \hat{V}_{ud}^R }$	
$ \omega $	$\frac{\text{Re}(A_2 A_0^*)}{ A_0 ^2}$	$\frac{ \hat{V}_{us}^L \text{Re } \hat{\mathcal{B}}_{ud}}{ \hat{V}_{ud}^L } - \frac{ \hat{V}_{ud}^L \text{Re } \hat{\mathcal{B}}_{us}}{ \hat{V}_{us}^L } \lesssim 10^{-3}$
$\text{Re } \frac{\epsilon'}{\epsilon}$	$\frac{\text{Im}(A_2 A_0^*)}{\sqrt{2} \epsilon A_0 ^2}$	$\left \frac{ \hat{V}_{us}^L \text{Im } \hat{\mathcal{B}}_{ud}}{ \hat{V}_{ud}^L } + \frac{ \hat{V}_{ud}^L \text{Im } \hat{\mathcal{B}}_{us}}{ \hat{V}_{us}^L } \right \lesssim 6 \times 10^{-7}$
$d_{n,p}$	eqs. (8.58) and (8.59)	$\frac{\text{Im } \hat{\mathcal{B}}_{ud}}{ \hat{V}_{ud}^L }, \frac{\text{Im } \hat{\mathcal{B}}_{us}}{ \hat{V}_{us}^L } \lesssim 3 - 6 \times 10^{-6}$
$\Gamma(Z \rightarrow \text{had})$	eq. (8.61)	$\hat{F}_{\alpha\alpha}^u, \hat{F}_{ii}^d \lesssim 5 \times 10^{-3}$ (for $\alpha \neq t$)
$ \epsilon , \Delta m_K$	$\frac{ \text{Im}(M_{12} A_0 \bar{A}_0^*) }{\sqrt{2} (\Delta m_K) A_0 \bar{A}_0 }, 2 M_{12} $	$\frac{ \hat{\mathcal{T}}_{ds,u} }{ \hat{V}_{ud}^L \hat{V}_{us}^L } < 6 \times 10^{-7} - 2 \times 10^{-4}$
$\text{Br}(K_L \rightarrow \pi^0 \nu \bar{\nu})$	$\lambda_{\pi\nu\bar{\nu}}, \text{eq. (8.82)}$	$\frac{ \text{Im } \hat{\mathcal{T}}_{ds,u} }{ \hat{V}_{ud}^L \hat{V}_{us}^L } < 2 \times 10^{-5}$
$\text{Br}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$		$\frac{ \hat{\mathcal{T}}_{ds,u} }{ \hat{V}_{ud}^L \hat{V}_{us}^L } < (3 - 8) \times 10^{-6}$

Table 6: Summary table of constraints on rephasing invariant mixing elements in presence of one VLQ doublet of quarks (assuming no cancellations).

write a bound to the non-standard contributions to ϵ'/ϵ as:

$$\left| \frac{\text{Im} \left[|\hat{V}_{us}^L|^2 \hat{\mathcal{B}}_{ud} + |\hat{V}_{ud}^L|^2 \hat{\mathcal{B}}_{us} - \hat{\mathcal{T}}_{u,ds} \cos^2 \theta_W \right]}{|\hat{V}_{us}^L \hat{V}_{ud}^L|} \right| \lesssim 5.6 \times 10^{-7}. \quad (8.89)$$

Let us also notice that when $z_u \neq 0$ this limit can be written as

$$\left| \frac{\text{Im} \left[|\hat{V}_{us}^L|^2 \hat{\mathcal{B}}_{ud} + |\hat{V}_{ud}^L|^2 \hat{\mathcal{B}}_{us} - \hat{\mathcal{B}}_{ud}^* \hat{\mathcal{B}}_{us} \frac{\cos^2 \theta_W}{\hat{F}_{uu}^2} \right]}{|\hat{V}_{us}^L \hat{V}_{ud}^L|} \right| \lesssim 5.6 \times 10^{-7}. \quad (8.90)$$

8.5 CAA-motivated scenarios

8.5.1 One VLQ doublet ($N = 1$)

Here, we examine in more detail the scenarios with vector-like doublets relevant for the Cabibbo angle anomalies, using the rephasing invariant formalism described in this work. As already recalled, it was shown in Ref. [49] that a single vector-like doublet mixing with the up, the down and the strange SM quarks cannot explain all the tensions, since the required couplings are excluded by limits on flavour-changing neutral currents (see also eq. (8.76)). We start by reviewing two scenarios with one extra doublet, in one case exhibiting large mixing with the up and the down SM quarks ($z_u \neq 0$ and $z_d \neq 0$), which can resolve the tension between V_{ud} and V_{us} (CAA1), and in a second case coupling predominantly to the up and strange quarks $z_u \neq 0$ and $z_s \neq 0$, causing of the tension between the $K\ell 3$ and $K\mu 2/\pi\mu 2$ determinations of V_{us} (CAA2). Then we study the scenario with the presence of two vector-like doublets with the same pattern of couplings respectively.

In the one doublet scenarios, the only non-zero matrix elements of the corresponding charge and neutral current matrices are either (CAA1)

$$\hat{V}_{ud}^R = \frac{v^2}{M_Q^2} z_u^* z_d, \quad \hat{F}_{dd} = \frac{v^2}{M_Q^2} |z_d|^2 > 0, \quad \hat{F}_{uu} = \frac{v^2}{M_Q^2} |z_u|^2 > 0, \quad (8.91)$$

or (CAA2)

$$\hat{V}_{us}^R = \frac{v^2}{M_Q^2} z_u^* z_s, \quad \hat{F}_{ss} = \frac{v^2}{M_Q^2} |z_s|^2 > 0, \quad \hat{F}_{uu} = \frac{v^2}{M_Q^2} |z_u|^2 > 0. \quad (8.92)$$

By performing a χ^2 fit of the determinations of $|V_{ud}|_\beta$, $|V_{us}|_{K\ell 3}$, $|V_{us}/V_{ud}|_{K\mu 2/\pi\mu 2}$ using the values in eq. (8.5) and the parameters in eq. (8.7) ($|\hat{V}_{us}^L|$ and $\text{Re}(\hat{V}_{ud}^{L*} \hat{V}_{ud}^R)/|\hat{V}_{ud}^L|$ or $\text{Re}(\hat{V}_{us}^{L*} \hat{V}_{us}^R)/|\hat{V}_{us}^L|$ in the two cases respectively) one finds for CAA1

$$\frac{\text{Re}(\hat{V}_{ud}^{L*} \hat{V}_{ud}^R)}{|\hat{V}_{ud}^L|} = -0.55(29) \times 10^{-3}, \quad |\hat{V}_{us}^L| = 0.22451(38), \quad (8.93)$$

with $\hat{V}_{ud}^L = 0.97447(9)$ obtained from unitarity of \hat{V}_L , or for CAA2

$$\frac{\text{Re}(\hat{V}_{us}^{L*} \hat{V}_{us}^R)}{|\hat{V}_{us}^L|} = -1.09(36) \times 10^{-3}, \quad |\hat{V}_{us}^L| = 0.22453(34), \quad (8.94)$$

with $\hat{V}_{ud}^L = 0.97446(8)$ from unitarity.

In these scenarios very clean and distinct predictions can be made about non-standard flavour-conserving neutral currents, since the mixing matrices of the RH charged currents \hat{V}_R and neutral currents \hat{F}_q originate from the same couplings (see




Figure 5: Neutral current space corresponding to eq. (8.98), one doublet case, and eq. (8.102), two doublets case, together with the excluded region at 95% CL level considering $Z \rightarrow \text{had}$ plus atomic parity violation (see text for details).

e.g. eqs. (8.91) and (8.92)). Hence, by assuming the presence of RH charged currents one is predicting a non-zero shift in RH neutral currents, which can be tested for example in Z -pole observables. In particular, in these two cases we have

$$\hat{F}_{dd}\hat{F}_{uu} = |\hat{V}_{ud}^R|^2 = \frac{[\text{Re } \hat{\mathcal{B}}_{ud}]^2 + [\text{Im } \hat{\mathcal{B}}_{ud}]^2}{|\hat{V}_{ud}^L|^2}, \quad (8.95)$$

or

$$\hat{F}_{ss}\hat{F}_{uu} = |\hat{V}_{us}^R|^2 = \frac{[\text{Re } \hat{\mathcal{B}}_{us}]^2 + [\text{Im } \hat{\mathcal{B}}_{us}]^2}{|\hat{V}_{us}^L|^2}. \quad (8.96)$$

In order to explain the Cabibbo angle anomalies we can substitute the values of $\text{Re } \hat{\mathcal{B}}_{ud(s)}/|\hat{V}_{ud(s)}^L|$ in eqs. (8.93) and (8.94). Then, one finds the relations

$$\hat{F}_{dd}\hat{F}_{uu} \geq \frac{(\text{Re } \hat{\mathcal{B}}_{ud})^2}{|\hat{V}_{ud}^L|^2} = (0.55(29) \times 10^{-3})^2, \quad (8.97)$$

or

$$\hat{F}_{ss}\hat{F}_{uu} \geq \frac{(\text{Re } \hat{\mathcal{B}}_{us})^2}{|\hat{V}_{us}^L|^2} = (1.09(36) \times 10^{-3})^2. \quad (8.98)$$

However, eqs. (8.95) and (8.96) should also be confronted with the strong experimental limits on CP-violating processes, which constrain the CP-odd rephasing invariants $\text{Im } \hat{\mathcal{B}}_{ud(s)}$ (see eqs. (8.54) and (8.60)). After taking into account the CP-violating probes, the product of couplings $\hat{F}_{uu}\hat{F}_{dd,ss}$ can be sharply determined and the relations in eqs. (8.97) and (8.98) become equalities at the per mille level.




Figure 6: Parameter space in the scenario with two vector-like quark doublets coupling with up and down and up and strange quark respectively, resolving CAA1 and CAA2 (see text for details).

Moreover, these relations should also include the other experimental constraints, from electroweak and Z-boson observables (see e.g. eq. (8.64) and Ref. [45]). Then, after considering all phenomenological effects, these scenarios lead to precise predictions on each coupling \hat{F}_{uu} , \hat{F}_{dd} (or \hat{F}_{ss}).

By performing a χ^2 fit for CAA2 using the parameters \hat{F}_{uu} , \hat{F}_{ss} , V_{Lus} , including the Cabibbo angle determinations of eq. (8.5) and the observables in section 8.3.4 (i.e. $\Gamma(Z \rightarrow \text{had})$, $Q_W(p)$, $Q_W(Cs)$) we obtain an improvement of $\chi_{\text{SM}}^2 - \chi_{\text{min}}^2 = 10.1$ which is independent of the mass of the vector-like doublet. The remaining discrepancy is due to the difference between the determinations of Cabibbo angle from kaon and β decays. In Figure 5a we show the 1σ (green) region ($\chi_{\text{min}}^2 + 1$). The constraints at 2σ CL are also displayed, for the Z boson decay to hadrons (magenta), the weak charge of the proton (red). In addition, we illustrate the relation (8.98) in Figure 5a. In particular, we show the result of the fit of the anomalies, namely eq. (8.5), as the blue curves, indicating 1σ region ($\chi_{\text{min}}^2 + 1$). By assuming that the charged current couplings assume the value needed for the anomaly, that is eq. (8.94), CP violation limits forbid the striped cyan region (which would be allowed by the CAA solution) and the neutral current couplings lie inside the blue band.

8.5.2 Two VLQ doublets ($N = 2$)

We now examine a scenario with two vector-like doublets, which could be the source of all the tensions in the first row of the CKM matrix while also not contradicting the stringent limits from flavour changing neutral currents and other precision electroweak observables. We assume that one doublet couples predominantly with the up-quark and the down quark while a second doublet has large couplings with the up quark and the strange quark. The mass matrices related to this scenario are given in eq. (6.37), with the mixing matrices V_R , F_u , F_d given in eqs. (6.40), (6.41) and (6.44). Then, the relevant




Figure 7: Neutral current space in the scenario with two vector-like quark doublets coupling with up and down and up and strange quark respectively, as predicted by resolving CAA1 and CAA2 (see text for details).

matrix elements of the corresponding charged and neutral current matrices read

$$\begin{aligned}
\hat{V}_{ud}^R &= \frac{v^2}{M_Q^2} z_{1u}^* z_{1d}, & \hat{V}_{us}^R &= \frac{v^2}{a^2 M_Q^2} z_{2u}^* z_{2s}, \\
\hat{F}_{dd} &= \frac{v^2}{M_Q^2} |z_{1d}|^2, & \hat{F}_{ss} &= \frac{v^2}{a^2 M_Q^2} |z_{2s}|^2, & \hat{F}_{uu} &= \frac{v^2}{M_Q^2} \left(|z_{1u}|^2 + \frac{|z_{2u}|^2}{a^2} \right),
\end{aligned} \tag{8.99}$$

where again it is clear that there exists a direct connection between neutral and charged mixing elements. In particular, as right handed charged currents are assumed in order to explain the anomalies, then modified neutral current couplings are predicted with the peculiar relation

$$\hat{F}_{uu} = \frac{|\hat{V}_{ud}^R|^2}{\hat{F}_{dd}} + \frac{|\hat{V}_{us}^R|^2}{\hat{F}_{ss}}. \tag{8.100}$$

In order to explain the Cabibbo angle anomalies the charged currents mixing $\text{Re } \hat{\mathcal{B}}_{ud} / |\hat{V}_{ud}^L|$ and $\text{Re } \hat{\mathcal{B}}_{us} / |\hat{V}_{us}^L|$ should assume the values in eq. (8.8). Then, one gets the relation between charged- and neutral-current couplings:

$$1 = \frac{|\hat{V}_{ud}^R|^2}{\hat{F}_{uu} \hat{F}_{dd}} + \frac{|\hat{V}_{us}^R|^2}{\hat{F}_{uu} \hat{F}_{ss}} \geq \frac{(\text{Re } \hat{\mathcal{B}}_{ud})^2}{|\hat{V}_{ud}^L|^2} \frac{1}{\hat{F}_{uu} \hat{F}_{dd}} + \frac{(\text{Re } \hat{\mathcal{B}}_{us})^2}{|\hat{V}_{us}^L|^2} \frac{1}{\hat{F}_{uu} \hat{F}_{ss}}. \tag{8.101}$$

Hence, the presence of right handed currents generating the Cabibbo angle anomalies would imply flavour-conserving neutral currents with couplings above the hyperbola of axes $\hat{F}_{uu} \hat{F}_{dd}$ and $\hat{F}_{uu} \hat{F}_{ss}$. After taking into account experimental bounds on the corresponding CP violating phases, the product of flavour-conserving neutral-current couplings lie *on* the hyperbola at the per mille level:

$$\hat{F}_{uu} = \frac{(0.79(27) \times 10^{-3})^2}{\hat{F}_{dd}} + \frac{(1.24(37) \times 10^{-3})^2}{\hat{F}_{ss}}, \tag{8.102}$$

where we inserted in eq. (8.101) the values in eq. (8.8). This relation is shown in Figure 5b (blue curves, corresponding to $\chi^2 = 1$ region). While the anomalies can be addressed in the region above the curves, CP-violation bounds force the couplings to remain in the blue band once the charged current couplings acquire the values needed for CAA1 and CAA2.

The RH charged currents explaining the anomalies hence imply the presence of anomalous Z couplings because of the connection between charged- and neutral-current couplings. Neutral-current couplings are also subject to constraints from electroweak and Z -boson observables (see e.g. eq. (8.64)) which should be included in the phenomenological analysis of the Cabibbo angle anomalies. Then, after considering all phenomenological effects, the prediction of each coupling \hat{F}_{uu} , \hat{F}_{dd} , \hat{F}_{ss} become quite accurate and testable for instance by improving precision on Z -decay observables.

We show this connection and the corresponding parameter space in Figures 6 and 7. In Figure 6 we illustrate the result of a χ^2 fit using the parameters $-z_u v/M_Q$, $-z_s v/M_Q$, $-z_d v/M_Q$ and V_{Lus} including the observables: determinations of the Cabibbo angle (8.5), $\Gamma(Z \rightarrow \text{had})$, $Q_W(p)$ and $Q_W(Cs)$ (see section 8.3.4). We receive an improvement over the SM $\chi_{\text{SM}}^2 - \chi_{\text{min}}^2 = 18.1$. The 1σ and 2σ regions ($\chi_{\text{min}}^2 + 1$, $\chi_{\text{min}}^2 + 4$) are displayed in green and light-green, marginalizing over the other variables. We also show the constraints at 2σ CL for the Z boson decay to hadrons (magenta), the weak charge of the proton (red), the atomic parity violation in ^{133}Cs (purple). The fit including only the anomalies, namely eq. (8.5), is indicated by the blue bands, which indicate the 1σ region ($\chi_{\text{min}}^2 + 1$) corresponding to eqs. (8.8).

The analogous χ^2 fit can be performed using the neutral currents couplings as parameters, i.e. \hat{F}_{uu} , \hat{F}_{ss} , \hat{F}_{dd} (and V_{Lus}) including the same observables in (8.5), and section 8.3.4. In Figure 7 we illustrate the results of the fit on the \hat{F}_{uu} - \hat{F}_{ss} , \hat{F}_{uu} - \hat{F}_{dd} and \hat{F}_{dd} - \hat{F}_{ss} planes, respectively, marginalizing over the other variables. The 1σ and 2σ regions of the fit ($\chi_{\text{min}}^2 + 1$, $\chi_{\text{min}}^2 + 4$) are presented in green and light green. The relation in eq. (8.102) emerges in the stand-alone fit of the anomalies, which at one sigma gives the cyan curves in the plots. CAA1 and CAA2 are resolved in the whole region above the curves.

Besides being the potential resolution of the Cabibbo angle anomalies, the Yukawa textures in eq. (6.37) are compatible with all other experimental constraints, in particular the stringent limit on flavour changing neutral currents. In fact, as it can be inferred from eqs. (6.40) and (6.41), there are not flavour changing neutral currents at tree level. There still is contribution at loop level e.g. to neutral kaon mixing:

$$M_{12,\text{NP}}^K \simeq \frac{1}{3} m_{K^0} f_K^2 \frac{G_F^2 m_W^2}{4\pi^2} \left(\frac{\lambda_u}{|\lambda_u|} s_{45} c_{45} z_{1d} z_{2s}^* \frac{v^2}{M_Q^2} \right)^2 \times \left[F \left(\frac{a^2 M_Q^2}{m_W^2} \right) + F \left(\frac{M_Q^2}{m_W^2} \right) - 2 F \left(\frac{a^2 M_Q^2}{m_W^2}, \frac{a^2 M_Q^2}{m_W^2} \right) \right], \quad (8.103)$$

where $F(x)$ are Inami-Lim functions [134]. This contribution disappears in the limit of mass degeneracy of the two heavy species (corresponding to $a = 1$). In the general

case on which the masses of the vector-like doublets are not equal, the combination of functions $F(x)$ increases with the mass M_Q^2 as $\sim x f(a)$, where $x = M_Q^2/m_W^2$ and $f(a)$ is a number which depends on the value of a . Let us emphasize that the combination of mixing elements in eq. (8.103) is rephasing invariant.

We want to test this effect in the scenario explaining the Cabibbo angle anomalies. Then, we should consider that there exist an upper limit on the vector-like doublet masses. In fact, $|z_u^* z_{d(s)}| v^2/M_Q^2 \simeq 10^{-3}$, assuming the couplings $z_{u,d,s}$ to be at most 1, we have $M_Q < 6.2$ TeV, $a M_Q < 4.9$ TeV. On the other hand, there is a lower bound on the mass given by the LHC limits, $M_Q, a M_Q \gtrsim 1$ TeV. Then, the mass M_Q and parameter a are constrained as

$$1 \text{ TeV} \lesssim M_Q \lesssim 6.2 \text{ TeV}, \quad 4.9 > a > 0.16, \quad (8.104)$$

with $a = 0.16$ when $M_Q = 6.2$ TeV and $a = 4.9$ when $M_Q = 1$ TeV. With these values, $f(a)$ is a number smaller than 1 ($f(0.16) \sim 0.1$, $f(5) \sim 0.5$).

The new contribution should be compared to the SM one in eq. (8.66) and confronted with experimental constraints on Δm_K and ϵ_K . Bounds on the new physics contribution can be estimated as $|M_{12,\text{NP}}^K| < |M_{12,\text{SM}}^K| \Delta_K$, $|\text{Im} M_{12,\text{NP}}^K| < |\text{Im} M_{12,\text{SM}}^K| \Delta_{\epsilon_K}$, with $\Delta_K = 1$ and, using the results in Ref. [141], $\Delta_{\epsilon_K} = 0.3$. Then, by comparing the new contribution in eq. (8.103) to the SM one in eq. (8.66) we have

$$\begin{aligned} \text{Im} \left[\frac{\lambda_u^2}{|\lambda_u|^2} (s_{45} c_{45} z_{1d} z_{2s}^*)^2 \right] \frac{v^4}{M_Q^4} \frac{M_Q^2 f(a)}{m_W^2} &\lesssim 1.6 \times 10^{-7}, \\ \left| \frac{\lambda_u^2}{|\lambda_u|^2} (s_{45} c_{45} z_{1d} z_{2s}^*)^2 \right| \frac{v^4}{M_Q^4} \frac{M_Q^2 f(a)}{m_W^2} &\lesssim 2.3 \times 10^{-5}. \end{aligned} \quad (8.105)$$

In the limit $|a^2 - 1|/a \gg |z_{1u} z_{2u}| v^2/M_Q^2$ and $|a^2 - 1| \gg ||z_{1u}|^2 - |z_{2u}|^2| v^2/M_Q^2$, s_{45} is given by eq. (6.43) and we have

$$\left| \frac{\lambda_u^2}{|\lambda_u|^2} \left(z_{1u}^* z_{1d} z_{2u} z_{2s}^* \frac{v^4}{M_Q^4} \right)^2 \right| \frac{af(a)}{a^2 - 1} \frac{M_Q^2}{m_W^2} \simeq (10^{-6})^2 \frac{af(a)}{a^2 - 1} \frac{M_Q^2}{m_W^2} \lesssim 10^{-10}, \quad (8.106)$$

which is well below the experimental constraints. Here in the first parentheses the charged-current couplings $\hat{V}_{ud}^R, \hat{V}_{us}^R$ responsible for the CAA appear and we substituted their values, see eq. (8.8). We can wonder if there is a peculiar intermediate scenario, in which the combination of Inami-Lim functions may not disappear while the mixing angle can be large, as $\theta_{45} \simeq \pi/4$, with $1 - |a| \lesssim 10^{-3}$. In this case we would have

$$\left| \frac{\lambda_u^2}{|\lambda_u|^2} (s_{45} c_{45} z_{1d} z_{2s}^*)^2 \right| \frac{v^4}{M_Q^4} \frac{M_Q^2 f(a)}{m_W^2} \simeq \frac{(10^{-6})^2}{4} \frac{M_Q^4}{v^4} \frac{M_Q^2 f(a)}{m_W^2} \lesssim 10^{-10}, \quad (8.107)$$

where the last inequality follows since for $|1 - |a|| \lesssim 10^{-3}$ the combination of Inami-Lim functions gives a value $< 5 \times 10^{-4}$ for $M_Q = 6.2$ TeV.

Let us close this section with two remarks. If Cabibbo anomalies were to be confirmed and addressed by this scenario, two new puzzles would emerge: 1. Why does each VLQ (mostly) couple to only one light SM quark family in each sector (needed to evade FCNC constraints)? 2. A priori CP phases in nature are arbitrary. Why does one have $|\arg \hat{\mathcal{B}}_{ud(s)}| = |\arg(\hat{V}_{ud(s)}^{L*} \hat{V}_{ud(s)}^R) - \pi| \lesssim 10^{-3}$? Perhaps a symmetry mechanism, beyond the reach of this work, could shed some light on both issues. Let us also notice that, in an alternative scenario in which e.g. $\text{Re} \hat{\mathcal{B}}_{ud} \sim \text{Im} \hat{\mathcal{B}}_{ud}$, CP-violating probes on the invariant $|\text{Im} \hat{\mathcal{B}}_{ud}| \lesssim 10^{-6}$ are naturally testing possible scenarios with VLQ doublets at $M_Q \sim 200 \text{ TeV}$, far beyond the direct reach of current and near future high-energy colliders, thus becoming one of the highest-scale BSM scenarios currently being tested and not violating any of the SM symmetries (accidental or not).

9 Summary and conclusions

In this work, we studied models of VLQ doublets with standard charges, which are minimal extensions of the SM and highly motivated from a model-building point of view. Notably, these models provide favoured solutions to the Cabibbo angle anomalies (CAAs).

We emphasized that a prominent tool in the analysis of these models is obtained through a weak-basis invariant (WBI) description. In an extensive survey, we explained how to construct WBIs for VLQ doublet models and how to relate these to physical parameters.³¹

An important and differentiating aspect of WBIs constructed for VLQ doublet models — and in contrast with the SM or VLQ singlet extensions — is that, due to the presence of exotic right-handed currents new and simpler rephasing invariants and WBIs can be obtained, which can also be related to Lagrangian parameters and observables. This extra character of doublet representations allows for lower-mass-dimension ($M = 6$) CP-odd WB invariants. This could represent a significant enhancement of CPV with respect to the SM or even to the VLQ-singlet extensions.

For the one-doublet case, we found a special weak basis, referred to as the “stepladder” WB, which in a simple and comprehensive way enables us to extract all physical parameters step-by-step, without having to solve complicated equations. We identified a set of WBIs totally characterizing the single VLQ doublet scenario ($N = 1$). An alternative procedure, using instead the minimal weak basis of section 3.1.1, is also used to generalize this characterization to an arbitrary number of doublets ($N > 1$). Moreover, we presented useful approximate and exact parametrizations for the quark mixing in section 4, for arbitrary N .

We then focused on studying rephasing invariants and CP-odd WBIs, whose imaginary parts signal the presence of CPV. In the doublet scenario, new kinds of rephasing invariants arise, which result from the existence of right-handed charged and neutral currents mixing matrices V_R , F_u and F_d . The interplay between the two types of charged currents, right-handed and left-handed, generates bilinears, of the form $V_{\alpha i}^L V_{\alpha i}^{R*}$, involving only two quarks. Flavour changing neutral currents give rise to trilinear rephasing invariants of the form $F_{\alpha\beta}^u V_{\alpha i}^{L*} V_{\beta i}^L$ and $F_{ij}^d V_{\alpha i}^L V_{\alpha j}^{L*}$, involving three quarks.

We found that the mass dimension and number of insertions of the mixing elements in WBIs play a crucial role. In fact, we uncover a direct connection between the type of *effective* rephasing invariants contained in different WBIs and their structure, i.e. the number of Hermitian blocks from which the WBIs are built. We denote by *effective* the rephasing invariants that involve only the three SM quarks and that emerge after taking into account the large hierarchy between the EW and the VLQ mass scale. These features then provide a deeper understanding of the links between WBIs, rephasing invariants and observables. We also find a complete set of WBIs which describe the CP properties of the one VLQ doublet scenario and whose vanishing would ensure CP invariance. We present this set in table 3. Moreover, we briefly studied the extreme chiral limit for the

³¹The physical parameter count for an arbitrary number N of VLQ doublets is given in Table 2.

one- and two-VLQ doublet scenarios. We show that in this regime of extremely high energies, even though the lighter quark masses can be taken as vanishing, CPV can still be achieved, in contrast to what happens in the SM.

After revisiting the EFT form of these models at energies below the VLQ masses, we carried out a phenomenological analysis, using the WBI description previously developed. Such description in terms of invariants provides unambiguous constraints on the model, which is particularly attractive when describing CP-violating phenomena. The crucial connection between WBIs and effective rephasing invariants hints at the strength of different CP-violating VLQ contributions — the different structures of invariants emerging in the presence of VLQ doublets suggest which processes one can expect to be most affected, particularly when involving CP-violation, prior to any explicit computation.

We studied the phenomenological implications in terms of the new effective rephasing invariants and their interplay with CPV, with special attention to the bilinears induced by the new effective charged right-handed currents and trilinears induced by the new neutral right-handed currents. They lead to significant contributions for e.g. direct CP violation, kaon mixing, rare kaon decays, electric dipole moments, low-energy EW flavour-conserving observables and Z decays. We calculated these and imposed limits on the corresponding invariant quantities allowed by the model. The most relevant phenomenological results are summarized in Table 6.

Finally, we analyzed specific scenarios addressing the Cabibbo Anomalies with VLQ doublets, explicitly showing, using rephasing invariant quantities, the remarkable relations between RH neutral- and charged-current observables involving both CP-even and CP-odd probes.

Acknowledgements

This work was partially supported by Fundação para a Ciência e a Tecnologia (FCT, Portugal) through the projects CERN/FIS-PAR/0002/2021, 2024.02004.CERN and CFTP-FCT Unit UIDB/00777/2020 and UIDP/00777/2020, which are partially funded through POCTI (FEDER), COMPETE, QREN and EU. The work of B.B. was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under grant 396021762 - TRR 257. A.R.S. was supported by the Generalitat Valenciana (Spain) through the plan GenT program (CIDEIG/2023/12) and by the Spanish Government (Agencia Estatal de Investigación MCIN/AEI/10.13039/501100011033) Grant No. PID2023-146220NB-I00.

A Notation

Table 7: Notations introduced in the text and equations where they are defined.

Notation	Description ($q = u, d; \chi = L, R$)	Definition
N	number of VLQ doublets added to the SM	eq. (2.1)
\mathcal{M}_q	$(3 + N)$ -dimensional mass matrix of q -type quarks	eq. (2.5)
Y_q	3×3 standard q -type Yukawa matrix	eq. (2.2)
Z_q	$N \times 3$ matrix of q -type Yukawa couplings to VLQs	eq. (2.2)
\overline{M}	$3 \times N$ matrix of bare mass terms involving SM quarks	eq. (2.2)
M_Q	$N \times N$ matrix of VLQ-only bare mass terms	eq. (2.2)
D_Q	M_Q in a basis in which it is diagonal and non-negative	eq. (3.5)
\mathcal{Y}_q	$(3 + N) \times 3$ matrix of all q -type Yukawa couplings	eq. (2.6)
m_q	the product $v \mathcal{Y}_q$, i.e. all masses terms or EW origin	eq. (2.6)
M	$(3 + N) \times 3$ matrix of all bare masses terms	eq. (2.6)
\mathcal{V}_χ^q	$(3 + N)$ -dimensional unitary rotations bidiagonalizing \mathcal{M}_q	eq. (2.8)
\mathcal{D}_q	$(3 + N)$ -dim. diagonal matrix of physical q -type masses	eq. (2.10)
M^q	new heavy physical masses (M_1^q, \dots, M_N^q)	eq. (2.10)
A_χ^q	matrix containing the first 3 rows of \mathcal{V}_χ^q	eq. (2.12)
B_χ^q	matrix containing the last N rows of \mathcal{V}_χ^q	eq. (2.12)
V_L	$(3 + N)$ -dim. LH charged-current mixing matrix (unitary)	eq. (2.16)
V_R	$(3 + N)$ -dim. RH charged-current mixing matrix (non-unit.)	eq. (2.16)
V_{CKM}	3×3 non-unitary CKM mixing matrix	eq. (2.18)
F_q	$(3 + N)$ -dim. Hermitian matrix controlling q -sector FCNCs	eq. (2.21)
\hat{V}_L	V_{CKM} at leading order, part of a parameterization of V_L	eq. (4.23)
\hat{V}_R	3×3 upper-left block of V_R at leading order, cf. eq. (4.19)	eq. (4.29)
\hat{F}_q	3×3 upper-left block of F_q at leading order, cf. eq. (4.20)	eq. (4.29)
\mathcal{W}_L	weak-basis rotation of LH fields	eq. (3.1)
W_R	weak-basis rotation of RH VLQ fields	eq. (3.1)
W_R^q	weak-basis rotation of RH standard q -type quarks	eq. (3.1)
\hat{Y}_q	Y_q in a basis in which it is diagonal and non-negative	eq. (3.7)

Table 7: (continued)

Notation	Description ($q = u, d; \chi = L, R$)	Definition
$\hat{U}_{q\chi}$	3×3 unitary rotations bidiagonalizing Y_q	eq. (3.7)
\hat{V}	3×3 CKM-like matrix in the definition of the minimal WB	eq. (3.9)
\mathbf{z}_q	rotated Z_q in the definition of the minimal WB	eq. (3.10)
$\hat{\mathbf{z}}_q$	rephased \mathbf{z}_q , with N less phases	eq. (3.12)
\tilde{V}	3×3 unitary matrix in the definition of the stepladder WB	eq. (3.13)
\mathcal{B}	rephasing-invariant bilinears, defined in terms of V_χ and F_q	eq. (3.26)
\mathcal{T}	rephasing-invariant trilinears, defined in terms of V_χ and F_q	eq. (3.27)
\mathcal{Q}	rephasing-invariant quartets, defined in terms of V_χ and F_q	eq. (3.28)
$\hat{\mathcal{B}}$	rephasing-invariant bilinears, defined in terms of \hat{V}_χ and \hat{F}_q	eq. (4.29)
$\hat{\mathcal{T}}$	rephasing-invariant trilinears, defined in terms of \hat{V}_χ and \hat{F}_q	eq. (4.29)
$\hat{\mathcal{Q}}$	rephasing-invariant quartets, defined in terms of \hat{V}_χ and \hat{F}_q	eq. (4.29)
$\hat{\mathcal{Q}}$	rephasing-invariant quartets generalizing the SM ones	eq. (4.31)
\mathcal{H}_q	LH Hermitian matrix $\mathcal{M}_q \mathcal{M}_q^\dagger = h_q + H$	eq. (5.1)
h_q	LH Hermitian matrix $m_q m_q^\dagger$	eq. (3.3)
H	LH Hermitian matrix $M M^\dagger$	eq. (3.3)
\mathcal{H}_{qR}	RH Hermitian matrix $\mathcal{M}_q^\dagger \mathcal{M}_q$	eq. (5.1)
h_{qR}	RH Hermitian matrix $m_q^\dagger m_q$	eq. (5.9)
H_R	RH Hermitian matrix $M^\dagger M$	eq. (5.10)

B Effective CKM angles vs. Lagrangian parameters

The effective matrices $D^q = \text{diag}(m_1^q, m_2^q, m_3^q)$ and \hat{V}_L in eqs. (7.15) and (7.16) are not directly \hat{Y}_q and \hat{V} of the minimal WB, eq. (3.12). Let us explicitly relate them to the initial (minimal) parameters in \hat{Y}_u, \hat{Y}_d and \hat{V} . From eqs. (7.13) and (7.14) one has, up to $\mathcal{O}(1/M_Q^4)$ corrections,

$$(D^q)_{ii} = m_i^q = v \hat{Y}_{qi} \left(\mathbf{1} - \frac{1}{2} \hat{F}_{ii}^q \right), \quad (\text{B.1})$$

and

$$\begin{aligned} i \alpha_i^u &\equiv i (\delta L_q)_{ii} = i (\delta R_u)_{ii}, \\ i \alpha_i^d &\equiv i (\delta L_d)_{ii} = i (\delta R_d)_{ii}, \end{aligned} \quad (\text{B.2})$$

where we have defined $i \delta L_d \equiv \mathbf{1} - \hat{V}_L^\dagger (\mathbf{1} - i \delta L_q) \hat{V}$. For $i \neq j$, one has instead

$$\begin{aligned} i (\delta L_q)_{ij} &\stackrel{i \neq j}{=} \frac{m_i^u m_j^u}{(m_i^u)^2 - (m_j^u)^2} \hat{F}_{ij}^u, & i (\delta L_d)_{ij} &\stackrel{i \neq j}{=} \frac{m_i^d m_j^d}{(m_i^d)^2 - (m_j^d)^2} \hat{F}_{ij}^d, \\ i (\delta R_u)_{ij} &\stackrel{i \neq j}{=} \frac{1}{2} \frac{(m_i^u)^2 + (m_j^u)^2}{(m_i^u)^2 - (m_j^u)^2} \hat{F}_{ij}^u, & i (\delta R_d)_{ij} &\stackrel{i \neq j}{=} \frac{1}{2} \frac{(m_i^d)^2 + (m_j^d)^2}{(m_i^d)^2 - (m_j^d)^2} \hat{F}_{ij}^d. \end{aligned} \quad (\text{B.3})$$

Recall that \hat{V}_L and \hat{V} have the same functional form, with $\hat{V}_L = V(\vec{\theta}_{\hat{V}_L}) \simeq V_{\text{CKM}}$ and $\hat{V} = V(\vec{\theta}_{\hat{V}})$. Thus, $\vec{\theta}_{\hat{V}_L} = (\theta_{12}, \theta_{13}, \theta_{23}, \delta_{13})$, while $\vec{\theta}_{\hat{V}} \equiv (\theta_{12}^{\hat{V}}, \theta_{13}^{\hat{V}}, \theta_{23}^{\hat{V}}, \delta_{13}^{\hat{V}})$. Expanding $V(\vec{\theta})$ in the neighbourhood of $\vec{\theta}_{\hat{V}_L}$ and using the definition of δL_d , one finds

$$\frac{\partial V}{\partial \vec{\theta}} \Big|_{\vec{\theta} = \vec{\theta}_{\hat{V}_L}} \cdot (\vec{\theta}_{\hat{V}_L} - \vec{\theta}_{\hat{V}}) = i \left[\hat{V} \delta L_d - \delta L_q \hat{V} \right], \quad (\text{B.4})$$

which can be used to extract the five unphysical external phase differences $\alpha_i^u - \alpha_j^d$ in the field redefinitions and, crucially, $(\theta_{12}, \theta_{13}, \theta_{23}, \delta_{13})$. The exact solution is relatively cumbersome and offers limited insight. Replacing instead the CKM angles and phase and quark masses by their experimental values (correct up to neglected second-order $\mathcal{O}(1/M_Q^4)$ corrections) and keeping only the largest numerical coefficients, one finds

$$\begin{aligned} \theta_{12} &\simeq \theta_{12}^{\hat{V}} - 0.050 \text{ Re } \hat{F}_{12}^d, \\ \theta_{23} &\simeq \theta_{23}^{\hat{V}} - 0.018 \text{ Re } \hat{F}_{23}^d, \\ \theta_{13} &\simeq \theta_{13}^{\hat{V}} - 0.00033 \text{ Re } \hat{F}_{13}^d - 0.0015 \text{ Re } \hat{F}_{23}^d + 0.00086 \text{ Re } \hat{F}_{13}^d + 0.0040 \text{ Re } \hat{F}_{23}^d, \\ \delta_{13} &\simeq \delta_{13}^{\hat{V}} + 1.1 \text{ Re } \hat{F}_{23}^d. \end{aligned} \quad (\text{B.5})$$

Notice how the leading prefactors in the corrections of the parameters happen to roughly scale as the values of the parameters themselves. In the limit of all CKM angles equal to

zero, the (typically largest) numerical prefactors in front of $\text{Re } \hat{F}_{ij}^d$ for the θ_{ij} corrections are approximately $-m_i^d/m_j^d$, which accidentally happen to be of the same order as the θ_{ij} themselves. More importantly, the rotation from the parameters in \hat{V} to the ones in \hat{V}_L is always suppressed by FCNC factors. Thus, for phenomenological purposes — and unlike for the mass-like parameters $D^q \neq v \hat{Y}_q$, see eq. (B.1) — the approximate identification of the angles and phase in $\vec{\theta}_{\hat{V}_L}$ with the ones of the original Lagrangian in the minimal weak basis, $\vec{\theta}_{\hat{V}_L} \simeq \vec{\theta}_{\hat{V}}$, is legitimate.

C Conditions for CP Invariance

The vanishing of all physical phases in a given parameterization gives sufficient conditions for CP invariance (CPI). However, that is not strictly necessary. A less restrictive set of sufficient conditions consists in imposing that, given a parameterization, the imaginary part of all rephasing invariants one can build vanish. For instance, in the case of the minimal WB of section 3.1.1, CPI is achieved in a one doublet scenario if³²

$$\begin{aligned}
I_{\alpha i} &\equiv \text{Im} \left(z_\alpha \hat{V}_{\alpha i} z_i^* \right) = \text{Im} \left(z_u P_\alpha \hat{V} P_i z_d^\dagger \right) = 0, \\
I_{i\alpha j} &\equiv \text{Im} \left(z_i \hat{V}_{\alpha i}^* \hat{V}_{\alpha j} z_j^* \right) = \text{Im} \left(z_d P_i \hat{V}^\dagger P_\alpha \hat{V} P_j z_d^\dagger \right) = 0, \\
I_{\alpha i \beta} &\equiv \text{Im} \left(z_\alpha \hat{V}_{\alpha i} \hat{V}_{\beta i}^* z_\beta^* \right) = \text{Im} \left(z_u P_\alpha \hat{V} P_i \hat{V}^\dagger P_\beta z_u^\dagger \right) = 0, \\
I_{\alpha i \beta j} &\equiv \text{Im} \left(z_\alpha \hat{V}_{\alpha i} \hat{V}_{\beta i}^* \hat{V}_{\beta j} z_j^* \right) = \text{Im} \left(z_u P_\alpha \hat{V} P_i \hat{V}^\dagger P_\beta \hat{V} P_j z_d^\dagger \right) = 0, \\
I'_{\alpha i \beta j} &\equiv \text{Im} \left(\hat{V}_{\alpha i} \hat{V}_{\beta i}^* \hat{V}_{\beta j} \hat{V}_{\alpha j}^* \right) = \text{Im} \text{Tr} \left[P_\alpha \hat{V} P_i \hat{V}^\dagger P_\beta \hat{V} P_j \hat{V}^\dagger \right] = 0,
\end{aligned} \tag{C.1}$$

where $(P_\alpha)_{ij} = \delta_{i\alpha} \delta_{j\alpha}$ and $\alpha, \beta, i, j = 1, 2, 3$.

In turn, assuming non-degeneracy, one can always write the projectors as

$$P_\alpha = \sum_{n=0,1,2} c_{n,\alpha}^u \hat{D}_u^{2n+1}, \quad P_i = \sum_{n=0,1,2} c_{n,i}^d \hat{D}_d^{2n+1}, \tag{C.2}$$

where $c_{n,\alpha}^u$ and $c_{n,i}^d$ are non-zero coefficients which depend solely on the elements of $\hat{D}_u \equiv v \hat{Y}_u$ and $\hat{D}_d \equiv v \hat{Y}_d$, respectively. Thus, we have

$$\mathcal{I}_{nm} \equiv \text{Im} \text{Tr} \left[z_u \hat{D}_u^{2n+1} \hat{V} \hat{D}_d^{2m+1} z_d^\dagger \right] = 0 \quad \Rightarrow \quad I_{\alpha i} = 0. \tag{C.3}$$

Alternatively we can write the projectors in terms of different combinations such as

$$P_\alpha = \sum_{n=1,2,3} c_{n,\alpha}^u \hat{D}_u^{2n}, \quad P_i = \sum_{n=1,2,3} c_{n,i}^d \hat{D}_d^{2n}, \tag{C.4}$$

so that using both eq. (C.2) and eq. (C.4) we have

$$\begin{aligned}
\mathcal{I}_{nmp}^d &\equiv \text{Im} \text{Tr} \left[z_d \hat{D}_d^{2n+1} \hat{V}^\dagger \hat{D}_u^{2m} \hat{V} \hat{D}_d^{2p+1} z_d^\dagger \right] = 0 \quad \Rightarrow \quad I_{i\alpha j} = 0, \\
\mathcal{I}_{nmp}^u &\equiv \text{Im} \text{Tr} \left[z_u \hat{D}_u^{2n+1} \hat{V} \hat{D}_d^{2m} \hat{V}^\dagger \hat{D}_u^{2p+1} z_u^\dagger \right] = 0 \quad \Rightarrow \quad I_{\alpha i \beta} = 0, \\
\mathcal{I}_{nmpq} &\equiv \text{Im} \text{Tr} \left[z_u \hat{D}_u^{2n+1} \hat{V} \hat{D}_d^{2m} \hat{V}^\dagger \hat{D}_u^{2p} \hat{V} \hat{D}_d^{2q+1} z_d^\dagger \right] = 0 \quad \Rightarrow \quad I_{\alpha i \beta j} = 0, \\
\mathcal{I}'_{nmpq} &\equiv \text{Im} \text{Tr} \left[\hat{D}_u^4 \hat{V} \hat{D}_d^4 \hat{V}^\dagger \hat{D}_u^2 \hat{V} \hat{D}_d^2 \hat{V}^\dagger \right] = 0 \quad \Rightarrow \quad I'_{\alpha i \beta j} = 0.
\end{aligned} \tag{C.5}$$

³²Using the fact that \hat{V} is a 3×3 unitary matrix, one can show that any other rephasing invariant can be constructed by combining these 5 types of invariants.

In order to connect this new set of conditions to the WBIs discussed in section 6.1 we note that

$$\mathcal{H}_u = \begin{pmatrix} \hat{D}_u^2 & v\hat{D}_u\mathbf{z}_u^\dagger \\ v\mathbf{z}_u\hat{D}_u & v^2\mathbf{z}_u\mathbf{z}_u^\dagger + M_Q^2 \end{pmatrix}, \quad \mathcal{H}_d = \begin{pmatrix} \hat{V}\hat{D}_d^2\hat{V}^\dagger & v\hat{V}\hat{D}_d\mathbf{z}_d^\dagger \\ v\mathbf{z}_d\hat{D}_d\hat{V}^\dagger & v^2\mathbf{z}_d\mathbf{z}_d^\dagger + M_Q^2 \end{pmatrix}, \quad (\text{C.6})$$

so that each one of the conditions in eq. (C.1) corresponds, in the same order, to the following conditions in terms of the Hermitian building blocks \mathcal{H}_u and \mathcal{H}_d ,

$$\begin{aligned} \mathcal{I}_{nm} &= \text{Im Tr} \left[P_4 \mathcal{H}_u [(\mathbb{1} - P_4) \mathcal{H}_u]^n [(\mathbb{1} - P_4) \mathcal{H}_d]^{m+1} \right] = 0, \\ \mathcal{I}_{nmp}^d &= \text{Im Tr} \left[P_4 \mathcal{H}_d [(\mathbb{1} - P_4) \mathcal{H}_d]^n [(\mathbb{1} - P_4) \mathcal{H}_u]^m [(\mathbb{1} - P_4) \mathcal{H}_d]^{p+1} \right] = 0, \\ \mathcal{I}_{nmp}^u &= \text{Im Tr} \left[P_4 \mathcal{H}_u [(\mathbb{1} - P_4) \mathcal{H}_u]^n [(\mathbb{1} - P_4) \mathcal{H}_d]^m [(\mathbb{1} - P_4) \mathcal{H}_u]^{p+1} \right] = 0, \\ \mathcal{I}_{nmpq} &= \text{Im Tr} \left[P_4 \mathcal{H}_u [(\mathbb{1} - P_4) \mathcal{H}_u]^n [(\mathbb{1} - P_4) \mathcal{H}_d]^m [(\mathbb{1} - P_4) \mathcal{H}_u]^p [(\mathbb{1} - P_4) \mathcal{H}_d]^{q+1} \right] = 0, \\ \mathcal{I}'_{nmpq} &= \text{Im Tr} \left[[(\mathbb{1} - P_4) \mathcal{H}_u]^2 [(\mathbb{1} - P_4) \mathcal{H}_d]^2 (\mathbb{1} - P_4) \mathcal{H}_u (\mathbb{1} - P_4) \mathcal{H}_d \right] = 0, \end{aligned} \quad (\text{C.7})$$

where $P_4 = \mathbb{1} - P_1 - P_2 - P_3$ and $\mathbb{1}$ is the 4×4 identity matrix.

To ultimately achieve a direct connection between these conditions and the WBI studied in the main text we must now remove the many P_4 that feature in these expressions, for which a useful property is

$$\text{Tr} (P_4 A P_4 B) = \text{Tr} (P_4 A) \text{Tr} (P_4 B). \quad (\text{C.8})$$

Using this property one can show that the form of \mathcal{I}_{nm} in eq. (C.7) can be expanded into a convoluted combination of traces that are exclusively of three types:

$$\text{Tr} (P_4 \mathcal{H}_u^a), \quad \text{Tr} (P_4 \mathcal{H}_d^a), \quad \text{Tr} (P_4 \mathcal{H}_u^a \mathcal{H}_d^b). \quad (\text{C.9})$$

Thus, since only the last type of invariant is in general CP-odd, the vanishing of \mathcal{I}_{nm} can be achieved if

$$\text{Im Tr} \left[P_4 \mathcal{H}_u^a \mathcal{H}_d^b \right] = 0, \quad (\text{C.10})$$

for all $1 \leq a \leq n+1$ and $1 \leq b \leq m+1$. Likewise, one can show that the expansion of \mathcal{I}_{nmp}^d in eq. (C.7), depends solely on four types of traces: the ones in eq. (C.9) plus $\text{Tr} (P_4 \mathcal{H}_d^a \mathcal{H}_u^b \mathcal{H}_d^c)$, meaning that

$$\text{Im Tr} \left[P_4 \mathcal{H}_d^a \mathcal{H}_u^b \mathcal{H}_d^c \right] = 0, \quad (\text{C.11})$$

for all $1 \leq a \leq n+1$, $1 \leq b \leq m$ and $1 \leq c \leq p+1$, along with the condition of eq. (C.10) leads to the vanishing of \mathcal{I}_{nmp}^d .

In the same way, one can show that the vanishing of \mathcal{I}_{nmp}^u can be achieved if the conditions of eq. (C.10) and

$$\text{Im Tr} \left[P_4 \mathcal{H}_u^a \mathcal{H}_d^b \mathcal{H}_u^c \right] = 0 \quad (\text{C.12})$$

are both true, while the vanishing of \mathcal{I}_{nmpq} is achieved if the conditions of eqs. (C.10) to (C.12) and

$$\text{Im Tr} \left[P_4 \mathcal{H}_u^a \mathcal{H}_d^b \mathcal{H}_u^c \mathcal{H}_d^d \right] = 0 \quad (\text{C.13})$$

are fulfilled. Finally, the last invariant vanishes if all conditions in eqs. (C.10) to (C.13) and

$$\text{Im Tr} \left[\mathcal{H}_u^2 \mathcal{H}_d^2 \mathcal{H}_u \mathcal{H}_d \right] = 0 \quad (\text{C.14})$$

are verified. Hence, using the fact that $H = M_Q^2 P_4$ in the minimal WB, we can conclude that the conditions

$$\begin{aligned} \text{Im Tr} \left[\mathcal{H}_u^n \mathcal{H}_d^m H \right] &= 0, \quad (n, m = 1, 2, 3), \\ \text{Im Tr} \left[\mathcal{H}_d^n \mathcal{H}_u^m \mathcal{H}_d^p H \right] &= 0, \quad (n > p, n, p = 1, 2, 3, m = 1, 2), \\ \text{Im Tr} \left[\mathcal{H}_u^n \mathcal{H}_d^m \mathcal{H}_u^p H \right] &= 0, \quad (n > p, n, p = 1, 2, 3, m = 1, 2), \\ \text{Im Tr} \left[\mathcal{H}_u^n \mathcal{H}_d^m \mathcal{H}_u^p \mathcal{H}_d^q H \right] &= 0, \quad (n > p, n, q = 1, 2, 3, m, p = 1, 2), \\ \text{Im Tr} \left[\mathcal{H}_u^2 \mathcal{H}_d^2 \mathcal{H}_u \mathcal{H}_d \right] &= 0 \end{aligned} \quad (\text{C.15})$$

necessarily imply CPI in a one-doublet model. These WBIs are the ones studied in section 6.1.1. Here we used the Cayley-Hamilton theorem to obtain a limited set of conditions. A more systematic application of this theorem, together with some Group Theory techniques can be used to give a minimal set of conditions also valid for arbitrary mass degeneracies, as recently done in Ref. [60] for the singlet case. We anticipate that this set is $\mathcal{O}(\lesssim 10)$ larger for the doublet case. Nonetheless, the set in eq. (C.15) still amounts to a very large number of conditions, and since the phase content of a one VLQ doublet model corresponds to at most 6 phases we expect the set of WBI conditions that lead to CPI to be much smaller. In fact, as we have shown in section 5.2.1, when no z_α or z_i coupling vanishes and there exist 6 physical phases, we only need 9 conditions to ensure CPI. In what follows we will analyze all of the remaining possible cases described in section 6.3 and presented in Table 3 and, for each one of them, identify a limited number of WBI conditions that can lead to CPI.

C.1 Case-by-case analysis

Here we will demonstrate the results of Table 3. To that end, we make use of specific WBs for which these proofs are more straightforward. For some cases, we make use of the minimal WB of section 3.1.1. However, at the end of this section we switch to the stepladder WB, since it is much more convenient to address the scenarios where all $z_{i/\alpha} \neq 0$ in one sector, just as it proved to be more useful when considering the case of all $z_{i/\alpha} \neq 0$ in section 5.2.1.

Note that the goal is to achieve a set of WBI conditions for CPI. Therefore, the final result is independent of the particular choice of WB.

Using the minimal WB:

We start by assume that all but one of the physical phases are contained in z_α and z_i , with the remaining one, which we denote by δ_0 , being contained in \hat{V} , which can take

the form

$$\hat{V} = \begin{pmatrix} \hat{V}_{11}^r & \hat{V}_{12}^r & \hat{V}_{13}^r \\ \hat{V}_{21}^r & \hat{V}_{22}^r & \hat{V}_{23}^r \\ \hat{V}_{31}^r & \hat{V}_{32}^r & \hat{V}_{33}^r \end{pmatrix} \quad (\text{C.16})$$

in some version of the minimal WB. Here the index r identifies an entry as real. Note that the complex entries can be parametrized in the following way

$$\hat{V}_{ij} = a_{ij} + b_{ij}e^{i\delta_0}, \quad (\text{C.17})$$

where a_{ij} and b_{ij} are real quantities.

- **One $z_i = 0$ and one $z_\alpha = 0$:**

Here we take $z_u = z_d = 0$. In this case 4 physical phases are present, δ_0 and the phases of z_t, z_s and z_b . The simplest type of invariant which is sensitive to all these phases is \mathcal{I}_{nm} of eq. (C.3) which takes the form

$$\begin{aligned} \mathcal{I}_{nm} = & \hat{m}_c^{2n+1} \left[\hat{m}_s^{2m+1} \text{Im} \left(\hat{V}_{22} z_s^* z_c \right) + \hat{m}_b^{2m+1} \text{Im} \left(\hat{V}_{23} z_b^* z_c \right) \right] \\ & + \hat{m}_t^{2n+1} \left[\hat{m}_s^{2m+1} \text{Im} \left(\hat{V}_{32} z_s^* z_t \right) + \hat{m}_b^{2m+1} \text{Im} \left(\hat{V}_{33} z_b^* z_t \right) \right], \end{aligned} \quad (\text{C.18})$$

where the 4 phases appear in 4 distinct rephasing invariant combinations. Here, the $\hat{m}_i \equiv v \hat{y}_i$ denote the eigenvalues of the \hat{D}_q matrices.

Hence, CPI can be achieved if we guarantee the simultaneous vanishing of four invariants of this type, such as

$$\mathcal{I}_{00} = \mathcal{I}_{10} = \mathcal{I}_{01} = \mathcal{I}_{11} = 0, \quad (\text{C.19})$$

provided that these conditions form a non-singular system.

Now, note that

$$\mathcal{I}_{00} = \frac{1}{M_Q^2} \text{Im Tr} [\mathcal{H}_u \mathcal{H}_d H] \quad (\text{C.20})$$

vanishes when

$$\text{Im Tr} [\mathcal{H}_u \mathcal{H}_d H] = 0. \quad (\text{C.21})$$

Assuming that is the case, we then have

$$\begin{aligned} \mathcal{I}_{10} &= \frac{1}{M_Q^2} \text{Im Tr} [\mathcal{H}_u^2 \mathcal{H}_d H], \\ \mathcal{I}_{01} &= \frac{1}{M_Q^2} \text{Im Tr} [\mathcal{H}_u \mathcal{H}_d^2 H], \end{aligned} \quad (\text{C.22})$$

and their vanishing can be achieved via

$$\text{Im Tr} [\mathcal{H}_u^2 \mathcal{H}_d H] = \text{Im Tr} [\mathcal{H}_u \mathcal{H}_d^2 H] = 0, \quad (\text{C.23})$$

which then leads to

$$\mathcal{I}_{11} = \frac{1}{M_Q^2} \text{Im Tr} [\mathcal{H}_u^2 \mathcal{H}_d^2 H] . \quad (\text{C.24})$$

Therefore, the conditions in eq. (C.19) can be achieved if

$$\text{Im Tr} [\mathcal{H}_u^{n+1} \mathcal{H}_d^{m+1} H] = 0 , \quad (\text{C.25})$$

with $(n, m) = (0, 0), (1, 0), (0, 1), (1, 1)$.

In turn, the system of linear equations in eq. (C.19) implies

$$\text{Im} \left(\hat{V}_{22} z_s^* z_c \right) = \text{Im} \left(\hat{V}_{23}^r z_b^* z_c \right) = \text{Im} \left(\hat{V}_{32} z_s^* z_t \right) = \text{Im} \left(\hat{V}_{33}^r z_b^* z_t \right) = 0 , \quad (\text{C.26})$$

assuming a non-singular system, i.e.

$$\hat{m}_c^2 \hat{m}_t^2 \hat{m}_s^2 \hat{m}_b^2 (\hat{m}_b^2 - \hat{m}_s^2)^2 (\hat{m}_t^2 - \hat{m}_c^2)^2 \neq 0 . \quad (\text{C.27})$$

Thus, in a non-degenerate scenario, we can eliminate four distinct rephasing invariant combinations of the four physical phases and, therefore achieve CPI if we require the vanishing of the four 4-block WBI of the form of eq. (C.25).

• **Two $z_{i/\alpha} = 0$ in one sector and one $z_{\alpha/i} = 0$ in the other sector :**

We assume the case with $z_u = z_d = z_s = 0$, where there are 3 phases present: δ_0 , the phase of z_t and the phase of z_b .

The simplest type of CP-odd invariants acquires the form

$$\mathcal{I}_{nm} = \hat{m}_b^{2m+1} \left[\hat{m}_c^{2n+1} \text{Im} \left(\hat{V}_{23}^r z_b^* z_c \right) + \hat{m}_t^{2n+1} \text{Im} \left(\hat{V}_{33}^r z_b^* z_t \right) \right] , \quad (\text{C.28})$$

which is sensitive to the phases of z_t and z_b . The vanishing of all invariants of this type only requires

$$\text{Im Tr} [\mathcal{H}_u \mathcal{H}_d H] = \text{Im Tr} [\mathcal{H}_u^2 \mathcal{H}_d H] = 0 , \quad (\text{C.29})$$

since it would imply $\text{Im } z_t = \text{Im } z_b = 0$, that is if the corresponding system of linear equations is non-singular, i.e.

$$\hat{m}_t \hat{m}_c \hat{m}_b^2 (\hat{m}_t^2 - \hat{m}_c^2) \neq 0 , \quad (\text{C.30})$$

which again amounts to having non-degeneracy.

The next simplest and non-vanishing type of CP-odd invariant that is sensible to the remaining phase δ_0 is now of the form³³

$$\begin{aligned} \mathcal{I}_{nmp}^u &= A_{nmp} \text{Im}(\hat{V}_{21} \hat{V}_{31}^* z_c z_t) + B_{nmp} \text{Im}(\hat{V}_{22} \hat{V}_{32}^* z_c z_t) \\ &= A'_{nmp} \text{Im}(\hat{V}_{21} \hat{V}_{31}^* z_c z_t) + B'_{nmp} \text{Im}(\hat{V}_{23}^r \hat{V}_{33}^r z_c z_t) \\ &= A'_{nmp} \text{Im}(\hat{V}_{21} \hat{V}_{31}^* z_c z_t) \\ &= A'_{nmp} \frac{z_c z_t}{V_{23}^r V_{33}^r} \text{Im } Q_{2133} , \end{aligned} \quad (\text{C.31})$$

³³Here we make use of the unitarity condition $V_{22} V_{32}^* = -V_{21} V_{31}^* - V_{23} V_{33}^*$ and the fact that V_{23} and V_{33} are real in the WB we are using.

where

$$A'_{nmp} = \hat{m}_c^{2n+1} \hat{m}_t^{2n+1} \left(\hat{m}_t^{2(p-n)} - \hat{m}_c^{2(p-n)} \right) \left(\hat{m}_d^{2m} - \hat{m}_s^{2m} \right), \quad (\text{C.32})$$

and assuming $V_{23}^r, V_{33}^r \neq 0$.³⁴

Given that, at this point, all $z_{i/\alpha}$ couplings are real, we have $\mathcal{I}_{nm} = 0$ for all n and m , so that

$$\mathcal{I}_{011}^u = \frac{1}{M_Q^2} \text{Im Tr} [\mathcal{H}_u \mathcal{H}_d \mathcal{H}_u^2 H], \quad (\text{C.33})$$

and the vanishing of this 4-block WBI, i.e.

$$\text{Im Tr} [\mathcal{H}_u \mathcal{H}_d \mathcal{H}_u^2 H] = 0, \quad (\text{C.34})$$

leads to $\mathcal{I}_{110}^u = 0$. Assuming no degeneracy we must have $A_{011} \neq 0$ so that the vanishing of this invariant ensures $\text{Im } Q_{2133} = 0$ and similarly for all other quartets, meaning there is no CPV induced by δ_0 in \hat{V} . Therefore, we need three conditions, such as

$$\text{Im Tr} [\mathcal{H}_u \mathcal{H}_d H] = \text{Im Tr} [\mathcal{H}_u^2 \mathcal{H}_d H] = \text{Im Tr} [\mathcal{H}_u \mathcal{H}_d \mathcal{H}_u^2 H] = 0, \quad (\text{C.35})$$

in order to achieve CPI.

- **Three $z_{i/\alpha} = 0$ in one sector and one $z_{\alpha/i} = 0$ in the other sector :**

Here we take $z_u = z_d = z_s = z_b = 0$ and only the phase δ_0 and the phase of z_t are physical. In this scenario we automatically have

$$\mathcal{I}_{nm} = \mathcal{I}_{nmp}^d = \mathcal{I}_{nmpq} = 0, \quad (\text{C.36})$$

while

$$\mathcal{I}_{nmp}^u = A_{nmp} \text{Im} \left(\hat{V}_{22}^* \hat{V}_{32} z_c z_t^* \right) + B_{nmp} \text{Im} \left(\hat{V}_{23}^r \hat{V}_{33}^r z_c z_t^* \right) \quad (\text{C.37})$$

is sensitive to both physical phases and where

$$A_{nmp} = \hat{m}_c^{2n+1} \hat{m}_t^{2n+1} \left(\hat{m}_t^{2(p-n)} - \hat{m}_c^{2(p-n)} \right) \left(\hat{m}_s^{2m} - \hat{m}_d^{2m} \right), \quad (\text{C.38})$$

$$B_{nmp} = \hat{m}_c^{2n+1} \hat{m}_t^{2n+1} \left(\hat{m}_t^{2(p-n)} - \hat{m}_c^{2(p-n)} \right) \left(\hat{m}_b^{2m} - \hat{m}_d^{2m} \right). \quad (\text{C.39})$$

The fact that all $\mathcal{I}_{nm} = 0$ then allows us to write

$$\mathcal{I}_{011}^u = \frac{1}{M_Q^2} \text{Im Tr} [\mathcal{H}_u \mathcal{H}_d \mathcal{H}_u^2 H], \quad (\text{C.40})$$

and the vanishing of this quantity can be achieved if

$$\text{Im Tr} [\mathcal{H}_u \mathcal{H}_d \mathcal{H}_u^2 H] = 0, \quad (\text{C.41})$$

³⁴Both these conditions are required to ensure that the final step in eq. (C.31) is valid. Still, if either one of these conditions is not fulfilled, then phase δ_0 is no longer an internal physical phase and can be rephased out of \hat{V} , meaning that $\mathcal{I}_{nmp}^u = 0$, for all n, m and p , and eq. (C.29) is sufficient to achieve CPI when there is no mass degeneracy.

which in turn leads to

$$\mathcal{I}_{021}^u = \frac{1}{M_Q^2} \text{Im Tr} [\mathcal{H}_u \mathcal{H}_d^2 \mathcal{H}_u^2 H] . \quad (\text{C.42})$$

Hence, the conditions $\mathcal{I}_{011} = \mathcal{I}_{021} = 0$ are fulfilled by having

$$\text{Im Tr} [\mathcal{H}_u \mathcal{H}_d \mathcal{H}_u^2 H] = \text{Im Tr} [\mathcal{H}_u \mathcal{H}_d^2 \mathcal{H}_u^2 H] = 0 . \quad (\text{C.43})$$

At the same time, from eq. (C.37), these conditions corresponds to the following linear system of equations

$$\begin{pmatrix} A_{011} & B_{011} \\ A_{021} & B_{021} \end{pmatrix} \begin{pmatrix} \text{Im}(\hat{V}_{22}^* \hat{V}_{32} z_c z_t^*) \\ \text{Im}(\hat{V}_{23}^r \hat{V}_{33}^r z_c z_t^*) \end{pmatrix} = 0 , \quad (\text{C.44})$$

which is non-singular if

$$\hat{m}_c^2 \hat{m}_t^2 (\hat{m}_t^2 - \hat{m}_c^2)^2 (\hat{m}_s^2 - \hat{m}_d^2) (\hat{m}_b^2 - \hat{m}_d^2) (\hat{m}_b^2 - \hat{m}_s^2) \neq 0 , \quad (\text{C.45})$$

and since we assume no mass degeneracy we conclude that the vanishing of these two invariants must imply

$$\text{Im}(\hat{V}_{22}^* \hat{V}_{32} z_c z_t^*) = \text{Im}(\hat{V}_{23}^r \hat{V}_{33}^r z_c z_t^*) = 0 , \quad (\text{C.46})$$

which leads to CPI.

- **Two $z_{i/\alpha} = 0$ in one sector and two $z_{\alpha/i} = 0$ in the other sector :**

Without loss of generality we consider the case $z_d = z_s = z_u = z_c = 0$. In this case we only have two physical phases: δ_0 and the phase of z_b . Also one can check that

$$\mathcal{I}_{nmp}^u = \mathcal{I}_{nmp}^d = 0 . \quad (\text{C.47})$$

As for invariants of the \mathcal{I}_{nm} type one has

$$\mathcal{I}_{nm} = \hat{m}_t^{2n+1} \hat{m}_b^{2m+1} \text{Im} \left(z_t z_b^* \hat{V}_{33}^r \right) . \quad (\text{C.48})$$

Now, if one 3-block WBI vanishes then z_b must be real. That is the case, for instance, if $\text{Im Tr} [\mathcal{H}_u \mathcal{H}_d H] = 0$ because, with z_t and V_{33}^r being real we have

$$\mathcal{I}_{00} = \text{Im Tr} [\mathcal{H}_u \mathcal{H}_d H] = 0 \quad \Rightarrow \quad \text{Im } z_b = 0 , \quad (\text{C.49})$$

and the only phase remaining is now δ_0 .

This phase is captured by invariants of the type

$$\mathcal{I}_{nmpq} = \hat{m}_t^{2n+1} \hat{m}_b^{2q+1} (\hat{m}_u^{2p} - \hat{m}_c^{2p}) (\hat{m}_d^{2m} - \hat{m}_s^{2m}) \text{Im} \left(\hat{V}_{11}^r \hat{V}_{13}^r \hat{V}_{31} z_b^* z_t \right) \propto \sin \delta_0 \quad (\text{C.50})$$

which are proportional $\sin \delta_0$ due to now z_b being real along with z_t , V_{11}^r and V_{13}^r . At this point we have

$$\mathcal{I}_{nm} = \mathcal{I}_{nmp}^u = \mathcal{I}_{nmp}^d = 0 , \quad (\text{C.51})$$

so that

$$\mathcal{I}_{0110} = \frac{1}{M_Q^2} \text{Im Tr} [\mathcal{H}_u \mathcal{H}_d^2 \mathcal{H}_u^2 \mathcal{H}_d H] . \quad (\text{C.52})$$

Thus, the vanishing of the remaining phase can be ensured if this 5-block WBI vanishes, i.e.

$$\mathcal{I}_{0110} = \frac{1}{M_Q^2} \text{Im} [\mathcal{H}_u \mathcal{H}_d \mathcal{H}_u \mathcal{H}_d H] = 0 . \quad (\text{C.53})$$

With no mass degeneracy we then must have $\sin \delta_0 = 0$. Therefore, we can conclude that

$$\text{Im Tr} [\mathcal{H}_u \mathcal{H}_d H] = \text{Im Tr} [\mathcal{H}_u \mathcal{H}_d \mathcal{H}_u \mathcal{H}_d H] = 0 \quad (\text{C.54})$$

ensures CPI.

Using the stepladder WB:

For the remaining two cases we drop the minimal WB of section 3.1.1 and instead focus on the stepladder WB. Recall from the discussion in section 3.1.2, that there is a direct connection between the number of $z_{i/\alpha}$ couplings vanishing in the former WB and the number of off-diagonal r_q couplings vanishing in the latter.

- **One** $z_{i/\alpha} = 0$:

Consider now the case of $z_d = 0$. To analyze this case we switch to the stepladder WB in the $r_4^d = 0$ limit. Here we have 5 physical phases, which we choose to parametrize \tilde{V} , making all r_q couplings real, just like we did in section 5.2.1.

Now, using eqs. (5.36) to (5.40), we can compute seven of the CP-odd invariants $\text{Im}(J_{ij})$, while the two remaining ones are identically zero ($J_{13}, J_{33} \propto r_4^d = 0$). In this way we can obtain the imaginary parts of all entries of \tilde{V} , except that of \tilde{V}_{31} (\tilde{V}_{11} can be taken as real). However, using the fact that \tilde{V} is unitary one can always extract the value of $\text{Im} \tilde{V}_{31}$ from simply having computed all the seven non-zero $\text{Im}(J_{ij})$.

In fact, even in an extreme scenario where all seven WBIs vanish because \tilde{V} has a form such as

$$\tilde{V} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ e^{i\alpha} & 0 & 0 \end{pmatrix} , \quad (\text{C.55})$$

making it is impossible to determine the complex phase α via unitarity relations, there is still no CP violation due to $r_4^d = 0$, since the phase α can be eliminated by rephasing the RH down-type quark fields.

Hence, CPI is achieved if seven 3-block WBIs vanish. For instance,

$$\text{Im Tr} [\mathcal{H}_u^n \mathcal{H}_d^m H] = 0 , \quad (\text{C.56})$$

with $(n, m) = (1, 1), (1, 2), (2, 1), (2, 2), (2, 3), (3, 1), (3, 2)$, ensures CPI.

• **Two $z_{i/\alpha} = 0$ in the same sector :**

Consider the limit of $r_2^d = 0$, which is analogous to having two $z_i = 0$ in the down sector. In this scenario we can perform rephasing of the quark fields that lead to the following form for the matrix \tilde{V}

$$\tilde{V} = \begin{pmatrix} \tilde{V}_{11}^r & \tilde{V}_{12}^r & \tilde{V}_{13}^r \\ \tilde{V}_{21} & \tilde{V}_{22} & \tilde{V}_{23} \\ \tilde{V}_{31} & \tilde{V}_{32} & \tilde{V}_{33} \end{pmatrix}, \quad (\text{C.57})$$

where the r -label identifies real entries. Additionally, we can identify $r_5^d = m_d$ and $r_3^d = m_s$.

In that case we can compute three of the CP-odd invariants in eqs. (5.36) to (5.40), J_{11} , J_{21} and J_{31} , while the rest are trivially zero in this limit. These two will vanish if

$$\text{Im Tr} [\mathcal{H}_u \mathcal{H}_d H] = \text{Im Tr} [\mathcal{H}_u^2 \mathcal{H}_d H] = \text{Im Tr} [\mathcal{H}_u^3 \mathcal{H}_d H] = 0, \quad (\text{C.58})$$

implying $\text{Im } \tilde{V}_{i3} = 0$, for $i = 1, 2, 3$. At this point we have

$$\tilde{V} = \begin{pmatrix} \tilde{V}_{11}^r & \tilde{V}_{12}^r & \tilde{V}_{13}^r \\ \tilde{V}_{21} & \tilde{V}_{22} & \tilde{V}_{23}^r \\ \tilde{V}_{31} & \tilde{V}_{32} & \tilde{V}_{33}^r \end{pmatrix}, \quad (\text{C.59})$$

and the presence of CPV hinges on having

$$\text{Im} \left(\tilde{V}_{21} \tilde{V}_{32} \tilde{V}_{22}^* \tilde{V}_{31}^* \right) = -\text{Im} \left(\tilde{V}_{21} \tilde{V}_{31}^* \tilde{V}_{23}^r \tilde{V}_{33}^r \right) = -\text{Im} \left(\tilde{V}_{21} \tilde{V}_{31}^* \right) \tilde{V}_{23}^r \tilde{V}_{33}^r \neq 0. \quad (\text{C.60})$$

Assuming the conditions in eq. (C.58) are verified, we then have

$$\text{Im Tr} [\mathcal{H}_u \mathcal{H}_d \mathcal{H}_u^2 H] = (r_0^u r_1^u)^2 r_2^u r_3^u (m_d^2 - m_s^2) \text{Im} \left(\tilde{V}_{31} \tilde{V}_{32} \right). \quad (\text{C.61})$$

If there is no mass degeneracy, we then have

$$\text{Im Tr} [\mathcal{H}_u \mathcal{H}_d \mathcal{H}_u H] = 0, \quad (\text{C.62})$$

so that $\text{Im} \left(\tilde{V}_{31} \tilde{V}_{32} \right) = 0$ which from eq. (C.60) implies CPI. Hence CPI is achieved if

$$\text{Im Tr} [\mathcal{H}_u \mathcal{H}_d H] = \text{Im Tr} [\mathcal{H}_u^2 \mathcal{H}_d H] = \text{Im Tr} [\mathcal{H}_u^3 \mathcal{H}_d H] = \text{Im Tr} [\mathcal{H}_u \mathcal{H}_d \mathcal{H}_u^2 H] = 0. \quad (\text{C.63})$$

• **Three $z_{i/\alpha} = 0$ in one sector :**

Now we choose to work the limit of $r_0^d = 0$, which is analogous to having all down-type couplings $z_i = 0$. Here we are interested in computing 4-block CP-odd WBIs, i.e.

$$J_{nmp} \equiv \frac{1}{M_Q^{2(1+n+m+p)}} \text{Im Tr} [\mathcal{H}_u^n \mathcal{H}_d^m \mathcal{H}_u^p H], \quad (\text{C.64})$$

with $n \neq p$, as all other types vanish. Note that in this limit we will have

$$\mathcal{H}_d = \begin{pmatrix} \tilde{V} & \\ & 1 \end{pmatrix} \text{diag}(m_d, m_s, m_b, M_{B'}) \begin{pmatrix} \tilde{V}^\dagger & \\ & 1 \end{pmatrix}, \quad (\text{C.65})$$

where in this scenario we have $M_{B'} = M_Q$.

Let us now consider the simplest invariants, given by

$$J_{1m2} = \frac{(r_0^u r_1^u)^2 r_2^u r_3^u}{M_Q^{2m}} \left\{ (m_s^{2m} - m_d^{2m}) \text{Im} \left(\tilde{V}_{32} \tilde{V}_{22}^* \right) + (m_b^{2m} - m_d^{2m}) \text{Im} \left(\tilde{V}_{33} \tilde{V}_{23}^* \right) \right\}. \quad (\text{C.66})$$

If two of these vanish, let us say $J_{112} = J_{122} = 0$, we will have the following system

$$\begin{pmatrix} m_s^2 - m_d^2 & m_b^2 - m_d^2 \\ m_s^4 - m_d^4 & m_b^4 - m_d^4 \end{pmatrix} \begin{pmatrix} \text{Im} \left(\tilde{V}_{32} \tilde{V}_{22}^* \right) \\ \text{Im} \left(\tilde{V}_{33} \tilde{V}_{23}^* \right) \end{pmatrix} = 0, \quad (\text{C.67})$$

which is non-singular for non-degenerate masses, in which case we must have

$$\text{Im} \left(\tilde{V}_{3i} \tilde{V}_{2i}^* \right) = 0 \quad (\text{C.68})$$

and $J_{1m2} = 0$ for all values of m .

Now considering the next simplest WBI of the form

$$J_{1m3} = \frac{(r_0^u r_1^u)^2 r_2^u r_3^u r_4^u r_5^u}{M_Q^{2m}} \left\{ (m_s^{2m} - m_d^{2m}) \text{Im} \left(\tilde{V}_{32} \tilde{V}_{12}^* \right) + (m_b^{2m} - m_d^{2m}) \text{Im} \left(\tilde{V}_{33} \tilde{V}_{13}^* \right) \right\}, \quad (\text{C.69})$$

where we assume not only that all J_{1m2} computed above vanish, but also no mass degeneracy is present. Thus, if for instance $J_{113} = J_{123} = 0$, then we must have

$$\text{Im} \left(\tilde{V}_{3i} \tilde{V}_{1i}^* \right) = 0. \quad (\text{C.70})$$

Finally, we may consider

$$J_{2m3} = \frac{(r_0^u r_1^u r_2^u r_3^u)^2 r_4^u r_5^u}{M_Q^{2m}} \left\{ (m_s^{2m} - m_d^{2m}) \text{Im} \left(\tilde{V}_{22} \tilde{V}_{12}^* \right) + (m_b^{2m} - m_d^{2m}) \text{Im} \left(\tilde{V}_{23} \tilde{V}_{13}^* \right) \right\}, \quad (\text{C.71})$$

and having for instance $J_{213} = J_{223} = 0$ will imply

$$\text{Im} \left(\tilde{V}_{2i} \tilde{V}_{1i}^* \right) = 0. \quad (\text{C.72})$$

At this point it is safe to say that all quartets vanish and there is no internal phase in \tilde{V} . Any other phases present in \tilde{V} have also to be common to all non-zero elements in a given column of this matrix, so that, given that \mathcal{M}_d is diagonal, they can be factored out via a rephasing of the RH down-type quark fields.

Hence the vanishing of six 4-block WBIs is sufficient (although perhaps not minimal) to achieve CPI.

References

- [1] F. Gursev, P. Ramond and P. Sikivie, *A Universal Gauge Theory Model Based on E_6* , *Phys. Lett. B* **60** (1976) 177.
- [2] Y. Achiman and B. Stech, *Quark Lepton Symmetry and Mass Scales in an E_6 Unified Gauge Model*, *Phys. Lett. B* **77** (1978) 389.
- [3] Z. G. Berezhiani and G. R. Dvali, *Possible solution of the hierarchy problem in supersymmetrical grand unification theories*, *Bull. Lebedev Phys. Inst.* **5** (1989) 55.
- [4] R. Barbieri, G. R. Dvali, A. Strumia, Z. Berezhiani and L. J. Hall, *Flavor in supersymmetric grand unification: A Democratic approach*, *Nucl. Phys. B* **432** (1994) 49 [[hep-ph/9405428](#)].
- [5] Z. Berezhiani, *SUSY $SU(6)$ GIFT for doublet-triplet splitting and fermion masses*, *Phys. Lett. B* **355** (1995) 481 [[hep-ph/9503366](#)].
- [6] L. Randall and R. Sundrum, *A Large mass hierarchy from a small extra dimension*, *Phys. Rev. Lett.* **83** (1999) 3370 [[hep-ph/9905221](#)].
- [7] N. Arkani-Hamed, A. G. Cohen and H. Georgi, *Electroweak symmetry breaking from dimensional deconstruction*, *Phys. Lett. B* **513** (2001) 232 [[hep-ph/0105239](#)].
- [8] N. Arkani-Hamed, A. G. Cohen, T. Gregoire and J. G. Wacker, *Phenomenology of electroweak symmetry breaking from theory space*, *JHEP* **08** (2002) 020 [[hep-ph/0202089](#)].
- [9] M. Perelstein, M. E. Peskin and A. Pierce, *Top quarks and electroweak symmetry breaking in little Higgs models*, *Phys. Rev. D* **69** (2004) 075002 [[hep-ph/0310039](#)].
- [10] T. Han, H. E. Logan, B. McElrath and L.-T. Wang, *Phenomenology of the little Higgs model*, *Phys. Rev. D* **67** (2003) 095004 [[hep-ph/0301040](#)].
- [11] S. Fajfer, A. Greljo, J. F. Kamenik and I. Mustac, *Light Higgs and Vector-like Quarks without Prejudice*, *JHEP* **07** (2013) 155 [[1304.4219](#)].
- [12] Z. G. Berezhiani, *The Weak Mixing Angles in Gauge Models with Horizontal Symmetry: A New Approach to Quark and Lepton Masses*, *Phys. Lett. B* **129** (1983) 99.
- [13] S. Dimopoulos, *Natural Generation of Fermion Masses*, *Phys. Lett. B* **129** (1983) 417.
- [14] Z. G. Berezhiani, *Horizontal Symmetry and Quark-Lepton Mass Spectrum: The $SU(5) \otimes SU(3)_H$ Model*, *Phys. Lett. B* **150** (1985) 177.

- [15] Z. G. Berezhiani and R. Rattazzi, *Universal seesaw and radiative quark mass hierarchy*, *Phys. Lett. B* **279** (1992) 124.
- [16] Z. G. Berezhiani and R. Rattazzi, *Inverse hierarchy approach to fermion masses*, *Nucl. Phys. B* **407** (1993) 249 [hep-ph/9212245].
- [17] Z. Berezhiani and A. Rossi, *Predictive grand unified textures for quark and neutrino masses and mixings*, *Nucl. Phys. B* **594** (2001) 113 [hep-ph/0003084].
- [18] Z. G. Berezhiani and J. L. Chkareuli, *Low-energy horizontal symmetry of $SU(3)_H \times U(1)_H$ and B anti- B oscillation. (In Russian)*, *Sov. J. Nucl. Phys.* **52** (1990) 383.
- [19] Z. Berezhiani, *Unified picture of the particle and sparticle masses in SUSY GUT*, *Phys. Lett. B* **417** (1998) 287 [hep-ph/9609342].
- [20] A. Anselm and Z. Berezhiani, *Weak mixing angles as dynamical degrees of freedom*, *Nucl. Phys. B* **484** (1997) 97 [hep-ph/9605400].
- [21] Z. Berezhiani and A. Rossi, *Flavor structure, flavor symmetry and supersymmetry*, *Nucl. Phys. B Proc. Suppl.* **101** (2001) 410 [hep-ph/0107054].
- [22] J. E. Kim, *Weak Interaction Singlet and Strong CP Invariance*, *Phys. Rev. Lett.* **43** (1979) 103.
- [23] Z. G. Berezhiani and M. Y. Khlopov, *Cosmology of Spontaneously Broken Gauge Family Symmetry*, *Z. Phys. C* **49** (1991) 73.
- [24] A. E. Nelson, *Naturally Weak CP Violation*, *Phys. Lett. B* **136** (1984) 387.
- [25] S. M. Barr, *Solving the Strong CP Problem Without the Peccei-Quinn Symmetry*, *Phys. Rev. Lett.* **53** (1984) 329.
- [26] K. S. Babu and R. N. Mohapatra, *A Solution to the Strong CP Problem Without an Axion*, *Phys. Rev. D* **41** (1990) 1286.
- [27] Z. G. Berezhiani, *On the possibility of a solution to the strong CP problem without axion in a $SU(3)_H$ family symmetry model*, *Mod. Phys. Lett. A* **6** (1991) 2437.
- [28] Z. G. Berezhiani, R. N. Mohapatra and G. Senjanovic, *Planck scale physics and solutions to the strong CP problem without axion*, *Phys. Rev. D* **47** (1993) 5565 [hep-ph/9212318].
- [29] G. C. Branco and L. Lavoura, *On the Addition of Vector Like Quarks to the Standard Model*, *Nucl. Phys. B* **278** (1986) 738.
- [30] F. del Aguila, L. Ametller, G. L. Kane and J. Vidal, *Vector Like Fermion and Standard Higgs Production at Hadron Colliders*, *Nucl. Phys. B* **334** (1990) 1.

- [31] Y. Nir and D. J. Silverman, *Z Mediated Flavor Changing Neutral Currents and Their Implications for CP Asymmetries in D^0 Decays*, *Phys. Rev. D* **42** (1990) 1477.
- [32] F. del Aguila, M. Perez-Victoria and J. Santiago, *Observable contributions of new exotic quarks to quark mixing*, *JHEP* **09** (2000) 011 [[hep-ph/0007316](#)].
- [33] G. Barenboim, F. J. Botella and O. Vives, *Constraining Models with Vector-Like Fermions from FCNC in K and B Physics*, *Nucl. Phys. B* **613** (2001) 285 [[hep-ph/0105306](#)].
- [34] G. Cacciapaglia, A. Deandrea, D. Harada and Y. Okada, *Bounds and Decays of New Heavy Vector-like Top Partners*, *JHEP* **11** (2010) 159 [[1007.2933](#)].
- [35] F. J. Botella, G. C. Branco and M. Nebot, *The Hunt for New Physics in the Flavour Sector with up vector-like quarks*, *JHEP* **12** (2012) 040 [[1207.4440](#)].
- [36] K. Ishiwata, Z. Ligeti and M. B. Wise, *New Vector-Like Fermions and Flavor Physics*, *JHEP* **10** (2015) 027 [[1506.03484](#)].
- [37] W. Wang, Z.-H. Xiong and X.-Y. Zhao, *General scan in flavor parameter space in models with vector quark doublets and an enhancement in the $B \rightarrow X_s \gamma$ process*, *Chin. Phys. C* **40** (2016) 093102 [[1603.05756](#)].
- [38] A. Biekötter, J. L. Hewett, J. S. Kim, M. Krämer, T. G. Rizzo, K. Rolbiecki et al., *Complementarity of Resonant Scalar, Vector-Like Quark and Superpartner Searches in Elucidating New Phenomena*, *Int. J. Mod. Phys. A* **32** (2017) 1750032 [[1608.01312](#)].
- [39] C. Bobeth, A. J. Buras, A. Celis and M. Jung, *Patterns of Flavour Violation in Models with Vector-Like Quarks*, *JHEP* **04** (2017) 079 [[1609.04783](#)].
- [40] F. J. Botella, G. C. Branco, M. Nebot, M. N. Rebelo and J. I. Silva-Marcos, *Vector-like Quarks at the Origin of Light Quark Masses and Mixing*, *Eur. Phys. J. C* **77** (2017) 408 [[1610.03018](#)].
- [41] G. Cacciapaglia, A. Deandrea, N. Gaur, D. Harada, Y. Okada and L. Panizzi, *The LHC potential of Vector-like quark doublets*, *JHEP* **11** (2018) 055 [[1806.01024](#)].
- [42] B. Belfatto, R. Beradze and Z. Berezhiani, *The CKM unitarity problem: A trace of new physics at the TeV scale?*, *Eur. Phys. J. C* **80** (2020) 149 [[1906.02714](#)].
- [43] G. C. Branco, J. T. Penedo, P. M. F. Pereira, M. N. Rebelo and J. I. Silva-Marcos, *Addressing the CKM unitarity problem with a vector-like up quark*, *JHEP* **07** (2021) 099 [[2103.13409](#)].

- [44] F. J. Botella, G. C. Branco, M. N. Rebelo, J. I. Silva-Marcos and J. F. Bastos, *Decays of the heavy top and new insights on ϵ_K in a one-VLQ minimal solution to the CKM unitarity problem*, *Eur. Phys. J. C* **82** (2022) 360 [2111.15401], [Erratum: *Eur.Phys.J.C* 82, 423 (2022)].
- [45] B. Belfatto and S. Trifinopoulos, *Cabibbo angle anomalies and oblique corrections: The remarkable role of the vectorlike quark doublet*, *Phys. Rev. D* **108** (2023) 035022 [2302.14097].
- [46] J. M. Alves, G. C. Branco, A. L. Cherchiglia, C. C. Nishi, J. T. Penedo, P. M. F. Pereira et al., *Vector-like singlet quarks: A roadmap*, *Phys. Rept.* **1057** (2024) 1 [2304.10561].
- [47] F. del Aguila, M. Perez-Victoria and J. Santiago, *Effective description of quark mixing*, *Phys. Lett. B* **492** (2000) 98 [hep-ph/0007160].
- [48] J. A. Aguilar-Saavedra, R. Benbrik, S. Heinemeyer and M. Pérez-Victoria, *Handbook of vectorlike quarks: Mixing and single production*, *Phys. Rev. D* **88** (2013) 094010 [1306.0572].
- [49] B. Belfatto and Z. Berezhiani, *Are the CKM anomalies induced by vector-like quarks? Limits from flavor changing and Standard Model precision tests*, *JHEP* **10** (2021) 079 [2103.05549].
- [50] A. Crivellin, M. Kirk, T. Kitahara and F. Mescia, *Global fit of modified quark couplings to EW gauge bosons and vector-like quarks in light of the Cabibbo angle anomaly*, *JHEP* **03** (2023) 234 [2212.06862].
- [51] V. Cirigliano, W. Dekens, J. de Vries, E. Mereghetti and T. Tong, *Anomalies in global SMEFT analyses. A case study of first-row CKM unitarity*, *JHEP* **03** (2024) 033 [2311.00021].
- [52] J. Bernabeu, G. C. Branco and M. Gronau, *CP Restrictions on Quark Mass Matrices*, *Phys. Lett. B* **169** (1986) 243.
- [53] M. Gronau, A. Kfir and R. Loewy, *Basis Independent Tests of CP Violation in Fermion Mass Matrices*, *Phys. Rev. Lett.* **56** (1986) 1538.
- [54] M. Olechowski and S. Pokorski, *Some Useful Invariants of Quark Mass Matrices*, *Phys. Lett. B* **231** (1989) 165.
- [55] F. del Aguila and J. A. Aguilar-Saavedra, *Invariant formulation of CP violation for four quark families*, *Phys. Lett. B* **386** (1996) 241 [hep-ph/9605418].
- [56] F. del Aguila, J. A. Aguilar-Saavedra and G. C. Branco, *CP violation from new quarks in the chiral limit*, *Nucl. Phys. B* **510** (1998) 39 [hep-ph/9703410].
- [57] G. C. Branco and J. I. Silva-Marcos, *Invariants, Alignment and the Pattern of Fermion Masses and Mixing*, *Phys. Lett. B* **715** (2012) 315 [1112.1631].

- [58] F. Albergaria, G. C. Branco, J. F. Bastos and J. I. Silva-Marcos, *CP-odd and CP-even weak-basis invariants in the presence of vector-like quarks*, *J. Phys. G* **50** (2023) 055001 [2210.14248].
- [59] M. P. Bento, J. P. Silva and A. Trautner, *The basis invariant flavor puzzle*, *JHEP* **01** (2024) 024 [2308.00019].
- [60] E. L. F. de Lima and C. C. Nishi, *Flavor invariants for the SM with one singlet vector-like quark*, *JHEP* **11** (2024) 157 [2408.10325].
- [61] G. C. Branco, L. Lavoura and J. P. Silva, *CP Violation*, vol. 103. Oxford University Press, 1999.
- [62] C.-Y. Chen, S. Dawson and Y. Zhang, *Higgs CP Violation from Vectorlike Quarks*, *Phys. Rev. D* **92** (2015) 075026 [1507.07020].
- [63] A. Santamaria, *Masses, mixings, Yukawa couplings and their symmetries*, *Phys. Lett. B* **305** (1993) 90 [hep-ph/9302301].
- [64] J. Berger and Y. Grossman, *Parameter counting in models with global symmetries*, *Phys. Lett. B* **675** (2009) 365 [0811.1019].
- [65] C. Jarlskog, *Commutator of the Quark Mass Matrices in the Standard Electroweak Model and a Measure of Maximal CP Nonconservation*, *Phys. Rev. Lett.* **55** (1985) 1039.
- [66] C. Jarlskog, *A Basis Independent Formulation of the Connection Between Quark Mass Matrices, CP Violation and Experiment*, *Z. Phys. C* **29** (1985) 491.
- [67] PARTICLE DATA GROUP collaboration, R. L. Workman et al., *Review of Particle Physics*, *PTEP* **2022** (2022) 083C01.
- [68] CMS collaboration, A. Tumasyan et al., *Search for a heavy resonance decaying into a top quark and a W boson in the lepton+jets final state at $\sqrt{s} = 13$ TeV*, *JHEP* **04** (2022) 048 [2111.10216].
- [69] L.-L. Chau and W.-Y. Keung, *Comments on the Parametrization of the Kobayashi-Maskawa Matrix*, *Phys. Rev. Lett.* **53** (1984) 1802.
- [70] F. J. Botella and L.-L. Chau, *Anticipating the Higher Generations of Quarks from Rephasing Invariance of the Mixing Matrix*, *Phys. Lett. B* **168** (1986) 97.
- [71] J. de Blas, J. C. Criado, M. Perez-Victoria and J. Santiago, *Effective description of general extensions of the Standard Model: the complete tree-level dictionary*, *JHEP* **03** (2018) 109 [1711.10391].
- [72] A. Crivellin, M. Kirk, T. Kitahara and F. Mescia, *Large $t \rightarrow cZ$ as a sign of vectorlike quarks in light of the W mass*, *Phys. Rev. D* **106** (2022) L031704 [2204.05962].

- [73] L. Allwicher et al., *Computing tools for effective field theories: SMEFT-Tools 2022 Workshop Report, 14–16th September 2022, Zürich*, *Eur. Phys. J. C* **84** (2024) 170 [2307.08745].
- [74] A. Carmona, A. Lazopoulos, P. Olgoso and J. Santiago, *Matchmakereft: automated tree-level and one-loop matching*, *SciPost Phys.* **12** (2022) 198 [2112.10787].
- [75] J. Fuentes-Martín, M. König, J. Pagès, A. E. Thomsen and F. Wilsch, *A proof of concept for matchete: an automated tool for matching effective theories*, *Eur. Phys. J. C* **83** (2023) 662 [2212.04510].
- [76] E. E. Jenkins, A. V. Manohar and P. Stoffer, *Low-Energy Effective Field Theory below the Electroweak Scale: Operators and Matching*, *JHEP* **03** (2018) 016 [1709.04486].
- [77] S. L. Glashow, J. Iliopoulos and L. Maiani, *Weak Interactions with Lepton-Hadron Symmetry*, *Phys. Rev. D* **2** (1970) 1285.
- [78] S. L. Glashow and S. Weinberg, *Natural Conservation Laws for Neutral Currents*, *Phys. Rev. D* **15** (1977) 1958.
- [79] E. A. Paschos, *Diagonal Neutral Currents*, *Phys. Rev. D* **15** (1977) 1966.
- [80] C. Y. Seng, M. Gorchtein and M. J. Ramsey-Musolf, *Dispersive evaluation of the inner radiative correction in neutron and nuclear β decay*, *Phys. Rev. D* **100** (2019) 013001 [1812.03352].
- [81] C.-Y. Seng, M. Gorchtein, H. H. Patel and M. J. Ramsey-Musolf, *Reduced Hadronic Uncertainty in the Determination of V_{ud}* , *Phys. Rev. Lett.* **121** (2018) 241804 [1807.10197].
- [82] J. C. Hardy and I. S. Towner, *Superallowed $0^+ \rightarrow 0^+$ nuclear β decays: 2020 critical survey, with implications for V_{ud} and CKM unitarity*, *Phys. Rev. C* **102** (2020) 045501.
- [83] M. Gorchtein, *γW Box Inside Out: Nuclear Polarizabilities Distort the Beta Decay Spectrum*, *Phys. Rev. Lett.* **123** (2019) 042503 [1812.04229].
- [84] UCN τ collaboration, F. M. Gonzalez et al., *Improved neutron lifetime measurement with UCN τ* , *Phys. Rev. Lett.* **127** (2021) 162501 [2106.10375].
- [85] V. F. Ezhov et al., *Measurement of the neutron lifetime with ultra-cold neutrons stored in a magneto-gravitational trap*, *JETP Lett.* **107** (2018) 671 [1412.7434].
- [86] R. W. Pattie, Jr. et al., *Measurement of the neutron lifetime using a magneto-gravitational trap and in situ detection*, *Science* **360** (2018) 627 [1707.01817].

- [87] A. P. Serebrov et al., *Neutron lifetime measurements with a large gravitational trap for ultracold neutrons*, *Phys. Rev. C* **97** (2018) 055503 [1712.05663].
- [88] S. Arzumanov, L. Bondarenko, S. Chernyavsky, P. Geltenbort, V. Morozov, V. V. Nesvizhevsky et al., *A measurement of the neutron lifetime using the method of storage of ultracold neutrons and detection of inelastically up-scattered neutrons*, *Phys. Lett. B* **745** (2015) 79.
- [89] A. Steyerl, J. M. Pendlebury, C. Kaufman, S. S. Malik and A. M. Desai, *Quasielastic scattering in the interaction of ultracold neutrons with a liquid wall and application in a reanalysis of the Mambo I neutron-lifetime experiment*, *Phys. Rev. C* **85** (2012) 065503.
- [90] A. Pichlmaier, V. Varlamov, K. Schreckenbach and P. Geltenbort, *Neutron lifetime measurement with the UCN trap-in-trap MAMBO II*, *Phys. Lett. B* **693** (2010) 221.
- [91] A. Serebrov et al., *Measurement of the neutron lifetime using a gravitational trap and a low-temperature Fomblin coating*, *Phys. Lett. B* **605** (2005) 72 [nucl-ex/0408009].
- [92] D. Mund, B. Maerkisch, M. Deissenroth, J. Krempel, M. Schumann, H. Abele et al., *Determination of the Weak Axial Vector Coupling from a Measurement of the Beta-Asymmetry Parameter A in Neutron Beta Decay*, *Phys. Rev. Lett.* **110** (2013) 172502 [1204.0013].
- [93] UCNA collaboration, M. A. P. Brown et al., *New result for the neutron β -asymmetry parameter A_0 from UCNA*, *Phys. Rev. C* **97** (2018) 035505 [1712.00884].
- [94] B. Märkisch et al., *Measurement of the Weak Axial-Vector Coupling Constant in the Decay of Free Neutrons Using a Pulsed Cold Neutron Beam*, *Phys. Rev. Lett.* **122** (2019) 242501 [1812.04666].
- [95] FLAVOUR LATTICE AVERAGING GROUP (FLAG) collaboration, Y. Aoki et al., *FLAG Review 2021*, *Eur. Phys. J. C* **82** (2022) 869 [2111.09849].
- [96] FLAVOUR LATTICE AVERAGING GROUP (FLAG) collaboration, Y. Aoki et al., *FLAG Review 2024*, 2411.04268.
- [97] C.-Y. Seng, D. Galviz, W. J. Marciano and U.-G. Meißner, *Update on $|V_{us}|$ and $|V_{us}/V_{ud}|$ from semileptonic kaon and pion decays*, *Phys. Rev. D* **105** (2022) 013005 [2107.14708].
- [98] C.-Y. Seng, D. Galviz, M. Gorchtein and U.-G. Meißner, *Complete theory of radiative corrections to $K_{\ell 3}$ decays and the V_{us} update*, *JHEP* **07** (2022) 071 [2203.05217].

- [99] M. Moulson, *Experimental determination of V_{us} from kaon decays*, *PoS CKM2016* (2017) 033 [1704.04104].
- [100] W. J. Marciano, *Precise determination of $|V_{us}|$ from lattice calculations of pseudoscalar decay constants*, *Phys. Rev. Lett.* **93** (2004) 231803 [hep-ph/0402299].
- [101] V. Cirigliano and H. Neufeld, *A note on isospin violation in $P_{\ell 2(\gamma)}$ decays*, *Phys. Lett. B* **700** (2011) 7 [1102.0563].
- [102] M. Di Carlo, D. Giusti, V. Lubicz, G. Martinelli, C. T. Sachrajda, F. Sanfilippo et al., *Light-meson leptonic decay rates in lattice QCD+QED*, *Phys. Rev. D* **100** (2019) 034514 [1904.08731].
- [103] P. Boyle et al., *Isospin-breaking corrections to light-meson leptonic decays from lattice simulations at physical quark masses*, *JHEP* **02** (2023) 242 [2211.12865].
- [104] Y. Grossman, E. Passemar and S. Schacht, *On the Statistical Treatment of the Cabibbo Angle Anomaly*, *JHEP* **07** (2020) 068 [1911.07821].
- [105] A. M. Coutinho, A. Crivellin and C. A. Manzari, *Global Fit to Modified Neutrino Couplings and the Cabibbo-Angle Anomaly*, *Phys. Rev. Lett.* **125** (2020) 071802 [1912.08823].
- [106] K. Cheung, W.-Y. Keung, C.-T. Lu and P.-Y. Tseng, *Vector-like Quark Interpretation for the CKM Unitarity Violation, Excess in Higgs Signal Strength, and Bottom Quark Forward-Backward Asymmetry*, *JHEP* **05** (2020) 117 [2001.02853].
- [107] A. Crivellin and M. Hoferichter, *β Decays as Sensitive Probes of Lepton Flavor Universality*, *Phys. Rev. Lett.* **125** (2020) 111801 [2002.07184].
- [108] M. Endo and S. Mishima, *Muon $g - 2$ and CKM unitarity in extra lepton models*, *JHEP* **08** (2020) 004 [2005.03933].
- [109] B. Capdevila, A. Crivellin, C. A. Manzari and M. Montull, *Explaining $b \rightarrow s\ell^+\ell^-$ and the Cabibbo angle anomaly with a vector triplet*, *Phys. Rev. D* **103** (2021) 015032 [2005.13542].
- [110] A. Crivellin, F. Kirk, C. A. Manzari and M. Montull, *Global Electroweak Fit and Vector-Like Leptons in Light of the Cabibbo Angle Anomaly*, *JHEP* **12** (2020) 166 [2008.01113].
- [111] M. Kirk, *Cabibbo anomaly versus electroweak precision tests: An exploration of extensions of the Standard Model*, *Phys. Rev. D* **103** (2021) 035004 [2008.03261].
- [112] C. A. Manzari, A. M. Coutinho and A. Crivellin, *Modified lepton couplings and the Cabibbo-angle anomaly*, *PoS LHCP2020* (2021) 242 [2009.03877].

- [113] A. K. Alok, A. Dighe, S. Gangal and J. Kumar, *The role of non-universal Z couplings in explaining the V_{us} anomaly*, *Nucl. Phys. B* **971** (2021) 115538 [2010.12009].
- [114] A. Crivellin, C. A. Manzari, M. Alguero and J. Matias, *Combined Explanation of the $Z \rightarrow bb^-$ Forward-Backward Asymmetry, the Cabibbo Angle Anomaly, and $\tau \rightarrow \mu\nu\nu$ and $b \rightarrow sl + \ell^-$ Data*, *Phys. Rev. Lett.* **127** (2021) 011801 [2010.14504].
- [115] A. Crivellin, F. Kirk, C. A. Manzari and L. Panizzi, *Searching for lepton flavor universality violation and collider signals from a singly charged scalar singlet*, *Phys. Rev. D* **103** (2021) 073002 [2012.09845].
- [116] A. Crivellin, M. Hoferichter and C. A. Manzari, *Fermi Constant from Muon Decay Versus Electroweak Fits and Cabibbo-Kobayashi-Maskawa Unitarity*, *Phys. Rev. Lett.* **127** (2021) 071801 [2102.02825].
- [117] D. Marzocca and S. Trifinopoulos, *Minimal Explanation of Flavor Anomalies: B -Meson Decays, Muon Magnetic Moment, and the Cabibbo Angle*, *Phys. Rev. Lett.* **127** (2021) 061803 [2104.05730].
- [118] O. Fischer et al., *Unveiling hidden physics at the LHC*, *Eur. Phys. J. C* **82** (2022) 665 [2109.06065].
- [119] R. Dcruz and K. S. Babu, *Resolving W boson mass shift and CKM unitarity violation in left-right symmetric models with a universal seesaw mechanism*, *Phys. Rev. D* **108** (2023) 095011 [2212.09697].
- [120] A. Pich and A. Rodríguez-Sánchez, *$SU(3)$ analysis of four-quark operators: $K \rightarrow \pi\pi$ and vacuum matrix elements*, *JHEP* **06** (2021) 005 [2102.09308].
- [121] T. Blum et al., *Lattice determination of the $K \rightarrow (\pi\pi)_{I=2}$ Decay Amplitude A_2* , *Phys. Rev. D* **86** (2012) 074513 [1206.5142].
- [122] RBC, UKQCD collaboration, Z. Bai et al., *Standard Model Prediction for Direct CP Violation in $K \rightarrow \pi\pi$ Decay*, *Phys. Rev. Lett.* **115** (2015) 212001 [1505.07863].
- [123] RBC, UKQCD collaboration, R. Abbott et al., *Direct CP violation and the $\Delta I = 1/2$ rule in $K \rightarrow \pi\pi$ decay from the standard model*, *Phys. Rev. D* **102** (2020) 054509 [2004.09440].
- [124] S. Bertolini, A. Maiezza and F. Nesti, *$K \rightarrow \pi\pi$ hadronic matrix elements of left-right current-current operators*, *Phys. Rev. D* **88** (2013) 034014 [1305.5739].
- [125] V. Cirigliano, W. Dekens, J. de Vries and E. Mereghetti, *An ϵ' improvement from right-handed currents*, *Phys. Lett. B* **767** (2017) 1 [1612.03914].
- [126] P. Chen, H. Ke and X. Ji, *Direct CP Violation in K -decay and Minimal Left-Right Symmetry Scale*, *Phys. Lett. B* **677** (2009) 157 [0810.2576].

- [127] PARTICLE DATA GROUP collaboration, S. Navas et al., *Review of particle physics*, *Phys. Rev. D* **110** (2024) 030001.
- [128] V. Cirigliano, H. Gisbert, A. Pich and A. Rodríguez-Sánchez, *Theoretical status of ϵ'/ϵ* , *J. Phys. Conf. Ser.* **1526** (2020) 012011 [1912.04736].
- [129] V. Cirigliano, H. Gisbert, A. Pich and A. Rodríguez-Sánchez, *Isospin-violating contributions to ϵ'/ϵ* , *JHEP* **02** (2020) 032 [1911.01359].
- [130] C. Abel et al., *Measurement of the Permanent Electric Dipole Moment of the Neutron*, *Phys. Rev. Lett.* **124** (2020) 081803 [2001.11966].
- [131] B. K. Sahoo, *Improved limits on the hadronic and semihadronic CP violating parameters and role of a dark force carrier in the electric dipole moment of ^{199}Hg* , *Phys. Rev. D* **95** (2017) 013002 [1612.09371].
- [132] S. Alioli, V. Cirigliano, W. Dekens, J. de Vries and E. Mereghetti, *Right-handed charged currents in the era of the Large Hadron Collider*, *JHEP* **05** (2017) 086 [1703.04751].
- [133] V. Cirigliano, W. Dekens, J. de Vries, E. Mereghetti and T. Tong, *Beta-decay implications for the W-boson mass anomaly*, *Phys. Rev. D* **106** (2022) 075001 [2204.08440].
- [134] T. Inami and C. S. Lim, *Effects of Superheavy Quarks and Leptons in Low-Energy Weak Processes $K_L \rightarrow \mu\bar{\mu}$, $K^+ \rightarrow \pi^+\nu\bar{\nu}$ and $K_0 \leftrightarrow \bar{K}_0$* , *Prog. Theor. Phys.* **65** (1981) 297 [Erratum: *Prog.Theor.Phys.* 65, 1772 (1981)].
- [135] G. Buchalla, A. J. Buras and M. E. Lautenbacher, *Weak decays beyond leading logarithms*, *Rev. Mod. Phys.* **68** (1996) 1125 [hep-ph/9512380].
- [136] Z. Bai, N. H. Christ, T. Izubuchi, C. T. Sachrajda, A. Soni and J. Yu, *$K_L - K_S$ Mass Difference from Lattice QCD*, *Phys. Rev. Lett.* **113** (2014) 112003 [1406.0916].
- [137] Z. Bai, N. H. Christ and C. T. Sachrajda, *The $K_L - K_S$ Mass Difference*, *EPJ Web Conf.* **175** (2018) 13017.
- [138] RBC/UKQCD collaboration, N. Garron, R. J. Hudspith and A. T. Lytle, *Neutral Kaon Mixing Beyond the Standard Model with $n_f = 2 + 1$ Chiral Fermions Part 1: Bare Matrix Elements and Physical Results*, *JHEP* **11** (2016) 001 [1609.03334].
- [139] A. J. Buras, S. Jager and J. Urban, *Master formulae for $\Delta F = 2$ NLO QCD factors in the standard model and beyond*, *Nucl. Phys. B* **605** (2001) 600 [hep-ph/0102316].

- [140] C. Bobeth, A. J. Buras, A. Celis and M. Jung, *Yukawa enhancement of Z-mediated new physics in $\Delta S = 2$ and $\Delta B = 2$ processes*, *JHEP* **07** (2017) 124 [1703.04753].
- [141] M. Bona et al., *Unitarity Triangle global fits beyond the Standard Model: UTfit 2021 NP update*, *PoS EPS-HEP2021* (2022) 500.
- [142] L. S. Littenberg, *The CP Violating Decay $K_L^0 \rightarrow \pi^0 \nu \bar{\nu}$* , *Phys. Rev. D* **39** (1989) 3322.
- [143] B. Kayser, *CP violation in the K and B systems*, in *ICTP Summer School in High-energy Physics and Cosmology*, pp. 432–478, 11, 1996, hep-ph/9702264.
- [144] Y. Grossman and Y. Nir, *$K_L \rightarrow \pi^0 \nu \nu$ beyond the standard model*, *Phys. Lett. B* **398** (1997) 163 [hep-ph/9701313].
- [145] G. Buchalla and G. Isidori, *The CP conserving contribution to $K_L \rightarrow \pi^0 \nu \bar{\nu}$ in the standard model*, *Phys. Lett. B* **440** (1998) 170 [hep-ph/9806501].
- [146] A. J. Buras, D. Buttazzo, J. Girrbach-Noe and R. Knegjens, *$K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 \nu \bar{\nu}$ in the Standard Model: status and perspectives*, *JHEP* **11** (2015) 033 [1503.02693].
- [147] A. Pich, *Effective Field Theory with Nambu-Goldstone Modes*, *Les Houches Lect. Notes* **108** (2020) 137 [1804.05664].