

# An oscillating Rastall universe crossing the phantom divide line

Nasr Ahmed<sup>1,2</sup>, Anirudh Pradhan<sup>3</sup>, Archana Dixit<sup>4</sup>

<sup>1</sup> Mathematics Department, Faculty of Science, Taibah University, Saudi Arabia.

<sup>2</sup> Astronomy Department, National Research Institute of Astronomy and Geophysics, Helwan, Cairo, Egypt

<sup>3</sup> Centre for Cosmology, Astrophysics and Space Science (CCASS), GLA University, Mathura-281 406, Uttar Pradesh, India

<sup>4</sup> Department of Mathematics, Gurugram University, Gurugram-122003, Haryana, India

<sup>1,2</sup>E-mail: nasr.ahmed@nriag.sci.eg

<sup>3</sup>E-mail: pradhan.anirudh@gmail.com

<sup>4</sup>E-mail: archana.ibs.maths@gmail.com

## Abstract

A cyclic flat universe with quintom behaviour and future big rip has been presented in the framework of Rastall gravity, which is an extension of the standard  $\Lambda$ CDM model. The Hubble parameter oscillates periodically between positive and negative values from one cycle to the next. Cosmic transit has been simulated through an oscillating time-dependent deceleration parameter, and is expected to occur at approximately 8.7 Gyr. The causality is satisfied all the time except near the initial singularity and the future Big Rip singularity. The apparent horizon, entropy and other thermodynamical quantities associated to the current model have been analyzed. Energy conditions have been investigated.

*Keywords:* Cyclic universe; Modified gravity; Causality; Phantom crossing

PACS number: 04.50+h, 04.20Jb, 95.35+d, 98.80.Es

## 1 Introduction and Motivation

One of the most significant and fascinating phenomena in contemporary cosmology is the universe's accelerating expansion. Our understanding of the composition and development of the cosmos underwent a paradigm change when it was discovered in the late 1990s through the observation of Type Ia Supernovae [1, 2]. Numerous independent observations, including as large-scale structural surveys and measurements of cosmic microwave background radiation, have supported this ground-breaking discovery [3, 4, 5]. At both large scales and strong curvature, Einstein theory of General Relativity is inconsistent with the observations. Cosmic acceleration has been a basic motivation behind modified gravity theories since it can't be explained by GR without assuming dark energy [2, 6]. Examples of such modified gravity theories are  $f(R)$  gravity [7], Gauss-Bonnet gravity [8, 9] where the Gauss-Bonnet term  $\mathcal{G} = R^2 - 4R^{\mu\nu}R_{\mu\nu} + R^{\mu\nu\rho\delta}R_{\mu\nu\rho\delta}$  has been used in the action instead of the Ricci scalar  $R$ , and  $f(T)$  gravity [10] where the torsion scalar ( $T$ ) has been

used. the generalized  $f(R, T)$  gravity where  $T$  is the trace of the energy-momentum tensor. More examples include  $\kappa(R, T)$  gravity [11] and extra-dimensional gravity theories [12, 13, 14, 15].

Bouncing scenarios (also named as cyclic cosmology) have been firstly suggested as an alternative way to solve some of the problems of the standard model of cosmology such as flatness and initial singularity. The universe in the singularity-free Big Bounce emerges from a previous contracting stage [16, 17, 18, 19] (see [20] for a review) where the contraction-expansion cycle is supposed to continue forever. There have been an extensive interest in cyclic cosmological models in the literature. they have been studied in many modified gravity theories [18, 21, 22, 23, 24, 25, 19, 26]. A general cyclic model in  $f(R)$  gravity has been constructed in [27]. A unitary version of Conformal Cyclic Cosmology (CCC) has been suggested in [28]. An M-theory inspired cyclic Model with two branes has been presented in [29] where the EoS parameter satisfies  $\omega \gg 1$  during the contraction phase. A new novel cyclic theory of the universe has been presented in [30] where the scale factor grows exponentially from one cycle to another. This recent cyclic model resolves many problems and produces a nearly scale invariant spectrum of density perturbations. An interesting singularity-free oscillatory model has been discussed in [31] where the late-time evolution was probed in the context of Quasi Steady State cosmology QSSC [32, 33, 34] to alleviate the Hubble tension. The cosmic scale factor parametrization has been chosen according to the description in [32, 33, 34, 35, 36] as

$$a(t) = e^{t/P} [1 + \eta \cos(2\pi t/Q)]. \quad (1)$$

Here  $\eta$  is a dimensionless constant while  $P$  and  $Q$  have dimensions of time. In the limit when the dimensionless constant  $\eta$  goes to zero, A de Sitter-like evolution occurs. When time goes to zero, the scale factor tends to the value  $1 + \eta$  where singularity doesn't exist.

Rastall gravity is one such modification that was proposed by Peter Rastall in 1972 [37]. Due to its presentation as a modified theory of gravity with a non-conserved stress energy tensor and a peculiar non-minimal coupling between geometry and matter, Rastall gravity (RG) has emerged as a leading theory of gravity in modern times. The conservation law that demonstrated a divergence free energy momentum tensor in general relativity (GR) has been accepted; that is,  $\nabla_\nu T_{\mu\nu} = 0$ , where  $\nabla_\nu$  is the covariant derivative. Rastall, on the other hand, adopted a new and distinct conservation law that conjectured the GR concepts. In this theory, the energy momentum tensor does not a conserved quantity. Rastall theorem is defined as  $\nabla_\mu = \left(\frac{\kappa}{16\pi}\right) \nabla_\mu R$ , where  $\kappa$  is the coupling constant and  $R$  be the Ricci scalar. The second Bianchi identity in this theory remains the same i.e.,  $\nabla^\mu G_{\nu\mu} = 0$ , where  $G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu}$  be the Einstein tensor. One significant aspect of RG is that, while the mathematical portion of the theory remains invariant, modifications have only been made to the matter portion [38, 39, 40]. Das et al. [41] demonstrated that RG is equal to Einstein gravity and offered some cosmological implications in the structure of modified RG. Holographic dark energy in RG was introduced by Ghaffari et al. [42], taking into account vacuum energy, which serves as DE. They accepted the idea that the current accelerating cosmos is caused by the sum of this energy and the Rastall term. With the assumption that the universe is made up of interacting/non-interacting dark energy (DE) and dark matter (DM), Saleem and Shahnila [43] investigated the phenomenon of cosmic evolution using curved FLRW space-time bounded by an apparent horizon with a particular holographic cut-off in the framework of Rastall gravity. Saleem et al. [44] recently introduced constant-roll warm inflation, a novel method for determining the

precise inflationary solutions to the Friedman equations within the framework of Rastall theory of gravity (RTG). Using a linear parametrization of the Equation of State (EoS) in FLRW background, Singh et al. [45] examined the evolution of the universe within the framework of RTG. In the framework of RTG, Saleem and Hassan [46] studied the dynamics of warm inflation induced by vector fields and concluded that the modified theory was compatible with the 2018 Planck observational data. In various contexts, several researchers have examined the cosmological scenario within the framework of Rastall gravity [47, 48, 49, 50, 51, 52, 53, 54, 55].

Due to the ad hoc periodic parametrization of the deceleration parameter  $q$  with the correct sign flipping as  $q = m \cos kt - 1$  [56], the universe accelerates after an epoch of deceleration (for every single cycle) which agrees with recent observations. Such periodic form of  $q$  leads to the following forms for the Hubble parameter and scale factor

$$H = \frac{k}{m \sin kt + c}, \quad a = a_0 \left[ \tan\left(\frac{1}{2}kt\right) \right]^{\frac{1}{m}}, \quad (2)$$

where  $m > 0$ ,  $k > 0$  and  $k$  acts as a cosmic frequency parameter. This specific form of the deceleration parameter has firstly been introduced in [56] where a new oscillating Quintom Model has been constructed. Using the well known redshift relation  $z = \frac{1}{a} - 1$  we get

$$t = \frac{2}{k} \tan^{-1} \frac{1}{a_0^m (z+1)^m}. \quad (3)$$

Observations suggests that the signature change of  $q$  occurs at  $z = 0.64$  for  $m = 1.55$  [57, 58, 59, 60]. Since cosmic transit happens when  $q = 0$  (*i.e.*  $\ddot{a} = 0$ ), we have

$$t_{q=0} = \frac{1}{k} \cos^{-1} \frac{1}{m} \approx 8.7 \text{ Gyr} \quad \text{for } m = 1.55, \quad k = 0.1 \quad (4)$$

The scale factor (2) suffers from future Big Rip singularity where the slope of  $a(t)$  increases hugely and goes to infinity  $\frac{da}{dt} \rightarrow \infty \Rightarrow$  which eventually tearing apart the space-time fabric .

## 2 Framework of the Model

Our goal is to obtain solutions by carefully taking into consideration the matter source and the gravitational background. The basic idea behind Rastall gravity [61] is that the energy-momentum conservation in GR can't be always valid in curved space-time and, instead of  $T_{;\mu}^{\mu\nu} = 0$ , we should have

$$T_{;\mu}^{\mu\nu} = \lambda R^{,\nu}, \quad (5)$$

which leads to the generalized field equations

$$G_{\mu\nu} + K \lambda g_{\mu\nu} R = K T_{\mu\nu}. \quad (6)$$

GR is recovered for  $\lambda = 0$ .  $G_{\mu\nu}$  is the Einstein tensor and  $K$  is a coupling constant. The FRW metric given by

$$ds^2 = -dt^2 + a^2(t) \left[ \frac{dr^2}{1 - \kappa r^2} + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \right], \quad (7)$$

where  $a(t)$  is the scale factor and  $\kappa$  equals 0 for a flat universe. Applying equation (6) to the metric (7) we obtain the cosmological equations as

$$3(1 - 4K\lambda)H^2 - 6K\lambda\dot{H} + 3(1 - 2K\lambda)\frac{\kappa}{a^2} = K\rho, \quad (8)$$

$$3(1 - 4K\lambda)H^2 + 2(1 - 3K\lambda)\dot{H} + (1 - 6K\lambda)\frac{\kappa}{a^2} = -Kp. \quad (9)$$

In the current work, we will be interested only in the observationally supported flat case where  $\kappa = 0$  [62, 63, 64, 65]. The energy density  $\rho$ , cosmic pressure  $p$  and EoS parameter  $\omega$  are written as

$$\rho = -\frac{1}{K} \left( 12\lambda KH^2 + 6\lambda K\dot{H} - 3H^2 \right), \quad (10)$$

$$p = \frac{1}{K} \left( 12\lambda KH^2 + 6\lambda K\dot{H} - 3H^2 - 2\dot{H} \right), \quad (11)$$

where the dot denotes differentiation w.r.t time and the EoS parameter  $\omega = p/\rho$ . Equations (10) and (11) show that

$$p = -\rho - \frac{2}{K}\dot{H}, \quad (12)$$

where  $\dot{H}$  expresses the rate of change of the Hubble parameter as

$$\dot{H} = -\frac{K}{2}(\rho + p) = -\frac{K}{2}(\gamma\rho). \quad (13)$$

We have considered the EoS  $p = (\gamma - 1)\rho$  with  $\frac{2}{3} \leq \gamma \leq 2$ . As has been pointed out in [88], inserting this equation for  $\dot{H}$  in (8) for the flat case results in .

$$H^2 = \frac{K\rho(3K\lambda\gamma - 1)}{3(4k\lambda - 1)}. \quad (14)$$

The physical behavior of energy density has been plotted in figure (1). The pressure changes sign from positive in a decelerating era to negative in an accelerating era dominated by dark energy with negative pressure. As a function of redshift, The EoS parameter  $\omega(z)$  meets  $-1$  at  $z = 0$ . The time evolution of the EoS parameter verses reveals a Quintom behavior as it passes the phantom divide line at far future. The Hubble parameter  $H$  is decreasing during the expansion as  $\dot{H} < 0$ , while it is increasing during the contraction where  $\dot{H} > 0$ . The general dynamical behavior can be deduced from the negative values of  $q$ . In each cycle, the EoS parameter  $\omega(t)$  lies between  $-2.25$  and  $\frac{1}{3}$  which agrees with observations [67, 68]. It starts from a positive value (radiation-like era), keeps decreasing to zero (dust era  $\omega = 0$ ) and passes to the negative values. After reaching the DE-dominated era, it crosses the cosmological constant boundary to the phantom era where  $\omega < -1$  possessing a Quintom feature. The Quintom dynamics associated with the crossing of the phantom divide line leads to  $\omega < -1$  today is also observationally supported [69]. A cosmological bouncing Quintom model has been studied in [70].  $\omega$  is an increasing function in the second half of the cosmic cycle starting from  $-2.25$ . Some observations favor DE with  $\omega$  less than  $-1$  [71, 72, 73]. The idea whether DE can evolve to the phantom era or not has been extensively investigated. In [71] it has been shown that, in a DE dominated universe, the transition of DE from  $\omega \geq -1$  to  $\omega < -1$  can't be explained by models of classical scalar fields dynamics unless more complicated physics is included [74, 75, 76].

### 3 Work Density and Entropy

In this section, we calculate and analyze some thermodynamical quantities. The  $(n+1)$ -dimensional FRW metric is written as

$$ds^2 = -dt^2 + a^2(t) \left( \frac{dr^2}{1 - \kappa r^2} + r^2 d\Omega_{n-1}^2 \right), \quad (15)$$

where the line element of an  $(n-1)$ -dimensional unit sphere is denoted by  $\Omega_{n-1}^2$ . This metric can also be written as [86]

$$ds^2 = h_{ab} dx^a dx^b + \tilde{r}^2 \Omega_{n-1}^2, \quad (16)$$

where  $\tilde{r} = a(t)r$ ,  $x^0 = t$ ,  $x^1 = r$  and the 2-dimensional metric  $h_{ab} = \text{diag}(-1, \frac{a^2}{1-\kappa r^2})$ . The apparent horizon is defined geometrically as an imaginary surface beyond which null geodesic congruences recede from the observer [66]. It is a dynamical structure which evolves with time and is determined by the relation  $h_{ab} \partial_a \tilde{r} \partial_b \tilde{r} = 0$ . This leads to the expression of the apparent horizon's radius in terms of the Hubble parameter as

$$\tilde{r}_A = \frac{1}{\sqrt{H^2 + \frac{\kappa}{a^2}}}. \quad (17)$$

For a flat universe (the case we are considering here), The apparent horizon's radius is the inverse of the Hubble parameter  $1/H$  which is the same definition of the Hubble horizon  $\tilde{r}_H$ . That means

$$\tilde{r}_A = \tilde{r}_H = \frac{1}{H}, \quad \text{for } \kappa = 0. \quad (18)$$

Consequently, the rate of change of the horizon radius  $\dot{\tilde{r}}_A = -\frac{\dot{H}}{H^2} = -\tilde{r}_A^2 \dot{H}$  or can be written in general as

$$\dot{\tilde{r}}_A = -\tilde{r}_A^3 H \left( \dot{H} - \frac{\kappa}{a^2} \right). \quad (19)$$




Figure 1: Evolution of the model parameters against the redshift  $z$  (a) The DP  $q(z)$  changes sign for  $m > 1$  and equals  $-1$  at  $z = 0$ . (b) The Hubble parameter  $H(z)$ . (c) The physical behaviour of energy density  $\rho(z)$ . (d) A sign flipping in the evolution of cosmic pressure  $p(z)$  (e) The EoS parameter  $\omega(z)$  equals to  $-1$  at  $z = 0$ . (f)  $\omega(t)$  evolution shows a Quantum behaviour where it crosses the phantom divide line at very late times. GR stands for General Relativity and RG for Rastall Gravity. (g) The rate of change of the Hubble parameter  $\dot{H} < 0$  during the expansion which means a decreasing  $H$ , while this rate is  $> 0$  during the contraction. (h) The Hubble parameter  $H > 0$  during the expansion, and  $H < 0$  during the contraction. (i) The evolution of scale factor shows an enormous increase over a short period of time (a future big rip). The unit of time  $t$  is taken in Gyr. Here  $m = 1.55$ ,  $\lambda = 1.4$ ,  $K = 0.01$ ,  $k = 0.1$  and  $a_0 = 1$ .

The work density  $W$  is defined as the work done by the cosmic volume change due to the change of the radius of the apparent horizon. For the current model we get

$$W = -\frac{1}{2}T^{ab}h_{ab} = \frac{1}{2}(\rho - p) \quad (20)$$

$$= -\frac{1}{K} \left( 12\lambda KH^2 + 6\lambda K\dot{H} - 3H^2 - \dot{H} \right). \quad (21)$$

In the case of the FRW spacetime with a perfect fluid. The surface gravity  $\kappa_{sg}$  is also defined as

$$\kappa_{sg} = \frac{1}{2\sqrt{-h}}\partial_a(\sqrt{-h}h^{ab}\partial_b\tilde{r}_A) = -\frac{1}{\tilde{r}_A} \left( 1 - \frac{1}{2} \frac{\dot{\tilde{r}}_A}{H\tilde{r}_A} \right), \quad (22)$$

where  $h$  is the determinant of  $h_{ab}$ . In terms of  $\kappa_{sg}$ , the temperature on the horizon is given as  $\frac{|\kappa_{sg}|}{2\pi}$

$$T_A = \frac{1}{2\pi\tilde{r}_A} \left| 1 - \frac{1}{2} \frac{\dot{\tilde{r}}_A}{H\tilde{r}_A} \right|. \quad (23)$$

So, the temperature depends on the Hubble parameter  $H$ , The radius of the apparent horizon  $\tilde{r}_A$  and its rate of change during evolution  $\dot{\tilde{r}}_A$  which is a result of the dynamical nature of the apparent horizon. If the change in the apparent horizon is very slow, then  $\frac{\dot{\tilde{r}}_A}{H\tilde{r}_A} \ll 1$  and we obtain the expression for Hawking temperature  $T_H = \frac{1}{2\pi\tilde{r}_A}$  which resembles the temperature of a spherically symmetric black hole with horizon radius  $\tilde{r}_A$  [87].

The modified Bekenstein-Hawking entropy in Rastall gravity on the apparent horizon is given as [88, 89]

$$\tilde{S} = \left( 1 + \frac{2\gamma}{4\gamma - 1} \right) S_0, \quad (24)$$

with  $S_0 = \frac{A}{4}$  is the Bekenstein-Hawking entropy on the apparent horizon and  $A = 4\pi\tilde{r}_A^2$  is the area. The normal Bekenstein-Hawking entropy in GR is recovered for  $\gamma \rightarrow 0$ . So, the area  $A = 4\pi\tilde{r}_A^2$  is related to the modified area  $\tilde{A}$  by  $\tilde{A} = \left( 1 + \frac{2\gamma}{4\gamma - 1} \right) A$  where the units  $c = G = \hbar = 1$  has been considered. As a function of cosmic time  $t$ , the entropy should always be an increasing function where the Universe evolves to the equilibrium state of maximum entropy. That means the two conditions

$$\dot{S} \geq 0 \quad \text{and} \quad \ddot{S} \leq 0, \quad (25)$$

should be satisfied. We recall that for a system at equilibrium, entropy has the maximum value which means that it can't increase anymore. On the other hand, any decrease is not possible as it violates the second law of thermodynamics which states that

$$d(S_m + S_h) > 0, \quad (26)$$

where  $S_h$  is the horizon entropy, and  $S_m$  the entropy of the entire matter fields. At the apparent horizon, the FRW equations can be written as  $dE = TdS + WdV$  where  $E$  is the total energy and  $W = \frac{1}{2}(\rho - p)$  is the work density [77, 78]. The cosmological work density has been related to the cosmological constant in [79]. The thermodynamical quantities  $w$ ,  $\tilde{r}_A$ ,  $T_A =$  and  $S$  can

be simply expressed as a function of redshift  $z$  to probe their behavior. Figure 2 shows that the work density  $W$  has the same physical behavior of energy density  $\rho$  versus  $z$ . In terms of cosmic time, it's a decreasing function during expansion which means that the work done by the cosmic volume change due to the change of the apparent horizon radius decreases. The first derivative of the event horizon radius is positive during the expansion  $\dot{\tilde{r}}_A > 0$  and negative during the contraction  $\dot{\tilde{r}}_A < 0$ . A comparison between the evolution of Hawking temperature  $T_H$  and the temperature on the horizon  $T_A$  versus  $z$  has been plotted in Figure 2(c). Entropy  $\tilde{S}$  is always positive. While the condition  $\dot{\tilde{S}} > 0$  is satisfied, the non-positivity of the equilibrium condition  $\ddot{\tilde{S}}$  exists only for the second half of cosmic cycle.

## 4 Causality and Energy Conditions

One way to investigate the physical acceptability is testing the classical linear energy conditions (ECs) [80, 81], as the quantum corrections are ignored in the present model. The null, weak, strong and dominant ECs are respectively:  $\rho + p \geq 0$ ;  $\rho \geq 0$ ,  $\rho + p \geq 0$ ;  $\rho + 3p \geq 0$  and  $\rho \geq |p|$ . These linear conditions can't be satisfied in the existence of quantum effects [82]. The strong energy condition (SEC), for example, implies that gravity should always be attractive which is not realistic when describing cosmic acceleration or inflation [83, 84, 85]. For the present model,

$$\begin{aligned}\rho + p &= -\frac{2}{K}\dot{H}, \\ \rho - p &= -\frac{2}{K}(12\lambda KH^2 + 6\lambda K\dot{H} - 3H^2 - \dot{H}), \\ \rho + 3p &= \frac{6}{K}(4\lambda KH^2 + 2\lambda K\dot{H} - H^2 - \dot{H}).\end{aligned}$$

The SEC is not expected to be valid during the negative pressure-dominated accelerating era as the negative pressure represents a repulsive gravity effect. The DEC is valid all the time for this model which is expected as it implies the non-negativity of energy density. The SEC is valid only with the domination of attractive gravity in the decelerating era (at the first half of each cycle), and then becomes invalid with the domination of repulsive gravity during the accelerating era.

The causality condition for the adiabatic square sound speed  $0 \leq \frac{dp}{d\rho} \leq 1$  should be satisfied through cosmic evolution. For the current model, using (10) and (11), we get

$$c_s^2 = -\frac{1}{3} \frac{12\lambda KH\dot{H} + 3\lambda K\ddot{H} - 3H\dot{H} - \ddot{H}}{4\lambda KH\dot{H} + \lambda K\ddot{H} - H\dot{H}}. \quad (27)$$

Figure 2(h) shows that the causality is satisfied except near the initial and the future Big Rip singularities






Figure 2: (a) Work density verses redshift. (b) Radius of the apparent horizon verses redshift. (c) Temperature on dynamical horizon verses Hawking temperature of a spherically symmetric black hole. (d) & (e) evolution of entropy verses  $t$  and  $z$  shows that  $\tilde{S} \geq 0$ . (f)  $\dot{\tilde{S}}(t)$  and  $\ddot{\tilde{S}}(t)$ , while the condition  $\dot{\tilde{S}} > 0$  is satisfied, the non-positivity of the equilibrium condition  $\ddot{\tilde{S}} < 0$  exists only for the second half of cosmic cycle. (g)  $\dot{\tilde{S}}(z)$  and  $\ddot{\tilde{S}}(z)$ . A violation of  $\dot{\tilde{S}} < 0$  starts at  $z \gtrsim 0.4$  (h) The causality condition  $0 \leq \frac{dp}{d\rho} \leq 1$  is satisfied except near the initial and the future Big Rip singularities (i) Linear energy conditions. Here  $m = 1.55$ ,  $\lambda = 1.4$ ,  $K = 0.01$ ,  $k = 0.1$  and  $a_0 = 1$ .

## 5 Conclusion

We have presented a cyclic cosmological model of a flat universe with future big rip singularity in the framework of Rastall gravity. The model possesses a quintom behavior where the EoS parameter crosses the phantom divide line (cosmological constant boundary) at  $\omega = -1$ . In such oscillatory cosmic evolution, the deceleration-to-acceleration Cosmic transit happens at around 8.7 Gyr. A recent analysis that included data from cosmic chronometers and the Pantheon compilation of supernovae, calculated the transition redshift to be  $z_t = 0.61_{-0.16}^{+0.24}$  [90]. Given that the universe is currently 13.8 Gyr old, this correlates to a transition period of roughly 7.5 to 9.8 billion years ago. A transition redshift of  $z_t = 0.60_{-0.12}^{+0.21}$  was discovered by another study that looked at a parametrization of the deceleration parameter [91]. This estimate, which places the transition era between 7.7 and 9.6 Gyr ago, is in good agreement with the previous one. A cosmic deceleration-to-acceleration transition that took place approximately 8.7 Gyr ago is within the observed range based on these observational estimations. Thus, our conclusion so seems to be in accordance with recent evidence from observations.

The EoS parameter  $\omega$  meets  $-1$  at redshift  $z = 0$ . The model exhibits the correct physical evolution of energy density and a sign flipping of cosmic pressure in compatible with cosmic transit. Observational evidence supports that the EoS parameter  $\omega$  is very close to  $-1$  at the present epoch (redshift  $z = 0$ ) [92, 93].

The first derivative of the Hubble parameter  $\dot{H}$  is negative during the expansion which means a decreasing  $H$ , and positive during contraction which means an increasing  $H$ . The EoS parameter  $\omega$  lies in the range  $-2.25 \leq \omega(t) \lesssim \frac{1}{3}$ . It begins at the radiation-like era then passes through the dust era, the current dark energy-dominated era before crossing the cosmological constant boundary to the phantom era. In the same time, plotting the  $\omega$  evolution in Rastall Gravity (RG) and in General relativity (GR) together shows a tiny difference between the two curves.

The evolution of the temperature on dynamical horizon and Hawking temperature has been plotted versus redshift. The entropy in the model is positive all the time. It starts as an increasing function of time in the first expanding half cycle until it reaches maximum value corresponding to an equilibrium state. The rate of change of the entropy along with the equilibrium condition has been calculated and plotted. The causality is satisfied all the time except near the initial singularity and the future Big Rip singularity.

It is pertinent to investigate an oscillating Rastall universe that crosses the phantom division line because it provides a theoretical foundation for investigating dynamic dark energy models and possible gravity alterations. Therefore, our research is useful and pertinent to the oscillating Rastall cosmos.

## Acknowledgments

The authors (A. Pradhan & A. Dixit) are grateful for the resources and support provided during a visit to the Inter-University Centre for Astronomy & Astrophysics (IUCAA), Pune, India, as part of their Associateship program.

## ORCID

Nasr Ahmed <https://orcid.org/0000-0003-1544-9970>

Anirudh Pradhan <https://orcid.org/0000-0002-1932-8431>

Archana Dixit <https://orcid.org/0000-0003-4285-4162>

## References

- [1] S. Perlmutter *et al.* [Supernova Cosmology Project], Measurements of  $\Omega$  and  $\Lambda$  from 42 high redshift supernovae, *Astrophys. J.* **517** (1999), 565-586, [arXiv:astro-ph/9812133 [astro-ph]].
- [2] A. G. Riess *et al.* [Supernova Search Team], Observational evidence from supernovae for an accelerating universe and a cosmological constant, *Astron. J.* **116** (1998), 1009-1038, [arXiv:astro-ph/9805201 [astro-ph]].
- [3] D. N. Spergel *et al.* [WMAP], First year Wilkinson Microwave Anisotropy Probe (WMAP) observations: Determination of cosmological parameters, *Astrophys. J. Suppl.* **148** (2003), 175-194, [arXiv:astro-ph/0302209 [astro-ph]].
- [4] E. Komatsu *et al.* [WMAP], Five-Year Wilkinson Microwave Anisotropy Probe observations: Cosmological interpretation, *Astrophys. J. Suppl.* **180**, 330-376 (2009).
- [5] P. Astier *et al.* [SNLS], The Supernova Legacy Survey: measurement of  $\Omega_M$ ,  $\Omega_\Lambda$  and  $w$  from the first year data set *Astron. Astrophys.* **447**, 31-48 (2006).
- [6] E. J. Copeland, M. Sami and S. Tsujikawa, Dynamics of dark energy, *Int. J. Mod. Phys. D* **15** (2006) 1753.
- [7] S. Nojiri and S. D. Odintsov, Modified  $f(R)$  gravity consistent with realistic cosmology: From a matter dominated epoch to a dark energy universe, *Phys. Rev. D* **74** (2006) 086005.
- [8] S. Nojiri, S. D. Odintsov and P. V. Tretyakov, From inflation to dark energy in the non-minimal modified gravity, *Prog. Theor. Phys. Suppl.* **172** (2008) 81-89.

- [9] N. Ahmed, A. Alfreedi and A. Alzulaibani, Some cosmological features of 4D Gauss-Bonnet gravity with varying cosmological constant. *Int. J. Geom. Meth. Mod. Phys.* **20** (2003) 2350053.
- [10] R. Ferraro and F. Fiorini, Modified teleparallel gravity: Inflation without an inflaton, *Phys. Rev. D* **75** (2007) 084031.
- [11] N. Ahmed and A. Pradhan, Probing cosmic acceleration in  $\kappa(R, T)$  gravity, *Indian J. Phys.* **96** (2022) 301-307.
- [12] J. M. Overduin and P. S. Wesson, Kaluza-Klein gravity, *Phys. Rept.* **283** (1997) 303-380.
- [13] T. P. Sotiriou, Horava-Lifshitz gravity: a status report, *J. Phys. Conf. Ser.* **283** (2011) 012034.
- [14] N. Ahmed, K. Bamba and F. Salama, The possibility of a stable flat dark energy-dominated Swiss-cheese brane-world universe, *Int. J. Geom. Meth. Mod. Phys.* **17** (2020) 2050075.
- [15] N. Ahmed and H. Rafat, Topological Origin of Holographic Principle, application to worm-holes. *Int. J. Geom. Methods Mod. Phys.* Vol. 15, No. 8 – 1850131 (2018).
- [16] M. Novello and S. E. P. Bergliaffa, Bouncing Cosmologies. *Phys. Rept.* **463** (2008) 127-213.
- [17] A. Ijjas and P. J Steinhardt, Bouncing cosmology made simple, *Class. Quant. Grav.* **35** (2018) 135004
- [18] P. Sahoo, S. Bhattacharjee, S. K. Tripathy and P. K. Sahoo, Bouncing scenario in  $f(R, T)$  gravity, *Mod. Phys. Lett. A* **35** (2020) 2050095.
- [19] S. Nojiri, S. D. Odintsov, V. K. Oikonomou and T. Paul, Nonsingular bounce cosmology from Lagrange multiplier  $f(R, T)$  gravity, *Phys. Rev. D* **100** (2019) 084056.
- [20] R. Brandenberger and P. Peter, Bouncing cosmologies: Progress and problems, *Found. Phys.* **47** 797 (2017).
- [21] K. Bamba, A. N. Makarenko, A. N. Myagky, S. Nojiri and S. D. Odintsov, Bounce cosmology from  $F(R)$  gravity and  $F(R)$  bigravity, *J. Cosmol. Astropart. Phys.* **1401** (2014) 008.
- [22] K. Bamba, A. N. Makarenko, A. N. Myagky and S. D. Odintsov, Bouncing cosmology in modified Gauss-Bonnet gravity, *Phys. Lett. B* **732** (2014) 349-355.
- [23] K. Bamba, A. N. Makarenko, A. N. Myagky and S. D. Odintsov, Bounce universe from string-inspired Gauss-Bonnet gravity, *J. Cosmol. Astropart. Phys.*, **1504** (2015) 001.
- [24] S. K. Tripathy, R. K. Khuntia, P. Parida, Bouncing cosmology in an extended theory of gravity, *Eur. Phys. J Plus* **134** (2019) 504.
- [25] A. de la Cruz-Dombriz, G. Farrugia, J. L. Said and D. S.-C. Gomez, Cosmological bouncing solutions in extended teleparallel gravity theories, *Phys. Rev. D* **97** (2018) 104040.

- [26] N. Ahmed, T. M Kamel and M. I. Nouh, A non-singular closed bouncing universe without violation of null energy condition, *Rev. Mex. Astron. Astrofis.* **58 (2)** (2022) 245-255, arXiv:2204.11854v1 [gr-qc].
- [27] P. Pavlovic and M. Sossich, Cyclic cosmology in modified gravity, *Phys. Rev. D* **95** (2017) 103519.
- [28] P. H. Frampton, Cyclic cosmology and holographic entanglement entropy, *Int. J. Mod. Phys. A* **33** (2018) 1844028.
- [29] P. J. Steinhardt, N. Turok, Cosmic evolution in a cyclic universe *Phys. Rev. D* **65**, 126003 (2002).
- [30] A. Ijjasa, P. J. Steinhardt, A new kind of cyclic universe, *Phys. Lett. B* **795** (2019) 666–672.
- [31] M K. Sharma et. al, The oscillatory universe, phantom crossing and the Hubble tension. *Annals of Physics V* 454-169345 (2023).
- [32] F. Hoyle, G. Burbidge, J. V. Narlikar, A Quasi–Steady State cosmological model with creation of matter, *Astrophys. J.* **410** (1993) 437.
- [33] F. Hoyle, G. Burbidge, J. V. Narlikar, Astrophysical deductions from the quasi-steady-state cosmology, *MNRAS* **267(4)** (1994) 1007.
- [34] F. Hoyle, G. Burbidge, J. V. Narlikar, Further astrophysical quantities expected in a quasi-steady state Universe, *A&A* **289(3)** (1994) 729.
- [35] F. Hoyle, G. Burbidge, J. V. Narlikar, The basic theory underlying the quasi-steady-state cosmology, *Proc. R. Soc. A* **448** (1995) 191.
- [36] R. Sachs, J. V. Narlikar, F. Hoyle, The quasi-steady state cosmology: analytical solutions of field equations and their relationship to observations, *A&A* **313** (1996) 703.
- [37] P. Rastall, Generalization of the Einstein theory, *Phys. Rev. D* **6** (1972) 3357-3359.
- [38] W.A.G. De Moraes and A.F. Santos, Lagrangian formalism for Rastall theory of gravity and Gödel-type universe, *Gen. Rel. Grav.* **51** (2019) 167-184.
- [39] V. Majernik and L. Richterek, Rastall’s gravity equations and Mach’s Principle, arXiv:gr-qc/0610070 (2006).
- [40] P. Rastall, A theory of gravity, *Can. J. Phys.* **54** (1976) 66-75.
- [41] D. Das, S. Dutta and S. Chakraborty, Cosmological consequences in the framework of generalized Rastall theory of gravity, *Eur. Phys. J. C* **78** (2018) 810-818.
- [42] S. Ghaffari, A.A Mamon, H. Moradpour and A.H. Ziaie, Holographic dark energy in Rastall theory, *Mod. Phys. Lett. A* **35** (2020) 2050276-2050290.

- [43] R. Saleem and Shahnila, Cosmological evolution interacting/non-interaction holographic dark energy model for curved FLRW space-time in Rastall gravity, *Phys. Dark Univ.* **32** (2021) 100808.
- [44] R. Saleem, I. Shahid, M. I. Aslam and A. Wahab, Constant-roll warm inflation within Rastall gravity, *Nucl. Phys. B* **1009** (2024) 116728.
- [45] J. K. Singh, A. Singh, S. G. Ghosh and S. D. Maharaj, Evolution of the universe with quintessence model in Rastall gravity, arXiv:2410.12012[gr-qc] (2024).
- [46] R. Saleem and J. Hassan, Confronting the warm vector inflation in Rastall theory of gravity with Planck 2018 data, *Phys. Dark Univ.* **28** (2020) 100515.
- [47] A. Banerjee, T. Tangphati, S. Hansraj and A. Pradhan, Strange quark star models from Rastall gravity, *Ann. Phys.* **451** (2023), 169267.
- [48] A. Banerjee, T. Tangphati and A. Pradhan, Possible existence of quark stars in Rastall gravity, *Int. J. Mod. Phys. D* **32** (2023), 2350026.
- [49] T. Tangphati, A. Banerjee, S. Hansraj and A. Pradhan, The criteria of the anisotropic quark star models in Rastall gravity, *Ann. Phys.* **452** (2023) 169285.
- [50] T. Tangphati, A. Banerjee, I. Sakalli and A. Pradhan, Quark stars in Rastall gravity with recent astrophysical observations, *Chin. J. Phys.* **90** (2024) 422-433.
- [51] A. Banerjee, S. Islam, A. Dixit and A. Pradhan, Wormholes in Rastall gravity and nonvacuum spacetime, *Gravitation & Cosmology* **30** (2024) 496-505.
- [52] Shaily, A. Singh, J. K. Singh, S. Hussain and R. Myrzakulov, Stability analysis of a dark energy model in Rastall gravity, arXiv:2402.08709 [gr-qc] (2024).
- [53] A. Singh, R. Raushan and R. Chaubey, Qualitative aspects of Rastall gravity with barotropic fluid, *Can. J. Phys.* **99** (2021) 1073-1081
- [54] A. Singh, G. P. Singh and A. Pradhan, Cosmic dynamics and qualitative study of Rastall model with spatial curvature, *Int. J. Mod. Phys. A* **37** (2022) 2250104.
- [55] O Akarsu, N. Katirci, S. Kumar, R. C. Nunes, B. Ozturk and S. Sharma, Rastall gravity extension of the standard  $\Lambda$ CDM model: theoretical features and observational constraints, *Eur. Phys. J. C* **80** (2020) 1050.
- [56] M. Shen and L. Zhao, Oscillating quintom model with time periodic varying deceleration parameter, *Chin. Phys. Lett.* **31** (2014) 010401.
- [57] S. Capozziello, O. Farooq, O. Luongo and B. Ratra, Cosmographic bounds on the cosmological deceleration-acceleration transition redshift in  $f(R)$  gravity, *Phys. Rev. D* **90** (2014) 044016.
- [58] S. Capozziello, O. Luongo and E. N. Saridakis, Transition redshift in  $f(T)$  cosmology and observational constraints, *Phys. Rev. D* **91** (2015) 124037.

- [59] O. Farooq, F. Madiyar, S. Crandall and B. Ratra, Hubble parameter measurement constraints on the redshift of the deceleration-acceleration transition, dynamical dark energy, and space curvature, *Astrophys. J.* **835** (2017) 26.
- [60] P. H. R. S. Moraes, R. A. C. Correa and G. Ribeiro, Comment on "Cosmological inviability of f(R,T) gravity, *Astrophys Space Sci.* **361** (2016) 227.
- [61] P. Rastall, Generalization of the Einstein Theory, *Phys. Rev. D* **6** (1972) 3357.
- [62] M. Tegmark et al., Cosmological parameters from SDSS and WMAP, *Phys. Rev. D* **69** 103501 (2004).
- [63] C.L. Bennett et al., First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Preliminary Maps and Basic Results, *Astrophys. J. Suppl.* **148** 1 (2003).
- [64] D.N. Spergel et al., First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Determination of Cosmological Parameters, *Astrophys. J. Suppl.* **148** 175 (2003).
- [65] N. Ahmed and S. Z. Alamri, A stable flat entropy-corrected FRW universe. *Int. J. Geom. Methods Mod. Phys.* **16** (2019) 1950159.
- [66] S. Saha, *Elements of cosmological thermodynamics*, Springer 2018.
- [67] E. Komatsu, K. M. Smith, J. Dunkley, et. al., Seven-Year Wilkinson Microwave Anisotropy Probe (WMAP) observations: Cosmological interpretation, *Astrophys. J. Suppl. Ser.* **192** (2011) 18.
- [68] D. J. Eisenstein, I. Zehavi, D. W. Hogg et al., Detection of the Baryon Acoustic Peak in the Large-Scale Correlation Function of SDSS Luminous Red Galaxies, *Astrophys. J.* **633** (2005) 560.
- [69] G.-B. Zhao, J.-Q. Xia, H. Li, C. Tao, J. M. Virey, Z. H. Zhu and X. Zhang, Probing for dynamics of dark energy and curvature of universe with latest cosmological observations, *Phys. Lett. B* **648** (2007) 8-13.
- [70] Yi-Fu Cai, T. Qiu, X. Zhang, Yun-Song Piao and M. Li, Bouncing universe with Quintom matter, *JHEP* **0710** 071 (2007).
- [71] A. Vikman, Can dark energy evolve to the Phantom?, *Phys.Rev. D* **71** (2005) 023515.
- [72] S. W. Allen, R. W. Schmidt, H. Ebeling, et al., Constraints on dark energy from Chandra observations of the largest relaxed galaxy clusters, *Mon. Not.Roy. Astron. Soc.* **353** (2004) 457.
- [73] A. G. Riess et al., Type Ia Supernova Discoveries at  $z > 1$  from the Hubble Space Telescope: Evidence for Past Deceleration and Constraints on Dark Energy Evolution\*, *Astrophys. J.* **607** (2004) 665-687.
- [74] C. Armendariz-Pion, Could dark energy be vector-like?, *JCAP* **0407** 007 (2004), astro-ph/0405267.

- [75] Bo Feng, X. Wang and X. Zhang, Dark Energy Constraints from the Cosmic Age and Supernova, *Phys. Lett. B* **607** (2005) 35-41, astro-ph/0404224.
- [76] V. Sahni and Y. Shtanov, Braneworld models of dark energy, *JCAP* **0311** (2003) 014.
- [77] R. G. Cai and S. P. Kim, First law of thermodynamics and Friedmann equations of Friedmann-Robertson-Walker universe, *JHEP* **2005(02)** (2005) 050.
- [78] M. Akbar and R. G. Cai, Thermodynamic behavior of the Friedmann equation at the apparent horizon of the FRW universe, *Phys. Rev. D* **75** (2007) 084003.
- [79] N. Ahmed and S. Z. Alamri, A cyclic universe with varying cosmological constant in  $f(R, T)$  gravity, *Can. J. Phys.* **999** (2019) 1-8.
- [80] S. W. Hawking and G. F. R. Ellis, *The large scale structure of spacetime*, (Cambridge University Press, England 1973).
- [81] R. M. Wald, *General Relativity* (University of Chicago Press, Chicago 1984).
- [82] P. Martín-Moruno and M. Visser, Semiclassical energy conditions for quantum vacuum states, *JHEP* **1309** (2013) 050.
- [83] M. Visser, Energy Conditions in the Epoch of Galaxy Formation, *Science* **276** (1997) 88-90.
- [84] M. Visser, General relativistic energy conditions: The Hubble expansion in the epoch of galaxy formation, *Phys. Rev. D* **56** (1997) 7578.
- [85] M. Visser, Energy conditions and galaxy formation, gr-qc/9710010.
- [86] D. Bak and S. J. Rey, Cosmic Holography, *Class. Quant. Grav.* **17** (2000) L83, [arXiv:hep-th/9902173].
- [87] L. Sanchez and H. Quevedo, Thermodynamics of the FLRW apparent horizon, *Phys. Lett. B* **839** (10) 137778 (2023).
- [88] K. Bamba, A. Jawad, S. Rafique and H. Moradpour, Thermodynamics in Rastall gravity with entropy corrections, *Eur. Phys. J. C* **78** (2018) 986.
- [89] H. Moradpour and I. G. Salako, Thermodynamic analysis of the static spherically symmetric field equations in Rastall theory, *Adv. High Ener. Phys.* **2016** (2016) 3492796.
- [90] Y. Yang and Y. Gong, The evidence of cosmic acceleration and observational constraints, *JCAP* **2000(06)** (2020) 059, arXiv:1912.07375 [astro-ph.CO].
- [91] S. Arora and P. K. Sahoo, Revisiting kink-like parametrization and constraints using OHD/Pantheon+/BAO samples, *Phys. Dark Univ.* **45** (2024) 101510, arXiv:2312.06464 [astro-ph.CO].
- [92] A. Tripathi, A. Sangwan and H. K. Jassal, Dark energy equation of state parameter and its variation at low redshifts, *JCAP* **2017(06)** (2017) 012, arXiv:1611.01899 [astro-ph.CO].



- [93] B. R. Dinda and N. Banerjee, A comprehensive data-driven odyssey to explore the equation of state of dark energy, *Eur. Phys. J. C* **84** (2024) 688, arXiv:2403.14223 [astro-ph.CO].