

Universal Wilson loop Bound of Quantum Geometry: Z_2 Bound and Physical Consequences

Jiabin Yu,^{1,2,*} Jonah Herzog-Arbeitman,² and B. Andrei Bernevig^{2,3,4}

¹*Department of Physics, University of Florida, Gainesville, FL, USA*

²*Department of Physics, Princeton University, Princeton, New Jersey 08544, USA*

³*Donostia International Physics Center, P. Manuel de Lardizabal 4, 20018 Donostia-San Sebastian, Spain*

⁴*IKERBASQUE, Basque Foundation for Science, Maria Diaz de Haro 3, 48013 Bilbao, Spain*

We define the absolute Wilson loop winding and prove that it bounds the quantum metric from below. This Wilson loop lower bound naturally reproduces the known Chern and Euler bounds of the quantum metric, and provides an explicit lower bound of the quantum metric due to the time-reversal protected Z_2 index, answering a hitherto open question. In general, the Wilson loop lower bound can be applied to any other topological invariants characterized by Wilson loop winding, such as the particle-hole Z_2 index. As physical consequences of the Z_2 bound, we show that the time-reversal Z_2 index bounds superfluid weight and optical conductivity from below, and bounds the direct gap of a band insulator from above.

I. INTRODUCTION

Quantum geometry is a fundamental framework for the geometric properties of Bloch wavefunctions, which underpins a wide range of physical phenomena. A key quantity that characterizes quantum geometry is the quantum metric [1–4],

$$[g_{\mathbf{k}}]_{ij} = \frac{1}{2} \text{Tr} [\partial_{k_i} P_{\mathbf{k}} \partial_{k_j} P_{\mathbf{k}}] , \quad (1)$$

where i, j denote spatial components, \mathbf{k} is Bloch momentum, and $P_{\mathbf{k}}$ is the projector for the isolated set of bands of interest. The quantum metric plays a pivotal role in diverse physics, including optical conductivity [5–11], the superfluid weight of superconductors [12–21], electron-phonon coupling [22, 23], and correlated charge fluctuations [24, 25]. Exploring the physical implications of the quantum geometry has become a central focus in condensed matter physics, with efforts directed toward identifying more and more physical quantities influenced by the quantum metric. (See Ref. [26] for a review.)

In this work, we will not focus on finding new physical responses influenced by the quantum metric; instead, we will focus on the topological lower bounds of the quantum metric. Such bounds provide important insights on the physical phenomena they are tied to (such as the existence of superconductivity in flat band systems). It is natural that the quantum metric cannot vanish in systems with nontrivial band topology, as a vanishing quantum metric implies an absence of \mathbf{k} -dependence in the periodic part of the Bloch states (in certain gauge), inevitably leading to trivial topology. Nevertheless, the key question is to find the explicit expression for the lower bound given by various topological invariants. Explicit lower bounds have been established for the Chern number [27], Euler number [16, 19, 28, 29], and some 2D

obstructed atomic limits and fragile topological states [30]; however, determining the explicit lower bound for the quantum metric associated with the Kane-Mele time-reversal (TR) Z_2 invariant [31–34] remains a long-term unresolved challenge. A broader question is how nontrivial Wilson loop (WL) winding [35], which captures the Chern number [36, 37], Euler number [38–40], TR Z_2 index [35], and other topological invariants [41, 42], rigorously enforces lower bounds on the quantum metric.

In this work, we resolve these two questions. We first prove a explicit expression for the lower bound of the quantum metric due to nontrivial WL winding. Specifically, we define the “absolute WL winding”, which is the sum of the absolute phase changes of all WL eigenvalues without cancellation. Then, with the non-abelian Stokes theorem [43], we can prove that the integration of $\text{Tr}[g_{\mathbf{k}}]$ is bounded from below by the absolute WL winding. As a consequence, we derive the TR- Z_2 lower bound of the quantum metric in two spatial dimensions, which reads

$$\frac{1}{2\pi} \int_{BZ} d^2k \text{Tr}[g_{\mathbf{k}}] \geq 2Z_2 , \quad (2)$$

where the integration is over the Brillouin zone (BZ) and the TR- Z_2 index takes the values 0 in the trivial case and 1 in the topological case, characterized by helical edge states [31, 33, 44] and an obstruction to local Kramers pair Wannier functions [45]. It is protected by spinful time-reversal obeying $\mathcal{T}^2 = -1$. Eq. (2) also holds for the particle-hole (PH) protected Z_2 index, which, as Ref. [46] points out, is equivalent to the TR- Z_2 index. We further describe how the inequality Eq. (2) provides lower bounds for superfluid weight and optical conductivity and an upper bound for the direct gap of a band insulator.

II. GENERAL WILSON LOOP BOUND OF THE QUANTUM METRIC

We start by sketching the proof of the general WL lower bound of the quantum metric. The setup of the

* yujiabin@ufl.edu




FIG. 1. (a) shows the WL of an infinitesimal $\hat{D}_{\mathbf{k}} = [k_x, k_x + dk_x] \times [k_y, k_y + dk_y]$, which equals $\exp[-iF_{\mathbf{k}}dk_xdk_y]$ with $F(\mathbf{k})$ the non-abelian Berry curvature. (b) shows that the multiplication of WLs $W(\partial\hat{D}_{(k_x, k_y)})$ and $W(\partial\hat{D}_{(k_x, k_y + dk_y)})$ is not necessarily $W(\partial[\hat{D}_{(k_x, k_y)} \cup \hat{D}_{(k_x, k_y + dk_y)}])$ owing to the non-abelian nature of the WL. (c) shows the WL of an infinitesimal $\hat{D}_{\mathbf{k}} = [k_x, k_x + dk_x] \times [k_y, k_y + dk_y]$ combined with with dressing $h_{\mathbf{k}}$ and $h_{\mathbf{k}}^{-1}$, where $h_{\mathbf{k}}$ is the red Wilson line, and the inverse $h_{\mathbf{k}}^{-1}$ is the blue Wilson line. Here $\tilde{F}(\mathbf{k}) = h_{\mathbf{k}}F_{\mathbf{k}}h_{\mathbf{k}}^{-1}$, the dressed non-abelian Berry curvature. (d) shows the merging of two dressed WLs into one dressed WL on a larger loop.

problem is an isolated set of N bands of a 2D noninteracting lattice system, and we label the periodic parts of the relevant Bloch states as $|u_{n, \mathbf{k}}\rangle$ with $n = 1, 2, \dots, N$. The general Wilson line for the isolated set of bands along path γ is defined as [35]

$$W(\gamma) = \lim_{L \rightarrow +\infty} \langle u_{\mathbf{k}_0} | P_{\mathbf{k}_1} P_{\mathbf{k}_2} \cdots P_{\mathbf{k}_{L-1}} P_{\mathbf{k}_L} | u_{\mathbf{k}_L} \rangle, \quad (3)$$

where $P_{\mathbf{k}} = \sum_n |u_{n, \mathbf{k}}\rangle \langle u_{n, \mathbf{k}}|$, $\mathbf{k}_0, \mathbf{k}_1, \mathbf{k}_2, \dots, \mathbf{k}_L$ are sequential momenta on γ with \mathbf{k}_0 at the beginning and \mathbf{k}_L at the end. Eq. (3) becomes a WL when γ forms a loop. Meanwhile, the non-abelian Berry curvature for the isolated set of bands reads

$$F_{\mathbf{k}} = \partial_{k_x} A_{\mathbf{k}, y} - \partial_{k_y} A_{\mathbf{k}, x} - i[A_{\mathbf{k}, x}, A_{\mathbf{k}, y}], \quad (4)$$

where

$$[A_{\mathbf{k}, i}]_{mn} = i \langle u_{m, \mathbf{k}} | \partial_{k_i} | u_{n, \mathbf{k}} \rangle. \quad (5)$$

The first step of the proof is to relate the WL to the non-abelian Berry curvature, which can be done through

a known theorem called non-abelian Stokes theorem [43]. The theorem holds for any simply connected region D ; but for simplicity, we, in the main text, consider a rectangular region $D = [0, K_x] \times [0, K_y]$, and we label the boundary (counter-clockwise) of D as ∂D . For the WL along ∂D (labeled as $W(\partial D)$ according to Eq. (3)) with starting momentum $\mathbf{k}_0 = (0, 0)$, the non-abelian Stokes theorem states that

$$\begin{aligned} W(\partial D) &= \mathcal{P} \exp \left[-i \int_D \tilde{F}_{\mathbf{k}} dk_x dk_y \right] \\ &= \prod_{i=L_1-1, \dots, 1, 0} \prod_{j=0, 1, \dots, L_2-1} \exp \left[-i \tilde{F}_{\left(\frac{iK_x}{L_1}, \frac{jK_y}{L_2}\right)} \frac{K_x}{L_1} \frac{K_y}{L_2} \right], \end{aligned} \quad (6)$$

where L_1 and L_2 limit to infinity, \mathcal{P} is the path ordering that moves larger k_x and smaller k_y to the left as shown by the second equality. $\tilde{F}_{\mathbf{k}}$ is the dressed non-abelian Berry curvature

$$\tilde{F}_{\mathbf{k}} = h_{\mathbf{k}} F_{\mathbf{k}} h_{\mathbf{k}}^{-1}, \quad (7)$$

where $h_{\mathbf{k}} = W_{\mathbf{k}_0 \rightarrow (k_x, 0)} W_{(k_x, 0) \rightarrow \mathbf{k}}$ with $\mathbf{k}_0 \rightarrow \mathbf{k}_1$ denoting the straight path from \mathbf{k}_0 to \mathbf{k}_1 .

To understand why Eq. (6) uses the dressed non-abelian Berry curvature $\tilde{F}_{\mathbf{k}}$ instead of the non-abelian Berry curvature $F_{\mathbf{k}}$, let us first note that the WL along the boundary of an infinitesimal plaquette is just the non-abelian Berry curvature (to leading order in $dk_x dk_y$) as shown in Fig. 1(a). Specifically, given an infinitesimal plaquette $\hat{D}_{(k_x, k_y)} = [k_x, k_x + dk_x] \times [k_y, k_y + dk_y]$, we have

$$W_{\partial\hat{D}_{(k_x, k_y)}} = \exp[-iF_{(k_x, k_y)} dk_x dk_y]. \quad (8)$$

Then, if we want to directly add the non-abelian Berry curvature for two nearby plaquettes, say $\hat{D}_{(k_x, k_y)}$ and $\hat{D}_{(k_x, k_y + dk_y)}$, we would have

$$\begin{aligned} &\exp[-iF_{(k_x, k_y)} dk_x dk_y] \exp[-iF_{(k_x, k_y + dk_y)} dk_x dk_y] \\ &= W_{\partial\hat{D}_{(k_x, k_y)}} W_{\partial\hat{D}_{(k_x, k_y + dk_y)}}, \end{aligned} \quad (9)$$

which does not necessarily equal to the Wilson loop around $\partial[\hat{D}_{(k_x, k_y)} \cup \hat{D}_{(k_x, k_y + dk_y)}]$ (Fig. 1(b)), a general feature of its non-abelian nature. This means that if we multiply $\exp[-iF_{(k_x, k_y)} dk_x dk_y]$ for all plaquettes in D , no matter how we select the path, we cannot necessarily arrive at the $W_{\partial D}$. The Wilson line dressing in Eq. (7) resolve this issue. With the dressing, multiplying $\exp[-i\tilde{F}_{(k_x, k_y)} dk_x dk_y]$ for two nearby plaquettes yields

$$\begin{aligned} &\exp[-i\tilde{F}_{(k_x, k_y)} dk_x dk_y] \exp[-i\tilde{F}_{(k_x, k_y + dk_y)} dk_x dk_y] \\ &= h_{(k_x, k_y)} W_{\partial[\hat{D}_{(k_x, k_y)} \cup \hat{D}_{(k_x, k_y + dk_y)}]} h_{(k_x, k_y)}^{-1}, \end{aligned} \quad (10)$$

which contains the Wilson loop along $\partial \left[\hat{D}_{(k_x, k_y)} \cup \hat{D}_{(k_x, k_y + dk_y)} \right]$, as shown in Fig. 1(c,d). As we multiply the $\exp \left[-i \tilde{F}_{(k_x, k_y)} dk_x dk_y \right]$ for all plaquettes in D , the dressing fully cancels, and leads to Eq. (6). The detailed proof is presented in App. [B1].

With the non-abelian Stokes theorem, we now consider the region D_s that continuously and monotonically depends on $s \in [0, s_f]$ with D_0 having zero area. Here by ‘‘monotonically’’, we require that $D_s \subset D_{s'}$ for any $s \leq s'$. The WL $W(\partial D_s)$ now depends continuously on s and has eigenvalues $e^{i\lambda_{i,s}}$ with $i = 1, 2, \dots, N$; we also require the starting point \mathbf{k}_0 of $W(\partial D_s)$ is the same for all $s \in [0, s_f]$. Without loss of generality, we will always choose $\lambda_{i,s}$ to continuously depend on s . We can then prove (in App. [C]) that

$$\begin{aligned} \int_0^{s_f} ds \sum_i \left| \frac{d\lambda_{i,s}}{ds} \right| &\leq \int_0^{s_f} ds \rho[-i\partial_s \log W(\partial D_s)] \\ &\leq \int_{D_{s_f}} d^2k \rho(F_{\mathbf{k}}), \end{aligned} \quad (11)$$

where $\rho(A)$ is the sum of the absolute values of all eigenvalues of the matrix A , which is called the Schatten 1-norm [47], and we have used the fact that $\tilde{F}_{\mathbf{k}}$ and $F_{\mathbf{k}}$ have the same eigenvalues. The proof of Eq. (11) exploits the properties (especially the triangle inequality) of the Schatten 1-norm, which is elaborated in App. [C]. Since $\rho(F_{\mathbf{k}}) \leq \text{Tr}[g_{\mathbf{k}}]$ as shown in App. [C], we arrive at the WL lower bound of the quantum metric

$$\mathcal{N} \equiv \int_0^{s_f} ds \sum_i \left| \frac{d\lambda_{i,s}}{ds} \right| \leq \int_{D_{s_f}} d^2k \text{Tr}[g_{\mathbf{k}}]. \quad (12)$$

We refer to \mathcal{N} defined in Eq. (12) as the ‘‘absolute WL winding’’, because it counts the winding of the WL eigenvalues without cancellation, as shown in Fig. 2(a). We emphasize that the absolute WL winding does not depend on how we rank $\lambda_{i,s}$ as long as they are continuous; it is because $\lambda_{i,s}$ generally touch or cross at measure-zero points of s which can be neglected for the integral. Naturally, the absolute WL winding is no smaller than the absolute value of the total WL winding, which is $\left| \int_0^{s_f} ds \sum_i \frac{d\lambda_{i,s}}{ds} \right|$.

One special yet useful case for the WL winding bound is when $D_s = \{k_1 \mathbf{b}_1 / (2\pi) + k_2 \mathbf{k}_2 / (2\pi) | k_1 \in [0, s], k_2 \in [0, 2\pi]\}$. Here \mathbf{b}_1 and \mathbf{b}_2 are two basis reciprocal lattice vectors, which means $D_{2\pi} = \text{BZ}$ is the entire first Brillouin zone. In this case, we can define a new WL along k_2 at a fixed k_1 :

$$\mathcal{W}_{k_1} = W(k_1 \mathbf{b}_1 / (2\pi) \rightarrow k_1 \mathbf{b}_1 / (2\pi) + \mathbf{b}_2). \quad (13)$$

Then, the absolute winding of $W(\partial D_s)$ in Eq. (12) can be directly replaced by the absolute winding of $\mathcal{W}_s = \mathcal{W}_{k_1=s}$, resulting in

$$\int_0^{s_f} ds \sum_l \left| \frac{d\phi_l(s)}{ds} \right| \leq \int_{D_{s_f}} d^2k \text{Tr}[g_{\mathbf{k}}], \quad (14)$$

where $\phi_l(s)$ is the phase of the l th eigenvalue of \mathcal{W}_s and is always chosen to be continuous. The validity of the replacement comes from the fact that

$$\begin{aligned} W(\partial D_s) &= W(\Gamma \rightarrow s\mathbf{b}_1 / (2\pi)) \mathcal{W}_s W^\dagger(\Gamma \rightarrow s\mathbf{b}_1 / (2\pi)) \\ &\quad \times W(\mathbf{b}_2 \rightarrow \Gamma) \end{aligned} \quad (15)$$

with $\Gamma = (0, 0)$, which can be directly derived from the definition of $W(\partial D_s)$. As $W(\mathbf{b}_2 \rightarrow \Gamma)$ is independent of s , we have

$$\begin{aligned} \rho[-i\partial_s \log W(\partial D_s)] &= \rho[-i\partial_s \log W(\Gamma \rightarrow s\mathbf{b}_1 / (2\pi)) \mathcal{W}_s W^{-1}(\Gamma \rightarrow s\mathbf{b}_1 / (2\pi))] \\ &\geq \int_0^{s_f} ds \sum_l \left| \frac{d\phi_l(s)}{ds} \right|, \end{aligned} \quad (16)$$

where we have made use of the fact that $W(\Gamma \rightarrow s\mathbf{b}_1 / (2\pi)) \mathcal{W}_s W^\dagger(\Gamma \rightarrow s\mathbf{b}_1 / (2\pi))$ has the same eigenvalues as \mathcal{W}_s . Combined with Eq. (12), we arrive at Eq. (14).

From the WL lower bound of the quantum metric in Eq. (14), we can directly derive the known Chern bound [27] by choosing $s_f = 2\pi$. In this case, the total WL winding of \mathcal{W}_s is just $2\pi \text{Ch}$ (with Ch the Chern number), which directly gives the Chern bound $|\text{Ch}| \leq \frac{1}{2\pi} \int_{\text{BZ}} \text{Tr}[g_{\mathbf{k}}] dk_x dk_y$. For Euler number $e_2 \in \mathbb{Z}$ defined for an isolated set of two bands with combination of time-reversal and inversion (or two-fold rotational) symmetries, the absolute WL winding would be no smaller than $4\pi|e_2|$, as we can always choose a Chern gauge where we can view the two components of the basis as two Chern bands with Chern numbers $\pm e_2$. Therefore, we can reproduce the known Euler bound [16, 19, 28, 29], which is $2|e_2| \leq \frac{1}{2\pi} \int_{\text{BZ}} \text{Tr}[g_{\mathbf{k}}] dk_x dk_y$.

III. Z_2 BOUND OF THE QUANTUM METRIC

Besides reproducing the known Chern and Euler bounds, Eq. (12) can also provide the TR-protected Z_2 bound of the quantum metric, which has remained an open question up to now.

To prove the bound, let us first review how to calculate TR-protected Z_2 index from WL winding. The setup is an isolated set of $2N$ bands of a non-interacting system with non-negligible spin-orbit coupling. The number of bands is necessarily even due to Kramers degeneracy of spinful TR symmetry \mathcal{T} . To calculate the Z_2 index of the isolated set of bands, we will use the WL \mathcal{W}_{k_1} in Eq. (13), with eigenvalues $e^{i\phi_l(k_1)}$ with $l = 1, 2, \dots, 2N$. Without loss of generality, we can choose $\phi_l(k_1)$ to be continuous for $k_1 \in [0, \pi]$ (see Fig. 2(b) for an example and see App. [D] for details). We can fix $\phi_l(0) \in [0, 2\pi)$. In this case, Z_2 reads [35]

$$Z_2 = \sum_{l=1}^{2N} M_l \text{ mod } 2, \quad (17)$$




FIG. 2. (a) is a schematic plot of a 2-band WL $W(\partial D_s)$, where D_s is a disk-like region that continuously depends on $s \in [0, s_f]$ with D_0 an measure-zero region (such as a point). The absolute WL winding of $W(\partial D_s)$ is equal to $|\lambda_{2,s_1} - \lambda_{2,0}| + |\lambda_{2,s_f} - \lambda_{2,s_1}| + |\lambda_{1,s_f} - \lambda_{1,0}|$. (b) is a schematic plot of a 2-band WL $W(k_1)$ in Eq. (13) for $Z_2 = 1$. The absolute WL winding of $W(k_1)$ is equal to $|\phi_2(p_1) - \phi_2(0)| + |\phi_2(p_2) - \phi_2(p_1)| + |\phi_2(\pi) - \phi_2(p_2)| + |\phi_1(\pi) - \phi_1(0)|$, which is no smaller than $\phi_2(\pi) - \phi_1(\pi) = 2\pi Z_2$.

where $2\pi M_l = \phi_l(\pi) - [\phi_l(\pi) \bmod 2\pi]$ with $[x \bmod 2\pi] \in [0, 2\pi)$ and $M_l \in \mathbb{Z}$.

Now we discuss how Z_2 bounds the quantum geometry from below. As exemplified in Fig. 2(b) and elaborated in App. [D], the Z_2 index in Eq. (17) suggests that the absolute WL winding of W_{k_1} over $k_1 \in [0, \pi]$ must be no smaller than $2\pi Z_2$, *i.e.*,

$$\frac{1}{2\pi} \int_0^\pi dk_1 \sum_l \left| \frac{d\phi_l(k_1)}{dk_1} \right| \geq Z_2, \quad (18)$$

where $\phi_l(k_1)$ is the phase of the l th eigenvalue of W_{k_1} , and we choose $\phi_l(k_1)$ to be continuous. Combined with Eq. (14) and TR symmetry, we arrive at the inequality in Eq. (2). This inequality can be saturated, as two TR-related lowest Landau levels with opposite spins serve as one example.

Eq. (2) also holds for the PH-protected Z_2 index, where the PH symmetry is antiunitary and squares to -1 . It comes from the equivalence between the TR- Z_2 and PH- Z_2 indices, which was pointed out in [46], as reviewed in App. [D].

IV. PHYSICAL CONSEQUENCES OF THE Z_2 BOUND

The TR Z_2 bound can provide bounds on various physical quantities. The first example is the superfluid weight of 2D superconductors. Consider a set of isolated two normal-state bands that preserve TR symmetry. Assuming a mean-field treatment in the flat-band limit where the pairing potential Δ is uniform over the BZ (see App. [E1]), the superfluid weight is proportional to the minimal integrated Fubini-Study metric [12, 48, 49].

Hence the Z_2 bound yields

$$\text{Tr}[D_{SW}(T=0)] \geq \frac{4|\Delta|}{\pi} \sqrt{f(1-f)} \frac{Z_2}{\pi}, \quad (19)$$

where $D_{SW}(T=0)$ is the zero-temperature superfluid weight tensor, $|\Delta|$ is the strength of the pairing, and $f \in [0, 1]$ is the occupation fraction of the two bands ($f = 1$ means the two bands are fully filled). Here and henceforth we take $\hbar = e = 1$ with electron having charge $-e$.

Besides superfluid weight, the optical conductivity of a band insulator has been shown to be related to its quantum geometry [5]

$$\int_0^\infty \frac{d\omega}{\omega} \sum_i \text{Re}[\sigma_{ii}(\omega + i0^+)] = \frac{1}{4\pi} \int d^2k \text{Tr}[g(\mathbf{k})], \quad (20)$$

where $\sigma_{ij}(\omega + i0^+)$ is the optical conductivity tensor, and 0^+ is an infinitesimal real positive number. Combined with the sum rule derived from the Kubo formula [50], it has been shown that the direct band gap E_g is bounded above by the quantum metric [51, 52]

$$E_g \leq \frac{n\pi}{m} \frac{1}{\frac{1}{4\pi} \int d^2k \text{Tr}[g(\mathbf{k})]}, \quad (21)$$

where n is the electron density, and m is the electron mass. Combining these known results with the TR Z_2 bound in Eq. (2), we can immediately arrive at the lower bound of integrated optical conductivity and the upper bound of the direct band gap for a 2D TR-invariant band insulator, *i.e.*,

$$\int_0^\infty \frac{d\omega}{\omega} \sum_i \text{Re}[\sigma_{ii}(\omega + i0^+)] \geq Z_2$$

$$E_g \leq \frac{n\pi}{mZ_2}. \quad (22)$$

More details are given in App. [E2].

V. CONCLUSION

We show that the absolute WL winding bounds the quantum metric from below, which provides the TR- Z_2 and PH- Z_2 lower bound of the quantum metric, as well as recovering the previously known Chern and Euler bounds. As a result, the TR- Z_2 index provides lower bounds for superfluid weight and optical conductivity and an upper bound for the band gap of a band insulator. The WL bound is applicable to any topological invariant that can be characterized by WL winding.

VI. ACKNOWLEDGMENT

J. Y.'s work at University of Florida is supported by startup funds at University of Florida. J. Y.'s work at

Princeton University is supported by the Gordon and Betty Moore Foundation through Grant No. GBMF8685 towards the Princeton theory program. J. H.-A. is supported by a Hertz Fellowship, with additional support from DOE Grant No. DE-SC0016239. B.A.B. was supported by the Gordon and Betty Moore Foundation through Grant No. GBMF8685 towards the Princeton theory program, the Gordon and Betty Moore Foundation's EPiQS Initiative (Grant No. GBMF11070), the

Office of Naval Research (ONR Grant No. N00014-20-1-2303), the Global Collaborative Network Grant at Princeton University, the Simons Investigator Grant No. 404513, the BSF Israel US foundation No. 2018226, the NSF-MERSEC (Grant No. MERSEC DMR 2011750), the Simons Collaboration on New Frontiers in Superconductivity, and the Schmidt Foundation at the Princeton University.

-
- [1] J. Provost and G. Vallee, Riemannian structure on manifolds of quantum states, *Communications in Mathematical Physics* **76**, 289 (1980).
- [2] G. Fubini, Sulle metriche definite da una forma hermitiana, *Atti del Reale istituto Veneto di Scienze, Lettere ed Arti* **63**, 502 (1904).
- [3] E. Study, Kürzeste Wege im komplexen Gebiet, *Mathematische Annalen* **60**, 321 (1905).
- [4] I. Souza, T. Wilkens, and R. M. Martin, Polarization and localization in insulators: Generating function approach, *Physical Review B* **62**, 1666 (1999), [cond-mat/9911007](#).
- [5] I. Souza, T. Wilkens, and R. M. Martin, Polarization and localization in insulators: Generating function approach, *Phys. Rev. B* **62**, 1666 (2000).
- [6] R. Resta, Polarization Fluctuations in Insulators and Metals: New and Old Theories Merge, *Physical Review Letters* **96**, 137601 (2006), [cond-mat/0512247](#).
- [7] R. M. Martin, *Electronic Structure* (2004).
- [8] C. Aebischer, D. Baeriswyl, and R. M. Noack, Dielectric Catastrophe at the Mott Transition, *Physical Review Letters* **86**, 468 (2001), [cond-mat/0006354](#).
- [9] N. Verma and R. Queiroz, Instantaneous Response and Quantum Geometry of Insulators, [arXiv 10.48550/arxiv.2403.07052](#) (2024), [2403.07052](#).
- [10] Y. Onishi and L. Fu, Universal relation between energy gap and dielectric constant, [arXiv 10.48550/arxiv.2401.04180](#) (2024), [2401.04180](#).
- [11] I. Souza, R. M. Martin, and M. Stengel, Optical bounds on many-electron localization, [arXiv 10.48550/arxiv.2407.17908](#) (2024), [2407.17908](#).
- [12] S. Peotta and P. Törmä, Superfluidity in topologically nontrivial flat bands, *Nature Communications* **6**, 8944 (2015).
- [13] A. Julku, S. Peotta, T. I. Vanhala, D.-H. Kim, and P. Törmä, Geometric origin of superfluidity in the Lieb-lattice flat band, *Phys. Rev. Lett.* **117**, 045303 (2016).
- [14] L. Liang, T. I. Vanhala, S. Peotta, T. Siro, A. Harju, and P. Törmä, Band geometry, berry curvature, and superfluid weight, *Phys. Rev. B* **95**, 024515 (2017).
- [15] X. Hu, T. Hyart, D. I. Pikulin, and E. Rossi, Geometric and conventional contribution to the superfluid weight in twisted bilayer graphene, *Phys. Rev. Lett.* **123**, 237002 (2019).
- [16] F. Xie, Z. Song, B. Lian, and B. A. Bernevig, Topology-bounded superfluid weight in twisted bilayer graphene, *Phys. Rev. Lett.* **124**, 167002 (2020).
- [17] A. Julku, T. J. Peltonen, L. Liang, T. T. Heikkilä, and P. Törmä, Superfluid weight and berezinskii-kosterlitz-thouless transition temperature of twisted bilayer graphene, *Phys. Rev. B* **101**, 060505 (2020).
- [18] E. Rossi, Quantum metric and correlated states in two-dimensional systems, *Current Opinion in Solid State and Materials Science* **25**, 100952 (2021).
- [19] J. Yu, M. Xie, F. Wu, and S. Das Sarma, Euler-obstructed nematic nodal superconductivity in twisted bilayer graphene, *Phys. Rev. B* **107**, L201106 (2023).
- [20] P. Törmä, Essay: Where Can Quantum Geometry Lead Us?, *Physical Review Letters* **131**, 240001 (2023).
- [21] H. Tian, X. Gao, Y. Zhang, S. Che, T. Xu, P. Cheung, K. Watanabe, T. Taniguchi, M. Randeria, F. Zhang, C. N. Lau, and M. W. Bockrath, Evidence for dirac flat band superconductivity enabled by quantum geometry, *Nature* **614**, 440 (2023).
- [22] J. Yu, C. J. Ciccarino, R. Bianco, I. Errea, P. Narang, and B. A. Bernevig, Non-trivial quantum geometry and the strength of electron-phonon coupling, *Nature Physics* **20**, 1262 (2024).
- [23] P. Zhu and A. Alexandradinata, Anomalous shift and optical vorticity in the steady photovoltaic current, *Phys. Rev. B* **110**, 115108 (2024).
- [24] P. M. Tam, J. Herzog-Arbeitman, and J. Yu, Quantum geometry and entanglement in two-dimensional insulators: A view from the corner charge fluctuation, [arXiv preprint arXiv:2406.17023](#) (2024).
- [25] X.-C. Wu, K.-L. Cai, M. Cheng, and P. Kumar, *Corner charge fluctuations and many-body quantum geometry* (2024), [arXiv:2408.16057 \[cond-mat.str-el\]](#).
- [26] J. Yu, B. A. Bernevig, R. Queiroz, E. Rossi, P. Törmä, and B.-J. Yang, To appear.
- [27] R. Roy, Band geometry of fractional topological insulators, *Physical Review B* **90**, 165139 (2014).
- [28] S. Kwon and B.-J. Yang, Quantum geometric bound and ideal condition for euler band topology, *Phys. Rev. B* **109**, L161111 (2024).
- [29] W. J. Jankowski, A. S. Morris, A. Bouhon, F. N. Ünal, and R.-J. Slager, *Optical manifestations of topological euler class* (2024), [arXiv:2311.07545 \[cond-mat.mes-hall\]](#).
- [30] J. Herzog-Arbeitman, V. Peri, F. Schindler, S. D. Huber, and B. A. Bernevig, Superfluid weight bounds from symmetry and quantum geometry in flat bands, *Phys. Rev. Lett.* **128**, 087002 (2022).
- [31] C. L. Kane and E. J. Mele, Z_2 topological order and the quantum spin hall effect, *Phys. Rev. Lett.* **95**, 146802 (2005).
- [32] B. A. Bernevig and S.-C. Zhang, Quantum spin hall effect, *Phys. Rev. Lett.* **96**, 106802 (2006).
- [33] C. L. Kane and E. J. Mele, Quantum spin hall effect in graphene, *Phys. Rev. Lett.* **95**, 226801 (2005).
- [34] B. A. Bernevig, T. L. Hughes, and S.-C. Zhang, Quantum spin hall effect and topological phase transition in hgte

- quantum wells, *Science* **314**, 1757 (2006).
- [35] R. Yu, X. L. Qi, A. Bernevig, Z. Fang, and X. Dai, Equivalent expression of z_2 topological invariant for band insulators using the non-abelian berry connection, *Phys. Rev. B* **84**, 075119 (2011).
- [36] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs, Quantized hall conductance in a two-dimensional periodic potential, *Phys. Rev. Lett.* **49**, 405 (1982).
- [37] A. Alexandradinata, X. Dai, and B. A. Bernevig, Wilson-loop characterization of inversion-symmetric topological insulators, *Phys. Rev. B* **89**, 155114 (2014).
- [38] J. Ahn, D. Kim, Y. Kim, and B.-J. Yang, Band topology and linking structure of nodal line semimetals with Z_2 monopole charges, *Phys. Rev. Lett.* **121**, 106403 (2018).
- [39] Z. Song, Z. Wang, W. Shi, G. Li, C. Fang, and B. A. Bernevig, All magic angles in twisted bilayer graphene are topological, *Phys. Rev. Lett.* **123**, 036401 (2019).
- [40] J. Ahn, S. Park, and B.-J. Yang, Failure of nielsen-ninomiya theorem and fragile topology in two-dimensional systems with space-time inversion symmetry: Application to twisted bilayer graphene at magic angle, *Phys. Rev. X* **9**, 021013 (2019).
- [41] A. Bouhon, A. M. Black-Schaffer, and R.-J. Slager, Wilson loop approach to fragile topology of split elementary band representations and topological crystalline insulators with time-reversal symmetry, *Phys. Rev. B* **100**, 195135 (2019).
- [42] B. Bradlyn, Z. Wang, J. Cano, and B. A. Bernevig, Disconnected elementary band representations, fragile topology, and wilson loops as topological indices: An example on the triangular lattice, *Phys. Rev. B* **99**, 045140 (2019).
- [43] I. Y. Aref'eva, Non-abelian stokes formula, *Theoretical and Mathematical Physics* **43**, 353 (1980).
- [44] L. Fu and C. L. Kane, Time reversal polarization and a Z_2 adiabatic spin pump, *Phys. Rev. B* **74**, 195312 (2006).
- [45] A. A. Soluyanov and D. Vanderbilt, Wannier representation of z_2 topological insulators, *Phys. Rev. B* **83**, 035108 (2011).
- [46] Z.-D. Song, B. Lian, N. Regnault, and B. A. Bernevig, Twisted bilayer graphene. ii. stable symmetry anomaly, *Phys. Rev. B* **103**, 205412 (2021).
- [47] R. Bhatia, *Positive definite matrices* (Princeton university press, 2009).
- [48] P. Törmä, S. Peotta, and B. A. Bernevig, Superconductivity, superfluidity and quantum geometry in twisted multilayer systems, *Nature Reviews Physics* **4**, 528 (2022).
- [49] K.-E. Huhtinen, J. Herzog-Arbeitman, A. Chew, B. A. Bernevig, and P. Törmä, Revisiting flat band superconductivity: Dependence on minimal quantum metric and band touchings, *Phys. Rev. B* **106**, 014518 (2022).
- [50] R. Kubo, Statistical-mechanical theory of irreversible processes. i. general theory and simple applications to magnetic and conduction problems, *Journal of the physical society of Japan* **12**, 570 (1957).
- [51] S. Kivelson, Wannier functions in one-dimensional disordered systems: Application to fractionally charged solitons, *Phys. Rev. B* **26**, 4269 (1982).
- [52] Y. Onishi and L. Fu, Fundamental bound on topological gap, arXiv [10.48550/arxiv.2306.00078](https://arxiv.org/abs/10.48550/arxiv.2306.00078) (2023), [2306.00078](https://arxiv.org/abs/2306.00078).
- [53] T. Fukui, Y. Hatsugai, and H. Suzuki, Chern numbers in discretized brillouin zone: efficient method of computing (spin) hall conductances, *Journal of the Physical Society of Japan* **74**, 1674 (2005).
- [54] M. Tovmasyan, S. Peotta, P. Törmä, and S. D. Huber, Effective theory and emergent $su(2)$ symmetry in the flat bands of attractive hubbard models, *Physical Review B* **94**, 245149 (2016).
- [55] J. Herzog-Arbeitman, A. Chew, K.-E. Huhtinen, P. Törmä, and B. A. Bernevig, Many-body superconductivity in topological flat bands, arXiv preprint arXiv:2209.00007 (2022).

CONTENTS

I. Introduction	1
II. General Wilson Loop Bound of the quantum Metric	1
III. Z_2 Bound of the quantum Metric	3
IV. Physical Consequences of the Z_2 Bound	4
V. Conclusion	4
VI. Acknowledgment	4
References	5
A. Quantum Geometric Tensor, Quantum Metric, and Berry Curvature	7
B. Mathematical Concepts	8
1. Non-abelian Stokes Theorem	8
2. Schatten Norms	11
a. General Definition	11
b. Schatten 1-norm for Hermitian Matrices	12
C. Wilson loop Lower Bound of the quantum Metric	13
D. Z_2 Lower Bound of the quantum Metric	15
1. Review of TR Z_2 index from Wilson Loop	15
2. Review of PH Z_2 index	16
3. Proof of the Z_2 Lower Bound of the quantum Metric	16
E. Physical Consequences of the TR Z_2 Bound	17
1. TR Z_2 Lower Bound for Superfluid Weight	17
2. Z_2 Bounds for Optical Conductivity and Band Gap	22

Appendix A: Quantum Geometric Tensor, Quantum Metric, and Berry Curvature

In this appendix, we review the key concepts used in our work following Ref. [26]. Consider an isolated set of N bands, and we label the periodic part of the Bloch states as $|u_{\mathbf{k},n}\rangle$ with $n = 1, 2, \dots, N$. The quantum geometric tensor for the set of bands reads

$$[Q_{ij}(\mathbf{k})]_{mn} = \langle \partial_i u_{\mathbf{k},m} | (1 - P_{\mathbf{k}}) | \partial_j u_{\mathbf{k},n} \rangle , \quad (\text{A1})$$

where $P_{\mathbf{k}} = \sum_n |u_{\mathbf{k},n}\rangle \langle u_{\mathbf{k},n}|$, and $\partial_i = \partial_{k_i}$. The non-abelian Berry curvature reads

$$[F_{ij,\mathbf{k}}]_{mn} = i[Q_{ij}(\mathbf{k}) - Q_{ji}(\mathbf{k})]_{mn} , \quad (\text{A2})$$

and the non-abelian quantum metric reads

$$[G_{ij,\mathbf{k}}]_{mn} = \frac{1}{2} [Q_{ij}(\mathbf{k}) + Q_{ji}(\mathbf{k})]_{mn} . \quad (\text{A3})$$

Both $F_{ij,\mathbf{k}}$ and $G_{ij,\mathbf{k}}$ are Hermitian matrix for fixed i, j :

$$F_{ij,\mathbf{k}}^\dagger = F_{ij,\mathbf{k}} , \quad G_{ij,\mathbf{k}}^\dagger = G_{ij,\mathbf{k}} . \quad (\text{A4})$$

The quantum metric reads

$$[g_{\mathbf{k}}]_{ij} = \text{Tr}[G_{ij,\mathbf{k}}] = \frac{1}{2} \text{Tr} [(\partial_i P_{\mathbf{k}}) \partial_j P_{\mathbf{k}}] . \quad (\text{A5})$$

$[Q_{ij}(\mathbf{k})]_{mn}$ is positive semi-definite. To see this, let us consider a generic U_{im} , which gives

$$\sum_{ij,mn} [Q_{ij}(\mathbf{k})]_{mn} U_{im}^* U_{jn} = \sum_{ij,mn} U_{im}^* \langle \partial_i u_{\mathbf{k},m} | (1 - P_{\mathbf{k}}) | \partial_j u_{\mathbf{k},n} \rangle U_{jn} = \langle X | (1 - P_{\mathbf{k}}) | X \rangle , \quad (\text{A6})$$

where $|X\rangle = \sum_{jn} |\partial_j u_{\mathbf{k},n}\rangle U_{jn}$. Since $1 - P_{\mathbf{k}}$ only has nonnegative eigenvalues, we know

$$\sum_{ij,mn} [Q_{ij}(\mathbf{k})]_{mn} U_{im}^* U_{jn} \geq 0 \quad (\text{A7})$$

for any U_{jn} , meaning that $[Q_{ij}(\mathbf{k})]_{mn}$ is positive semi-definite.

The positive semi-definiteness of the quantum geometric tensor provides an inequality between quantum metric and the non-abelian Berry curvature. To show it, let us choose U_{im} in Eq. (A7) as $U_{im} = (\delta_{i,i_0} + \text{si}\delta_{i,j_0})v_m$ for $s = \pm 1$ and arbitrary v_m . In this case, we have

$$\begin{aligned} \sum_{mn} v_m^* [Q_{i_0 i_0}(\mathbf{k}) + Q_{j_0 j_0}(\mathbf{k}) + \text{si}Q_{i_0 j_0}(\mathbf{k}) - \text{si}Q_{j_0 i_0}(\mathbf{k})]_{mn} v_n &\geq 0 \\ \Rightarrow v^\dagger [G_{i_0 i_0, \mathbf{k}} + G_{j_0 j_0, \mathbf{k}} + sF_{i_0 j_0, \mathbf{k}}] v &\geq 0 \end{aligned} \quad (\text{A8})$$

for any i_0, j_0 and vector v . In 2D, we define

$$F_{\mathbf{k}} = F_{xy, \mathbf{k}} , \quad (\text{A9})$$

which gives

$$v^\dagger \left[\sum_i G_{ii, \mathbf{k}} + sF_{\mathbf{k}} \right] v \geq 0 \Rightarrow v^\dagger \left[\sum_i G_{ii, \mathbf{k}} \right] v \geq |v^\dagger F_{\mathbf{k}} v| \quad (\text{A10})$$

for any vector v . Eventually, we arrive at, in 2D,

$$\text{Tr}[g_{\mathbf{k}}] \geq \sum_l |v_l^\dagger F_{\mathbf{k}} v_l| \quad (\text{A11})$$

for any choice of a complete set of orthonormal v_l . A special choice of v_l is the eigenbasis of $F_{\mathbf{k}}$.

Appendix B: Mathematical Concepts

In this appendix, we discuss two mathematical concepts: non-abelian Stokes theorem and Schatten Norms.

1. Non-abelian Stokes Theorem

In this part, we review the non-abelian Stokes theorem following [43]. Before considering the generic case, let us first discuss the case of a rectangular region D parametrized by (x, y) with $x \in [0, X]$ and $y \in [0, Y]$. Suppose we have N vector fields (just like the periodic parts of N Bloch states) defined on D , labeled as $v_{n,x,y}$ with $n = 1, 2, 3, \dots, N$. Here we require $v_{n,x,y}^\dagger v_{n',x,y} = \delta_{nn'}$, and we use x, y instead of \mathbf{k} to show that the derivation holds for any parametrization x, y , not just the Bloch momentum. The corresponding projector reads

$$P_{x,y} = v_{x,y} v_{x,y}^\dagger , \quad (\text{B1})$$

where

$$v_{x,y} = (v_{1,x,y} \ v_{2,x,y} \ v_{3,x,y} \ \cdots \ v_{N,x,y}) . \quad (\text{B2})$$

Clearly, we can define the Wilson line for v , which reads

$$W(\gamma) = \lim_{L \rightarrow \infty} v_{x_0, y_0}^\dagger P_{x_1, y_1} P_{x_2, y_2} P_{x_3, y_3} \cdots P_{x_{L-1}, y_{L-1}} v_{x_L, y_L} , \quad (\text{B3})$$

where $(x_0, y_0), (x_1, y_1), \dots, (x_{L-1}, y_{L-1}), (x_L, y_L)$ are arranged sequentially on the path γ , and (x_0, y_0) and (x_L, y_L) are the initial and final points of γ , respectively. We can also define the non-abelian Berry connection

$$\mathbf{A}(x, y) = i(v_{x,y}^\dagger \partial_x v_{x,y}, v_{x,y}^\dagger \partial_y v_{x,y}) \quad (\text{B4})$$

and the non-abelian Berry curvature

$$F_{x,y} = \partial_x A_2(x, y) - \partial_y A_1(x, y) - i[A_1(x, y), A_2(x, y)] . \quad (\text{B5})$$

To prove the relation between the Wilson loop and non-abelian Berry curvature, let us split the $[0, X]$ into L_1 equal parts and $[0, Y]$ into L_2 equal parts, resulting in area into $L_1 \times L_2$ rectangular plaquettes for D . A generic plaquette is labeled as

$$\tilde{D}_{x_i, y_j} = \{(x, y) | x \in [x_i, x_i + dx], y \in [y_j, y_j + dy]\} , \quad (\text{B6})$$

where $x_i = iX/L_1$ with $i = 0, 1, 2, \dots, L_1 - 1$, and $y_j = jY/L_2$ with $j = 0, 1, 2, \dots, L_2 - 1$, $dx = X/L_1$, and $dy = Y/L_2$. Eventually, we will take L_1 and L_2 to infinity, which means dx and dy are infinitesimal quantities.

For the infinitesimal \tilde{D}_{x_i, y_j} , the Wilson loop around the boundary of \tilde{D}_{x_i, y_j} is naturally connected to the non-abelian Berry curvature [53] via

$$\begin{aligned} w_{i,j} &= v_{x_i, y_j}^\dagger v_{x_i+dx, y_j} v_{x_i+dx, y_j}^\dagger v_{x_i+dx, y_j+dy} v_{x_i+dx, y_j+dy}^\dagger \left(v_{x_i, y_j+dy}^\dagger v_{x_i+dx, y_j+dy} \right)^{-1} \left(v_{x_i, y_j}^\dagger v_{x_i, y_j+dy} \right)^{-1} \\ &= \left[1 - iA_1(x_i, y_j)dx + \frac{1}{2} v_{x_i, y_j}^\dagger \partial_{x_i}^2 v_{x_i, y_j} dx^2 \right] \\ &\quad \times \left[1 - iA_2(x_i, y_j)dy - i\partial_{x_i} A_2(x_i, y_j)dx dy + \frac{1}{2} v_{x_i, y_j}^\dagger \partial_{y_j}^2 v_{x_i, y_j} dy^2 \right] \\ &\quad \times \left[1 - iA_1(x_i, y_j)dx - i\partial_{y_j} A_1(x_i, y_j)dx dy + \frac{1}{2} v_{x_i, y_j}^\dagger \partial_{x_i}^2 v_{x_i, y_j} dx^2 \right]^{-1} \\ &\quad \times \left[1 - iA_2(x_i, y_j)dy + \frac{1}{2} v_{x_i, y_j}^\dagger \partial_{y_j}^2 v_{x_i, y_j} dy^2 \right]^{-1} + O(dx^3, dx^2 dy, dx dy^2, dy^3) \\ &= 1 - iA_1(x_i, y_j)dx - iA_2(x_i, y_j)dy + iA_1(x_i, y_j)dx + iA_2(x_i, y_j)dy \\ &\quad - A_1(x_i, y_j)A_2(x_i, y_j)dx dy + A_1(x_i, y_j)A_2(x_i, y_j)dx dy \\ &\quad + A_2(x_i, y_j)A_1(x_i, y_j)dx dy - A_1(x_i, y_j)A_2(x_i, y_j)dx dy \\ &\quad - i\partial_{x_i} A_2(x_i, y_j)dx dy + i\partial_{y_j} A_1(x_i, y_j)dx dy + O(dx^3, dx^2 dy, dx dy^2, dy^3) \\ &= 1 - i\partial_{x_i} A_2(x_i, y_j)dx dy + i\partial_{y_j} A_1(x_i, y_j)dx dy - [A_1(x_i, y_j), A_2(x_i, y_j)]dx dy \\ &\quad + O(dx^3, dx^2 dy, dx dy^2, dy^3) \\ &= 1 - iF_{x_i, y_j} dx dy + O(dx^3, dx^2 dy, dx dy^2, dy^3) \\ &= \exp \left[-iF_{x_i, y_j} dx dy + O(dx^3, dx^2 dy, dx dy^2, dy^3) \right] , \end{aligned} \quad (\text{B7})$$

where we note that we have used the matrix inverse rather than the Hermitian conjugate in the first line.

To recover the non-abelian Berry curvature on the entire D from the infinitesimal result in Eq. (B7), one naïve way is to (matrix) multiply all $w_{i,j}$ together. But as discussed in the Main Text, the non-abelian nature is incompatible with this guess. The solution is to dress $w_{i,j}$:

$$\tilde{w}_{i,j} = h_{i,j} w_{i,j} h_{i,j}^{-1} , \quad (\text{B8})$$

where

$$h_{i,j} = v_{x_0, y_0} v_{x_0, y_0}^\dagger v_{x_1, y_0} v_{x_1, y_0}^\dagger v_{x_2, y_0} \cdots v_{x_{i-1}, y_0} v_{x_i, y_0} v_{x_i, y_0}^\dagger v_{x_i, y_1} v_{x_i, y_1}^\dagger v_{x_i, y_2} v_{x_i, y_2}^\dagger \cdots v_{x_i, y_{j-1}} v_{x_i, y_j} , \quad (\text{B9})$$

which has the following properties:

$$h_{i,j+1} = h_{i,j} v_{x_i, y_j}^\dagger v_{x_i, y_{j+1}} \quad (\text{B10})$$

and

$$h_{i+1,0} = h_{i,0} v_{x_i, y_0}^\dagger v_{x_{i+1}, y_0} . \quad (\text{B11})$$

Now we would like to derive the multiplication of all $\tilde{w}_{i,j}$'s step by step. We first consider the multiplication of $\tilde{w}_{i,j}$ and $\tilde{w}_{i,j+1}$:

$$\begin{aligned}
& \tilde{w}_{i,j} \tilde{w}_{i,j+1} \\
&= h_{i,j} v_{x_i, y_j}^\dagger v_{x_{i+1}, y_j} v_{x_{i+1}, y_{j+1}}^\dagger v_{x_{i+1}, y_{j+1}} \left(v_{x_i, y_{j+1}}^\dagger v_{x_{i+1}, y_{j+1}} \right)^{-1} \left(v_{x_i, y_j}^\dagger v_{x_{i+1}, y_{j+1}} \right)^{-1} h_{i,j}^{-1} \\
&\quad \times h_{i,j} v_{x_i, y_j}^\dagger v_{x_{i+1}, y_j} v_{x_{i+1}, y_{j+1}}^\dagger v_{x_{i+1}, y_{j+1}} v_{x_{i+1}, y_{j+1}}^\dagger v_{x_{i+1}, y_{j+1}} \left(v_{x_i, y_{j+2}}^\dagger v_{x_{i+1}, y_{j+2}} \right)^{-1} \left(v_{x_i, y_{j+1}}^\dagger v_{x_{i+1}, y_{j+2}} \right)^{-1} \left[v_{x_i, y_j}^\dagger v_{x_{i+1}, y_{j+1}} \right]^{-1} h_{i,j}^{-1} \\
&= h_{i,j} v_{x_i, y_j}^\dagger v_{x_{i+1}, y_j} \left(v_{x_{i+1}, y_j}^\dagger v_{x_{i+1}, y_{j+1}} v_{x_{i+1}, y_{j+1}}^\dagger v_{x_{i+1}, y_{j+2}} \right) \\
&\quad \times \left(v_{x_i, y_{j+2}}^\dagger v_{x_{i+1}, y_{j+2}} \right)^{-1} \left(v_{x_i, y_j}^\dagger v_{x_{i+1}, y_{j+1}} v_{x_{i+1}, y_{j+1}}^\dagger v_{x_{i+1}, y_{j+2}} \right)^{-1} h_{i,j}^{-1} ;
\end{aligned} \tag{B12}$$

then the multiplication of all $\tilde{w}_{i,j}$'s with fixed i reads

$$\begin{aligned}
\tilde{u}_i &= \tilde{w}_{i,0} \tilde{w}_{i,1} \cdots \tilde{w}_{i, L_1 - 1} \\
&= h_{i,0} \left[v_{x_i, y_0}^\dagger v_{x_{i+1}, y_0} \right] \left[v_{x_{i+1}, y_0}^\dagger v_{x_{i+1}, y_1} \cdots v_{x_{i+1}, y_{L_2-2}}^\dagger v_{x_{i+1}, y_{L_2-1}} v_{x_{i+1}, y_{L_2-1}}^\dagger v_{x_{i+1}, y_{L_2}} \right] \\
&\quad \times \left[v_{x_i, y_{L_2}}^\dagger v_{x_{i+1}, y_{L_2}} \right]^{-1} \left[v_{x_i, y_0}^\dagger v_{x_{i+1}, y_1} \cdots v_{x_i, y_{L_2-2}}^\dagger v_{x_{i+1}, y_{L_2-1}} v_{x_{i+1}, y_{L_2-1}}^\dagger v_{x_{i+1}, y_{L_2}} \right]^{-1} h_{i,0}^{-1} .
\end{aligned} \tag{B13}$$

Now we multiply \tilde{u}_i by \tilde{u}_{i+1} , leading to

$$\begin{aligned}
& \tilde{u}_{i+1} \tilde{u}_i \\
&= h_{i,0} v_{x_i, y_0}^\dagger v_{x_{i+1}, y_0} \left[v_{x_{i+1}, y_0}^\dagger v_{x_{i+2}, y_0} \right] \left[v_{x_{i+2}, y_0}^\dagger v_{x_{i+2}, y_1} \cdots v_{x_{i+2}, y_{L_2-2}}^\dagger v_{x_{i+2}, y_{L_2-1}} v_{x_{i+2}, y_{L_2-1}}^\dagger v_{x_{i+2}, y_{L_2}} \right] \\
&\quad \times \left[v_{x_{i+1}, y_{L_2}}^\dagger v_{x_{i+2}, y_{L_2}} \right]^{-1} \left[v_{x_{i+1}, y_0}^\dagger v_{x_{i+1}, y_1} \cdots v_{x_{i+1}, y_{L_2-2}}^\dagger v_{x_{i+1}, y_{L_2-1}} v_{x_{i+1}, y_{L_2-1}}^\dagger v_{x_{i+1}, y_{L_2}} \right]^{-1} \left[v_{x_i, y_0}^\dagger v_{x_{i+1}, y_0} \right]^{-1} h_{i,0}^{-1} \\
&\quad \times h_{i,0} \left[v_{x_i, y_0}^\dagger v_{x_{i+1}, y_0} \right] \left[v_{x_{i+1}, y_0}^\dagger v_{x_{i+1}, y_1} \cdots v_{x_{i+1}, y_{L_2-2}}^\dagger v_{x_{i+1}, y_{L_2-1}} v_{x_{i+1}, y_{L_2-1}}^\dagger v_{x_{i+1}, y_{L_2}} \right] \\
&\quad \times \left[v_{x_i, y_{L_2}}^\dagger v_{x_{i+1}, y_{L_2}} \right]^{-1} \left[v_{x_i, y_0}^\dagger v_{x_{i+1}, y_1} \cdots v_{x_i, y_{L_2-2}}^\dagger v_{x_{i+1}, y_{L_2-1}} v_{x_{i+1}, y_{L_2-1}}^\dagger v_{x_{i+1}, y_{L_2}} \right]^{-1} h_{i,0}^{-1} \\
&= h_{i,0} \left[v_{x_i, y_0}^\dagger v_{x_{i+1}, y_0} v_{x_{i+1}, y_0}^\dagger v_{x_{i+2}, y_0} \right] \left[v_{x_{i+2}, y_0}^\dagger v_{x_{i+2}, y_1} \cdots v_{x_{i+2}, y_{L_2-2}}^\dagger v_{x_{i+2}, y_{L_2-1}} v_{x_{i+2}, y_{L_2-1}}^\dagger v_{x_{i+2}, y_{L_2}} \right] \\
&\quad \times \left[v_{x_i, y_{L_2}}^\dagger v_{x_{i+1}, y_{L_2}} v_{x_{i+1}, y_{L_2}}^\dagger v_{x_{i+2}, y_{L_2}} \right]^{-1} \left[v_{x_i, y_0}^\dagger v_{x_{i+1}, y_1} \cdots v_{x_i, y_{L_2-2}}^\dagger v_{x_{i+1}, y_{L_2-1}} v_{x_{i+1}, y_{L_2-1}}^\dagger v_{x_{i+1}, y_{L_2}} \right]^{-1} h_{i,0}^{-1} .
\end{aligned} \tag{B14}$$

Then, we can define the path-ordered multiplication of all $\tilde{w}_{i,j}$'s:

$$\mathcal{P} \prod_{i,j} \tilde{w}_{i,j} = \prod_{i=L_1-1, \dots, 2, 1, 0} \prod_{j=0, 1, 2, \dots, L_2-1} \tilde{w}_{i,j} , \tag{B15}$$

which in the limit of $L_1, L_2 \rightarrow \infty$ reads

$$\begin{aligned}
& \lim_{L_1, L_2 \rightarrow \infty} \mathcal{P} \prod_{i,j} \tilde{w}_{i,j} \\
&= \tilde{u}_{L_1-1} \tilde{u}_{L_1-2} \cdots \tilde{u}_0 \\
&= \lim_{L_1, L_2 \rightarrow \infty} h_{0,0} \left[v_{x_0, y_0}^\dagger v_{x_1, y_0} \cdots v_{x_{L_1-2}, y_0}^\dagger v_{x_{L_1-1}, y_0} v_{x_{L_1-1}, y_0}^\dagger v_{x_{L_1}, y_0} \right] \\
&\quad \times \left[v_{x_{L_1}, y_0}^\dagger v_{x_{L_1}, y_1} v_{x_{L_1}, y_1}^\dagger v_{x_{L_1}, y_2} \cdots v_{x_{L_1}, y_{L_2-1}}^\dagger v_{x_{L_1}, y_{L_2}} \right] \\
&\quad \times \left[v_{x_0, y_{L_2}}^\dagger v_{x_1, y_{L_2}} \cdots v_{x_{L_1-2}, y_{L_2}}^\dagger v_{x_{L_1-1}, y_{L_2}} v_{x_{L_1-1}, y_{L_2}}^\dagger v_{x_{L_1}, y_{L_2}} \right]^{-1} \\
&\quad \times \left[v_{x_0, y_0}^\dagger v_{x_0, y_1} v_{x_0, y_1}^\dagger v_{x_0, y_2} \cdots v_{x_0, y_{L_2-1}}^\dagger v_{x_0, y_{L_2}} \right]^{-1} h_{0,0}^{-1} \\
&= W_{\partial D} ,
\end{aligned} \tag{B16}$$

where we used $h_{0,0} = 1$.

On the other hand, Eq. (B7) gives us

$$\begin{aligned} \tilde{w}_{i,j} &= h_{ij} \exp \left[-i F_{x_i, y_j} dx dy + O(dx^3, dx^2 dy, dx dy^2, dy^3) \right] h_{ij}^{-1} \\ &= \exp \left[-i h_{ij} F_{x_i, y_j} h_{ij}^{-1} dx dy + O(dx^3, dx^2 dy, dx dy^2, dy^3) \right], \end{aligned} \quad (\text{B17})$$

which means

$$\begin{aligned} \lim_{L_1, L_2 \rightarrow \infty} \mathcal{P} \prod_{i,j} \tilde{w}_{i,j} &= \lim_{L_1, L_2 \rightarrow \infty} \prod_{i=L_1-1, \dots, 2, 1, 0} \prod_{j=0, 1, 2, \dots, L_2-1} \tilde{w}_{i,j} \\ &= \lim_{L_1, L_2 \rightarrow \infty} \prod_{i=L_1-1, \dots, 2, 1, 0} \prod_{j=0, 1, 2, \dots, L_2-1} \exp \left[-i h_{ij} F_{x_i, y_j} h_{ij}^{-1} dx dy + O(dx^3, dx^2 dy, dx dy^2, dy^3) \right] \\ &\equiv \mathcal{P} \exp \left[-i \int_D dx dy \tilde{F}_{x,y} \right], \end{aligned} \quad (\text{B18})$$

where

$$\tilde{F}_{x,y} = h_{x,y} F_{x,y} h_{x,y}^{-1}, \quad (\text{B19})$$

and

$$h_{x,y} = W((x_0, y_0) \rightarrow (x, y_0)) W((x, y_0) \rightarrow (x, y)) \quad (\text{B20})$$

with $(x, y) \rightarrow (x', y')$ being the straight path from (x, y) to (x', y') . Note that we use the indices to avoid clashing notation with Eq. (B9). Eventually, we arrive at the non-abelian stokes theorem

$$W(\partial D) = \mathcal{P} \exp \left[-i \int_D \tilde{F}_{x,y} dx dy \right]. \quad (\text{B21})$$

Eq. (B21) holds for any simply connected D , since it is homeomorphic to a unit disk, which can be parameterized by two parameters with fixed range.

2. Schatten Norms

In this part, we briefly review the definition of Schatten norms, which will be used in the proof of bounds, and provide some useful properties for them.

a. General Definition

Given a generic complex matrix A , its Schatten p -norm is defined as [47]

$$\|A\|_p = \left[\sum_i s_i(A)^p \right]^{1/p} \quad (\text{B22})$$

with $p \in [1, \infty]$, where $s_i(A)$ is the i th singular value of A , and the summation is over all singular values of A . Recall that the singular values of a square matrix A are the square roots of the eigenvalues of AA^\dagger . The Schatten p -norm is a matrix norm [47], which means it must have the following properties.

- $\|A\|_p \geq 0$;
- $\|A\|_p = 0 \Leftrightarrow A = 0$;
- $\|\alpha A\|_p = |\alpha| \|A\|_p$ for any $\alpha \in \mathbb{C}$;
- $\|A + B\|_p \leq \|A\|_p + \|B\|_p$ for any matrix B that has the same dimension as A .

b. Schatten 1-norm for Hermitian Matrices

In particular, we will use the Schatten 1-norm for hermitian matrix A . For convenience, we define

$$\rho(A) = \|A\|_1 . \quad (\text{B23})$$

For Hermitian matrices, singular values are the same as the absolute values of the eigenvalues, and thus the summation of the singular values in the norm can be replaced by the summation of the absolute values of the eigenvalues, *i.e.*,

$$\rho(A) = \|A\|_1 = \sum_i s_i(A) = \sum_i |\lambda_i| , \quad (\text{B24})$$

where λ_i is the i th eigenvalue of the Hermitian A .

There are two useful properties of the Schatten 1-norm ρ for Hermitian matrix A . First, given a generic set of orthonormal set of vectors u_α , then we have

$$\rho(A) \geq \sum_\alpha |u_\alpha^\dagger A u_\alpha| . \quad (\text{B25})$$

To see that, suppose $\{v_i\}$ is the complete set of orthonormal eigenvectors of A for eigenvalue λ_i , and we have

$$C_{i\alpha} = v_i^\dagger u_\alpha \quad (\text{B26})$$

with $\sum_\alpha |C_{i,\alpha}|^2 \leq 1$ where the equality holds when u_α is complete. Then,

$$\sum_\alpha |u_\alpha^\dagger A u_\alpha| = \sum_\alpha \left| \sum_i C_{i\alpha}^* C_{i\alpha} \lambda_i \right| = \sum_\alpha \left| \sum_i |C_{i\alpha}|^2 \lambda_i \right| \leq \sum_\alpha \sum_i |C_{i\alpha}|^2 |\lambda_i| \leq \sum_i |\lambda_i| = \rho(A) . \quad (\text{B27})$$

Second, consider a hermitian matrix that smoothly depends on s , *i.e.*, $A(s)$. Suppose $\{v_{i,s}\}$ is the complete set of orthonormal eigenvectors of $A(s)$ with eigenvalues labeled by $\lambda_{i,s}$, we have

$$\rho \left(\frac{dA(s)}{ds} \right) \geq \sum_i \left| v_{i,s}^\dagger \frac{dA(s)}{ds} v_{i,s} \right| = \sum_i \left| \frac{d(v_{i,s}^\dagger A(s) v_{i,s})}{ds} \right| = \sum_i \left| \frac{d\lambda_{i,s}}{ds} \right| , \quad (\text{B28})$$

where we consider the case that $\lambda_{i,s}$ is isolated in a neighborhood of s . Here we used the Feynman-Hellman theorem $v_{i,s}^\dagger \frac{dA(s)}{ds} v_{i,s} = \frac{d}{ds} (v_{i,s}^\dagger A(s) v_{i,s})$ which is locally valid since we can always choose a smooth gauge for v_i in the neighborhood of s . Note that the final inequality is completely gauge-invariant. To make contact with the Wilson loop, we will need an exponentiated form of the inequality [B28](#):

$$\rho \left(-i \frac{d}{ds} \left[e^{iA(s)} \right] e^{-iA(s)} \right) \geq \sum_i \left| -i v_{i,s}^\dagger \frac{d}{ds} \left[e^{iA(s)} \right] e^{-iA(s)} v_{i,s} \right| = \sum_i \left| -i v_{i,s}^\dagger \frac{d}{ds} \left[e^{iA(s)} \right] v_{i,s} e^{-i\lambda_{i,s}} \right| = \sum_i \left| \frac{d\lambda_{i,s}}{ds} \right| \quad (\text{B29})$$

Moreover, owing to

$$\frac{d}{ds} e^{X(s)} = \int_0^1 e^{\alpha X(s)} \frac{dX(s)}{ds} e^{(1-\alpha)X(s)} d\alpha , \quad (\text{B30})$$

we have

$$\rho \left(-i \frac{d}{ds} \left[e^{iA(s)} \right] e^{-iA(s)} \right) = \rho \left(\int_0^1 e^{i\alpha A(s)} \frac{dA(s)}{ds} e^{-i\alpha A(s)} d\alpha \right) \leq \int_0^1 \rho \left(e^{i\alpha A(s)} \frac{dA(s)}{ds} e^{-i\alpha A(s)} \right) d\alpha = \rho \left(\frac{dA(s)}{ds} \right) \quad (\text{B31})$$

where we have used the triangle inequality of the Schatten norm. In sum, we have

$$\sum_i \left| \frac{d\lambda_{i,s}}{ds} \right| \leq \rho \left(-i \frac{d}{ds} \left[e^{iA(s)} \right] e^{-iA(s)} \right) \leq \rho \left(\frac{dA(s)}{ds} \right) , \quad (\text{B32})$$

for the eigenvalues $\lambda_{i,s}$ of $A(s)$, which means the exponentiated form is a tighter bound than Eq. [\(B28\)](#).




FIG. 3. These is two examples of the D_s with D_0 . In (a), D_0 is the black point, and $\partial D_{s>0}$ are the black solid lines. In this case, D_0 is the starting point for $W(\partial D_s)$ for all s . In (b), $D_s = [0, s] \times [0, Y]$. The black dot in this case is not D_0 but still the starting point for $W(\partial D_s)$ for all s .

Immediately we have, given two Hermitian matrices $A(s)$ and $B(s)$ that smoothly depend on s , we have

$$\begin{aligned}
\rho\left(-i\frac{d}{ds}\left[e^{iA(s)}e^{iB(s)}\right]e^{-iB(s)}e^{-iA(s)}\right) &= \rho\left(-i\left[\frac{d}{ds}e^{iA(s)}\right]e^{-iA(s)}-ie^{iA(s)}\left[\frac{d}{ds}e^{iB(s)}\right]e^{-iB(s)}e^{-iA(s)}\right) \\
&\leq \rho\left(-i\left[\frac{d}{ds}e^{iA(s)}\right]e^{-iA(s)}\right) + \rho\left(-ie^{iA(s)}\left[\frac{d}{ds}e^{iB(s)}\right]e^{-iB(s)}e^{-iA(s)}\right) \\
&= \rho\left(-i\left[\frac{d}{ds}e^{iA(s)}\right]e^{-iA(s)}\right) + \rho\left(-i\left[\frac{d}{ds}e^{iB(s)}\right]e^{-iB(s)}\right) \\
&\leq \rho\left(\frac{dA(s)}{ds}\right) + \rho\left(\frac{dB(s)}{ds}\right),
\end{aligned} \tag{B33}$$

where we, for the third line, use the fact that unitary transformations cannot change eigenvalues.

Appendix C: Wilson loop Lower Bound of the quantum Metric

In this appendix, we prove the WL lower bound (Eq. (12) and Eq. (14)) of the quantum metric for an isolated set of N bands in 2D.

Consider a simply connected region that depend smoothly on a continuous parameter $s \in [0, s_f]$, labeled as D_s , with D_0 having zero area. We require that $D_s \subset D_{s'}$ for any $s \leq s'$, and require the starting point \mathbf{k}_0 of $W(\partial D_s)$ is the same for all $s \in [0, s_f]$. (See two examples of D_s in Fig. 3.) The eigendecomposition of $W(\partial D_s)$ reads $W(\partial D_s) = V_s e^{i\Lambda(s)} V_s^\dagger$ with real diagonal continuous $\Lambda(s) = \text{diag}(\dots, \lambda_{i,s}, \dots)$ and unitary V_s . We choose $\Lambda(s=0) = 0$ without loss of generality, since $W(\partial D_0) = 1$ always holds. Then, we have

$$\begin{aligned}
\int_0^{s_f} ds \rho\left([-i\partial_s W(\partial D_s)] W(\partial D_s)^\dagger\right) &= \int_0^{s_f} ds \rho\left(W(\partial D_s)^\dagger [-i\partial_s W(\partial D_s)]\right) = \\
&\int_0^{s_f} ds \rho\left(\partial_s \left[-ie^{iV_s\Lambda(s)}V_s^\dagger\right]e^{-iV_s\Lambda(s)}V_s^\dagger\right) \geq \int_0^{s_f} ds \sum_i \left|\frac{d\lambda_{i,s}}{ds}\right|,
\end{aligned} \tag{C1}$$

where we have used Eq. (B32). We call $\int_0^{s_f} ds \sum_i \left|\frac{d\lambda_{i,s}}{ds}\right|$ the absolute winding of the WL $W(\partial D_s)$. We emphasize that the absolute winding does not depend on how we rank $\lambda_{i,s}$ as long as they are continuous. We note that since $\lambda_{i,s}$ at most touches at measure zero points, we don't need to consider them when deriving Eq. (C1), allowing us to use Eq. (B32) safely.

The next step is to show the absolute WL winding bounds the quantum metric from below. To show this, we first note that

$$[-i\partial_s W(\partial D_s)] W(\partial D_s)^\dagger = -i\frac{1}{ds} [W(\partial D_{s+ds}) - W(\partial D_s)] W(\partial D_s)^\dagger = -i\frac{1}{ds} [W(\partial D_{s+ds})W(\partial D_s)^\dagger - 1]. \tag{C2}$$

As shown in Fig. 3, $W(\partial D_{s+ds})W(\partial D_s)^\dagger$ is a WL along the boundary of $D_{s+ds} - D_s$. To simplify $W(\partial D_{s+ds})W(\partial D_s)^\dagger$, let us parametrize ∂D_s as $\{\mathbf{k}(s, t) | t \in [0, 1]\}$, where $\mathbf{k}(s, 0) = \mathbf{k}(s, 1) = \mathbf{k}_0$ and $\mathbf{k}(s, t)$ goes through ∂D_s smoothly as

t increases from 0 to 1. In this case, $D_{s+ds} - D_s = \{\mathbf{k}(s', t) | t \in [0, 1], s' \in [s, s + ds]\}$. $W(\partial D_{s+ds})W(\partial D_s)^\dagger$ is then Wilson loop along the boundary of $(D_{s+ds} - D_s)$, as

$$W(\partial D_{s+ds})W(\partial D_s)^\dagger = W((s, 0) \rightarrow (s + ds, 0))W((s + ds, 0) \rightarrow (s + ds, 1))W((s + ds, 1) \rightarrow (s, 1))W((s, 1) \rightarrow (s, 0)) , \quad (\text{C3})$$

where we have used that $W((1, s + ds) \rightarrow (1, s)) = W((0, s) \rightarrow (0, s + ds)) = 1$. To use the non-abelian stokes theorem in Eq. (B21), we first note that the starting point of $W(\partial D_{s+ds})W(\partial D_s)^\dagger$ is $(s, 0)$. Define

$$h_{t,s'} = W((s, 0) \rightarrow (s', 0))W((s', 0) \rightarrow (s', t)) = W((0, 0) \rightarrow (s', 0))W((s', 0) \rightarrow (s', t)) , \quad (\text{C4})$$

and

$$\tilde{F}_{s',t} = h_{s',t} F_{s',t} h_{s',t}^{-1} , \quad (\text{C5})$$

where $F_{s',t}$ is given by replacing (x, y) in Eq. (B5) by (s', t) , and we have used

$$W((s, 0) \rightarrow (s', 0)) = W((0, 0) \rightarrow (s', 0)) = 1 . \quad (\text{C6})$$

Then, we have

$$\begin{aligned} & W(\partial D_{s+ds})W(\partial D_s)^\dagger \\ &= \mathcal{P} \exp \left[-i \int_s^{s+ds} ds' \int_0^1 dt \tilde{F}_{s',t} \right] \\ &= \lim_{L_2 \rightarrow \infty} \prod_{j=0}^{L_2-1} \exp \left[-ids \frac{1}{L_2} \tilde{F}_{s,j/L_2} \right] + O(ds^2) \\ &= 1 - ids \int_0^1 dt \tilde{F}_{(s,t)} + O(ds^2) , \end{aligned} \quad (\text{C7})$$

leading to

$$[-i\partial_s W(\partial D_s)] W(\partial D_s)^\dagger = - \int_0^1 dt \tilde{F}_{(s,t)} . \quad (\text{C8})$$

Substituting Eq. (C8) into the left-most term in Eq. (C1), we obtain

$$\int_0^{s_f} ds \rho \left([-i\partial_s W(\partial D_s)] W(\partial D_s)^\dagger \right) = \int_0^{s_f} ds \rho \left(\int_0^1 dt \tilde{F}_{s,t} \right) \leq \int_0^{s_f} ds \int_0^1 dt \rho \left(\tilde{F}_{s,t} \right) = \int_0^{s_f} ds \int_0^1 dt \rho(F_{s,t}) , \quad (\text{C9})$$

where we use the fact that $F_{s,t}$ and $\tilde{F}_{s,t}$ have the same eigenvalues.

Previously, we choose the parametrize (s, t) to make sure the region of the parameters is rectangular, allowing us to directly use Eq. (B21). We now convert it back to the Bloch momentum, which is what we normally use. Define $F_{\mathbf{k}}$ as Eq. (B5) with $(x, y) = (k_x, k_y)$. Then,

$$F_{s,t} = \det \left[\frac{\partial(s, t)}{\partial(k_x, k_y)} \right] F_{\mathbf{k}} , \quad (\text{C10})$$

where $\frac{\partial(s, t)}{\partial(k_x, k_y)}$ is the Jacobian matrix. Then,

$$\begin{aligned} \int_0^{s_f} ds \int_0^1 dt \rho(F_{s,t}) &= \int_0^{s_f} ds \int_0^1 dt \rho \left(\det \left[\frac{\partial(s, t)}{\partial(k_x, k_y)} \right] F_{\mathbf{k}} \right) = \int_0^{s_f} ds \int_0^1 dt \left| \det \left[\frac{\partial(s, t)}{\partial(k_x, k_y)} \right] \right| \rho(F_{\mathbf{k}}) \\ &= \int_0^{s_f} ds \int_0^1 dt \left| \det \left[\frac{\partial(s, t)}{\partial(k_x, k_y)} \right] \right| \rho(F_{\mathbf{k}}) = \int_{D_{s_f}} d^2k \rho(F_{\mathbf{k}}) , \end{aligned} \quad (\text{C11})$$

which means

$$\int_0^{s_f} ds \rho \left([-i\partial_s W(\partial D_s)] W(\partial D_s)^\dagger \right) \leq \int_{D_{s_f}} d^2k \rho(F_{\mathbf{k}}) . \quad (\text{C12})$$

As Eq. (A11) suggests that $\rho(F_{\mathbf{k}}) \leq \text{Tr}[g_{\mathbf{k}}]$, we arrive at

$$\frac{1}{2\pi} \int_{D_{s_f}} d^2k \text{Tr}[g_{\mathbf{k}}] \geq \frac{1}{2\pi} \int_0^{s_f} ds \rho \left([-i\partial_s W(\partial D_s)] W(\partial D_s)^\dagger \right) \geq \int_0^{s_f} ds \sum_i \left| \frac{d\lambda_{i,s}}{ds} \right|. \quad (\text{C13})$$

One special yet useful case for the WL winding bound is when $D_s = \{k_1 \mathbf{b}_1/(2\pi) + k_2 \mathbf{b}_2/(2\pi) | k_1 \in [0, s], k_2 \in [0, 2\pi]\}$. (See Fig. 3(b) with $Y = 2\pi$.) Here \mathbf{b}_1 and \mathbf{b}_2 are two basis reciprocal lattice vectors, which means $D_{2\pi} = \text{BZ}$ is the entire first Brillouin zone. In this case, we can define a new WL which is

$$\mathcal{W}_{k_1} = W(k_1 \mathbf{b}_1/(2\pi) \rightarrow k_1 \mathbf{b}_1/(2\pi) + \mathbf{b}_2), \quad (\text{C14})$$

which is the WL along k_2 at a fixed k_1 . Then, the absolute winding of $W(\partial D_s)$ in Eq. (C13) can be directly replaced by the absolute winding of \mathcal{W}_s , resulting in

$$\int_0^{s_f} ds \sum_l \left| \frac{d\phi_l(s)}{ds} \right| \leq \int_{D_{s_f}} d^2k \text{Tr}[g_{\mathbf{k}}], \quad (\text{C15})$$

where $\phi_l(s)$ is the phase of the l th eigenvalue of \mathcal{W}_s and is always chosen to be continuous. To prove this, we first note that

$$W(\partial D_s) = W(\Gamma \rightarrow s\mathbf{b}_1/(2\pi)) \mathcal{W}_s W^\dagger(\Gamma \rightarrow s\mathbf{b}_1/(2\pi)) W(\mathbf{b}_2 \rightarrow \Gamma), \quad (\text{C16})$$

where $\Gamma = (0, 0)$. As $W(\mathbf{b}_2 \rightarrow \Gamma)$ is independent of s , we have

$$\begin{aligned} & \rho[-iW^\dagger(\partial D_s)\partial_s W(\partial D_s)] \\ &= \rho[-iW(\Gamma \rightarrow s\mathbf{b}_1/(2\pi)) \mathcal{W}_s^\dagger W^{-1}(\Gamma \rightarrow s\mathbf{b}_1/(2\pi)) \partial_s \{W(\Gamma \rightarrow s\mathbf{b}_1/(2\pi)) \mathcal{W}_s W^{-1}(\Gamma \rightarrow s\mathbf{b}_1/(2\pi))\}] \\ &\geq \int_0^{s_f} ds \sum_l \left| \frac{d\phi_l(s)}{ds} \right|, \end{aligned} \quad (\text{C17})$$

where we use the fact that $W(\Gamma \rightarrow s\mathbf{b}_1/(2\pi)) \mathcal{W}_s W^\dagger(\Gamma \rightarrow s\mathbf{b}_1/(2\pi))$ has the same eigenvalues as \mathcal{W}_s and use the second inequality in Eq. (C13). Combined with Eq. (C13), we arrive at Eq. (C15).

Appendix D: Z_2 Lower Bound of the quantum Metric

In this appendix, we derive the Z_2 lower bound of the quantum metric. We will discuss two Z_2 indices, one protected by the TR symmetry and the other protected by the particle-hole (PH) symmetry, which are in fact equivalent.

1. Review of TR Z_2 index from Wilson Loop

Before discussing the lower bound, let us first review how to calculate TR Z_2 index from WL. Since the TR Z_2 index is defined for 2D systems with spinful TR symmetry [31], we consider an isolated set of $2N$ bands owing to Kramer's degeneracy. The WL of interest is \mathcal{W}_{k_1} in Eq. (C14). Because of TR symmetry, we have

$$\mathcal{W}_{k_1} = U_{\frac{k_1}{2\pi}\mathbf{b}_1}^T \mathcal{W}_{k_1}^T U_{\frac{k_1}{2\pi}\mathbf{b}_1}^* \quad \text{for } k_1 = 0, \pi, \quad (\text{D1})$$

where $[U_{\mathbf{k}}]_{mn} = \langle u_{-\mathbf{k},m} | \mathcal{T} | u_{\mathbf{k},n} \rangle$. It means that \mathcal{W}_0 and \mathcal{W}_π have Kramer's degeneracy.

As discussed in Ref. [35], we track the phase of the eigenvalues of \mathcal{W}_{k_1} from $k_1 = 0$ to $k_1 = \pi$ to determine Z_2 index. Specifically, \mathcal{W}_{k_1} has eigenvalues $e^{i\phi_l(k_1)}$ with $l = 1, 2, \dots, 2N$, and we can choose $\phi_l(k_1)$ to be continuous for $k_1 \in [0, \pi]$ which is always allowed. Without loss of generality, we fix $\phi_l(0) \in [0, 2\pi)$; furthermore, we have in general

$$\phi_l(\pi) = [\phi_l(\pi) \bmod 2\pi] + 2\pi M_l, \quad (\text{D2})$$

where

$$[x \bmod 2\pi] \in [0, 2\pi), \quad (\text{D3})$$

and $M_l \in \mathbb{Z}$. Finally, Z_2 reads

$$Z_2 = \sum_{l=1}^{2N} M_l \bmod 2. \quad (\text{D4})$$




FIG. 4. (a) A representative Wilson loop for an isolated set of two bands with $Z_2 = 1$. (b) A representative Wilson loop for an isolated set of two bands with $Z_2 = 0$. (c) A representative Wilson loop given by coupling (a) to (b), which is an isolated set of four bands with $Z_2 = 1$.

2. Review of PH Z_2 index

As discussed in Ref. [46], the PH symmetry also protects a Z_2 index in a same way as the TR symmetry, which we review in this section.

Consider the PH matrix $U_{\mathcal{P}}(\mathbf{k})$, which satisfies $U_{\mathcal{P}}(\mathbf{k})h^*(\mathbf{k})U_{\mathcal{P}}^\dagger(\mathbf{k}) = -h(-\mathbf{k})$ and $U_{\mathcal{P}}(\mathbf{k})U_{\mathcal{P}}^*(-\mathbf{k}) = -1$, where $h(\mathbf{k})$ is the matrix Hamiltonian. Suppose we have an isolated set of $2N$ bands of $h(\mathbf{k})$ that preserves the PH symmetry, and we note the corresponding eigenvectors as $v_{n,\mathbf{k}}$ with $n = 1, 2, 3, \dots, 2N$, which satisfies

$$U_{\mathcal{P}}(\mathbf{k})v_{n,\mathbf{k}}^* = \sum_{m=1}^{2N} v_{m,-\mathbf{k}} [U_{\mathbf{k}}]_{mn} \quad (\text{D5})$$

with unitary $U_{\mathbf{k}}$. Since Eq. (D5) has the same form as the TR symmetry acting on the eigenstates, the PH-preserving $2N$ bands must be able to have the same Z_2 topology as the TR-preserved bands, characterized by the same WL winding as Eq. (D4).

3. Proof of the Z_2 Lower Bound of the quantum Metric

We will only show the derivation for the TR Z_2 , since the derivation for the PH Z_2 is exactly the same.

According to Eq. (C15), we can find a Z_2 lower bound of the quantum metric as long as we can bound the absolute winding of \mathcal{W}_{k_1} . In the following, we will prove

$$\sum_l \left| \int_0^\pi dk_1 \frac{d}{dk_1} \phi_l(k_1) \right| \geq 2\pi Z_2, \quad (\text{D6})$$

where $\phi_l(k_1)$ are the phases of the eigenvalues of \mathcal{W}_{k_1} , which we choose to be continuous. Of course, Eq. (D6) is trivially true for $Z_2 = 0$. In the following, we will consider the case where $Z_2 = 1$.

Before proving the general case, let us look at an example. As shown by Ref. [35], for an isolated set of two bands with $Z_2 = 1$, the two WL bands are connected as shown in Fig. 4(a). Since Z_2 is a stable topology, it means that if we couple an isolated set of two bands with $Z_2 = 1$ to an isolated set of two bands with $Z_2 = 0$ (Fig. 4(b)), we obtain an isolated set of four bands with $Z_2 = 1$. As shown in Fig. 4(c), the isolated set of four bands with $Z_2 = 1$ still have fully connected WL bands. Therefore, this example suggests an intuitive argument that for an isolated set of any number of bands with $Z_2 = 1$, the WL bands are fully connected. Such argument was presented in Ref. [46].

Now we prove the connectivity for the WL bands in the case of $Z_2 = 1$, which is essential for the proof of the Z_2 bound. Since the absolute WL winding does not care about how we rank $\phi_l(k_1)$, we choose

$$[\phi_l(k_1) - \phi_{l'}(k_1) + 2\pi n][\phi_l(k'_1) - \phi_{l'}(k'_1) + 2\pi n] \geq 0 \text{ for any } k_1, k'_1 \in [0, \pi], n \in \mathbb{Z} \text{ and different } l \text{ and } l' \quad (\text{D7})$$

and $\phi_l(0) \in [0, 2\pi)$. In other words, we make sure $\phi_l(k_1)$ and $\phi_{l'}(k_1)$ do not cross each other for any different l and l' and for any 2π shifts, though $\phi_l(k_1)$ and $\phi_{l'}(k_1)$ may still have touching that is either accidental or protected by extra symmetries for any $k_1 \in (0, \pi)$. Eq. (D7) is always allowed. Without loss of generality, we choose $\phi_1(k_1) \leq \phi_2(k_1) \leq \phi_3(k_1) \leq \dots \leq \phi_{2N}(k_1)$. What we want to show is that it is impossible to have a direct gap among

$\phi_l(k_1)$'s, *i.e.*, it is impossible to have l_0 that satisfies $\phi_{l_0+1}(k_1) > \phi_{l_0}(k_1)$ for all $k_1 \in [0, \pi]$. First note that $e^{i\phi_l(\pi)}$ form Kramers pairs owing to TR symmetry. In principle, given a Kramers pair $e^{i\phi_{l_1}(\pi)}$ and $e^{i\phi_{l_2}(\pi)}$ with $l_1 \neq l_2$, only $\phi_{l_1}(\pi) = \phi_{l_2}(\pi) \bmod 2\pi$ is required, whereas it is allowed to have $\phi_{l_1}(\pi) \neq \phi_{l_2}(\pi)$. It is impossible to have $\phi_{l_1}(\pi) = \phi_{l_2}(\pi)$ for all Kramers pair $e^{i\phi_{l_1}(\pi)}$ and $e^{i\phi_{l_2}(\pi)}$ with $l_1 \neq l_2$, since otherwise $M_{l_1} = M_{l_2}$ in Eq. (D4) for all pairs and Z_2 must be zero. Therefore, we must have at least one Kramers pair $e^{i\phi_{l_1}(\pi)}$ and $e^{i\phi_{l_2}(\pi)}$ with $l_1 \neq l_2$ such that $\phi_{l_1}(\pi) \neq \phi_{l_2}(\pi)$, which we call nontrivial Kramers pair. Consider a generic nontrivial Kramers pair $e^{i\phi_{l_1}(\pi)}$ and $e^{i\phi_{l_2}(\pi)}$ with $l_1 \neq l_2$. We can assume $l_1 < l_2$ without loss of generality, which means $\phi_{l_2}(\pi) = \phi_{l_1}(\pi) + 2\pi n$ with integer $n > 0$. If $\phi_{l_2}(\pi) < \phi_{2N}(\pi)$, then $\phi_{l_1}(k_1) + 2\pi n$ crosses $\phi_{2N}(k_1)$ since $\phi_{l_1}(0) + 2\pi n > \phi_{2N}(0)$ and $\phi_{l_1}(\pi) + 2\pi n < \phi_{2N}(\pi)$, which contradicts Eq. (D7). Similarly, if $\phi_{l_1}(\pi) > \phi_{l_2}(\pi)$, then $\phi_{l_2}(k_1) - 2\pi n$ crosses $\phi_{l_1}(k_1)$ since $\phi_{l_2}(0) - 2\pi n < \phi_{l_1}(0)$ and $\phi_{l_2}(\pi) - 2\pi n > \phi_{l_1}(\pi)$, which again contradicts Eq. (D7). Therefore, for any nontrivial Kramers pair $e^{i\phi_{l_1}(\pi)}$ and $e^{i\phi_{l_2}(\pi)}$ with $l_1 < l_2$, we have $\phi_{2N}(\pi) = \phi_{l_2}(\pi) = \phi_{l_1}(\pi) + 2\pi n = \phi_{l_1}(\pi) + 2\pi n$ with integer $n > 0$, meaning that $M_{l_1} + M_{l_2} = 2M_{l_1} + n$ and n is the same for different nontrivial Kramers pair.

In other words, suppose we have x nontrivial Kramers pair, the Z_2 index is just $xn \bmod 2$, according to Eq. (D4). We then know that we can only have x being odd, since an even x will make $Z_2 = 0$. Now suppose there is a gap between ϕ_{l_0} and ϕ_{l_0+1} . l_0 cannot be odd, since odd l_0 will make $\phi_{l_0+1}(k_1)$ and $\phi_{l_0}(k_1)$ touching at $k_1 = 0$. However, l_0 cannot be even either. Because for each nontrivial Kramers pairs $e^{i\phi_{l_1}(\pi)}$ and $e^{i\phi_{l_2}(\pi)}$ with $l_1 < l_2$, we must have $l_1 \leq l_0$ and $l_2 \geq l_0 + 1$, which means $(l_0 - x)$ WL bands with $l \leq l_0$ must all form trivial Kramers pairs to realize the gap and thus $(l_0 - x)$ is even. However, if l_0 is even, then $(l_0 - x)$ must be odd, leading to contradiction. Therefore, l_0 cannot be even either, which means l_0 does not exist, *i.e.*, it is impossible to have a direct gap among $\phi_l(k_1)$'s

The connectivity (*i.e.*, the absence of direct gap) and the no-crossing condition (Eq. (D7)) tell us that $\phi_l(0) = \phi_{l+1}(0)$ for odd l and $\phi_l(\pi) = \phi_{l+1}(\pi)$ for even l . Thus,

$$\begin{aligned} \sum_l \left| \int_0^\pi dk_1 \frac{d}{dk_1} \phi_l(k_1) \right| &\geq \left[\sum_{l=1,3,5,\dots,2N-1} \phi_l(0) - \phi_l(\pi) \right] + \left[\sum_{l=2,4,6,\dots,2N} \phi_l(\pi) - \phi_l(0) \right] \\ &= \sum_{l=1,3,5,\dots,2N-1} [\phi_l(0) - \phi_{l+1}(0)] + \sum_{l=2,4,6,\dots,2N-2} [\phi_l(\pi) - \phi_{l+1}(\pi)] + \phi_{2N}(\pi) - \phi_1(\pi) = \phi_{2N}(\pi) - \phi_1(\pi) = 2\pi n \geq 2\pi. \end{aligned} \quad (\text{D8})$$

Eq. (D6) is proven for $Z_2 = 1$.

Combining Eq. (D6) with Eq. (C15), we arrive at

$$2\pi Z_2 \leq \int_{\text{half BZ}} d^2k \text{Tr}[g_{\mathbf{k}}]. \quad (\text{D9})$$

Combined with the fact that $\text{Tr}[g_{\mathbf{k}}] = \text{Tr}[g_{-\mathbf{k}}]$ owing to TR symmetry, we obtain

$$\frac{1}{2\pi} \int_{\text{BZ}} d^2k \text{Tr}[g] \geq 2Z_2. \quad (\text{D10})$$

We note that the PH Z_2 index provides the same bound as Eq. (D10).

Appendix E: Physical Consequences of the TR Z_2 Bound

In this section, we will discuss three physical consequences, superfluid weight, optical conductivity and band gap, that can be bounded by the Z_2 bound in Eq. (D10). We always choose

$$\hbar = e = 1 \quad (\text{E1})$$

with electron having charge $-e$.

1. TR Z_2 Lower Bound for Superfluid Weight

We first discuss the mean-field superfluid weight in flat bands with a nontrivial Z_2 index, which we show is bounded below according to the inequality proved in the Main Text. The derivation that relates superfluid weight to quantum geometry has been worked out in Ref. [12–21] and is included here for completeness. The new result here is the new bound of superfluid weight due to the Z_2 index.

We consider a normal-phase 2D single-particle Hamiltonian H_0 that can realize a exactly-flat doubly degenerate Z_2 band at energy E_0 . Such a Hamiltonian is allowed in the continuous Hamiltonian (such as moiré Hamiltonians) but may have long-range hoppings in a tight-binding formalism. In general, H_0 reads

$$H_0 = \sum_{\mathbf{k}, i, j} c_{\mathbf{k}, i}^\dagger [h_0(\mathbf{k})]_{ij} c_{\mathbf{k}, j} , \quad (\text{E2})$$

Suppose when the Z_2 band is partially filled, we will have superconductivity with order parameter

$$H_\Delta = \frac{1}{2} \Delta \sum_{\mathbf{k}} c_{\mathbf{k}}^\dagger U_{\mathcal{T}} \left(c_{-\mathbf{k}}^\dagger \right)^T + h.c. , \quad (\text{E3})$$

where

$$c_{\mathbf{k}}^\dagger = (\dots, c_{\mathbf{k}, i}^\dagger, \dots) , \quad (\text{E4})$$

and $U_{\mathcal{T}}$ is the matrix representation of the TR symmetry, *i.e.*,

$$\mathcal{T} c_{\mathbf{k}}^\dagger \mathcal{T}^{-1} = c_{-\mathbf{k}}^\dagger U_{\mathcal{T}} , \quad (\text{E5})$$

where $U_{\mathcal{T}}$ is a unitary matrix that satisfies

$$U_{\mathcal{T}} U_{\mathcal{T}}^* = -1 \Rightarrow U_{\mathcal{T}} = -U_{\mathcal{T}}^T . \quad (\text{E6})$$

Then, the mean-field Hamiltonian reads

$$H_{MF} = \sum_{\mathbf{k}} c_{\mathbf{k}}^\dagger [h_0(\mathbf{k}) - \mu] c_{\mathbf{k}} + \frac{1}{2} \Delta \sum_{\mathbf{k}} c_{\mathbf{k}}^\dagger U_{\mathcal{T}} \left(c_{-\mathbf{k}}^\dagger \right)^T + \frac{1}{2} \Delta^* \sum_{\mathbf{k}} c_{-\mathbf{k}}^T U_{\mathcal{T}}^\dagger c_{\mathbf{k}} . \quad (\text{E7})$$

For examples of microscopic Hamiltonians where the above mean-field description of uniform pairing described by a single number Δ is accurate, we refer to reader to [54, 55].

We now focus on the superfluid weight [48]. For that purpose, we will add thread flux in the mean-field Hamiltonian, and obtain

$$H_{MF}(\mathbf{A}) = \sum_{\mathbf{k}} c_{\mathbf{k}}^\dagger [h_0(\mathbf{k} + \mathbf{A}) - \mu] c_{\mathbf{k}} + \frac{1}{2} \Delta \sum_{\mathbf{k}} c_{\mathbf{k}}^\dagger U_{\mathcal{T}} \left(c_{-\mathbf{k}}^\dagger \right)^T + \frac{1}{2} \Delta^* \sum_{\mathbf{k}} c_{-\mathbf{k}}^T U_{\mathcal{T}}^\dagger c_{\mathbf{k}} . \quad (\text{E8})$$

Let us consider the case that the flat Z_2 band is well isolated, and thus we can project the mean-field Hamiltonian to that specific band. To do so, we use $U_{n, \mathbf{k}}$ to label the eigenvectors for the flat Z_2 band ($n = 1, 2$). The creation operator for the Z_2 band reads

$$\psi_{n, \mathbf{k}, \mathbf{A}}^\dagger = c_{\mathbf{k}}^\dagger U_{n, \mathbf{k} + \mathbf{A}} , \quad (\text{E9})$$

and thus the projected mean-field Hamiltonian reads

$$\begin{aligned} \bar{H}_{MF}(\mathbf{A}) &= \sum_{\mathbf{k}} \psi_{\mathbf{k}, \mathbf{A}}^\dagger [E_0 - \mu] \psi_{\mathbf{k}, \mathbf{A}} + \frac{1}{2} \sum_{\mathbf{k}} \psi_{\mathbf{k}, \mathbf{A}}^\dagger \Delta_{\mathbf{k}, \mathbf{A}} \left(\psi_{-\mathbf{k}, \mathbf{A}}^\dagger \right)^T + \frac{1}{2} \sum_{\mathbf{k}} \psi_{-\mathbf{k}, \mathbf{A}}^T \Delta_{\mathbf{k}, \mathbf{A}}^\dagger \psi_{\mathbf{k}, \mathbf{A}} \\ &= \frac{1}{2} \sum_{\mathbf{k}} \left(\psi_{\mathbf{k}, \mathbf{A}}^\dagger \quad \psi_{-\mathbf{k}, \mathbf{A}}^T \right) h_{\text{BdG}}(\mathbf{k}, \mathbf{A}) \begin{pmatrix} \psi_{\mathbf{k}, \mathbf{A}} \\ \left(\psi_{-\mathbf{k}, \mathbf{A}}^\dagger \right)^T \end{pmatrix} . \end{aligned} \quad (\text{E10})$$

where $\psi_{\mathbf{k}, \mathbf{A}}^\dagger = (\psi_{1, \mathbf{k}, \mathbf{A}}^\dagger, \psi_{2, \mathbf{k}, \mathbf{A}}^\dagger)$,

$$\Delta_{\mathbf{k}, \mathbf{A}} = \Delta U_{\mathbf{k} + \mathbf{A}}^\dagger U_{\mathcal{T}} U_{-\mathbf{k} + \mathbf{A}}^* , \quad (\text{E11})$$

$$h_{\text{BdG}}(\mathbf{k}, \mathbf{A}) = \begin{pmatrix} [E_0 - \mu] \sigma_0 & \Delta_{\mathbf{k}, \mathbf{A}} \\ \Delta_{\mathbf{k}, \mathbf{A}}^\dagger & -[E_0 - \mu] \sigma_0 \end{pmatrix} , \quad (\text{E12})$$

and

$$U_{\mathbf{k}} = (U_{1, \mathbf{k}} \quad U_{2, \mathbf{k}}) . \quad (\text{E13})$$

The order parameter in Eq. (E11) naturally obeys the requirement from the fermion statistics:

$$\Delta_{\mathbf{k},\mathbf{A}}^T = \Delta U_{-\mathbf{k}+\mathbf{A}}^\dagger U_{\mathcal{T}}^T U_{\mathbf{k}+\mathbf{A}}^* = \Delta_{-\mathbf{k},\mathbf{A}}. \quad (\text{E14})$$

The TR symmetry provides

$$U_{\mathcal{T}} U_{\mathbf{k}}^* = U_{-\mathbf{k}} D_{\mathcal{T}}(\mathbf{k}) \quad (\text{E15})$$

with unitary $D_{\mathcal{T}}(\mathbf{k})$ satisfying

$$D_{\mathcal{T}}(-\mathbf{k}) D_{\mathcal{T}}^*(\mathbf{k}) = -1. \quad (\text{E16})$$

As a result, we have

$$\Delta_{\mathbf{k},\mathbf{A}} = \Delta U_{\mathbf{k}+\mathbf{A}}^\dagger U_{\mathcal{T}} U_{-\mathbf{k}+\mathbf{A}}^* = \Delta U_{\mathbf{k}+\mathbf{A}}^\dagger U_{\mathbf{k}-\mathbf{A}} D_{\mathcal{T}}(-\mathbf{k} + \mathbf{A}) \quad (\text{E17})$$

The superfluid weight is derived from the free energy, which reads

$$\begin{aligned} \Omega(\mathbf{A}) &= \sum_{\mathbf{k}} \left(\frac{1}{2} \text{Tr}[(E_0 - \mu)\sigma_0] - \frac{1}{\beta} \sum_n^{E_{\mathbf{k},n}(\mathbf{A}) > 0} \log(2 \cosh(\beta E_{\mathbf{k},n}(\mathbf{A})/2)) \right) \\ &= \sum_{\mathbf{k}} \left((E_0 - \mu) - \frac{1}{\beta} \sum_n^{E_{\mathbf{k},n}(\mathbf{A}) > 0} \log(2 \cosh(\beta E_{\mathbf{k},n}(\mathbf{A})/2)) \right), \end{aligned} \quad (\text{E18})$$

where $E_{\mathbf{k},n}(\mathbf{A})$ is the eigenvalue of $h_{\text{BdG}}(\mathbf{k}, \mathbf{A})$. To derive the analytical expression for $E_{\mathbf{k},n}(\mathbf{A})$, let us consider $h_{\text{BdG}}^2(\mathbf{k}, \mathbf{A})$, which reads

$$h_{\text{BdG}}^2(\mathbf{k}, \mathbf{A}) = [E_0 - \mu]^2 + \begin{pmatrix} \Delta_{\mathbf{k},\mathbf{A}} \Delta_{\mathbf{k},\mathbf{A}}^\dagger & \\ & \Delta_{\mathbf{k},\mathbf{A}}^\dagger \Delta_{\mathbf{k},\mathbf{A}} \end{pmatrix}. \quad (\text{E19})$$

The TR symmetry provides

$$\begin{aligned} \Delta_{\mathbf{k},\mathbf{A}} \Delta_{\mathbf{k},\mathbf{A}}^\dagger &= |\Delta|^2 U_{\mathbf{k}+\mathbf{A}}^\dagger U_{\mathbf{k}-\mathbf{A}} D_{\mathcal{T}}(-\mathbf{k} + \mathbf{A}) D_{\mathcal{T}}^\dagger(-\mathbf{k} + \mathbf{A}) U_{\mathbf{k}-\mathbf{A}}^\dagger U_{\mathbf{k}+\mathbf{A}} \\ &= |\Delta|^2 U_{\mathbf{k}+\mathbf{A}}^\dagger U_{\mathbf{k}-\mathbf{A}} U_{\mathbf{k}-\mathbf{A}}^\dagger U_{\mathbf{k}+\mathbf{A}}. \end{aligned} \quad (\text{E20})$$

Combining the TR symmetry with Eq. (E14), we arrive at

$$\begin{aligned} &D_{\mathcal{T}}^\dagger(-\mathbf{k} - \mathbf{A}) \Delta_{\mathbf{k},\mathbf{A}} \Delta_{\mathbf{k},\mathbf{A}}^\dagger D_{\mathcal{T}}(-\mathbf{k} - \mathbf{A}) \\ &= |\Delta|^2 D_{\mathcal{T}}^\dagger(-\mathbf{k} - \mathbf{A}) U_{\mathbf{k}+\mathbf{A}}^\dagger U_{\mathbf{k}-\mathbf{A}} U_{\mathbf{k}-\mathbf{A}}^\dagger U_{\mathbf{k}+\mathbf{A}} D_{\mathcal{T}}(-\mathbf{k} - \mathbf{A}) \\ &= |\Delta|^2 U_{-\mathbf{k}-\mathbf{A}}^T U_{\mathcal{T}}^\dagger U_{\mathcal{T}} U_{-\mathbf{k}+\mathbf{A}}^* U_{-\mathbf{k}+\mathbf{A}}^T U_{\mathcal{T}}^\dagger U_{\mathcal{T}} U_{-\mathbf{k}-\mathbf{A}}^* \\ &= |\Delta|^2 U_{-\mathbf{k}-\mathbf{A}}^T U_{-\mathbf{k}+\mathbf{A}}^* U_{-\mathbf{k}+\mathbf{A}}^T U_{-\mathbf{k}-\mathbf{A}}^* \\ &= D_{\mathcal{T}}^*(\mathbf{k} + \mathbf{A}) \left[\Delta_{-\mathbf{k},\mathbf{A}}^\dagger \Delta_{-\mathbf{k},\mathbf{A}} \right]^T D_{\mathcal{T}}^T(\mathbf{k} + \mathbf{A}) \\ &= D_{\mathcal{T}}^*(\mathbf{k} + \mathbf{A}) \Delta_{\mathbf{k},\mathbf{A}}^\dagger \Delta_{\mathbf{k},\mathbf{A}} D_{\mathcal{T}}^T(\mathbf{k} + \mathbf{A}), \end{aligned} \quad (\text{E21})$$

Therefore, the eigenvalues of $\Delta_{\mathbf{k},\mathbf{A}} \Delta_{\mathbf{k},\mathbf{A}}^\dagger$ are equal to those of $\Delta_{\mathbf{k},\mathbf{A}}^\dagger \Delta_{\mathbf{k},\mathbf{A}}$, which we labeled as $\lambda_{0,\mathbf{k},\mathbf{A}}$ and $\lambda_{1,\mathbf{k},\mathbf{A}}$. Furthermore, based on the TR symmetry, we have

$$\begin{aligned} \Delta_{\mathbf{k},\mathbf{A}}^\dagger \Delta_{\mathbf{k},\mathbf{A}} &= |\Delta|^2 D_{\mathcal{T}}^\dagger(-\mathbf{k} + \mathbf{A}) U_{\mathbf{k}-\mathbf{A}}^\dagger U_{\mathbf{k}+\mathbf{A}} U_{\mathbf{k}+\mathbf{A}}^\dagger U_{\mathbf{k}-\mathbf{A}} D_{\mathcal{T}}(-\mathbf{k} + \mathbf{A}) \\ &= D_{\mathcal{T}}^\dagger(-\mathbf{k} + \mathbf{A}) \Delta_{\mathbf{k},-\mathbf{A}} \Delta_{\mathbf{k},-\mathbf{A}}^\dagger D_{\mathcal{T}}(-\mathbf{k} + \mathbf{A}), \end{aligned} \quad (\text{E22})$$

which means we can always choose (and we will choose) $\lambda_{\alpha,\mathbf{k},\mathbf{A}} = \lambda_{\alpha,\mathbf{k},-\mathbf{A}}$ for $\alpha = 0, 1$. The eigenvalues of $h_{\text{BdG}}(\mathbf{k}, \mathbf{A})$ are just

$$E_{s,\alpha}(\mathbf{k}) = (-1)^s \sqrt{(E_0 - \mu)^2 + \lambda_{\alpha,\mathbf{k},\mathbf{A}}} \quad (\text{E23})$$

for $s = 0, 1$ and $\alpha = 0, 1$. Clearly, the positive $E_{+, \alpha}(\mathbf{k})$ are non-negative, which gives

$$\Omega(\mathbf{A}) = \sum_{\mathbf{k}} \left((E_0 - \mu) - \frac{1}{\beta} \sum_{\alpha} \log \left(2 \cosh \left(\frac{\beta}{2} \sqrt{(E_0 - \mu)^2 + \lambda_{\alpha, \mathbf{k}, \mathbf{A}}} \right) \right) \right). \quad (\text{E24})$$

The superfluid weight just reads

$$[D_s(T)]_{ij} = \frac{1}{\mathcal{V}} \frac{\partial^2 \Omega(\mathbf{A})}{\partial A_i \partial A_j} \Big|_{\mathbf{A} \rightarrow 0}, \quad (\text{E25})$$

where \mathcal{V} is the sample volume. In a fully self-consistent treatment, the partial derivatives must be replaced with total derivatives. Nevertheless, it was shown in [49] that TR symmetry guarantees the same result holds as long as $g_{\mathbf{k}}$ is the minimal quantum metric.

We are particularly interested in the zero-temperature case, where the free energy becomes

$$\Omega_0(\mathbf{A}) = \sum_{\mathbf{k}} \left((E_0 - \mu) - \frac{1}{2} \sum_{\alpha} \sqrt{(E_0 - \mu)^2 + \lambda_{\alpha, \mathbf{k}, \mathbf{A}}} \right). \quad (\text{E26})$$

As a result, the zero-temperature superfluid weight reads

$$\begin{aligned} \mathcal{V} [D_{SW}(0)]_{ij} &= \frac{\partial^2}{\partial A_i \partial A_j} \sum_{\mathbf{k}} \left((E_0 - \mu) - \frac{1}{2} \sum_{\alpha} \sqrt{(E_0 - \mu)^2 + \lambda_{\alpha, \mathbf{k}, \mathbf{A}}} \right) \Big|_{\mathbf{A} \rightarrow 0} \\ &= -\frac{1}{2} \sum_{\mathbf{k}} \sum_{\alpha} \frac{\partial^2}{\partial A_i \partial A_j} \sqrt{(E_0 - \mu)^2 + \lambda_{\alpha, \mathbf{k}, \mathbf{A}}} \Big|_{\mathbf{A} \rightarrow 0} \\ &= -\frac{1}{2} \sum_{\mathbf{k}} \sum_{\alpha} \frac{\partial}{\partial A_i} \frac{1}{2\sqrt{(E_0 - \mu)^2 + \lambda_{\alpha, \mathbf{k}, \mathbf{A}}}} \frac{\partial}{\partial A_j} \lambda_{\alpha, \mathbf{k}, \mathbf{A}} \Big|_{\mathbf{A} \rightarrow 0} \\ &= -\frac{1}{2} \sum_{\mathbf{k}} \sum_{\alpha} \left[-\frac{1}{4[(E_0 - \mu)^2 + \lambda_{\alpha, \mathbf{k}, \mathbf{A}}]^{3/2}} \frac{\partial \lambda_{\alpha, \mathbf{k}, \mathbf{A}}}{\partial A_i} \frac{\partial \lambda_{\alpha, \mathbf{k}, \mathbf{A}}}{\partial A_j} + \frac{1}{2\sqrt{(E_0 - \mu)^2 + \lambda_{\alpha, \mathbf{k}, \mathbf{A}}}} \frac{\partial^2}{\partial A_i \partial A_j} \lambda_{\alpha, \mathbf{k}, \mathbf{A}} \right] \Big|_{\mathbf{A} \rightarrow 0}. \end{aligned} \quad (\text{E27})$$

From Eq. (E20), we know

$$\mathcal{V} \Delta_{\mathbf{k}, 0} \Delta_{\mathbf{k}, 0}^{\dagger} = |\Delta|^2 \sigma_0, \quad (\text{E28})$$

which means

$$\lambda_{0, \mathbf{k}, 0} = \lambda_{1, \mathbf{k}, 0} = |\Delta|^2. \quad (\text{E29})$$

Therefore, the zero-temperature superfluid weight becomes

$$\begin{aligned} \mathcal{V} [D_{SW}(0)]_{ij} &= -\frac{1}{2} \sum_{\mathbf{k}} \sum_{\alpha} \left[-\frac{1}{4[(E_0 - \mu)^2 + |\Delta|^2]^{3/2}} \frac{\partial \lambda_{\alpha, \mathbf{k}, \mathbf{A}}}{\partial A_i} \frac{\partial \lambda_{\alpha, \mathbf{k}, \mathbf{A}}}{\partial A_j} + \frac{1}{2\sqrt{(E_0 - \mu)^2 + |\Delta|^2}} \frac{\partial^2}{\partial A_i \partial A_j} \lambda_{\alpha, \mathbf{k}, \mathbf{A}} \right] \Big|_{\mathbf{A} \rightarrow 0} \\ &= -\frac{1}{2} \sum_{\mathbf{k}} \sum_{\alpha} \left[-\frac{1}{4[(E_0 - \mu)^2 + |\Delta|^2]^{3/2}} \frac{\partial \lambda_{\alpha, \mathbf{k}, \mathbf{A}}}{\partial A_i} \frac{\partial \lambda_{\alpha, \mathbf{k}, \mathbf{A}}}{\partial A_j} + \frac{1}{2\sqrt{(E_0 - \mu)^2 + |\Delta|^2}} \frac{\partial^2}{\partial A_i \partial A_j} \lambda_{\alpha, \mathbf{k}, \mathbf{A}} \right] \Big|_{\mathbf{A} \rightarrow 0}. \end{aligned} \quad (\text{E30})$$

Since $\lambda_{\alpha, \mathbf{k}, \mathbf{A}} = \lambda_{\alpha, \mathbf{k}, -\mathbf{A}}$, we have

$$\frac{\partial \lambda_{\alpha, \mathbf{k}, \mathbf{A}}}{\partial A_i} \Big|_{\mathbf{A} \rightarrow 0} = 0. \quad (\text{E31})$$

Then,

$$\begin{aligned}
\mathcal{V}[D_{SW}(0)]_{ij} &= -\frac{1}{4\sqrt{(E_0 - \mu)^2 + |\Delta|^2}} \sum_{\mathbf{k}} \sum_{\alpha} \left[\frac{\partial^2}{\partial A_i \partial A_j} \lambda_{\alpha, \mathbf{k}, \mathbf{A}} \right]_{\mathbf{A} \rightarrow 0} \\
&= -\frac{1}{4\sqrt{(E_0 - \mu)^2 + |\Delta|^2}} \sum_{\mathbf{k}} \left[\frac{\partial^2}{\partial A_i \partial A_j} \text{Tr}[\Delta_{\mathbf{k}, \mathbf{A}} \Delta_{\mathbf{k}, \mathbf{A}}^\dagger] \right]_{\mathbf{A} \rightarrow 0} \\
&= -\frac{|\Delta|^2}{4\sqrt{(E_0 - \mu)^2 + |\Delta|^2}} \sum_{\mathbf{k}} \left[\frac{\partial^2}{\partial A_i \partial A_j} \text{Tr}[P_{\mathbf{k}-\mathbf{A}} P_{\mathbf{k}+\mathbf{A}}] \right]_{\mathbf{A} \rightarrow 0} \\
&= -\frac{|\Delta|^2}{4\sqrt{(E_0 - \mu)^2 + |\Delta|^2}} \sum_{\mathbf{k}} \left[\frac{\partial^2}{\partial A_i \partial A_j} \text{Tr}[P_{\mathbf{k}-\mathbf{A}} P_{\mathbf{k}+\mathbf{A}}] \right]_{\mathbf{A} \rightarrow 0} \\
&= -\frac{|\Delta|^2}{4\sqrt{(E_0 - \mu)^2 + |\Delta|^2}} \sum_{\mathbf{k}} \{ \text{Tr}[\partial_{k_i} \partial_{k_j} P_{\mathbf{k}} P_{\mathbf{k}}] - \text{Tr}[\partial_{k_i} P_{\mathbf{k}} \partial_{k_j} P_{\mathbf{k}}] - \text{Tr}[\partial_{k_j} P_{\mathbf{k}} \partial_{k_i} P_{\mathbf{k}}] + \text{Tr}[P_{\mathbf{k}} \partial_{k_i} \partial_{k_j} P_{\mathbf{k}}] \} \\
&= \frac{|\Delta|^2}{\sqrt{(E_0 - \mu)^2 + |\Delta|^2}} \sum_{\mathbf{k}} \text{Tr}[\partial_{k_i} P_{\mathbf{k}} \partial_{k_j} P_{\mathbf{k}}] = \frac{2|\Delta|^2}{\sqrt{(E_0 - \mu)^2 + |\Delta|^2}} \sum_{\mathbf{k}} [g_{\mathbf{k}}]_{ij}
\end{aligned} \tag{E32}$$

where

$$P_{\mathbf{k}} = U_{\mathbf{k}} U_{\mathbf{k}}^\dagger, \tag{E33}$$

the second last equality uses

$$\text{Tr}[\partial_{k_i} \partial_{k_j} P_{\mathbf{k}} P_{\mathbf{k}}] = \partial_{k_i} (\text{Tr}[\partial_{k_j} P_{\mathbf{k}} P_{\mathbf{k}}]) - \text{Tr}[\partial_{k_j} P_{\mathbf{k}} \partial_{k_i} P_{\mathbf{k}}] = -\text{Tr}[\partial_{k_j} P_{\mathbf{k}} \partial_{k_i} P_{\mathbf{k}}] \tag{E34}$$

which comes from

$$\text{Tr}[\partial_{k_j} P_{\mathbf{k}} P_{\mathbf{k}}] = \frac{1}{2} (\text{Tr}[\partial_{k_j} P_{\mathbf{k}} P_{\mathbf{k}}] + \text{Tr}[P_{\mathbf{k}} \partial_{k_j} P_{\mathbf{k}}]) = \frac{1}{2} \partial_{k_j} \text{Tr}[P_{\mathbf{k}} P_{\mathbf{k}}] = 0, \tag{E35}$$

and

$$[g_{\mathbf{k}}]_{ij} = \frac{1}{2} \text{Tr}[\partial_{k_i} P_{\mathbf{k}} \partial_{k_j} P_{\mathbf{k}}]. \tag{E36}$$

From the expression of Eq. (E26), we can obtain the zero-temperature average value of the electron number, which reads

$$\langle N_e \rangle = -\partial_i \Omega_0(0) = -\partial_i \sum_{\mathbf{k}} \left((E_0 - \mu) - \frac{1}{2} \sum_{\alpha} \sqrt{(E_0 - \mu)^2 + |\Delta|^2} \right) = N \left(1 - \frac{E_0 - i}{\sqrt{(E_0 - \mu)^2 + |\Delta|^2}} \right), \tag{E37}$$

which means the filled portion of the normal-state bands is

$$f = \langle N_e \rangle / (2N) = \frac{1}{2} \left[1 - \frac{E_0 - i}{\sqrt{(E_0 - \mu)^2 + |\Delta|^2}} \right]. \tag{E38}$$

As

$$f(1-f) = \frac{1}{4} - \left(\frac{1}{2} - f \right)^2 = \frac{1}{4} - \frac{1}{4} \frac{(E_0 - \mu)^2}{(E_0 - \mu)^2 + |\Delta|^2} = \frac{1}{4} \frac{|\Delta|^2}{(E_0 - \mu)^2 + |\Delta|^2}, \tag{E39}$$

we arrive at

$$[D_{SW}(0)]_{ij} = 4|\Delta| \sqrt{f(1-f)} \int \frac{d^2 k}{(2\pi)^2} [g_{\mathbf{k}}]_{ij} \tag{E40}$$

Thus, at the mean-field level, the zero-temperature superfluid weight is bounded from below by the Z_2 index, *i.e.*

$$\text{Tr}[D_s(0)] = 4|\Delta| \sqrt{f(1-f)} \int \frac{d^2 k}{(2\pi)^2} \text{Tr}[g_{\mathbf{k}}] \geq \frac{4|\Delta|}{\pi} \sqrt{f(1-f)} \frac{Z_2}{\pi}. \tag{E41}$$

2. Z_2 Bounds for Optical Conductivity and Band Gap

In this part, we discuss how the Z_2 bound affects the optical conductivity. We will consider a 2D non-interacting band insulator with TR symmetry. The optical conductivity tensor reads [5, 50]

$$\sigma_{ij}(\omega + i0^+) = -i \int \frac{d^2k}{(2\pi)^2} \sum_{m,n} \frac{(\varepsilon_{n,\mathbf{k}} - \varepsilon_{m,\mathbf{k}})[A_{\mathbf{k},i}]_{nm}[A_{\mathbf{k},j}]_{mn}}{\omega + \varepsilon_{n,\mathbf{k}} - \varepsilon_{m,\mathbf{k}} + i0^+} (f_n - f_m) , \quad (\text{E42})$$

where 0^+ is an infinitesimal positive real number, $\varepsilon_{n,\mathbf{k}}$ is the n th energy band,

$$f_n = \theta(\varepsilon_n(\mathbf{k} - \mu)) \quad (\text{E43})$$

with μ the chemical potential in the band gap, and

$$[A_{\mathbf{k},i}]_{nm} = i \langle u_{n,\mathbf{k}} | \partial_{k_i} | u_{m,\mathbf{k}} \rangle \quad (\text{E44})$$

with $|u_{n,\mathbf{k}}\rangle$ the periodic part of the Bloch state. It is known that [5]

$$\int_0^\infty d\omega \frac{\text{Re}[\sigma_{ii}(\omega + i0^+)]}{\omega} = \frac{1}{4\pi} \int d^2k g_{ii}(\mathbf{k}) , \quad (\text{E45})$$

where $g_{ij}(\mathbf{k})$ is the quantum metric for the occupied band, and we have use the insulating property of the system. Combined with our Z_2 bound for quantum geometry (Eq. (D10)), we have a Z_2 lower bound for the optical conductivity:

$$\int_0^\infty \frac{d\omega}{\omega} \sum_i \text{Re}[\sigma_{ii}(\omega + i0^+)] = \frac{1}{4\pi} \int d^2k \text{Tr}[g(\mathbf{k})] \geq Z_2 . \quad (\text{E46})$$

In general, the non-interacting Hamiltonian for solids have the following form

$$H = \int d^d r \sum_{s,s'} c_{\mathbf{r},s}^\dagger \left[-\frac{\nabla^2}{2m} \delta_{ss'} - i\nabla \cdot \mathbf{Y}_{ss'}(\mathbf{r}) - i\mathbf{Y}_{ss'}(\mathbf{r}) \cdot \nabla + V(\mathbf{r})\delta_{ss'} \right] c_{\mathbf{r},s'} , \quad (\text{E47})$$

where \mathbf{r} is the position, s is the spin, and

$$\{c_{\mathbf{r},s}^\dagger, c_{\mathbf{r}',s'}\} = \delta(\mathbf{r} - \mathbf{r}')\delta_{ss'} . \quad (\text{E48})$$

$\mathbf{Y}_{ss'}$ accounts for spin-orbit coupling. The form of the Hamiltonian implies a sum rule for the optical conductivity, which reads [50, 52]

$$\int_0^\infty d\omega \text{Re}[\sigma_{ii}(-\omega + i0^+)] = \frac{n\pi}{2m} , \quad (\text{E49})$$

where n is the electron density. Combined with

$$\int_0^\infty \frac{d\omega}{\omega} \sum_i \text{Re}[\sigma_{ii}(\omega + i0^+)] \leq \frac{1}{E_g} \int_0^\infty d\omega \sum_i \text{Re}[\sigma_{ii}(\omega + i0^+)] \quad (\text{E50})$$

with E_g the direct band gap, we arrive at a Z_2 upper bound for the gap

$$E_g \leq \frac{\int_0^\infty d\omega \sum_i \text{Re}[\sigma_{ii}(\omega + i0^+)]}{\int_0^\infty \frac{d\omega}{\omega} \sum_i \text{Re}[\sigma_{ii}(\omega + i0^+)]} = \frac{n\pi}{m} \frac{1}{\frac{1}{4\pi} \int d^2k \text{Tr}[g(\mathbf{k})]} \leq \frac{n\pi}{mZ_2} . \quad (\text{E51})$$

We note that the relation between the gap and the quantum metric was derived in Ref. [51, 52].