

Article Bayesian analysis of hybrid neutron star EOS constraints within a nonlocal color superconducting quark matter model

Alexander Ayriyan^{1,2}, David Blaschke^{1,3,4}, Juan Pablo Carlomagno^{5,6}, Gustavo A. Contrera^{5,6} and Ana Gabriela Grunfeld^{5,7}

- ¹ Institute for Theoretical Physics, University of Wroclaw, Max Born Pl. 9, 50-204, Wroclaw, Poland
- ² A. Alikhanyan National Science Lab, Alikhanyan Brothers street 2, 0036 Yerevan, Armenia
- ³ Helmholtz-Zentrum Dresden-Rossendorf (HZDR), Bautzner Landstrasse 400, 01328 Dresden, Germany
- ⁴ Center for Advanced Systems Understanding (CASUS), Untermarkt 20, 02826 Görlitz, Germany
- ⁵ CONICET, Godoy Cruz 2290, Buenos Aires, Argentina
- ⁶ IFLP, UNLP, CONICET, Facultad de Ciencias Exactas, Diagonal 113 entre 63 y 64, La Plata 1900, Argentina
- ⁷ Departamento de Física, Comisión Nacional de Energía Atómica, Av. Libertador 8250, (1429) Buenos Aires, Argentina
- * Correspondence: alexander.ayriyan@uwr.edu.pl

Abstract: We present a physics-informed Bayesian analysis of equation of state constraints using observational data for masses, radii and tidal deformability of pulsars and a generic class of hybrid neutron star equation of state with color superconducting quark matter on the basis of a recently developed nonlocal chiral quark model. The nuclear matter phase is described within a relativistic density functional model of the DD2 class and the phase transition is obtained by a Maxwell construction. We find the region in the two-dimensional parameter space spanned by the vector meson coupling and the scalar diquark coupling, where three conditions are fulfilled: 1) the Maxwell construction can be performed, 2) the maximum mass of the hybrid neutron star is not smaller than 2.0 M_{\odot} and 3) the onset density of the phase transition is not below the nuclear saturation density $n_0 = 0.15$ fm⁻³. The result of this study shows that the favorable neutron star equation of state has low onset masses for the occurrence of a color superconducting quark matter core between $0.5 - 0.7 M_{\odot}$ and maximum masses in the range $2.15 - 2.22 M_{\odot}$. In the typical mass range of $1.2 - 2.0 M_{\odot}$, the radii of these stars are between 11.9 and 12.4 km, almost independent of the mass. In principle, hybrid stars would allow for larger maximum masses than provided by the hadronic reference equation of state.

Keywords: Bayesian analysis; hybrid neutron stars; color superconductivity; quark deconfinement

1. Introduction

Exploring the phases of matter under extreme conditions, like those in heavy-ion collisions [1] or within neutron stars [2], is a topic that has attracted significant attention over the past decades [3]. We are particularly interested in studying transitions between different phases of strongly interacting matter, like quark-gluon plasma, color superconducting quark matter, and hadronic matter. These transitions have a big impact on astrophysical phenomena like the formation of eccentric binaries and isolated millisecond pulsars (MSPs) [4], binary neutron star mergers [5,6], and even trigger supernova explosions of massive blue supergiant stars [7,8] to mention just a few recent applications.

The status of quark matter as a state of matter in the inner core of neutron stars is still controversial. In particular, since the available data on masses and radii of pulsars do not allow for a firm model-independent conclusion about a quark deconfinement transition, it has been tried to apply agnostic Bayesian analysis methods to check the evidence for a strong phase transition, see [9] and references therein. In their study, the authors of [9] found evidence against a strong first-order (deconfinement) phase transition in neutron stars. However, since the lightest neutron star to be formed as a result of a supernova explosion according to state-of-the-art simulations has a mass of at least $1.1 M_{\odot}$, and there

Citation: Ayriyan, A.; Blaschke, D.; Carlomagno, J.P.; Contrera, G.A.; Grunfeld, A.G Bayesian analysis of hybrid neutron star EOS. *Universe* **2024**, 1,0. https://doi.org/

> Received: Revised: Accepted: Published:

Copyright: © 2025 by the authors. Submitted to <u>Universe</u> for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). is no scenario available that would explain the existence of a lighter neutron star, the recent measurement of a low mass and radius for HESS J1731-347 [10] has been considered very skeptical in the community. Also because the authors themselves could not present an idea about the origin of a neutron star in the subsolar mass region. Therefore, direct evidence for a strong phase transition from mass and radius measurements in the subsolar mass range is presently lacking the observational basis.

In this situation, we want to suggest a physics-informed Bayesian analysis of modern multi-messenger mass and radius measurements along the lines of our earlier work [11], which can now be made with the new, presently available set of observational data. Such a physics-informed study will be based on a class of hybrid equations of state (EOS) within the two-phase formalism that uses separate models for hadronic and quark matter phases joined by a phase transition construction. Given an appropriate hadronic EOS and the choice of the phase transition construction, there are two free parameters of the color superconducting quark matter model to be determined which cannot be reliably fixed with vacuum properties of hadrons. These are the couplings of the vector meson (G_V) and diquark (G_D) currents to the corresponding mean fields. As has been shown in [12] using the three-flavor color superconducting Nambu–Jona-Lasinio (NJL) model [13], that with such a setting the required maximum mass $M_{\rm max} \sim 2 \ M_{\odot}$ can be achieved by a sufficiently large G_V , while an early onset of deconfinement in the range of typical or even subsolar neutron star masses is achieved by a large value of G_D . In a first systematic study employing this setting, in Ref. [14], the regions in the two-dimensional plane of these two coupling strengths have been identified, where no stable hybrid stars were possible and where the onset of deconfinement is below a certain density. In between these two limits, the lines were shown along which the constraint on the minimal value of the maximum mass could be fulfilled. In these early studies the Dirac-Brueckner-Hartree-Fock (DBHF) hadronic EOS [15] was used together with a Maxwell construction for the phase transition. An alternative study by Baym et al. [16] came to similar conclusions for the preferable region of parameter values in the plane of vector meson and diquark coupling constants using, however, a softer APR-like hadronic EOS and a crossover construction [17], see also the discussion in the review [18]. Investigations based on the NJL model of quark matter are plagued with the absence of confinement in that model which becomes problematic when extending its application to the range of finite temperatures, as in supernova explosions and binary neutron star mergers, where it would predict deconfinement at a too low temperature of about 50 - 70 MeV.

In order to overcome this problem, two main routes have been suggested: (A) a confining density functional approach [19] as a relativistic formulation of the string-flip model [20,21] which was recently generalized to include diquark condensation and chiral symmetry [22,23]; and (B) density-dependent quark mass models (see, e.g., [24–31]), in which the bag pressure is interpreted as a density-dependent scalar mean field that can be reinterpreted as a contribution to the dynamical quark mass. The confining models of class (A) can be mapped by a Taylor expansion of the confining density functional w.r.t. the underlying quark bilinears up to second order onto an effective NJL model with medium dependent masses. Equivalently, one can also fit the pressure as a thermodynamic potential of the confining density functional approach by adding a medium dependent bag pressure and vector mean field to the nonlocal NJL model [32,33]. In the present work, we will employ the latter approach to a confining quark matter EOS as the basis for the physics-informed Bayesian analysis to be performed.

2. Hybrid neutron star EOS and fitting results

There are two main strategies to construct hybrid neutron star EOS which would fulfill the maximum mass constraint that requires the most massive hybrid star configuration to have at least 2 M_{\odot} at a radius around 12 km. One can start with a nuclear matter EOS that is relatively soft at supranuclear densities, like the well-known variational EOS by Akmal, Pandharipande and Ravenhall (APR) [34] or its updated versions by Togashi and coworkers [35–37]. Then, a crossover construction like the one by Masuda et al. [17] is required to facilitate the transition to a stiffer quark matter EoS at high densities in a thermodynamically consistent way. This strategy has been applied, e.g., by Baym et al. [16] and by Ayriyan et al. [11], who realized the two-zone interpolation scheme (TZIS) that was described in [18]. See also [23] for the extension of the TZIS to finite temperature and multiple critical endpoints.

The alternative is to start from a stiff nuclear EOS, for which a Maxwell construction to a sufficiently stiff quark matter EOS can be performed, so that despite a softening of the EOS due to the phase transition nevertheless the maximum mass constraint can be fulfilled. As examples for such stiff hadronic EOS we mention the DBHF [15] which was used in [12,14] and the relativistic density functional EOS with density-dependent nucleon-meson couplings, denoted as DD2 [38]which was recently used, e.g., in [22,23,33,39–41]. This latter strategy will be applied in the present work.

2.1. Nuclear matter phase

We employ the DD2 relativistic density functional EOS with the parametrization described in Ref. [38]. This EOS describes well all known nuclear matter saturation properties as well as the nuclear symmetry energy and its slope at saturation density $n_{\text{sat}} = 0.15 \text{ fm}^{-3}$.

2.2. Color superconducting 2SC quark matter

In this work, we consider the scenario of two-flavor color-superconducting quark matter appearing in the cores of NSs at sufficiently high densities. For its description, we employ the 3DFF nonlocal chiral quark model [33] defined by an effective Euclidean action functional that in the case of two light flavors is given by

$$S_{E} = \int d^{4}x \left\{ \bar{\psi}(x) \left(-i\partial \!\!\!/ + \hat{m} - \gamma_{0}\hat{\mu} \right) \psi(x) - \frac{G_{S}}{2} \left[j_{S}^{f}(x) j_{S}^{f}(x) + \eta_{D} \left[j_{D}^{a}(x) \right]^{\dagger} j_{D}^{a}(x) - \eta_{V} j_{V}^{\mu}(x) j_{V}^{\mu}(x) \right] \right\}.$$
(1)

Here $\hat{m} = \text{diag}(m_u, m_d)$ is the current quark mass matrix with the masses for u and d quarks, which are assumed to coincide $m_u = m_d = 2.3$ MeV, whereas $\hat{\mu} = \text{diag}(\mu_u, \mu_d)$ is the diagonal matrix for the chemical potentials of u and d quarks, which are to be adjusted so as to fulfill charge neutrality and β -equilibrium constraints relevant for neutron star matter. As a result remains the baryon chemical potential $\mu_B = 3\mu = 3(\mu_u + \mu_d)/2$ as independent thermodynamic variable. The coupling strength in the scalar meson channel is $G_S = 9.92 \text{ GeV}^{-2}$ and $\eta_V = G_V/G_S(\eta_D = G_D/G_S)$ is the relative coupling in the vector meson (scalar diquark) channel. The currents are given by nonlocal operators based on a separable approximation to the effective one gluon exchange (OGE) model of QCD [42,43],

$$\begin{split} j_{S}^{f}(x) &= \int d^{4}z \, g_{S}(z) \, \bar{\psi}(x + \frac{z}{2}) \, \Gamma_{f} \, \psi(x - \frac{z}{2}) \,, \\ j_{D}^{a}(x) &= \int d^{4}z \, g_{D}(z) \, \bar{\psi}_{C}(x + \frac{z}{2}) \, i \gamma_{5} \tau_{2} \lambda_{a} \, \psi(x - \frac{z}{2}) \,, \\ j_{V}^{\mu}(x) &= \int d^{4}z \, g_{V}(z) \, \bar{\psi}(x + \frac{z}{2}) \, i \gamma^{\mu} \, \psi(x - \frac{z}{2}) \,, \end{split}$$

$$\end{split}$$

where we defined $\psi_C(x) = \gamma_2 \gamma_4 \bar{\psi}^T(x)$ and $\Gamma_f = (\mathbf{1}, i\gamma_5 \tau)$, while τ and λ_a , with a = 2, 5, 7, stand for Pauli and Gell-Mann matrices acting on flavor and color spaces, respectively. The functions $g_i(z)$, i = S, D, V, in Eqs. (2) are nonlocal "instantaneous" form factors (3D-FF) characterizing the effective quark interaction, which depend on the spatial components of the position 4-vector z only. For the nonlocality a Gaussian ansatz is employed which after Fourier transformation to the momentum space reads

$$g_i(\vec{p}) = \exp(-\vec{p}^2/\Lambda_i^2), \ i = S, D,$$
 (3)

with the 3-momentum vector \vec{p} and the effective range $\Lambda_S = \Lambda_D = 885.47$ MeV fitted to Coulomb-gauge lattice QCD data [44]. Note that following [33], in the present work we consider the vector current in Eq. (2) to be local, $g_V(z) = \delta(z)$. This restriction could be easily relaxed, see [45]. The equation of state $p(T, \mu) = T \ln Z(T, \mu)$ for this model follows from the path-integral representation of the partition function $Z(T, \mu)$ which is evaluated at the mean-field level and results in the pressure $p_{MF}(\mu_B)$ which, following [33], is augmented by a μ_B -dependent bag pressure as in Ref. [32], given by the equation

$$B(\mu_B) = B_0 f_{<}(\mu_B)$$
 (4)

with

$$f_{<}(\mu_{B}) = \frac{1}{2} \left[1 - \tanh\left(\frac{\mu_{B} - \mu_{<}}{\Gamma_{<}}\right) \right], \tag{5}$$

where we use $\mu_{<} = 895$ MeV, $\Gamma_{<} = 180$ MeV and $B_0 = 35$ MeV/fm³ as the optimal values to reproduce the astrophysical constraints from multi-messenger astronomy of pulsars. The resulting zero-temperature pressure $p(\mu_B) = p_{MF}(\mu_B) - B(\mu_B)$ leaves the two couplings η_V and η_D as free parameters of the 3DFF color superconducting quark matter model EOS which forms the basis of the present work. Each EOS model of the present study can be characterized as a point in the two-dimensional parameter space spanned by the pair (η_V, η_D) which will therefore play a central role.

In order to perform a Bayesian Analysis, we fit the above introduced quark matter EoS to the constant speed of sound (CSS) form of the EoS [46]

$$p = A \left(\frac{\mu}{\mu_x}\right)^{1+1/c_s^2} - B \tag{6}$$

where the $\mu = \mu_B/3$ is the quark chemical potential and $\mu_x = 1$ GeV sets a scale. Before, such a fit has been provided in [46] for the covariant nonlocal chiral quark model EoS. Here, we present the fit (6) of the 3DFF nonlocal, color superconducting quark matter model [33] in the form

$$A(\eta_D, \eta_V) = a_1 \eta_D^2 + b_1 \eta_V^2 + c_1 \eta_D \eta_V + d_1 \eta_D + e_1 \eta_V + f_1,$$
(7)

$$c_s^2(\eta_D, \eta_V) = a_2\eta_D^2 + b_2\eta_V^2 + c_2\eta_D\eta_V + d_2\eta_D + e_2\eta_V + f_2,$$
(8)

$$B(\eta_D, \eta_V) = a_3\eta_D^2 + b_3\eta_V^2 + c_3\eta_D\eta_V + d_3\eta_D + e_3\eta_V + f_3,$$
(9)

The fitted parameter values are given in Tab. 1.

Table 1. Values of the parameters of Eqs. (7) - (9).

i	parameter	units	a _i	b _i	c _i	d _i	e _i	f_i
1	Α	$\frac{\text{MeV}}{\text{fm}^3}$	88.637	1.861	-30.368	-76.654	2.891	95.324
2	В	$\frac{\text{MeV}}{\text{fm}^3}$	-192.28	-10.180	30.455	390.92	-28.613	-100.95
3	c_s^2		0.142	-0.071	0.066	-0.166	0.134	0.427

2.3. Phase transition construction

We employ the Maxwell construction of the nuclear-to-quark matter phase transition. The resulting EoS is characterized by a critical pressure $P_c = P_h(\mu_c) = P_q(\mu_c)$ for the onset of the transition from the hadronic phase $P_h(\mu)$ for $\mu \le \mu_c$ to the 2SC quark matter phase $P_q(\mu)$ for $\mu \ge \mu_c$ with a jump in energy density $\Delta \varepsilon = \varepsilon_q(\mu_c) - \varepsilon_h(\mu_c)$. In Fig. 1, we show the hybrid neutron star EoS after the Maxwell construction has been performed. In the left

panel, we show the whole range of energy densities considered here, while in the right panel a zoom into the region relevant for the phase transition is shown.



Figure 1. Diagram of pressure vs. energy density, which shows the jump in energy density at the phase transition from hadronic DD2 EoS (green solid line) to the color superconducting quark matter EoS in its CSS parametrization (left panel). The right panel shows a closeup of the phase transition region at lower energy densities than that for which the maximum mass configuration of the hadronic DD2 EoS is obtained (indicated by a plus sign on the left panel).

For some values of the parameters (η_D, η_V) there is no crossing of the hadronic and quark-matter pressure curves so that no Maxwell construction can be performed. In Fig. 2, we depict the jump in baryon density $\Delta n = n_q(\mu_c) - n_h(\mu_c)$, where $n_i = dP_i(\mu)/d\mu$ for i = h, q as a contour plot in the $\eta_D - \eta_V$ plane.



Figure 2. Diagram of the coupling constants $\eta_D - \eta_V$, which shows jump in baryon density Δn at the phase transition from the hadronic DD2 to the color superconducting 3DFF quark matter EoS. In the parameter region filled with greyscale dots a Maxwell construction is possible. The white spots in the middle of that region stand for parameter sets for which the transition would degenerate to a crossover with $\Delta n = 0$.

3. Mass, radius and tidal deformability fom TOV solutions

In Fig. 3 we show the dependence of the neutron star mass on its radius (left panel) and on its central energy density (right panel) for the EoS parameterizations provided in the previous section. We note that there are cases of hybrid star sequences which have a maximum mass exceeding that of the hadronic DD2 reference EOS shown by the green

solid line in both panels. On the right panel of that figure we indicate by a green "X" the location of the maximum mass configuration of the purely hadronic case which has $M_{\text{max}} = 2.4 M_{\odot}$ and a central energy density $\varepsilon_c^{\text{max}} = 1150 \text{ MeV/fm}^3$.



Figure 3. Mass-radius relation (left panel) and Mass versus central energy density (right panel) for the hybrid EoS with a phase transition from the hadronic DD2 EoS to color superconducting quark matter described in the nonlocal 3DFF model. The green lines in both panels highlight the results for the hadronic neutron star sequences obtained with the DD2 EOS.

In Fig. 4 we show the tidal deformability as a function of the neutron star mass for the hadronic DD2 reference EOS (red solid line) and the hybrid star sequences (grey lines) of the present study. The data point with error bars corresponds to the gravitational wave analysis of the LIGO Virgo Collaboration for the inspiral phase of the binary neutron star merger GW170817 [47]. The comparison indicates that hybrid star sequences with an early onset of quark deconfinement at masses well below $1.4 M_{\odot}$ lead to a softening of the EOS at the onset energy density which entails a reduction of the value of the tidal deformability at that mass and thus leads to a fulfillment of the tidal deformability constraint while the DD2 EOS was slightly too stiff to match that constraint.



Figure 4. Tidal deformability vs. mass for the set of hybrid EoS introduced in this work. The green line highlights the result for the hadronic DD2 EOS and the green cross indicates the maximum mass configuration of that sequence. The data point with error bars corresponds to the gravitational wave analysis of the LIGO Virgo Collaboration for the inspiral phase of the binary neutron star merger GW170817 [47].

4. Bayesian analysis for optimal EOS

Bayesian analysis was performed using modern and highly selective observational constraints. The lower limit for the maximum mass was taken as $M = 2.10^{+0.04}_{-0.04} M_{\odot}$ [48] for PSR J0348+0432 and $M = 2.35^{+0.17}_{-0.17} M_{\odot}$ [49] for the "black widow" pulsar PSR J0952-0607. For mass-radius constraints, the following values were used:

- (A) PSR J0030+0451: $M = 1.44^{+0.07}_{-0.07} M_{\odot}$ and $R = 13.7^{+2.6}_{-1.5}$ km [50]; (B) PSR J0740+6620: $M = 2.08^{+0.07}_{-0.07} M_{\odot}$ and $R = 13.7^{+2.6}_{-1.5}$ km [51]; (C) PSR J0437-4715: $M = 1.418^{+0.037}_{-0.037} M_{\odot}$ and $R = 11.36^{+0.95}_{-0.63}$ km [52]; (D) HESS J1731-347: $M = 0.77^{+0.20}_{-0.17} M_{\odot}$ and $R = 10.04^{+0.86}_{-0.78}$ km [10].

Additionally, the tidal deformability constraint $\Lambda_{1.4} = 290^{+290}_{-120}$, obtained from the binary neutron star merger GW170817 [47] was applied to the analysis.

The result of the Bayesian analysis with modern observational constraints as described above is shown in Fig. 5. We note that the 60% credible region forms a triangular shaped area very similar to the almost cigar-shaped region that was obtained before, e.g., in Refs. [16], [40] and in [41], but not from a Bayesian analysis. The most favorable parameter set can be read-off the peak positions for the marginalized probability distributions in the η_V and η_D -directions, respectively. These peak positions are very close to the best choice values $\eta_V = 0.5$ and $\eta_D = 1.1$ that were found heuristically in [33], where the present class of hybrid EOS was introduced.



Figure 5. Bayesian analysis of the most likely parameter values in the $\eta_V - \eta_D$ plane under modern observational constraints for masses, radii and the tidal deformability of neutron stars.

5. Results

In Fig. 6 we show the different cases of phase transitions, again overlaid to the credible regions of the Bayesian analysis with the following color coding:

- White: No Maxwell transition,

- Gray: Maxwell transition occurs after the DD2 instability value (no stable hybrid stars),

- Orange: Maxwell transition starts before the DD2 instability value, with a jump to the DD2 instability (no stable hybrid stars),

- Green: Maxwell transition ends before the DD2 instability value (stable hybrid stars exist)

- Blue: Maxwell transition ends before the DD2 instability value and satisfies the Seidov criterion [53] (extremely light twin stars).

In Fig. 7 we display observables in the plane of the EOS parameters η_V and η_D as color-bubble diagrams overlaid to the Bayesian likelihood contours for the 60% and 90% credibility regions. On the left panel we show the difference in maximum mass between hybrid stars and the DD2 hadronic reference EOS with a maximum mass of 2.4 M_{\odot} (the



Figure 6. Phase transition categories in the $\eta_V - \eta_D$ diagram, overlaid with the 60% (blue dashed line) and 90% (red dashed lines) credible regions of the Bayesian analysis. For details, see text.

black solid neutrality line). The yellow bubble region indicates that no stable hybrid stars are possible there.

We want to mention that in [16], the maximum possible mass was at 2.35 M_{\odot} and in [40,41] at 2.42 M_{\odot} . Thus the corresponding parameter sets of the latter works would lie in the corner of the green region, between the 60% and 90% confidence regions, while the present class of hybrid EOS has the parameter sets with the largest maximum masses at the very corner of the green region, outside even the 90% confidence region.

The reason for this region of highest-mass hybrid EOS to be disfavored lies in the fact that they predict hybrid star sequences that are not sufficiently compact in the mass region of 1.4 M_{\odot} , where the tidal deformability was measured for the binary neutron star merger GW170817 and requires radii in the range $R_{1.4} = 10.94 - 12.61$ km at 90% confidence [54]. On the right panel we show the tidal deformability constraint: $70 < \Lambda < 580$ and can observe that the 60% credible region (blue dashed line) encloses the pairs of quark matter parameter values for which the tidal deformability constraint is fulfilled.



Figure 7. Color-bubble diagrams for observables in the $\eta_V - \eta_D$ plane of EOS parameters overlaid to the Bayesian likelihood contours for the 60% and 90% credibility regions. Left panel: Difference in maximum mass between hybrid stars and the DD2 hadronic reference EOS with a maximum mass of 2.4 M_{\odot} (the black solid neutrality line). Right panel: Tidal deformability constraint: 70 < Λ < 580.

In Fig. 8 we show the squared speed of sound as a function of the energy density for the hadronic DD2 reference EOS (green solid line), the hybrid star EOS with color superconducting 3DFF quark matter fitted to $c_S^2 = const$ (grey lines) and the cases favored after Bayesian analysis shown in the legend. In the first-order phase transition region holds $c_S^2 = 0$. The maximum energy density that can be reached in the center of a neutron star made of DD2 matter for its maximum mass configuration (vertical green dashed line) gets reduced (vertical green dotted line) when a deconfinement transition is possible, but does not lead to stable hybrid stars.



Figure 8. Squared speed of sound as a function of the energy density for the hadronic DD2 reference EOS (green solid line), the hybrid star EOS with color superconducting 3DFF quark matter fitted to $c_s^2 = const$ (grey lines) and the cases favored after Bayesian analysis shown in the legend. In the first-order phase transition region holds $c_s^2 = 0$. The vertical green dashed (dotted) line indicates the maximum energy density in neutron stars of DD2 matter without (with) the possibility of quark deconfinement.

The final result of this Bayesian analysis study for the class of hybrid EOS with the hadronic DD2 RDF reference EOS and the nonlocal 3DFF color superconducting quark matter EOS fitted to the CSS form of EOS is shown in Fig. 9. We note that the EOS parametrisations favored by the Bayesian analysis with the displayed set of observational constraints listed in section 4 provide evidence for an early onset of deconfinement below 0.7 M_{\odot} and maximum masses not exceeding 2.25 M_{\odot} . In the region of observed pulsar masses, $M \sim 1.1 - 2.1 M_{\odot}$, the radii are approximately mass-independent in the range of $R \sim 12$ km. This feature has been discussed in Ref. [55] as an indicator for maximum masses in the range $M_{\text{max}} \sim 2.3 - 2.4 M_{\odot}$ within a model-independent study. This is consistent with our result shown in Fig. 9.

6. Conclusions

In this work, we have performed for the first time a physics-informed Bayesian analysis of the EOS constraints from modern observational data for masses, radii and the tidal deformability of pulsars within the class of hybrid EOS with a stiff nuclear matter phase, color superconducting (two-flavor) quark matter and a Maxwell construction for the phase transition. It supersedes an earlier physics-informed Bayesian analysis study [11] which was performed within the alternative scheme that uses a soft nuclear matter EOS



Figure 9. Mass-radius diagram showing the hadronic DD2 reference EOS (green solid line) and the hybrid star sequences correspoding to the hybrid EOS Maxwell-constructed from it with the CSS fitted form of the nonlocal 3DFF color superconducting quark matter EOS (grey lines). The EOS parametrizations favored by the Bayesian analysis with the modern mass-radius constraints (colored and labelled regions) are shown by the colored lines explained in the legend.

together with a two-zone interpolation scheme for the transition to color superconducting quark matter. The main result are constrained regions of 60% and 90% credibility in the two-dimensional EOS parameter space of the vector meson and diquark couplings ($\eta_V - \eta_D$ plane) which allow to further constrain the triangular-shaped region that was found earlier in heuristic studies without employing Bayesian analysis methods [14,16,40,41]. We found that the upper right corner of that triangle which corresponds to hybrid stars with maximum masses above the purely hadronic maximum mass of 2.41 M_{\odot} are incompatible with the 60% credibility region of the Bayesian analysis! This appears to be a consequence of the tidal deformability constraint which disfavors these high-maximum mass hybrid sequences because of their insufficient compactness.

The main result of the study is the delineation of the generic structures in the $\eta_V - \eta_D$ plane:

- the rightmost line of proportionality $\eta_V \propto \eta_D$ with the early onset of deconfinement, eventually accompanied with the mass twin phenomenon (blue dots on Fig. 6),
- the region of stable hybrid stars (green dots on Fig. 6),
- the upper right corner of this region, where the hybrid stars with highest maximum mass are found which are eventually disfavored because of missing compactness. In the present work, this corner is at $\eta_V = 1.0$, $\eta_D = 1.2$, corresponding to $M_{\text{max}} = 2.57 M_{\odot}$ at $R_{\text{max}} = 12.6$ km.
- the region parameter region, where the onset of deconfinement is before the maximum central energy density of purely hadronic neutron stars, $\varepsilon_{c, \text{ onset}} < \varepsilon_{c, \text{ max}}$ (yellow dots on Fig. 6), where there are no stable hybrid stars but due to the presence of a deconfinement transition in the EOS the maximum mass is lowered relative to the purely hadronic case,
- the region, where the onset of deconfinement occurs after the maximum central energy density of purely hadronic neutron stars, $\varepsilon_{c, \text{ onset}} > \varepsilon_{c, \text{ max}}$ (grey dots on Fig. 6)

Outside the region highlighted by colored dots in Fig. 6, there is no Maxwell construction of a phase transition possible so that these parameter values of the 3DFF color superconducting quark matter EOS have no influence on the mass-radius relation of the neutron stars.

The next step in the development of this physics-informed Bayesian analysis method is the implementation of the requirement that the high-density asymptotics of the family of hybrid EOS should be compatible with the benchmark from perturbative QCD at densities $n \ge 40 n_0$, see [56,57]. A promising and straightforward strategy to reach this goal is the relaxation of the locality assumption for the vector mean field. This has recently been demonstrated by Ivanytskyi [45]. Work in this direction is in preparation.

Funding: A.A. and D.B. were supported by NCN under grant No. 2021/43/P/ST2/03319. A.G.G., G.A.C., and J.P.C. would like to acknowledge CONICET, ANPCyT and UNLP (Argentina) for financial support under grants No. PIP 2022-2024 GI - 11220210100150CO, PICT19-00792, PICT22-03-00799 and X960, respectively.

Acknowledgments: We thank Dr. Oleksii Ivanytskyi for providing us with the template figure of present observational mass and radius constraints for pulsars that we used in preparing our figure 9.

References

- Sorensen, A.; et al. Dense nuclear matter equation of state from heavy-ion collisions. Prog. Part. Nucl. Phys. 2024, 134, 104080, [arXiv:nucl-th/2301.13253]. https://doi.org/10.1016/j.ppnp.20 23.104080.
- 2. Blaschke, D.; Schaffner-Bielich, J.; Schulze, H.J., Eds. <u>Exotic Matter in Neutron Stars</u>; Vol. 52, European Physical Journal A, Springer, 2016.
- Rezzolla, L.; Pizzochero, P.; Jones, D.I.; Rea, N.; Vidaña, I., Eds. <u>The Physics and Astrophysics of Neutron</u> Vol. 457, <u>Astrophysics and Space Science Library</u>, Springer, 2018. <u>https://doi.org/10.1007/97</u> 8-3-319-97616-7.
- Chanlaridis, S.; Ohse, D.; Antoniadis, J.; Blaschke, D.; Alvarez-Castillo, D.E.; Danchev, V.; Misra, D.; Langer, N. Formation of twin compact stars in low-mass X-ray binaries: Implications on eccentric and isolated millisecond pulsar populations 2024. [arXiv:astro-ph.HE/2409.04755].
- Bauswein, A.; Bastian, N.U.F.; Blaschke, D.B.; Chatziioannou, K.; Clark, J.A.; Fischer, T.; Oertel, M. Identifying a first-order phase transition in neutron star mergers through gravitational waves. <u>Phys. Rev. Lett.</u> 2019, <u>122</u>, 061102, [arXiv:astro-ph.HE/1809.01116]. https://doi.org/10 .1103/PhysRevLett.122.061102.
- Bauswein, A.; Blacker, S.; Vijayan, V.; Stergioulas, N.; Chatziioannou, K.; Clark, J.A.; Bastian, N.U.F.; Blaschke, D.B.; Cierniak, M.; Fischer, T. Equation of state constraints from the threshold binary mass for prompt collapse of neutron star mergers. <u>Phys. Rev. Lett.</u> 2020, <u>125</u>, 141103, [arXiv:astro-ph.HE/2004.00846]. https://doi.org/10.1103/PhysRevLett.125.141103.
- Fischer, T.; Bastian, N.U.F.; Wu, M.R.; Baklanov, P.; Sorokina, E.; Blinnikov, S.; Typel, S.; Klähn, T.; Blaschke, D.B. Quark deconfinement as a supernova explosion engine for massive blue supergiant stars. <u>Nature Astron.</u> 2018, 2, 980–986, [arXiv:astro-ph.HE/1712.08788]. https: //doi.org/10.1038/s41550-018-0583-0.
- Bauswein, A.; Blaschke, D.; Fischer, T., Effects of a strong phase transition on supernova explosions, compact stars and their mergers; 2022; [arXiv:nucl-th/2203.17188]. https://doi.org/ 10.1142/9789811220944_0008.
- Brandes, L.; Weise, W.; Kaiser, N. Evidence against a strong first-order phase transition in neutron star cores: Impact of new data. <u>Phys. Rev. D</u> 2023, <u>108</u>, 094014, [arXiv:nucl-th/2306.06218]. https://doi.org/10.1103/PhysRevD.108.094014.
- 10. Doroshenko, V.; Suleimanov, V.; Pühlhofer, G.; Santangelo, A. A strangely light neutron star within a supernova remnant. Nat. Astron. **2022**, *6*, 1444–1451.
- Ayriyan, A.; Blaschke, D.; Grunfeld, A.G.; Alvarez-Castillo, D.; Grigorian, H.; Abgaryan, V. Bayesian analysis of multimessenger M-R data with interpolated hybrid EoS. <u>Eur. Phys. J. A</u> 2021, 57, 318, [arXiv:astro-ph.HE/2102.13485].
- Klahn, T.; Blaschke, D.; Sandin, F.; Fuchs, C.; Faessler, A.; Grigorian, H.; Ropke, G.; Trumper, J. Modern compact star observations and the quark matter equation of state. <u>Phys. Lett. B</u> 2007, 654, 170–176, [nucl-th/0609067]. https://doi.org/10.1016/j.physletb.2007.08.048.
- Blaschke, D.; Fredriksson, S.; Grigorian, H.; Oztas, A.M.; Sandin, F. The Phase diagram of three-flavor quark matter under compact star constraints. <u>Phys. Rev. D</u> 2005, <u>72</u>, 065020, [hep-ph/0503194]. https://doi.org/10.1103/PhysRevD.72.065020.
- 14. Klähn, T.; Łastowiecki, R.; Blaschke, D.B. Implications of the measurement of pulsars with two solar masses for quark matter in compact stars and heavy-ion collisions: A Nambu–Jona-

Lasinio model case study. Phys. Rev. D **2013**, <u>88</u>, 085001, [arXiv:nucl-th/1307.6996]. https://doi.org/10.1103/PhysRevD.88.085001.

- 15. Fuchs, C. The Relativistic Dirac-Brueckner approach to nuclear matter. Lect. Notes Phys. 2004, 641, 119–146, [nucl-th/0309003]. https://doi.org/10.1007/978-3-540-39911-7_4.
- Baym, G.; Furusawa, S.; Hatsuda, T.; Kojo, T.; Togashi, H. New Neutron Star Equation of State with Quark-Hadron Crossover. <u>Astrophys. J.</u> 2019, <u>885</u>, 42, [arXiv:astro-ph.HE/1903.08963]. https://doi.org/10.3847/1538-4357/ab441e.
- 17. Masuda, K.; Hatsuda, T.; Takatsuka, T. Hadron–quark crossover and massive hybrid stars. PTEP **2013**, 2013, 073D01, [arXiv:nucl-th/1212.6803]. https://doi.org/10.1093/ptep/ptt045.
- Baym, G.; Hatsuda, T.; Kojo, T.; Powell, P.D.; Song, Y.; Takatsuka, T. From hadrons to quarks in neutron stars: a review. <u>Rept. Prog. Phys.</u> 2018, <u>81</u>, 056902, [arXiv:astro-ph.HE/1707.04966]. https://doi.org/10.1088/1361-6633/aaae14.
- Kaltenborn, M.A.R.; Bastian, N.U.F.; Blaschke, D.B. Quark-nuclear hybrid star equation of state with excluded volume effects. Phys. Rev. D 2017, <u>96</u>, 056024, [arXiv:astro-ph.HE/1701.04400]. https://doi.org/10.1103/PhysRevD.96.056024.
- Horowitz, C.J.; Moniz, E.J.; Negele, J.W. Hadron structure in a simple model of quark/nuclear matter. Phys. Rev. D 1985, 31, 1689–1699. https://doi.org/10.1103/PhysRevD.31.1689.
- Ropke, G.; Blaschke, D.; Schulz, H. Pauli Quenching Effects in a Simple String Model of Quark / Nuclear Matter. Phys. Rev. D 1986, 34, 3499–3513. https://doi.org/10.1103/PhysRevD.34.3499.
- 22. Ivanytskyi, O.; Blaschke, D. Density functional approach to quark matter with confinement and color superconductivity. <u>Phys. Rev. D</u> 2022, <u>105</u>, 114042, [arXiv:nucl-th/2204.03611]. https://doi.org/10.1103/PhysRevD.105.114042.
- Ivanytskyi, O.; Blaschke, D. A new class of hybrid EoS with multiple critical endpoints for simulations of supernovae, neutron stars and their mergers. <u>Eur. Phys. J. A</u> 2022, <u>58</u>, 152, [arXiv:nucl-th/2205.03455]. https://doi.org/10.1140/epja/s10050-022-00808-5.
- 24. Fowler, G.N.; Raha, S.; Weiner, R.M. Confinement and Phase Transitions. <u>Z. Phys. C</u> 1981, <u>9</u>, 271. https://doi.org/10.1007/BF01410668.
- Plumer, M.; Raha, S.; Weiner, R.M. Effect of confinement on the velocity of sound in quark-gluon plasma. Phys. Lett. B 1984, 139, 198–202. https://doi.org/10.1016/0370-2693(84)91244-9.
- Wen, X.J.; Zhong, X.H.; Peng, G.X.; Shen, P.N.; Ning, P.Z. Thermodynamics with density and temperature dependent particle masses and properties of bulk strange quark matter and strangelets. <u>Phys. Rev. C</u> 2005, <u>72</u>, 015204, [hep-ph/0506050]. https://doi.org/10.1103/ PhysRevC.72.015204.
- Yin, S.y.; Su, R.K. Consistent thermodynamic treatment for a quark-mass density-dependent model. <u>Phys. Rev. C</u> 2008, <u>77</u>, 055204, [arXiv:nucl-th/0801.2813]. https://doi.org/10.1103/ PhysRevC.77.055204.
- Xia, C.J.; Peng, G.X.; Chen, S.W.; Lu, Z.Y.; Xu, J.F. Thermodynamic consistency, quark mass scaling, and properties of strange matter. <u>Phys. Rev. D</u> 2014, <u>89</u>, 105027, [arXiv:hep-ph/1405.3037]. https://doi.org/10.1103/PhysRevD.89.105027.
- Li, A.; Zuo, W.; Peng, G.X. Massive hybrid stars with a first order phase transition. Phys. Rev. C 2015, 91, 035803, [arXiv:astro-ph.SR/1503.02739]. https://doi.org/10.1103/PhysRevC.91.035 803.
- Lugones, G.; Grunfeld, A.G. Cold dense quark matter with phenomenological medium effects: A self-consistent formulation of the quark-mass density-dependent model. Phys. Rev. D 2023, 107, 043025, [arXiv:nucl-th/2209.03455]. https://doi.org/10.1103/PhysRevD.107.043025.
- 31. Issifu, A.; da Silva, F.M.; Menezes, D.P. Proto-strange quark stars from density-dependent quark mass model. <u>Eur. Phys. J. C</u> 2024, 84, 463, [arXiv:nucl-th/2311.12511]. https://doi.org/10.114 0/epjc/s10052-024-12828-0.
- 32. Alvarez-Castillo, D.E.; Blaschke, D.B.; Grunfeld, A.G.; Pagura, V.P. Third family of compact stars within a nonlocal chiral quark model equation of state. <u>Phys. Rev. D</u> 2019, <u>99</u>, 063010, [arXiv:hep-ph/1805.04105]. https://doi.org/10.1103/PhysRevD.99.063010.
- 33. Contrera, G.A.; Blaschke, D.; Carlomagno, J.P.; Grunfeld, A.G.; Liebing, S. Quark-nuclear hybrid equation of state for neutron stars under modern observational constraints. <u>Phys. Rev. C</u> 2022, 105, 045808, [arXiv:nucl-th/2201.00477]. https://doi.org/10.1103/PhysRevC.105.045808.
- 34. Akmal, A.; Pandharipande, V.R.; Ravenhall, D.G. The Equation of state of nucleon matter and neutron star structure. <u>Phys. Rev. C</u> **1998**, <u>58</u>, 1804–1828, [nucl-th/9804027]. https://doi.org/10.1103/PhysRevC.58.1804.

- 35. Togashi, H.; Takano, M. Variational study for the equation of state of asymmetric nuclear matter at finite temperatures. <u>Nucl. Phys. A</u> 2013, <u>902</u>, 53–73, [arXiv:nucl-th/1302.4261]. https://doi.org/10.1016/j.nuclphysa.2013.02.014.
- 36. Togashi, H.; Hiyama, E.; Yamamoto, Y.; Takano, M. Equation of state for neutron stars with hyperons by the variational method. <u>Phys. Rev. C</u> 2016, <u>93</u>, 035808, [arXiv:nucl-th/1602.08106]. https://doi.org/10.1103/PhysRevC.93.035808.
- Togashi, H.; Nakazato, K.; Takehara, Y.; Yamamuro, S.; Suzuki, H.; Takano, M. Nuclear equation of state for core-collapse supernova simulations with realistic nuclear forces. <u>Nucl. Phys. A</u> 2017, 961, 78–105, [arXiv:nucl-th/1702.05324]. https://doi.org/10.1016/j.nuclphysa.2017.02.010.
- Typel, S.; Ropke, G.; Klähn, T.; Blaschke, D.; Wolter, H.H. Composition and thermodynamics of nuclear matter with light clusters. Phys. Rev. 2010, C81, 015803, [arXiv:nucl-th/0908.2344]. https://doi.org/10.1103/PhysRevC.81.015803.
- Carlomagno, J.P.; Contrera, G.A.; Grunfeld, A.G.; Blaschke, D. Thermal twin stars within a hybrid equation of state based on a nonlocal chiral quark model compatible with modern astrophysical observations. <u>Phys. Rev. D</u> 2024, <u>109</u>, 043050, [arXiv:nucl-th/2312.01975]. https: //doi.org/10.1103/PhysRevD.109.043050.
- 40. Gärtlein, C.; Ivanytskyi, O.; Sagun, V.; Blaschke, D. Hybrid star phenomenology from the properties of the special point. <u>Phys. Rev. D</u> 2023, <u>108</u>, 114028, [arXiv:nucl-th/2301.10765]. https://doi.org/10.1103/PhysRevD.108.114028.
- 41. Gärtlein, C.; Sagun, V.; Ivanytskyi, O.; Blaschke, D.; Lopes, I. Fastest spinning millisecond pulsars: indicators for quark matter in neutron stars? **2024**. [arXiv:nucl-th/2412.07758].
- Blaschke, D.B.; Gomez Dumm, D.; Grunfeld, A.G.; Klahn, T.; Scoccola, N.N. Hybrid stars within a covariant, nonlocal chiral quark model. <u>Phys. Rev.</u> 2007, <u>C75</u>, 065804, [arXiv:nucl-th/nuclth/0703088]. https://doi.org/10.1103/PhysRevC.75.065804.
- Gomez Dumm, D.; Blaschke, D.B.; Grunfeld, A.G.; Scoccola, N.N. Phase diagram of neutral quark matter in nonlocal chiral quark models. <u>Phys. Rev.</u> 2006, <u>D73</u>, 114019, [arXiv:hep-ph/hepph/0512218]. https://doi.org/10.1103/PhysRevD.73.114019.
- 44. Burgio, G.; Schrock, M.; Reinhardt, H.; Quandt, M. Running mass, effective energy and confinement: the lattice quark propagator in Coulomb gauge. <u>Phys. Rev. D</u> 2012, <u>86</u>, 014506, [arXiv:hep-lat/1204.0716]. https://doi.org/10.1103/PhysRevD.86.014506.
- Ivanytskyi, O. Asymptotically conformal CFL quark matter within a nonlocal chiral quark model 2024. [arXiv:hep-ph/2409.05859].
- Shahrbaf, M.; Antić, S.; Ayriyan, A.; Blaschke, D.; Grunfeld, A.G. Constraining free parameters of a color superconducting nonlocal Nambu–Jona-Lasinio model using Bayesian analysis of neutron stars mass and radius measurements. <u>Phys. Rev. D</u> 2023, <u>107</u>, 054011, [arXiv:nuclth/2105.00029]. https://doi.org/10.1103/PhysRevD.107.054011.
- 47. Abbott, B.P.; et al. GW170817: Measurements of neutron star radii and equation of state. <u>Phys.</u> Rev. Lett. **2018**, 121, 161101, [arXiv:gr-qc/1805.11581].
- Antoniadis, J.; Freire, P.C.C.; Wex, N.; Tauris, T.M.; Lynch, R.S.; van Kerkwijk, M.H.; Kramer, M.; Bassa, C.; Dhillon, V.S.; Driebe, T.; et al. A Massive Pulsar in a Compact Relativistic Binary. Science 2013, <u>340</u>, 448, [arXiv:astro-ph.HE/1304.6875]. https://doi.org/10.1126/science.123323
- Romani, R.W.; Kandel, D.; Filippenko, A.V.; Brink, T.G.; Zheng, W. PSR J0952–0607: The Fastest and Heaviest Known Galactic Neutron Star. <u>Astrophys. J. Lett.</u> 2022, <u>934</u>, L17, [arXiv:astroph.HE/2207.05124]. https://doi.org/10.3847/2041-8213/ac8007.
- Miller, M.C.; Lamb, F.K.; Dittmann, A.J.; Bogdanov, S.; Arzoumanian, Z.; Gendreau, K.C.; Guillot, S.; Harding, A.K.; Ho, W.C.G.; Lattimer, J.M.; et al. PSR J0030+0451 Mass and Radius from NICER Data and Implications for the Properties of Neutron Star Matter. <u>The Astrophysical</u> Journal Letters 2019, <u>887</u>, L24. <u>https://doi.org/10.3847/2041-8213/ab50c5</u>.
- Miller, M.C.; Lamb, F.K.; Dittmann, A.J.; Bogdanov, S.; Arzoumanian, Z.; Gendreau, K.C.; Guillot, S.; Ho, W.C.G.; Lattimer, J.M.; Loewenstein, M.; et al. The Radius of PSR J0740+6620 from NICER and XMM-Newton Data. <u>The Astrophysical Journal Letters</u> 2021, <u>918</u>, L28. https: //doi.org/10.3847/2041-8213/ac089b.
- Choudhury, D.; Salmi, T.; Vinciguerra, S.; Riley, T.E.; Kini, Y.; Watts, A.L.; Dorsman, B.; Bogdanov, S.; Guillot, S.; Ray, P.S.; et al. A NICER View of the Nearest and Brightest Millisecond Pulsar: PSR J0437–4715. <u>The Astrophysical Journal Letters</u> 2024, <u>971</u>, L20. https://doi.org/10.3847/20 41-8213/ad5a6f.
- 53. Seidov, Z.F. The Stability of a Star with a Phase Change in General Relativity Theory. <u>Soviet</u> Astronomy **1971**, <u>15</u>, 347.

- 54. Dietrich, T.; Coughlin, M.W.; Pang, P.T.H.; Bulla, M.; Heinzel, J.; Issa, L.; Tews, I.; Antier, S. Multimessenger constraints on the neutron-star equation of state and the Hubble constant. Science 2020, <u>370</u>, 1450–1453, [arXiv:astro-ph.HE/2002.11355]. https://doi.org/10.1126/science. abb4317.
- 55. Ferreira, M.; Providência, C. Constraining neutron star matter from the slope of the mass-radius curves. <u>Phys. Rev. D</u> 2024, <u>110</u>, 063018, [arXiv:nucl-th/2406.12582]. https://doi.org/10.1103/PhysRevD.110.063018.
- Komoltsev, O.; Kurkela, A. How Perturbative QCD Constrains the Equation of State at Neutron-Star Densities. <u>Phys. Rev. Lett.</u> 2022, <u>128</u>, 202701, [arXiv:nucl-th/2111.05350]. https://doi.org/ 10.1103/PhysRevLett.128.202701.
- 57. Gorda, T.; Komoltsev, O.; Kurkela, A. Ab-initio QCD Calculations Impact the Inference of the Neutron-star-matter Equation of State. <u>Astrophys. J.</u> **2023**, <u>950</u>, 107, [arXiv:nucl-th/2204.11877]. https://doi.org/10.3847/1538-4357/acce3a.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.