

The Lie symmetry algebra of the Longstaff-Schwartz model

OUKNINE Anas*

Univ Rouen Normandie, CNRS, Normandie Univ, LMRS UMR 6085, F-76000 Rouen
France

Email: anas.ouknine1@univ-rouen.fr

January 3, 2025

Abstract - This study uses Lie's theory of symmetries to compute the symmetry group of a class of partial differential equations parameterized by four constants: $u_t = -((a - bx)u_x + (d - ey)u_y + \frac{x}{2}u_{xx} + \frac{y}{2}u_{yy})$; under the various conditions on the constants a, b, d and e , we deduce the largest and smallest Lie algebra of symmetries, and we also determined the structure of these algebras.

Keywords - Stochastic differential equation, Longstaff-Schwartz model, Kolmogorov backward equation, Itô formula, Lie algebra, symmetry group, prolongation formula, determining system, infinitesimal criterion of symmetry(Lie condition).

AMS Subject classifications - 34A26, 91B28 ,60H10

1 Introduction

The Longstaff-Schwartz model is a financial pricing model used to value American-style options. It was developed by Francis Longstaff and Eduardo Schwartz in 1992[13]. The model is widely used in the field of quantitative finance and is particularly effective for valuing options on assets that have complex underlying dynamics.

The Longstaff-Schwartz model is base on a two-factor model based on Cox Ingersoll and Ross (CIR) process

$$\begin{cases} dX_t = (a - bX_t)dt + \sqrt{X_t}dB_t^1 \\ dY_t = (d - eY_t)dt + \sqrt{Y_t}dB_t^2 \end{cases} \quad (1.1)$$

with B^1 and B^2 are two independent Brownian motions.

The weighted sum of these factors is considered to form the short-term interest rate:

$$r_t = \alpha X_t + \beta Y_t$$

with α and β nonnegative constants.

One of the main motivations for developing a multifactor model of the term structure is that single-factor models imply perfect correlation in the instantaneous returns on bonds of all maturities, a characteristic clearly inconsistent with reality.

The purpose of this study is to determine the symmetry groups of the class of (1+2) parabolic linear second-order partial differential equations:

$$u_t = -\mathcal{L}u \quad (1.2)$$

where

$$\mathcal{L}u = (a - bx)u_x + (d - ey)u_y + \frac{x}{2}u_{xx} + \frac{y}{2}u_{yy}$$

is the infinitesimal generator of equation (1.1), where a, b, d and e are constants. (1.1).

It follows from the Ito formula that for each $u \in \mathcal{C}^{1,2}(\mathbb{R}^+ \times \mathbb{R}^2)$, $u(t, X_t, Y_t)$ is a martingale if and only if $u_t + \mathcal{L}u = 0$. With a final condition at time T ,

equation (1.2) is known as the Kolmogorov backward equation, describing the backward time evolution of the probability density of the SDE (1.1), the density ρ verifies $\rho(s, x, y)dy = \mathbb{P}(X_s \in dy / X_T = x)$, $s \leq T$ and $\rho(T - t, x, y)$ is a solution of the Kolmogorov backward equation.

As we proceed in the computations, we must distinguish between cases for the constants a, b, d and e to identify the largest groups of symmetries that leave the equation invariant.

Before starting our work, we observed in various articles that certain software programs, such as Maple, Mathematica, and Python are capable of computing Lie symmetry groups for partial differential equations. However, a challenge arises in the Longstaff Schwartz model, that involves four constant parameters in the drift function. Existing software tools lack the ability to differentiate between these constants or impose necessary conditions (if needed) to identify symmetry groups. This limitation serves as our motivation to initiate computations, seeking to determine these conditions independently so that we can systematically ascertain symmetry groups in all cases and compare our findings with those obtained using Maple.

The remainder of this paper is organized as follows. In the next section, we recall the basic results of the Lie point symmetries for partial differential equations with an illustration of the heat equation. Section 3 is dedicated to identifying and analyzing the system of partial differential equations that determines the final structure of the symmetries of the Longstaff-Schwartz equation. Next, in Section 4, we solve the determining system and calculate the associated symmetries, and we give the class of solutions in every case. In the last section, we determine the structure of the Lie algebras linked to each Lie symmetry group and establish an isomorphism between these Lie algebras.

2 Prerequisites and preliminaries

2.1 Symmetry groups

In this section, we provide a concise overview of the theory of Lie symmetry groups for differential equations, closely following the presentation in Craddock and Platen [3].

The theory of Lie symmetry groups offers a systematic method to determine the Lie symmetry group connected to a given differential equation. This involves finding the infinitesimal generators of the group, which are vector fields that generate the symmetry transformations. The generators encode information regarding the symmetries of the equation and can be used to derive exact solutions or reduce the order of the equation. Lie symmetry groups have been applied in various fields of mathematics and physics. They are particularly important in the study of partial differential equations, where they can be used to classify and solve equations in areas such as fluid dynamics, quantum mechanics, general relativity, and finance. Lie symmetry groups are also connected to other branches of mathematics, including group theory, differential geometry, and representation theory.

In Lie symmetry theory, there are at least two methods to find symmetry groups. First, the method of Olver's prolongation, which we use in this study. Olver's book [14] provides a comprehensive treatment of Lie symmetry groups and their application to differential equations. It explores various mathematical techniques, such as Lie algebras, Lie derivatives, and group actions, to characterize and analyze symmetries. The book also discusses practical algorithms and

methods for finding the symmetries of specific differential equations, as well as the connection between symmetries and conservation laws.

Second, the method of isovectors introduced by B. Kent Harrison and Frank B. Estabrook in 1971, given a system of partial differential equations after a change in variables and/or possible unknowns, they can be expressed as the vanishing of a family of differential forms. An isovector is then a vector field of all variables preserving the differential ideal generated by the forms; see [6] for more details on this method.

The symmetry of a partial differential equation is a modification or transformation that maps the solutions of an equation to new solutions. More precisely, symmetries allow the construction of a complex solution from simple solutions. If \mathcal{C}_Δ denotes the set of all solutions of the PDE

$$\Delta(x, u^{(n)}) = 0 \quad (2.1)$$

Then we can see the symmetry \mathcal{S} as an automorphism of \mathcal{C}_Δ i.e $\mathcal{S} : \mathcal{C}_\Delta \rightarrow \mathcal{C}_\Delta$ means if $u \in \mathcal{C}_\Delta$ it implies that $\mathcal{S}u \in \mathcal{C}_\Delta$.

In the late XIX century, Lie developed a method to determine all symmetry groups for a system of differential equations, providing an essential and powerful tool for the analysis of differential equations. Olver's book [14] provides a comprehensive description of this theory, along with the works of Bluman and Kumei [1], [15] by Ovsiannikov, and the studies of Lescot and Zambrini [12, 9], who explored the symmetries of the Hamilton-Jacobi-Bellman equation and the Black-Scholes equation used in finance using the isovectors method [6].

The theory of symmetries is summarized by Lie's theorem, which will be discussed later. For the motivation of our work, we consider a PDE of order n involving p independent variables and q dependent variables, defined on a connected subset $M \subset X \times U$. The PDE takes the form (2.1) where $\Delta(x, y)$ is a C^∞ function on $M \subset X \times U$ where X and U contain independent and dependent variables, respectively.

$$u_J^{(n)} = \frac{\partial^k u}{\partial x_1^{j_1} \dots \partial x_k^{j_k}}$$

Here, $J = (j_1, \dots, j_k)$ is a multi-index with $j_i \in \mathbb{N}$ for $i \in \{1, 2, \dots, k\}$, the order of such a multi-index, which we denote by $\#J = k \leq n$, indicates how many derivatives are being taken. For more details, we refer to Olver's book [14] Chapter 2, which we describe below.

We begin by introducing the notion of vector fields in the form of

$$\mathbf{v} = \sum_{k=1}^p \xi_k(x, u) \frac{\partial}{\partial x_k} + \sum_{i=1}^q \phi_i(x, u) \frac{\partial}{\partial u_i}, \quad (2.2)$$

defined on $M \subset X \times U$.

We can view \mathbf{v} as a first-order differential operator: The vector field (2.2) is referred to as the infinitesimal generator of a one parameter local Lie group called the flow of \mathbf{v} , that transforms elements $(x, u) \in M \subset X \times U$. This group is denoted as \mathcal{G} . Our goal is to determine the conditions for ξ_k and ϕ_i which will guarantee that \mathcal{G} will be a symmetry group of (2.1).

Before stating Lie's theorem, we must introduce the notion of the **nth prolongation** of \mathbf{v} , denoted $pr^{(n)}(\mathbf{v})$ is defined on the space $M^{(n)} \subset X \times U^{(n)}$ where $X \times U^{(n)}$ is the space of independent variables, dependent variables and the derivatives of the dependent variables up to order n , we call it the **n-th order jet space** of the underlying space $X \times U$.

First, we define the n th prolongation of group \mathcal{G} , which is the natural extension of the action of \mathcal{G} from (x, u) to all derivatives of u up to order n , denoted by $pr^n \mathcal{G}$ [14, p. 100]. The infinitesimal generator of $pr^n \mathcal{G}$ is called the n th prolongation of \mathbf{v} , and is denoted $pr^n \mathbf{v}$. It is possible to obtain an explicit formula for $pr^n \mathbf{v}$.

Theorem 2.1. *Olver[14, p. 113]*

Let \mathbf{v} be a vector field of the form (2.2) on an open subset $M \subset X \times U$. Then, the n -th prolongation

of \mathbf{v} is the vector field:

$$\text{pr}^{(n)}\mathbf{v} = \mathbf{v} + \sum_{\alpha=1}^q \sum_J \phi_\alpha^J(x, u^{(n)}) \frac{\partial}{\partial u_J^\alpha}$$

defined on the corresponding jet space $M^{(n)} \subset X \times U^{(n)}$, the second summation is over all multi-indices $J = (j_1, \dots, j_k)$, with $1 \leq j_k \leq p, 1 \leq k \leq n$. The coefficient function ϕ_α^J of $\text{pr}^{(n)}\mathbf{v}$ is given by the following formula:

$$\phi_\alpha^J(x, u^{(n)}) = D_J \left(\phi_\alpha - \sum_{i=1}^p \xi^i u_{J,i}^\alpha \right) + \sum_{i=1}^p \xi^i u_{J,i}^\alpha,$$

where $u_i^\alpha = \partial u^\alpha / \partial x^i$, and $u_{J,i}^\alpha = \partial u_J^\alpha / \partial x^i = \frac{\partial^{k+1} u^\alpha}{\partial x^i \partial x^{j_1} \dots \partial x^{j_k}}$. Here D_J denotes the total derivative operator D .

Proposition 2.2. Given $P(x, u^{(n)})$ a smooth function of x, u and derivatives of u up to order n , defined on an open subset $M^{(n)} \subset X \times U^{(n)}$, the total derivative of P with respect to x^i has the form

$$D_i P = \frac{\partial P}{\partial x^i} + \sum_{\alpha=1}^p \sum_J u_{J,i}^\alpha \frac{\partial P}{\partial u_J^\alpha}$$

For example, suppose that $q = 1$ (one dependent variable u) and $p = 2$ (2 independent variables x, t). Let $J = (1, 1)$ be a multi-index; then, $u_J = u_{xt} = \frac{\partial^2 u}{\partial x \partial t}$, $u_{J,t} = u_{xtt}$. It is clear that ϕ^x can be written for ϕ^J when $J = (1, 0)$, $\phi^{xx} = \phi^J$ when $J = (2, 0)$ etc.

Now, we will present a version of Lie's theorem, which we call the infinitesimal criterion of symmetry, which is the key to the theory of Lie symmetry groups for differential equations. This allows the necessary and sufficient conditions for the vector field (2.2) to generate symmetries of the differential equation.

Theorem 2.3. *Lie*[14, p. 165] Let $\Delta(x, u^{(n)}) = 0$ be an n -th order partial differential equation as defined above. A local group of transformations \mathcal{G} acting on an open subset $M \subset X \times U$ is a symmetry group of the system if and only if $\text{pr}^n \mathbf{v}[\Delta(x, u^{(n)})] = 0$ whenever $\Delta(x, u^{(n)}) = 0$, for every infinitesimal generator of \mathcal{G} .

When Theorem (2.3) is applied to a partial differential equation, it leads us to what we refer to as the determining system for the functions ξ_k and ϕ . These determining systems are often solved using heuristic methods. Finally, we derive vector fields that generate symmetry groups. What is particularly interesting about these vector fields is that they form Lie algebra under the typical Lie bracket. We obtained the following results owing to Lie.

Corollary 2.4. Let $\Delta(x, u^{(n)}) = 0$ be an n -th order partial differential equation. The set of all infinitesimal symmetries of this system forms a Lie subalgebra of the algebra of C^∞ -vector fields on M . Moreover, if this subalgebra is finite-dimensional, the symmetry group of the system is a local Lie group of transformations acting on M .

The procedure to find the group transformation, which is generated by the infinitesimal symmetries that we already find, is known as **exponentiating** the vector field. To exponentiate the infinitesimal symmetry \mathbf{v}_k , we solve the system of first-order ordinary differential equations:

$$\frac{d\tilde{x}}{d\varepsilon} = \xi(\tilde{x}, \tilde{t}, \tilde{y}), \quad \frac{d\tilde{t}}{d\varepsilon} = \tau(\tilde{x}, \tilde{t}, \tilde{y}), \quad \frac{d\tilde{u}}{d\varepsilon} = \phi(\tilde{x}, \tilde{t}, \tilde{y}).$$

with initial conditions $\tilde{x}(0) = x, \tilde{t}(0) = t, \tilde{u}(0) = u$.

We define $u^{\varepsilon, v}(t, x) := \tilde{u}_\varepsilon(\tilde{t}_{-\varepsilon}, \tilde{x}_{-\varepsilon})$.

Theorem 2.5. For every infinitesimal generator that satisfies Lie's condition (2.3), we have that $u^{\varepsilon, v}$ is a solution of PDE (2.1) for every ε sufficiently small.

For more details on the transformed solution of the PDE, see [14, p. 94].

Remark 2.6.

- $u^{\varepsilon, v}$ is the action of the symmetry generated by v on u .
- ε is the parameter of the symmetry group.

Example 2.7 (Heat equation). We illustrate here the example of the well known one dimensional heat equation,

$$u_{xx} = u_t. \quad (2.3)$$

Here $p=2$ and $q=1$, so $X = \mathbb{R}^2$ and $U = \mathbb{R}$.

To find the symmetries of (2.3), we set

$$v = \xi(x, t, u) \frac{\partial}{\partial x} + \tau(x, t, u) \frac{\partial}{\partial t} + \phi(x, t, u) \frac{\partial}{\partial u}$$

and compute the second prolongation of \mathbf{v} . Thanks to Lie's theorem \mathbf{v} generates a symmetry group of the heat equation if and only if

$$pr^2 \mathbf{v}(u_{xx} - u_t) = 0$$

whenever $u_{xx} - u_t = 0$,

where

$$pr^2 \mathbf{v} = \mathbf{v} + \phi^x \frac{\partial}{\partial u_x} + \phi^t \frac{\partial}{\partial u_t} + \phi^{xx} \frac{\partial}{\partial u_{xx}} + \phi^{xt} \frac{\partial}{\partial u_{xt}} + \phi^{tt} \frac{\partial}{\partial u_{tt}}.$$

This condition provides the determining system for ξ , τ and ϕ . For more details see Olver [14, p. 120].

Solving the determining system gives the expression of ξ , τ and ϕ , then we determine the Lie algebra spanned by the infinitesimal symmetries. Thus, the basis of the Lie algebra of symmetry is

$$\begin{aligned} v_1 &= \frac{\partial}{\partial x} \\ v_2 &= \frac{\partial}{\partial t} \\ v_3 &= u \frac{\partial}{\partial u}, \\ v_4 &= x \frac{\partial}{\partial x} + 2t \frac{\partial}{\partial t} - \frac{u}{2} \frac{\partial}{\partial u} \\ v_5 &= 2t \frac{\partial}{\partial x} - xu \frac{\partial}{\partial u}, \\ v_6 &= 4xt \frac{\partial}{\partial x} + 4t^2 \frac{\partial}{\partial t} - (x^2 + 2t)u \frac{\partial}{\partial u}. \end{aligned}$$

Additionally, the infinite dimensional Lie subalgebra spanned by $v_\beta = \beta(x, t) \frac{\partial}{\partial u}$ where β is an arbitrary solution of the heat equation, which corresponds to the linearity of the equation.

Exponentiating v_k and using theorem 2.5 gives the symmetry transformations below:

$$u^{\varepsilon, v_1}(t, x) = u(x - \varepsilon, t), \quad (2.4)$$

$$u^{\varepsilon, v_2}(x, t) = u(x, t - \varepsilon), \quad (2.5)$$

$$u^{\varepsilon, v_3}(x, t) = e^\varepsilon u(x, t), \quad (2.6)$$

$$u^{\varepsilon, v_4}(x, t) = e^{-\frac{\varepsilon}{2}} u(e^{-\varepsilon} x, e^{-2\varepsilon} t), \quad (2.7)$$

$$u^{\varepsilon, v_5}(x, t) = e^{-\varepsilon x + \varepsilon^2 t} u((x - 2\varepsilon t), t), \quad (2.8)$$

$$u^{\varepsilon, v_6}(x, t) = \frac{1}{\sqrt{1 + 4\varepsilon t}} \exp\left(\frac{-\varepsilon x^2}{1 + 4\varepsilon t}\right) u\left(\frac{x}{1 + 4\varepsilon t}, \frac{t}{1 + 4\varepsilon t}\right). \quad (2.9)$$

This implies that if $u(x, t)$ is a solution of the heat equation, then the right-hand side of equations ((2.4), ..., (2.9)) is also a solution for ε to be sufficiently small. For an application, let us examine the infinitesimal symmetry (2.9). When $u(x, t) = 1$ is a solution of the heat equation, then by symmetry:

$$u^{\varepsilon, v_6}(x, t) = \frac{1}{\sqrt{1 + 4\varepsilon t}} \exp\left(\frac{-\varepsilon x^2}{1 + 4\varepsilon t}\right) \text{ is also a solution.}$$

Allow $t \rightarrow t - \frac{1}{4\varepsilon}$ and $\varepsilon = \pi$ then $g(x, t) = \frac{1}{\sqrt{4\pi t}} \exp\left(\frac{-x^2}{4t}\right)$ which is the fundamental solution for the heat equation that we found from $u = 1$ by a group transformation.

Lescot and Zambrini [12] found the same symmetries of the heat equation using the isovectors method [6].

2.2 A Lie algebra

We define $\mathcal{M} = \langle E, F, H, X, Y, Z \rangle$ with $[E, F] = H$, $[H, F] = -2F$, $[H, E] = 2E$, $[X, Z] = 0$, $[Y, Z] = 0$ and $[X, Y] = Z$. \mathcal{M} is a Lie algebra. From the commutator relations, we have that, $\langle E, F, H \rangle \cong \mathfrak{sl}_2$ and $\langle X, Y, Z \rangle \cong \mathfrak{h}_3$, the Heisenberg algebra of dimension 3

Table 1: Lie brackets of the Algebra of heat equation.

$[\cdot, \cdot]$	E	F	H	X	Y	Z
E	0	H	$-2E$	0	X	0
F	$-H$	0	$2F$	Y	0	0
H	$2E$	$-2F$	0	X	$-Y$	0
X	0	$-Y$	$-X$	0	Z	0
Y	$-X$	0	Y	$-Z$	0	0
Z	0	0	0	0	0	0

From the commutator table, we notice that $\langle X, Y, Z \rangle$ is an ideal of \mathcal{M} , and $\langle E, F, H \rangle$ is a subalgebra of \mathcal{M} , therefore, \mathcal{M} is a semidirect product $\mathfrak{sl}_2 \ltimes \mathfrak{h}_3$. More precisely, from [11] and Theorem 4.1, in [10] $\mathcal{M} \cong \mathcal{H}_{0,0}$, where $\mathcal{H}_{0,0}$ is the Lie symmetry algebra (of dimension 6) of the heat equation

$$\frac{\partial u}{\partial t} = -\frac{\partial^2 u}{\partial q^2}.$$

Also, from the commutator table, one can see that $\mathcal{N} := \{E, F, H, Z\}$ is a subalgebra of \mathcal{M} , and $\mathcal{N} \cong \mathfrak{sl}_2 \times \mathbb{R}$. Using Theorem (9.1) in [10], for each $C \neq 0$ and D , \mathcal{N} is isomorphic to the symmetry algebra of the equation

$$\frac{\partial u}{\partial t} = -\frac{\partial^2 u}{\partial q^2} + \left(\frac{C}{q^2} + Dq^2\right) u.$$

3 Equations Defining Infinitesimal Symmetries.

In this section, we aim to determine the Lie algebra of symmetries called $\mathcal{V}_{a,b,d,e}$ for the PDE:

$$\Delta(x, y, t, u^{(2)}) = (a - bx)u_x + (d - ey)u_y + \frac{x}{2}u_{xx} + \frac{y}{2}u_{yy} + u_t = 0. \quad (3.1)$$

Here we have three independent variables x, y and t , and one dependent variable u , resulting in $p = 3$ and $q = 1$ in our notation. Equation (3.1) is a second-order PDE with $n = 2$. We can identify the set of its solutions with the linear subvariety in $X \times U^{(2)}$ determined by the vanishing of $\Delta(x, y, t, u^{(2)})$.

We look for vector fields on $X \times U$ of the form

$$v = \xi(x, y, t, u) \frac{\partial}{\partial x} + \gamma(x, y, t, u) \frac{\partial}{\partial y} + \tau(x, y, t, u) \frac{\partial}{\partial t} + \phi(x, y, t, u) \frac{\partial}{\partial u}.$$

Our goal is to determine all coefficient functions ξ, γ, τ and ϕ such that the corresponding one-parameter group $\exp(\varepsilon v)$ is a symmetry group of (3.1). Otherwise, we search for symmetry groups that leave the subvariety $\mathcal{S}_\Delta = \{(x, y, t, u^{(2)}) : \Delta(x, y, t, u^{(2)}) = 0\}$ invariant.

According to Theorem (2.4), we need to calculate the second prolongation

$$pr^2 \mathbf{v} = \mathbf{v} + \phi^x \frac{\partial}{\partial u_x} + \phi^y \frac{\partial}{\partial u_y} + \phi^t \frac{\partial}{\partial u_t} + \phi^{xx} \frac{\partial}{\partial u_{xx}} + \phi^{yy} \frac{\partial}{\partial u_{yy}} + \phi^{tt} \frac{\partial}{\partial u_{tt}} + \phi^{xy} \frac{\partial}{\partial u_{xy}} + \phi^{xt} \frac{\partial}{\partial u_{xt}} + \phi^{yt} \frac{\partial}{\partial u_{yt}}$$

of \mathbf{v} whose coefficients are

$$\phi^x = \phi_x + (\phi_u - \xi_x)u_x - \xi_u u_x^2 - \gamma_x u_y - \tau_x u_t - \gamma_u u_x u_y - \tau_u u_x u_t,$$

$$\phi^y = \phi_y + (\phi_u - \gamma_y)u_y - \gamma_u u_y^2 - \xi_y u_x - \xi_u u_x u_y - \tau_y u_t - \tau_u u_t u_y,$$

$$\phi^t = \phi_t + (\phi_u - \tau_t)u_t - \xi_t u_x - \xi_u u_t u_x - \gamma_t u_y - \gamma_u u_t u_y - \tau_u u_t^2,$$

$$\begin{aligned} \phi^{xx} = & \phi_{xx} + (2\phi_{xu} - \xi_{xx})u_x + (\phi_u - 2\xi_x)u_{xx} + (\phi_{uu} - 2\xi_{xu})u_x^2 - 3\xi_u u_x u_{xx} - 2\gamma_{xu} u_x u_y - 2\gamma_x u_{xy} - 2\gamma_u u_x u_{xy} \\ & - \xi_{uu} u_x^3 - \gamma_{xx} u_y - \gamma_u u_{xx} u_y - \gamma_{uu} u_x^2 u_y - 2\tau_x u_{xt} - 2\tau_u u_{xt} u_x - \tau_{xx} u_t - 2\tau_{xu} u_x u_t - \tau_u u_{xx} u_t - \tau_{uu} u_x^2 u_t, \end{aligned}$$

$$\begin{aligned} \phi^{yy} = & \phi_{yy} + (2\phi_{yu} - \gamma_{yy})u_y + (\phi_u - 2\gamma_y)u_{yy} + (\phi_{uu} - 2\gamma_{yu})u_y^2 - 3\gamma_u u_y u_{yy} - 2\xi_{yu} u_{xy} - 2\xi_u u_{xy} u_y - \xi_{yy} u_x \\ & - 2\xi_{yu} u_x u_y - \xi_x u_{yy} - \xi_{uu} u_x u_y^2 - \gamma_{uu} u_y^3 - 2\tau_y u_{yt} - 2\tau_u u_y u_{yt} - \tau_{yy} u_t - 2\tau_{uy} u_t u_y - \tau_u u_t u_y - \tau_{uu} u_t u_y^2. \end{aligned}$$

Remark 3.1. In this context, we refrained from calculating $\phi^{tt}, \phi^{yt}, \phi^{xt}$ and ϕ^{xy} because the terms u_{tt}, u_{xy}, u_{yt} and u_{xt} are not present in Equation (3.1). Consequently, in the prolongation formula, the derivatives of Δ with respect to u_{tt}, u_{xy}, u_{yt} and u_{xt} vanish.

Applying $pr^{(2)}\mathbf{v}$ to $\Delta(x, y, t, u^{(2)})$, $pr^{(2)}\mathbf{v}(\Delta) = 0$ knowing that $\Delta = 0$, with

$$\begin{aligned}
pr^{(2)}\mathbf{v}(\Delta) &= -b\xi u_x + \frac{\xi}{2}u_{xx} - e\gamma u_y + \frac{\gamma}{2}u_{yy} \\
&+ (a - bx) [\phi_x + (\phi_u - \xi_x)u_x - \xi_u u_x^2 - \gamma_x u_y - \tau_x u_t - \gamma_u u_x u_y - \tau_u u_x u_t] \\
&+ (d - ey) [\phi_y + (\phi_u - \gamma_y)u_y - \gamma_u u_y^2 - \xi_y u_x - \xi_u u_x u_y - \tau_y u_t - \tau_u u_t u_y] \\
&+ \frac{x}{2} \left[\phi_{xx} + (2\phi_{xu} - \xi_{xx})u_x + (\phi_u - 2\xi_x)u_{xx} + (\phi_{uu} - 2\xi_{xu})u_x^2 - 3\xi_u u_x u_{xx} - 2\gamma_{xu} u_x u_y - 2\gamma_x u_{xy} - 2\gamma_u u_x u_{xy} \right. \\
&\quad \left. - \xi_{uu} u_x^3 - \gamma_{xx} u_y - \gamma_u u_{xx} u_y - \gamma_{uu} u_x^2 u_y - 2\tau_x u_{xt} - 2\tau_u u_{xt} u_x - \tau_{xx} u_t - 2\tau_{xu} u_x u_t - \tau_u u_{xx} u_t - \tau_{uu} u_x^2 u_t \right] \\
&+ \frac{y}{2} \left[\phi_{yy} + (2\phi_{yu} - \gamma_{yy})u_y + (\phi_u - 2\gamma_y)u_{yy} + (\phi_{uu} - 2\gamma_{yu})u_y^2 - 3\gamma_u u_y u_{yy} - 2\xi_y u_{xy} - 2\xi_u u_{xy} u_y - \xi_{yy} u_x \right. \\
&\quad \left. - 2\xi_{yu} u_x u_y - \xi_x u_{yy} - \xi_{uu} u_x u_y^2 - \gamma_{uu} u_y^3 - 2\tau_y u_{yt} - 2\tau_u u_y u_{yt} - \tau_{yy} u_t - 2\tau_{uy} u_t u_y - \tau_u u_t u_y - \tau_{uu} u_t u_y^2 \right] = 0
\end{aligned}$$

and replace u_t with $-(a - bx)u_x - (d - ey)u_y - \frac{x}{2}u_{xx} - \frac{y}{2}u_{yy}$ wherever it occurs. The infinitesimal criterion of symmetry in (2.3) results in a polynomial equation for the derivatives of u . The coefficients of this polynomial equation are constructed from ξ , γ , τ and ϕ along with their derivatives. However, because ξ , γ , τ and ϕ depend only on x, y, t and u , the coefficients must all be zero. Hence, the determining system is:

Monomials	Coefficients
u_x	$-b\xi - (a - bx)\xi_x + (a - bx)^2\tau_x - (d - ey)\xi_y + (a - bx)(d - ey)\tau_y - \xi_t$ $+(a - bx)\tau_t + \frac{x}{2}(2\phi_{xu} - \xi_{xx}) + \frac{x}{2}(a - bx)\tau_{xx} - \frac{y}{2}\xi_{yy} + \frac{y}{2}(a - bx)\tau_{yy} = 0$ (3.2)
u_y	$-e\gamma - (d - ey)\gamma_y + (d - ey)^2\tau_y - (a - bx)\gamma_x + (a - bx)(d - ey)\tau_x - \gamma_t$ $+(d - ey)\tau_t + \frac{y}{2}(2\phi_{yu} - \gamma_{yy}) + \frac{y}{2}(d - ey)\tau_{yy} - \frac{x}{2}\gamma_{xx} + \frac{x}{2}(d - ey)\tau_{xx} = 0$ (3.3)
$u_x u_y$	$-x\gamma_{xu} + x(d - ey)\tau_{xu} - y\xi_{yu} - y(a - bx)\tau_{uy} = 0$ (3.4)
u_x^2	$-(a - bx)\xi_u + (a - bx)^2\tau_u + \frac{x}{2}(\phi_{uu} - 2\xi_{xu}) + x(a - bx)\tau_{xu} = 0$ (3.5)
u_y^2	$-(d - ey)\gamma_u + (d - ey)^2\tau_u + \frac{y}{2}(\phi_{uu} - 2\gamma_{yu}) + y(d - ey)\tau_{uy} = 0$ (3.6)
u_{xx}	$\frac{\xi}{2} + \frac{x}{2}(a - bx)\tau_x + \frac{y}{2}(d - ey)\tau_y + \frac{x}{2}(\phi_u - 2\xi_x) + \frac{x^2}{4}\tau_{xx} + \frac{y^2}{4}\tau_{yy} = 0$ (3.7)
u_{yy}	$\frac{\gamma}{2} + \frac{y}{2}(d - ey)\tau_y + \frac{y}{2}(a - bx)\tau_x + \frac{y}{2}(\phi_u - 2\gamma_y) + \frac{y^2}{4}\tau_{yy} + \frac{x^2}{4}\tau_{xx} = 0$ (3.8)
$u_x u_{xx}$	$-x\xi_u + \frac{x^2}{2}\tau_{xu} = 0$ (3.9)
$u_y u_{yy}$	$-y\gamma_u + \frac{y^2}{2}\tau_{yu} = 0$ (3.10)
$u_{yy} u_x$	$\frac{xy}{2}\tau_{xu} = 0$ (3.11)
$u_{xx} u_y$	$\frac{xy}{2}\tau_{uy} = 0$ (3.12)
$u_{xx} u_{yy}$	$-xy\tau_u = 0$ (3.13)
u_{yy}^2	$\frac{y^2}{4}\tau_u = 0$ (3.14)
u_{xx}^2	$\frac{x^2}{4}\tau_u = 0$ (3.15)
u_{xy}	$x\gamma_x + y\xi_y = 0$ (3.16)
$u_x u_{xy}$	$-x\gamma_u = 0$ (3.17)
$u_x^2 u_y$	$-\frac{x}{2}\gamma_{uu} + \frac{x}{2}(d - ey)\tau_{uu} = 0$ (3.18)
u_x^3	$-\frac{x}{2}\xi_{uu} - \frac{x}{2}(a - bx)\tau_{uu} = 0$ (3.19)
u_y^3	$-\frac{y}{2}\gamma_{uu} - \frac{y}{2}(d - ey)\tau_{uu} = 0$ (3.20)
u_{xt}	$-x\tau_x = 0$ (3.21)
$u_{xt} u_x$	$-x\tau_u = 0$ (3.22)

Monomes	Coefficients	
$u_x u_y^2$	$-\frac{y}{2}\xi_{uu} + \frac{y}{2}(a - bx)\tau_{uu} = 0$	(3.23)
$u_x^2 u_t$	$\frac{x}{2}\tau_{uu} = 0$	(3.24)
$u_y^2 u_t$	$\frac{y}{2}\tau_{uu} = 0$	(3.25)
$u_{xy} u_y$	$-y\xi_u = 0$	(3.26)
u_{yt}	$y\tau_y = 0$	(3.27)
$u_y u_{yt}$	$-y\tau_u = 0$	(3.28)
.....	$\frac{y}{2}(d - ey)\tau_{uu} = 0$	(3.29)
$u_y^2 u_{yy}$	$\frac{y^2}{4}\tau_{uu} = 0$	(3.30)
$u_x^2 u_{xx}$	$\frac{x^2}{4}\tau_{uu} = 0$	(3.31)
constant term	$(a - bx)\phi_x + (d - ey)\phi_y + \frac{x}{2}\phi_{xx} + \frac{y}{2}\phi_{yy} + \phi_t = 0$	(3.32)

First, Equations (3.21),(3.22) and (3.27) require that τ be solely a function of t . Subsequently, Equations(3.17) and (3.26) demonstrate that ξ and γ are independent of u .

Furthermore, according to (3.5) and (3.6), ϕ is affine in u , implying that

$$\phi(x, y, t, u) = \lambda(x, y, t)u + \mu(x, y, t)$$

for certain functions λ and μ . Substituting ϕ into equation (3.32) reveals that λ and μ satisfy equation (3.1).

Since ϕ is affine on u , we define $\mathcal{L}_{a,b,d,e} = \{v \in \mathcal{V}_{a,b,d,e} \mid \mu_v = 0\}$, a subspace of $\mathcal{V}_{a,b,d,e}$ where $\mu_v := \mu$.

Lemma 3.2. $\mathcal{L}_{a,b,d,e}$ is a subalgebra of $\mathcal{V}_{a,b,d,e}$.

Proof. Consider $v, v' \in \mathcal{L}_{a,b,d,e}$ implying $\mu_v = \mu_{v'} = 0$. We aim to show that $[v, v'] \in \mathcal{L}_{a,b,d,e}$. By the definition of $\mathcal{L}_{a,b,d,e}$, it suffices to prove that $\mu_{[v,v']} = 0$. Expressing ϕ as $\phi_v = v(u) = \lambda_v u + \mu_v$ we have $\phi_{[v,v']} = \lambda_{[v,v']}u + \mu_{[v,v']}$.

We compute as follows:

$$\begin{aligned} \phi_{[v,v']} &= [v, v'](u) = v(v'(u)) - v'(v(u)) \\ &= v(\lambda_{v'}u + \mu_{v'}) - v'(\lambda_v u + \mu_v) \\ &= (v(\lambda_{v'}) - v'(\lambda_v))u + (v(\mu_{v'}) - v'(\mu_v) + \lambda_{v'}\mu_v - \lambda_v\mu_{v'}) \end{aligned}$$

Since ξ, γ, τ do not depend on x, y, t , the terms $v(\lambda_{v'}) - v'(\lambda_v)$ and $v(\mu_{v'}) - v'(\mu_v) + \lambda_{v'}\mu_v - \lambda_v\mu_{v'}$ depend only on x, y, t . Thus, $\mu_{[v,v']} = v(\mu_{v'}) - v'(\mu_v) + \lambda_{v'}\mu_v - \lambda_v\mu_{v'}$ and $\lambda_{[v,v']} = v(\lambda_{v'}) - v'(\lambda_v)$. Hence, $\mu_{[v,v']} = 0$, providing that $[v, v'] \in \mathcal{L}_{a,b,d,e}$. \square

Remark 3.3. $\mathcal{L}_{a,b,d,e}$ represent the space of non trivial symmetries, that is, symmetries that do not arise from the linearity of the equation.

Now we seek symmetries of the form:

$$v = \xi(x, y, t)\frac{\partial}{\partial x} + \gamma(x, y, t)\frac{\partial}{\partial y} + \tau(t)\frac{\partial}{\partial t} + (\lambda(x, y, t)u + \mu(x, y, t))\frac{\partial}{\partial u}.$$

Because τ does not depend on x, y or u , conditions (3.7) and (3.8) become:

$$\begin{cases} \frac{\xi}{2} + \frac{x}{2}\tau_t - x\xi_x = 0, \\ \frac{\gamma}{2} + \frac{y}{2}\tau_t - y\gamma_y = 0; \end{cases}$$

Solving those two equations gives:

$$\begin{cases} \xi = \tau_t x + \sqrt{x}f(t, y), \\ \gamma = \tau_t y + \sqrt{y}g(x, t), \end{cases}$$

for certain functions f and g .

Substituting ξ and γ into equation (3.16) gives

$$2(x\gamma_x + y\xi_y) = 2x(\sqrt{y}g_x) + 2y(\sqrt{x}f_y) = 2\sqrt{x}g_x + 2\sqrt{y}f_y = 0,$$

thus there exists a function h such that $-2\sqrt{x}g_x = 2\sqrt{y}f_y = h(t)$.

$$\begin{aligned} \text{Then } & \begin{cases} g_x = -\frac{1}{2\sqrt{x}}h(t), \\ f_y = \frac{1}{2\sqrt{y}}h(t), \end{cases} \\ \text{so } & \begin{cases} g(x, t) = -\sqrt{x}h(t) + k(t), \\ f(y, t) = \sqrt{y}h(t) + l(t), \end{cases} \end{aligned}$$

for certain functions k and l . We remind that the unknown functions are $h(t), l(t), k(t), \tau(t)$.

$$\text{We have } \begin{cases} \xi = \tau_t x + \sqrt{x}(\sqrt{y}h(t) + l(t)), \\ \gamma = \tau_t y + \sqrt{y}(-\sqrt{x}h(t) + k(t)). \end{cases}$$

Replacing ξ and γ in the equations (3.2) and (3.3) gives

$$\begin{cases} -b(\tau_t x + \sqrt{x}(\sqrt{y}h + l)) - (a - bx) \left(\frac{h}{2} \frac{\sqrt{y}}{\sqrt{x}} + \frac{l}{2} \frac{1}{\sqrt{x}} \right) - (d - ey) \left(\frac{h}{2} \frac{\sqrt{x}}{\sqrt{y}} \right) \\ - \tau_{tt}x - h_t \sqrt{x} \sqrt{y} - \sqrt{x}l_t + x\lambda_x + \frac{x}{2} \left(\frac{h}{4} \sqrt{y}x^{-\frac{3}{2}} + \frac{l}{4}x^{-\frac{3}{2}} \right) + \frac{y}{2} \left(\frac{h}{4} \sqrt{x}y^{-\frac{3}{2}} \right) = 0, \\ -e(\tau_t y + \sqrt{y}(-\sqrt{x}h + k)) - (d - ey) \left(-\frac{h}{2} \frac{\sqrt{x}}{\sqrt{y}} + \frac{k}{2} \frac{1}{\sqrt{y}} \right) - (a - bx) \left(-\frac{h}{2} \frac{\sqrt{y}}{\sqrt{x}} \right) \\ - \tau_{tt}y + \sqrt{x}\sqrt{y}h_t - k_t \sqrt{y} + y\lambda_y - \frac{y}{2} \left(\frac{h}{4} \sqrt{x}y^{-\frac{3}{2}} - \frac{k}{4}y^{-\frac{3}{2}} \right) - \frac{x}{2} \left(\frac{h}{4} \sqrt{y}x^{-\frac{3}{2}} \right) = 0, \end{cases}$$

simplifying we have

$$\left\{ \begin{aligned} \lambda(x, y, t) &= (\tau_{tt} + b\tau_t)x + 2 \left(h_t + \frac{h}{2}(b - e) \right) \sqrt{x}\sqrt{y} + h \left(\frac{1}{4} - a \right) \frac{\sqrt{y}}{\sqrt{x}} + h \left(d - \frac{1}{4} \right) \frac{\sqrt{x}}{\sqrt{y}} \\ &+ 2 \left(l_t + \frac{bl}{2} \right) \sqrt{x} + l \left(\frac{1}{4} - a \right) \frac{1}{\sqrt{x}} + \theta(y, t), \end{aligned} \right. \quad (3.33)$$

$$\left\{ \begin{aligned} \lambda(x, y, t) &= (\tau_{tt} + e\tau_t)y + 2 \left(-h_t + \frac{h}{2}(b - e) \right) \sqrt{x}\sqrt{y} + h \left(\frac{1}{4} - a \right) \frac{\sqrt{y}}{\sqrt{x}} + h \left(d - \frac{1}{4} \right) \frac{\sqrt{x}}{\sqrt{y}} \\ &+ 2 \left(k_t + \frac{ek}{2} \right) \sqrt{y} + k \left(\frac{1}{4} - d \right) \frac{1}{\sqrt{y}} + \beta(x, t), \end{aligned} \right. \quad (3.34)$$

for some functions θ and β .

For a fixed t and y , utilizing (3.33), we observe that $\sqrt{x}\lambda(x, y, t) \xrightarrow{x \rightarrow 0} 0$ and

$\sqrt{x}\lambda(x, y, t) \xrightarrow{x \rightarrow 0} h(\frac{1}{4} - a)\sqrt{y} + l(\frac{1}{4} - a)$. From the uniqueness of the limit, we obtain

$h(\frac{1}{4} - a)\sqrt{y} + l(\frac{1}{4} - a) = 0$. Similarly, for a fixed t and x , we find that $h(\frac{1}{4} - d)\sqrt{x} + k(\frac{1}{4} - d) = 0$.

Thus, for every x and y :

$$\begin{aligned} h(\frac{1}{4} - a)\sqrt{y} + l(\frac{1}{4} - a) &= 0 \\ h(\frac{1}{4} - d)\sqrt{x} + k(\frac{1}{4} - d) &= 0. \end{aligned}$$

Since h, k and l depend only on t , we have

$$h\left(\frac{1}{4} - a\right) = h\left(\frac{1}{4} - d\right) = l\left(\frac{1}{4} - a\right) = k\left(\frac{1}{4} - d\right) = 0 \text{ for every } t. \quad (3.35)$$

Remark 3.4. • If $a \neq \frac{1}{4}$:

From 3.35 we deduce that $h = l = 0$.

• If $d \neq \frac{1}{4}$:

From 3.35 we deduce that $h = k = 0$.

Thus equations (3.33) and (3.34) becomes

$$\begin{cases} \lambda(x, y, t) = (\tau_{tt} + b\tau_t)x + 2\left(h_t + \frac{h}{2}(b - e)\right)\sqrt{x}\sqrt{y} + 2\left(l_t + \frac{bl}{2}\right)\sqrt{x} + \theta(y, t), \\ \lambda(x, y, t) = (\tau_{tt} + e\tau_t)y + 2\left(-h_t + \frac{h}{2}(b - e)\right)\sqrt{x}\sqrt{y} + 2\left(k_t + \frac{ek}{2}\right)\sqrt{y} + \beta(x, t), \end{cases} \quad (3.36)$$

$$\begin{cases} \lambda(x, y, t) = (\tau_{tt} + e\tau_t)y + 2\left(-h_t + \frac{h}{2}(b - e)\right)\sqrt{x}\sqrt{y} + 2\left(k_t + \frac{ek}{2}\right)\sqrt{y} + \beta(x, t), \\ \lambda(x, y, t) = (\tau_{tt} + b\tau_t)x + 2\left(h_t + \frac{h}{2}(b - e)\right)\sqrt{x}\sqrt{y} + 2\left(l_t + \frac{bl}{2}\right)\sqrt{x} + \theta(y, t), \end{cases} \quad (3.37)$$

From (3.36) and (3.37) we have

$$\begin{aligned} & \lim_{x \rightarrow 0} \left[(\tau_{tt} + b\tau_t)x + 2\left(h_t + \frac{h}{2}(b - e)\right)\sqrt{x}\sqrt{y} + 2\left(l_t + \frac{bl}{2}\right)\sqrt{x} + \theta(y, t) \right] \\ &= \lim_{x \rightarrow 0} \left[(\tau_{tt} + e\tau_t)y + 2\left(-h_t + \frac{h}{2}(b - e)\right)\sqrt{x}\sqrt{y} + 2\left(k_t + \frac{ek}{2}\right)\sqrt{y} + \beta(x, t) \right] \end{aligned}$$

and

$$\begin{aligned} & \lim_{y \rightarrow 0} \left[(\tau_{tt} + b\tau_t)x + 2\left(h_t + \frac{h}{2}(b - e)\right)\sqrt{x}\sqrt{y} + 2\left(l_t + \frac{bl}{2}\right)\sqrt{x} + \theta(y, t) \right] \\ &= \lim_{y \rightarrow 0} \left[(\tau_{tt} + e\tau_t)y + 2\left(-h_t + \frac{h}{2}(b - e)\right)\sqrt{x}\sqrt{y} + 2\left(k_t + \frac{ek}{2}\right)\sqrt{y} + \beta(x, t) \right]. \end{aligned}$$

Therefore

$$\begin{cases} \theta(y, t) = (\tau_{tt} + e\tau_t)y + 2\left(k_t + \frac{ek}{2}\right)\sqrt{y} + s(t), \\ \beta(x, t) = (\tau_{tt} + b\tau_t)x + 2\left(l_t + \frac{bl}{2}\right)\sqrt{x} + s(t). \end{cases} \quad (3.38)$$

where $s(t) := \beta(0, t) = \theta(0, t)$.

Remark 3.5. Using (3.36), (3.37), and (3.38), we obtain:

$$\begin{aligned} \lambda(x, y, t) &= (\tau_{tt} + b\tau_t)x + 2\left(h_t + \frac{h}{2}(b - e)\right)\sqrt{x}\sqrt{y} + 2\left(l_t + \frac{bl}{2}\right)\sqrt{x} + (\tau_{tt} + e\tau_t)y + 2\left(k_t + \frac{ek}{2}\right)\sqrt{y} + s(t) \\ &= (\tau_{tt} + e\tau_t)y + 2\left(-h_t + \frac{h}{2}(b - e)\right)\sqrt{x}\sqrt{y} + 2\left(k_t + \frac{ek}{2}\right)\sqrt{y} + (\tau_{tt} + b\tau_t)x + 2\left(l_t + \frac{bl}{2}\right)\sqrt{x} + s(t). \end{aligned}$$

So, $4h_t\sqrt{x}\sqrt{y} = 0$ for every x and y . Therefore, $h_t = 0$.

Because λ satisfies (3.1), and from equation (3.35) and remark 3.5, then

$$\begin{aligned} & (a - b)x \left[\tau_{tt} + b\tau_t + \frac{h}{2}(b - e)\frac{\sqrt{y}}{\sqrt{x}} + \left(l_t + \frac{bl}{2}\right)\frac{1}{\sqrt{x}} \right] + (d - ey) \left[\tau_{tt} + e\tau_t + \frac{h}{2}(b - e)\frac{\sqrt{x}}{\sqrt{y}} + \left(k_t + \frac{ek}{2}\right)\frac{1}{\sqrt{y}} \right] \\ &+ \frac{x}{2} \left[-\frac{h}{4}(b - e)\sqrt{y}x^{-\frac{3}{2}} - \frac{1}{2}\left(l_t + \frac{bl}{2}\right)x^{-\frac{3}{2}} \right] + \frac{y}{2} \left[-\frac{h}{4}(b - e)\sqrt{x}y^{-\frac{3}{2}} - \frac{1}{2}\left(k_t + \frac{ek}{2}\right)y^{-\frac{3}{2}} \right] \\ &+ \left[(\tau_{ttt} + b\tau_{tt})x + 2\left(h_{tt} + \frac{h_t}{2}(b - e)\right)\sqrt{x}\sqrt{y} + 2\left(l_{tt} + \frac{bl_t}{2}\right)\sqrt{x} + (\tau_{ttt} + e\tau_{tt})y + 2\left(k_{tt} + \frac{ek_t}{2}\right)\sqrt{y} + s_t \right] = 0. \end{aligned}$$

Therefore

$$\begin{aligned}
& (\tau_{ttt} - b^2\tau_t)x + (\tau_{ttt} - e^2\tau_t)y - \frac{h}{2}(b^2 - e^2)\sqrt{x}\sqrt{y} + \left(2l_{tt} - b^2\frac{l}{2}\right)\sqrt{x} + \left(2k_{tt} - e^2\frac{k}{2}\right)\sqrt{y} \\
& + ((a+d)\tau_{tt} + (ab+de)\tau_t + s_t) = 0 \text{ for every } x \text{ and } y.
\end{aligned}$$

Then

$$\begin{cases} h_t = 0, & (3.39a) \end{cases}$$

$$\begin{cases} (b^2 - e^2)\tau_t = 0, & (3.39b) \end{cases}$$

$$\begin{cases} (b^2 - e^2)h = 0, & (3.39c) \end{cases}$$

$$\begin{cases} l_{tt} - \frac{b^2}{4}l = 0, & (3.39d) \end{cases}$$

$$\begin{cases} k_{tt} - \frac{e^2}{4}k = 0, & (3.39e) \end{cases}$$

$$\begin{cases} (a+d)\tau_{tt} + (ab+de)\tau_t + s_t = 0. & (3.39f) \end{cases}$$

and μ a solution of equation (3.1).

Recall that the unknowns are τ , h , l , k , s , and μ . Thus, the coefficients of the vector field are given by:

$$\xi(t, x, y) = \tau_t x + \sqrt{x}(\sqrt{y}h(t) + l(t)),$$

$$\gamma(t, x, y) = \tau_t y + \sqrt{y}(-\sqrt{x}h(t) + k(t)),$$

$$\phi(t, x, y, u) = \left[(\tau_{tt} + b\tau_t)x + h(b-e)\sqrt{x}\sqrt{y} + 2\left(l_t + \frac{bl}{2}\right)\sqrt{x} + (\tau_{tt} + e\tau_t)y + 2\left(k_t + \frac{ek}{2}\right)\sqrt{y} + s(t) \right] u + \mu(t, x, y).$$

This provides the final structure of the symmetry group. To proceed further, it is essential to distinguish between cases based on conditions involving the constants a , b , d and e .

4 Computing the Infinitesimal Symmetries.

Now, we determine the Lie algebras of symmetries and compute their Lie brackets. From equations 3.35 and 3.39c, we split our analysis into (4×4) cases:

$$(b = e, b \neq 0), (b = -e, b \neq 0), (b = e = 0), (b \neq \pm e)$$

and

$$(a = \frac{1}{4}, d = \frac{1}{4}), (a = \frac{1}{4}, d \neq \frac{1}{4}), (a \neq \frac{1}{4}, d = \frac{1}{4}), (a \neq \frac{1}{4}, d \neq \frac{1}{4}).$$

Remark 4.1. We observe that the determining equations (3.39b), (3.39c), (3.39d), (3.39e), and (3.39f) always have three particular solutions.

$$\xi = \gamma = \tau = \mu = 0, \lambda = cte \text{ means } h = l = k = \tau = 0, s = cte,$$

$$\xi = \gamma = \lambda = \mu = 0, \tau = cte \text{ means } h = l = k = s = 0, \tau = cte,$$

$$\xi = \gamma = \tau = 0, \lambda = 0 \text{ and } \mu \text{ is a solution of the PDE (3.1) means } h = l = k = \tau = 0, s = 0 \text{ and } \mu \text{ is a solution,}$$

These solutions provides us with multiples of the following symmetries:

$$v_u = u \frac{\partial}{\partial u}, \quad v_t = \frac{\partial}{\partial t}, \quad v_\mu = \mu(x, y, t) \frac{\partial}{\partial u} \text{ where } \mu \text{ is a solution of (3.1).}$$

v_μ corresponds to the linear superposition principle, which generates infinite-dimensional subalgebra. v_u illustrates that we can multiply the solutions by constants, and v_t reflects the invariance of the equation under translation in time.

4.1 Case 1 ($a = \frac{1}{4}, d = \frac{1}{4}$)

4.1.1 Subcase 1.1 ($b \neq \pm e, a = \frac{1}{4}, d = \frac{1}{4}$)

From equations (3.39b), (3.39c), (3.39d), (3.39e), and (3.39f), we obtain:

$$\tau_t = 0, \quad h = 0, \quad l_{tt} - b^2 \frac{l}{4} = 0, \quad k_{tt} - e^2 \frac{k}{4} = 0 \quad \text{and} \quad s_t = 0,$$

solving these equations yields:

$$\begin{cases} \xi = \sqrt{x} \left(c_3 \exp\left(\frac{b}{2}t\right) + c_4 \exp\left(-\frac{b}{2}t\right) \right), \\ \gamma = \sqrt{y} \left(c_5 \exp\left(\frac{e}{2}t\right) + c_6 \exp\left(-\frac{e}{2}t\right) \right), \\ \tau = c_1, \\ \phi = \left(2b \exp\left(\frac{b}{2}t\right) c_3 \sqrt{x} + 2e \exp\left(\frac{e}{2}t\right) c_5 \sqrt{y} + c_2 \right) u + \mu(x, y, t). \end{cases}$$

where $c_1, c_2, c_3, c_4, c_5, c_6$ are arbitrary constants and $\mu(x, y, t)$ is an arbitrary solution of the PDE (3.1).

For $c_i = 1$ and $c_j = 0$ if $i \neq j$, so the basis of the Lie algebra is:

$$\begin{aligned} v_1 &= \frac{\partial}{\partial t}, \quad v_2 = u \frac{\partial}{\partial u}, \quad v_3 = \sqrt{x} \exp\left(\frac{b}{2}t\right) \frac{\partial}{\partial x} + 2b \exp\left(\frac{b}{2}t\right) \sqrt{x} u \frac{\partial}{\partial u}, \\ v_4 &= \sqrt{x} \exp\left(-\frac{b}{2}t\right) \frac{\partial}{\partial x} v_5 = \sqrt{y} \exp\left(\frac{e}{2}t\right) \frac{\partial}{\partial y} + 2e \exp\left(\frac{e}{2}t\right) \sqrt{y} u \frac{\partial}{\partial u}, \\ v_6 &= \sqrt{y} \exp\left(-\frac{e}{2}t\right) \frac{\partial}{\partial y}. \end{aligned}$$

The commutation relations between these vector fields are given by the commutator table of $\mathcal{V}_{a,b,d,e}$ as follows:

Table 2: Commutator table ($b \neq \pm e, a = \frac{1}{4}, d = \frac{1}{4}$).

$[v_i, v_j]$	v_1	v_2	v_3	v_4	v_5	v_6	v_μ
v_1	0	0	$-\frac{b}{2}v_3$	$-\frac{e}{2}v_4$	$\frac{b}{2}v_5$	$\frac{e}{2}v_6$	v_{μ_t}
v_2	0	0	0	0	0	0	$-v_\mu$
v_3	$\frac{b}{2}v_3$	0	0	0	bv_2	0	v_{μ_1}
v_4	$\frac{e}{2}v_4$	0	0	0	0	ev_2	v_{μ_2}
v_5	$-\frac{b}{2}v_5$	0	$-bv_2$	0	0	0	v_{μ_3}
v_6	$-\frac{e}{2}v_6$	0	0	$-ev_2$	0	0	v_{μ_4}
v_{μ_t}	$-v_\mu$	v_μ	$-v_{\mu_1}$	$-v_{\mu_2}$	$-v_{\mu_3}$	$-v_{\mu_4}$	0

where $\mu_1 = \exp\left(-\frac{b}{2}t\right) \sqrt{x} \mu_x$, $\mu_2 = \exp\left(-\frac{e}{2}t\right) \sqrt{y} \mu_y$,

$$\mu_3 = -2 \exp\left(\frac{b}{2}t\right) \sqrt{x} b \mu + \exp\left(\frac{b}{2}t\right) \sqrt{x} \mu_x, \quad \mu_4 = -2 \exp\left(\frac{e}{2}t\right) \sqrt{y} e \mu + \exp\left(\frac{e}{2}t\right) \sqrt{y} \mu_y.$$

Remark 4.2. In all subsequent cases, the whole table will correspond to $\mathcal{V}_{a,b,d,e}$ and the table without the last line and column to $\mathcal{L}_{a,b,d,e}$.

Corollary (2.4) guarantees that the set of all infinitesimal symmetries is a Lie algebra. We can conclude that if $\mu(x, y, t)$ is any solution of equation (3.1), so are $\mu_1, \mu_2, \mu_3, \mu_4$; this is easily checked directly.

The one-parameter groups G_i generated by the v_i is given as follows: This gives the transformed point $\exp(\varepsilon v_i)(x, y, t, u) = (\tilde{x}, \tilde{y}, \tilde{t}, \tilde{u})$

$$\begin{aligned}
G_1 &: (x, y, t + \varepsilon, u), \\
G_2 &: (x, y, t, \exp(\varepsilon)u), \\
G_3 &: \left(\frac{(\varepsilon \exp(\frac{bt}{2}) + 2\sqrt{x})^2}{4}, y, t, u \exp\left(\frac{\exp(\frac{bt}{2})b\varepsilon(\varepsilon \exp(\frac{bt}{2}) + 4\sqrt{x})}{2}\right) \right), \\
G_4 &: \left(\frac{(2\exp(\frac{bt}{2})\sqrt{x} + \varepsilon)^2 \exp(-bt)}{4}, y, t, u \right), \\
G_5 &: \left(x, \frac{(\varepsilon \exp(\frac{et}{2}) + 2\sqrt{y})^2}{4}, t, u \exp\left(\frac{\exp(\frac{et}{2})e\varepsilon(\varepsilon \exp(\frac{et}{2}) + 4\sqrt{y})}{2}\right) \right), \\
G_6 &: \left(x, \frac{(2\exp(\frac{et}{2})\sqrt{y} + \varepsilon)^2 \exp(-et)}{4}, t, u \right).
\end{aligned}$$

Because each one-parameter group G_i is a symmetry group, using Theorem 2.5, if $u(x, y, t)$ is a solution of (3.1), so are the u^{ε, v_i} , $1 \leq i \leq 6$, given by:

$$\begin{aligned}
u^{\varepsilon, v_1}(x, y, t) &= u(x, y, t - \varepsilon), \\
u^{\varepsilon, v_2}(x, y, t) &= \exp(\varepsilon)u(x, y, t), \\
u^{\varepsilon, v_3}(x, y, t) &= \exp\left(\frac{b\varepsilon \exp(\frac{bt}{2})(4\sqrt{x} - \varepsilon \exp(\frac{bt}{2}))}{2}\right) u\left(\frac{(2\sqrt{x} - \varepsilon \exp(\frac{bt}{2}))^2}{4}, y, t\right), \\
u^{\varepsilon, v_4}(x, y, t) &= u\left(\frac{(2\exp(\frac{bt}{2})\sqrt{x} - \varepsilon)^2 \exp(-bt)}{4}, y, t\right), \\
u^{\varepsilon, v_5}(x, y, t) &= \exp\left(\frac{e\varepsilon \exp(\frac{et}{2})(4\sqrt{y} - \varepsilon \exp(\frac{et}{2}))}{2}\right) u\left(\frac{(2\sqrt{y} - \varepsilon \exp(\frac{et}{2}))^2}{4}, y, t\right), \\
u^{\varepsilon, v_6}(x, y, t) &= u\left(x, \frac{(2\exp(\frac{et}{2})\sqrt{y} - \varepsilon)^2 \exp(-et)}{4}, t\right).
\end{aligned}$$

4.1.2 Subcase 1.2 ($b = e$, $b \neq 0$, $a = \frac{1}{4}$, $d = \frac{1}{4}$)

$$\begin{cases}
\xi = (c_1 b \exp(bt) - c_2 b \exp(-bt))x + \sqrt{x} \left(c_4 \sqrt{y} + c_5 \exp(\frac{b}{2}t) + c_6 \exp(-\frac{b}{2}t) \right), \\
\gamma = (c_1 b \exp(bt) - c_2 b \exp(-bt))y + \sqrt{y} \left(-c_4 \sqrt{x} + c_7 \exp(\frac{b}{2}t) + c_8 \exp(-\frac{b}{2}t) \right), \\
\tau = c_1 \exp(bt) + c_2 \exp(-bt) + c_3, \\
\phi = \left((2b^2 c_1 \exp(bt))(x + y) + 2bc_5 \exp(\frac{b}{2}t)\sqrt{x} + 2bc_7 \exp(\frac{b}{2}t)\sqrt{y} - bc_1 \exp(bt) - \frac{1}{2}bc_3 + c_9 \right) u + \mu(x, y, t).
\end{cases}$$

$$\begin{aligned}
v_1 &= b \exp(bt)x \frac{\partial}{\partial x} + b \exp(bt)y \frac{\partial}{\partial y} + \exp(bt) \frac{\partial}{\partial t} + 2b \exp(bt)(b(x+y) - \frac{1}{2})u \frac{\partial}{\partial u}, \\
v_2 &= -b \exp(-bt)x \frac{\partial}{\partial x} - b \exp(-bt)y \frac{\partial}{\partial y} + \exp(-bt) \frac{\partial}{\partial t}, \quad v_3 = \frac{\partial}{\partial t} - \frac{b}{2}u \frac{\partial}{\partial u}, \quad v_4 = \sqrt{x}\sqrt{y} \frac{\partial}{\partial x} - \sqrt{x}\sqrt{y} \frac{\partial}{\partial y}, \\
v_5 &= \sqrt{x} \exp(\frac{b}{2}t) \frac{\partial}{\partial x} + 2b\sqrt{x} \exp(\frac{b}{2}t)u \frac{\partial}{\partial u}, \quad v_6 = \sqrt{x} \exp(-\frac{b}{2}t) \frac{\partial}{\partial x}, \\
v_7 &= \sqrt{y} \exp(\frac{b}{2}t) \frac{\partial}{\partial y} + 2b\sqrt{y} \exp(\frac{b}{2}t)u \frac{\partial}{\partial u}, \quad v_8 = \sqrt{y} \exp(-\frac{b}{2}t) \frac{\partial}{\partial y}, \quad v_9 = u \frac{\partial}{\partial u}.
\end{aligned}$$

Table 3: Commutator table($b = e$, $b \neq 0$, $a = \frac{1}{4}$, $d = \frac{1}{4}$).

$[v_i, v_j]$	v_1	v_2	v_3	v_4	v_5	v_6	v_7	v_8	v_9	v_μ
v_1	0	$-2bv_3$	$-bv_1$	0	0	$-bv_5$	0	$-bv_7$	0	v_{μ_1}
v_2	$2bv_3$	0	bv_2	0	bv_6	0	bv_8	0	0	v_{μ_2}
v_3	bv_1	$-bv_2$	0	0	$\frac{b}{2}v_5$	$-\frac{b}{2}v_6$	$\frac{b}{2}v_7$	$-\frac{b}{2}v_8$	0	v_{μ_t}
v_4	0	0	0	0	$\frac{1}{2}v_7$	$\frac{1}{2}v_8$	$-\frac{1}{2}v_5$	$-\frac{1}{2}v_6$	0	v_{μ_3}
v_5	0	$-bv_6$	$-\frac{b}{2}v_5$	$-\frac{1}{2}v_7$	0	$-bv_9$	0	0	0	v_{μ_4}
v_6	bv_5	0	$\frac{b}{2}v_6$	$-\frac{1}{2}v_8$	bv_9	0	0	0	0	v_{μ_5}
v_7	0	$-bv_8$	$-\frac{b}{2}v_7$	$\frac{1}{2}v_5$	0	0	0	$-bv_9$	0	v_{μ_6}
v_8	bv_7	0	$\frac{b}{2}v_8$	$\frac{1}{2}v_6$	0	0	bv_9	0	0	v_{μ_7}
v_9	0	0	0	0	0	0	0	0	0	$-v_\mu$
v_μ	$-v_{\mu_1}$	$-v_{\mu_2}$	$-v_{\mu_t}$	$-v_{\mu_3}$	$-v_{\mu_4}$	$-v_{\mu_5}$	$-v_{\mu_6}$	$-v_{\mu_7}$	v_μ	0

$$\mu_1 = b \exp(bt)x\mu_x + b \exp(bt)y\mu_y + \exp(bt)\mu_t + b \exp(bt)(1 - 2b(x+y))\mu,$$

$$\mu_2 = -b \exp(-bt)x\mu_x - b \exp(-bt)y\mu_y + \exp(bt)\mu_t,$$

$$\mu_3 = \sqrt{x}\sqrt{y}\mu_x - \sqrt{x}\sqrt{y}\mu_y, \quad \mu_4 = \sqrt{x} \exp(\frac{b}{2}t)\mu_x - 2b \exp(\frac{b}{2}t)\sqrt{x}\mu,$$

$$\mu_5 = \sqrt{x} \exp(-\frac{b}{2}t)\mu_x, \quad \mu_6 = \sqrt{y} \exp(\frac{b}{2}t)\mu_y - 2b \exp(\frac{b}{2}t)\sqrt{y}\mu, \quad \mu_7 = \sqrt{y} \exp(-\frac{b}{2}t)\mu_y.$$

The transformed point by a one-parameter group $\exp(\varepsilon v_i)(x, y, t, u) = (\tilde{x}, \tilde{y}, \tilde{t}, \tilde{u})$ are:

$$\begin{aligned}
G_1 &: \left(\frac{x}{1 - b\varepsilon \exp(bt)}, \frac{y}{1 - b\varepsilon \exp(bt)}, t - \frac{\ln(1 - b\varepsilon \exp(bt))}{b}, (1 - b\varepsilon \exp(bt))u \exp\left(\frac{2b^2\varepsilon \exp(bt)(x+y)}{1 - b\varepsilon \exp(bt)}\right) \right), \\
G_2 &: \left(\frac{x}{1 + b\varepsilon \exp(-bt)}, \frac{y}{1 + b\varepsilon \exp(-bt)}, t + \frac{\ln(1 + b\varepsilon \exp(-bt))}{b}, u \right), \\
G_3 &: (x, y, t + \varepsilon, u), \\
G_4 &: \left(\sqrt{x}\sqrt{y} \sin(\varepsilon) + \left(\frac{x-y}{2}\right) \cos(\varepsilon) + \left(\frac{x+y}{2}\right), -\sqrt{x}\sqrt{y} \sin(\varepsilon) - \left(\frac{x-y}{2}\right) \cos(\varepsilon) - \left(\frac{x+y}{2}\right), t, u \right), \\
G_5 &: \left(\frac{(\varepsilon \exp(\frac{bt}{2}) + 2\sqrt{x})^2}{4}, y, t, u \exp\left(\frac{\exp(\frac{bt}{2})b\varepsilon(\varepsilon \exp(\frac{bt}{2}) + 4\sqrt{x})}{2}\right) \right), \\
G_6 &: \left(\frac{(2 \exp(\frac{bt}{2})\sqrt{x} + \varepsilon)^2 \exp(-bt)}{4}, y, t, u \right), \\
G_7 &: \left(x, \frac{(\varepsilon \exp(\frac{bt}{2}) + 2\sqrt{y})^2}{4}, t, u \exp\left(\frac{\exp(\frac{bt}{2})b\varepsilon(\varepsilon \exp(\frac{bt}{2}) + 4\sqrt{y})}{2}\right) \right), \\
G_8 &: \left(x, \frac{(2 \exp(\frac{bt}{2})\sqrt{y} + \varepsilon)^2 \exp(-bt)}{4}, t, u \right), \\
G_9 &: (x, y, t, u e^\varepsilon).
\end{aligned}$$

Since each one parameter group G_i is a symmetry group, using Theorem 2.5 if $u(x, y, t)$ is a solution of (3.1), so are the u^{ε, v_i} , $1 \leq i \leq 9$ given by:

$$\begin{aligned}
u^{\varepsilon, v_1}(x, y, t) &= \frac{1}{1 + b\varepsilon \exp(bt)} \exp(2b^2\varepsilon \exp(bt)(x+y)) u\left(\frac{x}{1 + b\varepsilon \exp(bt)}, \frac{y}{1 + b\varepsilon \exp(bt)}, t - \frac{\ln(1 + \varepsilon b \exp(bt))}{b}\right), \\
u^{\varepsilon, v_2}(x, y, t) &= u\left(\frac{x}{1 - b\varepsilon \exp(-bt)}, \frac{y}{1 - b\varepsilon \exp(-bt)}, t + \frac{\ln(1 - b\varepsilon \exp(-bt))}{b}\right), \\
u^{\varepsilon, v_3}(x, y, t) &= u(x, y, t - \varepsilon), \\
u^{\varepsilon, v_4}(x, y, t) &= u\left(-\sqrt{x}\sqrt{y} \sin(\varepsilon) + \left(\frac{x-y}{2}\right) \cos(\varepsilon) + \left(\frac{x+y}{2}\right), \sqrt{x}\sqrt{y} \sin(\varepsilon) - \left(\frac{x-y}{2}\right) \cos(\varepsilon) - \left(\frac{x+y}{2}\right), t\right), \\
u^{\varepsilon, v_5}(x, y, t) &= \exp\left(\frac{b\varepsilon \exp(\frac{bt}{2})(4\sqrt{x} - \varepsilon \exp(\frac{bt}{2}))}{2}\right) u\left(\frac{(2\sqrt{x} - \varepsilon \exp(\frac{bt}{2}))^2}{4}, y, t\right), \\
u^{\varepsilon, v_6}(x, y, t) &= u\left(\frac{(2 \exp(\frac{bt}{2})\sqrt{x} - \varepsilon)^2 \exp(-bt)}{4}, y, t\right), \\
u^{\varepsilon, v_7}(x, y, t) &= \exp\left(\frac{b\varepsilon \exp(\frac{bt}{2})(4\sqrt{y} - \varepsilon \exp(\frac{bt}{2}))}{2}\right) u\left(\frac{(2\sqrt{y} - \varepsilon \exp(\frac{bt}{2}))^2}{4}, y, t\right), \\
u^{\varepsilon, v_8}(x, y, t) &= u\left(x, \frac{(2 \exp(\frac{bt}{2})\sqrt{y} - \varepsilon)^2 \exp(-bt)}{4}, t\right), \\
u^{\varepsilon, v_9}(x, y, t) &= \exp(\varepsilon)u(x, y, t).
\end{aligned}$$

4.1.3 Subcase 1.3 ($b = -e$, $b \neq 0$, $a = \frac{1}{4}$, $d = \frac{1}{4}$)

$$\begin{aligned}
v_1 &= b \exp(bt)x \frac{\partial}{\partial x} + b \exp(bt)y \frac{\partial}{\partial y} + \exp(bt) \frac{\partial}{\partial t} + \frac{b \exp(bt)(4bx - 1)}{2} u \frac{\partial}{\partial u}, \\
v_2 &= -b \exp(-bt)x \frac{\partial}{\partial x} - b \exp(-bt)y \frac{\partial}{\partial y} + \exp(-bt) \frac{\partial}{\partial t} + \frac{b \exp(-bt)(4by + 1)}{2} u \frac{\partial}{\partial u}, \\
v_3 &= \frac{\partial}{\partial t}, \quad v_4 = \sqrt{x}\sqrt{y} \frac{\partial}{\partial x} - \sqrt{x}\sqrt{y} \frac{\partial}{\partial y} + 2b\sqrt{x}\sqrt{y}u \frac{\partial}{\partial u}, \\
v_5 &= \sqrt{x} \exp\left(\frac{b}{2}t\right) \frac{\partial}{\partial x} + 2b\sqrt{x} \exp\left(\frac{b}{2}t\right) u \frac{\partial}{\partial u} \\
v_6 &= \sqrt{x} \exp\left(-\frac{b}{2}t\right) \frac{\partial}{\partial x}, \quad v_7 = \sqrt{y} \exp\left(-\frac{b}{2}t\right) \frac{\partial}{\partial y} - 2b\sqrt{y} \exp\left(-\frac{b}{2}t\right) u \frac{\partial}{\partial u}, \\
v_8 &= \sqrt{y} \exp\left(\frac{b}{2}t\right) \frac{\partial}{\partial y}, \quad v_9 = u \frac{\partial}{\partial u}.
\end{aligned}$$

Table 4: Commutator table ($b = -e$, $b \neq 0$, $a = \frac{1}{4}$, $d = \frac{1}{4}$).

$[v_i, v_j]$	v_1	v_2	v_3	v_4	v_5	v_6	v_7	v_8	v_9	v_μ
v_1	0	$-2bv_3$	$-bv_1$	0	0	$-bv_5$	$-bv_8$	0	0	v_{μ_1}
v_2	$2bv_3$	0	bv_2	0	bv_6	0	0	bv_7	0	v_{μ_2}
v_3	bv_1	$-bv_2$	0	0	$\frac{b}{2}v_5$	$-\frac{b}{2}v_6$	$-\frac{b}{2}v_7$	$\frac{b}{2}v_8$	0	v_{μ_t}
v_4	0	0	0	0	$\frac{1}{2}v_8$	$\frac{1}{2}v_7$	$-\frac{1}{2}v_6$	$-\frac{1}{2}v_5$	0	v_{μ_3}
v_5	0	$-bv_6$	$-\frac{b}{2}v_5$	$-\frac{1}{2}v_8$	0	$-bv_9$	0	0	0	v_{μ_4}
v_6	bv_5	0	$\frac{b}{2}v_6$	$-\frac{1}{2}v_7$	bv_9	0	0	0	0	v_{μ_5}
v_7	bv_8	0	$\frac{b}{2}v_7$	$\frac{1}{2}v_6$	0	0	0	bv_9	0	v_{μ_6}
v_8	0	$-bv_7$	$-\frac{b}{2}v_8$	$\frac{1}{2}v_5$	0	0	$-bv_9$	0	0	v_{μ_7}
v_9	0	0	0	0	0	0	0	0	0	$-v_\mu$
v_μ	$-v_{\mu_1}$	$-v_{\mu_2}$	$-v_{\mu_t}$	$-v_{\mu_3}$	$-v_{\mu_4}$	$-v_{\mu_5}$	$-v_{\mu_6}$	$-v_{\mu_7}$	v_μ	0

$$\begin{aligned}
\mu_1 &= b \exp(bt)x\mu_x + b \exp(bt)y\mu_y + \exp(bt)\mu_t + \frac{1}{2}b(1 - 4b) \exp(bt)\mu, \\
\mu_2 &= -b \exp(-bt)x\mu_x - b \exp(-bt)y\mu_y + \exp(-bt)\mu_t - \frac{1}{2}b(1 + by) \exp(bt)\mu, \\
\mu_3 &= \sqrt{x}\sqrt{y}\mu_x - \sqrt{x}\sqrt{y}\mu_y - 2b\sqrt{x}\sqrt{y}\mu, \quad \mu_4 = \sqrt{x} \exp\left(\frac{b}{2}t\right)\mu_x - 2b \exp\left(\frac{b}{2}t\right)\sqrt{x}\mu, \\
\mu_5 &= \sqrt{x} \exp\left(-\frac{b}{2}t\right)\mu_x, \quad \mu_6 = \sqrt{y} \exp\left(-\frac{b}{2}t\right)\mu_y - 2b \exp\left(-\frac{b}{2}t\right)\sqrt{y}\mu, \quad \mu_7 = \sqrt{y} \exp\left(-\frac{b}{2}t\right)\mu_y.
\end{aligned}$$

The transformed point by a one-parameter group $\exp(\varepsilon v_i)(x, y, t, u) = (\tilde{x}, \tilde{y}, \tilde{t}, \tilde{u})$ are:

$$\begin{aligned}
G_1 &: \left(\frac{x}{1 - b\varepsilon \exp(bt)}, \frac{y}{1 - b\varepsilon \exp(bt)}, t - \frac{\ln(1 - b\varepsilon \exp(bt))}{b}, \sqrt{1 - b\varepsilon \exp(bt)} u \exp\left(\frac{2b^2\varepsilon \exp(bt)x}{1 - b\varepsilon \exp(bt)}\right) \right), \\
G_2 &: \left(\frac{x}{1 + b\varepsilon \exp(-bt)}, \frac{y}{1 + b\varepsilon \exp(-bt)}, t + \frac{\ln(1 - b\varepsilon \exp(-bt))}{b}, \sqrt{1 + b\varepsilon \exp(-bt)} u \exp\left(\frac{2b^2\varepsilon \exp(-bt)y}{1 + b\varepsilon \exp(-bt)}\right) \right), \\
G_3 &: (x, y, t + \varepsilon, u), \\
G_4 &: \left(\sqrt{x}\sqrt{y} \sin(\varepsilon) + \left(\frac{x-y}{2}\right) \cos(\varepsilon) + \left(\frac{x+y}{2}\right), -\sqrt{x}\sqrt{y} \sin(\varepsilon) - \left(\frac{x-y}{2}\right) \cos(\varepsilon) + \left(\frac{x+y}{2}\right), t, \right. \\
&\quad \left. u \exp\left(b(2\sqrt{x}\sqrt{y} \sin(\varepsilon) + (x-y) \cos(\varepsilon) - (x-y))\right) \right), \\
G_5 &: \left(\frac{(\varepsilon \exp(\frac{bt}{2}) + 2\sqrt{x})^2}{4}, y, t, u \exp\left(\frac{\exp(\frac{bt}{2})b\varepsilon (\varepsilon \exp(\frac{bt}{2}) + 4\sqrt{x})}{2}\right) \right), \\
G_6 &: \left(\frac{(2\exp(\frac{bt}{2})\sqrt{x} + \varepsilon)^2 \exp(-bt)}{4}, y, t, u \right), \\
G_7 &: \left(x, \frac{(\varepsilon \exp(\frac{et}{2}) + 2\sqrt{y})^2}{4}, t, u \exp\left(\frac{\exp(\frac{et}{2})e\varepsilon (\varepsilon \exp(\frac{et}{2}) + 4\sqrt{y})}{2}\right) \right), \\
G_8 &: \left(x, \frac{(2\sqrt{y} + \varepsilon \exp(\frac{bt}{2}))^2}{4}, t, u \right), \\
G_9 &: (x, y, t, u \exp(\varepsilon)).
\end{aligned}$$

Since each one parameter group G_i is a symmetry group, using Theorem 2.5 if $u(x, y, t)$ is a solution of

(3.1), so are the u^{ε, v_i} , $1 \leq i \leq 9$ given by:

$$\begin{aligned}
u^{\varepsilon, v_1}(x, y, t) &= \frac{\exp\left(\frac{2b^2 \varepsilon x \exp(bt)}{1 + b\varepsilon \exp(bt)}\right)}{\sqrt{1 + b\varepsilon \exp(bt)}} u\left(\frac{x}{1 + b\varepsilon \exp(bt)}, \frac{y}{1 + b\varepsilon \exp(bt)}, t - \frac{\ln(1 + \varepsilon b \exp(bt))}{b}\right), \\
u^{\varepsilon, v_2}(x, y, t) &= \sqrt{1 + \frac{b\varepsilon \exp(-bt)}{1 + b\varepsilon \exp(-bt)}} \exp\left(\frac{2b^2 \varepsilon \exp(-bt)y}{(1 - b\varepsilon \exp(-bt))(1 + 2b\varepsilon \exp(-bt))}\right) \\
&\quad u\left(\frac{x}{(1 - \varepsilon b \exp(-bt))}, \frac{y}{(1 - \varepsilon b \exp(-bt))}, t + \frac{\ln(1 + b\varepsilon \exp(-bt))}{b}\right), \\
u^{\varepsilon, v_3}(x, y, t) &= u(x, y, t - \varepsilon), \\
u^{\varepsilon, v_4}(x, y, t) &= \exp\left[b\left(\sqrt{\left(-\sqrt{xy} \sin(\varepsilon) + \frac{(x-y)}{2} \cos(\varepsilon) + \frac{(x+y)}{2}\right)\left(\sqrt{xy} \sin(\varepsilon) - \frac{(x-y)}{2} \cos(\varepsilon) + \frac{(x+y)}{2}\right)}\right)} \sin(\varepsilon)\right. \\
&\quad \left. + (-2\sqrt{x}\sqrt{y} \sin(\varepsilon) + (x-y) \cos(\varepsilon))(\cos(\varepsilon) - 1)\right] \\
&\quad u\left(-\sqrt{x}\sqrt{y} \sin(\varepsilon) + \frac{(x-y)}{2} \cos(\varepsilon) + \frac{(x+y)}{2}, \sqrt{x}\sqrt{y} \sin(\varepsilon) - \frac{(x-y)}{2} \cos(\varepsilon) - \frac{(x+y)}{2}, t\right), \\
u^{\varepsilon, v_5}(x, y, t) &= \exp\left(\frac{b\varepsilon \exp(\frac{bt}{2})(4\sqrt{x} - \varepsilon \exp(\frac{bt}{2}))}{2}\right) u\left(\frac{(2\sqrt{x} - \varepsilon \exp(\frac{bt}{2}))^2}{4}, y, t\right), \\
u^{\varepsilon, v_6}(x, y, t) &= u\left(\frac{(2 \exp(\frac{bt}{2})\sqrt{x} - \varepsilon)^2}{4} \exp(-bt), y, t\right), \\
u^{\varepsilon, v_7}(x, y, t) &= \exp\left(-\frac{b\varepsilon e^{-\frac{bt}{2}}(4\sqrt{y} - \varepsilon \exp(-\frac{bt}{2}))}{2}\right) u\left(x, \frac{(2\sqrt{y} - \varepsilon \exp(-\frac{bt}{2}))^2}{4}, t\right), \\
u^{\varepsilon, v_8}(x, y, t) &= u\left(x, \frac{(2\sqrt{y} - \varepsilon \exp(\frac{bt}{2}))^2}{4}, t\right), \\
u^{\varepsilon, v_9}(x, y, t) &= \exp(\varepsilon)u(x, y, t).
\end{aligned}$$

4.1.4 Subcase 1.4 ($b = e = 0$, $a = \frac{1}{4}$, $d = \frac{1}{4}$)

$$\begin{aligned}
v_1 &= 2tx \frac{\partial}{\partial x} + 2ty \frac{\partial}{\partial y} + t^2 \frac{\partial}{\partial t} + (2(x+y) - \frac{t}{2})u \frac{\partial}{\partial u}, \\
v_2 &= x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} + t \frac{\partial}{\partial t}, \\
v_3 &= \frac{\partial}{\partial t}, \quad v_4 = \sqrt{x}\sqrt{y} \frac{\partial}{\partial x} - \sqrt{x}\sqrt{y} \frac{\partial}{\partial y}, \quad v_5 = \sqrt{xt} \frac{\partial}{\partial x} + 2\sqrt{x}u \frac{\partial}{\partial u}, \\
v_6 &= \sqrt{x} \frac{\partial}{\partial x}, \quad v_7 = \sqrt{yt} \frac{\partial}{\partial y} + 2\sqrt{y}u \frac{\partial}{\partial u}, \quad v_8 = \sqrt{y} \frac{\partial}{\partial y}, \quad v_9 = u \frac{\partial}{\partial u}.
\end{aligned}$$

Remark 4.3. $\{v_1, \dots, v_9\}$ is the largest Lie algebra of symmetries that we found.

Table 5: Commutator table($b = e = 0, a = \frac{1}{4}, d = \frac{1}{4}$).

$[v_i, v_j]$	v_1	v_2	v_3	v_4	v_5	v_6	v_7	v_8	v_9	v_μ
v_1	0	$-v_1$	$-2v_2 + \frac{1}{2}v_9$	0	0	$-v_5$	0	$-v_7$	0	v_{μ_1}
v_2	v_1	0	$-v_3$	0	$\frac{1}{2}v_5$	$-\frac{1}{2}v_6$	$\frac{1}{2}v_7$	$-\frac{1}{2}v_8$	0	v_{μ_2}
v_3	$2v_2 - \frac{1}{2}v_9$	v_3	0	0	v_6	0	v_8	0	0	v_{μ_3}
v_4	0	0	0	0	$\frac{1}{2}v_7$	$\frac{1}{2}v_8$	$-\frac{1}{2}v_5$	$-\frac{1}{2}v_6$	0	v_{μ_3}
v_5	0	$-\frac{1}{2}v_5$	$-v_6$	$-\frac{1}{2}v_7$	0	$-v_9$	0	0	0	v_{μ_4}
v_6	0	$\frac{1}{2}v_6$	0	$-\frac{1}{2}v_8$	v_9	0	0	0	0	v_{μ_5}
v_7	0	$-\frac{1}{2}v_7$	$-v_8$	$\frac{1}{2}v_5$	0	0	0	$-v_9$	0	v_{μ_6}
v_8	$-v_7$	$\frac{1}{2}v_8$	0	$\frac{1}{2}v_6$	0	0	v_9	0	0	v_{μ_7}
v_9	0	0	0	0	0	0	0	0	0	$-v_\mu$
v_μ	$-v_{\mu_1}$	$-v_{\mu_2}$	$-v_{\mu_3}$	$-v_{\mu_3}$	$-v_{\mu_4}$	$-v_{\mu_5}$	$-v_{\mu_6}$	$-v_{\mu_7}$	v_μ	0

$$\begin{aligned} \mu_1 &= 2tx\mu_x + 2ty\mu_y + t^2\mu_t + \left(\frac{t}{2} - 2(x+y)\right)\mu, \quad \mu_2 = x\mu_x + y\mu_y + t\mu_t, \quad \mu_3 = \mu_t, \\ \mu_4 &= \sqrt{x}\sqrt{y}\mu_x - \sqrt{x}\sqrt{y}\mu_y, \quad \mu_5 = \sqrt{x}t\mu_x - 2\sqrt{x}\mu, \quad \mu_6 = \sqrt{x}\mu_x, \quad \mu_7 = \sqrt{y}t\mu_y - 2\sqrt{y}\mu. \end{aligned}$$

The transformed point by a one-parameter group $\exp(\varepsilon v_i)(x, y, t, u) = (\tilde{x}, \tilde{y}, \tilde{t}, \tilde{u})$ are:

$$G_1 : \left(\frac{x}{(1-t\varepsilon)^2}, \frac{y}{(1-t\varepsilon)^2}, \frac{t}{1-t\varepsilon}, u \exp\left(\frac{2\varepsilon(x+y)}{1-t\varepsilon}\right) \sqrt{1-t\varepsilon} \right),$$

$$G_2 : (\exp(\varepsilon)x, \exp(\varepsilon)y, \exp(\varepsilon)t, u),$$

$$G_3 : (x, y, t + \varepsilon, u),$$

$$G_4 : \left(\sqrt{x}\sqrt{y}\sin(\varepsilon) + \left(\frac{x-y}{2}\right)\cos(\varepsilon) + \left(\frac{x+y}{2}\right), -\sqrt{x}\sqrt{y}\sin(\varepsilon) - \left(\frac{x-y}{2}\right)\cos(\varepsilon) + \left(\frac{x+y}{2}\right), t, u \right),$$

$$G_5 : \left(x + t\sqrt{x}\varepsilon + \frac{t^2\varepsilon^2}{4}, y, t, u \exp\left(\frac{\varepsilon(t\varepsilon + 4\sqrt{x})}{2}\right) \right),$$

$$G_6 : \left(\sqrt{x}\varepsilon + \frac{\varepsilon^2}{4} + x, y, t, u \right),$$

$$G_7 : \left(x, y + t\sqrt{y}\varepsilon + \frac{t^2\varepsilon^2}{4}, t, u \exp\left(\frac{\varepsilon(t\varepsilon + 4\sqrt{y})}{2}\right) \right),$$

$$G_8 : (x, \sqrt{y}\varepsilon + \frac{\varepsilon^2}{4} + y, t, u),$$

$$G_9 : (x, y, t, \exp(\varepsilon)u).$$

Since each one parameter group G_i is a symmetry group, using Theorem 2.5 if $u(x, y, t)$ is a solution of

(3.1), so are the u^{ε, v_i} , $1 \leq i \leq 9$ given by:

$$\begin{aligned}
u^{\varepsilon, v_1}(x, y, t) &= \frac{\exp\left(\frac{2\varepsilon(x+y)}{(1+t\varepsilon)^3}\right)}{\sqrt{1+t\varepsilon}} u\left(\frac{x}{(1+t\varepsilon)^2}, \frac{y}{(1+t\varepsilon)^2}, \frac{t}{1+t\varepsilon}\right), \\
u^{\varepsilon, v_2}(x, y, t) &= u(\exp(-\varepsilon)x, \exp(-\varepsilon)y, \exp(-\varepsilon)t), \\
u^{\varepsilon, v_3}(x, y, t) &= u(x, y, t - \varepsilon), \\
u^{\varepsilon, v_4}(x, y, t) &= u\left(-\sqrt{x}\sqrt{y}\sin(\varepsilon) + \left(\frac{x-y}{2}\right)\cos(\varepsilon) + \left(\frac{x+y}{2}\right), \sqrt{x}\sqrt{y}\sin(\varepsilon) - \left(\frac{x-y}{2}\right)\cos(\varepsilon) - \left(\frac{x+y}{2}\right), t\right), \\
u^{\varepsilon, v_5}(x, y, t) &= \exp\left(\frac{\varepsilon\left(\varepsilon t + 4\sqrt{x-t\sqrt{x}\varepsilon + \frac{t^2\varepsilon^2}{2}}\right)}{2}\right) u\left(x - t\sqrt{x}\varepsilon + \frac{t^2\varepsilon^2}{2}, y, t\right), \\
u^{\varepsilon, v_6}(x, y, t) &= u\left(\sqrt{x}\varepsilon - \frac{\varepsilon^2}{4} + x, y, t\right), \\
u^{\varepsilon, v_7}(x, y, t) &= \exp\left(\frac{\varepsilon\left(\varepsilon t + 4\sqrt{y-t\sqrt{y}\varepsilon + \frac{t^2\varepsilon^2}{2}}\right)}{2}\right) u\left(x, y - t\sqrt{y}\varepsilon + \frac{t^2\varepsilon^2}{2}, t\right), \\
u^{\varepsilon, v_8}(x, y, t) &= u\left(-\sqrt{x}\varepsilon + \frac{\varepsilon^2}{4} + x, y, t\right), \\
u^{\varepsilon, v_9}(x, y, t) &= \exp(\varepsilon)u(x, y, t).
\end{aligned}$$

4.2 Case 2 ($a \neq \frac{1}{4}, d \neq \frac{1}{4}$)

Because $a \neq \frac{1}{4}, d \neq \frac{1}{4}$ and from remark (3.35), then $h = k = l = 0$ in all the following subcases.

4.2.1 Subcase 2.1 ($b \neq \pm e, a \neq \frac{1}{4}, d \neq \frac{1}{4}$)

$$\tau = c_1, \quad \xi = \gamma = 0, \quad \phi = c_2u + \mu(x, y, t).$$

$$v_1 = \frac{\partial}{\partial t}, \quad v_2 = u \frac{\partial}{\partial u}$$

Remark 4.4. $\{v_1, v_2\}$ is the smallest Lie algebra of infinitesimal symmetries that we found.

Table 6: Commutator table($b \neq \pm e, a \neq \frac{1}{4}, d \neq \frac{1}{4}$).

$[v_i, v_j]$	v_1	v_2	v_μ
v_1	0	0	$v_{\mu t}$
v_2	0	0	v_μ
v_μ	$-v_{\mu t}$	$-v_\mu$	0

The transformed point by a one-parameter group $\exp(\varepsilon v_i)(x, y, t, u) = (\tilde{x}, \tilde{y}, \tilde{t}, \tilde{u})$ are:

$$G_1 : (x, y, t + \varepsilon, u), \quad G_2 : (x, y, t, \exp(\varepsilon)u).$$

Since each one parameter group G_i is a symmetry group, using Theorem 2.5 if $u(x, y, t)$ is a solution of (3.1), so are the u^{ε, v_i} , $1 \leq i \leq 2$ given by:

$$u^{\varepsilon, v_1}(x, y, t) = f(x, y, t - \varepsilon), \quad u^{\varepsilon, v_2}(x, y, t) = \exp(\varepsilon)f(x, y, t).$$

4.2.2 Subcase 2.2($b = e, b \neq 0, a \neq \frac{1}{4}, d \neq \frac{1}{4}$)

$$v_1 = b \exp(bt)x \frac{\partial}{\partial x} + b \exp(bt)y \frac{\partial}{\partial y} + b \exp(bt)2(b(x+y) - (a+d)u) \frac{\partial}{\partial u},$$

$$v_2 = -b \exp(-bt)x \frac{\partial}{\partial x} - b \exp(-bt)y \frac{\partial}{\partial y} + \exp(-bt) \frac{\partial}{\partial t}, v_3 = \frac{\partial}{\partial t} - b(a+d)u \frac{\partial}{\partial u}, v_4 = u \frac{\partial}{\partial u}.$$

Table 7: Commutator table($b = e, b \neq 0, a \neq \frac{1}{4}, d \neq \frac{1}{4}$).

$[v_i, v_j]$	v_1	v_2	v_3	v_4	v_μ
v_1	0	$-2bv_3$	$-bv_1$	0	v_{μ_1}
v_2	$2bv_3$	0	bv_2	0	v_{μ_2}
v_3	bv_1	$-bv_2$	0	0	v_{μ_3}
v_4	0	0	0	0	$-v_\mu$
v_μ	$-v_{\mu_1}$	$-v_{\mu_2}$	$-v_{\mu_3}$	v_μ	0

$$\mu_1 = b \exp(bt)x\mu_x + b \exp(bt)y\mu_y + \exp(bt)\mu_t + b(a+d-b(x+y)) \exp(bt)\mu,$$

$$\mu_2 = -b \exp(-bt)x\mu_x - b \exp(-bt)y\mu_y + \exp(-bt)\mu_t, \mu_3 = b(a+d)\mu + \mu_t.$$

The transformed point by a one-parameter group $\exp(\varepsilon v_i)(x, y, t, u) = (\tilde{x}, \tilde{y}, \tilde{t}, \tilde{u})$ are:

$$G_1 : \left(\frac{x}{1 - \varepsilon b \exp(bt)}, \frac{y}{1 - \varepsilon b \exp(bt)}, t - \frac{\ln(1 - \varepsilon b \exp(bt))}{b}, (1 - \varepsilon b \exp(bt))^{2(d+a)} u \exp\left(\frac{2b^2 \exp(bt)\varepsilon(x+y)}{1 - \varepsilon b \exp(bt)}\right) \right),$$

$$G_2 : \left(\frac{x}{1 + \varepsilon b \exp(-bt)}, \frac{y}{1 + \varepsilon b \exp(-bt)}, t + \frac{\ln(1 + b\varepsilon \exp(-bt))}{b}, u \right),$$

$$G_3 : (x, y, t + \varepsilon, \exp(-\varepsilon b(a+d))u),$$

$$G_4 : (x, y, t, \exp(\varepsilon)u).$$

Since each one parameter group G_i is a symmetry group, using Theorem 2.5 if $u(x, y, t)$ is a solution of (3.1), so are the $u^{\varepsilon, v_i}, 1 \leq i \leq 4$ given by:

$$u^{\varepsilon, v_1}(x, y, t) = \frac{\exp\left(\frac{2b^2 \varepsilon \exp(bt)(x+y)}{1 + b\varepsilon \exp(bt)}\right)}{(1 + b\varepsilon \exp(bt))^{2(a+d)}} u\left(\frac{x}{1 + \varepsilon b \exp(bt)}, \frac{y}{1 + \varepsilon b \exp(bt)}, t - \frac{\ln(1 + b\varepsilon \exp(bt))}{b}\right),$$

$$u^{\varepsilon, v_2}(x, y, t) = u\left(\frac{x}{1 - \varepsilon b \exp(-bt)}, \frac{y}{1 - \varepsilon b \exp(-bt)}, t + \frac{\ln(1 - \exp(-bt)b\varepsilon)}{b}\right),$$

$$u^{\varepsilon, v_3}(x, y, t) = \exp(-\varepsilon b(a+d))u(x, y, t - \varepsilon),$$

$$u^{\varepsilon, v_4}(x, y, t) = \exp(\varepsilon)u(x, y, t).$$

4.2.3 Subcase 2.3 ($b = -e$, $b \neq 0$, $a \neq \frac{1}{4}$, $d \neq \frac{1}{4}$)

$$\begin{aligned} v_1 &= b \exp(bt)x \frac{\partial}{\partial x} + b \exp(bt)y \frac{\partial}{\partial y} + \exp(bt) \frac{\partial}{\partial t} - 2b(-bx + a) \exp(bt)u \frac{\partial}{\partial u}, \\ v_2 &= -b \exp(-bt)x \frac{\partial}{\partial x} - b \exp(-bt)y \frac{\partial}{\partial y} + \exp(-bt) \frac{\partial}{\partial t} + 2b(by + d) \exp(-bt)u \frac{\partial}{\partial u}, \\ v_3 &= \frac{\partial}{\partial t} + b(db - ab)u \frac{\partial}{\partial u}, \quad v_4 = u \frac{\partial}{\partial u}. \end{aligned}$$

Table 8: Commutator table($b = -e$, $b \neq 0$, $a \neq \frac{1}{4}$, $d \neq \frac{1}{4}$).

$[v_i, v_j]$	v_1	v_2	v_3	v_4	v_μ
v_1	0	$-2bv_3$	$-bv_1$	0	v_{μ_1}
v_2	$2bv_3$	0	$2bv_2$	0	v_{μ_2}
v_3	bv_1	$-bv_2$	0	0	v_{μ^3}
v_4	0	0	0	0	$-v_\mu$
v_μ	$-v_{\mu_1}$	$-v_{\mu_2}$	$-v_{\mu_3}$	v_μ	0

$$\begin{aligned} \mu_1 &= b \exp(bt)x\mu_x + b \exp(bt)y\mu_y + \exp(bt)\mu_t + 2b(a - bx) \exp(bt)\mu, \\ \mu_2 &= -b \exp(-bt)x\mu_x - b \exp(-bt)y\mu_y + \exp(-bt)\mu_t - 2b(d + bx) \exp(-bt)\mu, \\ \mu_3 &= b^2(a - d)\mu + \mu_t. \end{aligned}$$

The transformed point by a one-parameter group $\exp(\varepsilon v_i)(x, y, t, u) = (\tilde{x}, \tilde{y}, \tilde{t}, \tilde{u})$ are:

$$\begin{aligned} G_1 &: \left(\frac{x}{1 - b\varepsilon \exp(bt)}, \frac{y}{1 - b\varepsilon \exp(bt)}, t - \frac{\ln(1 - b\varepsilon \exp(bt))}{b}, \frac{u \exp\left(\frac{2b^2\varepsilon \exp(bt)x}{1 - b\varepsilon \exp(bt)}\right)}{1 - b\varepsilon \exp(bt)} \right), \\ G_2 &: \left(\frac{x}{1 + b\varepsilon \exp(-bt)}, \frac{y}{1 + b\varepsilon \exp(-bt)}, t + \frac{\ln(1 + b\varepsilon \exp(-bt))}{b}, \frac{\exp\left(-\frac{2b(\exp(-bt)dt + b\varepsilon(dt - y))}{b\varepsilon + \exp(-bt)}\right)u}{(b\varepsilon + \exp(-bt))^{-2d}} \right), \\ G_3 &: (x, y, t + \varepsilon, b(db - ab)\varepsilon + u(x, y, t)), \\ G_4 &: (x, y, t, \exp(\varepsilon)u). \end{aligned}$$

Since each one parameter group G_i is a symmetry group, using Theorem 2.5 if $u(x, y, t)$ is a solution of

(3.1), so are the u^{ε, v_i} , $1 \leq i \leq 4$ given by:

$$u^{\varepsilon, v_1}(x, y, t) = (1 + b\varepsilon \exp(bt)) \exp\left(\frac{2b^2\varepsilon \exp(bt)x}{1 + b\varepsilon \exp(bt)}\right) u\left(\frac{x}{1 + b\varepsilon \exp(bt)}, \frac{y}{1 + b\varepsilon \exp(bt)}, t - \frac{\ln(1 + b\varepsilon \exp(bt))}{b}\right),$$

$$u^{\varepsilon, v_2}(x, y, t) = \frac{\exp\left(\frac{2b^2\varepsilon \exp(-bt)y}{1 - b\varepsilon \exp(-bt)}\right)}{(1 - b\varepsilon \exp(-bt))^{2d}} u\left(\frac{x}{1 - b\varepsilon \exp(-bt)}, \frac{y}{1 - b\varepsilon \exp(-bt)}, t + \frac{\ln(1 - b\varepsilon \exp(-bt))}{b}\right),$$

$$u^{\varepsilon, v_3}(x, y, t) = \exp(\varepsilon b(d - a))u(x, y, t - \varepsilon),$$

$$u^{\varepsilon, v_4}(x, y, t) = \exp(\varepsilon)u(x, y, t)$$

4.2.4 Subcase 2.4 ($b = e = 0$, $a \neq \frac{1}{4}$, $d \neq \frac{1}{4}$)

$$v_1 = 2tx \frac{\partial}{\partial x} + 2ty \frac{\partial}{\partial y} + t^2 \frac{\partial}{\partial t} + (2(x + y) - 2(a + d)t)u \frac{\partial}{\partial u}, \quad v_2 = x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} + t \frac{\partial}{\partial t}, \quad v_3 = \frac{\partial}{\partial t}, \quad v_4 = u \frac{\partial}{\partial u}.$$

Table 9: Commutator table ($b = e = 0$, $a \neq \frac{1}{4}$, $d \neq \frac{1}{4}$).

$[v_i, v_j]$	v_1	v_2	v_3	v_4	v_μ
v_1	0	$-v_1$	$-2v_2 + 2(a + d)v_4$	0	v_{μ_1}
v_2	v_1	0	$-v_3$	0	v_{μ_2}
v_3	$2v_2 - 2(a + d)v_4$	v_3	0	0	v_{μ_3}
v_4	0	0	0	0	$-v_\mu$
v_μ	$-v_{\mu_1}$	$-v_{\mu_2}$	$-v_{\mu_3}$	v_μ	0

$$\mu_1 = 2tx\mu_x + 2ty\mu_y + t^2\mu_t + (2t(a + d) - 2(x + y))\mu, \quad \mu_2 = x\mu_x + y\mu_y + t\mu_t, \quad \mu_3 = \mu_t.$$

The transformed point by a one-parameter group $\exp(\varepsilon v_i)(x, y, t, u) = (\tilde{x}, \tilde{y}, \tilde{t}, \tilde{u})$ are:

$$G_1 : \left(\frac{x}{(1 - t\varepsilon)^2}, \frac{y}{(1 - t\varepsilon)^2}, \frac{t}{1 - t\varepsilon}, u \exp\left(\frac{2\varepsilon(x + y)}{1 - t\varepsilon}\right) (1 - t\varepsilon)^{a+d} \right),$$

$$G_2 : (\exp(\varepsilon)x, \exp(\varepsilon)y, \exp(\varepsilon)t, u),$$

$$G_3 : (x, y, t + \varepsilon, u),$$

$$G_4 : (x, y, t, \exp(\varepsilon)u).$$

Since each one parameter group G_i is a symmetry group, using Theorem 2.5 if $u(x, y, t)$ is a solution of (3.1), so are the u^{ε, v_i} , $1 \leq i \leq 9$ given by:

$$\begin{aligned}
u^{\varepsilon, v_1}(x, y, t) &= \frac{\exp\left(\frac{2\varepsilon(x+y)}{(1+t\varepsilon)^3}\right)}{(1+t\varepsilon)^{a+d}} u\left(\frac{x}{(1+t\varepsilon)^2}, \frac{y}{(1+t\varepsilon)^2}, \frac{t}{(1+t\varepsilon)}\right), \\
u^{\varepsilon, v_2}(x, y, t) &= u(\exp(-\varepsilon)x, \exp(-\varepsilon)y, \exp(-\varepsilon)t), \\
u^{\varepsilon, v_3}(x, y, t) &= u(x, y, t - \varepsilon), \\
u^{\varepsilon, v_4}(x, y, t) &= \exp(\varepsilon)u(x, y, t).
\end{aligned}$$

4.3 Case 3 ($a = \frac{1}{4}$, $d \neq \frac{1}{4}$)

Since $d \neq \frac{1}{4}$ and from remark (3.35), using (3.39b) we find that $h = k = 0$ and $l_{tt} - \frac{b^2}{4}l = 0$ in all the following subcases.

4.3.1 Subcase 3.1 ($b \neq \pm e$, $a = \frac{1}{4}$, $d \neq \frac{1}{4}$)

Using (3.39b) we find out that $\tau_t = 0$. Thus

$$\xi = \sqrt{x} \left(c_3 \exp\left(\frac{b}{2}t\right) + c_4 \exp\left(-\frac{b}{2}t\right) \right), \quad \gamma = 0, \quad \tau = c_1, \quad \phi = \left(2bc_3 \exp\left(\frac{b}{2}t\right)\sqrt{x} + c_2 \right) u + \mu(x, y, t)$$

$$v_1 = \frac{\partial}{\partial t}, \quad v_2 = u \frac{\partial}{\partial u}, \quad v_3 = \sqrt{x} \exp\left(\frac{b}{2}t\right) \frac{\partial}{\partial x} + 2b \exp\left(\frac{b}{2}t\right) \sqrt{x} u \frac{\partial}{\partial u}, \quad v_4 = \sqrt{x} \exp\left(-\frac{b}{2}t\right) \frac{\partial}{\partial x}.$$

Table 10: Commutator table ($b \neq \pm e$, $a = \frac{1}{4}$, $d \neq \frac{1}{4}$).

$[v_i, v_j]$	v_1	v_2	v_3	v_4	v_μ
v_1	0	0	$\frac{b}{2}v_3$	$-\frac{b}{2}v_4$	v_{μ_t}
v_2	0	0	0	0	$-v_\mu$
v_3	$-\frac{b}{2}v_3$	0	0	$-bv_2$	v_{μ_1}
v_4	$\frac{b}{2}v_4$	0	bv_2	0	v_{μ_2}
v_μ	$-v_{\mu_t}$	v_μ	$-v_{\mu^1}$	$-v_{\mu^2}$	0

$$\mu_1 = -2 \exp\left(\frac{b}{2}t\right) \sqrt{x} b \mu + \exp\left(\frac{b}{2}t\right) \sqrt{x} \mu_x, \quad \mu_2 = \exp\left(-\frac{b}{2}t\right) \sqrt{x} \mu_x.$$

The transformed point by a one-parameter group $\exp(\varepsilon v_i)(x, y, t, u) = (\tilde{x}, \tilde{y}, \tilde{t}, \tilde{u})$ are:

$$\begin{aligned}
G_1 &: (x, y, t + \varepsilon, u), \\
G_2 &: (x, y, t, \exp(\varepsilon)u), \\
G_3 &: \left(\frac{(\varepsilon \exp\left(\frac{bt}{2}\right) + 2\sqrt{x})^2}{4}, y, t, u \exp\left(\frac{\exp\left(\frac{bt}{2}\right)b\varepsilon (\varepsilon \exp\left(\frac{bt}{2}\right) + 4\sqrt{x})}{2}\right) \right), \\
G_4 &: \left(\frac{(2 \exp\left(\frac{bt}{2}\right)\sqrt{x} + \varepsilon)^2 \exp(-bt)}{4}, y, t, u \right).
\end{aligned}$$

Since each one parameter group G_i is a symmetry group, using Theorem 2.5 if $u(x, y, t)$ is a solution of (3.1), so are the u^{ε, v_i} , $1 \leq i \leq 4$ given by:

$$\begin{aligned} u^{\varepsilon, v_1}(x, y, t) &= u(x, y, t - \varepsilon), \\ u^{\varepsilon, v_2}(x, y, t) &= \exp(\varepsilon)u(x, y, t), \\ u^{\varepsilon, v_3}(x, y, t) &= \exp\left(\frac{\exp(\frac{bt}{2})b\varepsilon(4\sqrt{x} - \varepsilon \exp(\frac{bt}{2}))}{2}\right) u\left(\frac{(2\sqrt{x} - \varepsilon \exp(\frac{bt}{2}))^2}{4}, y, t\right), \\ u^{\varepsilon, v_4}(x, y, t) &= u\left(\frac{(2\exp(\frac{bt}{2})\sqrt{x} - \varepsilon)^2 \exp(-bt)}{4}, y, t\right). \end{aligned}$$

4.3.2 Subcase 3.2 ($b = e$, $b \neq 0$, $a = \frac{1}{4}$, $d \neq \frac{1}{4}$)

$$\begin{aligned} v_1 &= b \exp(bt)x \frac{\partial}{\partial x} + b \exp(bt)y \frac{\partial}{\partial y} + \exp(bt) \frac{\partial}{\partial t} + 2b \exp(bt) \left(b(x+y) - \left(\frac{1}{4} + d\right) \right) u \frac{\partial}{\partial u}, \\ v_2 &= -b \exp(-bt)x \frac{\partial}{\partial x} - b \exp(-bt)y \frac{\partial}{\partial y} + \exp(-bt) \frac{\partial}{\partial t}, \quad v_3 = \frac{\partial}{\partial t} - b\left(\frac{1}{4} + d\right)u \frac{\partial}{\partial u} \\ v_4 &= \sqrt{x} \exp\left(\frac{b}{2}t\right) \frac{\partial}{\partial x} + 2b\sqrt{x} \exp\left(\frac{b}{2}t\right)u \frac{\partial}{\partial u}, \quad v_5 = \sqrt{x} \exp\left(-\frac{b}{2}t\right) \frac{\partial}{\partial x}, \quad v_6 = u \frac{\partial}{\partial u}. \end{aligned}$$

Table 11: Commutator table($b = e$, $b \neq 0$, $a = \frac{1}{4}$, $d \neq \frac{1}{4}$).

$[v_i, v_j]$	v_1	v_2	v_3	v_4	v_5	v_6	v_μ
v_1	0	$-2bv_3$	$-bv_1$	0	$-bv_4$	0	v_{μ_1}
v_2	$2bv_3$	0	bv_2	bv_5	0	0	v_{μ_2}
v_3	bv_1	$-bv_2$	0	$\frac{b}{2}v_4$	$-\frac{b}{2}v_5$	0	v_{μ_3}
v_4	0	$-bv_5$	$-\frac{b}{2}v_4$	0	$-bv_6$	0	v_{μ_4}
v_5	bv_4	0	$\frac{b}{2}v_5$	bv_6	0	0	v_{μ_5}
v_6	0	0	0	0	0	0	$-v_\mu$
v_μ	$-v_{\mu_1}$	$-v_{\mu_2}$	$-v_{\mu_3}$	$-v_{\mu_4}$	$-v_{\mu_5}$	v_μ	0

$$\begin{aligned} \mu_1 &= b \exp(bt)x\mu_x + b \exp(bt)y\mu_y + \exp(bt)\mu_t + 2b \exp(bt)\left(d + \frac{1}{4} - b(x+y)\right)\mu, \\ \mu_2 &= -b \exp(-bt)x\mu_x - b \exp(-bt)y\mu_y + \exp(-bt)\mu_t, \quad \mu_3 = b(a+d)\mu + \mu_t, \\ \mu_4 &= \sqrt{x} \exp\left(\frac{b}{2}t\right)\mu_x - 2b \exp\left(\frac{b}{2}t\right)\sqrt{x}\mu, \quad \mu_5 = \sqrt{x} \exp\left(-\frac{b}{2}t\right)\mu_x. \end{aligned}$$

The transformed point by a one-parameter group $\exp(\varepsilon v_i)(x, y, t, u) = (\tilde{x}, \tilde{y}, \tilde{t}, \tilde{u})$ are:

$$\begin{aligned}
G_1 &: \left(\frac{x}{1 - b\varepsilon \exp(bt)}, \frac{y}{1 - b\varepsilon \exp(bt)}, t - \frac{\ln(1 - b\varepsilon \exp(bt))}{b}, (1 - b\varepsilon \exp(bt))^{(2d+\frac{1}{2})} u \exp\left(\frac{2b^2\varepsilon \exp(bt)(x+y)}{1 - b\varepsilon \exp(bt)}\right) \right), \\
G_2 &: \left(\frac{x}{1 + b\varepsilon \exp(-bt)}, \frac{y}{1 + b\varepsilon \exp(-bt)}, t + \frac{\ln(1 + b\varepsilon \exp(-bt))}{b}, u \right), \\
G_3 &: \left(x, y, t + \varepsilon, \exp(-\varepsilon b(\frac{1}{4} + d))u \right), \\
G_4 &: \left(\frac{(\varepsilon \exp(\frac{bt}{2}) + 2\sqrt{x})^2}{4}, y, t, u \exp\left(\frac{\exp(\frac{bt}{2})b\varepsilon(\varepsilon \exp(\frac{bt}{2}) + 4\sqrt{x})}{2}\right) \right), \\
G_5 &: \left(\frac{(2 \exp(\frac{bt}{2})\sqrt{x} + \varepsilon)^2 \exp(-bt)}{4}, y, t, u \right) \\
G_6 &: (x, y, t, \exp(\varepsilon)u)
\end{aligned}$$

Since each one parameter group G_i is a symmetry group, using Theorem 2.5 if $u(x, y, t)$ is a solution of (3.1), so are the u^{ε, v_i} , $1 \leq i \leq 6$ given by:

$$\begin{aligned}
u^{\varepsilon, v_1}(x, y, t) &= \frac{\exp\left(\frac{2b^2\varepsilon \exp(bt)(x+y)}{1+b\varepsilon \exp(bt)}\right)}{(1 + b\varepsilon \exp(bt))^{2(\frac{1}{4}+d)}} u\left(\frac{x}{1 + \varepsilon b \exp(bt)}, \frac{y}{1 + \varepsilon b \exp(bt)}, t - \frac{\ln(1 + b\varepsilon \exp(bt))}{b}\right), \\
u^{\varepsilon, v_2}(x, y, t) &= u\left(\frac{x}{1 - \varepsilon b \exp(-bt)}, \frac{y}{1 - \varepsilon b \exp(-bt)}, t + \frac{\ln(1 - \exp(-bt)b\varepsilon)}{b}\right), \\
u^{\varepsilon, v_3}(x, y, t) &= \exp\left(-\varepsilon b\left(\frac{1}{4} + d\right)\right) u(x, y, t - \varepsilon), \\
u^{\varepsilon, v_4}(x, y, t) &= \exp\left(\frac{b\varepsilon \exp(\frac{bt}{2})(4\sqrt{x} - \varepsilon \exp(\frac{bt}{2}))}{2}\right) u\left(\frac{(2\sqrt{x} - \varepsilon \exp(\frac{bt}{2}))^2}{4}, y, t\right), \\
u^{\varepsilon, v_5}(x, y, t) &= u\left(\frac{(2 \exp(\frac{bt}{2})\sqrt{x} - \varepsilon)^2 \exp(-bt)}{4}, y, t\right), \\
u^{\varepsilon, v_6}(x, y, t) &= \exp(\varepsilon)u(x, y, t).
\end{aligned}$$

4.3.3 Subcase 3.3 ($b = -e$, $b \neq 0$, $a = \frac{1}{4}$, $d \neq \frac{1}{4}$)

$$\begin{aligned}
v_1 &= b \exp(bt)x \frac{\partial}{\partial x} + b \exp(bt)y \frac{\partial}{\partial y} + \exp(bt) \frac{\partial}{\partial t} + \frac{b(4bx - 1)}{2} \exp(bt)c_1 u \frac{\partial}{\partial u}, \\
v_2 &= -b \exp(-bt)x \frac{\partial}{\partial x} - b \exp(-bt)y \frac{\partial}{\partial y} + \exp(-bt) \frac{\partial}{\partial t} + 2b(by + d) \exp(-bt)u \frac{\partial}{\partial u}, \\
v_3 &= \frac{\partial}{\partial t} - b\left(\frac{1}{4} - d\right)u \frac{\partial}{\partial u}, \quad v_4 = \sqrt{x} \exp\left(\frac{b}{2}t\right) \frac{\partial}{\partial x} + 2b\sqrt{x} \exp\left(\frac{b}{2}t\right)u \frac{\partial}{\partial u}, \quad v_5 = \sqrt{x} \exp\left(-\frac{b}{2}t\right) \frac{\partial}{\partial x}, \quad v_6 = u \frac{\partial}{\partial u}.
\end{aligned}$$

Table 12: Commutator table($b = -e$, $b \neq 0$, $a = \frac{1}{4}$, $d \neq \frac{1}{4}$).

$[v_i, v_j]$	v_1	v_2	v_3	v_4	v_5	v_6	v_μ
v_1	0	$-2bv_3$	$-bv_1$	0	$-bv_4$	0	v_{μ_1}
v_2	$2bv_3$	0	bv_2	bv_5	0	0	v_{μ_2}
v_3	bv_1	$-bv_2$	0	$\frac{b}{2}v_4$	$-\frac{b}{2}v_5$	0	v_{μ_3}
v_4	0	$-bv_5$	$-\frac{b}{2}v_4$	0	$-bv_6$	0	v_{μ_4}
v_5	bv_4	0	$\frac{b}{2}v_5$	bv_6	0	0	v_{μ_5}
v_6	0	0	0	0	0	0	$-v_\mu$
v_μ	$-v_{\mu_1}$	$-v_{\mu_2}$	$-v_{\mu_3}$	$-v_{\mu_4}$	$-v_{\mu_5}$	v_μ	0

$$\mu_1 = b \exp(bt)x\mu_x + b \exp(bt)y\mu_y + \exp(bt)\mu_t + 2b\left(\frac{1}{4} - bx\right) \exp(bt)\mu,$$

$$\mu_2 = -b \exp(-bt)x\mu_x - b \exp(-bt)y\mu_y + \exp(-bt)\mu_t - 2b(d + by) \exp(-bt)\mu,$$

$$\mu_3 = b\left(\frac{1}{4} - d\right)\mu + \mu_t, \quad \mu_4 = \exp\left(\frac{bt}{2}\right)\sqrt{x}\mu_x - 2b \exp\left(\frac{bt}{2}\right)\sqrt{x}\mu, \quad \mu_5 = \exp\left(-\frac{bt}{2}\right)\sqrt{x}\mu_x.$$

The transformed point by a one-parameter group $\exp(\varepsilon v_i)(x, y, t, u) = (\tilde{x}, \tilde{y}, \tilde{t}, \tilde{u})$ are:

$$G_1 : \left(\frac{x}{1 - b\varepsilon \exp(bt)}, \frac{y}{1 - b\varepsilon \exp(bt)}, t - \frac{\ln(1 - b\varepsilon \exp(bt))}{b}, \sqrt{1 - b\varepsilon \exp(bt)}u \exp\left(\frac{2b^2\varepsilon \exp(bt)x}{1 - b\varepsilon \exp(bt)}\right) \right),$$

$$G_2 : \left(\frac{x}{1 + b\varepsilon \exp(-bt)}, \frac{y}{1 + b\varepsilon \exp(-bt)}, t + \frac{\ln(1 - b\varepsilon \exp(-bt))}{b}, (b\varepsilon + \exp(bt))^{2d}u \exp\left(\frac{-2b(dt \exp(bt) + b\varepsilon(dt - y))}{b\varepsilon + \exp(bt)}\right) \right)$$

$$G_3 : \left(x, y, t + \varepsilon, -b\left(\frac{1}{4} - d\right)\varepsilon + u(x, y, t) \right),$$

$$G_4 : \left(\frac{(\varepsilon \exp(\frac{bt}{2}) + 2\sqrt{x})^2}{4}, y, t, u \exp\left(\frac{\exp(\frac{bt}{2})b\varepsilon(\varepsilon \exp(\frac{bt}{2}) + 4\sqrt{x})}{2}\right) \right),$$

$$G_5 : \left(\frac{(2 \exp(\frac{bt}{2})\sqrt{x} + \varepsilon)^2 \exp(-bt)}{4}, y, t, u \right),$$

$$G_6 : (x, y, t, u \exp(\varepsilon)).$$

Since each one parameter group G_i is a symmetry group, using Theorem 2.5 if $u(x, y, t)$ is a solution of (3.1), so are the u^{ε, v_i} , $1 \leq i \leq 6$ given by:

$$\begin{aligned}
u^{\varepsilon, v_1}(x, y, t) &= \frac{\exp\left(\frac{2b^2\varepsilon x \exp(bt)}{1+b\varepsilon \exp(bt)}\right)}{\sqrt{1+b\varepsilon \exp(bt)}} u\left(\frac{x}{1+b\varepsilon \exp(bt)}, \frac{y}{1+b\varepsilon \exp(bt)}, t - \frac{\ln(1+\varepsilon b \exp(bt))}{b}\right), \\
u^{\varepsilon, v_2}(x, y, t) &= \frac{\exp\left(-\frac{b\varepsilon \exp(-bt)y}{1-b\varepsilon \exp(-bt)}\right)}{(1-b\varepsilon \exp(-bt))^2} u\left(\frac{x}{(1-\varepsilon b \exp(-bt))}, \frac{y}{(1-\varepsilon b \exp(-bt))}, t + \frac{\ln(1+b\varepsilon \exp(-bt))}{b}\right), \\
u^{\varepsilon, v_3}(x, y, t) &= \exp\left(-\varepsilon b\left(\frac{1}{4} - d\right)\right) u(x, y, t - \varepsilon), \\
u^{\varepsilon, v_4}(x, y, t) &= \exp\left(\frac{b\varepsilon \exp(\frac{bt}{2})(4\sqrt{x} - \varepsilon \exp(\frac{bt}{2}))}{2}\right) u\left(\frac{(2\sqrt{x} - \varepsilon \exp(\frac{bt}{2}))^2}{4}, y, t\right), \\
u^{\varepsilon, v_5}(x, y, t) &= u\left(\frac{(2 \exp(\frac{bt}{2})\sqrt{x} - \varepsilon)^2}{4} \exp(-bt), y, t\right), \\
u^{\varepsilon, v_6}(x, y, t) &= \exp(\varepsilon)u(x, y, t).
\end{aligned}$$

4.3.4 Subcase 3.4 ($b = e = 0, a = \frac{1}{4}, d \neq \frac{1}{4}$)

$$\begin{aligned}
v_1 &= 2tx \frac{\partial}{\partial x} + 2ty \frac{\partial}{\partial y} + t^2 \frac{\partial}{\partial t} + (2(x+y) - 2(\frac{1}{4} + d)t)u \frac{\partial}{\partial u}, \\
v_2 &= x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} + t \frac{\partial}{\partial t}, \quad v_3 = \frac{\partial}{\partial t}, \\
v_4 &= \sqrt{x}t \frac{\partial}{\partial x} + 2\sqrt{x}u \frac{\partial}{\partial u}, \quad v_5 = \sqrt{x} \frac{\partial}{\partial x}, \\
v_6 &= u \frac{\partial}{\partial u}.
\end{aligned}$$

Table 13: Commutator table ($b = e = 0, b \neq 0, a = \frac{1}{4}, d \neq \frac{1}{4}$).

$[v_i, v_j]$	v_1	v_2	v_3	v_4	v_5	v_6	v_μ
v_1	0	$-v_1$	$-2v_2 + 2(d + \frac{1}{4})v_6$	0	$-v_4$	0	v_{μ_1}
v_2	v_1	0	$-v_3$	$\frac{1}{2}v_4$	$-\frac{1}{2}v_5$	0	v_{μ_2}
v_3	$2v_2 - 2(d + \frac{1}{4})v_6$	v_3	0	v_5	0	0	v_{μ_3}
v_4	0	$-\frac{1}{2}v_4$	$-v_5$	0	$-v_6$	0	v_{μ_4}
v_5	v_4	$\frac{1}{2}v_5$	0	v_6	0	0	v_{μ_5}
v_6	0	0	0	0	0	0	$-v_\mu$
v_μ	$-v_{\mu_1}$	$-v_{\mu_2}$	$-v_{\mu_3}$	$-v_{\mu_4}$	$-v_{\mu_5}$	v_μ	0

$$\mu_1 = 2tx\mu_x + 2ty\mu_y + t^2\mu_t + (2t(\frac{1}{4} + d) - 2(x+y))\mu,$$

$$\mu_2 = x\mu_x + y\mu_y + t\mu_t, \quad \mu_3 = \mu_t, \quad \mu_4 = \sqrt{x}t\mu_x - 2\sqrt{x}\mu, \quad \mu_5 = \sqrt{x}\mu_x.$$

The transformed point by a one-parameter group $\exp(\varepsilon v_i)(x, y, t, u) = (\tilde{x}, \tilde{y}, \tilde{t}, \tilde{u})$ are:

$$\begin{aligned}
G_1 &: \left(\frac{x}{(1-t\varepsilon)^2}, \frac{y}{(1-t\varepsilon)^2}, \frac{t}{1-t\varepsilon}, u \exp\left(\frac{2\varepsilon(x+y)}{1-t\varepsilon}\right) (1-t\varepsilon)^{\frac{1}{4}+d} \right), \\
G_2 &: (\exp(\varepsilon)x, \exp(\varepsilon)y, \exp(\varepsilon)t, u), \\
G_3 &: (x, y, t + \varepsilon, u), \\
G_4 &: \left(x + t\sqrt{x}\varepsilon + \frac{t^2\varepsilon^2}{4}, y, t, u \exp\left(\frac{\varepsilon(t\varepsilon + 4\sqrt{x})}{2}\right) \right), \\
G_5 &: \left(\sqrt{x}\varepsilon + \frac{\varepsilon^2}{4} + x, y, t, u \right), \\
G_6 &: (x, y, t, \exp(\varepsilon)u).
\end{aligned}$$

Since each one parameter group G_i is a symmetry group, using Theorem 2.5 if $u(x, y, t)$ is a solution of (3.1), so are the u^{ε, v_i} , $1 \leq i \leq 6$ given by:

$$\begin{aligned}
u^{\varepsilon, v_1}(x, y, t) &= \frac{\exp\left(\frac{2\varepsilon(x+y)}{(1+t\varepsilon)^3}\right)}{(1+t\varepsilon)^{\frac{1}{4}+d}} u\left(\frac{x}{(1+t\varepsilon)^2}, \frac{y}{(1+t\varepsilon)^2}, \frac{t}{1+t\varepsilon}\right), \\
u^{\varepsilon, v_2}(x, y, t) &= u(\exp(-\varepsilon)x, \exp(-\varepsilon)y, \exp(-\varepsilon)t), \\
u^{\varepsilon, v_3}(x, y, t) &= u(x, y, t - \varepsilon), \\
u^{\varepsilon, v_4}(x, y, t) &= \exp\left(\frac{\varepsilon\left(\varepsilon t + 4\sqrt{x - t\sqrt{x}\varepsilon + \frac{t^2\varepsilon^2}{2}}\right)}{2}\right) u\left(x - t\sqrt{x}\varepsilon + \frac{t^2\varepsilon^2}{2}, y, t\right), \\
u^{\varepsilon, v_5}(x, y, t) &= u\left(\sqrt{x}\varepsilon - \frac{\varepsilon^2}{4} + x, y, t\right), \\
u^{\varepsilon, v_6}(x, y, t) &= \exp(\varepsilon)u(x, y, t).
\end{aligned}$$

4.4 Case 4 ($a \neq \frac{1}{4}$, $d = \frac{1}{4}$)

This case is symmetric to the last case (case 3), because $a \neq \frac{1}{4}$ and from remark (3.35), using (3.39e) we find out that $h = l = 0$ and $k_{tt} - \frac{e^2}{4}k = 0$ in all the following subcases.

4.4.1 Subcase 4,1 ($b \neq \pm e$, $a \neq \frac{1}{4}$, $d = \frac{1}{4}$)

$$\begin{aligned}
v_1 &= \frac{\partial}{\partial t}, \quad v_2 = u \frac{\partial}{\partial u}, \\
v_3 &= \sqrt{y} \exp\left(\frac{e}{2}t\right) \frac{\partial}{\partial y} + 2e \exp\left(\frac{e}{2}t\right) \sqrt{y} u \frac{\partial}{\partial u}, \\
v_4 &= \sqrt{y} \exp\left(-\frac{e}{2}t\right) \frac{\partial}{\partial y}.
\end{aligned}$$

Table 14: Commutator table($b \neq \pm e$, $a \neq \frac{1}{4}$, $d = \frac{1}{4}$).

$[v_i, v_j]$	v_1	v_2	v_3	v_4	v_μ
v_1	0	0	$\frac{\epsilon}{2}v_3$	$-\frac{\epsilon}{2}v_4$	v_{μ_t}
v_2	0	0	0	0	$-v_\mu$
v_3	$-\frac{\epsilon}{2}v_3$	0	0	$-ev_2$	v_{μ_1}
v_4	$\frac{\epsilon}{2}v_4$	0	ev_2	0	v_{μ^2}
v_μ	$-v_{\mu_t}$	v_μ	$-v_{\mu_1}$	$-v_{\mu_2}$	0

$$\mu_1 = -2 \exp\left(\frac{\epsilon}{2}t\right)\sqrt{y}e\mu + \exp\left(\frac{\epsilon}{2}t\right)\sqrt{y}\mu_y, \quad \mu_2 = \exp\left(-\frac{\epsilon}{2}t\right)\sqrt{y}\mu_y.$$

The transformed point by a one-parameter group $\exp(\epsilon v_i)(x, y, t, u) = (\tilde{x}, \tilde{y}, \tilde{t}, \tilde{u})$ are:

$$\begin{aligned} G_1 &: (x, y, t + \epsilon, u), \\ G_2 &: (x, y, t, \exp(\epsilon)u), \\ G_3 &: \left(x, \frac{(\epsilon \exp(\frac{\epsilon t}{2}) + 2\sqrt{y})^2}{4}, t, u \exp\left(\frac{\exp(\frac{\epsilon t}{2})e\epsilon(\epsilon \exp(\frac{\epsilon t}{2}) + 4\sqrt{y})}{2}\right) \right), \\ G_4 &: \left(x, \frac{(2\exp(\frac{\epsilon t}{2})\sqrt{y} + \epsilon)^2 \exp(-\epsilon t)}{4}, t, u \right). \end{aligned}$$

Since each one parameter group G_i is a symmetry group, using Theorem 2.5 if $u(x, y, t)$ is a solution of (3.1), so are the u^{ϵ, v_i} , $1 \leq i \leq 4$ given by:

$$\begin{aligned} u^{\epsilon, v_1}(x, y, t) &= u(x, y, t - \epsilon), \\ u^{\epsilon, v_2}(x, y, t) &= \exp(\epsilon)u(x, y, t), \\ u^{\epsilon, v_3}(x, y, t) &= \exp\left(\frac{\exp(\frac{\epsilon t}{2})e\epsilon(4\sqrt{y} - \epsilon \exp(\frac{\epsilon t}{2}))}{2}\right) u\left(\frac{(2\sqrt{y} - \epsilon \exp(\frac{\epsilon t}{2}))^2}{4}, y, t\right), \\ u^{\epsilon, v_4}(x, y, t) &= u\left(x, \frac{(2\exp(\frac{\epsilon t}{2})\sqrt{y} - \epsilon)^2 \exp(-\epsilon t)}{4}, t\right). \end{aligned}$$

4.4.2 Subcase 4.2 ($b = e$, $b \neq 0$, $a \neq \frac{1}{4}$, $d = \frac{1}{4}$)

$$\begin{aligned} v_1 &= b \exp(bt)x \frac{\partial}{\partial x} + b \exp(bt)y \frac{\partial}{\partial y} + \exp(bt) \frac{\partial}{\partial t} + 2b \exp(bt) \left(b(x+y) - (a + \frac{1}{4}) \right) u \frac{\partial}{\partial u}, \\ v_2 &= -b \exp(-bt)x \frac{\partial}{\partial x} - b \exp(-bt)y \frac{\partial}{\partial y} + \exp(-bt) \frac{\partial}{\partial t}, \quad v_3 = \frac{\partial}{\partial t} - b(\frac{1}{4} + a)u \frac{\partial}{\partial u} \\ v_4 &= \sqrt{y} \exp(\frac{b}{2}t) \frac{\partial}{\partial x} + 2b\sqrt{y} \exp(\frac{b}{2}t)u \frac{\partial}{\partial u}, \quad v_5 = \sqrt{y} \exp(-\frac{b}{2}t) \frac{\partial}{\partial y}, \quad v_6 = u \frac{\partial}{\partial u}. \end{aligned}$$

Table 15: Commutator table($b = e$, $a \neq \frac{1}{4}$, $d = \frac{1}{4}$).

$[v_i, v_j]$	v_1	v_2	v_3	v_4	v_5	v_6	v_μ
v_1	0	$-2bv_3$	$-bv_1$	0	$-bv_4$	0	v_{μ_1}
v_2	$2bv_3$	0	bv_2	bv_5	0	0	v_{μ_2}
v_3	bv_1	$-bv_2$	0	$\frac{b}{2}v_4$	$-\frac{b}{2}v_5$	0	v_{μ_3}
v_4	0	$-bv_5$	$-\frac{b}{2}v_4$	0	$-bv_6$	0	v_{μ_4}
v_5	bv_4	0	$\frac{b}{2}v_5$	bv_6	0	0	v_{μ_5}
v_6	0	0	0	0	0	0	$-v_\mu$
v_μ	$-v_{\mu_1}$	$-v_{\mu_2}$	$-v_{\mu_3}$	$-v_{\mu_4}$	$-v_{\mu_5}$	v_μ	0

$$\begin{aligned} \mu_1 &= b \exp(bt)x\mu_x + b \exp(bt)y\mu_y + \exp(bt)\mu_t + 2b \exp(bt)(d + \frac{1}{4} - b(x+y))\mu, \\ \mu_2 &= -b \exp(-bt)x\mu_x - b \exp(-bt)y\mu_y + \exp(-bt)\mu_t, \\ \mu_3 &= b(a+d)\mu + \mu_t, \quad \mu_4 = \sqrt{y} \exp(\frac{b}{2}t)\mu_y - 2b \exp(\frac{b}{2}t)\sqrt{x}\mu, \quad \mu_5 = \sqrt{y} \exp(-\frac{b}{2}t)\mu_y. \end{aligned}$$

The transformed point by a one-parameter group $\exp(\varepsilon v_i)(x, y, t, u) = (\tilde{x}, \tilde{y}, \tilde{t}, \tilde{u})$ are:

$$G_1 : \left(\frac{x}{1 - b\varepsilon \exp(bt)}, \frac{y}{1 - b\varepsilon \exp(bt)}, t - \frac{\ln(1 - b\varepsilon \exp(bt))}{b}, (1 - b\varepsilon \exp(bt))^{2a + \frac{1}{2}} u \exp\left(\frac{2b^2\varepsilon \exp(bt)(x+y)}{1 - b\varepsilon \exp(bt)}\right) \right),$$

$$G_2 : \left(\frac{x}{1 + b\varepsilon \exp(-bt)}, \frac{y}{1 + b\varepsilon \exp(-bt)}, t + \frac{\ln(1 + b\varepsilon \exp(-bt))}{b}, u \right),$$

$$G_3 : \left(x, y, t + \varepsilon, \exp(-\varepsilon b(\frac{1}{4} + d))u \right),$$

$$G_4 : \left(x, \frac{(\varepsilon \exp(\frac{bt}{2}) + 2\sqrt{y})^2}{4}, t, u \exp\left(\frac{\exp(\frac{bt}{2})b\varepsilon(\varepsilon \exp(\frac{bt}{2}) + 4\sqrt{y})}{2}\right) \right),$$

$$G_5 : \left(x, \frac{(2 \exp(\frac{bt}{2})\sqrt{y} + \varepsilon)^2 \exp(-bt)}{4}, t, u \right)$$

$$G_6 : (x, y, t, \exp(\varepsilon)u)$$

Since each one parameter group G_i is a symmetry group, using Theorem 2.5 if $u(x, y, t)$ is a solution of (3.1), so are the u^{ε, v_i} , $1 \leq i \leq 6$ given by:

$$\begin{aligned}
u^{\varepsilon, v_1}(x, y, t) &= \frac{\exp\left(\frac{2b^2\varepsilon \exp(bt)(x+y)}{1+b\varepsilon \exp(bt)}\right)}{(1+b\varepsilon \exp(bt))^{2(\frac{1}{4}+a)}} u\left(\frac{x}{1+\varepsilon b \exp(bt)}, \frac{y}{1+\varepsilon b \exp(bt)}, t - \frac{\ln(1+b\varepsilon \exp(bt))}{b}\right), \\
u^{\varepsilon, v_2}(x, y, t) &= u\left(\frac{x}{1-\varepsilon b \exp(-bt)}, \frac{y}{1-\varepsilon b \exp(-bt)}, t + \frac{\ln(1-\exp(-bt)b\varepsilon)}{b}\right), \\
u^{\varepsilon, v_3}(x, y, t) &= \exp(-\varepsilon b(\frac{1}{4}+a))f(x, y, t - \varepsilon), \\
u^{\varepsilon, v_4}(x, y, t) &= \exp\left(\frac{b\varepsilon \exp(\frac{bt}{2})(4\sqrt{y} - \varepsilon \exp(\frac{bt}{2}))}{2}\right) u\left(x, \frac{(2\sqrt{y} - \varepsilon \exp(\frac{bt}{2}))^2}{4}, t\right) \\
u^{\varepsilon, v_5}(x, y, t) &= u\left(x, \frac{(2\exp(\frac{bt}{2})\sqrt{y} - \varepsilon)^2}{4} \exp(-bt), t\right), \\
u^{\varepsilon, v_6}(x, y, t) &= \exp(\varepsilon)u(x, y, t).
\end{aligned}$$

4.4.3 Subcase 4.3($b = -e, b \neq 0, a \neq \frac{1}{4}, d = \frac{1}{4}$)

$$\begin{aligned}
v_1 &= b \exp(bt)x \frac{\partial}{\partial x} + b \exp(bt)y \frac{\partial}{\partial y} + \exp(bt) \frac{\partial}{\partial t} - 2b(-bx + a) \exp(bt)u \frac{\partial}{\partial u}, \\
v_2 &= -b \exp(-bt)x \frac{\partial}{\partial x} - b \exp(-bt)y \frac{\partial}{\partial y} + \exp(-bt) \frac{\partial}{\partial t} + \frac{b(4by + 1)}{2} \exp(-bt)u \frac{\partial}{\partial u}, \quad v_3 = \frac{\partial}{\partial t} + b(\frac{1}{4} - a)u \frac{\partial}{\partial u} \\
v_4 &= \sqrt{y} \exp(-\frac{b}{2}t) \frac{\partial}{\partial x} - 2b\sqrt{y} \exp(-\frac{b}{2}t)u \frac{\partial}{\partial u}, \quad v_5 = \sqrt{y} \exp(\frac{b}{2}t) \frac{\partial}{\partial y}, \quad v_6 = u \frac{\partial}{\partial u}.
\end{aligned}$$

Table 16: Commutator table($b = -e, a \neq \frac{1}{4}, d = \frac{1}{4}$).

$[v_i, v_j]$	v_1	v_2	v_3	v_4	v_5	v_6	v_μ
v_1	0	$-2bv_3$	$-bv_1$	$-bv_5$	0	0	v_{μ_1}
v_2	$2bv_3$	0	bv_2	0	bv_4	0	v_{μ_2}
v_3	bv_1	$-bv_2$	0	$-\frac{b}{2}v_4$	$\frac{b}{2}v_5$	0	v_{μ_3}
v_4	bv_5	0	$\frac{b}{2}v_4$	0	bv_6	0	v_{μ_4}
v_5	0	$-bv_4$	$-\frac{b}{2}v_5$	$-bv_6$	0	0	v_{μ_5}
v_6	0	0	0	0	0	0	$-v_\mu$
v_μ	$-v_{\mu_1}$	$-v_{\mu_2}$	$-v_{\mu_3}$	$-v_{\mu_4}$	$-v_{\mu_5}$	v_μ	0

$$\begin{aligned}
\mu_1 &= b \exp(bt)x\mu_x + b \exp(bt)y\mu_y + \exp(bt)\mu_t + 2b(a - bx) \exp(bt)\mu, \\
\mu_2 &= -b \exp(-bt)x\mu_x - b \exp(-bt)y\mu_y + \exp(-bt)\mu_t - 2b(\frac{1}{4} + by) \exp(-bt)\mu, \\
\mu_3 &= -b(\frac{1}{4} - a)\mu + \mu_t, \quad \mu_4 = \exp(-\frac{bt}{2})\sqrt{y}\mu_y + 2b \exp(-\frac{bt}{2})\sqrt{y}\mu, \quad \mu_5 = \exp(\frac{bt}{2})\sqrt{y}\mu_y.
\end{aligned}$$

The transformed point by a one-parameter group $\exp(\varepsilon v_i)(x, y, t, u) = (\tilde{x}, \tilde{y}, \tilde{t}, \tilde{u})$ are:

$$\begin{aligned}
G_1 &: \left(\frac{x}{1 - b\varepsilon \exp(bt)}, \frac{y}{1 - b\varepsilon \exp(bt)}, t - \frac{\ln(1 - b\varepsilon \exp(bt))}{b}, \frac{1}{1 - b\varepsilon \exp(bt)} u \exp\left(\frac{2 \exp(bt) b^2 \varepsilon x}{1 - b\varepsilon \exp(bt)}\right) \right), \\
G_2 &: \left(\frac{x}{1 + b\varepsilon \exp(-bt)}, \frac{y}{1 + b\varepsilon \exp(-bt)}, t + \frac{\ln(1 - b\varepsilon \exp(-bt))}{b}, \sqrt{1 + b\varepsilon \exp(-bt)} u \exp\left(\frac{2b^2 \varepsilon \exp(-bt) y}{1 + b\varepsilon \exp(-bt)}\right) \right), \\
G_3 &: \left(x, y, t + \varepsilon, b \left(\frac{b}{4} - ab \right) \varepsilon + u(x, y, t) \right), \\
G_4 &: \left(x, \frac{(\varepsilon \exp(-\frac{bt}{2}) + 2\sqrt{y})^2}{4}, t, u \exp\left(\frac{-b \exp(-\frac{bt}{2}) \varepsilon (\varepsilon \exp(-\frac{bt}{2}) + 4\sqrt{y})}{2}\right) \right), \\
G_5 &: \left(x, \frac{(2\sqrt{y} + \varepsilon \exp(\frac{bt}{2}))^2}{4}, t, u \right), \\
G_6 &: (x, y, t, u \exp(\varepsilon)).
\end{aligned}$$

Since each one parameter group G_i is a symmetry group, using Theorem 2.5 if $u(x, y, t)$ is a solution of (3.1), so are the u^{ε, v_i} , $1 \leq i \leq 6$ given by:

$$\begin{aligned}
u^{\varepsilon, v_1}(x, y, t) &= (1 + b\varepsilon \exp(bt)) \exp\left(\frac{2b^2 \varepsilon \exp(bt) x}{1 - b\varepsilon \exp(bt)}\right) u\left(\frac{x}{1 + b\varepsilon \exp(bt)}, \frac{y}{1 + b\varepsilon \exp(bt)}, t - \frac{\ln(1 + \varepsilon b \exp(bt))}{b}\right) \\
u^{\varepsilon, v_2}(x, y, t) &= \sqrt{1 + \frac{b\varepsilon \exp(-bt)}{1 + b\varepsilon \exp(-bt)}} \exp\left(\frac{2b^2 \varepsilon \exp(-bt) y}{(1 - b\varepsilon \exp(-bt))(1 + 2b\varepsilon \exp(-bt))}\right) \\
&\quad u\left(\frac{x}{(1 - \varepsilon b \exp(-bt))}, \frac{y}{(1 - \varepsilon b \exp(-bt))}, t + \frac{\ln(1 + b\varepsilon \exp(-bt))}{b}\right), \\
u^{\varepsilon, v_3}(x, y, t) &= \exp\left(\varepsilon b \left(\frac{1}{4} - a\right)\right) u(x, y, t - \varepsilon), \\
u^{\varepsilon, v_4}(x, y, t) &= \exp\left(-\frac{b\varepsilon \exp(-\frac{bt}{2}) (4\sqrt{y} - \varepsilon \exp(-\frac{bt}{2}))}{2}\right) u\left(x, \frac{(2\sqrt{y} - \varepsilon \exp(-\frac{bt}{2}))^2}{4}, t\right), \\
u^{\varepsilon, v_5}(x, y, t) &= u\left(x, \frac{(2\sqrt{y} - \varepsilon \exp(\frac{bt}{2}))^2}{4}, t\right), \\
u^{\varepsilon, v_6}(x, y, t) &= \exp(\varepsilon) u(x, y, t).
\end{aligned}$$

4.4.4 Subcase 4.4($b = e = 0$, $a \neq \frac{1}{4}$, $d = \frac{1}{4}$)

$$\begin{aligned}
v_1 &= 2tx \frac{\partial}{\partial x} + 2ty \frac{\partial}{\partial y} + t^2 \frac{\partial}{\partial t} + (2(x + y) - 2(\frac{1}{4} + d)t) u \frac{\partial}{\partial u}, \quad v_3 = \frac{\partial}{\partial t}, \\
v_2 &= x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} + t \frac{\partial}{\partial t}, \quad v_4 = \sqrt{y} t \frac{\partial}{\partial x} + 2\sqrt{y} u \frac{\partial}{\partial u}, \quad v_5 = \sqrt{y} \frac{\partial}{\partial y}, \quad v_6 = u \frac{\partial}{\partial u}.
\end{aligned}$$

Table 17: Commutator table($b = e = 0, a \neq \frac{1}{4}, d = \frac{1}{4}$).

$[v_i, v_j]$	v_1	v_2	v_3	v_4	v_5	v_6	v_μ
v_1	0	$-v_1$	$-2v_2 + 2(d + \frac{1}{4})v_6$	0	$-v_4$	0	v_{μ_1}
v_2	v_1	0	$-v_3$	$\frac{1}{2}v_4$	$-\frac{1}{2}v_5$	0	v_{μ_2}
v_3	$2v_2 - 2(d + \frac{1}{4})v_6$	$-bv_2$	0	v_5	0	0	v_{μ_3}
v_4	0	$-\frac{1}{2}v_4$	$-v_5$	0	$-v_6$	0	v_{μ_4}
v_5	v_4	$\frac{1}{2}v_5$	0	$-v_6$	0	0	v_{μ_5}
v_6	0	0	0	0	0	0	$-v_\mu$
v_μ	$-v_{\mu_1}$	$-v_{\mu_2}$	$-v_{\mu_3}$	$-v_{\mu_4}$	$-v_{\mu_5}$	v_μ	0

$$\mu_1 = 2tx\mu_x + 2ty\mu_y + t^2\mu_t + (2t(\frac{1}{4} + d) - 2(x + y))\mu,$$

$$\mu_2 = x\mu_x + y\mu_y + t\mu_t, \quad \mu_3 = \mu_t, \quad \mu_4 = \sqrt{y}t\mu_y - 2\sqrt{y}\mu, \quad \mu_5 = \sqrt{y}\mu_y.$$

The transformed point by a one-parameter group $\exp(\varepsilon v_i)(x, y, t, u) = (\tilde{x}, \tilde{y}, \tilde{t}, \tilde{u})$ are:

$$G_1 : \left(\frac{x}{(1-t\varepsilon)^2}, \frac{y}{(1-t\varepsilon)^2}, \frac{t}{1-t\varepsilon}, u \exp\left(\frac{2\varepsilon(x+y)}{1-t\varepsilon}\right) (1-t\varepsilon)^{a+\frac{1}{4}} \right),$$

$$G_2 : (\exp(\varepsilon)x, \exp(\varepsilon)y, \exp(\varepsilon)t, u),$$

$$G_3 : (x, y, t + \varepsilon, u),$$

$$G_4 : \left(x, y + t\sqrt{y}\varepsilon + \frac{t^2\varepsilon^2}{4}, t, u \exp\left(\frac{\varepsilon(t\varepsilon + 4\sqrt{y})}{2}\right) \right),$$

$$G_5 : \left(x, \sqrt{y}\varepsilon + \frac{\varepsilon^2}{4} + y, t, u \right),$$

$$G_6 : (x, y, t, \exp(\varepsilon)u).$$

Since each one parameter group G_i is a symmetry group, using Theorem 2.5 if $u(x, y, t)$ is a solution of (3.1), so are the $u^{\varepsilon, v_i}, 1 \leq i \leq 6$ given by:

$$u^{\varepsilon, v_1}(x, y, t) = \frac{\exp\left(\frac{2\varepsilon(x+y)}{(1+t\varepsilon)^3}\right)}{(1+t\varepsilon)^{a+\frac{1}{4}}} u\left(\frac{x}{(1+t\varepsilon)^2}, \frac{y}{(1+t\varepsilon)^2}, \frac{t}{1+t\varepsilon}\right),$$

$$u^{\varepsilon, v_2}(x, y, t) = u(\exp(-\varepsilon)x, \exp(-\varepsilon)y, \exp(-\varepsilon)t),$$

$$u^{\varepsilon, v_3}(x, y, t) = u(x, y, t - \varepsilon),$$

$$u^{\varepsilon, v_4}(x, y, t) = \exp\left(\frac{\varepsilon\left(\varepsilon t + 4\sqrt{y - t\sqrt{y}\varepsilon + \frac{t^2\varepsilon^2}{2}}\right)}{2}\right) u\left(x, y - t\sqrt{y}\varepsilon + \frac{t^2\varepsilon^2}{2}, t\right),$$

$$u^{\varepsilon, v_5}(x, y, t) = u\left(x, -\sqrt{y}\varepsilon + \frac{\varepsilon^2}{4} + y, t\right),$$

$$u^{\varepsilon, v_6}(x, y, t) = \exp(\varepsilon)u(x, y, t).$$

5 The structure of the Lie algebras

Recall $\mathcal{V}_{a,b,d,e}$ defined in Remark 4.2, the algebra generated by v_1, \dots, v_r and v_μ , where $r \in \{2, 4, 6, 9\}$. Similarly, $\mathcal{L}_{a,b,d,e}$ represent the finite-dimensional algebra generated by v_1, \dots, v_r $r \in \{2, 4, 6, 9\}$. Additionally, \mathcal{H} is the infinite-dimensional ideal of $\mathcal{V}_{a,b,d,e}$ consisting of the v_μ where μ is a solution of (3.1).

$[\mathcal{V}_{a,b,d,e}, \mathcal{H}] \subset \mathcal{H}$ therefore $\mathcal{V}_{a,b,d,e} = \mathcal{L}_{a,b,d,e} \ltimes \mathcal{H}$.

Now, we proceed to analyze the structure of $\mathcal{L}_{a,b,d,e}$ in each case.

5.1 Case 1 ($a = \frac{1}{4}, d = \frac{1}{4}$)

5.1.1 Subcase 1.1 ($b \neq \pm e, a = d = \frac{1}{4}$)

We have $\mathcal{Z}(\mathcal{L}_{a,b,d,e}) = \langle v_2 \rangle$ thus $\mathcal{L}_{a,b,d,e}/\mathcal{Z}(\mathcal{L}_{a,b,d,e}) \cong \mathbb{R}v_1 \ltimes \mathcal{A}$ where $\mathcal{A} = \langle \bar{v}_3, \bar{v}_4, \bar{v}_5, \bar{v}_6 \rangle$ is an abelian Lie algebra of dimension 4.

■ if $b = 0, e \neq 0$:(4.1.1)

From the commutator table, the center of $\mathcal{L}_{a,b,d,e}$ is $\mathcal{Z}(\mathcal{L}_{a,b,d,e}) = \langle v_2, v_3, v_5 \rangle$, which is an abelian Lie algebra of dimension 3. Then

$$\frac{\mathcal{L}_{a,b,d,e}}{\mathcal{Z}(\mathcal{L}_{a,b,d,e})} \cong \frac{\frac{\mathcal{L}_{a,b,d,e}}{\mathbb{R}v_2}}{\frac{\mathcal{Z}(\mathcal{L}_{a,b,d,e})}{\mathbb{R}v_2}} = \frac{\mathbb{R}v_1 \ltimes \langle \bar{v}_3, \bar{v}_4, \bar{v}_5, \bar{v}_6 \rangle}{\langle \bar{v}_3, \bar{v}_5 \rangle} \cong \mathbb{R}\bar{v}_1 \ltimes \langle \bar{v}_4, \bar{v}_6 \rangle \cong L(3, 2, x = -1) \cong \mathfrak{iso}_2$$

which is the Poincaré algebra, using the isomorphism defined below

$$\phi : \begin{cases} e_1 \longrightarrow \bar{v}_4, \\ e_2 \longrightarrow \bar{v}_6, \\ e_3 \longrightarrow \frac{2}{e}\bar{v}_1. \end{cases}$$

where $\{e_1, e_2, e_3\}$ are the basis of \mathfrak{iso}_2 , using the notation of [2, p. 16].

■ if $b \neq 0, e = 0$:(4.1.1)

This case is symmetric to the last one. We have that, $\mathcal{Z}(\mathcal{L}_{a,b,d,e}) = \langle \bar{v}_2, \bar{v}_4, \bar{v}_6 \rangle$, then

$$\frac{\mathcal{L}_{a,b,d,e}}{\mathcal{Z}(\mathcal{L}_{a,b,d,e})} \cong \frac{\frac{\mathcal{L}_{a,b,d,e}}{\mathbb{R}v_2}}{\frac{\mathcal{Z}(\mathcal{L}_{a,b,d,e})}{\mathbb{R}v_2}} = \frac{\mathbb{R}v_1 \ltimes \langle \bar{v}_3, \bar{v}_4, \bar{v}_5, \bar{v}_6 \rangle}{\langle \bar{v}_4, \bar{v}_6 \rangle} \cong \mathbb{R}\bar{v}_1 \ltimes \langle \bar{v}_3, \bar{v}_5 \rangle \cong L(3, 2, x = -1) \cong \mathfrak{iso}_2$$

using the isomorphism defined below

$$\psi : \begin{cases} e_1 \longrightarrow \bar{v}_5, \\ e_2 \longrightarrow \bar{v}_6, \\ e_3 \longrightarrow \frac{2}{b}\bar{v}_1. \end{cases}$$

where $\{e_1, e_2, e_3\}$ are the basis of \mathfrak{iso}_2 , using the notation of [2, p. 16].

5.1.2 Subcase 1.2 ($b = e, b \neq 0, a = \frac{1}{4}, d = \frac{1}{4}$)

We have $J = \langle v_4, v_5, v_6, v_7, v_8, v_9 \rangle$ is an ideal of $\mathcal{L}_{a,b,d,e}$ of dimension 6, then $\mathcal{L}_{a,b,d,e}/J = \langle \bar{v}_1, \bar{v}_2, \bar{v}_3 \rangle \cong \mathfrak{sl}_2$ via the isomorphism

$$\begin{cases} \bar{v}_1 \longrightarrow -bE, \\ \bar{v}_2 \longrightarrow bF, \\ \bar{v}_3 \longrightarrow \frac{b}{2}H, \end{cases} \quad (5.1a)$$

where $\{E, F, H\}$ is the canonical basis of \mathfrak{sl}_2 , then:

$$\mathcal{L}_{a,b,d,e} \cong \mathfrak{sl}_2 \ltimes J.$$

5.1.3 Subcase 1.3 ($b = -e, b \neq 0, a = \frac{1}{4}, d = \frac{1}{4}$)

Similar to the last case,

$$\mathcal{L}_{a,b,d,e} \cong \mathfrak{sl}_2 \times J$$

5.1.4 Subcase 1.4 ($b = e = 0, a = \frac{1}{4}, d = \frac{1}{4}$)

We have that $J = \langle \bar{v}_4, \bar{v}_5, \bar{v}_6, \bar{v}_7, \bar{v}_8, \bar{v}_9 \rangle$ is an ideal of $\mathcal{L}_{a,b,d,e}$ of dimension 6, and, $\mathcal{L}_{a,b,d,e}/J \cong \mathfrak{sl}_2$ via the isomorphism defined as follows:

$$\begin{cases} \bar{v}_1 \longrightarrow E, \\ \bar{v}_2 \longrightarrow \frac{H}{2}, \\ \bar{v}_3 \longrightarrow -F, \end{cases} \quad (5.2a)$$

Thus,

$$\mathcal{L}_{a,b,d,e} \cong \mathfrak{sl}_2 \times J.$$

5.2 Case 2 ($a \neq \frac{1}{4}, d \neq \frac{1}{4}$)

5.2.1 Subcase 2.1 ($b \neq \pm e, a \neq \frac{1}{4}, d \neq \frac{1}{4}$)

From the commutator table, we deduce that $\mathcal{L}_{a,b,d,e} = \langle v_1, v_2 \rangle$ is an abelian Lie algebra of dimension 2.

5.2.2 Subcase 2.2 ($b = e, b \neq 0, a \neq \frac{1}{4}, d \neq \frac{1}{4}$)

We have $I = \langle v_1, v_2, v_3 \rangle \cong \mathfrak{sl}_2$ via the isomorphism (5.3b), and $\mathcal{Z}(\mathcal{L}_{a,b,d,e}) = \langle v_4 \rangle$. Thus:

$$\mathcal{L}_{a,b,d,e} \cong \mathbb{R}v_4 \times \mathfrak{sl}_2 \cong \mathcal{N}.$$

5.2.3 Subcase 2.3 ($b = -e, b \neq 0, a \neq \frac{1}{4}, d \neq \frac{1}{4}$)

Same as last case:

$$\mathcal{L}_{a,b,d,e} \cong \mathbb{R}v_4 \times \mathfrak{sl}_2 \cong \mathcal{N}.$$

5.2.4 Subcase 2.4 ($b = e = 0, a \neq \frac{1}{4}, d \neq \frac{1}{4}$)

Again we have that:

$$\mathcal{L}_{a,b,d,e} \cong \mathbb{R}v_4 \times \mathfrak{sl}_2 \cong \mathcal{N}.$$

5.3 Case 3 ($a = \frac{1}{4}, d \neq \frac{1}{4}$)

5.3.1 Subcase 3.1 ($b \neq \pm e, a = \frac{1}{4}, d \neq \frac{1}{4}$)

Suppose that $b \neq 0$, because the case $b = 0$ $\mathcal{L}_{a,b,d,e}$ is an abelian Lie algebra of dimension 4. From the commutator table $\mathcal{Z}(\mathcal{L}_{a,b,d,e}) = \mathbb{R}v_2$. Thus:

$$\mathcal{L}_{a,b,d,e}/\mathbb{R}v_2 \cong L(3, 2, x = -1) \cong \mathfrak{iso}_2,$$

using the notations of [2, p. 16].

5.3.2 Subcase 3.2 ($b = e, b \neq 0, a = \frac{1}{4}, d \neq \frac{1}{4}$)

Using the table in paragraph 4.3.2, one sees that the linear mapping from $\mathcal{L}_{a,b,d,e}$ to \mathcal{M} defined by:

$$\begin{cases} v_1 \longrightarrow E, & (5.3a) \\ v_2 \longrightarrow -b^2 F & (5.3b) \\ v_3 \longrightarrow \frac{b}{2} H, & (5.3c) \\ v_4 \longrightarrow X, & (5.3d) \\ v_5 \longrightarrow -bY, & (5.3e) \\ v_6 \longrightarrow Z. & (5.3f) \end{cases}$$

It is a morphism of Lie algebra. Therefore, $\mathcal{L}_{a,b,d,e}$ is isomorphic to \mathcal{M} , Hence to $\mathcal{H}_{0,0}$, used in [11].

5.3.3 Subcase 3.3 ($b = -e, b \neq 0, a = \frac{1}{4}d \neq \frac{1}{4}$)

Similar to the previous case:

$$\mathcal{L}_{a,b,d,e} \cong \mathcal{M} \cong \mathcal{H}_{0,0}.$$

5.3.4 Subcase 3.4 ($b = e = 0, a = \frac{1}{4}, d \neq \frac{1}{4}$)

Again, in this case:

$$\mathcal{L}_{a,b,d,e} \cong \mathcal{M} \cong \mathcal{H}_{0,0}.$$

5.4 Case 4 ($a \neq \frac{1}{4}, d = \frac{1}{4}$)

This case is symmetric to case 3.

5.4.1 Subcase 4.1 ($b \neq \pm e, a \neq \frac{1}{4}, d = \frac{1}{4}$)

We assume $e \neq 0$, because the case $e = 0$ makes $\mathcal{L}_{a,b,d,e}$ an abelian algebra dimension 4. From the commutator table,

$$\mathcal{L}_{a,b,d,e}/\mathbb{R}v_2 \cong \mathfrak{iso}_2$$

5.4.2 Subcase 4.2 ($b = e, b \neq 0, a \neq \frac{1}{4}, d = \frac{1}{4}$)

$$\mathcal{L}_{a,b,d,e} \cong \mathcal{M} \cong \mathcal{H}_{0,0}.$$

5.4.3 Subcase 4.3 ($b = -e, b \neq 0, a \neq \frac{1}{4}, d = \frac{1}{4}$)

$$\mathcal{L}_{a,b,d,e} \cong \mathcal{M} \cong \mathcal{H}_{0,0}.$$

5.4.4 Subcase 4.4 ($b = e = 0, a \neq \frac{1}{4}, d = \frac{1}{4}$)

$$\mathcal{L}_{a,b,d,e} \cong \mathcal{M} \cong \mathcal{H}_{0,0}.$$

6 Isomorphism between Lie algebras.

In each case we designate the respective Lie algebras as follows:

$$\begin{aligned} \mathcal{L}_0^0 : b \neq \pm e \text{ and } a \neq \frac{1}{4}d \neq \frac{1}{4}, & \quad \mathcal{L}_0^1 : b = e \text{ and } a \neq \frac{1}{4}d \neq \frac{1}{4}, & \quad \mathcal{L}_0^2 : b = -e \text{ and } a \neq \frac{1}{4}d \neq \frac{1}{4}, & \quad \mathcal{L}_0^3 : b = e = 0 \text{ and } a \neq \frac{1}{4}d \neq \frac{1}{4}, \\ \mathcal{L}_1^0 : b \neq \pm e \text{ and } a = \frac{1}{4}d \neq \frac{1}{4}, & \quad \mathcal{L}_1^1 : b = e \text{ and } a = \frac{1}{4}d \neq \frac{1}{4}, & \quad \mathcal{L}_1^2 : b = -e \text{ and } a = \frac{1}{4}d \neq \frac{1}{4}, & \quad \mathcal{L}_1^3 : b = e = 0 \text{ and } a = \frac{1}{4}d \neq \frac{1}{4}, \\ \mathcal{L}_2^0 : b \neq \pm e \text{ and } a \neq \frac{1}{4}d = \frac{1}{4}, & \quad \mathcal{L}_2^1 : b = e \text{ and } a \neq \frac{1}{4}d = \frac{1}{4}, & \quad \mathcal{L}_2^2 : b = -e \text{ and } a \neq \frac{1}{4}d = \frac{1}{4}, & \quad \mathcal{L}_2^3 : b = e = 0 \text{ and } a \neq \frac{1}{4}d = \frac{1}{4}, \\ \mathcal{L}_3^0 : b \neq \pm e \text{ and } a = \frac{1}{4}d = \frac{1}{4}, & \quad \mathcal{L}_3^1 : b = e \text{ and } a = \frac{1}{4}d = \frac{1}{4}, & \quad \mathcal{L}_3^2 : b = -e \text{ and } a = \frac{1}{4}d = \frac{1}{4}, & \quad \mathcal{L}_3^3 : b = e = 0 \text{ and } a = \frac{1}{4}d = \frac{1}{4}. \end{aligned}$$

In all cases we noticed that

$$\begin{aligned} \mathcal{L}_0^0 \subset \mathcal{L}_1^0 \subset \mathcal{L}_3^0, & \quad \mathcal{L}_0^1 \subset \mathcal{L}_1^1 \subset \mathcal{L}_3^1, & \quad \mathcal{L}_0^2 \subset \mathcal{L}_1^2 \subset \mathcal{L}_3^2, \\ \mathcal{L}_0^0 \subset \mathcal{L}_2^0 \subset \mathcal{L}_3^0, & \quad \mathcal{L}_0^1 \subset \mathcal{L}_2^1 \subset \mathcal{L}_3^1, & \quad \mathcal{L}_0^2 \subset \mathcal{L}_2^2 \subset \mathcal{L}_3^2. \end{aligned}$$

$$\text{and } \mathcal{L}_0^1 \cong \mathcal{L}_2^0 \cong \mathcal{L}_3^0, \quad \mathcal{L}_1^1 \cong \mathcal{L}_2^1 \cong \mathcal{L}_3^1 \cong \mathcal{L}_2^2 \cong \mathcal{L}_3^2, \quad \mathcal{L}_1^0 \cong \mathcal{L}_2^0, \quad \mathcal{L}_3^0 \cong \mathcal{L}_3^2.$$

We summarize our results in the following table :

Table 18: Summarized results.

Cases	Subcases	Finite dimensional Lie algebra of symmetry	isomorphism between Lie algebras
$a = \frac{1}{4} d = \frac{1}{4}$	$b \neq \pm e$	$\mathcal{L}_3^0 : 6$ vector fields	$\mathcal{L}_3^0 \cong \mathfrak{iso}_2$
	$b = e, b \neq 0$	$\mathcal{L}_3^1 : 9$	$\mathcal{L}_3^1 \cong \mathcal{L}_3^2 \cong \mathcal{L}_3^3 \cong \mathfrak{sl}_2 \times J$
	$b = -e, b \neq 0$	$\mathcal{L}_3^2 : 9$	
	$b = e = 0$	$\mathcal{L}_3^3 : 9$	
$a \neq \frac{1}{4} d \neq \frac{1}{4}$	$b \neq \pm e$	$\mathcal{L}_0^0 : 2$ vector fields	$\mathcal{L}_0^1 \cong \mathcal{L}_0^2 \cong \mathcal{L}_0^3 \cong \mathbb{R}v_4 \times \mathfrak{sl}_2$
	$b = e, b \neq 0$	$\mathcal{L}_0^1 : 4$	
	$b = -e, b \neq 0$	$\mathcal{L}_0^2 : 4$	
	$b = e = 0$	$\mathcal{L}_0^3 : 4$	
$a = \frac{1}{4} d \neq \frac{1}{4}$	$b \neq \pm e$	$\mathcal{L}_1^0 : 4$ vector fields	$\mathcal{L}_1^0 \cong \mathfrak{iso}_2$
	$b = e, b \neq 0$	$\mathcal{L}_1^1 : 6$	$\mathcal{L}_1^1 \cong \mathcal{L}_1^2 \cong \mathcal{L}_1^3 \cong \mathfrak{sl}_2 \times \mathfrak{h}_3$
	$b = -e, b \neq 0$	$\mathcal{L}_1^2 : 6$	
	$b = e = 0$	$\mathcal{L}_1^3 : 6$	
$a \neq \frac{1}{4} d = \frac{1}{4}$	$b \neq \pm e$	$\mathcal{L}_2^0 : 4$ vector fields	$\mathcal{L}_2^0 \cong \mathfrak{iso}_2$
	$b = e, b \neq 0$	$\mathcal{L}_2^1 : 6$	$\mathcal{L}_2^1 \cong \mathcal{L}_2^2 \cong \mathcal{L}_2^3 \cong \mathfrak{sl}_2 \times \mathfrak{h}_3$
	$b = -e, b \neq 0$	$\mathcal{L}_2^2 : 6$	
	$b = e = 0$	$\mathcal{L}_2^3 : 6$	

Remark 6.1. From the beginning, we observed that the PDE class (1.2) admits the permutation of x and y as an equivalence transformation (symmetry group). Therefore, we might have anticipated that $\mathcal{L}_{a,b,d,e}$ and $\mathcal{L}_{d,e,a,b}$ would be isomorphic.

7 Conclusion

We examined the Lie point symmetries of the Kolmogorov backward equation (1.2) by, analyzing Lie algebras and their structures. Furthermore, we can extend this analysis to compute the Lie symmetries of stochastic differential equation (1.1) associated with partial differential equation (1.2). These symmetries are referred to as W-symmetries or random Lie point symmetries following the notation introduced by Gaeta [4]. For more comprehensive information, one can refer to Gaeta's works [5, 4] as well as the contributions by Kozlov [7], who has made additional advancements in this theory. For example, let us consider the Longstaff-Schwartz model (1.1) in the case ($b = e = 0$), and ($a = d = \frac{1}{4}$). Using Theorem 3.5 in [8], we find that the SDE has the following symmetries:

$$v_2 = x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} + \frac{\partial}{\partial t}, v_4 = \sqrt{x} \frac{\partial}{\partial x}, v_6 = \sqrt{y} \frac{\partial}{\partial y}, v_8 = \frac{\partial}{\partial t}.$$

Exponentiating the symmetry v_4 , gives the Lie group of transformation associated to v_4 which is $(\tilde{x}, \tilde{y}, \tilde{t}) = ((\sqrt{x + \frac{\varepsilon}{2}})^2, y, t)$, using Theorem 3.5 in [8], $\tilde{X}_t = (\sqrt{X_t + \frac{\varepsilon}{2}})^2$ must be a solution of the SDE

$dX_t = \frac{1}{4}dt + \sqrt{X_t}dB_t^1$. Using Ito formula, one finds that,

$$\begin{aligned} d\tilde{X}_t &= \left(1 + \frac{\varepsilon}{2\sqrt{X_t}}\right) (adt + \sqrt{X_t}dB_t^1) - \frac{1}{8} \left(\varepsilon X_t^{-\frac{1}{2}}\right) dt \\ &= adt + \frac{\varepsilon}{2\sqrt{X_t}} \left(a - \frac{1}{4}\right) + \left(\sqrt{X_t} + \frac{\varepsilon}{2}\right) dB_t^1 \\ d\tilde{X}_t &= \frac{1}{4}dt + \sqrt{\tilde{X}_t}dB_t^1 \end{aligned}$$

Whence, \tilde{X}_t is a solution of the SDE.

Acknowledgement

This paper and the research behind it would not have been possible without the exceptional support of Mr Paul LESCOT. His enthusiasm, knowledge and exacting attention to details have been an inspiration and kept my work on track.

References

- [1] George W Bluman and Sukeyuki Kumei. *Symmetries and differential equations*, volume 81. Springer Science & Business Media, 2013.
- [2] Adam Bowers. Classification of three-dimensional real lie algebras. *available online at the webpage math.ucsd.edu/abowers/downloads/survey/3d_Lie_alg_classify.pdf*, 2005.
- [3] Mark Craddock, Eckhard Platen, et al. *Symmetry group methods for fundamental solutions and characteristic functions*. Citeseer, 2003.
- [4] Giuseppe Gaeta. Lie-point symmetries and stochastic differential equations: Ii. *Journal of Physics A: Mathematical and General*, 33(27):4883, jul 2000.
- [5] Giuseppe Gaeta and Niurka Rodríguez Quintero. Lie-point symmetries and stochastic differential equations. *Journal of Physics A: Mathematical and General*, 32(48):8485, dec 1999.
- [6] B Kent Harrison and Frank B Estabrook. Geometric approach to invariance groups and solution of partial differential systems. *Journal of Mathematical Physics*, 12(4):653–666, 1971.
- [7] Roman Kozlov. Symmetries of systems of stochastic differential equations with diffusion matrices of full rank. *Journal of Physics A: Mathematical and Theoretical*, 43(24):245201, 2010.
- [8] Roman Kozlov. Symmetries of kolmogorov backward equation. *arXiv preprint arXiv:2010.16128*, 2020.
- [9] Paul Lescot. Symmetries of the Black-Scholes equation. *Methods Appl. Anal.*, 19(2):147–160, 2012.
- [10] Paul Lescot and Hélène Quintard. Symmetries of the backward heat equation with potential and interest rate models. *Comptes Rendus. Mathématique*, 352(6):525–528, 2014.
- [11] Paul Lescot, Helene Quintard, and Jean-Claude Zambrini. Solving stochastic differential equations with cartan’s exterior differential systems. *arXiv preprint arXiv:1404.4802*, 2014.
- [12] Paul Lescot and Jean-Claude Zambrini. Isovector for the Hamilton-Jacobi-Bellman equation, formal stochastic differentials and first integrals in Euclidean quantum mechanics. In *Seminar on stochastic analysis, random fields and applications IV, Centro Stefano Franscini, Ascona, Switzerland, May 20–24, 2002. Proceedings*, pages 187–202. Basel: Birkhäuser, 2004.
- [13] Francis A Longstaff and Eduardo S Schwartz. A two-factor interest rate model and contingent claims valuation. *The Journal of Fixed Income*, 2(3):16–23, 1992.
- [14] Peter J. Olver. *Applications of Lie groups to differential equations*, volume 107 of *Grad. Texts Math.* Springer, Cham, 1986.
- [15] Lev Vasil’evich Ovsiannikov. *Group analysis of differential equations*. Academic press, 2014.