

## ON THE MINIMAL DENOMINATOR PROBLEM IN FUNCTION FIELDS

NOY SOFFER ARANOV

ABSTRACT. We study the minimal denominator problem in function fields. In particular, we compute the probability distribution function of the random variable which returns the degree of the smallest denominator  $Q$ , for which the ball of a fixed radius around a point contains a rational function of the form  $\frac{p}{Q}$ . Moreover, we discuss the distribution of the random variable which returns the denominator of minimal degree, as well as higher dimensional and  $P$ -adic generalizations. This can be viewed as a function field generalization of a paper by Chen and Haynes.

## 1. INTRODUCTION

Meiss and Sanders [SM20] described an experiment in which a distance  $\delta > 0$  is fixed, and for randomly chosen  $x \in [0, 1)$ , they study the statistics of the function

$$q_{\min}(x, \delta) = \min \left\{ q : \exists \frac{p}{q} \in B(x, \delta), \gcd(p, q) = 1 \right\}. \quad (1.1)$$

Chen and Haynes [CH21] computed the probability that  $\mathbb{P}(q_{\min}(x, \delta) = q)$  for every  $\delta > 0$  and for every  $q \leq \lfloor \frac{1}{\delta} \rfloor$ . Moreover, they proved that  $\mathbb{E}[q_{\min}(\cdot, \delta)] = \frac{16}{\pi^2 \cdot \delta^{\frac{1}{2}}} + O(\log^2 \delta)$ . Markloff [Mar24] generalized the results of [CH21] to higher dimensions by studying the statistics of Farey fractions. The minimal denominator problem was investigated in the real setting in several other papers such as [DK77, Ste13], but it is not well studied over other fields.

In this paper, we use linear algebra and number theory to study the function field analogue of the function  $q_{\min}(x, \delta)$ , as well as its higher dimensional and  $P$ -adic analogues in the function field setting. In particular, we prove a function field analogue of the results of [CH21]. We note that unlike [CH21, Mar24], we do not study the distribution of Farey fractions, rather we use linear algebra and lattice point counting techniques, which work better in ultrametric spaces.

**1.1. Function Field Setting.** In this setting, we let  $q$  be a prime power and denote the ring of Laurent polynomials over  $\mathbb{F}_q$  by

$$\mathcal{R} = \left\{ \sum_{n=0}^N a_n x^n : a_n \in \mathbb{F}_q, N \in \mathbb{N} \cup \{0\} \right\}.$$

We let  $\mathcal{K}$  be the field of fractions of  $\mathcal{R}$ , and define an absolute value on  $\mathcal{K}$  by  $\left| \frac{f}{g} \right| = q^{\deg(f) - \deg(g)}$ , where  $f, g \in \mathcal{R}$  and  $g \neq 0$ . Then, the completion of  $\mathcal{K}$  with respect to  $|\cdot|$  is

$$\mathcal{K}_{\infty} = \left\{ \sum_{n=-N}^{\infty} a_n x^{-n} : a_n \in \mathbb{F}_q \right\}.$$

We let  $\mathcal{O} = \{\alpha \in \mathcal{K}_{\infty} : |\alpha| \leq 1\}$ , and let

$$\mathfrak{m} = x^{-1}\mathcal{O} = \{\alpha \in \mathcal{K}_{\infty} : |\alpha| \leq q^{-1}\}.$$

For  $\alpha \in \mathcal{K}_{\infty}$ , we write  $\alpha = [\alpha] + \{\alpha\}$ , where  $[\alpha] \in \mathcal{R}$  and  $\{\alpha\} \in \mathfrak{m}$ . In this paper, we define the Haar measure on  $\mathcal{K}_{\infty}$  to be the unique translation invariant measure  $\mu$ , such that  $\mu(\mathfrak{m}) = 1$ .

In  $\mathcal{K}_{\infty}^n$ , we define the supremum norm as  $\|(v_1, \dots, v_n)\| = \max_{i=1, \dots, n} \|v_i\|$ . Similarly, for  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathcal{K}_{\infty}^n$ , we let  $[\alpha] = ([\alpha_1], \dots, [\alpha_n])$  and  $\{\alpha\} = (\{\alpha_1\}, \dots, \{\alpha_n\})$ .

**1.2. Main Results.** We prove a function field analogue of the main results of [CH21]. Let  $n \in \mathbb{N}$ . For  $\delta > 0$  and  $\alpha \in \mathcal{K}_\infty^n$ , we define the minimal denominator degree by

$$\deg_{\min}(\alpha, \delta) = \min \left\{ d : \exists \frac{P}{Q}, \deg(Q) = d, \left| \alpha - \frac{P}{Q} \right| < \delta \right\}.$$

We say that  $Q$  is a minimal denominator for  $\alpha$  if  $\deg(Q) = \deg_{\min}(\alpha, \delta)$  and  $\left| \alpha - \frac{P}{Q} \right| < \delta$ . We note that if  $Q$  is a minimal denominator for  $\alpha$ , then, it is also a minimal denominator for  $\{\alpha\}$ . Hence, we only focus on  $\alpha \in \mathfrak{m}^n$ . Moreover, since the absolute value  $|\cdot|$  obtains values in  $\{0\} \cup \{q^k : k \in \mathbb{Z}\}$ , then, for every  $q^{-(k+1)} < \delta \leq q^{-k}$ , we have  $\deg_{\min}(\alpha, \delta) = \deg_{\min}(\alpha, q^{-k})$ . Hence, we only focus on  $\delta = q^{-k}$ , where  $k \in \mathbb{N}$ . We firstly compute the probability distribution function of  $\deg_{\min}(\cdot, q^{-k})$  when  $n = 1$ . From now on, we denote the probability distribution by  $\mathbb{P}$ .

**Theorem 1.1.** *Let  $k \in \mathbb{N}$ . Then, we have*

$$\mathbb{P}(\deg_{\min}(\alpha, q^{-1}) = d) = \begin{cases} \frac{1}{q} & d = 0, \\ \frac{q-1}{q} & d = 1 \end{cases},$$

and for every  $k \geq 2$ , we have

$$\mathbb{P}(\deg_{\min}(\alpha, q^{-k}) = d) = \begin{cases} q^{-k} & d = 0, \\ \frac{q-1}{q^{k-2d+1}} & d \leq \lceil \frac{k}{2} \rceil, d \in \mathbb{N}, \\ 0 & \text{else.} \end{cases} \quad (1.2)$$

**Corollary 1.2.** *We have*

$$\mathbb{E}[\deg_{\min}(\cdot, q^{-k})] = \begin{cases} \frac{q-1}{q} & k = 1, \\ \frac{q-1}{q^k} \left( \frac{q^{2\lceil \frac{k}{2} \rceil + 1} (\lceil \frac{k}{2} \rceil + 1) - (\lceil \frac{k}{2} \rceil + 2) q^{2\lceil \frac{k}{2} \rceil + 1}}{(q^2 - 1)^2} \right) & \text{else.} \end{cases} \quad (1.3)$$

*Proof.* When  $k = 1$ , the claim is immediate. Otherwise, by Theorem 1.1, we have

$$\begin{aligned} \mathbb{E}[\deg_{\min}(\alpha, q^{-k})] &= \sum_{d=0}^{\lceil \frac{k}{2} \rceil} d \frac{q-1}{q^k} q^{2d-1} = \frac{q-1}{q^k} \frac{d}{dt} \left( \sum_{d=0}^{\lceil \frac{k}{2} \rceil} t^d \right)_{t=q^2} \\ &= \frac{q-1}{q^k} \frac{d}{dt} \left( \frac{t^{\lceil \frac{k}{2} \rceil + 1} - 1}{t-1} \right)_{t=q^2} = \frac{q-1}{q^k} \left( \frac{q^{2\lceil \frac{k}{2} \rceil + 1} (\lceil \frac{k}{2} \rceil + 1) - (\lceil \frac{k}{2} \rceil + 2) q^{2\lceil \frac{k}{2} \rceil + 1}}{(q^2 - 1)^2} \right). \end{aligned} \quad (1.4)$$

□

Moreover, in every dimension, there is a unique monic polynomial which is a denominator of minimal degree.

**Lemma 1.3.** *For every  $\alpha \in \mathfrak{m}^n$  and for every  $k \geq 1$ , there exists a unique monic polynomial  $Q \in \mathcal{R}$ , such that  $\deg(Q) = \deg_{\min}(\alpha, q^{-k})$  and  $\|Q\alpha\| < q^{-k}$ .*

This motivates the following definition.

*Remark 1.4.* Due to Lemma 1.3, we denote the unique monic polynomial  $Q$  satisfying  $\deg(Q) = \deg_{\min}(\alpha, q^{-k})$  and  $\|Q\alpha\| < q^{-k}$  by  $Q_{\min}(\alpha, q^{-k})$ .

We also compute the distribution of  $Q_{\min}(\cdot, q^{-k})$ . To do so, we shall use some notations from number theory.

**Definition 1.5.** For a polynomial  $Q$ , we let  $d(Q)$  be the number of prime divisors of  $Q$ , we let  $D(Q)$  be the number of monic divisors of  $Q$ , and we let  $S(Q)$  be the set of divisors of  $Q$ . We define

$$\mu(Q) = \begin{cases} (-1)^{d(Q)} & Q \text{ is square free,} \\ 0 & \text{if there exists } P \text{ such that } P^2 \mid Q \end{cases}$$

**Definition 1.6.** For a polynomial  $Q \in \mathcal{R}$ , we define  $S_{\text{monic}}^{\mathbb{1}, \ell}(Q)$  to be the set of  $\ell$  tuples  $(a_1, \dots, a_\ell)$ , such that  $a_i$  are distinct monic polynomials which divide  $Q$ , and  $\deg(a_i) < \deg(Q)$ .

**Theorem 1.7.** Let  $Q$  be a monic polynomial with  $\deg(Q) \leq \lceil \frac{k}{2} \rceil$ . Then, for every  $k \geq 1$ , the probability that  $Q_{\min}(\alpha, q^{-k}) = Q$  is

$$\frac{1}{q^k} \left( |Q| + \sum_{N|Q, \deg(N) < \deg(Q)} |N| \sum_{\ell=1}^{D(N)} (-1)^\ell \left( \frac{D\left(\frac{Q}{N}\right)!}{\left(D\left(\frac{Q}{N}\right) - \ell\right)!} + \sum_{M \in S\left(\frac{Q}{N}\right): D\left(\frac{Q}{NM}\right) \geq \ell} \mu(M) \frac{D(M)!}{(D(M) - \ell)!} \right) \right). \quad (1.5)$$

In particular, if  $Q$  is an irreducible monic polynomial of degree  $d$ , then,

$$\mathbb{P}(Q_{\min}(\alpha, q^{-k}) = Q) = \frac{q^d - 1}{q^k}. \quad (1.6)$$

The higher dimensional setting discusses a simultaneous solution for the equations  $|\{Q\alpha_i\}| < q^{-k}$ .

**Lemma 1.8.** For every  $k \in \mathbb{N}$ , and for every  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathfrak{m}^{n-1}$ , we have

$$\deg_{\min}(\alpha, q^{-k}) \geq \max_{i=1, \dots, n} \deg_{\min}(\alpha_i, q^{-k}).$$

*Proof.* If there exists  $Q \in \mathcal{R}$  with  $\deg(Q) = \deg_{\min}(\alpha, q^{-k})$ , such that for every  $i = 1, \dots, n$ , we have  $\|Q\alpha_i\| < q^{-k}$ , then,  $\deg(Q) \geq \deg_{\min}(\alpha_i, q^{-k})$  for every  $i = 1, \dots, n$ . Hence,  $\deg_{\min}(\alpha, q^{-k}) \geq \max \deg_{\min}(\alpha_i, q^{-k})$ .  $\square$

Hence, it is natural to ask what is the probability that  $\deg_{\min}(\alpha, q^{-k}) = \max_{i=1, \dots, n} \deg_{\min}(\alpha_i, q^{-k})$ . This is an immediate corollary of Lemma 1.3 and Lemma 1.8

**Corollary 1.9.** Let  $\alpha \in \mathfrak{m}^n$  and let  $k \geq 1$ . Then, for  $Q \in \mathcal{R}$  with  $\deg(Q) = d$ , for  $0 \leq d \leq \lceil \frac{k}{2} \rceil$ , we have  $Q_{\min}(\alpha, q^{-k}) = Q$  if and only if

- (1) for every  $i = 1, \dots, n$ , we have  $Q_{\min}(\alpha_i, q^{-k})|Q$ ;
- (2) there exists  $i$  such that  $Q_{\min}(\alpha_i, q^{-k}) = Q$ .

*Proof of Corollary 1.9.* By Lemma 1.8, we have  $\deg_{\min}(\alpha, q^{-k}) \geq \deg_{\min}(\alpha_i, q^{-k})$  for every  $i = 1, \dots, n$ . For  $i = 1, \dots, n$ , let  $Q_i = Q_{\min}(\alpha_i, q^{-k})$ . If  $Q_i = Q$  for every  $i$ , then, there is nothing to prove. Otherwise, let  $i$  be such that  $Q_i \neq Q$ . Then, by Lemma 1.3, if  $\left| \alpha - \frac{P_i}{Q_i} \right| < q^{-k}$  and  $\left| \alpha - \frac{P}{Q} \right| < q^{-k}$ , then,  $\frac{P}{Q} = \frac{P_i}{Q_i}$ . Thus,  $Q_i$  divides  $Q$ .  $\square$

Furthermore, we bound the probability that  $\deg_{\min}(\alpha, q^{-k}) \leq d$  in higher dimensions.

**Lemma 1.10.** Let  $n, k \in \mathbb{N}$ . Then, for every  $\alpha \in \mathfrak{m}^n$ , we have  $\deg_{\min}(\alpha, q^{-k}) \leq \left\lceil \frac{nk}{n+1} \right\rceil$ .

Moreover, for every  $d \leq \left\lceil \frac{nk}{n+1} \right\rceil$ , we have

$$\mathbb{P}(\deg_{\min}(\alpha, q^{-k}) \leq d) \leq q^{-(kn - (n+1)d)}. \quad (1.7)$$

We shall also discuss a  $P$ -adic variant of the minimal denominator problem. Let  $P \in \mathcal{R}$  be an irreducible polynomial, let  $\alpha \in \mathfrak{m}$ , and let  $k \geq 1$ . We define

$$\deg_{\min, P}(\alpha, q^{-k}) = \min \left\{ d \geq 0 : \exists m \geq 0, \exists \frac{a}{b} : \deg(b) = d, \frac{a}{P^m b} \in B(\alpha, q^{-k}) \right\} = \inf_{m \geq 0} \deg_{\min}(P^m \alpha, q^{-k}).$$

**Theorem 1.11.** For every irreducible polynomial  $P$  and for almost every  $\alpha \in \mathfrak{m}$ , we have  $\deg_{\min, P}(\alpha, q^{-k}) = 0$ . Moreover, for every  $1 \leq d \leq \lceil \frac{k}{2} \rceil$ , we have

$$\dim_H \{ \alpha \in \mathfrak{m} : \deg_{\min, P}(\alpha, q^{-k}) \geq d \} = \frac{\log_q(|P|^k - |P|^{2(d-1)})}{k \deg(P)}.$$

Moreover, when  $d = 0$ , we have an equality.

*Remark 1.12.* A natural question pertains to the minimal denominator question in the infinite residue case, for example over  $\mathbb{Q}[x]$  or  $\mathbb{C}[x]$ . Since many of our results rely on counting techniques, these results do not hold for function fields with an infinite residue field.

**1.3. Acknowledgements.** I would like to thank Eran Igra and Albert Artiles who accidentally introduced me to the minimal denominator problem, and thus led to the birth of this paper.

## 2. HANKEL MATRICES

We first translate the minimal denominator problem to the language of linear algebra. For  $k, \ell \in \mathbb{N}$  and  $\alpha = \sum_{i=1}^{\infty} \alpha_i x^{-i} \in \mathfrak{m}$ , we define the Hankel matrix of  $\alpha$  of order  $(k, \ell)$  as

$$\Delta_{\alpha}(k, \ell) = \begin{pmatrix} \alpha_1 & \alpha_2 & \dots & \alpha_{\ell} \\ \alpha_2 & \alpha_3 & \dots & \alpha_{\ell+1} \\ \vdots & \dots & \ddots & \vdots \\ \alpha_k & \alpha_{k+1} & \dots & \alpha_{k+\ell-1} \end{pmatrix}.$$

Assume that  $\alpha \in \mathfrak{m}$  and  $\frac{P}{Q} \in \mathbb{F}_q(x)$ . Then,  $\left| \alpha - \frac{P}{Q} \right| < q^{-k}$  if and only if  $|Q\alpha - P| < q^{-(k - \deg(Q))}$ . Let  $d = \deg(Q)$ . Note that by Dirichlet's theorem,  $\deg_{\min}(\alpha, q^{-k}) \leq k$ . Hence, we can assume that  $d \leq k$ . Then,  $|\{Q\alpha\}| < q^{-(k-d)}$  if and only if

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \dots & \alpha_{d+1} \\ \alpha_2 & \alpha_3 & \dots & \alpha_{d+2} \\ \vdots & \dots & \ddots & \vdots \\ \alpha_{k-d} & \alpha_{k-d+1} & \dots & \alpha_k \end{pmatrix} \begin{pmatrix} Q_0 \\ Q_1 \\ \vdots \\ Q_d \end{pmatrix} = 0, \quad (2.1)$$

where  $Q = \sum_{i=0}^d Q_i x^i$ . We notice that if  $d+1 \geq k-d+1$ , that is if  $d \geq \frac{k}{2}$ , then, there exists a non-trivial solution to (2.1). Hence,  $\deg_{\min}(\alpha, q^{-k}) \leq \lceil \frac{k}{2} \rceil$ .

*Remark 2.1.* We note that  $\deg_{\min}(\alpha, q^{-k}) = d$ , for  $d \leq \lceil \frac{k}{2} \rceil$ , if and only if for every  $j < d$ , we have that the matrix  $\Delta_{\alpha}(k-j, j+1)$  has rank  $j+1$ , but the matrix  $\Delta_{\alpha}(k-d, d+1)$  has rank  $d$ .

**Lemma 2.2.** *For every  $0 \neq \alpha \in \mathfrak{m}$ , there exists a unique  $d \leq \frac{k}{2}$  for which there exist coprime  $P, Q$  with  $\deg(Q) = d$  such that  $\left| \alpha - \frac{P}{Q} \right| < q^{-k}$ .*

*Proof.* Assume that there exist  $P, Q, P', Q' \in \mathcal{R}$  such that

- (1)  $Q, Q' \neq 0$
- (2)  $\gcd(P, Q) = 1 = \gcd(P', Q')$ ,
- (3)  $\deg(Q') = d' < d = \deg(Q) \leq \frac{k}{2}$ ,
- (4)  $\left| \alpha - \frac{P}{Q} \right| < q^{-k}$ , and
- (5)  $\left| \alpha - \frac{P'}{Q'} \right| < q^{-k}$ .

Then, by the ultrametric inequality and the fact that these fractions are reduced, we have

$$\left| \frac{PQ' - P'Q}{QQ'} \right| = \left| \frac{P}{Q} - \frac{P'}{Q'} \right| < \frac{1}{q^k}. \quad (2.2)$$

Thus,  $|PQ' - P'Q| < \frac{q^{d+d'}}{q^k} \leq q^{2d} q^k = q^{-(k-2d)} \leq 1$ . Hence,  $\frac{P}{Q} = \frac{P'}{Q'}$ , which contradicts the assumption that  $\gcd(P', Q') = \gcd(P, Q) = 1$ .  $\square$

We first use this reinterpretation to prove Lemma 1.3. To do so we define for  $\alpha \in \mathfrak{m}^n$  the matrix

$$\Delta_{\alpha}(k, \ell) = \begin{pmatrix} \Delta_{\alpha_1}(k, \ell) \\ \vdots \\ \Delta_{\alpha_n}(k, \ell) \end{pmatrix}.$$

*Proof of Lemma 1.3.* We notice that if  $d = \deg_{\min}(\alpha, q^{-k})$ , then,

$$\text{rank}(\Delta_{\alpha}(k-d-1, d)) = d = \text{rank}(\Delta_{\alpha}(k-d, d+1)).$$

Hence,  $\dim \text{Ker}(\Delta_{\alpha}(k-d, d+1)) = 1$ . Let  $Q$  be a polynomial satisfying  $\deg(Q) = d$  and  $\|Q\alpha\| < q^{-k}$ . Without loss of generality we can assume that  $Q_d = 1$ . Thus,

$$\Delta_{\alpha}(k-d, d+1)(Q_0, Q_1, \dots, Q_{d-1}, 1)^t = 0.$$

Since  $\dim \text{Ker}(\Delta_{\alpha}(k-d, d+1)) = 1$ , then,  $(Q_0, Q_1, \dots, Q_{d-1}, 1)$  is the unique vector  $\mathbf{v} = (v_0, \dots, v_d)$  with  $v_d = 1$ , such that  $\Delta_{\alpha}(k-d, d+1)\mathbf{v} = 0$ . Thus, by (2.1),  $Q$  is the unique monic polynomial of minimal degree with  $\|Q\alpha\| < q^{-k}$ .  $\square$

We shall use several facts about ranks of Hankel matrices to prove Theorems 1.1 and 1.7.

**Theorem 2.3.** [GAGR11, Theorem 5.1] *Let  $r > 0$ . Then, the number of invertible  $h \times h$  Hankel matrices with entries in  $\mathbb{F}_q$  of rank  $r$ ,  $N(r, h; q)$ , is equal to*

$$N(r, h; q) = \begin{cases} 1 & r = 0 \\ q^{2r-2}(q^2 - 1) & 1 \leq r \leq h-1 \\ q^{2h-2}(q-1) & r = h \end{cases}. \quad (2.3)$$

We shall also use the following generalization of Theorem 2.3.

**Theorem 2.4.** [DG22, Theorem 1.1] *Let  $k, \ell \in \mathbb{N}$ , let  $F$  be a finite field with  $|F| = q$ , and let  $r \leq \min\{k, \ell\} - 1$ . Then, the number of Hankel matrices  $\Delta_{\alpha}(k, \ell)$  over  $F$  with rank at most  $r$  is  $q^{2r}$ .*

**Lemma 2.5.** [ANL21, Lemma 2.3] *Let  $m, n \in \mathbb{N}$ , and let  $k \leq \min\{m, n-1\}$ . Let  $H = \Delta_{\alpha}(m, n)$  be a Hankel matrix. If the first  $k$  columns of  $H$  are independent, but the first  $k+1$  columns of  $H$  are dependent, then,  $\det(\Delta_{\alpha}(k, k)) \neq 0$ .*

### 3. PROOFS IN THE ONE DIMENSIONAL CASE

*Proof of Theorem 1.1.* By using the reinterpretation in §2, we realize that if  $k = 1$ , then, there is a non-trivial solution for (2.1) when  $d \geq 1$ . Moreover, there exists a solution for (2.1) when  $k = 1$  and  $d = 0$  if and only if  $\alpha_1 = 0$ . Hence,

$$\mathbb{P}(\deg_{\min}(\alpha, q^{-1}) = 0) = \frac{1}{q}, \mathbb{P}(\deg_{\min}(\alpha, q^{-1}) = 1) = \frac{q-1}{q}.$$

If,  $k \geq 2$  and  $d \geq \frac{k}{2}$ , then, there exists a non-trivial solution to (2.1). Hence,  $\deg_{\min}(\alpha, q^{-k}) \leq \lceil \frac{k}{2} \rceil$ . Firstly, if  $d = 0$ , then,  $\alpha_1 = \dots = \alpha_k$ , and therefore,  $\mathbb{P}(\deg_{\min}(\alpha, q^{-k}) = 0) = q^{-k}$ .

Let  $1 \leq d \leq \frac{k}{2}$  and let  $\alpha \in \mathbb{F}_q((x^{-1}))$ . By Remark 2.1, we have  $d = \deg_{\min}(\alpha, q^{-k})$  if and only if the columns of the Hankel matrix  $\Delta_{\alpha}(k-d, d+1)$  are linearly dependent, but the columns of  $\Delta_{\alpha}(k-d-1, d)$  are linearly independent. Hence, by Lemma 2.5, the matrix  $\Delta_{\alpha}(d, d)$  is invertible. Hence, there exist unique  $a_1, \dots, a_d \in \mathbb{F}_q$ , such that

$$\begin{pmatrix} \alpha_{d+1} \\ \alpha_{d+2} \\ \vdots \\ \alpha_{2d} \end{pmatrix} = a_1 \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_d \end{pmatrix} + \dots + a_d \begin{pmatrix} \alpha_d \\ \alpha_{d+1} \\ \vdots \\ \alpha_{2d-1} \end{pmatrix}. \quad (3.1)$$

On the other hand, since the columns of  $\Delta_{\alpha}(k-d, d+1)$  are linearly dependent, and the columns of  $\Delta_{\alpha}(k-d+1, d)$  are linearly independent, there exist  $b_1, \dots, b_d \in \mathbb{F}_q$ , such that

$$\begin{pmatrix} \alpha_{d+1} \\ \alpha_{d+2} \\ \vdots \\ \alpha_k \end{pmatrix} = b_1 \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_{k-d} \end{pmatrix} + \dots + b_d \begin{pmatrix} \alpha_d \\ \alpha_{d+1} \\ \vdots \\ \alpha_{k-1} \end{pmatrix}. \quad (3.2)$$

Thus, by (3.1) and (3.2), we have  $a_i = b_i$  for every  $i = 1, \dots, d$ . Hence, given an invertible matrix  $d \times d$  Hankel matrix  $\Delta_\alpha(d, d)$  and some  $\alpha_{2d} \in \mathbb{F}_q$ , there is exactly one way to extend the word  $(\alpha_1, \dots, \alpha_{2d})$  to a Laurent sequence  $\sigma = \sum_{i=1}^{\infty} \alpha_i x^{-i}$  satisfying  $\deg_{\min}(\sigma, q^{-k}) = d$ . Therefore, by Theorem 2.3 (see also Theorem 2.4), we have

$$\mathbb{P}(\deg_{\min}(\alpha, q^{-k}) = d) = \frac{q^{2d-1}(q-1)}{q^k}.$$

□

To prove Theorem 1.7, we use the following fact from [CR19]: We have  $\|Q\alpha\| < q^{-(k-d)}$ , where  $Q = Q_0 + Q_1x + \dots + Q_{d-1}x^{d-1} + x^d$ , if and only if

$$\begin{pmatrix} Q_0 & \dots & Q_{d-1} & -1 & 0 & \dots & 0 \\ 0 & Q_0 & \dots & Q_{d-1} & -1 & 0 & \dots \\ \vdots & \dots & \ddots & \ddots & \dots & \ddots & \ddots \\ 0 & \dots & \dots & Q_0 & \dots & Q_{d-1} & -1 \end{pmatrix} \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_k \end{pmatrix} = \mathbf{0}. \quad (3.3)$$

We denote the matrix the right hand side of (3.3) by  $A_Q$ , and we denote  $\pi_k(\alpha) = (\alpha_1, \dots, \alpha_k)^t$ . Then,  $A_Q \in M_{k-d \times k}(\mathbb{F}_q)$ , and  $\dim(\text{Ker}(A_Q)) = d$ . Hence,  $|\text{Ker}(A_Q)| = q^d$ .

**Proposition 3.1.** *The number of primitive vectors in  $S_{\text{monic}}^{\mathfrak{A}, \ell}(Q)$  is*

$$|\widehat{S}_{\text{monic}}^{\mathfrak{A}, \ell}| = \begin{cases} \frac{D(Q)!}{(D(Q)-\ell)!} + \sum_{N \in S(Q): D(\frac{Q}{N}) \geq \ell} \mu(N) \frac{D(N)!}{(D(N)-\ell)!} & D(Q) \geq \ell, \\ 0 & \text{else.} \end{cases} \quad (3.4)$$

*Remark 3.2.* We use Proposition 3.1 to count the number of tuples whose greatest common denominator is 1, instead of the classical counting method, since this yields a more compact expression in the proof of Theorem 1.7

*Proof of Proposition 3.1.* We first note that

$$|S_{\text{monic}}^{\mathfrak{A}, \ell}(Q)| = \begin{cases} \frac{D(Q)!}{(D(Q)-\ell)!} & |D(Q)| \geq \ell, \\ 0 & \text{else.} \end{cases}$$

Hence, to count primitive vectors in  $S_{\text{monic}}^{\mathfrak{A}, \ell}$ , we use the exclusion inclusion principle and induction to obtain that

$$\begin{aligned} |\widehat{S}_{\text{monic}}^{\mathfrak{A}, \ell}| &= \left| S_{\text{monic}}^{\mathfrak{A}, \ell} \setminus \bigcup_{P \in S(Q) \text{ prime}} P S_{\text{monic}}^{\mathfrak{A}, \ell} \left( \frac{Q}{P} \right) \right| \\ &= |S_{\text{monic}}^{\mathfrak{A}, \ell}| - \sum_{P_1, \dots, P_i \in S(Q) \text{ prime}} (-1)^{i+1} \left| P_1 \dots P_i S_{\text{monic}}^{\mathfrak{A}, \ell} \left( \frac{Q}{P_1 \dots P_i} \right) \right| \\ &= \frac{D(Q)!}{(D(Q)-\ell)!} + \sum_{N \in S(Q)} \mu(N) \left| S_{\text{monic}}^{\mathfrak{A}, \ell} \left( \frac{Q}{N} \right) \right| \\ &= \frac{D(Q)!}{(D(Q)-\ell)!} + \sum_{N \in S(Q): D(\frac{Q}{N}) \geq \ell} \mu(N) \frac{D(N)!}{(D(N)-\ell)!}. \end{aligned} \quad (3.5)$$

□

*Proof of Theorem 1.7.* Let  $Q$  be a monic polynomial of degree at most  $\lceil \frac{k}{2} \rceil$ . By Lemma 1.3, if  $\pi_k(\alpha) \in \text{Ker}(A_D) \cap \text{Ker}(A_Q)$ , where  $\deg(D) < \deg(Q)$ , then,  $D|Q$ . Hence, by the exclusion

inclusion principle,

$$\begin{aligned}
\mathbb{P}(Q_{\min}(\alpha, q^{-k}) = Q) &= \mathbb{P}\left(\pi_k(\alpha) \in (\text{Ker}(A_Q)) \setminus \bigcup_{D|Q, \deg(D) < \deg(Q)} \text{Ker}(A_D)\right) \\
&= \mathbb{P}(\pi_k(\alpha) \in \text{Ker}(A_Q)) - \sum_{D_1, \dots, D_\ell | Q} (-1)^{\ell+1} \mathbb{P}\left(\bigcap_{i=1}^{\ell} \text{Ker}(A_{D_i})\right) \\
&= q^{-(k-\deg(Q))} + \sum_{D_1, \dots, D_\ell | Q} (-1)^\ell \mathbb{P}(\text{Ker}(A_{\gcd(D_1, \dots, D_\ell)})).
\end{aligned} \tag{3.6}$$

We notice that if  $N|Q$ , then,  $N = \gcd(D_1, \dots, D_\ell)$  if and only if for every  $i = 1, \dots, \ell$ , there exists a monic polynomial  $a_i \in \mathcal{R}$  such that  $D_i = a_i N$ ,  $a_i | \frac{Q}{N}$ , and  $\gcd(a_1, \dots, a_\ell) = 1$ . Hence,  $(a_1, \dots, a_\ell) \in \mathcal{R}^\ell$  is a primitive vector with distinct coordinates, which are all monic polynomials which divide  $\frac{Q}{N}$ , so that  $(a_1, \dots, a_\ell) \in S_{\text{monic}}^{\mathbb{N}, \ell}\left(\frac{Q}{N}\right)$ . Hence, by Proposition 3.1, we have

$$\begin{aligned}
&\sum_{D_1, \dots, D_\ell | Q} (-1)^{\ell+1} \mathbb{P}(\text{Ker}(A_{\gcd(D_1, \dots, D_\ell)})) \\
&= \frac{1}{q^k} \sum_{N|Q, \deg(N) < \deg(Q)} |N| \sum_{\ell=1}^{D(Q)} (-1)^{\ell+1} \#\{(D_1, \dots, D_\ell) : \gcd(D_1, \dots, D_\ell) = N\} \\
&= \frac{1}{q^k} \sum_{N|Q, \deg(N) < \deg(Q)} |N| \sum_{\ell=1}^{D(N)} (-1)^{\ell+1} \left| \widehat{S}_{\text{monic}}^{\mathbb{N}, \ell}\left(\frac{Q}{N}\right) \right| \\
&= \frac{1}{q^k} \sum_{N|Q, \deg(N) < \deg(Q)} |N| \sum_{\ell=1}^{D(N)} (-1)^{\ell+1} \left( \frac{D\left(\frac{Q}{N}\right)!}{\left(D\left(\frac{Q}{N}\right) - \ell\right)!} + \sum_{M \in S\left(\frac{Q}{N}\right): D\left(\frac{Q}{N}\right) \geq \ell} \mu(M) \frac{D(M)!}{(D(M) - \ell)!} \right).
\end{aligned} \tag{3.7}$$

Hence, the probability that  $Q_{\min}(\alpha, q^{-k}) = Q$ , for  $\deg(Q) \leq \lceil \frac{k}{2} \rceil$  is equal to

$$\frac{1}{q^k} \left( |Q| + \sum_{N|Q, \deg(N) < \deg(Q)} |N| \sum_{\ell=1}^{D(N)} (-1)^\ell \left( \frac{D\left(\frac{Q}{N}\right)!}{\left(D\left(\frac{Q}{N}\right) - \ell\right)!} + \sum_{M \in S\left(\frac{Q}{N}\right): D\left(\frac{Q}{N}\right) \geq \ell} \mu(M) \frac{D(M)!}{(D(M) - \ell)!} \right) \right).$$

□

#### 4. PROOF OF THEOREM 1.11

*Proof of Theorem 1.11.* We notice that  $|P^m b \alpha - a| < q^{-(k-d)}$ , where  $d = \deg(b)$ , if and only if

$$\begin{pmatrix} \alpha_m & \alpha_{m+1} & \cdots & \alpha_{m+d} \\ \alpha_{m+1} & \alpha_{m+2} & \cdots & \alpha_{m+d+1} \\ \vdots & \cdots & \ddots & \vdots \\ \alpha_{m+k-d} & \cdots & \cdots & \alpha_{m+k} \end{pmatrix} \begin{pmatrix} b_0 \\ b_1 \\ \vdots \\ b_d \end{pmatrix} = 0, \tag{4.1}$$

where  $\alpha = \sum_{i=1}^{\infty} P^{-i} \alpha_i$ , for some  $\alpha_i \in \mathbb{F}_q[x]/P\mathbb{F}_q[x]$ . Hence,  $\deg_{\min, P}(\alpha, q^{-k}) \leq \frac{k}{2}$ . Since almost every string is normal [Bor09], then, for every  $k$ , for every prime  $P$ , and for almost every  $\alpha \in \mathfrak{m}$ , the string  $0^k$  appears in the infinite word  $\{\alpha_i\}_{i \in \mathbb{N}}$ . Hence, if  $\alpha_m = \alpha_{m+1} = \cdots = \alpha_{m+k-1} = 0$ , then, there exists  $a$ , such that  $|P^m \alpha - a| < q^{-(k-d)}$ . Thus,  $\mathbb{P}(\deg_{\min, P}(\alpha, q^{-k}) = 0) = 1$ .

By Theorem 2.4, for every  $k$  and every  $d \leq k-d$ , the number of Hankel matrices  $\text{rank}(\Delta_\alpha(k-d-1, d)) = d$  with entries in  $\mathbb{F}_q[x]/P\mathbb{F}_q[x]$  of rank  $d$  is  $|P|^k - |P|^{2(d-1)}$ . Thus, for every  $1 \leq d \leq \frac{k}{2}$ ,

we have

$$\dim_H\{\alpha \in \mathfrak{m} : \deg_{\min,t}(\alpha, q^{-k}) \geq d\} = \frac{\log_q(|P|^k - |P|^{2(d-1)})}{k \deg(P)}.$$

□

## 5. PROOFS OF LEMMA 1.10

*Proof of Lemma 1.10.* We note that  $\|Q\alpha\| < q^{-k}$  if and only if

$$\Delta_\alpha(k-d, d+1)(Q_0, Q_1, \dots, Q_d)^t = 0. \quad (5.1)$$

Thus, if  $d \geq \frac{nk}{n+1}$ , there will always be a non-zero solution to (5.1). If  $d = 0$ , then, for every  $i = 1, \dots, n$ , we have  $(\alpha_{i1}, \dots, \alpha_{ik}) = 0$ , where  $\alpha_i = \sum_{j=1}^{\infty} \alpha_{ij} x^{-j}$ . Hence,

$$\mathbb{P}\left(\deg_{\min}(\alpha, q^{-k}) = 0\right) = q^{-nk}. \quad (5.2)$$

Let  $1 \leq d \leq \left\lceil \frac{nk}{n+1} \right\rceil$ , and let  $Q$  be a monic polynomial with  $\deg(Q) = d$ . Since  $|\text{Ker}(A_Q) \setminus \{0\}| = q^d$ , then, we have

$$\mathbb{P}\left(Q_{\min}(\alpha, q^{-k}) = Q\right) \leq \mathbb{P}\left(\bigcap_{i=1}^n (\alpha_i \in \text{Ker}(A_Q))\right) = q^{-n(k-d)} \quad (5.3)$$

Hence,

$$\mathbb{P}\left(\deg_{\min}(\alpha, q^{-k}) = d\right) \leq q^{-nk+(n+1)d}. \quad (5.4)$$

□

## REFERENCES

- [ANL21] Faustin Adiceam, Erez Nesharim, and Fred Lunnon, *On the  $t$ -adic littlewood conjecture*, Duke Mathematical Journal **170**(10) (2021), 2371–2419.
- [Bor09] E. Borel, *Les probabilités dénombrables et leurs applications arithmétiques.*, Rend. Circ. Mat. Palermo **27** (1909), 247–271 (French).
- [CH21] Huayang Chen and Alan Haynes, *Expected value of the smallest denominator in a random interval of fixed radius*, ArXiv **abs/2109.12668** (2021).
- [CR19] Sam Chow and Luka Rimanic, *Lonely runners in function fields*, Mathematika **65** (2019), 677–701.
- [DG22] Omesh Dhar Dwivedi and Darij Grinberg, *On the rank of hankel matrices over finite fields*, Linear Algebra and its Applications **641** (2022), 156–181.
- [DK77] H.G Meijer D Kruyswijk, *On small denominators and farey sequences*, Indagationes Mathematicae (Proceedings) **80** (1977), 332–337.
- [GAGR11] Mario García-Armas, Sudhir R. Ghorpade, and Samrith Ram, *Relatively prime polynomials and nonsingular hankel matrices over finite fields*, Journal of Combinatorial Theory, Series A **118** (2011), 819–828.
- [Mar24] J. Marklof, *Smallest denominators*, Bull. London Math. Soc. **56** (2024), 1920–1938.
- [SM20] Evelyn Sander and James .D. Meiss, *Birkhoff averages and rotational invariant circles for area-preserving maps*, Physica D: Nonlinear Phenomena **411** (2020).
- [Stel3] C. L. Stewart, *On the distribution of small denominators in the farey series of order  $n$* , Advances in combinatorics (2013), 275–286.

*Email address:* `noysofferaranov@math.utah.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY, UTAH, USA