

Minimal Models for RNA Simulations

D. Thirumalai, Naoto Hori, and Hung T. Nguyen

January 3, 2025

Department of Chemistry, The University of Texas at Austin, Austin, TX 78712, USA

School of Pharmacy, University of Nottingham, Nottingham, NG7 2RD, UK

Department of Chemistry, University at Buffalo, NY 14260, USA

The increasing importance of RNA as a prime player in biology can hardly be overstated. It is suspected that the functions of RNA are linked to their structures and dynamics. Many of the problems in RNA, such as folding and RNA-RNA interactions that drive phase separation even in the absence of proteins, require cations. Because experiments alone cannot directly reveal the dynamics of cation-RNA interactions, well calibrated theory and computations are needed to predict how ions control the behavior of RNA. In this review, we outline the development of coarse-grained models at different resolutions with application to phase separation in low complexity sequences. We also describe folding of ribozymes and riboswitches with a focus on the impact of monovalent and divalent cations. We outline major challenges that need to be overcome to simulate complex problems such as assembly of ribosomes.

1 Introduction

Although RNA sometimes referred to as the “dark matter”, the spotlight on RNA biology has never been brighter with the award of Nobel prizes in medicine in two consecutive years, one in 2023 for the COVID-19 vaccine based on nucleoside base modifications of RNA¹, and another in 2024 for the role of microRNA

in regulating gene expression^{2;3}. These are the latest high profile examples in RNA biology which took off after the demonstration that RNA could function as enzymes over forty years ago⁴⁻⁶. In the intervening years the functional repertoire of RNA has continuously expanded producing one surprise after another and with no end in sight. As is always the case in science, the discovery of new phenomena usually create a treasure trove of problems that need quantitative explanations, which in this field require both biophysical experiments⁷⁻⁹ and computational approaches¹⁰⁻¹⁴.

The focus of this review is to highlight the utility of coarse-grained (CG) models in the folding of RNA and phase separation in low complexity sequences. Although not a substitute for atomically detailed simulations in water with explicit ions, there is a great need for CG models in understanding the folding of ribozymes (RNA enzymes) as well in the simulations of multiple chains that span a range of time and length scales. Over the last two decades several studies have shown that CG models, beginning with the initial reports on the mechanical force-induced unfolding of a RNA hairpin^{10;15}, have been efficacious in making predictions that compare favorably with experiments. The CG models, at different resolutions, have been particularly important in elucidating the role of divalent cations in controlling RNA folding¹⁶⁻¹⁸.

For the purposes of this article, let us classify CG models into a few categories depending on the resolution of the RNA and the treatment of ions (Figure 1). (i) In the Single Site Interaction (SIS) model, each nucleotide is represented by a bead^{19;20}. Electrostatic interactions associated with phosphate groups and the cations are modeled implicitly using the Debye-Hückel (DH) interactions¹⁹⁻²¹. (ii) In the Three Interaction Site (TIS) model¹⁰, each nucleotide has three sites representing phosphate, sugar, and base. Interactions involving charged groups (phosphate and metal ions) are treated explicitly¹⁶ or implicitly²², depending on the applications. (iii) If the monovalent (such as Na⁺ or K⁺) cation concentration exceeds the concentration of divalent cations greatly, as is the case in several folding experiments, the electrostatic interactions associated with monovalent cations could be treated using the DH potential where as divalent cations (Mg²⁺ or Ca²⁺) have to be explicitly treated for accurate description of RNA folding. The mixed description of cations when combined with the TIS model for RNA is referred to as the TIS-CAT model²³. (iv) There are models that use higher resolution (six or seven beads per nucleotide) in which charged interactions are either neglected or treated in an *ad hoc* manner. Most of these models were not designed to investigate RNA folding²⁴ or RNA-RNA interactions. Due to space limitations, we only discuss models that have made predictions for experiments or compared simulation results directly with experimental findings. Models that have been developed for the purpose of structure predictions²⁵, while important, are beyond the scope of this review.




Figure 1: Coarse-grained models for RNA. (a) Hierarchy of models at different resolution. In the TIS model, each nucleotide consists of three sites representing phosphate (red), sugar (yellow), and base (blue), whereas in the SIS model, a single bead represents an entire nucleotide. (b) Treatment of ions in the TIS model framework. *Left*: When the solvent contains only monovalent ions, implicit ion treatment based on Debye-Hückel theory is appropriate. *Middle*: RNA in a solution with monovalent salts and divalent cations such as Mg^{2+} and Ca^{2+} . *Right*: In the TIS-CAT model, divalent ions are treated explicitly while monovalent ions are implicit.

2 Single Interaction Site (SIS) Model

The Single Interaction Site (SIS) model represents each nucleotide by single interaction center¹⁹. The SIS model for folded RNA retains chain connectivity and favorable attractive interactions between sites that stabilize the native fold. The initial formulation of the SIS model was primarily used to provide a theoretical framework to predict the outcomes of single molecule pulling experiments in which mechanical forces were to probe the folding landscapes of RNA constructs and ribozymes^{26–28}. We first describe applications to force-driven conformational changes in a riboswitch to demonstrate the versatility of the model and set the stage for phase separation studies.

Response of Riboswitches to Mechanical Force: Riboswitches, consisting of an aptamer domain (AD) and an expression platform, contain RNA elements in the untranslated region of mRNA that allosterically regulate translation and transcription in bacteria by binding to metabolites with exquisite specificity^{29;30}. Often transcription and riboswitch folding are coupled^{31–33}. The functional selectivity is best illustrated using the purine riboswitches as examples. For example, adenine (*add*) riboswitch³⁴ activates translation upon binding the metabolite purine, whereas the structurally similar *pbuE* adenine riboswitch²⁸ activates transcription in the presence of purine. Both of these are ON riboswitches, which means that translation or transcription is activated only upon binding of the metabolite. In contrast, OFF riboswitches shut down gene expression when the metabolite binds to the aptamer domain. The SAM-III riboswitch binds S-adenosylmethionine (SAM) and represses translation upon binding SAM which sequesters the Shine-Dalgarno ribosome binding sequence. Single molecule force spectroscopy using optical tweezers, performed either by applying constant force to the ends of the RNA or by increasing the force at a constant rate, have been used to assess the characteristics of the intermediates in riboswitches^{28;35}. To complement these studies and to describe the structures of the intermediates that the folding landscape of riboswitches could be quantitatively mapped using SIS model^{36;37}.

In conventional single-molecular force spectroscopy, performed in the force ramp or force clamp mode, only states that are populated in the accessible force range can be characterized. Experiments performing force pulses during stretch-quench cycles could uncover the hidden states through which RNA folds. Let us start from a fully unfolded riboswitch, prepared at a high stretching force, f_s , and then quench the force to f_q that is less than the critical force needed to unfold for a time period t_q to initiate refolding. If t_q is less than the folding time RNA would only access partially structured state. By repeating the process of

stretching and quenching the force for varying time period (t_q) folding can be interrupted, and the states that are difficult to capture in a single or step wise quenching of force could be resolved. The SIS model simulations with the metabolite SAM bound showed that a spectrum of states was populated when t_q was varied. Importantly, the simulations showed³⁶ that the refolding time as a function of force is accurately predicted using calculated free energy profile and Kramers rate theory^{38;39}.

Phase separation in repeat RNA sequences: More recently, the SIS model has been extended to probe phase separation in low complexity RNA repeat sequences, which cause several neurological and neuromuscular disorders, such as the Huntington disease⁴⁰. *In vitro* experiments⁴¹ showed that the trinucleotide repeat sequences such as $(CAG)_n$ and $(CUG)_n$ undergo phase separation in which a dense phase coexists with a dilute phase. Importantly, phase separation occurs only if the number of repeats n exceeds a critical value. To determine the mechanism of droplet formation using the SIS model, we adjusted the only parameter in the SIS model, corresponding to the energy scale for base-pair interactions, to reproduce the known structures of a short $(CAG)_2$ duplex. Multichain simulations recapitulate the length and concentration dependence of the phase separation, in quantitative agreement with the *in vitro* experiments (Figure 2 a and b). The driving force for coacervate formation is the intermolecular base-pair interactions that has both an enthalpic and entropic contribution. Unexpectedly, it was found that once RNA molecules are recruited in the droplets, they undergo large conformational change, from a hairpin-like conformation in isolation to a stretched state, to form an extensive network of intermolecular interactions (Figure 2c). This soft network, in turn, constrains the RNA conformational fluctuation and mobility, which was rationalized using reptation dynamics⁴². The SIS model has been adopted to simulate a number of repeat RNA sequences using the LAMMPS, which apparently enhances the computational efficiency⁴³. Like the previous work, simulations of a number of low complexity RNA sequences showed that the enhanced number of ways of forming Watson-Crick pairs render stability to the dense droplets.

Based on the SIS model, a theory was advanced for anticipating the propensity of RNA to self-associate. It was discovered that the free energy gap separating the ground state and low lying excited states in repeat sequences are small²¹. As a result, such sequences can readily unfold, thus exposing a stretch of nucleotides that can engage in inter molecular base pairing^{44;45}, resulting in phase separation. In contrast, the free energy gap in heterogeneous sequences in which the 5' and 3' ends are in proximity are sufficiently great that the excited state has negligible population, which prevents unwarranted RNA-RNA interactions.

The single bead representation of RNA has recently been used to investigate phase separations in mixture

of RNA and proteins.^{20;46;47} In these models, interactions of RNA residues are parameterized to reproduce condensate formation with proteins, but the capability of such models to fold RNA into complex secondary and tertiary structures has not been demonstrated. Understanding the impact of RNA in influencing phase separation in proteins will also require a consistent model of RNA as well as interactions with ions.

3 Three Interaction Site (TIS) Model

In the Three Interaction Site (TIS) model each nucleotide consists of three distinct interaction sites – phosphate (P), sugar (S), and nucleobase (B) (Fig. 1). This allows the TIS model to capture the chemically distinct behavior of these moieties, which is essential for describing the sequence dependent structural dynamics. Since its first application in 2005¹⁰, the TIS model has been widely used to study RNA folding in a variety of contexts. Examples include RNA folding induced by temperature jump¹⁵ or mechanical forces⁴⁹, folding pathways shifting with changes in ion concentration on the folding pathways⁵⁰, and effects of molecular crowding⁵¹, and solvent viscosity⁵². The three-bead representation has also been widely adopted in other RNA models^{25;53} and to investigate various aspects of DNA biophysics^{54;55}.

Over the years dramatic improvements have been within the TIS model by incorporating the nearest neighbor thermodynamics into the TIS representation²². This was validated by quantitative comparison with experiments on RNA thermodynamics, such as melting of RNA hairpins and pseudoknots (PKs) (Figure 3a), which are essential building blocks of larger RNA molecules^{22;49}. Recent work has focused on developing methods to accurately describe electrostatic interactions associated with monovalent and divalent cations and phosphate groups (Figure 1b). In most CG models of RNA, electrostatic interactions are modeled using the DH potential, which is sufficient for monovalent ions. In fact, for example, the TIS model with DH interactions accurately captured that folding pathways of a PK as a function of monovalent ion concentration, in accord with experiments⁵⁰. However, it was a long-standing limitation of CG models that the effects of divalent cations cannot be described accurately using the DH theory. This limitation was overcome in an important study that proposed a way to include both monovalent (Na^+ , K^+ , Cl^-) and divalent ions (Mg^{2+} , Ca^{2+}) in the TIS model explicitly¹⁶. Simulations using this model captures the complicated ion-RNA electrostatic interactions, enabling visualization of how binding of individual cations facilitate self-assembly of RNA molecules.

In a series of papers we illustrated the success of the TIS model with explicit ions by quantitatively re-




Figure 2: RNA phase separation at the molecular level using coarse-grained simulations. (a, b) The single interaction site model reproduces droplet formation (a) that occurs only when the RNA with ~ 30 CAG repeats and RNA concentration exceeds critical concentration (b)⁴⁸. (c) Snapshots showing incorporation event of a $(CAG)_{47}$ molecule (red) into a droplet. The dramatic conformational change that occurs can be seen in the accompanying secondary structure diagrams. *Left*: Initially, the isolated $(CAG)_{47}$ chain formed a hairpin shape due to self-complementary base pairs. *Middle*: The RNA is unwound from one end to form new base pairs with other $(CAG)_{47}$ chains present on the surface of droplet. *Right*: Once incorporated into the droplet, each RNA chain tends to elongate as formation of inter-chain base pairs dominate due to entropic gain.

producing experiments on the folding of the *Azoarcus* group I intron ribozyme^{16;17;56}. In particular, the simulations reproduced the Mg^{2+} concentration dependence of folding¹⁶. Furthermore, specific Mg^{2+} binding sites agreed with the experimentally determined positions of Mg^{2+} ^{16;56} (Figure 3b). This allowed the authors to confidently discuss how individual Mg^{2+} ions navigate the formation of secondary and tertiary structural motifs in order, which cannot be traced experimentally. The model also captured a subtle difference between Mg^{2+} and Ca^{2+} . Ca^{2+} can drive RNA folding to some extent similar to Mg^{2+} , but not enough to reach its enzymatically active conformation due to the slightly larger size of the ionic radius.

It has been shown experimentally that RNA of a certain length with a complex fold often gets caught in kinetic traps and takes a long time to fold or misfold⁵⁷. The TIS simulation of the *Azoarcus* ribozyme has recently been extended to a kinetics study that showed how such multi-step folding and misfolding of the RNA occurs upon sudden changes in solution conditions, such as Mg^{2+} concentration jump¹⁷. By simulating a series of Mg^{2+} -induced folding events, it has been shown that the fate of the RNA molecules – whether they rapidly fold to the catalytically active state, get trapped in intermediate states, or even misfold – can be determined at a very early stage when the collapsing chain arrangement is driven by Mg^{2+} ion binding to specific sites (Figure 3c). Notably, the major misfolded state that appeared in the simulation was similar to misfolded structures of *Tetrahymena* ribozyme recently observed by cryo-electron microscopy⁵⁸, giving further confidence to the simulation model.

In an important application of the TIS model, ions (Mg^{2+} , K^+ , and F^-) were included explicitly to investigate the Mg^{2+} -dependent transition of a fluoride riboswitch⁵⁹ folding in the presence of K^+ and F^- ions⁶⁰. Kumar and Reddy investigated the folding of the aptamer domain (AD), which led them to propose a novel mechanism for the function of the fluoride riboswitch and F^- encapsulation by the cationic pocket composed of three Mg^{2+} ions. Folding of the AD induces sequential binding of the two Mg^{2+} ions to the phosphate groups driven in part by a transition from an outer to an inner-shell coordination through dehydration (Figure 4a). The third Mg^{2+} and F^- ion bind to the riboswitch in two steps. Binding of Mg^{2+} results in trigonal cationic pocket. Subsequently, Mg^{2+} and F^- form a water-mediated ion pair (Mg-water-F).

Let us highlight couple of additional applications of the TIS model. Simulations of the central domain of the bacterial ribosome, performed over a broad range of Mg^{2+} concentration, reproduced a number of experimental findings quantitatively (Figure 3d)⁶¹. Strikingly, the TIS model simulations predicted not only the multi-pathway folding of G-quadruplex folding but also characterized the structures of the intermediates, thus complementing the experimental results⁶². The handful of applications show that the combination of

TIS model simulations and experiments could be used to fully determine the folding of large RNAs.

4 TIS-CAT

A technical challenge in explicitly treating both divalent and monovalent ions in simulations is that it dramatically decreases the computational efficiency. To overcome this problem the TIS-CAT hybrid model was developed in which monovalent ions are implicitly treated using the DH potential but divalent cations are explicitly included (Figure 1b)²³. The TIS-CAT model is valid when the concentration of monovalent ions (typically >50 mM) is higher than the divalent ion concentration, which is normally in the millimolar range in the folding experiments. We developed a comprehensive theoretical model combining the Reference Interaction Site Model (RISM)⁶⁴⁻⁶⁶, grounded in liquid state theory, with coarse-grained molecular simulations. The RISM theory enables detailed calculations of the potential of mean force (PMF), which is distance and the ionic concentration-dependent effective potential between divalent ions and the phosphate group (Figure 4a). The isotropic PMF has two distinct minima: one for inner-sphere coordination (direct binding to the phosphate) and another for outer-sphere coordination (binding through water mediation). This dual-mode binding is crucial for accurate prediction of the thermodynamics of ion-induced RNA folding²³.

Simulations using the hybrid model demonstrated that divalent ions, particularly Mg^{2+} , have a profound effect on RNA folding through both inner- and outer-sphere coordination with phosphate groups. In smaller RNAs, such as the Beet Western Yellow Virus (BWYV) pseudoknot, Mg^{2+} predominantly binds via outer-sphere interactions, where the ion remains hydrated and interacts indirectly with RNA through water molecules. In contrast, in larger RNAs like the 58-nucleotide ribosomal RNA fragment (58-nt rRNA, Figure 4b), there is a significant change toward inner-sphere binding, where Mg^{2+} directly interacts with phosphate groups after partially losing its hydration shell. This change is due to the increased electrostatic potential in the more densely packed phosphate regions of larger RNA molecules, which necessitates stronger binding to stabilize the folded structure.

The simulations also highlighted the role of bridging interactions on RNA stability⁶⁷. These occur when a single divalent ion binds simultaneously to two or more phosphate groups, acting as a bridge that stabilizes the RNA's tertiary structure. Bridging interactions are particularly important in larger RNAs where multiple phosphate groups are in proximity. The simulations showed that reducing the concentration of divalent ions reduces the likelihood of the bridging interactions, leading to RNA destabilization. For example, in the 58-nt




Figure 3: Ion-driven thermodynamics and kinetics of RNA folding driven by ions. (a) Melting profile of BWYV pseudoknot measured in experiment (black)⁶³ at 500 mM NaCl and TIS simulation (red) with implicit monovalent salt at the same concentration. (b) Nucleotide position dependent ion fingerprints in the folding of *Azoarcus* ribozyme, represented by local concentrations of Mg^{2+} (blue) and K^+ (green). Comparisons with the Mg^{2+} binding sites resolved in X-ray crystal structure are shown. (c) *Left*: Kinetic partitioning mechanism in the folding of ribozyme. Differences in partial formations of secondary and tertiary structures at an early stage, driven by specific binding of Mg^{2+} , lead to dramatic difference in the fate – folding to the native structure or misfolding. A major source of misfolding is caused by topological frustration where the position of J8/7 strand (red) relative to neighboring helices are opposite to that in the native state. *Right*: The same mechanism of misfolding was recently observed in cryo-EM experiments of a similar ribozyme from *Tetrahymena*⁵⁸. The images were adopted from¹⁷ under the CC-BY license. (d) *Left*: Conformational transitions driven by Mg^{2+} in the central domain of bacterial ribosome. *Right*: Comparison of the high density spots of Mg^{2+} around the central junction obtained in the simulation (grey) with the experimentally observed Mg^{2+} positions (red).

rRNA, specific nucleotides were identified as key binding sites for Mg^{2+} , with significant accumulation of ions around them²³. These nucleotides are located at structurally important positions, and the Mg^{2+} ions in these regions are crucial for maintaining the integrity of the folded RNA. This finding is particularly important for understanding the stability of larger RNA molecules, such as ribozymes and ribosomal RNAs, where multiple regions of the RNA come into proximity during folding.

The charge density of divalent cations plays a crucial role in determining their coordination mode with RNA. The high charge density of Mg^{2+} favors outer-sphere binding in most small and intermediate-sized RNAs because it maintains a stable hydration shell. However, in regions of highly negative electrostatic potential, such as in larger RNA molecules or deep pockets of compact structures, Mg^{2+} can partially dehydrate and engage in inner-sphere coordination, as in the fluoride riboswitch. In contrast, Ca^{2+} , which has a lower charge density and thus weaker interaction with its hydration layer, dehydrates more readily and binds directly to phosphate groups in the inner-sphere coordination, even in smaller RNAs. This difference in binding behavior is important for understanding how different divalent ions influence RNA folding and stability. The findings suggest that Mg^{2+} binds RNA in multiple ways depending on the RNA size and structure, while Ca^{2+} is more rigid in its preference for direct phosphate binding.

Recently, single molecule FRET experiments and simulations using the TIS-CAT model were combined to detect three distinct states that are populated at different Mg^{2+} concentrations in guanidine-II riboswitch that controls translation (Figure 4c)¹⁸. When guanidine is bound (unbound) translation is turned on (suppressed)⁶⁹. Remarkably, the calculated distributions of FRET efficiency as a function Mg^{2+} using the TIS-CAT model without any adjustable parameter were in excellent agreement with experiments (Figure 4d)¹⁸. The TIS-CAT simulations were used to determine the structures of the states that are populated at three values of Mg^{2+} concentration. Note that unlike in guanidine-II riboswitch it is only upon binding to the metabolite is translation is suppressed in the SAM riboswitch. Nevertheless, the CG models accurately predict the outcomes of experiments, albeit using different resolution for the RNA.

5 Concluding Remarks

Although coarse-grained models for nucleic acids were first introduced only twenty-five years ago¹⁰, a great deal of progress has been made in the intervening years, spurred largely by advances in the experimental fronts (such as single molecule studies, cryo-EM, time-resolved SAXS experiments, and ion-counting




Figure 4: (a) The effective potential between phosphate and Mg^{2+} used in the TIS-CAT model (red) is a combination of the PMF derived from the RISM theory for the short range (grey dashed line) interaction and the Debye-Hückel theory for the long range (cyan) potential. There are two minima corresponding to the direct (dehydrated) and water-mediated binding of Mg^{2+} to the phosphate. (b) Preferential interaction coefficient, Γ , depending on the bulk Mg^{2+} concentrations as a function of monovalent salt concentration. Γ values calculated from TIS-CAT simulations (symbols)²³ are in excellent agreement with the experimental data (dashed curves)⁶⁸. (c) Three distinct conformations of guanidine-II riboswitch from TIS-CAT simulations at different Mg^{2+} concentrations (upper) and secondary structures indicating the positions of experimental FRET dyes (lower)¹⁸. (d) Comparison of smFRET efficiency distribution between the simulations and experiments.¹⁸ The images in (c and d) were taken from¹⁸ under the CC-BY license © 2022 Fuks, Falkner, Schwierz and Hengesbach.

experiments). While these developments, some of which are sketched here, are encouraging, frontiers in RNA biology are rapidly moving to problems that invariably require coming to grips with more interactions involving multiple proteins, DNA, and RNA. The prime example is ribosome assembly, which has been extensively studied experimentally⁷⁰⁻⁷², thus producing accurate data. Although there are combination of a variety of simulation methods and experiments have provided invaluable insights⁷³, it is fair to say that computational methods alone are currently not capable of tackling the dynamics of ribosome assembly or similarly complicated problems. We believe that innovations in this context are sorely needed.

Acknowledgments: This work was supported by a grant from the National Science Foundation (CHE 2320256) and the Welch Foundation through the Collie-Welch Chair (F-0019). NH was supported by JST PRESTO (JPMJPR22EA).

References

- [1] K. Karikó, M. Buckstein, H. Ni, and D. Weissman, “Suppression of RNA recognition by Toll-like receptors: the impact of nucleoside modification and the evolutionary origin of RNA,” Immunity, vol. 23, no. 2, pp. 165–175, 2005.
- [2] R. C. Lee, R. L. Feinbaum, and V. Ambros, “The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*,” Cell, vol. 75, no. 5, pp. 843–854, 1993.
- [3] B. Wightman, I. Ha, and G. Ruvkun, “Posttranscriptional regulation of the heterochronic gene *lin-14* by *lin-4* mediates temporal pattern formation in *C. elegans*,” Cell, vol. 75, no. 5, pp. 855–862, 1993.
- [4] T. R. Cech, A. J. Zaug, and P. J. Grabowski, “*In vitro* splicing of the ribosomal-RNA precursor of *Tetrahymena*: involvement of a quinosine nucleotide in the excision of the intervening sequence,” Cell, vol. 27, pp. 487–496, 1981.
- [5] C. Guerrier-Takada, K. Gardiner, T. Marsh, N. Pace, and S. Altman, “The RNA moiety of Ribonuclease-P is the catalytic subunit of the enzyme,” Cell, vol. 35, pp. 849–857, 1983.
- [6] J. A. Doudna and T. R. Cech, “The chemical repertoire of natural ribozymes,” Nature, vol. 418, no. 6894, pp. 222–228, 2002.
- [7] S. A. Woodson, “Structure and assembly of group I introns,” Curr. Opin. Struct. Biol., vol. 15, pp. 324–330, 2005.

- [8] S. A. Woodson, “Compact intermediates in RNA folding,” Annu. Rev. Biophys., vol. 39, no. 1, pp. 61–77, 2010.
- [9] D. Herschlag, S. Bonilla, and N. Bisaria, “The story of RNA folding, as told in epochs,” Cold Spring Harb. Perspect. Biol., vol. 10, no. 10, p. a032433, 2018.
- [10] C. Hyeon and D. Thirumalai, “Mechanical unfolding of RNA hairpins,” Proc. Natl. Acad. Sci. U.S.A., vol. 102, no. 19, pp. 6789–6794, 2005.
- [11] R. L. Hayes, J. K. Noel, A. Mandic, P. C. Whitford, K. Y. Sanbonmatsu, U. Mohanty, and J. N. Onuchic, “Generalized manning condensation model captures the RNA ion atmosphere,” Phys. Rev. Lett., vol. 114, no. 25, p. 258105, 2015.
- [12] R. L. Hayes, J. K. Noel, U. Mohanty, P. C. Whitford, S. P. Hennelly, J. N. Onuchic, and K. Y. Sanbonmatsu, “Magnesium fluctuations modulate RNA dynamics in the SAM-I riboswitch,” J. Am. Chem. Soc., vol. 134, no. 29, pp. 12043–12053, 2012.
- [13] M. J. Boniecki, G. Lach, W. K. Dawson, K. Tomala, P. Lukasz, T. Soltysinski, K. M. Rother, and J. M. Bujnicki, “SimRNA: a coarse-grained method for RNA folding simulations and 3D structure prediction,” Nucl. Acids Res., vol. 44, no. 7, pp. e63–e63, 2016.
- [14] Z. Xia, D. P. Gardner, R. R. Gutell, and P. Ren, “Coarse-grained model for simulation of RNA three-dimensional structures,” J. Phys. Chem. B, vol. 114, no. 42, pp. 13497–13506, 2010.
- [15] C. Hyeon and D. Thirumalai, “Multiple probes are required to explore and control the rugged energy landscape of rna hairpins,” J. Am. Chem. Soc., vol. 130, no. 5, pp. 1538–1539, 2008.
- [16] N. A. Denesyuk and D. Thirumalai, “How do metal ions direct ribozyme folding?,” Nat. Chem., vol. 7, no. 10, pp. 793–801, 2015.
- [17] N. Hori and D. Thirumalai, “Watching ion-driven kinetics of ribozyme folding and misfolding caused by energetic and topological frustration one molecule at a time,” Nucl. Acids Res., vol. 51, no. 19, pp. 10737–10751, 2023.
- [18] C. Fuks, S. Falkner, N. Schwierz, and M. Hengesbach, “Combining Coarse-Grained Simulations and Single Molecule Analysis Reveals a Three-State Folding Model of the Guanidine-II Riboswitch,” Front. Mol. Biosci., vol. 9, p. 826505, 2022.
- [19] C. Hyeon, R. I. Dima, and D. Thirumalai, “Pathways and kinetic barriers in mechanical unfolding and refolding of RNA and proteins,” Structure, vol. 14, no. 11, pp. 1633–1645, 2006.
- [20] R. M. Regy, G. L. Dignon, W. Zheng, Y. C. Kim, and J. Mittal, “Sequence dependent phase separation

- of protein-polynucleotide mixtures elucidated using molecular simulations,” Nucl. Acids Res., vol. 48, no. 22, pp. gkaa1099–, 2020.
- [21] H. Maity, H. T. Nguyen, N. Hori, and D. Thirumalai, “Odd–even disparity in the population of slipped hairpins in RNA repeat sequences with implications for phase separation,” Proc. Natl. Acad. Sci. U.S.A., vol. 120, no. 24, p. e2301409120, 2023.
- [22] N. A. Denesyuk and D. Thirumalai, “Coarse-grained model for predicting RNA folding thermodynamics,” J. Phys. Chem. B, vol. 117, no. 17, pp. 4901–4911, 2013.
- [23] H. T. Nguyen, N. Hori, and D. Thirumalai, “Theory and simulations for RNA folding in mixtures of monovalent and divalent cations,” Proc. Natl. Acad. Sci. U.S.A., vol. 116, no. 42, pp. 21022–21030, 2019.
- [24] J. J. Uusitalo, H. I. Ingólfsson, S. J. Marrink, and I. Faustino, “Martini coarse-grained force field: extension to RNA,” Biophys. J., vol. 113, no. 2, pp. 246–256, 2017.
- [25] J. Li and S.-J. Chen, “RNA 3D structure prediction using coarse-grained models,” Front. Mol. Biosci., vol. 8, p. 720937, 2021.
- [26] M. T. Woodside, P. C. Anthony, W. M. Behnke-Parks, K. Larizadeh, D. Herschlag, and S. M. Block, “Direct measurement of the full, sequence-dependent folding landscape of a nucleic acid,” Science, vol. 314, no. 5801, pp. 1001–1004, 2006.
- [27] M. T. Woodside and S. M. Block, “Reconstructing folding energy landscapes by single-molecule force spectroscopy,” Annu. Rev. Biophys., vol. 43, no. 1, pp. 19–39, 2014.
- [28] W. J. Greenleaf, K. L. Frieda, D. A. Foster, M. T. Woodside, and S. M. Block, “Direct observation of hierarchical folding in single riboswitch aptamers,” Science, vol. 319, no. 5863, pp. 630–633, 2008.
- [29] A. Serganov and E. Nudler, “A decade of riboswitches,” Cell, vol. 152, no. 1, pp. 17–24, 2013.
- [30] W. C. Winkler and R. R. Breaker, “Regulation of bacterial gene expression by riboswitches,” Annu. Rev. Microbiol., vol. 59, pp. 487–517, 2005.
- [31] K. E. Watters, E. J. Strobel, A. M. Yu, J. T. Lis, and J. B. Lucks, “Cotranscriptional folding of a riboswitch at nucleotide resolution,” Nat. Struct. Mol. Biol., vol. 23, no. 12, pp. 1124–1131, 2016.
- [32] T.-t. Sun, C. Zhao, and S.-J. Chen, “Predicting cotranscriptional folding kinetics for riboswitch,” J. Phys. Chem. B, vol. 122, no. 30, pp. 7484–7496, 2018.
- [33] H. Uhm, W. Kang, K. S. Ha, C. Kang, and S. Hohng, “Single-molecule FRET studies on the cotran-

- scriptional folding of a thiamine pyrophosphate riboswitch,” Proc. Natl. Acad. Sci. U.S.A., vol. 115, no. 2, pp. 331–336, 2018.
- [34] A. Serganov, Y.-R. Yuan, O. Pikovskaya, A. Polonskaia, L. Malinina, A. T. Phan, C. Hobartner, R. Micura, R. R. Breaker, and D. J. Patel, “Structural basis for discriminative regulation of gene expression by adenine-and guanine-sensing mRNAs,” Chem. Biol., vol. 11, no. 12, pp. 1729–1741, 2004.
- [35] K. L. Frieda and S. M. Block, “Direct observation of cotranscriptional folding in an adenine riboswitch,” Science, vol. 338, no. 6105, pp. 397–400, 2012.
- [36] J.-C. Lin and D. Thirumalai, “Kinetics of allosteric transitions in s-adenosylmethionine riboswitch are accurately predicted from the folding landscape,” J. Am. Chem. Soc., vol. 135, no. 44, pp. 16641–16650, 2013.
- [37] J.-C. Lin and D. Thirumalai, “Relative Stability of Helices Determines the Folding Landscape of Adenine Riboswitch Aptamers,” J. Am. Chem. Soc., vol. 130, no. 43, pp. 14080–14081, 2008.
- [38] H. A. Kramers, “Brownian motion a field of force and the diffusion model of chemical reaction,” Physica, vol. 7, pp. 284–304, 1940.
- [39] P. Hanggi, P. Talkner, and M. Borkovec, “Reaction-rate theory: fifty years after Kramers,” Rev. Mod. Phys., vol. 62, pp. 251–341, 1990.
- [40] A. R. La Spada and J. P. Taylor, “Repeat expansion disease: progress and puzzles in disease pathogenesis,” Nat. Rev. Genet., vol. 11, no. 4, pp. 247–258, 2010.
- [41] A. Jain and R. D. Vale, “RNA phase transitions in repeat expansion disorders,” Nature, vol. 546, no. 7657, pp. 243–247, 2017.
- [42] P. G. de Gennes, “Reptation of a Polymer Chain in the Presence of Fixed Obstacles,” J. Chem. Phys., vol. 55, no. 2, pp. 572–579, 1971.
- [43] D. Aierken and J. A. Joseph, “Accelerated simulations reveal physicochemical factors governing stability and composition of RNA clusters,” J. Chem. Theory Comput., 2024. (in press).
- [44] C. Roden and A. S. Gladfelter, “RNA contributions to the form and function of biomolecular condensates,” Nat. Rev. Mol. Cell Biol., vol. 22, no. 3, pp. 183–195, 2021.
- [45] E. M. Langdon and A. S. Gladfelter, “A new lens for RNA localization: Liquid-liquid phase separation,” Annu. Rev. Microbiol., vol. 72, pp. 255–271, 2018.
- [46] J. A. Joseph, A. Reinhardt, A. Aguirre, P. Y. Chew, K. O. Russell, J. R. Espinosa, A. Garaizar, and

- R. Collepardo-Guevara, “Physics-driven coarse-grained model for biomolecular phase separation with near-quantitative accuracy,” Nat. Comput. Sci., vol. 1, no. 11, pp. 732–743, 2021.
- [47] G. Valdes-Garcia, L. Heo, L. J. Lapidus, and M. Feig, “Modeling concentration-dependent phase separation processes involving peptides and RNA via residue-based coarse-graining,” J. Chem. Theory Comput., vol. 19, no. 2, pp. 669–678, 2023.
- [48] H. T. Nguyen, N. Hori, and D. Thirumalai, “Condensates in RNA repeat sequences are heterogeneously organized and exhibit reptation dynamics,” Nat. Chem., vol. 14, no. 7, pp. 775–785, 2022.
- [49] N. Hori, N. A. Denesyuk, and D. Thirumalai, “Salt effects on the thermodynamics of a frameshifting RNA pseudoknot under tension,” J. Mol. Biol., vol. 428, no. 14, pp. 2847–2859, 2016.
- [50] J. Roca, N. Hori, S. Baral, Y. Velmurugu, R. Narayanan, P. Narayanan, D. Thirumalai, and A. Ansari, “Monovalent ions modulate the flux through multiple folding pathways of an rna pseudoknot,” Proc. Natl. Acad. Sci. U.S.A., vol. 115, no. 31, pp. E7313–E7322, 2018.
- [51] N. A. Denesyuk and D. Thirumalai, “Crowding promotes the switch from hairpin to pseudoknot conformation in human telomerase RNA,” J. Am. Chem. Soc., vol. 133, no. 31, pp. 11858–11861, 2011.
- [52] N. Hori, N. A. Denesyuk, and D. Thirumalai, “Frictional effects on RNA folding: Speed limit and Kramers turnover,” J. Phys. Chem. B, vol. 122, no. 49, pp. 11279–11288, 2018.
- [53] C. Tan, J. Jung, C. Kobayashi, D. U. L. Torre, S. Takada, and Y. Sugita, “Implementation of residue-level coarse-grained models in GENESIS for large-scale molecular dynamics simulations,” PLoS Comput. Biol., vol. 18, no. 4, p. e1009578, 2022.
- [54] T. A. Knotts, IV, N. Rathore, D. C. Schwartz, and J. J. de Pablo, “A coarse grain model for DNA,” J. Chem. Phys., vol. 126, no. 8, pp. 084901 – 13, 2007.
- [55] D. Chakraborty, N. Hori, and D. Thirumalai, “Sequence-dependent three interaction site model for single- and double-stranded DNA,” J. Chem. Theory Comput., vol. 14, no. 7, pp. 3763–3779, 2018.
- [56] N. Hori, N. A. Denesyuk, and D. Thirumalai, “Ion condensation onto ribozyme is site specific and fold dependent,” Biophys. J., vol. 116, no. 12, pp. 2400–2410, 2019.
- [57] D. Thirumalai and S. A. Woodson, “Kinetics of folding of proteins and RNA,” Acc. Chem. Res., vol. 29, no. 9, pp. 433–439, 1996.
- [58] S. L. Bonilla and K. Jang, “Challenges, advances, and opportunities in RNA structural biology by cryo-EM,” Curr. Opin. Struct. Biol., vol. 88, p. 102894, 2024.
- [59] L. M. Hertz, E. N. White, K. Kuznedelov, L. Cheng, A. M. Yu, R. Kakkaramadam, K. Severinov,

- A. Chen, and J. B. Lucks, “The effect of pseudoknot base pairing on cotranscriptional structural switching of the fluoride riboswitch,” Nucl. Acids Res., vol. 52, no. 8, pp. 4466–4482, 2024.
- [60] S. Kumar and G. Reddy, “Mechanism of fluoride ion encapsulation by magnesium ions in a bacterial riboswitch,” J. Phys. Chem. B, vol. 127, no. 43, pp. 9267–9281, 2023.
- [61] N. Hori, N. A. Denesyuk, and D. Thirumalai, “Shape changes and cooperativity in the folding of the central domain of the 16S ribosomal RNA,” Proc. Natl. Acad. Sci. U.S.A., vol. 118, no. 10, p. e2020837118, 2021.
- [62] M. Ugrina, I. Burkhart, D. Muller, H. Schwalbe, and N. Schwierz, “RNA G-quadruplex folding is a multi-pathway process driven by conformational entropy,” Nucl. Acids Res., vol. 52, no. 1, pp. 87–100, 2024.
- [63] A. M. Soto, V. Misra, and D. E. Draper, “Tertiary structure of an RNA pseudoknot is stabilized by “diffuse” Mg^{2+} ions,” Biochemistry, vol. 46, no. 11, pp. 2973 – 2983, 2007.
- [64] D. Chandler and H. C. Andersen, “Optimized cluster expansions for classical fluids. ii. theory of molecular liquids,” J. Chem. Phys., vol. 57, no. 5, pp. 1930–1937, 1972.
- [65] F. Hirata and P. J. Rossky, “An extended rism equation for molecular polar fluids,” Chem. Phys. Lett., vol. 83, no. 2, pp. 329–334, 1981.
- [66] F. Hirata, B. M. Pettitt, and P. J. Rossky, “Application of an extended rism equation to dipolar and quadrupolar fluids,” J. Chem. Phys., vol. 77, no. 1, pp. 509–520, 1982.
- [67] H. T. Nguyen and D. Thirumalai, “Charge density of cation determines inner versus outer shell coordination to phosphate in RNA,” J. Phys. Chem. B, vol. 124, no. 20, pp. 4114–4122, 2020.
- [68] D. Grilley, V. Misra, G. Caliskan, and D. E. Draper, “Importance of partially unfolded conformations for Mg^{2+} -induced folding of RNA tertiary structure: Structural models and free energies of Mg^{2+} interactions,” Biochemistry, vol. 46, no. 36, pp. 10266–10278, 2007.
- [69] M. E. Sherlock, S. N. Malkowski, and R. R. Breaker, “Biochemical validation of a second guanine riboswitch class in bacteria,” Biochemistry, vol. 56, no. 2, pp. 352–358, 2017.
- [70] M. L. Rodgers and S. A. Woodson, “A roadmap for rRNA folding and assembly during transcription,” Trends Biochem. Sci., vol. 46, no. 11, pp. 889–901, 2021.
- [71] Z. Shajani, M. T. Sykes, and J. R. Williamson, “Assembly of bacterial ribosomes,” Annu. Rev. Biochem., vol. 80, no. 1, pp. 501–526, 2011.

- [72] O. Duss, G. A. Stepanyuk, J. D. Puglisi, and J. R. Williamson, “Transient protein-RNA interactions guide nascent ribosomal RNA folding,” Cell, vol. 179, no. 6, pp. 1357–1369, 2019.
- [73] L. V. Bock, S. Gabrielli, M. H. Kolář, and H. Grubmüller, “Simulation of complex biomolecular systems: the ribosome challenge,” Annu. Rev. Biophys., vol. 52, no. 1, pp. 361–390, 2023.