

# Presentations for monoids of endomorphisms of a star graph

Vítor H. Fernandes\*, Jörg Koppitz and T. Musunthia

January 3, 2025

## Abstract

In this paper, we consider the monoids of all endomorphisms, of all weak endomorphisms, of all strong endomorphisms and of all strong weak endomorphisms of a star graph with a finite number of vertices. Our main objective is to exhibit a presentation for each of them.

2020 *Mathematics subject classification*: 20M20, 20M05, 05C12, 05C25.

*Keywords*: transformations, partial endomorphisms, star graphs, presentations.

## Introduction

For a set  $\Omega$ , denote by  $\mathcal{PT}(\Omega)$  the monoid (under composition) of all partial transformations on  $\Omega$ , by  $\mathcal{T}(\Omega)$  the submonoid of  $\mathcal{PT}(\Omega)$  of all full transformations on  $\Omega$  and by  $\mathcal{S}(\Omega)$  the *symmetric group* on  $\Omega$ , i.e. the subgroup of  $\mathcal{PT}(\Omega)$  of all permutations on  $\Omega$ .

Let  $G = (V, E)$  be a simple graph (i.e. undirected, without loops and without multiple edges). Let  $\alpha$  be a full transformation of  $V$ . We say that  $\alpha$  is:

- an *endomorphism* of  $G$  if  $\{u, v\} \in E$  implies  $\{u\alpha, v\alpha\} \in E$ , for all  $u, v \in V$ ;
- a *weak endomorphism* of  $G$  if  $\{u, v\} \in E$  and  $u\alpha \neq v\alpha$  imply  $\{u\alpha, v\alpha\} \in E$ , for all  $u, v \in V$ ;
- a *strong endomorphism* of  $G$  if  $\{u, v\} \in E$  if and only if  $\{u\alpha, v\alpha\} \in E$ , for all  $u, v \in V$ ;
- a *strong weak endomorphism* of  $G$  if  $\{u, v\} \in E$  and  $u\alpha \neq v\alpha$  if and only if  $\{u\alpha, v\alpha\} \in E$ , for all  $u, v \in V$ ;
- an *automorphism* of  $G$  if  $\alpha$  is a bijective strong endomorphism (i.e.  $\alpha$  is bijective and  $\alpha$  and  $\alpha^{-1}$  are both endomorphisms). For finite graphs, any bijective weak endomorphism is an automorphism.


Denote by:

- $\text{End}G$  the set of all endomorphisms of  $G$ ;
- $\text{wEnd}G$  the set of all weak endomorphisms of  $G$ ;
- $\text{sEnd}G$  the set of all strong endomorphisms of  $G$ ;
- $\text{swEnd}G$  the set of all strong weak endomorphisms of  $G$ ;
- $\text{Aut}G$  the set of all automorphisms of  $G$ .

---

\*This work is funded by national funds through the FCT - Fundação para a Ciência e a Tecnologia, I.P., under the scope of the (Center for Mathematics and Applications) projects UIDB/00297/2020 (<https://doi.org/10.54499/UIDB/00297/2020>) and UIDP/00297/2020 (<https://doi.org/10.54499/UIDP/00297/2020>).

Clearly,  $\text{End}G$ ,  $\text{wEnd}G$ ,  $\text{sEnd}G$ ,  $\text{swEnd}G$  and  $\text{Aut}G$  are monoids under composition of mappings with the identity mapping  $\text{id}$  as the identity element. Moreover,  $\text{Aut}G$  is also a group. It is also clear that  $\text{Aut}G \subseteq \text{sEnd}G \subseteq \text{End}G$ ,  $\text{swEnd}G \subseteq \text{wEnd}G$




(these inclusions may not be strict).

Monoids of endomorphisms of graphs have many important applications, particularly related to automata theory; see [21]. A large number of interesting results concerning graphs and algebraic properties of their endomorphism monoids have been obtained by several authors (see, for example, [3, 10, 17, 18, 22, 23, 25, 30]). In recent years, the first two authors together with Dimitrova and Quinteiro, also have studied such types of monoids, in particular by considering finite undirected paths and cycles; see [6, 7, 8].

For any non negative integer  $n$ , let  $\Omega_n = \{1, 2, \dots, n\}$  and  $\Omega_n^0 = \{0, 1, \dots, n\} = \{0\} \cup \Omega_n$ . Notice that,  $\Omega_0 = \emptyset$  and  $\Omega_0^0 = \{0\}$ . For an integer  $n \geq 1$ , consider the *star graph*

$$S_n = (\Omega_{n-1}^0, \{\{0, i\} \mid 1 \leq i \leq n-1\})$$

with  $n$  vertices.



These very basic graphs, which are special examples of *trees* and also of *complete bipartite graphs*, play a significant role in Graph Theory, for instance, through the notions of *star chromatic number* and *star arboricity*. We may also find important applications of star graphs in Computer Science. For example, in Distributed Computing the *star network* is one of the most common computer network topologies.

In 2023, Fernandes and Paulista [14] considered the monoid of all partial isometries of the star graph  $S_n$ . They determined the rank (i.e. the minimum size of a generating set) and the cardinality of this monoid as well as described its Green's relations and exhibited a presentation. On the other hand, recently, the first two authors together with Dimitrova considered in [4, 5] the *partial counterparts* of the monoids  $\text{sEnd}(S_n)$ ,  $\text{swEnd}(S_n)$ ,  $\text{End}(S_n)$  and  $\text{wEnd}(S_n)$ , studying their regularity, describing their Green's relations and computing their cardinalities and ranks.

In this paper, our main aim is to determine presentations for the monoids  $\text{End}S_n$ ,  $\text{swEnd}S_n$  and  $\text{wEnd}S_n$ , which we will carry out in Section 3. In the first section, we will look at some basic concepts and results about presentations. In Section 2, we describe the monoids  $\text{Aut}S_n$ ,  $\text{sEnd}S_n$ ,  $\text{End}S_n$ ,  $\text{swEnd}S_n$  and  $\text{wEnd}S_n$ , and recall some of their unpublished properties, namely cardinalities, regularity, generators and ranks, which were studied by the first author together with his undergraduate students Duarte Grilo and Samuel Medalha within the *Undergraduate Research Opportunity Program* of NOVA School of Science and Technology. For the sake of completeness, we present here brief proofs of these results.

For general background on Semigroup Theory and standard notations, we would like to refer the reader to Howie's book [19]. Regarding Algebraic Graph Theory, we refer to Knauer's book [22]. We would also like to point out that we made use of computational tools, namely GAP [16].

# 1 On presentations

Let  $X$  be a set and denote by  $X^*$  the free monoid generated by  $X$ . Often, in this context, the set  $X$  is called an *alphabet* and its elements are called *words*. A *monoid presentation* is an ordered pair  $\langle X \mid R \rangle$ , where  $X$  is an alphabet and  $R$  is a subset of  $X^* \times X^*$ . An element  $(u, v)$  of  $X^* \times X^*$  is called a *relation* of  $X^*$  and it is usually represented by  $u = v$ . A monoid  $M$  is said to be *defined by a presentation*  $\langle X \mid R \rangle$  if  $M$  is isomorphic to  $X^*/\sim_R$ , where  $\sim_R$  denotes the congruence on  $X^*$  generated by  $R$ , i.e.  $\sim_R$  is the smallest congruence on  $X^*$  containing  $R$ . Suppose that  $X$  is a generating set of a monoid  $M$  and let  $u = v$  be a relation of  $X^*$ . We say that a relation  $u = v$  of  $X^*$  is *satisfied* by  $X$  if  $u = v$  is an equality in  $M$ . For more details, see [24] or [29].

A well-known direct method to obtain a presentation for a monoid, that we might consider folklore, is given by the following result; see, for example, [29, Proposition 1.2.3].

**Proposition 1.1.** *Let  $M$  be a monoid generated by a set  $X$ . Then,  $\langle X \mid R \rangle$  is a presentation for  $M$  if and only if the following two conditions are satisfied:*

1. *The generating set  $X$  of  $M$  satisfies all the relations from  $R$ ;*
2. *If  $w_1, w_2 \in X^*$  are any two words such that the generating set  $X$  of  $M$  satisfies the relation  $w_1 = w_2$ , then  $w_1 \sim_R w_2$ .*

For finite monoids, a standard method to find a presentation is described by the following result, adapted for the monoid case from [29, Proposition 3.2.2].

**Theorem 1.2** (Guess and Prove method). *Let  $M$  be a finite monoid, let  $X$  be a generating set for  $M$ , let  $R \subseteq X^* \times X^*$  be a set of relations and let  $W \subseteq X^*$ . Assume that the following conditions are satisfied:*

1. *The generating set  $X$  of  $M$  satisfies all the relations from  $R$ ;*
2. *For each word  $w \in X^*$ , there exists a word  $w' \in W$  such that  $w \sim_R w'$ ;*
3.  $|W| \leq |M|$ .

*Then,  $M$  is defined by the presentation  $\langle X \mid R \rangle$ .*

Notice that, if  $W$  satisfies the above conditions, then, in fact,  $|W| = |M|$ .

Let  $X$  be an alphabet, let  $R \subseteq X^* \times X^*$  be a set of relations and let  $W$  be a subset of  $X^*$ . We say that  $W$  is a set of *canonical forms* for a finite monoid  $M$  if Conditions 2 and 3 of Theorem 1.2 are satisfied. Suppose that the empty word belongs to  $W$  and, for each letter  $x \in X$  and for each word  $w \in W$ , there exists a word  $w' \in W$  such that  $wx \sim_R w'$ . Then, it is easy to show that  $W$  satisfies Condition 2.

In this paper, we will mainly use the method given by Theorem 1.2 and the condition described above instead of Condition 2.

A presentation for the symmetric group  $\mathcal{S}(\Omega_n)$  was determined by Moore [26] by the end of the 19th century (1897). Nearly six decades later (1958), Aizenštāt [1] gave a presentation for full transformation monoid  $\mathcal{T}(\Omega_n)$ . A few years later (1961), presentations for the partial transformation monoid  $\mathcal{PT}(\Omega_n)$  and for the symmetric inverse monoid  $\mathcal{I}(\Omega_n)$  were found by Popova [27]. In 1962, Aizenštāt [2] and Popova [28] exhibited presentations for the monoids of all order-preserving transformations and of all order-preserving partial transformations of a finite chain, respectively, and from the sixties until our days several authors obtained presentations for many classes of monoids. Further examples can be found, for instance, in [9, 11, 12, 13, 15, 20, 29].

We finish this section by recalling the presentations of  $\mathcal{S}(\Omega_n)$ ,  $\mathcal{T}(\Omega_n)$  and  $\mathcal{PT}(\Omega_n)$  aforementioned; see [12].

Let  $n \geq 3$  and  $a = \begin{pmatrix} 1 & 2 & 3 & \cdots & n \\ 2 & 1 & 3 & \cdots & n \end{pmatrix}$ ,  $b = \begin{pmatrix} 1 & 2 & \cdots & n-1 & n \\ 2 & 3 & \cdots & n & 1 \end{pmatrix}$ ,  $c = \begin{pmatrix} 2 & 3 & \cdots & n \\ 2 & 3 & \cdots & n \end{pmatrix}$  and  $e = \begin{pmatrix} 1 & 2 & 3 & \cdots & n \\ 1 & 1 & 3 & \cdots & n \end{pmatrix}$ . Then, we have: for  $n \geq 3$ ,

$$\mathcal{S}(\Omega_n) = \langle a, b \mid a^2 = b^n = (ba)^{n-1} = (ab^{n-1}ab)^3 = (ab^{n-j}ab^j)^2 = 1 \ (2 \leq j \leq n-2); \rangle$$

for  $n \geq 4$ ,

$$\begin{aligned} \mathcal{T}(\Omega_n) = \langle a, b, e \mid a^2 = b^n = (ba)^{n-1} = (ab^{n-1}ab)^3 = (ab^{n-j}ab^j)^2 = 1 \ (2 \leq j \leq n-2), \\ ae = b^{n-2}ab^2eb^{n-2}ab^2 = bab^{n-1}abeb^{n-1}abab^{n-1} = (ebab^{n-1})^2 = e, \\ (b^{n-1}abe)^2 = eb^{n-1}abe = (eb^{n-1}ab)^2, (ebab^{n-2}ab)^2 = (bab^{n-2}abe)^2 \rangle \end{aligned}$$

and

$$\begin{aligned} \mathcal{PT}(\Omega_n) = \langle a, b, c, e \mid a^2 = b^n = (ba)^{n-1} = (ab^{n-1}ab)^3 = (ab^{n-j}ab^j)^2 = 1 \ (2 \leq j \leq n-2), \\ b^{n-1}abcb^{n-1}ab = bacab^{n-1} = c = c^2, (ca)^2 = cac = (ac)^2, \\ ae = b^{n-2}ab^2eb^{n-2}ab^2 = bab^{n-1}abeb^{n-1}abab^{n-1} = (ebab^{n-1})^2 = e, \\ (b^{n-1}abe)^2 = eb^{n-1}abe = (eb^{n-1}ab)^2, (ebab^{n-2}ab)^2 = (bab^{n-2}abe)^2, \\ ec = cac, ce = ca, eac = ea, eab^{n-1}abac = ab^{n-1}abacab^{n-1}abaeab^{n-1}aba \rangle; \end{aligned}$$

for  $n = 3$ ,

$$\begin{aligned} \mathcal{T}(\Omega_3) = \langle a, b, e \mid a^2 = b^3 = (ba)^2 = (ab^2ab)^3 = 1, ae = bab^2abeb^2abab^2 = (ebab^2)^2 = e, \\ (b^2abe)^2 = eb^2abe = (eb^2ab)^2 \rangle \end{aligned}$$

and

$$\begin{aligned} \mathcal{PT}(\Omega_3) = \langle a, b, c, e \mid a^2 = b^3 = (ba)^2 = (ab^2ab)^3 = 1, b^2abcb^2ab = bacab^2 = c = c^2, (ca)^2 = cac = (ac)^2, \\ ae = bab^2abeb^2abab^2 = (ebab^2)^2 = e, (b^2abe)^2 = eb^2abe = (eb^2ab)^2, \\ ec = cac, ce = ca, eac = ea, eab^2abac = ab^2abacab^2abaeab^2aba \rangle. \end{aligned}$$

## 2 Descriptions, regularity and ranks

Let  $\mathcal{PT}_{n-1}^0 = \{\alpha \in \mathcal{T}(\Omega_{n-1}^0) \mid 0\alpha = 0\}$ ,  $\mathcal{T}_{n-1}^0 = \{\alpha \in \mathcal{T}(\Omega_{n-1}^0) \mid 0\alpha = 0 \text{ and } \Omega_{n-1}\alpha \subseteq \Omega_{n-1}\}$  and let  $\mathcal{S}_{n-1}^0 = \mathcal{S}(\Omega_{n-1}^0) \cap \mathcal{T}_{n-1}^0$ . Then, it is clear that  $\mathcal{PT}_0^0 = \mathcal{T}_0^0 = \mathcal{T}(\Omega_0^0) = \{(\begin{smallmatrix} 0 \\ 0 \end{smallmatrix})\} = \mathcal{S}(\Omega_0^0) = \mathcal{S}_0^0$  and, for  $n \geq 2$ ,  $\mathcal{PT}_{n-1}^0$  is a submonoid of  $\mathcal{T}(\Omega_{n-1}^0)$  isomorphic to  $\mathcal{PT}(\Omega_{n-1})$ ,  $\mathcal{T}_{n-1}^0$  is a submonoid of  $\mathcal{T}(\Omega_{n-1}^0)$  isomorphic to  $\mathcal{T}(\Omega_{n-1})$  and  $\mathcal{S}_{n-1}^0$  is a subgroup of  $\mathcal{S}(\Omega_{n-1}^0)$  isomorphic to  $\mathcal{S}(\Omega_{n-1})$ . Moreover, for  $n \geq 1$ , it is a routine matter to check that:

- $\text{sEnd}S_n = \text{End}S_n = \mathcal{T}_{n-1}^0 \cup \{(\begin{smallmatrix} 0 & 1 & \cdots & n-1 \\ i & 0 & \cdots & 0 \end{smallmatrix}) \mid i \in \Omega_{n-1}\}$ ;
- $\text{swEnd}S_n = \text{End}S_n \cup \{(\begin{smallmatrix} 0 & 1 & \cdots & n-1 \\ i & i & \cdots & i \end{smallmatrix}) \mid i \in \Omega_{n-1}\}$ ;
- $\text{wEnd}S_n = \{\alpha \in \mathcal{T}(\Omega_{n-1}^0) \mid 0\alpha \neq 0 \text{ implies } \text{Im } \alpha \subseteq \{0, 0\alpha\}\}$ .

Therefore, we have:

- $|\text{End}S_n| = (n-1)^{n-1} + n - 1$ , for  $n \geq 1$ ;
- $|\text{swEnd}S_n| = (n-1)^{n-1} + 2n - 1$ , for  $n \geq 2$ ;
- $|\text{wEnd}S_n| = n^{n-1} + (n-1)2^{n-1}$ , for  $n \geq 1$ .

Observe that,  $\text{Aut}S_1 = \text{End}S_1 = \text{swEnd}S_1 = \text{wEnd}S_1 = \mathcal{T}(\Omega_0^0) = \{(\begin{smallmatrix} 0 \\ 0 \end{smallmatrix})\}$ ,  $\text{Aut}S_2 = \text{End}S_2 = \mathcal{S}(\Omega_1^0) = \{(\begin{smallmatrix} 0 & 1 \\ 0 & 1 \end{smallmatrix}), (\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix})\}$  and  $\text{swEnd}S_2 = \text{wEnd}S_2 = \mathcal{T}(\Omega_1^0) = \{(\begin{smallmatrix} 0 & 1 \\ 0 & 1 \end{smallmatrix}), (\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}), (\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}), (\begin{smallmatrix} 0 & 1 \\ 1 & 1 \end{smallmatrix})\}$ . Furthermore, as the automorphisms of a graph are its bijective endomorphisms, for  $n \geq 3$ , we have

- $\text{Aut}S_n = \mathcal{S}_{n-1}^0$  and so  $|\text{Aut}S_n| = (n-1)!$ .

It is obvious that, for  $n = 1, 2$ , the monoids  $\text{End}S_n$ ,  $\text{swEnd}S_n$  and  $\text{wEnd}S_n$  are all regular. In fact, it is easy to show that this is the case for any  $n \geq 1$ :

**Theorem 2.1.** For  $n \geq 1$ , the monoids  $\text{End}S_n$ ,  $\text{swEnd}S_n$  and  $\text{wEnd}S_n$  are all regular.

*Proof.* By the previous observation, we may consider  $n \geq 3$ . As  $\mathcal{T}_{n-1}^0$  is a submonoid of  $\mathcal{T}(\Omega_{n-1}^0)$  isomorphic to  $\mathcal{T}(\Omega_{n-1})$  and it is well known that this last monoid is regular, each element of  $\mathcal{T}_{n-1}^0$  is regular in  $\mathcal{T}_{n-1}^0$  and so also regular in  $\text{End}S_n$ ,  $\text{swEnd}S_n$  and  $\text{wEnd}S_n$ .

Let  $\alpha_i = \begin{pmatrix} 0 & 1 & \cdots & n-1 \\ i & 0 & \cdots & 0 \end{pmatrix}$  for  $1 \leq i \leq n-1$ . Then,  $\alpha_i^3 = \alpha_i$ , whence  $\alpha_i$  is regular in  $\text{End}S_n$  and so  $\alpha_i$  is also regular in  $\text{swEnd}S_n$  and  $\text{wEnd}S_n$ , for all  $1 \leq i \leq n-1$ . We can conclude right away that  $\text{End}S_n$  is a regular monoid.

Let  $\varepsilon_i = \begin{pmatrix} 0 & 1 & \cdots & n-1 \\ i & i & \cdots & i \end{pmatrix}$  for  $0 \leq i \leq n-1$ . Then,  $\varepsilon_i$  is an idempotent, whence  $\varepsilon_i$  is regular in  $\text{swEnd}S_n$  and so  $\varepsilon_i$  is also regular in  $\text{wEnd}S_n$ , for all  $1 \leq i \leq n-1$ . At this point, we can conclude that  $\text{swEnd}S_n$  is a regular monoid.

Now, let  $\alpha \in \text{wEnd}S_n$ .

First, suppose that  $0\alpha = 0$  and take  $\beta \in \mathcal{T}(\Omega_{n-1}^0)$  such that  $\alpha = \alpha\beta\alpha$ . Notice that such  $\beta$  exists since  $\mathcal{T}(\Omega_{n-1}^0)$  is a regular monoid. Define  $\bar{\beta} \in \mathcal{T}(\Omega_{n-1}^0)$  by  $0\bar{\beta} = 0$  and  $i\bar{\beta} = i\beta$  for  $i \in \Omega_{n-1}$ . Then,  $\bar{\beta} \in \text{wEnd}S_n$  and  $\alpha = \alpha\bar{\beta}\alpha$ , whence  $\alpha$  is regular in  $\text{wEnd}S_n$ .

Next, suppose that  $0\alpha \neq 0$ . If  $\text{Im } \alpha = \{0\alpha\}$ , then  $\alpha$  is an idempotent and so  $\alpha$  is a regular element. Hence, suppose that  $\text{Im } \alpha = \{0, 0\alpha\}$ . Take  $i \in 0\alpha^{-1}$  and let  $\beta = \begin{pmatrix} 0 & 1 & \cdots & n-1 \\ i & 0 & \cdots & 0 \end{pmatrix} \in \text{wEnd}S_n$ . Therefore,  $\alpha = \alpha\beta\alpha$  and so  $\alpha$  is regular in  $\text{wEnd}S_n$ , as required.  $\square$

From now on and throughout this paper we will always consider  $n \geq 3$ . Observe that, in this case, we have  $\text{Aut}S_n \subsetneq \text{End}S_n \subsetneq \text{swEnd}S_n \subsetneq \text{wEnd}S_n$ .

Let us define

$$a_0 = \begin{pmatrix} 0 & 1 & 2 & 3 & \cdots & n-1 \\ 0 & 2 & 1 & 3 & \cdots & n-1 \end{pmatrix}, b_0 = \begin{pmatrix} 0 & 1 & 2 & \cdots & n-2 & n-1 \\ 0 & 2 & 3 & \cdots & n-1 & 1 \end{pmatrix} \text{ and } e_0 = \begin{pmatrix} 0 & 1 & 2 & 3 & \cdots & n-1 \\ 0 & 1 & 1 & 3 & \cdots & n-1 \end{pmatrix}.$$

Then,  $\{a_0, b_0, e_0\}$  is a generating set with minimum size of the monoid  $\mathcal{T}_{n-1}^0$  (notice that,  $a_0 = b_0$  for  $n = 3$ ); see [12]. Let us also consider the following transformations of  $\mathcal{T}(\Omega_{n-1}^0)$ :

$$c_0 = \begin{pmatrix} 0 & 1 & 2 & \cdots & n-1 \\ 0 & 0 & 2 & \cdots & n-1 \end{pmatrix}, z = \begin{pmatrix} 0 & 1 & 2 & \cdots & n-1 \\ 1 & 0 & 0 & \cdots & 0 \end{pmatrix} \text{ and } z_0 = \begin{pmatrix} 0 & 1 & 2 & \cdots & n-1 \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix}.$$

Notice that,  $\{a_0, b_0, e_0, c_0\}$  is a generating set with minimum size of the monoid  $\mathcal{PT}_{n-1}^0$ ; see [12]. Moreover, it is easy to check that  $a_0, b_0, e_0, c_0, z, z_0 \in \text{wEnd}(S_n)$ . Furthermore, we have:

**Theorem 2.2.** For  $n \geq 4$ ,  $\{a_0, b_0, e_0, z\}$ ,  $\{a_0, b_0, e_0, z, z_0\}$  and  $\{a_0, b_0, e_0, c_0, z\}$  are generating sets with minimum size of  $\text{End}S_n$ ,  $\text{swEnd}S_n$  and  $\text{wEnd}S_n$ , respectively. Consequently,  $\text{End}S_n$ ,  $\text{swEnd}S_n$  and  $\text{wEnd}S_n$  have ranks 4, 5 and 5, respectively.

*Proof.* Since  $\{a_0, b_0, e_0\}$  is a generating set of the monoid  $\mathcal{T}_{n-1}^0$  and, for all  $i \in \Omega_{n-1}$ ,  $\begin{pmatrix} 0 & 1 & \cdots & n-1 \\ i & 0 & \cdots & 0 \end{pmatrix} = z \begin{pmatrix} 0 & 1 & \cdots & n-1 \\ 0 & i & \cdots & i \end{pmatrix}$  and  $\begin{pmatrix} 0 & 1 & \cdots & n-1 \\ i & i & \cdots & i \end{pmatrix} = z_0 \begin{pmatrix} 0 & 1 & \cdots & n-1 \\ i & 0 & \cdots & 0 \end{pmatrix}$ , it immediately follows that  $\{a_0, b_0, e_0, z\}$  and  $\{a_0, b_0, e_0, z, z_0\}$  are generating sets of  $\text{End}S_n$  and  $\text{swEnd}S_n$ , respectively.

Next, observe that  $\text{wEnd}S_n = \mathcal{PT}_{n-1}^0 \cup \{\alpha \in \mathcal{T}(\Omega_{n-1}^0) \mid 0\alpha \neq 0 \text{ and } \text{Im } \alpha \subseteq \{0, 0\alpha\}\}$ . Since  $\mathcal{PT}_{n-1}^0$  is generated by  $\{a_0, b_0, e_0, c_0\}$ , in order to show that  $\{a_0, b_0, e_0, c_0, z\}$  generates  $\text{wEnd}S_n$ , it suffices to show that  $\{\alpha \in \mathcal{T}(\Omega_{n-1}^0) \mid 0\alpha \neq 0 \text{ and } \text{Im } \alpha \subseteq \{0, 0\alpha\}\}$  is contained in the monoid generated by  $\{a_0, b_0, e_0, c_0, z\}$ . So, take  $\alpha \in \mathcal{T}(\Omega_{n-1}^0)$  such that  $0\alpha \neq 0$  and  $\text{Im } \alpha \subseteq \{0, 0\alpha\}$ . Then,  $\alpha = \begin{pmatrix} 0 & A & B \\ 0\alpha & 0\alpha & 0 \end{pmatrix}$ , for some (disjoint) subsets  $A$  and  $B$  of  $\Omega_{n-1}$  such that  $A \cup B = \Omega_{n-1}$ . Let  $\alpha_0 = \begin{pmatrix} 0 & A & B \\ 0 & 0 & 0\alpha \end{pmatrix}$  and  $\beta = \begin{pmatrix} 0 & 1 & \cdots & n-1 \\ 0 & 0\alpha & \cdots & 0\alpha \end{pmatrix}$ . Then,  $\alpha_0, \beta \in \mathcal{PT}_{n-1}^0$  and  $\alpha = \alpha_0 z \beta$ , which concludes the proof.

Finally, it is easy to conclude that any generating set of  $\text{End}S_n$  cannot have less than four elements and that both  $\text{swEnd}S_n$  and  $\text{wEnd}S_n$  cannot be generated by less than five elements.  $\square$

For  $n = 3$ , as  $b_0 = a_0$  and  $e_0 = z^2$ , it is a routine matter to conclude that  $\{a_0, z\}$ ,  $\{a_0, z, z_0\}$  and  $\{a_0, c_0, z\}$  are generating sets with minimum size of  $\text{End}S_3$ ,  $\text{swEnd}S_3$  and  $\text{wEnd}S_3$ , respectively. Therefore,  $\text{End}S_3$ ,  $\text{swEnd}S_3$  and  $\text{wEnd}S_3$  have ranks 2, 3 and 3, respectively.

### 3 Presentations

Recall that  $\mathcal{T}_{n-1}^0$  and  $\mathcal{PT}_{n-1}^0$  are monoids isomorphic to  $\mathcal{T}(\Omega_{n-1})$  and  $\mathcal{PT}(\Omega_{n-1})$ , respectively.

Let  $\langle a_0, b_0, e_0 \mid R \rangle$  be a presentation of  $\mathcal{T}_{n-1}^0$  on the generators  $a_0, b_0$  and  $e_0$ . Then, we have:

**Theorem 3.1.** For  $n \geq 4$ ,

$$\langle a_0, b_0, e_0, z \mid R, a_0z = b_0z = e_0z = z, z^2 = (e_0b_0)^{n-3}e_0 \rangle$$

is a presentation of  $\text{End}S_n$  on the generators  $a_0, b_0, e_0$  and  $z$ .

*Proof.* First, notice that, it is a routine matter to show that the set of generators  $\{a_0, b_0, e_0, z\}$  satisfies all the relations  $R, b_0z = a_0z = e_0z = z$  and  $z^2 = (e_0b_0)^{n-3}e_0$ .

Denote by  $\sim_R$  the congruence on  $\{a_0, b_0, e_0\}^*$  generated by  $R$  and by  $\sim$  the congruence on  $\{a_0, b_0, e_0, z\}^*$  generated by  $R \cup \{b_0z = a_0z = e_0z = z, z^2 = (e_0b_0)^{n-3}e_0\}$ .

Let  $w \in \{a_0, b_0, e_0, z\}^*$ . Then  $w = u_1zu_2z \cdots u_kzu$ , for some  $u_1, u_2, \dots, u_k, u \in \{a_0, b_0, e_0\}^*$  and  $k \geq 0$ . As a consequence of the relations  $b_0z = a_0z = e_0z = z$ , we have  $u_i z \sim z$ , for  $1 \leq i \leq k$ , and so  $w \sim z^k u$ . On the other hand, considering also the relation  $z^2 = (e_0b_0)^{n-3}e_0$ , we also get  $z^3 = z^2z \sim (e_0b_0)^{n-3}e_0z \sim z$ , whence  $z^k u \sim zu$ , if  $k$  is odd, and  $z^k u \sim z^2 u \sim (e_0b_0)^{n-3}e_0 u$ , if  $k$  is an even integer greater than zero. Therefore, we obtain  $w \sim zu'$  or  $w \sim u'$ , for some  $u' \in \{a_0, b_0, e_0\}^*$ .

Now, let  $w_1, w_2 \in \{a_0, b_0, e_0, z\}^*$  be such that  $w_1 = w_2$  (equality in  $\text{End}S_n$ ). Our goal is to show that  $w_1 \sim w_2$ , with which the proof is concluded.

Let  $u_1, u_2 \in \{a_0, b_0, e_0\}^*$  be such that  $w_i \sim u_i$  or  $w_i \sim zu_i$ , for  $i = 1, 2$ . Hence, we may consider four cases.

CASE 1:  $w_1 \sim u_1$  and  $w_2 \sim u_2$ .

Then,  $u_1 = w_1 = w_2 = u_2$  (equalities in  $\text{End}S_n$ ) and so  $u_1 = u_2$  (equality in  $\mathcal{T}_{n-1}^0$ ). Since  $\langle a_0, b_0, e_0 \mid R \rangle$  is a presentation of  $\mathcal{T}_{n-1}^0$ , we have  $u_1 \sim_R u_2$ , whence  $u_1 \sim u_2$  and so  $w_1 \sim w_2$ .

CASE 2:  $w_1 \sim u_1$  and  $w_2 \sim zu_2$ . In this case, we have

$$\begin{pmatrix} 0 & 1 & \cdots & n-1 \\ 0 & 1u_1 & \cdots & (n-1)u_1 \end{pmatrix} = u_1 = w_1 = w_2 = zu_2 = \begin{pmatrix} 0 & 1 & \cdots & n-1 \\ 1 & 0 & \cdots & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 & \cdots & n-1 \\ 0 & 1u_2 & \cdots & (n-1)u_2 \end{pmatrix} = \begin{pmatrix} 0 & 1 & \cdots & n-1 \\ 1u_2 & 0 & \cdots & 0 \end{pmatrix},$$

whence  $u_1 = \begin{pmatrix} 0 & 1 & \cdots & n-1 \\ 0 & 0 & \cdots & 0 \end{pmatrix} \notin \mathcal{T}_{n-1}^0$ , which is a contradiction. Thus, this case does not occur.

CASE 3:  $w_1 \sim zu_1$  and  $w_2 \sim u_2$ .

Similar to the previous case, this case does not occur.

CASE 4:  $w_1 \sim zu_1$  and  $w_2 \sim zu_2$ .

Then,  $zu_1 = w_1 = w_2 = zu_2$  (equalities in  $\text{End}S_n$ ), whence  $zu_1 = zu_2$  and so  $z^2u_1 = z^2u_2$ . Since  $z^2 = (e_0b_0)^{n-3}e_0$ , we get  $(e_0b_0)^{n-3}e_0u_1 = (e_0b_0)^{n-3}e_0u_2$ , which is an equality in  $\mathcal{T}_{n-1}^0$ . Therefore, as  $\langle a_0, b_0, e_0 \mid R \rangle$  is a presentation of  $\mathcal{T}_{n-1}^0$ , we have  $(e_0b_0)^{n-3}e_0u_1 \sim_R (e_0b_0)^{n-3}e_0u_2$ , whence  $(e_0b_0)^{n-3}e_0u_1 \sim (e_0b_0)^{n-3}e_0u_2$  and so  $z^2u_1 \sim z^2u_2$ , from which follows that  $z^3u_1 \sim z^3u_2$  and, as  $z^3 \sim z$ , finally, that  $zu_1 \sim zu_2$ . Thus  $w_1 \sim w_2$ , by transitivity, as required.  $\square$

Next, let  $\langle a_0, b_0, e_0, z \mid Q \rangle$  be a presentation of  $\text{End}S_n$  on the generators  $a_0, b_0, e_0$  and  $z$ . Hence, we get:

**Theorem 3.2.** For  $n \geq 4$ ,

$$\langle a_0, b_0, e_0, z, z_0 \mid Q, a_0z_0 = b_0z_0 = e_0z_0 = zz_0 = z_0^2 = z_0a_0 = z_0b_0 = z_0e_0 = z_0 \rangle$$

is a presentation of  $\text{swEnd}S_n$  on the generators  $a_0, b_0, e_0, z$  and  $z_0$ .

*Proof.* We begin by observing that we can routinely show that the set of generators  $\{a_0, b_0, e_0, z, z_0\}$  satisfies all the relations  $Q$  and  $a_0z_0 = b_0z_0 = e_0z_0 = zz_0 = z_0^2 = z_0a_0 = z_0b_0 = z_0e_0 = z_0$ .

Denote by  $\sim_Q$  the congruence on  $\{a_0, b_0, e_0, z\}^*$  generated by  $Q$  and by  $\sim$  the congruence on  $\{a_0, b_0, e_0, z, z_0\}^*$  generated by  $Q \cup \{a_0z_0 = b_0z_0 = e_0z_0 = zz_0 = z_0^2 = z_0a_0 = z_0b_0 = z_0e_0 = z_0\}$ .

Let  $w \in \{a_0, b_0, e_0, z, z_0\}^* \setminus \{a_0, b_0, e_0, z\}^*$ . Then,  $w = u_1 z_0 u_2 z_0 \cdots u_k z_0 u$ , for some  $u_1, u_2, \dots, u_k, u \in \{a_0, b_0, e_0, z\}^*$  and  $k \geq 1$ . As a consequence of the relations  $a_0 z_0 = b_0 z_0 = e_0 z_0 = z z_0 = z_0$ , we get  $u_i z_0 \sim z_0$ , for  $1 \leq i \leq k$ , and so  $w \sim z_0^k u \sim z_0 u$ , also in view of the relation  $z_0^2 = z_0$ . On the other hand, we showed in the proof of Theorem 3.1 that  $u \sim_Q z u'$  or  $u \sim_Q u'$ , for some  $u' \in \{a_0, b_0, e_0\}^*$ . Hence,  $w \sim z_0 z u'$  or  $w \sim z_0 u' \sim z_0$ , for some  $u' \in \{a_0, b_0, e_0\}^*$ , also taking into account the relations  $z_0 a_0 = z_0 b_0 = z_0 e_0 = z_0$ . Observe that, as transformation, we obtain  $w = \begin{pmatrix} 0 & 1 & \cdots & n-1 \\ i & i & \cdots & i \end{pmatrix}$ , for some  $i \in \Omega_{n-1}^0$ , and so  $w \notin \text{End}S_n$ .

Now, let  $w_1, w_2 \in \{a_0, b_0, e_0, z, z_0\}^*$  be such that  $w_1 = w_2$  (equality in  $\text{swEnd}S_n$ ). We aim to show that  $w_1 \sim w_2$ . By the last observation above, we can conclude that  $w_1, w_2 \in \{a_0, b_0, e_0, z\}^*$  or  $w_1, w_2 \in \{a_0, b_0, e_0, z, z_0\}^* \setminus \{a_0, b_0, e_0, z\}^*$ . If  $w_1, w_2 \in \{a_0, b_0, e_0, z\}^*$ , then  $w_1 \sim_Q w_2$  and so  $w_1 \sim w_2$ . So, suppose that  $w_1, w_2 \in \{a_0, b_0, e_0, z, z_0\}^* \setminus \{a_0, b_0, e_0, z\}^*$ . Hence, as transformation,  $w_1 = w_2 = \begin{pmatrix} 0 & 1 & \cdots & n-1 \\ i & i & \cdots & i \end{pmatrix}$ , for some  $i \in \Omega_{n-1}^0$ . If  $i = 0$ , then we must have  $w_1 \sim z_0 \sim w_2$ . On the other hand, suppose that  $i \in \Omega_{n-1}$ . Then,  $w_1 \sim z_0 z u'_1$  and  $w_2 \sim z_0 z u'_2$ , for some  $u'_1, u'_2 \in \{a_0, b_0, e_0\}^*$ , and (as transformations)  $z u'_1 = z u'_2 = \begin{pmatrix} 0 & 1 & \cdots & n-1 \\ i & 0 & \cdots & 0 \end{pmatrix}$ . Thus,  $z u'_1 \sim_Q z u'_2$  and so  $w_1 \sim z_0 z u'_1 \sim z_0 z u'_2 \sim w_2$ , as required.  $\square$

Now, let  $\langle a_0, b_0, e_0, c_0 \mid S \rangle$  be a presentation of  $\mathcal{PT}_{n-1}^0$  on the generators  $a_0, b_0, e_0$  and  $c_0$ . Thus, we obtain:

**Theorem 3.3.** For  $n \geq 4$ ,

$$\langle a_0, b_0, e_0, c_0, z \mid S, a_0 z = b_0 z = e_0 z = z, z^2 = (e_0 b_0)^{n-3} e_0, z^2 c_0 = z c_0 \rangle$$

is a presentation of  $\text{wEnd}S_n$  on the generators  $a_0, b_0, e_0, c_0$  and  $z$ .

*Proof.* First, observe that it is a routine matter to check that the set of generators  $\{a_0, b_0, e_0, c_0, z\}$  satisfies all the considered relations.

Next, recall that  $\text{wEnd}S_n$  is the disjoint union of  $\mathcal{PT}_{n-1}^0$  and

$$\left\{ \begin{pmatrix} 0 & A & B \\ i & i & 0 \end{pmatrix} \in \mathcal{T}(\Omega_{n-1}^0) \mid A \cup B = \Omega_{n-1}, A \cap B = \emptyset, i \in \Omega_{n-1} \right\}.$$

For each  $A, B \subseteq \Omega_{n-1}$  such that  $A \cup B = \Omega_{n-1}$  and  $A \cap B = \emptyset$ , let  $u_{A,B}$  be a word of  $\{a_0, b_0, e_0, c_0\}^*$  representing the transformation  $\begin{pmatrix} 0 & A & B \\ 0 & 0 & 1 \end{pmatrix} \in \mathcal{PT}_{n-1}^0$ . For each  $1 \leq i \leq n-1$ , let  $u_i$  be a word of  $\{a_0, b_0, e_0\}^*$  representing the transformation  $\begin{pmatrix} 0 & 1 & \cdots & n-1 \\ 0 & i & \cdots & i \end{pmatrix} \in \mathcal{T}_{n-1}^0$ . Notice that, as transformations, we have  $\begin{pmatrix} 0 & A & B \\ i & i & 0 \end{pmatrix} = u_{A,B} z u_i$ , for any disjoint subsets  $A$  and  $B$  of  $\Omega_{n-1}$  such that  $A \cup B = \Omega_{n-1}$  and  $1 \leq i \leq n-1$ . Let  $W_0$  be a set of canonical forms for the monoid  $\mathcal{PT}_{n-1}^0$  regarding the set of generators  $\{a_0, b_0, e_0, c_0\}$ . Let

$$W_1 = \{u_{A,B} z u_i \in \{a_0, b_0, e_0, c_0, z\}^* \mid A \cup B = \Omega_{n-1}, A \cap B = \emptyset, 1 \leq i \leq n-1\}$$

and let  $W = W_0 \cup W_1$ . Then, clearly,  $|W| = |\text{wEnd}S_n|$ .

Denote by  $\sim_S$  the congruence on  $\{a_0, b_0, e_0, c_0\}^*$  generated by  $S$  and by  $\sim$  the congruence on  $\{a_0, b_0, e_0, c_0, z\}^*$  generated by  $S \cup \{a_0 z = b_0 z = e_0 z = z, z^2 = (e_0 b_0)^{n-3} e_0, z^2 c_0 = z c_0\}$ .

Let  $w \in W_0$ . Since  $\langle a_0, b_0, e_0, c_0 \mid S \rangle$  is a presentation of  $\mathcal{PT}_{n-1}^0$ , for each  $x \in \{a_0, b_0, e_0, c_0\}$ , we have  $w x \sim_S w'$ , whence  $w x \sim w'$ , for some  $w' \in W_0$ . On the other hand, being  $A = \{k \in \Omega_{n-1} \mid (k)w = 0\}$  and  $B = \Omega_{n-1} \setminus A$ , we have  $w(e_0 b_0)^{n-3} e_0 = u_{A,B}$  and  $(e_0 b_0)^{n-3} e_0 = u_1$  (considering  $w, u_1$  and  $u_{A,B}$  as transformations of  $\mathcal{PT}_{n-1}^0$ ), whence  $w(e_0 b_0)^{n-3} e_0 \sim_S u_{A,B}$  and  $(e_0 b_0)^{n-3} e_0 \sim_S u_1$ , and so

$$w z \sim w(e_0 b_0)^{n-3} e_0 (e_0 b_0)^{n-3} e_0 z \sim u_{A,B} z^2 z = u_{A,B} z z^2 \sim u_{A,B} z (e_0 b_0)^{n-3} e_0 \sim u_{A,B} z u_1 \in W_1.$$

Now, take a subset  $A$  of  $\Omega_{n-1}$ ,  $B = \Omega_{n-1} \setminus A$  and  $1 \leq i \leq n-1$ . Let  $w = u_{A,B} z u_i \in W_1$ . For each  $x \in \{a_0, b_0, e_0\}$ , we have  $u_i x = u_{(i)x}$  (considering  $u_i$  and  $x$  as transformation of  $\mathcal{T}_{n-1}^0$ ), whence  $u_i x \sim_S u_{(i)x}$  and so  $w x = u_{A,B} z u_i x \sim u_{A,B} z u_{(i)x} \in W_1$ . On the other hand, (as transformations) we have  $u_1 c_0 = c_0 u_1 c_0$  and  $u_i c_0 = u_i$ , if  $2 \leq i \leq n-1$ , whence

$$w c_0 = u_{A,B} z u_1 c_0 \sim u_{A,B} z c_0 u_1 c_0 \sim u_{A,B} z^2 c_0 u_1 c_0 \sim u_{A,B} (e_0 b_0)^{n-3} e_0 c_0 u_1 c_0 \sim w',$$

for some  $w' \in W_0$ , and  $w c_0 = u_{A,B} z u_i c_0 \sim u_{A,B} z u_i = w \in W_1$ , if  $2 \leq i \leq n-1$ . Finally, since  $u_i \in \{a_0, b_0, e_0\}^*$ , we have  $u_i z \sim z$  and so  $w z = u_{A,B} z u_i z \sim u_{A,B} z^2 \sim u_{A,B} (e_0 b_0)^{n-3} e_0 \sim w'$ , for some  $w' \in W_0$ .

Thus, we proved that  $W$  is a set of canonical forms for  $\text{wEnd}S_n$ , which completes this proof.  $\square$

We finish this paper by exhibiting also presentations for  $\text{End}S_3$ ,  $\text{swEnd}S_3$  and  $\text{wEnd}S_3$ . These presentations can be verified by using GAP [16]:

$$\text{End}S_3 = \langle a_0, z \mid a_0^2 = 1, a_0z = z, z^3 = z \rangle;$$

$$\text{swEnd}S_3 = \langle a_0, z, z_0 \mid a_0^2 = 1, a_0z = z, z^3 = z, a_0z_0 = zz_0 = z_0^2 = z_0a_0 = z_0z^2 = z_0 \rangle;$$

$$\begin{aligned} \text{wEnd}S_3 = \langle a_0, c_0, z \mid a_0^2 = 1, a_0z = z, z^3 = z, c_0^2 = c_0, (c_0a_0)^2 = (a_0c_0)^2 = c_0a_0c_0, \\ z^2c_0 = c_0a_0c_0, c_0z^2 = c_0a_0, z^2a_0c_0 = z^2a_0, z^2c_0 = zc_0 \rangle. \end{aligned}$$

## References

- [1] A.Ya. Aĭzenštat, Defining relations of finite symmetric semigroups, *Mat. Sb. N. S.* 45 (1958), 261–280 (Russian).
- [2] A.Ya. Aĭzenštat, The defining relations of the endomorphism semigroup of a finite linearly ordered set, *Sibirsk. Mat.* 3 (1962), 161–169 (Russian).
- [3] M. Böttcher and U. Knauer, *Endomorphism spectra of graphs*, *Discrete Mathematics* 109 (1992), 45–57.
- [4] I. Dimitrova, V. H. Fernandes and J. Koppitz, *On partial endomorphisms of a star graph*, *Quaestiones Mathematicae* (2024), 1–21 (<https://doi.org/10.2989/16073606.2024.2374796>).
- [5] I. Dimitrova, V. H. Fernandes and J. Koppitz, *Presentations for monoids of partial endomorphisms of a star graph*, arXiv:2410.21084 [math.RA] (<https://doi.org/10.48550/arXiv.2410.21084>).
- [6] I. Dimitrova, V. H. Fernandes, J. Koppitz and T. M. Quinteiro, *Ranks of monoids of endomorphisms of a finite undirected path*, *Bull. Malaysian Math. Sci. Soc.* 43(2) (2020), 1623–1645.
- [7] I. Dimitrova, V. H. Fernandes, J. Koppitz and T. M. Quinteiro, *Partial Automorphisms and Injective Partial Endomorphisms of a Finite Undirected Path*, *Semigroup Forum* 103(1), (2021), 87–105.
- [8] I. Dimitrova, V. H. Fernandes, J. Koppitz and T. M. Quinteiro, *On monoids of endomorphisms of a cycle graph*, *Mathematica Slovaca* 74(5), (2024), 1071–1088.
- [9] J. East, Generators and relations for partition monoids and algebras, *J. Algebra* 339 (2011), 1–26.
- [10] S. Fan, *On End-regular graphs*, *Discrete Mathematics* 159 (1996), 95–102.
- [11] V.H. Fernandes, The monoid of all injective order preserving partial transformations on a finite chain, *Semigroup Forum* 62 (2001), 178–204.
- [12] V.H. Fernandes, *Presentations for some monoids of partial transformations on a finite chain: a survey*, *Semigroups, Algorithms, Automata and Languages*, eds. Gracinda M. S. Gomes & Jean-Éric Pin & Pedro V. Silva, World Scientific (2002), 363–378.
- [13] V.H. Fernandes, G.M.S. Gomes and M.M. Jesus, Presentations for some monoids of injective partial transformations on a finite chain, *Southeast Asian Bull. Math.* 28 (2004), 903–918.
- [14] V.H. Fernandes and T. Paulista, *On the monoid of partial isometries of a finite star graph*, *Commun. Algebra* 51 (2023), 1028–1048.
- [15] V.H. Fernandes and T.M. Quinteiro, Presentations for monoids of finite partial isometries, *Semigroup Forum* 93 (2016), 97–110.



- [16] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.13.0*; 2024.  
(<https://www.gap-system.org>)
- [17] R. Gu and H. Hou, *End-regular and End-orthodox generalized lexicographic products of bipartite graphs*, Open Math. 14 (2016), 229–236.
- [18] H. Hou, Y. Luo and S. Fan, *End-regular and End-orthodox joins of split graphs*, Ars Combinatoria 105 (2012), 305–318.
- [19] J.M. Howie, *Fundamentals of Semigroup Theory*, Clarendon Press, Oxford, 1995.
- [20] J.M. Howie and N. Ruškuc, *Constructions and presentations for monoids*, Commun. Algebra 22 (1994), 6209–6224.
- [21] A.V. Kelarev, *Graph Algebras and Automata*, Marcel Dekker, New York, 2003.
- [22] U. Knauer, *Algebraic graph theory: morphisms, monoids, and matrices*, De Gruyter, Berlin, 2011.
- [23] U. Knauer and A. Wanichsombat, *Completely Regular Endomorphisms of Split Graphs*, Ars Combinatoria 115 (2014), 357–366.
- [24] G. Lallement, *Semigroups and Combinatorial Applications*, John Wiley & Sons, New York, 1979.
- [25] W. Li, *Graphs with regular monoids*, Discrete Mathematics 265 (2003), 105–118.
- [26] E.H. Moore, *Concerning the abstract groups of order  $k!$  and  $\frac{1}{2}k!$  holohedrally isomorphic with the symmetric and the alternating substitution groups on  $k$  letters*, Proc. London Math. Soc. 28 (1897), 357–366.
- [27] L.M. Popova, *The defining relations of certain semigroups of partial transformations of a finite set*, Leningrad. Gos. Ped. Inst. Učen. Zap. 218 (1961), 191–212 (Russian).
- [28] L.M. Popova, *Defining relations of a semigroup of partial endomorphisms of a finite linearly ordered set*, Leningrad. Gos. Ped. Inst. Učen. Zap. 238 (1962), 78–88 (Russian).
- [29] N. Ruškuc, *Semigroup Presentations*, Ph. D. Thesis, University of St-Andrews, 1995.
- [30] E. Wilkeit, *Graphs with a regular endomorphism monoid*, Arch. Math. 66 (1996), 344–352.

VÍTOR H. FERNANDES, Center for Mathematics and Applications (NOVA Math) and Department of Mathematics, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Monte da Caparica, 2829-516 Caparica, Portugal; e-mail: [vhf@fct.unl.pt](mailto:vhf@fct.unl.pt).

JÖRG KOPPITZ, Institute of Mathematics and Informatics, Bulgarian Academy of Sciences, 1113 Sofia, Bulgaria; e-mail: [koppitz@math.bas.bg](mailto:koppitz@math.bas.bg).

TIWADEE MUSUNTHIA, Department of Mathematics, Faculty of Science, Silpakorn University, Nakorn Pathom, Thailand 73000; e-mail: [tiwadee.m@gmail.com](mailto:tiwadee.m@gmail.com).