

CRYSTALLINE LIFTS OF SEMISIMPLE G -VALUED GALOIS REPRESENTATIONS WITH FIXED DETERMINANT

KENSUKE AOKI

ABSTRACT. For a finite extension K/\mathbb{Q}_p and a split reductive group G over \mathcal{O}_K , let $\bar{\rho}: \text{Gal}_K \rightarrow G(\overline{\mathbb{F}}_p)$ be a continuous quasi-semisimple mod p G -valued representation of the absolute Galois group Gal_K . Let $\bar{\rho}^{\text{ab}}$ be the abelianization of $\bar{\rho}$ and fix a crystalline lift ψ of $\bar{\rho}^{\text{ab}}$. We show the existence of a crystalline lift ρ of $\bar{\rho}$ with regular Hodge-Tate weights such that the abelianization of ρ coincides with ψ . We also show analogous results in the case that G is a quasi-split tame group and $\bar{\rho}: \text{Gal}_K \rightarrow {}^L G(\overline{\mathbb{F}}_p)$ is a semisimple mod p L -parameter. These theorems are generalizations of those of Lin and Böckle-Iyengar-Paškūnas.

CONTENTS

1. Introduction	1
2. Notation	3
3. Crystalline extensions to Cartan subgroups	3
4. Crystalline lifts of quasi-semisimple representations with fixed abelianizations	6
5. Potentially crystalline lifts of semisimple L -parameters with fixed abelianizations	10
References	13

1. INTRODUCTION

Let K/\mathbb{Q}_p be a finite extension with its ring of integers \mathcal{O}_K , $\text{Gal}_K := \text{Gal}(\overline{K}/K)$ be the absolute Galois group of K , L/K be a finite extension with its ring of integers \mathcal{O}_L . Let $\mathfrak{m}_K, \mathfrak{m}_L$ be the maximal ideals of $\mathcal{O}_K, \mathcal{O}_L$, and k, k_L be the residue fields of $\mathcal{O}_K, \mathcal{O}_L$, respectively. Let \mathbb{C}_p be the p -adic completion of \overline{K} .

For a reductive group G over \mathcal{O}_L , a continuous G -valued representation $\rho: \text{Gal}_K \rightarrow G(L)$ is *Hodge-Tate* (resp. *crystalline*, *potentially crystalline*) if for any algebraic L -representation $\tau: G \rightarrow \text{GL}(V)$, the composite $\tau \circ \rho$ is a Hodge-Tate (resp. crystalline, potentially crystalline) representation. For a Hodge-Tate G -valued representation $\rho: \text{Gal}_K \rightarrow G(L)$, we say that ρ has *regular* Hodge-Tate weights if for each $\sigma \in \text{Hom}_{\mathbb{Q}_p}(L, \mathbb{C}_p)$, the associated cocharacter $\mathcal{HT}(\rho)^\sigma$ of $G_{\mathbb{C}_p}$ is regular, that is, not killed by any root of $G_{\mathbb{C}_p}$ (see Definition 3.6). Let G^{ab} be the abelianization of G . The composite of ρ with $G(L) \rightarrow G^{\text{ab}}(L)$ is written as ρ^{ab} . We call it the *abelianization* of ρ .

A continuous representation $\rho: \text{Gal}_K \rightarrow G(\mathcal{O}_L)$ is said to be *Hodge-Tate* (resp. *crystalline*, *potentially crystalline*) if the composite $\text{Gal}_K \xrightarrow{\rho} G(\mathcal{O}_L) \hookrightarrow G(L)$ is Hodge-Tate (resp. crystalline, potentially crystalline).

From here, let G be a connected split reductive group over \mathcal{O}_L and let T be a maximal torus of G . We say $\bar{\rho}: \text{Gal}_K \rightarrow G(\overline{\mathbb{F}}_p)$ is *quasi-semisimple* if there exists a maximal torus T' of $G_{\overline{\mathbb{F}}_p} := G \otimes_{\mathcal{O}_K} \overline{\mathbb{F}}_p$ such that $\bar{\rho}(I_K) \subset T'(\overline{\mathbb{F}}_p)$ and $\bar{\rho}(\text{Gal}_K) \subset$

$N_{G_{\overline{\mathbb{F}}_p}}(T')(\overline{\mathbb{F}}_p)$, where $I_K \subset \text{Gal}_K$ denotes the inertia subgroup and $N_{G_{\overline{\mathbb{F}}_p}}(T')$ denotes the normalizer of T' in $G_{\overline{\mathbb{F}}_p}$.

The first main theorem of this paper is as follows:

Theorem 1.1 (Theorem 4.4). *Assume that T is a split torus and the composite $T \hookrightarrow G \twoheadrightarrow G^{\text{ab}}$ has a right inverse, that is, there exists a homomorphism $G^{\text{ab}} \rightarrow T$ such that the composite $G^{\text{ab}} \rightarrow T \rightarrow G^{\text{ab}}$ is the identity. Let $\overline{\rho}: \text{Gal}_K \rightarrow G(k_L)$ be a quasi-semisimple G -valued representation such that $\overline{\rho}(I_K) \subset T(k_L)$ and $\overline{\rho}(\text{Gal}_K) \subset N_G(T)(k_L)$. Let $\overline{\rho}^{\text{ab}}: \text{Gal}_K \rightarrow G^{\text{ab}}(k_L)$ be the composite of $\overline{\rho}$ and the natural map $G(k_L) \rightarrow G^{\text{ab}}(k_L)$. Let $\psi: \text{Gal}_K \rightarrow G^{\text{ab}}(\mathcal{O}_L)$ be a crystalline lift of $\overline{\rho}^{\text{ab}}$. Then there exist a finite extension L'/L and a crystalline lift $\rho: \text{Gal}_K \rightarrow G(\mathcal{O}_{L'})$ of $\overline{\rho}$ with regular Hodge-Tate weights such that $\rho^{\text{ab}} = \psi$. Moreover, if we do not impose a crystalline lift ρ has regular Hodge-Tate weights, L'/L can be taken to be a finite unramified extension.*

Theorem 1.1 is a generalization of the results of Lin [9, Theorem 5, Theorem 6] and Böckle-Iyengar-Paškūnas [4, Proposition 2.6].

We also prove analogous results of Theorem 1.1 for potentially crystalline lifts of semisimple mod p L -parameters for quasi-split tame groups over K . Let G be a connected quasi-split reductive group over K . We assume that G is tame, that is, G splits over a tame Galois extension F of K . Write \widehat{G} for the Langlands dual group of G and ${}^L G = \widehat{G} \rtimes \text{Gal}(F/K)$ for the L -group of G . Let \widehat{T} be a maximal torus of \widehat{G} .

The second main theorem of this paper is as follows:

Theorem 1.2 (Theorem 5.5). *Let $\overline{\rho}: \text{Gal}_K \rightarrow {}^L G(\overline{\mathbb{F}}_p)$ be a semisimple mod p L -parameter in the sense of [10, Definition 3.2.5]. Fix a maximal torus S of G such that $\overline{\rho}$ factors through ${}^L S(\overline{\mathbb{F}}_p)$ (see Theorem 5.3). Assume that the composite $S \hookrightarrow G \twoheadrightarrow G^{\text{ab}}$ has a right inverse, which is written as $i: G^{\text{ab}} \rightarrow S$. For $A = \overline{\mathbb{F}}_p^*$ or $\overline{\mathbb{Z}}_p^*$ and an L -parameter $\tau: \text{Gal}_K \rightarrow {}^L S(A)$, write τ^{ab} for the composite of τ and the map ${}^L S(A) \rightarrow {}^L G^{\text{ab}}(A)$ induced by the composite $G^{\text{ab}} \xrightarrow{i} S \hookrightarrow G$ and we set $\overline{\psi} := \overline{\rho}^{\text{ab}}$. Let $\psi: \text{Gal}_K \rightarrow {}^L G^{\text{ab}}(\overline{\mathbb{Z}}_p)$ be a potentially crystalline lift of $\overline{\psi}$. Then $\overline{\rho}$ admits a potentially crystalline lift $\rho: \text{Gal}_K \rightarrow {}^L G(\overline{\mathbb{Z}}_p)$ which has regular Hodge-Tate weights and satisfies $\rho^{\text{ab}} = \psi$.*

The existence of crystalline lifts with fixed determinant in the case of $G = \text{GL}_m$ is proved by Böckle-Iyengar-Paškūnas [4, Proposition 2.6, Proposition 2.7]. In the paper, they firstly proved the case where $\overline{\rho}: \text{Gal}_K \rightarrow \text{GL}_m(\overline{\mathbb{F}}_p)$ is absolutely irreducible and the general case is deduced from the former case using the geometry of the Emerton-Gee stacks. Also, Lin [9, Theorem 5, Theorem 6] showed the existence of crystalline lifts with regular Hodge-Tate weights of quasi-semisimple mod p G -valued representations when G is a split reductive group over \mathcal{O}_K (see [10, Theorem 5.3.1] for semisimple mod p L -parameters for quasi-split tame groups). In this paper, we combine the methods of [4] and [9], [10] to prove the existence of crystalline lifts of $\overline{\rho}$ with fixed abelianization.

Remark 1.3. Let $\overline{\rho}: \text{Gal}_K \rightarrow G(k_L)$ be a continuous G -valued representation and $R_{\overline{\rho}}^{\square}$ be the universal framed deformation ring constructed by Balaji [2, Theorem 1.2.2]. Write $\mathcal{X}_{\overline{\rho}}^{\square} := \text{Spf}(R_{\overline{\rho}}^{\square})^{\text{rig}}$ for the rigid generic fiber of $\text{Spf}(R_{\overline{\rho}}^{\square})$ as constructed

in [8, Definition 7.1.3]. For $G = \mathrm{GSp}_{2n}$, Zariski density of crystalline points on an irreducible component of \mathcal{X}_p^\square which has a crystalline point is shown in [1] using the geometry of the trianguline deformation space for GSp_{2n} . To prove the full version of the Zariski density theorem for GSp_{2n} , we have to show that each irreducible component of \mathcal{X}_p^\square has a crystalline point. By the results of Paškūnas-Quast [11], it suffices to show that any mod p G -valued representation $\bar{\rho}$ has a crystalline lift with fixed abelianization. We expect that Theorem 1.1 is the first step for the full version of the Zariski density theorem for split reductive groups.

The strategies of the proofs of Theorem 1.1 and Theorem 1.2 are based on Lin's strategies in [9] and [10], respectively, where the existence of crystalline lifts of semisimple mod p G -valued representations is shown. In order to control the abelianizations of crystalline lifts, we use the calculations of abelianizations of Galois representations via the reciprocity map of local class field theory as in the work of Böckle-Iyengar-Paškūnas [4].

Acknowledgement. The author thanks his advisor Tetsushi Ito for valuable comments on the draft of the paper.

2. NOTATION

Let p be a prime number. For a group G and a subset $S \subset G$, we write $N_G(S)$ for the normalizer of S in G and $C_G(S)$ for the centralizer of S in G . We fix an algebraic closure $\overline{\mathbb{Q}_p}$ of \mathbb{Q}_p . Let K be a finite extension of \mathbb{Q}_p contained in $\overline{\mathbb{Q}_p}$. Let \mathbb{C}_p be the p -adic completion of $\overline{\mathbb{Q}_p}$. Let L be a sufficiently large finite extension of K such that $|\mathrm{Hom}_{\mathbb{Q}_p}(K, L)| = [K : \mathbb{Q}_p]$. Let k_L be the residue field of L and $\Sigma_L := \mathrm{Hom}_{\mathbb{Q}_p}(L, \mathbb{C}_p)$. Let $\mathrm{Gal}_K := \mathrm{Gal}(\overline{\mathbb{Q}_p}/K)$ be the absolute Galois group of K , and I_K be the inertia subgroup of Gal_K . Let W_K be the Weil group of K . For a finite Galois extension E/K , let $W_{E/K} := W_K/W_E^{\mathrm{der}}$, where W_E^{der} is the closure of the derived subgroup W_E^{der} of W_E , be the relative Weil group as defined in [12]. Write $\widehat{\mathbb{Z}} := \prod_{\ell: \text{prime}} \mathbb{Z}_\ell$. We fix a uniformizer $\varpi_K \in \mathcal{O}_K$ and normalize the reciprocity isomorphism $\mathrm{rec}_K: K^* \xrightarrow{\sim} W_K^{\mathrm{ab}}$ of local class field theory such that ϖ_K is mapped to the geometric Frobenius automorphism.

For an algebraic group G over a commutative ring R , write $G^{\mathrm{der}} := [G, G]$ for the derived subgroup of G , $G^{\mathrm{ab}} := G/G^{\mathrm{der}}$ for the abelianization of G , and G° for the identity component of G . Write $X^*(G) := \mathrm{Hom}_R(G, \mathbb{G}_{m,R})$ for the character group of G and $X_*(G) := \mathrm{Hom}_R(\mathbb{G}_{m,R}, G)$ for the cocharacter group of G .

Let A be a commutative topological R -algebra. For a continuous representation $\rho: \mathrm{Gal}_K \rightarrow G(A)$, let $\rho^{\mathrm{ab}}: \mathrm{Gal}_K \rightarrow G^{\mathrm{ab}}(A)$ be the composite of ρ with the abelianization map $G(A) \rightarrow G^{\mathrm{ab}}(A)$.

3. CRYSTALLINE EXTENSIONS TO CARTAN SUBGROUPS

In this section, let G be a connected split reductive group over \mathcal{O}_K and T be a maximal torus of G . We assume that the composite $T \hookrightarrow G \rightarrow G^{\mathrm{ab}}$ has a right inverse, that is, there exists a homomorphism $G^{\mathrm{ab}} \rightarrow T$ such that the composite $G^{\mathrm{ab}} \rightarrow T \rightarrow G^{\mathrm{ab}}$ is the identity. Note that $G^{\mathrm{ab}} \cong Z^\circ/(Z^\circ \cap G^{\mathrm{der}})$, where Z is the center of G , is a torus.

We recall the notation of [9, §4]. Let $\Phi_K \in \mathrm{Gal}_K$ be a lift of a topological generator of $\mathrm{Gal}_K/I_K \cong \widehat{\mathbb{Z}}$. Let $W(G, T)$ be the Weyl group scheme of (G, T) . For

the ease of notation, write $W(G, T)$ for the group $W(G, T)(\mathcal{O}_L) \cong W(G, T)(k_L)$, which is a subgroup of the finite group $W(G, T)(\overline{L})$.

Let $M_{T, \text{crys}}$ denote the set of continuous $T(\mathcal{O}_L)$ -valued representations $I_K \rightarrow T(\mathcal{O}_L)$ of the inertia group I_K , which can be extended to a crystalline representation $\text{Gal}_{K'} \rightarrow T(\mathcal{O}_L)$ for some finite unramified extension K'/K inside \overline{K} . Let M_{T, k_L} denote the set of continuous mod ϖ_L representations $I_K \rightarrow T(k_L)$ of I_K .

Both $M_{T, \text{crys}}$ and M_{T, k_L} are naturally equipped with the structures of $\mathbb{Z}[W(G, T)]$ -modules defined by $wv := (\sigma \mapsto wv(\sigma)w^{-1})$ for $w \in W(G, T)$ and $v \in M_{T, \text{crys}}$ (or $v \in M_{T, k_L}$). Similarly, they are naturally equipped with the structures of $\mathbb{Z}[\text{Gal}_K/I_K]$ -modules defined by $\alpha v := (\sigma \mapsto v(\alpha^{-1}\sigma\alpha))$ for $\alpha \in \text{Gal}_K$ and $v \in M_{T, \text{crys}}$ (or $v \in M_{T, k_L}$).

- Definition 3.1.** (1) A continuous G -valued representation $\overline{\rho}: \text{Gal}_K \rightarrow G(\overline{\mathbb{F}}_p)$ is said to be *elliptic* if it does not factor through any $P(\overline{\mathbb{F}}_p) \subset G(\overline{\mathbb{F}}_p)$ where P is a parabolic subgroup of $G_{\overline{\mathbb{F}}_p}$.
- (2) A subgroup $\Gamma \subset G(\overline{\mathbb{F}}_p)$ is said to be *G -completely reducible* if for any parabolic subgroup P of $G_{\overline{\mathbb{F}}_p}$ such that $\Gamma \subset P(\overline{\mathbb{F}}_p)$, a Levi subgroup L of P also satisfies $\Gamma \subset L(\overline{\mathbb{F}}_p)$. We also say that a continuous G -valued representation $\overline{\rho}: \text{Gal}_K \rightarrow G(\overline{\mathbb{F}}_p)$ is *G -completely reducible* if the image of $\overline{\rho}$ is G -completely reducible.
- (3) We say $\overline{\rho}: \text{Gal}_K \rightarrow G(\overline{\mathbb{F}}_p)$ is *quasi-semisimple* if there exists a maximal torus T' of $G_{\overline{\mathbb{F}}_p}$ such that $\overline{\rho}(I_K) \subset T'(\overline{\mathbb{F}}_p)$ and $\overline{\rho}(\text{Gal}_K) \subset N_{G_{\overline{\mathbb{F}}_p}}(T')(\overline{\mathbb{F}}_p)$.

Note that an elliptic continuous G -valued representation $\text{Gal}_K \rightarrow G(\overline{\mathbb{F}}_p)$ is G -completely reducible.

Theorem 3.2. *If a continuous G -valued representation $\text{Gal}_K \rightarrow G(\overline{\mathbb{F}}_p)$ is G -completely reducible, then $\overline{\rho}$ is quasi-semisimple.*

Proof. See [9, Theorem 4]. □

For showing the main theorem, we will reduce to the case where $\overline{\rho}: \text{Gal}_K \rightarrow G(\overline{\mathbb{F}}_p)$ is elliptic. By Theorem 3.2, it suffices to consider the existence of a lift $\rho: \text{Gal}_K \rightarrow N_G(T)(\mathcal{O}_L)$ of $\overline{\rho}$ satisfying $\rho(\Phi_K) = w^{-1}$ for an element $w \in N_G(T)(\mathcal{O}_L)$.

In the following, we generalize Lin's argument in [9, §4] to the case that $\rho(\Phi_K)$ is not of finite order and the homomorphism $\mathbb{Z} \rightarrow G(\mathcal{O}_L)$, $1 \mapsto \rho(\Phi_K)$ can be topologically extended to a homomorphism $\widehat{\mathbb{Z}} \rightarrow G(\mathcal{O}_L)$.

Definition 3.3. For a Hausdorff topological group G , we say that a homomorphism $\mathbb{Z} \rightarrow G$, $1 \mapsto g$ is *extendable* if it can be extended to a (unique) continuous homomorphism $\widehat{\mathbb{Z}} \rightarrow G$.

Lemma 3.4. *Let ℓ be a prime number, m be a positive integer, G be a locally profinite group, and $g \in G$.*

- (1) *If $\lim_{k \rightarrow \infty} g^{\ell^k} = e$, the homomorphism $\mathbb{Z} \rightarrow G$, $1 \mapsto g$ is extendable.*
- (2) *The homomorphism $1 \mapsto g$ is extendable if and only if the homomorphism $1 \mapsto g^m$ is extendable.*

Proof. (1) The homomorphism $\mathbb{Z} \rightarrow G$, $1 \mapsto g$ can be extended to $\mathbb{Z}_\ell \rightarrow G$ because there exists a system $\{U_i\}_{i \geq 1}$ of neighborhoods of the unity $e \in G$, where each U_i

is a profinite open subgroup of G . Then (1) follows by taking the composite of the projection $\widehat{\mathbb{Z}} \rightarrow \mathbb{Z}_\ell$ with a map $\mathbb{Z}_\ell \rightarrow G$.

(2) The “only if” part is obvious. If $1 \mapsto g^m$ is extendable, there exists a continuous homomorphism $\widehat{f}' : m\widehat{\mathbb{Z}} \rightarrow G$ which extends $m\mathbb{Z} \rightarrow G$, that is the restriction of $f : \mathbb{Z} \rightarrow G$, $1 \mapsto g$. For each $a \in [0, m-1]$, the map

$$\widehat{f}'_a : a + m\widehat{\mathbb{Z}} \rightarrow G, \quad a + mb \mapsto f(a)\widehat{f}'(mb)$$

is continuous. Since $\{a + m\widehat{\mathbb{Z}} \mid a \in [0, m-1]\}$ is an open covering of $\widehat{\mathbb{Z}}$, the map $\bigsqcup_{a \in [0, m-1]} \widehat{f}'_a : \widehat{\mathbb{Z}} \rightarrow G$ is a continuous homomorphism which extends f . \square

We prove the following lemma, which is proved by Lin [9, Lemma 4] when w has finite order.

Lemma 3.5. *Let $\zeta : N_G(T) \rightarrow W(G, T)$ be the quotient map.*

- (1) *Let $v : I_K \rightarrow T(\mathcal{O}_L)$ be a continuous representation such that $v \in M_{T, \text{crys}}$. For an element $w \in N_G(T)(\mathcal{O}_L)$, the representation v extends to a continuous representation $\rho : \text{Gal}_K \rightarrow N_G(T)(\mathcal{O}_L)$ satisfying $\rho(\Phi_K) = w^{-1}$ and $\rho|_{I_K} = v$ if and only if*

$$v \in \ker(M_{T, \text{crys}} \xrightarrow{\zeta(w) \otimes 1 - 1 \otimes \Phi_K} M_{T, \text{crys}}).$$

- (2) *Let $\bar{v} : I_K \rightarrow T(k_L)$ be a continuous representation such that $\bar{v} \in M_{T, k_L}$. For an element $\bar{w} \in N_G(T)(k_L)$, the representation \bar{v} extends to a continuous representation $\bar{\rho} : \text{Gal}_K \rightarrow N_G(T)(k_L)$ satisfying $\bar{\rho}(\Phi_K) = \bar{w}^{-1}$ and $\bar{\rho}|_{I_K} = \bar{v}$ if and only if*

$$\bar{v} \in \ker(M_{T, k_L} \xrightarrow{\zeta(\bar{w}) \otimes 1 - 1 \otimes \Phi_K} M_{T, k_L}).$$

Proof. (1) Assume that $v \in \ker(\zeta(w) \otimes 1 - 1 \otimes \Phi_K : M_{T, \text{crys}} \rightarrow M_{T, \text{crys}})$. By [9, Lemma 4], the representation v extends to a representation $v' : W_K \cong I_K \rtimes \mathbb{Z} \rightarrow G(\mathcal{O}_L)$. So it suffices to show that v' extends to a continuous representation of the absolute Galois group Gal_K , which is homeomorphic to the profinite set $I_K \times \widehat{\mathbb{Z}}$.

Since the centralizer $C_G(T)$ is the maximal torus T and the Weyl group $W(G, T)$ is finite, there exists a positive integer m such that $w^m \in T(\mathcal{O}_L)$. For a finite extension L'/L , the torus T splits over $\mathcal{O}_{L'}$ and the topology of the group $T(\mathcal{O}_L)$ is the restriction of that of $T(\mathcal{O}_{L'})$. Fix an isomorphism $i : T \cong \mathbb{G}_{m, \mathcal{O}_L}^d$ over $\mathcal{O}_{L'}$. Then there is also a positive integer m' such that the j -th component of $i(w^{mm'}) \in (\mathcal{O}_{L'}^*)^d$ is in $1 + \mathfrak{m}_{L'}$ for every $1 \leq j \leq d$. Then $\lim_{k \rightarrow \infty} (w^{mm'})^{p^k} = 1$ in $T(\mathcal{O}_{L'})$. The limit also exists and equals to 1 since $T(\mathcal{O}_L)$ is a closed subset of $T(\mathcal{O}_{L'})$. So $1 \mapsto w^{mm'}$ and then $1 \mapsto w$ is extendable by Lemma 3.4. The “only if” part is obvious.

- (2) See [9, Lemma 4 (2)] \square

We recall the notion of co-labeled Hodge-Tate characters following [9, §6, Definition 3].

Definition 3.6. Let G be a reductive group over L .

- (1) For a Hodge-Tate G -valued representation $\rho : \text{Gal}_K \rightarrow G(L)$, there is a cocharacter $\mathcal{HT}(\rho)^\sigma : \mathbb{G}_{m, \mathbb{C}_p} \rightarrow G_{\mathbb{C}_p}$ associated with ρ for each $\sigma \in \Sigma_L =$

$\mathrm{Hom}_{\mathbb{Q}_p}(L, \mathbb{C}_p)$ by Tannakian theory (cf. [9, Definition 3]). The *co-labeled Hodge-Tate character* associated with ρ is defined by

$$\mathcal{HT}(\rho) := (\mathcal{HT}(\rho)^\sigma)_{\sigma \in \Sigma_L} \in \prod_{\sigma \in \Sigma_L} X_*(G_{\mathbb{C}_p}).$$

- (2) A cocharacter of $G_{\mathbb{C}_p}$ is said to be *regular* if it is not killed by any roots of $G_{\mathbb{C}_p}$ (with respect to a fixed maximal torus of $G_{\mathbb{C}_p}$). For a Hodge-Tate G -valued representation $\rho: \mathrm{Gal}_K \rightarrow G(L)$, we say that ρ has *regular Hodge-Tate weights* if for each $\sigma \in \Sigma_L$, the cocharacter $\mathcal{HT}(\rho)^\sigma$ of $G_{\mathbb{C}_p}$ is regular.

4. CRYSTALLINE LIFTS OF QUASI-SEMISIMPLE REPRESENTATIONS WITH FIXED ABELIANIZATIONS

We prove the first main theorem in this section. In addition to Lin's argument [9, Proposition 2, Theorem 6], we use the calculation of the norms of Galois characters due to Böckle-Iyengar-Paškūnas [4, Lemma 2.1] in order to construct a crystalline lift with a fixed abelianization.

We call the homomorphism $K^* \rightarrow \mathcal{O}_K^*$ defined by $\varpi_K \mapsto 1$, $\mathcal{O}_K^* \ni a \mapsto a$ the *fundamental character (of K)*, and the composite of the fundamental character with the reduction map $\mathcal{O}_K^* \rightarrow k^*$ the *mod ϖ_K fundamental character (of K)*. Write $\omega_K: K^* \rightarrow \mathcal{O}_K^*$ for the fundamental character of K . Note that the fundamental character is the continuous character of K^* associated with the Lubin-Tate character, which is a crystalline character $\mathrm{Gal}_K \rightarrow \mathcal{O}_K^*$. Let $\mathrm{unr}(c): K^* \rightarrow \mathcal{O}_L^*$ be the unramified character which sends the uniformizer ϖ_K to an element $c \in \mathcal{O}_L^*$. We recall basic properties of crystalline characters of Gal_K .

Lemma 4.1. *The continuous character $K^* \rightarrow \mathcal{O}_L^*$ associated with a crystalline character $\mathrm{Gal}_K \rightarrow \mathcal{O}_L^*$ can be written as the twist of $\mathrm{unr}(c)$ for an element $c \in \mathcal{O}_L^*$ with a homomorphism $x \mapsto \prod_{\sigma: K \rightarrow L} \sigma(\omega_K(x))^{a_\sigma}$ where $a_\sigma \in \mathbb{Z}$ for each $\sigma \in \mathrm{Hom}_{\mathbb{Q}_p}(K, L)$.*

Proof. See [6, Proposition B.4]. □

Lemma 4.2. *Let $f := [k_L : k]$ and K_f be an unramified extension of K of degree f . Let G be a connected split reductive group over K and $T \subset G$ be a K -split maximal torus. Let $M_{T, \mathrm{crys}}^0 \subset M_{T, \mathrm{crys}}$ be the $\mathbb{Z}[W(G, T)] \otimes \mathbb{Z}[\mathrm{Gal}_K / I_K]$ -submodule which consists of continuous $T(\mathcal{O}_L)$ -valued representations of I_K that can be extended to a continuous representation $\mathrm{Gal}_{K_f} \rightarrow T(\mathcal{O}_L)$. Then the natural reduction map $M_{T, \mathrm{crys}}^0 \rightarrow M_{T, k_L}$ is surjective.*

Proof. The group M_{T, k_L} is generated by the mod ϖ_{K_f} fundamental character. Since it lifts to the fundamental character of K_f which extends to a crystalline character of Gal_{K_f} (i.e. the Lubin-Tate character of Gal_{K_f}), the assertion follows. □

The following lemma is used to take a crystalline lift which has regular Hodge-Tate weights.

Lemma 4.3. *Let G be a connected split reductive group over K and $T \subset G$ be a K -split maximal torus. Consider the map*

$$w - 1: X_*(T) \rightarrow X_*(T), \quad \chi \mapsto \mathrm{ad}(w)(\chi) \cdot \chi^{-1}$$

for every non-trivial element $w \in W(G, T)$. Let $X_*^{\text{ab}}: X_*(T) \rightarrow X_*(G^{\text{ab}})$ be the homomorphism of the cocharacter groups induced by the composite $T \hookrightarrow G \twoheadrightarrow G^{\text{ab}}$. Then the property

(R_w) : the rank of $\ker(X_*^{\text{ab}})/(\ker(X_*^{\text{ab}}) \cap \ker(w - 1))$ is positive

is satisfied for every non-trivial element $w \in W(G, T)$.

Proof. Let $G^{\text{ad}} := G/Z^\circ$, which is a split reductive group. If Z° is trivial, the condition (R_w) is satisfied for every non-trivial element $w \in W(G, T)$ of the Weyl group since w acts non-trivially on $X_*(T)$.

For general G , there exists a decomposition

$$X_*(T) \otimes_{\mathbb{Z}} \mathbb{Q} \cong (X_*(Z^\circ) \otimes_{\mathbb{Z}} \mathbb{Q}) \oplus (X_*(G^{\text{ad}}) \otimes_{\mathbb{Z}} \mathbb{Q})$$

since the composite $Z^\circ \hookrightarrow G \twoheadrightarrow G^{\text{ab}}$ is an isogeny. Since the center of G^{ad} is trivial, the lemma follows from the case where Z° is trivial. \square

We prove the first main theorem of this paper.

Theorem 4.4. *Let G be a connected split reductive group over \mathcal{O}_L and T be a split maximal torus of G over \mathcal{O}_L . Assume that the composite $T \hookrightarrow G \twoheadrightarrow G^{\text{ab}}$ has a right inverse.*

Let $\bar{\rho}: \text{Gal}_K \rightarrow G(k_L)$ be a quasi-semisimple G -valued representation such that $\bar{\rho}(I_K) \subset T(k_L)$ and $\bar{\rho}(\text{Gal}_K) \subset N_G(T)(k_L)$. Let $\bar{\rho}^{\text{ab}}: \text{Gal}_K \rightarrow G^{\text{ab}}(k_L)$ be the abelianization of $\bar{\rho}$ and $\psi: \text{Gal}_K \rightarrow G^{\text{ab}}(\mathcal{O}_L)$ be a crystalline lift of $\bar{\rho}^{\text{ab}}$. Then there exist a finite extension L'/L and a crystalline lift $\rho: \text{Gal}_K \rightarrow G(\mathcal{O}_{L'})$ with regular Hodge-Tate weights such that $\rho^{\text{ab}} = \psi$.

Moreover, if we do not impose a crystalline lift ρ has regular Hodge-Tate weights, L'/L can be taken to be a finite unramified extension.

Proof. Step 1.

First, we introduce notations in [9, Proposition 2] needed in the proof of the theorem. We naturally consider $M_{T, \text{crys}}$ and M_{T, k_L} as $\mathbb{Z}[W(G, T)] \otimes \mathbb{Z}[\text{Gal}_K / I_K]$ -modules. For an element $\tau \in W(G, T)$, write

$$\begin{aligned} M_{T, \tau, \text{crys}} &:= \ker(M_{T, \text{crys}} \xrightarrow{\tau \otimes 1 - 1 \otimes \Phi_K} M_{T, \text{crys}}), \\ M_{T, \tau, k_L} &:= \ker(M_{T, k_L} \xrightarrow{\tau \otimes 1 - 1 \otimes \Phi_K} M_{T, k_L}) \end{aligned}$$

for subgroups of $M_{T, \text{crys}}$ and M_{T, k_L} .

From here, let $\tau \in W(G, T)$ be the image of $\bar{\rho}(\Phi_K^{-1})$ by the quotient map $\zeta: N_G(T)(k_L) \rightarrow W(G, T)$. Then we have $\bar{\rho}|_{I_K} \in M_{T, \tau, k_L}$.

Let $f := [k_L : k]$ and K_f be an unramified extension of K of degree f . We may assume $\tau^f = 1$ after replacing L with its finite unramified extension. We put

$$\Xi := \sum_{i=0}^{f-1} \tau^i \otimes \Phi_K^{f-1-i}.$$

Let $M_{T, \text{crys}}^0 \subset M_{T, \text{crys}}$ be the $\mathbb{Z}[W(G, T)] \otimes \mathbb{Z}[\text{Gal}_K / I_K]$ -submodule which consists of continuous $T(\mathcal{O}_L)$ -valued representations of I_K that can be extended to a continuous representation $\text{Gal}_{K_f} \rightarrow T(\mathcal{O}_L)$. We put $M_{T, \tau, \text{crys}}^0 := M_{T, \text{crys}}^0 \cap M_{T, \tau, \text{crys}}$. Note that the natural reduction map $M_{T, \text{crys}}^0 \rightarrow M_{T, k_L}$ is surjective by Lemma 4.2.

Since $(\tau \otimes 1)^f = (1 \otimes \Phi_K)^f = \text{id}$ on both $M_{T,\text{crys}}$ and M_{T,k_L} , we have an injection $\Xi M_{T,\text{crys}}^0 \hookrightarrow M_{T,\tau,\text{crys}}^0$. Then it follows that the composite

$$\Xi M_{T,\text{crys}}^0 \hookrightarrow M_{T,\tau,\text{crys}}^0 \hookrightarrow M_{T,\tau,\text{crys}} \rightarrow M_{T,\tau,k_L}$$

is the same as the composite of the natural surjection $\Xi M_{T,\text{crys}}^0 \rightarrow \Xi M_{T,k_L}$ and the natural isomorphism

$$\Xi M_{T,k_L} \cong M_{T,\tau,k_L}$$

in the proof of [9, Proposition 2], and thus the composite $\Xi M_{T,\text{crys}}^0 \rightarrow M_{T,\tau,k_L}$ is surjective.

Step 2.

Next, we prove the existence of crystalline lifts with fixed abelianizations. Fix isomorphisms $\mathbb{G}_m^{d_0} \times \mathbb{G}_m^{d_1} \cong G^{\text{ab}} \times T' \cong T$ for a split torus T' by the assumption of the existence of a right inverse $G^{\text{ab}} \rightarrow T$.

For $A = k_L$ or \mathcal{O}_L and a continuous representation $r: H \rightarrow T(A)$ of a locally profinite group H , we write $r_i: H \rightarrow \mathbb{G}_m(A) = A^*$ for the i -th component of r via the isomorphism $\mathbb{G}_m^{d_0} \times \mathbb{G}_m^{d_1} \cong T$ for each $i \in [1, d_0 + d_1]$.

Let $\bar{\rho}' \in M_{T,k_L}$ be a continuous representation such that $\Xi \bar{\rho}'$ is mapped to $\bar{\rho}|_{I_K}$ by the natural isomorphism $\Xi M_{T,k_L} \cong M_{T,\tau,k_L}$ in Step 1. We construct an element $\rho' \in M_{T,\text{crys}}^0$ as follows:

- For $d_0 + 1 \leq i \leq d_0 + d_1$, take $\rho'_i \in M_{\mathbb{G}_m,\text{crys}}^0$ to be mapped to $\bar{\rho}'_i$ by the surjection $M_{\mathbb{G}_m,\text{crys}}^0 \rightarrow M_{\mathbb{G}_m,k_L}$ (see Lemma 4.2).
- For $1 \leq i \leq d_0$, let $\psi_i: \text{Gal}_K \rightarrow \mathcal{O}_L^*$ be the i -th component of ψ . The character $\psi'_i: K^* \rightarrow \mathcal{O}_L^*$ associated with ψ_i may be written as the twist of an unramified character $\text{unr}(c_i)$ with $x \mapsto \prod_{\sigma: K \rightarrow L} \sigma(\omega_K(x))^{a_{\sigma,i}}$, where $c_i \in \mathcal{O}_L^*$, $a_{\sigma,i} \in \mathbb{Z}$ for each $\sigma \in \text{Hom}_{\mathbb{Q}_p}(K, L)$ by Lemma 4.1. Then there exist distinct integers $b_{s,i}$ ($s \in \text{Hom}_{\mathbb{Q}_p}(K_f, L)$) such that the character

$$\theta: \mathcal{O}_{K_f}^* \rightarrow \mathcal{O}_L^*, \quad x \mapsto \prod_{s \in \text{Hom}_{\mathbb{Q}_p}(K_f, L)} s(x)^{b_{s,i}}$$

satisfies the following conditions by [4, Lemma 2.5].

- The integers $b_{s,i}$ ($s \in \text{Hom}_{\mathbb{Q}_p}(K_f, L)$) are pairwise distinct.
- The character θ lifts the character $\mathcal{O}_{K_f}^* \rightarrow k_L^*$ associated with the composite of $I_{K_f} \xrightarrow{\cong} I_K$ and $\bar{\rho}'_i$.
- $\theta(x) = \psi'_i(x)$ for all $x \in \mathcal{O}_K^*$.

Let $\rho'_i: I_K \rightarrow \mathcal{O}_L^*$ be the composite of the following maps

$$I_K = I_{K_f} \hookrightarrow W_{K_f} \twoheadrightarrow W_{K_f}^{\text{ab}} \xrightarrow{\cong} K_f^* \xrightarrow{\text{rec}_{K_f}^{-1}} K_f^* \xrightarrow{\prod_{s: K_f \rightarrow L} s(\omega_{K_f}(-))^{b_{s,i}}} L^*,$$

Then the continuous $T(\mathcal{O}_L)$ -valued representation $\rho' \in M_{T,\text{crys}}^0$ constructed as above satisfies the following properties:

- The image of ρ' by the surjection $M_{T,\text{crys}}^0 \rightarrow M_{T,k_L}$ is $\bar{\rho}'$.
- The abelianization $\rho'^{\text{ab}}: I_K \rightarrow G^{\text{ab}}(\mathcal{O}_L)$ is extendable to a crystalline representation $\psi': \text{Gal}_{K_f} \rightarrow G^{\text{ab}}(\mathcal{O}_L)$ such that $\psi|_{W_K}$ is the same as the composite

$$W_K \twoheadrightarrow W_K^{\text{ab}} \xrightarrow{\cong} K^* \hookrightarrow K_f^* \xrightarrow{\cong} W_{K_f}^{\text{ab}} \xrightarrow{\psi'|_{W_{K_f}}} G^{\text{ab}}(\mathcal{O}_L).$$

Let R' and $\Xi R'$ be the $T(\mathcal{O}_L)$ -valued crystalline representations $\text{Gal}_{K_f} \rightarrow T(\mathcal{O}_L)$ extended from ρ' and $\Xi\rho'$ respectively as above. By the calculation in the proof of [4, Lemma 2.1], we have

$$\begin{aligned} (\Xi R')^{\text{ab}}(\text{rec}_{K_f}(y)) &= \prod_{\sigma \in \text{Gal}(K_f/K)} R'^{\text{ab}}(\sigma \text{rec}_{K_f}(y) \sigma^{-1}) \\ &= \prod_{\sigma \in \text{Gal}(K_f/K)} R'^{\text{ab}}(\text{rec}_{K_f}(\sigma(y))) \\ &= R'^{\text{ab}}(\text{rec}_{K_f}(\text{Nm}_{K_f/K}(y))) \\ &= \psi|_{W_K}(\text{rec}_K(\text{Nm}_{K_f/K}(y))) \\ &= \psi|_{W_{K_f}}(\text{rec}_{K_f}(y)) \end{aligned}$$

for any $y \in K_f^*$. Here, the first equality follows from the fact that each τ^i acts trivially on the abelianization ρ'^{ab} , the fourth equality follows from the second bullet of the properties of the representation $\rho' \in M_{T, \text{crys}}^0$, and the last equality follows from $\text{Nm}_{K_f/K} \circ \text{rec}_{K_f}^{-1} = \text{rec}_K^{-1} \circ \lambda$ by local class field theory, where $\lambda: W_{K_f}^{\text{ab}} \rightarrow W_K^{\text{ab}}$ is induced by the inclusion $W_{K_f} \hookrightarrow W_K$. So we have $(\Xi R')^{\text{ab}}|_{W_{K_f}} = \psi|_{W_{K_f}}$, and by continuity $(\Xi R')^{\text{ab}} = \psi|_{\text{Gal}_{K_f}}$. Moreover, it follows that the image of $\Xi\rho'$ by the surjection $\Xi M_{T, \text{crys}}^0 \rightarrow M_{T, \tau, k_L}$ is $\Xi \bar{\rho}' = \bar{\rho}|_{I_K}$ by the construction.

Let $w \in N_G(T)(\mathcal{O}_L)$ be an element which is a lift of the element $\bar{\rho}(\Phi_K)^{-1}$. We may assume that the image of w by the composite $N_G(T)(\mathcal{O}_L) \hookrightarrow G(\mathcal{O}_L) \rightarrow G^{\text{ab}}(\mathcal{O}_L)$ is $\psi(\Phi_K)^{-1}$ by the assumption of the existence of a right inverse of the map $T \rightarrow G^{\text{ab}}$.

Since $\Xi M_{T, \text{crys}}^0 \subset M_{T, \tau, \text{crys}}^0$, there exists a continuous representation $\rho: \text{Gal}_K \rightarrow N_G(T)(\mathcal{O}_L) \subset G(\mathcal{O}_L)$ such that $\rho|_{\text{Gal}_{K_f}} = \Xi R'$ and $\rho^{\text{ab}} = \psi$ by Lemma 3.5 (1) where w is taken as above. Then ρ is a crystalline lift of $\bar{\rho}$ satisfying $\rho^{\text{ab}} = \psi$. But the Hodge-Tate weights of ρ are not necessarily regular.

Step 3.

Finally, we show the existence of a crystalline lift with fixed abelianization which has regular Hodge-Tate weights. We prove it by an argument of Lin [9, Lemma 12, Theorem 6].

Let $i: T \hookrightarrow G$ be the embedding and fix embeddings $K_f \subset L \subset \mathbb{C}_p$. For constructing such a crystalline lift with regular Hodge-Tate weights, we first show the existence of a crystalline representation $t: \text{Gal}_{K_f} \rightarrow T(\mathcal{O}_L)$ satisfying the following properties for each $\sigma \in \text{Hom}_{\mathbb{Q}_p}(L, \mathbb{C}_p)$:

- $\mathcal{HT}(i(t))^\sigma$ is regular if $\sigma|_{K_f}$ coincides with the fixed embedding $K_f \hookrightarrow \mathbb{C}_p$.
- $\mathcal{HT}(i(t))^\sigma$ is the trivial cocharacter if $\sigma|_{K_f}$ does not coincide with the fixed embedding $K_f \hookrightarrow \mathbb{C}_p$.
- t^{ab} is trivial.

The existence of such t is reduced to show $\ker(X_*^{\text{ab}}) \not\subset \bigcup_{w \in W(G, T) \setminus \{e\}} \ker(w - 1)$ by considering powers of the fundamental character of K_f for each coordinate of $T(\mathcal{O}_L) \cong (\mathcal{O}_L^*)^{d_0 + d_1}$ (see the proof of [9, Lemma 12]). In fact, let

$$x \in \ker(X_*^{\text{ab}}) \setminus \left(\bigcup_{w \in W(G, T) \setminus \{e\}} \ker(w - 1) \right).$$

Then the composite

$$W_{K_f} \twoheadrightarrow W_{K_f}^{\text{ab}} \xrightarrow[\cong]{\text{rec}_{K_f}^{-1}} K_f^* \hookrightarrow L^* \cong \mathbb{G}_m(L) \xrightarrow{x(L)} T(L)$$

extends to a required crystalline representation $t: \text{Gal}_{K_f} \rightarrow T(L)$. The assertion follows from Lemma 4.3.

For such t , let $v_0 \in M_{T, \text{crys}}^0$ be the composite of the following maps

$$I_K = I_{K_f} \xrightarrow{t|_{I_{K_f}}} T(\mathcal{O}_L).$$

Then we have $v_0^{\text{ab}} = 1$. For a sufficiently large positive integer M , the representation $v' := v \cdot (\Xi v_0)^{MN}$, where $v := \rho|_{I_K}$ and N is the cardinality of k_L^* , satisfies $v'^{\text{ab}} = (v \cdot (\Xi v_0)^{MN})^{\text{ab}} = v^{\text{ab}}$ and $\overline{v'} = \overline{v}$. Since $v, \Xi v_0 \in \Xi M_{T, \text{crys}}^0 \subset M_{T, \tau, \text{crys}}^0$, the representation v' extends to a continuous representation $\rho'': \text{Gal}_K \rightarrow G(\mathcal{O}_L)$, which is a crystalline lift of $\overline{\rho}$ by $K_f \subset L$ and [9, Corollary 2]. We have $\rho''^{\text{ab}}|_{I_K} = \rho^{\text{ab}}|_{I_K}$ by the construction. So replacing L with its finite extension, there is an unramified representation $u: \text{Gal}_K \rightarrow Z^\circ(\mathcal{O}_L)$ whose composite with the map $Z^\circ(\mathcal{O}_L) \rightarrow G^{\text{ab}}(\mathcal{O}_L)$ is $(\rho''^{\text{ab}})^{-1} \cdot \rho^{\text{ab}}$ since $Z^\circ \rightarrow G^{\text{ab}}$ is an isogeny of algebraic tori. Then $\rho_{\text{reg}} := u\rho''$ is a crystalline lift of $\overline{\rho}$ which satisfies $\rho_{\text{reg}}^{\text{ab}} = \rho^{\text{ab}} = \psi$ by the construction. In addition, the representation ρ_{reg} has regular Hodge-Tate weights since for each $\sigma \in \text{Hom}_{\mathbb{Q}_p}(L, \mathbb{C}_p)$, there exists an integer $i \in [0, f-1]$ and an element $\sigma' \in \text{Hom}_{K_f}(L, \mathbb{C}_p)$ such that $\sigma = \sigma' \circ \Phi_K^{f-1-i}$ and

$$\mathcal{HT}(\Xi v_0)^\sigma = \tau^i \mathcal{HT}(v_0)^{\sigma'} \tau^{-i}$$

by the calculation in the proof of [9, Theorem 6]. \square

5. POTENTIALLY CRYSTALLINE LIFTS OF SEMISIMPLE L -PARAMETERS WITH FIXED ABELIANIZATIONS

In this section, we also prove the existence of potentially crystalline lifts with a fixed potentially crystalline abelianization of semisimple mod p L -parameters for connected quasi-split tame groups.

First, we recall the definitions of L -groups and L -parameters following [10, §2, §3]. Let K/\mathbb{Q}_p be a finite extension and G be a connected quasi-split reductive group over K . Fix a K -pinning $(B, T, \{X_\alpha\})$ of G . The pinned group $(G, B, T, \{X_\alpha\})$ has a dual pinned group $(\widehat{G}, \widehat{B}, \widehat{T}, \{Y_\alpha\})$ defined over \mathbb{Z} . Let $E \subset \overline{K}$ be a splitting field of G , which is a finite Galois extension over K . The Galois action on the based root datum induces a Galois action $\text{Gal}(E/K) \rightarrow \text{Aut}(\widehat{G}, \widehat{B}, \widehat{T}, \{Y_\alpha\})$. Write ${}^L G := \widehat{G} \rtimes \text{Gal}_K$ (see [5] and [13, §1.1] for more details). Write ${}^L \{*\} \cong \text{Gal}_K$ for the L -group of the trivial group.

Definition 5.1 (cf. [10, Definition 3.1.1, Definition 3.1.3]). Let G, H be connected reductive groups over K and \widehat{G}, \widehat{H} be their Langlands dual groups.

- (1) For a finite ring A , a group homomorphism

$$\widehat{G}(A) \rtimes \text{Gal}(E/K) \rightarrow \widehat{H}(A) \rtimes \text{Gal}(E/K)$$

is said to be an L -homomorphism (with A -coefficients) if each element $g \times \sigma \in \widehat{G}(A) \rtimes \text{Gal}(E/K)$ is sent to $h \times \sigma \in \widehat{H}(A) \rtimes \text{Gal}(E/K)$.

- (2) For a finite ring
- A
- , a group homomorphism

$$\widehat{G}(A) \rtimes W_{E/K} \rightarrow \widehat{H}(A) \rtimes W_{E/K}$$

is said to be an L -homomorphism (with A -coefficients) if each element $g \times \sigma \in \widehat{G}(A) \rtimes \text{Gal}(E/K)$ is sent to $h \times \sigma \in \widehat{H}(A) \rtimes \text{Gal}(E/K)$.

- (3) For a finite ring
- A
- , a group homomorphism

$$\widehat{G}(A) \rtimes \text{Gal}_K = {}^L G(A) \rightarrow {}^L H(A) = \widehat{H}(A) \rtimes \text{Gal}_K$$

is said to be an L -homomorphism (with A -coefficients) if each element $g \times \sigma \in \widehat{G}(A) \rtimes \text{Gal}_K$ is sent to $h \times \sigma \in \widehat{H}(A) \rtimes \text{Gal}_K$, and there exists an open subgroup $\Gamma \subset \text{Gal}_K$ such that $1 \times \sigma \in \widehat{G}(A) \rtimes \text{Gal}_K$ is sent to $1 \times \sigma \in \widehat{H}(A) \rtimes \text{Gal}_K$ for every $\sigma \in \Gamma$.

- (4) An
- L
- parameter (of
- Gal_K
- (resp.
- $\text{Gal}(E/K)$
- ,
- $W_{E/K}$
-) for
- G
- with
- A
- coefficients) is an
- L
- homomorphism

$$\text{Gal}_K \simeq {}^L \{*\}(A) \rightarrow {}^L G(A),$$

$$\text{(resp. } \text{Gal}(E/K) \rightarrow \widehat{G}(A) \rtimes \text{Gal}(E/K), W_{E/K} \rightarrow \widehat{G}(A) \rtimes W_{E/K}),$$

defined up to $\widehat{G}(A)$ -conjugacy. Note that an L -parameter can be considered as an ${}^L G(A)$ -valued Galois representation $\text{Gal}_K \rightarrow {}^L G(A)$.

- (5) For a profinite ring R , an L -parameter (for G with R -coefficients) is a compatible system of L -parameters $\text{Gal}_K \rightarrow {}^L G(A)$, where A is a finite quotient of R .
- (6) For $A = \overline{\mathbb{F}}_p$ or $\overline{\mathbb{Z}}_p$, an L -parameter (for G with A -coefficients) is the composite of an L -parameter $\text{Gal}_K \rightarrow {}^L G(A')$ for a finitely generated \mathbb{Z}_p -algebra $A' \subset A$ and the inclusion ${}^L G(A') \hookrightarrow {}^L G(A)$.

Definition 5.2 (cf. [10, Definition 5.1.1]). Let $\Lambda \subset \overline{\mathbb{Z}}_p$ be a discrete valuation ring. For a connected quasi-split reductive group G over K which splits over a finite Galois extension E/K , an L -parameter $\rho: \text{Gal}_K \rightarrow \widehat{G}(\Lambda) \rtimes \text{Gal}(E/K)$ is said to be *crystalline* (resp. *potentially crystalline*) if for some closed embeddings of algebraic groups $\widehat{G} \rtimes \text{Gal}_{E/K} \hookrightarrow \text{GL}_d$ for some $d \geq 1$, the composite $\text{Gal}_K \rightarrow \widehat{G}(\Lambda) \rtimes \text{Gal}(E/K) \rightarrow \text{GL}_d(\Lambda)$ is crystalline (resp. *potentially crystalline*).

An L -parameter $\bar{\rho}: \text{Gal}_K \rightarrow {}^L G(\overline{\mathbb{F}}_p)$ is said to be *semisimple* if $\bar{\rho}$ is semisimple in the sense of [10, Definition 3.2.5].

The following Langlands-Shelstad factorizations of L -parameters are shown in [10].

Theorem 5.3 ([10, Theorem 3.4.1]). *For a semisimple L -parameter $\bar{\rho}: \text{Gal}_K \rightarrow {}^L G(\overline{\mathbb{F}}_p)$, there exists a tame K -torus S of G and a canonical L -homomorphism ${}^L j: {}^L S_{\overline{\mathbb{F}}_p} \rightarrow {}^L G_{\overline{\mathbb{F}}_p}$ such that $\bar{\rho}$ factors through ${}^L j$.*

For a semisimple L -parameter $\bar{\rho}: \text{Gal}_K \rightarrow {}^L G(\overline{\mathbb{F}}_p)$, we fix a tame K -torus $S \subset G$ satisfying the condition of Theorem 5.3 in the assertion of Theorem 5.5.

For $A = \overline{\mathbb{Z}}_p^*$ or $\overline{\mathbb{F}}_p^*$, note that the module $H_{\text{cont}}^1(W_{E/K}, X^*(T_E) \otimes_{\mathbb{Z}} A)$ (resp. $H_{\text{cont}}^1(\text{Gal}_K, X^*(T_E) \otimes_{\mathbb{Z}} A)$) is naturally bijective to the set of $\widehat{G}(A)$ -conjugacy classes of L -parameters $W_{E/K} \rightarrow \widehat{G}(A) \rtimes W_{E/K}$ (resp. $\text{Gal}_K \rightarrow {}^L G(A) \rtimes \text{Gal}_K$). We also use the p -adic local Langlands correspondence for tori with divisible coefficients.

Theorem 5.4 ([3]). *For a K -torus T with splitting field E , which is a finite Galois extension over K , and a divisible abelian topological group D , there exists a natural isomorphism*

$$H_{\text{cont}}^1(W_{E/K}, X^*(T_E) \otimes_{\mathbb{Z}} D) \xrightarrow{\sim} \text{Hom}_{\text{cont}}(T(K), D).$$

Write $T_E := T \times_K E$. Then the following natural maps

$$\begin{aligned} H_{\text{cont}}^1(\text{Gal}_K, X^*(T_E) \otimes_{\mathbb{Z}} \overline{\mathbb{F}}_p^*) &\rightarrow H_{\text{cont}}^1(W_{E/K}, X^*(T_E) \otimes_{\mathbb{Z}} \overline{\mathbb{F}}_p^*), \\ H_{\text{cont}}^1(\text{Gal}_K, X^*(T_E) \otimes_{\mathbb{Z}} \overline{\mathbb{Z}}_p^*) &\rightarrow H_{\text{cont}}^1(W_{E/K}, X^*(T_E) \otimes_{\mathbb{Z}} \overline{\mathbb{Z}}_p^*), \end{aligned}$$

are also isomorphisms.

Proof. For the first isomorphism, see [3], [10, Theorem 4.3.1]. Let A be a finite discrete ring. Then the set of L -homomorphisms $\rho: W_{E/K} \rightarrow \widehat{T}(A) \rtimes W_{E/K}$ is naturally bijective to the set of L -homomorphisms $\rho: W_K \rightarrow \widehat{T}(A) \rtimes W_K$ since W_E acts trivially on \widehat{T} .

By [10, Lemma 3.1.2 (2), (4)], an L -homomorphism $\rho: W_K \rightarrow \widehat{T}(A) \rtimes W_K$ naturally corresponds to an L -parameter $\rho: \text{Gal}_K \rightarrow {}^L G(A)$. So we have an isomorphism

$$H_{\text{cont}}^1(\text{Gal}_K, X^*(T_E) \otimes_{\mathbb{Z}} A) \xrightarrow{\sim} H_{\text{cont}}^1(W_{E/K}, X^*(T_E) \otimes_{\mathbb{Z}} A) \quad (5.1)$$

It follows that the last maps are isomorphisms by considering the (co)limits of the isomorphisms (5.1) for finite discrete coefficients A . \square

For a K -torus S and an L -parameter $\bar{\rho}: \text{Gal}_K \rightarrow {}^L S(\overline{\mathbb{F}}_p)$, let $[\bar{\rho}]: \text{Gal}_K \rightarrow {}^L S(\overline{\mathbb{Z}}_p)$ be the L -parameter induced by the isomorphisms in Theorem 5.4 for $D = \overline{\mathbb{F}}_p^*$ and the Teichmüller character $\overline{\mathbb{F}}_p^* \hookrightarrow \overline{\mathbb{Z}}_p^* (\subset W(\overline{\mathbb{F}}_p)^*)$ from $\bar{\rho}$.

Theorem 5.5. *Let G be a connected quasi-split tame group over K and $\bar{\rho}: \text{Gal}_K \rightarrow {}^L G(\overline{\mathbb{F}}_p)$ be a semisimple mod p L -parameter. Fix a tame K -torus S of G such that $\bar{\rho}$ factors through ${}^L S(\overline{\mathbb{F}}_p)$. Assume that the composite $S \hookrightarrow G \twoheadrightarrow G^{\text{ab}}$ has a right inverse, which is written as $i: G^{\text{ab}} \rightarrow S$.*

For $A = \overline{\mathbb{F}}_p^$ or $\overline{\mathbb{Z}}_p^*$ and an L -parameter $\tau: \text{Gal}_K \rightarrow {}^L S(A)$, write τ^{ab} for the composite of τ and the map ${}^L S(A) \rightarrow {}^L G^{\text{ab}}(A)$ induced by the composite $G^{\text{ab}} \xrightarrow{i} S \hookrightarrow G$ and we set $\bar{\psi} := \bar{\rho}^{\text{ab}}$.*

Let $\psi: \text{Gal}_K \rightarrow {}^L G^{\text{ab}}(\overline{\mathbb{Z}}_p)$ be a potentially crystalline lift of $\bar{\psi}$. Then $\bar{\rho}$ admits a potentially crystalline lift $\rho: \text{Gal}_K \rightarrow {}^L G(\overline{\mathbb{Z}}_p)$ which has regular Hodge-Tate weights and satisfies $\rho^{\text{ab}} = \psi$.

Proof. Let E be a finite Galois extension of K such that the image of $\bar{\rho}$ is contained in ${}^L G(E)$, S splits over E , $\psi|_{\text{Gal}_E}$ is crystalline, the character $\text{Gal}_K \rightarrow k^*$ associated with the mod ϖ_K fundamental character is trivial on Gal_E . We fix an embedding $E \hookrightarrow \overline{K}$ and a uniformizer $\varpi_E \in E$.

For a K -torus T which splits over E , let $\text{Nm}_{E/K}: T(E) \rightarrow T(K)$ be the norm map induced by $\text{Nm}_{E/K}: \text{Res}_{E/K} T_E \rightarrow T$, where $T_E := S \times_K E$. By Theorem 5.4, we have the following commutative diagram

$$\begin{array}{ccccccc} H_K^1(G^{\text{ab}}) & \xrightarrow{\cong} & \text{Hom}_{\text{cont}}(G^{\text{ab}}(K), \overline{\mathbb{Z}}_p^*) & \longrightarrow & \text{Hom}_{\text{cont}}(S(K), \overline{\mathbb{Z}}_p^*) & \xleftarrow{\cong} & H_K^1(S) \\ \downarrow \text{-}|_{\text{Gal}_E} & & \downarrow \text{-}\circ\text{Nm}_{E/K} & & \downarrow \text{-}\circ\text{Nm}_{E/K} & & \downarrow \text{-}|_{\text{Gal}_E} \\ H_E^1(G^{\text{ab}}) & \xrightarrow{\cong} & \text{Hom}_{\text{cont}}(G^{\text{ab}}(E), \overline{\mathbb{Z}}_p^*) & \longrightarrow & \text{Hom}_{\text{cont}}(S(E), \overline{\mathbb{Z}}_p^*) & \xleftarrow{\cong} & H_E^1(S) \end{array}$$

where $H_K^1(-) := H_{\text{cont}}^1(\text{Gal}_K, X^*(- \times_K \overline{K}) \otimes_{\mathbb{Z}} \overline{\mathbb{Z}}_p^*)$.

By the existence of the right inverse of the map $S \rightarrow G^{\text{ab}}$, we have $S \cong G^{\text{ab}} \times S'$ for some K -torus S' . Let $\overline{\rho}_{S'}: \text{Gal}_K \rightarrow {}^L S'(\overline{\mathbb{F}}_p)$ be the ${}^L S'(\overline{\mathbb{F}}_p)$ -component of $\overline{\rho}$. Then $[\overline{\rho}_{S'}]: \text{Gal}_K \rightarrow {}^L S'(\overline{\mathbb{Z}}_p)$ is a potentially crystalline L -parameter with its reduction $\overline{\rho}_{S'}$ since its image is finite. Let

$$\rho' := (\psi, [\overline{\rho}_{S'}]) \in H_K^1(G^{\text{ab}}) \oplus H_K^1(S') \cong H_K^1(S).$$

It follows that ρ' is a potentially crystalline lift of $\overline{\rho}$ and we have $\rho'^{\text{ab}} = \psi$ by the construction.

Note that $\text{Gal}(E/K)$ acts on $X_*(\widehat{S}_E) = X^*(S_E)$. For a character $x \in X^*(S_E)$, let $\widetilde{\chi}_x: S(E) \rightarrow \overline{\mathbb{Z}}_p^*$ be the composite of a character $x(E): S(E) \rightarrow E^*$, the fundamental character $E^* \rightarrow \mathcal{O}_E^*$, and a fixed inclusion $\mathcal{O}_E^* \hookrightarrow \overline{\mathbb{Z}}_p^*$.

By Step 3 of the proof of Theorem 4.4, there exists a cocharacter $x \in X_*(\widehat{S}_E)$ which is regular, i.e., not killed by any roots of \widehat{S}_E , and the trivial abelianization.

It follows that for any $\sigma \in \text{Gal}(E/K)$, the cocharacter $\sigma \cdot x \in X_*(\widehat{S}_E)$ is also regular, and the L -parameter corresponding to $\chi_x := \widetilde{\chi}_x|_{S(K)}$ has regular Hodge-Tate weights by [10, Lemma 5.2.6].

Then $\rho' \rho''^M$, where $\rho'' \in H_K^1(S)$ is the potentially crystalline representation corresponding to χ_x , has regular Hodge-Tate weights for a sufficiently large integer M . The L -parameter $\rho' \rho''^M [\overline{\rho}]^{-M}$ is what we want. \square

REFERENCES

- [1] K. Aoki, *Zariski density of crystalline points on GSp_{2n} -valued local deformation rings*, preprint, available at <https://arxiv.org/abs/2407.17005>.
- [2] S. Balaji, *G-valued potentially semi-stable deformation rings*, ProQuest LLC, Ann Arbor, MI, 2012, Thesis (Ph. D.)-The University of Chicago.
- [3] C. Birkbeck, *On the p -adic Langlands correspondence for algebraic tori*, J. Théor. Nombres Bordeaux **32** (2020), no. 1, 133-158.
- [4] G. Böckle, A. Iyengar, V. Paškūnas, *Zariski density of crystalline points*, Proc. Natl. Acad. Sci. USA **120** (2023), no. 13, Paper No. e2221042120, 7.
- [5] K. Buzzard, T. Gee, *The conjectural connections between automorphic representations and Galois representations*, Automorphic forms and Galois representations. Vol. 1, 135-187. London Math. Soc. Lecture Note Ser., **414**, Cambridge University Press, Cambridge, 2014.
- [6] B. Conrad, *Lifting global representations with local properties*, preprint, available at <https://math.stanford.edu/~conrad/papers/locchar.pdf>.
- [7] M. Emerton, T. Gee, *Moduli stacks of étale (φ, Γ) -modules and the existence of crystalline lifts*, Ann. of Math. Stud., **215**, Princeton University Press, Princeton, NJ, 2023.
- [8] A. J. de Jong, *Crystalline Dieudonné module theory via formal and rigid geometry*, Inst. Hautes Études Sci. Publ. Math. No. **82** (1995), 5-96.
- [9] Z. Lin, *Crystalline lifts and a variant of the Steinberg-Winter theorem*, Doc. Math. **27** (2022), 2441-2468.
- [10] Z. Lin, *A Deligne-Lustig type correspondence for tame p -adic groups*, preprint, available at <https://arxiv.org/abs/2306.02093>.
- [11] V. Paškūnas, J. Quast, *On local Galois deformation rings: generalised reductive groups*, preprint, available at <https://arxiv.org/abs/2404.14622>.
- [12] J. T. Tate, *Number theoretic background*, in Automorphic forms, representations and L -functions, Proceedings of Symposia in Pure Mathematics, vol. **33**, Part 2, American Mathematical Society, 1979, p. 3-26.
- [13] X. Zhu, *A note on Integral Satake isomorphisms*, available at <https://arxiv.org/abs/2005.13056>

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, KYOTO UNIVERSITY
Email address: aoki.kensuke.88s@st.kyoto-u.ac.jp