NEW WEIGHTED SPECTRAL GEOMETRIC MEAN AND QUANTUM DIVERGENCE

MIRAN JEONG, SEJONG KIM AND TIN-YAU TAM

ABSTRACT. A new class of weighted spectral geometric means has recently been introduced. In this paper, we present its inequalities in terms of the Löwner order, operator norm, and trace. Moreover, we establish a log-majorization relationship between the new spectral geometric mean, and the Rényi relative operator entropy. We also give the quantum divergence of the quantity, given by the difference of trace values between the arithmetic mean and new spectral geometric mean. Finally, we study the barycenter that minimizes the weighted sum of quantum divergences for given variables.

Mathematics Subject Classification (2020): 15B48, 81P17

Keywords: spectral geometric mean, Rényi relative operator entropy, log-majorization, quantum divergence, barycenter

1. INTRODUCTION

Kubo and Ando [16] introduced the concept of operator means for the set \mathbb{P} of positive invertible operators on a complex Hilbert space \mathcal{H} , associated with operator monotone functions. Their operator mean, denoted by σ , is defined for positive invertible operators A and B, as

$$A\sigma B = A^{1/2} f(A^{-1/2} B A^{-1/2}) A^{1/2},$$

where f is an operator monotone function on $(0, \infty)$. A fundamental example of Kubo-Ando's operator means is the (metric) geometric mean defined using $f(x) = x^t$ for $t \in [0, 1]$, and denoted by $A \sharp_t B$. In the finite-dimensional setting, \mathbb{P}_m , of $m \times m$ positive definite matrices, the geometric mean $A \sharp_t B$ has nice geometric meanings such as being the unique geodesic connecting A and B under the natural Riemannian trace distance

$$d_R(A,B) = \|\log(A^{-1/2}BA^{-1/2})\|_F,$$

where $\|\cdot\|_F$ denotes the Forbenius norm. In other words,

$$A\sharp_t B = \underset{X \in \mathbb{P}_m}{\operatorname{arg\,min}} (1-t) d_R^2(X, A) + t d_R^2(X, B).$$

This characterization allows the extension of two-variable geometric means to multivariable means, such as the Cartan mean.

On the other hand, various operator means that do not satisfy monotonicity under the Löwner partial order, known as non-Kubo-Ando means, have also been studied. A prominent example is the spectral geometric mean, whose weighted form is given in [18]:

$$A\natural_t B = (A^{-1}\sharp B)^t A (A^{-1}\sharp B)^t, \quad t \in [0,1].$$

Although the spectral geometric mean lacks monotonicity, it satisfies the in-betweenness property (a weaker version of monotonicity) with respect to the near-order [8]. Many results comparing the geometric mean and the spectral geometric mean have been established [7, 11, 12, 14, 15], including the following log-majorization relationship:

$$A \sharp_t B \prec_{\log} \exp((1-t)\log A + t\log B) \prec_{\log} A \natural_t B.$$

Recently, a new spectral geometric mean was introduced in [6]:

$$F_t(A,B) := (A^{-1} \sharp_t B)^{1/2} A^{2-2t} (A^{-1} \sharp_t B)^{1/2}, \quad t \in [0,1].$$

This is a path connecting A at t = 0 and B at t = 1, with $F_{1/2}(A, B) = A \natural_{1/2} B$. It shares properties with the spectral geometric mean $A \natural_t B$. In the context of positive invertible operators on \mathcal{H} , we derive interesting inequalities of this new mean in terms of Löwner order and operator norm. In the finite-dimensional setting \mathbb{P}_m , we establish a trace inequality and a log-majorization relationship between the new spectral geometric mean and Rényi relative operator entropy. These results allow us to define the quantity

$$\Phi(A, B) = \operatorname{tr} \left[A \nabla_t B - F_t(A, B) \right], \quad t \in [0, 1]$$

as a quantum divergence, analogous to the findings in [13]. Moreover, we study the barycenter as a multi-variable mean that minimizes the weighted sum of quantum divergences for given variables.

2. New weighted spectral geometric mean

Let $B(\mathcal{H})$ be the Banach space of all bounded linear operators on a Hilbert space \mathcal{H} over the complex field, and let $S(\mathcal{H}) \subset B(\mathcal{H})$ be the set of all self-adjoint operators. Let $\mathbb{P} \subset S(\mathcal{H})$ be the open convex cone of all positive definite operators. For the finitedimensional setting $\mathcal{H} = \mathbb{C}^m$, we denote by \mathbb{H}_m and \mathbb{P}_m the sets of all $m \times m$ Hermitian matrices and positive definite matrices, respectively.

The (metric) geometric mean of $A, B \in \mathbb{P}$ is defined as

$$A\sharp_t B := A^{1/2} (A^{-1/2} B A^{-1/2})^t A^{1/2}, \quad t \in [0, 1].$$

This mean represents the unique geodesic on \mathbb{P}_m joining A and B under the natural Riemannian trace metric. Note that $A \sharp B = A \sharp_{1/2} B$ is a unique positive definite solution of Riccati equation

$$XA^{-1}X = B.$$

The following is the well-known arithmetic-geometric-harmonic mean inequalities:

$$A!_t B \le A \sharp_t B \le A \nabla_t B, \quad t \in [0, 1],$$

where $A\nabla_t B = (1-t)A + tB$ and $A!_t B = [(1-t)A^{-1} + tB^{-1}]^{-1}$ are the arithmetic and harmonic means, respectively.

The spectral geometric mean of $A, B \in \mathbb{P}$ is defined as

$$A \natural_t B := (A^{-1} \sharp B)^t A (A^{-1} \sharp B)^t, \quad t \in [0, 1].$$

This mean has properties analogous to the (metric) geometric mean [11, 14, 15] and represents a geodesic on \mathbb{P} joining A and B under the semi-metric $d(A, B) = 2 || \log(A^{-1} \sharp B) ||$, where $|| \cdot ||$ denotes the operator norm. For simplicity, we write $A \natural B = A \natural_{\frac{1}{2}} B$.

Recently, a new weighted spectral geometric mean was introduced in [6]:

$$F_t(A,B) = (A^{-1} \sharp_t B)^{\frac{1}{2}} A^{2-2t} (A^{-1} \sharp_t B)^{\frac{1}{2}}, \quad t \in [0,1].$$

One can easily see that $F_0(A, B) = A$, $F_1(A, B) = B$, and $F_{\frac{1}{2}}(A, B) = A \natural B$. Moreover, $F_t(A, B)$ is a unique positive definite solution of the Riccati equation

$$(A^{-1}\sharp_t B)^{\frac{1}{2}} = A^{2t-2}\sharp X, \quad t \in [0,1].$$

The following properties of $F_t(A, B)$ were established in [6, Proposition 2.2] for the finite-dimensional setting \mathbb{P}_m , and the proofs extend to \mathbb{P} .

Proposition 2.1. Let $A, B \in \mathbb{P}$. For all $t \in [0, 1]$, the following properties hold.

- (1) $F_t(A, B) = A^{1-t}B^t$ if A, B commute.
- (2) $F_t(aA, bB) = a^{1-t}b^t F_t(A, B)$ for any a, b > 0.
- (3) $F_t(UAU^*, UBU^*) = UF_t(A, B)U^*$ for any unitary operator U.
- (4) $F_t(A^{-1}, B^{-1}) = F_t(A, B)^{-1}$.
- (5) $2(A\nabla_t B^{-1})^{-\frac{1}{2}} A^{2(t-1)} \le F_t(A, B) \le \left[2(A^{-1}\nabla_t B)^{-\frac{1}{2}} A^{2(1-t)}\right]^{-1}.$
- (6) $\lim_{t \to 0} F_t(A^s, B^s)^{1/s} = \exp((1-t)\log A + t\log B).$

3. Operator inequalities and eigenvalue relationships

The (metric) geometric mean $A \sharp_t B$ has a well-known property:

$$A\sharp_t B \le I \Longrightarrow A^p \sharp_t B^p \le I \quad \text{for all } p \ge 1.$$
(3.1)

This result, first discovered by Ando and Hiai [1], has been generalized to the Cartan mean for \mathbb{P}_n [21] and the Karcher mean for \mathbb{P} [17]. In the finite-dimensional case \mathbb{P}_m , the Ando-Hiai inequality can be expressed as the following log-majorization relationship:

$$(A^p \sharp_t B^p)^{1/p} \prec_{\log} (A^q \sharp_t B^q)^{1/q}, \quad 0 < q \le p \le 1.$$

Lemma 3.1. [10, Lemma 2.7] Let $S \in S(\mathcal{H})$ and $X \in \mathbb{P}$. Then $SXS \leq X$ implies $S \leq I$. Furthermore, SXS = X for $S, X \in \mathbb{P}$ if and only if S = I.

Using this result, we establish the following inequalities for the new weighted spectral geometric mean $F_t(A, B)$.

Proposition 3.2. Let $A, B \in \mathbb{P}$ and $t \in [0, 1]$. Then the following statements hold.

- (1) If $F_t(A^{-1}, B) \leq A^{2(t-1)}$, then $A^p \sharp_t B^p \leq I$ for all $p \geq 1$.
- (2) $F_t(A, B) = A^{2-2t}$ if and only if $B = A^{\frac{1}{t}-1}$ for $t \in (0, 1]$.

Proof. Note that $(A \sharp_t B)^{\frac{1}{2}} A^{2t-2} (A \sharp_t B)^{\frac{1}{2}} = F_t(A^{-1}, B) \leq A^{2(t-1)}$. From Lemma 3.1, we have $(A \sharp_t B)^{\frac{1}{2}} \leq I$, and thus $A \sharp_t B \leq I$ for $t \in [0, 1]$. By the Ando-Hiai inequality (3.1), $A^p \sharp_t B^p \leq I$ for all $p \geq 1$.

Furthermore, $F_t(A, B) = A^{2-2t}$ if and only if $A^{-1}\sharp_t B = I$ by Lemma 3.1, which is equivalent to $(A^{\frac{1}{2}}BA^{\frac{1}{2}})^t = A$. This yields $B = A^{\frac{1}{t}-1}$ for $t \in (0, 1]$.

Corollary 3.3. Let $A, B \in \mathbb{P}$ and $t \in (0, 1]$. Then $F_t(A, B) \leq A^{2-2t}$ for $A \leq I$ implies $B \leq A^{-1}$.

Proof. By Proposition 3.2, $F_t(A, B) \leq A^{2-2t}$ implies $(A^{-1}\sharp_t B)^{\frac{1}{2}} \leq I$ but not $(A^{-1}\sharp_t B) \leq I$. Since $A \leq I$, we have

$$(A^{\frac{1}{2}}BA^{\frac{1}{2}})^t \le A \le I,$$

which simplifies to $A^{\frac{1}{2}}BA^{\frac{1}{2}} \leq I$, that is, $B \leq A^{-1}$.

Theorem 3.4. Let $A, B \in \mathbb{P}$. Then for $t \in \mathbb{R}$,

$$F_t(A,B) \le I \iff (A^{\frac{1}{2}}BA^{\frac{1}{2}})^t \le A^{2t-1}.$$

Proof. Note that $F_t(A, B) \leq I$ if and only if $A^{2(1-t)} \leq (A^{-1}\sharp_t B)^{-1}$, which is equivalent to $A^{2(t-1)} \geq A^{-1}\sharp_t B$. By taking congruence transformation via $A^{\frac{1}{2}}$, we obtain $A^{2t-1} \geq (A^{\frac{1}{2}}BA^{\frac{1}{2}})^t$.

Corollary 3.5. Let $A, B \in \mathbb{P}$ such that $A \leq \alpha I$ and $B \leq \beta I$ for some $\alpha, \beta > 0$. Then $F_t(A, B) \leq \alpha^{1-t} \beta^t I$ for $t \in [0, 1]$.

Proof. We first assume that $A, B \leq I$. Then we have $A^{\frac{1}{2}}BA^{\frac{1}{2}} \leq A$ since $B \leq I$, and

$$(A^{\frac{1}{2}}BA^{\frac{1}{2}})^t \le A^t \le A^{2t-1}$$

The first inequality follows from the Löwner-Heinz inequality, and the last inequality follows from $A \leq I$ and $t \in [0, 1]$. Thus, $F_t(A, B) \leq I$ by Theorem 3.4.

For general $A \leq \alpha I$ and $B \leq \beta I$ for some $\alpha, \beta > 0$, let $A_1 := \alpha^{-1}A \leq I$ and $B_1 := \beta^{-1}B \leq I$, so $F_t(A_1, B_1) \leq I$. Using the joint homogeneity of F_t , we have

$$F_t(A_1, B_1) = \frac{1}{\alpha^{1-t}\beta^t} F_t(A, B) \le I.$$

It yields the desired inequality.

Remark 3.6. Since $A \leq ||A||I$ and $B \leq ||B||I$ for the operator norm $||\cdot||$, Corollary 3.5 yields $F_t(A, B) \leq ||A||^{1-t} ||B||^t I$ for any $t \in [0, 1]$, and hence,

$$||F_t(A, B)|| \le ||A||^{1-t} ||B||^t.$$

Switching to the finite-dimensional setting \mathbb{P}_m , we now provide an inequality relating the trace of the new weighted spectral geometric mean to the traces of the individual matrices.

Theorem 3.7. Let $A, B \in \mathbb{P}_m$. Then

$$\operatorname{tr} F_t(A, B) \le (\operatorname{tr} A)^{1-t} (\operatorname{tr} B)^t, \quad t \in [0, 1].$$

Proof. For any $A, B \in \mathbb{P}_m$ and $t \in [0, 1]$

$$\operatorname{tr} F_t(A,B) = \operatorname{tr} \left[A^{1-t} (A^{-1} \sharp_t B) A^{1-t} \right] = \operatorname{tr} \left[A^{\frac{1}{2}-t} (A^{\frac{1}{2}} B A^{\frac{1}{2}})^t A^{\frac{1}{2}-t} \right].$$
(3.2)

By the Araki-Lieb-Thirring inequality in [2],

$$\operatorname{tr} F_t(A, B) = \operatorname{tr} \left[A^{\frac{1}{2}-t} (A^{\frac{1}{2}} B A^{\frac{1}{2}})^t A^{\frac{1}{2}-t} \right]$$

$$\leq \operatorname{tr} \left[(A^{\frac{1}{2t}-1} \cdot A^{\frac{1}{2}} B A^{\frac{1}{2}} \cdot A^{\frac{1}{2t}-1})^t \right] = \operatorname{tr} \left[(A^{\frac{1-t}{2t}} B A^{\frac{1-t}{2t}})^t \right]$$

$$\leq (1-t) \operatorname{tr}(A) + t \operatorname{tr}(B).$$

The last inequality follows from [4, Theorem 10]. Replacing A, B by density matrices $\rho = \frac{A}{\text{tr}A}, \sigma = \frac{B}{\text{tr}B}$ in the above inequality, respectively, yields tr $F_t(\rho, \sigma) \leq 1$, and using the joint homogeneity of F_t in Proposition 2.1 (2) we obtain

$$\operatorname{tr} F_t(A, B) \le (\operatorname{tr} A)^{1-t} (\operatorname{tr} B)^t.$$

For $t \in [0, 1]$ and z > 0 the quantity

$$Q_{t,z}(A,B) = (A^{\frac{1-t}{2z}}B^{\frac{t}{z}}A^{\frac{1-t}{2z}})^z$$

is the matrix version of Rényi relative entropy. In particular, $Q_{t,t}(A, B)$ is known as the sandwiched Rényi relative entropy [19]. The quantity $Q_{t,z}$ can be defined for all $t \in \mathbb{R}$. The next theorem establishes a connection between the new weighted spectral geometric mean and Rényi relative entropy through log-majorization

Theorem 3.8. Let $A, B \in \mathbb{P}_m$. Then

$$F_t(A, B) \prec_{\log} Q_{t,z}(A, B), \quad 0 < z \le t \le 1.$$

Proof. Note that $Q_{t,z}(A, B)$ and $F_t(A, B)$ are homogeneous and are preserved by antisymmetric tensor power. Since

$$\det F_t(A,B) = (\det A)^{1-t} (\det B)^t = \det(A^{\frac{1-t}{2z}} B^{\frac{t}{z}} A^{\frac{1-t}{2z}})^z = \det Q_{t,z}(A,B),$$

it is sufficient to show that

$$Q_{t,z}(A,B) \le I \implies F_t(A,B) \le I.$$

Assume that $Q_{t,z}(A, B) \leq I$. Then $B^{\frac{t}{z}} \leq A^{\frac{t-1}{z}}$ because z > 0. Since $0 < \frac{z}{t} \leq 1$, $B \leq A^{1-\frac{1}{t}}$ by the Löwner-Heinz inequality. This yields $A^{1/2}BA^{1/2} \leq A^{\frac{2t-1}{t}}$, which implies $(A^{1/2}BA^{1/2})^t \leq A^{2t-1}$ by the Löwner-Heinz inequality. By Theorem 3.4 we obtain $F_t(A, B) \leq I$.

Corollary 3.9. Let $A, B \in \mathbb{P}_m$ and $\frac{1}{2} \leq t \leq 1$. Then

$$F_t(A, B) \prec_{\log} Q_{t,t}(A, B) \prec_{\log} A \natural_t B \prec_{w \log} A \diamond_t B.$$

Proof. The first log-majorization follows by Theorem 3.8 when z = t, and the last weak log-majorization follows by [12, Theorem 4.4]. From [7, Corollary 8],

$$Q_{t,z}(A,B) \prec_{\log} A \natural_t B$$

when $z \ge \max\{t, 1-t\}$ for $t \in [0, 1]$. Thus, the preceding log-majorization holds when $\frac{1}{2} \le t \le 1$ and $z \ge t$. The proof is complete by choosing z = t.

4. QUANTUM DIVERGENCE AND BARYCENTER

Bhatia, Gaubert and Jain [3] introduced a notion of quantum divergence on the Riemannian manifold \mathbb{P}_m . A map $\Phi : \mathbb{P}_m \times \mathbb{P}_m \to \mathbb{R}$ is called a *quantum divergence* if it satisfies the following properties:

- (I) $\Phi(A, B) \ge 0$, and equality holds if and only if A = B;
- (II) The derivative $D\Phi$ of Φ with respect to the second variable vanishes on the diagonal, that is,

$$D\Phi(A,B)|_{B=A} = 0;$$

(III) The second derivative is non-negative on the diagonal, that is,

$$D^2\Phi(A,B)|_{B=A}[Y,Y] \ge 0$$

for any Hermitian matrix Y.

Gan, Jeong and Kim showed in [13] that the map

$$\Psi(A,B) = \operatorname{tr}(A\nabla_t B - A\natural_t B)$$

is a quantum divergence for $t \in [0, \frac{1}{2}]$, and studied the barycenter minimizing the weighted sum of quantum divergences.

Theorem 4.1. The map $\Phi : \mathbb{P}_m \times \mathbb{P}_m \to \mathbb{R}$ defined by

$$\Phi(A,B) = \operatorname{tr}[A\nabla_t B - F_t(A,B)]$$
(4.3)

is a quantum divergence for $t \in [0, 1]$.

Proof. We verify the three conditions of quantum divergence.

From Theorem 3.7, we have

$$\operatorname{tr} F_t(A, B) \le (\operatorname{tr} A)^{1-t} (\operatorname{tr} B)^t \le (1-t) \operatorname{tr} A + t \operatorname{tr} B.$$

Therefore,

$$\Phi(A, B) = \operatorname{tr}[A\nabla_t B - F_t(A, B)] \ge 0$$

with equality if and only if A = B.

To prove positive definiteness of Φ , consider:

(a) Case 1: $t \in [0, \frac{1}{2}]$. If $\Phi(A, B) = 0$, then $(\operatorname{tr} A)^{1-t}(\operatorname{tr} B)^t = (1-t)\operatorname{tr} A + t \operatorname{tr} B$, and $\operatorname{tr} F_t(\rho, \sigma) = 1$ for positive definite density matrices $\rho = \frac{A}{\operatorname{tr} A}$ and $\sigma = \frac{B}{\operatorname{tr} B}$. By the definition of (metric) geometric mean on (3.2), we have

$$\operatorname{tr} F_t(\rho, \sigma) = \operatorname{tr} \left[\rho^{\frac{1-2t}{2}} (\rho^{\frac{1}{2}} \sigma \rho^{\frac{1}{2}})^t \rho^{\frac{1-2t}{2}} \right] = \operatorname{tr} \left[\rho^{1-2t} \left((\rho^{\frac{1}{2}} \sigma \rho^{\frac{1}{2}})^{\frac{1}{2}} \right)^{2t} \right]$$
$$\leq (1-2t) \operatorname{tr} \rho + 2t \operatorname{tr} (\rho^{\frac{1}{2}} \sigma \rho^{\frac{1}{2}})^{\frac{1}{2}}$$
$$\leq 1.$$

The first inequality follows from the fact that

$$\operatorname{tr}(A\sharp_t B) \le \operatorname{tr}(A^{1-t}B^t) \le (1-t)\operatorname{tr} A + t\operatorname{tr} B, \quad t \in [0,1],$$

and the second inequality follows from the property of quantum fidelity. Since tr $F_t(\rho, \sigma) = 1$, we have

$$\operatorname{tr}(\rho^{\frac{1}{2}}\sigma\rho^{\frac{1}{2}})^{\frac{1}{2}} = 1.$$

So tr A = tr B, and $\rho = \sigma$. Thus, A = B.

(b) Case 2: $t \in [\frac{1}{2}, 1]$. Note that

$$F_t(A,B) \prec_{\log} Q_{t,t}(A,B) \prec_{\log} A \natural_t B \prec_{w \log} A \diamond_t B \leq A \nabla_t B,$$

where the last inequality follows by [5, Theorem 6]. So we have

$$\operatorname{tr} F_t(A,B) \le \operatorname{tr} Q_{t,t}(A,B) \le \operatorname{tr}(A\natural_t B) \le \operatorname{tr}(A\diamond_t B) \le \operatorname{tr}(A\nabla_t B).$$

Thus, $\Phi(A, B) = 0$ implies $\operatorname{tr}(A\nabla_t B) = \operatorname{tr}(A \natural_t B) = \operatorname{tr}(A \diamond_t B)$. Since

$$\operatorname{tr}\left[A\nabla_t B - A \diamond_t B\right] = t(1-t)\operatorname{tr}\left[(A+B) - 2(A^{1/2}BA^{1/2})^{1/2}\right] = t(1-t)d_W^2(A,B),$$

where d_W denotes the Bures-Wasserstein distance, we obtain A = B.

(ii) One can see from (3.2) that

$$D\Phi(A,B)(B) = \operatorname{tr}(tI - A^{1/2-t}D(A^{1/2}BA^{1/2})^t(B)A^{1/2-t}).$$

Using the following integral representation:

$$A^{t} = \frac{\sin t\pi}{\pi} \int_{0}^{\infty} (\lambda A^{-1} + I)^{-1} \lambda^{t-1} d\lambda, \quad t \in (0, 1),$$

we have

$$D(A^{1/2}BA^{1/2})^{t}(B) = \frac{\sin t\pi}{\pi} \int_{0}^{\infty} (\lambda A^{-1/2}B^{-1}A^{-1/2} + I)^{-1}A^{-1/2}B^{-2}A^{-1/2}(\lambda A^{-1/2}B^{-1}A^{-1/2} + I)^{-1}\lambda^{t}d\lambda.$$

Thus, we obtain

$$D\Phi(A,B)(B)|_{B=A} = \operatorname{tr}\left(tI - A^{1/2-t}\frac{\sin t\pi}{\pi} \int_0^\infty (\lambda A^{-2} + I)^{-1} A^{-3} (\lambda A^{-2} + I)^{-1} \lambda^t d\lambda A^{1/2-t}\right)$$
$$= \operatorname{tr}\left(tI - A^{1-t}\frac{\sin t\pi}{\pi} \int_0^\infty (\lambda I + A^2)^{-2} \lambda^t d\lambda A^{1-t}\right)$$
$$= \operatorname{tr}\left(tI - tA^{2(1-t)+2t-2}\right) = 0,$$

where the last equality follows from the fact that for $t \in (0, 1)$

$$tA^{t-1} = D(A^t) = \frac{\sin t\pi}{\pi} \int_0^\infty (\lambda I + A)^{-2} \lambda^t d\lambda.$$

(iii) Let $Y \in \mathbb{H}_m$. Set $X := \lambda A^{-1/2} B^{-1} A^{-1/2} + I$. From the above observation, one can see that

$$D\Phi(A,B)(B)[Y] = \operatorname{tr} \left(tY - A^{1/2-t}D(A^{1/2}BA^{1/2})^t(B)[Y]A^{1/2-t}\right)$$

= $\operatorname{tr} \left(tY - A^{1/2-t}\frac{\sin t\pi}{\pi}\int_0^\infty X^{-1}A^{-1/2}B^{-1}YB^{-1}A^{-1/2}X^{-1}\lambda^t d\lambda A^{1/2-t}\right).$

Then, we see that

$$\begin{split} D^2 \Phi(A,B)(B,B)[Y,Y] \\ &= -\operatorname{tr} \left(A^{1/2-t} \frac{\sin t\pi}{\pi} \int_0^\infty Z A^{-1/2} B^{-1} Y B^{-1} A^{-1/2} X^{-1} \lambda^t d\lambda A^{1/2-t} \right) \\ &+ \operatorname{tr} \left(A^{1/2-t} \frac{\sin t\pi}{\pi} \int_0^\infty X^{-1} A^{-1/2} B^{-1} Y B^{-1} Y B^{-1} A^{-1/2} X^{-1} \lambda^t d\lambda A^{1/2-t} \right) \\ &+ \operatorname{tr} \left(A^{1/2-t} \frac{\sin t\pi}{\pi} \int_0^\infty X^{-1} A^{-1/2} B^{-1} Y B^{-1} Y B^{-1} A^{-1/2} X^{-1} \lambda^t d\lambda A^{1/2-t} \right) \\ &- \operatorname{tr} \left(A^{1/2-t} \frac{\sin t\pi}{\pi} \int_0^\infty X^{-1} A^{-1/2} B^{-1} Y B^{-1} A^{-1/2} Z \lambda^t d\lambda A^{1/2-t} \right), \end{split}$$

where $Z := D(X^{-1})(B)[Y]$. We denote the four terms in the above summation as I, II, III and IV, respectively. If B = A, then using the fact that

$$D(X^{-1})(B)[Y]|_{B=A} = \lambda(\lambda A^{-2} + I)^{-1} A^{-3/2} Y A^{-3/2} (\lambda A^{-2} + I)^{-1}$$

we obtain

$$I = IV$$

= $-\operatorname{tr}\left(A^{1-t}\frac{\sin t\pi}{\pi}\int_{0}^{\infty}(\lambda I + A^{2})^{-1}YA^{-1/2}(\lambda I + A^{2})^{-1}A^{-1/2}Y(\lambda I + A^{2})^{-1}\lambda^{t+1}d\lambda A^{1-t}\right)$
II = III
= $\operatorname{tr}\left(A^{1-t}\frac{\sin t\pi}{\pi}\int_{0}^{\infty}(\lambda I + A^{2})^{-1}YA^{-1}Y(\lambda I + A^{2})^{-1}\lambda^{t}d\lambda A^{1-t}\right).$

10

Thus, we have

$$\begin{split} D^2 \Phi(A,B)(B,B)[Y,Y]|_{B=A} \\ &= 2 \operatorname{tr} \left(\frac{\sin t\pi}{\pi} \int_0^\infty A^{1-t} (\lambda I + A^2)^{-1} Y A^{-1} Y (\lambda I + A^2)^{-1} A^{1-t} \lambda^t d\lambda \right) \\ &- 2 \operatorname{tr} \left(\frac{\sin t\pi}{\pi} \int_0^\infty A^{1-t} (\lambda I + A^2)^{-1} Y A^{-1/2} (\lambda I + A^2)^{-1} A^{-1/2} Y (\lambda I + A^2)^{-1} A^{1-t} \lambda^{t+1} d\lambda \right) \\ &= 2 \operatorname{tr} \left(\frac{\sin t\pi}{\pi} \int_0^\infty A^{1-t} (\lambda I + A^2)^{-1} Y H Y (\lambda I + A^2)^{-1} A^{1-t} \lambda^{t+1} d\lambda \right), \end{split}$$

where $H := A^{-1} - \lambda A^{-1/2} (\lambda I + A^2)^{-1} A^{-1/2}$. Since $I + \lambda^{-1} A^2 > I$ for any $\lambda > 0$, $I - (I + \lambda^{-1} A^2)^{-1} > 0$. Applying the congruence transformation via $A^{-1/2}$ on both sides yields that

$$A^{-1} - \lambda A^{-1/2} (\lambda I + A^2)^{-1} A^{-1/2} = A^{-1/2} \left\{ I - \left(I + \frac{1}{\lambda} A^2 \right)^{-1} \right\} A^{-1/2} > 0.$$

Hence, $D^2 \Phi(A, B)(B, B)[Y, Y]|_{B=A} \ge 0$ for any $Y \in \mathbb{H}_m$.

Proposition 4.2. The quantum divergence Φ given by (4.3) is invariant under unitary congruence transformation and tensor product with a density matrix.

Proof. By Proposition 2.1 (3),

$$\Phi(UAU^*, UBU^*) = \operatorname{tr}[U(A\nabla_t B - F_t(A, B))U^*] = \operatorname{tr}[A\nabla_t B - F_t(A, B)] = \Phi(A, B)$$

for any unitary matrix U. Since

$$F_t(A \otimes C, B \otimes D) = F_t(A, B) \otimes F_t(C, D)$$

for any $A, B, C, D \in \mathbb{P}_m$, the bi-linearity of the tensor product ensures

$$\Phi(A \otimes \rho, B \otimes \rho) = \operatorname{tr}[(A \nabla_t B - F_t(A, B)) \otimes \rho] = \operatorname{tr}[A \nabla_t B - F_t(A, B)] \operatorname{tr} \rho = \Phi(A, B)$$

for any density matrix ρ .

Theorem 4.3. For a given $A \in \mathbb{P}_m$ and $t \in (0,1)$, the map $f : \mathbb{P}_m \to \mathbb{R}$ defined by $f(X) = \operatorname{tr} F_t(A, X)$ is strictly concave.

Proof. From (3.2), $f(X) = \operatorname{tr} F_t(A, X) = \operatorname{tr} [A^{1-t}(A^{-1}\sharp_t X)A^{1-t}]$ for $X \in \mathbb{P}_m$ and $t \in (0, 1)$. Then for any $X, Y \in \mathbb{P}_m$ and $\lambda \in [0, 1]$,

$$f((1-\lambda)X + \lambda Y) = \operatorname{tr}[A^{1-t}(A^{-1}\sharp_t((1-\lambda)X + \lambda Y))A^{1-t}]$$

$$\geq \operatorname{tr}[A^{1-t}((1-\lambda)A^{-1}\sharp_tX + \lambda A^{-1}\sharp_tY)A^{1-t}]$$

$$= (1-\lambda)\operatorname{tr}[A^{1-t}(A^{-1}\sharp_tX)A^{1-t}] + \lambda\operatorname{tr}[A^{1-t}(A^{-1}\sharp_tY)A^{1-t}]$$

$$= (1-\lambda)f(X) + \lambda f(Y).$$

The inequality follows from the joint concavity of (metric) geometric mean:

$$A\sharp_t((1-\lambda)X + \lambda Y)) \ge (1-\lambda)A\sharp_t X + \lambda A\sharp_t Y.$$

The equality of the preceding argument holds if and only if X = Y. Thus, the map f is strictly concave.

Remark 4.4. The data processing inequality is an information-theoretic principle stating that the information content of a signal cannot increase under a local physical operation. For a quantum divergence Φ , this means that for any completely positive trace-preserving map Ψ and for $A, B \in \mathbb{P}_m$, the inequality

$$\Phi(\Psi(A), \Psi(B)) \le \Phi(A, B)$$

must hold.

According to [20, Theorem 5.16], if a map Φ is jointly convex and invariant under unitary congruence transformation and tensor product with density matrices, then it satisfies the data processing inequality. From Theorem 4.3, the quantum divergence Φ is convex with respect to the second variable, but not necessarily with respect to the first variable. Despite this, Φ satisfies the data processing inequality when t = 1/2.

Theorem 4.3 ensures that the quantum divergence Φ given by (4.3) is strictly convex with respect to the second variable. Consequently, the minimization problem for a positive probability vector $\omega = (w_1, \ldots, w_n)$ and $A_1, \ldots, A_n \in \mathbb{P}_m$

$$\underset{X \in \mathbb{P}_m}{\operatorname{arg\,min}} \sum_{j=1}^n w_j \Phi(A_j, X)$$

has a unique solution. We call it the barycenter of A_1, \ldots, A_n for the divergence Φ , denoted by $\mathfrak{B}_t(\omega; A_1, \ldots, A_n)$.

Theorem 4.5. The barycenter $\mathfrak{B}_t(\omega; A_1, \ldots, A_n)$ of $A_1, \ldots, A_n \in \mathbb{P}_m$ and $t \in (0, 1)$ for the quantum divergence Φ is the unique positive definite solution of the equation

$$tI = \sum_{j=1}^{n} w_j \frac{\sin t\pi}{\pi} \int_0^\infty A_j^{-t} (\lambda A_j^{-1} + X)^{-2} A_j^{-t} \lambda^t d\lambda, \qquad (4.4)$$

where $\omega = (w_1, \ldots, w_n)$ is a positive probability vector.

Proof. From the definition of the barycenter, $\mathfrak{B}_t(\omega; A_1, \ldots, A_n)$ minimizes the objective function

$$F(X) = \sum_{j=1}^{n} w_j \Phi(A_j, X),$$

where $\Phi(A_j, X) = \operatorname{tr}[A_j \nabla_t X - F_t(A_j, X)]$. From the proof of (ii) in Theorem 4.1, the gradient of F(X) with respect to X is given by

$$\nabla F(X) = \sum_{j=1}^{n} w_j \left[tI - A_j^{\frac{1}{2}-t} \frac{\sin t\pi}{\pi} \int_0^\infty (\lambda I + A_j^{\frac{1}{2}} X A_j^{\frac{1}{2}})^{-1} A_j (\lambda I + A_j^{\frac{1}{2}} X A_j^{\frac{1}{2}})^{-1} \lambda^t d\lambda A_j^{\frac{1}{2}-t} \right].$$

Since the objective function F(X) is strictly convex by Theorem 4.3, the barycenter $\mathfrak{B}_t(\omega; A_1, \ldots, A_n)$ coincides with the unique positive definition solution X of the equation obtained by setting $\nabla F(X) = 0$. Thus,

$$tI = \sum_{j=1}^{n} w_j A_j^{\frac{1}{2}-t} \frac{\sin t\pi}{\pi} \int_0^\infty (\lambda I + A_j^{\frac{1}{2}} X A_j^{\frac{1}{2}})^{-1} A_j (\lambda I + A_j^{\frac{1}{2}} X A_j^{\frac{1}{2}})^{-1} \lambda^t d\lambda A_j^{\frac{1}{2}-t}$$
$$= \sum_{j=1}^{n} w_j A_j^{-t} \frac{\sin t\pi}{\pi} \int_0^\infty (\lambda A_j^{-1} + X)^{-2} \lambda^t d\lambda A_j^{-t},$$

since $A_j^{\frac{1}{2}} (\lambda I + A_j^{\frac{1}{2}} X A_j^{\frac{1}{2}})^{-1} A_j^{\frac{1}{2}} = (\lambda A_j^{-1} + X)^{-1}$ for each *i*.

The gradient of F(X) with respect to X is given by:

$$\nabla F(X) = \sum_{j=1}^{n} w_j \left[tI - \sin(t\pi) \int_0^\infty A_j^{-t} (\lambda A_j^{-1} + X)^{-2} \lambda^t \, d\lambda \right].$$

Setting $\nabla F(X) = 0$ yields the necessary condition for the minimizer $B_t(\omega; A_1, \ldots, A_n)$, which satisfies:

$$tI = \sum_{j=1}^{n} w_j \sin(t\pi) \int_0^\infty A_j^{-t} (\lambda A_j^{-1} + X)^{-2} \lambda^t \, d\lambda.$$

Since F(X) is strictly convex (Theorem 4.3), this equation has a unique positive definite solution X, which corresponds to the barycenter $B_t(\omega; A_1, \ldots, A_n)$.

Theorem 4.6. The barycenter $\mathfrak{B}_t(\omega; A_1, \ldots, A_n)$ of $\mathbb{A} = (A_1, \ldots, A_n) \in \mathbb{P}_m^n$ and satisfies the following:

(1) if A_1, \ldots, A_n commute then

$$\mathfrak{B}_t(\omega;\mathbb{A}) = \left(\sum_{j=1}^n w_j A_j^{1-t}\right)^{\frac{1}{1-t}}, \quad t \in (0,1),$$

where $\omega = (w_1, \ldots, w_n)$ is a positive probability vector.

(2) The barycenter is invariant under permutation of the input matrices, that is,

$$\mathfrak{B}_t(\omega_{\sigma};\mathbb{A}_{\sigma})=\mathfrak{B}_t(\omega;\mathbb{A}),$$

for any permutation σ .

(3) The barycenter is equivariant under unitary congruence transformation, that is,

$$\mathfrak{B}_t(\omega; U\mathbb{A}U^*) = U\mathfrak{B}_t(\omega; \mathbb{A})U^*,$$

for any unitary matrix U.

Proof. For (1) it is enough to show that $X = \left(\sum_{j=1}^{n} w_j A_j^{1-t}\right)^{\frac{1}{1-t}}$ is a unique solution of (4.4) when A_1, \ldots, A_n . Assuming that X and A_j 's commute, the equation (4.4) reduces to

$$tI = \sum_{j=1}^{n} w_j A_j^{1-t} \frac{\sin t\pi}{\pi} \int_0^\infty (\lambda I + A_j X)^{-2} \lambda^t d\lambda A_j^{1-t}$$
$$= \sum_{j=1}^{n} w_j A_j^{1-t} t (A_j X)^{t-1} A_j^{1-t}$$
$$= t X^{t-1} \sum_{j=1}^{n} w_j A_j^{1-t}.$$

Solving for X, we get $X = \left(\sum_{j=1}^{n} w_j A_j^{1-t}\right)^{\frac{1}{1-t}}$.

Since the weights $\omega = (w_1, \ldots, w_n)$ and matrices A_1, \ldots, A_n can be permuted without affecting the sum in the barycenter equation, it follows that $\mathfrak{B}_t(\omega_{\sigma}; \mathbb{A}_{\sigma}) = \mathfrak{B}_t(\omega; \mathbb{A})$ for any permutation σ .

From Proposition 4.2, F_t is invariant under unitary congruence transformations. Hence we have (3), that is, $\mathfrak{B}_t(\omega; U\mathbb{A}U^*) = U\mathfrak{B}_t(\omega; \mathbb{A})U^*$.

Acknowledgement

There are no conflicts of interest associated with this work. The research of Sejong Kim was conducted during the research year of Chungbuk National University in 2024 at the University of Nevada, Reno. His work is also supported by the National Research Foundation of Korea funded by the Ministry of Science and ICT under grant no. NRF-2022R1A2C4001306.

References

- T. Ando and F. Hiai, Log majorization and complementary Golden-Thompson type inequalities, Linear Algebra Appl., 197/198 (1994), 113–131.
- [2] H. Araki, On an inequality of Lieb and Thirring, Lett. Math. Phys., 19 (1990), 167–170.
- [3] R. Bhatia, S. Gaubert and T. Jain, *Matrix versions of the Hellinger distance*, Lett. Math. Phys., 109 (2019), 1777–1804.
- [4] R. Bhatia, T. Jain and Y. Lim, Strong convexity of sandwiched entropies and related optimization problems, Rev. Math. Phys., 30 (2018), 1850014, 18 pp.
- R. Bhatia, T. Jain and Y. Lim, On the Bures-Wasserstein distance between positive definite matrices, Expo. Math. 37 (2019), 165–191.
- [6] T.-H. Dinh, T.-Y. Tam and T.-D. Vuong, A new weighted spectral geometric mean and properties, Electron. J. Linear Algebra, 40 (2024), 333–342.
- [7] R. Dumitru and J. A. Franco, Log-majorizations related to the spectral geometric and Rényi means, Acta Sci. Math., (2024), DOI: 10.1007/s44146-024-00128-8.
- [8] R. Dumitru and J. A. Franco, Near order and metric-like functions on the cone of positive definite matrices, Positivity, 28 (2024), Paper No. 2, 15 pp.
- [9] F. Hansen and G. K. Pedersen, Jensen's inequality for operators and Löwner's theorem, Math. Ann., 258 (1981/1982), 229–241.
- [10] J. Hwang and S. Kim, Two-variable Wasserstein means of positive definite operators, Mediterr. J. Math., 19 (2022), Paper No. 110, 16pp.
- [11] L. Gan and S. Kim, Revisit on spectral geometric mean, Linear Multilinear Algebra, 72 (2024), 944–955.

- [12] L. Gan and S. Kim, Weak log-majorization between the geometric and Wasserstein means, J. Math. Anal. Appl., 530 (2024), Paper No. 127711, 14 pp.
- [13] L. Gan, M. Jeong and S. Kim, Quantum divergence and barycenter associated with the spectral geometric mean, Linear Algebra Appl., to appear. DOI: 10.1016/j.laa.2024.05.004.
- [14] L. Gan and T.-Y. Tam, Inequalities and limits of weighted spectral geometric mean, Linear Multilinear Algebra, 72 (2024), 261–282.
- [15] S. Kim, Operator inequalities and gyrolines of the weighted geometric means, Math. Inequal. Appl., 24 (2021), 491–514.
- [16] F. Kubo and T. Ando, Means of positive linear operators, Math. Ann., 246 (1980), 205–224.
- [17] J. Lawson and Y. Lim, Karcher means and Karcher equations of positive definite operators, Trans. Am. Math. Soc. Ser. B, 1 (2014), 1–22.
- [18] H. Lee and Y. Lim, Metric and spectral geometric means on symmetric cones, Kyungpook Math. J., 47 (2007), 133–150.
- [19] M. Wilde, A. Winter and D. Yang, Strong converse for the classical capacity of entanglement-breaking and Hadamard channels via a sandwiched Renyi relative entropy, Comm. Math. Phys., 331 (2014), 593–622.
- [20] M. M. Wolf, Quantum channels & operations: guide tour, Lecture notes available at http://wwwm5.ma.tum.de/foswiki/pub/M5/A;;gemeines/MichaelWolf/QChannelLecture.pdf, July 2012
- [21] T. Yamazaki, The Riemannian mean and matrix inequalities related to the Ando-Hiai inequality and chaotic order, Oper. Matrices, 6 (2012), 577–588.

MIRAN JEONG, DEPARTMENT OF MATHEMATICS, CHUNGBUK NATIONAL UNIVERSITY, CHEONGJU 28644, KOREA

Email address: jmr4006@chungbuk.ac.kr

SEJONG KIM, DEPARTMENT OF MATHEMATICS, CHUNGBUK NATIONAL UNIVERSITY, CHEONGJU 28644, KOREA

Email address: skim@chungbuk.ac.kr

TIN-YAU TAM, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEVADA AT RENO, NV 89557, USA

Email address: ttam@unr.edu