

Liminal $\mathrm{SL}_2\mathbb{Z}_p$ -representations and odd-th cyclic covers of genus one two-bridge knots

Honami Sakamoto, Ryoto Tange, and Jun Ueki

ABSTRACT

Let p be a prime number and K a genus one two-bridge knot. We prove that if p divides the size of the 1st homology group of some odd-th cyclic branched cover of the knot K , then its group $\pi_1(S^3 - K)$ admits a liminal $\mathrm{SL}_2\mathbb{Z}_p$ -character. In addition, we discuss the existence of liminal $\mathrm{SL}_2\mathbb{Z}_p$ -representations. In the course of the argument, we also point out a constraint for prime numbers dividing certain Lucas-type sequences by using the Legendre symbols.

Contents

1	Introduction	1
2	Genus one two-bridge knots	2
3	Liminal $\mathrm{SL}_2\mathbb{Z}_p$-characters	3
4	Liminal representations	5
5	Lucas and Fibonacci type sequences	6
6	Cyclic covers	7
	References	8

1. Introduction

Let p be a prime number and \mathbb{Z}_p denote the ring of p -adic integers. In the context of Hida–Mazur theory and its analogue in low dimensional topology [MTTU17, KMTT18, TTU22, TU24, BTTU23], there are special interests in irreducible $\mathrm{SL}_2\mathbb{Z}_p$ -representations whose residual representations are reducible. In this paper, following Mazur [Maz11, Section 19], we aim to “go the other way”.

Let π be a group. A function $\chi : \pi \rightarrow \mathbb{Z}_p$ is called *an $\mathrm{SL}_2\mathbb{Z}_p$ -character* if there exists an SL_2 -representation ρ over an extension of \mathbb{Z}_p such that $\chi = \mathrm{tr} \rho$ holds. An $\mathrm{SL}_2\mathbb{Z}_p$ -representation (resp. character) is said to be *liminal* if it is absolutely reducible and its every open neighborhood contains an absolutely irreducible $\mathrm{SL}_2\mathbb{Z}_p$ -representation (resp. character).

We will prove the following.

THEOREM 1.1. *Let K be a genus one two-bridge knot in S^3 . If a prime number p divides the size of the 1st homology group of some odd-th cyclic branched cover of K , then its group $\pi_1(S^3 - K)$ admits a liminal $\mathrm{SL}_2\mathbb{Z}_p$ -character.*

To prove Theorem 1.1, we utilize the character varieties, the cyclic resultants of the Alexander polynomials, and the Legendre symbols, and establish a property of certain Fibonacci/Lucas-type

sequences as well.

The argument will be given as follows. In Section 2, we recollect properties of genus one two-bridge knots, namely, double twist knots of type $J(2k, 2l)$ with $(0, 0) \neq (k, l) \in \mathbb{Z}^2$ and calculate the intersection of the varieties of irreducible and reducible $\mathrm{SL}_2\mathbb{C}$ -characters (Proposition 2.1). In Section 3, by using Hensel's lemma several times, we prove that $J(2k, 2l)$ admits a liminal $\mathrm{SL}_2\mathbb{Z}_p$ -character iff (i) $p = 2$ and $r \equiv 1 \pmod{8}$ or (ii) $p \neq 2$ and the Legendre symbol satisfies $(\frac{r}{p}) = 1$, where r denotes the square-free part of $4k^2l^2 - kl$ (Theorem 3.1), and calculate several examples. In Section 4, we further discuss the existence of liminal $\mathrm{SL}_2\mathbb{Z}_p$ -representations (Proposition 4.2).

In Section 5, for a given $m \in \mathbb{Z}$, we write $t^2 - t + m = (t - a)(t - b)$ and define the Lucas-type sequence $(L_n)_n \in \mathbb{Z}^{\mathbb{N}}$ by $L_n = a^n + b^n$. We prove that $p \mid L_{2n+1}$ for some $n \in \mathbb{Z}_{\geq 0}$ implies $(\frac{4m^2 - m}{p}) = 1$ (Proposition 5.1) and attach examples. Note that the classical one with $m = -1$ is the case of the figure-eight knot $J(2, -2) = 4_1$.


In Section 6, we recall some properties of the cyclic covers $M_n \rightarrow S^3$ of a knot K , prove for a knot K with its Alexander polynomial $\Delta_K(t) = mt^2 - (2m - 1)t + m$ that $\#H_1(M_{2n+1}) = L_{2n+1}^2$ holds, yielding that $p \mid \#H_1(M_{2n+1})$ implies $(\frac{4m^2 - m}{p}) = 1$ (Theorem 6.1), and deduce Theorem 1.1.

It has been classically known by Burde and de Rham [Bur67, dR67] that such deformative characters correspond to the zeros of the Alexander polynomials (see also [HPSP01]), so our result has deep context in its background. We describe further queries and perspectives in Remark 6.4. We also remark that a p -adic analogue of Burde–de Rham theorem and its application to number theoretical settings are due to the second author and others [MTT24].

Acknowledgments. We are grateful to Léo Benard, Tomoki Mihara, Shin-ichiro Seki, Motoo Tange, Anh T. Tran, Yuji Terashima, Yoshikazu Yamaguchi, and Hyuga Yoshizaki for their useful comments. The third author has been partially supported by JSPS KAKENHI Grant Number JP23K12969.

2. Genus one two-bridge knots

It is known that every genus one two-bridge knot is realized as a *double twist knot* of type $J(2k, 2l)$ with $(0, 0) \neq (k, l) \in \mathbb{Z}^2$ defined by the following diagram. A basic reference is [Tra18].



Especially, $J(2, 2l)$ is known as a twist knot; $J(2, 0)$ is the unknot 0_1 , $J(2, 2)$ is the trefoil 3_1 , and $J(2, -2)$ is the figure-eight knot 4_1 .

The group of such a knot admits a presentation

$$\pi := \pi_1(S^3 - J(2k, 2l)) = \langle a, b \mid w^l a = b w^l \rangle$$

with a, b being meridians and $w = (ba^{-1})^k (b^{-1}a)^k$.

A Seifert matrix is given by $V = \begin{pmatrix} k & 1 \\ 0 & l \end{pmatrix}$, so its Alexander polynomial becomes

$$\Delta_{J(2k,2l)}(t) = \det(tV - V^\perp) = klt^2 + (1 - 2kl)t + kl.$$

For each $g \in \pi$, let $\mathrm{tr} g$ denote the map $\mathrm{Hom}(\pi, \mathrm{SL}_2\mathbb{C}) \rightarrow \mathbb{C}; \rho \mapsto \mathrm{tr} \rho(g)$. Then the conjugacy classes of $\mathrm{SL}_2\mathbb{C}$ -representations are parametrized by

$$x := \mathrm{tr} a, \quad y := \mathrm{tr} ab^{-1}.$$

(This y is different from that in [TU24] and has a strong advantage in our calculation. The old y coincides with $\mathrm{tr} ab = \mathrm{tr} a \mathrm{tr} b - \mathrm{tr} ab^{-1} = x^2 - y$.) We further put $z := \mathrm{tr} w$.

Each conjugacy class of non-abelian $\mathrm{SL}_2\mathbb{C}$ -representation is presented by *Riley's universal representation* $\rho^R : \pi \rightarrow \mathrm{SL}_2\mathbb{C}$ defined by

$$\rho^R(a) = \begin{pmatrix} s & 1 \\ 0 & s^{-1} \end{pmatrix}, \quad \rho^R(b) = \begin{pmatrix} s & 0 \\ 2 - y & s^{-1} \end{pmatrix},$$

where $x = s + s^{-1}$ and x, y satisfy Riley's polynomial equation $f_{k,l}(x, y) = \Phi_{k,l}(x, y - 2) = 0$.

There is a bijective correspondence between conjugacy classes of irreducible representations and points on $f_{k,l}(x, y) = 0$ with $y - 2 \neq 0$. Each point of $y - 2 = 0$ corresponds to a pair of the class of reducible representations and that of abelian representations.

PROPOSITION 2.1. *The intersection of $f_{k,l}(x, y) = 0$ and $y - 2 = 0$ is $(x, y) = (\pm\sqrt{4 - \frac{1}{kl}}, 2)$.*

For each $n \in \mathbb{Z}$, let $S_n(z) \in \mathbb{Z}[z]$ denote the n -th Chebyshev polynomial of the second type defined by $S_{n-1}(2 \cos \theta) = \frac{\sin n\theta}{\sin \theta}$ with $\theta \in \mathbb{R}$, $\sin \theta \neq 0$. This is equivalent to say that $S_{-1}(z) = 0$, $S_0(z) = 1$, and $S_{n+1}(z) - zS_n(z) + S_{n-1}(z) = 0$ for every $n \in \mathbb{Z}$. These polynomials satisfy $S_{-1-n}(z) = -S_{-1+n}(z)$ and $S_{n-1}(\pm 2) = (\pm 1)^{n-1}n$.

We have

$$z = \mathrm{tr} w = 2 + (y - 2)(-x^2 + y + 2)S_{m-1}^2(y).$$

By [Tra18, Subsection 2.2], the variety of irreducible characters is given by

$$f_{k,l}(x, y) = S_l(z) - (1 + (-x^2 + y + 2)S_{k-1}(y)(S_k(y) - S_{k-1}(y))S_{l-1}(z)).$$

Proof of Proposition 2.1. If $y - 2 = 0$, then we have $z = 2$, $0 = f_{k,l}(x, 2) = l - (1 + (-x^2 + 2 + 2)k((k + 1) - k))l = 1 - (4 - x^2)kl$, and hence $x = \pm\sqrt{4 - \frac{1}{kl}}$. \square

REMARK 2.2. In the old parameters, the intersection points become $(x, \eta) = (\pm\sqrt{4 - \frac{1}{kl}}, -\frac{1}{kl})$. Note that Proposition 2.1 persists over any algebraically closed field \mathbb{F} with $\mathrm{char} \mathbb{F} \nmid kl$.

3. Liminal $\mathrm{SL}_2\mathbb{Z}_p$ -characters

THEOREM 3.1. *Suppose $kl \neq 0$. Then the group of $J(2k, 2l)$ admits a liminal $\mathrm{SL}_2\mathbb{Z}_p$ -character iff*

- i) $p = 2$ and $4k^2l^2 - kl \equiv 1 \pmod{8}$ or
- ii) $p \neq 2$, $p \nmid kl$, and the Legendre symbol satisfies $\left(\frac{r}{p}\right) = 1$, where r denotes the square-free part of $4k^2l^2 - kl$.

Here, we define *the square-free part* of $0 \neq a \in \mathbb{Z}$ by a/b^2 , where b denotes the maximal integer with $b^2 \mid a$. (Note that some people use this term for other notions.) *The Legendre symbol* $(\frac{a}{p}) \in \{0, \pm 1\}$ of $a \in \mathbb{Z}$ over p is defined as follows: If $p \mid a$, we put $(\frac{a}{p}) = 0$. Suppose $p \nmid a$. If $a \equiv x^2 \pmod{p}$ for some $x \in \mathbb{Z}$, then we put $(\frac{a}{p}) = 1$. Otherwise, we put $(\frac{a}{p}) = -1$. The following lemmas are consequences of Hensel's lemma (cf. [Neu99, Chapter II (4.6)]).

LEMMA 3.2. *Let $0 \neq a \in \mathbb{Z}$ with $p^2 \nmid a$. Then, we have $\sqrt{a} \in \mathbb{Z}_p$ iff*

- i) $p = 2$ and $a \equiv 1 \pmod{8}$ or
- ii) $p \neq 2$ and $(\frac{a}{p}) = 1$.

Proof. (i) Suppose $p = 2$. If $\sqrt{a} \in \mathbb{Z}_2$, then $\sqrt{a} = \sum_i b_i 2^i$ with $b_i \in \{0, 1\}$, and $a \equiv (b_0 + 2b_1 + 4b_2)^2 \equiv b_0^2 + 4b_1(b_0 + b_1)$. Hence, by the assumption $f \nmid a$, we have $b_0 = 1$. In both cases $b_1 = 0, 1$, we have that $a \equiv 1 \pmod{8}$. Conversely, suppose that $a \equiv 1 \pmod{8}$ and write $a = 8b + 1$ with $b \in \mathbb{Z}$. If $X^2 - a = 0$ have a solution $X = \alpha$ in \mathbb{Z}_2 , then $\alpha = 2\beta + 1$ for some $\beta \in \mathbb{Z}_2$, and $\alpha^2 - a = (2\beta + 1)^2 - (8b + 1) = 4(\beta^2 + \beta - 2b)$. Since the polynomial $Y^2 + Y - 2b \pmod{2}$ has single roots in \mathbb{F}_2 , Hensel's lemma yields roots of $Y^2 + Y - 2b$ in \mathbb{Z}_2 , and hence those of $X^2 - a$ in \mathbb{Z}_2 .

ii) Suppose $p \neq 2$. If $p \mid a$, then $\sqrt{a} \notin \mathbb{Z}_p$. If $p \nmid a$, then $X^2 - a \pmod{p}$ has single roots in \mathbb{F}_p and they lift to a quadratic extension of \mathbb{Z}_p by Hensel's lemma. In addition, the following are equivalent; $(\frac{a}{p}) = 1$, the roots of $X^2 - a \pmod{p}$ belong to \mathbb{F}_p , the roots of $X^2 - a$ belong to \mathbb{Z}_p . \square

LEMMA 3.3. *There is a bijective correspondence between liminal $\mathrm{SL}_2\mathbb{Z}_p$ -characters of $\pi_1(S^3 - J(2k, 2l))$ and the intersection points of $f_{k,l}(x, y) = 0$ and $y - 2 = 0$ in \mathbb{Z}_p^2 .*

Proof. If ρ is a liminal representation with $\mathrm{Im} \mathrm{tr} \rho \subset \mathbb{Z}_p$, then $\mathrm{tr} \rho$ is on $y - 2 = 0$ in \mathbb{Z}_p^2 . In addition, for every $n \in \mathbb{Z}_{>0}$, there is an irreducible representation ρ' with $\rho \equiv \rho' \pmod{p^n}$, so $\mathrm{tr} \rho$ is on $f_{k,l}(x, y) \equiv 0 \pmod{p^n}$. Thus $\mathrm{tr} \rho$ is on $f_{k,l}(x, y) = 0$ in \mathbb{Z}_p^2 as well.

Let us prove the converse. Note that Riley's representation ρ^R in Section 2 may be seen as a representation over a quadratic extension of $\mathbb{Z}_p[x, y]$.

By Proposition 2.1, if $f_{k,l}(x, y) = 0$ and $y - 2 = 0$ has an intersection point in \mathbb{Z}_p^2 , then $f_{k,l}(x, 2) = (f_{k,l}(x, y) \pmod{(y - 2)})$ has a single root α in \mathbb{Z}_p , and Hensel's lemma yields the implicit function $x = x_f(y) \in \mathbb{Z}_p[[y - 2]]$ around $(x, y) = (\alpha, 2)$. By substituting $x = x_f(y)$ in Riley's representation, we obtain an irreducible representation ρ^R over a quadratic extension of $\mathbb{Z}_p[[y - 2]]$ with $\mathrm{Im} \mathrm{tr} \rho^R \subset \mathbb{Z}_p[[y - 2]]$.

Note that every element of $\mathbb{Z}_p[[y - 2]]$ converges in \mathbb{Z}_p at any $y \in \mathbb{Z}_p$ in the p -adic unit disc $|y - 2|_p < 1$. If we substitute $y = 2$, then this ρ^R yields a reducible representation ρ with $\mathrm{tr} \rho \subset \mathbb{Z}_p$ at $(\alpha, 2)$. If we instead substitute $y = \eta \in \mathbb{Z}_p$ with $0 \neq |2 - \eta|_p < 1$, namely, $2 \equiv \eta \pmod{p^n}$ for some $n \in \mathbb{Z}_{>0}$, then ρ^R yields an absolutely irreducible representation ρ_η with $\mathrm{tr} \rho_\eta \subset \mathbb{Z}_p$. Thus, $\mathrm{tr} \rho$ is a liminal $\mathrm{SL}_2\mathbb{Z}_p$ -character. \square

Proof of Theorem 3.1. Since $f_{k,l}(x, y) = 0$ and $y - 2 = 0$ have an intersection point in \mathbb{Z}_p^2 if and only if $p \nmid kl$ and $\sqrt{4k^2l^2 - kl} \in \mathbb{Z}_p$, Lemma 3.2 and Lemma 3.3 yield the assertion. \square

EXAMPLE 3.4. The condition for $\pi_1(S^3 - J(2, 2l))$ admitting a liminal $\mathrm{SL}_2\mathbb{Z}_p$ -representation becomes $(\frac{4l^2 - l}{p}) = 1$. By an elementary calculation, we obtain the following.

- (i) $J(2, 2) = 3_1$ (trefoil): $(\frac{3}{p}) = 1$, i.e., $p \equiv \pm 1 \pmod{12}$.

- (ii) $J(2, -2) = 4_1$ (figure-eight): $(\frac{5}{p}) = 1$, i.e., $p \equiv \pm 1 \pmod{5}$.
- (iii) $J(2, 4)$: $(\frac{14}{p}) = 1$, i.e., $p \equiv \pm 1, \pm 9, \pm 25, \pm 5, \pm 11, \pm 13 \pmod{56}$.
- (iv) $J(2, -4)$: $(\frac{18}{p}) = 1$, i.e., $p \equiv \pm 1 \pmod{8}$.
- (v) $J(2, 6)$: $p = 2$ or $(\frac{33}{p}) = 1$, i.e., $p \equiv \pm 1, \pm 2, \pm 4, \pm 8, \pm 16 \pmod{33}$.
- (vi) $J(2, -6)$: $(\frac{39}{p}) = 1$, i.e., $\pm p \equiv 1, 5, 7, 19, 23, 25, 26, 35, 41, 49, 61, 67 \pmod{156}$.

Proof. The condition for $p = 2$ is $4l^2 - l \equiv 1 \pmod{8}$, that is, $l \equiv 3 \pmod{8}$. Suppose $p \neq 2$.

(i) If $l = 1$, then $4l^2 - l = 3$, and the condition becomes $(\frac{3}{p}) = 1$. If $p \equiv 1 \pmod{4}$, then by the quadratic reciprocity law, we have $(\frac{p}{3}) = (\frac{3}{p})$. Since $x \equiv \pm 1$ implies $x^2 \equiv 1$, we have that $(\frac{3}{p}) = 1$ iff $p \equiv 1 \pmod{3}$. If instead $p \equiv -1 \pmod{4}$, then by $(\frac{p}{3}) = -(\frac{3}{p})$, we have that $(\frac{3}{p}) = 1$ iff $p \equiv -1 \pmod{3}$. Thus, the condition becomes $p \equiv \pm 1 \pmod{12}$.

(ii) If $l = -1$, then the condition becomes $(\frac{5}{p}) = 1$. By $5 \equiv 1 \pmod{4}$, we have $(\frac{5}{p}) = (\frac{p}{5})$. Since $x \equiv \pm 1, \pm 2$ implies $x^2 \equiv \pm 1 \pmod{5}$, the condition becomes $p \equiv \pm 1 \pmod{5}$.

(iii) If $l = 2$, then the condition becomes $(\frac{14}{p}) = (\frac{2}{p})(\frac{7}{p}) = 1$. The second supplement asserts that $(\frac{2}{p}) = 1$ iff $p \equiv \pm 1 \pmod{8}$. Note that $7 \equiv 3 \pmod{4}$ and that $x \equiv \pm 1, \pm 2, \pm 3$ implies $x^2 \equiv 1, 4, 2 \pmod{7}$. If $p \equiv 1 \pmod{4}$, then $(\frac{7}{p}) = (\frac{p}{7})$, so $(\frac{7}{p}) = 1$ iff $p \equiv 1, 2, 4 \pmod{7}$. If $p \equiv -1 \pmod{4}$, then $(\frac{7}{p}) = -(\frac{p}{7})$, so $(\frac{7}{p}) = 1$ iff $p \equiv -1, -2, -4 \pmod{7}$. Thus, the condition becomes “ $\pm p \equiv \pm 1 \pmod{8}$ and $\pm p \equiv 1, 2, 4 \pmod{7}$ ” or “ $\pm p \equiv 3 \pmod{8}$ and $\pm p \equiv 1, 2, 4 \pmod{7}$ ” (double sign in the same order), yielding the assertion.

(iv) If $l = -2$, then by $4l^2 - l = 18 = 2 \cdot 3^2$, the condition becomes $(\frac{2}{p}) = 1$. The second supplement asserts that $(\frac{2}{p}) = 1$ iff $p \equiv \pm 1 \pmod{8}$.

(v) If $l = 3$, then the condition becomes $1 = (\frac{33}{p}) = (\frac{3}{p})(\frac{11}{p}) = ((-1)^{\frac{p-1}{2}})^2 (\frac{p}{3})(\frac{p}{11}) = (\frac{p}{3})(\frac{p}{11})$. Since $x \equiv \pm 1$ implies $x^2 \equiv 1 \pmod{3}$, we have that $(\frac{p}{3}) = 1$ iff $p \equiv 1 \pmod{3}$. Since $x \equiv \pm 1, \pm 2, \pm 3, \pm 4, \pm 5$ implies $x^2 \equiv 1, 4, 9, 5, 3 \pmod{11}$, we have $(\frac{p}{11}) = 1$ iff $p \equiv 1, 4, 9, 5, 3 \pmod{11}$. Both values of $(\frac{p}{3})$ and $(\frac{p}{11})$ are 1 iff $p \equiv 1, 4, 31, 16, 25 \pmod{33}$. Both values are -1 iff $p \equiv 2 \pmod{3}$ and $p \equiv 2, 6, 7, 8, 10 \pmod{11}$, hence iff $p \equiv 2, 17, 29, 8, 32 \pmod{33}$. Thus, the condition becomes $p \equiv 1, 2, 4, 8, 16, 17, 25, 29, 31, 32 \pmod{33}$, hence the assertion.

(vi) If $l = -3$, then the condition becomes $(\frac{39}{p}) = (\frac{3}{p})(\frac{13}{p}) = 1$. As we saw in (i), we have $(\frac{3}{p}) = 1$ iff $p \equiv \pm 1 \pmod{12}$, and hence $(\frac{3}{p}) = -1$ iff $p \equiv \pm 5 \pmod{12}$. In addition, since $x \equiv \pm 1, \pm 2, \pm 3, \pm 4, \pm 5, \pm 6$ implies $x^2 \equiv 1, 4, 9, 3, 12, 10 \pmod{13}$, we have that $(\frac{13}{p}) = 1$ iff $p \equiv \pm 1, \pm 3, \pm 4, \pmod{13}$, and $(\frac{13}{p}) = -1$ iff $p \equiv \pm 2, \pm 5, \pm 6, \pmod{13}$. Thus, we have $(\frac{39}{p}) = 1$ iff $\pm p \equiv 1, 25, 133, 49, 121, 61, 41, 89, 5, 125, 149, 137 \pmod{156}$, and hence the assertion. \square

4. Liminal representations

Let ρ be a representation over a finite extension of \mathbb{Z}_p such that $\bar{\rho} := \rho \pmod{p}$ is absolutely irreducible and suppose that $\mathrm{Im} \mathrm{tr} \bar{\rho} \subset \mathbb{F}_p$. Since the Brauer group of a finite field is trivial, the Skolem–Noether theorem assures that such $\bar{\rho}$ is conjugate to some representation $\bar{\rho}'$ over \mathbb{F}_p (cf. [Mar16, Subsection 3.5]). In addition, Nyssen’s theorem [Nys96, Théorème 1] asserts that $\bar{\rho}'$ lifts to some ρ' over \mathbb{Z}_p such that $\mathrm{tr} \rho = \mathrm{tr} \rho'$, and such ρ' is unique up to conjugate.

One might wonder whether this argument extends to representations on the Zariski closure of the variety $X_{\mathrm{irr}}(S^3 - J(2k, 2l)) = \{f_{f,l}(x, y) = 0\} \setminus \{y - 2 = 0\}$ of irreducible characters. Nyssen’s proof uses the fact that being absolutely irreducible is equivalent to the surjectivity of

the corresponding algebra homomorphism, so the result does not necessarily extend to residually reducible representations. In addition, as pointed out in [KMTT18, Subsection 2.3], in the setting of knot group representations, the deformation problem is not unobstructed in the sense that the 2nd cohomology of the adjoint representation does not vanish. The following question seems subtle.

QUESTION 4.1. Given a liminal character over \mathbb{Z}_p , can we find a liminal $\mathrm{SL}_2\mathbb{Z}_p$ -representation?

We may say at least the following.

PROPOSITION 4.2. *Under the conditions of Theorem 3.1, if $p \neq 2$ and $(\frac{-kl}{p}) = 1$, then there exists a liminal $\mathrm{SL}_2\mathbb{Z}_p$ -representaion.*

Proof. Let ρ^R be the one in Lemma 3.3 and note that $s = \frac{x - \sqrt{x^2 - 4}}{2}$. By Proposition 2.1, we have $\sqrt{x^2 - 4}|_{y=2} = \sqrt{\frac{-1}{kl}}$. If $(\frac{-kl}{p}) = 1$, then $\sqrt{x^2 - 4} = \sqrt{\frac{-1}{kl}} \in \mathbb{Z}_p$, and $\rho^R \bmod y - 2$ becomes a liminal $\mathrm{SL}_2\mathbb{Z}_p$ -representation. \square

5. Lucas and Fibonacci type sequences

Let $m \in \mathbb{Z}$ and write $t^2 - t + m = (t - a)(t - b)$, so that we have $a + b = 1$ and $ab = m$. We define the Lucas-type sequence and the Fibonacci-type sequence by $L_n = a^n + b^n$ and $F_n = \frac{a^n - b^n}{a - b}$ respectively. Then, we have $L_0 = 2$, $L_1 = 1$, $L_2 = 1 - 2m$, $L_{n+2} = L_{n+1} - mL_n$, $F_0 = 0$, $F_1 = 1$, $F_2 = 1$, $F_{n+2} = F_{n+1} - mF_n$, and hence $L_n, F_n \in \mathbb{Z}$. We have

$$L_n^2 + (4m - 1)F_n^2 = 4m^n. \quad \cdots (\star)$$

Indeed, by $(a - b)^2 = (a + b)^2 - 4ab = 1 - 4m$, we have $(a^n + b^n)^2 + (4m - 1)(\frac{a^n - b^n}{a - b})^2 = (a^n + b^n)^2 - (a^n - b^n)^2 = 4a^n b^n = 4m^n$.

Now suppose that p divides L_{2n+1} for some $n \in \mathbb{Z}_{\geq 0}$. Then we have $(4m - 1)F_n^2 \equiv (2m^n)^2 m \pmod{p}$. This implies that $m(4m - 1)$ must be a square mod p . In addition, we have $p \nmid m(4m - 1)$. Indeed, if $p \mid m$, then the recurrence formula yields that $L_n \equiv 1 \pmod{p}$ for all $n \in \mathbb{Z}_{> 0}$, contradicting $p \mid L_{2n+1}$. If instead $p \mid 4m - 1$, then (\star) yields $p = 2$ or $p \mid m$, hence again contradiction. Thus, we have the following.

PROPOSITION 5.1. *If a prime number p divides L_{2n+1} for some $n \in \mathbb{Z}_{\geq 0}$, then the Legendre symbol satisfies $(\frac{4m^2 - m}{p}) = 1$.*

EXAMPLE 5.2. The following is comparable with Example 3.4.

(i) Let $m = 1$. Then the condition $(\frac{12}{p}) = 1$ becomes $p = 2$ or $p \equiv \pm 1 \pmod{12}$. By $L_{n+1} = L_n - L_{n-1}$, we see that

$$\frac{n \bmod 6}{L_n} \left\| \begin{array}{c|c|c|c|c|c} 1 & 2 & 3 & 4 & 5 & 6 \\ \hline 1 & -1 & -2 & -1 & 1 & 2 \end{array} \right.,$$

so only $p = 2$ actually appears.

(ii) Let $m = -1$. Then, L_n and F_n become the classical ones, and the condition $(\frac{5}{p}) = 1$ becomes $p = 2$ or $p \equiv \pm 1 \pmod{5}$. We actually have

$$\frac{n}{L_n} \left\| \begin{array}{c|c|c|c|c|c|c|c|c|c|c|c|c|c} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 & 14 \\ \hline 1 & 3 & 2^2 & 7 & 11 & 2 \cdot 3^2 & 29 & 47 & 2^2 \cdot 19 & 3 \cdot 41 & 199 & 2 \cdot 7 \cdot 23 & 521 & 3 \cdot 281 \end{array} \right.,$$

LIMINAL $SL_2\mathbb{Z}_p$ -REPRESENTATIONS

	15	16	17	18	19	20	21
	$2^2 \cdot 11 \cdot 31$	$13 \cdot 17$	3571	$2 \cdot 3 \cdot 13 \cdot 37$	9349	15127	$2^4 \cdot 29 \cdot 211$

We list the value of L_{2n+1} with $0 \leq n \leq 10$ for (ii) $m = -1$, (iii) $m = 2$, (iv) $m = -2$, (v) $m = 3$, (vi) $m = -3$:

$2n+1$	1	3	5	7	9	11	13	15	17	19	21
(ii)	1	2^2	11	29	$2^2 \cdot 19$	199	521	$2^2 \cdot 11 \cdot 31$	3571	9349	$2^4 \cdot 29 \cdot 211$
(iii)	1	5	11	13	5	67	181	$5^2 \cdot 11$	101	797	$5 \cdot 13 \cdot 43$
(iv)	1	7	31	127	$7 \cdot 73$	$23 \cdot 89$	8191	$7 \cdot 31 \cdot 151$	131071	524287	$7^2 \cdot 127 \cdot 337$
(v)	1	2^3	31	83	$2^3 \cdot 17$	67	1559	$2^3 \cdot 29 \cdot 31$	21929	44917	$2^3 \cdot 41 \cdot 83$
(vi)	1	$2 \cdot 5$	61	337	$2 \cdot 5 \cdot 181$	$23 \cdot 491$	51169	$2 \cdot 5^2 \cdot 61 \cdot 89$	$67 \cdot 21487$	7634353	$2 \cdot 5 \cdot 41 \cdot 293 \cdot 337$

We may often find a large prime number there.

REMARK 5.3. At least when $m = -1$ (the classical case), the converse of Proposition 5.1 does not hold. Indeed, Lagarias [Lag85] proved that the density of

$$S_A := \{p \mid p \equiv \pm 1 \pmod{5} \text{ and } p \text{ divides } L_n \text{ for some } n \in \mathbb{Z}_{>0}\}$$

in the set of prime numbers is $5/12$. So, the density of

$$\{p \mid p \equiv \pm 1 \pmod{5} \text{ and } p \text{ divides } L_{2n+1} \text{ for some } n \in \mathbb{Z}_{\geq 0}\}$$

is smaller than the density $1/2$ of the set $\{p \mid p \equiv \pm 1 \pmod{5}\}$.

6. Cyclic covers

Let $\Delta_K(t)$ denote the Alexander polynomial of a knot K in S^3 . Then Fox–Weber’s formula [Web79] asserts that the $\mathbb{Z}/n\mathbb{Z}$ -cover $M_n \rightarrow S^3$ branched over K satisfies

$$r_n := |H_1(M_n; \mathbb{Z})| = |\text{Res}(t^n - 1, \Delta_K(t))|$$

for each $n \in \mathbb{Z}_{>0}$. Here, if G is a finite group, then $|G|$ denotes the order $\#G$ of G , and if G is an infinite group, then we put $|G| = 0$. For polynomials $f(t), g(t) \in \mathbb{Z}[t]$, $\text{Res}(f(t), g(t)) \in \mathbb{Z}$ denotes their resultant. We remark that if K is a two-bridge knot of genus one, then the precise group structure of $H_1(M_n)$ is known by Fox and others ([Fox60], see also [Ily21]). Now, we assert the following.

THEOREM 6.1. *If $\Delta_K(t) = mt^2 + (1 - 2m)t + m$ with $m \in \mathbb{Z}$, then $p \mid r_n$ for some odd n implies that $\left(\frac{4m^2 - m}{p}\right) = 1$.*

Proof. Since Euler’s totient function satisfies $\varphi(m) \leq 2$ only for $m \leq 6$, we see that $r_n = 0$ happens only for $m = 1$ and $6 \mid n$, so we have $0 \neq r_n = \#H_1(M_n)$ for any odd n .

If $m = 0$, then $r_n = 1$ and $p \nmid r_n$. Suppose $m \neq 0$. Let us write $\Delta_K(t) = mt^2 + (1 - 2m)t + m = m(t - \alpha)(t - \beta)$ with $m \in \mathbb{Z}$. Then, we have $\alpha\beta = 1$, $\alpha + \beta = \frac{2m-1}{m}$, and $r_n = m^n(\alpha^n - 1)(\beta^n - 1) = m^n(2 - \alpha^n - \beta^n)$. In addition, let us write $t^2 - t + m = (t - a)(t - b)$, so that we have $a + b = 1$, $ab = m$, $a^2 + b^2 = 1 - 2m = -m(\alpha + \beta)$, $a^2b^2 = m^2 = (-m\alpha)(-m\beta)$, and hence $\{a^2, b^2\} = \{-m\alpha, -m\beta\}$. We may assume $a^2 = -m\alpha$, $b^2 = -m\beta$.

If n is odd, then we have $L_n^2 = (a^n + b^n)^2 = ((-m\alpha)^n + (-m\beta)^n + 2m^n) = m^n(2 - \alpha^n - \beta^n) = -\text{Res}(t^n - 1, \Delta_K(t)) = r_n$. Hence, the assertion follows from Proposition 5.1. \square

Proof of Theorem 1.1. Since $\Delta_{J(2k, 2l)}(t) = klt^2 + (1 - 2kl)t + kl$, by putting $m = kl$, Theorems 3.1 and 6.1 yield the assertion. \square

REMARK 6.2. If n is even, then we have $\text{Res}(t^n - 1, \Delta_k(t)) = m^n(\alpha^n + \beta^n - 2) = ((-m\alpha)^n + (-m\beta)^n - 2m^n) = (a^n - b^n)^2 = (a - b)^2 F_n^2 = (1 - 4m)F_n^2 = L_n^2 - (2m^{n/2})^2$. Thus, $p \mid r_n$ does not lead to a similar condition of the Legendre symbol.

EXAMPLE 6.3. If $4m^2 - m$ has a square factor, then the converse of Theorem 1.1 does not hold. For instance, if $m = 16$ and $p = 3$, then $4m^2 - m = 16 \cdot 63 = 7 \cdot 12^2$, $(7/3) = (1/3) = 1$. So $J(2, 32)$ admits a liminal representation over \mathbb{Z}_3 . In this case $3 \nmid r_{2n-1}$ for all $n \in \mathbb{Z}_{>0}$.

REMARK 6.4. The following are still in mystery.

(1) Does Theorem 1.1 extend to some wider class of knots? It is literally false in general; we find that $K = 6_3$ with $\Delta_K(t) = -t^4 + 3t^3 - 3t^2 + 3t - 1$ satisfies $r_5 = |\text{Res}(t^5 - 1, \Delta_K(t))| = 2^4$, while the intersection of two varieites is empty in \mathbb{F}_2^2 . However, such a prime number seems very rare. We expect that there is a correct formulation or interpretation in a more general setting.

(2) One may believe that the Burde–de Rham theory is in the background of Theorem 1.1. However, there is no obvious implication, and we rather expect that our result will shed a new light around there.

(3) The Burde–de Rham theory has been extended to higher-dimensional representations by Heusner–Porti [HP15]. We wonder if our study also extends to higher dimensional cases. The study of $\text{SL}_3\mathbb{C}$ -character variety of the figure-eight knot [HMP16] might be a clue.

(4) As we asked in Question 4.1, it seems to be a subtle question to ask if there always exists a liminal $\text{SL}_2\mathbb{Z}_p$ -representation under the condition of Theorem 3.1.

REFERENCES

- BTTU23 Léo Bénard, Ryoto Tange, Anh T. Tran, and Jun Ueki, *Multiplicity of non-acyclic SL_2 -representations and L -functions of the twisted Whitehead links*, preprint. arXiv:2303.15941, 2023.
- Bur67 Gerhard Burde, *Darstellungen von Knotengruppen*, Math. Ann. **173** (1967), 24–33. MR 212787
- dR67 Georges de Rham, *Introduction aux polynômes d’un nœud*, Enseign. Math. (2) **13** (1967), 187–194. MR 240804
- Fox60 Ralph H. Fox, *The homology characters of the cyclic coverings of the knots of genus one*, Ann. of Math. (2) **71** (1960), 187–196. MR 119210
- HMP16 Michael Heusener, Vicente Muñoz, and Joan Porti, *The $\text{SL}(3, \mathbb{C})$ -character variety of the figure eight knot*, Illinois J. Math. **60** (2016), no. 1, 55–98. MR 3665172
- HP15 Michael Heusener and Joan Porti, *Representations of knot groups into $\text{SL}_n(\mathbb{C})$ and twisted Alexander polynomials*, Pacific J. Math. **277** (2015), no. 2, 313–354. MR 3402353
- HPSP01 Michael Heusener, Joan Porti, and Eva Suárez Peiró, *Deformations of reducible representations of 3-manifold groups into $\text{SL}_2(\mathbb{C})$* , J. Reine Angew. Math. **530** (2001), 191–227. MR 1807271
- Ily21 Mednykh Ilya, *Homology group of branched cyclic covering over a 2-bridge knot of genus two*, preprint. arXiv:2111.04292, November 2021.
- KMTT18 Takahiro Kitayama, Masanori Morishita, Ryoto Tange, and Yuji Terashima, *On certain L -functions for deformations of knot group representations*, Trans. Amer. Math. Soc. **370** (2018), no. 5, 3171–3195. MR 3766846
- Lag85 J. C. Lagarias, *The set of primes dividing the Lucas numbers has density 2/3*, Pacific J. Math. **118** (1985), no. 2, 449–461. MR 789184
- Mar16 Julien Marché, *Character varieties in SL_2 and Kauffman Skein algebras*, Topology, Geometry and Algebra of low-dimensional manifolds, RIMS Kôkyûroku, no. 1991, Res. Inst. Math. Sci. (RIMS), Kyoto, 4 2916, pp. 27–42.

- Maz11 Barry Mazur, *How can we construct abelian Galois extensions of basic number fields?*, Bull. Amer. Math. Soc. (N.S.) **48** (2011), no. 2, 155–209. MR 2774089
- MTT24 Yasushi Mizusawa, Ryoto Tange, and Yuji Terashima, *On the Burde–de Rham theorem for finitely presented pro- p groups*, to appear in Int. Math. Res. Not. IMRN (2024), arXiv:2408.01270.
- MTTU17 Masanori Morishita, Yu Takakura, Yuji Terashima, and Jun Ueki, *On the universal deformations for SL_2 -representations of knot groups*, Tohoku Math. J. (2) **69** (2017), no. 1, 67–84. MR 3640015
- Neu99 Jürgen Neukirch, *Algebraic number theory*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 322, Springer-Verlag, Berlin, 1999, Translated from the 1992 German original and with a note by Norbert Schappacher, With a foreword by G. Harder. MR 1697859 (2000m:11104)
- Nys96 Louise Nyssen, *Pseudo-représentations*, Math. Ann. **306** (1996), no. 2, 257–283. MR 1411348 (98a:20013)
- Tra18 Anh T. Tran, *Twisted Alexander polynomials of genus one two-bridge knots*, Kodai Math. J. **41** (2018), no. 1, 86–97. MR 3777388
- TTU22 Ryoto Tange, Anh T. Tran, and Jun Ueki, *Non-acyclic SL_2 -representations of twist knots, -3 -Dehn surgeries, and L -functions*, Int. Math. Res. Not. IMRN (2022), no. 15, 11690–11731. MR 4458562
- TU24 Ryoto Tange and Jun Ueki, *Twisted Iwasawa invariants of knots*, Math. Nachr. **297** (2024), no. 4, 1519–1534. MR 4734983
- Web79 Claude Weber, *Sur une formule de R. H. Fox concernant l’homologie des revêtements cycliques*, Enseign. Math. (2) **25** (1979), no. 3-4, 261–272 (1980). MR 570312 (81d:57011)

Honami Sakamoto sakamo10ho73@gmail.com

Department of Mathematics, Faculty of Science, Ochanomizu University; 2-1-1 Otsuka, Bunkyo-ku, 112-8610, Tokyo, Japan

Ryoto Tange rtange.math@gmail.com

Department of Mathematics, School of Education, Waseda University; 1-104 Totsuka-cho, Shinjuku-ku, 169-8050, Tokyo, Japan

Jun Ueki uekijun46@gmail.com

Department of Mathematics, Faculty of Science, Ochanomizu University; 2-1-1 Otsuka, Bunkyo-ku, 112-8610, Tokyo, Japan