Superdiffusive Central Limit Theorem for the Stochastic Burgers Equation at the critical dimension

January 10, 2025

Giuseppe Cannizzaro¹ **, Quentin Moulard**² **, Fabio Toninelli**²

¹University of Warwick, UK

²Technical University of Vienna, Austria

Email: giuseppe.cannizzaro@warwick.ac.uk, quentin.moulard@tuwien.ac.at, fabio.toninelli@tuwien.ac.at

Abstract

The Stochastic Burgers Equation (SBE) is a singular, non-linear Stochastic Partial Differential Equation (SPDE) that describes, on mesoscopic scales, the fluctuations of stochastic driven diffusive systems with a conserved scalar quantity. In space dimension $d = 2$, the SBE is critical, being formally scale invariant under diffusive scaling. As such, it falls outside of the domain of applicability of the theories of Regularity Structures and paracontrolled calculus. In apparent contrast with the formal scale invariance, we fully prove the conjecture first appeared in [H. van Beijeren, R. Kutner, & H. Spohn, Phys. Rev. Lett., 1986] according to which the 2d-SBE is logarithmically superdiffusive, i.e. its diffusion coefficient diverges like $(\log t)^{2/3}$ as $t \to \infty$, thus removing subleading diverging multiplicative corrections in [D. De Gaspari & L. Haunschmid-Sibitz, Electron. J. Probab., 2024] and in [H.-T. Yau, Ann. of Math., 2004] for 2d-ASEP. We precisely identify the constant prefactor of the logarithm and show it is proportional to $\lambda^{4/3}$, for $\lambda > 0$ the coupling constant, which, intriguingly, turns out to be exactly the same as for the *one-dimensional* Stochastic Burgers/KPZ equation. More importantly, we prove that, under super-diffusive space-time rescaling, the SBE has an explicit Gaussian fixed point in the Renormalization Group sense, by deriving a superdiffusive central limit-type theorem for its solution.

This is the first scaling limit result for a critical singular SPDE, beyond the weak coupling regime, and is obtained via a refined control, on all length-scales, of the resolvent of the generator of the SBE. We believe our methods are well-suited to study other out-of-equilibrium driven diffusive systems at the critical dimension, such as 2d-ASEP, which, we conjecture, have the same large-scale Fixed Point as SBE.

Key words and phrases. Stochastic Partial Differential equations, critical dimension, Stochastic Burgers equation, super-diffusion, diffusion matrix, Asymmetric Simple Exclusion Process

Contents

1 Introduction

In recent years, the interest in singular Stochastic Partial Differential Equations (SPDEs) has grown dramatically and the field has witnessed spectacular progress. Virtually any *subcritical equation*, i.e. whose non-linearity becomes weaker when *zooming in at small scales*, can be given a rigorous path-wise meaning via the theory of Regularity Structures [\[Hai14\]](#page-71-0), paracontrolled cal-culus [\[GIP15\]](#page-71-1), or the Flow approach [\[Kup16,](#page-72-0) [Duc22\]](#page-70-0)^{[1](#page-1-1)}. Notable examples include the dynamical Φ_d^4 model for $d < 4$ [\[Hai14\]](#page-71-0), the Yang-Mills-Higgs equation in $d = 2, 3$ [\[CCHS22,](#page-69-0) [CCHS24\]](#page-69-1) and the KPZ equation [\[Hai13,](#page-71-2) [GP17\]](#page-71-3), just to mention a few. On the other extreme, the understanding of *supercritical equations*, i.e. whose non-linearity becomes weaker when *zooming out at large scales*, is much less systematic but some progress has been recently made. The large-scale be-haviour for the (non-)linear multiplicative stochastic heat equation [\[GRZ18,](#page-71-4) [GL20,](#page-71-5) [GHL23\]](#page-71-6), the KPZ equation [\[CCM24,](#page-69-2) [DGRZ20,](#page-70-1) [CNN22,](#page-70-2) [LZ22\]](#page-72-1) and the Stochastic Burgers equation [\[CGT24\]](#page-70-3), all in dimension $d \geq 3$, has been identified and convergence to the linear Stochastic Heat Equation (SHE) under diffusive scaling, proved.

In contrast, *critical SPDEs*, i.e. those which are formally scale-invariant, meaning that the non-linearity is left unchanged when *zooming either in or out*, are way less understood.

¹Let us also mention the energy solution approach of [\[GJ14,](#page-71-7) [GP18,](#page-71-8) [GPP24\]](#page-71-9) which is though not path-wise and needs the explicit knowledge of the invariant measure, but is less technically demanding and is better suited to determine scaling-limits of interacting particle systems.

Paradigmatic examples in dimension $d = 2$ are the 2d KPZ equation, the Anisotropic KPZ equation and the 2d Stochastic Burgers Equation (SBE). While, in view of scale-invariance, naive scaling arguments fail to predict whether the effect of non-linearity is relevant or not, nonrigorous Renormalization Group techniques suggest that at large scales their solution approaches a "fixed point", whose nature and scaling exponents depend crucially on the symmetries of the equation at hand. For instance, for AKPZ and SBE, [\[Wol91\]](#page-73-0) and [\[vBKS85\]](#page-73-1) respectively predict that the diffusive scaling exponents should be modified by logarithmic corrections but a Gaussian fixed point is still expected, while for the $2d$ KPZ equation the fixed point should be non-Gaussian [\[BS95\]](#page-69-3) (and the scaling exponents are not known). Up to now, however, we are not aware of any singular, critical SPDE whose large-scale behaviour has been rigorously identified, except in the *weak coupling regime*, where the strength of the non-linearity is artificially tuned and is forced to vanish as the scale of the system grows (see Section [1.2](#page-8-0) for a discussion).

In this work, we obtain the *first full scaling limit result* for a singular critical SPDE *beyond* the weak coupling regime: we show that the 2d SBE (with coupling constant of order 1) converges *under logarithmically superdiffusive scaling*, to a linear Stochastic Heat Equation with explicit renormalised coefficients.

Before precisely stating our theorems, let us introduce the equation of interest and put it into context. In any dimension $d \geq 1$, the SBE is formally defined on $\mathbb{R}_+ \times \mathbb{R}^d$ by

$$
\partial_t u = \frac{1}{2} \Delta u + \lambda (\mathfrak{w} \cdot \nabla) u^2 + \nabla \cdot \vec{\xi}, \qquad (1.1)
$$

where $\vec{\xi} = (\xi_1, \dots, \xi_d)$ is a d-dimensional vector-valued space-time white noise, and $\lambda > 0$ is the so-called *coupling constant*. Here, $\mathfrak w$ is a unit vector in $\mathbb R^d$, but, by rotational invariance, the specific choice of $\mathfrak w$ is irrelevant.

The SBE was introduced in [\[vBKS85\]](#page-73-1) as an effective continuous model for the mesoscopic fluctuations of driven diffusive systems with a single conserved scalar quantity, such as the Asymmetric Simple Exclusion Process (ASEP), the Zero-Range Process with asymmetric rates and other driven, conservative, interacting particle systems. In fact, the three terms at the righthand side of (1.1) represent respectively a diffusive term, the non-linear effect of the driving field acting in direction w, and a microscopic noise of conservative type. On the basis of the so-called mode-coupling theory, [\[vBKS85\]](#page-73-1) predicts the large-scale behaviour of SBE in any dimension $d \geq 1$: in the subcritical dimension $d = 1$, the authors conjecture strong superdiffusion with a $t^{1/3}$ divergence of the diffusion coefficient for large time t; in the critical dimension $d = 2$, logarithmic enhancement of diffusion of order $(\log t)^{2/3}$; and, in the super-critical dimensions $d > 3$, classical diffusive behaviour. Let us point out that in $d = 1$, the SBE is nothing but the space derivative of the KPZ equation which, by now, is mathematically well-understood (see [\[QS15\]](#page-72-2), for a review): not only the $t^{1/3}$ behaviour was confirmed [\[BQS11\]](#page-69-4), but the solution was shown to have a non-Gaussian and universal fixed point (the "KPZ fixed point" [\[MQR21\]](#page-72-3)), that also describes the large-scale behaviour of a large class of one-dimensional driven diffusive processes such as ASEP [\[QS23\]](#page-72-4) and directed polymers in random environment [\[Vir20\]](#page-73-2). In dimension $d \geq 3$, the series of works [\[LY97,](#page-72-5) [CLO01,](#page-70-4) [LOY04\]](#page-72-6) has confirmed the conjectured normal diffusion of ASEP and the convergence of the particle density fluctuations to a linear

Stochastic Heat Equation at large scales. A counterpart on the SPDE side is the recent [\[CGT24\]](#page-70-3), where the same result is obtained for the SBE in $d \geq 3$. As for the critical dimension $d = 2$, the breakthrough work [Yauo4] proved $(\log t)^{2/3}$ diffusivity for ASEP, while in [\[GHS24\]](#page-71-10) the analogous was shown for the SBE. Let us stress though that in both cases, the upper and lower bounds on the diffusivity feature *diverging subleading corrections* and their statement is phrased in the *Tauberian sense*, i.e. in terms of Laplace transform.

The problem when analysing the SPDE [\(1.1\)](#page-2-0) is that it is severely ill-defined due to the presence of the non-linearity. Since dimension $d = 2$ is critical, it falls outside of the scope of any of the path-wise theories mentioned above which successfully addressed subcritical equations and, in fact, a local solution theory is not even expected.[2](#page-3-1) Our main results are the following. First, we determine the *sharp large-time asymptotic behaviour of the diffusion matrix*, including the explicit constant prefactor, thus significantly improving over both [\[Yau04,](#page-73-3) [GHS24\]](#page-71-10) and fully addressing the long-standing conjecture of [\[vBKS85\]](#page-73-1). Second, we move well beyond that and obtain the (Gaussian) *large-scale fixed point of the regularised SBE*, when space-time is rescaled in a suitable, logarithmically superdiffusive way.

In addition to being the first such result for critical SPDEs, our work opens intriguing perspectives for other driven diffusive systems, including interacting particle systems, as we will discuss more in detail at the end of Section [1.2.](#page-8-0) In a broader perspective, this paper can be seen as an out-of-equilibrium counterpart of the celebrated work [\[ADC21\]](#page-69-5) that proves that the large-scale limit of the Ising and Φ_d^4 $\frac{4}{d}$ equilibrium measures, in the critical dimension $d = 4$, is Gaussian, despite the presence of logarithmic corrections to the mean field critical exponents.

1.1 The main results: asymptotic superdiffusion and large-scale limit

To give a meaning to the formal equation (1.1) , we regularise it with a smooth compactly supported, non-negative, even-symmetric function ρ whose mass equals 1. Then, we consider

$$
\partial_t u = \frac{1}{2} \Delta u + \lambda \mathcal{N}^1(u) + \nabla \cdot \vec{\xi}^1, \qquad (1.2)
$$

where $u = u(t, x)$, for $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^2$, $\vec{\xi}^1 \stackrel{\text{def}}{=} \varrho * \vec{\xi} \stackrel{\text{def}}{=} (\varrho * \xi_1, \varrho * \xi_2)$, and

$$
\mathcal{N}^1(f) \stackrel{\text{def}}{=} (\mathfrak{w} \cdot \nabla) \varrho^{*2} * f^2.
$$
 (1.3)

Note that, thanks to the convolution(s), the noise $\bar{\xi}^1$ is smooth and, as we will see in Proposition [2.3,](#page-14-0) the dynamics is globally well-posed and the solution u smooth in space. Moreover, it is a Markov process (with respect to a suitable filtration) and stationary. Indeed, a fundamental property of [\(1.1\)](#page-2-0) is that, formally, it admits the law of a spatial white noise η on \mathbb{R}^2 , denoted by \mathbb{P} , as a stationary measure, which is the continuum analog of the stationarity of Bernoulli i.i.d. measures for ASEP. The regularisation in (1.2) has been chosen similarly to [\[FQ15\]](#page-70-5) (which is, however, in $d = 1$, in such a way that stationarity is preserved, which means that the law of the mollified space white noise $\eta^1 \stackrel{\text{def}}{=} \varrho * \eta$, denoted by \mathbb{P}^{ϱ} , is stationary for the solution to [\(1.2\)](#page-3-2).

²In a direction orthogonal to ours, the noteworthy work [\[GPP24\]](#page-71-9) determines a local solution theory for a scalingcritical SPDE, the fractional SBE in $d = 1$ but their approach does not apply to $d = 2$ and, contrary to ours, their focus in on small-scales.

INTRODUCTION 5

In order to study the large-scale behaviour of u , a convenient tool is the so-called diffusion matrix, which, for $t \geq 0$, is the 2×2 matrix whose entries are given by

$$
D_{i,j}(t) = \frac{1}{t} \int_{\mathbb{R}^2} x_i x_j \left[S(t, x) - S(0, x) \right] dx, \quad S(t, x) \stackrel{\text{def}}{=} \mathbf{E}(u(0, 0) u(t, x)), \quad (1.4)
$$

with E denoting the expectation with respect to the product measure of η^1 and the space-time noise $\vec{\xi}^1$, and u being the stationary solution of [\(1.2\)](#page-3-2) (i.e. u_0 has the same law as η^1). Intuitively, if $e \in \mathbb{R}^2$ is an unit vector, one can interpret $\langle D(t) e, e \rangle$ as the ratio $(L_e(t))^2/t$, with $L_e(t)$ the time-dependent correlation length in the direction e at time t , defined as the distance beyond which the correlation $S(t, x)$ is negligible. While it is useful to keep this interpretation in mind, we will not try to make this connection rigorous.

The integral defining $D(t)$ is convergent because $x \mapsto S(t, x)$ decays fast enough as $|x| \rightarrow$ $+\infty$, as is proven in Section [2.](#page-13-0) Further, upon taking $\lambda = 0$ in [\(1.2\)](#page-3-2), which reduces the SBE to the linear stochastic heat equation, one has

$$
S(t,x)|_{\lambda=0} = \left(\varrho^{*2} * \mathcal{P}_t\right)(x)
$$

with \mathcal{P}_t the usual heat semigroup, and therefore

$$
D(t)|_{\lambda=0} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.
$$

In particular, if for large t the diffusion matrix of a given system has $O(1)$ entries, then its large-scale behaviour is said to be diffusive, while if it grows with t it is *superdiffusive* (and if it decreases, subdiffusive). Even though the formal scale invariance of (1.1) could at first sight suggest the former, this turns out not to be the case. Our first main theorem determines the precise asymptotic behaviour of $D(t)$ as $t \to \infty$.

Theorem 1.1 *Let* $\lambda > 0$ *and, without loss of generality, take* $\mathfrak{w} = \mathfrak{e}_1$ *, where* $(\mathfrak{e}_1, \mathfrak{e}_2)$ *stands for* the canonical basis of \mathbb{R}^2 . Then as $t \to +\infty$ it holds

$$
D(t) = \begin{pmatrix} C_{\text{eff}}(\lambda) (\log t)^{2/3} (1 + o(1)) & 0 \\ 0 & 1 \end{pmatrix},
$$
 (1.5)

where the constant $C_{\text{eff}}(\lambda)$ *is explicit and given by*

$$
C_{\rm eff}(\lambda) \stackrel{\rm def}{=} \left(\frac{3}{2\pi}\right)^{2/3} \lambda^{4/3} \,. \tag{1.6}
$$

As mentioned above, Theorem [1.1](#page-4-0) substantially improves on the previously known bounds [\[GHS24\]](#page-71-10) (and the analog for 2d ASEP in [Yauo4]). Our result is for D itself and not for the Laplace transform of $t \mapsto tD(t)$, as was the case in the above-mentioned references. More importantly, it eliminates spurious multiplicative, subdominant corrections of order (log log log t)^C in [\[GHS24\]](#page-71-10) (respectively, of order exp[($log log log t$)²] for ASEP in [Yauo4]) and it identifies the precise constant $C_{\text{eff}}(\lambda)$. Further, the "o(1)" in the first entry is explicit and our estimates actually show that $D_{1,1}(t) = C_{\text{eff}}(\lambda) \log(t)^{2/3} + O(\log(t)^{1/2 + o(1)})$ (see [\(6.20\)](#page-51-2)).

Let us also underline two important implications of (1.5) and (1.6) . First, the effective diffusivity in direction w is *independent of the microscopic diffusivity in the same direction*, which means that it is *fully produced by the non-linearity*. To see this, take $\mathfrak{w} = \mathfrak{e}_1, \nu_1, \nu_2 > 0$ and Q to be the 2 × 2 diagonal matrix such that $Q_{i,j} = \nu_i \delta_{i,j}, i, j = 1, 2$. Consider the solution v of

$$
\partial_t v = \frac{1}{2} \nabla \cdot Q \nabla v + \lambda \tilde{\mathcal{N}}^1(v) + \nabla \cdot \sqrt{Q} \,\vec{\xi}^1
$$

where $\tilde{\mathcal{N}}^1$ coincides with \mathcal{N}^1 in [\(1.3\)](#page-3-3) but with the mollifier ϱ replaced by $(\nu_1 \nu_2)^{-1/2} \varrho (Q^{-1/2})$. Then, easy scaling arguments show that $u(t, x) \stackrel{\text{law}}{=} (v_1v_2)^{1/4} v(t, \sqrt{Q}x)$, for u the solution to [\(1.2\)](#page-3-2) with λ replaced by $\lambda \nu_1^{-3/4} \nu_2^{-1/4}$, and the diffusion matrix associated to v, D^v satisfies

$$
D_{i,j}^v(t) = (\nu_i \nu_j)^{1/2} D_{i,j}(t).
$$
\n(1.7)

Therefore, as $t \to \infty$, [\(1.5\)](#page-4-1) and [\(1.6\)](#page-4-2) imply that $D_{1,2}^v(t) = D_{2,1}^v(t) = 0$, $D_{2,2}^v = \nu_2$ and especially that $D_{1,1}^v(t) = C(\nu_1, \nu_2, \lambda) (\log t)^{2/3} (1 + o(1))$ with

$$
C(\nu_1, \nu_2, \lambda) = \nu_2^{-1/3} C_{\text{eff}}(\lambda)
$$

so that overall D^v is *independent of* ν_1 .

Remark 1.2 The scaling argument above can be reversed, to show that the lack of dependence of the diffusion matrix on the parameter ν_1 forces the $\lambda^{4/3}$ dependence of $C_{\text{eff}}(\lambda)$. In fact, by [\(1.7\)](#page-5-0), $C(\nu_1, \nu_2, \lambda) = \nu_1 C(1, 1, \lambda \nu_1^{-3/4} \nu_2^{-1/4})$, which means that, assuming $C(\nu_1, \nu_2, \lambda)$ is independent of ν_1 , we get that $C_{\text{eff}}(\lambda) = C(1, 1, \lambda)$ is proportional to $\lambda^{4/3}$.

The proportionality $C_{\text{eff}}(\lambda) \propto \lambda^{4/3}$ is quite intriguing, especially because exactly the same holds for the *one-dimensional* Stochastic Burgers equation^{[3](#page-5-1)}, despite the fact that the t-dependence of the diffusion coefficients are completely different in the two cases (of order $t^{1/3}$ in one dimension, in contrast with $(\log t)^{2/3}$ in two dimensions). The equality of the exponents of λ in $d = 1, 2$ is *not* a coincidence, but rather a reflection of the fact that the diffusion coefficient is asymptotically (for large t) independent of the microscopic diffusion coefficient in the direction where the non-linearity acts.

The second implication of Theorem [1.1](#page-4-0) is the *precise identification of the scaling* under which one can expect [\(1.2\)](#page-3-2) to approach its Fixed Point. If one naively tries the classical diffusive rescaling, i.e. $u_{\text{diff}}^{\tau}(t, x) \stackrel{\text{def}}{=} \tau^{1/2} u(\tau t, \tau^{1/2} x)$, then an argument similar to that used to deduce [\(1.7\)](#page-5-0), shows that $D_{\text{diff}}^{\tau}(t) = D(\tau t)$ which, for $t > 0$ fixed and $\tau \to \infty$, blows up as $(\log \tau)^{2/3}$ by [\(1.5\)](#page-4-1). Thus, the diffusive scaling must be adjusted logarithmically and the above statement strongly suggests that one should instead look at

$$
u^{\tau}(t,x) \stackrel{\text{def}}{=} \tau^{1/2} (\log \tau)^{1/6} u(\tau t, \tau^{1/2} R_{\tau}^{-1/2} x), \quad R_{\tau} \stackrel{\text{def}}{=} \begin{pmatrix} (\log \tau)^{-2/3} & 0\\ 0 & 1 \end{pmatrix}, \tag{1.8}
$$

³This follows immediately from elementary scaling properties of the 1d SBE, plus the fact that its diffusion coefficent grows proportionally to the cube root of time

as, under such scaling, the entries of the diffusion matrix are asymptotically $O(1)$ (see [\(6.1\)](#page-46-2)). In principle, Theorem [\(1.1\)](#page-4-0) does not fix the size of the fluctuations, i.e. the prefactor in the definition of u^{τ} , but this needs to be chosen in such a way that the law \mathbb{P}^{τ} of the mollified white noise $\varrho_{\tau} * \eta$, with $\varrho_{\tau}(\cdot) \stackrel{\text{def}}{=} \tau (\log \tau)^{1/3} \varrho(\tau^{1/2} R_{\tau}^{-1/2} \cdot)$, is a stationary measure for u^{τ} .

The equation solved by u^{τ} then becomes

$$
\partial_t u^\tau = \frac{1}{2} \nabla \cdot R_\tau \nabla u + \frac{\lambda}{\sqrt{\log \tau}} \mathcal{N}^\tau(u^\tau) + \nabla \cdot \sqrt{R_\tau} \vec{\xi}^\tau, \tag{1.9}
$$

with $\vec{\xi}^{\tau} \stackrel{\text{law}}{=} \varrho_{\tau} * \vec{\xi}$ and $\mathcal{N}^{\tau}(f) \stackrel{\text{def}}{=} \mathcal{N}^{\varrho_{\tau}}(f)$, see [\(1.3\)](#page-3-3). Note that, for $\tau \to \infty$, ϱ_{τ} is converging to a δ-function so that, morally, identifying the large-scale limit of u^{τ} coincides with making sense of the singular critical SPDE [\(1.1\)](#page-2-0).

Let us also stress that all the terms at the right hand side of (1.9) describing the dynamics in the direction e_2 , i.e. the Laplacian $\partial_2^2 u^\tau$ and the noise $\partial_2 \xi_2^\tau$, are unaffected by the scaling, meaning that their coefficients are constant in τ , and those in \mathfrak{e}_1 , i.e. $\partial_1^2 u^{\tau}$, the non-linearity and $\partial_1 \xi_1^{\tau}$, are all multiplied by τ -dependent quantities that vanish as $\tau \to \infty$. The main challenge we have to face is that, even if the vanishing coefficients should tame the growth caused by the non-linearity, they also lessen the regularising properties of the linear part of the equation, making it apriori unclear what to expect in the limit.

This is precisely what is addressed by our second main result, which more broadly provides a full *central limit theorem for* u^{τ} . It shows that as $\tau \to \infty$, u^{τ} converges in finite-dimensional distributions to a Gaussian process, given by the solution of an *anisotropic linear Stochastic Heat Equation* with *explicit renormalised* coefficients.

Theorem 1.3 *Let* Q *be a probability measure, absolutely continuous with respect to* P *and for* $\tau > 0$, let $(u^{\tau}(t))_{t \geq 0}$ be the solution of [\(1.9\)](#page-6-0) with $\mathfrak{w} = \mathfrak{e}_1$ and initial condition $u^{\tau}(0) = \varrho_{\tau} * u_0$ *for* $u_0 \sim \mathbb{Q}$. *Then, for any* $k \in \mathbb{N}$, $0 \le t_1 \le \cdots \le t_k$ *and* $\varphi_1, ..., \varphi_k \in \mathcal{S}(\mathbb{R}^2)$,

$$
(\langle u^{\tau}(t_1), \varphi_1 \rangle, \cdots, \langle u^{\tau}(t_k), \varphi_k \rangle) \xrightarrow{\tau \to +\infty} (\langle u^{\text{eff}}(t_1), \varphi_1 \rangle, \cdots, \langle u^{\text{eff}}(t_k), \varphi_k \rangle)
$$

where the convergence is in law and $(u^{\text{eff}}(t))_{t\geq0}$ is the process that starts at $u^{\text{eff}}(0) = u_0$ and *solves the anisotropic linear Stochastic Heat Equation given by*

$$
\partial_t u^{\text{eff}} = \frac{1}{2} \, \nabla \cdot D_{\text{eff}} \, \nabla u^{\text{eff}} + \nabla \cdot \sqrt{D_{\text{eff}}} \, \vec{\xi}, \quad \text{for} \quad D_{\text{eff}} \stackrel{\text{def}}{=} \begin{pmatrix} C_{\text{eff}}(\lambda) & 0 \\ 0 & 1 \end{pmatrix}, \tag{1.10}
$$

where $\vec{\xi}$ *is a space-time white noise and the constant* $C_{\text{eff}}(\lambda)$ *is that in* [\(1.5\)](#page-4-1)*.*

The previous theorem fully characterises the large-scale behaviour of (1.2) and, more generally, it strongly indicates what to expect for other out-of-equilibrium driven diffusive systems with one conserved quantity at the critical dimension. Phenomenologically, it says that in the \mathfrak{e}_2 direction the original diffusion is unaltered (with the mollifier replaced by the δ function), while, in the \mathfrak{e}_1 direction, the linear dynamics vanishes and the non-linear one produces *a new noise and a new Laplacian* whose strength explicitly depends on the coupling constant λ , i.e. on the microscopic strength of the non-linearity.

Even though the creation of new Laplacian/noise from the non-linearity has already been observed in the context of critical SPDEs (e.g. [\[CSZ20\]](#page-70-6) for KPZ, [\[CET23b\]](#page-69-6) for AKPZ or, more explicitly, [\[CC22\]](#page-69-7) for the multiplicative SHE), this is the first time, up to the authors' knowledge, where this is coupled with a vanishing microscopic diffusivity.

Theorem [1.3](#page-6-1) is a corollary of Theorem [1.4](#page-7-0) below, which concerns the convergence of the semigroup of the solution u^{τ} of [\(1.9\)](#page-6-0) to that of u^{eff} . To properly state it, we need a few preliminary definitions. Let $P_{u_0}^{\tau}$ and $P_{u_0}^{\text{eff}}$ be the laws (with respective expectations $\mathbf{E}_{u_0}^{\tau}$ and $\mathbf{E}_{u_0}^{\text{eff}}$) of u^{τ} and u^{eff} given their initial condition $u_0^{\tau} = u_0 * \varrho_{\tau}$ and u_0 , and denote by $(P_t^{\tau})_{t\geq 0}$ and $(P_t^{\text{eff}})_{t\geq 0}$ their Markov semigroups.

Our analysis of P^{τ} and P^{eff} will be carried out on the spaces $\mathbb{L}^{\theta}(\mathbb{P}^{\tau})$ (to deal with u^{τ}) and $\mathbb{L}^{\theta}(\mathbb{P})$ (to deal with u^{eff}), for $\theta \in [1,\infty)$, where we recall that \mathbb{P} and \mathbb{P}^{τ} stand for the law of the spatial white noise η and its mollified counterpart $\eta^{\tau} = \varrho_{\tau} * \eta$. To relate $\mathbb{L}^{\theta}(\mathbb{P}^{\tau})$ and $\mathbb{L}^{\theta}(\mathbb{P})$, we define the isometric embedding $\iota_{\tau} : \mathbb{L}^{\theta}(\mathbb{P}^{\tau}) \to \mathbb{L}^{\theta}(\mathbb{P})$ which sends the functional $(\eta^{\tau} \to F(\eta^{\tau}))$ to the functional $(\eta \to F(\rho_{\tau} * \eta))$ (for more on it see Lemma [1.7](#page-12-0) below). With a slight abuse of notation, we will say that a sequence $(F^{\tau})_{\tau>0}$, with $F^{\tau} \in \mathbb{L}^{\theta}(\mathbb{P}^{\tau})$, converges to $F \in \mathbb{L}^{\theta}(\mathbb{P})$ if $\iota_{\tau} F^{\tau} \to F$ in $\mathbb{L}^{\theta}(\mathbb{P})$ as $\tau \to +\infty$.

Theorem 1.4 $For any $\theta \in [1, +\infty)$ and sequence $(F^{\tau})_{\tau>0}$ converging to F in $\mathbb{L}^{\theta}(\mathbb{P})$, $(P_t^{\tau} F^{\tau})_{\tau>0}$$ *converges to* $P_t^{\text{eff}}F$ *in* $\mathbb{L}^{\theta}(\mathbb{P})$ *uniformly over* $t \geq 0$ *, that is to say*

$$
\sup_{t\geq 0} \mathbb{E}\left[|\iota_{\tau} P_t^{\tau} F^{\tau} - P_t^{\text{eff}} F|^{\theta}\right] \xrightarrow{\tau \to +\infty} 0. \tag{1.11}
$$

As a consequence, for every $k \geq 1$, $(F_1^{\tau})_{\tau > 0}$, ..., $(F_k^{\tau})_{\tau > 0}$ converging to $F_1, ..., F_k$ in $\mathbb{L}^{\theta}(\mathbb{P})$ for *all* $\theta \in [1, +\infty)$ *, and* $0 \le t_1 \le ... \le t_k$ *, we have*

$$
\mathbf{E}_{u_0}^{\tau}[F_1^{\tau}(u^{\tau}(t_1))\dots F_k^{\tau}(u^{\tau}(t_k))] \xrightarrow{\tau \to +\infty} \mathbf{E}_{u_0}^{\text{eff}}[F_1(u^{\text{eff}}(t_1))\dots F_k(u^{\text{eff}}(t_k))], \qquad (1.12)
$$

where the convergence holds in $\mathbb{L}^{\theta}(\mathbb{P}(\mathrm{d}u_0))$ for every $\theta \in [1, +\infty)$.

The above theorem is substantially stronger than Theorem [1.3.](#page-6-1) In fact, (1.12) says that the finite dimensional distributions of the process u^{τ} converge to those of the limit process not just in average, but in $\mathbb{L}^{\theta}(\mathbb{P})$ with respect to the initial condition. In this sense, the convergence stated in [\(1.12\)](#page-7-1) is stronger than both annealed and stable convergence (and that in [\[KLO12,](#page-72-7) Definition 2.6]), and is closer in spirit to a quenched central limit theorem, although we make no claim of almost-sure convergence. Note also that, because of the supremum in time in (1.11) due to the ergodicity of the limit process, we have a uniform control of the law of the process u^{τ} , *over all time scales*, which rules out pathological deviations from the scaling limit. Moreover, while we formulated Theorem [1.4](#page-7-0) for λ of order 1, this being the most interesting scenario, our techniques would allow to obtain the scaling limit of the process in the whole *crossover window* where $\lambda = \lambda(\tau)$ tends to 0 with τ in any desired way, between the weak-coupling regime (see next section) and the case $\lambda = O(1)$.

The proof of Theorems [1.1](#page-4-0) and [1.4](#page-7-0) relies on a careful analysis of the resolvent of u^{τ} . We will present the overall strategy and provide a thorough account of the novelties of the present work

in Section [2.3,](#page-21-0) after deriving the main properties of u^{τ} for fixed τ and studying the action of its generator \mathcal{L}^{τ} on $\mathbb{L}^2(\mathbb{P}^{\tau})$. For now, let us only point out that our arguments give a *scale-dependent control of the effective diffusion operator*, in the spirit of Renormalization Group. Indeed, what our proof shows is that, for $F \in \mathbb{L}^2(\mathbb{P}^{\tau})$ sufficiently regular (think of $F = \eta(\varphi)$ for some φ smooth), $(1 - \mathcal{L}^{\tau})^{-1}F$ approaches $(1 - \mathcal{D}_{eff}^{\tau})^{-1}F$, which is the resolvent of the solution to the non-homogeneous Stochastic Heat Equation given by

$$
\partial_t u^{\tau}_{\rm eff} = \tfrac{1}{2} \nabla \cdot D_{\rm eff}^{\tau} \nabla u^{\tau}_{\rm eff} + \nabla \cdot \sqrt{D_{\rm eff}^{\tau}} \vec{\xi}
$$

and D_{eff}^{τ} is the operator whose Fourier multiplier is

$$
D_{\text{eff}}^{\tau}(k) = \begin{pmatrix} (\log \tau)^{-2/3} + g^{\tau} (L^{\tau} (\frac{1}{2} |\sqrt{R_{\tau}} k|^2)) & 0\\ 0 & 1 \end{pmatrix}.
$$
 (1.13)

Above, R^{τ} is the matrix in [\(1.8\)](#page-5-2), L^{τ} is a function proportional to $x \mapsto \log(1 + \tau/x)/\log \tau$ (see [\(3.14\)](#page-29-1) for the precise definition) and, interestingly, g^{τ} is the solution of the ODE

$$
\dot{g}^{\tau}(x) = \frac{1}{\sqrt{\nu_{\tau} + g^{\tau}(x)}}, \qquad g^{\tau}(0) = 0.
$$
 (1.14)

Inspecting [\(1.13\)](#page-8-1) we see that for momenta k of order 1 and τ large, one has $D_{\text{eff}}^{\tau}(k) \approx D_{\text{eff}}$ with the latter defined as in [\(1.10\)](#page-6-2), which in turn implies the proximity of u^{τ} and u^{eff} . But more than that, since $L^{\tau}(x)$ *is small when* x *is large* and g^{τ} satisfies an *initial value* problem, the effective diffusivity $g^{\tau}(L^{\tau}(\frac{1}{2}))$ $\frac{1}{2}$ $\sqrt{R_{\tau}}k$ ²)) at scale k *encodes the cumulative effect of momenta larger than* k, that is, *of smaller spatial scales*. In other words, the ODE [\(1.14\)](#page-8-2) corresponds to what would be called in Renormalization Group a *flow equation for the scale-dependent coefficient* g^{τ} .

1.2 Relation with the literature and future perspectives

As alluded to earlier, most of the work on critical SPDEs (or more general out-of-equilibrium critical systems) concerns the so-called *weak coupling* or *intermediate disorder* scaling, which, in a sense, provides a bridge between subcritical and critical equations and can be thought of as a way to zoom into the singular part of the equation at hand. More precisely, starting from a given SPDE (say [\(1.2\)](#page-3-2)), this amounts to introduce *a vanishing coupling constant* multiplying the nonlinearity, the noise or the initial condition (for [\(1.2\)](#page-3-2), set $\lambda = \lambda_{\tau} = \bar{\lambda}/\sqrt{\log \tau}$), and then take the usual diffusive scaling (i.e. $u \mapsto \sqrt{\tau}u(\tau t, \sqrt{\tau}x)$). Even though there is currently no systematic way to analyse such setting, several interesting results in a variety of different contexts have recently appeared. This is the case e.g. for the KPZ equation [\[CD20,](#page-69-8) [Gu20,](#page-71-11) [CSZ20\]](#page-70-6), the AKPZ equation [\[CES21,](#page-69-9) [CET23b\]](#page-69-6), the SBE [\[CGT24\]](#page-70-3), (self-)interacting diffusions [\[CG24,](#page-69-10) [YY24\]](#page-73-4) the Allen-Cahn equation with weakly critical random initial data [\[GRZ24\]](#page-71-12), and, especially, the multiplicative (non-)linear Stochastic Heat Equation in $d = 2$ [\[BC98,](#page-69-11) [CSZ17,](#page-70-7) [DG22,](#page-70-8) [Tao24\]](#page-72-8). The latter in particular culminated in the ground-breaking work [\[CSZ23\]](#page-70-9), which first constructed the scaling limit of directed polymers in random environment in the intermediate disorder regime at the critical value of the disorder strength, a process that now goes under the name of critical "2d-Stochastic Heat Flow" (see also [\[Tsa24\]](#page-72-9), for a more recent characterisation). As already stressed, the present work goes beyond weak coupling as the vanishing coefficients at the right

hand side of [\(1.9\)](#page-6-0) are not put there "by hand" but come from simply *zooming out at the correct scale*, which for the SBE corresponds to (1.8) .

Another critical, out-of-equilibrium model which has attracted significant attention due to its physical importance in connection with turbulence, is the DCGFF, i.e. a singular SDE describing the motion of a Brownian particle in two dimensions, subject to a divergence-free drift given by the curl of a log-correlated random field (e.g. the $2d$ GFF). The analysis of its mean-square displacement was initiated in [\[TV12\]](#page-72-10) and later continued first in [\[CHST22\]](#page-70-10), using functional analytical techniques similar in flavour to those in the present work, and then in [\[CMOW22\]](#page-70-11), via stochastic homogeneisation arguments, where the conjectured annealed logarithmic superdiffusivity (see [\[TV12\]](#page-72-10)) was established. The remarkable recent work [\[ABRK24\]](#page-69-12), which proves the analog of Theorem [1.3](#page-6-1) in this context, obtains the full conjecture of [\[BCGL87\]](#page-69-13) as it derives a quenched invariance principle under a logarithmically adjusted scaling (that said, the tools exploited therein are completely different from ours and it is unclear whether they could be adapted to the SPDE setting).

Ultimately, the major achievement of the present paper is that it establishes a rigorous framework in which the large-scale behaviour of stationary space-time systems at criticality can be analysed. Moreover, in relation to the literature, we believe it raises a number of intriguing questions:

- the convergence, under similar logarithmically adjusted scaling, of the anisotropic selfrepelling polymer of $[TV_12]$, of the density fluctuation profile of 2d ASEP and of other critical interacting particle systems, to a non-trivial Gaussian limit process. We are not aware of any rigorous statement or even explicit conjecture in this sense;
- the investigation of the role and universality of the $\lambda^{4/3}$ dependence [\(1.6\)](#page-4-2) of the diffusion coefficient on the coupling constant (see Remark [1.2\)](#page-5-3) for the above-mentioned systems at *and above* the critical dimension;
- the study of large-scale behaviour of other critical SPDEs with (non-)quadratic nonlinearity, as the AKPZ equation or the stochastic Navier-Stokes equation, and, in the long-run, away from stationarity.

At last, a very fascinating line of works [\[OW24,](#page-72-11) [MOW24\]](#page-72-12) has displayed compelling features of the DCGFF such as intermittency properties and an unexpected connection with a diffusion on the Lie group $SL(2)$, and it would be interesting to see if analogous relations can be found in the present context or even more broadly for other systems at the critical dimension.

Organization of the paper

The rest of the paper is organised as follows. In subsection [1.3,](#page-11-0) we recall basic tools from Wiener space analysis. Section [2.1](#page-13-1) states basic results of the SBE [\(1.9\)](#page-6-0) for fixed value of $\tau > 0$ such as global well-posedness, Markovianity and stationarity. Moreover, the two-point correlation function is shown to decay polynomially fast and a Green-Kubo formula for the diffusion matrix in [\(1.4\)](#page-4-3) is rigorously derived, which we believe is of independent interest. The semigroup and the generator of [\(1.9\)](#page-6-0), together with the action of the latter on Fock spaces, are analysed in Section [2.2](#page-19-0) where we further recall basic properties of the limit equation [\(1.10\)](#page-6-2). In Section [2.3,](#page-21-0) we provide a high-level roadmap that on the one hand outlines the overall strategy of proofs, and

should thus guide the reader throughout, while on the other introduces the main novelties of the paper. Some of these are then detailed in Section [3](#page-24-0) such as: a new splitting of the antisymmetric part of the generator that allows to single out the singularity in Fourier and growth in the "Wiener chaos index" (Section [3.2\)](#page-26-0); a (refined version of the) Replacement Lemma (Lemma [3.6\)](#page-30-0) and the Recursive Replacement Lemma (Lemma [3.9\)](#page-32-0). Section [4](#page-35-0) applies these tools to derive the needed estimates on the resolvent, as summarised in its main statement Theorem [4.1.](#page-35-1) Theorems [1.3](#page-6-1) and [1.4,](#page-7-0) concerning the superdiffusive central limit theorem are proven in Section [5,](#page-43-0) while Theorem [1.1](#page-4-0) about the diffusion matrix in Section [6.](#page-46-0) At last, Appendices [A](#page-51-0) and [B](#page-61-0) contain the proofs of the statements in Section [2,](#page-13-0) while technical bounds and a general replacement estimate (which we deem interesting in its own right) can be found in Appendix [C.](#page-64-0)

Notations and Function spaces

Let e_1 and e_2 be the canonical basis vectors of \mathbb{R}^2 and $|\cdot|$ be the usual Euclidean norm. For $p_1,...,p_n \in \mathbb{R}^2$, we set $p_{1:n} \stackrel{\text{def}}{=} (p_1,...,p_n) \in \mathbb{R}^{2n}$ and, given a 2×2 matrix M , $Mp_{1:n} \stackrel{\text{def}}{=}$ $(Mp_1, ..., Mp_n)$. We write (for $e \in \mathbb{R}^2$)

$$
|p_{1:n}| \stackrel{\text{def}}{=} \left(\sum_{i=1}^n |p_i|^2\right)^{\frac{1}{2}}, \qquad |e \cdot p_{1:n}| \stackrel{\text{def}}{=} \left(\sum_{i=1}^n |e \cdot p_i|^2\right)^{\frac{1}{2}}.
$$
 (1.15)

For $\tau > 0$ let the matrix R_{τ} and $\nu_{\tau} > 0$ be respectively given according to

$$
R_{\tau} \stackrel{\text{def}}{=} \begin{pmatrix} \nu_{\tau} & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad \nu_{\tau} \stackrel{\text{def}}{=} \frac{1}{\left(1 \vee \log \tau\right)^{2/3}}, \tag{1.16}
$$

and note that it holds

$$
|\sqrt{R_{\tau}}p_{1:n}|^2 = \nu_{\tau}|\mathfrak{e}_1 \cdot p_{1:n}|^2 + |\mathfrak{e}_2 \cdot p_{1:n}|^2.
$$
 (1.17)

Let $\mathcal{S}(\mathbb{R}^d)$ be the Schwartz space of smooth functions whose derivatives decay faster then any polynomial and $\mathcal{S}'(\mathbb{R}^d)$ be its dual. For $\varphi \in \mathcal{S}(\mathbb{R}^d)$, we define its Fourier transform by

$$
\mathcal{F}(\varphi)(p) = \hat{\varphi}(p) \stackrel{\text{def}}{=} \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^n} \varphi(x) e^{-\iota p \cdot x} dx,
$$

while, for $f \in \mathcal{S}'(\mathbb{R}^d)$, it is given via duality.

Throughout the paper we fix a mollifier $\varrho \in \mathcal{S}(\mathbb{R}^2)$ (but the results are independent of it) which is smooth, of mass 1, even and compactly supported in a ball of radius 1. For $\tau > 0$ and $n \in \mathbb{N}$, we set

$$
\varrho_{\tau}(\cdot) \stackrel{\text{def}}{=} \tau \nu_{\tau}^{-1/2} \varrho(\tau^{1/2} R_{\tau}^{-1/2} \cdot), \quad \text{and} \quad \Theta_{\tau}(\cdot) \stackrel{\text{def}}{=} (2\pi \hat{\varrho}_{\tau}(\cdot))^2 = (2\pi \hat{\varrho}(\tau^{-1/2} R_{\tau}^{1/2} \cdot))^2, \quad (1.18)
$$

and define the measure Ξ_n^{τ} on \mathbb{R}^{2n} according to

$$
\Xi_n^{\tau}(\mathrm{d}p_{1:n}) \stackrel{\text{def}}{=} \prod_{i=1}^n \Theta_{\tau}(p_i) \,\mathrm{d}p_{1:n} \,, \qquad p_{1:n} \in \mathbb{R}^{2n} \tag{1.19}
$$

with the convention that, for $\tau = \infty$, $\Theta_{\tau} \equiv 1$ and $\Xi_n = \Xi_n^{\infty}$ is the Lebesgue measure.

For $\tau > 0$ and $\gamma \in \mathbb{R}$, we introduce the τ -dependent (anisotropic) Sobolev spaces with respect to Ξ_n^{τ} , which are given by the completion of $\mathcal{S}(\mathbb{R}^{2n})$ under the (semi-)norm^{[4](#page-11-1)}

$$
||f||_{H^{\gamma}_{\tau}(\mathbb{R}^{2n})}^{2} \stackrel{\text{def}}{=} \int \left(1 + \frac{1}{2} |\sqrt{R_{\tau}} p_{1:n}|^{2} \right)^{\gamma} |\hat{f}(p_{1:n})|^{2} \Xi_{n}^{\tau}(\mathrm{d}p_{1:n}). \tag{1.20}
$$

For $\gamma = 0$, this corresponds to the space $L^2(\mathbb{R}^{2n}) \stackrel{\text{def}}{=} L^2(\mathbb{R}^{2n}, \Xi_n^{\tau})$, whose scalar product will be denoted by $\langle \cdot, \cdot \rangle_{L^2_{\tau}(\mathbb{R}^{2n})}$.

Finally, we will write $a \leq b$ if there exists a constant $C > 0$ independent of any quantity relevant for the result, such that $a \leq Cb$ and $a \approx b$ if $a \leq b$ and $b \leq a$. If we want to highlight the dependence of the constant C on a specific quantity Q, we write instead \leq_{Q} .

1.3 Gaussian white noise and Fock spaces

Let η be a space white noise on \mathbb{R}^2 , i.e. an isonormal Gaussian process on $L^2(\mathbb{R}^2)$ (see [Nuao6, Definition 1.1.1]) and, for $\tau \ge 0$, set $\eta^{\tau} \stackrel{\text{def}}{=} \varrho_{\tau} * \eta$ (with $\eta^{\infty} \equiv \eta$). Then, η^{τ} is a centred Gaussian field with covariance

$$
\mathbb{E}^{\tau}[\eta^{\tau}(\varphi)\eta^{\tau}(\psi)] = \langle \varphi, \psi \rangle_{L_{\tau}^{2}(\mathbb{R}^{2})} = \int_{\mathbb{R}^{2}} \hat{\varphi}(p) \overline{\hat{\psi}(p)} \Xi_{1}^{\tau}(dp), \qquad \varphi, \psi \in L_{\tau}^{2}(\mathbb{R}^{2}).
$$

We will denote its law by \mathbb{P}^{τ} and by $\mathbb{L}^{\theta}(\mathbb{P}^{\tau})$ the space of θ -integrable random variable with respect to \mathbb{P}^{τ} . A random variable $F \in \mathbb{L}^{\theta}(\mathbb{P}^{\tau})$ is said to be *cylinder* if it is of the form $F = f(\eta^{\tau}(\varphi_1), \ldots, \eta^{\tau}(\varphi_n))$ for $f: \mathbb{R}^n \to \mathbb{R}$ smooth and growing at most polynomially at infinity, and $\varphi_1,\ldots,\varphi_n\in \mathcal{S}(\mathbb{R}^2)$. The set of cylinder random variables is dense in $\mathbb{L}^{\theta}(\mathbb{P}^{\tau})$, for $\theta < \infty$.

According to [Nuao6, Theorem 1.1.1], $\mathbb{L}^2(\mathbb{P}^{\tau})$ admits the orthogonal decomposition $\mathbb{L}^2(\eta^{\tau}) =$ $\bigoplus_{n\geq 0} \mathcal{H}_n^{\tau}$, where \mathcal{H}_n^{τ} is the *n-th homogeneous Wiener chaos*, i.e. the closure in $\mathbb{L}^2(\mathbb{P}^{\tau})$ of $\text{Span}_{\mathcal{H}}\{H_n(\eta^\tau(h)) \,|\, h \in \mathcal{S}(\mathbb{R}^2), \|h\|_{L^2_\tau(\mathbb{R}^2)} = 1\}$ and H_n is the *n*-th Hermite polynomial. Further, [Nuao6, Theorem 1.1.2] ensures the existence of a canonical isometry I^{τ} onto $\mathbb{L}^2(\mathbb{P}^{\tau})$ which is the unique extension of the map that, for every $n \ge 0$, assigns $h^{\otimes n}$ to $n! H_n(\eta^\tau(h)) \in \mathcal{H}_n^\tau$ for $h \in \mathcal{S}(\mathbb{R}^2)$ with $L^2(\mathbb{R}^2)$ -norm equal to 1. The space on which I^{τ} is defined is the Fock space ΓL^2_τ , and is given by the closure of $\oplus_{n\geq 0} \Gamma_n L^2_\tau$, where $\Gamma_n L^2_\tau$ is the subset of $L^2_\tau(\mathbb{R}^{2n})$ made of functions which are symmetric with respect to permutations of their variables, endowed with the norm

$$
||f||_{\tau}^{2} = ||f||_{\Gamma L_{\tau}^{2}}^{2} \stackrel{\text{def}}{=} \sum_{n\geq 0} n! ||f_{n}||_{L_{\tau}^{2}(\mathbb{R}^{2n})}^{2}, \qquad (1.21)
$$

for $f = (f_n)_{n \geq 0} \in \bigoplus_{n \geq 0} \Gamma_n L_\tau^2$. Note that, by construction, the restriction I_n^{τ} of I^{τ} to $\Gamma_n L_\tau^2$ is an isometry with values in \mathcal{H}_n^{τ} and, again by [Nuao6, Theorem 1.1.2], for any $F \in \mathbb{L}^2(\mathbb{P}^{\tau})$ there exists $f = (f_n)_n \in \Gamma L_\tau^2$ such that $F = \sum_n I_n(f_n)$ and $\mathbb{E}^\tau[F^2] = ||f||_\tau^2$.

The bulk of our analysis will focus on various operators acting on (subspaces of) $\mathbb{L}^2(\mathbb{P}^{\tau})$. In view of the isometry I^{τ} , we will abuse notation and identify them with the corresponding operator acting on ΓL_{τ}^2 . An important notion is that of *diagonal* operator that we now define.

⁴As the zero-set of Θ_{τ} is of empty interior since Θ_{τ} is analytic, these are actually norms.

INTRODUCTION **13**

Definition 1.5 An operator \mathcal{D} on ΓL^2_τ is said to be *diagonal* if there exists a family of measurable kernels $\mathbf{d} = (\mathbf{d}_n)_{n \geq 0}$ such that for all $n \geq 0$ and $\varphi \in \Gamma_n L^2_\tau \cap \mathcal{S}(\mathbb{R}^{2n})$, it holds $\mathcal{FD}\varphi(p_{1:n}) =$ $\mathbf{d}_n(p_{1:n})\hat{\varphi}(p_{1:n})$ for (Lebesgue-almost-) every $p_{1:n} \in \mathbb{R}^{2n}$. If $\mathfrak{D}^{(1)}$ and $\mathfrak{D}^{(2)}$ are diagonal, we write $\mathfrak{D}^{(1)} \leq \mathfrak{D}^{(2)}$ if their kernels satisfy $\mathbf{d}^{(1)} \leq \mathbf{d}^{(2)}$ and say that $\mathfrak{D}^{(1)}$ is positive if $\mathfrak{D}^{(1)} \geq 0$. Given a diagonal operator \mathfrak{D} with kernel $\mathbf{d} = (\mathbf{d}_n)_{n \geq 0}$ and a measurable scalar function f, we define $f(\mathfrak{D})$ to be the diagonal operator whose kernel is $f(\mathbf{d}) = (f \circ \mathbf{d}_n)_{n \geq 0}$.

Definition 1.6 We define the number operator N to be the diagonal operator acting on ψ = $(\psi_n)_n \in \Gamma L^2_\tau$ such that $\|\psi_n\|_\tau$ decays faster than any polynomial in n, as $\mathcal{N}\psi_n \stackrel{\text{def}}{=} n\psi_n$.

In order to discuss the domain and well-posedness of the operators we will encounter, it is convenient to introduce, for $\gamma \in \mathbb{R}$, the Sobolev space corresponding to ΓL^2_τ , which we denote by ΓH^{γ} and whose norm is the same as that in [\(1.21\)](#page-11-2) but with $\|\cdot\|_{H^{\gamma}(\mathbb{R}^{2n})}^2$ replacing $\|\cdot\|_{L^2(\mathbb{R}^{2n})}^2$.

To conclude this preliminary section, let us clarify the relation between $\mathbb{L}^{\theta}(\mathbb{P}^{\tau})$ and $\mathbb{L}^{\theta}(\mathbb{P}) =$ $\mathbb{L}^{\theta}(\mathbb{P}^{\infty})$, for $\theta \in [1,\infty)$. Recall that the operator $\iota^{\tau} \colon \mathbb{L}^{\theta}(\mathbb{P}^{\tau}) \to \mathbb{L}^{\theta}(\mathbb{P})$ which maps the functional $\eta^{\tau} \mapsto F(\eta^{\tau})$ to the functional $\eta \mapsto F(\varrho_{\tau} * \eta)$, is an isometric embedding. In the next basic lemma we show that ι^{τ} is an isometry which admits an inverse j^{τ} .

Lemma 1.7 *For every* $\theta \in [1, +\infty)$ *, the map* $\iota^{\tau} : \mathbb{L}^{\theta}(\mathbb{P}^{\tau}) \to \mathbb{L}^{\theta}(\mathbb{P})$ *is an isometry which admits an inverse, that we set to be* $j^{\tau} \stackrel{\text{def}}{=} (i^{\tau})^{-1}$ *. Furthermore, for* $\theta = 2$ *,* i^{τ} *and* j^{τ} *are respectively the operators from* ΓL^2_τ to $\Gamma L^2 = \Gamma L^2_\infty$ and viceversa, that for every $\varphi \in \Gamma_n L^2_\tau$, $\psi \in \Gamma_n L^2$, $n \in \mathbb{N}$, *satisfy*

$$
\mathcal{F}(\iota^{\tau}(\varphi))(p_{1:n}) = \hat{\varphi}(p_{1:n}) \prod_{i=1}^{n} \Theta_{\tau}(p_i)^{1/2}, \qquad \mathcal{F}(j^{\tau}(\psi))(p_{1:n}) = \frac{\hat{\psi}(p_{1:n})}{\prod_{i=1}^{n} \Theta_{\tau}(p_i)^{1/2}}.
$$
 (1.22)

Proof. As ι^{τ} is an isometric embedding, to show that it is an isometry, it suffices to prove that it is surjective. To do so, we will build its inverse, i.e. a map $j^{\tau} \colon \mathbb{L}^{\theta}(\mathbb{P}) \to \mathbb{L}^{\theta}(\mathbb{P}^{\tau})$ such that $\iota^{\tau} \circ j^{\tau} =$ Id. First, let $(\varrho_{\tau}*)^{-1} : L^2(\mathbb{R}^2) \to L^2_{\tau}(\mathbb{R}^2)$ be given as

$$
\mathcal{F}((\varrho_\tau)^{-1}f)(p) \stackrel{\text{def}}{=} \frac{\hat{f}(p)}{\Theta_\tau(p)^{1/2}},
$$

which is well defined, as ρ is compactly supported, its Fourier transform is analytic and thus the set of points at which Θ_{τ} vanishes has zero Lebesgue measure. Furthermore, (ϱ_{τ}) and $(\varrho_{\tau})^{-1}$ are both isometries respectively from $L^2(\mathbb{R}^2)$ to $L^2(\mathbb{R}^2)$ and viceversa, and are inverses of one another. Now, note that for $\varphi \in L^2(\mathbb{R}^2)$, $\eta(\varphi) = \eta^{\tau}((\varrho_{\tau}*)^{-1}\varphi)$, that is to say $\eta = \eta^{\tau} \circ (\varrho_{\tau}*)^{-1}$. Hence, the map j^{τ} : $\mathbb{L}^{\theta}(\mathbb{P}) \ni F(\eta) \mapsto F(\eta^{\tau} \circ (\varrho_{\tau}*)^{-1}) \in \mathbb{L}^{\theta}(\mathbb{P}^{\tau})$ is also well-defined, is an isometry and it is the inverse of ι^{τ} .

At last, [\(1.22\)](#page-12-1) can be immediately deduced by the definitions of $(\varrho_{\tau})^{-1}$ and j^{τ} . \Box

2 Preliminaries, novelties and roadmap

In this section, we present some preliminary results concerning the solution to the regularised scaled SBE in [\(1.9\)](#page-6-0). The statements are essential, since they represent the foundations on which our analysis relies, but basic, in that their proof uses classical techniques from PDE theory and functional analysis. This is the reason why we preferred to postpone most of the arguments (some of which quite involved) to Appendices [A](#page-51-0) and [B,](#page-61-0) and could be skipped at first read.

After studying basic properties of the solution (Section [2.1\)](#page-13-1), its semigroup and generator (Section [2.2\)](#page-19-0), in Section [2.3](#page-21-0) we provide a more detailed heuristic as to how the proof of the main result works and what are the novel ideas behind it.

Throughout this section, the regularisation parameter $\tau > 0$ is *fixed* and, unless otherwise specified, in the estimates herein no uniformity in its value is claimed (or true).

2.1 The regularised stochastic Burgers equation

To rigorously discuss the equation of interest, let $(0, \mathcal{F}, P)$ be a probability space supporting a couple of independent spatial white noise η on \mathbb{R}^2 and 2-dimensional space-time white noise $\vec{\xi} = (\xi_1, \xi_2)$ on $\mathbb{R} \times \mathbb{R}^2$, whose laws are respectively denoted by \mathbb{P} and P , and $P \stackrel{\text{def}}{=} \mathbb{P} \otimes P$. Let $(\mathcal{F}_t)_{t\geq0}$ be the filtration in which \mathcal{F}_t is the (completed) σ -algebra generated by η and $\vec{\xi}|_{(-\infty,t]}$. Let $\overline{\varrho} \in \mathcal{S}(\mathbb{R}^2)$ be the mollifier (i.e. smooth, even-symmetric, compactly supported in a ball of radius 1 and with total mass equal to 1) introduced above and, for $\tau > 0$, ρ_{τ} be as in [\(1.18\)](#page-10-0). Let $\eta^{\tau} \stackrel{\text{def}}{=} \eta * \varrho_{\tau}$ and $\vec{\xi}^{\tau} \stackrel{\text{def}}{=} \vec{\xi} * \varrho_{\tau} = (\xi_1 * \varrho_{\tau}, \xi_2 * \varrho_{\tau})$ the rescaled white noises, whose joint law is $\mathbf{P}^{\tau} \stackrel{\text{def}}{=} \mathbb{P}^{\tau} \otimes P^{\tau}.$

Now, we want to study the SPDE in [\(1.9\)](#page-6-0) given by

$$
\partial_t u^\tau = \frac{1}{2} \nabla \cdot R_\tau \nabla u^\tau + \lambda \nu_\tau^{3/4} \mathcal{N}^\tau [u^\tau] + \nabla \cdot \sqrt{R_\tau} \vec{\xi}^\tau, \qquad u_0^\tau = \eta^\tau, \tag{2.1}
$$

where $u^{\tau} = u^{\tau}(t, x) = u_t^{\tau}(x)$ for $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^2$ and \mathcal{N}^{τ} is given as in [\(1.3\)](#page-3-3) but with ϱ_{τ} in place of ϱ , and R_{τ} , ν_{τ} as in [\(1.16\)](#page-10-1). To do so, we need to introduce a space in which u^{τ} lives and a suitable notion of solution. The former is $C_{\text{loc}}^{\beta}(\mathbb{R}_+, \mathscr{C}^{\alpha}(p_a))$ for $\beta, \alpha, a > 0$, i.e. the space of locally β-Hölder continuous functions in time with values in the weighted Besov-Hölder space $\mathscr{C}^{\alpha}(\mathfrak{p}_a)$ of regularity α . The precise definition of $\mathscr{C}^{\alpha}(\mathfrak{p}_a)$ will be given in Appendix [A.1](#page-51-1) but, for now, the reader can think of the space of α -Hölder continuous functions whose norm restricted to a ball of radius x grows at most as $p_{-a}(x) = (1+|x|)^a$ (in particular this is true for all derivatives of order smaller than $|\alpha|$). Concerning the notion of solution, it will be convenient for us to work with mild solutions.

Definition 2.1 We say that u^{τ} is a mild solution of [\(2.1\)](#page-13-2) driven by $(\eta^{\tau}, \vec{\xi}^{\tau})$ if u^{τ} is $(\mathcal{F}_t)_{t\geq 0}$ adapted, it belongs to $C_{loc}(\mathbb{R}_+, \mathcal{C}^{\alpha}(\mathbf{p}_a))$ **P**^{*r*}-a.s., and for every $t \geq 0$ it holds

$$
u_t^{\tau} = K_t^{\tau} \eta^{\tau} + \lambda \nu_{\tau}^{3/4} \int_0^t K_{t-s}^{\tau} \mathcal{N}^{\tau} [u_s^{\tau}] ds + \int_0^t K_{t-s}^{\tau} \nabla \cdot R_{\tau} \vec{\xi}^{\tau} (ds), \qquad (2.2)
$$

where the last summand is to be interpreted as a stochastic convolution in the sense of Walsh [\[Wal86\]](#page-73-5), and $K_t^{\tau} \stackrel{\text{def}}{=} e^{\frac{t}{2} \nabla \cdot R_{\tau} \cdot \nabla}$ is the heat semigroup associated to $\frac{1}{2} \nabla \cdot R_{\tau} \nabla$.

Remark 2.2 In previous works (and especially [\[CGT24\]](#page-70-3)), the notion of solution considered is that of analytically weak solution, which is obtained by multiplying both sides of (2.1) by a function φ , integrating in time and space, and requiring the equality to hold for a sufficiently large class of φ 's. In general, the notion of mild solution is stronger than that of weak but, given the regularity of the noise (and the initial condition) for τ fixed, the two can be easily shown to be equivalent arguing as in [\[DPZ08,](#page-70-12) Theorem 5.4].

The next proposition summarises the main properties of u^{τ} we are concerned with, i.e. global well-posedness, Markovianity and stationarity. See Appendix [A.3,](#page-55-0) for the proof.

Proposition 2.3 *Let* $\tau > 0$ *be fixed. For any* $\alpha > 0$ *and* $a > 0$ *, the regularised stochastic* Burgers equation [\(2.1\)](#page-13-2) driven by $(\eta^{\tau}, \vec{\xi}^{\tau})$ admits a unique mild solution whose paths belong *to* $C_{\text{loc}}(\mathbb{R}_+,\mathcal{C}^{\alpha}(\mathbf{p}_a))$ *(actually to* $C_{\text{loc}}^{\beta}(\mathbb{R}_+,\mathcal{C}^{\alpha}(\mathbf{p}_a))$ *for any* $\beta \in [0,1/2)$ *) and path-by-path uniqueness holds. Moreover, for every* $\alpha, a, p, T > 0$ *and* $\beta \in [0, 1/2)$ *, there exists a positive constant* $C = C(\tau, \alpha, \beta, a, p, T) < \infty$ *such that*

$$
\mathbf{E}^{\tau}\Big[\|u^{\tau}\|_{C^{\beta}([0,T],\mathcal{C}^{\alpha}(\mathbf{p}_a))}^p\Big] \leq C\,. \tag{2.3}
$$

At last, $(u_t^\tau)_{t\geq 0}$ defines a $(\mathcal{F}_t)_{t\geq 0}$ -Markov process on $\mathscr{C}^\alpha(\mathrm{p}_a)$, which is stationary, that is, for any $t \geq 0$, $u_t^{\tau} \stackrel{law}{=} \eta^{\tau}$; and skew-reversible, in the sense that, for any $T \geq 0$, $\{u_{T-t}^{\tau}: t \in [0,T]\} \stackrel{law}{=}$ $\{v_t^\tau : t \in [0,T]\}$ where v^τ is the unique mild solution to [\(2.1\)](#page-13-2) driven by $(\tilde{\eta}^\tau, \tilde{\xi}^\tau) \stackrel{law}{=} (\eta^\tau, \tilde{\xi}^\tau)$ with λ *replaced by* $-\lambda$ *.*

In light of the previous proposition, we not only know that (2.1) admits a unique mild solution u^{τ} driven by $(\eta^{\tau}, \vec{\xi}^{\tau})$ but also that u^{τ} is smooth (and stationary). Therefore, we can define its two-point correlation function $S^{\tau} \colon \mathbb{R}_+ \times \mathbb{R}^2 \to \mathbb{R}$ according to

$$
S_t^{\tau}(x) = \mathbf{E}^{\tau}[u_t^{\tau}(x) u_0^{\tau}(0)], \qquad (t, x) \in \mathbb{R}_+ \times \mathbb{R}^2.
$$
 (2.4)

Note that, since u^{τ} is centred, S^{τ} is indeed the covariance of $u_t^{\tau}(x)$ and $u_0^{\tau}(0)$. Further, it is smooth and its derivatives, which, for $\mathbf{k} \in \mathbb{N}^2$, will be denoted by $\partial_{\mathbf{k}}^{|\mathbf{k}|} S^{\tau} = \partial_1^{k_1} \partial_2^{k_2} S^{\tau}$ (with the convention that if $k_i = 0$ then there is no derivative in the *i*-th direction), can be obtained by commuting the expectation and the derivative operator (this is allowed as, locally, the sup-norm of the derivatives of u_t^{τ} have finite moments of all orders since, in law, u_t^{τ} equals η^{τ} by stationarity).

In the next proposition, we continue our analysis of u^{τ} and show that its correlations, encoded via S^{τ} , and their spatial derivatives decay in space faster than any polynomial.

Proposition 2.4 *Let* S^{τ} *be defined according to* [\(2.4\)](#page-14-1)*. Then, for every* $T, \gamma > 0$ *and* $\mathbf{k} \in \mathbb{N}^2$ *, there exists a positive constant* $C = C(\tau, \gamma, T, \mathbf{k}) < \infty$ *such that for every* $t \leq T$ *and* $x \in \mathbb{R}^2$ *.*

$$
|\partial_{\mathbf{k}}^{|\mathbf{k}|} S_t^{\tau}(x) | \leqslant C |x|^{-\gamma}.
$$

Similarly, for $\mathbf{k} \in \mathbb{N}^2$, we have

$$
\left| \mathbf{E}^{\tau} [\partial_{\mathbf{k}}^{\vert \mathbf{k} \vert} \mathcal{M}^{\tau} [u_t^{\tau}] (x) u_0^{\tau} (0)] \right| \vee \left| \mathbf{E}^{\tau} [\partial_{\mathbf{k}}^{\vert \mathbf{k} \vert} \mathcal{M}^{\tau} [u_t^{\tau}] (x) \mathcal{M}^{\tau} [u_0^{\tau}] (0)] \right| \leqslant C |x|^{-\gamma}, \tag{2.6}
$$

where ^M^τ [f] *is given by*

$$
\mathcal{M}^{\tau}[f] \stackrel{\text{def}}{=} \varrho_{\tau}^{*2} * f^2 - \mathbb{E}^{\tau}[\varrho_{\tau}^{*2} * f^2(0)]. \tag{2.7}
$$

Proof. Throughout the proof, since τ is fixed, without loss of generality, we take it to be equal to 1 and remove it from the notation. Note that this gives $\nu_{\tau} = 1$ and $R_{\tau} = Id$, the identity matrix. The result for general τ can then be deduced by scaling. Further, we will denote by $C > 0$ a constant only depending on the stated quantities, and whose value may change from line to line.

Let us begin with a few preliminary definitions and observations. Let $\varphi : \mathbb{R}^2 \to [0, 1]$ be a compactly supported smooth function such that $\varphi \equiv 1$ on $B_0(1/8) \subset \mathbb{R}^2$, i.e. the ball centred at the origin of radius 1/8, and $\varphi \equiv 0$ on $B_0(1/4)^c$, and set, for $x_0 \in \mathbb{R}^2$, $|x_0| > 4$, $\varphi_{x_0}(y) = \varphi(y/|x_0|)$. Let η' be a space white noise on \mathbb{R}^2 independent of η and define $\tilde{\eta}$ according to

$$
\tilde{\eta} = \sqrt{\varphi_{x_0}} \eta' + \sqrt{1 - \varphi_{x_0}} \eta.
$$

Note that $\tilde{\eta}$ is distributed according to a spatial white noise on \mathbb{R}^2 and, setting as usual $\tilde{\eta}^1 \stackrel{\text{def}}{=} \tilde{\eta} * \varrho$, we have that $\tilde{\eta}^1$ coincides with η^1 on $B_0(R)^c$ where $R \stackrel{\text{def}}{=} |x_0|/4 + 1$ (because ϱ has support in the ball of radius 1), and $\tilde{\eta}^1|_{B_0(|x_0|/8-1)}$ is independent of $\eta^1|_{B_0(|x_0|/8-1)}$.

Let \tilde{u} be the unique mild solution of [\(2.1\)](#page-13-2) driven by $(\tilde{\eta}^1, \vec{\xi})$ and u that driven by $(\eta^1, \vec{\xi}^1)$, so that $\tilde{u} \equiv u$. Set $v \stackrel{\text{def}}{=} u - \tilde{u}$ and $w \stackrel{\text{def}}{=} u + \tilde{u}$ and note that they satisfy [\(A.19\)](#page-57-0) with $X = K_t v_0 = K_t(\eta^1 - \tilde{\eta}^1)$. Then, thanks to the previous observations, the **k**-th derivative, for $\mathbf{k} \in \mathbb{N}^2$, of the two-point correlation function S at x_0 can be written as

$$
\partial_{\mathbf{k}}^{|\mathbf{k}|} S_t(x_0) = \mathbf{E}[\partial_{\mathbf{k}}^{|\mathbf{k}|} \tilde{u}_t(x_0) u_0(0)] + \mathbf{E}[\partial_{\mathbf{k}}^{|\mathbf{k}|} v_t(x_0) u_0(0)] = \mathbf{E}[\partial_{\mathbf{k}}^{|\mathbf{k}|} v_t(x_0) u_0(0)] \tag{2.8}
$$

where we used that $\partial_{\mathbf{k}}^{|\mathbf{k}|} \tilde{u}$ is independent of $\partial_{\mathbf{k}}^{|\mathbf{k}|} \eta^1|_{B_0(|x_0|/8-1)}$.

At this point, we need to control $\partial_{\bf k}^{|{\bf k}|} v_t(x_0)$. To do so, we will first take advantage of [\(2.3\)](#page-14-2) to get rid of the randomness of w, η^1 and $\tilde{\eta}^1$ so to be able to estimate it with the solution to deterministic PDEs and ultimately prove that these decay sufficiently fast. For $\delta > 0$ and R as above, define the event

$$
E = \left\{ \sup_{(t,x)\in[0,T]\times\mathbb{R}^2} \left| \mathbf{p}_{\delta}(x) \partial_{\mathbf{k}}^{|\mathbf{k}|} w_t(x) \right| \vee \sup_{x\in B_0(R)} |\partial_{\mathbf{k}}^{|\mathbf{k}|} v_0(x)| \le R^{\delta} \right\}
$$
(2.9)

and notice that thanks to [\(2.3\)](#page-14-2) and [\(A.9\)](#page-53-1) (the latter with $t = 0$), the weighted norm of w and the local sup-norm of $v_0 = \eta^1 - \tilde{\eta}^1$ have finite moments of all orders, which immediately implies that $P(E^c) \lesssim R^{-\nu}$ for any given $\nu > 0$. Therefore,

$$
|\partial_{\mathbf{k}}^{|\mathbf{k}|} S_t(x_0)| \le \mathbf{E}[|\partial_{\mathbf{k}}^{|\mathbf{k}|} v_t(x_0)u_0(0)|\mathbb{1}_{E^c}] + \mathbf{E}[|\partial_{\mathbf{k}}^{|\mathbf{k}|} v_t(x_0)u_0(0)|\mathbb{1}_E] \n\lesssim R^{-\nu} \sqrt{\mathbf{E}[|\partial_{\mathbf{k}}^{|\mathbf{k}|} v_t(x_0)|^2 |u_0(0)|^2]} + \sqrt{\mathbf{E}[|\partial_{\mathbf{k}}^{|\mathbf{k}|} v_t(x_0)|^2 \mathbb{1}_E]\mathbf{E}[|u_0(0)|^2]} \n\lesssim R^{-\nu} + \sqrt{\mathbf{E}[|\partial_{\mathbf{k}}^{|\mathbf{k}|} v_t(x_0)|^2 \mathbb{1}_E]}
$$
\n(2.10)

where, in the first step we applied Cauchy-Schwarz, while, in the last, again Cauchy-Schwarz on the first expectation and then stationarity and finiteness of moments of $u_0(0) = \eta^1(0)$ and of

 $\partial_{\bf k}^{|{\bf k}|}\eta^1(0)$, to conclude that the bound is uniform in t and x_0 . Thus, we are left to show that, on $E, |\partial_{\mathbf{k}}^{\mathbf{|k|}} v_t(x_0)|$ can be estimated by the right hand side of [\(2.5\)](#page-14-3).

Recall the definition of $\mathcal{N}^1[\cdot, \cdot]$ in [\(A.20\)](#page-57-1), that v solves [\(A.19\)](#page-57-0) and that, by construction, v_0 is compactly supported in a ball of radius R. Then, on E, for every $(t, x) \in [0, T] \times \mathbb{R}^2$, we have

$$
|\partial_{\mathbf{k}}^{|\mathbf{k}|} v_t(x)| \le K_t |\partial_{\mathbf{k}}^{|\mathbf{k}|} v_0|(x) + \int_0^t K_{t-s} |\partial_{\mathbf{k}}^{|\mathbf{k}|} \mathcal{N}^1[v_s, w_s]|(x) ds
$$
\n
$$
\le K_t (R^\delta \mathbb{1}_{B_0(R)})(x_0) + R^\delta \int_0^t K_{t-s} * |\partial_{\mathbf{k}}^{|\mathbf{k}|} (\mathfrak{w} \cdot \nabla) \varrho^{*2}| * (|v_s| \mathfrak{p}_{-\delta})(x) ds.
$$
\n(2.11)

Let us consider first the case of $|\mathbf{k}| = 0$. As ϱ^{*2} has compact support and $p_{-\delta}$ is a weight, [\(A.1\)](#page-51-3) ensures the existence of a constant $c_{\delta} > 0$ such that the convolution in the second summand above can be estimated by

$$
\begin{aligned} |(\mathfrak{e}_1 \cdot \nabla)\varrho^{*2}| * (|v_s| \mathfrak{p}_{-\delta})(z) &= \int |\partial_1 \varrho^{*2}|(z-y)|v_s(y)|\mathfrak{p}_{-\delta}(y)dy \\ &\le c_\delta p_{-\delta}(z) \int |\partial_1 \varrho^{*2}|(z-y)|v_s(y)|dy \\ &= p_{-\delta}(z) \int \Phi(z-y)[|v_s(y)| - |v_s(z)|]dy + m_\Phi p_{-\delta}(z)|v_s(z)| \end{aligned} \tag{2.12}
$$

where $\Phi \stackrel{\text{def}}{=} c_{\delta} |\partial_1 \varrho^{*2}|$ (which is compactly supported, positive and even) and m_{Φ} is its total mass. Therefore, [\(2.11\)](#page-16-0) and [\(2.12\)](#page-16-1) imply that, uniformly in $(t, x) \in [0, T] \times \mathbb{R}^2$, v is bounded by the deterministic function $Z = Z_t(x)$ which solves

$$
\partial_t Z_t(x) = \left(\frac{1}{2}\Delta + L_{\Phi, R^{\delta}p - \delta}\right) Z_t(x) + m_{\Phi} R^{\delta} p_{-\delta}(x) Z_t(x), \quad Z_0(x) = R^{\delta} 1_{B_0(R)}(x), \quad (2.13)
$$

for $L_{\Phi,R\delta p-\delta}$ the operator acting on functions $u: \mathbb{R}^2 \to \mathbb{R}$ as

$$
L_{\Phi, R^{\delta} p_{-\delta}} u(x) \stackrel{\text{def}}{=} R^{\delta} p_{-\delta}(x) \int \Phi(x - y) [u(y) - u(x)] dy.
$$
 (2.14)

As detailed in Appendix [A.4,](#page-60-0) the advantage of singling out the operator $L_{\Phi,R\delta p_{-\delta}}$ is that $\frac{1}{2}\Delta$ + $L_{\Phi, R^{\delta} p_{-\delta}}$ is the generator of a Markov process on \mathbb{R}^2 , and this allows to derive an explicit expression for the solution to [\(2.13\)](#page-16-2) via a Feynman-Kac formula, which in turn implies the (exponential) decay. Indeed, it is not hard to see that Lemma [A.6](#page-60-1) applies to Z with the choices of Φ as above, $f = R^{\delta}p_{-\delta}$, $g = m_{\Phi}R^{\delta}p_{-\delta}$ and $h = R^{\delta}1_{B_0(R)}$. Hence, recalling that $R = |x_0|/4 + 1$ (so that, since h is supported in a ball of radius $r_h = R$, we have $|x_0| \ge 2r_h$ as $|x_0| > 4$, as required by Lemma [A.6\)](#page-60-1), we deduce that on the event E

$$
|v_t(x_0)| \le Z_t(x_0) \lesssim e^{-C|x_0|}
$$

which, once plugged into [\(2.10\)](#page-15-0) with $|\mathbf{k}| = 0$, implies [\(2.5\)](#page-14-3).

For $|\mathbf{k}| \geq 0$, we go back to [\(2.11\)](#page-16-0) and upper-bound |v| at the right hand side by Z. Then, arguing as in [\(2.12\)](#page-16-1), we deduce that $|\partial_{\bf k}^{|{\bf k}|}v|$ is upper bounded by the solution $Z^{\bf k}$ of the nonhomogeneous deterministic heat equation given by

$$
\partial_t Z_t^{\mathbf{k}}(x) = \frac{1}{2} \Delta Z_t^{\mathbf{k}}(x) + c_\delta R^\delta p_{-\delta}(x) \Big(|\partial_{\mathbf{k}}^{|{\mathbf{k}}|} \partial_1 \varrho^{*2} | * Z_t \Big)(x) \,, \qquad Z_0(x) = R^\delta 1\!\!1_{B_0(R)}(x) \,.
$$

Since the initial condition is compactly supported and, by Lemma [A.6,](#page-60-1) the second summand decays exponentially (outside the ball of radius $2R = |x_0|/2 + 2$), it can be easily checked that also $Z_t^{\mathbf{k}}$ $t^{k}(x_0)$ decays exponentially in $|x_0|$ from which [\(2.5\)](#page-14-3) follows at once.

The proof of [\(2.6\)](#page-14-4) requires only minor changes. For instance, arguing as in [\(2.8\)](#page-15-1), the first term at the right hand side can be rewritten as

$$
\mathbf{E}\Big[\partial_{\mathbf{k}}^{\vert\mathbf{k}\vert}(\mathcal{M}^1[u_t](x_0)-\mathcal{M}^1[\tilde{u}_t](x_0))u_0(0)\Big].
$$

On the other hand,

$$
\partial_{\mathbf{k}}^{|\mathbf{k}|}(\mathcal{M}^{1}[u_{t}](x_{0}) - \mathcal{M}^{1}[\tilde{u}_{t}](x_{0})) = \partial_{\mathbf{k}}^{|\mathbf{k}|} \varrho^{*2} * (u_{t}^{2} - \tilde{u}_{t}^{2})(x_{0}) = \partial_{\mathbf{k}}^{|\mathbf{k}|} \varrho^{*2} * (v_{t}w_{t})(x_{0}).
$$

At this point, one proceeds exactly as above, removing $|w|$ as in [\(2.10\)](#page-15-0) and then bounding $|v|$ by
Z. We omit obvious details Z. We omit obvious details.

A first consequence of Proposition [2.4](#page-14-5) is that the two-point correlation function S^{τ} has total mass equal to 1, i.e.

$$
\int S_t^{\tau}(x) dx = \int S_0^{\tau}(x) dx = \int \varrho_{\tau}^{*2}(x) dx = 1
$$
\n(2.15)

independently of $t \geq 0$. To see this, note first that, since u^{τ} is a (spatially smooth) mild solution to [\(2.1\)](#page-13-2), then it is standard to show that for every $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^2$ we have

$$
u_t^\tau(x) = \eta^\tau(x) + \frac{1}{2} \int_0^t \nabla \cdot R_\tau \nabla u_s^\tau(x) \, ds + \lambda \nu_\tau^{3/4} \int_0^t \mathcal{N}^\tau [u_s^\tau](x) \, ds + \int_0^t \nabla \cdot \sqrt{R_\tau} \vec{\xi}^\tau(\mathrm{d}s, x) \tag{2.16}
$$

which, by multiplying both sides by $u_0^{\tau}(0) = \eta^{\tau}(0)$ and taking expectation, gives

$$
S_t^{\tau}(x) - S_0^{\tau}(x) = \frac{1}{2} \int_0^t \nabla \cdot R_{\tau} \nabla S_s^{\tau}(x) \, ds + \lambda \nu_{\tau}^{3/4} \int_0^t \mathbf{E} \Big[\partial_1 \mathcal{M}^{\tau}[u_s^{\tau}](x) u_0^{\tau}(0) \Big] ds \,, \quad (2.17)
$$

where we used that the initial condition and the driving noise are independent. Integrating with respect to x , all terms in the right-hand side vanish, as follows by integration by parts, which is allowed by the decay properties in Proposition [2.4.](#page-14-5) A similar argument delivers the second important consequence, namely a Green-Kubo formula for the diffusivity matrix D^{τ} defined as in [\(1.4\)](#page-4-3) but with u replaced by the rescaled process u^{τ} in [\(1.8\)](#page-5-2).

Proposition 2.5 Let S^{τ} be the two-point correlation function in [\(2.4\)](#page-14-1). Then, for every $t \geq 0$, *the diffusivity matrix* D^{τ} *in* [\(1.4\)](#page-4-3) *with* S^{τ} *in place of S is well-defined and admits the following Green-Kubo representation*

$$
D_{1,1}^{\tau}(t) = \nu_{\tau} + \frac{2\lambda^2 \nu_{\tau}^{3/2}}{t} \int_0^t \int_0^s \int_{\mathbb{R}^2} \mathbf{E} \left[\mathcal{M}^{\tau}[u_{r}^{\tau}](x) \mathcal{M}^{\tau}[u_{0}^{\tau}](0) \right] dx dr ds , \qquad (2.18)
$$

where M^{τ} *is defined in* [\(2.7\)](#page-15-2)*, while* $D_{2,2}^{\tau}(t) = 1$ *and* $D_{1,2}^{\tau}(t) = D_{2,1}^{\tau}(t) = 0$ *.*

In previous works [\[GHS24,](#page-71-10) [CET23a\]](#page-69-14), the right hand side of [\(2.18\)](#page-18-0) (or the corresponding one for the AKPZ equation) with $\tau = \nu_{\tau} = 1$ was adopted as *definition* of bulk diffusion coefficient, but the equivalence with an expression similar to that in [\(1.4\)](#page-4-3) was only argued heuristically.

Proof. First, by $(2,5)$, the integral at the right hand side of (1.4) is absolutely convergent and thus the diffusion matrix D^{τ} is well-defined. On the other hand, the proof of [\(2.18\)](#page-18-0) is essentially the same as in [\[GHS24,](#page-71-10) App. B], except that the authors consider the bulk diffusion coefficient, which is a scalar quantity, and the argument therein is formal, since it involves an unjustified integration by parts, which holds in view of the decay of correlations shown in Proposition [2.4.](#page-14-5)

Start from [\(2.17\)](#page-17-0), multiply by x_ix_j/t , for $i, j \in \{1, 2\}$, and integrate with respect to x, to obtain

$$
D_{i,j}^{\tau}(t) = \frac{1}{2t} \int_0^t \int_{\mathbb{R}^2} x_i x_j \nabla \cdot R_{\tau} \nabla S_s^{\tau}(x) dx ds + \frac{\lambda \nu_{\tau}^{3/4}}{t} \int_0^t \int_{\mathbb{R}^2} x_i x_j \mathbf{E}^{\tau} \left[\partial_1 \mathcal{M}^{\tau} [u_s^{\tau}] (x) (u_0^{\tau}(0) - u_s^{\tau}(0)) \right] dx ds
$$
\n(2.19)

where we used the fact that $\mathcal{M}^{\tau}[u_s]$ is quadratic while $u_s^{\tau}(0)$ is linear in u^{τ} , and that, for every $s \geq 0$, $u_s^{\tau} \stackrel{\text{law}}{=} \eta^{\tau}$ with η^{τ} Gaussian, so that $\mathbf{E}[\mathcal{M}^{\tau}[u_s^{\tau}](x)u_s^{\tau}(0)] = \mathbb{E}[\mathcal{M}^{\tau}[\eta^{\tau}](x)\eta^{\tau}(0)] = 0$. For the first term at the right hand side of [\(2.19\)](#page-18-1), we apply integration by parts twice and use that $\nabla \cdot R_{\tau} \nabla x_i x_j = 2\nu_{\tau} \mathbb{1}_{i=j=1} + 2 \mathbb{1}_{i=j=2}$ together with [\(2.15\)](#page-17-1). As for the second, we use translation invariance, the tower property and stationarity to write

$$
\mathbf{E}^{\tau}\Big[\partial_1\mathcal{M}^{\tau}[u_s^{\tau}](x)(u_0^{\tau}(0) - u_s^{\tau}(0))\Big] = \mathbb{E}^{\tau}\Big[\partial_1\mathcal{M}^{\tau}[\tilde{\eta}^{\tau}](0)\mathbf{E}^{\tau}_{\tilde{\eta}^{\tau}}[\tilde{u}_s^{\tau}(x) - \tilde{u}_0^{\tau}(x)]\Big]
$$

where $\tilde{\eta}^{\tau} \stackrel{\text{def}}{=} u_s^{\tau} (= \tilde{u}_0^{\tau})$ is a regularised spatial white noise and \tilde{u} is the time-reversed process, i.e. $\tilde{u}_r \stackrel{\text{def}}{=} u_{s-r}$. Thanks to the skew-reversibility of u stated in Proposition [2.3,](#page-14-0) in law, \tilde{u} satisfies the same equation as [\(2.16\)](#page-17-2) with λ replaced by $-\lambda$ and a (different) space-time white noise which is independent of $\tilde{\eta}$. Hence, by plugging in such expression in place of the increment of \tilde{u} , integrating by parts and using the independence of initial condition and driving noise, we obtain

$$
-\frac{\lambda^2 \nu_{\tau}^{3/2}}{t} \int_0^t \int_0^s \int_{\mathbb{R}^2} x_i x_j \mathbf{E} \Big[\partial_1 \mathcal{M}^{\rho}[u_r^{\tau}](x) \partial_1 \mathcal{M}^{\rho}[u_0^{\tau}](0) \Big] dr ds dx = \frac{\lambda^2 \nu_{\tau}^{3/2}}{t} \int_0^t \int_0^s \int_{\mathbb{R}^2} x_i x_j \mathbf{E} \Big[\partial_1^2 \mathcal{M}^{\rho}[u_r](x) \mathcal{M}^{\rho}[u_0](0) \Big] dr ds dx.
$$

Another integration by parts, allowed by [\(2.6\)](#page-14-4), ensures that the above coincides with the second summand in [\(2.18\)](#page-18-0) if $i = j = 1$ and is 0 otherwise. Putting first and second summands together, the statement follows. \Box

2.2 The semigroup and the generator of SBE

In Proposition [2.3,](#page-14-0) we showed that, for every $\tau > 0$, [\(2.1\)](#page-13-2) admits a unique solution u^{τ} and that such solution is a stationary Markov process. As such, it possesses both a semigroup P^{τ} and a generator \mathcal{L}^{τ} , and all of the analysis detailed in the next sections hinges on determining their behaviour as τ goes to $+\infty$. Prior to that, we need to ensure that, for fixed $\tau > 0$, P^{τ} satisfies suitable basic properties and to study the action of \mathcal{L}^{τ} on its domain.

For $t \geq 0$ and $F \in \mathbb{L}^1(\mathbb{P}^{\tau})$, the semigroup P^{τ} of u^{τ} is given by

$$
P_t^\tau F \stackrel{\text{def}}{=} \mathbb{E}^\tau [F(u_t^\tau) | \mathcal{F}_0], \qquad (2.20)
$$

which is clearly well-defined.

Lemma 2.6 *Let* $\theta \in [1,\infty)$ *. Then,* $(P_t^{\tau})_{t\geq 0}$ *in* [\(2.20\)](#page-19-1) *defines a contractive strongly continuous* semigroup on $\mathbb{L}^{\theta}(\mathbb{P}^{\tau})$ such that, for every $F \in \mathbb{L}^{\theta}(\mathbb{P}^{\tau})$,

$$
||P_t^\tau F||_{\mathbb{L}^\theta(\mathbb{P}^\tau)} \le ||F||_{\mathbb{L}^\theta(\mathbb{P}^\tau)}
$$
\n(2.21)

holds uniformly in τ *and also for* $\theta = \infty$. In particular, the generator \mathscr{L}^{τ} : Dom (\mathscr{L}^{τ}) $\mathbb{L}^{\theta}(\mathbb{P}^{\tau}) \to \mathbb{L}^{\theta}(\mathbb{P}^{\tau})$ associated to P^{τ} is closed and, for every $\mu > 0$, its resolvent $(\mu - \mathcal{L}^{\tau})^{-1}$ is *bounded uniformly in* τ *by* μ^{-1} *and has range given by* Dom(\mathcal{L}^{τ}).

Remark 2.7 Even though for most of the paper we will focus on the case $\theta = 2$, we stated the previous result for generic $\theta \in [1,\infty)$ as the contractivity in [\(2.21\)](#page-19-2) (which is uniform in $\tau > 0$) will be essential to upgrade the convergence of the *two*-point distribution of u^{τ} (i.e. of random vectors of the form $(u_{t_1}^T(\varphi_1), u_{t_2}^T(\varphi_2))$ to that of *finite* dimensional distributions (see Section [5](#page-43-0) for more details).

Proof. The contractivity bound [\(2.21\)](#page-19-2) follows by a standard application of Jensen's inequality for conditional expectations and stationarity of u^{τ} . To see that P^{τ} is a strongly continuous semigroup on $\mathbb{L}^{\theta}(\mathbb{P}^{\tau})$ for $\theta \in [1,\infty)$, it suffices to verify that, for a given bounded cylinder random variable F (or F in any dense subset of $\mathbb{L}^{\theta}(\mathbb{P}^{\tau})$), $P_t^{\tau}F$ converges to F as $t \to 0$ in $\mathbb{L}^{\theta}(\mathbb{P}^{\tau})$. This in turn is a consequence the continuity of u^{τ} in time and smoothness of F. At last, the properties of the generator listed in the statement follow by classical Hille-Yosida theory (see e.g. [\[EK86,](#page-70-13) Theorem 1.2.6]). \Box

We now want to study the action of \mathcal{L}^{τ} on $\mathbb{L}^{2}(\mathbb{P}^{\tau})$, which, by abusing notation, is identified with the Fock space ΓL_{τ}^2 . To obtain an explicit expression for such action, let us introduce the subset ΓH_{τ} of ΓL_{τ}^2 consisting of elements $f = (f_n)_{n \geq 0} \in \Gamma H_{\tau}^2$ such that the sequence $(\|f_n\|_{\Gamma_n H^2_\tau})_{n\geq 0}$ decays faster than any inverse power. Then, the following lemma, whose proof is postponed to Appendix [B,](#page-61-0) holds.

Lemma 2.8 *Let* $\tau > 0$ *be fixed. Let* S^{τ} , \mathcal{A}_{+}^{τ} *and* \mathcal{A}_{-}^{τ} *be the operators such that* $S^{\tau}(\Gamma_{0}L_{\tau}^{2}) =$ $\mathcal{A}_+^{\tau}(\Gamma_0 L_\tau^2) = \mathcal{A}_-^{\tau}(\Gamma_1 L_\tau^2) = 0$, and that, for any $n \in \mathbb{N}$, map $\Gamma_n L_\tau^2 \cap \mathcal{S}(\mathbb{R}^{2n})$ to $\Gamma_n L_\tau^2, \Gamma_{n+1} L_\tau^2$ and $\Gamma_{n-1}L^2$, respectively, and whose action on $\psi\in\Gamma_nL^2_\tau\cap\mathcal{S}(\mathbb{R}^{2n})$ is given by

$$
\mathcal{F}(\mathcal{S}^\tau \psi)(p_{1:n}) \stackrel{\text{def}}{=} \mathcal{F}((\nu_\tau \mathcal{L}_0^{\mathfrak{e}_1} + \mathcal{L}_0^{\mathfrak{e}_2})\psi)(p_{1:n}) = -\frac{1}{2} |\sqrt{R_\tau} p_{1:n}|^2 \hat{\psi}(p_{1:n}) \tag{2.22}
$$

$$
\mathcal{F}(\mathcal{A}_{+}^{\tau}\psi)(p_{1:n+1}) \stackrel{\text{def}}{=} \lambda \nu_{\tau}^{3/4} \frac{2}{n+1} \sum_{1 \leq i < j \leq n+1} \mathcal{K}_{p_i, p_j}^{+,\tau} \hat{\psi}(p_i + p_j, p_{1:n+1} \setminus \{i, j\}), \tag{2.23}
$$

$$
\mathcal{F}(\mathfrak{A}_{-}^{T}\psi)(p_{1:n-1}) \stackrel{\text{def}}{=} \lambda \nu_{\tau}^{3/4} n \sum_{i=1}^{n-1} \int_{\mathbb{R}^2} \mathfrak{X}_{q,p_i-q}^{-,\tau} \hat{\psi}(q, p_i-q, p_{1:n-1}\setminus \{i\}) \, \mathrm{d}q. \tag{2.24}
$$

where

$$
\mathcal{F}(\mathcal{L}_0^{\mathfrak{e}_j} \psi)(p_{1:n}) \stackrel{\text{def}}{=} -\frac{1}{2} |\mathfrak{e}_j \cdot p_{1:n}|^2 \hat{\psi}(p_{1:n}), \qquad j = 1, 2, \qquad (2.25)
$$

$$
\mathcal{K}_{q,q'}^{+,\tau} \stackrel{\text{def}}{=} -\frac{\iota}{2\pi} \left[\mathfrak{e}_1 \cdot (q+q') \right] \Theta_\tau(q+q') , \quad \mathcal{K}_{q,q'}^{-,\tau} \stackrel{\text{def}}{=} -\frac{\iota}{2\pi} \left[\mathfrak{e}_1 \cdot (q+q') \right] \Theta_\tau(q) \Theta_\tau(q') , \text{(2.26)}
$$

 ∂_{τ} *is as in* [\(1.18\)](#page-10-0)*.* Then, \mathcal{L}^{τ} coincides with $S^{\tau} + \mathcal{A}_{+}^{\tau} + \mathcal{A}_{-}^{\tau}$ on ΓH_{τ} *, which is contained in* Dom(\mathcal{L}^{τ})*. Further, on* ΓH_{τ} , S^{τ} *and* $\mathcal{A}^{\tau} \stackrel{\text{def}}{=} \mathcal{A}_{+}^{\tau} + \mathcal{A}_{-}^{\tau}$ *are respectively the symmetric and* s kew-symmetric parts of \mathscr{L}^{τ} and $(\mathscr{A}_{\pm}^{\tau})^* = -\mathscr{A}_{\mp}^{\tau}$. At last, $\Gamma \mathbf{H}_{\tau}$ is a core for \mathscr{L}^{τ} .

Remark 2.9 The importance of $\mathbf{r} \mathbf{H}_{\tau}$ is that, compared to the domain of \mathcal{L}^{τ} , which is only implicitly given, its elements possess an explicit description and, since it is a core, any property or statement regarding \mathcal{L}^{τ} only need to be checked on it. Such a feature will be used throughout the paper.

Before proceeding let us comment on the nature of the operators S^{τ} , \mathcal{A}_{+}^{τ} and \mathcal{A}_{-}^{τ} , and provide some intuition regarding their action on ΓL^2_τ . The Fock space $\Gamma_n L^2_\tau$ can be seen as the span of "*n*-particle states", each particle carrying a momentum p_i , $i \in \{1, ..., n\}$. In this basis, while S^{τ} is diagonal and its action simply consists of multiplying by $((-1/2)$ times) the sum of squared scaled momenta, \mathcal{A}_{+}^{τ} and \mathcal{A}_{-}^{τ} are not. Indeed, \mathcal{A}_{+}^{τ} takes one of the *n* particles, say that of momentum $p_i + p_j$, and *creates* two particles, whose momenta are p_i and p_j . Dually, \mathcal{A}_{-}^{τ} takes two particles, say with momenta p_i and p_j , and merges (or *annihilates*) them into one of momentum $p_i + p_j$. The kernels $\mathcal{K}^{+, \tau}, \mathcal{K}^{-, \tau}$ determine the rates at which these transitions occur.

The difficulty in analysing \mathcal{A}^{τ} (and more specifically \mathcal{A}_{+}^{τ}) lies precisely in the fact that, even though $p_i + p_j$ might be small, there is no mechanism, uniform in τ , that prevents p_i and p_j from being arbitrarily large. To see this, let φ be as regular as it gets, say in $\mathcal{S}(\mathbb{T}^2)$, and consider the ΓH_{τ}^{-1} -norm of $\mathcal{A}^{\tau}\varphi = \mathcal{A}_{+}^{\tau}\varphi$ (the importance of such norm will become apparent in the next lemma). Then, it is not hard to see that

$$
\|\mathcal{A}^\tau \varphi\|_{\Gamma H_\tau^{-1}}^2 = \frac{\lambda^2}{4\pi^2} \int_{\mathbb{R}^2} dp \, |\mathfrak{e}_1 \cdot p|^2 |\hat{\varphi}(p)|^2 \left(\nu_\tau^{3/2} \int dq \frac{\Theta_\tau(p) \Theta_\tau(q) \Theta_\tau(p-q)}{1 + \frac{1}{2} |\sqrt{R_\tau}(q, p-q)|^2}\right)
$$
\n
$$
\gtrsim \nu_\tau \log \tau \int_{\mathbb{R}^2} dp \, |\mathfrak{e}_1 \cdot p|^2 |\hat{\varphi}(p)|^2 \qquad (2.27)
$$

which explodes as $\tau \to \infty$ since $\nu_{\tau} \log \tau \sim (\log \tau)^{1/3}$.

Finally, let us conclude with the following statement which will allow us to perform apriori estimates on the resolvent and approximations thereof.

Lemma 2.10 *Let* $\tau > 0$ *be fixed. Then* $Dom(\mathcal{L}^{\tau}) \subset \Gamma H_{\tau}^1$ *and for* $\psi \in Dom(\mathcal{L}^{\tau})$ *it holds*

$$
\|\psi\|_{\Gamma H_{\tau}^{1}}^{2} = \langle (1 - \mathcal{L}^{\tau}) \psi, \psi \rangle_{\tau}.
$$
 (2.28)

Proof. Thanks to Lemma [2.8,](#page-19-3) [\(2.28\)](#page-21-1) is satisfied for $\psi \in \mathbf{F}$ H_T. Indeed, using the skew-symmetry of A^{τ} , it holds

$$
\|\psi\|_{\Gamma H_{\tau}^{1}}^{2} = \langle (1 - \mathcal{S}^{\tau}) \psi, \psi \rangle_{\tau} = \langle (1 - \mathcal{L}^{\tau}) \psi, \psi \rangle_{\tau}, \qquad (2.29)
$$

which, by Cauchy-Schwarz, implies

$$
\|\psi\|_{\Gamma H^1_\tau} \le \| (1 - \mathcal{L}^\tau) \psi \|_{\Gamma H^{-1}_\tau} \,. \tag{2.30}
$$

Now, let $\psi \in \text{Dom}(\mathcal{L}^{\tau})$, and $(\psi^n)_{n\geq 0} \subset \text{TH}_{\tau}$ be such that $\psi^n \to \psi$ and $(1 - \mathcal{L}^{\tau}) \psi^n \to$ $(1 - \mathcal{L}^{\tau}) \psi$, in ΓL_{τ}^{2} (the existence of such a sequence is due to ΓH_{τ} being a core). Then, applying [\(2.30\)](#page-21-2) to $\psi_n - \psi_m$, we deduce that $(\psi_n)_{n \geq 0}$ is Cauchy in ΓH^1_τ . As it already converges to ψ in ΓL^2_τ , the convergence must necessarily occur in ΓH^1_τ which implies that $\psi \in \Gamma H^1_\tau$. Hence, Dom $(\mathcal{L}^{\tau}) \subset \Gamma H_{\tau}^{\frac{1}{2}}$. Upon applying [\(2.29\)](#page-21-3) to ψ_n and taking the limit in n, [\(2.28\)](#page-21-1) follows. \Box

Let us conclude this section by collecting a few facts on the semigroup, P^{eff} , and the generator, \mathcal{L}^{eff} , of the solution u^{eff} of the limiting (effective) equation in [\(1.10\)](#page-6-2), which we here recall

$$
\partial_t u^{\text{eff}} = \frac{1}{2} \nabla \cdot D_{\text{eff}} \nabla u^{\text{eff}} + \nabla \cdot \sqrt{D_{\text{eff}}} \vec{\xi}, \quad D_{\text{eff}} \stackrel{\text{def}}{=} \begin{pmatrix} C_{\text{eff}}(\lambda) & 0 \\ 0 & 1 \end{pmatrix}, \tag{2.31}
$$

where $C_{\text{eff}}(\lambda)$ is as in [\(1.6\)](#page-4-2).

Lemma 2.11 *The semigroup* $(P_t^{\text{eff}})_{t\geq0}$ *of the solution* u^{eff} *to* [\(2.31\)](#page-21-4) *is strongly continuous on* $\mathbb{L}^{\theta}(\mathbb{P})$ *for* $\theta \in [1,\infty)$ and contractive on $\mathbb{L}^{\theta}(\mathbb{P})$ *for* $\theta \in [1,\infty]$ *. Its generator* \mathscr{L}^{eff} *is closed and has bounded resolvent. Further,* $\mathscr{L}^{eff} = C_{eff}(\lambda) \mathscr{L}_0^{\mathfrak{e}_1} + \mathscr{L}_0^{\mathfrak{e}_2}$, where $C_{eff}(\lambda)$ is the constant given *in* [\(1.6\)](#page-4-2) *and the operators* $\mathcal{L}_0^{\mathfrak{e}_1}$, $\mathcal{L}_0^{\mathfrak{e}_2}$ *were defined in* [\(2.25\)](#page-20-0)*.*

At last, u^{eff} *is* $\mathbb{L}^2(\mathbb{P})$ *-mixing, in the sense that for all* $F \in \mathbb{L}^2(\mathbb{P})$ *, we have*

$$
\lim_{t \to \infty} ||P_t^{\text{eff}} F - \mathbb{E}[F]|| = 0. \tag{2.32}
$$

Proof. Basic properties of the semigroup and generator of u^{eff} are standard and can be proved, e.g., using the same procedure as in Lemmas [2.6](#page-19-4) and [2.8.](#page-19-3) The $\mathbb{L}^2(\mathbb{P})$ -mixing property is wellknown and straightforward to verify (a possible way to do so is by using the explicit expression for the action of the semigroup on ΓL^2). \Box

2.3 Heuristics, novelties and roadmap

Before delving into the details of the proof, we give here a high-level roadmap that outlines the strategy and the main novelties, and should guide the reader through the next sections.

As stated in Theorem [1.4,](#page-7-0) our goal is to prove that the semigroup $(P_t^{\tau})_{t\geq 0}$ of the solution to the regularised SBE in [\(2.1\)](#page-13-2) converges strongly to that of the effective limit equation [\(2.31\)](#page-21-4), $(P_t^{\text{eff}})_{t\geq0}$. Via a suitable version of the classical Trotter-Kato theorem, this follows if we are able

to show strong convergence of the resolvent $(\mu - \mathcal{L}^{\tau})^{-1}$ to $(\mu - \mathcal{L}^{\text{eff}})^{-1}$ for some $\mu > 0$, which amounts to prove that for every $\varphi \in \mathbb{L}^2(\mathbb{P})$, we have

$$
(\mu - \mathcal{L}^{\tau})^{-1} \varphi \xrightarrow[\tau \to \infty]{\mathbb{L}^2(\mathbb{P})} (\mu - \mathcal{L}^{\text{eff}})^{-1} \varphi
$$
 (2.33)

(see Proposition [5.1](#page-43-1) below for a more precise statement). Establishing [\(2.33\)](#page-22-0) requires a thorough understanding of its left-hand side, whose structure is highly non-trivial, and a way to see how, as τ goes to ∞ , the relatively simple right hand side arises. Let us fix $\mu = 1$ and, for the sake of the present informal discussion, a smooth element in the first chaos, i.e. take $\varphi \in \mathcal{S}(\mathbb{R}^2)$, and set $r^{\tau} \stackrel{\text{def}}{=} (1 - \mathcal{L}^{\tau})^{-1} \varphi.$

A crucial observation is that the behaviour of the resolvent is intimately related to the following operator fixed-point-equation

$$
\hbar^{\tau} = -\mathcal{A}_{-}^{\tau} \mathcal{H}_{\tau} \mathcal{A}_{+}^{\tau}, \qquad \text{with} \quad \mathcal{H}_{\tau} \stackrel{\text{def}}{=} (1 - \mathcal{S}^{\tau} + \hbar_{\tau})^{-1}.
$$
 (2.34)

(For a heuristic as to where (2.34) comes from see [\[CT24,](#page-70-14) Section 5.1].) Indeed, assume that [\(2.34\)](#page-22-1) admits a solution h_{τ} and that h_{τ} is diagonal in chaos, i.e. it maps $\Gamma_n L_{\tau}^2$ to itself. Now, define \tilde{r}^{τ} formally as

$$
\tilde{r}^{\tau} \stackrel{\text{def}}{=} \sum_{n\geq 0} (\mathcal{H}_{\tau} \mathcal{A}_{+}^{\tau})^n \mathcal{H}_{\tau} \varphi , \qquad (2.35)
$$

which means that $\tilde{r}_0^{\tau} = 0$, $\tilde{r}_1^{\tau} = \mathcal{H}_{\tau} \varphi$ and, when projected onto the *n*-th chaos component for $n > 1$, $\tilde{r}_n^{\tau} = (\mathcal{H}_{\tau} \mathcal{A}_{+}^{\tau})^{n-1} \mathcal{H}_{\tau} \varphi = \mathcal{H}_{\tau} \mathcal{A}_{+}^{\tau} \tilde{r}_{n-1}^{\tau}$. In particular, it satisfies

$$
\tilde{r}^{\tau} = \mathcal{H}_{\tau} \mathcal{A}_{+}^{\tau} \tilde{r}^{\tau} + \mathcal{H}_{\tau} \varphi.
$$
\n(2.36)

We claim that \tilde{r}^{τ} coincides with r^{τ} . Indeed, applying $(1 - \mathcal{L}^{\tau})$ to \tilde{r}^{τ} we obtain

$$
(1 - \mathcal{L}^{\tau})\tilde{r}^{\tau} = (1 - \mathcal{S}^{\tau} + \hbar_{\tau})\tilde{r}^{\tau} - \mathcal{A}_{+}^{\tau}\tilde{r}^{\tau} - \mathcal{A}_{-}^{\tau}\tilde{r}^{\tau} - \hbar_{\tau}\tilde{r}^{\tau}
$$

$$
= [\mathcal{H}_{\tau}^{-1}\tilde{r}^{\tau} - \mathcal{A}_{+}^{\tau}\tilde{r}^{\tau}] + [-\mathcal{A}_{-}^{\tau}\mathcal{H}_{\tau}\mathcal{A}_{+}^{\tau} - \hbar_{\tau}]\tilde{r}^{\tau} - \mathcal{A}_{-}^{\tau}\mathcal{H}_{\tau}\varphi
$$

where in the first step we added and subtracted $h_{\tau}\tilde{r}^{\tau}$ and in the second we used the definition of \mathcal{H}_{τ} and plugged [\(2.36\)](#page-22-2) into $-\mathcal{A}^{\tau}\tilde{r}^{\tau}$. Now, by (2.36) the first term coincides with φ , the second is 0 thanks to [\(2.34\)](#page-22-1), and the last vanishes since, by Lemma [2.8,](#page-19-3) $\mathcal{A}^{\tau}(\Gamma_1 L^2_{\tau}) \equiv 0$. In other words, $(1 - \mathcal{L}^{\tau})\tilde{r}^{\tau} = \varphi = (1 - \mathcal{L}^{\tau})r^{\tau}$ which, by applying the resolvent at both sides, proves the claim.

What the previous shows is that the operator \hbar_{τ} is the building block at the core of $(1 - \mathcal{L}^{\tau})^{-1}$ as it provides a (semi-)*explicit representation* for r^{τ} . With it in hands, [\(2.33\)](#page-22-0) would follow provided we could prove that

(1) $(r_n^{\tau})_{n\geq 2}$ vanishes as $\tau \to 0$ in ΓL^2_{τ} , (2) $r_1^{\tau} = \mathcal{H}_{\tau} \varphi \in \Gamma_1 L_{\tau}^2$ is close to $(1 - \mathcal{L}^{\text{eff}})^{-1} \varphi$.

The problem with the above is that, even assuming existence and uniqueness of the fixed point, h^{τ} is likely to be a very complicated operator to which we have *no direct access*. The way out, first explored in [\[CGT24\]](#page-70-3) (and [\[CET23b,](#page-69-6) [CG24\]](#page-69-10)) in the so-called weak coupling scaling is

to identify an operator \mathcal{C}^{τ} that, on one hand, is an *approximate* solution of [\(2.34\)](#page-22-1), and on the other has a sufficiently simple form (in particular, it is diagonal in the sense of Definition [1.5\)](#page-12-2) that allows to rigorously show (1) and (2) in an appropriate sense. To move beyond weak-coupling though, this is still not enough.

The difficulty lies in the fact that the error in the approximate fixed point equation (referred to as the "Replacement Lemma", [\[CGT24,](#page-70-3) Lemma 2.5]), while small in τ in each given Fock space component $\Gamma_n L_\tau^2$, grows polynomially in the chaos index n. Such growth is caused by the "off-diagonal terms" (see [\(3.5\)](#page-25-0)) in scalar products involving A^{τ} as explained in the proof of Lemma [3.1.](#page-25-1) Let us remark that this problem is *structural* and somewhat intrinsic to stationary non-reversible Markov processes (see [\[KLO12\]](#page-72-7)). It lead to the introduction of the so-called *graded sector condition* (see [\[HTVo12\]](#page-71-13)) in the study of Gaussian fluctuations for such systems and is responsible for the sub-dominant corrections to the $(\log t)^{2/3}$ -divergence of the diffusion coefficient $D(t)$ of 2d-ASEP [Yauo4] and of the 2d-SBE [\[GHS24\]](#page-71-10), as well as those to the (log t)^{1/2} divergence of $D(t)$ of the AKPZ equation [\[CET23a\]](#page-69-14).

The main novelty of the present work is the identification of a way that allows *both* to determine a diagonal approximate fixed point for [\(2.34\)](#page-22-1) *and* to tame the growth in chaos, thereby establishing [\(2.33\)](#page-22-0) in the strong coupling regime (thus beyond the range of applicability of the graded sector condition, see Remarks [3.5](#page-28-0) and [B.3\)](#page-62-0), and obtaining the precise asymptotic for $D(t)$, as well as a (superdiffusive) central limit theorem. To understand how the strategy came about, there are two aspects of the antisymmetric part of the generator that should be taken into account, namely, the *magnitude* and the *direction* of the momenta it produces.

To see the former, note that \mathcal{A}^{τ} (and more specifically \mathcal{A}_{+}^{τ}) is "dangerous" for two very distinct reasons. As was discussed after Lemma [2.8,](#page-19-3) the operator \mathcal{A}^{τ}_{+} , even when acting on smooth functions with localised momenta, creates *new* momenta which can range from low to arbitrarily high. The creation of high momenta causes the output to be very irregular in real space (or spread out in momentum space) and is the mechanism responsible for the growth of $D(t)$ and, under the correct scaling, for the origin of \mathcal{L}^{eff} . When instead the new momenta are of the same magnitude as the original ones, then new and old "resonate" and generate contributions which are lower order in τ but grow polynomially in the number of variables (i.e. the chaos number).

For the latter, note that even though A^{τ} takes as an input momenta from every direction, the singularity it produces is concentrated *only in the direction* \mathfrak{e}_1 , which is the direction of the "drift", i.e. the non-linearity, of SBE (2.1) . To witness, the right hand side of (2.27) (but also of every upper bound we will obtain on \mathcal{A}^{τ}) only involves the *anisotropic* H^1 -norm of φ , i.e. the H^1 -norm in the direction \mathfrak{e}_1 , which is $\|(-\mathcal{L}_0^{\mathfrak{e}_1})^{1/2}\varphi\|_{\tau} = \frac{1}{2}$ $\frac{1}{2} || (e_1 \cdot \nabla) \varphi ||_{\tau}$. Further, in such direction, the linear part of the dynamics is vanishing at an order $\nu_{\tau} \ll \nu_{\tau}^{3/4}$, the latter being the coefficient of N^{τ} (and unsurprisingly the rate of convergence we derive is $\nu_{\tau}^{1/4}$).

To translate the previous observations in mathematical terms, we introduce two novel tools. First, we identify a splitting of \mathcal{A}_{+}^{τ} (and, by duality, \mathcal{A}_{-}^{τ}) which decouples the singularity in Fourier and the growth in chaos (see Section [3.2\)](#page-26-0). More specifically, we write it as the sum of $\mathcal{A}_+^{\sharp,\tau}$ and $\mathcal{A}_+^{b,\tau}$, with $\mathcal{A}_+^{\sharp,\tau}$ only producing high modes, and no growth in the chaos number, while $\mathcal{A}_{+}^{\flat,\tau}$ keeping track of the latter and regular (actually, small) in Fourier. Second, we introduce a

Recursive Replacement Lemma (see Lemma [3.9\)](#page-32-0) which (morally) allows to recursively estimate each of the summands at the right hand side of [\(2.35\)](#page-22-3) and precisely pins down the contribution of the anisotropic ΓH_{τ}^1 -norm of the resolvent r^{τ} , i.e. the ΓH_{τ}^1 -norm in the direction \mathfrak{e}_1 .

With the previous at hand, the strategy of proof becomes

- (I) find an approximate fixed point q^{τ} to [\(2.34\)](#page-22-1) but with $\mathcal{A}^{\sharp,\tau}_{+}$ and $\mathcal{A}^{\sharp,\tau}_{-}$ in place of \mathcal{A}^{τ}_{+} and \mathcal{A}^{τ}_{-} (so to avoid growth in the chaos in the estimates);
- (II) define the Ansatz for the resolvent s^{τ} as the right hand side of [\(2.35\)](#page-22-3) but with $\mathcal{C}^{\tau} \stackrel{\text{def}}{=}$ $(\mu - \mathcal{S}^{\tau} + q^{\tau})^{-1}$ and $\mathcal{A}_{+}^{\sharp,\tau}$ instead of \mathcal{H}_{τ} and \mathcal{A}_{+}^{τ} ;
- (III) control the anisotropic ΓH_{τ}^1 -norm of s^{τ} , i.e. $\|(-\nu_{\tau}\mathcal{L}_0^{\mathfrak{e}_1})^{1/2} s^{\tau}\|_{\tau}$, with the Recursive Replacement Lemma and show that it vanishes in the limit for $\tau \to \infty$. This will show both that the contribution of $\mathcal{A}^{\flat,\tau} s^{\tau}$ is small and that s^{τ} is sufficiently close to the resolvent applied to φ ;
- (IV) verify that r^{τ} satisfies [\(1\)](#page-22-4) and [\(2\)](#page-22-5) above, thus proving [\(2.33\)](#page-22-0).

The proof of these points will be carried out in Sections [3](#page-24-0) and [4.](#page-35-0) In the former, we will introduce: the splitting $\mathcal{A}^{\sharp,\tau}/\mathcal{A}^{\flat,\tau}$, the approximate fixed point (point [\(I\)\)](#page-24-2); the Recursive Replacement Lemma; and the main estimates needed in the next. Section [4](#page-35-0) puts these tools to practice and addresses points [\(II\)](#page-24-3)[-\(IV\).](#page-24-4)

3 Estimates on the generator acting in Fock spaces

This section is devoted to the analysis of the antisymmetric part A^{τ} of the generator \mathcal{L}^{τ} . We will rigorously present the novel tools mentioned in Section [2.3](#page-21-0) and obtain the bounds we will need in order to establish [\(2.33\)](#page-22-0).

3.1 Diagonal and off-diagonal contributions

As should be clear by [\(2.34\)](#page-22-1), a crucial role will be played by operators of the form $-\mathcal{A}^{\top}\mathcal{D}\mathcal{A}^{\top}$ with $\mathfrak D$ a non-negative diagonal operator on ΓL^2_τ according to Definition [1.5,](#page-12-2) with kernel $\mathbf d = (\mathbf d_n)_{n\geq 0}$ and that we assume to be well-defined on ΓH_{τ} (this is the case, e.g., if \mathfrak{D} is bounded). Notice first that, since by Lemma [2.8](#page-19-3) $-\mathcal{A}_{-}^{\tau} = (\mathcal{A}_{+}^{\tau})^*$ on $\Gamma \mathbf{H}_{\tau}$ and diagonal operators are always self-adjoint (on their natural domain), it holds

$$
\langle -\mathcal{A}_-^T \mathfrak{D} \mathcal{A}_+^T \varphi, \psi \rangle_{\tau} = \langle \mathfrak{D} \mathcal{A}_+^T \varphi, \mathcal{A}_+^T \psi \rangle_{\tau} \quad \text{and} \quad \langle -\mathcal{A}_-^T \mathfrak{D} \mathcal{A}_+^T \varphi, \varphi \rangle_{\tau} = ||\mathfrak{D}^{1/2} \mathcal{A}_+^T \varphi||_{\tau}^2 \tag{3.1}
$$

for $\varphi, \psi \in \mathbf{\Gamma} \mathbf{H}_{\tau}$.

Let us now introduce the *diagonal* and *off-diagonal terms* of operators $-\mathcal{A}^{\top} \mathcal{D} \mathcal{A}^{\top}_+$, with \mathcal{D} as above. For $\varphi, \psi \in \Gamma \mathbf{H}_{\tau}$, the former is given by

$$
\langle -\mathcal{A}_{-}^{\tau}\mathfrak{D}\mathcal{A}_{+}^{\tau}\varphi, \psi \rangle_{\tau, \text{diag}} \stackrel{\text{def}}{=} \sum_{n} n! \int \Xi_{n}^{\tau}(\mathrm{d}p_{1:n}) \hat{\varphi}_{n}(p_{1:n}) \overline{\hat{\psi}}_{n}(p_{1:n})
$$

$$
\times \sum_{i=1}^{n} (\mathfrak{e}_{1} \cdot p_{i})^{2} \frac{\lambda^{2} \nu_{\tau}^{3/2}}{2\pi^{2}} \int_{\mathbb{R}^{2}} \mathrm{d}q \,\mathbf{d}_{n+1}(q, p_{i} - q, p_{1:n \setminus \{i\}}) \mathcal{J}_{q, p_{1} - q}^{\tau}
$$
(3.2)

for $\int_{q,q'}^{\tau} \triangleq \Theta_{\tau}(q)\Theta_{\tau}(q')\Theta_{\tau}(q+q'),$ while the others, referred to as the *off-diagonal terms* of type 1 and 2 respectively, are

$$
\langle -\mathcal{A}^{\tau}_{-}\mathfrak{D}\mathcal{A}^{\tau}_{+}\varphi, \psi \rangle_{\tau, \text{off}_1} \stackrel{\text{def}}{=} \sum_{n} n! c_{\text{off}_1}(n) \lambda^2 \nu_{\tau}^{3/2} \int \Xi_{n+1}^{\tau}(dp_{1:n+1}) \mathbf{d}_{n+1}(p_{1:n+1}) \mathcal{K}_{p_1, p_2}^{+,\tau} \overline{\mathcal{K}_{p_1, p_3}^{+,\tau}}
$$

$$
\times \langle \hat{d}(p_1 + p_2, p_1, \ldots, p_{n+1}) \rangle \overline{\hat{d}}(p_1 + p_2, p_1, \ldots, p_{n+1}) \langle q_1, q_2 \rangle
$$

$$
\times \hat{\varphi}(p_1 + p_2, p_{1:n+1 \setminus \{1,2\}}) \hat{\psi}(p_1 + p_3, p_{1:n+1 \setminus \{1,3\}}), \qquad (3.3)
$$

$$
\langle -\mathcal{A}_{-}^{T} \mathfrak{D} \mathcal{A}_{+}^{T} \varphi, \psi \rangle_{\tau, \text{off}_2} \stackrel{\text{def}}{=} \sum_{n} n! c_{\text{off}_2}(n) \lambda^2 \nu_{\tau}^{3/2} \int \Xi_{n+1}^{\tau}(\text{d}p_{1:n+1}) \,\text{d}_{n+1}(p_{1:n+1}) \mathcal{K}_{p_1, p_2}^{+,\tau} \overline{\mathcal{K}_{p_3, p_4}^{+,\tau}}
$$

$$
\times \hat{\varphi}(p_1 + p_2, p_{1:n+1} \setminus \{1, 2\}) \overline{\hat{\psi}}(p_3 + p_4, p_{1:n+1} \setminus \{3, 4\})
$$
(3.4)

where $c_{off_1}(n) \stackrel{\text{def}}{=} 4n(n-1)$ and $c_{off_2}(n) \stackrel{\text{def}}{=} n(n-1)(n-2)$, and $\mathcal{K}_{q,q'}^{+,\tau}$ is as in [\(2.26\)](#page-20-2). In the next lemma, we show that [\(3.2\)](#page-24-5)-[\(3.4\)](#page-25-2) indeed provide a decomposition of $-\mathcal{A}^T_-\mathcal{D}\mathcal{A}^T_+$.

Lemma 3.1 Let \mathfrak{D} be a non-negative diagonal operator on ΓL^2_τ which is well-defined on $\Gamma \mathbf{H}_\tau$. *Then, the operator* $-\mathfrak{A}_{-}^{T} \mathfrak{D} \mathfrak{A}_{+}^{T}$ *preserves the chaos, in the sense that it commutes with the number operator* N, and for every $\varphi, \psi \in \Gamma H_{\tau}$ we have

$$
\langle -\mathcal{A}_-^{\tau} \mathfrak{D} \mathcal{A}_+^{\tau} \varphi, \psi \rangle_{\tau} = \langle -\mathcal{A}_-^{\tau} \mathfrak{D} \mathcal{A}_+^{\tau} \varphi, \psi \rangle_{\tau, \text{diag}} + \sum_{i=1}^2 \langle -\mathcal{A}_-^{\tau} \mathfrak{D} \mathcal{A}_+^{\tau} \varphi, \psi \rangle_{\tau, \text{off}_i} \tag{3.5}
$$

where the summands at the right hand side are respectively given by [\(3.2\)](#page-24-5)*,* [\(3.3\)](#page-25-3) *and* [\(3.4\)](#page-25-2)*.*

As a consequence, for every $n \in \mathbb{N}$ *, we have*

$$
\left| \langle -\mathcal{A}_-^{\tau} \mathfrak{D} \mathcal{A}_+^{\tau} \varphi_n, \psi_n \rangle_{\tau, \text{off}_i} \right| \lesssim n^2 \langle -\mathcal{A}_-^{\tau} \mathfrak{D} \mathcal{A}_+^{\tau} \varphi_n, \varphi_n \rangle_{\tau, \text{diag}}^{1/2} \langle -\mathcal{A}_-^{\tau} \mathfrak{D} \mathcal{A}_+^{\tau} \psi_n, \psi_n \rangle_{\tau, \text{diag}}^{1/2}, \qquad (3.6)
$$

for all $\varphi, \psi \in \Gamma H_{\tau}$ *.*

Proof. The fact that $-\mathcal{A}^T \mathcal{D} \mathcal{A}^T +$ preserves the chaos follows immediately from Lemma [2.8,](#page-19-3) while (3.5) can be argued as in [\[CET23a,](#page-69-14) Lemma 3.6] so that we will only sketch the proof. Consider the first equality in [\(3.1\)](#page-24-6). By [\(2.23\)](#page-20-3), the Fourier kernels of $\mathcal{A}^{\tau}_{+}\varphi$ and $\mathcal{A}^{\tau}_{+}\psi$ contain a sum over indices $i < j$, so that, when multiplying them and expanding the product, we obtain a sum over $i < j$, $i' < j'$. The diagonal part then corresponds to the sum over $i = i' < j = j'$, the off-diagonal of type 1 to that over $|\{i, j\} \cap \{i', j'\}| = 1$ while the off-diagonal of type 2 to that over $|\{i, j\} \cap \{i', j'\}| = 0$. Equations [\(3.3\)](#page-25-3) and [\(3.4\)](#page-25-2) are then obtained via a change of variables and are a consequence of the fact that φ and ψ are symmetric with respect to permutation of their variables. The constants $c_{\text{off}_i}(n)$, $i = 1, 2$, are then simple combinatorial factors arising from counting the number of possible choices of $i < j$, $i' < j'$ satisfying $|\{i, j\} \cap \{i', j'\}| = 1$ or $|\{i, j\} \cap \{i', j'\}| = 0$. At last, [\(3.6\)](#page-25-4) is a straightforward application of Cauchy-Schwarz and the bound $c_{\text{off}_i}(n) = O(n^3)$. \Box

The expressions [\(3.2\)](#page-24-5), [\(3.3\)](#page-25-3) and [\(3.4\)](#page-25-2) are quite suggestive in regards to the expected contributions from these terms. An inspection of the first hints at where the diffusivity in the direction e_1 comes from. Indeed, its right hand side features an integral in q which *involves neither* φ *nor*

 ψ , so that its convergence as $\tau \to \infty$ is totally independent of them. In particular, their decay in Fourier plays no role, as we saw in [\(2.27\)](#page-20-1) for $\mathcal{D} = (1 - \mathcal{S}^{\tau})^{-1}$. Further, if we assume for a moment that, for the right choice of \mathcal{D} , such integral were a constant D independent of $p_{1:n}$, then we would have $\langle -\mathcal{A}^T \mathcal{D} \mathcal{A}^T_+ \varphi, \psi \rangle_{\tau,\text{diag}} = \langle (-D \mathcal{L}_0^{\epsilon_1}) \varphi, \psi \rangle_{\tau}$. Our goal is to show that for the (approximate) fixed point to [\(2.34\)](#page-22-1), in the limit for $\tau \to \infty$ such integral indeed converges to a constant and that such constant coincides with the first entry of D_{eff} in [\(1.10\)](#page-6-2).

On the other hand, the value of $c_{off_i}(\mathcal{N})$, $i = 1, 2$, in [\(3.3\)](#page-25-3) and [\(3.4\)](#page-25-2), and the trivial bound [\(3.6\)](#page-25-4) say that off-diagonal terms can be controlled by the diagonal one, times a factor growing like the square of the chaos index. This bound alone is way too brutal for our purposes since, as hinted at above, we expect the effective diffusivity to arise from the diagonal terms alone. Thus, the off-diagonal terms should actually be *negligible* in the large-scale limit but, as we will discuss in next section, to see this we need to perform a much more refined analysis of the operators $-\mathcal{A}^{\tau}_{-}\mathfrak{D}\mathcal{A}^{\tau}_{+}.$

3.2 Singling out momentum and chaos singularities

In order to set apart the issues coming from the singularity in Fourier and the chaos-growth, we now introduce a splitting of the operator A^{τ} which is one of the main novelties of the present work. For any $n \in \mathbb{N}$ and $\psi \in \Gamma_n L^2_\tau \cap \mathcal{S}(\mathbb{R}^{2n})$, we define the operators $\mathcal{A}_+^{\sharp,\tau}$ and $\mathcal{A}_-^{\sharp,\tau}$ according to

$$
\mathcal{F}(\mathcal{A}_{+}^{\sharp,\tau}\psi)(p_{1:n+1}) \stackrel{\text{def}}{=} \lambda \nu_{\tau}^{3/4} \frac{2}{n+1} \sum_{1 \leq i < j \leq n+1} \mathcal{K}_{p_i,p_j}^{+,\tau} \chi_{p_i,p_j}^{\sharp,\tau}(p_i+p_j,p_{1:n+1} \setminus \{i,j\}) \times \hat{\psi}(p_i+p_j,p_{1:n+1} \setminus \{i,j\}) \tag{3.7}
$$

$$
\mathcal{F}(\mathfrak{A}_{-}^{\sharp,\tau}\psi)(p_{1:n-1}) \stackrel{\text{def}}{=} \lambda \nu_{\tau}^{3/4} n \sum_{i=1}^{n-1} \int_{\mathbb{R}^2} \chi_{q,p_i-q}^{\sharp,\tau}(p_{1:n-1}) \mathfrak{X}_{q,p_i-q}^{-,\tau} \hat{\psi}(q,p_i-q,p_{1:n-1}\setminus\{i\}) \mathrm{d}q \tag{3.8}
$$

and set $\mathcal{A}^{\sharp,\tau} \stackrel{\text{def}}{=} \mathcal{A}_+^{\sharp,\tau} + \mathcal{A}_-^{\sharp,\tau}$, where χ^{\sharp} is given by

$$
\chi_{q,q'}^{\sharp,\tau}(p_{1:N}) \stackrel{\text{def}}{=} \mathbb{1}_{|\sqrt{R_{\tau}}q| \wedge |\sqrt{R_{\tau}}q'| > 2|\sqrt{R_{\tau}}p_{1:N}|}.
$$
\n(3.9)

Upon setting $\chi^{\flat,\tau} \stackrel{\text{def}}{=} 1 - \chi^{\sharp,\tau}$, we also introduce the operator $\mathcal{A}^{\flat,\tau} \stackrel{\text{def}}{=} \mathcal{A}^{\flat,\tau}_+ + \mathcal{A}^{\flat,\tau}_-$, where $\mathcal{A}^{\flat,\tau}_+$ and $\mathfrak{A}^{\flat,\tau}$ are given as in [\(3.7\)](#page-26-1) and [\(3.8\)](#page-26-2) but with χ^{\flat} in place of χ^{\sharp} . Before discussing their features, let us state in the next lemma their basic properties.

Lemma 3.2 *Let us fix* $\tau > 0$ *. Then the operators* $\mathbb{R}^{\sharp,\tau}$ *and* $\mathbb{R}^{\flat,\tau}$ *are continuous on* ΓH_{τ} *and their sum coincides with* \mathcal{A}^{τ} *. Further, on* ΓH_{τ} *, they are skew-symmetric, and* $(\mathcal{A}_{+}^{\sharp,\tau})^* = -\mathcal{A}_{-}^{\sharp,\tau}$ *,* $(\mathfrak{A}^{\flat,\tau}_{+})^* = -\mathfrak{A}^{\flat,\tau}_{-}$ (in particular [\(3.1\)](#page-24-6) remains true upon replacing $\mathfrak{A}^{\tau}_{\pm}$ by $\mathfrak{A}^{\sharp,\tau}_{\pm}$ or $\mathfrak{A}^{\flat,\tau}_{\pm}$). At *last, for any bounded diagonal operator* $\mathcal D$ *satisfying the assumptions of Lemma* [3.1,](#page-25-1) *both* $-\mathfrak{A}^{\sharp,\tau}_{-}\mathfrak{D}\mathfrak{A}^{\sharp,\tau}_{+}$ and $-\mathfrak{A}^{b,\tau}_{-}\mathfrak{D}\mathfrak{A}^{b,\tau}_{+}$ preserve the chaos, and the same decomposition into diagonal and off-diagonal terms given in [\(3.5\)](#page-25-0) holds with χ^\sharp and χ^\flat appropriately placed at the right hand *sides of* [\(3.2\)](#page-24-5)*-*[\(3.4\)](#page-25-2)*.*

Proof. Continuity can be argued as in Lemma [B.2,](#page-62-1) while the other properties follow immediately by the definition of the operators and straightforward computations. \Box

The advantage of the previous decomposition is that $\mathcal{A}^{\sharp,\tau}$ and $\mathcal{A}^{\flat,\tau}$, while singular in Fourier and chaos respectively, are "regular" in the other. To substantiate such claims, we first show that $\mathcal{A}^{\sharp,\tau}$ does not produce any off-diagonal contribution (and thus its bounds will not display any dependence in N).

Lemma 3.3 *Let* \mathfrak{D} *satisfy the assumptions of Lemma* [3.1.](#page-25-1) *Then, the operator* $-\mathfrak{A}^{\sharp,\tau}_{-} \mathfrak{D} \mathfrak{A}^{\sharp,\tau}_{+}$ *is a non-negative diagonal operator according to Definition [1.5](#page-12-2) on* ΓH^τ *. In other words, for any* $\varphi \in \Gamma \mathbf{H}_{\tau}$ and $\psi \in \Gamma L^2_{\tau}$ it holds

$$
\langle -\mathcal{A}^{\sharp,\tau}_{-}\mathfrak{D}\mathcal{A}^{\sharp,\tau}_{+}\varphi,\psi\rangle_{\tau} = \langle -\mathcal{A}^{\sharp,\tau}_{-}\mathfrak{D}\mathcal{A}^{\sharp,\tau}_{+}\varphi,\psi\rangle_{\tau,\text{diag}}.
$$
 (3.10)

Proof. Let us begin by showing [\(3.10\)](#page-27-0). Thanks to Lemma [3.1](#page-25-1) (and Lemma [3.2\)](#page-26-3), it follows provided $\langle -\mathcal{A}_-^{\sharp,\tau}\mathcal{D}\mathcal{A}_+^{\sharp,\tau}\varphi,\psi\rangle_{\tau,\text{off}_i}=0$ for $i=1,2$. We will focus on the case $i=1$ as the other is completely analogous. By definition, we have

$$
\langle -\mathcal{A}_{-}^{\sharp,\tau}\mathfrak{D}\mathcal{A}_{+}^{\sharp,\tau}\varphi,\psi\rangle_{\tau,\text{off}_1} = \sum_{n} n! c_{\text{off}_1}(n) \lambda^2 \nu_{\tau}^{3/2} \int \Xi_{n+1}^{\tau}(\mathrm{d}p_{1:n+1}) \,\mathrm{d}_{n+1}(p_{1:n+1}) \mathfrak{X}_{p_1,p_2}^{+,\tau} \overline{\mathfrak{X}_{p_1,p_3}^{+,\tau}} \times \chi_{p_1,p_2}^{\sharp,\tau}(p_1+p_2,p_{1:n+1}\setminus\{1,2\}) \chi_{p_1,p_3}^{\sharp,\tau}(p_1+p_3,p_{1:n+1}\setminus\{1,3\}) \times \hat{\varphi}(p_1+p_2,p_{1:n+1}\setminus\{1,2\}) \overline{\hat{\psi}}(p_1+p_3,p_{1:n+1}\setminus\{1,3\}) .
$$

Now, the product of the χ^{\sharp} 's can be different from 0 only if they are both equal to 1. By [\(3.9\)](#page-26-4), this holds provided that

$$
|\sqrt{R_{\tau}}p_1| \wedge |\sqrt{R_{\tau}}p_2| > 2|\sqrt{R_{\tau}}(p_1 + p_2, p_{1:n+1 \setminus \{1,2\}})| \ge |\sqrt{R_{\tau}}p_3|
$$

and

$$
|\sqrt{R_{\tau}}p_2| \wedge |\sqrt{R_{\tau}}p_3| > 2|\sqrt{R_{\tau}}(p_2 + p_3, p_{1:n+1 \setminus \{2,3\}})| \ge |\sqrt{R_{\tau}}p_1|
$$

i.e. $|\sqrt{R_{\tau}}p_1| > |\sqrt{R_{\tau}}p_3| > |\sqrt{R_{\tau}}p_1|$ which is impossible. Hence, [\(3.10\)](#page-27-0) follows.

Since the second line at right hand side of [\(3.2\)](#page-24-5) defines a non-negative Fourier multiplier (as it only depends on $p_{1:n}$) and $\mathcal{A}^{\sharp,\tau}_{+}$, $\mathcal{A}^{\sharp,\tau}_{-}$ and \mathcal{D} are well-defined on $\Gamma \mathbf{H}_{\tau}$, we conclude that $-\mathcal{A}_+^{\sharp,\tau}\mathcal{D}\mathcal{A}_+^{\sharp,\tau}$ defines a diagonal operator according to Definition [1.5](#page-12-2) on $\Gamma\mathbf{H}_\tau$, so that the proof of the statement is complete. \Box

On the other hand, the operator norm of $\mathcal{A}^{b,\tau}$ from ΓH_{τ}^1 to ΓH_{τ}^{-1} is bounded *uniformly in* τ (but the bound depends polynomially on the number operator N), which, together with [\(2.27\)](#page-20-1), says that the singularity in Fourier is fully captured by $\mathcal{A}^{\sharp,\tau}$. The next proposition states an even more refined estimate.

Proposition 3.4 *There exists a constant* $C > 0$ *independent of* τ *such that for every* $\varphi \in \Gamma H_{\tau}$ *it holds*

$$
\|\mathcal{A}_{\pm}^{\flat,\tau}\varphi\|_{\Gamma H_{\tau}^{-1}} \leq C\lambda \|\mathcal{N}(-\nu_{\tau}\mathcal{L}_0^{\mathfrak{e}_1})^{1/2}\varphi\|_{\tau}.
$$
 (3.11)

Remark 3.5 The bound [\(3.11\)](#page-27-1) on $\mathcal{A}^{b,\tau}$ is reminiscent of the so-called Graded Sector Condition (GSC) of [\[KLO12,](#page-72-7) Section 2.7.4] but with three differences. First, the GSC refers to the *full* operator \mathcal{A}_{\pm}^{τ} , while in [\(3.11\)](#page-27-1) only $\mathcal{A}_{\pm}^{\nu,\tau}$ appears. Second, the GSC would require the power of $\mathcal N$ to be strictly less than 1 or λ to be sufficiently small (see [\[HTVo12\]](#page-71-13)). Arguing as in [\[CGT24,](#page-70-3) Lemma 2.4], it should be possible to improve (3.11) and show that instead of N one has \sqrt{N} at its right hand side. On the other hand, the GSC will play no role in our analysis, because anyway it *does not hold independently of* τ *for the full operator* \mathfrak{A}^τ *, but at best only for* $\mathfrak{A}^{\flat,\tau}$ *(see* also Remark [B.3\)](#page-62-0). For this reason, we refrained from optimising [\(3.11\)](#page-27-1) and limited ourselves to use the trivial bound [\(3.6\)](#page-25-4). Third, and more importantly, the operator at the right hand side of [\(3.11\)](#page-27-1) is *not* the symmetric part of \mathcal{L}^{τ} , i.e. \mathcal{S}^{τ} , as is usual for GSC, *but only its component in the* \mathfrak{e}_1 -direction, i.e. $(-\nu_\tau \mathcal{L}_0^{\mathfrak{e}_1})$, that is in a sense much smaller than $-\delta^\tau$. As mentioned in Section [2.3,](#page-21-0) this fact will be crucial for us (see Section [4.2\)](#page-39-0). Note that, if φ is smooth, say in $\mathcal{S}(\mathbb{R}^{2n})$ for some $n \in \mathbb{N}$, then $\|(-\nu_\tau \mathcal{L}_0^{\epsilon_1})^{1/2}\varphi\|_{\tau} \lesssim_{\varphi,n} \sqrt{\nu_\tau} \sim (\log \tau)^{-1/3}$ which vanishes as $\tau \to \infty$ (see [\(2.27\)](#page-20-1) for comparison).

Proof. Let us consider the diagonal bounded operators $\mathcal{D}^{(1)} = (1 - \mathcal{S}^{\tau})^{-1}$ and, for $\kappa > 0$, $\mathfrak{D}^{(2)} = (\kappa - \nu_\tau \mathcal{L}_0^{\mathfrak{e}_1})^{-1}$. A simple duality argument connected to the fact that $\mathfrak{A}^{\flat,\tau}_{-} = -(\mathfrak{A}^{\flat,\tau}_{+})^*$ ensures that it suffices to prove that for all $\varphi \in \mathbf{FH}_{\tau}$ and $(i, j) \in \{(1, 2), (2, 1)\}\)$, uniformly over $\kappa > 0$ we have

$$
\|(\mathfrak{D}^{(i)})^{1/2} \mathfrak{A}^{\flat,\tau}_{+}(\mathfrak{D}^{(j)})^{1/2} \varphi\|_{\tau} \lesssim \|\mathcal{N}\varphi\|_{\tau}.
$$
 (3.12)

Since both $\mathfrak{D}^{(1)}$ and $\mathfrak{D}^{(2)}$ are bounded operators, they are well-defined on $\Gamma \mathbf{H}_{\tau}$, which implies that (3.1) holds. Hence, by (3.5) and (3.6) we get

$$
\begin{split}\n\|(\mathfrak{D}^{(i)})^{1/2} \mathfrak{A}_{+}^{b,\tau} (\mathfrak{D}^{(j)})^{1/2} \varphi\|_{\tau}^{2} &\leq \langle -\mathfrak{A}_{-}^{b,\tau} \mathfrak{D}^{(i)} \mathfrak{A}_{+}^{b,\tau} \mathcal{N} (\mathfrak{D}^{(j)})^{1/2} \varphi, \mathcal{N} (\mathfrak{D}^{(j)})^{1/2} \varphi \rangle_{\tau, \text{diag}} \\
&= \frac{\lambda^{2}}{2\pi^{2}} \sum_{n} n! \, n^{2} \int \Xi_{n}^{\tau} (\mathrm{d}p_{1:n}) \, |\hat{\varphi}_{n}(p_{1:n})|^{2} \times \\
&\times \mathbf{d}_{n}^{(j)}(p_{1:n}) \nu_{\tau}^{3/2} \sum_{i=1}^{n} (\mathfrak{e}_{1} \cdot p_{i})^{2} \int_{\mathbb{R}^{2}} \mathrm{d}q \, \mathbf{d}_{n+1}^{(i)}(q, p_{i} - q, p_{1:n} \setminus \{i\}) \chi_{q, p_{i} - q}^{b}(p_{1:n}) \mathcal{J}_{q, p_{1} - q}^{\tau}\n\end{split} \tag{3.13}
$$

where we further used that $\mathcal N$ and $-\mathfrak{A}^{b,\tau}_{-}\mathfrak{D}\mathfrak{A}^{b,\tau}_{+}$ commute and the definition of the diagonal part in [\(3.2\)](#page-24-5). To estimate the last line above, note that χ^{\flat} is bounded above by the sum of the indicator functions over $\{\sqrt{R_{\tau}}q| < 2|\sqrt{R_{\tau}}p_{1:n}|\}$ and $\{\sqrt{R_{\tau}}(p_i - q)| < 2|\sqrt{R_{\tau}}p_{1:n}|\}$, but the two resulting summands coincide as can be shown by a simple change of variables. Further, the triangle inequality easily gives

$$
\mathbf{d}_{n+1}^{(i)}(q, p_i-q, p_{1:n\setminus\{i\}}) \lesssim \mathbf{d}_n^{(i)}(p_{1:n}).
$$

Putting these observations together, we obtain

$$
\mathbf{d}_{n}^{(j)}(p_{1:n})\nu_{\tau}^{3/2} \sum_{i=1}^{n} (\mathfrak{e}_{1} \cdot p_{i})^{2} \int_{\mathbb{R}^{2}} d q \, \mathbf{d}_{n+1}^{(i)}(q, p_{i} - q, p_{1:n \setminus \{i\}}) \chi_{q, p_{i} - q}^{\flat}(p_{1:n}) \mathcal{F}_{q, p_{1} - q}^{\tau}
$$
\n
$$
\lesssim \mathbf{d}_{n}^{(j)}(p_{1:n}) \mathbf{d}_{n}^{(i)}(p_{1:n}) \nu_{\tau}^{3/2} \sum_{i=1}^{n} (\mathfrak{e}_{1} \cdot p_{i})^{2} \int_{\mathbb{R}^{2}} d q \, \mathbf{1}_{|\sqrt{R_{\tau}}q| < 2|\sqrt{R_{\tau}}p_{1:n}|}
$$

$$
\lesssim \mathbf{d}_n^{(j)}(p_{1:n})\mathbf{d}_n^{(i)}(p_{1:n})\nu_\tau |\mathfrak{e}_1 \cdot p_{1:n}|^2 |\sqrt{R_\tau}p_{1:n}|^2 \lesssim 1\,,
$$

the second to last step being a consequence of the fact that, for any $r > 0$, the volume of $\{|\sqrt{R_{\tau}}q| \leq r\}$ is $r^2 \nu_{\tau}^{-1/2}$, while the last follows from the definition of $\mathfrak{D}^{(1)}$, $\mathfrak{D}^{(2)}$, recalling that ${i, j} = {1, 2}$. Plugging the above back into [\(3.13\)](#page-28-1), [\(3.12\)](#page-28-2) and thus the statement follow. \Box

3.3 The approximate fixed point \mathcal{G}_M^{τ} and the Replacement Lemmas

In this section, we focus on operators involving $\mathcal{A}^{\sharp,\tau}$ only and aim at refining our understanding of operators of the form $-\mathcal{A}_-^{\sharp,\tau} \mathfrak{D} \mathcal{A}_+^{\sharp,\tau}$ for carefully chosen diagonal \mathfrak{D} . More precisely, our goals are to first identify an approximate fixed point \mathcal{G}^{τ} to [\(2.34\)](#page-22-1) (with $\mathcal{A}^{\sharp,\tau}_{\pm}$ in place of \mathcal{A}^{τ}_{\pm}) which is diagonal and bounded, and then to obtain suitable estimates on $\mathcal{G}^\tau\mathcal{A}_+^{\sharp,\tau}$ as this will appear in our ansatz for the solution to the resolvent equation (see [\(II\)](#page-24-3) and $(2.36)/(4.7)$ $(2.36)/(4.7)$).

Let us begin by introducing the candidate approximate fixed point \mathcal{G}^{τ}_{M} , whose definition requires a few notations. Let L^{τ} be the function on \mathbb{R}_+ given by

$$
L^{\tau}(x) \stackrel{\text{def}}{=} \frac{\lambda^2 \nu_{\tau}^{3/2}}{\pi} \log \left(1 + \frac{\tau}{1+x} \right). \tag{3.14}
$$

and, for a measurable function $f: \mathbb{R}_+ \to \mathbb{R}_+$, set $-\mathcal{L}_0^{e_1}[f]$ to be the diagonal operator

$$
-\mathcal{L}_0^{\mathfrak{e}_1}[f] \stackrel{\text{def}}{=} -f(L^{\tau}(-\mathcal{S}^{\tau}))\mathcal{L}_0^{\mathfrak{e}_1},\tag{3.15}
$$

whose Fourier multiplier is $f(L^{\tau}(\frac{1}{2}))$ $\frac{1}{2}$ $\sqrt{R_{\tau}}p_{1:n}|^{2})\frac{1}{2}$ $\frac{1}{2}|\mathfrak{e}_1 \cdot p_{1:n}|^2$ for any $p_{1:n} \in \mathbb{R}^{2n}$ and $n \in \mathbb{N}$.

For $M \geq 0$, let g_M^{τ} be the function on \mathbb{R}_+ defined as

$$
g_M^{\tau}(x) \stackrel{\text{def}}{=} (M\nu_{\tau}) \vee g^{\tau}(x) \tag{3.16}
$$

where

$$
g^{\tau}(x) \stackrel{\text{def}}{=} \left(\frac{3}{2}x + \nu_{\tau}^{3/2}\right)^{2/3} - \nu_{\tau},\tag{3.17}
$$

and q_M^{τ} and \mathcal{G}_M^{τ} be the diagonal operators

$$
q_M^{\tau} \stackrel{\text{def}}{=} -\mathcal{L}_0^{\epsilon_1}[g_M^{\tau}], \quad \text{and} \quad \mathcal{L}_M^{\tau} \stackrel{\text{def}}{=} (1 - \mathcal{S}^{\tau} + g_M^{\tau})^{-1}, \quad (3.18)
$$

which are the approximate fixed point candidates alluded to earlier.

Before stating the lemma that will make the previous claim rigorous, let us comment on the structure (and the advantages) of the operators q_M^{τ} and \mathcal{G}_M^{τ} . What they provide is a *scale-byscale* description of how the limiting operator arises. Indeed, the function g^T appearing in their definition is what determines the size of the diffusivity and, as we will see in the proof of the next statement (see also [\[CGT24,](#page-70-3) Remark 2.6]), it must be chosen in such a way that it solves the **ODE**

$$
\dot{y}(\ell) = \frac{1}{\sqrt{\nu_{\tau} + y(\ell)}}, \qquad y(0) = 0.
$$
\n(3.19)

If $p_{1:n} \in \mathbb{R}^{2n}$ and $M \ge 0$ are given, then the Fourier multiplier of g_M^{τ} satisfies

$$
\mathcal{F}(q_M^{\tau})(p_{1:n}) = g_M^{\tau}(L^{\tau}(\frac{1}{2}|\sqrt{R_{\tau}}p_{1:n}|^2))\frac{1}{2}|\mathfrak{e}_1 \cdot p_{1:n}|^2
$$

$$
= \left\{ \left[\left(\frac{3}{2} L^{\tau} \left(\frac{1}{2} |\sqrt{R_{\tau}} p_{1:n}|^2 \right) + \nu_{\tau}^{3/2} \right)^{2/3} - \nu_{\tau} \right] \vee (M \nu_{\tau}) \right\} \frac{1}{2} |\mathfrak{e}_1 \cdot p_{1:n}|^2
$$

$$
\xrightarrow{\tau \to \infty} \frac{C_{\text{eff}}(\lambda)}{2} |\mathfrak{e}_1 \cdot p_{1:n}|^2
$$
 (3.20)

with $C_{\text{eff}}(\lambda)$ as in [\(1.6\)](#page-4-2), so that the right hand side coincides with the Fourier multiplier of the limiting operator in the direction \mathfrak{e}_1 . As will be made more precise in the next section, this last observation addresses point [\(2\)](#page-22-5) in Section [2.3](#page-21-0)

On the other hand, if $-\mathcal{S}^{\tau} \gtrsim \tau$ then $L^{\tau}(-\mathcal{S}^{\tau})$ (i.e. ℓ in [\(3.19\)](#page-29-2)) is vanishingly small and $g^{\tau}(L^{\tau}(-S^{\tau}))$ is smaller than ν_{τ} so that it does not provide any relevant (or new) information. The parameter M provides an extra degree of freedom which controls the ratio between ν_{τ} and g_M^{τ} for very small ℓ , that is, for very small spatial scales. We will see in Lemma [3.9](#page-32-0) that its role is to artificially force the approximate fixed point to enter some perturbative regime, which will ultimately make our analysis easier. In the spirit of Renormalization Group, the somewhat arbitrary parameter M , while technically useful, does not influence the large-scale properties of the system, notably the diffusion matrix for momenta of order 1, that is independent of M .

We are ready for the first of the two main results of this section, which precisely quantifies how close q_M^{τ} and \mathcal{G}_M^{τ} are from solving [\(2.34\)](#page-22-1).

Lemma 3.6 (The Replacement Lemma) For $M \geq 0$ and $\tau > 0$, let q_M^{τ} and \mathcal{G}_M^{τ} be the *operators defined in* [\(3.18\)](#page-29-3). Then, there exists a constant $C = C(\lambda, M) > 0$ such that for every $\varphi \in \Gamma H_{\tau}$ *it holds*

$$
\| - \mathcal{A}_-^{\sharp,\tau} \mathcal{G}_M^{\tau} \mathcal{A}_+^{\sharp,\tau} \varphi - \mathcal{G}_M^{\tau} \varphi \|_{\Gamma H_\tau^{-1}} \le C \, \| (-\nu_\tau \mathcal{L}_0^{\mathfrak{e}_1})^{1/2} \varphi \|_{\tau} \,.
$$

Remark 3.7 In the proof, the dependence of the constant on λ and M is explicitly tracked and it turns out to be proportional to $\lambda^2 \frac{1 \vee \log(M)}{1 \vee M^{1/2}} + M$.

Remark 3.8 Compared to previous works in which statements as the above have been exploited, Lemma [3.6](#page-30-0) displays two main differences. The first lies in the fact that the norm of φ at the right hand side is *not* the whole of the ΓH_{τ}^1 -norm but only part of it, and more precisely the part in the \mathfrak{e}_1 direction (i.e. the anisotropic ΓH_τ^1 -norm). Second, [\(3.21\)](#page-30-1) does not display any growth in chaos, i.e. its right hand side is *independent* of the number operator N . In all cases previously considered, the error in the Replacement Lemma contained an exploding polynomial factor of the form \mathcal{N}^{γ} , for some $\gamma > 0$, which limited its applicability to the weak-coupling regime.

Proof. Using orthogonality of chaoses and the fact that $-\mathcal{A}^{\sharp,\tau}_{-} \mathcal{G}_{M}^{\tau} \mathcal{A}_{+}^{\sharp,\tau}$ and \mathcal{G}_{M}^{τ} preserve the chaos, it is easy to see that it is enough to prove [\(3.21\)](#page-30-1) for $\varphi \in \Gamma_n L^2_\tau \cap \Gamma \mathbf{H}_\tau$ and any $n \geq 1$. Moreover, the duality between ΓH_{τ}^{-1} and ΓH_{τ}^{1} ensures that we only need to establish the existence of $C = C(\lambda, M) > 0$ such that for all $\varphi, \psi \in \Gamma_n L^2_\tau \cap \Gamma \mathbf{H}_\tau$ it holds

$$
\left| \langle -\mathcal{A}_-^{\sharp,\tau} \mathcal{G}_M^\tau \mathcal{A}_+^{\sharp,\tau} \varphi, \psi \rangle_\tau - \langle \mathcal{g}_M^\tau \varphi, \psi \rangle_\tau \right| \le C \left\| (-\nu_\tau \mathcal{L}_0^{\mathfrak{e}_1})^{1/2} \varphi \right\|_\tau \left\| (-\nu_\tau \mathcal{L}_0^{\mathfrak{e}_1})^{1/2} \psi \right\|_\tau. \tag{3.22}
$$

Now, by Lemma [3.3](#page-27-2) the first summand reduces to its diagonal part (see [\(3.2\)](#page-24-5)) so that, recalling the definitions of q_M^{τ} and \mathcal{G}_M^{τ} in [\(3.18\)](#page-29-3), we have

$$
\left| \langle -\mathcal{A}^{\sharp,\tau}_{-} \mathcal{L}_{M}^{\tau} \mathcal{A}^{\sharp,\tau}_{+} \varphi, \psi \rangle_{\tau} - \langle \mathcal{Q}_{M}^{\tau} \varphi, \psi \rangle_{\tau} \right| \leq n! \int \Xi_{n}^{\tau}(\mathrm{d}p_{1:n}) \left| \hat{\varphi}(p_{1:n}) \right| \hat{\psi}(p_{1:n}) \times \tag{3.23}
$$

$$
\times \sum_{i=1}^n \frac{1}{2} |\mathfrak{e}_1 \cdot p_i|^2 \left| \frac{\lambda^2 \nu_\tau^{3/2}}{\pi^2} \int_{\mathbb{R}^2} \frac{\mathcal{J}_{q, p_i - q}^\tau \chi_{q, p_i - q}^\sharp(p_{1:n})}{1 + \Gamma_i^\tau + g_M^\tau (L^\tau(\Gamma_i^\tau)) \Gamma_i^{\mathfrak{e}_1}} \mathrm{d}q - g_M^\tau (L^\tau(\frac{1}{2} |\sqrt{R_\tau} p_{1:n}|^2)) \right|,
$$

where we adopted the same notations as in [C.1,](#page-64-1) i.e. for $i \in \{1, \ldots, n\}$,

$$
\Gamma_i^{\tau} \stackrel{\text{def}}{=} \frac{1}{2} |\sqrt{R_{\tau}}(q, p_i - q, p_{1:n \setminus \{i\}})|^2, \quad \Gamma_i^{\mathfrak{e}_1} \stackrel{\text{def}}{=} \frac{1}{2} |\mathfrak{e}_1 \cdot (q, p_i - q, p_{1:n \setminus \{i\}})|^2.
$$

By Remark [C.2,](#page-64-2) we can apply Lemma [C.1](#page-64-3) with the function Φ in its statement given by $\Phi(\gamma, \gamma', \ell) \stackrel{\text{def}}{=} (\gamma + g_M^{\tau}(\ell)\gamma')^{-1}$, so that [\(C.4\)](#page-64-4) gives

$$
\left| \frac{\lambda^2 \nu_{\tau}^{3/2}}{\pi^2} \int_{\mathbb{R}^2} \frac{\mathcal{J}_{q,p_i-q}^{\tau} \chi_{q,p_i-q}^{\sharp}(p_{1:n})}{1 + \Gamma_i^{\tau} + g_M^{\tau} (L^{\tau}(\Gamma_i^{\tau})) \Gamma_i^{\epsilon_1}} dq - \int_0^{L^{\tau}(\frac{1}{2}|\sqrt{R_{\tau}}p_{1:n}|^2)} \int_0^{\nu_{\tau}^{-1}} \frac{1}{\pi \sqrt{\varsigma(1 - \nu_{\tau}\varsigma)}} \frac{1}{1 + g_M^{\tau}(\ell) \varsigma} d\varsigma d\ell \right| \lesssim \lambda^2 \frac{1 \vee \log(M)}{1 \vee M^{1/2}} \nu_{\tau} .
$$
\n(3.24)

Let us now focus on the second term at the left hand side and, to abbreviate the notations, set $\ell^{\tau} \stackrel{\text{def}}{=} L^{\tau}(\frac{1}{2})$ $\frac{1}{2}$ $\sqrt{R_{\tau}}p_{1:n}|^2$). We first compute the integral in ς using the change of variables $\varsigma = (\nu_{\tau}(1+t^2))^{-1}$, so that

$$
\int_0^{\nu_\tau^{-1}} \frac{1}{\pi \sqrt{\varsigma(1 - \nu_\tau \varsigma)}} \frac{1}{1 + g_M^{\tau}(\ell) \varsigma} d\varsigma = \frac{1}{\sqrt{1 + g_M^{\tau}(\ell)}},\tag{3.25}
$$

and we are left to control the difference between the integral on $[0, \ell^{\tau}]$ of the latter, and $g_M^{\tau}(\ell^{\tau})$. To do so, we will separately upper and lower bound [\(3.25\)](#page-31-0) in terms of $g_M^{\tau}(\ell^{\tau})$. Recall that g^{τ} in [\(3.17\)](#page-29-4) is clearly pointwise smaller than g_M^{τ} in [\(3.16\)](#page-29-5) and was chosen so that it solves the ODE [\(3.19\)](#page-29-2). Hence,

$$
\int_0^{\ell^{\tau}} \frac{\mathrm{d}\ell}{\sqrt{1+g^{\tau}_M(\ell)}} \leq \int_0^{\ell^{\tau}} \frac{\mathrm{d}\ell}{\sqrt{1+g^{\tau}(\ell)}} = g^{\tau}(\ell^{\tau}) \leq g^{\tau}_M(\ell^{\tau}).
$$

On the other hand, let $\bar{\ell} \ge 0$ be such that $g^{\tau}(\bar{\ell}) = M \nu_{\tau}$. Then, $g_M^{\tau} \equiv g^{\tau}$ on $[\bar{\ell}, +\infty)$, and we can lower bound (3.25) as

$$
\int_0^{\ell^\tau} \frac{\mathrm{d}\ell}{\sqrt{1+g_M^\tau(\ell)}} \ge \int_{\bar{\ell}\wedge\ell^\tau}^{\ell^\tau} \frac{\mathrm{d}\ell}{\sqrt{1+g^\tau(\ell)}} = g^\tau(\ell^\tau) - g^\tau(\bar{\ell}\wedge\ell^\tau) \ge g_M^\tau(\ell^\tau) - M\nu_\tau \tag{3.26}
$$

As a consequence, we deduce

$$
\bigg|\int_0^{L^{\tau}\left(\frac{1}{2}|\sqrt{R_{\tau}}p_{1:n}|^2\right)}\frac{\mathrm{d}\ell}{\sqrt{1+g_M^{\tau}(\ell)}}-g_M^{\tau}(L^{\tau}\left(\frac{1}{2}|\sqrt{R_{\tau}}p_{1:n}|^2\right))\bigg|\leq M\nu_{\tau}\,.
$$

Plugging the latter, together with (3.25) and (3.24) , into (3.23) , (3.22) follows by a simple application of Cauchy-Schwarz, and the proof of the statement is thus complete. \Box

What the previous lemma gives is a quantitative bound on the distance between the operators $-\mathcal{A}^{\sharp,\tau}_{\perp} \mathcal{B}^{\tau}_{M} \mathcal{A}^{\sharp,\tau}_{\perp}$ and \mathcal{G}^{τ}_{M} . For [\(3.21\)](#page-30-1) to be effective, we need its right hand side to go to 0 and, as can be seen from Remark [3.7,](#page-30-4) such smallness cannot come from the constant C (which instead grows linearly in M). Thus, we must hope that, for the functions of interest to us, namely the ansatz anticipated in point [\(II\),](#page-24-3) the anisotropic ΓH_{τ}^1 -norm vanishes in τ for M sufficiently large.

In the next lemma, we state the second main result of this section which is the core of the proof of Proposition [4.8,](#page-39-1) that in turn shows that this is indeed the case.

Lemma 3.9 (The Recursive Replacement Lemma) Let $f: \mathbb{R}_+ \to \mathbb{R}_+$ be a locally Lipschitz *function such that for every* x > 0 *where its derivative exists it holds*

$$
f(x) \le C_f \left(\nu_\tau + g_M^\tau(x)\right) \quad \text{and} \quad |f'(x)| \le C_f \frac{\nu_\tau + g_M^\tau(x)}{x},\tag{3.27}
$$

for some $C_f > 0$. Then, there exists a constant $C_M^{(1)} = C_M^{(1)}(\lambda) > 0$ independent of f and such *that* $\lim_{M \to \infty} C_M^{(1)} = 0$, for which for every $\varphi \in \mathbf{r}$ \mathbf{H}_{τ} we have

$$
\left\| \left(-\mathcal{L}_0^{\mathfrak{e}_1}[f] \right)^{1/2} \mathcal{G}_M^{\tau} \mathcal{A}_+^{\sharp,\tau} \varphi \right\|_{\tau}^2 \leq \left\| \left(-\mathcal{L}_0^{\mathfrak{e}_1}[\mathcal{F}_\tau(f) + \nu_\tau C_f C_M^{(1)}] \right)^{1/2} \varphi \right\|_{\tau}^2, \tag{3.28}
$$

where \mathfrak{G}_M^{τ} is as in [\(3.18\)](#page-29-3) and \mathcal{F}_{τ} is the linear operator acting on $\mathscr{C}(\mathbb{R}_+,\mathbb{R}_+)$ as

$$
\mathcal{F}_{\tau}(f)(x) \stackrel{\text{def}}{=} \frac{1}{2} \int_0^x \frac{f(\ell)}{(\nu_{\tau} + g^{\tau}(\ell))^{3/2}} \, \mathrm{d}\ell \,. \tag{3.29}
$$

Remark 3.10 In the proof, the dependence of the constant $C_M^{(1)}$ on λ and M is explicitly tracked and it turns out to be proportional to $\lambda^2 \frac{1 \vee \log M}{1 \vee M^{1/2}}$ $\frac{1 \vee \log M}{1 \vee M^{1/2}}$

Remark 3.11 The $1/2$ in the expression of \mathcal{I}_{τ} in [\(3.29\)](#page-32-1), and more precisely the fact that it is strictly less than 1, will play a crucial role in our analysis (see Remark [4.11\)](#page-40-0). As we will see in the proof, it comes from differentiating [\(3.25\)](#page-31-0) with respect to g_M^{τ} , and the square-root therein is a direct consequence of the anisotropy of the SBE. Indeed, for "isotropic" models, as the AKPZ equation, the power would be one (as shown in $[CET23b, Eq. (3.29)]$).

Proof. Arguing as in Lemma [3.6,](#page-30-0) it suffices to prove the statement for $\varphi \in \Gamma_n L^2_\tau \cap \Gamma \mathbf{H}_\tau$ and any $n \geq 1$. Furthermore, by Lemma [3.3](#page-27-2) and the fact that both \mathcal{C}_M^{τ} and $-\mathcal{L}_0^{\epsilon_1}[f]$ are non-negative diagonal (see (3.15) for the latter), we have

$$
\begin{split} \Vert (-\mathcal{L}_0^{\mathfrak{e}_1} [f])^{1/2} \, \mathcal{G}_M^{\tau} \mathcal{A}_+^{\sharp,\tau} \varphi \Vert_{\tau}^2 & = \langle -\mathcal{L}_0^{\mathfrak{e}_1} [f] \, (\mathcal{G}_M^{\tau})^2 \, \mathcal{A}_+^{\sharp,\tau} \varphi , \mathcal{A}_+^{\sharp,\tau} \varphi \rangle_{\tau, \text{diag}} \\ & = n! \, \int \Xi_n^{\tau} (\mathrm{d} p_{1:n}) \, |\hat{\varphi}_n(p_{1:n})|^2 \, \times \\ & \times \sum_{i=1}^n \frac{1}{2} |\mathfrak{e}_1 \cdot p_i|^2 \, \frac{\lambda^2 \nu_{\tau}^{3/2}}{\pi^2} \int_{\mathbb{R}^2} \mathrm{d} q \, \frac{f(L^{\tau}(\Gamma_i^{\tau})) \Gamma_i^{\mathfrak{e}_1}}{1 + \Gamma_i^{\tau} + g_M^{\tau}(L^{\tau}(\Gamma_i^{\tau})) \Gamma_i^{\mathfrak{e}_1}} \, \mathfrak{F}_{q, p_i - q}^{\tau} \chi_{q, p_i - q}^{\sharp}(p_{1:n}) \end{split}
$$

where we adopted the same notations as in $(C.1)$.

Once again, we apply Lemma [C.1,](#page-64-3) this time with the function Φ given by $\Phi(\gamma, \gamma', \ell) =$ $f(\ell) \gamma'$ $\frac{f(t)\gamma}{(1+g_M^T(\ell)\gamma')^2}$, which, by Remark [C.2,](#page-64-2) satisfies [\(C.2\)](#page-64-5) and [\(C.3\)](#page-64-6). Hence, we get

$$
\begin{aligned} &\left|\frac{\lambda^2\nu_\tau^{3/2}}{\pi^2}\int_{\mathbb{R}^2}\mathrm{d}q\,\frac{f(L^\tau(\Gamma_i^\tau))\Gamma_i^{\mathfrak{e}_1}}{1+\Gamma_i^\tau+\Gamma_i^{\mathfrak{e}_1}g_M^\tau(L^\tau(\Gamma_i^\tau))} \mathfrak{F}^\tau_{q,p_i-q}\chi_{q,p_i-q}^\sharp(p_{1:n})\right.\\ &\left.-\int_0^{L^\tau\left(\tfrac{1}{2}|\sqrt{R_\tau}p_{1:n}|^2\right)}\int_0^{\nu_\tau^{-1}}\frac{1}{\pi\sqrt{\varsigma(1-\nu_\tau\varsigma)}}\,\frac{f(\ell)\varsigma}{(1+g_M^\tau(\ell)\varsigma)^2}\mathrm{d}\varsigma\mathrm{d}\ell\right|\leq \nu_\tau\,C_f\lambda^2\frac{1\vee\log(M)}{1\vee M^{1/2}}\,. \end{aligned}
$$

We are left to compute the second integral at the left hand side. Set $\ell^{\tau} \stackrel{\text{def}}{=} L^{\tau}(\frac{1}{2})$ $\frac{1}{2}$ $\sqrt{R_{\tau}}p_{1:n}|^2$). The integral in ς can be immediately deduced by [\(3.25\)](#page-31-0) by differentiating with respect to $g_M^{\tau}(\ell)$ (which is a fixed number for given ℓ), so that we obtain

$$
\int_0^{\ell^{\tau}} \int_0^{\nu_{\tau}^{-1}} \frac{1}{\pi \sqrt{\varsigma(1-\nu_{\tau}\varsigma)}} \frac{f(\ell)\varsigma}{(1+g_M^{\tau}(\ell)\varsigma)^2} d\varsigma d\ell = \frac{1}{2} \int_0^{\ell^{\tau}} \frac{f(\ell)}{(1+g_M^{\tau}(\ell))^{3/2}} d\ell
$$

$$
\leq \frac{1}{2} \int_0^{\ell^{\tau}} \frac{f(\ell)}{(1+g^{\tau}(\ell))^{3/2}} d\ell = \mathcal{F}(f)(\ell^{\tau}),
$$

and the proof is concluded.

The reason why we dubbed the previous Lemma *Recursive* is because we will use [\(3.28\)](#page-32-2) *recursively* on the chaos components of our ansatz (see the proof of Proposition [4.8\)](#page-39-1), while *Replacement* is motivated by the fact that to leading order we expect the quantity at the left hand side of [\(3.28\)](#page-32-2) to behave as that at the right hand side *without* the correction $\nu_{\tau} C_f C_M^{(1)}$, and thus we would like to replace the former by the latter. That said, a matching lower bound, which would fully justify the name "replacement", is only true at the cost of a larger (negative) additive error (in the spirit of [\(3.26\)](#page-31-2)) and, in our setting, an upper bound suffices.

An important observation is that while $C_M^{(1)}$ vanishes as $M \to \infty$, the constant C in [\(3.21\)](#page-30-1) explodes (see Remark [3.7\)](#page-30-4). For our purposes though, it will be enough to choose M sufficiently large so that $C_M^{(1)} < 1/2$ (as this will help stabilize the iteration $f \mapsto \mathcal{F}_{\tau}(f) + \nu_{\tau} C_f C_M^{(1)}$) but *fixed*, thus not creating problems when applying Lemma [3.6.](#page-30-0)

We conclude this section with a lemma whose goal is to control the norm of operators of the form $\mathcal{C}_M^{\pi} \mathcal{A}_+^{\sharp,\tau}$ and $\mathcal{A}_-^{\sharp,\tau} \mathcal{C}_M^{\tau}$ as these play a crucial role in addressing point [\(1\)](#page-22-4) in Section [2.3.](#page-21-0)

Proposition 3.12 *There exist a constant* $C_M^{(2)} = C_M^{(2)}(\lambda) > 0$ *which converges to* 0 *as* $M \to \infty$ *, such that for every* $\varphi \in \Gamma H_{\tau}$ *it holds*

$$
\|\mathcal{G}_M^\tau \mathcal{A}_+^{\sharp,\tau} \varphi\|_\tau \le C_M^{(2)} \|\varphi\|_\tau. \tag{3.30}
$$

Furthermore, for any $\alpha \in [0, 1)$ *there exists a constant* $C = C(\lambda, \alpha) > 0$ *, for which*

$$
\|\mathcal{G}_M^{\tau}\mathcal{A}_+^{\sharp,\tau}\mathbb{1}_{-\delta^{\tau}\leq\tau^{\alpha}}\varphi\|_{\tau}\vee\|\mathcal{A}_-^{\sharp,\tau}\mathcal{G}_M^{\tau}\mathbb{1}_{-\delta^{\tau}\leq\tau^{\alpha}}\varphi\|_{\tau}\leq C\,\nu_{\tau}^{3/4}\,\|\varphi\|_{\tau}\,,\tag{3.31}
$$

where $1\!\!1_{-8^{\tau}\leq \tau^{\alpha}}$ *stands for the diagonal operator whose kernel is* $1\!\!1_{|\sqrt{R_{\tau}}p_{1:n}|^2/2\leq \tau^{\alpha}}$ *.*

 \Box

Remark 3.13 As for the replacement lemmas, the dependence of $C_M^{(2)}$ and C on M, λ and τ can be explicitly tracked and they can respectively be taken to be proportional to $\lambda/(1 \vee M^{3/4})$ and $(1 - \beta)^{-1/2} + \lambda \nu_{\tau}^{-3/2} \tau^{-(\beta - \alpha)/2}$, for any $0 \le \alpha < \beta < 1$.

Proof. As in the proof of Replacement Lemmas [3.6](#page-30-0) and [3.9,](#page-32-0) it suffices to prove the statement for $\varphi \in \Gamma_n L^2_\tau \cap \Gamma \mathbf{H}_\tau$ and any $n \geq 1$. Furthermore, by Lemma [3.3,](#page-27-2) the fact that both \mathcal{L}_M^{τ} and $\mathscr{L}_0^{\mathfrak{e}_1}[f]$ are diagonal (see [\(3.15\)](#page-29-6) for the latter), and formula [\(3.2\)](#page-24-5), we have

$$
\|\mathcal{G}_M^{\pi}\mathcal{A}_+^{\sharp,\tau}\varphi\|_{\tau}^2 = \langle -\mathcal{A}_-^{\sharp,\tau}(\mathcal{G}_M^{\tau})^2 \mathcal{A}_+^{\sharp,\tau}\varphi, \varphi \rangle_{\tau,\text{diag}} \tag{3.32}
$$

$$
\lesssim n! \int \Xi_n^{\tau}(\mathrm{d}p_{1:n}) |\hat{\varphi}(p_{1:n})|^2 \lambda^2 \sum_{i=1}^n |\mathfrak{e}_1 \cdot p_i|^2 \nu_{\tau}^{3/2} \int_{\mathbb{R}^2} \frac{\mathrm{d}q}{(1 + \Gamma_i^{\tau} + g_M^{\tau}(L^{\tau}(\Gamma_i^{\tau})) \Gamma_i^{\mathfrak{e}_1})^2}
$$

where we adopted the same notations as in $(C.1)$. Let us now focus on the inner integral which will be bounded in two different ways to obtain [\(3.30\)](#page-33-0) and [\(3.31\)](#page-33-1), respectively.

For the former, since $g_M^{\tau} \ge M \nu_{\tau}$ (see [\(3.17\)](#page-29-4)) we have

$$
\begin{aligned}\int_{\mathbb{R}^2}\frac{\mathrm{d} q}{\left(1+\Gamma_i^\tau+g_M^\tau(L^\tau(\Gamma_i^\tau))\,\Gamma_i^{\mathfrak{e}_1}\right)^2}&\lesssim \int_{\mathbb{R}^2}\frac{\mathrm{d} q}{(\nu_\tau(M\vee 1)|\mathfrak{e}_1\cdot p_{1:n}|^2+|\sqrt{R_\tau}q|^2+M\nu_\tau|\mathfrak{e}_1\cdot q|^2)^2}\\&\lesssim \frac{1}{\nu_\tau^{3/2}(1\vee M^{3/2})|\mathfrak{e}_1\cdot p_{1:n}|^2}\end{aligned}
$$

where the last estimate follows by $(C.5)$ in Lemma [C.3.](#page-65-1) Then, plugging the previous into [\(3.32\)](#page-34-0), we obtain (3.30) .

Let us now turn to [\(3.31\)](#page-33-1). The bound on the term containing $\mathcal{A}^{\sharp,\tau}_{-}$ follows from that on $\mathcal{A}_+^{\sharp,\tau}$ and a duality argument. Indeed, duality ensures that it suffices to control the ΓL^2_τ -norm of $\mathbb{1}_{-8^{\tau} \leq \tau^{1/2}} \mathcal{G}_M^{\tau} \mathcal{A}_+^{\sharp,\tau} \varphi$ via the right hand side of [\(3.31\)](#page-33-1). But, thanks to the definition of χ^{\sharp} in [\(3.9\)](#page-26-4), we have

$$
1_{-\delta^\tau \leq \tau^\alpha} \mathcal{G}_M^\tau \mathcal{A}_+^{\sharp,\tau} = 1_{-\delta^\tau \leq \tau^\alpha} \mathcal{G}_M^\tau \mathcal{A}_+^{\sharp,\tau} 1_{-\delta^\tau \leq \tau^\alpha}
$$

so that we can apply [\(3.31\)](#page-33-1) for $\mathcal{A}_+^{\sharp,\tau}$ to conclude.

To prove the estimate on $\mathcal{A}_+^{\sharp,\tau}$, we split the integral in [\(3.32\)](#page-34-0) in two regions, corresponding to $|q|^2 \leq \tau^{\beta}$ and $|q|^2 > \tau^{\beta}$, for some given β such that $\alpha < \beta < 1$. In the first, we have $\Gamma_{i}^{\tau}=\frac{1}{2}|$ $\sqrt{R_{\tau}}(q, p_i - q, p_{2:n})|^2 \lesssim \tau^{\beta}$, which implies that $g_M^{\tau}(L^{\tau})$ $\frac{1}{2}$ $\sqrt{R_{\tau}}(q, p_i-q, p_{2:n})|^2) \geq$ $\lambda^{4/3} (1 - \beta)^{2/3}$. Hence,

$$
\begin{aligned} \int \frac{1_{\{|q|^2\leq\tau^\beta\}}\mathrm{d}q}{\left(1+\Gamma_i^\tau+g_M^\tau(L^\tau(\Gamma_i^\tau))\,\Gamma_i^{\mathfrak{e}_1}\right)^2} &\lesssim \int \frac{\mathrm{d}q}{\left(|\sqrt{R_\tau}q|^2+\lambda^{4/3}(1-\beta)^{2/3}(|\mathfrak{e}_1\cdot q|^2+|\mathfrak{e}_1\cdot p_{1:n}|^2)\right)^2}\\ &\lesssim \frac{1}{\lambda^2(1-\beta)|\mathfrak{e}_1\cdot p_{1:n}|^2} \end{aligned}
$$

where in the last step we used $(C.5)$ from Lemma [C.3](#page-65-1) with A and M therein respectively given by $A^2 = \lambda^{4/3} (1 - \beta)^{2/3} |\mathfrak{e}_1 \cdot p_{1:n}|^2$ and $M = \nu_\tau^{-1} \lambda^{4/3} (1 - \beta)^{2/3}$. For the other instead, we keep only the term $|\sqrt{R_T}q|^4 \gtrsim \nu_T^2|q|^4$ at the denominator and obtain

$$
\int \frac{1_{\{|q|^2>\tau^\beta\}}{\rm d}q}{(1+\Gamma_i^\tau+g_M^\tau(L^\tau(\Gamma_i^\tau))\,\Gamma_i^{\mathfrak e_1})^2} \lesssim \int \frac{1_{\{|q|^2>\tau^\beta\}}{\rm d}q}{\nu_\tau^2|q|^4} \lesssim \frac{1}{\nu_\tau^2\tau^\beta} \leq \frac{1}{\nu_\tau^3\tau^{\beta-\alpha}|\mathfrak e_1\cdot p_{1:n}|^2}
$$

the last bound being a consequence of $\tau^{\alpha} \ge |\sqrt{R_{\tau}} p_{1:n}|^2 \gtrsim \nu_{\tau} |\mathfrak{e}_1 \cdot p_{1:n}|^2$. Therefore, [\(3.31\)](#page-33-1) follows by estimating [\(3.32\)](#page-34-0) via the two equations above and thus, the proof of the proposition is complete. \Box

4 The Resolvent and its ansatz

Now that we collected all the necessary bounds, we turn to the analysis of the resolvent and aim at obtaining the estimates needed to prove Theorems [1.1](#page-4-0) and [1.3.](#page-6-1) To properly state them, let φ be an element of $\Gamma_{n_0} L_\tau^2$ for some fixed $n_0 \in \mathbb{N}$ and r^τ be the resolvent applied to φ , i.e.

$$
r^{\tau} \stackrel{\text{def}}{=} (1 - \mathcal{L}^{\tau})^{-1} \varphi \,. \tag{4.1}
$$

Note that, if $n_0 = 0$, then r^{τ} coincides with φ (and there is thus nothing to prove) so that in what follows we will focus on the case of $n_0 > 0$. The main result of the present section is summarised in the following theorem.

Theorem 4.1 Let $n_0 \in \mathbb{N}$, $n_0 \geq 1$, and M be sufficiently large so that the constants $C_M^{(1)}$ and C (2) ^M *in* [\(3.28\)](#page-32-2) *and* [\(3.30\)](#page-33-0) *satisfy*

$$
C_M^{(1)} \vee C_M^{(2)} \le \frac{1}{2} \,. \tag{4.2}
$$

Then, there exists a constant $C = C(\lambda, M, n_0) > 0$ *such that for all* $\varphi \in \Gamma_{n_0} L^2_{\tau}$ *, we have*

$$
||r^{\tau}||_{\tau} \le C \Big(||\varphi||_{\tau,M} + ||\mathcal{G}_M^{\tau}\varphi||_{\tau} \Big), \tag{4.3}
$$

$$
\left| \|\boldsymbol{r}^{\tau}\|_{\boldsymbol{\Gamma} H_{\tau}^{1}}^{2} - \|(\mathcal{E}_{M}^{\tau})^{1/2} \varphi\|_{\tau}^{2} \right| \leq C \left\| \varphi \right\|_{\tau,M} \left(\|\varphi\|_{\tau,M} + \|(\mathcal{E}_{M}^{\tau})^{1/2} \varphi\|_{\tau} \right) \tag{4.4}
$$

where r τ *is defined according to* [\(4.1\)](#page-35-2) *and the (semi-)norm* ||| · ||| *is given by*

$$
\|\varphi\|_{\tau,M} \stackrel{\text{def}}{=} \nu_{\tau}^{1/4} \sqrt{|\log \nu_{\tau}|} \| (-\mathcal{L}_0^{\mathfrak{e}_1})^{1/2} \mathcal{L}_M^{\tau} \varphi\|_{\tau} + \|\mathcal{A}_-^{\sharp,\tau} \mathcal{L}_M^{\tau} \varphi\|_{\tau} . \tag{4.5}
$$

Furthermore, if there exists α *such that* $\text{Supp}(\mathcal{F}\varphi) \subset \{-\mathcal{S}^\tau \leq \tau^\alpha\}$, then there exists a constant $C = C(\lambda, \alpha) > 0$ for which

$$
\|r^{\tau} - \mathcal{G}_M^{\tau} \varphi\| \le C \, \nu_{\tau}^{1/4} \sqrt{|\log \nu_{\tau}|} \, \|\varphi\|_{\tau} \,. \tag{4.6}
$$

As outlined in Section [2.3,](#page-21-0) the proof of the above statement is based on a approximation of the resolvent with a suitably chosen ansatz s^{τ} (point [\(II\)](#page-24-3) of the roadmap). The latter is introduced in Section [4.1](#page-36-0) where its main properties together with a control over the ΓH_{τ}^1 -norm of the difference between r^{τ} and s^{τ} are derived. Most of the estimates we obtain therein are in terms of $\|(-\nu_\tau \mathcal{L}_0^{\epsilon_1})^{1/2} s^\tau\|_{\tau}$ whose control is only possible via the Recursive Replacement Lemma [3.9.](#page-32-0) This is detailed in Section [4.2](#page-39-0) (point [\(III\)\)](#page-24-7), at the end of which all the results derived are put together and the proof of Theorem [4.1](#page-35-1) is completed.
4.1 The ansatz and its properties

As explained in Section [2.3,](#page-21-0) the resolvent is a complicated object whose structure intimately depends on the fixed point operator to [\(2.34\)](#page-22-0). The ansatz that we will now introduce shares the same structure, i.e. that in [\(2.36\)](#page-22-1), but with two twists: we replace \mathcal{H}^{τ} with \mathcal{G}_{M}^{τ} given in [\(3.18\)](#page-29-0), and the whole of \mathcal{A}_{+}^{τ} with $\mathcal{A}_{+}^{\sharp,\tau}$ in [\(3.7\)](#page-26-0).

Let φ be an element of $\Gamma_{n_0} L_\tau^2$ for some fixed $n_0 \geq 1$. We define the ansatz s^τ for the resolvent equation in [\(4.1\)](#page-35-0) according to

$$
s^{\tau} \stackrel{\text{def}}{=} \sum_{k=0}^{+\infty} (\mathcal{G}_M^{\tau} \mathcal{A}_+^{\sharp,\tau})^k \mathcal{G}_M^{\tau} \varphi.
$$
 (4.7)

Equation [\(4.7\)](#page-36-0) provides the full chaos decomposition of s^{τ} as, for any k, $(\mathcal{C}_{M}^{\tau}\mathcal{A}_{+}^{\sharp,\tau})^{k}\mathcal{C}_{M}^{\tau} \in$ $\Gamma_{n_0+k} L_\tau^2$. Let us first show that, for fixed $\tau > 0$, the series at the right hand side is absolutely convergent in ΓH_{τ} so that s^{τ} is indeed a well-defined element of the core, and derive an expression for the action of the generator on it.

Lemma 4.2 *Let* $\varphi \in \Gamma_{n_0} L^2_{\tau}$ *for some* $n_0 \geq 1$ *, and assume that* [\(4.2\)](#page-35-1) *holds. Then, for* $\tau > 0$ *fixed, the sum at the right hand side of* [\(4.7\)](#page-36-0) *is absolutely convergent in* $\Gamma \textbf{H}_{\tau}$ *, so that* s^{τ} *is indeed well defined. Moreover, it satisfies*

$$
s^{\tau} = \mathcal{G}_M^{\tau} \mathcal{A}_+^{\sharp,\tau} s^{\tau} + \mathcal{G}_M^{\tau} \varphi , \qquad (4.8)
$$

and

$$
(1 - \mathcal{L}^{\tau})s^{\tau} = \varphi + (-\mathcal{A}^{\sharp,\tau}_{-}\mathcal{G}_{M}^{\tau}\mathcal{A}^{\sharp,\tau}_{+} - q^{\tau}_{M})s^{\varepsilon} - \mathcal{A}^{\flat,\tau}s^{\varepsilon} - \mathcal{A}^{\sharp,\tau}_{-}\mathcal{G}_{M}^{\tau}\varphi.
$$
(4.9)

Proof. Assume first that the sum in [\(4.7\)](#page-36-0) indeed converges absolutely in ΓH_{τ} , and let us prove [\(4.8\)](#page-36-1) and [\(4.9\)](#page-36-2). For [\(4.8\)](#page-36-1), notice that

$$
s^\tau=\mathcal{G}^\tau_M\varphi+\sum_{k\geq 0}(\mathcal{G}^\tau_Mg_+^{\sharp,\tau})^{k+1}\,\mathcal{G}^\tau_M\varphi=\mathcal{G}^\tau_M\varphi+\mathcal{G}^\tau_Mg_+^{\sharp,\tau}s^\tau\,,
$$

where in the last step we are able to collect $\mathcal{C}_M^{\tau} \mathcal{A}_+^{\sharp,\tau}$ from the infinite sum, which is allowed as this operator is continuous on ΓH_{τ} and the sum is absolutely convergent therein.

For [\(4.9\)](#page-36-2), by adding and subtracting $q_M^{\tau} s^{\tau}$ from $(1 - \mathcal{L}^{\tau}) s^{\tau}$, we obtain

$$
(1 - \mathcal{L}^{\tau}) s^{\tau} = (1 - \mathcal{S}^{\tau} + q^{\tau}_{M}) s^{\tau} - \mathcal{A}_{+}^{\sharp, \tau} s^{\tau} - \mathcal{A}_{-}^{\sharp, \tau} s^{\tau} - q^{\tau}_{M} s^{\tau} - \mathcal{A}^{\flat, \tau} s^{\varepsilon}
$$

= $(\mathcal{G}_{M}^{\tau})^{-1} s^{\tau} - \mathcal{A}_{+}^{\sharp, \tau} s^{\tau} - \mathcal{A}_{-}^{\sharp, \tau} s^{\tau} - q^{\tau}_{M} s^{\tau} - \mathcal{A}^{\flat, \tau} s^{\tau}$
= $\varphi + (-\mathcal{A}_{-}^{\sharp, \tau} \mathcal{G}_{M}^{\tau} \mathcal{A}_{+}^{\sharp, \tau} - q^{\tau}_{M}) s^{\tau} - \mathcal{A}_{-}^{\sharp, \tau} \mathcal{G}_{M}^{\tau} \varphi - \mathcal{A}^{\flat, \tau} s^{\tau},$

the second step being a consequence of the definition of \mathcal{C}_{M}^{τ} in [\(3.18\)](#page-29-0) and the last of two applications of [\(4.8\)](#page-36-1), one for the first two terms and the other for the third and fourth.

Let us now argue that the series at right hand side of [\(4.7\)](#page-36-0) converges absolutely in $\mathbf{r} \mathbf{H}_{\tau}$. First, by Proposition [3.12](#page-33-0) and [\(4.2\)](#page-35-1), $\|\mathcal{G}_{M}^{\tau}\mathcal{A}_{+}^{\sharp,\tau}\|_{\Gamma L_{\tau}^{2}\to\Gamma L_{\tau}^{2}} \leq 1/2$. Hence, a k-times iteration of this bound gives

$$
||s_{n_0+k}^\tau||_\tau = ||(\mathcal{G}_M^\tau \mathcal{A}_+^{\sharp,\tau})^k \mathcal{G}_M^\tau \varphi||_\tau \le 2^{-k} ||\mathcal{G}_M^\tau \varphi||_\tau \le 2^{-k} ||\varphi||_\tau. \tag{4.10}
$$

Now, since, by definition, it can be directly checked that $s_{n_0+k+1}^{\tau} = \mathcal{C}_{\varepsilon} \mathcal{A}_{+}^{\sharp,\tau} s_{n_0+k}^{\tau}$, we have

$$
||s_{n_0+k+1}^{\tau}||_{\Gamma H^2_{\tau}} = ||\mathcal{G}_{\varepsilon} \mathcal{A}_{+}^{\sharp,\tau} s_{n_0+k}^{\tau}||_{\Gamma H^2_{\tau}} \le ||\mathcal{A}_{+}^{\sharp,\tau} s_{n_0+k}^{\tau}||_{\tau} \lesssim_{\lambda,\tau} (n_0+k)^{3/2} ||s_{n_0+k}^{\tau}||_{\tau}, \quad (4.11)
$$

where in the last step we exploited [\(B.1\)](#page-62-0) which, even though stated (and proved) for \mathcal{A}^{τ}_{\pm} , also holds for $\mathcal{A}_{\pm}^{\sharp,\tau}$ (with the same proof). Combining [\(4.10\)](#page-36-3) and [\(4.11\)](#page-37-0) concludes the proof.

The (rather) explicit structure of s^{τ} allows to efficiently estimate both its ΓL_{τ}^2 and ΓH_{τ}^1 -norms, as the next proposition shows.

Proposition 4.3 *Assumte that* [\(4.2\)](#page-35-1) *holds. Then, there exists a constant* $C = C(\lambda, M) > 0$ *such that for all* $\varphi \in \Gamma_{n_0} L_\tau^2$ *for some* $n_0 \in \mathbb{N}$ *, we have*

$$
||s^{\tau}||_{\tau} \le 2 ||\mathcal{C}_M^{\tau} \varphi||_{\tau}, \qquad (4.12)
$$

$$
\left| \|s^{\tau}\|_{\Gamma H^1_{\tau}}^2 - \|(\mathcal{G}_M^{\tau})^{1/2} \varphi\|_{\tau}^2 \right| \le C \|(-\nu_{\tau} \mathcal{L}_0^{\mathfrak{e}_1})^{1/2} s^{\tau} \|_{\tau}^2 \tag{4.13}
$$

where s^{τ} is defined according to [\(4.7\)](#page-36-0). Furthermore, if φ is such that $\text{Supp}(\mathcal{F}\varphi) \subset \{-8^{\tau} \leq \tau^{\alpha}\}\$ *for some* $\alpha \in [0, 1)$ *, then there exists a constant* $C = C(\lambda, \alpha) > 0$ *independent of* φ *for which*

$$
||s^{\tau} - \mathcal{G}_M^{\tau} \varphi||_{\tau} \le \nu_{\tau}^{3/4} C ||\varphi||_{\tau}, \qquad (4.14)
$$

Remark 4.4 As the proof shows, the constant C in (4.13) is the same at that of the Replacement Lemma, i.e. the one for which (3.21) holds, and that in (4.14) coincides with that of (3.31) .

Proof. As noted in the proof of Lemma [4.2,](#page-36-4) Proposition [3.12](#page-33-0) and [\(4.2\)](#page-35-1) imply that $||\mathcal{G}_M^{\pi} \mathcal{A}_+^{\sharp,\tau}||_{\Gamma} \mathcal{L}_\tau^2 \to \Gamma \mathcal{L}_\tau^2$ 1/2. Hence,

$$
\|s^\tau\|_\tau \leq \sum_k \|(\mathfrak{E}^\tau_M \mathfrak{A}_+^{\sharp,\tau})^k \mathfrak{E}^\tau_M \varphi\|_\tau \leq \sum_k 2^{-k} \|\mathfrak{E}^\tau_M \varphi\|_\tau \leq 2 \|\mathfrak{E}^\tau_M \varphi\|_\tau\,,
$$

from which [\(4.12\)](#page-37-3) follows.

For [\(4.13\)](#page-37-1), Lemma [4.2](#page-36-4) ensures that $s^{\tau} \in \mathbf{FH}_{\tau} \subset \text{Dom}(\mathcal{L}^{\tau})$, so that [\(2.28\)](#page-21-1) gives

$$
||s^{\tau}||_{\Gamma H_{\tau}^{1}}^{2} = \langle (1 - \mathcal{L}^{\tau})s^{\tau}, s^{\tau} \rangle_{\tau}.
$$
\n(4.15)

Plugging [\(4.9\)](#page-36-2) in the above formula, we obtain

$$
\|s^{\tau}\|_{\Gamma H^1_{\tau}}^2 = \langle \varphi, s^{\tau} \rangle_{\tau} + \langle (-\mathcal{A}^{\sharp,\tau}_{-} \mathcal{G}_{M}^{\tau} \mathcal{A}^{\sharp,\tau}_{+} - \mathcal{G}_{M}^{\tau}) s^{\tau}, s^{\tau} \rangle_{\tau} - \langle \mathcal{A}^{\flat,\tau} s^{\tau}, s^{\tau} \rangle_{\tau} - \langle \mathcal{A}^{\sharp,\tau}_{-} \mathcal{G}_{M}^{\tau} \varphi, s^{\tau} \rangle_{\tau}.
$$

Now, the third and fourth terms are 0: the third since $\mathcal{A}^{\flat,\tau}$ is skew-symmetric on $\mathbf{\Gamma} \mathbf{H}_{\tau}$ by Lemma [3.2;](#page-26-1) the fourth since $s^{\tau} \in \overline{\bigoplus_{n \ge n_0} \Gamma_n L_{\tau}^2}$ while $\mathscr{A}_{-}^{\sharp,\tau} \mathscr{G}_{M}^{\tau} \varphi \in \Gamma_{n_0-1} L_{\tau}^2$ and these spaces are orthogonal. For the first, we use again the orthogonality of Fock spaces with different indices and [\(4.7\)](#page-36-0), and deduce

$$
\langle \varphi, s^{\tau} \rangle_{\tau} = \langle \varphi, s_{n_0}^{\tau} \rangle_{\tau} = \langle \varphi, \mathfrak{E}_M^{\tau} \varphi \rangle_{\tau} = \|(\mathfrak{E}_M^{\tau})^{1/2} \varphi\|_{\tau}^2 \,.
$$

Thus we are left to control the second term, on which we apply (3.21) in the Replacement Lemma [3.6,](#page-30-1) and get [\(4.13\)](#page-37-1).

At last, we turn to [\(4.14\)](#page-37-2) whose proof is similar to that of [\(4.12\)](#page-37-3). By definition

$$
\|s^{\tau}-\mathcal{G}_M^{\tau}\varphi\|_{\tau}\leq \sum_{k\geq 1}\|(\mathcal{G}_M^{\tau}\mathcal{A}_+^{\sharp,\tau})^k\mathcal{G}_M^{\tau}\varphi\|_{\tau}\leq \|(\mathcal{G}_M^{\tau}\mathcal{A}_+^{\sharp,\tau})\mathcal{G}_M^{\tau}\varphi\|_{\tau}
$$

where we used that $||\mathcal{G}_{M}^{\tau} \mathcal{A}_{+}^{\sharp,\tau}||_{\Gamma}L_{\tau}^{2} \to \Gamma L_{\tau}^{2} \leq 1/2$. Now, since \mathcal{G}_{M}^{τ} is diagonal, the Fourier transforms of φ and $\mathcal{G}_{M}^{\tau}\varphi$ have the same support, so that [\(3.31\)](#page-33-1) holds. Further exploiting that $\mathscr{C}_M^{\tau} \leq 1$, we obtain

$$
\|(\mathcal{G}^\tau_M \mathcal{A}_+^{\sharp,\tau}) \mathcal{G}^\tau_M \varphi\|_{\tau} \lesssim \nu_\tau^{3/4} \|\mathcal{G}^\tau_M \varphi\|_{\tau} \leq \nu_\tau^{3/4} \|\varphi\|_{\tau}
$$

which concludes the proof.

Remark 4.5 The bound [\(4.14\)](#page-37-2) shows in particular that, for φ smooth, s^{τ} is asymptotically equal to $\mathcal{L}_{M}^{\tau} \varphi$ in ΓL_{τ}^{2} , which means that s^{τ} is well-approximated by the $k = 0$ summand at the right hand side of [\(4.7\)](#page-36-0). In ΓH_{τ}^1 , the situation is drastically different: while the ΓH_{τ}^1 of the whole s^{τ} *does* converge to a positive constant for $\tau \to \infty$, and this (as can be seen from [\(4.13\)](#page-37-1); and this will play a crucial role in obtaining the limiting diffusion matrix), it can be shown that the ΓH_{τ}^1 norm of each individual summand in [\(4.7\)](#page-36-0) tends to zero with $\tau \to \infty$. In fact, one can prove that the summands that indeed contribute are those whose k diverges with τ as log log τ .

To conclude this section, we derive an apriori estimate on the difference of s^{τ} and r^{τ} in ΓH_{τ}^{1} (and thus also in ΓL_{τ}^2).

Proposition 4.6 *There exists a constant* $C = C(\lambda, M) > 0$ *such that for any* $\varphi \in \Gamma_{n_0} L^2_{\tau}$ *for some* $n_0 \geq 1$ *, we have*

$$
||r^{\tau} - s^{\tau}||_{\Gamma H^{1}_{\tau}} \leq C ||\mathcal{N}(-\nu_{\tau}\mathcal{L}_{0}^{\mathfrak{e}_{1}})^{1/2} s^{\tau}||_{\tau} + ||\mathcal{A}_{-}^{\sharp,\tau}\mathcal{L}_{M}^{\tau}\varphi||_{\tau}
$$
(4.16)

where r ^τ *and* s ^τ *are respectively given as in* [\(4.1\)](#page-35-0) *and* [\(4.7\)](#page-36-0)*, and* ^N *is the number operator in Definition [1.6.](#page-12-0)*

Remark 4.7 The constant C in [\(4.16\)](#page-38-0) can be taken to be proportional to $\lambda + \lambda^2 + M$.

Proof. Note that $r^{\tau} - s^{\tau} \in Dom(\mathcal{L}^{\tau})$, so that Lemma [2.10](#page-21-2) gives

$$
||r^{\tau} - s^{\tau}||^2_{\Gamma H^1_{\tau}} = \langle (1 - \mathcal{L}^{\tau})(r^{\tau} - s^{\tau}), r^{\tau} - s^{\tau} \rangle_{\tau}.
$$
 (4.17)

Thanks to the definition of r^{τ} and [\(4.9\)](#page-36-2), we can compute $(1 - \mathcal{L}^{\tau})(r^{\tau} - s^{\tau})$ and plug it into the above expression. What we obtain is the sum of three terms

$$
-\langle[(-\mathscr{A}_-^{\sharp,\tau})\mathscr{G}_M^\tau\mathscr{A}_+^{\sharp,\tau}-g_M^\tau]s^\tau,r^\tau-s^\tau\rangle_\tau+\langle\mathscr{A}^{\flat,\tau}s^\tau,r^\tau-s^\tau\rangle_\tau+\langle\mathscr{A}_-^{\sharp,\tau}\mathscr{G}_M^\tau\varphi,r^\tau-s^\tau\rangle_\tau
$$

 \Box

that we denote by (I) , (II) and (III) , and separately estimate. For the first, we apply the Replacement Lemma [3.6](#page-30-1) which gives

$$
|(\mathbf{I})| \le C \|(- \nu_\tau \mathcal{L}_0^{\mathfrak{e}_1})^{1/2} s^\tau \|_\tau \| r^\tau - s^\tau \|_{\Gamma H^1_\tau} \,. \tag{4.18}
$$

Proposition [3.4](#page-27-0) allows to control (II) as

$$
|(\text{II})| \lesssim \lambda \left\| \mathcal{N}(-\nu_\tau \mathcal{L}_0^{\epsilon_1})^{1/2} s^\tau \right\|_{\tau} \left\| r^\tau - s^\tau \right\|_{\Gamma H_\tau^1}.
$$
 (4.19)

At last, we simply use Cauchy-Schwarz on (III), thus obtaining

$$
|(\text{III})| \le ||\mathcal{A}_-^{\sharp,\tau}\mathcal{C}_M^{\tau}\varphi||_{\tau} ||r^{\tau} - s^{\tau}||_{\tau} \le ||\mathcal{A}_-^{\sharp,\tau}\mathcal{C}_M^{\tau}\varphi||_{\tau} ||r^{\tau} - s^{\tau}||_{\Gamma H^1_{\tau}}.
$$
 (4.20)

As a consequence, [\(4.16\)](#page-38-0) follows upon plugging [\(4.18\)](#page-39-0), [\(4.19\)](#page-39-1) and [\(4.20\)](#page-39-2) at the right hand side of [\(4.17\)](#page-38-1), and dividing both sides by $||r^{\tau} - s^{\tau}||_{\Gamma H^{1}_{\tau}}$. \Box

4.2 Estimating the anisotropic norm of the ansatz and proof of Theorem [4.1](#page-35-2)

For the estimates in the previous section to be effective and allow us to deduce Theorem [4.1,](#page-35-2) we need to control the $\|w(N)(-\nu_{\tau}\mathcal{L}_0^{\epsilon_1})^{1/2} \cdot \|_{\tau}$ -norm of the ansatz s^{τ} in [\(4.7\)](#page-36-0), for suitable weights $w: \mathbb{N} \to \mathbb{R}_+$, and prove that it is small as $\tau \to +\infty$. This is the content of the next proposition.

Proposition 4.8 *Assume that* [\(4.2\)](#page-35-1) *holds.* Let $w: \mathbb{N} \to \mathbb{R}_+$ be such that, for every $k \geq 0$ *,* $w(k + 1) \leq 2w(k)$. Then, for every $\varphi \in \Gamma_{n_0} L^2_{\tau}$ for some $n_0 \geq 1$,

$$
\| [w(\mathcal{N} - n_0)]^{1/2} \left(-\nu_\tau \mathcal{L}_0^{\mathfrak{e}_1} \right)^{1/2} s^{\tau} \|_{\tau} \leq 2 \| (-\mathcal{L}_0^{\mathfrak{e}_1} [f_w^{\tau}])^{1/2} \mathcal{L}_{M}^{\tau} \varphi \|_{\tau}, \tag{4.21}
$$

where

$$
f_w^{\tau}(x) \stackrel{\text{def}}{=} \nu_{\tau} \sum_{k \ge 0} w(k) \frac{1}{k!} \left(\log \sqrt{\frac{\nu_{\tau} + g^{\tau}(x)}{\nu_{\tau}}} \right)^k \tag{4.22}
$$

and g^{\dagger} is the function in [\(3.17\)](#page-29-1).

Remark 4.9 The reason why we imposed $w(\cdot + 1) \le 2w(\cdot)$ is to ensure that f_w in [\(4.22\)](#page-39-3) satisfies the assumptions of the Recursive Replacement Lemma [3.9](#page-32-0) with constant $C_{f_w} = w(0)$ as can be immediately checked via a direct computation. Notice also that, if w is non-decreasing then $\mathcal{I}_{\tau}(f_w) + w(0) \nu_{\tau} \leq f_w$ where \mathcal{I}_{τ} is the map given in [\(3.29\)](#page-32-1).

Remark 4.10 Even though for the sake of the present paper, it would have been sufficient to consider w given by the identity, we preferred to state Proposition [4.8](#page-39-4) in broader generality as it reveals deeper properties of the ansatz and partially, by [\(4.16\)](#page-38-0), of the resolvent [\(4.1\)](#page-35-0). Indeed, the bound [\(4.21\)](#page-39-5) shows that the anisotropic ΓH_{τ}^1 -norm of the former *decays in chaos exponentially fast* as can be checked by taking w to be $w(k) = c^k$ for $c \le 2$, in which case

$$
f_w^{\tau}(x) = \nu_{\tau}^{1-\frac{c}{2}} \left(\nu_{\tau} + g^{\tau}(x) \right)^{\frac{c}{2}}
$$
 (4.23)

and that, upon choosing $c < 2$, the weighted norm *vanishes as* $\tau \to \infty$, as one sees from [\(4.23\)](#page-39-6). Even though [\(4.16\)](#page-38-0) per se does not allow us to deduce the same at the level of the resolvent (and we will not need it), we suspect that this is indeed the case. Note that weighted (in \mathcal{N}) apriori bounds on the resolvent are highly non-trivial and, the only available ones, strongly rely on the graded sector condition (see [\[KLO12\]](#page-72-0)), which, as pointed out in Remark [3.5,](#page-28-0) does not hold in our context.

The proof of the above proposition is based on two main ingredients: the explicit iterative structure of the Ansatz s^{τ} and the Recursive Replacement Lemma [3.9.](#page-32-0) Before delving into the details, let us provide a brief heuristic that on the one hand shows how they come into play and on the other sheds some light on the role of the map \mathcal{I}_{τ} in [\(3.29\)](#page-32-1).

Thanks to [\(4.7\)](#page-36-0), for every $N \ge 0$, the $(N + n_0)$ -chaos component of s^{τ} is given by

$$
s^{\tau}_{N+n_0} = (\mathcal{G}^{\tau}_M \mathcal{A}^{\sharp,\tau}_+)^N \mathcal{G}^{\tau}_M \varphi \,,
$$

which implies

$$
\|(-\nu_{\tau}\mathcal{L}_{0}^{\epsilon_{1}})^{1/2}s_{N+n_{0}}^{\tau}\|_{\tau}^{2} = \|(-\nu_{\tau}\mathcal{L}_{0}^{\epsilon_{1}})^{1/2}(\mathcal{G}_{M}^{\tau}\mathcal{A}_{+}^{\sharp,\tau})^{N}\mathcal{G}_{M}^{\tau}\varphi\|_{\tau}^{2}
$$

$$
= \left\|\left[(-\mathcal{L}_{0}^{\epsilon_{1}}[\nu_{\tau}])^{1/2}\mathcal{G}_{M}^{\tau}\mathcal{A}_{+}^{\sharp,\tau}\right](\mathcal{G}_{M}^{\tau}\mathcal{A}_{+}^{\sharp,\tau})^{N-1}\mathcal{G}_{M}^{\tau}\varphi\right\|_{\tau}^{2} (4.24)
$$

where we have taken $w \equiv 1$ and used that, by [\(3.15\)](#page-29-2), $-\mathcal{L}_0^{\mathfrak{e}_1}[\nu_{\tau}] = -\nu_{\tau}\mathcal{L}_0^{\mathfrak{e}_1}$. To estimate the norm at the right hand side we apply [\(3.28\)](#page-32-2) with $f \equiv \nu_{\tau}$ and, neglecting the error for the sake of this heuristic, we obtain

$$
\|(-\nu_\tau \mathcal{L}_0^{\mathfrak{e}_1})^{1/2} s_{N+n_0}^{\tau} \|_\tau^2 \le \|(-\mathcal{L}_0^{\mathfrak{e}_1} [\mathcal{F}_\tau(\nu_\tau)])^{1/2} (\mathcal{G}_M^\tau \mathcal{A}_+^{\sharp,\tau})^{N-1} \mathcal{G}_M^\tau \varphi\|_\tau^2,
$$

where \mathcal{F}_{τ} is as in [\(3.29\)](#page-32-1) and the right hand side has precisely the same shape of [\(4.24\)](#page-40-0) but with $\mathcal{I}_{\tau}(\nu_{\tau})$ and $N-1$ in place of ν_{τ} and N. In other words, we can iterate the above procedure $N - 1$ times, ultimately getting

$$
\|(-\nu_{\tau}\mathcal{L}_{0}^{\epsilon_{1}})^{1/2}s_{N+n_{0}}^{\tau}\|_{\tau}^{2} \leq \|(-\mathcal{L}_{0}^{\epsilon_{1}}[\mathcal{J}_{\tau}^{\circ N}(\nu_{\tau})])^{1/2}\mathcal{G}_{M}^{\tau}\varphi\|_{\tau}^{2}
$$
(4.25)

for $\mathcal{F}_{\tau}^{\circ N}$ the N-fold composition of the map \mathcal{F}_{τ} with itself. On the other hand, $\mathcal{F}_{\tau}^{\circ N}(\nu_{\tau})$ is nothing but the N-step Picard iteration for the ODE

$$
\dot{y}(\ell) = \frac{1}{2} \frac{y(\ell)}{(\nu_{\tau} + g^{\tau}(\ell))^{3/2}}, \qquad y(0) = \nu_{\tau}, \qquad (4.26)
$$

(which is the linearisation of (3.19)) whose unique solution is

$$
y(x) = \nu_{\tau} \sqrt{\frac{g^{\tau}(x) + \nu_{\tau}}{\nu_{\tau}}} = \sqrt{\nu_{\tau}} \sqrt{g^{\tau}(x) + \nu_{\tau}}.
$$
 (4.27)

Remark 4.11 Here we see how crucial is the fact that the coefficient of the integral at the right hand side of (3.29) is $1/2 < 1$. Indeed, the $1/2$ therein corresponds to the square-root inside the logarithm in (4.22) which in turn results in the square-root in the second term of (4.27) . Since $g^{\tau}(x) + \nu_{\tau} \lesssim 1$, we get $y(x) \lesssim \nu_{\tau}/\sqrt{\nu_{\tau}} = \sqrt{\nu_{\tau}} \to 0$ thus suggesting that the $\|(-\nu_{\tau}\mathcal{L}_0^{\mathfrak{e}_1})^{1/2} \cdot \|_{\tau}$ norm of the $(N + n_0)$ -chaos component of the ansatz s^{τ} vanishes at rate $\nu_{\tau}^{1/4}$.

The Resolvent and its ansatz **42**

To make the previous argument rigorous, we need to be able to sum over N the bound (4.25) , introduce the dependence on the number operator via the weight w and take into account the error in [\(3.28\)](#page-32-2). This boils down to a stability type result for the Picard iteration of [\(4.26\)](#page-40-3) which reduces to understand how \mathcal{F}_{τ} iterates. This is precisely what the following lemma explains.

Lemma 4.12 *For* $k \in \mathbb{N}$, *let* $p_k^{\tau}: \mathbb{R}_+ \to \mathbb{R}_+$ *be the function given by*

$$
p_k^{\tau}(x) \stackrel{\text{def}}{=} \nu_{\tau} \frac{1}{k!} \left(\log \sqrt{\frac{\nu_{\tau} + g^{\tau}(x)}{\nu_{\tau}}} \right)^k, \tag{4.28}
$$

where g^{τ} is defined according to [\(3.17\)](#page-29-1). Then, p_k^{τ} is smooth, non-negative and non-decreasing *and, for all* $x > 0$ *, it holds*

$$
\sum_{k\geq 0} 2^k p_k^{\tau}(x) \leq \nu_{\tau} + g^{\tau}(x), \quad \sum_{k\geq 0} 2^k (p_k^{\tau})'(x) \leq \frac{2}{3} \frac{\nu_{\tau} + g^{\tau}(x)}{x}.
$$
 (4.29)

Furthermore, for every $k \geq 0$ *, we have*

$$
\mathcal{F}_{\tau}(p_k^{\tau}) = p_{k+1}^{\tau} \,. \tag{4.30}
$$

The statement follows by basic computations, and thus its proof is omitted. We now have all the elements we need in order to prove Proposition [4.8.](#page-39-4)

Proof of Proposition [4.8.](#page-39-4) We claim that for every $0 \le m \le N$ it holds

$$
\sum_{k=m}^{N} w(k) \| (-\nu_{\tau} \mathcal{L}_0^{\epsilon_1})^{1/2} s_{n_0+k}^{\tau} \|^2 \le \| (-\mathcal{L}_0^{\epsilon_1} [f_{m,N}])^{1/2} s_{n_0+m}^{\tau} \|^2 , \tag{4.31}
$$

where $f_{m,N} \stackrel{\text{def}}{=} 2 \sum_{k=0}^{N-m} w(k+m) p_k^{\tau}$ and p_k^{τ} is as in [\(4.28\)](#page-41-0). Given the claim, then [\(4.21\)](#page-39-5) follows by taking $m = 0$ and $N \to \infty$.

To prove the claim, we fix $N > 0$ and proceed by induction from $m = N$ to $m = 0$. The case $m = N$ is obvious, so assume [\(4.31\)](#page-41-1) holds for $m + 1$, with $m \in \{0, \ldots, N - 1\}$. Then, by the induction hypothesis, we have

$$
\sum_{k=m}^{N} w(k) \| (-\nu_{\tau} \mathcal{L}_{0}^{\epsilon_{1}})^{1/2} s_{n_{0}+k}^{\tau} \|^{2}
$$
\n
$$
\leq w(m) \| (-\nu_{\tau} \mathcal{L}_{0}^{\epsilon_{1}})^{1/2} s_{n_{0}+m}^{\tau} \|^{2} + \| (-\mathcal{L}_{0}^{\epsilon_{1}} [f_{m+1,N}])^{1/2} s_{n_{0}+m+1}^{\tau} \|^{2}
$$
\n
$$
= w(m) \| (-\nu_{\tau} \mathcal{L}_{0}^{\epsilon_{1}})^{1/2} s_{n_{0}+m}^{\tau} \|^{2} + \| \left[(-\mathcal{L}_{0}^{\epsilon_{1}} [f_{m+1,N}])^{1/2} \mathcal{L}_{M}^{\tau} s_{n}^{\tau} + \left| s_{n_{0}+m}^{\tau} \right| \right]^{2}.
$$
\n(4.32)

where in the last step we used [\(4.7\)](#page-36-0). In order to use the Recursive Replacement Lemma [3.9](#page-32-0) to bound the second term, we need to check that $f_{m+1,N}$ satisfies its assumptions. Since $w(\cdot + 1) \leq 2 w(\cdot)$, we have

$$
f_{m+1,N}(x) \le \sum_{k\ge 0} 2^{k+1} w(m) p_k^{\tau}(x) \le 2w(m) (\nu_{\tau} + g^{\tau}(x)), \quad \text{for all } x > 0,
$$

the last bound being a consequence of [\(4.29\)](#page-41-2), and similarly for $f'_{m+1,N}$. Hence, [\(3.27\)](#page-32-3) holds for $f_{m+1,N}$ with $C_{f_{m+1,N}} = 2w(m)$. Thus, plugging [\(3.28\)](#page-32-2) into [\(4.32\)](#page-41-3), we deduce

$$
\sum_{k=m}^{N} w(k) \| (-\nu_{\tau} \mathcal{L}_{0}^{\mathfrak{e}_{1}})^{1/2} s_{n_{0}+k}^{\tau} \|^{2}
$$
\n
$$
\leq w(m) \| (-\nu_{\tau} \mathcal{L}_{0}^{\mathfrak{e}_{1}})^{1/2} s_{n_{0}+m}^{\tau} \|^{2} + \| (-\mathcal{L}_{0}^{\mathfrak{e}_{1}} [\mathcal{J}_{\tau}(f_{m+1,N}) + \nu_{\tau} w(m)])^{1/2} s_{n_{0}+m}^{\tau} \|^{2}
$$
\n
$$
= \| (-\mathcal{L}_{0}^{\mathfrak{e}_{1}} [\mathcal{J}_{\tau}(f_{m+1,N}) + 2\nu_{\tau} w(m)])^{1/2} s_{n_{0}+m}^{\tau} \|^{2},
$$

where we also used [\(4.2\)](#page-35-1) to bound $\nu_{\tau} C_{f_{m+1,N}} C_{M,\tau}^{(2)} \le \nu_{\tau} w(m)$. But now, thanks to [\(4.30\)](#page-41-4), $\mathcal{I}_{\tau}(f_{m+1,N}) + 2\nu_{\tau}w(m) = f_{m,N}$, which concludes the proof of the induction and therefore of $(4.21).$ $(4.21).$ \Box

Thanks to the previous proposition and the results in the previous subsection, we can now prove Theorem [4.1.](#page-35-2)

Proof of Theorem [4.1.](#page-35-2) We will first prove [\(4.3\)](#page-35-3) and [\(4.4\)](#page-35-4), for which we take $\varphi \in \Gamma_{n_0} L^2_{\tau}$, for some $n_0 \geq 1$. Notice first that

$$
||r^{\tau}||_{\tau} \leq ||r^{\tau} - s^{\tau}||_{\Gamma H_{\tau}^{1}} + ||s^{\tau}||_{\tau} \lesssim ||\mathcal{N}(-\nu_{\tau}\mathcal{L}_{0}^{\mathfrak{e}_{1}})^{1/2}s^{\tau}||_{\tau} + ||\mathcal{A}_{-}^{\sharp,\tau}\mathcal{L}_{M}^{\tau}\varphi||_{\tau} + ||\mathcal{L}_{M}^{\tau}\varphi||_{\tau}
$$
(4.33)

where in the last step we used [\(4.16\)](#page-38-0) and [\(4.12\)](#page-37-3). The triangle inequality together with the basic estimate $a \le \sqrt{|a^2 - b^2|} + b$, which holds for any $a, b > 0$, implies

$$
\begin{aligned} \left| \|r^{\tau}\|_{\Gamma H^1_{\tau}}^2 - \|(\mathfrak{E}^{\tau}_M)^{1/2}\varphi\|_{\tau}^2 \right| &\leq \left| \|r^{\tau}\|_{\Gamma H^1_{\tau}}^2 - \|s^{\tau}\|_{\Gamma H^1_{\tau}}^2 \right| + \left| \|s^{\tau}\|_{\Gamma H^1_{\tau}}^2 - \|(\mathfrak{E}^{\tau}_M)^{1/2}\varphi\|_{\tau}^2 \right| \\ &\leq \|r^{\tau} - s^{\tau}\|_{\Gamma H^1_{\tau}} (\|r^{\tau} - s^{\tau}\|_{\Gamma H^1_{\tau}} + 2\|s^{\tau}\|_{\Gamma H^1_{\tau}}) + \left| \|s^{\tau}\|_{\Gamma H^1_{\tau}}^2 - \|(\mathfrak{E}^{\tau}_M)^{1/2}\varphi\|_{\tau}^2 \right| \\ &\lesssim \left(\|r^{\tau} - s^{\tau}\|_{\Gamma H^1_{\tau}} + \sqrt{|\|s^{\tau}\|_{\Gamma H^1_{\tau}}^2 - \|(\mathfrak{E}^{\tau}_M)^{1/2}\varphi\|_{\tau}^2}| \right)^2 + \|r^{\tau} - s^{\tau}\|_{\Gamma H^1_{\tau}} \|(\mathfrak{E}^{\tau}_M)^{1/2}\varphi\|_{\tau} \end{aligned}
$$

and thus, by (4.16) and (4.13) , we deduce

$$
\left| \|r^{\tau}\|_{\Gamma H^1_{\tau}}^2 - \|(\mathcal{E}_M^{\tau})^{1/2} \varphi\|_{\tau}^2 \right| \lesssim \left(\|\mathcal{N}(-\nu_{\tau}\mathcal{L}_0^{\mathfrak{e}_1})^{1/2} s^{\tau}\|_{\tau} + \|\mathcal{A}_-^{\sharp,\tau}\mathcal{E}_M^{\tau}\varphi\|_{\tau} \right) \times \left(4.34 \right) \times \left(\|\mathcal{N}(-\nu_{\tau}\mathcal{L}_0^{\mathfrak{e}_1})^{1/2} s^{\tau}\|_{\tau} + \|\mathcal{A}_-^{\sharp,\tau}\mathcal{E}_M^{\tau}\varphi\|_{\tau} + \|(\mathcal{E}_M^{\tau})^{1/2} \varphi\|_{\tau} \right).
$$

As a consequence, we are left to estimate the weighted norm of s^{τ} . Upon taking $w : \mathbb{N} \to \mathbb{R}_+$ to be defined as $w(k) = n_0 + k$, Proposition [4.8](#page-39-4) gives

$$
\| [w(\mathcal{N} - n_0)]^{1/2} \left(-\nu_\tau \mathcal{L}_0^{\mathfrak{e}_1} \right)^{1/2} s^{\tau} \|_{\tau} \leq 2 \| (-\mathcal{L}_0^{\mathfrak{e}_1} [f_w^{\tau}] \right)^{1/2} \mathcal{L}_{M}^{\tau} \varphi \|_{\tau}
$$

$$
\lesssim \nu_\tau^{1/4} \sqrt{|\log \nu_\tau|} \| (-\mathcal{L}_0^{\mathfrak{e}_1})^{1/2} \mathcal{L}_{M}^{\tau} \varphi \|_{\tau}
$$
 (4.35)

which holds since the function $f_w^{\tau}(L^{\tau})$, for f_w^{τ} as in [\(4.22\)](#page-39-3) can be bounded uniformly in $x > 0$ as

$$
f_w^{\tau}(L^{\tau}(x)) \lesssim \nu_{\tau}^{1/2} \sqrt{\nu_{\tau} + g^{\tau}(L^{\tau}(x))} \max \left\{ n_0 \vee \log \sqrt{\frac{\nu_{\tau} + g^{\tau}(L^{\tau}(x))}{\nu_{\tau}}} \right\}
$$

The superdiffusive Central Limit Theorem **44**

$$
\lesssim \nu_\tau^{1/2} \max\left\{n_0 \vee \log \nu_\tau^{-1/2}\right\} \lesssim_{n_0} \nu_\tau^{1/2} |\log \nu_\tau|
$$

where we used that L^{τ} is bounded uniformly in x, and consequently so is $g^{\tau}(L^{\tau})$, and, in the last step, that n_0 is fixed. By plugging [\(4.35\)](#page-42-0) into both [\(4.33\)](#page-42-1) and [\(4.34\)](#page-42-2) and recalling the definition of the seminorm in (4.5) , (4.3) and (4.4) follow.

We now turn to (4.6) , for which we proceed similarly to what was done in (4.33) . Assume φ is such that Supp(φ) $\subset \{-8^\tau \leq \tau^\alpha\}$ for some $\alpha \in (0,1)$. Then, Proposition [4.6](#page-38-2) and equations (4.35) and (4.14) give

$$
\begin{aligned}\|r^{\tau}-\mathcal{G}_M^{\tau}\varphi\|_{\tau}&\leq \|r^{\tau}-s^{\tau}\|_{\Gamma H^1_{\tau}}+\|s^{\tau}-\mathcal{G}_M^{\tau}\varphi\|_{\tau}\\ &\lesssim \nu_{\tau}^{1/4}\sqrt{|\log \nu_{\tau}|}\|(-\mathcal{L}_0^{\mathfrak{e}_1})^{1/2}\mathcal{G}_M^{\tau}\varphi\|_{\tau}+\|\mathcal{A}_-^{\sharp,\tau}\mathcal{G}_M^{\tau}\varphi\|_{\tau}+\nu_{\tau}^{3/4}\|\varphi\|_{\tau}\\ &\lesssim \nu_{\tau}^{1/4}\sqrt{|\log \nu_{\tau}|}\|(-\mathcal{L}_0^{\mathfrak{e}_1})^{1/2}\mathcal{G}_M^{\tau}\varphi\|_{\tau}+\nu_{\tau}^{3/4}\|\varphi\|_{\tau}\end{aligned}
$$

where the last step follows by [\(3.31\)](#page-33-1). To conclude, notice that, since \mathcal{C}_{M}^{τ} is diagonal, then $\text{Supp}(\mathcal{F}(\mathcal{G}_M^{\tau}\varphi)) = \text{Supp}(\hat{\varphi}) \subset \{-\mathcal{S}^{\tau} \leq \tau^{\alpha}\}\$ and on the latter set we have

$$
g_M^{\tau}(L^{\tau}(-S^{\tau})) \ge g_M^{\tau}(L^{\tau}(\tau^{\alpha})) \gtrsim_{\alpha} 1.
$$

Therefore, $(-\mathcal{L}_0^{\mathfrak{e}_1})^{1/2}\mathcal{G}_M^{\tau} \lesssim_{\alpha} (\mathcal{G}_M^{\tau})^{1/2} \leq 1$, from which we get [\(4.6\)](#page-35-6) and thus conclude the proof of the statement.

5 The superdiffusive Central Limit Theorem

The goal of this and the next section is to show the main results of this paper, which, as we will soon see, are direct consequences of the analysis carried out in the previous.

Here, we focus on Theorems [1.4](#page-7-0) and [1.3.](#page-6-0) As mentioned before, for the former we will exploit (a version of) the Trotter-Kato theorem which requires to show convergence of the resolvent $(1 - \mathcal{L}^{\tau})^{-1}$ to $(1 - \mathcal{L}^{\text{eff}})^{-1}$. This is the content of the next Theorem which addresses point [\(IV\)](#page-24-0) of Section [2.3.](#page-21-0) Before stating it, recall that, as mentioned in Section [2,](#page-13-0) the two resolvents are defined on different spaces, namely $\mathbb{L}^{\theta}(\mathbb{P}^{\tau})$ and $\mathbb{L}^{\theta}(\mathbb{P})$, $\theta \in [1,\infty)$, which can be mapped to one another thanks to the surjective isometries ι^{τ} , j^{τ} introduced in Lemma [1.7.](#page-12-1)

Theorem 5.1 *For every* $F \in \mathbb{L}^2(\mathbb{P})$ *, we have*

$$
\iota^{\tau} (1 - \mathcal{L}^{\tau})^{-1} j^{\tau} F \xrightarrow[\tau \to \infty]{\mathbb{L}^2(\mathbb{P})} (1 - \mathcal{L}^{\text{eff}})^{-1} F , \qquad (5.1)
$$

where $ι^τ$ *and* $j^τ$ *are the maps defined in Lemma* [1.7](#page-12-1) *and* \mathcal{L}^{eff} *is the operator introduced in Lemma [2.11.](#page-21-3)*

Proof. By Lemmas [1.7](#page-12-1) and [2.6,](#page-19-0) and Lemma [2.11,](#page-21-3) the operators $\iota^{\tau} (1 - \mathcal{L}^{\tau})^{-1} j^{\tau}$ and $(1 - \mathcal{L}^{\text{eff}})^{-1}$ are bounded, uniformly in τ , from $\mathbb{L}^2(\mathbb{P})$ to itself, so that it suffices to verify [\(5.1\)](#page-43-0) for F belonging to a subset spanning a dense subspace of $\mathbb{L}^2(\mathbb{P})$, e.g. the set of F of the form $I_n^{\tau}(\varphi)$ with $n \in \mathbb{N}$ and $\varphi \in \Gamma_n L^2$ such that $\mathcal{F}\varphi$ has compact support (recall the notation I^{τ} from Section [1.3](#page-11-0) for

The superdiffusive Central Limit Theorem **45**

the isometry between Fock space and $L^2(\mathbb{P})$). Hence, we are left to verify that for such n and φ , we have

$$
\iota^{\tau} (1 - \mathcal{L}^{\tau})^{-1} j^{\tau} \varphi \xrightarrow[\tau \to \infty]{\mathbb{L}^{2}(\mathbb{P})} (1 - \mathcal{L}^{\text{eff}})^{-1} \varphi.
$$
 (5.2)

Notice that, by [\(1.22\)](#page-12-2), for any (bounded) diagonal operator \mathfrak{D} acting on ΓL^2_τ (see Definition [1.5\)](#page-12-3), the diagonal operator $\iota^T \mathcal{D} j^T$ acting on ΓL^2_{∞} has the same kernel as \mathcal{D} , so that, with a slight abuse of notations, we denote it by $\iota^T \mathfrak{D} j^T = \mathfrak{D}$. Then we have

$$
\begin{aligned} \|\iota^{\tau}(1-\mathcal{L}^{\tau})^{-1}j^{\tau}\varphi-(1-\mathcal{L}^{\text{eff}})^{-1}\varphi\|\\ \leq\|\iota^{\tau}(1-\mathcal{L}^{\tau})^{-1}j^{\tau}\varphi-\iota^{\tau}\mathcal{G}_{M}^{\tau}j^{\tau}\varphi\|+\|\mathcal{G}_{M}^{\tau}\varphi-(1-\mathcal{L}^{\text{eff}})^{-1}\varphi\|\,. \end{aligned}
$$

By [\(4.14\)](#page-37-2), the first term goes to 0 as $\tau \to +\infty$. From the explicit expression of the kernels of \mathfrak{G}_M^{τ} and \mathcal{L}^{eff} , it is easy to see that also the second term vanishes as $\tau \to +\infty$, so that the proof is complete.

Thanks to the previous result, we are now ready to prove Theorems [1.4](#page-7-0) and [1.3.](#page-6-0)

Proof of Theorems [1.4](#page-7-0) and [1.3.](#page-6-0) Let us immediately say that, given Theorem [1.4,](#page-7-0) Theorem [1.3](#page-6-0) follows immediately by taking $F_i^{\tau}(\eta^{\tau}) \stackrel{\text{def}}{=} f_i(\eta^{\tau}(\varphi_i))$, for $i = 1, \ldots, k, \varphi_1, \ldots, \varphi_k \in \mathcal{S}(\mathbb{R}^2)$ and f_1, \ldots, f_n bounded continuous functions on \mathbb{R} .

Turning to Theorem [1.4,](#page-7-0) we split its proof in two parts. First, we will show strong convergence of the semigroup, and more precisely that [\(1.11\)](#page-7-1) holds for any $\theta \in [1,\infty)$. Then, we will use the latter to deduce the convergence in [\(1.12\)](#page-7-2).

Part I: Convergence of the semigroup. Thanks to Lemma [1.7](#page-12-1) and Theorem [5.1,](#page-43-1) we are precisely in the setting of the Trotter-Kato theorem [\[IK02,](#page-72-1) Theorem 4.2], which implies that for every $F \in \mathbb{L}^2(\mathbb{P})$ and $T > 0$, it holds

$$
\sup_{t\in[0,T]}\|t^{\tau}P_t^{\tau}j^{\tau}F-P_t^{\text{eff}}F\|_{\mathbb{L}^2(\mathbb{P})}\xrightarrow[\tau\to+\infty]{}0.
$$

This can be rewritten in the following form. Let $(F_\tau)_\tau$ be sequence such that $F_\tau \in L^2(\mathbb{P}^\tau)$ and $||\iota^\tau F_\tau - F||_{\mathbb{L}^2(\mathbb{P})} \to 0$ as $\tau \to +\infty$. Then noticing that $||\iota^\tau F_\tau - F||_{\mathbb{L}^2(\mathbb{P})} = ||F^\tau - j^\tau F||_{\mathbb{L}^2(\mathbb{P}^\tau)}$ and using the contractivity of the semi-groups stated in Lemmas [2.6](#page-19-0) and [2.11\)](#page-21-3), we also deduce that

$$
\sup_{t \in [0,T]} \|t^{\tau} P_t^{\tau} F^{\tau} - P_t^{\text{eff}} F\|_{\mathbb{L}^2(\mathbb{P})} \xrightarrow[\tau \to +\infty]{} 0. \tag{5.3}
$$

Hence, we are left to extend the above to $T = \infty$ and any value of $\theta \in [1, \infty)$. Let us begin with the former.

Without loss of generality, we can take F^{τ} and F with mean zero. Let $T > 0$ and note that, for any $t \geq T$, the contractivity of P^{τ} and P^{eff} implies

$$
\| \iota^{\tau} P_t^{\tau} F^{\tau} - P_t^{\text{eff}} F \|_{\mathbb{L}^2(\mathbb{P})} \leq \| P_t^{\tau} F^{\tau} \|_{\mathbb{L}^2(\mathbb{P}^{\tau})} + \| P_t^{\text{eff}} F \|_{\mathbb{L}^2(\mathbb{P})}
$$

\n
$$
\leq \| P_T^{\tau} F^{\tau} \|_{\mathbb{L}^2(\mathbb{P}^{\tau})} + \| P_T^{\text{eff}} F \|_{\mathbb{L}^2(\mathbb{P})}
$$

\n
$$
\leq \| \iota^{\tau} P_T^{\tau} F^{\tau} - P_T^{\text{eff}} F \|_{\mathbb{L}^2(\mathbb{P})} + 2 \| P_T^{\text{eff}} F \|_{\mathbb{L}^2(\mathbb{P})},
$$

where we also used that ι^{τ} is an isometry. Therefore,

$$
\sup_{t\geq 0} \| \iota^\tau P_t^\tau F^\tau - P_t^{\text{eff}} F \|_{\mathbb{L}^2(\mathbb{P})} \leq \sup_{t\in[0,T]} \| \iota^\tau P_t^\tau F^\tau - P_t^{\text{eff}} F \|_{\mathbb{L}^2(\mathbb{P})} + 2 \| P_T^{\text{eff}} F \|_{\mathbb{L}^2(\mathbb{P})}.
$$

We can now take the limsup as $\tau \to +\infty$ at both sides so that, at the right hand side, the first term vanishes and what is left goes to 0 in the $T \to \infty$ limit by [\(2.32\)](#page-21-4). Hence, [\(1.11\)](#page-7-1) holds for $\theta = 2.$

To prove that it also holds for $\theta \in [1,\infty)$, let $(F^{\tau})_{\tau}$ be such that $F^{\tau} \in \mathbb{L}^{\theta}(\mathbb{P}^{\tau})$ and $\|\iota^{\tau} F^{\tau} F\|_{\mathbb{F}^{\theta(\mathbb{P})}} \to 0$ as $\tau \to +\infty$. Applying [\(2.21\)](#page-19-1) with θ therein equal to $+\infty$, it is immediate to see that [\(1.11\)](#page-7-1) for $\theta = 2$ extends to $\theta \in [1, \infty)$ if F^{τ} (and F) is bounded by a constant, uniformly in τ . For general F^{τ} and F , let $M > 0$ and $\chi_M(x) \stackrel{\text{def}}{=} (-M) \vee (M \wedge x)$. Clearly, $\chi^M(F^{\tau})$ and $\chi^M(F)$ are bounded and satisfy $\| \iota^\tau \chi^M(F^\tau) - \chi^M(F) \|_{\mathbb{L}^\theta(\mathbb{P})} \to 0$ as $\tau \to +\infty$. Using once again the contractivity of the semigroups, we obtain the bound

$$
\| \iota^{\tau} P_t^{\tau} F^{\tau} - P_t^{\text{eff}} F \|_{\mathbb{L}^{\theta}(\mathbb{P})} \leq \| \iota^{\tau} P_t^{\tau} \chi^M(F^{\tau}) - P_t^{\text{eff}} \chi^M(F) \|_{\mathbb{L}^{\theta}(\mathbb{P})} + 2 \| F - \chi^M(F) \|_{\mathbb{L}^{\theta}(\mathbb{P})} + \| \iota^{\tau} F^{\tau} - F \|_{\mathbb{L}^{\theta}(\mathbb{P})},
$$

and we take the supremum in $t \geq 0$. Now, we can pass to the limit first in $\tau \to +\infty$, so that the first and last terms vanish (the former since $\chi^M(F^{\tau})$ and $\chi^M(F)$ are bounded and the latter by construction) and then in $M \to \infty$ which ensures that also the second term converges to 0. Therefore, [\(1.11\)](#page-7-1) holds for any $\theta \in [1, \infty)$.

Part II: Convergence of the finite-dimensional laws. We first rewrite the conditional expectations in [\(1.12\)](#page-7-2). To do so, let $0 = t_0 \le t_1 \le \cdots \le t_k$ and set $\Delta t_i \stackrel{\text{def}}{=} t_i - t_{i-1}$ for $i \in \{1, ..., k\}$. Define recursively the functionals G_i^{τ} and G_i^{eff} , $i \in \{1, ..., k+1\}$, according to $G_{k+1}^{\tau} = G_{k+1}^{\text{eff}} \stackrel{\text{def}}{=} 1$ and for every $i \in \{1, ..., k\}$

$$
G_i^{\tau} = P_{\Delta t_i}^{\tau} (F_i^{\tau} G_{i+1}^{\tau}) \qquad \text{and} \qquad G_i^{\text{eff}} = P_{\Delta t_i}^{\text{eff}} (F_i G_{i+1}^{\text{eff}}).
$$
 (5.4)

Note that both $\bigcap_{\theta \geq 1} \mathbb{L}^{\theta}(\mathbb{P}^{\tau})$ and $\bigcap_{\theta \geq 1} \mathbb{L}^{\theta}(\mathbb{P})$ are stable under multiplication and the action of the respective semigroups. Thus, the above recursion is well-defined and, for every $i \in \{1, \ldots, k\}$, $G_i^{\tau} \in \bigcap_{\theta \geq 1} \mathbb{L}^{\theta}(\mathbb{P}^{\tau})$ and $G_i^{\text{eff}} \in \bigcap_{\theta \geq 1} \mathbb{L}^{\theta}(\mathbb{P})$. The Markov property implies that the two terms in (1.12) can be written as

$$
\mathbf{E}_{u_0}^{\tau}[F_1^{\tau}(u_{t_1}^{\tau})...F_k^{\tau}(u_{t_k}^{\tau})] = \iota^{\tau}G_1^{\tau}(u_0) \text{ and } \mathbf{E}_{u_0}^{\text{eff}}[F_1(u_{t_1}^{\text{eff}})...F_k(u_{t_k}^{\text{eff}})] = G_1^{\text{eff}}(u_0). \quad (5.5)
$$

Therefore, if we show that for every $i \in \{1,\ldots,k+1\}$ it holds $||\iota^\tau G_i^\tau - G_i^{\text{eff}}||_{\mathbb{L}^{\theta}(\mathbb{P})} \to 0$ as $\tau \to +\infty$, for every $\theta \in [1,\infty)$, then [\(1.12\)](#page-7-2) follows at once. To prove it, we proceed by (backward) induction. The result is obvious for $i = k + 1$, since $\iota^{\tau} G_{k+1}^{\tau} = 1 = G_{k+1}^{\text{eff}}$. Assume now it holds for $i + 1$, and let us verify the same is true for i. Since by assumption $||\iota^\tau F_i^\tau - F_i||_{\mathbb{L}^{\theta}(\mathbb{P})} \to 0$ for all $\theta \in [1, \infty)$, and, by induction, the same holds with G_{i+1}^τ and G_{i+1}^{eff} in place of F_i^{τ} and F_i , the same must be true for their product. By applying [\(1.11\)](#page-7-1), we obtain

$$
G_i^{\tau} = P_{\Delta t_i}^{\tau}(F_i^{\tau} G_{i+1}^{\tau}) \xrightarrow[\tau \to \infty]{\mathbb{L}^{\theta}(\mathbb{P})} P_{\Delta t_i}^{\text{eff}}(F_i G_{i+1}^{\text{eff}}) = G_i^{\text{eff}}
$$

which completes the proof.

6 The large-scale behaviour of the Diffusion Matrix

In this final section, we show Theorem [1.1,](#page-4-0) whose proof amounts to determine the asymptotic behaviour of the diffusion matrix $t \mapsto D(t)$ in [\(1.4\)](#page-4-1) as $t \to \infty$.

To do so, let us first make a couple of remarks. By Proposition [2.5](#page-18-0) (for $\tau = 1$), the only non-trivial entry of D is $D_{1,1}$ which can be written in terms of the rescaled process u^{τ} as

$$
D_{1,1}^{\tau}(s) = \nu_{\tau} D_{1,1}(\tau s), \qquad s \ge 0.
$$
 (6.1)

In particular, this means that Theorem [1.1](#page-4-0) follows provided we show that, for $s = 1$, the left hand side converges as $\tau \to \infty$ to $C_{\text{eff}}(\lambda)$ in [\(1.6\)](#page-4-2). Now, before that, we need to manipulate the Green-Kubo formula in [\(2.18\)](#page-18-1) so to unveil the connection with the results in the previous section.

6.1 The diffusivity as the energy associated to a parabolic problem

In order to simplify our analysis and ensure that all the quantities under study are well-defined, we approximate the spatial integral at the right hand side of [\(2.18\)](#page-18-1) in such a way that the functional \mathcal{M}^{τ} in [\(2.7\)](#page-15-0) is written in terms of the operator \mathcal{A}^{τ}_+ . Loosely speaking, $\mathcal{M}^{\tau}[\eta^{\tau}]$ is the antiderivative in the \mathfrak{e}_1 direction of the nonlinearity $\mathcal{N}^\tau[\eta^\tau]$, which, when tested against a test function $\psi \in \mathcal{S}(\mathbb{R}^2) \cap \Gamma_1 L^2_\tau$, satisfies $\mathcal{N}^\tau[\eta^\tau](\psi) = \mathcal{A}^\tau_+\eta^\tau(\psi)$. Thus, we would like to say that $M^{\tau}[\eta^{\tau}](\psi)$ is $\mathcal{A}_{+}^{\tau}\eta^{\tau}(\partial_{\mathfrak{e}_1}^{-1}\psi)$, but this only makes sense provided $\text{Supp}(\hat{\psi}) \cap (\mathfrak{e}_2 \mathbb{R}) = \emptyset$.

To do so, let us fix an even symmetric function $\psi \in \mathcal{S}(\mathbb{R}^2)$ such that $\|\psi\|_{L^2(\mathbb{R}^2)} = 1$ and whose Fourier transform $\hat{\psi}$ has compact support in a ball of radius 1 around the origin contained in $(\mathfrak{e}_2 \mathbb{R})^c$, with $\mathfrak{e}_2 \mathbb{R}$ denoting the vertical axis. For $\delta > 0$, set $\psi^{\delta}(\cdot) \stackrel{\text{def}}{=} \delta \psi(\delta \cdot)$, so that the convolution of ψ^{δ} with itself converges to 1 as $\delta \to 0$. Then, we hope that

$$
\int_{\mathbb{R}^2} \mathbf{E}[\mathcal{M}^\tau[u_r^\tau](x) \mathcal{M}^\tau[u_0^\tau](0)] (\psi^\delta * \psi^\delta)(x) dx \stackrel{\delta \to 0}{\sim} \int_{\mathbb{R}^2} \mathbf{E}[\mathcal{M}^\tau[u_r^\tau](x) \mathcal{M}^\tau[u_0^\tau](0)] dx. (6.2)
$$

The advantage of the left hand side is that, by translation invariance and the symmetry of ψ^{δ} , it equals

$$
\int_{\mathbb{R}^4} \mathbf{E}[\mathcal{M}^\tau[u_r^\tau](x - y)\mathcal{M}^\tau[u_0^\tau](-y)] \psi^\delta(x - y)\psi^\delta(y) \mathrm{d}x \mathrm{d}y
$$
\n
$$
= \mathbf{E}[\mathcal{M}^\tau[u_r^\tau](\psi^\delta)\mathcal{M}^\tau[u_0^\tau](\psi^\delta)] = \frac{1}{\lambda^2 \nu_\tau^{3/2}} \langle P_r^\tau \mathfrak{m}^{\tau,\delta}, \mathfrak{m}^{\tau,\delta} \rangle_\tau \tag{6.3}
$$

where in the last step we denoted by $\mathfrak{m}^{\tau,\delta} \in \Gamma_2 L_\tau^2 \cap \mathbf{\Gamma} \mathbf{H}_\tau$ the kernel of $\lambda \nu_\tau^{3/4} \mathcal{M}^{\tau,\delta}$, and P^τ is the semigroup of the rescaled SBE. Now, since by construction $\text{Supp}(\hat{\psi}^{\delta}) \subset (\mathfrak{e}_2 \mathbb{R})^c$, $\partial_{\mathfrak{e}_1}^{-1} \psi^{\delta}$ is well-defined and a direct computation shows that

$$
\mathfrak{m}^{\tau,\delta} = \mathfrak{A}_+^{\tau} \partial_{\mathfrak{e}_1}^{-1} \psi^{\delta} \,. \tag{6.4}
$$

In the next lemma we show that the Green-Kubo formula [\(2.18\)](#page-18-1) can be expressed in the form of the energy of the solution of the parabolic equation given by

$$
(\partial_t - \mathcal{L}^\tau) \varphi^{\tau,\delta} = \mathfrak{m}^{\tau,\delta}, \qquad \varphi^{\tau,\delta}|_{t=0} = 0 \tag{6.5}
$$

which is explicit and given by

$$
\varphi_t^{\tau,\delta} = \int_0^t P_s^{\tau} \mathfrak{m}^{\tau,\delta} \, \mathrm{d}s \,, \qquad t \ge 0. \tag{6.6}
$$

Proposition 6.1 *For any* $t, \tau > 0$ *fixed, the following equality holds*

$$
D_{1,1}^{\tau}(t) = \nu_{\tau} + \lim_{\delta \to 0} \frac{\mathcal{E}_t^{\tau}(\varphi^{\tau,\delta})}{t}.
$$
 (6.7)

where, for any $\delta > 0$, $\varphi^{\delta, \tau}$ is the unique solution to (6.5) and, for any $(f_s)_{s \geq 0} \in C^1([0, t], \Gamma L^2_\tau) \cap$ $C([0, t], Dom(\mathcal{L}^{\tau}))$, the positive quadratic form \mathcal{E}^{τ}_t is defined as

$$
\mathcal{E}_t^{\tau}(f) \stackrel{\text{def}}{=} \|f_t\|_{\tau}^2 + 2 \int_0^t \|(-\mathcal{S}^{\tau})^{1/2} f_s\|_{\tau}^2 ds = \|f_0\|_{\tau}^2 + 2 \int_0^t \langle (\partial_s - \mathcal{L}^{\tau}) f_s, f_s \rangle_{\tau} ds. \tag{6.8}
$$

Proof. Let us first briefly argue the second equality in [\(6.8\)](#page-47-0). A proof similar to that of Lemma [2.10,](#page-21-2) gives that for every $\varphi \in \text{Dom}(\mathscr{L}^{\tau})$, $\|(-\mathscr{S}^{\tau})^{1/2}\varphi\|_{\tau}^{2} = \langle (-\mathscr{L}^{\tau})\varphi, \varphi \rangle_{\tau}$. Therefore,

$$
\int_0^t \langle (\partial_s - \mathcal{L}^\tau) f_s, f_s \rangle_\tau \, \mathrm{d} s = \int_0^t \langle \partial_s f_s, f_s \rangle_\tau \, \mathrm{d} s + \int_0^t \langle (-\mathcal{L}^\tau) f_s, f_s \rangle_\tau \, \mathrm{d} s
$$

from which the conclusion follows by integration by parts.

Next, we address [\(6.7\)](#page-47-1), which essentially boils down to show that [\(6.2\)](#page-46-1) holds. For this, note that $|\psi^{\delta} * \psi^{\delta}(x) - 1| = |\psi * \psi(\delta x) - \psi * \psi(0)| \lesssim \delta |x|$ and, thus, the polynomial decay of correlations in [\(2.6\)](#page-14-0) implies

$$
\left|\mathbf{E}[\langle \mathcal{M}^{\tau}[u_{r}^{\tau}], \psi^{\delta} \rangle \langle \mathcal{M}^{\tau}[u_{0}^{\tau}], \psi^{\delta} \rangle] - \int_{\mathbb{R}^{2}} \mathbf{E}[\mathcal{M}^{\tau}[u_{r}^{\tau}](x) \mathcal{M}^{\tau}[u_{0}^{\tau}](0)] dx \right| \lesssim_{\tau} \delta.
$$

Together with (2.18) and (6.3) , the previous gives

$$
D_{1,1}^{\tau}(t) = \nu_{\tau} + \lim_{\delta \to 0} \frac{2\lambda^2 \nu_{\tau}^{3/2}}{t} \int_0^t ds \int_0^s dr \mathbf{E}[\langle \mathcal{M}^{\tau}[u_{r}^{\tau}], \psi^{\delta} \rangle \langle \mathcal{M}^{\tau}[u_{0}^{\tau}], \psi^{\delta} \rangle]
$$

= $\nu_{\tau} + \lim_{\delta \to 0} \frac{2}{t} \int_0^t ds \int_0^s dr \langle P_r^{\tau} \mathbf{m}^{\tau, \delta}, \mathbf{m}^{\tau, \delta} \rangle_{\tau} = \nu_{\tau} + \lim_{\delta \to 0} \frac{2}{t} \int_0^t ds \langle \varphi^{\tau, \delta}, \mathbf{m}^{\tau, \delta} \rangle_{\tau}$

where in the last step we used [\(6.6\)](#page-47-2). Since by [\(6.5\)](#page-46-0), $\mathfrak{m}^{\tau,\delta} = (\partial_t - \mathcal{L}^{\tau}) \varphi^{\tau,\delta}$, the second equality of [\(6.8\)](#page-47-0) implies [\(6.7\)](#page-47-1) and the proof of the statement is complete. \Box

To combine the previous proposition with the results from Section [4,](#page-35-7) we need to understand the relation between the parabolic equation in (6.5) and the elliptic one given by

$$
(1 - \mathcal{L}^{\tau}) r^{\tau, \delta} = \mathfrak{m}^{\tau, \delta}.
$$
 (6.9)

This is the content of the next (rather straightforward) lemma^{[5](#page-47-3)}.

⁵See also [\[CMOW22,](#page-70-0) Section 9], for a similar a similar analysis in a different context.

Lemma 6.2 *Let* $\varphi^{\tau,\delta}$ *and* $r^{\tau,\delta}$ *be respectively the solutions of* [\(6.5\)](#page-46-0) *and* [\(6.9\)](#page-47-4)*. Then, for every* $T > 0$, there exists a constant $C = C(T) > 0$ such that

$$
\sup_{t \in [0,T]} \mathcal{E}_t^{\tau} (\varphi^{\tau,\delta} - r^{\tau,\delta}) \le C \|r^{\tau,\delta}\|_{\tau}^2.
$$
\n(6.10)

Proof. First, recall that $(\partial_t - \mathcal{L}^{\tau}) \varphi^{\tau,\delta} = \mathfrak{m}^{\tau,\delta}$ while $(\partial_t - \mathcal{L}^{\tau}) r^{\tau,\delta} = -\mathcal{L}^{\tau} r^{\tau,\delta} = \mathfrak{m}^{\tau,\delta} - r^{\tau,\delta}$, since $r^{\tau,\delta}$ does not depend on t. Thus, by [\(6.8\)](#page-47-0) and Cauchy-Schwarz, we deduce

$$
\mathcal{E}_t^{\tau}(\varphi^{\tau,\delta} - r^{\tau,\delta}) = \|r^{\tau,\delta}\|_{\tau}^2 + 2 \int_0^t \langle r^{\tau,\delta}, \varphi_s^{\tau,\delta} - r^{\tau,\delta} \rangle_{\tau} ds
$$

$$
\leq (1+t) \|r^{\tau,\delta}\|_{\tau}^2 + \int_0^t \|\varphi_s^{\tau,\delta} - r^{\tau,\delta}\|_{\tau}^2 ds
$$

$$
\leq (1+t) \|r^{\tau,\delta}\|_{\tau}^2 + \int_0^t \mathcal{E}_{s,\tau}(\varphi^{\tau,\delta} - r^{\tau,\delta}) ds,
$$

where the last step follows by the first equality in (6.8) . Then, Grönwall's lemma gives

$$
\mathscr{E}_t^{\tau}(\varphi^{\tau,\delta} - r^{\tau,\delta}) \le (1+t) e^t \, \|r^{\tau,\delta}\|_{\tau}^2 \,,
$$

which concludes the proof.

6.2 Proof of Theorem [1.1](#page-4-0)

In view of Proposition [6.1](#page-47-5) and Lemma [6.2,](#page-48-0) we are left to identify the limit as first $\delta \to 0$ and then $\tau \to \infty$ of $\mathscr{E}_t^{\tau}(\tau^{\tau,\delta})$ for $r^{\tau,\delta}$ the solution to the elliptic equation [\(6.9\)](#page-47-4). This is the point at which Theorem [4.1](#page-35-2) comes into play. To apply it, we need to estimate the right hand sides of [\(4.3\)](#page-35-3) and [\(4.4\)](#page-35-4) for φ given by $\mathfrak{m}^{\tau,\delta}$ in [\(6.4\)](#page-46-3). We want to control the (semi-)norms appearing therein using the results of Section [3.3,](#page-29-4) which are though stated in terms of the sharp operators $\mathscr{A}_\pm^{\sharp,\tau}$. Hence, let us consider the following slight modification of $\mathfrak{m}^{\tau,\delta}$

$$
\tilde{\mathfrak{m}}^{\tau,\delta} \stackrel{\text{def}}{=} \mathfrak{A}_{+}^{\sharp,\tau} \partial_{\mathfrak{e}_1}^{-1} \psi^{\delta} \tag{6.11}
$$

and show that its resolvent is close to $r^{\tau,\delta}$ in ΓH^1_τ .

Lemma 6.3 *Let* $r^{\tau,\delta}$ *and* $\tilde{r}^{\tau,\delta}$ *be respectively the solutions to* [\(6.9\)](#page-47-4) *and*

$$
(1 - \mathcal{L}^{\tau})\tilde{r}^{\tau,\delta} = \tilde{\mathfrak{m}}^{\tau,\delta}.
$$
\n(6.12)

Then, there exists a constant $C = C(\lambda) > 0$ *, independent of* δ *such that*

$$
||r^{\tau,\delta} - \tilde{r}^{\tau,\delta}||_{\Gamma H^1_\tau} \leq C \nu_\tau^{1/2} \delta. \tag{6.13}
$$

Proof. By [\(2.28\)](#page-21-1) and the equations [\(6.9\)](#page-47-4) and [\(6.12\)](#page-48-1), we have

$$
\begin{split} \|\boldsymbol{r}^{\tau,\delta}-\tilde{\boldsymbol{r}}^{\tau,\delta}\|_{\Gamma H_{\tau}^{1}}^{2} &= \langle (1-\mathcal{L}^{\tau})(\boldsymbol{r}^{\tau,\delta}-\tilde{\boldsymbol{r}}^{\tau,\delta}), \boldsymbol{r}^{\tau,\delta}-\tilde{\boldsymbol{r}}^{\tau,\delta}\rangle_{\tau} \\ &= \langle \mathfrak{m}^{\tau,\delta}-\tilde{\mathfrak{m}}^{\tau,\delta}, \boldsymbol{r}^{\tau,\delta}-\tilde{\boldsymbol{r}}^{\tau,\delta}\rangle_{\tau} \leq \|\mathfrak{m}^{\tau,\delta}-\tilde{\mathfrak{m}}^{\tau,\delta}\|_{\Gamma H_{\tau}^{-1}} \|\boldsymbol{r}^{\tau,\delta}-\tilde{\boldsymbol{r}}^{\tau,\delta}\|_{\Gamma H_{\tau}^{1}} \end{split}
$$

 \Box

so that we are left to control the ΓH_{τ}^{-1} -norm of the difference between $\mathfrak{m}^{\tau,\delta}$ and $\tilde{\mathfrak{m}}^{\tau,\delta}$. Their definition in [\(6.4\)](#page-46-3) and [\(6.11\)](#page-48-2) together with the fact that $\mathcal{A}_{+}^{\tau} = \mathcal{A}_{+}^{\sharp,\tau} + \mathcal{A}_{+}^{\flat,\tau}$, implies that $\mathfrak{m}^{\tau,\delta}$ – $\tilde{\mathfrak{m}}^{\tau,\delta} = \mathfrak{A}^{\flat,\tau} \partial_{\mathfrak{e}_1}^{-1} \psi^{\delta}$. Thus,

$$
\|{\mathscr A}^{\flat,\tau} \partial_{\mathfrak e_1}^{-1} \psi^\delta\|_{\Gamma H_\tau^{-1}}^2 \leq \|{\mathscr A}^{\flat,\tau} \partial_{\mathfrak e_1}^{-1} \psi^\delta\|_\tau^2 \lesssim \lambda^2 \nu_\tau^{3/2} \int_{\mathbb R^2} {\rm d} p\, |\hat\psi^\delta(p)|^2 \int_{\mathbb R^2} {\rm d} q\, 1\!\!1_{|\sqrt{R^\tau}q|\leq 2\delta} \lesssim \lambda^2 \nu_\tau \delta^2
$$

where we used that $\|\psi^{\delta}\|_{L^2(\mathbb{R}^2)} = 1$ and that the volume of $\{q : |\sqrt{R'}q| \leq 2\delta\}$ is of order $\nu_{\tau}^{-1/2} \delta^2$. Hence, [\(6.13\)](#page-48-3) follows at once. \Box

Let us now analyse $\tilde{m}^{\tau,\delta}$ and estimate all the (semi-)norms needed to conclude our proof.

Lemma 6.4 *Let* $\tilde{m}^{\tau,\delta}$ *be defined according to* [\(6.11\)](#page-48-2). Then, there exists a constant C = $C(\lambda, M) > 0$ *such that*

$$
\limsup_{\delta \to 0} \|\tilde{\mathfrak{m}}^{\tau,\delta}\|_{\tau,M} \vee \|\mathcal{G}_M^{\tau}\tilde{\mathfrak{m}}^{\tau,\delta}\|_{\tau} \leq C\nu_{\tau}^{1/4} |\log \nu_{\tau}| \tag{6.14}
$$

$$
\limsup_{\delta \to 0} \left| \| (\mathcal{C}_M^{\tau})^{1/2} \tilde{\mathfrak{m}}^{\tau,\delta} \|_{\tau}^2 - \frac{1}{2} C_{\text{eff}}(\lambda) \right| \leq C \nu_{\tau}
$$
\n(6.15)

where $\|\cdot\|_{\tau,M}$ *is the (semi-)norm in* [\(4.5\)](#page-35-5).

Proof. Let us begin by controlling $\|\mathcal{G}_{\mathcal{M}}^{\tau} \tilde{m}^{\tau,\delta}\|_{\tau}$. Since the support of $\hat{\psi}$ is contained in a ball of radius 1 away from $e_2\mathbb{R}$, that of $\mathfrak{F}(\partial_{e_1}^{-1}\psi^{\delta})$ is contained in a ball of radius δ which is independent of τ . Hence, we can apply [\(3.31\)](#page-33-1), which gives

$$
\|\mathcal{G}_M^\tau \tilde{\mathfrak{m}}^{\tau,\delta}\|_\tau = \|\mathcal{G}_M^\tau \mathcal{A}_+^{\sharp,\tau} \partial_{\mathfrak{e}_1}^{-1} \psi^\delta\|_\tau \lesssim \nu_\tau^{3/4} \|\partial_{\mathfrak{e}_1}^{-1} \psi^\delta\|_\tau \lesssim \nu_\tau^{3/4} \|\partial_{\mathfrak{e}_1}^{-1} \psi\|_{L^2(\mathbb{R}^2)} \lesssim \nu_\tau^{3/4}
$$

and in the third step we used that $\Theta^{\tau} \lesssim 1$ and in the last last step that $\hat{\psi}$ is supported away from $\mathfrak{e}_2\mathbb{R}$ and that ψ is smooth, so that $\|\partial_{\mathfrak{e}_1}^{-1}\psi\|_{L^2(\mathbb{R}^2)} < \infty$. Note that, for τ large enough, the right hand side is controlled by the right hand side of [\(6.14\)](#page-49-0).

Concerning $\|\tilde{\mathfrak{m}}^{\tau,\delta}\|_{\tau,M}$, we will separately bound the two summands in its definition. For the second, the definition of $\mathfrak{A}^{\sharp,\tau}_+$ in [\(3.7\)](#page-26-0) ensures that $\mathfrak{Fm}^{\tau,\delta}$ is supported on modes (q,q') satisfying $|q + q'| \leq \delta$, for which the kernel $\mathcal{K}_{q,q'}^{-,\tau}$ in [\(2.26\)](#page-20-0) of $\mathcal{A}^{\sharp,\tau}_{-}$ can be bounded by δ . Hence, at a price which might blow up as $\tau \to \infty$, we have

$$
\| {\mathscr A}_{-}^{\sharp,\tau} {\mathscr G}_{M}^\tau \tilde{{\mathfrak m}}^{\tau,\delta} \|_{{\tau}}^2 \lesssim_{\lambda,M,\tau} \int_{{\mathbb R}^2} d p \, \delta^2 \, |\hat{\psi}^{\delta}(p)|^2 \int_{{\mathbb R}^2} d q \, \Theta_{\tau}(q) \Theta_{\tau}(p-q) \lesssim_{\lambda,M,\tau} \delta^2 \,,
$$

which means that, since we are taking $\delta \to 0$ first, the limsup_{$\delta \to 0$} of the previous vanishes. For the first, we first apply the Recursive Replacement Lemma [3.9](#page-32-0) with $f = p_0^{\tau} \equiv \nu_{\tau}$ and p_0^{τ} as in [\(4.28\)](#page-41-0) (meaning $C_f = 1$), and get

$$
\begin{aligned} \|(-\nu_\tau \mathcal{L}_0^{\mathfrak{e}_1})^{1/2} \mathcal{G}_M^\tau \tilde{\mathfrak{m}}^{\tau,\delta}\|_\tau &= \|(-\nu_\tau \mathcal{L}_0^{\mathfrak{e}_1})^{1/2} \mathcal{G}_M^\tau \mathcal{A}_+^{\sharp,\tau} \partial_{\mathfrak{e}_1}^{-1} \psi^\delta\|_\tau \\ &\leq \|(-\mathcal{L}_0^{\mathfrak{e}_1} [p_1^\tau + \nu_\tau C_M^{(1)}])^{1/2} \partial_{\mathfrak{e}_1}^{-1} \psi^\delta\|_\tau \end{aligned}
$$

$$
\lesssim \nu_{\tau}^{1/2} \sqrt{|\log \nu_{\tau}|} \| (-\mathcal{L}_0^{\mathfrak{e}_1})^{1/2} \partial_{\mathfrak{e}_1}^{-1} \psi^{\delta} \|_{\tau}
$$

$$
\lesssim \nu_{\tau}^{1/2} \sqrt{|\log \nu_{\tau}|} \| \psi^{\delta} \|_{\tau} \leq \nu_{\tau}^{1/2} \sqrt{|\log \nu_{\tau}|}
$$
(6.16)

where we used [\(4.30\)](#page-41-4) together with the definition of $\mathcal{L}_0^{\mathfrak{e}_1}[\cdot]$ in [\(3.15\)](#page-29-2) and the fact that

$$
p_1^{\tau}(L^{\tau}(-\mathcal{S}^{\tau})) + \nu_{\tau}C_M^{(1)} \leq \nu_{\tau} \left(\log \sqrt{\frac{\nu_{\tau} + g^{\tau}(L^{\tau}(-\mathcal{S}^{\tau}))}{\nu_{\tau}}} + 1 \right) \lesssim \nu_{\tau} |\log \nu_{\tau}|.
$$

Simplifying the $\nu_{\tau}^{1/2}$ at the right and left hand sides of [\(6.16\)](#page-50-0), we deduce that the first summand in [\(4.5\)](#page-35-5) is bounded by the right hand side of [\(6.14\)](#page-49-0) uniformly in $\delta > 0$, and thus, the analysis of $\|\tilde{\mathfrak{m}}^{\tau,\delta}\|_{\tau,M}$ is concluded.

At last, we turn to [\(6.15\)](#page-49-1). Let us first note that, by [\(3.1\)](#page-24-1) and the definition of $\tilde{m}^{\tau,\delta}$ in [\(6.11\)](#page-48-2), we have

$$
\|(\mathcal{C}_M^{\tau})^{1/2}\tilde{\mathfrak{m}}^{\tau,\delta}\|_{\tau}^2 = \langle -\mathcal{A}_-^{\sharp,\tau}\mathcal{C}_M^{\tau}\mathcal{A}_+^{\sharp,\tau}\partial_{\mathfrak{e}_1}^{-1}\psi^{\delta}, \partial_{\mathfrak{e}_1}^{-1}\psi^{\delta}\rangle_{\tau}.
$$

and the Replacement Lemma [3.6](#page-30-1) gives

$$
\begin{aligned} \left| \langle - \mathcal{A}_-^{\sharp,\tau} \mathcal{G}_M^\tau \mathcal{A}_+^{\sharp,\tau} \partial_{\mathfrak{e}_1}^{-1} \psi^\delta, \partial_{\mathfrak{e}_1}^{-1} \psi^\delta \rangle_\tau - \langle \mathcal{g}_M^\tau \partial_{\mathfrak{e}_1}^{-1} \psi^\delta, \partial_{\mathfrak{e}_1}^{-1} \psi^\delta \rangle_\tau \right| \\ \lesssim \| (-\nu_\tau \mathcal{L}_0^{\mathfrak{e}_1})^{1/2} \partial_{\mathfrak{e}_1}^{-1} \psi^\delta \|_\tau^2 \leq \nu_\tau \| \psi^\delta \|_{L^2(\mathbb{R}^2)} = \nu_\tau \,. \end{aligned}
$$

As a consequence, we are left to analyse

$$
\|(q_M^{\tau})^{1/2}\partial_{\mathfrak{e}_1}^{-1}\psi^{\delta}\|_{\tau}^2 = \|g_M^{\tau}(L^{\tau}(-\mathcal{S}^{\tau}))^{1/2}(-\mathcal{L}_0^{\mathfrak{e}_1})^{1/2}\partial_{\mathfrak{e}_1}^{-1}\psi^{\delta}\|_{\tau}^2 = \frac{1}{2}\|g_M^{\tau}(L^{\tau}(-\mathcal{S}^{\tau}))^{1/2}\psi^{\delta}\|_{\tau}^2,
$$

where the prefactor $1/2$ comes from the definition of $\mathcal{L}_0^{\mathfrak{e}_1}$ in [\(2.22\)](#page-19-2). Now, by construction, $|\hat{\psi}^{\delta}|^2$ converges to a Dirac mass at 0 as $\delta \to 0$, which means that, as $\delta \to 0$ the right hand side converges to

$$
g_M^{\tau}(L^{\tau}(0)) = \left(\frac{3\lambda^2}{2\pi}\nu_{\tau}^{3/2}\log(1+\tau) + \nu_{\tau}^{3/2}\right)^{2/3} - \nu_{\tau} = C_{\text{eff}}(\lambda) + O(\nu_{\tau})
$$

from which (6.15) follows at once.

Thanks to the previous lemmas, we can now complete the proof of Theorem [1.1.](#page-4-0)

Proof of Theorem [1.1.](#page-4-0) The bulk of the proof consists of establishing that, as $\tau \to \infty$, $D_{1,1}^{\tau}(1)$ converges to $C_{\text{eff}}(\lambda)$, which, by [\(6.7\)](#page-47-1), follows provided

$$
\lim_{\tau \to \infty} \lim_{\delta \to 0} \mathcal{E}_1^{\tau}(\varphi^{\tau,\delta}) = C_{\text{eff}}(\lambda), \qquad (6.17)
$$

where $\varphi^{\tau,\delta}$ is the solution of the parabolic equation [\(6.5\)](#page-46-0).

Let us first show that, in [\(6.17\)](#page-50-1), we can replace $\varphi^{\tau,\delta}$ with the solution $\tilde{r}^{\tau,\delta}$ of [\(6.12\)](#page-48-1) at a cost which is negligible with respect to δ and τ . Recall the definition of $r^{\tau,\delta}$ in [\(6.9\)](#page-47-4), and note that

$$
\mathcal{E}_1^{\tau}(\varphi^{\tau,\delta} - \tilde{r}^{\tau\delta}) \lesssim \mathcal{E}_1^{\tau}(\varphi^{\tau,\delta} - r^{\tau,\delta}) + \mathcal{E}_1^{\tau}(r^{\tau,\delta} - \tilde{r}^{\tau,\delta}) \lesssim \|r^{\tau,\delta}\|_{\tau}^2 + \|r^{\tau,\delta} - \tilde{r}^{\tau,\delta}\|_{\Gamma H_{\tau}^1}^2 \lesssim \|\tilde{r}^{\tau,\delta}\|_{\tau}^2 + \|r^{\tau,\delta} - \tilde{r}^{\tau,\delta}\|_{\Gamma H_{\tau}^1}^2
$$
\n(6.18)

 \Box

where in the last step we simply applied the triangle inequality to bound $||r^{\tau,\delta}||_{\tau}$ by the right hand side, and in the second, we used [\(6.10\)](#page-48-4) and fact that, for a time independent $\varphi \in Dom(\mathcal{L}^{\tau})$, [\(6.8\)](#page-47-0) implies that $\mathscr{E}_1^{\tau}(\varphi) \lesssim ||\varphi||_{\Gamma H_{\tau}^{\tau}}^2$. Now, the second term at the right hand side is of order $\nu_{\tau} \delta^2$ in view of [\(6.13\)](#page-48-3), while for the first we invoke [\(4.3\)](#page-35-3) (with $r^{\tau,\delta}$ replaced by $\tilde{r}^{\tau,\delta}$), which gives

$$
\limsup_{\delta \to 0} \|\tilde{r}^{\tau,\delta}\|_{\tau} \lesssim \limsup_{\delta \to 0} \left(\|\tilde{\mathfrak{m}}^{\tau,\delta}\|_{\tau,M} + \|\mathcal{G}_M^{\tau}\tilde{\mathfrak{m}}^{\tau,\delta}\|_{\tau} \right) \lesssim \nu_{\tau}^{1/4} |\log \nu_{\tau}| \tag{6.19}
$$

thanks to (6.14) .

As a consequence, [\(6.18\)](#page-50-2) vanishes in the double limit $\lim_{\tau\to\infty} \lim_{\delta\to 0}$ and we are left to study

$$
\mathscr{E}_1^{\tau}(\tilde{r}^{\tau,\delta}) = \|\tilde{r}^{\tau,\delta}\|_{\tau}^2 + 2\int_0^1 \|(-\mathcal{S}^{\tau})^{1/2}\tilde{r}^{\tau,\delta}\|_{\tau}^2 ds = \|\tilde{r}^{\tau,\delta}\|_{\tau}^2 + 2\|\tilde{r}^{\tau,\delta}\|_{\Gamma H_{\tau}^1}^2
$$

The $\limsup_{\delta \to 0} \|\tilde{r}^{\tau,\delta}\|_{\tau}^2$ can be controlled as in [\(6.19\)](#page-51-0), while for the ΓH_{τ}^1 -norm, by [\(4.4\)](#page-35-4) (with $r^{\tau,\delta}$ replaced by $\tilde{r}^{\tau,\delta}$), we get

$$
\begin{aligned} \limsup_{\delta \to 0} \left| \| {\tilde r}^{\tau,\delta} \|_{\Gamma H^1_\tau}^2 - \tfrac{1}{2} C_{\rm eff}(\lambda) \right| \\ & \quad \le \limsup_{\delta \to 0} \left| \| {\tilde r}^{\tau,\delta} \|_{\Gamma H^1_\tau}^2 - \| (\mathcal{G}^\tau_M)^{1/2} \tilde{\mathfrak{m}}^{\tau,\delta} \|_\tau^2 \right| + \left| \| (\mathcal{G}^\tau_M)^{1/2} \tilde{\mathfrak{m}}^{\tau,\delta} \|_\tau^2 - \tfrac{1}{2} C_{\rm eff}(\lambda) \right| \\ & \quad \lesssim \limsup_{\delta \to 0} \| \tilde{\mathfrak{m}}^{\tau,\delta} \|_{\tau,M} \left(\| \tilde{\mathfrak{m}}^{\tau,\delta} \|_{\tau,M} + \| (\mathcal{G}^\tau_M)^{1/2} \tilde{\mathfrak{m}}^{\tau,\delta} \|_\tau \right) + \nu_\tau \lesssim \nu_\tau^{1/4} |\log \nu_\tau| \end{aligned}
$$

where we used both (6.14) and (6.15) . By putting the previous estimates together, (6.17) follows at once.

In order to complete the proof of (1.5) , it suffices to exploit the scaling relation in (6.1) and the definition of ν_{τ} in [\(1.16\)](#page-10-0), which give

$$
D_{1,1}(\tau) = C_{\text{eff}}(\lambda) \nu_{\tau}^{-1} + O(\nu_{\tau}^{-3/4} |\log \nu_{\tau}|) = C_{\text{eff}}(\lambda) (\log \tau)^{2/3} \left(1 + O\left(\frac{\log \log \tau}{(\log \tau)^{1/6}}\right) \right). \tag{6.20}
$$

Appendix A The regularised SBE: proofs

In this appendix, we collect some of the proofs of the statements concerning the solution to the regularised SBE in [\(2.1\)](#page-13-1). Since $\tau > 0$ is fixed, we take it to be equal to 1 and omit the corresponding index so to lighten the notation. In particular, $\nu_{\tau} = 1$, R_{τ} is the identity matrix and [\(1.9\)](#page-6-1) reduces to [\(1.2\)](#page-3-0). The case of general τ can be recovered by scaling.

We begin by recalling the definition and main properties of weighted Besov spaces (for a thorough exposition we refer to [\[ET96,](#page-70-1) Ch. 4]).

A.1 Weighted Besov spaces

Since we want to construct a solution to (2.1) in infinite volume, we will need to consider weighted norms. A function $w: \mathbb{R}^2 \to \mathbb{R}_+$ is said to be a *weight* if there exists a positive constant $C > 0$ such that

$$
C^{-1} \le \sup_{|x-y| \le 1} \frac{w(x)}{w(y)} \le C.
$$
 (A.1)

.

The weights we will be concerned with are of polynomial or exponential type, namely

$$
p_a(x) \stackrel{\text{def}}{=} (1+|x|)^{-a}, \qquad e_b^{\sigma}(x) \stackrel{\text{def}}{=} \exp(-b(1+|x|)^{\sigma})
$$
 (A.2)

for $a, b \in \mathbb{R}_+$, $\sigma \in (0, 1)$ and $x \in \mathbb{R}^2$.

We now turn to Besov spaces, whose definition is based on a *dyadic partition of unity*, i.e. a family $(\varphi_j)_{j \geq -1} \subset \mathcal{C}_c^{\infty}(\mathbb{R}^2)$ of non-negative radial functions such that

- supp φ_{-1} and supp φ_0 are respectively contained in a ball and in an annulus around the origin,
- for $j \geq 0$, $\varphi_j(\cdot) = \varphi_0(2^{-j} \cdot)$
- for all $x \in \mathbb{R}^2$, $\sum_{j \geq -1} \varphi_j(x) = 1$,
- if $|j j'| > 1$, then supp $\varphi_j \cap \text{supp } \varphi_{j'} = \emptyset$.

We denote by Φ_j the inverse Fourier transform of φ_j and set $\Delta_j f \stackrel{\text{def}}{=} \Phi_j * f$ to be the j-th Littlewood-Paley block.

Definition A.1 Let w be a weight as in [\(A.1\)](#page-51-1), $\alpha \in \mathbb{R}$ and $p, q \in [1, \infty]$. We define the weighted Besov space $\mathfrak{B}_{p,q}^{\alpha}(w)$ as the space of distributions (if the weight is polynomial, and of ultra-distributions^{[6](#page-52-0)} if exponential) $f \in \mathscr{S}'(\mathbb{R}^2)$ (or $\mathscr{S}'_{\omega}(\mathbb{R}^2)$) such that

$$
||f||_{\mathcal{B}_{p,q}^{\alpha}(w)} \stackrel{\text{def}}{=} \left\| \left(2^{\alpha j} ||w \Delta_j f||_{L^p(\mathbb{R}^2)} \right)_{j \geq -1} \right\|_{\ell^q} = \left(\sum_{j \geq -1} 2^{\alpha j q} ||w \Delta_j f||_{L^p(\mathbb{R}^2)}^q \right)^{\frac{1}{q}} < \infty. \quad (A.3)
$$

For $p = q = \infty$, we adopt the notation $\mathcal{C}^{\alpha}(w) \stackrel{\text{def}}{=} \mathcal{B}_{\infty,\infty}^{\alpha}(w)$.

As given above, weighted Besov spaces are Banach and, for suitable choices of α , p , q and w they coincide with spaces the reader might be more familiar with. In particular, if $p = q = \infty$, $\alpha \in \mathbb{R} \setminus \mathbb{N}$ and $w \equiv 1$, $\mathcal{C}^{\alpha}(1)$ is the space of α -Hölder continuous functions. If $p = q = 2$, $\mathcal{B}_{2,2}^{\alpha}(w) = H^{\alpha}(w)$ is the usual (weighted) Sobolev space and, if $\alpha = 0$ and $w \equiv 1$, $\mathfrak{B}_{2,2}^0(1) = L^2(\mathbb{R}^2).$

Besov spaces satisfy a number of properties that we will need in what follows. Let us summarise them without proof.

• *(Besov embedding)* (see e.g. [\[ET96,](#page-70-1) Thm. 4.2.3]) for any weights $w_2 \leq w_1$ such that $\lim_{|x|\to\infty} w_2(x)/w_1(x) = 0, 1 \le p_1 \le p_2 \le \infty, 1 \le q_1 \le q_2 \le \infty, \alpha_1 \in \mathbb{R}$ and $\alpha_2 \leq \alpha_1 - 2(1/p_1 + 1/p_2)$, we have the continuous embedding

$$
\mathfrak{B}_{p_1,q_1}^{\alpha_1}(\mathbf{w}_1) \subseteq \mathfrak{B}_{p_2,q_2}^{\alpha_2}(\mathbf{w}_2).
$$
 (A.4)

• *(Semigroup estimates)* (see e.g. [\[MP19,](#page-72-2) Prop. 3.6]) Let $K_t \stackrel{\text{def}}{=} e^{\frac{t}{2}\Delta}$ be the semigroup associated to $\frac{1}{2}\Delta$. Then, for any $\alpha \in \mathbb{R}$, $\beta \ge 0$ and weight w, we have

$$
||K_t f||_{\mathscr{C}^{\alpha+\beta}(\mathbf{w})} \lesssim t^{-\frac{\beta}{2}} ||f||_{\mathscr{C}^{\alpha}(\mathbf{w})}, \tag{A.5}
$$

where $K_t f \stackrel{\text{def}}{=} K_t * f$.

⁶For the definition of ultra-distributions and their need in the case of exponential weights, see e.g. [\[MP19,](#page-72-2) Sec. 2.2]

• *(Product rule)* (see e.g. [\[MP19,](#page-72-2) Lem. 4.2]) For any $\alpha_1, \alpha_2 \in \mathbb{R}$ and weights w₁, w₂, if $\alpha_1 + \alpha_2 > 0$, then

$$
||fg||_{\mathscr{C}^{\alpha_1 \wedge \alpha_2}(w_1 w_2)} \lesssim ||f||_{\mathscr{C}^{\alpha_1}(w_1)} ||g||_{\mathscr{C}^{\alpha_2}(w_2)}.
$$
 (A.6)

A.2 Approximation and convergence for the linear equation

Before turning to Burgers, we consider the stationary solution to the linear stochastic heat equation (SHE) obtained by dropping the non-linearity in [\(2.1\)](#page-13-1).

Let $(\eta, \vec{\xi})$ be a couple of independent space and space-time Gaussian white noises on \mathbb{R}^2 and $\mathbb{R} \times \mathbb{R}^2$ respectively. In order to ensure measurability of the solution (both for SHE and SBE), we want to construct our approximating sequence in such a way that all of its elements are coupled together using the same noises. For this, starting from $(\eta, \vec{\xi})$ and given $M \in \mathbb{N}$, we set

$$
\eta_M \stackrel{\text{def}}{=} \sum_{y \in M\mathbb{Z}^2} \tau_y(\eta \mathbb{1}_{Q(M)}) \quad \text{and} \quad \vec{\xi}_M(\text{d}s) \stackrel{\text{def}}{=} \sum_{y \in M\mathbb{Z}^2} \tau_y(\vec{\xi} \mathbb{1}_{Q(M)}) \quad (A.7)
$$

where $Q(M) \stackrel{\text{def}}{=} [-M/2, M/2]^2$ and, for $y \in \mathbb{R}^2$, we denoted by τ_y the translation operator on $S'(\mathbb{R}^2)$ (and defined by duality from $\tau_y f(\cdot) = f(\cdot - y)$ for $f \in S(\mathbb{R}^2)$). Note that, thanks to spatial independence, $(\eta_M, \vec{\xi}_M)$ coincide in law to *periodic* independent space and space-time white noises with period $M \in \mathbb{N}$.

According to Definition [2.1,](#page-13-2) the unique mild solutions X and X^M of [\(2.1\)](#page-13-1) with $\lambda = 0$ driven by $(\eta^1, \vec{\xi}^1)$ and by $(\eta^1_M, \vec{\xi}^1_M)$ are respectively given by

$$
X_t = K_t \eta^1 + \int_0^t K_{t-s} \operatorname{div} \vec{\xi}^1(\mathrm{d}s) \quad \text{and} \quad X_t^M = K_t \eta^1_M + \int_0^t K_{t-s} \operatorname{div} \vec{\xi}^1_M(\mathrm{d}s), \tag{A.8}
$$

for K the usual heat kernel. Then, we have the following lemma.

Lemma A.2 In the setting outlined above, let X and X^M be as in [\(A.8\)](#page-53-0). Then, for any $\alpha, a, p, T > 0$ and $\beta \in [0, 1/2)$, X and X^M almost surely belong to $C_{\text{loc}}^{\beta}(\mathbb{R}_+, \mathcal{C}^{\infty}(\mathbb{R}^2))$ and *there exists a constant* $C = C(\rho) > 0$ *such that*

$$
\mathbf{E}\Big[\|X\|_{C^{\beta}([0,T],\mathcal{C}^{\alpha}(p_a))}^p\Big] \vee \sup_M \mathbf{E}\Big[\|X^M\|_{C^{\beta}([0,T],\mathcal{C}^{\alpha}(p_a))}^p\Big] \leq C\,. \tag{A.9}
$$

Furthermore,

$$
\lim_{M \to \infty} \mathbf{E} \Big[\|X - X^M\|_{C^{\beta}([0,T], \mathcal{C}^{\alpha}(p_a))}^p \Big] = 0. \tag{A.10}
$$

In the proof of the lemma (and in Lemma [A.4\)](#page-55-0) we will need the following technical result.

Lemma A.3 *For any* $\gamma, \kappa > 0$, $\mathbf{k} \in \mathbb{N}^2$ *and* $m \in \{1, 2\}$ *, there exists a constant* $C > 0$ *such that for every* $j \geq -1$, every $r \geq 0$ and all $x \in \mathbb{R}^2$, we have

$$
||\mathbf{p}_{-2\kappa}(\Delta_j K_r \partial_{\mathbf{k}}^{\|\mathbf{k}\|} \varrho^{*m})(x-\cdot)||_{L^2(\mathbb{R}^2)} \le C2^{-\gamma j}
$$
\n(A.11)

where Δ_j *denotes the j-th Littlewood-Paley block,* $p_{-\kappa}$ *is as in* [\(A.2\)](#page-52-1) *and* $\partial_{\bf k}^{|{\bf k}|} = \partial_1^{k_1} \partial_2^{k_2}$. *Moreover, for any* $0 \leq s \leq t$ *, we have*

$$
\|\mathbf{p}_{-\kappa}(\Delta_j(K_t - K_s)\partial_{\mathbf{k}}^{\|\mathbf{k}\|}\varrho)(x - \cdot)\|_{L^2(\mathbb{R}^2)} \le C(t - s)2^{-\gamma j} \,. \tag{A.12}
$$

The regularised SBE: proofs **55**

Proof. Notice that, upon taking $\kappa' > \kappa$ so that $\kappa' \in \mathbb{N}$, we can interpret the weight $p_{-\kappa'}$ as the inverse Fourier transform of a derivative, thus the left hand side of [\(A.11\)](#page-53-1) is bounded above by

$$
\|\mathbf{p}_{-\kappa'}(\Delta_j K_r\partial_\mathbf{k}^{|\mathbf{k}|}\varrho^{*m})(x-\cdot)\|_{L^2(\mathbb{R}^2)}\lesssim \|\mathcal{F}^{-1}((\Delta_j K_r\partial_\mathbf{k}^{|\mathbf{k}|}\varrho^{*m})(x-\cdot))\|_{H^{\kappa'}(\mathbb{R}^2)}
$$

where, for any $p = (p_1, p_2) \in \mathbb{R}^2$,

$$
\mathcal{F}^{-1}((\Delta_j K_r \partial_{\mathbf{k}}^{|\mathbf{k}|} \varrho^{*m})(x - \cdot))(p) = \varphi_j(p) e^{-r|p|^2} p_1^{k_1} p_2^{k_2} \hat{\varrho}^m(p) e^{-2\pi i p \cdot x}
$$

A similar argument holds for the left hand side of [\(A.12\)](#page-53-2), with the inverse Fourier transform of the argument given by

$$
\mathcal{F}^{-1}(\Delta_j (K_t - K_s) \partial_{\mathbf{k}}^{\vert \mathbf{k} \vert} \varrho(x - \cdot))(p) = \varphi_j(p) \left(e^{-t|p|^2} - e^{-s|p|^2} \right) p_1^{k_1} p_2^{k_2} \, \hat{\varrho}(p) \, e^{-2\pi i p \cdot x} \, .
$$

Hence, $(A.11)$ and $(A.12)$ follow by the fact that every factor in the above formulas is (uniformly) smooth, that $\hat{\varrho}$ and all its derivatives decay faster than any polynomial, and that φ_i is supported on modes p of order 2^j . \Box

Proof of Lemma [A.2.](#page-53-3) Since the proof of [\(A.9\)](#page-53-4) and [\(A.10\)](#page-53-5) follows the same steps with the second being slightly more involved, we only focus on the latter.

We proceed as usual, namely, setting $\tilde{X}^M \stackrel{\text{def}}{=} X - X^M$, we control the time increments of \tilde{X}^M (and \tilde{X}_0^M , but the proof is similar and thus omitted), apply first Besov's embedding [\(A.4\)](#page-52-2) and then Kolmogorov's criterion.

As a first step, we need to consider the Littlewood-Paley blocks of these increments and, using the coupling of $(\eta, \vec{\xi})$ and $(\eta^M, \vec{\xi}^M)$ provided by the construction in [\(A.7\)](#page-53-6), we write them purely in terms of $(\eta, \vec{\xi})$. To do so, let K_t^j $\frac{d}{dt} \stackrel{\text{def}}{=} \Delta_j K_t \varrho, j \geq -1$, and notice that

$$
\Delta_j \tilde{X}_t^M(x) = \eta(\tilde{K}_{t,x}^{j,M}) + \int_0^t \vec{\xi}(\mathrm{d}s, \nabla \tilde{K}_{t-s,x}^{j,M})
$$
\n(A.13)

where

$$
\tilde{K}_{t,x}^{j,M}(z) \stackrel{\text{def}}{=} \sum_{y \in M \mathbb{Z}_0^2} K_t^j(x-z) (\tau_y 1\!\!1_{Q(M)})(z) - \sum_{y \in M \mathbb{Z}_0^2} \tau_{-y} \Big(K_t^j(x-z) (\tau_y 1\!\!1_{Q(M)})(z) \Big)
$$

for $\mathbb{Z}_0^2 \stackrel{\text{def}}{=} \mathbb{Z}^2 \setminus \{0\}$. The absence of the summand corresponding to $y = 0$ is due to the difference. The right hand sides of [\(A.13\)](#page-54-0) are well-defined as, even though $\tilde{K}_{t,x}^{j,M}$ is not smooth (because of the behaviour on $M\mathbb{Z}^2$), it belongs to $L^2(\mathbb{R}^2)$ for all $x \in \mathbb{R}^2$. Indeed, for any $\mathbf{k} \in \mathbb{N}^2$, we have

$$
\|\partial_{\mathbf{k}}^{\mathbf{k}}\tilde{K}_{t,x}^{j,M}\|_{L^{2}(\mathbb{R}^{2})} \leq 2 \sum_{y\in M\mathbb{Z}_{0}^{2}} \|\partial_{\mathbf{k}}^{\mathbf{k}}K_{t}^{j}(x-\cdot)(\tau_{y}\mathbb{1}_{Q(M)})\|_{L^{2}(\mathbb{R}^{2})}
$$

$$
\lesssim \sum_{y\in M\mathbb{Z}_{0}^{2}} p_{\kappa}(y)\|p_{-\kappa}\partial_{\mathbf{k}}^{\mathbf{k}}K_{t}^{j}(x-\cdot)\|_{L^{2}(\mathbb{R}^{2})} \lesssim M^{-\kappa}2^{-j\gamma}
$$
(A.14)

where we have chosen $\kappa > 0$ large enough so that p_{κ} is summable and used [\(A.11\)](#page-53-1) in the last bound.

.

In view of [\(A.14\)](#page-54-1), we can now estimate to the time increments of $\Delta_i \tilde{X}^M$. Let $0 < a' < a$, $p > \max\{2/a', (1/2 - \beta)^{-1}\}\$ and $\alpha' \ge \alpha + 2/p$. Then, for any $0 \le s \le t$, we have

$$
\mathbf{E} \Big[\| \mathbf{p}_{a'} \Delta_{j} (\tilde{X}_{t}^{M} - \tilde{X}_{s}^{M}) \|_{L^{p}(\mathbb{R}^{2})}^{p} \Big] \n= \mathbf{E} \Big[\Big\| \mathbf{p}_{a'} \Big(\eta (\tilde{K}_{t,\cdot}^{j,M} - \tilde{K}_{s,\cdot}^{j,M}) + \int_{s}^{t} \vec{\xi} (\mathrm{d}r, \nabla \tilde{K}_{t-r,\cdot}^{j,M}) + \int_{0}^{s} \vec{\xi} (\mathrm{d}s, \nabla (\tilde{K}_{t-r,\cdot}^{j,M} - \tilde{K}_{s-r,\cdot}^{j,M})) \Big\|_{L^{p}(\mathbb{R}^{2})}^{p} \n= \int_{\mathbb{R}^{2}} \mathbf{p}_{a'}(x)^{p} \mathbf{E} \Big[\Big(\eta (\tilde{K}_{t,x}^{j,M} - \tilde{K}_{s,x}^{j,M}) + \int_{s}^{t} \vec{\xi} (\mathrm{d}r, \nabla \tilde{K}_{t-r,x}^{j,M}) + \int_{0}^{s} \vec{\xi} (\mathrm{d}s, \nabla (\tilde{K}_{t-r,x}^{j,M} - \tilde{K}_{s-r,x}^{j,M})) \Big)^{p} \Big] \mathrm{d}x \n\lesssim \int_{\mathbb{R}^{2}} \mathbf{p}_{a'}(x)^{p} \Big(\mathbb{E} [\eta (\tilde{K}_{t,x}^{j,M} - \tilde{K}_{0,x}^{j,M})^{2}] + \mathbf{E} \Big[\Big(\int_{s}^{t} \vec{\xi} (\mathrm{d}s, \nabla \tilde{K}_{t-s,x}^{j,M}) \Big)^{2} \Big] \Big) \Big\}^{p} \n+ \mathbf{E} \Big[\Big(\int_{0}^{s} \vec{\xi} (\mathrm{d}s, \nabla (\tilde{K}_{t-r,x}^{j,M} - \tilde{K}_{s-r,x}^{j,M})) \Big)^{2} \Big] \Big)^{\frac{p}{2}} \mathrm{d}x \n= \int_{\mathbb{R}^{2}} \mathbf{p}_{a'}(x)^{p} \Big(\|\tilde{K}_{t,x}^{j,M} - \tilde{K}_{s,x}^{j,M} \|_{L^{2}(\mathbb{R}^{2})}^{2} + \int_{s}^{t} \|\nabla \tilde
$$

where in the third step we used Gaussian hypercontractivity and the independence of η and $\vec{\xi}$, in the fourth Itô's isometry and in the last $(A.14)$ for the second summand, and the analogous bound for the first and third (which can be obtained arguing as in $(A.14)$ using $(A.12)$ in place of [\(A.11\)](#page-53-1)).

Now, using Besov's embeddings [\(A.4\)](#page-52-2) together with our assumption on a', α', p and $\gamma > \alpha'$, we get

$$
\mathbf{E}\Big[\| (\tilde{X}_t^M - \tilde{X}_s^M) \|_{\mathcal{C}^{\alpha}(\mathbf{p}_a)}^p \Big] \lesssim \mathbf{E}\Big[\| (\tilde{X}_t^M - \tilde{X}_s^M) \|_{\mathcal{B}^{\alpha'}_{p,p}(\mathbf{p}_{a'})}^p \Big]
$$
\n
$$
= \sum_{j \geq -1} 2^{j\alpha'^p} \mathbf{E}\Big[\| \mathbf{p}_{a'} \Delta_j \left[(\tilde{X}_t^M - \tilde{X}_s^M) \right] \|_{L^p(\mathbb{R}^2)}^p \Big] \lesssim_{\kappa} (t-s)^{p/2} M^{-\kappa}, \tag{A.15}
$$

from which, an application of Kolmogorov's criterion gives [\(A.10\)](#page-53-5).

 \Box

A.3 The regularised SBE: Proposition [2.3](#page-14-1)

This section is the core of the present appendix as it contains the proof of Proposition [2.3.](#page-14-1) The latter is based on two main ingredients: the derivation of suitable weighted estimates for a stationary periodic approximation to the solution of (2.1) (Lemma [A.4\)](#page-55-0) and a deterministic apriori bound which allows to control the difference of two solutions driven by possibly different noises (Lemma [A.5\)](#page-57-0). We begin with the former.

Lemma A.4 *For* $M \in \mathbb{N}$ *, let* η_M *and* $\vec{\xi}_M$ *be as in* [\(A.7\)](#page-53-6)*. Then, the equation* [\(2.1\)](#page-13-1) *driven by* $(\eta^1_M, \tilde{\xi}^1_M)$ admits a unique mild solution u^M according to Definition [2.1](#page-13-2) which is strong Markov, stationary, i.e. for every $t \geq 0$, $u_t^M \stackrel{\text{law}}{=} \eta_M^1$, and skew-reversible. Furthermore, for every

 $\alpha, a, p, T > 0$ and $\beta \in [0, 1/2)$, there exists a positive constant $C = C(\rho, \alpha, \beta, a, p, T) < \infty$ *such that*

$$
\sup_{M} \mathbf{E}\Big[\|u^M\|_{C^{\beta}([0,T], \mathcal{C}^{\alpha}(\mathbf{p}_a))}^p \Big] \leq C \,. \tag{A.16}
$$

Proof. Existence, uniqueness (of weak solutions in $C_{\text{loc}}(\mathbb{R}_+, \mathcal{S}'(\mathbb{T}_M^2))$, \mathbb{T}_M^2 being the twodimensional torus of side M), strong Markovianity, stationarity and skew-reversibility for fixed M (essentially) follow from [\[CGT24,](#page-70-2) Lemma 2.1]. In the afore-mentioned reference, it is shown that such properties hold for \tilde{u}^M which is the solution to

$$
\partial_t \tilde{u}^M = \frac{1}{2} \Delta \tilde{u}^M + \lambda \tilde{N}^1 [\tilde{u}^M] + (-\Delta)^{1/2} \xi_M , \qquad \tilde{u}_0^M \stackrel{\text{def}}{=} \eta_M
$$
 (A.17)

on \mathbb{T}_M^2 , with η_M and ξ_M space and space-time white noises on \mathbb{T}_M^2 , and

$$
\tilde{\mathcal{N}}^1[f](x) \stackrel{\text{def}}{=} [\varrho * (\mathfrak{e}_1 \cdot \nabla)(\varrho * f)^2](x) = [\varrho * \partial_1(\varrho * f)^2](x).
$$
 (A.18)

Compared to our setting, the ρ chosen therein is a cut-off function in Fourier space (so that in particular it is not compactly supported in real space) and the noise is formally different. That said, $(-\Delta)^{1/2} \xi_M \stackrel{\text{law}}{=}$ div $\vec{\xi}_M$ and, standard references (e.g. [\[DPZ08\]](#page-70-3)) guarantee existence and uniqueness of local in time weak solutions and that these solutions are adapted and strong Markov. On the other hand, the proof of global in time existence, stationarity and skew-reversibility given in [\[CGT24\]](#page-70-2) (see also [\[CT24,](#page-70-4) Lemma 3.3]) applies verbatim.

Therefore, setting $u^M \stackrel{\text{def}}{=} \varrho * \tilde{u}^M$, u^M is clearly smooth in space (and thus belongs to the unweighted space of spatially periodic functions with period M , $C_{loc}(\mathbb{R}_+, \mathcal{C}^\alpha_{per}_M)$, for any $\alpha > 0$), and is the (necessarily unique) weak solution to (1.2) (which is also mild, see Remark [2.2\)](#page-14-2) driven by $(\eta_M^1, \vec{\xi}_M^1)$. Further, it is a strong Markov process, has the law of η_M^1 as invariant measure and is skew-reversible.

Thus, identifying u^M with its periodic extension to the whole of \mathbb{R}^2 , we are left to show that, for every $T > 0$, it belongs to the weighted space $C^{\beta}([0,T], \mathcal{C}^{\alpha}(\mathbf{p}_a))$ uniformly over M, i.e. that [\(A.16\)](#page-56-0) holds. First, let us write the mild formulation as

$$
u_t^M = K_t \eta_M^1 + \lambda \int_0^t K_{t-s} \mathcal{N}^1[u_s^M] \, ds + \int_0^t K_{t-s} \, \text{div} \, \bar{\xi}_M^1(\text{d}s) = X_t^M + N_t^M
$$

in which X^M collects the first and the last summand, i.e. the terms containing the noises, and N^M the second, i.e. that containing the nonlinearity.

Let a' , p and α' be such that $0 < a' < a$, $p > max\{2/a', (1/2 - \beta)^{-1}\}$ and $\alpha' \ge \alpha + 2/p$. Let us begin by controlling the norm of the Littlewood-Paley block of N^M at a fixed time $t \geq 0$. Then,

$$
\mathbf{E}\Big[\| \mathbf{p}_{a'} \Delta_j N_t^M \|^p_{L^p(\mathbb{R}^2)}\Big] = \lambda^p \int_{\mathbb{R}^2} \mathbf{p}_{a'}(x)^p \mathbf{E}\Big[\Big(\int_0^t \Delta_j K_{t-r} \mathcal{N}^1[u_r^M](x) \, \mathrm{d}r\Big)^p\Big] \, \mathrm{d}x
$$

$$
\lesssim t^p \, \mathbb{E}\Big[\sup_{r \in [0,t]} \Big|\Delta_j K_r \mathcal{N}^1[\eta_M^1](0)\Big|^p\Big]
$$

where we first applied Jensen's inequality to the time integral, then we used stationarity of u^M to replace it with $\eta_M^{\bar{1}}$, translation invariance to remove the spatial dependence from the quantity in the expectation and the fact that $a'p > 2$ to bound the spatial integral. Now, for $\kappa > 0$ we have

$$
\left|\Delta_j K_r \mathcal{N}^1[\eta_M^1](0)\right| = \left|\langle \Delta_j K_r \partial_1 \varrho^{*2}, (\eta_M^1)^2 \rangle\right| \leq \|p_{-\kappa}\Delta_j K_r \partial_1 \varrho^{*2}\|_{L^2(\mathbb{R}^2)} \|(\eta_M^1)^2 p_{\kappa}\|_{L^2(\mathbb{R}^2)} \lesssim \|p_{-\kappa}\Delta_j K_r \partial_1 \varrho^{*2}\|_{L^2(\mathbb{R}^2)} \|\eta_M^1\|_{\mathscr{C}^{\alpha}(p_{\kappa})}^2 \lesssim 2^{-j\gamma} \| \eta_M^1\|_{\mathscr{C}^{\alpha}(p_{\kappa})}^2
$$

where the third step follows by [\(A.4\)](#page-52-2) and [\(A.6\)](#page-53-7) (up to take α sufficiently large) while the last step holds for any $\gamma > 0$ and $r \in [0, t]$, and is a consequence of [\(A.11\)](#page-53-1). Therefore, applying Besov's embedding once again, combining the last two estimates and choosing $\gamma > \alpha'$, we obtain

$$
\begin{aligned} \mathbf{E}\Big[\|N^M_t\|^p_{{\mathcal{C}^\alpha}({\mathbf{p}}_a)}\Big] &\lesssim \mathbf{E}\Big[\|N^M_t\|^p_{{\mathcal{B}^{\alpha'}_{{\mathcal{P}},{\mathcal{P}}}}({\mathbf{p}}_{a'})}\Big] = \sum_{j\geq -1} 2^{j\alpha'p}\,\mathbf{E}\Big[\Big\|{\mathbf{p}}_{a'}\,\Delta_j\,N^M_t\Big\|^p_{L^p({\mathbb{R}}^2)}\Big] \\ &\lesssim t^p\,\mathbb{E}\Big[\|\eta^1_M\|_{{\mathcal{C}^\alpha}({\mathbf{p}}_\kappa)}^{2p}\Big]\sum_{j\geq -1} 2^{-j(\gamma-\alpha')p}\lesssim t^p \end{aligned}
$$

in which we also used [\(A.9\)](#page-53-4) with $t = 0$.

Since u^M is Markov, stationary and has stationary increments, a bound on any time increment from s to t, with $0 \le s \le t$, follows provided we have one for $s = 0$. By the previous bound and [\(A.9\)](#page-53-4), we get

$$
\mathbf{E}\Big[\|u_t^M-u_0^M\|_{\mathscr{C}^{\alpha}(\mathbf{p}_a)}^p\Big]^{\frac{1}{p}}\leq \mathbf{E}\Big[\|X_t^M-X_0^M\|_{\mathscr{C}^{\alpha}(\mathbf{p}_a)}^p\Big]^{\frac{1}{p}}+\mathbf{E}\Big[\|N_t^M\|_{\mathscr{C}^{\alpha}(\mathbf{p}_a)}^p\Big]^{\frac{1}{p}}\lesssim_\beta t^\beta\,,
$$

for any $\beta \in [0, 1/2)$. By Kolmogorov's criterion, this implies [\(A.16\)](#page-56-0) so that the proof of the lemma is complete. lemma is complete.

At this point we can move to the second step, namely show that [\(2.1\)](#page-13-1) has a unique solution and that the sequence $\{u^M\}_M$ converges to that solution. As we will soon see, both the results follow from the next lemma.

Lemma A.5 *Let* $a, b, \alpha > 0, \sigma \in (0, 1)$ *such that* $a < \sigma$. *Let* $v \in C([0, T], \mathcal{C}^{\alpha}(e_b^{\sigma}))$ *be such that there exist* $X, w \in C([0, T], \mathcal{C}^{\alpha}(\mathfrak{p}_a))$ *for which*

$$
v_t = X_t + \lambda \int_0^t K_{t-s} \mathcal{N}^1[v_s, w_s] \, \mathrm{d}s \,, \qquad \forall \, t \in [0, T] \,, \tag{A.19}
$$

where ^N¹ [f, g] *is defined according to*

$$
\mathcal{N}^1[f,g] \stackrel{\text{def}}{=} (\partial_1 \varrho^{*2}) * (fg). \tag{A.20}
$$

Then, there exists a positive constant $C = C(\rho, \lambda, a, b, \sigma, T) < \infty$ *such that* v *satisfies*

$$
\sup_{t\in[0,T]}\|v_t\|_{\mathcal{C}^{\alpha}(\mathbf{e}_{b+t}^{\sigma})}\leq C\|X\|_{C([0,T],\mathcal{C}^{\alpha}(\mathbf{p}_a))}\exp\left(C\|w\|_{C([0,T],\mathcal{C}^{\alpha}(\mathbf{p}_a))}^{\frac{1}{1-a/\sigma}}\right) \tag{A.21}
$$

Proof. The proof follows the strategy implemented in $[HL15]$ in the context of the parabolic Anderson equation (which, contrary to our setting, is rough) and consists of taking advantage of the different weights imposed on v and w . The crucial ingredient is the (trivial) estimate

$$
\frac{\mathbf{e}_{b+t}(x)}{\mathbf{p}_a(x)\,\mathbf{e}_{b+s}(x)} = \frac{\left[(t-s)(1+|x|)^\sigma\right]^{\frac{a}{\sigma}}}{(t-s)^{\frac{a}{\sigma}}}e^{-(t-s)(1+|x|)^\sigma} \le e^{-\frac{a}{\sigma}\left(\frac{a}{\sigma}\right)^{\frac{a}{\sigma}}},\tag{A.22}
$$

which holds uniformly over $x \in \mathbb{R}^2$, and where in the last step we used that $y \mapsto y^{a/\sigma}e^{-y}$ is bounded by $e^{-a/\sigma} (a/\sigma)^{a/\sigma}$.

From [\(A.19\)](#page-57-1), we deduce

$$
||v_t||_{\mathscr{C}^{\alpha}(e_{b+t}^{\sigma})} \le ||X_t||_{\mathscr{C}^{\alpha}(e_{b+t}^{\sigma})} + \lambda \int_0^t ||K_{t-s} \mathcal{N}^1[v_s, w_s]||_{\mathscr{C}^{\alpha}(e_{b+t}^{\sigma})} ds
$$

\n
$$
\lesssim ||X_t||_{\mathscr{C}^{\alpha}(p_a)} + \lambda \int_0^t ||v_s w_s||_{\mathscr{C}^{\alpha}(e_{b+t}^{\sigma})} ds
$$

\n
$$
\lesssim ||X_t||_{\mathscr{C}^{\alpha}(p_a)} + \lambda \int_0^t (t-s)^{-a/\sigma} ||v_s||_{\mathscr{C}^{\alpha}(e_{b+s}^{\sigma})} ||w_s||_{\mathscr{C}^{\alpha}(p_a)} ds
$$

\n
$$
\lesssim ||X_t||_{\mathscr{C}^{\alpha}(p_a)} + \lambda ||w||_{\mathscr{C}([0,T],\mathscr{C}^{\alpha}(p_a))} \int_0^t (t-s)^{-a/\sigma} ||v_s||_{\mathscr{C}^{\alpha}(e_{b+s}^{\sigma})} ds,
$$

\n(A.23)

where in the second step we used the convolution bound (A, ς) while in the third the product rule [\(A.6\)](#page-53-7) and [\(A.22\)](#page-58-0). Now, recalling that $a/\sigma < 1$, Grönwall's lemma [\[Hen81,](#page-71-1) Lemma 7.1.1] implies the existence of $C = C(\rho, \lambda, a, b, \sigma) > 0$ for which [\(A.21\)](#page-57-2) holds, thus the proof is concluded. \Box

We now have all the elements to complete the proof of Proposition [2.3.](#page-14-1)

Proof of Proposition [2.3.](#page-14-1) The proof of the statement is divided into three steps which amount to show that 1) the sequence ${u^M}_{M}$ converges to a unique limit u which is a mild solution to [\(2.1\)](#page-13-1) and satisfies [\(2.3\)](#page-14-3), 2) [\(2.1\)](#page-13-1) admits a unique solution in $C_{loc}(\mathbb{R}_+,\mathcal{C}^{\alpha}(\mathfrak{p}_a))$, for $a, \alpha > 0$ (so that it is necessarily given by u) and 3) that u is a Markov process and is stationary. Let us fix a number of parameters which will be used throughout, namely $T > 0$, α , $a, b, \sigma > 0$, $\beta \in [0, 1/2)$ such that $a < \sigma < 1$, and $\alpha' > \alpha$, $\beta' \in (\beta, 1/2)$ and $a' \in (0, a)$.

1) Convergence of $\{u^M\}_M$. The result follows upon proving that the sequence is Cauchy in the right (Banach) space. Adopting the same notations as in the proof of Lemma [A.4,](#page-55-0) for $M, M' \in \mathbb{N}$ we have

$$
u^M_t - u^{M'}_t = X^M_t - X^{M'}_t + \lambda \int_0^t K_{t-s} \, \mathcal{N}^1[u^M_s - u^{M'}_s, u^M_s + u^{M'}_s] \, \mathrm{d} s \, \mathrm{d} s
$$

where $\mathcal{N}^1[\cdot,\cdot]$ is defined according to [\(A.20\)](#page-57-3). Hence, applying Lemma [A.5](#page-57-0) with $v = v^{M,M'} \stackrel{\text{def}}{=}$ $u^M - u^{M'}$, $X = X^{M,M'} = X^M - X^{M'}$ and $w = w^{M,M'} \stackrel{\text{def}}{=} u^M + u^{M'}$ leads to

$$
\sup_{t\in[0,T]}\|v^{M,M'}\|_{\mathscr{C}^{\alpha}(\mathrm{e}^{\sigma}_{b+t})}\leq C\|X^{M,M'}\|_{C([0,T],\mathscr{C}^{\alpha}(\mathrm{p}_a))}\exp\left(C\,\|w^{M,M'}\|_{C([0,T],\mathscr{C}^{\alpha}(\mathrm{p}_a))}^{\frac{1}{1-a/\sigma}}\right)
$$

for some constant $C = C(\rho, \lambda, a, b, \sigma, T)$, which, together with [\(A.16\)](#page-56-0) and [\(A.10\)](#page-53-5), easily implies

$$
\lim_{M,M'\to+\infty} \mathbf{E}\Big[1 \wedge \sup_{t\in[0,T]} \|u^M - u^M'\|_{\mathcal{C}^{\alpha}(e_{b+t}^{\sigma})}\Big] = 0.
$$
 (A.24)

At this point, it remains to show that the convergence takes place in a space with polynomial (instead of exponential) weight. To do so, note that our choices of $\alpha, \alpha', \beta, \beta', a$ and b ensure that the following embeddings hold

$$
C^{\beta'}([0,T],\mathcal{C}^{\alpha'}(p_{\alpha'}))\overset{c}{\hookrightarrow}C^{\beta}([0,T],\mathcal{C}^{\alpha}(p_{a}))\hookrightarrow C([0,T],\mathcal{C}^{\alpha}(e^{\sigma}_{b+T}))
$$

the first even being compact and where the last space contains those functions such that the norm at the left hand side of [\(A.21\)](#page-57-2) is finite. Note that, for generic Banach spaces $X_1 \subset X_2 \subset X_3$ such that the first embedding is compact, it is easy to see that for every $\delta > 0$, there exists a constant $A_{\delta} > 0$ such that

$$
||x||_{X_2} \le \delta ||x||_{X_1} + A_\delta ||x||_{X_3}.
$$

In particular, the above applies to $X_1 = C^{\beta'}([0,T], \mathcal{C}^{\alpha'}(p_{\alpha})), X_2 = C^{\beta}([0,T], \mathcal{C}^{\alpha}(p_a))$ and $X_3 = C([0, T], \mathcal{C}^{\alpha}(e_{b+T}^{\sigma}))$. Therefore, invoking Lemma [A.4,](#page-55-0) we deduce

$$
\mathbf{E}\Big[1\wedge\|u^M-u^{M'}\|_{C^{\beta}([0,T],\mathcal{C}^{\alpha}(\mathbf{p}_a))}\Big]\leq \delta \mathbf{E}[\|u^M-u^{M'}\|_{C^{\beta'}([0,T],\mathcal{C}^{\alpha'}(\mathbf{p}_a))}]+A_{\delta}\mathbf{E}[1\wedge\|u^M-u^{M'}\|_{C([0,T],\mathcal{C}^{\alpha}(e_{b+T}))}]\lesssim \delta+A_{\delta}\mathbf{E}[1\wedge\|u^M-u^{M'}\|_{C([0,T],\mathcal{C}^{\alpha}(e_{b+T}))}],
$$

which, using [\(A.24\)](#page-59-0) and that δ can be taken arbitrarily small, leads to

$$
\lim_{M,M'\to+\infty} \mathbf{E}\Big[1\wedge\|u^M-u^{M'}\|_{C^{\beta}([0,T],\mathcal{C}^{\alpha}(\mathbf{p}_a))}\Big]=0\,.
$$
 (A.25)

The latter implies that the sequence $(u^M)_M$ converges in probability in $C^\beta([0,T], \mathscr{C}^\alpha(p_a))$ to a unique limit u which, by $(A.16)$ and Fatou's lemma, satisfies (2.3) , and, up to passing to a subsequence, the convergence is almost sure. A diagonal argument then guarantees that we can find an event of full **P**-measure in which the convergence takes place in $C^{\beta}([0,T], \mathcal{C}^{\alpha}(\mathbf{p}_a))$ for every $T > 0$, i.e. in $C_{\text{loc}}^{\beta}(\mathbb{R}_+, \mathcal{C}^{\alpha}(\mathbf{p}_a))$, so that **P**-a.s. $u \in C_{\text{loc}}^{\beta}(\mathbb{R}_+, \mathcal{C}^{\alpha}(\mathbf{p}_a))$. Further, u is $(\mathcal{F}_t)_{t\geq 0}$ -adapted because each u^M is, and it can be easily seen to satisfy [\(2.2\)](#page-13-3) thanks to [\(A.10\)](#page-53-5) and the continuity of the map $C_{\text{loc}}^{\beta}(\mathbb{R}_+,\mathcal{C}^{\alpha}(\mathbf{p}_a)) \ni f \mapsto \int_0^t K_{t-s}\mathcal{N}^1[f_s]ds$. In other words, u is a mild solution to [\(2.1\)](#page-13-1) driven by $(n, \vec{\xi})$.

2) Path by path uniqueness. It is an immediate consequence of the fact that **P**-a.s. $X, u \in$ $C(\mathbb{R}_+,\mathcal{C}^{\alpha}(\mathfrak{p}_a))$ and Lemma [A.5](#page-57-0) applied with $v = u_1 - u_2$, $X \equiv 0$ and $w = u_1 + u_2$.

3) Skew-reversibility, stationarity and Markovianity. Since for every $t \geq 0$, u_t^M converges in probability to u_t in $\mathcal{C}^\alpha(\mathbf{p}_a)$ and skew-reversibility holds for u^M it also holds for the limit. Further, convergence in probability implies convergence in law, so that, since by Lemma [A.4,](#page-55-0) $u_t^M \stackrel{\text{law}}{=} \eta_M^1$, and η_M^1 converges in law to η^1 (because of Lemma [A.10,](#page-53-5) noticing that $X_0^M = \eta_M^1$ and $X_0 = \eta^1$), we deduce that u_t and η^1 have the same law, which implies stationarity.

For the Markov property, note that u is a measurable function of $(\eta, \vec{\xi})$, so that it can be written as $u = \Phi(\eta, \vec{\xi})$. Now, defining the translation $\theta_{\tau} f \stackrel{\text{def}}{=} (f_{t+\tau})_{t\geq 0}$ for $\tau \geq 0$, it is immediate to see that $\theta_\tau \Phi(\eta, \vec{\xi})$ and $\Phi(u(\tau), \theta_\tau \vec{\xi})$ are both mild solutions to [\(2.1\)](#page-13-1) driven by $(u_\tau, \theta_\tau \vec{\xi}^1)$ so that, by uniqueness, they coincide. In particular, since u_{τ} is \mathscr{F}_{τ} -measurable while $\theta_{\tau}\vec{\xi}^{\,1}$ is independent of \mathcal{F}_{τ} and has the same law as $\vec{\xi}^1$, for bounded functionals F it holds

$$
\mathbf{E}[F(\theta_\tau \Phi(\eta, \vec{\xi})) | \mathcal{F}_\tau] = \mathbf{E}[F(\Phi(u_\tau, \vec{\xi}))].
$$

which means that the process is Markov, hence the proof is completed.

A.4 Decay properties of a non-local PDE

In this appendix, we determine decay properties of the non-local PDE [\(2.13\)](#page-16-0) derived in the proof of Proposition [2.4.](#page-14-4) Let us take a slightly more general approach, namely consider the PDE

$$
\partial_t Z = \left(\frac{1}{2}\Delta + L_{\Phi,f}\right)Z + gZ, \qquad Z_0 = h, \qquad (A.26)
$$

where $Z = Z_t(x)$ for $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^2$, $f \ge 0$, $g \ge 0$, $h \ge 0$ are locally bounded functions on \mathbb{R}^2 , and L is the operator acting on functions $u: \mathbb{R}^2 \to \mathbb{R}$ as

$$
L_{\Phi,f}u(x) \stackrel{\text{def}}{=} f(x) \int \Phi(x-y)[u(y)-u(x)] \mathrm{d}y \tag{A.27}
$$

for Φ a non-negative, compactly supported and even function. Under suitable assumptions on f, g, h , the solution Z decays exponentially fast, as the next lemma shows.

Lemma A.6 Let Φ , f, g, h be functions on \mathbb{R}^2 such that Φ is non-negative, compactly supported and even, f,g are non-negative and such that there exists $\delta\in[1,1/2)$ for which $\sup_x|f(x)p_\delta(x)|\vee$ $\sup_x |g(x) \mathsf{p}_{\delta}(x)| < \infty$, and h is non-negative, bounded and compactly supported in a ball of *radius* $r_h > 0$. Then, for every $T \geq 0$ and $\nu > 0$ there exists a positive constant $C =$ $C(\Phi, \delta, T, \nu) < \infty$ *such that for all* $|x| \geq 2r_h$ *and* $t \leq T$ *, we have*

$$
Z_t(x) \le C \|h\|_{\infty} e^{-C|x|}.
$$
\n(A.28)

Proof. Throughout the proof we will denote by C a generic constant depending only on the parameters in the statement and whose value might change from line to line.

As mentioned in the proof of Proposition [2.4,](#page-14-4) $L_{\Phi, f}$ is the generator of a pure jump Markov process on \mathbb{R}^2 that has rate $f(x)\Phi(y-x)dy$ of jumping from x to dy. Therefore, $(1/2)\Delta + L_{\Phi,f}$ is the generator of a Markov process Y on \mathbb{R}^2 that has a Brownian and a jump part, the Brownian noise being independent from the Poisson jump noise. We call \mathbb{P}_{x}^{Y} the law of Y with initial condition x . The reader can then easily check that the following Feynman-Kac formula holds

$$
Z_t(x) = \mathbb{E}_x^Y \left[h(Y_t) e^{\int_0^t g(Y_s) \, ds} \right]
$$

which, by Cauchy-Schwarz, implies

$$
Z_t(x)^2 \le ||h||_{\infty}^2 \mathbb{P}_x^Y(|Y_t| \le r_h) \mathbb{E}_x^Y \left[e^{2 \int_0^t g(Y_s) ds} \right] \tag{A.29}
$$

 \Box

and we will separately bound the last two factors at the right hand side starting from the first.

Let $x \in \mathbb{R}^2$ be such that $|x| \ge 2r_h$. As Y starts at x, on the event $|Y_t| \le r_h$, Y has to move at least at distance $|x|/2$ in the time-interval [0, t]. Let N be the number of jumps that the process does in [0, t] starting from positions smaller than $|x|+r_{\Phi}$ and E be the event that $N > |x|/(4r_{\Phi})$, r_{Φ} being the radius of the support of Φ , which is finite by assumption. We claim that

$$
\mathbb{P}_x^Y(E) \lesssim e^{-C|x|},\tag{A.30}
$$

for C some constant depending on all the parameters in the statement, but independent of x . To see why [\(A.30\)](#page-61-0) holds, note that N is dominated by a Poisson random variable X of parameter $\lambda \lesssim t \sup\{f(y), |y| \leq |x| + r_{\Phi}\} \lesssim t|x|^{2\delta}$. Now, an elementary estimate then gives that the probability of $X > k$ is upper bounded by $\lambda^k/k!$, which, via an application of Stirling's formula, implies [\(A.30\)](#page-61-0) since $2\delta < 1$. On the other hand, Y's jumps are at most of size r_{Φ} , hence, on E^c , the contribution to $|Y_t - x|$ coming from the jumps counted by N is at most $Nr_{\Phi} \leq |x|/4$ and the Brownian noise must do the rest. Thus,

$$
\mathbb{P}_x^Y(|Y_t| \le r_h \; ; \; E^c) \le \mathbb{P}_0^B \Big(\sup_{0 < s_1 < s_2 < t} |B_{s_2} - B_{s_1}| \ge |x|/4 \Big) \lesssim e^{-C|x|^2} \,. \tag{A.31}
$$

We now turn to the second factor in $(A.29)$ for which, in view of $(A.30)$ and $(A.31)$, it suffices to show that

$$
\mathbb{E}_x^Y \left[e^{2 \int_0^t g(Y_s) \, ds} \right] \lesssim e^{C|x|^\gamma} \tag{A.32}
$$

for some γ < 1. Denote by M the maximum of |Y| in the interval [0, t]. Then, using that q grows at most as $|x|^{2\delta}$, we have

$$
\mathbb{E}_x^Y\Big[e^{2\int_0^t g(Y_s)ds}\Big] \lesssim \mathbb{E}_x^Y\Big[e^{CtM^{2\delta}}\Big] \leq e^{Ct|x|^{2\delta}} + \sum_{n>2|x|} \mathbb{P}_x^Y(M \in [n-1,n]) e^{Ctn^{2\delta}}.
$$

To control the probability of the event $\{M \in [n-1,n]\}$, we argue as above. If $M \in [n-1,n]$, then either Y has $n/(4r_{\Phi})$ jumps from positions with norm smaller than $n + r_{\varphi}$, or the Brownian noise induces a displacement of at least $n/2$ in time t. The latter event has probability \lesssim $\exp(-n^2/2t)$. The probability of the former event is upper bounded by the probability that a Poisson random variable of parameter $\lambda \lesssim tn^{2\delta}$ is larger than $n/(4r_{\Phi})$, which decays superexponentially in *n*, as mentioned above. Therefore, $\mathbb{P}_{x_0}^Y(M \in [n-1,n])$ decays at least exponentially in n and the sum is small since $2\delta < 1$. It follows that [\(A.32\)](#page-61-2) holds for $\gamma = 2\delta < 1$, and the proof of the statement is complete. \Box

Appendix B The action of \mathcal{L}^{τ} : proofs

In this appendix we provide the proof of Lemma [2.8,](#page-19-3) which will be split in two steps. Let us begin with the following.

Lemma B.1 *For any* $n \in \mathbb{N}$ *and* $\psi \in \mathcal{S}(\mathbb{R}^{2n}) \cap \Gamma_n L_\tau^2$, *it holds that* $\mathcal{L}^\tau \psi = (\mathcal{S}^\tau + \mathcal{A}_+^\tau + \mathcal{A}_-^\tau)\psi$, *where the operators at the right hand side are defined according to formulas* [\(2.22\)](#page-19-2)*-*[\(2.24\)](#page-20-1)*.*

Proof. The proof follows a (by now) standard procedure which has been implemented in a number of references for slightly different equations (see e.g. [\[GP20,](#page-71-2) [CES21,](#page-69-0) [CGT24\]](#page-70-2)), so we refrain from carrying out the (long but) straightforward computations. The main difference with respect to the literature listed above is that our equation is defined on the full space and the Gaussian analysis is done with respect to a regularised measure but the bulk of the argument is exactly the same. At first, one considers functionals of the form $I_n^{\tau}(h^{\otimes n}) = n! H_n(\eta^{\tau}(h))$ (with H_n the $n-th$ Hermite polynomial) for $h \in \mathcal{S}(\mathbb{R}^2)$ and applies Itô's formula to $H_n(u_t^\tau(h))$. Then, one reads off the drift of the latter from which one deduces the action of \mathcal{L}^{τ} on $I_n^{\tau}(h^{\otimes n})$. By basic calculus on the *n*-th Wiener-Itô integrals I_n^{τ} (and in particular the product formula [Nuao6, Proposition 1.1.3]) it is then possible to single out the contributions which respectively give rise to S^{τ} , \mathcal{A}_{-}^{τ} and \mathcal{A}_{-}^{τ} . Hence, the equality $\mathcal{L}^{\tau} = S^{\tau} + \mathcal{A}_{+}^{\tau} + \mathcal{A}_{-}^{\tau}$ holds on functionals of the form $h^{\otimes n}$ which then can be extended to $\mathcal{S}(\mathbb{R}^{2n}) \cap \Gamma_n L^2_\tau$ by polarisation.

In order to extend the definition of the operators S^{τ} , \mathcal{A}_{+}^{τ} and \mathcal{A}_{-}^{τ} to the whole of $\Gamma \mathbf{H}_{\tau}$, show that they satisfy the stated symmetry properties and verify that ΓH_{τ} is indeed a core, we will need the following lemma which holds for $\tau > 0$ *fixed*.

Lemma B.2 *The operators* S^{τ} , A^{τ}_+ *and* A^{τ}_- *are well-defined and continuous on* $\Gamma \mathbf{H}_{\tau}$ *and there exists a constant* $C = C(\tau) > 0$ *for all* $\psi \in \Gamma \mathbf{H}_{\tau} \cap \Gamma_n L^2_{\tau}$

$$
\|\mathcal{A}_{\pm}^{\tau}\psi\|_{\tau} \leq C n^{3/2} \|\psi\|_{\tau} . \tag{B.1}
$$

Furthermore, the graded sector condition [\[KLO12,](#page-72-0) Section 2.7.4] holds for fixed τ, i.e. there exists a constant $C = C(\tau) > 0$ *such that, for all* $\psi \in \Gamma_n H^1_\tau$ *, we have*

$$
\|\mathcal{A}^{\tau}_{\pm}\psi\|_{\Gamma H_{\tau}^{-1}} \le C\sqrt{n} \|\psi\|_{\Gamma H_{\tau}^{1}}.
$$
 (B.2)

Remark B.3 The dependence on τ in the graded sector condition [\(B.2\)](#page-62-1) can be explicitly tracked and it can be shown that $C = C(\tau) \sim (\log \tau)^{1/2}$. That is, the graded sector condition is *not satisfied uniformly in* τ (see also the discussion in Remark [3.5\)](#page-28-0).

Proof. Concerning well-posedness and continuity, they are obvious for S^{τ} , while for A^{τ}_+ and $A^{\tau}_$ the argument is similar so we will only focus on the former. Let $\gamma \ge 0$ and $\psi \in \mathcal{S}(\mathbb{R}^{2n}) \cap \Gamma_n L^2_{\tau}$. By Cauchy-Schwarz, we get

$$
\| \mathcal{A}_{+}^{\tau} \psi \|_{\Gamma H_{\tau}^{\gamma}}^{2} \lesssim_{\tau} (n+1) n^{2} n! \int \Xi_{n}^{\tau} (\mathrm{d} p_{1:n}) |\hat{\psi}(p_{1:n})|^{2} \times \\ \times \int |p_{1}|^{2} (1 + |\sqrt{R_{\tau}} (p_{1} - q, q, p_{2:n})|^{2})^{\gamma} \Theta_{\tau}(q) \Theta_{\tau}(p_{1} - q) \mathrm{d}q. \tag{B.3}
$$

Now, bounding $|p_1|^2 \lesssim |q|^2 + |p_1 - q|^2$ and using that Θ_τ decays fast (recall its definition in [\(1.18\)](#page-10-1)), the integral over q can be bounded above by a τ -dependent constant times $(1 + |\sqrt{R_{\tau}}p_{1:n}|^2)^{\gamma}$, from which we deduce

$$
\|\mathcal{A}_{+}^{\tau}\psi\|_{\Gamma H_{\tau}^{\gamma}} \lesssim_{\tau} n^{3/2} \|\psi\|_{\Gamma H_{\tau}^{\gamma}}^2.
$$
 (B.4)

In particular, by taking $\gamma = 2$, we conclude that the operator \mathcal{A}_{+}^{τ} can be extended by continuity to a map from $\Gamma \mathbf{H}_{\tau}$ to $\Gamma H_{\tau}^{\gamma} \subset \Gamma L_{\tau}^2$, and, by taking $\gamma = 0$, satisfies [\(B.1\)](#page-62-0).

At last it remains to argue [\(B.2\)](#page-62-1). The same bound was obtained in [\[CGT24,](#page-70-2) Lemma 2.4] in the periodic setting, but the estimate can be similarly shown in the present context (for τ fixed!) and the proof is thus omitted. \Box

We are now ready to complete the proof of Lemma [2.8.](#page-19-3)

Proof of Lemma [2.8.](#page-19-3) We have shown above that on $\mathcal{S}(\mathbb{R}^{2n}) \cap \Gamma_n L_\tau^2$, \mathcal{L}^τ coincides with \mathcal{S}^τ + $\mathcal{A}_{+}^{\tau} + \mathcal{A}_{-}^{\tau}$. Lemma [B.2](#page-62-2) guarantees that the operators \mathcal{S}^{τ} , \mathcal{A}_{+}^{τ} and \mathcal{A}_{-}^{τ} are continuous from $\Gamma \mathbf{H}_{\tau}$ to ΓL^2_τ so that, since by Lemma [2.6](#page-19-0) \mathcal{L}^τ is closed, we deduce that $\Gamma \mathbf{H}_\tau \subset \text{Dom}(\mathcal{L}^\tau)$ and that, on ΓH_{τ} , $\mathcal{L}^{\tau} = \mathcal{S}^{\tau} + \mathcal{A}_{+}^{\tau} + \mathcal{A}_{-}^{\tau}$. The symmetry properties can be straightforwardly checked from the formulas [\(2.22\)](#page-19-2)-[\(2.24\)](#page-20-1) so that we only need to prove that ΓH_{τ} is a core for \mathcal{L}^{τ} , which amounts to show that the closure of \mathcal{L}^{τ} on $\Gamma \mathbf{H}_{\tau}$ coincides with \mathcal{L}^{τ} .

At first, we claim that the closure of the range of $(1 - \mathcal{L}^{\tau})$ restricted to $\Gamma \mathbf{H}_{\tau}$ is dense in ΓL_{τ}^2 . We will prove such claim at the end, but first we argue that it implies the result. Let $x \in \text{Dom}(\mathcal{L}^{\tau})$ (so that $(1 - \mathcal{L}^{\tau})x \in \Gamma L_{\tau}^2$) and let $(x_n)_{n \geq 1}$ be a sequence in $\Gamma \mathbf{H}_{\tau}$ such that $(1 - \mathcal{L}^{\tau})x_n \to (1 - \mathcal{L}^{\tau})x$ in ΓL^2_{τ} , whose existence is guaranteed by the claim above. To conclude, it only remains to show that $(x_n)_{n\geq 1}$ converges to x in ΓL^2_τ . Notice that by [\(2.30\)](#page-21-5) (which is applicable since, for every $n, x_n \in \overline{\mathbf{\Gamma}} \mathbf{H}_{\tau} \subset \Gamma H_{\tau}^1 \cap \text{Dom}(\mathcal{L}^{\tau})$), we have

$$
||x_n - x_m||_{\tau} \lesssim ||(1 - \mathcal{L}^\tau)x_n - (1 - \mathcal{L}^\tau)x_m||_{\tau},
$$

so that $(x_n)_{n\geq 1}$ is Cauchy in ΓL^2_τ . Let $\tilde{x} \in \Gamma L^2_\tau$ be the corresponding limit. As \mathcal{L}^τ is closed, $\tilde{x} \in \text{dom}(\mathcal{L}^{\tau})$ and we have $(1 - \mathcal{L}^{\tau})\tilde{x} = (1 - \mathcal{L}^{\tau})x$. Hence, applying the resolvent to both sides we obtain $\tilde{x} = x$ as desired.

We are left to prove the claim. For $n \ge 1$, let us define the operator \mathcal{L}_n^{τ} acting on $\Gamma_{\le n} L_{\tau}^2 \stackrel{\text{def}}{=} \bigoplus_{k=0}^n \Gamma_k L_{\tau}^2$ as $_{k=0}^{n}$ $\Gamma_{k}L_{\tau}^{2}$ as

$$
\mathcal{L}_n^{\tau} \stackrel{\text{def}}{=} P^n(\mathcal{S}^{\tau} + \mathcal{A}^{\tau})P^n,
$$

where P^n is the orthogonal projection of ΓL^2_τ onto $\Gamma_{\leq n} L^2_\tau$. Now, $-\delta^\tau$ is positive self-adjoint while P^n of $T P^n$ is skew-symmetric and bounded for τ fixed (see [\(B.4\)](#page-62-3)), therefore $(1 - \mathcal{L}_n^{\tau})^{-1}$ is a bounded operator on $\Gamma_{\leq n} L^2_\tau$, taking values in $\Gamma_{\leq n} H^2_\tau (\subset \Gamma \mathbf{H}_\tau)$. Note that the inclusion holds because of the projection: any element in $\Gamma_{\leq n} H_{\tau}^{\overline{2}}$ is compactly supported in chaos so that, in particular, it decays sufficiently fast. Let us fix $\varphi \in \Gamma L^2_\tau$ and set $r^{(n)} \stackrel{\text{def}}{=} (1 - \mathcal{L}_n^{\tau})^{-1} P^n \varphi \in \Gamma L^2_\tau$ $\Gamma_{\leq n}H_{\tau}^2$. A consequence of [\[CGT24,](#page-70-2) Proposition 2.8] is that for any $\alpha \geq 0$, there exists a constant $C = C(\tau) > 0$ such that $r^{(n)}$ satisfies

$$
\sup_{k,n} (1 + k^{\alpha}) \|r^{(n)}\|_{\tau} \le C.
$$
 (B.5)

Now, observe that

$$
\|(1-\mathcal{L}^{\tau})r^{(n)}-\varphi\|_{\tau}\leq \|(Id-P^n)\varphi\|_{\tau}+\|\mathcal{A}_+^{\tau}r_n^{(n)}\|_{\tau}\lesssim \|(Id-P^n)\varphi\|_{\tau}+n^{3/2}\|r_n^{(n)}\|_{\tau},
$$

where in the second bound we exploited $(B.1)$. But now, the two terms at the right hand side vanish as $n \to \infty$ - the first since $\varphi \in \Gamma L^2_\tau$, and the second by the polynomial decay in n given by [\(B.5\)](#page-63-0). Thus, the proof is concluded. \Box

Appendix C The Replacement Lemmas: integral approximations estimates

In this appendix, we detail the main technical ingredient which is at the heart of the proof of the Replacement Lemmas in Section [3.3.](#page-29-4) Before stating it, let us introduce some useful notations which will be used throughout the section. For $p_{1:n} \in \mathbb{R}^{2n}$, $q \in \mathbb{R}^2$, and $i \in \{1, ..., n\}$, we denote by $q' \stackrel{\text{def}}{=} p_i - q$ and set

$$
\Gamma_i^{\tau} \stackrel{\text{def}}{=} \frac{1}{2} |\sqrt{R_{\tau}} (q, q', p_{1:n} \setminus \{i\})|^2, \qquad \tilde{\Gamma}^{\tau} \stackrel{\text{def}}{=} |\sqrt{R_{\tau}} q|^2 + \frac{1}{2} |\sqrt{R_{\tau}} p_{1:n}|^2, \Gamma_i^{\epsilon_1} \stackrel{\text{def}}{=} \frac{1}{2} |\epsilon_1 \cdot (q, q', p_{1:n} \setminus \{i\})|^2.
$$
\n(C.1)

Notice that for any $a, b, c \in \mathbb{R}^2$ such that $a + b = c$, then $|\sqrt{R_{\tau}}a|^2 + |\sqrt{R_{\tau}}b|^2 \approx |\sqrt{R_{\tau}}a|^2 + |\sqrt{R_{\tau}}b|^2$ | $\sqrt{R_{\tau}}c|^2 \approx |\sqrt{R_{\tau}}b|^2 + |\sqrt{R_{\tau}}c|^2$, and the same holds replacing the scaled norm on \mathbb{R}^2 with $|\cdot|$. In particular, this ensures that, for any $i \in \{1, ..., n\}$, $\Gamma_i^{\tau} \asymp \tilde{\Gamma}^{\tau}$.

Lemma C.1 *Let* Φ : $\mathbb{R}^3 \to \mathbb{R}$ *be a locally Lipschitz function such that there exists* $C_{\Phi} > 0$ *for which for every* $t > 0$, $i \in \{1, 2, 3\}$ *and* $x_1, x_2, x_3 > 0$ *where the partial derivatives of* Φ *are well-defined, it holds*

$$
\Phi(tx_1, tx_2, x_3) = t^{-1} \Phi(x_1, x_2, x_3)
$$
\n(C.2)

$$
|\Phi(x_1, x_2, x_3)| \le C_\Phi \frac{1 \vee \nu_\tau \frac{x_2}{x_1}}{x_1 + M \nu_\tau x_2}, \quad |\partial_i \Phi(x_1, x_2, x_3)| \le C_\Phi \frac{1 \vee \nu_\tau \frac{x_2}{x_1}}{(x_1 + M \nu_\tau x_2) x_i}.
$$
 (C.3)

Then, adopting the notations in [\(C.1\)](#page-64-0)*, uniformly over* $p_{1:n} \in \mathbb{R}^{2n}$, $i \in \{1, ..., n\}$ and $M \geq 0$, *we have*

$$
\left|\frac{\lambda^2 \nu_\tau^{3/2}}{\pi^2} \int_{\mathbb{R}^2} \Phi(1 + \Gamma_i^{\tau}, \Gamma_i^{\mathfrak{e}_1}, L^{\tau}(\Gamma_i^{\tau})) \mathcal{J}_{q,q'}^{\tau} \chi_{q,q'}^{\sharp} (p_{1:n}) \, \mathrm{d}q - \int_0^{L^{\tau}(\frac{1}{2}|\sqrt{R_{\tau}}p_{1:n}|^2)} \int_0^{\nu_\tau^{-1}} \frac{\Phi(1,\varsigma,\ell)}{\pi \sqrt{\varsigma(1 - \nu_\tau \varsigma)}} \mathrm{d}\varsigma \mathrm{d}\ell \right| \lesssim C_{\Phi} \lambda^2 \frac{1 \vee \log(1 + M)}{1 \vee M^{1/2}} \nu_\tau,
$$
\n(C.4)

where the map L^{τ} is given according to [\(3.14\)](#page-29-5), and $\mathcal{J}_{q,q'}^{\tau}$ was introduced in [\(3.2\)](#page-24-2).

Remark C.2 We will apply the previous statement to two specific functions, namely

$$
\Phi(\gamma, \gamma', \ell) = \frac{1}{\gamma + g_M^{\tau}(\ell) \gamma'}, \quad \text{and} \quad \Phi(\gamma, \gamma', \ell) = \frac{f(\ell) \gamma'}{(\gamma + g_M^{\tau}(\ell) \gamma')^2},
$$

for some function f on \mathbb{R}_+ . The former will appear in the Replacement Lemma [3.6](#page-30-1) and the latter in the Recursive Replacement Lemma [3.9.](#page-32-0) It is not hard to see that $(C.2)$ and $(C.3)$ hold for the first with $C_{\Phi} = 1$, and the same is true for the second provided f satisfies the assumptions of Lemma [3.9,](#page-32-0) with $C_{\Phi} = 6 C_f$ and C_f such that [\(3.27\)](#page-32-3) is verified.

In the proof of the above (and in that of Proposition [3.12\)](#page-33-0), we will exploit the basic integral estimates summarised in the next lemma.

Lemma C.3 *Let* γ , $A > 0$ *and* $M > 0$ *, then for every* $\tau > 0$ *it holds*

$$
\int_{\mathbb{R}^2} \frac{dq}{(A^2 + |\sqrt{R_\tau}q|^2 + M\nu_\tau |\mathfrak{e}_1 \cdot q|^2)^2} \lesssim \nu_\tau^{-1/2} \frac{1}{A^2} \frac{1}{1 \vee M^{1/2}} \tag{C.5}
$$

$$
\int_{\mathbb{R}^2} \frac{1_{\{\nu_\tau^{1/2} \mid \mathfrak{e}_1 \cdot q \mid \le A\}} dq}{A^2 + |\sqrt{R_\tau} q|^2 + M\nu_\tau |\mathfrak{e}_1 \cdot q|^2} \lesssim \nu_\tau^{-1/2} \frac{1 \vee \log(1+M)}{1 \vee M^{1/2}},
$$
\n(C.6)

$$
\int_{\mathbb{R}^2} \frac{dq}{|\sqrt{R_\tau}q|^2 + M\nu_\tau |\mathfrak{e}_1 \cdot q|^2} \left[\left(\frac{|\mathfrak{e}_1 \cdot q|}{A} \right)^\gamma \wedge \left(\frac{|\mathfrak{e}_1 \cdot q|}{A} \right)^{-\gamma} \right] \lesssim_\gamma \nu_\tau^{-1/2} \frac{1}{1 \vee M^{1/2}}, \quad (C.7)
$$

$$
\int_{\mathbb{R}^2} \frac{\mathrm{d}q}{|\sqrt{R_\tau}q|^2 + M\nu_\tau |\mathfrak{e}_1 \cdot q|^2} \Big[\Big(\frac{|\sqrt{R_\tau}q|}{A} \Big)^\gamma \wedge \Big(\frac{|\sqrt{R_\tau}q|}{A} \Big)^{-\gamma} \Big] \lesssim_\gamma \nu_\tau^{-1/2} \frac{1}{1 \vee M^{1/2}}. \tag{C.8}
$$

Proof. For the first three bounds, write $q = (x, y) \in \mathbb{R}^2$ and use the estimate $\sqrt{R_\tau}q|^2 + M\nu_\tau |\mathfrak{e}_1 \cdot \mathfrak{e}_2|$ $|q|^2 = \frac{\nu_\tau}{2}x^2 + \frac{1}{2}y^2 + M\nu_\tau x^2 \gtrsim y^2 + (1 \vee M)\nu_\tau x^2$. Then, for [\(C.5\)](#page-65-0), we have

$$
\begin{aligned} &\int_{\mathbb{R}^2}\frac{\mathrm{d} q}{(A^2+|\sqrt{R_\tau}q|^2+M\nu_\tau|\mathfrak{e}_1\cdot q|^2)^2} \lesssim \int_{\mathbb{R}}\int_{\mathbb{R}}\frac{1}{A^4+y^4+(1\vee M)^2\nu_\tau^2 x^4}\mathrm{d} x\mathrm{d} y\\ &\lesssim \frac{\nu_\tau^{-1/2}}{1\vee M^{1/2}}\int_{\mathbb{R}}\frac{\mathrm{d} y}{(A^4+y^4)^{3/4}} \lesssim \frac{\nu_\tau^{-1/2}}{1\vee M^{1/2}}\int_{\mathbb{R}}\frac{\mathrm{d} y}{A^3+y^3} \lesssim \frac{\nu_\tau^{-1/2}}{A^2(1\vee M^{1/2})} \end{aligned}
$$

where in the second and the last step we used the elementary bound

$$
\int_{\mathbb{R}} \frac{\mathrm{d}u}{a|u|^{\alpha} + b} \lesssim_{\alpha} \frac{1}{a^{1/\alpha} b^{1-1/\alpha}} \tag{C.9}
$$

which holds for any $\alpha > 1$, uniformly over $a, b > 0$.

Concerning [\(C.6\)](#page-65-1), we proceed similarly, i.e.

$$
\begin{aligned} &\int_{\mathbb{R}^2}\frac{1_{\{\nu_{\tau}^{1/2}\mid\mathfrak{e}_{1}:q\mid\leq A\}}\,\mathrm{d}q}{A^2+|\sqrt{R_\tau}q|^2+M\nu_{\tau}|\mathfrak{e}_{1}\cdot q|^2}&\lesssim \int_{\mathbb{R}}\int_{\mathbb{R}}\frac{1_{\{\nu_{\tau}^{1/2}\mid x\mid\leq A\}}}{A^2+y^2+(1\vee M)\nu_{\tau}x^2}\mathrm{d}x\mathrm{d}y}\\ &\lesssim \int_{\mathbb{R}}\frac{1_{\{\nu_{\tau}^{1/2}\mid x\mid\leq A\}}}{\sqrt{A^2+(1\vee M)\nu_{\tau}x^2}}\mathrm{d}x\lesssim \int_{\mathbb{R}}\frac{1_{\{\nu_{\tau}^{1/2}\mid x\mid\leq A\}}}{A+(1\vee M^{1/2})\nu_{\tau}^{1/2}|x|}\mathrm{d}x\lesssim \nu_{\tau}^{-1/2}\frac{1\vee\log(1+M)}{1\vee M^{1/2}} \end{aligned}
$$

the integral in y being controlled via [\(C.9\)](#page-65-2) with $\alpha = 2$ and the last via a direct computation. For [\(C.7\)](#page-65-3), we use once again [\(C.9\)](#page-65-2) with $\alpha = 2$ on the integral in y, so that

$$
\int_{\mathbb{R}^2} \frac{dq}{|\sqrt{R_{\tau}}q|^2 + M\nu_{\tau}|\mathfrak{e}_1 \cdot q|^2} \left[\left(\frac{|\mathfrak{e}_1 \cdot q|}{A} \right)^{\gamma} \wedge \left(\frac{|\mathfrak{e}_1 \cdot q|}{A} \right)^{-\gamma} \right] \leq \int_{\mathbb{R}^2} \frac{(|x|/A)^{\gamma} \wedge (|x|/A)^{-\gamma}}{y^2 + (M \vee 1)\nu_{\tau}x^2} dy dx \lesssim \frac{\nu_{\tau}^{-1/2}}{1 \vee M^{1/2}} \int_{\mathbb{R}_+} \frac{(x/A)^{\gamma} \wedge (x/A)^{-\gamma}}{x} dx
$$
 (C.10)

from which the bound follows by splitting the cases $x < A$ and $x > A$.

At last, in the integral at the left hand side of [\(C.8\)](#page-65-4) we first apply the change of variables $\tilde{q} = \sqrt{R_{\tau}} q$ and then pass to polar coordinates, so that it equals

$$
\nu_{\tau}^{-1/2} \int_{\mathbb{R}^2} \frac{(|q|/A)^{\gamma} \wedge (|q|/A)^{-\gamma}}{|q|^2 + M |\mathfrak{e}_1 \cdot q|^2} dq
$$

The Replacement Lemmas: integral approximations estimates **67**

$$
\lesssim \nu_{\tau}^{-1/2} \int_0^{2\pi} \frac{d\theta}{1 + (M \vee 1) \cos(\theta)^2} \int_{\mathbb{R}_+} \frac{(r/A)^{\gamma} \wedge (r/A)^{-\gamma}}{r} dr \lesssim_{\gamma} \frac{\nu_{\tau}^{-1/2}}{1 \vee M^{1/2}} :
$$

the integral over the radial component was bounded as in $(C.10)$, while that of the angular one can be obtained by a direct computation. \Box

We are now ready to prove Lemma [C.1.](#page-64-3)

Proof of Lemma [C.1.](#page-64-3) The bulk of the proof consists of showing that

$$
\frac{\lambda^2 \nu_{\tau}^{3/2}}{\pi^2} \bigg| \int_{\mathbb{R}^2} \Phi(1 + \Gamma_i^{\tau}, \Gamma_i^{\mathfrak{e}_1}, L^{\tau}(\Gamma_i^{\tau})) \mathcal{J}_{q,q'}^{\tau} \chi_{q,q'}^{\sharp} (p_{1:n}) \, \mathrm{d}q
$$
\n
$$
- \int_{\mathbb{R}^2} \frac{\Phi(1 + \tilde{\Gamma}^{\tau}, \frac{1 + \tilde{\Gamma}^{\tau}}{|\sqrt{R_{\tau}q}|^2} |\mathfrak{e}_1 \cdot q|^2, L^{\tau}(\tilde{\Gamma}^{\tau}))}{1 + (1 + \tilde{\Gamma}^{\tau})\tau^{-1}} \mathrm{d}q \bigg| \lesssim C_{\Phi} \lambda^2 \frac{1 \vee \log(1 + M)}{1 \vee M^{1/2}} \nu_{\tau} \,.
$$
\n(C.11)

To see how the latter implies [\(C.4\)](#page-64-4), let us manipulate the second term above, which we denote by I_{Φ} . Using the scaling relation [\(C.2\)](#page-64-1) and the symmetry of the integrand with respect to the inversion $q \mapsto -q$, we see that I_{Φ} equals

$$
\frac{\lambda^2\nu_\tau^{3/2}}{\pi^2}\int_{\mathbb{R}^2}\frac{\Phi(1,\frac{|\mathfrak{e}_1\cdot q|^2}{|\sqrt{R_\tau}q|^2},L^\tau(\tilde{\Gamma}^\tau))}{(1+\tilde{\Gamma}^\tau)(1+(1+\tilde{\Gamma}^\tau)\tau^{-1})}\,\mathrm{d}q = \frac{4\lambda^2\nu_\tau^{3/2}}{\pi^2}\int_{\mathbb{R}_+^2}\frac{\Phi(1,\frac{|\mathfrak{e}_1\cdot q|^2}{|\sqrt{R_\tau}q|^2},L^\tau(\tilde{\Gamma}^\tau))}{(1+\tilde{\Gamma}^\tau)(1+(1+\tilde{\Gamma}^\tau)\tau^{-1})}\,\mathrm{d}q\,.
$$

Then, we perform two change of variables, first $(\sigma, \varsigma) \stackrel{\text{def}}{=} (\tilde{\Gamma}^{\tau}, |\mathfrak{e}_1 \cdot q|^2 / |\sqrt{R_{\tau}} q|^2)$, for which the determinant of the Jacobian is $d\sigma d\zeta = 4\sqrt{\zeta(1 - \nu_\tau \zeta)} dq$, and then $\ell \stackrel{\text{def}}{=} L^\tau(\sigma)$, for which instead $d\ell = \frac{\lambda^2 \nu_\tau^3/2}{\pi} [(1+\sigma)(1+(1+\sigma)\tau^{-1})]^{-1} d\sigma$. As a consequence we obtain

$$
I_{\Phi} = \frac{\lambda^2 \nu_{\tau}^{3/2}}{\pi^2} \int_0^{\nu_{\tau}^{-1}} \int_{|\sqrt{R_{\tau}} p_{1:n}|^2/2}^{+\infty} \frac{\Phi(1,\varsigma,L^{\tau}(\sigma))}{(1+\sigma)(1+(1+\sigma)/\tau)} d\sigma \frac{d\varsigma}{\sqrt{\varsigma(1-\nu_{\tau}\varsigma)}}
$$

=
$$
\int_0^{L^{\tau}(\frac{1}{2}|\sqrt{R_{\tau}} p_{1:n}|^2)} \int_0^{\nu_{\tau}^{-1}} \frac{\Phi(1,\varsigma,\ell)}{\pi\sqrt{\varsigma(1-\nu_{\tau}\varsigma)}} d\varsigma d\ell,
$$
 (C.12)

which indeed coincides with the second integral at the left hand side of $(C.4)$.

We are thus left to show $(C.11)$. Heuristically, the reason why we expect it to hold is that only the regions $|e_1 \cdot q|, |e_1 \cdot q'| \gg 1 \vee |p_{1:n}|$ and $|q|, |q'| \ll \tau^{-1}$ should contribute and, on such regions, the integrands are indeed close. To formalise this idea, let $Q = \frac{def}{=} \{ \nu_{\tau}^{1/2} | \mathfrak{e}_1 \cdot q \}$ $10 (1 \vee \sqrt{R_{\tau}} p_{1:n}|)$ and Q_+ be its complement. Assume the following estimates hold

$$
\int_{Q_{-}} dq \frac{C}{1 + |\sqrt{R_{\tau}}q|^{2} + |\sqrt{R_{\tau}}p_{1:n}|^{2} + M\nu_{\tau}|\mathfrak{e}_{1} \cdot q|^{2}} \lesssim C_{\Phi} \frac{1 \vee \log(M)}{1 \vee M^{1/2}} \nu_{\tau}^{-1/2}, \tag{C.13}
$$

$$
\int_{Q_+} dq \frac{C}{|\sqrt{R_\tau}q|^2 + M\nu_\tau |\mathfrak{e}_1 \cdot q|^2} \left| \mathfrak{F}^\tau_{q,q'} - \frac{1}{1 + (1 + \tilde{\Gamma}^\tau)\tau^{-1}} \right| \lesssim C_\Phi \frac{\nu_\tau^{-1/2}}{1 \vee M^{1/2}},\tag{C.14}
$$

The Replacement Lemmas: integral approximations estimates **68**

$$
\int_{Q_+} dq \left| \Phi(1 + \Gamma_i^{\tau}, \Gamma_i^{\mathfrak{e}_1}, L^{\tau}(\Gamma_i^{\tau})) - \Phi(1 + \tilde{\Gamma}^{\tau}, \frac{1 + \tilde{\Gamma}^{\tau}}{|\sqrt{R_{\tau}}q|^2} |\mathfrak{e}_1 \cdot q|^2, L^{\tau}(\tilde{\Gamma}^{\tau})) \right| \lesssim C_{\Phi} \frac{\nu_{\tau}^{-1/2}}{1 \vee M^{1/2}}. (C.15)
$$

In order to deduce $(C.11)$, we begin by splitting the first integral al the left hand side of $(C.11)$ in the regions $Q_$ and $Q_$. For the integral over $Q_$, we bound Φ via the first inequality in [\(C.3\)](#page-64-2), use that $1 \vee \nu_{\tau} \frac{\Gamma_i^{\epsilon_1}}{1+\Gamma_i^{\tau}} \lesssim 1$, and control what is left with [\(C.13\)](#page-66-1). For the integral over Q_+ , note that, on Q_+ , the definition of χ^{\sharp} in [\(3.9\)](#page-26-2) ensures that $\chi^{\sharp}_{q,q'}(p_{1:n}) = 1$. Then, we replace $\mathcal{J}^{\tau}_{q,q'}$ with $[1 + (1 + \tilde{\Gamma}^{\tau}) \tau^{-1}]^{-1}$, by adding and subtracting the corresponding term, and estimate the error by treating Φ as above and exploiting [\(C.14\)](#page-66-2). At last, we want to substitute (Γ_i^{τ} , $\Gamma_i^{\epsilon_1}$ $i^{(1)}$) with $(\tilde{\Gamma}^{\tau}, \frac{1+\tilde{\Gamma}^{\tau}}{1+\sqrt{P_{-a}}})$ $\frac{1+\tilde{\Gamma}^{\tau}}{|\sqrt{R_{\tau}q}|^2}|\mathfrak{e}_1 \cdot q|^2$). Once again, we add and subtract the corresponding term, and control the difference by bounding $[1 + (1 + \tilde{\Gamma}^{\tau}) \tau^{-1}]^{-1} \le 1$ and using [\(C.15\)](#page-67-0). These last two steps are instrumental to reducing the integrand to a form in which the change of variables in $(C.12)$ is possible.

Therefore, we are left to show $(C.13)-(C.15)$ $(C.13)-(C.15)$. The first is the easiest since it can be deduced by simply removing $|\sqrt{R_{\tau}}p_{1:n}|^2$ from the denominator and then applying [\(C.6\)](#page-65-1) from Lemma $C.3.$

For [\(C.14\)](#page-66-2), recall the definition of ϱ_{τ} and Θ_{τ} in [\(1.18\)](#page-10-1), and that $\mathcal{J}_{q,q'}^{\tau} = \Theta_{\tau}(q)\Theta_{\tau}(q')\Theta_{\tau}(p_i)$. Since $\hat{\varrho}_{\tau} \in \mathcal{S}(\mathbb{R}^2)$ and $2\pi \hat{\varrho}_{\tau}(0) = 1$, we have

$$
|\mathcal{F}_{q,q'}^{\tau}-1| \lesssim \tau^{-1/2} \max\{|\sqrt{R_{\tau}}q|,|\sqrt{R_{\tau}}q'|,|\sqrt{R_{\tau}}p_i|\} \lesssim \tau^{-1/2} \sqrt{R_{\tau}}q|, \mathcal{F}_{q,q'}^{\tau} \lesssim \frac{1}{\tau^{-1/2} \max\{|\sqrt{R_{\tau}}q|,|\sqrt{R_{\tau}}q'|,|\sqrt{R_{\tau}}p_i|\}} \lesssim \frac{1}{\tau^{-1/2} |\sqrt{R_{\tau}}q|},
$$

where we also used that on Q_+ , $|\sqrt{R_\tau}q| \asymp \max\{|\sqrt{R_\tau}q|, |\sqrt{R_\tau}q'|, |\sqrt{R_\tau}p_i|\}.$ Therefore,

$$
\left| \mathcal{J}_{q,q'}^{\tau} - \frac{1}{1 + (1 + \tilde{\Gamma})\tau^{-1}} \right| \leq \mathbb{1}_{\{\tau^{-1/2}|\sqrt{R_{\tau}}q| \leq 1\}} \left(|\mathcal{J}_{q,q'}^{\tau} - 1| + \left| 1 - \frac{1}{1 + (1 + \tilde{\Gamma})\tau^{-1}} \right| \right) + \mathbb{1}_{\{\tau^{-1/2}|\sqrt{R_{\tau}}q| > 1\}} \left(\mathcal{J}_{q,q'}^{\tau} + \frac{1}{1 + (1 + \tilde{\Gamma})\tau^{-1}} \right) < \lesssim \tau^{-1/2} |\sqrt{R_{\tau}}q| \, \mathbb{1}_{\{\tau^{-1/2}|\sqrt{R_{\tau}}q| \leq 1\}} + \frac{\mathbb{1}_{\{\tau^{-1/2}|\sqrt{R_{\tau}}q| > 1\}}}{\tau^{-1/2} |\sqrt{R_{\tau}}q|}.
$$

Plugging the above estimate into the left hand side of $(C.14)$, we can upper bound it (modulo an absolute constant) by

$$
\int_{\mathbb{R}^2} \mathrm{d}q \left(\frac{\tau^{-1/2} |\sqrt{R_\tau} q| 1\!\!1_{\{\tau^{-1/2} |\sqrt{R_\tau} q| \leq 1\}}}{|\sqrt{R_\tau} q|^2 + M\nu_\tau |\mathfrak{e}_1 \cdot q|^2} + \frac{(\tau^{-1/2} |\sqrt{R_\tau} q|)^{-1} 1\!\!1_{\{\tau^{-1/2} |\sqrt{R_\tau} q| > 1\}}}{|\sqrt{R_\tau} q|^2 + M\nu_\tau |\mathfrak{e}_1 \cdot q|^2} \right) \lesssim \frac{\nu_\tau^{-1/2}}{1 \vee M^{1/2}} \,,
$$

the last step being a consequence of [\(C.8\)](#page-65-4), applied with $\gamma = 1$.

It remains to prove $(C.15)$. Let us first control the integrand as

$$
|\Phi(1+\Gamma_i^{\tau},\Gamma_i^{\epsilon_1},L^{\tau}(\Gamma_i^{\tau}))-\Phi(1+\tilde{\Gamma}^{\tau},\frac{1+\tilde{\Gamma}^{\tau}}{|\sqrt{R_{\tau}}q|^2}|\mathfrak{e}_1\cdot q|^2,L^{\tau}(\tilde{\Gamma}^{\tau}))|
$$

The Replacement Lemmas: integral approximations estimates **69**

$$
\leq |\Phi(1+\Gamma_i^{\tau}, \Gamma_i^{\epsilon_1}, L^{\tau}(\Gamma_i^{\tau})) - \Phi(1+\tilde{\Gamma}^{\tau}, \Gamma_i^{\epsilon_1}, L^{\tau}(\tilde{\Gamma}^{\tau}))| + |\Phi(1+\tilde{\Gamma}^{\tau}, \Gamma_i^{\epsilon_1}, L^{\tau}(\tilde{\Gamma}^{\tau})) - \Phi(1+\tilde{\Gamma}^{\tau}, \frac{\tilde{\Gamma}^{\tau}}{|\sqrt{R_{\tau}q}|^2} |\mathfrak{e}_1 \cdot q|^2, L^{\tau}(\tilde{\Gamma}^{\tau}))| + |\Phi(1+\tilde{\Gamma}^{\tau}, \frac{\tilde{\Gamma}^{\tau}}{|\sqrt{R_{\tau}q}|^2} |\mathfrak{e}_1 \cdot q|^2, L^{\tau}(\tilde{\Gamma}^{\tau})) - \Phi(1+\tilde{\Gamma}^{\tau}, \frac{1+\tilde{\Gamma}^{\tau}}{|\sqrt{R_{\tau}q}|^2} |\mathfrak{e}_1 \cdot q|^2, L^{\tau}(\tilde{\Gamma}^{\tau}))| \n\leq \frac{C_{\Phi}}{|\sqrt{R_{\tau}q}|^2 + M\nu_{\tau}|\mathfrak{e}_1 \cdot q|^2} \left(\frac{|\Gamma_i^{\tau} - \tilde{\Gamma}^{\tau}|}{|\sqrt{R_{\tau}q}|^2} + \frac{|\Gamma_i^{\epsilon_1} - \frac{\tilde{\Gamma}^{\tau}}{|\sqrt{R_{\tau}q}|^2} |\mathfrak{e}_1 \cdot q|^2|}{|\mathfrak{e}_1 \cdot q|^2} + \frac{1}{|\sqrt{R_{\tau}q}|^2} \right),
$$

where in the last step we used mean value theorem, assumption $(C, 3)$ on the gradient of Φ , which in particular gives

$$
\begin{split} |\partial_{\gamma}\Phi(\gamma,\gamma',L^{\tau}(\gamma))| &\leq |\partial_{1}\Phi(\gamma,\gamma',L^{\tau}(\gamma))| + |\partial_{3}\Phi(t,\gamma',L^{\tau}(\gamma))| \times |\partial_{\gamma}L^{\tau}(\gamma)| \\ &\leq \frac{C_{\Phi}(1\vee \nu_{\tau}\frac{\gamma'}{\gamma})}{(\gamma + M\nu_{\tau}\gamma')\gamma} + \frac{C_{\Phi}(1\vee \nu_{\tau}\frac{\gamma'}{\gamma})}{(\gamma + M\nu_{\tau}\gamma')L^{\tau}(\gamma)} \frac{L^{\tau}(\gamma)}{\gamma} \lesssim C_{\Phi}\frac{1\vee \nu_{\tau}\frac{\gamma'}{\gamma}}{(\gamma + M\nu_{\tau}\gamma')\gamma}, \end{split}
$$

that $\Gamma_i^{\tau} \asymp \tilde{\Gamma}^{\tau} \gtrsim |\sqrt{R_{\tau}}q|^2$ and that, on Q_+ , $\Gamma_i^{\mathfrak{e}_1} \asymp$ $\tilde{\Gamma}^\tau$ $\frac{\Gamma^{\tau}}{|\sqrt{R_{\tau}q}|^2}|\mathfrak{e}_1\cdot q|^2 \gtrsim |\mathfrak{e}_1\cdot q|^2$. Now,

$$
|\Gamma_i^{\tau} - \tilde{\Gamma}^{\tau}| = |\langle \sqrt{R_{\tau}} p_i, \sqrt{R_{\tau}} q \rangle| \leq |\sqrt{R_{\tau}} p_i| |\sqrt{R_{\tau}} q|,
$$

$$
|\Gamma_i^{\mathfrak{e}_1} - |\mathfrak{e}_1 \cdot q|^2| \leq |\mathfrak{e}_1 \cdot p_i| |\mathfrak{e}_1 \cdot q| + \frac{1}{2} |\mathfrak{e}_1 \cdot p_{1:n}|^2,
$$

$$
||\mathfrak{e}_1 \cdot q|^2 - \frac{1 + \tilde{\Gamma}}{|\sqrt{R_{\tau}} q|^2} |\mathfrak{e}_1 \cdot q|^2| \leq \frac{(1 + |\sqrt{R_{\tau}} p_{1:n}|^2) |\mathfrak{e}_1 \cdot q|^2}{|\sqrt{R_{\tau}} q|^2},
$$

so that, putting everything together, we obtain

$$
\int_{Q_{+}} dq \left| \Phi(1 + \Gamma_{i}^{\tau}, \Gamma_{i}^{\epsilon_{1}}, L^{\tau}(\Gamma_{i}^{\tau})) - \Phi(1 + \tilde{\Gamma}^{\tau}, \frac{1 + \tilde{\Gamma}^{\tau}}{|\sqrt{R_{\tau}}q|^{2}} | \mathfrak{e}_{1} \cdot q |^{2}, L^{\tau}(\tilde{\Gamma}^{\tau})) \right|
$$
\n
$$
\lesssim \int_{Q_{+}} \frac{C_{\Phi} dq}{|\sqrt{R_{\tau}}q|^{2} + M\nu_{\tau}| \mathfrak{e}_{1} \cdot q|^{2}} \left(\frac{|\sqrt{R_{\tau}}p_{i}|}{|\sqrt{R_{\tau}}q|} + \frac{|\mathfrak{e}_{1} \cdot p_{i}|}{|\mathfrak{e}_{1} \cdot q|} + \frac{|\mathfrak{e}_{1} \cdot p_{1:n}|^{2}}{|\mathfrak{e}_{1} \cdot q|^{2}} + \frac{1 + |\sqrt{R_{\tau}}p_{1:n}|^{2}}{|\sqrt{R_{\tau}}q|^{2}} \right)
$$
\n
$$
\lesssim C_{\Phi} \frac{\nu_{\tau}^{-1/2}}{1 \vee M^{1/2}}
$$

which holds thanks to $(C.7)$ and $(C.8)$ in Lemma [C.3.](#page-65-6) Therefore, the proof of $(C.15)$ (and of the lemma) is complete. \Box

Acknowledgments

We wish to thank Scott Armstrong, Harry Giles, Massimiliano Gubinelli, Felix Otto and Peter Morfe for enlightnening discussions. G. C. gratefully acknowledges financial support via the UKRI FL fellowship "Large-scale universal behaviour of Random Interfaces and Stochastic Operators" MR/W008246/1. The research of Q. M. and F. T. was funded by the Austrian Science Fund (FWF) 10.55776/F1002. For open access purposes, the authors have applied a CC BY public copyright license to any author accepted manuscript version arising from this submission.

References

- [ABRK24] S. Armstrong, A. Bou-Rabee, and T. Kuusi. Superdiffusive central limit theorem for a Brownian particle in a critically-correlated incompressible random drift. *arXiv preprint arXiv:2404.01115* (2024).
- [ADC21] M. AIZENMAN and H. DUMINIL-Copin. Marginal triviality of the scaling limits of critical 4d Ising and ϕ 4 ⁴ models. *Ann. Math. (2)* **194**, no. 1, (2021), 163–235. [http://dx.doi.org/](doi:10.4007/annals.2021.194.1.3) [10.4007/annals.2021.194.1.3](doi:10.4007/annals.2021.194.1.3).
- [BC98] L. Bertini and N. Cancrini. The two-dimensional stochastic heat equation: renormalizing a multiplicative noise. *J. Phys. A, Math. Gen.* **31**, no. 2, (1998), 615–622. [http://dx.doi.](doi:10.1088/0305-4470/31/2/019) [org/10.1088/0305-4470/31/2/019](doi:10.1088/0305-4470/31/2/019).
- [BCGL87] BOUCHAUD, J.P., COMTET, A., GEORGES, A., and LE DOUSSAL, P. Anomalous diffusion in random media of any dimensionality. *J. Phys. France* **48**, no. 9, (1987), 1445–1450. [http://](doi:10.1051/jphys:019870048090144500) [dx.doi.org/10.1051/jphys:019870048090144500](doi:10.1051/jphys:019870048090144500).
- $[BOS11]$ M. Balázs, J. Quastel, and T. Sepp λ in Equation exponent of the KPZ/stochastic Burgers equation. *J. Am. Math. Soc.* **24**, no. 3, (2011), 683–708. [http://dx.doi.org/10.](doi:10.1090/S0894-0347-2011-00692-9) [1090/S0894-0347-2011-00692-9](doi:10.1090/S0894-0347-2011-00692-9).
- [BS95] A.-L. BARABÁSI and H. E. STANLEY. *Fractal concepts in surface growth*. Cambridge University Press, Cambridge, 1995. [http://dx.doi.org/10.1017/CBO9780511599798](doi:10.1017/CBO9780511599798).
- [CC₂₂] F. CARAVENNA and F. COTTINI. Gaussian limits for subcritical chaos. *Electron. J. Probab.* **27**, (2022), 35. Id/No 81. [http://dx.doi.org/10.1214/22-EJP798](doi:10.1214/22-EJP798).
- [CCHS22] A. CHANDRA, I. CHEVYREV, M. HAIRER, and H. SHEN. Langevin dynamic for the 2d Yang-Mills measure. *Publ. Math., Inst. Hautes Etud. Sci. ´* **136**, (2022), 1–147. [http://dx.doi.](doi:10.1007/s10240-022-00132-0) [org/10.1007/s10240-022-00132-0](doi:10.1007/s10240-022-00132-0).
- [CCHS24] A. CHANDRA, I. CHEVYREV, M. HAIRER, and H. SHEN. Stochastic quantisation of Yang-Mills-Higgs in 3d. *Invent. Math.* **237**, no. 2, (2024), 541–696. [http://dx.doi.org/10.](doi:10.1007/s00222-024-01264-2) [1007/s00222-024-01264-2](doi:10.1007/s00222-024-01264-2).
- [CCM24] F. Comers, C. Cosco, and C. Mukherjee. Space-time fluctuation of the Kardar-Parisi-Zhang equation in $d \geq 3$ and the Gaussian free field. *Ann. Inst. Henri Poincaré, Probab. Stat.* **60**, no. 1, (2024), 82–112. [http://dx.doi.org/10.1214/22-AIHP1272](doi:10.1214/22-AIHP1272).
- [CD₂₀] S. CHATTERJEE and A. DUNLAP. Constructing a solution of the $(2 + 1)$ -dimensional KPZ equation. *Ann. Probab.* **48**, no. 2, (2020), 1014–1055. [http://dx.doi.org/10.1214/](doi:10.1214/19-AOP1382) [19-AOP1382](doi:10.1214/19-AOP1382).
- [CES21] G. CANNIZZARO, D. ERHARD, and P. SCHÖNBAUER. 2d anisotropic KPZ at stationarity: scaling, tightness and nontriviality. *Ann. Probab.* **49**, no. 1, (2021), 122–156. [http://dx.](doi:10.1214/20-AOP1446) [doi.org/10.1214/20-AOP1446](doi:10.1214/20-AOP1446).
- [CET23a] G. Cannizzaro, D. Erhard, and F. Toninelli. The Stationary AKPZ equation: logarithmic superdiffusivity. *Commun. Pure Appl. Math.* **76**, no. 11, (2023), 3044–3103. [http://dx.](doi:10.1002/cpa.22108) [doi.org/10.1002/cpa.22108](doi:10.1002/cpa.22108).
- [CET23b] G. Cannizzaro, D. Erhard, and F. Toninelli. Weak coupling limit of the anisotropic KPZ equation. *Duke Math. J.* **172**, no. 16, (2023), 3013–3104. [http://dx.doi.org/10.1215/](doi:10.1215/00127094-2022-0094) [00127094-2022-0094](doi:10.1215/00127094-2022-0094).
- [CG24] G. CANNIZZARO and H. GILES. An invariance principle for the 2d weakly self-repelling Brownian polymer, 2024. URL <https://arxiv.org/abs/2403.06730>.
- [CGT24] G. Cannizzaro, M. Gubinelli, and F. Toninelli. Gaussian fluctuations for the stochastic Burgers equation in dimension $d > 2$. *Commun. Math. Phys.* **405**, no. 4, (2024), 60. Id/No 89. [http://dx.doi.org/10.1007/s00220-024-04966-z](doi:10.1007/s00220-024-04966-z).
- [CHST22] G. CANNIZZARO, L. HAUNSCHMID-SIBITZ, and F. TONINELLI. $\sqrt{\log t}$ -superdiffusivity for a Brownian particle in the curl of the 2d GFF. *Ann. Probab.* **50**, no. 6, (2022), 2475–2498. [http://dx.doi.org/10.1214/22-AOP1589](doi:10.1214/22-AOP1589).
- [CLO01] C.-C. CHANG, C. LANDIM, and S. OLLA. Equilibrium fluctuations of asymmetric simple exclusion processes in dimension $d \geq 3$. *Probab. Theory Related Fields* **119**, no. 3, (2001), 381–409. [http://dx.doi.org/10.1007/PL00008764](doi:10.1007/PL00008764).
- [CMOW22] G. CHATZIGEORGIOU, P. MORFE, F. OTTO, and L. WANG. The gaussian free-field as a stream function: asymptotics of effective diffusivity in infra-red cut-off. *arXiv preprint arXiv:2212.14244* (2022).
- [CNN22] C. Cosco, S. Nakajima, and M. Nakashima. Law of large numbers and fluctuations in the sub-critical and L^2 regions for she and kpz equation in dimension $d \geq 3$. *Stochastic Processes Appl.* **151**, (2022), 127–173. [http://dx.doi.org/10.1016/j.spa.2022.05.](doi:10.1016/j.spa.2022.05.010) [010](doi:10.1016/j.spa.2022.05.010).
- [CSZ17] F. CARAVENNA, R. SUN, and N. ZYGOURAS. Universality in marginally relevant disordered systems. *Ann. Appl. Probab.* **27**, no. 5, (2017), 3050–3112. [http://dx.doi.org/10.](doi:10.1214/17-AAP1276) [1214/17-AAP1276](doi:10.1214/17-AAP1276).
- [CSZ20] F. Caravenna, R. Sun, and N. Zygouras. The two-dimensional KPZ equation in the entire subcritical regime. *Ann. Probab.* **48**, no. 3, (2020), 1086–1127. [http://dx.doi.org/10.](doi:10.1214/19-AOP1383) [1214/19-AOP1383](doi:10.1214/19-AOP1383).
- [CSZ23] F. Caravenna, R. Sun, and N. Zygouras. The critical 2d stochastic heat flow. *Invent. Math.* **233**, no. 1, (2023), 325–460. [http://dx.doi.org/10.1007/s00222-023-01184-7](doi:10.1007/s00222-023-01184-7).
- [CT24] G. Cannizzaro and F. Toninelli. Lecture notes on stationary critical and super-critical SPDEs, 2024. URL <https://arxiv.org/abs/2403.15006>.
- [DG22] A. DUNLAP and Y. GU. A forward-backward SDE from the 2d nonlinear stochastic heat equation. *Ann. Probab.* **50**, no. 3, (2022), 1204–1253. [http://dx.doi.org/10.1214/](doi:10.1214/21-AOP1563) [21-AOP1563](doi:10.1214/21-AOP1563).
- [DGRZ20] A. DUNLAP, Y. GU, L. RYZHIK, and O. ZEITOUNI. Fluctuations of the solutions to the KPZ equation in dimensions three and higher. *Probab. Theory Relat. Fields* **176**, no. 3-4, (2020), 1217–1258. [http://dx.doi.org/10.1007/s00440-019-00938-w](doi:10.1007/s00440-019-00938-w).
- [DPZ08] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions.*, vol. 44 of *Encycl. Math. Appl.* Cambridge: Cambridge University Press, reprint of the 1992 hardback ed. ed., 2008. [http://dx.doi.org/10.1017/CBO9780511666223](doi:10.1017/CBO9780511666223).
- [Duc22] P. Duch. Renormalization of singular elliptic stochastic PDEs using flow equation (2022).
- [EK86] S. N. Ethier and T. G. Kurtz. *Markov processes*. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons, Inc., New York, 1986. Characterization and convergence.
- [ET96] D. E. Edmunds and H. Triebel. *Function spaces, entropy numbers, differential operators*, vol. 120 of *Camb. Tracts Math.* Cambridge: Cambridge Univ. Press, 1996.
- [FQ15] T. Funaki and J. Quastel. KPZ equation, its renormalization and invariant measures. *Stoch. Partial Differ. Equ., Anal. Comput.* **3**, no. 2, (2015), 159–220. [http://dx.doi.org/10.](doi:10.1007/s40072-015-0046-x) [1007/s40072-015-0046-x](doi:10.1007/s40072-015-0046-x).
- [GHL23] L. GEROLLA, M. HAIRER, and X.-M. Li. Fluctuations of stochastic PDEs with long-range correlations. Preprint, arXiv:2303.09811 [math.PR] (2023), 2023. URL [https://arxiv.](https://arxiv.org/abs/2303.09811) [org/abs/2303.09811](https://arxiv.org/abs/2303.09811).
- [GHS24] D. D. GASPARI and L. HAUNSCHMID-SIBITZ. $(\log t)^{2/3}$ -superdiffusivity for the 2d stochastic Burgers equation. *Electronic Journal of Probability* **29**, no. none, (2024), 1 – 34. [http://](doi:10.1214/24-EJP1249) [dx.doi.org/10.1214/24-EJP1249](doi:10.1214/24-EJP1249).
- [GIP15] M. GUBINELLI, P. IMKELLER, and N. PERKOWSKI. Paracontrolled distributions and singular PDEs. *Forum Math. Pi* **3**, (2015), e6, 75. [http://dx.doi.org/10.1017/fmp.2015.2](doi:10.1017/fmp.2015.2).
- [GJ14] P. Goncalves and M. Jara. Nonlinear fluctuations of weakly asymmetric interacting particle systems. *Arch. Ration. Mech. Anal.* **212**, no. 2, (2014), 597–644. [http://dx.doi.org/10.](doi:10.1007/s00205-013-0693-x) [1007/s00205-013-0693-x](doi:10.1007/s00205-013-0693-x).
- [GL20] Y. Gu and J. Li. Fluctuations of a nonlinear stochastic heat equation in dimensions three and higher. *SIAM J. Math. Anal.* **52**, no. 6, (2020), 5422–5440. [http://dx.doi.org/10.](doi:10.1137/19M1296380) [1137/19M1296380](doi:10.1137/19M1296380).
- [GP17] M. Gubinelli and N. Perkowski. KPZ reloaded. *Commun. Math. Phys.* **349**, no. 1, (2017), 165–269. [http://dx.doi.org/10.1007/s00220-016-2788-3](doi:10.1007/s00220-016-2788-3).
- [GP18] M. Gubinelli and N. Perkowski. Energy solutions of KPZ are unique. *J. Am. Math. Soc.* **31**, no. 2, (2018), 427–471. [http://dx.doi.org/10.1090/jams/889](doi:10.1090/jams/889).
- [GP20] M. Gubinelli and N. Perkowski. The infinitesimal generator of the stochastic Burgers equation. *Probab. Theory Relat. Fields* **178**, no. 3-4, (2020), 1067–1124. [http://dx.doi.](doi:10.1007/s00440-020-00996-5) [org/10.1007/s00440-020-00996-5](doi:10.1007/s00440-020-00996-5).
- [GPP₂₄] L. GrÄFNER, N. PERKOWSKI, and S. POPAT. Energy solutions of singular SPDEs on Hilbert spaces with applications to domains with boundary conditions. Preprint, arXiv:2411.07680 [math.PR] (2024), 2024. URL <https://arxiv.org/abs/2411.07680>.
- [GRZ18] Y. Gu, L. RYZHIK, and O. ZEITOUNI. The Edwards-Wilkinson limit of the random heat equation in dimensions three and higher. *Commun. Math. Phys.* **363**, no. 2, (2018), 351–388. [http://dx.doi.org/10.1007/s00220-018-3202-0](doi:10.1007/s00220-018-3202-0).
- [GRZ24] S. Gabriel, T. Rosati, and N. Zygouras. The Allen-Cahn equation with weakly critical random initial datum. *Probab. Theory Relat. Fields* (2024). [http://dx.doi.org/10.](doi:10.1007/s00440-024-01312-1) [1007/s00440-024-01312-1](doi:10.1007/s00440-024-01312-1).
- [Gu20] Y. Gu. Gaussian fluctuations from the 2D KPZ equation. *Stoch. Partial Differ. Equ. Anal. Comput.* **8**, no. 1, (2020), 150–185. [http://dx.doi.org/10.1007/](doi:10.1007/s40072-019-00144-8) [s40072-019-00144-8](doi:10.1007/s40072-019-00144-8).
- [Hai13] M. Hairer. Solving the KPZ equation. *Ann. Math. (2)* **178**, no. 2, (2013), 559–664. [http://](doi:10.4007/annals.2013.178.2.4) [dx.doi.org/10.4007/annals.2013.178.2.4](doi:10.4007/annals.2013.178.2.4).
- [Hai14] M. Hairer. A theory of regularity structures. *Invent. Math.* **198**, no. 2, (2014), 269–504. [http://dx.doi.org/10.1007/s00222-014-0505-4](doi:10.1007/s00222-014-0505-4).
- [Hen81] D. Henry. *Geometric theory of semilinear parabolic equations*, vol. 840 of *Lect. Notes Math.* Springer, Cham, 1981.
- [HL₁₅] M. Hairer and C. Labbé. A simple construction of the continuum parabolic Anderson model on R 2 . *Electron. Commun. Probab.* **20**, (2015), 11. Id/No 43. [http://dx.doi.org/](doi:10.1214/ECP.v20-4038) [10.1214/ECP.v20-4038](doi:10.1214/ECP.v20-4038).
- [HTVo12] I. Horvárn, B. Tórn, and B. VET" o. Relaxed sector condition. *Bull. Inst. Math. Acad. Sin. (N.S.)* **7**, no. 4, (2012), 463-476.
- [IK02] K. Ito and F. Kappel. *Evolution equations and approximations*, vol. 61 of *Ser. Adv. Math. Appl. Sci.* Singapore: World Scientific, 2002.
- [KLO12] T. Komorowski,C. Landim, and S. Olla. *Fluctuations in Markov processes. Time symmetry and martingale approximation.*, vol. 345 of *Grundlehren Math. Wiss.* Berlin: Springer, 2012. [http://dx.doi.org/10.1007/978-3-642-29880-6](doi:10.1007/978-3-642-29880-6).
- [Kup16] A. Kupiainen. Renormalization group and stochastic PDEs. *Ann. Henri Poincare´* **17**, no. 3, (2016), 497–535. [http://dx.doi.org/10.1007/s00023-015-0408-y](doi:10.1007/s00023-015-0408-y).
- [LOY04] C. LANDIM, S. OLLA, and H. YAU. Diffusive behaviour of the equilibrium fluctuations in the asymmetric exclusion processes. *Adv. Stud. Pure Math.* **39**, (2004), 307–324. [http://dx.](doi:10.2969/aspm/03910307) [doi.org/10.2969/aspm/03910307](doi:10.2969/aspm/03910307).
- [LY97] C. LANDIM and H. T. YAU. Fluctuation-dissipation equation of asymmetric simple exclusion processes. *Probab. Theory Relat. Fields* **108**, no. 3, (1997), 321–356. [http://dx.doi.](doi:10.1007/s004400050112) [org/10.1007/s004400050112](doi:10.1007/s004400050112).
- [LZ22] D. LYGKONIS and N. ZYGOURAS. Edwards–Wilkinson fluctuations for the directed polymer in the full L^2 -regime for dimensions $d \geq 3$. *Ann. Inst. H. Poincaré (B): Prob. Stat.* **58(1)**, $(2022), 65 - 104.$
- [MOW24] P. Morfe, F. Otto, and C. WAGNER. A Critical Drift-Diffusion Equation: Connections to the Diffusion on **SL**(2). Preprint, arXiv:2410.15983 [math.PR] (2024), 2024. URL [https://](https://arxiv.org/abs/2410.15983) arxiv.org/abs/2410.15983.
- [MP19] J. Martin and N. Perkowski. Paracontrolled distributions on Bravais lattices and weak universality of the 2d parabolic Anderson model. *Ann. Inst. Henri Poincare, Probab. Stat. ´* **55**, no. 4, (2019), 2058–2110. [http://dx.doi.org/10.1214/18-AIHP942](doi:10.1214/18-AIHP942).
- [MQR21] K. MATETSKI, J. QUASTEL, and D. REMENIK. The KPZ fixed point. *Acta Math.* 227, no. 1, (2021), 115–203. [http://dx.doi.org/10.4310/ACTA.2021.v227.n1.a3](doi:10.4310/ACTA.2021.v227.n1.a3).
- [Nua06] D. Nualart. *The Malliavin calculus and related topics*. Probability and its Applications (New York). Springer-Verlag, Berlin, second ed., 2006.
- [OW24] F. Orro and C. WAGNER. A critical drift-diffusion equation: intermittent behavior. Preprint, arXiv:2404.13641 [math.PR] (2024), 2024. URL [https://arxiv.org/abs/](https://arxiv.org/abs/2404.13641) [2404.13641](https://arxiv.org/abs/2404.13641).
- [QS15] J. QUASTEL and H. SPOHN. The one-dimensional KPZ equation and its universality class. *J. Stat. Phys.* **160**, no. 4, (2015), 965–984. [http://dx.doi.org/10.1007/](doi:10.1007/s10955-015-1250-9) [s10955-015-1250-9](doi:10.1007/s10955-015-1250-9).
- [QS23] J. QUASTEL and S. SARKAR. Convergence of exclusion processes and the KPZ equation to the KPZ fixed point. *J. Am. Math. Soc.* **36**, no. 1, (2023), 251–289. [http://dx.doi.org/](doi:10.1090/jams/999) [10.1090/jams/999](doi:10.1090/jams/999).
- [Tao24] R. Tao. Gaussian fluctuations of a nonlinear stochastic heat equation in dimension two. *Stoch. Partial Differ. Equ., Anal. Comput.* **12**, no. 1, (2024), 220–246. [http://dx.doi.](doi:10.1007/s40072-022-00282-6) [org/10.1007/s40072-022-00282-6](doi:10.1007/s40072-022-00282-6).
- [Tsa24] L.-C. Tsai. Stochastic heat flow by moments. Preprint, arXiv:2410.14657 [math.PR] (2024), 2024. URL <https://arxiv.org/abs/2410.14657>.
- [TV₁₂] B. Tórn and B. VALKÓ. Superdiffusive bounds on self-repellent Brownian polymers and diffusion in the curl of the Gaussian free field in $d = 2$. *J. Stat. Phys.* **147**, no. 1, (2012), 113–131. [http://dx.doi.org/10.1007/s10955-012-0462-5](doi:10.1007/s10955-012-0462-5).
- [vBKS85] H. van Beijeren, R. Kutner, and H. Spohn. Excess noise for driven diffusive systems. *Phys. Rev. Lett.* **54**, (1985), 2026–2029. [http://dx.doi.org/10.1103/PhysRevLett.](doi:10.1103/PhysRevLett.54.2026) [54.2026](doi:10.1103/PhysRevLett.54.2026).
- [Vir20] B. Virag. The heat and the landscape I. *arXiv preprint* (2020). [http://arxiv.org/abs/](arXiv:2008.07241) [2008.07241](arXiv:2008.07241).
- [Wal86] J. B. WALSH. An introduction to stochastic partial differential equations. École d'été de probabilités de Saint-Flour XIV - 1984, Lect. Notes Math. 1180, 265-437 (1986)., 1986.
- [Wol91] E. Wolf, DIETRICH. Kinetic roughening of vicinal surfaces. *Phys. Rev. Lett.* **67**, (1991), 1783–1786.
- [Yauo4] H.-T. YAU. $(\log t)^{2/3}$ law of the two dimensional asymmetric simple exclusion process. Ann. *of Math. (2)* **159**, no. 1, (2004), 377–405. [http://dx.doi.org/10.4007/annals.2004.](doi:10.4007/annals.2004.159.377) [159.377](doi:10.4007/annals.2004.159.377).
- [YY24] H. Yang and Z. Yang. Weak coupling limit of a Brownian particle in the curl of the 2D GFF. *arXiv preprint arXiv:2405.05778* (2024).