# TORSION OF ELLIPTIC CURVES WITH RATIONAL *j*-INVARIANT OVER QUARTIC NUMBER FIELDS

LUCAS HAMADA

ABSTRACT. Let E be an elliptic curve, defined over a quartic extension K of  $\mathbb{Q}$ , with  $j(E) \in \mathbb{Q}$ . In this paper, we classify the possible torsion subgroup structures  $E(K)_{\text{tors}}$ .

## INTRODUCTION

Let E be an elliptic curve defined over a number field K. According to the Mordell-Weil theorem, the Mordell-Weil group E(K) of K-rational points of E is a finitely generated abelian group, which can be expressed as:

$$E(K) \cong E(K)_{\text{tors}} \oplus \mathbb{Z}^{\oplus r},$$

where  $E(K)_{\text{tors}}$  denotes the torsion subgroup. The first major breakthrough regarding the torsion subgroup of an elliptic curve was achieved by B. Mazur, who classified, up to isomorphism, all possible Q-torsion subgroups of elliptic curves defined over Q. This result is presented in the following theorem:

THEOREM 0.1 (B. Mazur [14]). Let E be an elliptic curve defined over  $\mathbb{Q}$ . Then,  $E(\mathbb{Q})_{\text{tors}}$  is isomorphic to one of the following 15 groups:

$$\mathbb{Z}/N_1\mathbb{Z},$$
  
 $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2N_2\mathbb{Z},$ 
 $N_1 = 1, \dots, 10, 12$   
 $N_2 = 1, \dots, 4.$ 

Several analogues of Theorem 0.1 were proven for different classes of elliptic curves: Let  $\Phi(d)$  denote the set of all isomorphism classes of torsion subgroup  $E(K)_{\text{tors}}$ , where K runs through all extensions of  $\mathbb{Q}$  of degree d and E runs through all elliptic curves defined over K. The classification of  $\Phi(d)$  have been obtained for various values of d:

 $-\Phi(2)$ : S. Kamienny in [12], and M. Kenku and F. Momose in [13].

- Φ(3): D. Jeon, C. H. Kim, and A. Schweizer in [11], and M. Derickx, A. Etropolski,
 J. Morrow, M. van Hoeij, and D. Zureick-Brown in [4].

 $-\Phi(4)$ : M. Derickx and F. Najman in [3].

Moreover, let  $\Phi_{\mathbb{Q}}(d)$  denote the set of all isomorphism classes of torsion subgroups  $E(K)_{\text{tors}}$ , where E runs through all elliptic curves defined over  $\mathbb{Q}$  and K runs through all extensions of  $\mathbb{Q}$  of degree d. The following classifications are known:

<sup>2020</sup> Mathematics Subject Classification. Primary 11G05; Secondary 12F05.

Key words and phrases. Elliptic curves, Torsion points, number fields.

- $-\Phi_{\mathbb{Q}}(2)$  and  $\Phi_{\mathbb{Q}}(3)$ : F. Najman in [15],
- $-\Phi_{\mathbb{Q}}(4)$ :M. Chou in [1], and E. González-Jiménez and F. Najman in [7]
- $-\Phi_{\mathbb{Q}}(p)$  for primes  $p \geq 5$ : González-Jiménez and F. Najman in [6] and [7].

In this paper, we study the torsion subgroup structure of elliptic curves with *j*-invariant belonging to  $\mathbb{Q}$ . Let  $\Phi_{j\in\mathbb{Q}}(d)$  denote the set of all isomorphism classes of torsion subgroups  $E(K)_{\text{tors}}$ , where K runs through all extensions of  $\mathbb{Q}$  of degree d and E runs through all elliptic curves E defined over K satisfying  $j(E) \in \mathbb{Q}$ . The first significant work on this problem was conducted by T. Gužvić in [8], where he determined  $\Phi_{j\in\mathbb{Q}}(p)$  for all prime numbers p. Following this, J. E. Cremona and F. Najman in [16] extended the results to determine  $\Phi_{j\in\mathbb{Q}}(d)$  for all d not divisible by primes less than 11.

The goal of this paper is to determine the set  $\Phi_{j\in\mathbb{Q}}(4)$ . Specifically, we provide a complete description as follows:

THEOREM 0.2. Let E be an elliptic curve, defined over a quartic extension K of  $\mathbb{Q}$ , with  $j(E) \in \mathbb{Q}$ . Then, the torsion subgroup  $E(K)_{\text{tors}}$  is isomorphic to one of the following groups, and each group occurs as a torsion subgroup for some elliptic curve E:

 $\mathbb{Z} / N_1 \mathbb{Z}, \qquad N_1 = 1, \dots, 10, 12, 13, 15, 16, 17, 20, 21, 24, \\ \mathbb{Z} / 2 \mathbb{Z} \oplus \mathbb{Z} / 2 N_2 \mathbb{Z}, \qquad N_2 = 1, \dots, 6, 8, \\ \mathbb{Z} / 3 \mathbb{Z} \oplus \mathbb{Z} / 3 N_3 \mathbb{Z}, \qquad N_3 = 1, 2, \\ \mathbb{Z} / 4 \mathbb{Z} \oplus \mathbb{Z} / 4 N_4 \mathbb{Z}, \qquad N_4 = 1, 2, \\ \mathbb{Z} / 5 \mathbb{Z} \oplus \mathbb{Z} / 5 \mathbb{Z}, \\ \mathbb{Z} / 6 \mathbb{Z} \oplus \mathbb{Z} / 6 \mathbb{Z}.$ 

**Outline of the paper.** The paper is organized as follows: In Section 1, we review fundamental properties of torsion points on elliptic curves defined over  $\mathbb{Q}$ . Section 2 establishes two necessary lemmas regarding number fields, which will be instrumental for the proof of Theorem 0.2. In Section 3, we discuss a prior result concerning the torsion points of elliptic curves with complex multiplication (CM), which allows us to focus exclusively on elliptic curves without CM in subsequent sections. Section 4 presents key results concerning elliptic curves E with  $j(E) \in \mathbb{Q} \setminus \{0, 1728\}$ . Finally, in Section 5, we prove Theorem 0.2 and provide examples of elliptic curves corresponding to each torsion subgroup structure listed in the theorem.

Acknowledgments. The author deeply appreciates Yuichiro Taguchi and Yuto Nagashima for their insightful discussions and constructive comments on the early version of this paper.

#### 1. Generalities on torsion points of rational elliptic curves

In this section, we recall some properties of torsion points of elliptic curves defined over  $\mathbb{Q}$ . The main result is the list determined by E. González-Jimenéz and F. Najman about the possible extension degrees of the field of definition of torsion points of prime orders.

THEOREM 1.1 (E. González-Jimenéz and F. Najman [7], Proposition 5.7). Let E be an elliptic curve defined over  $\mathbb{Q}$ , p a prime number and  $P \in E(\overline{\mathbb{Q}})$  a point of order p. Then, for  $p \leq 13$  or p = 37 all of the cases in the table below occur, and they are the only possible ones. For p = 17, the cases in the table below are the only possible ones (it is still unknown if the cases  $[\mathbb{Q}(P) : \mathbb{Q}] = 96$  or 192 might occur).

p	$[\mathbf{Q}(P):\mathbf{Q}]$
2	1,2,3
3	1, 2, 3, 4, 6, 8
5	1, 2, 4, 5, 8, 10, 16, 20, 24
7	1, 2, 3, 6, 7, 9, 12, 14, 18, 21, 24, 36, 42, 48
11	5, 10, 20, 40, 55, 80, 100, 110, 120
13	3, 4, 6, 12, 24, 39, 48, 52, 72, 78, 96, 144, 156, 168
17	$8, 16, 32, \boldsymbol{96}, 136, \boldsymbol{192}, 256, 272, 288$
37	12, 36, 72, 444, 1296, 1332, 1368

When the elliptic curve admits a cyclic isogeny (see Silverman [18]), we get stronger results as the following lemma:

LEMMA 1.2 (L. Hamada [9], Lemmas 2.6 and 2.7). Let E be an elliptic curve defined over  $\mathbb{Q}$ , and K a finite Galois extension of  $\mathbb{Q}$ .

- (1) If  $P_n \in E(K)$  has order n and  $\langle P_n \rangle$  is  $\operatorname{Gal}(K/\mathbb{Q})$ -invariant, then  $[\mathbb{Q}(P_n) : \mathbb{Q}]$  divides  $\varphi(n)$ , where  $\varphi$  is the Euler function.
- (2) If  $E(K)[p^n] = \langle P_{p^n}, Q_p \rangle \cong (\mathbb{Z}/p^n \mathbb{Z}) \oplus (\mathbb{Z}/p \mathbb{Z})$ , where p is a prime number,  $P_{p^n} \in E(K)[p^n] - E(\mathbb{Q})$  is a point of order  $p^n$  that is not defined in  $\mathbb{Q}$ , and  $Q_p \in E(\mathbb{Q})[p]$  is a point of order p. If the degree  $[K : \mathbb{Q}]$  is coprime to p, then  $[\mathbb{Q}(P_{p^n}) : \mathbb{Q}]$  divides p - 1.

In [1], M. Chou provided a useful criterion for determining when an elliptic curve admits a cyclic isogeny.

LEMMA 1.3 (M. Chou. Adapted from the proof of Proposition 4.1 in [1]). Let K be a Galois extension of  $\mathbb{Q}$ , and let E be an elliptic curve defined over  $\mathbb{Q}$  such that  $E(K)[nm] = (\mathbb{Z}/nm\mathbb{Z}) \oplus (\mathbb{Z}/m\mathbb{Z})$  for some positive integers n and m. Then, E has a cyclic n-isogeny.

PROOF. This follows from the fact that [m](E(K)[nm]) is a  $\operatorname{Gal}(K/\mathbb{Q})$ -invariant cyclic subgroup of E(K) of order n.

#### 2. Lemmas on number fields

This section establishes two lemmas on number fields, to be used in the proof of Theorem 0.2. First, we prove a condition under which a quadratic extension of  $\mathbb{Q}$  is contained within a cyclic quartic extension of  $\mathbb{Q}$ . Specifically, we prove the following:

LEMMA 2.1. Let K be a cyclic quartic extension of  $\mathbb{Q}$  and F be its unique non-trivial subfield. Then,  $F = \mathbb{Q}(\sqrt{n})$  for some positive integer  $n = 2^a p_1 p_2 \cdots p_k$  where  $a \in \{0, 1\}$ and  $p_i$  are primes that satisfy  $p_i \equiv 1 \pmod{4}$ . In particular, F is not equal to neither  $\mathbb{Q}(\sqrt{-1}), \mathbb{Q}(\sqrt{-3})$  nor  $\mathbb{Q}(\sqrt{3})$ .

To prove this, we need the following theorem by A.-M. Legendre.

THEOREM 2.2 (A.-M. Legendre. See J.-P. Serre [17]). Let a, b, c be three integers, not all of the same sign and such that abc is square-free. Then, the equation

$$aX^2 + bY^2 + cZ^2 = 0$$

has a solution in integers  $(X, Y, Z) \neq (0, 0, 0)$  if and only if -bc, -ca, -ab are quadratic residues modulo |a|, modulo |b| and modulo |c|, respectively.

Proof of Lemma 2.1. By [1, Lemma 3.1], a quadratic extension F of  $\mathbb{Q}$  is an intermediate field of a quartic cyclic extension of  $\mathbb{Q}$  if and only if  $F = \mathbb{Q}(\sqrt{n})$  for some positive square-free integer n such that there exist non-zero integers a, b and c satisfying

$$a^2 - nb^2 = nc^2.$$

Since n is square-free, this is equivalent to the existence of a non-zero solution of the diophantine equation

$$nX^2 - Y^2 - Z^2 = 0.$$

By Theorem 2.2, this is equivalent to -1 be a quadratic residuo modulo n which is equivalent to -1 be a quadratic residue for each prime divisor of n. Therefore  $n = 2^a p_1 p_2 \cdots p_k$  for  $a \in \{0, 1\}$  and  $p_i \equiv 1 \pmod{4}$ .

The second lemma is a technical result that addresses the non-existence of cyclic extensions of degree 3, under the conditions considered in the proof of Theorem 0.2.

LEMMA 2.3. Let  $K/\mathbb{Q}$  be a Galois extension with Galois group  $\operatorname{Gal}(K/\mathbb{Q})$  isomorphic to  $S_4$ , or  $D_4$ . Let  $L = K(\sqrt{\alpha})$  be a quadratic extension of K with  $\alpha \in K$ , and let  $\widehat{L}$  be its Galois closure over  $\mathbb{Q}$ . Then, there does not exist a cyclic extension F of  $\mathbb{Q}$  of degree 3 that is contained in  $\widehat{L}$ .

PROOF. First, we claim that  $[\widehat{L} : K]$  is a power of 2. Indeed, let  $\sigma$  be an element of  $\operatorname{Gal}(\widehat{L}/\mathbb{Q})$ . The Galois conjugate of L by  $\sigma$  is  $K(\sqrt{\sigma(\alpha)})$ , also a quadratic extension of

K. Therefore, since we know that  $\hat{L}$  is the compositum of all Galois conjugates of L and each of them are Galois extensions over K, we have the equality

$$[\widehat{L}:K] = \prod_{\sigma'} [L^{\sigma'}:K],$$

where the product is taken over a maximal set of linearly disjoint Galois conjugates of L. This is a product of powers of 2, where we get the claim.

Finally, suppose that  $F/\mathbb{Q}$  is a cyclic extension of degree 3 contained in L. Looking at the subgroup lattice of  $S_4$  and  $D_4$  we see that  $K \cap F = \mathbb{Q}$ . Since both K and F are Galois extension of  $\mathbb{Q}$ , there are linearly disjoint and we conclude that their compositum KF satisfies [KF:K] = 3, contradicting the first claim.  $\Box$ 

# 3. TORSION OF ELLIPTIC CURVES WITH COMPLEX MULTIPLICATION

In this section, we briefly review torsion subgroups of elliptic curves with complex multiplication. P. L. Clark, P. Corn, A. Rice, and J. Stankewicz in [2] classified, up to isomorphism, the torsion subgroups of elliptic curves E with complex multiplication defined over number fields of various degrees. In particular, they determined the possible torsion subgroups for E defined over quartic extensions of  $\mathbb{Q}$ :

THEOREM 3.1 (P. L. Clark, P. Corn, A. Rice, and J. Stankewicz [2]). Let E be an elliptic curve defined over a quartic extension K of  $\mathbb{Q}$ , with complex multiplication. Then, the torsion subgroup  $E(K)_{\text{tors}}$  is isomorphic to one of the following groups:

$\mathbb{Z}/N_1\mathbb{Z},$	$N_1 = 1, \dots, 8, 10, 12, 13, 21,$
$\mathbb{Z}/2\mathbb{Z}\oplus\mathbb{Z}/2N_2\mathbb{Z},$	$N_2 = 1, \dots, 5,$
$\mathbb{Z}/3\mathbb{Z}\oplus\mathbb{Z}/3N_3\mathbb{Z},$	$N_3 = 1, 2,$
$\mathbb{Z}/4\mathbb{Z}\oplus\mathbb{Z}/4\mathbb{Z}$ .	

As a result, throughout the remainder of this paper, we may assume that all elliptic curves under consideration do not have complex multiplication. In particular, this implies that their j-invariants are neither 0 nor 1728, allowing us to apply the theory presented in the next section.

# 4. On elliptic curves with rational j-invariant

In this section, we begin by briefly reviewing the theory of quadratic twists of elliptic curves. We then focus on the specific case where the elliptic curve has *j*-invariant in  $\mathbb{Q} \setminus \{0, 1728\}$ . Let *E* be an elliptic curve defined over a field *K*, of characteristic zero, of the form

$$E: y^2 = x^3 + Ax + B \quad (A, B \in K).$$

Let D be a non-zero element of K and let  $E^D$  be the elliptic curve of the form:

$$E^D: y^2 = x^3 + AD^2x + BD^3.$$

It is known (see J. H. Silverman [18]) that there exists an isomorphism  $\phi$ , defined over the field  $K(\sqrt{D})$ , from E to  $E^D$  as follows:

(1) 
$$\phi \colon E \xrightarrow{\sim} E^{D}$$
$$(x, y) \longmapsto \left(xD, yD\sqrt{D}\right).$$

This  $E^D$  is called the quadratic twist of E by D. From the format of the isomorphism, we get the following lemma:

LEMMA 4.1 (L. Hamada [9], Lemma 3.3). If E(K) has a point  $P_n$  of order n, then its image  $\phi(P_n)$  in  $E^D(K(\sqrt{D}))$  belongs to  $E^D(K)$  if and only if  $\sqrt{D} \in K$  or n = 2.

Now, suppose the *j*-invariant j(E) is a rational number not equal to neither 0 nor 1728, and define E' to be the elliptic curve

(2) 
$$E': y^2 = x^3 - \frac{27j(E)}{j(E) - 1728}x + \frac{54j(E)}{j(E) - 1728}x$$

Then E' is defined over  $\mathbb{Q}$  and has the same *j*-invariant as E, that is, j(E') = j(E). For simplicity, denote the coefficients of E' by

$$A' := -\frac{27j(E)}{j(E) - 1728},$$
$$B' := \frac{54j(E)}{j(E) - 1728}.$$

Since  $j(E) = j(E') \notin \{0, 1728\}$ , by the definition of the *j*-invariant, all the coefficients A, B, A' and B' are non-zero. It is known that E' is a quadratic twist of E by  $D := BA'/AB' \in K$  and, in particular, there exists an isomorphism  $\phi$ , defined over the field  $L := K(\sqrt{D})$ , as in (1)

Now, observe that

$$A = A'/D^2,$$
  
$$B = B'/D^3,$$

so E is actually defined over the field  $\mathbb{Q}(D^2, D^3) = \mathbb{Q}(D)$ . During the proof of Theorem 0.2, we will get L to be equal to  $K(\sqrt{m})$  for some  $m \in \mathbb{Q}$ . We prove that in this case, E is actually K-isomorphic to an elliptic curve  $\widetilde{E}$  defined over  $\mathbb{Q}$ . More generally, we have:

THEOREM 4.2. Let E be an elliptic curve defined over a field K of characteristic zero, with  $j(E) \in \mathbb{Q} \setminus \{0, 1728\}$ , and let  $E' = E^D$  be its quadratic twist by D as defined by the formula (2). If  $L := K(\sqrt{D}) = K(\sqrt{m})$  for some  $m \in K$ , then there exists an elliptic curve  $\tilde{E}$  defined over  $\mathbb{Q}(m)$  that is K-isomorphic to E. PROOF. Since  $K(\sqrt{D}) = K(\sqrt{m})$ , by Kummer theory, there exists an element  $b \in K^{\times}$  such that

$$D = b^2 m$$

This implies that

$$Ab^4 = \frac{Ab^4m^2}{m^2} = \frac{AD^2}{m^2} = \frac{A'}{m^2} \in \mathbb{Q}(m)$$

In the same way, we can prove that  $Bb^6$  also belongs to  $\mathbb{Q}(m)$ . Now, consider the quadratic twist  $\widetilde{E}$  of E by  $b^2$  which, by the above, is defined over  $\mathbb{Q}(m)$ :

$$\widetilde{E}: y^2 = x^3 + Ab^4x + Bb^6.$$

By the argument at the beginning of this section, we know that there exists an isomorphism

$$\begin{split} \psi \colon E & \stackrel{\sim}{\longrightarrow} \widetilde{E} \\ (x,y) & \longmapsto \left( xb^2, yb^3 \right), \end{split}$$

and that this isomorphism  $\psi$  is defined over  $K(\sqrt{b^2}) = K$ . Therefore, E is K-isomorphic to the elliptic curve  $\widetilde{E}$ , which is defined over  $\mathbb{Q}(m)$ .

NOTATION 4.3. From now, if E is an elliptic curve defined over a field K of characteristic zero, satisfying  $j(E) \in \mathbb{Q} \setminus \{0, 1728\}$ , then E' is the elliptic curve defined by the formula in (2), D is the element of K such that  $E' = E^D$ ,  $L = L_{K,D} := K(\sqrt{D})$  is the field of definition of the isomorphism  $\phi$ , as defined in (1), and  $\hat{L}$  is the Galois closure of L over  $\mathbb{Q}$ , in case  $L/\mathbb{Q}$  is an algebraic extension.

The following theorem is the main result that relates the torsion points of E(K) with the torsion points of E'(K):

THEOREM 4.4 (E. González-Jimenéz [5], Theorem 3). Let E be an elliptic curve over a field K of characteristic zero, D a non-square element of K,  $L = K(\sqrt{D})$  and  $E^D$  a quadratic twist of E by D. There exist a pair of homomorphisms

$$E(L) \xrightarrow{\Psi} E(K) \times E^D(L) \xrightarrow{\overline{\Psi}} E(L)$$

such that  $\overline{\Psi} \circ \Psi = [2]$  and  $\Psi \circ \overline{\Psi} = [2] \times [2]$ . Moreover, for n odd, there exists an isomorphism

$$E(L)[n] \cong E(K)[n] \times E^D(K)[n].$$

During the proof of Theorem 0.2, the extension  $L/\mathbb{Q}$  may not be Galois. In such cases, to derive a contradiction, we employ the following two lemmas:

LEMMA 4.5 (L. Hamada [9]. Lemma 2.4). Let E be an elliptic curve defined over  $\mathbb{Q}$ , L an extession of  $\mathbb{Q}$ ,  $\widehat{L}$  its Galois closure over  $\mathbb{Q}$ , and  $\sigma \in \operatorname{Gal}(\widehat{L}/\mathbb{Q})$  an automorphism of  $\widehat{L}$ . Then, we have a group isomorphism:

$$E(L) \xrightarrow{\sim} E(L^{\sigma})$$
$$(x, y) \longmapsto (\sigma(x), \sigma(y))$$

NOTATION 4.6. Let G and H be groups. For the sake of simplicity, we write  $G \subseteq E(K)$ and  $E(K) \subseteq H$  when E(K) has a subgroup isomorphic to G and E(K) is isomorphic to a subgroup of H, respectively.

LEMMA 4.7 (L. Hamada [9], Lemmas 3.7 to 3.9). Let E be an elliptic curve defined over a Galois extension K of  $\mathbb{Q}$  satisfying  $j(E) \in \mathbb{Q} \setminus \{0, 1728\}$ .

(1) If  $\mathbb{Z}/n\mathbb{Z} \subseteq E(K)$  and  $(\mathbb{Z}/n\mathbb{Z}) \oplus (\mathbb{Z}/n\mathbb{Z}) \subseteq E'(L)$  for some integer  $n \geq 3$ , then  $L/\mathbb{Q}$  is a Galois extension.

Now, suppose  $L/\mathbb{Q}$  is not a Galois extension and let  $\widehat{L}$  be its Galois closure over  $\mathbb{Q}$ :

- (2) If  $\mathbb{Z}/p^n \mathbb{Z} \subseteq E(K)$  and  $\mathbb{Z}/p \mathbb{Z} \not\subseteq E'(K)$  for some odd prime p and some positive integer n, then  $[\widehat{L}:K] = 4$ , and  $(\mathbb{Z}/p^n \mathbb{Z}) \oplus (\mathbb{Z}/p^n \mathbb{Z}) \subseteq E'(\widehat{L})$ .
- (3) If  $\mathbb{Z}/2^n \mathbb{Z} \subseteq E(K)$  for some integer  $n \ge 2$ ,  $E'(K)[2^{\infty}] \subseteq (\mathbb{Z}/2\mathbb{Z}) \oplus (\mathbb{Z}/2\mathbb{Z})$ , then  $[\widehat{L}:K] = 4$ , and  $(\mathbb{Z}/2^{n-1}\mathbb{Z}) \oplus (\mathbb{Z}/2^{n-1}\mathbb{Z}) \subseteq E'(\widehat{L})$ .

# 5. Proof of the main theorem

Finally, in this section, we prove Theorem 0.2. Here, we use the notation fixed in the last section, that is, K is a quartic extension of  $\mathbb{Q}$  and E is an elliptic curve, **without** complex multiplication, defined over K with  $j(E) \in \mathbb{Q} \setminus \{0, 1728\}$ . E' is the elliptic curve over  $\mathbb{Q}$  as defined in 2, and L the quadratic extension of K such that there exists an isomorphism  $\phi$ , defined over L, between E and E'.

Fortunately, since M. Derickx and F. Najman classified  $\Phi(4)$  in [3], we already have a finite list of possible torsion subgroup structure for  $E(K)_{\text{tors}}$ :

THEOREM 5.1 (M. Derickx and F. Najman [3]). Let E be an elliptic curve, defined over a quartic extension K of  $\mathbb{Q}$ . Then, the torsion subgroup  $E(K)_{\text{tors}}$  is isomorphic to one of the following groups:

$\mathbb{Z}/N_1\mathbb{Z},$	$N_1 = 1, \dots, 18, 20, 21, 22, 24,$
$\mathbb{Z}/2\mathbb{Z}\oplus\mathbb{Z}/2N_2\mathbb{Z},$	$N_2 = 1, \ldots, 9,$
$\mathbb{Z}/3\mathbb{Z}\oplus\mathbb{Z}/3N_3\mathbb{Z},$	$N_3 = 1, 2, 3$
$\mathbb{Z}/4\mathbb{Z}\oplus\mathbb{Z}/4N_4\mathbb{Z},$	$N_4 = 1, 2,$
$\mathbb{Z} / 5 \mathbb{Z} \oplus \mathbb{Z} / 5 \mathbb{Z},$	
$\mathbb{Z} \ / 6 \mathbb{Z} \oplus \mathbb{Z} \ / 6 \mathbb{Z}$ .	

Now, comparing the list above with the list in the statement of Theorem 0.2, we see that we just need to prove that the following groups do not occur as torsion subgroups of elliptic curves with  $j(E) \in \mathbb{Q} \setminus \{0, 1728\}$ :

(3) 
$$\mathbb{Z} / N_1 \mathbb{Z}, \qquad N_1 = 11, 14, 18, 22,$$
$$\mathbb{Z} / 2 \mathbb{Z} \oplus \mathbb{Z} / 2 N_2 \mathbb{Z}, \qquad N_2 = 7, 9,$$
$$\mathbb{Z} / 3 \mathbb{Z} \oplus \mathbb{Z} / 9 \mathbb{Z}.$$

This will require the following three lemmas:

LEMMA 5.2.  $E(K)_{\text{tors}}$  has no points of order 11.

PROOF. Suppose  $P_{11} \in E(K)_{\text{tors}}$  is a point of order 11, and let  $\phi(P_{11})$  be its image in  $E'(L)_{\text{tors}}$ . By Theorem 1.1, we know that  $[\mathbb{Q}(\phi(P_{11})) : \mathbb{Q}]$  does not divide 8, hence it is impossible that  $\mathbb{Q}(P_{11}) \subseteq L$ , a contradiction.

LEMMA 5.3. If  $E(K)_{\text{tors}}$  has a point of order 7, then L = K or  $L = K(\sqrt{m})$  for some  $m \in \mathbb{Q}$ .

PROOF. Let  $P_7 \in E(K)_{\text{tors}}$  be a point of order 7, and let  $\phi(P_7)$  be its image in  $E'(L)_{\text{tors}}$ . Then, by Theorem 1.1 and the fact that  $[L : \mathbb{Q}]$  divides 8, we see that  $[\mathbb{Q}(\phi(P)) : \mathbb{Q}] = 1$ or 2. Since, by Lemma 4.1,  $L = K(\phi(P))$ , we conclude that L = K or  $L = K(\sqrt{m})$  for some  $m \in \mathbb{Q}$ .

LEMMA 5.4. If  $E(K)_{\text{tors}}$  has a point of order 9, then L = K or  $L = K(\sqrt{m})$  for some  $m \in \mathbb{Q}$ .

To prove this Lemma, we need the following consequence of the Weil pairing.

LEMMA 5.5 (L. Hamada [9], Lemma 5.3). Let E be an elliptic curve defined over a field K of characteristic zero. If  $E(K)_{\text{tors}}$  contains E[n], then K contains  $\mathbb{Q}(\zeta_n)$ . In particular, if the extension  $K/\mathbb{Q}$  is finite, then  $\varphi(n)$  divides  $[K:\mathbb{Q}]$ .

Proof of Lemma 5.4. Suppose  $E(K)_{\text{tors}}$  has a point of order 9. By Lemma 5.5 and the fact that  $\varphi(9) = 6$  does not divide 8, we can assume  $E'(L)[9] \subseteq (\mathbb{Z}/3\mathbb{Z}) \oplus (\mathbb{Z}/9\mathbb{Z})$ . Consequently, E'(K)[3] is isomorphic to either  $\{\mathcal{O}\}$  or  $\mathbb{Z}/3\mathbb{Z}$ . We now consider two cases, depending on whether  $K/\mathbb{Q}$  is a Galois extension or not.

Case 1:  $K/\mathbb{Q}$  is a Galois extension.

- (1.1) Suppose E'(K)[3] = 0. Then, by Theorem 4.4, we have  $E'(L)[9] \subseteq (\mathbb{Z}/3\mathbb{Z}) \oplus (\mathbb{Z}/9\mathbb{Z})$ .
  - If  $L/\mathbb{Q}$  is a Galois extension, then by Lemma 1.3, the elliptic curve E' admits a 3-isogeny. Let  $P_3 \in E'(L)$  be a point of order 3. By Lemma 1.2, the degree  $[\mathbb{Q}(P_3) : \mathbb{Q}]$  divides 2. Therefore, L = K or  $L = K(\sqrt{n})$  for some rational number n.

- If  $L/\mathbb{Q}$  is not a Galois extension, then by Lemma 4.7, we have  $\mathbb{Z}/9\mathbb{Z} \oplus \mathbb{Z}/9\mathbb{Z} \subseteq E'(\widehat{L})$ , where  $[\widehat{L}:\mathbb{Q}] = 16$ . However, since  $[\widehat{L}:\mathbb{Q}]$  must also be divisible by  $\varphi(9) = 6$ , this leads to a contradiction. Hence,  $L/\mathbb{Q}$  cannot be a non-Galois extension in this case.
- (1.2) Suppose  $E'(K)[3] = \langle Q_3 \rangle \cong \mathbb{Z}/3\mathbb{Z}$ . By Lemma 4.7,  $L/\mathbb{Q}$  is a Galois extension. Additionally, by Lemma 1.2, the degree  $[\mathbb{Q}(Q_3) : \mathbb{Q}]$  divides 2. If  $\mathbb{Q}(Q_3) = \mathbb{Q}(\sqrt{n}) \subseteq K$ , then we take the quadratic twist of E' by n. This produces an elliptic curve  $E'^n$  defined over  $\mathbb{Q}$  that is L-isomorphic to E and satisfies  $E'^n(\mathbb{Q})[3] \neq \{\mathcal{O}\}$ . Therefore, we can assume, without loss of generality, that  $Q_3 \in E'(\mathbb{Q})[3]$ . Now, let  $P_9 \in E(K)[9]$  be a point of order 9, and let  $\phi(P_9)$  denote its image in E'(L)[9]. By Lemma 1.2, the degree  $[\mathbb{Q}(\phi(P_9)) : \mathbb{Q}]$  divides 2. Consequently, by Lemma 4.1, we conclude that L = K or  $L = K(\sqrt{n})$  for some rational n.

# Case 2: $K/\mathbb{Q}$ is not a Galois extension.

Let  $\widehat{K}$  be the Galois closure of K over  $\mathbb{Q}$ . Suppose that the Galois group  $\operatorname{Gal}(\widehat{K}/\mathbb{Q})$  is isomorphic to either  $A_4$  or  $S_4$ :

- (2.1) Suppose  $E'(L)[3] \cong (\mathbb{Z}/3\mathbb{Z}) \oplus (\mathbb{Z}/9\mathbb{Z})$ . Then, by Lemma 5.5,  $\mathbb{Q}(\sqrt{3}) \subseteq L$ . Since  $K/\mathbb{Q}$  has no nontrivial intermediate field, we conclude that  $L = K(\sqrt{3})$ .
- (2.2) Suppose  $E'(K)[3] \cong \{\mathcal{O}\}$  and  $E(K) \cong E'(L)[9] \cong \mathbb{Z}/9\mathbb{Z}$ . Let  $P_9 \in E(K)$  be a point of order 9, and  $\phi(P_9)$  its image in E'(L).

**Subcase 1:** Suppose  $\operatorname{Gal}(\widehat{K}/\mathbb{Q}) \cong S_4$ . We claim that  $\mathbb{Q}([3]P_3)/\mathbb{Q}$  is a Galois extension:

- Indeed, suppose it is not and let  $\sigma \in \operatorname{Gal}(\widehat{L}/\mathbb{Q})$  be such that  $\mathbb{Q}([3]\phi(P_9))^{\sigma} \neq \mathbb{Q}([3]\phi(P_9))$ . Then, by Theorem 4.5,  $\phi(P_9)^{\sigma} \in E'(L^{\sigma})$  also has order 9. We claim that  $E'[9] = \langle \phi(P_9), \phi(P_9)^{\sigma} \rangle \subseteq E'(\widehat{L})$ . Indeed, if a and b are integers such that  $a \cdot \phi(P_9) + b \cdot \phi(P_9)^{\sigma} = \mathcal{O}$ , then both  $a \cdot \phi(P_9)$  and  $b \cdot \phi(P_9)^{\sigma}$  belong to  $E'(L \cap L^{\sigma})$ . However,  $\mathbb{Q}([3]\phi(P_9))$  is not contained in  $L \cap L^{\sigma}$ , hence  $E'(L \cap L^{\sigma})[3] = \{\mathcal{O}\}$  and we conclude that  $a \equiv b \equiv 0 \pmod{9}$ . With this, we get that  $E'[9] \subseteq E'(\widehat{L})$ , and by Lemma 5.5, the 9-th cyclotomic extension  $\mathbb{Q}(\zeta_9)/\mathbb{Q}$  is contained in  $\widehat{L}$ . However, by Lemma 2.3 this is not possible.

Finally, by Theorem 1.2,  $[\mathbb{Q}([3]\phi(P_9)):\mathbb{Q}]$  divides 2, and we conclude, by Lemma 4.1, that  $L = K(\sqrt{m})$  for some  $m \in \mathbb{Q}$ .

**Subcase 2:** Suppose  $\operatorname{Gal}(\widehat{K}/\mathbb{Q}) \cong A_4$ . Let  $[3]P_9 = (x,y) \in E(K)$ . By the definition of the isomorphism  $\phi$ , the x-coordinate,  $x(\phi([3]P_9))$ , of  $\phi([3]P_9)$  belongs to K. Since K has no nontrivial subfields, either  $x(\phi([3]P_9)) \in \mathbb{Q}$  or  $K = \mathbb{Q}(x(\phi([3]P_9)))$ . In the first case,  $\mathbb{Q}(\phi([3]P_9))/\mathbb{Q}$  is a quadratic extension, which implies  $L = K(\sqrt{m})$  for some  $m \in \mathbb{Q}$ . In the second case,  $L = \mathbb{Q}(\phi([3]P_9))$ , which further implies  $\widehat{K} \subseteq \mathbb{Q}(E'[3])$ . By [7, Theorem 3.2],  $\operatorname{Gal}(\mathbb{Q}(E'[3])/\mathbb{Q})$  is isomorphic to one of  $GL_2(\mathbb{F}_3)$ , 3Ns, 3B, or 3Nn (observe that this is the only step

we used that E do not have CM). However, none of these groups have a quotient isomorphic to  $A_4$ .

- At last, suppose the Galois group  $\operatorname{Gal}(\widehat{K}/\mathbb{Q})$  is isomorphic to  $D_4$ :
  - (2.3) Suppose  $E'(K)[3] \cong \{\mathcal{O}\}$ . If  $E'(L)[9] \cong \mathbb{Z}/9\mathbb{Z}$ , then we can use the same proof of the corresponding case where  $\operatorname{Gal}(\widehat{K}/\mathbb{Q})$  is isomorphic to  $S_4$ . If  $E'(L)[9] \cong$  $(\mathbb{Z}/3\mathbb{Z}) \oplus (\mathbb{Z}/9\mathbb{Z})$ , then, by Lemma 5.5,  $\mathbb{Q}(\sqrt{3})$  is the unique nontrivial intermediate field of the extension  $K/\mathbb{Q}$  and  $\mathbb{Q}(E'[3]) \subseteq L$ . If  $[\mathbb{Q}(E'[3]) : \mathbb{Q}]$  is equal to either 1, 2 or 8, then  $L = K(\sqrt{m})$  for some  $m \in \mathbb{Q}$ . If  $\mathbb{Q}(E'[3])/\mathbb{Q}$  is a Galois quartic extension, then it is an abelian extension, isomorphic to either  $(\mathbb{Z}/4\mathbb{Z})$  or  $(\mathbb{Z}/2\mathbb{Z}) \oplus (\mathbb{Z}/2\mathbb{Z})$ . Since, by Lemma 2.1,  $\mathbb{Q}(\sqrt{3})$  is not contained in any cyclic quartic extension of  $\mathbb{Q}$ , we conclude that there exists a quadratic number field  $\mathbb{Q}(\sqrt{m}) \subseteq L$ , with  $m \in \mathbb{Q}$ , other than  $\mathbb{Q}(\sqrt{3})$ , concluding that  $L = K(\sqrt{m})$ .
  - (2.4) Suppose  $E'(K)[3] = \langle Q_3 \rangle \cong \mathbb{Z}/3\mathbb{Z}$ . If  $K = \mathbb{Q}(Q_3)$ , then, by Theorem 4.4,  $L = \mathbb{Q}(E'[3])$  is a Galois extension of  $\mathbb{Q}$ , hence, by the lattice of subgroups of  $D_4$ , we conclude that  $L = K(\sqrt{m})$  for some  $m \in \mathbb{Q}$ . Next, suppose that  $[\mathbb{Q}(Q_3) : \mathbb{Q}]$  divides 2. Let  $\mathbb{Q}(Q_3) = \mathbb{Q}(\sqrt{m}) \subseteq K$ , with  $m \in \mathbb{Q}$ . By Theorem 4.4, we see that up to a twist by m, we can suppose without loss of generality that  $Q_3 \in E'(\mathbb{Q})[3]$ . Now, let  $P_3 \in E(K)$  be a point of order 3, and let  $\phi(P_3)$  be its image in E'(L). Observe that  $\mathbb{Q}(\phi(P_3)) = \mathbb{Q}(E'[3])$  is a Galois extension of  $\mathbb{Q}$ . Using Lemma 1.2, we conclude that  $[\mathbb{Q}(\phi(P_3) : \mathbb{Q}]$  divides 2, and  $L = K(\sqrt{n})$  for some  $n \in \mathbb{Q}$ .

Finally, using Theorem 4.2, we conclude that if  $E(K)_{\text{tors}}$  has points of order 7 or 9, then E is K-isomorphic to an elliptic curve  $\tilde{E}$  defined over  $\mathbb{Q}$ . Therefore,  $E(K)_{\text{tors}}$  is isomorphic to one of the groups of  $\Phi_{\mathbb{Q}}(4)$ , which, as stated in the introduction, was already found by M. Chou, E. González-Jiménez and F. Najman:

THEOREM 5.6 (M. Chou [1], E. González-Jiménez and F. Najman [7]). Let E be an elliptic curve defined over  $\mathbb{Q}$  and let K be a quartic extension of  $\mathbb{Q}$ . Then, the torsion subgroup  $E(K)_{\text{tors}}$  is isomorphic to one of the following groups:

 $\mathbb{Z} / N_1 \mathbb{Z}, \qquad N_1 = 1, \dots, 10, 12, 13, 15, 16, 20, 24, \\ \mathbb{Z} / 2 \mathbb{Z} \oplus \mathbb{Z} / 2 N_2 \mathbb{Z}, \qquad N_2 = 1, \dots, 6, 8, \\ \mathbb{Z} / 3 \mathbb{Z} \oplus \mathbb{Z} / 3 N_3 \mathbb{Z}, \qquad N_3 = 1, 2, \\ \mathbb{Z} / 4 \mathbb{Z} \oplus \mathbb{Z} / 4 N_4 \mathbb{Z}, \qquad N_4 = 1, 2, \\ \mathbb{Z} / 5 \mathbb{Z} \oplus \mathbb{Z} / 5 \mathbb{Z}, \\ \mathbb{Z} / 6 \mathbb{Z} \oplus \mathbb{Z} / 6 \mathbb{Z}.$ 

From this, we conclude that no group in (3) belongs to  $\Phi_{j\in\mathbb{Q}}(4)$ .

5.1. Examples of elliptic curves for each torsion structure. To complete the proof of Theorem 0.2, we need to show elliptic curves having each torsion structure listed in Theorem 0.2. Except for  $\mathbb{Z}/17\mathbb{Z}$  and  $\mathbb{Z}/21\mathbb{Z}$ , all the other groups have already appeared as torsion subgroups of elliptic curves defined over  $\mathbb{Q}$ , as shown in Theorem 5.6.

To illustrate an example of an elliptic curve with torsion subgroup isomorphic to  $\mathbb{Z}/17\mathbb{Z}$ , we present the result found by D. Jeon in [10]. In this case, we are fortunate that the corresponding elliptic curve has a *j*-invariant in  $\mathbb{Q}$ .

EXAMPLE 5.7 (Elliptic curve with a 17-torsion, D. Jeon [10]). Consider the quartic cyclic number field  $K = \mathbb{Q}[X]/(f(X))$ , where  $f(X) = 2X^4 - 5X^3 - 7X^2 + 10X + 8$ , and let  $\alpha$  be one of the roots of f(X) in K. Define the constants b and c as follows:

$$\begin{split} c &= -\frac{1337}{32\alpha^3} + \frac{3233}{64\alpha^2} + \frac{12285}{64\alpha} + \frac{1135}{16},\\ b &= -\frac{59}{16\alpha^3} + \frac{147}{32\alpha^2} + \frac{535}{32\alpha} + \frac{45}{8}. \end{split}$$

Consider the elliptic curve E = E(b, c), given by the equation

$$E: y^{2} + (1 - c)xy - by = x^{3} - bx.$$

Using MAGMA, it can be verified that j(E) = -882216989/131072, and its torsion subgroup  $E(K)_{\text{tors}}$  is isomorphic to  $\mathbb{Z}/17\mathbb{Z}$ .

Analogously, P. L. Clark, P. Corn, A. Rice and J. Stankewicz already found in [2] an elliptic curve, in this case with complex multiplication, with torsion subgroup isomorphic to  $\mathbb{Z}/21\mathbb{Z}$ :

EXAMPLE 5.8 (Elliptic curve with a 21-torsion, P. L. Clark, P. Corn, A. Rice and J. Stankewicz [2]). Consider the quartic number field  $K = \mathbb{Q}[X]/(f(X))$ , where  $f(X) = X^4 - X^3 + 2X + 1$ , and let  $\alpha$  be one of the roots of f(X) in K. Consider the elliptic curve E, given by the equation

$$E: y^2 = x^3 - (371952\alpha^3 + 3373488\alpha^2 + 3777840\alpha + 1228608).$$

Then, j(E) = 0 and  $E(K)_{\text{tors}} \cong \mathbb{Z}/21\mathbb{Z}$ .

## References

- M. CHOU, Torsion of rational elliptic curves over quartic Galois number fields, J. Number Theory, 160 (2016), 603–628.
- [2] P. L. CLARK, P. CORN, A. RICE AND J. STANKEWICZ, Computation on elliptic curves with complex multiplication, LMS J. of Comp. and Math., 1 (2014), 509–535.
- [3] M. DERICKX AND F. NAJMAN, Classification of torsion of elliptic curves over quartic fields, arXiv:2412.16016 [math.NT].
- [4] M. DERICKX, A. ETROPOLSKI, M. HOEIJ, J. S. MORROW, AND D. ZUREICK-BROWN, Sporadic cubic torsion, Algebra Number Theory, 15 (2021), no. 7, 1837–1864.

- [5] E. GONZÁLEZ-JIMÉNEZ AND J. TORNERO, Torsion of rational elliptic curves over quadratic fields, Rev. R. Acad. Cienc. Exactas, Fuís. Nat. (Esp.), 108 (2014), issue 2, 923–934.
- [6] E. GONZÁLEZ-JIMÉNEZ, Complete classification of the torsion structures of rational elliptic curves over quintic number fields, J. Algebra, 478 (2016), 484–505.
- [7] E. GONZÁLEZ-JIMÉNEZ AND F. NAJMAN, Growth of torsion groups of elliptic curves upon base change, Math. Comput., 89 (2016), 1457–1485.
- [8] T. GUŽIĆ, Torsion of elliptic curves with rational *j*-invariant defined over number fields of prime degree, Proc. Amer. Math. Soc., 149 (2021), 3261–3275.
- [9] L. HAMADA, Torsion of elliptic curves with rational *j*-invariant over the maximal elementary abelian 2-extension of  $\mathbb{Q}$ , arXiv:2412.11202 [math.NT].
- [10] D. JEON, Tetraelliptic modular curves  $X_1(N)$ , arXiv:2305.05851 [math.NT]
- [11] D. JEON, C. H. KIM, AND A. SCHWEIZER, On the torsion of elliptic curves over cubic number fields, Acta Arith., 113 (2004), 291–301.
- [12] S. KAMIENNY, Torsion points on elliptic curves over all quadratic fields, Duke Math. J., 53 (1986), 157–162.
- [13] M. A. KENKU AND F. MOMOSE, Torsion points on elliptic curves defined over quadratic fields, Nagoya Math. J., 109 (1988), 125–149.
- [14] B. MAZUR, Modular curves and the eisenstein ideal, Publ. Math. Inst. Hautes Études Sci., 47 (1977), 33–186.
- [15] F. NAJMAN, Torsion of rational elliptic curves over cubic fields and sporadic points on  $X_1(n)$ , Math. Res. Lett., 23 (2016), no. 1, 245–272.
- [16] J. E. CREMONA AND F. NAJMAN, Q-curves over odd degree number fields, Res. Number Theory, 7 (2021).
- [17] J.-P. SERRE, A course in arithmetic, Grad. texts in math., Springer, 1973.
- [18] J. H. SILVERMAN, The Arithmetic of Elliptic Curves, Grad. texts in math., Springer, 2009.

DEPARTMENT OF MATHEMATICS INSTITUTE OF SCIENCE TOKYO 2-12-1, O-OKAYAMA, MEGURO-KU, TOKYO 152-8551 JAPAN *Email address*: lucas.h.r.hamada@gmail.com