

# On the Wiedemann–Franz law violation in Graphene and quark-gluon plasma systems

Ashutosh Dwivedi<sup>1</sup>, Subhalaxmi Nayak<sup>1</sup>, Sathe Subodh Kiran<sup>2</sup>, Sabyasachi Ghosh<sup>1</sup>, Sesa Vempati<sup>1</sup>

<sup>1</sup>*Department of Physics, Indian Institute of Technology Bhilai,  
Kutalabhata, Durg, 491002, Chhattisgarh, India and*

<sup>2</sup>*Department of Physics, Sardar Vallabhbhai National Institute of Technology, Ichchhanath, Surat, 395007, Gujarat, India*

A comparative study of the thermodynamic and transport properties of the ultra-relativistic quark-gluon plasma (QGP) produced in Heavy ion collisions (HIC) with the "quasi-relativistic" massless electron-hole plasma in graphene sample has been performed. We observe that the enthalpy per net carrier density emerges as a useful physical quantity determining the hydrodynamic domain's transport variables. Lorenz ratio is defined as thermal to electrical conductivity ratio, normalized by temperature. In searching whether the Wiedemann-Franz (WF) law is obeyed or violated by checking the Lorenz ratio as one or deviated from one, we find that the Lorenz ratio determined from the fluid-based framework will always be responsible for the violation of the WF law. The reason is the proportional relation between Lorenz ratio and enthalpy per particle in the fluid. Based on the experimental observation, graphene, and QGP, both systems at low net charge density, exhibit WF law violation due to their fluid nature. However, graphene at high net charge density obeys the WF law, followed by metals with high Fermi energy or density. It indicates a fluid to the non-fluid transition of the graphene system from low to high-density domain. In this regard, the fluid or non-fluid aspect of QGP at high density is yet to be explored by future facilities like Compressed Baryonic matter (CBM) and Nuclotron-based Ion Collider Facility (NICA) experiments.

## I. INTRODUCTION

Understanding the charge carrier/quasi-particle transport under the influence of external perturbation such as electric field and/or thermal gradient is fundamentally vital in the context of device physics. Their concerning transport coefficients are electrical ( $\sigma$ ) and thermal ( $\kappa$ ) conductivities. Although  $\sigma$  and  $\kappa$  appear to be intrinsic properties, they crucially depend on various other parameters such as carrier and defect density, energy dispersion relation, Coulombic, and other scatterers, mean free path with reference to the size of the sample, temperature ( $T$ ) etc [1–35]. Notably, the  $\sigma$  and  $\kappa$  can be connected *via* the WF law, which yields a constant for conventional metals and known as Lorentz number ( $L_0$ ) [36–38]. Violations to the WF law are observed in graphene [39–42] and QGP systems [43–47], where the transport behavior of the carrier is not similar to that of conventional metals. Although these two systems are at the extreme ends of the chemical potential ( $\mu$ ) spectrum, the constituents share an exotic common property *viz* 'hydrodynamic transport' [48] that causes the violation of WF law. Furthermore, various physical characteristics such as masslessness, linear dispersion of energy-momentum relation ( $E_{\pm} = \pm\hbar|\vec{k}|v_F \equiv \pm|\vec{p}|v_F$  and  $E = |\vec{p}|c$  for graphene and QGP respectively) of charge carriers are notably similar. For graphene, the effective mass of the charge carriers vanish, and they, in many aspects, behave like ultra-relativistic Dirac particles [49, 50] similar to the constituents of QGP at the core of neutron star or in heavy ion collision experiments. On the other hand, in graphene, the  $\mu$  can be tuned by using doping techniques [1] while that of in QGP baryonic- $\mu$  can be altered by changing the energy of the collision-beam [51]. In the context of hydrodynamics behavior,  $\mu - T$  plane can be grossly classified into Dirac fluid (DF) or Fermi liquid (FL) regions [52–54]. For the domain  $\frac{\mu}{k_B T} \gg 1$ , the electrons in graphene obey the well-known Fermi liquid theory [52–54], whereas, in QGP, the same may be expected at the core of neutron stars. Here  $k_B$  is Boltzmann constant and  $T$  is temperature. For  $\frac{\mu}{k_B T} \ll 1$ , both QGP and lightly doped graphene are identified to follow fluidic behavior. In the case of QGP, ultra-relativistic hydrodynamics is highly successful in explaining the particle spectra and anisotropic flow coefficients extracted from the experimental data, see Ref. [51] and also the references therein. In the context of graphene various studies [55–64] have explored the experimental aspects of electron hydrodynamics. From theoretical viewpoints the electron hydrodynamics has also been explored focusing on various aspects like shear, hall viscosities and corresponding KSS bound [65–67]. More recently there is also experiment [63] on the determination of  $L \equiv \kappa/(\sigma T)$  whose results show a strong deviation of Lorentz ratio (LR)  $L/L_0$  from one in the DF regime of graphene. This leads to intensive theoretical investigation seeking the explanation of enhanced LR near the Dirac point of graphene. The authors of Ref. [39] have used the relativistic fluid dynamics formalism in the DF regime to explain the experimentally observed enhancement in LR, whereas in Ref. [40], the authors developed a "disorder enabled hydrodynamics" to explain the same. In Ref. [42], on the other hand, the authors explained the enhancement in LR with the inclusion of bipolar diffusion effect and a band gap at the Dirac point.

In this paper, by considering the similarity between QGP and graphene, we have developed a microscopic theory for the latter that takes care of hydrodynamic transport in the domain of low charge impurity densities/charge

puddles. We calculated the  $\kappa$  and  $\sigma$  and then Lorentz ratio for graphene (QGP) systems by explicitly considering the contribution to  $L$  from electrons and holes (quarks and antiquarks). LR for graphene with respect to net carrier density are compared with the experimental data [63] where a good agreement in the DF domain is obtained.

The article is arranged as follows. In Sec. II A and Sec. II B, we derive respectively the LR for graphene and QGP starting from the Boltzmann transport equation (BTE) in relaxation time approximation (RTA). Next, we compare the results of graphene and QGP with the plots of net number density, energy density, pressure, enthalpy, electrical conductivity, thermal conductivity, and LR in Sec. III. In Sec. IV, we conclude by summarizing our investigations.

## II. FORMALISM

### A. Thermoelectric transports in Graphene

The carrier transport in the materials can be described with the help of the BTE, which determines the fate of charge carriers in different energy bands [37, 38]. In graphene, the low-energy electron excitations can be modeled by a two-band electronic system comprised of the valence band and conduction band. The BTE for the two-band electronic system in the presence of electric field  $\vec{E}$  can be written as [68, 69]:

$$\frac{\partial f_\lambda}{\partial t} + \vec{v}_\lambda \cdot \frac{\partial f_\lambda}{\partial \vec{r}} - e\vec{E} \cdot \frac{\partial f_\lambda}{\partial \vec{p}_\lambda} = C_\lambda[f_\lambda], \quad (1)$$

where the band index  $\lambda = +$  for conduction band and  $-$  for valence band. The microscopic variables quasi-momentum (or crystal momentum), energy, and the group velocity of the electrons are respectively defined as  $\vec{p}_\lambda \equiv \hbar\vec{k}_\lambda$ ,  $E_\lambda \equiv E_\lambda(\vec{k}_\lambda)$ , and  $\vec{v}_\lambda \equiv \frac{1}{\hbar} \frac{\partial E_\lambda}{\partial \vec{k}_\lambda}$  with  $\vec{k}_\lambda$  being the wave vector in the reciprocal space. The collision kernel  $C_\lambda[f_\lambda] = \left(\frac{\partial f_\lambda}{\partial t}\right)_{coll}$  gives rise to changes in the distribution function due to random incessant collisions. The  $C_\lambda[f_\lambda]$  contains all the information about the momentum-conserving and momentum-non-conserving interaction of the electrons with other electrons, phonons, and lattice defects [70]. We will consider a temperature window where the momentum non-conserving scatterings are negligible compared to momentum-conserving electron-electron scatterings. In this temperature regime, the hydrodynamical electron flows are expected contrary to the diffusive flow of electrons observed in metals in normal conditions [70]. Therefore, in the temperature window under consideration, the collision kernel  $C_\lambda[f_\lambda]$  ensures energy, momentum, and charge conservation. Using the dispersion relation  $E_\lambda = \lambda v_F \hbar |\vec{k}_\lambda|$  the group velocity of the electrons in the vicinity of Dirac point is obtained as  $\vec{v}_\lambda = \frac{1}{\hbar} \frac{\partial E_\lambda}{\partial \vec{k}_\lambda} = \lambda v_F \frac{\vec{k}_\lambda}{|\vec{k}_\lambda|}$ . Rewriting Eq. (1) with the substitution of the group velocity  $\vec{v}_\lambda = \lambda v_F \frac{\vec{k}_\lambda}{|\vec{k}_\lambda|}$  we have,

$$\frac{E_\lambda}{v_F^2} \frac{\partial f_\lambda}{\partial t} + \hbar\vec{k}_\lambda \cdot \frac{\partial f_\lambda}{\partial \vec{r}} - e\vec{E} \cdot \left( \hbar\vec{k}_\lambda \frac{\partial f_\lambda}{\partial E_\lambda} + \frac{E_\lambda}{v_F^2} \frac{\partial f_\lambda}{\partial \vec{p}_\lambda} \right) = \frac{E_\lambda}{v_F^2} C[f_\lambda]. \quad (2)$$

Eq. (2) describes the dynamics of both the conduction band and valence band electrons in the graphene near the Dirac cone. Since it is customary to describe the valence band carriers by holes, we will write down the BTE for valence band carriers with the following change of variables  $f_h \equiv 1 - f_-$ ,  $E_h \equiv -E_-$  and  $\vec{k}_h \equiv -\vec{k}_-$  as,

$$\frac{E_h}{v_F^2} \frac{\partial f_h}{\partial t} + \hbar\vec{k}_h \cdot \frac{\partial f_h}{\partial \vec{r}} + e\vec{E} \cdot \left( \hbar\vec{k}_h \frac{\partial f_h}{\partial E_h} + \frac{E_h}{v_F^2} \frac{\partial f_h}{\partial \vec{p}_h} \right) = -\frac{E_h}{v_F^2} C_h[1 - f_h], \quad (3)$$

where we defined  $C_-[1 - f_h] \equiv C_h[1 - f_h]$ . Similarly, the BTE for the conduction band electrons in graphene can be rewritten by calling  $f_+ \equiv f_e$ ,  $E_+ = E_e$ ,  $\vec{k}_+ = \vec{k}_e$  and  $C_+[f_+] \equiv C_e[f_e]$  as:

$$\frac{E_e}{v_F^2} \frac{\partial f_e}{\partial t} + \hbar\vec{k}_e \cdot \frac{\partial f_e}{\partial \vec{r}} + e\vec{E} \cdot \left( \hbar\vec{k}_e \frac{\partial f_e}{\partial E_e} + \frac{E_e}{v_F^2} \frac{\partial f_e}{\partial \vec{p}_e} \right) = \frac{E_e}{v_F^2} C_e[f_e]. \quad (4)$$

Eq. (4) and (3) form the basis of the dynamics of the electrons and holes in the graphene.

The relations:  $\vec{p}_{e,h} = \hbar\vec{k}_{e,h}$ ,  $E_{e,h} = (\hbar k_{e,h})v_F$ , and  $\vec{v}_{e,h} = v_F \frac{\vec{k}_{e,h}}{k_{e,h}}$  is similar to a massless relativistic (ultra-relativistic) particle in a world where the limiting speed is  $v_F$ . Moreover, there exists numerous literature [39, 67, 70–76] in the field of graphene where the relativistic behavior of electrons in graphene has been explored regarding  $v_F$  as the limiting speed in the graphene world. Recently, in Ref. [77], the authors have calculated the ratio of shear viscosity to entropy density of the fluid in graphene by drawing an analogy with the relativistic hydrodynamics used

in the literature of QGP. The similarities and differences of the hydrodynamics practiced in the literature QGP and graphene were described in detail there. Here we will follow the same prescription to write the Eq. (4) and (3) in a covariant form where the role of universal speed  $c$  is replaced with the limiting speed  $v_F$  in the graphene world. By defining the four position vector  $x^\mu$ , four-momentum of electrons  $p_e^\mu$  and four-momentum of holes  $p_h^\mu$  (with  $\mu = 0$  to 2) as  $x^\mu \equiv (v_F t, x^i)$ ,  $p_e^\mu \equiv (\frac{E_e}{v_F}, p_e^i = \hbar k_e^i)$  and  $p_h^\mu \equiv (\frac{E_h}{v_F}, p_h^i = \hbar k_h^i)$ , Eq. (4) and (3) can be restated as:

$$p_e^\mu \partial_\mu f_e - e F^{\mu\nu} p_{e\nu} \frac{\partial f_e}{\partial p_e^\mu} = \tilde{C}_e[f_e], \quad (5)$$

$$p_h^\mu \partial_\mu f_h + e F^{\mu\nu} p_{h\nu} \frac{\partial f_h}{\partial p_h^\mu} = \tilde{C}_h[f_h], \quad (6)$$

where we adopted the relativistic notation  $\partial_\mu \equiv \frac{\partial}{\partial x^\mu}$  and redefined the collision terms as  $\tilde{C}_e[f_e] \equiv \frac{E_e}{v_F^2} C_e[f_e]$  and  $\tilde{C}_h[f_h] \equiv -\frac{E_h}{v_F^2} C_h[1 - f_h]$ . The  $F^{\mu\nu} = \frac{1}{v_F^2} (\tilde{E}^\mu u^\nu - \tilde{E}^\nu u^\mu)$  is the Faraday tensor with the fluid four-velocity  $u^\mu = \gamma_u (v_F, \vec{u})$ , where  $\gamma_u = 1/\sqrt{1 - u^2/v_F^2}$ .  $\tilde{E}^\mu$  is the comoving electric field perpendicular to  $u^\mu$  i.e.,  $\tilde{E}^\mu u_\mu = 0$ . One can observe the similarity of the electron-hole BTE of graphene given in Eq. (5) and (6) with the BTE electron-positron plasma [78–80] found in the literature of astrophysics or the quark-antiquark plasma [81–83] created in heavy ion collision experiments performed in Large Hadron Collider (LHC) or Relativistic Heavy Ion Collider (RHIC). This similarity will be used to define particle flow, stress-energy tensor, and the corresponding thermodynamic variables in a covariant manner. The net particle four flow  $N^\mu$  for the electron hydrodynamics in graphene is defined as the electron flow minus the hole flow:

$$N^\mu \equiv N_e^\mu - N_h^\mu = 4v_F^2 \left[ \int \frac{d^2 \vec{p}_e}{h^2 E_e} p_e^\mu f_e - \int \frac{d^2 \vec{p}_h}{h^2 E_h} p_h^\mu f_h \right]. \quad (7)$$

where 4 is the degeneracy factor, which is the product of spin and valley degeneracies, i.e., total degeneracy = spin degeneracy (2)  $\times$  valley degeneracy (2) = 4. Similarly, the total stress-energy tensor for the electron hydrodynamics in graphene is defined as:

$$T^{\mu\nu} \equiv T_e^{\mu\nu} + T_h^{\mu\nu} = 4v_F^2 \left[ \int \frac{d^2 \vec{p}_e}{h^2 E_e} p_e^\mu p_e^\nu f_e + \int \frac{d^2 \vec{p}_h}{h^2 E_h} p_h^\mu p_h^\nu f_h \right]. \quad (8)$$

In the out-of-equilibrium situation, the stress-energy tensor and the net particle flow can be split into two parts:  $T^{\mu\nu} = T_0^{\mu\nu} + T_D^{\mu\nu}$  and  $N^\mu = N_0^\mu + N_D^\mu$ . The  $T_0^{\mu\nu}$  and  $N_0^\mu$  are the ideal parts that give information about the thermodynamics of the system.  $T_D^{\mu\nu}$  and  $N_D^\mu$  are the dissipative parts containing information about the various dissipative flows like shear flow, energy diffusion, particle diffusion, etc. The ideal parts  $N_0^\mu$  and  $T_0^{\mu\nu}$  can be expressed as integrals over the local equilibrium distribution function  $f_e^0 = 1/(e^{(u_\mu p_e^\mu - \mu_e)/k_B T} + 1)$  and  $f_h^0 = 1/(e^{(u_\mu p_h^\mu - \mu_h)/k_B T} + 1)$  with  $\mu_e = \mu$  and  $\mu_h = -\mu$ :

$$N_0^\mu \equiv 4v_F^2 \left[ \int \frac{d^2 \vec{p}_e}{h^2 E_e} p_e^\mu f_e^0 - \int \frac{d^2 \vec{p}_h}{h^2 E_h} p_h^\mu f_h^0 \right] = n u^\mu, \quad (9)$$

$$T_0^{\mu\nu} \equiv 4v_F^2 \left[ \int \frac{d^2 \vec{p}_e}{h^2 E_e} p_e^\mu p_e^\nu f_e^0 + \int \frac{d^2 \vec{p}_h}{h^2 E_h} p_h^\mu p_h^\nu f_h^0 \right] = v_F^{-2} \mathcal{E} u^\mu u^\nu - P \Delta^{\mu\nu}, \quad (10)$$

where  $\Delta^{\mu\nu} = \eta^{\mu\nu} - v_F^{-2} u^\mu u^\nu$  and  $n$ ,  $\mathcal{E}$  and  $P$  are respectively the net number density, energy density, and pressure of the electron-hole fluid system. It is easy to see from Eq. (9) and (10) that the thermodynamic variables  $n$ ,  $\mathcal{E}$  and  $P$  can be expressed as:

$$n = v_F^{-2} u_\mu N_0^\mu = n_e - n_h = 4 \left[ \int \frac{d^2 \vec{p}_e}{h^2} f_e^0 - \int \frac{d^2 \vec{p}_h}{h^2} f_h^0 \right], \quad (11)$$

$$\mathcal{E} = v_F^{-2} u_\mu u_\nu T_0^{\mu\nu} = \mathcal{E}_e + \mathcal{E}_h = 4 \left[ \int \frac{d^2 \vec{p}_e}{h^2} E_e f_e^0 + \int \frac{d^2 \vec{p}_h}{h^2} E_h f_h^0 \right], \quad (12)$$

$$P = -\frac{1}{2} \Delta_{\mu\nu} T_0^{\mu\nu} = P_e + P_h = 4v_F^2 \left[ \int \frac{d^2 \vec{p}_e}{h^2} \frac{p_e^2}{2E_e} f_e^0 + \int \frac{d^2 \vec{p}_h}{h^2} \frac{p_h^2}{2E_h} f_h^0 \right]. \quad (13)$$

The above phase space integrals for the thermodynamic variables have been evaluated in Appendix VIB in terms of fermi integral functions. Now, the dissipative part of the particle flow  $N_D^\mu$  and stress-energy tensor  $T_D^{\mu\nu}$  can be

expressed as integral over  $\delta f_e \equiv f_e - f_e^0$  and  $\delta f_h \equiv f_h - f_h^0$  as,

$$N_D^\mu \equiv 4v_F^2 \left[ \int \frac{d^2 \vec{p}_e}{h^2 E_e} p_e^\mu \delta f_e - \int \frac{d^2 \vec{p}_h}{h^2 E_h} p_h^\mu \delta f_h \right], \quad (14)$$

$$T_D^{\mu\nu} \equiv 4v_F^2 \left[ \int \frac{d^2 \vec{p}_e}{h^2 E_e} p_e^\mu p_e^\nu \delta f_e + \int \frac{d^2 \vec{p}_h}{h^2 E_h} p_h^\mu p_h^\nu \delta f_h \right]. \quad (15)$$

The thermoelectric coefficients connected with the electron hydrodynamics in graphene can be obtained from the knowledge of  $N_D^\mu$ . For the evaluation of  $N_D^\mu$  from Eq. (14) one needs to determine the unknowns  $\delta f_e$  and  $\delta f_h$  with the help of Eq. (5) and (6). For the determination of  $\delta f_e$  and  $\delta f_h$  we will rewrite Eq. (5) and (6) in the RTA of Anderson-Witting type [84] i.e.  $C_e[f_e] = -\frac{u_\mu p_e^\mu}{v_F^2} \frac{\delta f_e}{\tau_c}$  and  $C_h[f_h] = -\frac{u_\mu p_h^\mu}{v_F^2} \frac{\delta f_h}{\tau_c}$ , where we assumed same collision time  $\tau_c$  for electrons and holes. At this juncture, it is necessary to discuss our choice of hydrodynamic frame and the energy dependence of the relaxation time. In the theory of relativistic dissipative hydrodynamics, the choice of hydrodynamic frame is of primary importance in which one defines the fluid velocity and thermodynamic variables through the use of matching conditions [85–87]. In Anderson-Witting type RTA models one usually works from the Landau-Lifshitz hydrodynamic frame<sup>1</sup> with energy independent relaxation time. Nevertheless, the actual relaxation time for a system is energy-dependent, and its energy dependence can be calculated for given interactions using quantum mechanical or quantum field theoretical methods. Since the present article is planned to see the effect of the fluid dynamic nature of the electron flow on the structure of thermoelectric coefficients, our assumption of a constant relaxation time is justifiable. The Eq. (5) and (6) in the RTA with the substitution of  $f_{e,h} = f_{e,h}^0 + \delta f_{e,h}$  become,

$$p_e^\mu \partial_\mu f_e^0 - \frac{e}{v_F^2} (\tilde{E}^\mu u^\nu - \tilde{E}^\nu u^\mu) p_{e\nu} \frac{\partial f_e^0}{\partial p_e^\mu} = -\frac{u_\mu p_e^\mu}{v_F^2} \frac{\delta f_e}{\tau_c}, \quad (16)$$

$$p_h^\mu \partial_\mu f_h^0 + \frac{e}{v_F^2} (\tilde{E}^\mu u^\nu - \tilde{E}^\nu u^\mu) p_{h\nu} \frac{\partial f_h^0}{\partial p_h^\mu} = -\frac{u_\mu p_h^\mu}{v_F^2} \frac{\delta f_h}{\tau_c}, \quad (17)$$

where we neglected the space-time gradients of  $\delta f_{e,h}$  from the LHS of the equations [90]. Evaluating the space-time gradients of  $f_{e,h}^0$  by using  $f_e^0 = 1/(e^{(u_\mu p_e^\mu - \mu_e)/k_B T} + 1)$  and  $f_h^0 = 1/(e^{(u_\mu p_h^\mu - \mu_h)/k_B T} + 1)$  in Eq. (16) and (17) with  $\mu \equiv \mu_e = -\mu_h$  we have,

$$\begin{aligned} \delta f_e = & \frac{\tau_c v_F^2}{u_\mu p_e^\mu} \left[ v_F^{-2} (u_\alpha p_e^\alpha) \left( (u_\beta p_e^\beta) D \frac{1}{k_B T} + \frac{p_e^\alpha D u_\alpha}{k_B T} - D \frac{\mu}{k_B T} \right) + \frac{p_e^\alpha p_e^\beta}{k_B T} \nabla_\alpha u_\beta + (u_\alpha p_e^\alpha) p_e^\beta \nabla_\beta \frac{1}{k_B T} \right. \\ & \left. - p_e^\alpha \nabla_\alpha \frac{\mu}{k_B T} + \frac{e p_e^\alpha}{k_B T} \tilde{E}_\alpha \right] f_e^0 (1 - f_e^0), \end{aligned} \quad (18)$$

$$\begin{aligned} \delta f_h = & \frac{\tau_c v_F^2}{u_\mu p_h^\mu} \left[ v_F^{-2} (u_\alpha p_h^\alpha) \left( (u_\beta p_h^\beta) D \frac{1}{k_B T} + \frac{p_h^\alpha D u_\alpha}{k_B T} + D \frac{\mu}{k_B T} \right) + \frac{p_h^\alpha p_h^\beta}{k_B T} \nabla_\alpha u_\beta + (u_\alpha p_h^\alpha) p_h^\beta \nabla_\beta \frac{1}{k_B T} \right. \\ & \left. + p_h^\alpha \nabla_\alpha \frac{\mu}{k_B T} - \frac{e p_h^\alpha}{k_B T} \tilde{E}_\alpha \right] f_h^0 (1 - f_h^0), \end{aligned} \quad (19)$$

where the spatial and temporal derivative operators  $\nabla^\mu$  and  $D$  are defined as  $\nabla^\mu \equiv \Delta^{\mu\nu} \partial_\nu \xrightarrow{LRF} (0, -\vec{\nabla})$  and  $D \equiv u^\mu \partial_\mu \xrightarrow{LRF} \frac{\partial}{\partial t}$ . Eq. (18) along with Eq. (19) can be readily used for the calculation of the thermoelectric coefficients of graphene. We will consider a full dynamical scenario in which the fluid velocity profile possesses a space-time gradient. Since the dissipative fluxes like shear flow, thermal flow, etc., are proportional to spatial gradients, the temporal derivatives of  $1/T$ ,  $\mu/T$ , and  $u^\mu$  occurring in Eq. (18) and (19) should be eliminated with the help of the conservation equation of ideal electron hydrodynamics in graphene,

$$Dn = -n \nabla_\mu u^\mu, \quad (20)$$

$$D u^\mu = \frac{v_F^2}{\mathcal{E} + P} \left[ \nabla^\mu P + \rho \tilde{E}^\mu \right], \quad (21)$$

$$D\mathcal{E} = -(\mathcal{E} + P) \nabla_\mu u^\mu, \quad (22)$$

<sup>1</sup> In Landau-Lifshitz frame the dissipative part of the energy flow  $W^\lambda$  vanishes [88, 89] i.e.,  $W^\lambda \equiv \Delta_\mu^\lambda T^{\mu\nu} u_\nu = 0$ .

where  $\rho \equiv -en = -e(n_e - n_h)$  is the charge density. The Eq. (20) ensures charge conservation, and Eq. (21) and (22) provide the energy-momentum conservation for the electron hydrodynamics. After all the temporal derivatives have been eliminated from Eq. (18) and (19) the  $\delta f_{e,h}$  can be easily expressed as a linear combination of three independent spatial gradient terms:  $\nabla_\mu u^\mu$ ,  $\frac{1}{2}(\nabla^\mu u^\nu + \nabla^\nu u^\mu) - \frac{1}{2}(\nabla_\alpha u^\alpha)\Delta^{\mu\nu}$ , and  $-\nabla^\alpha \frac{\mu}{k_B T} + \frac{e\tilde{E}^\alpha}{k_B T}$ . The first ( $\nabla_\mu u^\mu$ ) and second term ( $\frac{1}{2}(\nabla^\mu u^\nu + \nabla^\nu u^\mu) - \frac{1}{2}(\nabla_\alpha u^\alpha)\Delta^{\mu\nu}$ ) give rise to bulk and shear stresses in the fluid (see appendix VI A). They are of primary importance for the calculation of bulk and shear viscosity. Since the present article is structured for the calculation of thermoelectric coefficients, we will neglect viscous stresses to write,

$$\delta f_e = -\frac{\tau_c v_F^2}{u_\mu p_e^\mu} p_e^\alpha \left[ \frac{n}{\mathcal{E} + P} u_\beta p_e^\beta - 1 \right] \left( -\nabla_\alpha \frac{\mu}{k_B T} + \frac{e\tilde{E}_\alpha}{k_B T} \right) f_e^0 (1 - f_e^0), \quad (23)$$

$$\delta f_h = -\frac{\tau_c v_F^2}{u_\mu p_h^\mu} p_h^\alpha \left[ \frac{n}{\mathcal{E} + P} u_\beta p_h^\beta + 1 \right] \left( -\nabla_\alpha \frac{\mu}{k_B T} + \frac{e\tilde{E}_\alpha}{k_B T} \right) f_h^0 (1 - f_h^0), \quad (24)$$

The expression for the current density (dissipative part of charge flow)  $J^\mu$  can be written as,

$$J^\mu \equiv -e\Delta_\nu^\mu N_D^\nu = -4e v_F^2 \Delta_\nu^\mu \left[ \int \frac{d^2 \vec{p}}{h^2 E_e} p_e^\nu \delta f_e - \int \frac{d^2 \vec{p}_h}{h^2 E_h} p_h^\nu \delta f_h \right],$$

substituting  $u^\mu = (v_F, \vec{0})$  in the above expression we have,

$$J^i = -4e v_F^4 \tau_c \left[ \int \frac{d^2 \vec{p}_e}{h^2} \frac{p_e^i p_e^j}{E_e^2} \left( \frac{n E_e}{\mathcal{E} + P} - 1 \right) f_e^0 (1 - f_e^0) - \int \frac{d^2 \vec{p}_h}{h^2} \frac{p_h^i p_h^j}{E_h^2} \left( \frac{n E_h}{\mathcal{E} + P} + 1 \right) f_h^0 (1 - f_h^0) \right] X^j, \quad (25)$$

where we defined  $X^i \equiv \partial_i \frac{\mu}{k_B T} + \frac{e\tilde{E}^i}{k_B T}$ . We can simplify Eq. (25) (see Appendix VI B) to obtain,

$$J^i = \frac{4e\pi\tau_c}{h^2} \frac{nk_B T}{\mathcal{E} + P} \left[ 2(k_B T)^2 (f_2(A^{-1}) - f_2(A)) + \frac{\mathcal{E} + P}{n} k_B T (f_1(A^{-1}) + f_1(A)) \right] X^i$$

or,  $J^\alpha = \frac{4e\pi\tau_c}{h^2} \frac{nk_B T}{\mathcal{E} + P} \left[ 2(k_B T)^2 (f_2(A^{-1}) - f_2(A)) + \frac{\mathcal{E} + P}{n} k_B T (f_1(A^{-1}) + f_1(A)) \right] \left( -\nabla^\alpha \frac{\mu}{k_B T} + \frac{e\tilde{E}^\alpha}{k_B T} \right). \quad (26)$

The heat flow  $q^\mu$  for relativistic fluid is defined as the difference between dissipative part of energy flow  $W^\mu \equiv \Delta_\alpha^\mu T^{\alpha\beta} u_\beta$  and enthalpy flow  $h^\mu \equiv \mathfrak{h} \Delta_\alpha^\mu N_D^\alpha$  [88, 89] i.e.,  $q^\mu = W^\mu - h^\mu = \Delta_\alpha^\mu (T^{\alpha\beta} u_\beta - \mathfrak{h} N_D^\alpha)$ , where  $\mathfrak{h} \equiv \frac{\mathcal{E} + P}{n}$  is the enthalpy per particle. In Anderson-Witting type of RTA model one works in the Landau-Lifshitz hydrodynamic frame where the dissipative part of energy flow vanishes [88, 89] and the expression of heat flow becomes:  $q^\mu = -h^\mu = -\frac{\mathcal{E} + P}{n} \Delta_\alpha^\mu N_D^\alpha = -(\mathcal{E} + P) \frac{J^\mu}{\rho}$ . The preceding definition of heat flow with Eq. (26) give the following expression for  $q^\alpha$ :

$$q^\alpha = \frac{4\pi\tau_c k_B T}{h^2} \left[ 2(k_B T)^2 (f_2(A^{-1}) - f_2(A)) + \frac{\mathcal{E} + P}{n} k_B T (f_1(A^{-1}) + f_1(A)) \right] \left( -\nabla^\alpha \frac{\mu}{k_B T} + \frac{e\tilde{E}^\alpha}{k_B T} \right). \quad (27)$$

We can rewrite the expression of current density and heat flow with the help of Gibbs-Duhem relation  $n d(\frac{\mu}{k_B T}) = \frac{1}{k_B T} dP + (\mathcal{E} + P) d(\frac{1}{k_B T})$  as:

$$J^\mu = \frac{4e\pi\tau_c}{h^2} (k_B T)^2 \left[ 2(f_2(A^{-1}) - f_2(A)) + \frac{\mathcal{E} + P}{nk_B T} (f_1(A^{-1}) + f_1(A)) \right] \left[ -\frac{\rho}{\mathcal{E} + P} \tilde{\mathcal{E}}^\mu + \frac{1}{T} \nabla^\mu T \right], \quad (28)$$

$$q^\mu = \frac{4\pi\tau_c}{h^2} (k_B T)^2 \frac{\mathcal{E} + P}{n} \left[ 2(f_2(A^{-1}) - f_2(A)) + \frac{\mathcal{E} + P}{nk_B T} (f_1(A^{-1}) + f_1(A)) \right] \left[ -\frac{\rho}{\mathcal{E} + P} \tilde{\mathcal{E}}^\mu + \frac{1}{T} \nabla^\mu T \right], \quad (29)$$

where we used the definition  $\tilde{\mathcal{E}}^\mu \equiv \tilde{E}^\mu + \frac{1}{\rho} \nabla^\mu P$ . The electrical and thermal conductivity of the electron hydrodynamics can be identified by comparing the microscopically derived expressions of vectorial dissipative flows (charge and heat) in Eq. (28) and (29) with the macroscopic expressions:

$$J^\mu = a_{11} \tilde{\mathcal{E}}^\mu + a_{12} \nabla^\mu T$$

and,  $q^\mu = a_{21} \tilde{\mathcal{E}}^\mu + a_{22} \nabla^\mu T.$

In this paper, we will focus only on the diagonal components of the  $\mathbf{a}$ -matrix since the electrical conductivity  $\sigma$  and thermal conductivity  $\kappa$  are identified with the diagonal elements, i.e.,  $\sigma = a_{11}$  and  $\kappa = a_{22}$ . The off-diagonal element  $a_{12}$  gives rise to the electric current due to the spatial variation in the temperature  $T$ . Similarly, the element  $a_{21}$  gives rise to the heat current due to the presence of an electric field  $\tilde{E}^\mu$ . The electrical and thermal conductivity of graphene is given by,

$$\sigma = 4\pi\tau_c e^2 \left(\frac{k_B T}{h}\right)^2 \frac{n}{\mathcal{E} + P} \left[ 2(f_2(A^{-1}) - f_2(A)) + \frac{\mathcal{E} + P}{nk_B T} (f_1(A^{-1}) + f_1(A)) \right], \quad (30)$$

$$\kappa = 4\pi\tau_c k_B \left(\frac{k_B T}{h}\right)^2 \frac{\mathcal{E} + P}{nk_B T} \left[ 2(f_2(A^{-1}) - f_2(A)) + \frac{\mathcal{E} + P}{nk_B T} (f_1(A^{-1}) + f_1(A)) \right], \quad (31)$$

The  $L$  is given by,

$$L = \frac{\kappa}{\sigma T} = \left(\frac{\mathcal{E} + P}{nk_B T}\right)^2 \frac{k_B^2}{e^2} = \left(\frac{\hbar}{k_B T}\right)^2 \frac{k_B^2}{e^2}. \quad (32)$$

Now, we will move to the next section to address the transport coefficients of the QGP.

## B. Thermoelectric transport in QGP

The inherent properties of the quantum chromo dynamics (QCD) make two distinct phases of the quark-matter possible: confined quark-matter as we observed in hadrons and deconfined quark matter which is supposed to be present in the early stages of universe and the core of neutron stars owing to the infrared slavery and asymptotic freedom of the QCD interactions respectively [91–93]. The deconfined stage of the quark matter known as QGP can be created in the HIC experiments in RHIC and LHC. In the early stage of HIC collision experiments, the QGP formed can be effectively modeled by relativistic fluid dynamics with the transport coefficients obtained from BTE [51]. For simplicity we will take a single flavored quark-anti quark ( $q, \bar{q}$ ) system and write down the BTE as [82, 83],

$$p^\mu \partial_\mu f_{q,\bar{q}} - Q_{q,\bar{q}} F^{\mu\nu} p_\nu \frac{\partial f_{q,\bar{q}}}{\partial p^\mu} = -(u_\alpha p^\alpha) \frac{f_{q,\bar{q}} - f_{q,\bar{q}}^0}{\tau}, \quad (33)$$

where for notational convenience we ignored subscripts on momentum variables.  $F^{\mu\nu}$  and  $\tau$  are the electromagnetic Faraday tensor and average collision time, respectively. Here, we will identify the quark-antiquark system with up-quark and anti-up-quark, i.e.,  $(q, \bar{q}) = (u, \bar{u})$ .  $f_q^0 = 1/(e^{(u_\mu p_q^\mu - \mu_q)/k_B T} + 1)$  and  $f_{\bar{q}}^0 = 1/(e^{(u_\mu p_{\bar{q}}^\mu + \mu_q)/k_B T} + 1)$  are the local equilibrium distribution functions for the quarks and anti-quarks, where  $u^\mu$  is the fluid four-velocity and  $\mu_q$  is the chemical potential of quark. The charge  $Q_q = Q_u = \frac{2e}{3}$  and  $Q_{\bar{q}} = Q_{\bar{u}} = -\frac{2e}{3}$ . The derivation of thermoelectrical transport coefficients of this  $(u, \bar{u})$  plasma runs similar to Sec. II A. The expressions of thermodynamic variables of this system are also similar to Sec. II A. The main differences in the expressions of QGP and graphene arise because of the following two reasons. The universal speed  $c$  in QGP and fermi velocity  $v_F$  in graphene and the dimensionality of QGP (3D system) and graphene (2D system) are different. Keeping these two differences in mind, one can easily write down the expressions for QGP as follows:

$$\tilde{n} = 2 \left[ \int \frac{d^3 \vec{p}}{h^3} f_q^0 - \int \frac{d^3 \vec{p}}{h^3} f_{\bar{q}}^0 \right] = 16\pi \left(\frac{k_B T}{hc}\right)^3 (f_3(A) - f_3(A^{-1})), \quad (A \equiv e^{\mu_q/k_B T}), \quad (34)$$

$$\tilde{\mathcal{E}} = 3P = 2 \left[ \int \frac{d^3 \vec{p}}{h^3} E f_q^0 + \int \frac{d^3 \vec{p}}{h^3} E f_{\bar{q}}^0 \right] = 48\pi \frac{(k_B T)^4}{(hc)^3} (f_4(A) + f_4(A^{-1})), \quad (35)$$

$$\tilde{\sigma} = \frac{16\pi\tau Q_u^2}{3c} \left(\frac{k_B T}{h}\right)^3 \frac{\tilde{n}}{\tilde{\mathcal{E}} + \tilde{P}} \left[ 3(f_3(A^{-1}) - f_3(A)) + \frac{\tilde{\mathcal{E}} + \tilde{P}}{\tilde{n}k_B T} (f_2(A) + f_2(A^{-1})) \right], \quad (36)$$

$$\tilde{\kappa} = \frac{16\pi\tau k_B}{3c} \left(\frac{k_B T}{h}\right)^3 \frac{\tilde{\mathcal{E}} + \tilde{P}}{\tilde{n}k_B T} \left[ 3(f_3(A^{-1}) - f_3(A)) + \frac{\tilde{\mathcal{E}} + \tilde{P}}{\tilde{n}k_B T} (f_2(A) + f_2(A^{-1})) \right], \quad (37)$$

$$\tilde{L} = \frac{\tilde{\kappa}}{\tilde{\sigma} T} = \left(\frac{\tilde{\mathcal{E}} + \tilde{P}}{\tilde{n}k_B T}\right)^2 \frac{k_B^2}{Q_u^2} = \left(\frac{\tilde{\hbar}}{k_B T}\right)^2 \frac{k_B^2}{Q_u^2}, \quad (38)$$

where for the ease of the presentation, we used the same notations as that of Sec. II A but with a tilde to distinguish QGP variables from graphene variables.




FIG. 1: Representation of different physical systems starting from CMP system, graphene, and metals to HEP system, QGP and Neutron star in  $\mu - T$  plane

### III. RESULTS AND DISCUSSION

Let us first qualitatively discuss the Condensed Matter Physics (CMP) and High Energy Physics (HEP) systems in  $\mu - T$  plane to get acquainted with the order of magnitude of the  $\mu$  and  $k_B T$  values in different physical systems like graphene, metals, QGP and Neutron stars. For this purpose, we have first displayed the domain of CMP and HEP in the  $\mu - T$  plane in Fig. 1. We have separated the whole plane in two parts by the line  $\mu/k_B T = 1$ , where the regions,  $\mu/k_B T < 1$  and  $\mu/k_B T > 1$  may be identified with the DF and FL region, respectively. From the figure, it is apparent that the chemical potential  $\mu$  for CMP lies within 0 – 10 eV. For a typical graphene sample, it varies in the meV scale, whereas for metals, it can be 2 – 10 eV. Similarly, the temperature scale for the CMP systems is of the few eVs, whereas in HEP, it is of the order of a few MeVs. Now, coming to the FL and DF behavior of the electrons, it can be seen in Fig. 1 that the electrons in metals show FL behavior, whereas the electrons in the graphene sheet can have both DF and FL behavior. Similarly, for HEP-HIC systems, we may assume the existence of DF and FL regimes depending on the energy of the collisions. In very high-energy HICs with the center of energy of the colliding nuclei of the order of TeV, the QGP formed is a DF comprising ultrarelativistic quarks. On the other hand, the HIC energies of the CBM and NICA, which try to reproduce the Neutron star environment, produce quark matter which are in the FL region.

Now let us discuss the quantitative magnitude of the thermodynamic and transport variables of the graphene and QGP with the help of the expressions obtained in Sec. II A and II B in terms of fermi integral functions. Here, we will show their variation with respect to  $\mu/k_B T$  and compare the numerical magnitudes of corresponding graphene and QGP variables. For a gated graphene sample, the experimental procedure of increasing the gate voltage with positive polarity at a constant device temperature can be effectively mapped by showing the variation of different sample characteristics, i.e., carrier density, energy density, conductivity, etc, by changing  $\mu/k_B T$  along  $X$ -axis keeping  $T$  constant. Similarly, the thermodynamic characteristics of the QGP produced in HIC experiments as one moves from the high energy collisions of LHC and RHIC to CBM and NICA can be mapped by changing  $\mu_q/k_B T$  along the horizontal axis, but we have to do this by taking the trajectory of constant  $T$ , varying  $\mu$  in the accessible  $\mu - T$  domain of those HIC experiment setups (One can also visualize the same in terms of baryon chemical potential  $\mu_B$  and baryon density  $n_B$  by using the link,  $\mu_q = \mu_B/3$ , and  $\tilde{n} = 3n_B$ ). At first, the variation of the surface density of electrons  $n_e$ , holes  $n_h$  and net carriers  $n = n_e - n_h$  is shown in the left panel of Fig. 2 at a fixed temperature  $T = 60$  K. For this purpose, Eq. (11) can be used to get the numerical magnitudes of  $n_e$ ,  $n_h$  and  $n$ . The result shows that




FIG. 2: (a)The number density (b)energy density and pressure of graphene with respect to  $\mu/k_B T$  at  $T=60$  K .




FIG. 3: (a)The number density (b)energy density and pressure of QGP with respect to  $\mu_q/k_B T$  for QGP at  $T = 10^{12}$  K ( $\equiv 100$  MeV) .

at a constant temperature the number density of electron  $n_e$  (blue line) rises with a rise in  $\mu/k_B T$  or equivalently  $\mu$ , whereas the number density of holes  $n_h$  (red line) falls exponentially with increase in  $\mu/k_B T$  in accordance with Eq. (59). The net carrier density  $n$  (green line) also shows an increasing trend as  $\mu$  increases. At  $\mu/k_B T \geq 3$ , the net density almost align with the total electron density  $n_e$  and  $n_h \rightarrow 0$ . This suggests that in gated graphene devices at  $T = 60$  K ( $k_B T \sim 5$  meV), for  $\mu > 15$  meV, the major charge carriers are electrons, and their density is of the order of  $10^{10}$   $\text{cm}^{-2}$ . Therefore, at  $T = 60$  K, we have an electron-hole DF for  $\mu \ll 15$  meV and an electron FL for  $\mu > 15$  meV. In the right panel of Fig. 2, we display the energy density  $\mathcal{E}$  and pressure  $P$  as a function of  $\mu/k_B T$  at  $T = 60$  K. We employ Eqs. (12) and (13) to get  $\mathcal{E}$  and  $P$  as a function of  $\mu/T$ . The total energy density (green line) increases monotonically with a change in  $\mu$ . At  $T = 60$  K, the estimated energy density of carriers lies in the range  $\sim 0.1$  to  $1$   $\text{GeV}/\text{cm}^2$  for  $\mu$  in the range  $5$  to  $25$  meV. The curve for pressure  $P$  (red line) follows the same qualitative trend as energy density  $\mathcal{E}$ , with its magnitude being exactly half of  $\mathcal{E}$  in agreement with Eq. (62). Now, let us shift our attention to see the changes in the thermodynamic variables of QGP by varying  $\mu/k_B T$  at a fixed temperature  $T \approx 10^{12}$  K or  $\approx 100$  MeV. The net quark density  $\tilde{n}$  (blue line) of QGP as a function of  $\mu_q/k_B T$  has been presented in the left panel of Fig. 3. We apply the formula (34) laid down in the Sec. IIB to get the numerical magnitude of  $\tilde{n}$  for different  $\mu_q/k_B T$ . The net quark density shows a monotonic rise with an increase in  $\mu_q$ . A straight line (red dot-dashed line) corresponds to nuclear saturation density  $n = 0.16$   $\text{fm}^{-3}$  has been drawn as a reference point. The results show that for  $\mu$  in the range  $0$  to  $500$  MeV, the net quark density lies below nuclear saturation, whereas, for  $\mu > 500$  MeV, the net quark density surpasses the nuclear saturation density. The right panel of Fig. 3 portrays the change in energy density  $\tilde{\mathcal{E}}$  (blue line) and pressure  $\tilde{P}$  (orange line) in relation to  $\mu/k_B T$  at a constant  $T = 10^{12}$  K. Both energy density and pressure increases monotonically with change in  $\mu$  consistent with the Eq. (35). The magnitude of energy density for QGP lies between  $0.01$  to  $0.21$   $\text{GeV}/\text{fm}^3$  for the range of  $\mu$  between  $0$  to  $500$  MeV.

In the left panel of Fig. 4, we show the variation of enthalpy per particle normalized by the chemical potential  $\frac{h}{\mu}$






FIG. 4: (a) Enthalpy density per particle vs  $\mu/k_B T$  (b) normalised  $\kappa$  and  $\sigma$  in graphene vs  $n$  .




FIG. 5: (a) Enthalpy density per particle vs  $\mu_q/k_B T$  (b) normalised  $\kappa$  and  $\sigma$  vs  $\tilde{n}$  for QGP at  $T = 10^{12}$  K ( $\equiv 100$  MeV) .

(blue line) with respect to  $\frac{\mu}{k_B T}$  at a fixed temperature  $T = 60$  K. Enthalpy per particle  $\mathfrak{h} = \frac{\mathcal{E}+P}{n}$  is an important thermodynamic parameter in the hydrodynamic regime of transport and the expression of electrical conductivity, thermal conductivity and LR contain the factor  $\mathfrak{h}$  in their formula. Analysing the  $\mathcal{E}$  and  $P$  curves in Fig. 2(b) and net electron density  $n$  in Fig. 2(a), one can notice that towards  $\frac{\mu}{k_B T} \rightarrow 0$  domain,  $\mathcal{E}$  and  $P$  saturate towards finite values but  $n$  tends to zero. This is the reason for  $\mathfrak{h}$  or  $\frac{\mathfrak{h}}{\mu}$  tends to diverge at  $\frac{\mu}{k_B T} \rightarrow 0$ . Since  $\mu$  can be identified with Gibb's free energy per particle, the vertical axis in the plot represents the ratio of enthalpy to Gibb's free energy. The plot displays a decreasing trend of  $\frac{\mathfrak{h}}{\mu}$  with  $\frac{\mu}{k_B T}$  and almost aligns with the line (red dash-dotted line) at unity after  $\mu > 30$  meV (or  $\frac{\mu}{k_B T} > 6$ ). This suggests that the enthalpy of the graphene significantly differs from Gibb's free energy in the DF region and almost matches with Gibb's free energy in the FL domain. In terms of Euler thermodynamic relation  $Ts = \mathcal{E} + P - \mu n$ , we can also understand the fact when we go towards  $\frac{\mu}{k_B T} \gg 1$  or large  $\mu$  and small  $T$ , we can write the thermodynamic relation as,

$$\begin{aligned} \mathcal{E} + P - \mu n &\approx 0 \\ \implies \frac{\mathcal{E} + P}{n} &= \mu \end{aligned}$$

This fact is reflected in Fig .4(a).

Next, the variation of normalized electrical conductivity ( $\sigma'$ ) and thermal conductivity ( $\kappa'$ ) of graphene with respect to  $n$  at constant  $T = 60$  K is presented in the right panel of Fig. 4. The dimensionless ( $\kappa'$ ) and ( $\sigma'$ ) are defined with




FIG. 6: (a) The dependence of  $\frac{L}{L_0}$  with net carrier density  $n$  in graphene at  $T=60$  K (b) The dependence of  $\frac{\tilde{L}}{L_0}$  with net carrier density  $\tilde{n}$  in QGP at  $T = 100$  MeV.

the help of Eq. (30) and (31) as follows:

$$\sigma' = \frac{\sigma h^2}{4e^2 \pi \tau_c k_B T} = \frac{2nk_B T}{\mathcal{E} + P} (f_2(A^{-1}) - f_2(A)) + (f_1(A^{-1}) + f_1(A)) \quad (39)$$

$$\kappa' = \frac{\kappa h^2}{4\pi \tau_c k_B^3 T^2} = \left( \frac{\mathcal{E} + P}{nk_B T} \right)^2 \left[ \frac{2nk_B T}{\mathcal{E} + P} (f_2(A^{-1}) - f_2(A)) + (f_1(A^{-1}) + f_1(A)) \right] \quad (40)$$

The magnitude of  $\kappa'$  (red line) displayed in the right panel of Fig. 4 shows interesting behavior as one changes the net carrier density  $n$ . As one moves from right to left along the horizontal axis in the FL domain ( $\frac{\mu}{k_B T} \gg 1$ ) it first decreases and then increases significantly in the DF regime ( $\frac{\mu}{k_B T} \ll 1$ ) with  $\kappa' \rightarrow \infty$  as  $\mu \rightarrow 0$ . This singular behavior can be understood from Eq. (40), where the factor  $\frac{\mathcal{E} + P}{nk_B T}$  which diverges as one approaches  $n \rightarrow 0$ . We also plot the normalized electrical conductivity (green line). It can be observed that in the same range of  $n$ ,  $\sigma'$  is almost constant, and its magnitude lies way below  $\kappa'$ . In Fig. 5 we show the variation of  $\frac{\tilde{\mu}}{\mu}$  (blue line) with change in  $\mu_q/k_B T$  at a fixed  $T = 10^{12}$  K in the left panel. Similar to the graphene case, in the domain of  $\mu/k_B T \ll 1$ ,  $\frac{\tilde{\mu}}{\mu}$  of quark matter significantly differs from one (shown in red dash-dotted line). Similar to the plots of normalized electrical and thermal conductivity of graphene, we also portray the normalized  $\tilde{\kappa}'$  and  $\tilde{\sigma}'$  for QGP in the right panel of Fig. 5 with respect to  $\mu_q/k_B T$  by using the formula:

$$\tilde{\sigma}' = \frac{3h^3 c \tilde{\sigma}}{16Q_u^2 \pi \tau k_B^2 T^2} = \frac{3\tilde{n} k_B T}{\tilde{\mathcal{E}} + \tilde{P}} (f_3(A^{-1}) - f_3(A)) + (f_2(A^{-1}) + f_2(A)) \quad (41)$$

$$\tilde{\kappa}' = \frac{3h^3 c \tilde{\kappa}}{16\pi \tau k_B^4 T^3} = \left( \frac{\tilde{\mathcal{E}} + \tilde{P}}{\tilde{n} k_B T} \right)^2 \left[ \frac{3\tilde{n} k_B T}{\tilde{\mathcal{E}} + \tilde{P}} (f_3(A^{-1}) - f_3(A)) + (f_2(A^{-1}) + f_2(A)) \right]. \quad (42)$$

The qualitative trends of both  $\tilde{\kappa}'$  and  $\tilde{\sigma}'$  are same as that of the graphene with  $\tilde{\kappa}'$  being significantly higher than  $\tilde{\sigma}'$  at low net quark densities.

Finally in Fig. 6, we depicted the normalized LR  $\frac{L}{L_0}$  of graphene (in the left panel) as a function of net carrier density  $n$  at  $T = 60$  K and  $\frac{\tilde{L}}{L_0}$  of QGP (in the right panel) as a function of net quark density  $\tilde{n}$  at  $T = 10^{12}$  K. In the left panel of Fig. 6 we also present the experimental data of Ref. [63] corresponds to  $S_1$  (green dots) and  $S_2$  (red dots), where  $S_1$  is more cleaner graphene sample than  $S_2$ . We divide the graph into three regions: fluidic, mixed, and ohmic. The fluidic region corresponds to the DF regime where  $\mu \ll k_B T$  and hydrodynamics is applicable here. In contrast, the ohmic region corresponds to the FL domain of  $\mu \gg k_B T$  where carrier transport is diffusive, and hydrodynamics is not applicable. Between the fluidic and ohmic regions, a domain exists where carrier dynamics are neither completely hydrodynamic nor diffusive; this region is named mixed. It can be observed that the experimental data corresponds to  $S_1$  and  $S_2$  lie far away from the line  $L/L_0 = 1$  (black dashed) in the domain where  $n < 10^{10}$   $\text{cm}^{-2}$  and almost align with it when  $n > 2 \times 10^{10}$   $\text{cm}^{-2}$ . We can see that LR obtained with the help of Eq. (32) of Sec. II A show a similar kind of divergence tendency as experimentally observed by Crossno et al. [63]. We notice that experimental data of  $S_1$  and  $S_2$  both have divergence tendencies in the fluidic domain, and our fluid-base theoretical

curve is in good agreement with  $S_1$  data. These  $S_1$  and  $S_2$  correspond to different levels of cleanness of experimental samples. Since our present theory does not carry such quantitative inputs, so instead of quantitative matching of experimental data, we should highlight the qualitative divergence tendency in the fluid domain, noticed in both theory and experiment.

Interestingly, we have identified three gross domain of Fig. 6(a) along  $\frac{\mu}{k_B T}$  - axis or  $n$  - axis at  $T = 60$  K. They are (1) fluidic domain ( $n < 8 \times 10^9 \text{cm}^{-2}$ ), (2) mixed domain ( $8 \times 10^9 \text{cm}^{-2} < n < 1.5 \times 10^{10} \text{cm}^{-2}$ ) and (3) non fluidic or Ohmic domain ( $n > 1.5 \times 10^{10} \text{cm}^{-2}$ ). Large net carrier density ( $n > 1.5 \times 10^{10} \text{cm}^{-2}$ ) or large  $\frac{\mu}{k_B T}$  (*i.e.*  $\frac{\mu}{k_B T} \gg 1$ ), we will get our traditional Ohmic domain, where the Lorenz ratio become (universal) constant and equal to Lorenz number  $L_0$ . Metals as 3-dimensional non-relativistic matter [36] follow this constant Lorenz ratio rule, known as WF law because their Fermi energy remains within  $\mu = 2 - 10$  eV, which is located in Ohmic domain ( $\frac{\mu}{k_B T} \gg 1$ ). It is 2-dimensional graphene system, where Fermi energy  $\mu$  or net charge carrier density can be reduced from  $\frac{\mu}{k_B T} \gg 1$  to  $\frac{\mu}{k_B T} \ll 1$  domains. Recent experimental measurements [48, 55–64] suggest that ( $\frac{\mu}{k_B T} \ll 1$ ) domain show fluid property (popularly called DF). So, we used fluid-based LR expression for the fluidic domain ( $n < 8 \times 10^9 \text{cm}^{-2}$ ), and we get an increasing trend as we decrease the  $n$ . Experimental data in  $n < 8 \times 10^9 \text{cm}^{-2}$  and  $n > 1.5 \times 10^9 \text{cm}^{-2}$  follow respectively fluid-based and non-fluid/ Ohmic based theories. It indicates a mixed phenomenon or a transition from non-fluid to fluid phenomena within  $8 \times 10^9 \text{cm}^{-2} < n < 1.5 \times 10^{10} \text{cm}^{-2}$  domain. One can use switching functions [36] to show a smooth transition from the non-fluid to the fluid domain, but digging into the actual mechanism of transition may be considered interesting for future research.

Readers can go through earlier Refs. [39, 41, 42], which are focused on explaining this WF law violation data [63]. In Ref. [42], both the band gap and the case without a band gap in graphene are considered, along with bipolar diffusion. Additionally, Ref. [41] employs thermodynamic aspects through the poly-logarithmic function to address the enhancement of LR near the Dirac point. Among them, Ref. [39] has provided a more quantitative matching of WF law violation data, and they have grossly marked that a fluid aspect is a possible reason for the violation. In this regard, we are also pointing out similar physics qualitatively instead of quantitatively matching exactly Crossno et al.'s data. The present work is intended to explore similar hydrodynamical structures in graphene and QGP systems in relation to WF law violation.

In the QGP system, Refs. [43–47] have found a similar kind of WF law violation at  $\mu = 0$  region, expected to be produced in RHIC or LHC experiments. Fig. 6(b) demonstrates  $\frac{L}{L_0}$  of massless quark matter, which also supports the WF law violation due to the fluid aspect of quark matter. At high density  $n > n_0 = 0.16 \text{fm}^{-3}$  ( $2n_0 - 4n_0$ ), which is expected to be produced in CBM or NICA experiments and expected in neutron star environment, one may explore the  $\frac{L}{L_0}$  calculations, which is poorly known till now. So, future experimental data from CBM and NICA may play an important role in deciding the requirement of a hydro-dynamical or non-fluid framework for dense sector quark or hadronic matter. Our simple massless quark matter calculation suggests that if high-density QGP carries a fluid aspect, it will still violate WF law. However, a systemic review of earlier research and future research may be required to conclude something on WF law in high-density QGP systems.

#### IV. SUMMARY AND CONCLUSIONS

In summary, we have explored the thermoelectric transport in graphene and its equivalence with quark-gluon plasma both qualitatively and quantitatively. To obtain the thermoelectric transport coefficients in graphene in the Dirac fluid regime, we have used a covariant Boltzmann transport equation with the light speed  $c$  replaced by Fermi velocity  $v_F$ . The electric field and thermal gradients in the graphene sample give rise to dissipative flows like electrical and thermal flows, which have been evaluated with the help of the Boltzmann transport equation. We observe that the enthalpy per particle in graphene plays a decisive role in determining the numerical magnitude of the thermal and electrical conductivity. The enthalpy per particle significantly differs from the chemical potential at the Dirac point and affects the charge transport in the DF regime. Our calculation suggests a blowing up of the Lorenz ratio at low carrier density, which is consistent with the recent experimental measurements. The qualitative pattern obtained for the thermodynamic variables and transport coefficients for graphene fluid and quark-gluon plasma is similar. Two factors that make the difference in the quantitative magnitude of the thermodynamic and transport coefficients of quark-gluon plasma and graphene fluid are the dimensionality (three for quark-gluon plasma and two for graphene fluid) and the universal speed ( $c$  for quark-gluon plasma and  $v_F$  for graphene fluid). Now, when we follow the experimental data of the Lorenz ratio for the graphene system, we notice that low and high-density data prefer fluid and non-fluid theoretical calculation of the Lorenz ratio. Since these two theoretical estimations do not merge in the intermediate density, so a mixed phenomenon or transition from non-fluid to fluid phenomena is expected here.

Similar to the graphene case, we can get indirect experimental data of the Lorenz ratio for the quark-gluon plasma at low and high density. Relativistic heavy ion collider and Large hadron collider experiments already indicated the

fluid aspect of quark-gluon plasma at low density. So, the Wiedemann-Franz law violation can also be expected here, which is also confirmed by some recent references. Future experiments at Compressed Baryonic matter and Nuclotron-based Ion Collider fAcility (NICA) may provide a better knowledge of quark matter created at high density from which the correct framework to determine Lorentz ratio and validity or violation of Wiedemann-Franz law and can be inferred for the dense quark-gluon plasma.

## V. ACKNOWLEDGEMENT

SN and AD gratefully acknowledge the Ministry of Education (MoE), Govt. of India. The authors extend their thanks to the other members of the electron hydrodynamics club - Cho Win Aung, Thandar Zaw Win, and Narayan Prasad.

## VI. APPENDIX

### A. Calculation of $\delta f_{e,h}$ as sum of bulk, shear and thermoelectrical parts

In this appendix we will briefly show the transition from Eq. (18) and (19) to Eq. (23) and (24) via using the conservation Eqs. (20) to (22). The net number density  $n$  and total energy density  $\mathcal{E}$  being function of  $\frac{\mu}{k_B T}$  and  $\frac{1}{k_B T}$  can be differentiated with the help of operator  $D$  as follows:

$$Dn = \frac{\partial n}{\partial \left(\frac{\mu}{k_B T}\right)} D \frac{\mu}{k_B T} + \frac{\partial n}{\partial \left(\frac{1}{k_B T}\right)} D \frac{1}{k_B T}, \quad (43)$$

$$D\mathcal{E} = \frac{\partial \mathcal{E}}{\partial \left(\frac{\mu}{k_B T}\right)} D \frac{\mu}{k_B T} + \frac{\partial \mathcal{E}}{\partial \left(\frac{1}{k_B T}\right)} D \frac{1}{k_B T}. \quad (44)$$

Now using the conservation laws given by the Eq. (20) and Eq. (22) in the Eqs. (43) and (44) respectively we have,

$$\frac{\partial n}{\partial \left(\frac{\mu}{k_B T}\right)} D \frac{\mu}{k_B T} + \frac{\partial n}{\partial \left(\frac{1}{k_B T}\right)} D \frac{1}{k_B T} = -n \nabla_\mu u^\mu, \quad (45)$$

$$\frac{\partial \mathcal{E}}{\partial \left(\frac{\mu}{k_B T}\right)} D \frac{\mu}{k_B T} + \frac{\partial \mathcal{E}}{\partial \left(\frac{1}{k_B T}\right)} D \frac{1}{k_B T} = -(\mathcal{E} + P) \nabla_\mu u^\mu. \quad (46)$$

The Eq. (45) and (46) are two linear equation and can be easily solved for  $D \frac{1}{k_B T}$  and  $D \frac{\mu}{k_B T}$ . The solution of these equations are of the form:  $D \frac{1}{k_B T} = x_1 \nabla_\mu u^\mu$  and  $D \frac{\mu}{k_B T} = x_2 \nabla_\mu u^\mu$ . The quantities  $x_1$  and  $x_2$  are functions of thermodynamical variables, and their exact expressions are irrelevant to our purpose. Now we are ready to eliminate all the temporal derivative terms from Eq. (18) and Eq. (19) by substituting  $D \frac{1}{k_B T} = x_1 \nabla_\mu u^\mu$ ,  $D \frac{\mu}{k_B T} = x_2 \nabla_\mu u^\mu$ , and Eq. (21). Rewriting Eq. (18) in terms of spatial derivatives we have,

$$\begin{aligned} \delta f_e = & \frac{\tau_c v_F^2}{u_\mu p_e^\mu} \left[ v_F^{-2} (x_1 (u_\alpha p_e^\alpha)^2 - x_2 (u_\alpha p_e^\alpha)) \nabla_\beta u^\beta + \frac{p_e^\alpha p_e^\beta}{k_B T} \nabla_\alpha u_\beta + \frac{(u_\beta p_e^\beta) p_e^\alpha}{k_B T} \frac{\nabla_\alpha P}{\mathcal{E} + P} + \frac{(u_\beta p_e^\beta) p_e^\alpha}{k_B T} \frac{\rho}{\mathcal{E} + P} \tilde{E}_\alpha \right. \\ & \left. + (u_\beta p_e^\beta) p_e^\alpha \nabla_\alpha \frac{1}{k_B T} - p_e^\alpha \nabla_\alpha \frac{\mu}{k_B T} + \frac{e p_e^\alpha}{k_B T} \tilde{E}_\alpha \right] f_e^0 (1 - f_e^0). \end{aligned} \quad (47)$$

From the Gibbs-Duhem relation, we have,

$$\begin{aligned} n d \frac{\mu}{k_B T} &= \frac{1}{k_B T} dP + (\mathcal{E} + P) d \frac{1}{k_B T} \\ \implies n \nabla^\alpha \frac{\mu}{k_B T} &= \frac{1}{k_B T} \nabla^\alpha P + (\mathcal{E} + P) \nabla^\alpha \frac{1}{k_B T} \\ \implies \frac{(u_\beta p_e^\beta) p_e^\alpha}{k_B T} \frac{\nabla_\alpha P}{\mathcal{E} + P} &= (u_\beta p_e^\beta) p_e^\alpha \left[ \frac{n}{\mathcal{E} + P} \nabla_\alpha \frac{\mu}{k_B T} - \nabla_\alpha \frac{1}{k_B T} \right]. \end{aligned} \quad (48)$$

Using Eq. (48) in Eq. (47) we have,

$$\begin{aligned} \delta f_e &= \frac{\tau_c v_F^2}{u_\mu p_e^\mu} \left[ v_F^{-2} (x_1 (u_\alpha p_e^\alpha)^2 - x_2 (u_\alpha p_e^\alpha)) \nabla_\beta u^\beta + \frac{p_e^\alpha p_e^\beta}{k_B T} \nabla_\alpha u_\beta \right. \\ &\quad \left. - p_e^\alpha \left[ \frac{n}{\mathcal{E} + P} u_\beta p_e^\beta - 1 \right] \left( -\nabla_\alpha \frac{\mu}{k_B T} + \frac{e \tilde{E}_\alpha}{k_B T} \right) \right] f_e^0 (1 - f_e^0). \end{aligned} \quad (49)$$

Rewriting Eq. (49) using the result:  $p_e^\alpha p_e^\beta \nabla_\alpha u_\beta = p_e^\alpha p_e^\beta \sigma_{\alpha\beta} + \frac{1}{2} \Delta_{\alpha\nu} p_e^\alpha p_e^\nu (\nabla_\beta u^\beta)$ , where  $\sigma_{\alpha\beta} \equiv \frac{1}{2} (\nabla_\alpha u_\beta + \nabla_\beta u_\alpha) - \frac{1}{2} \Delta_{\alpha\beta} (\nabla_\mu u^\mu)$ , we have,

$$\begin{aligned} \delta f_e &= \frac{\tau_c v_F^2}{u_\mu p_e^\mu} \left[ v_F^{-2} (x_1 (u_\alpha p_e^\alpha)^2 - x_2 (u_\alpha p_e^\alpha)) + \frac{1}{2k_B T} \Delta_{\alpha\nu} p_e^\alpha p_e^\nu \nabla_\beta u^\beta + \frac{p_e^\alpha p_e^\beta}{k_B T} \sigma_{\alpha\beta} \right. \\ &\quad \left. - p_e^\alpha \left[ \frac{n}{\mathcal{E} + P} u_\beta p_e^\beta - 1 \right] \left( -\nabla_\alpha \frac{\mu}{k_B T} + \frac{e \tilde{E}_\alpha}{k_B T} \right) \right] f_e^0 (1 - f_e^0). \end{aligned} \quad (50)$$

Using similar steps one easily finds the out-of-equilibrium hole distribution  $\delta f_h$  in terms of spatial gradients as:

$$\begin{aligned} \delta f_h &= \frac{\tau_c v_F^2}{u_\mu p_h^\mu} \left[ v_F^{-2} (x_1 (u_\alpha p_h^\alpha)^2 - x_2 (u_\alpha p_h^\alpha)) + \frac{1}{2k_B T} \Delta_{\alpha\nu} p_h^\alpha p_h^\nu \nabla_\beta u^\beta + \frac{p_h^\alpha p_h^\beta}{k_B T} \sigma_{\alpha\beta} \right. \\ &\quad \left. - p_h^\alpha \left[ \frac{n}{\mathcal{E} + P} u_\beta p_h^\beta + 1 \right] \left( -\nabla_\alpha \frac{\mu}{k_B T} + \frac{e \tilde{E}_\alpha}{k_B T} \right) \right] f_h^0 (1 - f_h^0). \end{aligned} \quad (51)$$

In Eq. (50) and (51), the terms proportional to  $\nabla_\beta u^\beta$  and  $\sigma_{\alpha\beta}$  give rise to bulk and shear viscosity in the graphene. One can readily verify that upon ignoring the terms that correspond to the viscosity in the graphene, the out-of-equilibrium distribution functions for electron and hole matches with Eq. (23) and Eq. (24).

## B. Evaluation of thermodynamic variables and current density in terms of fermi integral functions

Here, we will define the fermi integral functions for electrons and holes used in the main text and briefly describe their properties. Finally, we will express the thermodynamic variables in terms of these integrals. Let us define the fermi integrals for electrons as:

$$f_j(A) = \frac{1}{\Gamma(j)} \int_0^\infty \frac{x^{j-1} dx}{e^x A^{-1} + 1}, \quad (A \equiv e^{\mu/k_B T}). \quad (52)$$

In the domain  $j > 0$  and  $-\infty < \frac{\mu}{k_B T} \leq 0$ , one can represent  $f_j(A)$  by the following series,

$$f_j(A) = A - \frac{A^2}{2j} + \frac{A^3}{3j} - \dots = e^{\mu/k_B T} - \frac{e^{2\mu/k_B T}}{2j} + \frac{e^{3\mu/k_B T}}{3j} - \dots \quad (53)$$

For the region  $j > 0$  and  $\frac{\mu}{k_B T} \gg 1$ , one can express  $f_j(A)$  in the form of sommerfeld's series as follows:

$$f_j(A) = \frac{1}{\Gamma(j+1)} \left( \frac{\mu}{k_B T} \right)^j + \frac{1}{\Gamma(j-1)} \frac{\pi^2}{6} \left( \frac{\mu}{k_B T} \right)^{j-2} + \dots \quad (54)$$

The fermi integrals for holes can be defined in a similar manner as follows:

$$f_j(A^{-1}) = \frac{1}{\Gamma(j)} \int_0^\infty \frac{x^{j-1} dx}{e^x A + 1}, \quad (A \equiv e^{\mu/k_B T}). \quad (55)$$

Different limiting expressions for the function  $f_j(A^{-1})$  can be obtained from  $f_j(A)$  by the replacement  $\mu \rightarrow -\mu$ . We will put down these expressions explicitly,

$$f_j(A^{-1}) = A^{-1} - \frac{A^{-2}}{2j} + \frac{A^{-3}}{3j} - \dots = e^{-\mu/k_B T} - \frac{e^{-2\mu/k_B T}}{2j} + \frac{e^{-3\mu/k_B T}}{3j} - \dots \quad (56)$$

where  $j > 0$  and  $0 \leq \frac{\mu}{k_B T} < \infty$ . Similarly, the Sommerfeld's series expression for the hole distribution is,

$$f_j(A^{-1}) = \frac{1}{\Gamma(j+1)} \left( \frac{-\mu}{k_B T} \right)^j + \frac{1}{\Gamma(j-1)} \frac{\pi^2}{6} \left( \frac{-\mu}{k_B T} \right)^{j-2} + \dots, \quad (57)$$

where  $j > 0$  and  $\frac{-\mu}{k_B T} \gg 1$ .

Now, we will move on to express the thermodynamic variables in terms of fermi integrals. The thermodynamic integral for  $n$  specified in Eqs. (11) can be simplified as,

$$\begin{aligned} n &= 4 \left[ \int \frac{d^2 \vec{p}_e}{h^2} f_e^0 - \int \frac{d^2 \vec{p}_h}{h^2} f_h^0 \right] \\ \Rightarrow n &= 4 \left[ \int_0^\infty \frac{d^2 \vec{p}_e}{h^2} \frac{1}{e^{(E_e - \mu)/k_B T} + 1} - \int_0^\infty \frac{d^2 \vec{p}_h}{h^2} \frac{1}{e^{(E_h + \mu)/k_B T} + 1} \right] \\ \Rightarrow n &= \frac{8\pi}{h^2} \left( \frac{k_B T}{v_F} \right)^2 \left[ \frac{1}{\Gamma(2)} \int_0^\infty \frac{x^{2-1} dx}{e^x A^{-1} + 1} - \frac{1}{\Gamma(2)} \int_0^\infty \frac{x^{2-1} dx}{e^x A + 1} \right], \text{ (where } x \equiv \frac{p_e v_F}{k_B T} \text{)} \\ \Rightarrow n &= \frac{8\pi}{h^2} \left( \frac{k_B T}{v_F} \right)^2 (f_2(A) - f_2(A^{-1})) . \end{aligned} \quad (58)$$

It can be easily seen from Eq. (58) that in the limit  $\frac{\mu}{k_B T} \rightarrow 0$  we have  $n = 0$  and for  $\frac{\mu}{k_B T} \gg 1$  we get,

$$n = \frac{8\pi\mu^2}{(hv_F)^2} \left[ \frac{1}{6} + \frac{\pi^2}{6} \left( \frac{k_B T}{\mu} \right)^2 - \left( \frac{k_B T}{\mu} \right)^2 e^{-\mu/k_B T} + \frac{1}{4} \left( \frac{k_B T}{\mu} \right)^2 e^{-2\mu/k_B T} \right]. \quad (59)$$

A quick look at Eq. (59) suggests that for  $\mu \gg k_B T$ , one can safely ignore the hole contribution (3rd and 4th term in the square bracket) to the net number density. We can also simplify the energy density  $\mathcal{E}$  given in Eq. (12) as,

$$\begin{aligned} \mathcal{E} &= 4 \left[ \int \frac{d^2 \vec{p}_e}{h^2} E_e f_e^0 + \int \frac{d^2 \vec{p}_h}{h^2} E_h f_h^0 \right] \\ \Rightarrow \mathcal{E} &= 4 \left[ \int_0^\infty \frac{d^2 \vec{p}_e}{h^2} \frac{E_e}{e^{(E_e - \mu)/k_B T} + 1} + \int_0^\infty \frac{d^2 \vec{p}_h}{h^2} \frac{E_h}{e^{(E_h + \mu)/k_B T} + 1} \right] \\ \Rightarrow \mathcal{E} &= \frac{16\pi}{h^2} \frac{(k_B T)^3}{v_F^2} \left[ \frac{1}{\Gamma(3)} \int_0^\infty \frac{x^{2-1} dx}{e^x A^{-1} + 1} + \frac{1}{\Gamma(3)} \int_0^\infty \frac{x^{2-1} dx}{e^x A + 1} \right], \text{ (where } x \equiv \frac{p_e v_F}{k_B T} \text{)} \\ \Rightarrow \mathcal{E} &= \frac{16\pi}{h^2} \frac{(k_B T)^3}{v_F^2} (f_3(A) + f_3(A^{-1})) . \end{aligned} \quad (60)$$

In the limit  $\frac{\mu}{k_B T} \rightarrow 0$  we have  $\mathcal{E} = \frac{32\pi}{h^2} \frac{(k_B T)^3}{v_F^2} \eta(3)$ , where the Dirichlet eta function  $\eta(j) = f_j(1) = \sum_{n=1}^\infty \frac{(-1)^{n-1}}{n^j}$  and in the limit  $\frac{\mu}{k_B T} \gg 1$  we have,

$$\mathcal{E} = \frac{16\pi}{(hv_F)^2} \mu^3 \left[ \frac{1}{6} + \frac{\pi^2}{6} \left( \frac{k_B T}{\mu} \right)^2 + \left( \frac{k_B T}{\mu} \right)^3 e^{-\mu/k_B T} - \frac{1}{8} \left( \frac{k_B T}{\mu} \right)^3 e^{-2\mu/k_B T} \right]. \quad (61)$$

From Eq. (61), it can be easily inferred that in the parameter range  $\mu \gg k_B T$ , the hole contribution to the energy density is negligible. Similarly, pressure  $P$  can be expressed as,

$$\begin{aligned} P &= 4 \left[ \int \frac{d^2 \vec{p}_e}{h^2} \frac{p_e^2}{2E_e} f_e^0 + \int \frac{d^2 \vec{p}_h}{h^2} \frac{p_h^2}{2E_h} f_h^0 \right] \\ \Rightarrow P &= 2 \left[ \int \frac{d^2 \vec{p}_e}{h^2} E_e f_e^0 + \int \frac{d^2 \vec{p}_h}{h^2} E_h f_h^0 \right] = \frac{\mathcal{E}}{2}, \end{aligned} \quad (62)$$

where we used  $p_{e,h} = \frac{E_{e,h}}{v_F}$ . Now, we will move on to give a step-wise derivation of the expression of current density provided in Eq. (26). By using the result of an integral of type  $\int d^2 \vec{p} G(|\vec{p}|) p^i p^j = \int d^2 p G(|\vec{p}|) \frac{p^2}{2} \delta^{ij}$ , ( $d^2 p \equiv 2\pi p dp$ )

where  $G(|\vec{p}|)$  is any arbitrary function, we can rewrite Eq. (25) as follows:

$$\begin{aligned}
J^i &= -4e v_F^4 \tau_c \left[ \int \frac{d^2 p_e}{h^2} \frac{p_e^2}{2E_e^2} \delta^{ij} \left( \frac{nE_e}{\mathcal{E} + P} - 1 \right) f_e^0(1 - f_e^0) - \int \frac{d^2 p_h}{h^2} \frac{p_h^2}{2E_h^2} \delta^{ij} \left( \frac{nE_h}{\mathcal{E} + P} + 1 \right) f_h^0(1 - f_h^0) \right] X^j, \\
&= -\frac{4\pi e \tau_c}{h^2} \left[ \int_0^\infty E_e \left( \frac{nE_e}{\mathcal{E} + P} - 1 \right) f_e^0(1 - f_e^0) dE_e - \int_0^\infty E_h \left( \frac{nE_h}{\mathcal{E} + P} + 1 \right) f_h^0(1 - f_h^0) dE_h \right] X^i \\
&= -\frac{4\pi e \tau_c}{h^2} \left[ \frac{n}{\mathcal{E} + P} \left( \int_0^\infty E_e^2 f_e^0(1 - f_e^0) dE_e - \int_0^\infty E_h^2 f_h^0(1 - f_h^0) dE_h \right) \right. \\
&\quad \left. - \left( \int_0^\infty E_e f_e^0(1 - f_e^0) dE_e + \int_0^\infty E_h f_h^0(1 - f_h^0) dE_h \right) \right] X^i. \tag{63}
\end{aligned}$$

We can easily evaluate the integrals occurring in Eq. (63) by using the definition of fermi integrals and the identity  $\frac{\partial f_j(A)}{\partial(\mu/k_B T)} = f_{j-1}(A)$  as,

$$\begin{aligned}
\int_0^\infty E_e^2 f_e^0(1 - f_e^0) dE_e &= \frac{\partial}{\partial(\mu/k_B T)} \int_0^\infty E_e^2 f_e^0 dE_e = 2(k_B T)^3 \frac{\partial f_3(A)}{\partial(\mu/k_B T)} = 2(k_B T)^3 f_2(A) \\
\text{and, } \int_0^\infty E_e f_e^0(1 - f_e^0) dE_e &= \frac{\partial}{\partial(\mu/k_B T)} \int_0^\infty E_e f_e^0 dE_e = (k_B T)^2 \frac{\partial f_2(A)}{\partial(\mu/k_B T)} = (k_B T)^2 f_1(A).
\end{aligned}$$

Similarly, the integrals corresponding to hole distributions can be obtained by just replacing  $\mu$  by  $-\mu$ . The final expression for Eq. (63) can be written as,

$$\begin{aligned}
J^i &= -\frac{4\pi e \tau_c}{h^2} \left[ \frac{n}{\mathcal{E} + P} 2(k_B T)^3 (f_2(A) - f_2(A^{-1}) - (k_B T)^2 (f_1(A) + f_1(A^{-1}))) \right] X^i \\
&= \frac{4\pi e \tau_c}{h^2} \frac{nk_B T}{\mathcal{E} + P} \left[ 2(k_B T)^2 (f_2(A^{-1}) - f_2(A) + \frac{\mathcal{E} + P}{n} k_B T (f_1(A) + f_1(A^{-1}))) \right] X^i,
\end{aligned}$$

which exactly matches with Eq. (26).

- 
- [1] K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, Y. Zhang, S. V. Dubonos, I. V. Grigorieva, and A. A. Firsov, Electric field effect in atomically thin carbon films, *Science* **306**, 666 (2004), <https://www.science.org/doi/pdf/10.1126/science.1102896>.
  - [2] C. Berger, Z. Song, T. Li, X. Li, A. Y. Ogbazghi, R. Feng, Z. Dai, A. N. Marchenkov, E. H. Conrad, P. N. First, and W. A. de Heer, Ultrathin epitaxial graphite: 2d electron gas properties and a route toward graphene-based nanoelectronics, *The Journal of Physical Chemistry B* **108**, 19912 (2004).
  - [3] K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, M. I. Katsnelson, I. V. Grigorieva, S. V. Dubonos, and A. A. Firsov, Two-dimensional gas of massless dirac fermions in graphene, *nature* **438**, 197 (2005).
  - [4] Y. Zhang, Y.-W. Tan, H. L. Stormer, and P. Kim, Experimental observation of the quantum hall effect and berry's phase in graphene, *nature* **438**, 201 (2005).
  - [5] Y.-W. Tan, Y. Zhang, K. Bolotin, Y. Zhao, S. Adam, E. H. Hwang, S. Das Sarma, H. L. Stormer, and P. Kim, Measurement of scattering rate and minimum conductivity in graphene, *Phys. Rev. Lett.* **99**, 246803 (2007).
  - [6] J.-H. Chen, M. Ishigami, C. Jang, D. Hines, M. Fuhrer, and E. Williams, Printed graphene circuits, *Advanced Materials* **19**, 3623 (2007), <https://onlinelibrary.wiley.com/doi/pdf/10.1002/adma.200701059>.
  - [7] C. Jang, S. Adam, J.-H. Chen, E. D. Williams, S. Das Sarma, and M. S. Fuhrer, Tuning the effective fine structure constant in graphene: Opposing effects of dielectric screening on short- and long-range potential scattering, *Phys. Rev. Lett.* **101**, 146805 (2008).
  - [8] J.-H. Chen, C. Jang, S. Adam, M. S. Fuhrer, E. D. Williams, and M. Ishigami, Charged-impurity scattering in graphene, *Nature Physics* **4**, 377–381 (2008).
  - [9] S. Cho and M. S. Fuhrer, Charge transport and inhomogeneity near the minimum conductivity point in graphene, *Phys. Rev. B* **77**, 081402 (2008).
  - [10] J. Martin, N. Akerman, G. Ulbricht, T. Lohmann, J. v. Smet, K. Von Klitzing, and A. Yacoby, Observation of electron-hole puddles in graphene using a scanning single-electron transistor, *Nature physics* **4**, 144 (2008).
  - [11] L. A. Ponomarenko, R. Yang, T. M. Mohiuddin, M. I. Katsnelson, K. S. Novoselov, S. V. Morozov, A. A. Zhukov, F. Schedin, E. W. Hill, and A. K. Geim, Effect of a high- $\kappa$  environment on charge carrier mobility in graphene, *Phys. Rev. Lett.* **102**, 206603 (2009).

- [12] M. Monteverde, C. Ojeda-Aristizabal, R. Weil, K. Bennaceur, M. Ferrier, S. Guéron, C. Glattli, H. Bouchiat, J. N. Fuchs, and D. L. Maslov, Transport and elastic scattering times as probes of the nature of impurity scattering in single-layer and bilayer graphene, *Phys. Rev. Lett.* **104**, 126801 (2010).
- [13] N. Shon and T. Ando, Quantum transport in two-dimensional graphite system, *Journal of the Physical Society of Japan* **67**, 2421 (1998), <https://doi.org/10.1143/JPSJ.67.2421>.
- [14] T. Ando, Screening effect and impurity scattering in monolayer graphene, *Journal of the Physical Society of Japan* **75**, 074716 (2006), <https://doi.org/10.1143/JPSJ.75.074716>.
- [15] K. Nomura and A. H. MacDonald, Quantum hall ferromagnetism in graphene, *Phys. Rev. Lett.* **96**, 256602 (2006).
- [16] K. Nomura and A. H. MacDonald, Quantum transport of massless dirac fermions, *Phys. Rev. Lett.* **98**, 076602 (2007).
- [17] S. Adam, E. Hwang, and S. Das Sarma, Scattering mechanisms and boltzmann transport in graphene, *Physica E: Low-dimensional Systems and Nanostructures* **40**, 1022 (2008).
- [18] N. M. R. Peres, F. Guinea, and A. H. Castro Neto, Electronic properties of disordered two-dimensional carbon, *Phys. Rev. B* **73**, 125411 (2006).
- [19] K. Ziegler, Robust transport properties in graphene, *Phys. Rev. Lett.* **97**, 266802 (2006).
- [20] P. M. Ostrovsky, I. V. Gornyi, and A. D. Mirlin, Electron transport in disordered graphene, *Phys. Rev. B* **74**, 235443 (2006).
- [21] M. Trushin and J. Schliemann, Minimum electrical and thermal conductivity of graphene: A quasiclassical approach, *Phys. Rev. Lett.* **99**, 216602 (2007).
- [22] S. Adam, E. Hwang, V. Galitski, and S. Das Sarma, A self-consistent theory for graphene transport, *Proceedings of the National Academy of Sciences* **104**, 18392 (2007).
- [23] E. H. Hwang, S. Adam, and S. D. Sarma, Carrier transport in two-dimensional graphene layers, *Phys. Rev. Lett.* **98**, 186806 (2007).
- [24] N. M. R. Peres, J. M. B. Lopes dos Santos, and T. Stauber, Phenomenological study of the electronic transport coefficients of graphene, *Phys. Rev. B* **76**, 073412 (2007).
- [25] D. S. Novikov, Numbers of donors and acceptors from transport measurements in graphene, *Applied Physics Letters* **91**, 102102 (2007), <https://pubs.aip.org/aip/apl/article-pdf/doi/10.1063/1.2779107/14374674/102102.1.online.pdf>.
- [26] M. I. Katsnelson and A. K. Geim, Electron scattering on microscopic corrugations in graphene, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* **366**, 195 (2008).
- [27] H. Kumazaki and D. S. Hirashima, Effects of impurities in two-dimensional graphite, *Journal of the Physical Society of Japan* **75**, 053707 (2006), <https://doi.org/10.1143/JPSJ.75.053707>.
- [28] J. Nilsson, A. H. C. Neto, F. Guinea, and N. M. R. Peres, Electronic properties of graphene multilayers, *Phys. Rev. Lett.* **97**, 266801 (2006).
- [29] D. K. Efetov and P. Kim, Controlling electron-phonon interactions in graphene at ultrahigh carrier densities, *Phys. Rev. Lett.* **105**, 256805 (2010).
- [30] S. Yiğen, V. Tayari, J. O. Island, J. M. Porter, and A. R. Champagne, Electronic thermal conductivity measurements in intrinsic graphene, *Phys. Rev. B* **87**, 241411 (2013).
- [31] E. H. Hwang and S. Das Sarma, Acoustic phonon scattering limited carrier mobility in two-dimensional extrinsic graphene, *Phys. Rev. B* **77**, 115449 (2008).
- [32] E. Muñoz, Phonon-limited transport coefficients in extrinsic graphene, *Journal of Physics: Condensed Matter* **24**, 195302 (2012).
- [33] C.-H. Park, N. Bonini, T. Sohler, G. Samsonidze, B. Kozinsky, M. Calandra, F. Mauri, and N. Marzari, Electron-phonon interactions and the intrinsic electrical resistivity of graphene, *Nano Letters* **14**, 1113 (2014), pMID: 24524418, <https://doi.org/10.1021/nl402696q>.
- [34] T. Sohler, M. Calandra, C.-H. Park, N. Bonini, N. Marzari, and F. Mauri, Phonon-limited resistivity of graphene by first-principles calculations: Electron-phonon interactions, strain-induced gauge field, and boltzmann equation, *Phys. Rev. B* **90**, 125414 (2014).
- [35] T. Y. Kim, C.-H. Park, and N. Marzari, The electronic thermal conductivity of graphene, *Nano Letters* **16**, 2439 (2016), pMID: 26907524, <https://doi.org/10.1021/acs.nanolett.5b05288>.
- [36] T. Z. Win, C. W. Aung, G. Khandal, and S. Ghosh, Wiedemann-franz law violation domain for graphene and nonrelativistic systems (2024), arXiv:2404.01061 [cond-mat.mes-hall].
- [37] N. Ashcroft and N. Mermin, *Solid State Physics* (Cengage Learning, 2011).
- [38] J. Ziman, *Electrons and Phonons: The Theory of Transport Phenomena in Solids*, International series of monographs on physics (OUP Oxford, 2001).
- [39] A. Lucas, J. Crossno, K. C. Fong, P. Kim, and S. Sachdev, Transport in inhomogeneous quantum critical fluids and in the dirac fluid in graphene, *Phys. Rev. B* **93**, 075426 (2016).
- [40] M. Zarenia, A. Principi, and G. Vignale, Disorder-enabled hydrodynamics of charge and heat transport in monolayer graphene, *2D Materials* **6**, 035024 (2019).
- [41] A. Rycerz, Wiedemann-franz law for massless dirac fermions with implications for graphene, *Materials* **14**, 10.3390/ma14112704 (2021).
- [42] Y.-T. Tu and S. Das Sarma, Wiedemann-franz law in graphene, *Physical Review B* **107**, 10.1103/physrevb.107.085401 (2023).
- [43] A. Jaiswal, B. Friman, and K. Redlich, Relativistic second-order dissipative hydrodynamics at finite chemical potential, *Physics Letters B* **751**, 548 (2015).
- [44] P. Sahoo, R. Sahoo, and S. K. Tiwari, Wiedemann-Franz law for hot QCD matter in a color string percolation scenario,



- Phys. Rev. D **100**, 051503 (2019), arXiv:1904.06961 [hep-ph].
- [45] R. Rath, S. Tripathy, B. Chatterjee, R. Sahoo, S. Kumar Tiwari, and A. Nath, Violation of Wiedemann-Franz Law for Hot Hadronic Matter created at NICA, FAIR and RHIC Energies using Non-extensive Statistics, *Eur. Phys. J. A* **55**, 125 (2019), arXiv:1902.07922 [hep-ph].
- [46] K. Singh, J. Dey, R. Sahoo, and S. Ghosh, Effect of time-varying electromagnetic field on wiedemann-franz law in a hot hadronic matter, *Physical Review D* **108**, 094007 (2023).
- [47] K. K. Pradhan, D. Sahu, R. Scaria, and R. Sahoo, Conductivity, diffusivity, and violation of the wiedemann-franz law in a hadron resonance gas with van der waals interactions, *Physical Review C* **107**, 014910 (2023).
- [48] D. A. Bandurin, I. Torre, R. K. Kumar, M. B. Shalom, A. Tomadin, A. Principi, G. H. Auton, E. Khestanova, K. S. Novoselov, I. V. Grigorieva, L. A. Ponomarenko, A. K. Geim, and M. Polini, Negative local resistance caused by viscous electron backflow in graphene, *Science* **351**, 1055 (2016), <https://www.science.org/doi/pdf/10.1126/science.aad0201>.
- [49] A. H. Castro Neto, F. Guinea, N. M. R. Peres, K. S. Novoselov, and A. K. Geim, The electronic properties of graphene, *Reviews of Modern Physics* **81**, 109–162 (2009).
- [50] S. Das Sarma, S. Adam, E. H. Hwang, and E. Rossi, Electronic transport in two-dimensional graphene, *Rev. Mod. Phys.* **83**, 407 (2011).
- [51] A. Jaiswal and V. Roy, Relativistic hydrodynamics in heavy-ion collisions: general aspects and recent developments, *Adv. High Energy Phys.* **2016**, 9623034 (2016), arXiv:1605.08694 [nucl-th].
- [52] L. D. Landau., The Theory of a Fermi Liquid, *J. Exptl. Theoret. Phys.* **3**, 920 (1957).
- [53] L. D. Landau., Oscillations in a Fermi Liquid, *J. Exptl. Theoret. Phys.* **5**, 101 (1957).
- [54] L. D. Landau., On the theory of Fermi Liquid, *J. Exptl. Theoret. Phys.* **35** (8), 70 (1959).
- [55] L. Levitov and G. Falkovich, Electron viscosity, current vortices and negative nonlocal resistance in graphene, *Nature Physics* **12**, 672 (2016).
- [56] R. Krishna Kumar, D. Bandurin, F. Pellegrino, Y. Cao, A. Principi, H. Guo, G. Auton, M. Ben Shalom, L. Ponomarenko, G. Falkovich, et al., Superballistic flow of viscous electron fluid through graphene constrictions, *Nature Physics* **13**, 1182 (2017).
- [57] D. A. Bandurin, A. V. Shytov, L. S. Levitov, R. K. Kumar, A. I. Berdyugin, M. Ben Shalom, I. V. Grigorieva, A. K. Geim, and G. Falkovich, Fluidity onset in graphene, *Nat. Commun.* **9**, 4533 (2018).
- [58] P. Gallagher, C.-S. Yang, T. Lyu, F. Tian, R. Kou, H. Zhang, K. Watanabe, T. Taniguchi, and F. Wang, Quantum-critical conductivity of the dirac fluid in graphene, *Science* **364**, 158 (2019), <https://www.science.org/doi/pdf/10.1126/science.aat8687>.
- [59] J. A. Sulpizio, L. Ella, A. Rozen, J. Birkbeck, D. J. Perello, D. Dutta, M. Ben-Shalom, T. Taniguchi, K. Watanabe, T. Holder, et al., Visualizing poiseuille flow of hydrodynamic electrons, *Nature* **576**, 75 (2019).
- [60] A. I. Berdyugin, S. G. Xu, F. M. D. Pellegrino, R. K. Kumar, A. Principi, I. Torre, M. B. Shalom, T. Taniguchi, K. Watanabe, I. V. Grigorieva, M. Polini, A. K. Geim, and D. A. Bandurin, Measuring hall viscosity of graphene's electron fluid, *Science* **364**, 162 (2019), <https://www.science.org/doi/pdf/10.1126/science.aau0685>.
- [61] L. Ella, A. Rozen, J. Birkbeck, M. Ben-Shalom, D. Perello, J. Zultak, T. Taniguchi, K. Watanabe, A. K. Geim, S. Ilani, et al., Simultaneous voltage and current density imaging of flowing electrons in two dimensions, *Nat. Nanotechnol.* **14**, 480 (2019).
- [62] M. J. Ku, T. X. Zhou, Q. Li, Y. J. Shin, J. K. Shi, C. Burch, L. E. Anderson, A. T. Pierce, Y. Xie, A. Hamo, et al., Imaging viscous flow of the dirac fluid in graphene, *Nature* **583**, 537 (2020).
- [63] J. Crossno, J. K. Shi, K. Wang, X. Liu, A. Harzheim, A. Lucas, S. Sachdev, P. Kim, T. Taniguchi, K. Watanabe, et al., Observation of the dirac fluid and the breakdown of the wiedemann-franz law in graphene, *Science* **351**, 1058 (2016).
- [64] A. Block, A. Principi, N. C. Hesp, A. W. Cummings, M. Liebel, K. Watanabe, T. Taniguchi, S. Roche, F. H. Koppens, N. F. van Hulst, et al., Observation of giant and tunable thermal diffusivity of a dirac fluid at room temperature, *Nature Nanotechnology* **16**, 1195 (2021).
- [65] M. Müller, J. Schmalian, and L. Fritz, Graphene: A nearly perfect fluid, *Phys. Rev. Lett.* **103**, 025301 (2009).
- [66] B. N. Narozhny, I. V. Gornyi, M. Titov, M. Schütt, and A. D. Mirlin, Hydrodynamics in graphene: Linear-response transport, *Phys. Rev. B* **91**, 035414 (2015).
- [67] S. A. Hartnoll, P. K. Kovtun, M. Müller, and S. Sachdev, Theory of the nernst effect near quantum phase transitions in condensed matter and in dyonic black holes, *Physical Review B* **76**, 10.1103/physrevb.76.144502 (2007).
- [68] B. N. Narozhny, Hydrodynamic approach to two-dimensional electron systems, *Riv. Nuovo Cim.* **45**, 661 (2022), arXiv:2207.10004 [cond-mat.mes-hall].
- [69] B. N. Narozhny, Electronic hydrodynamics in graphene, *Annals Phys.* **411**, 167979 (2019), arXiv:1905.09686 [cond-mat.mes-hall].
- [70] A. Lucas and K. C. Fong, Hydrodynamics of electrons in graphene, *Journal of Physics: Condensed Matter* **30**, 053001 (2018).
- [71] M. Müller and S. Sachdev, Collective cyclotron motion of the relativistic plasma in graphene, *Phys. Rev. B* **78**, 115419 (2008).
- [72] L. Fritz, J. Schmalian, M. Müller, and S. Sachdev, Quantum critical transport in clean graphene, *Phys. Rev. B* **78**, 085416 (2008).
- [73] M. Müller, L. Fritz, and S. Sachdev, Quantum-critical relativistic magnetotransport in graphene, *Phys. Rev. B* **78**, 115406 (2008).
- [74] M. S. Foster and I. L. Aleiner, Slow imbalance relaxation and thermoelectric transport in graphene, *Phys. Rev. B* **79**, 085415 (2009).

- [75] M. Mendoza, H. J. Herrmann, and S. Succi, Hydrodynamic model for conductivity in graphene, *Scientific reports* **3**, 1052 (2013).
- [76] A. Lucas, Sound waves and resonances in electron-hole plasma, *Physical Review B* **93**, 10.1103/physrevb.93.245153 (2016).
- [77] C. W. Aung, T. Z. Win, G. Khandal, and S. Ghosh, Shear viscosity expression for a graphene system in relaxation time approximation, *Phys. Rev. B* **108**, 235172 (2023), arXiv:2306.14747 [nucl-th].
- [78] L. Janicke, Non-linear electromagnetic waves in a relativistic plasma, *Plasma Physics* **19**, 209 (1977).
- [79] M. Palmroth, U. Ganse, Y. Pfau-Kempf, M. Battarbee, L. Turc, T. Brito, M. Grandin, S. Hoilijoki, A. Sandroos, and S. von Althaus, Vlasov methods in space physics and astrophysics, *Living Reviews in Computational Astrophysics* **4**, 10.1007/s41115-018-0003-2 (2018).
- [80] Y. Elskens and M. K.-H. Kiessling, Microscopic foundations of kinetic plasma theory: The relativistic vlasov–maxwell equations and their radiation-reaction-corrected generalization, *Journal of Statistical Physics* **180**, 749–772 (2020).
- [81] K. Kushwah and G. S. Denicol, Relativistic dissipative magnetohydrodynamics from the Boltzmann equation for a two-component gas, *Phys. Rev. D* **109**, 096021 (2024), arXiv:2402.01597 [nucl-th].
- [82] A. K. Panda, A. Dash, R. Biswas, and V. Roy, Relativistic non-resistive viscous magnetohydrodynamics from the kinetic theory: a relaxation time approach, *JHEP* **03**, 216, arXiv:2011.01606 [nucl-th].
- [83] A. K. Panda, A. Dash, R. Biswas, and V. Roy, Relativistic resistive dissipative magnetohydrodynamics from the relaxation time approximation, *Phys. Rev. D* **104**, 054004 (2021), arXiv:2104.12179 [nucl-th].
- [84] J. Anderson and H. Witting, A relativistic relaxation-time model for the boltzmann equation, *Physica* **74**, 466 (1974).
- [85] G. S. Rocha, D. Wagner, G. S. Denicol, J. Noronha, and D. H. Rischke, Theories of Relativistic Dissipative Fluid Dynamics, *Entropy* **26**, 189 (2024), arXiv:2311.15063 [nucl-th].
- [86] G. S. Rocha and G. S. Denicol, Transient fluid dynamics with general matching conditions: A first study from the method of moments, *Phys. Rev. D* **104**, 096016 (2021), arXiv:2108.02187 [nucl-th].
- [87] A. Dwibedi, N. Padhan, A. Chatterjee, and S. Ghosh, Transport Coefficients of Relativistic Matter: A Detailed Formalism with a Gross Knowledge of Their Magnitude, *Universe* **10**, 132 (2024), arXiv:2404.01421 [nucl-th].
- [88] S. R. De Groot, Relativistic Kinetic Theory. Principles and Applications, edited by W. A. Van Leeuwen and C. G. Van Weert (1980).
- [89] C. Cercignani and G. M. Kremer, Thermomechanics of relativistic fluids, in The Relativistic Boltzmann Equation: Theory and Applications (Birkhäuser Basel, Basel, 2002) pp. 99–110.
- [90] A. Jaiswal, Relativistic dissipative hydrodynamics from kinetic theory with relaxation time approximation, *Phys. Rev. C* **87**, 051901 (2013), arXiv:1302.6311 [nucl-th].
- [91] N. Itoh, Hydrostatic Equilibrium of Hypothetical Quark Stars, *Progress of Theoretical Physics* **44**, 291 (1970), <https://academic.oup.com/ptp/article-pdf/44/1/291/5357142/44-1-291.pdf>.
- [92] T. D. Lee and G. C. Wick, Vacuum stability and vacuum excitation in a spin-0 field theory, *Phys. Rev. D* **9**, 2291 (1974).
- [93] J. C. Collins and M. J. Perry, Superdense matter: Neutrons or asymptotically free quarks?, *Phys. Rev. Lett.* **34**, 1353 (1975).