Semi-Quenched Invariance Principle for the Random Lorentz Gas – Beyond the Boltzmann-Grad Limit

Bálint Tóth

Rényi Institute, Budapest, HU and University of Bristol, UK

January 3, 2025

Abstract

By synchronously coupling multiple Lorentz trajectories exploring the same environment consisting of randomly placed scatterers in \mathbb{R}^3 we upgrade the annealed invariance principle proved in [6] to quenched setting (that is, valid for almost all realizations of the environment) along sufficiently fast extractor sequences.

MSC2010: 60F17; 60K35; 60K37; 60K40; 82C22; 82C31; 82C40; 82C41

KEY WORDS AND PHRASES: Lorentz-gas; invariance principle; scaling limit; coupling; almost sure convergence

> Révész Pali emlékére (Dedicated to the memory of Pali Révész)

1 Introduction

Let $(\Omega, \mathcal{F}, \mathbf{P})$ be a sufficiently large probability space which supports (inter alia) a Poisson Point Process (PPP) of intensity 1 on \mathbb{R}^d , denoted ϖ . Other, independent random elements jointly defined on $(\Omega, \mathcal{F}, \mathbf{P})$ will also be considered later. Therefore, it is best to think about $(\Omega, \mathcal{F}, \mathbf{P})$ as a product space which in one of its factors supports the PPP ϖ and on the other factor (or factors) many other random elements, independent of ϖ , to be introduced later. To keep notation simple we do not denote explicitly this product structure of $(\Omega, \mathcal{F}, \mathbf{P})$. However, as this note is about *quenched* laws, that is about laws and limits conditioned on *typical* ϖ , we denote

$$\mathbf{P}_{\varpi}(\cdot) := \mathbf{P}(\cdot \mid \mathcal{F}(\varpi)), \qquad \mathbf{E}_{\varpi}(\cdot) := \mathbf{E}(\cdot \mid \mathcal{F}(\varpi)),$$

where $\mathcal{F}(\varpi) \subset \mathcal{F}$ is the sigma algebra generated by the PPP ϖ .

Given

$$\varepsilon > 0, \qquad r = r_{\varepsilon} := \varepsilon^{d/(d-1)}$$

and

$$v \in S^{d-1} := \{ u \in \mathbb{R}^d, |u| = 1 \}$$

let

$$t \mapsto X_{\varepsilon}(t) \in \mathbb{R}^d$$

be the Lorentz trajectory among fixed spherical scatterers of radius r centred at the points of the rescaled PPP

$$\varpi_{\varepsilon} := \{ \varepsilon q : q \in \varpi, \ |q| > \varepsilon^{-1} r = \varepsilon^{1/(d-1)} \}, \tag{1}$$

with initial conditions

$$X_{\varepsilon}(0) = 0, \qquad \dot{X}_{\varepsilon}(0) = v.$$

In plain words: $t \mapsto X_{\varepsilon}(t)$ is the trajectory of a point particle starting from the origin with velocity v, performing free flight in the complement of the scatterers and scattering elastically on them.

Notes: (1) In order to define the Lorentz trajectory we have to disregard those points of the rescaled PPP ϖ_{ε} within distance r from the origin. However, this will not effect whatsoever our arguments and conclusions since, with probability 1, for ε sufficiently small there are no points like this.

(2) Given ε and the initial velocity v, the trajectory $t \mapsto X_{\varepsilon}(t)$ is almost surely well defined for $t \in [0, \infty)$. That is: almost surely all scatterings will happen on an unique scatterer, the singular sets at the intersection of more than one scatterers will be a.s. avoided.

In order to properly (and, comparably) formulate our invariance principles first we recall the relevant function spaces. Let

$$\mathcal{C} := \mathcal{C}([0,\infty), \mathbb{R}^d) := \{\mathfrak{z} : [0,\infty) \to \mathbb{R}^d : \mathfrak{z} \text{ continuous}\},\$$

endowed with the topology of uniform convergence on compact subintervals of $[0.\infty)$, which is metrizable and C complete and separable. For details see, e.g., [12]. Further on, let

$$\mathcal{C}(\mathcal{C}) := \mathcal{C}(\mathcal{C}([0,1],\mathbb{R}^d),\mathbb{R}) := \{F : \mathcal{C} \to \mathbb{R} : F \text{ continuous, } \|F\|_{\infty} := \sup_{\mathfrak{z} \in \mathcal{C}} |F(\mathfrak{z})| < \infty\},\$$
$$\mathcal{C}_0(\mathcal{C}) := \mathcal{C}_0(\mathcal{C}([0,1],\mathbb{R}^d),\mathbb{R}) := \{F \in \mathcal{C}(\mathcal{C}) : \forall \delta > 0, \exists K \Subset \mathcal{C} : \sup_{\mathfrak{z} \in \mathcal{C} \setminus K} |F(\mathfrak{z})| < \delta\}.$$

 $(\mathcal{C}_0(\mathcal{C}), \|\cdot\|_{\infty})$ is a separable Banach space. We will also denote by $t \mapsto W(t)$ a standard Brownian motion in \mathbb{R}^d , and recall from [1], [11], [12] criteria for weak convergence of probability measures on \mathcal{C} .

In [6] the following *annealed* invariance principle was proved.

Theorem 1. ([6] Theorem 1) Let d = 3, $\varepsilon \to 0$, $r_{\varepsilon} = \varepsilon^{d/(d-1)}$ and $T_{\varepsilon} \to \infty$ be such that

$$\lim_{\varepsilon \to 0} r_{\varepsilon} T_{\varepsilon} = 0.$$
 (2)

Let $t \mapsto X_{\varepsilon}(t)$ be the sequence of Lorentz trajectories among the spherical scatterers of radius r_{ε} centred at the points ϖ_{ε} cf. (1), and with deterministic initial velocities $v_{\varepsilon} \in S^{d-1}$. For any $F \in \mathcal{C}_0(\mathcal{C})$,

$$\lim_{\varepsilon \to 0} \left| \mathbf{E} \left(F(T_{\varepsilon}^{-1/2} X_{\varepsilon}(T_{\varepsilon} \cdot)) \right) - \mathbf{E} \left(F(W(\cdot)) \right) \right| = 0.$$
(3)

Remarks: (R1) (On dimension.) Although some crucial elements of the proofs in [6], on which the present note is based, are worked out in full detail in dimension d = 3 only, we prefer to use the generic notation d for dimension with the explicit warning that in the actual results and proofs d = 3 is meant. See Remark (R7) below and the paragraph "remarks on dimension" in section 1 of [6] for comments on possible extensions to dimensions other than d = 3.

(R2) Theorem 1 is an *annealed* invariance principle in the sense that on the left-hand side of (3) the probability distribution of the rescaled Lorentz trajectory is provided by the random environment ϖ . The proofs in [6] rely on a genuinely annealed argument: a simultaneous realization of the PPP ϖ and the trajectory $t \mapsto X_{\varepsilon}(t)$.

(R3) The main result in [6] (Theorem 2 of that paper) is actually stronger, assuming

$$\lim_{\varepsilon \to 0} (r_{\varepsilon} |\log \varepsilon|)^2 T_{\varepsilon} = 0$$

rather than (2). However, the *semi-quenched* invariance principle of this note, Theorem 2 below, is directly comparable to this weaker version.

Theorem 2. Let d=3, $\varepsilon \to 0$, $r_{\varepsilon} = \varepsilon^{d/(d-1)}$, $T_{\varepsilon} \to \infty$ and $\beta_{\varepsilon} \in (0,1]$ be such that $\lim_{\varepsilon \to 0} r_{\varepsilon}(T_{\varepsilon} + \beta_{\varepsilon}^{-1}) = 0, \tag{4}$

and define the solid angle domains

$$B_{\varepsilon} := \{ u \in S^{d-1} : 2 \arcsin \sqrt{(1 - u \cdot v_{\varepsilon})/2} \le \beta_{\varepsilon} \}, \qquad v_{\varepsilon} \in S^{d-1} \quad deterministic.$$

Let $t \mapsto X_{\varepsilon}(t)$ be the sequence of Lorentz trajectories among the spherical scatterers of radius r_{ε} centred at the points ϖ_{ε} cf. (1), and with initial velocities $v_{\varepsilon} \sim \text{UNI}(B_{\varepsilon})$ sampled independently of the PPP ϖ . For any $F \in C_0(\mathcal{C})$,

$$\lim_{\varepsilon \to 0} \mathbf{E} \left(\left| \mathbf{E}_{\varpi} \left(F(T_{\varepsilon}^{-1/2} X_{\varepsilon}(T_{\varepsilon} \cdot)) \right) - \mathbf{E} \left(F(W(\cdot)) \right) \right| \right) = 0$$

Remarks ctd.: (R4) Theorem 2 is an invariance principle valid *in probability* with respect to the random environment ϖ . An equivalent formulation is that under the stated conditions, for any $\delta > 0$

$$\lim_{\varepsilon \to 0} \mathbf{P}\big(\{\varpi : D_{\mathrm{LP}}\big(\mathsf{law-of}(T_{\varepsilon}^{-1/2}X_{\varepsilon}(T_{\varepsilon}\cdot) \,|\, \mathcal{F}(\varpi)), \mathsf{law-of}(W(\cdot))\big) > \delta\}\big) = 0,$$

where $D_{LP}(\cdot, \cdot)$ denotes the Lévy-Prohorov distance between probability measures on \mathcal{C} .

We will actually prove a stronger statement from which Theorem 2 follows as a corollary: In the setting of Theorem 2, for almost all realizations of the PPP ϖ , along (precisely quantified) sufficiently fast converging subsequences $\varepsilon_n \to 0$, the invariance principle holds:

Theorem 3. Let d = 3, $\varepsilon_n \to 0$, $r_n := \varepsilon_n^{d/(d-1)}$, $T_n \to \infty$ and $\beta_n \in (0, 1]$ be such that

$$\sum_{n} \left(\log n \, r_n T_n + (\log n)^2 \left(r_n \beta_n^{-1} \right)^{(d-1)/d} \right) < \infty,\tag{5}$$

and define the solid angle domains

$$B_n := \{ u \in S^{d-1} : 2 \arcsin \sqrt{(1 - u \cdot v_n)/2} \le \beta_n \}, \qquad v_n \in S^{d-1} \ deterministic.$$

Let $t \mapsto X_n(t)$ be the sequence of Lorentz trajectories among the spherical scatterers of radius r_n centred at the points $\varpi_n := \varpi_{\varepsilon_n}$ cf. (1), and with initial velocities $v_n \sim \text{UNI}(B_n)$ sampled independently of the PPP ϖ . For almost all realizations of the PPP ϖ , for any $F \in \mathcal{C}_0(\mathcal{C})$,

$$\lim_{n \to \infty} \left| \mathbf{E}_{\varpi} \left(F(T_n^{-1/2} X_n(T_n \cdot)) \right) - \mathbf{E} \left(F(W(\cdot)) \right) \right| = 0.$$

Remarks ctd.: (R5) Theorem 2 is a corollary of Theorem 3, as under condition (4) from any sequence $\varepsilon_n \to 0$ a subsequence $\varepsilon_{n'}$ can be extracted that satisfies condition (5). On the other hand, Theorem 3 is genuinely stronger than Theorem 2, as the former provides an explicit quantitative characterization of the sequences $\varepsilon_n \to 0$ along which the quenched (i.e., almost sure) invariance principle holds.

(R6) For a comprehensive historical survey of the invariance principle for the random Lorentz gas we refer to the monograph [10] and to section 1 on [6]. We just mention here that the main milestones preceding [6] are [4, 5], [9], and [2]. The new result of this note (i.e., Theorems 2 and 3) is to be compared with that in [2] where a fully quenched invariance principle is proved for the 2-dimensional random Lorentz gas in the Boltzmann-Grad limit, on kinetic time scales. The weakness of our result (compared with [2]) is that the limit theorem is semi-quenched, in the sense that a.s. invariance principle is proved along sufficiently fast converging sequences ε_n only. On the other hand the strengths are twofold. (\star) The proof works in dimension d = 3 and it is "hands-on", not relying on the heavy computational details of [2] (performable only in d = 2). See remark (R7) below for possible extensions to dimensions other than d = 3. ($\star\star$) The time-scale of validity is much longer, $T_{\varepsilon} = o(\varepsilon^{-d/(d-1)})$ rather than $T_{\varepsilon} = \mathcal{O}(1)$, as in [2].

(R7) The results of [6] are stated and the proofs are fully spelled out for dimension d = 3. Therefore the new results of this note (which rely on those of [6]) are also valid in d = 3 only. However, as noted in the paragraph "remarks on dimension" in [6], extension to other dimensions is possible, on the expense of more involved details due

partly to recurrence (in d = 2) and partly to the non-uniform scattering cross section (in all dimensions other than d = 3). For arguments in d = 2 see [7], [8].

The strategy of proof in [6] (also extended to [7], [8]) was based on a *coupling* of the mechanical/Hamiltonian Lorentz trajectory within the environment consisting of randomly placed scatterers and the Markovian random flight trajectory. The coupling is realized as an *exploration* of the random environment along the trajectory of the tagged particle. This construction is *par excellence* annealed, as the environment and the trajectory of the moving particle are constructed synchronously (rather than first sampling the environment and consequently letting the particle move in the fully sampled environment). However, this exploration process can be realized synchronously with multiple (actually, many) moving particles, which, as long as they explore disjoint areas of the environment, are independent in the annealed sense (due to the Poisson character of the environment). Applying a Strong Law of Large Numbers to tests of these trajectories will provide the quenched invariance principle - valid for typical realizations of the environment. A somewhat similar exploration strategy is used in the very different context of random walks on sparse random graphs, [3].

Acknowledgements.

I thank Pietro Caputo and Justin Salez for drawing my attention to their paper [3]. This work was supported by the grant no. K-143468 of the Hungarian National Research, Development and Innovation Office (NKFIH).

2 Construction and Quenched Coupling

2.1 Synchronous Lorentz trajectories

Beside ε and $r = \varepsilon^{d/(d-1)}$ let $N \in \mathbb{N}$, and

$$v_j \in S^{d-1}, \qquad 1 \le j \le N.$$

Given these we define *jointly* N synchronous Lorentz trajectories

$$t \mapsto X_j(t) \in \mathbb{R}^d, \qquad 1 \le j \le N,$$

among fixed spherical scatterers of radius r centred at the points of the rescaled PPP ϖ_{ε} cf. (1), with initial conditions

$$X_j(0) = 0,$$
 $X_j(0) = v_j,$ $1 \le j \le N.$

(Given the parameters and the initial velocities, the trajectories $t \mapsto X_j(t)$, $1 \le j \le N$, are almost surely well defined for $t \in [0, \infty)$.)

We will consider the càdlàg version of the velocity processes

$$V_j(t) := X_j(t). \qquad 1 \le j \le N,$$

and use the notation $X := \{X_j : 1 \le j \le N\}.$

In order to construct the *quenched coupling* with Markovian flight processes (in the next subsection) we have to define some further variables in terms of the Lorentz processes $t \mapsto X(t)$.

First the collision times $\tau_{j,k}$, $1 \leq j \leq N$, $k \geq 0$:

$$\tau_{j,0} := 0, \qquad \qquad \tau_{j,k+1} := \inf\{t > \tau_{j,k} : V_j(t) \neq V_j(\tau_{j,k})\}.$$

In plain words: $\tau_{j,k}$ is the time of the k-th scattering of the Lorentz trajectory $X_j(\cdot)$. We will use the notation

$$X_{j,k} := X_j(\tau_{j,k}), \qquad V_{j,k+1} := V_j(\tau_{j,k}). \qquad X'_{j,k} := X_{j,k} + r \frac{V_{j,k} - V_{j,k+1}}{|V_{j,k} - V_{j,k+1}|}$$

That is: $X_{j,k}$ is the position of the Lorentz trajectory at the instant of its k-th collision, $V_{j,k+1}$ is its velocity right after this collision, and $X'_{j,k}$ is the position of the centre of the fixed scatterer which had caused this collision. Altogether, the continuous-time trajectory is written

$$X_{j}(t) = X_{j,k} + (t - \tau_{j,k})V_{j,k+1}, \qquad \text{for} \quad t \in [\tau_{j,k}, \tau_{j,k+1})$$

Next, the *indicators of freshness*:

$$a_{j,0} := 1, \qquad a_{j,k} := \begin{cases} 1 & \text{if } \forall \delta > 0 : \min_{\substack{1 \le i \le N \\ 0 \le s \le \tau_{j,k} - \delta}} |X_i(s) - X'_{j,k}| > r \\ 0 & \text{otherwise} \end{cases} \quad (k \ge 1).$$

In plain words, $a_{j,k}$ indicates whether the *j*-th trajectory at its *k*-th collision encounters a fresh scatterer, never seen in the past by either one of the *N* Lorentz trajectories. Finally, the *shadow indicators* $b_j(t, v), t \in [0, \infty), v \in S^{d-1}$:

$$b_j(t,v) := \begin{cases} 0 & \text{if } \forall \delta > 0 : \min_{\substack{1 \le i \le N \\ 0 \le s \le t - \delta}} \left| X_i(s) - X_j(t) + r \frac{v - V_j(t)}{|v - V_j(t)|} \right| > r, \\ 1 & \text{otherwise} \end{cases}$$

In plain words, $b_j(t, v)$ indicates whether at time t a virtual scatterer (virtually) causing new velocity v would be *mechanically inconsistent* with the past of the paths.

2.2 Quenched coupling with independent Markovian flight processes

On the same probability space $(\Omega, \mathcal{F}, \mathbf{P})$ and jointly with the Lorentz trajectories X, we construct N independent Markovian flight processes

$$t \mapsto Y_j(t) \in \mathbb{R}^d, \qquad 1 \le j \le N,$$

with initial conditions identical to those of the Lorentz trajectories

$$Y_j(0) = 0,$$
 $Y_j(0) = v_j,$ $1 \le j \le N.$

The processes $\{Y_j(\cdot): 1 \leq j \leq N\}$, are independent, and consist of i.i.d. EXP(1)distributed free flights with independent $UNI(S^{d-1})$ -distributed velocities. See [6] for a detailed exposition of the Markovian flight processes. We will again consider the càdlàg version of their velocity processes

$$U_j(t) := Y_j(t), \qquad 1 \le j \le N.$$

and use the notation $Y := \{Y_j : 1 \le j \le N\}.$

The construction of the coupling goes as follows. Assume that the probability space $(\Omega, \mathcal{F}, \mathbf{P})$, besides and independently of the PPP ϖ supports the fully independent random variables

$$\widetilde{\xi}_{j,k} \sim \text{EXP}(1), \qquad \qquad \widetilde{U}_{j,k+1} \sim \text{UNI}(S^{d-1}), \qquad \qquad j = 1, \dots, N, \quad k \ge 1,$$

and let

$$\widetilde{\theta}_{j,k} := \sum_{l=1}^{k} \widetilde{\xi}_{j,l}, \qquad \qquad b_{j,k} := b_j(\widetilde{\theta}_{j,k}, \widetilde{U}_{j,k+1}).$$

We construct the piecewise constant càdlàg velocity processes $U_j(\cdot)$ successively on the time intervals $[\tau_{j,k}, \tau_{j,k+1}), k = 0, 1, \ldots$

• At $\tau_{j,k}$:

• If
$$a_{j,k} = 0$$
 then let $U_j(\tau_{j,k}) = U_j(\tau_{j,k})$.
• If $a_{j,k} = 1$ then let $U_j(\tau_{j,k}) = V_{j,k+1}$.

- At any $\tilde{\theta}_{j,l} \in (\tau_j(k), \tau_j(k+1))$
 - If $b_{j,l} = 0$ then let $U_j(\theta_{j,l}) = U_j(\theta_{j,l}^-)$.
 - If $b_{j,l} = 1$ then let $U_j(\theta_{j,l}) = \widetilde{U}_{j,l+1}$.
- In the open subintervals of $(\tau_{j,k}, \tau_{j,k+1})$ determined by the times $\{\tilde{\theta}_{j,l} : l \geq 1\} \cap (\tau_{j,k}, \tau_{j,k+1})$ keep the value of $U_j(t)$ constant.

It is true - and not difficult to see - that the velocity processes $\{U_j(t): 1 \leq j \leq N\}$ constructed in this way are independent between them, and distributed as required. That is, they consist of i.i.d. EXP(1)-distributed intervals where their values are i.i.d. $\text{UNI}(S^{d-1})$. This is due to the fact that each Lorentzian scatterer is taken into account exactly once, when first explored by a Lorentz particle, and missing scatterings (due to areas shadowed by past trajectories) are compensated for by the auxiliary events at times $\tilde{\theta}_{j,l}$. Consistently with the notation introduced for the Lorentz trajectories, we write

$$\theta_{j,0} := 0,$$
 $\theta_{j,k+1} := \inf\{t > \theta_{j,k} : U_j(t) \neq U_j(\theta_{j,k})\},$ (6)

and

$$Y_{j,k} := Y_j(\theta_{j,k}), \qquad U_{j,k+1} := U_j(\theta_{j,k}), \qquad Y'_{j,k} := Y_{j,k} + r \frac{U_{j,k} - U_{j,k+1}}{|U_{j,k} - U_{j,k+1}|}.$$
 (7)

That is, $Y_{j,k}$ is the position of the Markovian flight trajectory at the instant of its k-th scattering, $U_{j,k+1}$ is its velocity right after this scattering, and $Y'_{j,k}$ would be the position of the centre of a spherical scatterer of radius r which could have caused this scattering. Altogether, the continuous time Markovian flight trajectory is written as

$$Y_j(t) = Y_{j,k} + (t - \theta_{j,k})U_{j,k+1} \qquad \text{for} \quad t \in [\theta_{j,k}, \theta_{j,k+1}).$$

Note that

$$\{\theta_{j,k}: k \ge 0\} \subseteq \{\tau_{j,k}: k \ge 0\} \cup \{\widetilde{\theta}_{j,k}: k \ge 0\}.$$

This coupling between Lorentz trajectories and Markovian flight processes has the same joint distribution as the one presented in [6]. However, it is realized in a different way. While in [6] first we constructed the Markovian flight processs Y and conditionally on this we constructed the coupled Lorentz exploration process X, here we do this in reverse order: first we realize the N Lorentz exploration processes $X = \{X_1, \ldots, X_N\}$ and given these we realize the N independent Markovian flight processes $Y = \{Y_1, \ldots, Y_N\}$ coupled to them.

2.3 Control of tightness of the coupling

We quantify the tightness of the coupling.

The relevant filtrations are

$$\begin{aligned} \mathcal{F}_t^X &:= \sigma(\{X_j(s) : 1 \le j \le N, \ 0 \le s \le t\}) \\ \mathcal{F}_t^Y &:= \sigma(\{Y_j(s) : 1 \le j \le N, \ 0 \le s \le t\}) \\ \mathcal{F}_t^{X,Y} &:= \mathcal{F}_t^X \lor \mathcal{F}_t^Y \end{aligned}$$

Next we define some relevant stopping times, indicating explicitly the filtration with respect to which they are adapted

$$\begin{aligned}
\sigma_{1} &:= \min\{\tau_{j,k} : a_{j,k} = 0\} & \text{stopping time w.r.t. } \mathcal{F}_{t}^{X} \\
\sigma_{2} &:= \min\{\theta_{j,l} : b_{j,l} = 1\} & \text{stopping time w.r.t. } \mathcal{F}_{t}^{X,Y} \\
\sigma_{3} &:= \inf\{t > 0 : \min\{|Y_{j}(t) - Y_{i,k}'| : \theta_{i,k} < t\} < r\} & \text{stopping time w.r.t. } \mathcal{F}_{t}^{Y} \\
\sigma_{4} &:= \min\{\theta_{i,k} : \inf\{|Y_{j}(s) - Y_{i,k}'| : 0 \le s \le \theta_{i,k}\} < r\} & \text{stopping time w.r.t. } \mathcal{F}_{t}^{Y} \\
\sigma &:= \inf\{t : X(t) \ne Y(t)\} = \min\{\sigma_{1}, \sigma_{2}\} & \text{stopping time w.r.t. } \mathcal{F}_{t}^{X,Y}
\end{aligned}$$

In plain words:

- σ_1 is the first time an already explored scatterer is re-encountered by one of N Lorentz particles. We'll call it the time of the first recollision. This is a stopping time with respect to the filtration \mathcal{F}_t^X
- σ_2 is the first time when in the construction of the Markovian flight processes a compensating scattering occurs. We'll call it the time of the first shadowed scattering. This is a stopping time with respect to the largest filtration $\mathcal{F}_t^{X,Y}$
- σ_3 is the first time when a Markovian flight trajectory encounters a virtual scatterer which would have caused an earlier scattering event of one of the Markovian flight processes. This is a stopping time with respect to the filtration \mathcal{F}_t^Y
- σ_4 is the first scattering of one of the Markovian flight trajectories encounters a virtual scatterer which would have caused an earlier scattering event of one of the Markovian flight processes. This is a stopping time with respect to the filtration \mathcal{F}_t^Y
- σ is the time of the first mismatch between the Lorentz trajectories X(t) and the coupled Markovian flight trajectories Y(t). This is (a priori) a stopping time with respect to the largest filtration $\mathcal{F}_t^{X,Y}$

Although these are stopping times with respect to different filtrations it clearly follows from the construction of the coupling that

$$\sigma_1 \mathbb{1}_{\{\sigma_1 < \sigma_2\}} = \sigma_3 \mathbb{1}_{\{\sigma_3 < \sigma_4\}} \quad \text{and} \quad \sigma_2 \mathbb{1}_{\{\sigma_2 < \sigma_1\}} = \sigma_4 \mathbb{1}_{\{\sigma_4 < \sigma_3\}}.$$

Hence, $\min\{\sigma_1, \sigma_2\} = \min\{\sigma_3, \sigma_4\}$ and thus actually

$$\sigma = \min\{\sigma_3, \sigma_4\}.\tag{8}$$

Although by definition σ is a priori adapted to the joint filtration $\mathcal{F}_t^{X,Y}$, due to the particularities of the coupling construction, according to (8) it is actually a stopping time with respect to the filtration of the Markovian flight trajectories \mathcal{F}_t^Y which makes it suitable to purely probabilistic control. In what follows we will use the expression (8) as definition of the first mismatch time σ .

Proposition 1. There exists an absolute constant $C < \infty$, such that for any $N, T < \infty$

$$\mathbf{P}(\sigma < T) \le Cr(NT + N^2 w^{-1}),\tag{9}$$

where

$$w := 2 \min_{1 \le i < j \le N} \arcsin \sqrt{(1 - v_i \cdot v_j)/2} \tag{10}$$

is the minimum angle between any two of the starting velocities.

Proof. Given $N, T < \infty$, let

$$A_{i} := \{ \min\{|Y_{i}(t) - Y_{i,k}| : 0 < t < T, 0 < \theta_{i,k} < T \} < 2r \}, \qquad 1 \le i \le N,$$

$$B_{i,j} := \{ \min\{|Y_{i}(t) - Y_{j,k}| : 0 < t < T, 0 < \theta_{j,k} < T \} < 2r \}, \qquad 1 \le i \ne j \le N,$$

$$\subseteq \{ \min\{|Y_{i}(t) - Y_{j,k}| : 0 < t < \infty, 1 \le k < \infty \} < 2r \} =: \widetilde{B}_{i,j}$$

Obviously,

$$\{\min\{\sigma_3, \sigma_4\} < T\} \subseteq \big(\bigcup_{1 \le i \le N} A_i\big) \bigcup \big(\bigcup_{1 \le i \ne j \le N} B_{i,j}\big) \subseteq \big(\bigcup_{1 \le i \le N} A_i\big) \bigcup \big(\bigcup_{1 \le i \ne j \le N} \widetilde{B}_{i,j}\big).$$
(11)

By careful application of the Green function estimates of of section 3 in [6] we get the bounds

$$\mathbf{P}(A_i) \le CrT, \qquad \mathbf{P}(\widetilde{B}_{i,j}) \le Crw^{-1}.$$
(12)

The first one of these is explicitly stated in Corollary 1 of Lemma 4 (on page 608) of of [6]. When proving the second one, one hase to take into account that the directions of the first flights of Y_i and Y_j are deterministic, v_i , respectively, v_j and the angle between these two directions determines the probability of interference between the two trajectories during the first free flights. We omit the computational details fully based on the aforementioned Green function estimates and very similar to those in [6].

Finally (9) follows from (11) and (12) by a straightforward union bound. \Box

3 Proof of Theorem 3

3.1 Triangular array of processes

Let now $\varepsilon_n \to 0$, $r_n = \varepsilon_n^{d/(d-1)}$, $T_n \to \infty$, $\beta_n \in (0, 1]$ be as in (5), and choose an increasing sequence N_n such that

$$(\log n)^{-1} N_n \to \infty \tag{13}$$

and the stronger summability

$$\sum_{n} \left(N_n r_n T_n + N_n^2 \left(r_n \beta_n^{-1} \right)^{(d-1)/d} \right) < \infty$$
(14)

still holds. (Given (5) this can be done.)

Assume that the probability space $(\Omega, \mathcal{F}, \mathbf{P})$ supports a *triangular array* of processes

$$\{ \{ (X_{n,j}(\cdot), Y_{n,j}(\cdot)) : 1 \le j \le N_n \} : n \ge 1 \}$$

row-wise constructed as in section 2, with parameters ε_n , r_n , β_n , and with i.i.d. initial velocities

$$v_{n,j} \sim \text{UNI}(B_n), \qquad 1 \le j \le N_n,$$

which are also independent of all other randomness in the row.

Note that

- The row-wise construction, and thus the joint distribution of $\{(X_{n,j}(\cdot), Y_{n,j}(\cdot)) : 1 \le j \le N_n\}$ is prescribed.
- The PPP $\varpi_n := \varpi_{\varepsilon_n}$ are obtained by rescaling the same realisation of the PPP ϖ . This makes the sequence of couplings *quenched*.
- The joint distribution of the probabilistic ingredients apart of ϖ in different rows is irrelevant.

Lemma 1. Let the sequence $N_n \in \mathbb{N}$ be as in (13) and $\{\{\Upsilon_{n,j} : 1 \leq j \leq N_n\} : n \geq 1\}$ a jointly defined triangular array of real valued, uniformly bounded, row-wise i.i.d. zeromean random variables:

$$\mathbf{P}(|\Upsilon_{n,j}| \le M) = 1, \qquad \mathbf{E}(\Upsilon_{n,j}) = 0.$$

Then,

$$\mathbf{P}\big(\lim_{n\to\infty}N_n^{-1}\sum_{j=1}^{N_n}\Upsilon_{n,j}\to 0\big)=1.$$

Proof. This is a direct (and straightforward) consequence of Hoeffding's inequality and the Borel-Cantelli lemma. By Hoeffding's inequality, for any $\delta > 0$

$$\mathbf{P}\left(\pm N_n^{-1}\sum_{j=1}^{N_n}\Upsilon_{n,j} > \delta\right) \le e^{-\delta^2 N_n/(4M^2)}.$$

Hence, due to (13) and Borel-Cantelli, for any $\delta > 0$

$$\mathbf{P}\Big(\lim_{n\to\infty}\pm N_n^{-1}\sum_{j=1}^{N_n}\Upsilon_{n,j}>\delta\Big)=0.$$

Proposition 2. Almost surely, for any $F \in C_0(\mathcal{C})$,

$$\lim_{n \to \infty} \left(N_n^{-1} \sum_{j=1}^n F(T_n^{-1/2} Y_{n,j}(T_n \cdot)) - \mathbf{E} \left(F(T_n^{-1/2} Y_{n,1}(T_n \cdot)) \right) \right) = 0$$
(15)

$$\lim_{n \to \infty} \left(N_n^{-1} \sum_{j=1}^n F(T_n^{-1/2} X_{n,j}(T_n \cdot)) - \mathbf{E}_{\varpi} \left(F(T_n^{-1/2} X_{n,1}(T_n \cdot)) \right) \right) = 0$$
(16)

Proof. The same statement with " for any $F \in \mathcal{C}_0(\mathcal{C})$, almost surely" follows from Lemma 1, noting that the triangular array of *annealed* random variables

$$\Upsilon_{n,j} := F(T_n^{-1/2}Y_{n,j}(T_n \cdot)) - \mathbf{E} \big(F(T_n^{-1/2}Y_{n,j}(T_n \cdot)) \big), \qquad 1 \le j \le N_n, \quad n \ge 1$$

respectively, the triangular array of quenched random variables

$$\widetilde{\Upsilon}_{n,j,\varpi} := F(T_n^{-1/2}X_{n,j}(T_n \cdot)) - \mathbf{E}_{\varpi} \big(F(T_n^{-1/2}X_{n,j}(T_n \cdot)) \big), \qquad 1 \le j \le N_n, \quad n \ge 1$$

meet the conditions of the lemma.

Going from "for any $F \in \mathcal{C}_0(\mathcal{C})$, almost surely" to "almost surely, for any $F \in \mathcal{C}_0(\mathcal{C})$ " we rely on separability of the Banach space $(\mathcal{C}_0(\mathcal{C}), \|\cdot\|_{\infty})$.

Proposition 3. For any $F \in C_0(\mathcal{C})$,

$$\lim_{n \to \infty} \mathbf{E} \left(F(T_n^{-1/2} Y_{n,1}(T_n \cdot)) \right) = \mathbf{E} \left(W(\cdot) \right) \right).$$
(17)

Proof. This is Donsker's theorem.

Proposition 4.

$$\mathbf{P}\big(\max\{n:\sigma_n < T_n\} < \infty\big) = 1. \tag{18}$$

That is: almost surely, for all but finitely many n

$$X_{n,j}(t) = Y_{n,j}(t), \quad 1 \le j \le N_n, \quad 0 \le t \le T_n.$$
 (19)

Proof. Let

$$\alpha_n := r_n^{1/d} \beta_n^{(d-1)/d}$$

With this choice

$$r_n \alpha_n^{-1} = (\alpha_n \beta_n^{-1})^{d-1} = (r_n \beta_n^{-1})^{(d-1)/d}$$

As in (10), denote

$$w_n := 2 \min_{1 \le i < j \le N_n} \arcsin \sqrt{\left(1 - v_{n,i} \cdot v_{n,j}\right)/2}$$

the minimum angle between any two of the starting velocities. Then, using elementary geometry and the stopping time bound (9) of Proposition 1,

$$\mathbf{P}(\sigma_n < T_n) \leq \mathbf{P}(w_n < \alpha_n) + \mathbf{P}(\{\sigma_n < T_n\} \cap \{w_n \ge \alpha_n\})$$

$$\leq CN_n^2(\alpha_n\beta_n^{-1})^{d-1} + C(N_nr_nT_n + N_n^2r_n\alpha_n^{-1})$$

$$\leq C(N_nr_nT_n + N_n^2(r_n\beta_n^{-1})^{(d-1)/d}).$$

The claim of Proposition 4 follows from Borel-Cantelli, using (14).

Finally, putting together (16), (15) of Proposition 2, (17) of Proposition 3 and (18)/(19) of Proposition 4 we get: Almost surely, for any $F \in \mathcal{C}_0(\mathcal{C})$,

$$\lim_{n \to \infty} \mathbf{E}_{\varpi} \left(F(T_n^{-1/2} X_{n,1}(T_n \cdot)) \right) = \mathbf{E} \left(F(W(\cdot)) \right)$$

which concludes the proof of Theorem 3.

References

- [1] P. Billingsley: Convergence of Probability Measures. Wiley, New York (1968)
- [2] C. Boldrighini, L.A. Bunimovich, Y.G. Sinai: On the Boltzmann equation for the Lorentz gas. J. Stat. Phys. 32: 477-501 (1983)
- [3] P. Caputo, J. Salez: Random walk on sparse random digraphs. Probab. Theory Relat. Fields 170: 933-960 (2018)
- [4] G. Gallavotti: Rigorous theory of the Boltzmann equation in the Lorentz gas. Nota Interna Univ di Roma 358 (1970)
- [5] G. Gallavotti: Statistical Mechanics. A Short Treatise. Theoretical and Mathematical Physics Series. Springer, Berlin (1999)
- [6] C. Lutsko, B. Tóth: Invariance principle for the random Lorentz gas beyond the Boltzmann-Grad limit. Commun. Math. Phys. 379 589-632 (2020)
- [7] C. Lutsko, B. Tóth: Invariance principle for the random Wind-Tree Process. Ann. H. Poincaré 22: 3357-3389 (2021)
- [8] C. Lutsko, B. Tóth: Diffusion of the random Lorentz process in a magnetic field. arXiv:2411.03984 (2024)
- [9] H. Spohn: The Lorentz process converges to a random flight process. Comm. Math. Phys. 60: 277-290 (1978)
- [10] H. Spohn: Large Scale Dynamics of Interacting Particles Springer, Berlin-Heidelberg, 1991
- [11] C. Stone: Weak convergence of stochastic processes defined on a semi-infinite time interval. Proc. Amer. Math. Soc. 14: 694-696 (1963)
- [12] W. Whitt: Weak convergence of probability measures on the function space $C[0, \infty)$ Ann. Math. Statist. **41**: 939-944 (1970)

AUTHOR'S ADDRESS: Alfréd Rényi Institute of Mathematics Reáltanoda utca 13-14 Budapest 1053, Hungary toth.balint@renyi.hu