

Polynomial time sampling from log-smooth distributions in fixed dimension under semi-log-concavity of the forward diffusion with application to strongly dissipative distributions

Adrien Vacher¹, Omar Chehab¹, Anna Korba¹

¹CREST, ENSAE, IP Paris

January 3, 2025

Abstract

In this article we provide a stochastic sampling algorithm with polynomial complexity in fixed dimension that leverages the recent advances on diffusion models where it is shown that under mild conditions, sampling can be achieved via an accurate estimation of intermediate scores across the marginals $(p_t)_{t \geq 0}$ of the standard Ornstein-Uhlenbeck process started at the density we wish to sample from. The heart of our method consists into approaching these scores via a computationally cheap estimator and relating the variance of this estimator to the smoothness properties of the forward process. Under the assumption that the density to sample from is L -log-smooth and that the forward process is semi-log-concave: $-\nabla^2 \log(p_t) \succeq -\beta I_d$ for some $\beta \geq 0$, we prove that our algorithm achieves an expected ϵ error in KL divergence in $O(d^7 L^{d+2} \epsilon^{-2(d+3)} (L + \beta)^2 d^{2(d+1)})$ time. In particular, our result allows to fully transfer the problem of sampling from a log-smooth distribution into a regularity estimate problem. As an application, we derive an exponential complexity improvement for the problem of sampling from a L -log-smooth distribution that is α -strongly log-concave distribution outside some ball of radius R : after proving that such distributions verify the semi-log-concavity assumption, a result which might be of independent interest, we recover a $\text{poly}(R, L, \alpha^{-1}, \epsilon^{-1})$ complexity in fixed dimension which exponentially improves upon the previously known $\text{poly}(e^{RL^2}, L, \alpha^{-1}, \log(\epsilon^{-1}))$ complexity in the low precision regime.

1 Introduction

Given some potential $V : \mathbb{R}^d \mapsto \mathbb{R}$ such that $\int e^{-V(x)} dx < \infty$, the sampling problem consists into obtaining a sample from some distribution p such that p is ϵ -close to $\mu \propto e^{-V}$ with respect to some divergence while maintaining the complexity, a.k.a. the number of queries to V and possibly to its derivatives, as low as possible with respect to ϵ and to the other parameters of the problem. As in Euclidean optimization, appealing complexity bounds can be obtained when μ is assumed to be L -log-smooth and α -strongly log-concave or equivalently, when the potential V is assumed to have an L -Lispchitz gradient and to be α -strongly convex. Namely, denoting the condition number $\kappa = L/\alpha$, it has been recently shown that the Underdamped Langevin Algorithm (ULA) [14, 49], one of the most popular sampling algorithm, could provide a sample $X \sim p$ such that $\text{KL}(p, \mu) \lesssim \epsilon$ in $\tilde{O}(d^{1/3} \kappa \epsilon^{-2/3})$ time under minor modifications [43, 20, 8]. Similarly, it was shown that the Metropolis Adjusted Langevin Algorithm (MALA) [48, 4], a debiased version of Langevin, could achieve a complexity of $\tilde{O}(d^{1/2} \kappa \log^4(1/\epsilon))$ for a well-chosen warm start [2]; in both cases, the dependence is sub-linear in the dimension. However, because strong log-concavity implies unimodality, this framework is far from satisfactory when it comes to obtaining guarantees for real-world applications, where the distributions to sample from can be highly multimodal [30, 17]. One natural relaxation is to confine multimodality within some fixed ball, also known as the *strong dissipativity* assumption [13, 31, 32, 19] where we assume that the distribution is strongly log-concave outside this ball. Note that this assumption is much more realistic as it encompasses several types of Gaussian mixtures and other popular Bayesian mixture models [34, 33, 19]. Yet, this extra flexibility currently comes at a heavy price as known complexity bounds now degrade exponentially with the size of the non-convex region: if we assume that μ is α -strongly log-concave outside the centered ball of radius R , ULA achieves a complexity of $\tilde{O}(e^{32RL^2} \kappa^2 \frac{d}{\epsilon^2})$ and MALA achieves a complexity $O(\frac{e^{40LR^2}}{\alpha} \kappa^{3/2} d^{1/2} (d \log(\kappa) + \log(1/\epsilon))^{3/2})$ [31]. Note that even if those are indeed worst case bounds, it

has been well observed in practice that the aforementioned samplers suffer from *metastability*: as the distance between the modes grows, samples get stuck in one local mode, leaving the other ones unexplored [44, 26] (see also Chehab et al. [9] for formal lower bounds in continuous time). While there is no hope for a general polynomial bound as it was shown in Lee et al. [29] that even sampling from a 2-modes distribution could require up to 2^{cd} queries to V with c a universal constant, sampling from a multimodal density in low dimension with provable polynomial bounds is still an open problem. This raises the following informal question:

”In fixed dimension, can we sample from a multimodal distribution in polynomial time?”

1.1 Related work

Sampling from multimodal distributions has been an active area of research for several decades now [37, 15, 38]. Indeed, it has long been observed that Monte Carlo Markov Chain (MCMC) based on local exploration of the space, such as ULA and MALA, are prone to *metastability* and remain stuck in local modes [47, 21, 42]. We review in this paragraph the main alternatives to Langevin based samplers that have been developed and the guarantees they provide.

Histogram methods Given some compact set $S \subset \mathbb{R}^d$, histogram methods build a regular n -grid over S and sample a point s_i of the grid according to the probability $e^{-s_i} / \sum_{i=1}^n e^{-V(s_i)}$. If V is assumed to be α -strongly convex and L -smooth for instance, then, if S is taken to be a ball of radius r centered at x^* the global minimum of V , the typical error of this method scales in

$$O(L(r/n)^{1/d}r + e^{-\alpha r^2}),$$

thus inducing a typical complexity of $\tilde{O}((L/\epsilon)^d)$ which is indeed polynomial in fixed dimension. However, if instead of being centered at the global minimum x^* , the ball is centered at some point x , the error remains $O(1)$ as long as $r \leq \|x - x^*\|$ as only the tail of μ is now discretized. When V is smooth and strongly convex, x^* can be approximated up to arbitrary precision yet in the case where V is assumed to be strongly convex only outside of some *unknown* ball, one cannot localize the global minimum x^* even in one dimension. In other terms, unless we have prior knowledge on the localization of the mass of μ , histogram methods can induce an arbitrary large error.

Importance Sampling (IS) Instead of focusing on the problem of sampling from μ , IS focuses on the related, yet slightly simpler problem of approximation of integrals under μ : given some bounded function f , IS methods approximate the integral

$$\mu(f) := \int_x f(x) d\mu(x).$$

For any W such that $\int e^{-W(x)} dx < +\infty$, denoting $\nu \propto e^{-W}$, the integral above can be re-written as an expectation under ν as $\mu(f) = \mathbb{E}_{Y \sim \nu}[f(Y) \frac{d\mu}{d\nu}(Y)]$. In particular, if we can access n i.i.d. samples $(x_i)_{1 \leq i \leq n}$ from the proposal ν , this quantity can be approximated as $\hat{\mu}(f) = \frac{\sum_{i=1}^n f(x_i) e^{-V(x_i)+W(x_i)}}{\sum_{i=1}^n e^{-V(x_i)+W(x_i)}}$. It was shown in Robins et al. [41] that the average quadratic error of this estimator scales as

$$\sup_{\|f\| \leq 1} \mathbb{E}[(\hat{\mu}(f) - \mu(f))^2] = O\left(\frac{\chi^2(\mu, \nu)}{n}\right).$$

Yet, unless ν is carefully chosen, this bound may blow up to infinity even for simple densities: if $\mu \sim \mathcal{N}(\mu_1, \sigma_1^2)$ and $\nu \sim \mathcal{N}(\mu_2, \sigma_2^2)$, it holds that

$$\chi^2(\mu, \nu) = \frac{1}{2} \left(\frac{\sigma_2^2}{\sigma_1 \sqrt{2\sigma_2^2 - \sigma_1^2}} e^{\frac{(\mu_1 - \mu_2)^2}{2\sigma_2^2 - \sigma_1^2}} - 1 \right).$$

In particular, not only this bound blows up when the variance of ν is not at least half the variance of μ but also, this bound always degrades exponentially with the quadratic distance between the mode of the proposal and the mode of the target μ . Hence, even in this one dimensional strongly log-concave toy example, IS requires a precise *a priori* knowledge of μ to tailor a good proposal ν (and also requires to be able to sample from ν).

Tempering heuristics Instead of directly sampling from $\mu \propto e^{-V}$, we start to sample from a flattened version of μ given by $\mu^\beta \propto e^{-\beta V}$ for β small and we progressively increase β to 1. When μ is assumed to be a finite mixture of the same shifted α -strongly log-concave and L -log-smooth distribution, Lee et al. [29] proved that for a well-chosen (stochastic) sequence of flattened distributions μ^{β_i} , sampling up to precision ϵ in KL can be achieved in $\text{poly}(L, \alpha^{-1}w_{\min}, d, \epsilon^{-1}, R)$ time with R the diameter of the shifts and w_{\min} the minimum weight in the mixture. Yet, despite the strong guarantees it offers, this framework excludes many practical cases such as mixtures of Gaussians with different variances for instance: in that case, the tempering strategy even fails in low dimension as it attributes biased weights to the different modes.

Reverse Diffusion (RD) methods As we shall review in details in Section 2, recent works show that, under milder and milder assumptions [12, 10, 16, 6], the problem of sampling can be transferred into the problem of approximating the score in L^2 distance along the standard Ornstein-Uhlenbeck (OU) process initialized at the targeted distribution μ . Over the past year, several works sought to leverage this result and focused on the task of estimating these intermediate scores given an oracle access to V and eventually to its gradient. Their guarantees can be summarized as follows:

- in Ding et al. [18], denoting $\pi^{\eta, \Sigma}$ the d -dimensional Gaussian density with mean η and covariance matrix Σ , the authors proved that if there existed η, Σ such that $d\mu/d\pi^{\eta, \Sigma}$ is globally L -Lipschitz and strictly bounded from below by ξ , then their algorithm achieved a complexity $\text{poly}(L, \xi^{-1}, d, \epsilon^{-1})$. We show in Appendix C that even for the simple mixture $\mu = \frac{1}{2}\mathcal{N}(1, 1) + \frac{1}{2}\mathcal{N}(-1, 1)$, the previous conditions do not hold.
- in Huang et al. [25], it is shown that under the assumption that the OU process remains L -log-smooth along the trajectory, denoting $m_2(\mu) = \int \|x\|^2 d\mu(x)$ the second moment of μ , their algorithm could reach a complexity of $e^{O(L^3 \log^3((Ld+m_2(\mu))/\epsilon))}$.
- in He et al. [22], it is shown that under a sub-quadratic growth assumption on V (which is weaker than log-smoothness), *i.e.* there exists $L > 0$ such that $V(x) - V(x^*) \leq L\|x - x^*\|^2$, with x^* a global minimizer of V , their algorithm could reach at best a complexity of $O(L^{d/2}\epsilon^{-d}e^{(L\|x^*\|^2 + \|x_N\|^2)})$, where x_N is the sample output by their algorithm. Under the reasonable (and desirable) assumption that $\mathbb{E}[\|x_N\|^2] \asymp m_2(\mu)$, Jensen inequality yields that their complexity scales at best as $O(L^{d/2}\epsilon^{-d}e^{(L\|x^*\|^2 + m_2(\mu))})$.

Hence, we observe that previous approaches based on reverse diffusion suffer at best from exponential complexity either with respect to L or to $m_2(\mu)$.

1.2 Our contributions

Polynomial time sampling under semi-log-concavity of the Ornstein-Uhlenbeck (OU) process As in the aforementioned work, we rely on the recent advances on reverse diffusion and focus on the task of estimating the intermediate scores $\nabla \log(p_t)$ with $(p_t)_{t \geq 0}$ the marginals of the standard OU process initialized at μ . Yet, unlike all previous works, we provide fully polynomial time guarantees in fixed dimension when μ is assumed to be log-smooth and when the forward process has semi-log-concave marginals.

Theorem 1 (Main result, informal) *Denoting p_t^V the density at time t of the forward OU process started at $\mu \propto e^{-V}$, then under the assumption that V is C^2 and L -smooth, that μ has finite second moment $m_2(\mu)$ and that $-\nabla^2 \log(p_t^V), -\nabla^2 \log(p_t^{2V}) \succeq -\beta I_d$ uniformly over \mathbb{R}^d , then there exists a stochastic algorithm that outputs a sample $X \sim \hat{p}$ such that $\mathbb{E}[\text{KL}(\mu, \hat{p})] \lesssim \epsilon$ in $O(d^7 L^{d+2} \epsilon^{-2(d+3)} (L + \beta)^2 (m_2(\mu) + d)^{2(d+1)})$ time where \lesssim hides a universal constant as well as log quantities in $d, \epsilon^{-1}, L, \beta, m_2(\mu)$.*

Hence, when the dimension d is fixed, we obtain a polynomial time guarantee. Note that this result pushes one step further the works of Chen et al. [12, 10], Conforti et al. [16], Benton et al. [6] who showed that the sampling problem could be transferred into a score approximation problem yet leaving this crucial question open. Using our estimator of the intermediate scores, we show that in fixed dimension, sampling can be framed as a regularity estimate problem: for any class of smooth potentials V such that $\nabla^2 \log(p_t^V), \nabla^2 \log(p_t^{2V})$ are bounded from above, we can sample from $\mu \propto e^{-V}$ in polynomial time.

Polynomial time sampling for smooth and strongly dissipative distributions Deriving classes of potentials V that allow quantitative bounds on $\nabla^2 \log(p_t^V)$ is an area of research of its own as it allows to recover transport maps from the standard Gaussian to measures of the form $\mu \propto e^{-V}$ with quantitative Lipschitz bounds. Such bounds were originally proven for the class of smooth and strongly convex potentials [27] and later derived for the classes of compactly supported distributions and mixtures of Gaussians with the same variance respectively [35]. In this work, we show that such bounds still hold for smooth potentials that are only assumed to be strongly convex outside some ball. Using Theorem 1, this allows us to recover polynomial complexity bounds for sampling from this class of distributions.

Theorem 2 (Application to smooth and strongly dissipative potentials) *Assume that V is C^2 , L -smooth and α -strongly convex outside some ball of radius R , then there exists a stochastic algorithm that that returns a sample $X \sim \hat{p}$ such that $\mathbb{E}[\text{KL}(\mu, \hat{p})] \lesssim \epsilon$ in $O(d^7 L^{d+8} \epsilon^{-2(d+3)} (R^2 \kappa^2 + \frac{d}{\alpha})^{2(d+4)})$ time.*

As previously mentioned, this bound exponentially improves upon the $O(e^{32RL^2} d \kappa^2 \epsilon^{-2})$ complexity of ULA [31] and the $O(\frac{e^{40LR^2}}{\alpha} \kappa^{3/2} d^{1/2} (d \log(\kappa) + \log(1/\epsilon))^{3/2})$ complexity of MALA [31] in the low dimension and low precision regime. We also show in Section 4 that our results exponentially improve upon Huang et al. [25] and He et al. [22].

1.3 Technical overview

In Section 2, we provide precise details about the reverse diffusion framework and formally state the result of Chen et al. [10]. Then we show how we derived our estimator of the intermediate scores and notably how it differs from the ones derived in [24] and [22]. In Section 3, we prove our main result. This proof can be decomposed in two major steps: first, we provide a non-asymptotic quadratic error for our estimator which is generic and does not rely on any particular assumption on μ . We observe that, once this quadratic error is integrated with respect to p_t^V , our upper bound scales as:

$$\int_{\mathbb{R}^d} (1 + \|\nabla \log(p_t^V)\|^2(z) + \|\nabla \log(p_t^{2V})\|^2(z)) \frac{p_t^{2V}(z)}{p_t^V(z)} dz.$$

Under the assumptions that $\nabla \log(p_t^V)$ and $\nabla \log(p_t^{2V})$ are Lipschitz, the previous quantity can be controlled by a bound that only depends on the zeroth and second order moments of the ratio $\Phi_t = \frac{p_t^{2V}}{p_t^V}$. Hence a first natural assumption is to assume that both p_t^V, p_t^{2V} are log-smooth at $t = 0$.

Assumption 1 (Smoothness) *The potential V is $C^2(\mathbb{R}^d)$ and has an L -Lipschitz gradient.*

We then rely on the results of Mikulincer and Shenfeld [36] who show that under Assumption 1, one can control the difference of Hessians of log-densities as

$$\nabla^2 \log(\pi) - \nabla^2 \log(p_t^V) \preceq \frac{e^{-2t}(L-1)}{(1-e^{-2t})(L-1)+1} I_d, \quad (1)$$

where π refers to the density of the standard Gaussian. This already provides the following upper-bound

$$\nabla^2 \log(p_t^V) \succeq -\frac{L}{L(1-e^{-2t})+e^{-2t}} I_d \succeq -(L+1)I_d. \quad (2)$$

Hence, in order to ensure $\nabla \log(p_t^V), \nabla \log(p_t^{2V})$ are Lipschitz, it suffices to assume the converse inequality *i.e.* the semi-log-concavity of the forward processes.

Assumption 2 (Semi-log-concavity of the forward processes) *There exists $\beta \geq 0$ such that $-\nabla^2 \log(p_t^V) \succeq -\beta I_d$ and $-\nabla^2 \log(p_t^{2V}) \succeq -\beta I_d$ uniformly over \mathbb{R}^d .*

We refer to this assumption as semi-log-concavity, in reference with semi-convexity as common in the Optimal Transport literature [3] and semi-log-concavity in the sampling literature [35]; note that this is significantly weaker than classical log-concavity which implies the tighter lower-bound $\beta \leq 0$. Hence, under Assumptions 1-2, it suffices to control the zeroth and the second order moments of Φ_t to recover a quantitative error bound. We shall again rely on the previous bound Eq. 1 that we combine with Grönwall's lemma to recover a $O(e^{td} L^d)$ bound on the zeroth order moment of Φ_t . Crucially, note that we recover a polynomial dependence with respect to L instead of an exponential dependence because whenever $t > 0$, the bound Eq. 1 remains bounded with respect to L . The second moment being more

delicate to handle, we briefly explain the main additional ingredients it requires and leave its complete proof in Appendix. We conclude the section by formally stating our main theorem.

In Section 4, we apply our result to the class of smooth, strongly convex potentials outside some ball and make the following assumption.

Assumption 3 (Strong Dissipativity) *The potential V is α -strongly convex outside some ball of radius R centered at $m \in \mathbb{R}^d$ denoted $B_R(m)$.*

We prove that under Assumptions 1-3, Assumption 2 is met with $\beta \lesssim L^2(R^2L^2\alpha^{-2} + d\alpha^{-1})$. Our proof relies on a bound of the variance of log-smooth and strongly log-concave measures outside some ball that we believe to be a novel one, as it is both polynomial and does not depend on the gradient of V and the center of the ball. The upper bound on β combined with our Theorem 1 allows us to recover a polynomial complexity in fixed dimension (Theorem 2) for the task of sampling from log-smooth, strongly dissipative distributions. We conclude the section by comparing in details our complexity with the ones yielded by previous works under strong dissipativity.

2 Preliminaries

In this section, we introduce the reverse diffusion framework and review the main existing bounds on how the resulting sampling error is affected by the error on the intermediate scores. Then, we show that these intermediate scores can be re-written as a ratio of expectations and detail how this re-writing differs from previous works.

2.1 Reverse diffusion: from sampling to score estimation

One of the core tool for sampling from $\mu \propto e^{-V}$ is the Langevin diffusion process

$$dX_t = -\nabla V(X_t)dt + \sqrt{2}dB_t, \quad (3)$$

where B_t is the standard Brownian motion. When discretized with a Euler-Maruyama scheme, this forward equation becomes the well-known Underdamped Langevin Algorithm (ULA, [40]) which is readily implementable whenever we can access ∇V . One of the fundamental question to determine the quality of the sample output by ULA is the speed of convergence of the process Eq. 3 towards the equilibrium μ . When μ is α -strongly log-concave, and more generally when μ verifies an α -log-Sobolev inequality [5], exponential convergence in KL occurs at an α rate [46]. Yet, if μ is multimodal, strong log-concavity no longer holds and the log-sobolev constant α is either no longer defined either scales exponentially with the distance between the modes [11, 9]. This exponential dependence is in correspondence with a phenomenon known as *metastability* where the process X_t gets stuck in a local mode of μ .

Reverse diffusion methods emerged as an alternative to Langevin based samplers in order to overcome metastability and were first introduced to the ML community in Song and Ermon [45]. Instead of targeting μ in Eq. 3, one targets the standard Gaussian measure with a Langevin diffusion initialized at μ . We thus obtain the so-called forward process

$$\begin{cases} dX_t = -X_t dt + dB_t, \\ X_0 \sim \mu, \end{cases} \quad (4)$$

which corresponds to the standard Ornstein-Uhlenbeck (OU) process initialized at μ . Note that since we are now targeting the standard Gaussian which is 1-strongly log-concave, the resulting process converges exponentially fast to the equilibrium. In order to sample from μ we consider the semi-discretized *backward process*: given a horizon T that we discretize as $0 = t_0 \leq t_1 \leq \dots \leq t_{N-1} \leq t_N = T$, the discretized process reads

$$\begin{cases} dY_t = Y_t dt + 2\nabla \log(p_{t_{N-k}})(Y_t)dt + \sqrt{2}dB_t, & t \in]T - t_{N-k}, T - t_{N-(k+1)}], \\ Y_0 \sim p_T, \end{cases} \quad (5)$$

where p_t is the distribution of the forward process Eq. 4 at time t . Note that this reverse process cannot be readily implemented for two reasons: first, it requires the knowledge of the intermediate scores $\nabla \log(p_{t_k})$ which are not available in closed form. Second, it requires sampling from the distribution p_T . Nevertheless, if one can access a proxy s_{t_k} of the scores $\nabla \log(p_{t_k})$, and considering T large enough so that $p_T \approx \pi$, we can implement instead

$$\begin{cases} dY_t = Y_t dt + 2s_{t_k}(Y_k)dt + \sqrt{2}dB_t, & t \in]T - t_{N-k}, T - t_{N-(k+1)}], \\ Y_0 \sim \pi, \end{cases} \quad (6)$$

where all iterations can be solved in closed form. Because the forward process Eq. 4 converges exponentially fast, we can expect the initialization error $Y_0 \sim \pi$ instead of $Y_0 \sim p_T$ to be small for a large time T . Furthermore, if the proxies s_{t_k} are sufficiently accurate, one can expect that the process output by the approximate scheme Eq. 6 has a distribution that is close to the target μ . Over the past three years, several works provided quantitative bounds of the error induced by the discretization, the use of an approximate score and the initialization error with respect to different divergences and under various assumptions [7, 28, 12, 10, 16]. Yet, we shall rely exclusively on the following theorem as it is the most suited to our framework.

Theorem 3 (Chen et al. [10]) *Assume that μ is L -log-smooth and has finite second order moment $m_2(\mu) = \int \|x\|^2 d\mu(x)$ and choose the following (backward) discretization: $t_{k-1} = t_k - c \min(\max(t_k, 1/L), 1)$ for $k \geq 2$ with $c = (\log(L) + T)/N$. Then, if c verifies $c \leq \frac{1}{2d}$, denoting p the distribution of the sample Y_T output by Eq. 6, it holds that*

$$\text{KL}(\mu, p) \lesssim (d + m_2(\mu))e^{-T} + \sum_{k=1}^N (t_k - t_{k-1}) \|\nabla \log(p_{t_k}) - s_{t_k}\|_{L^2(p_{t_k})}^2 + \frac{d^2(\log(L) + T)^2}{N},$$

where \lesssim hides a universal constant.

Remark 4 *In its original statement, Chen et al. [10, Theorem 5] assumes that*

$$\frac{1}{T} \sum_{k=1}^N (t_k - t_{k-1}) \|\nabla \log(p_{t_k}) - s_{t_k}\|_{L^2(p_{t_k})}^2 \leq \epsilon^2,$$

and directly provides the upper bound with the $T\epsilon^2$ term instead. Yet, an inspection of their Proposition 8 shows that we can safely use $\sum_{k=1}^N (t_k - t_{k-1}) \|\nabla \log(p_{t_k}) - s_{t_k}\|_{L^2(p_{t_k})}^2$. Similarly, it assumes that c should be lower than $\frac{1}{Kd}$ with K some universal constant. Yet a careful inspection of Lemma 13 and Lemma 17 shows that we can take $K = 2$.

The previous theorem shows that under mild assumptions that notably allow for multimodality, the problem of sampling from μ can be transferred into a score approximation problem along the forward process. In the next subsection, we build an estimator for these intermediate scores that is tractable given the knowledge of the unnormalized density $\mu \propto e^{-V}$, and show how it differs from previous works.

2.2 Construction of our estimator

The key observation to derive an estimator of the intermediate scores is that the forward process Eq. 4 is nearly available in closed form. Indeed, Eq. 4 integrates to

$$X_t = e^{-t} X_0 + B_{1-e^{-2t}},$$

where B_s is the standard d -dimensional Brownian motion evaluated at time s . In particular, conditionally on X_0 , X_t has a normal distribution $X_t | X_0 \sim \mathcal{N}(e^{-t} X_0, (1 - e^{-2t}) I_d)$. By Bayes rule, the distribution p_t^V of the forward process X_t initialized at $\mu \propto e^{-V}$ thus integrates to

$$p_t^V(z) = \frac{1}{(2\pi(1 - e^{-2t}))^{d/2} Z_V} \int e^{-\frac{\|e^{-t}x - z\|^2}{2(1 - e^{-2t})}} e^{-V(x)} dx,$$

with Z_V the normalizing constant $Z_V = \int e^{-V(x)} dx$. We thus recover that the score of the forward process re-writes as

$$\nabla \log(p_t^V)(z) = \frac{\int (z - e^{-t}x) e^{-\frac{\|e^{-t}x - z\|^2}{2(1 - e^{-2t})}} e^{-V(x)} dx}{(1 - e^{-2t}) \int e^{-\frac{\|e^{-t}x - z\|^2}{2(1 - e^{-2t})}} e^{-V(x)} dx} = \mathbb{E}_{Y \sim q_{t,z}} \left[\frac{z - e^{-t}Y}{1 - e^{-2t}} \right],$$

where we denoted $q_{t,z}(x) \propto e^{-\frac{\|e^{-t}x - z\|^2}{2(1 - e^{-2t})}} e^{-V(x)}$. As noted in Huang et al. [24], Huang et al. [25] and He et al. [22], if samples $y_i \sim q_{t,z}$ are available, one can recover an efficient empirical estimator of the score given by $\frac{-1}{n(1 - e^{-2t})} \sum_{i=1}^n y_i$ ready to be plugged in the reverse diffusion process Eq. 6. However, by doing so, the problem of sampling from e^{-V} is simply transferred to the problem of sampling from $q_{t,z}$,

which becomes almost as hard as t grows. In Huang et al. [24], the intermediate sampling problems are solved using the ULA algorithm and in He et al. [22], the authors use a rejection sampling scheme which is similar in spirit to importance sampling. Hence, it is not surprising that their final upper-bounds do not improve over standard methods as they employ them to solve intermediate problems that are progressively as hard as the original problem.

Instead, we observe that after applying the change of variable $y_t = ze^{-t} - x$ both on the numerator and on the denominator, we obtain a score formula which re-writes as a *ratio* of expectations under standard Gaussians

$$\nabla \log(p_t^V)(z) = \frac{-1}{1 - e^{-2t}} \frac{\mathbb{E}[Y_t e^{-V(e^t(z - Y_t))}]}{\mathbb{E}[e^{-V(e^t(z - Y_t))}]}, \quad (7)$$

with $Y_t \sim \mathcal{N}(0, (1 - e^{-2t})I_d)$. Unlike the previous identity, which involves an expectation under $q_{t,z}$, the expectations above are computationally cheap as we can easily sample from Y_t . While conventional statistical wisdom would suggest both the numerator and the denominator to be estimated with independent samples, we voluntarily choose to correlate them and implement instead

$$\hat{s}_{t,n}(z) := \frac{-1}{1 - e^{-2t}} \frac{\sum_{i=1}^n y_i e^{-V(e^t(z - y_i))}}{\sum_{i=1}^n e^{-V(e^t(z - y_i))}}, \quad (8)$$

where the y_i are independent Gaussians such that $y_i \sim \mathcal{N}(0, (1 - e^{-2t})I_d)$; we shall refer to this estimator as *self-normalized* as common in the sampling literature [1]. Depending on the locations of the modes of V we expect that this estimator behaves reasonably well in some region of the space and poorly in others. Yet, the key property of self-normalized estimators is that they remain nearly bounded: in our case, it holds uniformly in z, t that:

$$\|\hat{s}_{t,n}(z)\| \leq \frac{\max_i \|y_i\|}{1 - e^{-2t}} \sim \sqrt{d \log(n)}.$$

In particular, even in the regions where it behaves poorly, $\hat{s}_{t,n}(z)$ nevertheless remains bounded. Quantitatively, as recalled in the previous subsection, it suffices to control the average error of $\hat{s}_{t,n}$ in $L^2(p_t^V)$ error. In the next section, under the assumption that V is L -smooth and that the forward processes p_t^V, p_t^{2V} are semi-log-concave a.k.a. they verify $\nabla^2 \log(p_t^V), \nabla^2 \log(p_t^{2V}) \preceq \beta I_d$ for some $\beta \geq 0$, we shall prove that

$$\mathbb{E} \left[\int \|\hat{s}_{t,n}(z) - \nabla \log(p_t^V)(z)\|^2 dp_t^V(z) \right] \lesssim \frac{e^{2t(d+1)} L^{d+2} (L + \beta)^2 m_2(\mu)}{n(1 - e^{-2t})^2},$$

where \lesssim hides a universal constant and log factors.

Remark 5 *The estimator used in Ding et al. [18] is very similar to ours and simply differs by the fact that the authors discretize a Föllmer flow which corresponds to a reparametrized in time version of the standard OU process and where an arbitrary Gaussian may be targeted. The authors obtain strong guarantees yet under assumptions that exclude almost all distributions (see Appendix C). As a consequence, the authors avoid most of the technical difficulties we were confronted with.*

3 Proof of the main result

In this section, we prove a non-asymptotic bound on the variance of the estimator presented above. First, we derive a general non-asymptotic bound that does not rely on any particular assumptions. Using the estimates provided in Mikulincer and Shenfeld [36] and our additional semi-log-concavity assumption, this upper bound becomes fully explicit in the zeroth and second order moments of the ratio $\Phi_t = \frac{p_t^{2V}}{p_t^V}$. We then provide upper-bounds for these moments that again heavily leverage the estimates of Mikulincer and Shenfeld [36]. Finally, we combine our results with those of Chen et al. [10] to state our main theorem.

3.1 Non asymptotical upper-bound of the variance

Before studying our ratio estimator defined in Eq. 8, we start by separately upper-bounding the variances of its numerator and denominator as a function of the forward process.

Proposition 6 (Variance of estimators) *Let y_1, \dots, y_n i.i.d. distributed as $\mathcal{N}(0, (1 - e^{-2t})I_d)$ and denote $\hat{N}(z)$ and $\hat{D}(z)$ the numerator and denominator of the estimator defined in Eq. 8, we have:*

$$\begin{aligned} \mathbb{E}[\|\hat{N}(z) - N(z)\|^2] &\leq \frac{p_t^{2V} Z_{2V} e^{-td}}{n} \left(\Delta \log(p_t^{2V})(z) - \frac{\Delta \log(\pi)(z)}{1 - e^{-2t}} + \|\nabla \log(p_t^{2V})\|^2(z) \right), \\ \mathbb{E}[\|\hat{D}(z) - D(z)\|^2] &\leq \frac{p_t^{2V}(z) Z_{2V} e^{-td}}{n}. \end{aligned}$$

The proof is deferred to Appendix A.2. We can now derive a non asymptotic quadratic error for our estimator.

Proposition 7 (Non-asymptotic bound of the quadratic error) *For all $z \in \mathbb{R}^d$, n and $t > 0$, it holds that*

$$\mathbb{E} \left[\|\hat{s}_{t,n}(z) - \nabla \log(p_t^V)(z)\|^2 \right] \leq \frac{32e^2(d + \log(n))}{n(1 - e^{-2t})} \left(1 + \theta_t(z) \frac{p_t^{2V}(z) Z_{2V} e^{td}}{(p_t^V(z) Z_V)^2} \right),$$

with $\theta_t(z) = \Delta \log(p_t^{2V})(z) - \frac{\Delta \log(\pi)(z)}{1 - e^{-2t}} + \|\nabla \log(p_t^V)(z)\|^2 + \|\nabla \log(p_t^{2V})(z)\|^2 + 1$.

The complete proof is left to Appendix A.3 yet we briefly sketch the main arguments. We split the expectation on the event A where the empirical denominator \hat{D} (resp. the empirical numerator \hat{N}) is not too small (resp. not too large) with respect to its expectation D (resp. $\|N\|$) and on the complementary \bar{A} . Across A , we use a Taylor expansion of order 2 to make the variance of the numerator and of the denominator appear and we then use the previous proposition. Conversely, the estimator remains almost bounded on \bar{A} . Hence, to control the remaining error, we must simply upper-bound $\mathbb{P}(\bar{A})$ together with a Hölder inequality with a well-chosen exponent. We use Tchebychev inequality to upper-bound $\mathbb{P}(\bar{A})$ and make the variances of the numerator and of the denominator appear again, which concludes the proof.

Remark 8 *Generic results on self-normalized estimators were derived in Agapiou et al. [1, Theorem 2.3]. Yet, it would have involved an additional $1/(p_t^V)^4$ term which would have caused the integrated error $\int \delta(z)^2 dp_t^V(z)$ to diverge.*

Hence, we observe that the average quadratic error is related to the properties of the forward process Eq. 4 started at $\mu \propto e^{-V}$ and $\mu^2 \propto e^{-2V}$ respectively. Under our semi-log-concavity assumption, *i.e.* there exists $\beta \geq 0$ such that $\nabla^2 \log(p_t^{2V})$ is uniformly bounded from above by βI_d , the Laplacian terms can be upper-bounded as

$$\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1 - e^{-2t}} \leq d \left(\beta + \frac{1}{1 - e^{-2t}} \right). \quad (9)$$

In order to handle the gradient terms, we need an additional bound from below on the Hessian. Fortunately, such bounds were derived in Mikulincer and Shenfeld [36] when the potential V is assumed to be L -smooth.

Proposition 9 (Mikulincer and Shenfeld [36]) *Under the assumption that μ is L -log smooth and C^2 , it holds for all $z \in \mathbb{R}^d$ that*

$$\nabla^2 \log(\pi)(z) - \nabla^2 \log(p_t^V)(z) \preceq \frac{(L - 1)e^{-2t}}{(1 - e^{-2t})(L - 1) + 1} I_d. \quad (10)$$

The proof is left to Appendix A.4 and is mainly an identification of terms. As a result, the following lower bound holds:

$$\nabla^2 \log(p_t^V) \succeq -(L + 1)I_d.$$

In particular, under the assumption that both p_t^V, p_t^{2V} are semi-log-concave, we recover that

$$\begin{cases} \|\nabla \log(p_t^V)(z)\| \leq (L + \beta + 1)\|z\| + \|\nabla \log(p_t^V)(0)\|, \\ \|\nabla \log(p_t^{2V})(z)\| \leq (2L + \beta + 1)\|z\| + \|\nabla \log(p_t^{2V})(0)\|. \end{cases} \quad (11)$$

Hence, in order to control the average quadratic error integrated w.r.t. p_t^V , it suffices to control the zeroth and the second order moments of the ratio $\Phi_t(z) := \frac{p_t^{2V}}{p_t^V}$. This will be the object of the next paragraph.

3.2 Estimates on the moments of the ratio

In this subsection, we provide our upper bounds on the moments of Φ_t needed to obtain a interpretable upper bound from Proposition 7, starting with a bound on its zeroth moment. By slight abuse of notations, we shall denote $m_i(\Phi_t) = \int \|x\|^i \Phi_t(z) dz$ the i -th moment of Φ_t .

Lemma 10 (Bound on the zeroth moment of the ratio) *Assume that $\mu \propto e^{-V}$ is L -log-smooth and C^2 . It holds that*

$$m_0(\Phi_t) \leq \frac{(Z_V)^2}{Z_{2V}} e^{td} (L(1 - e^{-2t}) + e^{-2t})^d.$$

The proof of Lemma 10 is provided in Appendix A.5 and relies on applying Grönwall's lemma for the evolution of the zeroth moment, using an intermediate upper bound given by Proposition 9. Crucially, note that this bound is only polynomial in L when d is fixed instead of exponential. As mentioned earlier, this is a consequence of the bound of Mikulincer and Shenfeld [36] that remains bounded with respect to L whenever $t > 0$.

There remains to bound the second moment of Φ_t . This part is more delicate and will require in fact to bound the first moment of Φ_t . The full proof is deferred to Appendix A.8 yet we state the key intermediate lemmas to achieve this result. First we show that the maximum of the ratio Φ_t decreases.

Lemma 11 (Decrease of the maximum of Φ) *The maximum of the ratio Φ_t decreases with t .*

The proof is left in Appendix and is simply a combination of Lemma 25 with the implicit function theorem. Then, the second key lemma is a control on the integrated gradient of the forward process at 0.

Lemma 12 (Upper-bound integrated gradient) *Let $\mu \propto e^{-V}$ be a L -log smooth measure. Denoting $\mu(0)$ the density of μ with respect to the Lebesgue measure at 0, it holds that*

$$\int_0^t \|\nabla \log(p_s^V)\|^2(0) ds \leq -\log(\mu(0)) + \frac{d}{2} \log\left(\frac{L}{2\pi}\right).$$

The proof is left in Appendix and is essentially the consequence of Lemma 25 deriving the evolution of Φ_t , combined with the estimate in Proposition 9 in addition with a general result on the maximum of a log-smooth density. These two lemmas allow to recover a bound on the first moment of Φ_t which in turns allows to bound the second moment of Φ_t .

Lemma 13 *Let μ be L -log-smooth, C^2 and with finite second moment $m_2(\mu)$. It holds that*

$$m_2(\Phi_t) \leq \frac{2e^{t(d+2)}(Z_V)^2}{Z_{2V}} (L(1 - e^{-2t}) + e^{-2t})^{d+2} \left[m_2(\mu) + d(L+1) + 2d \log\left(\frac{L\mu(0)^{-2/d}}{2\pi}\right) \right].$$

We have now all the ingredients to state and prove our main result.

3.3 Assembling the pieces

We first derive the average integrated squared error of our estimator $\hat{s}_{n,t}$.

Theorem 14 (Integrated average squared error of the estimator) *Under Assumptions 1-2 and the additional assumptions that $\|\nabla \log(p_t^V)(0)\|, \|\nabla \log(p_t^{2V})(0)\| \leq \xi$ for some $\xi \geq 0$ and that μ has finite second moment $m_2(\mu)$, it holds that*

$$\int \mathbb{E}[\|\hat{s}_{n,t}(z) - \nabla \log(p_t)(z)\|^2] dp_t^V(z) \lesssim \frac{e^{2t(d+1)}(d + \log(n))(L+1)^d}{n(1 - e^{-2t})^2} (K_1 + K_2),$$

where we defined

$$\begin{cases} K_1 = (L + \beta + 1)^2 (L + 1)^2 \left[m_2(\mu) + d(L + 1) + 2d \log\left(\frac{L\mu(0)^{-2/d}}{2\pi}\right) \right], \\ K_2 = d\beta + \xi^2 + 1. \end{cases} \quad (12)$$

The proof is left in Appendix. Combined with the result of Chen et al. [10], we recover a polynomial time bound in fixed dimension.

Corollary 15 (Polynomial time sampling) *Under the same assumptions as in Theorem 14, if we run algorithm Eq. 6 for an appropriate (explicit) set of parameters and with the stochastic score estimators \hat{s}_{n_k, t_k} defined in Eq. 8, then, denoting \hat{p} the stochastic distribution of the output Y_N , it holds that*

$$\begin{cases} \mathbb{E}[\text{KL}(\mu, \hat{p})] \lesssim \epsilon, \\ \sum_{i=1}^N n_k \lesssim d^7 L^{d+2} \epsilon^{-2(d+3)} (m_2(\mu) + d)^{2(d+1)} (K_1 + K_2) \end{cases}$$

where \lesssim hides a universal constant as well as log factors with respect to K_1, K_2 that are defined in Eq. 12.

The proof is left in Appendix and provides the appropriate set of parameters to use.

Remark 16 *As the proof shows, one could have relaxed the uniform-in-time upper-bound on $\nabla^2 \log(p_t^V)$, $\nabla^2 \log(p_t^{2V})$ into a time-dependent version. This relaxation would simply have replaced the term β in our bounds with $\max_{1 \leq k \leq N} \beta_{t_k}$.*

4 Application to smooth and strongly dissipative potentials

We prove in this section that the results above apply for the class of smooth, strongly convex potentials outside a ball: namely we show that if Assumption 1 and Assumption 3 hold, then Assumption 2 holds with $\beta \asymp L^2(R^2(L/\alpha)^2 + d/\alpha)$. The proof is decomposed into two time regimes: first, when t is sufficiently small, the following result ensures that the forward process remains log-smooth, and *a fortiori* semi-log-concave, when μ is.

Proposition 17 (Chen et al. [10]) *If V is C^2 and L -smooth and if t is such that $1 - e^{-2t} \leq \frac{e^{-t}}{L}$, then it holds that $\|\nabla^2 \log(p_t^V)\| \leq 2Le^t$.*

Hence, we only need to control $\nabla^2 \log(p_t^V)$ when $t \gtrsim 1$. Tweedie's formula (see Appendix A.1) yields the identity

$$\nabla^2 \log(p_t^V)(x) - \nabla^2 \log(\pi)(x) = \frac{e^{-2t}}{1 - e^{-2t}} \left(\frac{\text{Cov}_{q_{t,x}}(X)}{1 - e^{-2t}} - I_d \right), \quad (13)$$

where $q_{t,x} \propto e^{-V(z)} e^{-\frac{e^{-2t}\|z-x\|^2}{2(1-e^{-2t})}}$. Hence, in order to control $\nabla^2 \log(p_t^V)$ from above, it suffices to upper-bound $\text{Cov}_{q_{t,x}}$ independently of x . Now remark that if V is α -strongly convex outside some ball of radius R with condition number $\kappa := L/\alpha$, then $q_{t,x}$ remains at least α -strongly log-concave outside the same ball of radius R and has most condition number κ . We thus derived the following result which upper-bounds the variance (and *a fortiori* the covariance matrix) of log-smooth and strongly log-concave measures outside some ball of radius R .

Proposition 18 *Let V be a potential that is L -smooth, α -strongly convex outside some ball of radius R . Then, denoting $\mu \propto e^{-V}$ and $\kappa = L/\alpha$, it holds for any critical point x^* that*

$$\text{Var}_\mu(X) \leq \mathbb{E}_\mu[\|X - x^*\|^2] \leq 8R^2(\kappa + 1)^2 + \frac{4d}{\alpha}.$$

The proof is deferred to Appendix B.1 and is based on the following estimate that we believe to be novel: for smooth and strongly dissipative potentials, we show that for any $x \in \mathbb{R}^d$ and any critical point x^* of V

$$\langle \nabla V(x), x - x^* \rangle \geq \alpha \|x - x^*\|^2 - 2R(L + \alpha) \|x - x^*\|.$$

Integrating this inequality with respect to μ and using $\mathbb{E}_\mu[\|X - x^*\|] \leq \mathbb{E}_\mu[\|X - x^*\|^2]^{1/2}$ then yields

$$\mathbb{E}_\mu[\|X - x^*\|^2] \leq 2R(\kappa + 1) \mathbb{E}_\mu[\|X - x^*\|^2]^{1/2} + \frac{d}{\alpha}$$

and recovers the result of Proposition 18.

Remark 19 *We note that Ma et al. [31] and Chehab et al. [9] also derived bounds for the second moment of smooth and (weakly) dissipative potentials which *a fortiori* provides bounds for the variance. Yet the former suffers an exponential dependence in LR^2 and the latter implicitly depends on the the gradient of the potential at 0 which would have induced a dependence in x when applied to $q_{t,x}$.*

Algorithm	Complexity
ULA [31]	$O(e^{32R^2L\kappa^2\frac{d}{\epsilon^2}})$
MALA [31]	$O(\frac{e^{40LR^2}}{\alpha}\kappa^{3/2}d^{1/2}(d\log(\kappa) + \log(1/\epsilon))^{3/2})$
RD + ULA [25]	$e^{O(L^3\log^3(L^2(R^2\kappa^2+d/\alpha)/\epsilon))}$
RD + Rejection Sampling [22]	$O(L^{d/2}\epsilon^{-d}e^{L\ x^*\ ^2+L^2(\kappa^2R^2+d/\alpha)})$
RD + Self-normalized IS (ours)	$O(d^7L^{d+8}\epsilon^{-2(d+3)}(R^2\kappa^2 + \frac{d}{\alpha})^{2(d+4)})$

Table 1: Complexity of sampling algorithms in the smooth, strongly dissipative case. We denoted x^* the global minimum of V .

We use the previous Proposition to recover the following uniform bound:

$$-\frac{e^{-2t}}{(1-e^{-2t})^2} \left(8R^2(\kappa+1)^2 + \frac{4d}{\alpha} \right) I_d \leq \nabla^2 \log(\pi) - \nabla^2 \log(p_t^V). \quad (14)$$

Combined with Proposition 9 and Proposition 17, we can prove that the forward process p_t^V is uniformly smooth along the trajectory.

Corollary 20 *Assume V is C^2 , L -smooth and α -strongly convex outside some ball of radius R . Then, denoting the condition number $\kappa = L/\alpha$, it holds that*

$$\|\nabla^2 \log(p_t^V)\| \leq 4(L+1)^2 \left(8R^2\alpha(\kappa+1)^2 + \frac{4d}{\alpha} + 1 \right).$$

The proof can be found in Appendix B.2. Note that numerous works in the reverse diffusion literature make the *a priori* assumption that the forward process is L -log-smooth [12, 24, 25]. The result above provides a non-trivial setting where this assumption does hold.

Remark 21 *Note that our bounds extend the results of Mikulincer and Shenfeld [36] and could be used in a similar fashion to recover estimates on the Log-Sobolev constants of strongly dissipative potentials. Yet, because of our residual dependence in d , we expect our resulting bounds to worsen those found in Ma et al. [31].*

In order to apply our Theorem 14, it remains now to bound the second moment $m_2(\mu)$, the gradients of the forward process $\|\nabla \log(p_t^V)\|$, $\|\nabla \log(p_t^{2V})\|$ and the density μ at 0 in the case where V is L -smooth and α -strongly convex outside some ball $B_R(m)$. In what follows, to ease notations, we shall assume that $\nabla V(0) = 0$. Note that this extra assumption is benign since we can always compute any critical point x^* up to arbitrary precision in polynomial time [23] and then sample from $\tilde{V} = V(\cdot + x^*)$. Using Proposition 18, this extra assumption readily implies $m_2(\mu) \leq 8R^2(\kappa+1)^2 + \frac{4d}{\alpha}$. In the next Proposition, we provide the needed upper bounds for the gradient and the density.

Proposition 22 *Let V be a C^2 , L -smooth potential outside some ball of radius R such that $\nabla V(0) = 0$. Then it holds that*

$$\begin{cases} \|\nabla \log(p_t^V)(0)\|^2 \leq (L+1)^2 m_2(\mu), \\ 2d \log(L\mu(0)^{-2/d}/2\pi) \leq 4(R^2\alpha(\kappa+1)^2 + \log(2\kappa)). \end{cases}$$

The proof is left in Appendix. We can now apply Corollary 15 and prove that strongly dissipative distributions can be sampled from in polynomial time in fixed dimension.

Corollary 23 *Under Assumptions 1-3 and if we further assume $\nabla V(0) = 0$, there exists a stochastic algorithm to sample from $\mu \propto e^{-V}$ that outputs a sample $Y_N \sim \hat{p}$ such that $\mathbb{E}[\text{KL}(\mu, \hat{p})] \lesssim \epsilon$ that runs in (deterministic) time $O(d^7L^{d+8}\epsilon^{-2(d+3)}(R^2\kappa^2 + \frac{d}{\alpha})^{2(d+4)})$ where \lesssim hides a universal constant and log factors in d, L, α, R .*

The proof is left in Appendix. We summarize in Table 1 how our bounds compare to those of previous works after plugging our estimate of $m_2(\mu)$ in the smooth, strongly dissipative case. Assuming that our estimate is tight, we observe that while our bound exhibits a worse dependence in ϵ^{-1}, d than in Ma et al. [31] and in He et al. [22], we provide an exponential improvement with respect to R, L over all previous works where we lowered the complexity to polynomial.

5 Conclusion

In this article, we pushed one step further the reduction from sampling to intermediate score estimation initiated over the past two years by Chen et al. [12, 10], Conforti et al. [16], Benton et al. [6]. Using our self-normalized estimator of the scores, we proved that in fixed dimension, the sampling problem could be further reduced to a regularity problem along the forward OU process. We then applied this result to derive polynomial time guarantees for the task of sampling from a log-smooth, strongly dissipative distribution thus providing an exponential improvement over previously known bounds. Yet we believe that our contribution could pave the way for numerous other non-log concave sampling problems in low dimensions. In future works, we plan to apply our results to Gaussian mixtures which are not fully covered by the strongly dissipative framework.

Acknowledgments

This work was funded by ANR JCJC WOS. We thank Pierre Monmarché for his help in proving Lemma 28.

References

- [1] Agapiou, S., Papaspiliopoulos, O., Sanz-Alonso, D., and Stuart, A. M. (2017). Importance sampling: Intrinsic dimension and computational cost. *Statistical Science*.
- [2] Altschuler, J. M. and Chewi, S. (2024). Faster high-accuracy log-concave sampling via algorithmic warm starts. *Journal of the ACM*.
- [3] Ambrosio, L., Gigli, N., and Savaré, G. (2008). *Gradient flows: in metric spaces and in the space of probability measures*. Springer Science & Business Media.
- [4] Atchadé, Y. F. (2006). An adaptive version for the metropolis adjusted langevin algorithm with a truncated drift. *Methodology and Computing in applied Probability*.
- [5] Bakry, D., Gentil, I., Ledoux, M., et al. (2014). *Analysis and geometry of Markov diffusion operators*. Springer.
- [6] Benton, J., Bortoli, V. D., Doucet, A., and Deligiannidis, G. (2024). Nearly d-linear convergence bounds for diffusion models via stochastic localization. In *ICLR*.
- [7] Bortoli, V. D., Thornton, J., Heng, J., and Doucet, A. (2021). Diffusion schrödinger bridge with applications to score-based generative modeling. In *NeurIPS*.
- [8] Bou-Rabee, N. and Marsden, M. (2025). Unadjusted hamiltonian mcmc with stratified monte carlo time integration. *Annals of Applied Probability*.
- [9] Chehab, O., Korba, A., Stromme, A., and Vacher, A. (2024). Provable convergence and limitations of geometric tempering for langevin dynamics. *arXiv preprint arXiv:2410.09697*.
- [10] Chen, H., Lee, H., and Lu, J. (2023a). Improved analysis of score-based generative modeling: User-friendly bounds under minimal smoothness assumptions. In *ICML*.
- [11] Chen, H.-B., Chewi, S., and Niles-Weed, J. (2021). Dimension-free log-sobolev inequalities for mixture distributions. *Journal of Functional Analysis*.
- [12] Chen, S., Chewi, S., Li, J., Li, Y., Salim, A., and Zhang, A. R. (2023b). Sampling is as easy as learning the score: theory for diffusion models with minimal data assumptions. In *ICLR*.
- [13] Cheng, X., Chatterji, N. S., Abbasi-Yadkori, Y., Bartlett, P. L., and Jordan, M. I. (2018a). Sharp convergence rates for langevin dynamics in the nonconvex setting. *arXiv preprint arXiv:1805.01648*.
- [14] Cheng, X., Chatterji, N. S., Bartlett, P. L., and Jordan, M. I. (2018b). Underdamped langevin mcmc: A non-asymptotic analysis. In *COLT*.
- [15] Chopin, N., Lelièvre, T., and Stoltz, G. (2012). Free energy methods for bayesian inference: efficient exploration of univariate gaussian mixture posteriors. *Statistics and Computing*.

- [16] Conforti, G., Durmus, A., and Silveri, M. G. (2025). Score diffusion models without early stopping: finite fisher information is all you need. *SIAM Journal on Mathematics of Data Science (SIMODS)*.
- [17] Dasgupta, S. and Hsu, D. (2008). Hierarchical sampling for active learning. In *ICML*.
- [18] Ding, Z., Jiao, Y., Lu, X., Yang, Z., and Yuan, C. (2023). Sampling via follmer flow. *arXiv preprint arXiv:2311.03660*.
- [19] Erdogdu, M. A., Hosseinzadeh, R., and Zhang, S. (2022). Convergence of langevin monte carlo in chi-squared and rényi divergence. In *AISTATS*.
- [20] Foster, J., Lyons, T., and Oberhauser, H. (2021). The shifted ode method for underdamped langevin mcmc. *arXiv preprint arXiv:2101.03446*.
- [21] Gelman, A., Shirley, K., et al. (2011). Inference from simulations and monitoring convergence. *Handbook of markov chain monte carlo*.
- [22] He, Y., Rojas, K., and Tao, M. (2024). Zeroth-order sampling methods for non-log-concave distributions: Alleviating metastability by denoising diffusion. In *NeurIPS*.
- [23] Hollender, A. and Zampetakis, E. (2023). The computational complexity of finding stationary points in non-convex optimization. In *COLT*.
- [24] Huang, X., Dong, H., Yifan, H., Ma, Y., and Zhang, T. (2023). Reverse diffusion monte carlo. In *ICLR*.
- [25] Huang, X., Zou, D., Dong, H., Ma, Y.-A., and Zhang, T. (2024). Faster sampling without isoperimetry via diffusion-based monte carlo. In *COLT*.
- [26] Huisinga, W., Meyn, S., and Schütte, C. (2004). Phase transitions and metastability in markovian and molecular systems. *The Annals of Applied Probability*.
- [27] Kim, Y.-H. and Milman, E. (2012). A generalization of caffarelli’s contraction theorem via (reverse) heat flow. *Mathematische Annalen*.
- [28] Lee, H., Lu, J., and Tan, Y. (2022). Convergence for score-based generative modeling with polynomial complexity. In *NeurIPS*.
- [29] Lee, H., Risteski, A., and Ge, R. (2018). Beyond log-concavity: Provable guarantees for sampling multi-modal distributions using simulated tempering langevin monte carlo. In *NeurIPS*.
- [30] Lei, M., Zavodszky, M. I., Kuhn, L. A., and Thorpe, M. F. (2004). Sampling protein conformations and pathways. *Journal of Computational Chemistry*.
- [31] Ma, Y.-A., Chen, Y., Jin, C., Flammarion, N., and Jordan, M. I. (2019). Sampling can be faster than optimization. *Proceedings of the National Academy of Sciences*.
- [32] Majka, M. B., Mijatović, A., and Szpruch, L. (2019). Non-asymptotic bounds for sampling algorithms without log-concavity. *The Annals of Applied Probability*.
- [33] Marin, J.-M., Mengersen, K., and Robert, C. P. (2005). Bayesian modelling and inference on mixtures of distributions. *Handbook of statistics*.
- [34] McLachlan, G. and Peel, D. (2004). *Finite mixture models*. Wiley.
- [35] Mikulincer, D. and Shenfeld, Y. (2023). *On the Lipschitz Properties of Transportation Along Heat Flows*. Springer International Publishing.
- [36] Mikulincer, D. and Shenfeld, Y. (2024). The brownian transport map. *Probability Theory and Related Fields*.
- [37] Neal, R. M. (1996). Sampling from multimodal distributions using tempered transitions. *Statistics and computing*.
- [38] Oliver, D. S. (2017). Metropolized randomized maximum likelihood for improved sampling from multimodal distributions. *SIAM/ASA Journal on Uncertainty Quantification*.

- [39] Robbins, H. E. (1992). An empirical bayes approach to statistics. In *Breakthroughs in Statistics: Foundations and basic theory*. Springer.
- [40] Roberts, G. O. and Tweedie, R. L. (1996). Exponential convergence of Langevin distributions and their discrete approximations. *Bernoulli*.
- [41] Robins, J. M., Holmes, C. C., McGree, J. M., Mengersen, K., Richardson, S., and Ryan, E. G. (2017). Importance sampling: Intrinsic dimension and computational cost. *Statistical Science*.
- [42] Roy, V. (2020). Convergence diagnostics for markov chain monte carlo. *Annual Review of Statistics and Its Application*.
- [43] Shen, R. and Lee, Y. T. (2019). The randomized midpoint method for log-concave sampling. In *NeurIPS*.
- [44] Shteto, I., Linares, J., and Varret, F. (1997). Monte carlo entropic sampling for the study of metastable states and relaxation paths. *Physical Review E*.
- [45] Song, Y. and Ermon, S. (2019). Generative modeling by estimating gradients of the data distribution. In *NeurIPS*.
- [46] Vempala, S. and Wibisono, A. (2019). Rapid convergence of the unadjusted Langevin algorithm: Isoperimetry suffices. In *NeurIPS*.
- [47] Welling, M. (2009). Herding dynamical weights to learn. In *ICML*.
- [48] Xifara, T., Sherlock, C., Livingstone, S., Byrne, S., and Girolami, M. (2014). Langevin diffusions and the metropolis-adjusted langevin algorithm. *Statistics And Probability Letters*.
- [49] Zhang, S., Chewi, S., Li, M., Balasubramanian, K., and Erdogdu, M. A. (2023). Improved discretization analysis for underdamped langevin monte carlo. In *COLT*.

A Proofs of Section 3

A.1 Identities of the forward process

Before starting our proofs, we recall useful identities on the density and the log hessian of the forward process known as the *Tweedie's formulas* [39].

Proposition 24 (Tweedie's formulas) *Denoting p_t^V the density of the forward process X_t initialized at $\mu \propto e^{-V}$, and $Y_t \sim \mathcal{N}(0, (1 - e^{-2t})I_d)$, it holds for all $z \in \mathbb{R}^d$ that*

$$p_t^V(z) = \frac{e^{td}}{Z^V} \mathbb{E}[e^{-V((z-Y_t)e^t)}], \quad (15)$$

and that

$$\nabla^2 \log(p_t^V)(z) = \frac{\mathbb{E}[Y_t Y_t^\top e^{-V((z-Y_t)e^t)}]}{(1 - e^{-2t})^2 \mathbb{E}[e^{-V((z-Y_t)e^t)}]} - \frac{I_d}{(1 - e^{-2t})} - (\nabla \log(p_t^V)(z))(\nabla \log(p_t^V)(z))^\top. \quad (16)$$

Proof. Recall that p_t^V is the law of the variable

$$X_t = e^{-t}X_0 + B_{1-e^{-2t}},$$

with $X_0 \sim \mu$ and B_s the standard Brownian motion evaluated at time s . Hence, using Bayes formula, we have

$$p_t^V(z) = \int p_t(z|x) dp_0(x) = \frac{1}{Z^V(1 - e^{-2t})^{d/2}(2\pi)^{d/2}} \int_{\mathbb{R}^d} e^{-\frac{\|z - xe^{-t}\|^2}{2(1 - e^{-2t})}} e^{-V(x)} dx.$$

After taking the logarithm and differentiating with respect to z , we obtain

$$\nabla \log(p_t^V)(z) = \frac{\int_{\mathbb{R}^d} -(z - xe^{-t}) e^{-\frac{\|z - xe^{-t}\|^2}{2(1 - e^{-2t})}} e^{-V(x)} dx}{(1 - e^{-2t}) \int_{\mathbb{R}^d} e^{-\frac{\|z - xe^{-t}\|^2}{2(1 - e^{-2t})}} e^{-V(x)} dx}.$$

To obtain the Hessian, we differentiate the formula above. The Jacobian of the numerator is given by

$$-I_d \int_{\mathbb{R}^d} e^{-\frac{\|z-xe^{-t}\|^2}{2(1-e^{-2t})}} e^{-V(x)} dx + \frac{1}{1-e^{-2t}} \int_{\mathbb{R}^d} (z-xe^{-t})(z-xe^{-t})^\top e^{-\frac{\|z-xe^{-t}\|^2}{2(1-e^{-2t})}} e^{-V(x)} dx,$$

from which we can deduce

$$\begin{aligned} \nabla^2 \log(p_t^V)(z) &= -\frac{I_d}{(1-e^{-2t})} + \frac{\int_{\mathbb{R}^d} (z-xe^{-t})(z-xe^{-t})^\top e^{-\frac{\|z-xe^{-t}\|^2}{2(1-e^{-2t})}} e^{-V(x)} dx}{(1-e^{-2t})^2 \int_{\mathbb{R}^d} e^{-\frac{\|z-xe^{-t}\|^2}{2(1-e^{-2t})}} e^{-V(x)} dx} \\ &\quad - \frac{(\int_{\mathbb{R}^d} (z-xe^{-t}) e^{-\frac{\|z-xe^{-t}\|^2}{2(1-e^{-2t})}} e^{-V(x)} dx)(\int_{\mathbb{R}^d} (z-xe^{-t}) e^{-\frac{\|z-xe^{-t}\|^2}{2(1-e^{-2t})}} e^{-V(x)} dx)^\top}{(1-e^{-2t})^2 (\int_{\mathbb{R}^d} e^{-\frac{\|z-xe^{-t}\|^2}{2(1-e^{-2t})}} e^{-V(x)} dx)^2} \\ &= -\frac{I_d}{(1-e^{-2t})} + \frac{\int_{\mathbb{R}^d} (z-xe^{-t})(z-xe^{-t})^\top e^{-\frac{\|z-xe^{-t}\|^2}{2(1-e^{-2t})}} e^{-V(x)} dx}{(1-e^{-2t})^2 \int_{\mathbb{R}^d} e^{-\frac{\|z-xe^{-t}\|^2}{2(1-e^{-2t})}} e^{-V(x)} dx} \\ &\quad - (\nabla \log(p_t^V)(z))(\nabla \log(p_t^V)(z))^\top. \end{aligned}$$

In order to rewrite the quantities above as expectations, we make the change of variable $y = z - xe^{-t}$ so that $x = (z - y)e^t$ and we obtain for the density p_t^V :

$$p_t^V(z) = \frac{e^{td}}{Z^V} \mathbb{E}[e^{-V((z-Y_t)e^t)}],$$

where $Y_t \sim \mathcal{N}(0, I_d(1 - e^{-2t}))$. Conversely, the score rewrites as

$$\nabla \log(p_t^V)(z) = \frac{\mathbb{E}[-Y_t e^{-V((z-Y_t)e^t)}]}{(1 - e^{-2t}) \mathbb{E}[e^{-V((z-Y_t)e^t)}]},$$

and the Hessian rewrites as

$$\nabla^2 \log(p_t^V)(z) = \frac{\mathbb{E}[Y_t Y_t^\top e^{-V((z-Y_t)e^t)}]}{(1 - e^{-2t})^2 \mathbb{E}[e^{-V((z-Y_t)e^t)}]} - \frac{I_d}{(1 - e^{-2t})} - (\nabla \log(p_t^V)(z))(\nabla \log(p_t^V)(z))^\top.$$

□

A.2 Proof of Proposition 6

Proof. For the numerator, we have

$$\hat{N}(z) - N(z) = \frac{-1}{n} \sum_{i=1}^n \frac{y_i e^{-V((z-y_i)e^t)}}{1 - e^{-2t}} + N,$$

hence, since the y_i , $i = 1, \dots, n$ are i.i.d. distributed as $Y_t \sim \mathcal{N}(0, (1 - e^{-2t})I_d)$,

$$\mathbb{E}[\|\hat{N}(z) - N(z)\|^2] = \frac{1}{n} \mathbb{E} \left[\left\| \frac{Y_t e^{-V((z-Y_t)e^t)}}{1 - e^{-2t}} - N(z) \right\|^2 \right] \leq \frac{1}{n} \mathbb{E} \left[\frac{\|Y_t\|^2 e^{-2V((z-Y_t)e^t)}}{(1 - e^{-2t})^2} \right].$$

Taking in the trace in the log hessian identity in Proposition 24 yields

$$\mathbb{E} \left[\frac{\|Y_t\|^2 e^{-2V((z-Y_t)e^t)}}{(1 - e^{-2t})^2} \right] = \left(\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1 - e^{-2t}} + \|\nabla \log(p_t^{2V})\|^2 \right) p_t^{2V} Z_{2V} e^{-td}.$$

Similarly, we have

$$\hat{D}(z) - D(z) = \frac{1}{n} \sum_{i=1}^n e^{-V((z-y_i)e^t)} - D,$$

hence we get using again Proposition 24,

$$\mathbb{E}[(\hat{D}(z) - D(z))^2] = \frac{1}{n} \left(\mathbb{E}[e^{-2V((z-Y_t)e^t)}] - \mathbb{E}[e^{-2((z-Y_t)e^t)}]^2 \right) \leq \frac{p_t^{2V}(z) Z_{2V} e^{-td}}{n}. \quad (17)$$

□

A.3 Proof of Proposition 7

Proof. To ease notations, we shall drop the dependence in z throughout the proof. Define the event $A = (\hat{D} \geq \eta D) \cap (\|\hat{N}\| \leq \kappa \|N\|)$ where $\eta \leq 1, \kappa \geq 1$ are positive scalars to be chosen later. We start to decompose the quadratic error as:

$$\mathbb{E} \left[\left\| \frac{\hat{N}}{\hat{D}} - \frac{N}{D} \right\|^2 \right] = \mathbb{E} \left[\left\| \frac{\hat{N}}{\hat{D}} - \frac{N}{D} \right\|^2 \mathbf{1}_A \right] + \mathbb{E} \left[\left\| \frac{\hat{N}}{\hat{D}} - \frac{N}{D} \right\|^2 \mathbf{1}_{\bar{A}} \right].$$

We now separately analyze the first and the second term. For the first term, define

$$\begin{cases} \theta : \mathbb{R}^d \times \mathbb{R}^* \rightarrow \mathbb{R} \\ (x, p) \mapsto \left\| \frac{x}{p} - \frac{N}{D} \right\|^2. \end{cases}$$

The gradient and Hessian of θ are given by

$$\begin{cases} \nabla \theta(x, p) = \frac{-2}{p} \left(\frac{N}{D} - \frac{x}{p}, \left\| \frac{x}{p} \right\|^2 - \left\langle \frac{x}{p}, \frac{N}{D} \right\rangle \right), \\ \nabla^2 \theta(x, p) = \frac{-2}{p^2} \begin{pmatrix} -I_d & \left(\frac{2x}{p} - \frac{N}{D} \right)^\top \\ \left(\frac{2x}{p} - \frac{N}{D} \right) & -3 \left\| \frac{x}{p} \right\|^2 + 2 \left\langle \frac{x}{p}, \frac{N}{D} \right\rangle \end{pmatrix}. \end{cases}$$

We thus make a Taylor expansion of order 2 of $\theta(\hat{N}, \hat{D})$ around (N, D) : there exists (a random) $\hat{t} \in [0, 1]$ such that

$$\begin{aligned} \theta(\hat{N}, \hat{D}) &= \theta(N, D) + \nabla \theta(N, D)^\top (\hat{N} - N, \hat{D} - D) \\ &\quad + \frac{1}{2} (\hat{N} - N, \hat{D} - D)^\top \nabla^2 \theta(\hat{N}_{\hat{t}}, \hat{D}_{\hat{t}}) (\hat{N} - N, \hat{D} - D). \end{aligned}$$

where we denoted $\hat{N}_{\hat{t}} = \hat{t}\hat{N} + (1-\hat{t})N$ and $\hat{D}_{\hat{t}} = \hat{t}\hat{D} + (1-\hat{t})D$. The two first terms in the expansion are null and we are left with

$$\begin{aligned} \theta(\hat{N}, \hat{D}) &= \frac{1}{\hat{D}_{\hat{t}}^2} \left(\|\hat{N} - N\|^2 - 2 \left\langle 2 \frac{\hat{N}_{\hat{t}}}{\hat{D}_{\hat{t}}} - \frac{N}{D}, \hat{N} - N \right\rangle (\hat{D} - D) \right) + \left(3 \frac{\|\hat{N}_{\hat{t}}\|^2}{\hat{D}_{\hat{t}}^2} - 2 \left\langle \frac{\hat{N}_{\hat{t}}}{\hat{D}_{\hat{t}}}, \frac{N}{D} \right\rangle \right) (\hat{D} - D)^2 \\ &\leq \frac{1}{\hat{D}_{\hat{t}}^2} \left(\|\hat{N} - N\|^2 + 2 \left\| 2 \frac{\hat{N}_{\hat{t}}}{\hat{D}_{\hat{t}}} - \frac{N}{D} \right\| \|\hat{N} - N\| |\hat{D} - D| + \left(3 \frac{\|\hat{N}_{\hat{t}}\|^2}{\hat{D}_{\hat{t}}^2} + 2 \left\| \frac{\hat{N}_{\hat{t}}}{\hat{D}_{\hat{t}}} \right\| \left\| \frac{N}{D} \right\| \right) (\hat{D} - D)^2 \right). \end{aligned}$$

Hence, almost surely over A

$$\left\| \frac{\hat{N}}{\hat{D}} - \frac{N}{D} \right\|^2 \leq \frac{1}{\eta^2 D^2} \left(\|\hat{N} - N\|^2 + \frac{6\kappa}{\eta} \left\| \frac{N}{D} \right\| \|\hat{N} - N\| |\hat{D} - D| + 5 \left\| \frac{N}{D} \right\|^2 \left(\frac{\kappa}{\eta} \right)^2 (\hat{D} - D)^2 \right).$$

Hence, after taking the expectation and applying Cauchy-Schwarz, we obtain

$$\begin{aligned} \mathbb{E} \left[\left\| \frac{\hat{N}}{\hat{D}} - \frac{N}{D} \right\|^2 \mathbf{1}_A \right] &\leq \frac{1}{\eta^2 D^2} \left(\mathbb{E}[\|\hat{N} - N\|^2] + \frac{6\kappa}{\eta} \left\| \frac{N}{D} \right\| \mathbb{E}[\|\hat{N} - N\|^2]^{1/2} \mathbb{E}[(\hat{D} - D)^2]^{1/2} \right. \\ &\quad \left. + 5 \left\| \frac{N}{D} \right\|^2 \left(\frac{\kappa}{\eta} \right)^2 \mathbb{E}[(\hat{D} - D)^2] \right). \end{aligned}$$

Now recall that $D = p_t^V Z_V e^{-td}$ and that $\frac{N}{D} = \nabla \log(p_t^V)$ which, combined with Proposition 6 yields for the first term:

$$\frac{\mathbb{E}[\|\hat{N} - N\|^2]}{\eta^2 D^2} \leq \frac{p_t^{2V} Z_{2V} e^{td}}{\eta^2 (p_t^V)^2 (Z_V)^2 n} \left(\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1 - e^{-2t}} + \|\nabla \log(p_t^{2V})\|^2 \right),$$

for the second term:

$$\frac{6\kappa}{\eta^3 D^2} \left\| \frac{N}{D} \right\| \|\hat{N} - N\| |\hat{D} - D| \leq \frac{6\kappa p_t^{2V} Z_{2V} e^{td}}{\eta^3 (p_t^V)^2 (Z_V)^2 n} \|\nabla \log(p_t^V)\| \left(\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1 - e^{-2t}} + \|\nabla \log(p_t^{2V})\|^2 \right)^{1/2}$$

and for the last term:

$$5 \left\| \frac{N}{\hat{D}} \right\|^2 \left(\frac{\kappa}{\eta^2 D} \right)^2 (\hat{D} - D)^2 \leq \frac{5\kappa^2 p_t^{2V} Z_{2V} e^{td}}{\eta^4 (p_t^V)^2 (Z_V)^2 n} \|\nabla \log(p_t^V)\|^2.$$

Hence we finally obtain

$$\begin{aligned} \mathbb{E} \left[\left\| \frac{\hat{N}}{\hat{D}} - \frac{N}{D} \right\|^2 \mathbf{1}_A \right] &\leq \frac{p_t^{2V} Z_{2V} e^{td}}{n\eta^2 (p_t^V)^2 (Z_V)^2} \left(\frac{6\kappa}{\eta} \|\nabla \log(p_t^V)\| \left[\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1 - e^{-2t}} + \|\nabla \log(p_t^{2V})\|^2 \right]^{1/2} \right. \\ &\quad \left. + \Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1 - e^{-2t}} + 6 \left(\frac{\kappa}{\eta} \right)^2 \|\nabla \log(p_t^V)\|^2 \right). \end{aligned} \quad (18)$$

Let us now handle the quadratic error of the estimator on the complementary \bar{A} . We have, using Young's inequality $\|a - b\|^2 \leq 2(\|a\|^2 + \|b\|^2)$,

$$\mathbb{E} \left[\left\| \frac{\hat{N}}{\hat{D}} - \frac{N}{D} \right\|^2 \mathbf{1}_{\bar{A}} \right] \leq 2 \left\| \frac{N}{D} \right\|^2 \mathbb{P}(\bar{A}) + 2\mathbb{E} \left[\mathbf{1}_{\bar{A}} \frac{\max_i \|y_i\|^2}{(1 - e^{-2t})^2} \right].$$

Now, recall that $X_i = \|y_i\|^2 (1 - e^{-2t})^{-1}$ are n independent variables such that for all i , $X_i \sim \chi^2(d)$. Using Hölder inequality for some $p \geq 1$, the second term can be upper-bounded as

$$\begin{aligned} \mathbb{E} \left[\mathbf{1}_{\bar{A}} \frac{\max_i \|y_i\|^2}{(1 - e^{-2t})^2} \right] &\leq \frac{1}{1 - e^{-2t}} \mathbb{E}[\max_i X_i^{p-1/p} \mathbb{P}(\bar{A})^{1-1/p}] \\ &\leq \frac{1}{1 - e^{-2t}} (n\mathbb{E}[X_1^p])^{1/p} \mathbb{P}(\bar{A})^{1-1/p} \leq \frac{1}{1 - e^{-2t}} n^{1/p} (d + 2p) \mathbb{P}(\bar{A})^{1-1/p}, \end{aligned}$$

where we used in the penultimate inequality that the max is smaller than the sum, and in the last one that $\mathbb{E}[X_1^p] = \prod_{i=0}^{p-1} (d + 2i)$ when $X_1 \sim \chi^2(d)$ combined with the fact that the geometric mean is lower than the arithmetic mean.

We now upper bound the probability of the event $\bar{A} = (\hat{D} < \eta D) \cup (\|\hat{N}\| > \kappa \|N\|)$. By Tchebychev inequality, using $\eta < 1$, it holds that

$$\mathbb{P}(\hat{D} < \eta D) \leq \frac{\mathbb{E}[(\hat{D} - D)^2]}{D^2(\eta - 1)^2} \leq \frac{p_t^{2V} Z_{2V} e^{td}}{n(p_t^V)^2 (Z_V)^2 (\eta - 1)^2} := \frac{U}{n(\eta - 1)^2}.$$

Similarly, recalling that $\|N\| = D \|\nabla \log(p_t^V)\|$, we have

$$\begin{aligned} \mathbb{P}(\|\hat{N}\| > \kappa \|N\|) &\leq \frac{\mathbb{E}[\|\hat{N} - N\|^2]}{\|N\|^2 (\kappa - 1)^2} \\ &\leq \frac{U}{n \|\nabla \log(p_t^V)\|^2 (\kappa - 1)^2} \left(\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1 - e^{-2t}} + \|\nabla \log(p_t^V)\|^2 \right). \end{aligned}$$

We now make a disjunction of cases: if $\|\nabla \log(p_t^V)\| \geq 1$, we pick $\eta = 1/2$ and $\kappa = 3/2$ so we recover

$$\begin{aligned} \mathbb{E} \left[\left\| \frac{\hat{N}}{\hat{D}} - \frac{N}{D} \right\|^2 \mathbf{1}_{\bar{A}} \right] &\leq \frac{8U}{n} \left[\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1 - e^{-2t}} + 2\|\nabla \log(p_t^V)\|^2 \right] \\ &\quad + \frac{n^{1/p} (d + 2p)}{1 - e^{-2t}} \left[\frac{8U}{n} \left(\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1 - e^{-2t}} + \|\nabla \log(p_t^V)\|^2 + 1 \right) \right]^{1-1/p} \\ &\leq \frac{8U}{n} \left[\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1 - e^{-2t}} + 2\|\nabla \log(p_t^V)\|^2 \right] \\ &\quad + \frac{n^{1/p} (d + 2p)}{1 - e^{-2t}} \left[\frac{8U}{n} \left(\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1 - e^{-2t}} + \|\nabla \log(p_t^V)\|^2 + 1 \right) + 1 \right]. \end{aligned}$$

We thus pick $p = \log(n)$ to get

$$\mathbb{E} \left[\left\| \frac{\hat{N}}{\hat{D}} - \frac{N}{D} \right\|^2 \mathbf{1}_{\bar{A}} \right] \leq \frac{16e^2 (d + 2 \log(n))}{n(1 - e^{-2t})} \left[U \left(\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1 - e^{-2t}} + \|\nabla \log(p_t^V)\|^2 + 1 \right) + 1 \right].$$

Combining this with the bound Eq. 18 eventually yields

$$\delta^2 \leq \frac{32e^2(d+2\log(n))}{n(1-e^{-2t})} \left[U \left(\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1-e^{-2t}} + \|\nabla \log(p_t^V)\|^2 + \|\nabla \log(p_t^{2V})\|^2 + 1 \right) + 1 \right],$$

where we used the inequality

$$\begin{aligned} & \|\nabla \log(p_t^V)\| \left[\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1-e^{-2t}} + \|\nabla \log(p_t^{2V})\|^2 \right]^{1/2} \\ & \leq \frac{1}{2} \left(\|\nabla \log(p_t^V)\|^2 + \Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1-e^{-2t}} + \|\nabla \log(p_t^{2V})\|^2 \right). \end{aligned}$$

In the case where $\|\nabla \log(p_t)\|^2 < 1$, we instead pick $\eta = 1/2$ and $\kappa = 1 + \frac{1}{2\|\nabla \log(p_t^V)\|}$. We obtain that

$$\mathbb{P}(\bar{A}) \leq \frac{4U}{n} \left(\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1-e^{-2t}} + \|\nabla \log(p_t^V)\|^2 + 1 \right)$$

and as previously, for $p = \log(n)$ we get

$$\mathbb{E} \left[\left\| \frac{\hat{N}}{\hat{D}} - \frac{N}{D} \right\|^2 \mathbf{1}_{\bar{A}} \right] \leq \frac{16e^2(d+2\log(n))}{n(1-e^{-2t})} \left[U \left(\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1-e^{-2t}} + \|\nabla \log(p_t^V)\|^2 + 1 \right) + 1 \right].$$

For this choice of κ, η , the bound on A becomes

$$\begin{aligned} \mathbb{E} \left[\left\| \frac{\hat{N}}{\hat{D}} - \frac{N}{D} \right\|^2 \mathbf{1}_A \right] & \leq \frac{4U}{n} \left(6 \left[\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1-e^{-2t}} + \|\nabla \log(p_t^{2V})\|^2 \right]^{1/2} + \Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1-e^{-2t}} + 6 \right) \\ & \leq \frac{4U}{n} \left(7 \left(\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1-e^{-2t}} \right) + 6(\|\nabla \log(p_t^{2V})\|^2 + 1) \right). \end{aligned}$$

Thus, we obtain as previously

$$\delta^2 \leq \frac{32e^2(d+2\log(n))}{n(1-e^{-2t})} \left[U \left(\Delta \log(p_t^{2V}) - \frac{\Delta \log(\pi)}{1-e^{-2t}} + \|\nabla \log(p_t^V)\|^2 + \|\nabla \log(p_t^{2V})\|^2 + 1 \right) + 1 \right],$$

□

A.4 Proof of Proposition 9

Proof. Define the Ornstein-Uhlenbeck semi-group Q_t as

$$\begin{aligned} Q_t(g)(z) & = \int g(ze^{-t} + \sqrt{1-e^{-2t}}y) e^{-\frac{\|y\|^2}{2}} (2\pi)^{-d/2} dy \\ & = \frac{1}{\sqrt{1-e^{-2t}}} \int g(u) e^{-\frac{\|u-e^{-t}z\|^2}{2(1-e^{-2t})}} (2\pi)^{-d/2} du, \end{aligned}$$

for all function g integrable w.r.t. the standard Gaussian measure. Taking g as $f = \frac{d\mu}{d\pi}$ with π the standard Gaussian, we obtain that

$$\begin{aligned} Q_t(f)(z) & = \frac{1}{Z_V \sqrt{1-e^{-2t}}} \int e^{-V(u)} e^{\frac{\|u\|^2}{2}} e^{-\frac{\|u-e^{-t}z\|^2}{2(1-e^{-2t})}} du \\ & = \frac{1}{Z_V \sqrt{1-e^{-2t}}} \int e^{-V(u)} e^{\frac{\|u\|^2(1-e^{-2t}) - \|u\|^2 + (z, ue^{-t}) - e^{-2t}\|z\|^2}{2(1-e^{-2t})}} du \\ & = \frac{1}{Z_V \sqrt{1-e^{-2t}}} \int e^{-V(u)} e^{\frac{-\|ue^{-t}-z\|^2 + \|z\|^2 - e^{-2t}\|z\|^2}{2(1-e^{-2t})}} du \\ & = \frac{e^{\frac{\|z\|^2}{2}}}{Z_V \sqrt{1-e^{-2t}}} \int e^{-V(u)} e^{-\frac{\|ue^{-t}-z\|^2}{2(1-e^{-2t})}} du. \end{aligned}$$

In particular, we remark that $\nabla \log(Q_t(f)) = \nabla \log(p_t^V) - \nabla \log(\pi)$. Now, the quantity $\nabla \log(Q_t(f))$ was studied in Mikulincer and Shenfeld [35] and they prove in Lemma 5 that for all z

$$\nabla^2 \log(Q_t(f))(z) \succeq \frac{(1-L)e^{-2t}}{(1-e^{-2t})(L-1)+1} I_d,$$

which is equivalent to

$$\nabla^2 \log(\pi)(z) - \nabla^2 \log(p_t^V)(z) \preceq \frac{(L-1)e^{-2t}}{(1-e^{-2t})(L-1)+1} I_d.$$

□

A.5 Proof of Lemma 10

Before proving Lemma 10, we introduce this preliminary result on the evolution of Φ_t .

We start by deriving the evolution of the ratio Φ_t .

Lemma 25 (Evolution of the ratio) *Let $t > 0$, it holds that*

$$\partial \Phi_t = \Phi_t (\Delta \log(\Phi_t) - \langle \nabla \log(\Phi_t), \nabla \log(\pi) \rangle + \|\nabla \log(\Phi_t)\|^2 + 2\langle \nabla \log(p_t^V), \nabla \log(\Phi_t) \rangle).$$

Proof. Recall that the log-density $\log(p_t^V)$ evolves as

$$\partial \log(p_t^V) = \Delta \log(p_t^V) + \|\nabla \log(p_t^V)\|^2 - \langle \nabla \log(p_t^V), \nabla \log(\pi) \rangle - \Delta \log(\pi).$$

Hence, we deduce that $\log(\Phi_t)$ evolves as

$$\partial \log(\Phi_t) = \Delta \log(\Phi_t) - \langle \nabla \log(\Phi_t), \nabla \log(\pi) \rangle + \|\nabla \log(p_t^{2V})\|^2 - \|\nabla \log(p_t^V)\|^2.$$

The difference of quadratic terms can be expressed as

$$\begin{aligned} \|\nabla \log(p_t^{2V})\|^2 - \|\nabla \log(p_t^V)\|^2 &= \|\nabla \log(\Phi_t) + \nabla \log(p_t^V)\|^2 - \|\nabla \log(p_t^V)\|^2 \\ &= \|\nabla \log(\Phi_t)\|^2 + 2\langle \nabla \log(p_t^V), \nabla \log(\Phi_t) \rangle, \end{aligned}$$

which allows to recover

$$\partial \log(\Phi_t) = \Delta \log(\Phi_t) - \langle \nabla \log(\Phi_t), \nabla \log(\pi) \rangle + \|\nabla \log(\Phi_t)\|^2 + 2\langle \nabla \log(p_t^V), \nabla \log(\Phi_t) \rangle.$$

□

We now provide the proof of Lemma 10.

Proof. Until the rest of the proof, the dependence on z of the integrand shall be implied unless expressed explicitly. We start by differentiating $m_0(\Phi_t)$ with respect to t :

$$\begin{aligned} \partial m_0(\Phi_t) &= \int \partial \Phi_t dz \\ &= \int \Phi_t (\Delta \log(\Phi_t) - \langle \nabla \log(\Phi_t), \nabla \log(\pi) \rangle + \|\nabla \log(\Phi_t)\|^2 + 2\langle \nabla \log(p_t^V), \nabla \log(\Phi_t) \rangle) dz, \end{aligned}$$

where we used Lemma 25 to compute $\partial \Phi_t$. Using integration by parts, the first term reads

$$\int \Phi_t \Delta \log(\Phi_t) dz = - \int \langle \nabla \Phi_t, \nabla \log(\Phi_t) \rangle dz = - \int \Phi_t \|\nabla \log(\Phi_t)\|^2 dz,$$

hence the first and the third terms cancel and we recover

$$\partial m_0(\Phi_t) = 2 \int \langle \nabla \log(p_t^V), \nabla \Phi_t \rangle dz - \int \langle \nabla \log(\pi), \nabla \Phi_t \rangle dz.$$

Using integration by parts again, we recover

$$\begin{aligned}\partial m_0(\Phi_t) &= \int \Delta \log(\pi) \Phi_t dz - 2 \int \Delta \log(p_t^V) \Phi_t dz \\ &= 2 \int (\Delta \log(\pi) - \Delta \log(p_t^V)) \Phi_t dz - \int \Delta \log(\pi) \Phi_t dz \\ &= dm_0(\Phi_t) + 2 \int (\Delta \log(\pi) - \Delta \log(p_t^V)) \Phi_t dz.\end{aligned}$$

Using Proposition 9, since V is L -smooth, the term $(\Delta \log(\pi) - \Delta \log(p_t^V))$ can be upper-bounded uniformly by $\frac{d(L-1)e^{-2t}}{(1-e^{-2t})(L-1)+1}$ so we eventually get

$$\partial m_0(\Phi_t) \leq dm_0(\Phi_t) \left(1 + \frac{2(L-1)e^{-2t}}{(1-e^{-2t})(L-1)+1} \right).$$

Hence we can use Gronwall lemma which yields

$$m_0(\Phi_t) \leq m_0(\Phi_0) \exp \left(d \int_0^t \left(1 + \frac{2(L-1)e^{-2s}}{(1-e^{-2s})(L-1)+1} \right) ds \right).$$

Denoting by Z_V (resp. Z_{2V}) the normalizing constant of e^{-V} (resp. e^{-2V}), the term $m_0(\Phi_0)$ reads

$$m_0(\Phi_0) = \int \frac{p_0^{2V}(z)}{p_0^V(z)} dz = \frac{Z_V}{Z_{2V}} \int \frac{e^{-2V(z)}}{e^{-V(z)}} dz = \frac{(Z_V)^2}{Z_{2V}}.$$

Finally, let us compute the integral above. Making the change of variable $u = e^{-2s}(L-1)$ we have $du = -2(L-1)e^{-2s} ds$ which yields

$$\begin{aligned}\int_0^t \frac{2(L-1)e^{-2s}}{(1-e^{-2s})(L-1)+1} ds &= - \int_{L-1}^{(L-1)e^{-2t}} \frac{1}{L-u} du \\ &= [\log(L-u)]_{L-1}^{(L-1)e^{-2t}} \\ &= \log(L - (L-1)e^{-2t}) \\ &= \log(L(1 - e^{-2t}) + e^{-2t}).\end{aligned}$$

Hence we recover

$$m_0(\Phi_t) \leq \frac{e^{td}(Z_V)^2}{Z_{2V}} (L(1 - e^{-2t}) + e^{-2t})^d.$$

□

A.6 Proof of Lemma 11

Proof. Let z_t be a point where Φ_t attains its maximum and denote $M_t = \log(\Phi_t)(z_t)$. By the implicit function theorem, z_t is differentiable hence we can compute ∂M_t as

$$\begin{aligned}\partial M_t &= \partial \log(\Phi_t)(z_t) + \langle \partial z_t, \nabla \log(\Phi_t)(z_t) \rangle \\ &= \Delta \log(\Phi_t)(z_t).\end{aligned}$$

Since z_t is a maximum, we have in particular $\Delta \Phi_t(z_t) \leq 0$ which implies that M_t decreases. □

A.7 Proof of Lemma 12

Before starting the proof, we need the following intermediate result.

Proposition 26 (Upper-bound of the maximum) *Let V be a C^2 potential such that $\nabla^2 V \preceq LI_d$. Denoting $\mu \propto e^{-V}$, it holds that $\frac{d\mu}{dz} \leq \left(\frac{\beta}{2\pi}\right)^{\frac{d}{2}}$.*

Proof. Recall that the density of μ can be re-written as

$$\frac{d\mu}{dz} = \frac{e^{-(V(z)-V_*)}}{\int_z e^{-(V(z)-V_*)} dz},$$

where V_* the minimum of V attained for some z_* . By definition $e^{-(V(z)-V_*)} \leq 1$ for all z . Furthermore, since V verifies $\nabla^2 V \preceq LL_d$, we are ensured that

$$V(z) - V_* \leq L \frac{\|z - z_*\|^2}{2},$$

which implies in particular that

$$\frac{1}{\int_z e^{-(V(z)-V_*)} dz} \leq \left(\frac{L}{2\pi}\right)^{\frac{d}{2}}.$$

□

We can now start the proof of the lemma.

Proof. Denoting π the density of the standard d dimensional Gaussian, recall that the density p_t^V evolves as

$$\partial p_t^V = \nabla \cdot \left(p_t^V \nabla \log \left(\frac{p_t^V}{\pi} \right) \right),$$

which can also be re-written as

$$\begin{aligned} \partial \log(p_t^V) &= \frac{\Delta p_t^V}{p_t^V} - \langle \nabla \log(p_t^V), \nabla \log(\pi) \rangle - \Delta \log(\pi) \\ &= \Delta \log(p_t^V) + \|\nabla \log(p_t^V)\|^2 - \langle \nabla \log(p_t^V), \nabla \log(\pi) \rangle - \Delta \log(\pi). \end{aligned}$$

In particular, for $z = 0$ this yields

$$\partial \log(p_t^V)(0) = \Delta \log(p_t^V)(0) + \|\nabla \log(p_t^V)\|^2(0) - \Delta \log(\pi)(0),$$

which implies

$$\int_0^t \|\nabla \log(p_s^V)\|^2(0) ds = \log(p_t^V)(0) - \log(p_0^V)(0) + \int_0^t \Delta \log(\pi)(0) - \Delta \log(p_t^V)(0) ds.$$

Using the uniform upper-bound of Proposition 9, the second term can upper-bounded as

$$\begin{aligned} \int_0^t \Delta \log(\pi)(0) - \Delta \log(p_t^V)(0) ds &\leq d \int_0^t \frac{(L-1)e^{-2s}}{(1-e^{-2s})(L-1)+1} ds \\ &= \frac{d}{2}(L-1) \int_{e^{-2t}}^1 \frac{1}{L-u(L-1)} du \\ &= \frac{d}{2} \log(L(1-e^{-2t}) + e^{-2t}). \end{aligned}$$

Furthermore, Proposition 9 shows that $\nabla^2 \log(p_t^V) \preceq \frac{L}{L(1-e^{-2t})+e^{-2t}}$. Thus, using Proposition 26, we recover that $\log(p_t^V)(0) \leq \frac{d}{2} \log\left(\frac{L}{L(1-e^{-2t})+e^{-2t}}\right) - \frac{d}{2} \log(2\pi)$. In particular, we recover

$$\int_0^t \|\nabla \log(p_s^V)\|^2(0) ds \leq -\log\left(\frac{d\mu}{dx}(0)\right) + \frac{d}{2} \log(L) - \frac{d}{2} \log(2\pi).$$

□

A.8 Proof of Lemma 13

Before starting our proof we first need to derive an upper-bound for the first moment of the ratio.

Lemma 27 *Let V be a C^2 , L -smooth potential such that $\mu \propto e^{-V}$ has finite second moment $m_2(\mu)$. It holds that*

$$m_1(\Phi_t) \leq \frac{(Z_V)^2}{Z_{2V}} e^{t(d+1)} (L(1 - e^{-2t}) + e^{-2t})^d \left(\sqrt{m_2(\mu)} + \sqrt{-2 \log(\mu(0)) + d \log\left(\frac{L}{2\pi}\right) + 2\sqrt{dL}} \right).$$

Proof. We differentiate $m_1(\Phi_t)$ and we recover

$$\partial m_1(\Phi_t) = \int \Phi_t (\Delta \log(\Phi_t) - \langle \nabla \log(\Phi_t), \nabla \log(\pi) \rangle + \|\nabla \log(\Phi_t)\|^2 + 2\langle \nabla \log(p_t^V), \nabla \log(\Phi_t) \rangle) \|z\| dz.$$

Integration by parts of the first term yields

$$\int \Phi_t \Delta \log(\Phi_t) \|z\| dz = - \int \Phi_t \|\nabla \log(\Phi_t)\|^2 \|z\| + \langle \nabla \Phi_t, \frac{z}{\|z\|} \rangle dz,$$

hence the squared gradients terms cancel and we recover

$$\partial m_1(\Phi_t) = \int \langle 2\nabla \log(p_t^V) - \nabla \log(\pi), \nabla \Phi_t \rangle \|z\| dz - \int \langle \nabla \Phi_t, \frac{z}{\|z\|} \rangle dz.$$

Let us denote by A the first term above. Integration by parts yields:

$$\begin{aligned} A &= \int \Phi_t (\Delta \log(\pi) - 2\Delta \log(p_t^V)) \|z\| dz + \int \Phi_t \langle \nabla \log(\pi) - 2\nabla \log(p_t^V), \frac{z}{\|z\|} \rangle dz \\ &= - \int \Phi_t \Delta \log(\pi) \|z\| dz + 2 \int \Phi_t (\Delta \log(\pi) - \Delta \log(p_t^V)) \|z\| dz \\ &\quad - \int \Phi_t \langle \nabla \log(\pi), \frac{z}{\|z\|} \rangle dz + 2 \int \Phi_t \langle \nabla \log(\pi) - \nabla \log(p_t^V), \frac{z}{\|z\|} \rangle dz \\ &= (d+1)m_1(\Phi_t) + 2 \int \Phi_t (\Delta \log(\pi) - \Delta \log(p_t^V)) \|z\| dz + 2 \int \Phi_t \langle \nabla \log(\pi) - \nabla \log(p_t^V), \frac{z}{\|z\|} \rangle dz. \end{aligned}$$

Using the upper bound given in Proposition 9, we get $2 \int \Phi_t (\Delta \log(\pi) - \Delta \log(p_t^V)) \|z\| dz \leq 2d \frac{(L-1)e^{-2t}}{(L-1)(1-e^{-2t})+1} m_1(\Phi_t)$. Similarly, we re-write the second term as

$$\begin{aligned} 2 \int \Phi_t \langle \nabla \log(\pi) - \nabla \log(p_t^V), \frac{z}{\|z\|} \rangle dz &= 2 \int \Phi_t \langle \nabla \log\left(\frac{\pi}{p_t^V}\right)(z) - \nabla \log\left(\frac{\pi}{p_t^V}\right)(0), \frac{z}{\|z\|} \rangle dz \\ &\quad + 2 \int \Phi_t \langle \nabla \log\left(\frac{\pi}{p_t^V}\right)(0), \frac{z}{\|z\|} \rangle dz \\ &\leq 2m_1(\Phi_t) \frac{(L-1)e^{-2t}}{(L-1)(1-e^{-2t})+1} + 2\|\nabla \log(p_t^V)(0)\| m_0(\Phi_t). \end{aligned}$$

Let us now handle the term $B = - \int \langle \nabla \Phi_t, \frac{z}{\|z\|} \rangle dz$. In one dimension, $B = 2\Phi_t(0) \leq \max(\Phi_t)$ and for $d \geq 2$, we have

$$\begin{aligned} B &= \int \frac{\Phi_t(d-1)}{\|z\|} dz \\ &= \int_{B_R} \frac{\Phi_t(d-1)}{\|z\|} dz + \int_{\overline{B_R}} \frac{\Phi_t(d-1)}{\|z\|} dz \\ &\leq \max(\Phi_t) \int_{B_R} \frac{d-1}{\|z\|} dz + \frac{d-1}{R} m_0(\Phi_t) \\ &= \max(\Phi_t) \frac{2\pi^{d/2}}{\Gamma(d/2)} R^{d-1} + \frac{d-1}{R} m_0(\Phi_t). \end{aligned}$$

Using Lemma 11 and Proposition 26, we have that $\max(\Phi_t) \leq \max(\Phi_0) = \frac{(Z_V)^2}{Z_{2V}} \max\left(\frac{d\mu}{dx}\right) \leq \frac{(Z_V)^2}{Z_{2V}} \left(\frac{L}{2\pi}\right)^{d/2}$. Hence, if we pick $R = (Z_{2V} m_0(\Phi_t) \Gamma(d/2) / Z_V^2)^{1/d} L^{-1/2} 2^{1/2-1/d}$, we get as an upper-bound for B :

$$\begin{aligned} B &\leq \frac{Z_V^2}{Z_{2V}} d\sqrt{L} 2^{1/d-1/2} (m_0(\Phi_t) Z_{2V} / Z_V^2)^{\frac{d-1}{d}} \Gamma(d/2)^{-1/d} \\ &\leq \frac{2Z_V^2}{Z_{2V}} \sqrt{dL} (m_0(\Phi_t) Z_{2V} / Z_V^2)^{\frac{d-1}{d}}. \end{aligned}$$

In particular, we recover that

$$\partial m_1(\Phi_t) \leq (d+1) \left(1 + 2 \frac{(L-1)e^{-2t}}{(L-1)(1-e^{-2t})+1}\right) m_1(\Phi_t) + 2 \|\nabla \log(p_t^V)(0)\| m_0(\Phi_t) + \frac{2Z_V^2}{Z_{2V}} \sqrt{dL} (m_0(\Phi_t) Z_{2V} / Z_V^2)^{\frac{d-1}{d}},$$

hence using Gronwall lemma, we have that

$$\begin{aligned} m_1(\Phi_t) &\leq e^{t(d+1)} (L(1-e^{-2t}) + e^{-2t})^{d+1} m_1(\Phi_0) \\ &\quad + 2 \int_0^t \left[\|\nabla \log(p_s^V)(0)\| m_1(\Phi_s) + \sqrt{dL} (m_0(\Phi_s) Z_{2V} / Z_V^2)^{\frac{d-1}{d}} \right] \frac{e^{t(d+1)} (L(1-e^{-2t}) + e^{-2t})^{d+1}}{e^{s(d+1)} (L(1-e^{-2s}) + e^{-2s})^{d+1}} ds. \end{aligned}$$

Using Lemma 10, we have that $m_0(\Phi_s) \leq \frac{Z_V^2}{Z_{2V}} e^{sd} (L(1-e^{-2s}) + e^{-2s})^d$ hence the first term of the integral is upper-bounded as:

$$\int_0^t \frac{e^{-s(d+1)} m_0(\Phi_s) \|\nabla \log(p_s^V)(0)\|}{(L(1-e^{-2s}) + e^{-2s})^{d+1}} ds \leq \frac{Z_V^2}{Z_{2V}} \int_0^t \|\nabla \log(p_s^V)(0)\| \frac{e^{-s}}{L(1-e^{-2s}) + e^{-2s}} ds.$$

By Cauchy-Schwarz it holds that

$$\int_0^t \frac{\|\nabla \log(p_s^V)(0)\| e^{-s}}{L(1-e^{-2s}) + e^{-2s}} ds \leq \sqrt{\int_0^t \|\nabla \log(p_s^V)(0)\|^2 ds} \sqrt{\int_0^t \frac{e^{-2s}}{(L(1-e^{-2s}) + e^{-2s})^2} ds}.$$

The integral term is given by

$$\begin{aligned} \int_0^t \frac{e^{-2s}}{(L(1-e^{-2s}) + e^{-2s})^2} ds &= \frac{1}{2} \int_{e^{-2t}}^1 \frac{1}{(L(1-u) + u)^2} du \\ &= \frac{1}{2(1-L)} \left[-\frac{1}{L(1-u) + u} \right]_{e^{-2t}}^1 \\ &= \frac{1}{2(1-L)} \left(\frac{1}{L(1-e^{-2t}) + e^{-2t}} - 1 \right) \\ &= \frac{1 - e^{-2t}}{2} \end{aligned}$$

Similarly,

$$\begin{aligned} \int_0^t m_0(\Phi_s)^{\frac{d-1}{d}} \frac{e^{-s(d+1)}}{(L(1-e^{-2s}) + e^{-2s})^{d+1}} ds &\leq \int_0^t \frac{e^{-2s}}{(L(1-e^{-2s}) + e^{-2s})^2} ds \\ &= \frac{1 - e^{-2t}}{2} \end{aligned}$$

Hence, we obtain

$$m_1(\Phi_t) \leq \frac{Z_V^2}{Z_{2V}} e^{t(d+1)} (L(1-e^{-2t}) + e^{-2t})^{d+1} \left(\frac{Z_{2V}}{Z_V^2} m_1(\Phi_0) + \sqrt{-2 \log(\mu(0)) + d \log\left(\frac{L}{2\pi}\right) + \sqrt{dL}} \right).$$

Finally, $m_1(\Phi_0) = \frac{(Z_V)^2}{Z_{2V}} \int \|z\| \frac{e^{-V(z)}}{\int e^{-V(z)}} dz = \frac{(Z_V)^2}{Z_{2V}} \sqrt{m_2(\mu)}$. \square

We can now prove Lemma 13.

Proof. We start by differentiating $m_2(\Phi_t)$:

$$\begin{aligned}
\partial m_2(\Phi_t) &= \int \|z\|^2 \partial \Phi_t dz \\
&= \int \|z\|^2 (\operatorname{div}(\nabla \Phi_t) - \langle \nabla \Phi_t, \nabla \log(\pi) \rangle + 2 \langle \nabla \log(p_t^V), \nabla \Phi_t \rangle) dz \\
&= - \int 2 \langle z, \nabla \Phi_t \rangle dz + \int (\Delta \log(\pi) - 2 \Delta \log(p_t^V)) \|z\|^2 \Phi_t dz + 2 \int \langle \nabla \log(\pi) - 2 \nabla \log(p_t^V), z \rangle \Phi_t dz \\
&= -2 \int \Delta \log(\pi) \Phi_t dz - \int \Delta \log(\pi) \|z\|^2 \Phi_t dz - 2 \int \langle \nabla \log(\pi), z \rangle \Phi_t dz \\
&\quad + 2 \int (\Delta \log(\pi) - \Delta \log(p_t^V)) \|z\|^2 \Phi_t dz + 4 \int \langle \nabla \log(\pi) - \nabla \log(p_t^V), z \rangle \Phi_t dz \\
&= 2dm_0(\Phi_t) + dm_2(\Phi_t) + 2m_2(\Phi_t) + 2 \int (\Delta \log(\pi) - \Delta \log(p_t^V)) \|z\|^2 \Phi_t dz + 4 \int \langle \nabla \log(\pi) - \nabla \log(p_t^V), z \rangle \Phi_t dz.
\end{aligned}$$

The first term $\int (\Delta \log(\pi) - \Delta \log(p_t^V)) \|z\|^2 \Phi_t dz$ is upper-bounded by $\frac{(L-1)e^{-2t}}{(1-e^{-2t})(L-1)+1} dm_2(\Phi_t)$ and for the second term we have

$$\begin{aligned}
\int \langle \nabla \log(\pi) - \nabla \log(p_t^V), z \rangle \Phi_t dz &= \int \langle \nabla \log\left(\frac{\pi}{p_t^V}\right)(z) - \nabla \log\left(\frac{\pi}{p_t^V}\right)(0), z \rangle \Phi_t dz \\
&\quad + \int \langle \nabla \log\left(\frac{\pi}{p_t^V}\right)(0), z \rangle \Phi_t dz \\
&\leq \int \frac{(L-1)e^{-2t}}{(1-e^{-2t})(L-1)+1} \|z\|^2 \Phi_t dz + \|\log(p_t^V)(0)\| \int \|z\| \Phi_t dz \\
&= \frac{(L-1)e^{-2t}}{(1-e^{-2t})(L-1)+1} m_2(\Phi_t) + \|\nabla \log(p_t^V)(0)\| m_1(\Phi_t).
\end{aligned}$$

Hence we recover

$$\partial m_2(\Phi_t) \leq (d+2) \left(1 + \frac{2(L-1)e^{-2t}}{(1-e^{-2t})(L-1)+1}\right) m_2(\Phi_t) + 4\|\nabla \log(p_t^V)(0)\| m_1(\Phi_t) + 2dm_0(\Phi_t).$$

We now use the Gronwall lemma to obtain

$$m_2(\Phi_t) \leq e^{t(d+2)} (L(1-e^{-2t}) + e^{-2t})^{d+2} \left(m_2(\Phi_0) + 2 \int_0^t \frac{e^{-s(d+2)} (2m_1(\Phi_s) \|\nabla \log(p_s^V)(0)\| + dm_0(\Phi_s))}{(L(1-e^{-2s}) + e^{-2s})^{d+2}} ds \right)$$

Recalling the upper-bound $m_1(\Phi_t) \leq e^{t(d+1)} (L(1-e^{-2t}) + e^{-2t})^{d+1} C$ where C is defined in Lemma 27, we upper-bound the integral term as

$$\begin{aligned}
\int_0^t \frac{e^{-s(d+2)} m_1(\Phi_s) \|\nabla \log(p_s^V)(0)\|}{(L(1-e^{-2s}) + e^{-2s})^{d+2}} ds &\leq C \int_0^t \frac{e^{-s}}{L(1-e^{-2s}) + e^{-2s}} \|\nabla \log(p_s^V)(0)\| ds \\
&\leq C \sqrt{\int_0^t \frac{e^{-2s}}{(L(1-e^{-2s}) + e^{-2s})^2} ds} \sqrt{\int_0^t \|\nabla \log(p_s^V)(0)\|^2 ds} \\
&= C \sqrt{\frac{1-e^{-2t}}{2}} \sqrt{\int_0^t \|\nabla \log(p_s^V)(0)\|^2 ds}.
\end{aligned}$$

Similarly,

$$\begin{aligned}
\int_0^t \frac{e^{-s(d+2)}}{(L(1-e^{-2s}) + e^{-2s})^{d+2}} m_0(\Phi_s) ds &\leq \int_0^t \frac{e^{-2s}}{(L(1-e^{-2s}) + e^{-2s})^2} ds \\
&= \frac{1-e^{-2t}}{2}.
\end{aligned}$$

Hence we recover that

$$m_2(\Phi_t) \leq e^{t(d+2)} (L(1-e^{-2t}) + e^{-2t})^{d+2} \left(m_2(\Phi_0) + 2C \sqrt{-2 \log(\mu(0)) + d \log\left(\frac{L}{2\pi}\right) + d} \right).$$

Using the expression of C , we recover eventually that

$$m_2(\Phi_t) \leq \frac{2e^{t(d+2)}Z_V^2}{Z_{2V}}(L(1 - e^{-2t}) + e^{-2t})^{d+2} \left[m_2(\mu) + d(L+1) - 4\log(\mu(0)) + 2d\log\left(\frac{L}{2\pi}\right) \right].$$

□

A.9 Proof of Theorem 14

Proof. Using Proposition 7 as well as the estimates Eq. 9 and Eq. 11, the integrated error reads

$$\begin{aligned} \int \delta(z)^2 dp_t^V(z) &\lesssim e^{td} \left[d\left(\beta + \frac{1}{1 - e^{-2t}}\right) + \xi^2 + 1 \right] \frac{(d + \log(n))Z_{2V}}{n(1 - e^{-2t})(Z_V)^2} \int \Phi_t(z) dz \\ &\quad + \frac{e^{td}Z_{2V}(d + \log(n))(L + \beta + 1)^2}{n(Z_V)^2(1 - e^{-2t})} \int \|z\|^2 \Phi_t(z) dz. \end{aligned}$$

We use Lemma 10 for the first term and Lemma 13 for the second term and we obtain

$$\begin{aligned} \int \delta(z)^2 dp_t^V(z) &\lesssim \frac{e^{2td}(d + \log(n))(L + 1)^d}{n(1 - e^{-2t})} \left(\left[d\left(\beta + \frac{1}{1 - e^{-2t}}\right) + \xi^2 + 1 \right] \right. \\ &\quad \left. + (L + \beta + 1)^2(L + 1)^2 e^{2t} \left[m_2(\mu) + d(L + 1) + 2d\log\left(\frac{L\mu(0)^{-2/d}}{2\pi}\right) \right] \right) \\ &\lesssim \frac{e^{2t(d+1)}(d + \log(n))(L + 1)^d}{n(1 - e^{-2t})^2} \left([d\beta + \xi^2 + 1] \right. \\ &\quad \left. + (L + \beta + 1)^2(L + 1)^2 \left[m_2(\mu) + d(L + 1) + 2d\log\left(\frac{L\mu(0)^{-2/d}}{2\pi}\right) \right] \right). \end{aligned}$$

□

A.10 Proof of Corollary 15

Proof. We run the recursion Eq. 6 with $T = \log((d + m_2(\mu))/\epsilon)$, $N = \frac{d^2(\log(L)+T)}{\epsilon}$, t_k as indicated in Theorem 3, the approximate score $\hat{s}_{n_k, t_k}(z) = \frac{-1}{(1 - e^{-2t_k})} \frac{\sum_{i=1}^{n_k} y_i^k e^{-V((z - y_i^k)e^{t_k})}}{\sum_{i=1}^{n_k} e^{-V((z - y_i^k)e^{t_k})}}$ with $y_i^k \sim \mathcal{N}(0, (1 - e^{-2t_k})I_d)$ and

$$n_k = \frac{e^{2t_k(d+1)}d(L+1)^d}{\epsilon(1 - e^{-2t_k})^2} (K_1 + K_2) := C \frac{e^{2t_k(d+1)}}{(1 - e^{-2t_k})^2}.$$

By definition of n_k and the other parameters, Theorem 3 yields

$$\mathbb{E}[\text{KL}(\mu, \hat{p})] \lesssim \epsilon,$$

where \lesssim hides a universal constant and logarithmic factors in $\epsilon, L, \beta, d, m_2(\mu), \log(\mu(0))$. The total running time of the algorithm is given by

$$\begin{aligned} \sum_{k=1}^N n_k &= C \sum_{k=1}^N \frac{e^{2t_k(d+1)}}{(1 - e^{-2t_k})^2} \\ &\leq CN \frac{e^{2t_N(d+1)}}{(1 - e^{-2t_1})^2} \\ &= \frac{CN}{(1 - e^{-2t_1})^2} (m_2(\mu) + d)^{2(d+1)} \epsilon^{-2(d+1)} \\ &\lesssim \frac{CN}{t_1^2} (m_2(\mu) + d)^{2(d+1)} \epsilon^{-2(d+1)}. \end{aligned}$$

Using the recursion in Theorem 3, we have that $t_1 = \frac{c}{L}$ where we recall $c = (\log(L) + T)/N$. This yields $t_1 = \epsilon \log((m_2(\mu) + d)L/\epsilon)/(Ld^2)$ and a total complexity of

$$\begin{aligned} \sum_{i=1}^N n_k &\lesssim d^7 L^{d+2} \epsilon^{-2(d+3)} (m_2(\mu) + d)^{2(d+1)} [(L + \beta)^2 L^2 (m_2(\mu) + dL + 2d\log\left(\frac{L\mu(0)^{-2/d}}{2\pi}\right)) \\ &\quad + (d\beta + \xi^2)]. \end{aligned}$$

□

B Proofs of Section 4

B.1 Proof of Proposition 18

Before proving the result, we need the following intermediate result.

Lemma 28 *Let V be an L -smooth, α -strongly convex potential outside some ball centered $B_R(m)$. Then for any critical point x^* of V , it holds*

$$\langle \nabla V(x), x - x^* \rangle \geq \alpha \|x - x^*\|^2 - 2R(L + \alpha) \|x - x^*\|.$$

Proof. For all $x \in \mathbb{R}^d$, the following identity holds

$$\langle \nabla V(x), x - x^* \rangle = \int \langle x - x^*, \nabla^2 V(tx + (1-t)x^*)(x - x^*) \rangle. \quad (19)$$

Using the equation above, we split the analysis in the case where the critical point x^* belongs to $B_R(m)$ and in the case where it does not. In the first case, if $\|x - x^*\| \leq 2R$, we can coarsely lower bound as

$$\langle \nabla V(x), x - x^* \rangle \geq -L \|x - x^*\|^2.$$

In the case where $\|x - x^*\| > 2R$, note that whenever $t \geq \frac{2R}{\|x - x^*\|}$, it holds that

$$\begin{aligned} \|tx + (1-t)x^* - m\| &= \|x^* - m + t(x - x^*)\| \\ &\geq t\|x - x^*\| - \|x^* - m\| \\ &> R, \end{aligned}$$

in other words, $tx + (1-t)x^* \notin B_R(m)$. This allows to recover the lower bound

$$\begin{aligned} \langle \nabla V(x), x - x^* \rangle &\geq -L \|x - x^*\|^2 \frac{2R}{\|x - x^*\|} + \alpha \|x - x^*\|^2 \left(1 - \frac{2R}{\|x - x^*\|}\right) \\ &= \alpha \|x - x^*\|^2 - 2R(L + \alpha) \|x - x^*\|. \end{aligned}$$

Hence we recover that

$$\begin{aligned} \langle \nabla V(x), x - x^* \rangle &\geq -\mathbf{1}_{x \in B_R(m)} L \|x - x^*\|^2 + \mathbf{1}_{x \notin B_R(m)} (\alpha \|x - x^*\|^2 - 2R(L + \alpha) \|x - x^*\|) \\ &\geq \alpha \|x - x^*\|^2 - 2R(L + \alpha) \|x - x^*\|. \end{aligned}$$

In the case where $x^* \notin B_R(m)$, let us introduce the cone

$$C = \{x \in \mathbb{R}^d \mid \exists t \in [0, 1], tx + (1-t)x^* \in B_R(m)\};$$

note that we indeed have $B_R(m) \subset C$. By definition of C , if $x \notin C$, it holds that

$$\langle \nabla V(x), x - x^* \rangle \geq \alpha \|x - x^*\|^2.$$

We now split the analysis in the case where $x \in C \cap B_R(m)$ and where $x \in C \cap \overline{B_R(m)}$. Denoting $x(t) = tx + (1-t)x^*$ we define $t_1 = \inf\{t \in [0, 1] \mid x(t) \in B_R(m)\}$; remark that $t_1 = \frac{\|x(t_1) - x^*\|}{\|x - x^*\|}$. By definition of t_1 , it holds for any $t < t_1$ that $x(t) \notin B_R(m)$. Thus, if $x \in C \cap B_R(m)$, we get the following lower-bound

$$\begin{aligned} \langle \nabla V(x), x - x^* \rangle &\geq \alpha \|x - x^*\|^2 \frac{\|x(t_1) - x^*\|}{\|x - x^*\|} - L \|x - x^*\|^2 \left(1 - \frac{\|x(t_1) - x^*\|}{\|x - x^*\|}\right) \\ &\geq \alpha \|x - x^*\|^2 \frac{\|x - x^*\| - \|x - x(1)\|}{\|x - x^*\|} - L \|x - x^*\|^2 \left(1 - \frac{\|x - x^*\| - \|x - x(1)\|}{\|x - x^*\|}\right) \\ &\geq \alpha \|x - x^*\|^2 - 2R(L + \alpha) \|x - x^*\|. \end{aligned}$$

Now, if $x \in C$, note that $t < \frac{\|m - x^*\| - R}{\|x - x^*\|}$, then it holds that

$$\begin{aligned} \|tx + (1-t)x^* - m\| &\geq \|x^* - m\| - t\|x - x^*\| \\ &> R. \end{aligned}$$

Furthermore, introducing for $x \in \mathbb{R}^d$ the projection $p_B(x)$ onto the ball $B_R(m)$ we recall that $p_B(x) - x$ is (positively) colinear to $m - p_B(x)$. Thus it holds that

$$\begin{aligned} \frac{\|m - x^*\| - R}{\|x - x^*\|} &= \frac{\|p_B(x^*) - x^*\| + \|p_B(x^*) - m\| - R}{\|x - x^*\|} \\ &= \frac{\|p_B(x^*) - x^*\|}{\|x - x^*\|}. \end{aligned}$$

Ne now prove that $\langle x - p_B(x^*), p_B(x^*) - x^* \rangle \geq 0$. By the absurd, assume that $\langle x - p_B(x^*), p_B(x^*) - x^* \rangle < 0$, then

$$\langle tx + (1-t)x^* - p_B(x^*), p_B(x^*) - x^* \rangle < 0.$$

In particular, for all $t \in [0, 1]$, it holds that

$$\begin{aligned} \|tx + (1-t)x^* - m\|^2 &= \|m - p_B(x^*) + p_B(x^*) - (tx + (1-t)x^*)\|^2 \\ &= \|m - p_B(x^*)\|^2 + 2\langle m - p_B(x^*), p_B(x^*) - (tx + (1-t)x^*) \rangle \\ &\quad + \|p_B(x^*) - (tx + (1-t)x^*)\|^2 \\ &= \|m - p_B(x^*)\|^2 + \|p_B(x^*) - (tx + (1-t)x^*)\|^2 \\ &\quad + 2\frac{\|m - p_B(x^*)\|}{\|p_B(x^*) - x^*\|} \langle p_B(x^*) - x^*, p_B(x^*) - (tx + (1-t)x^*) \rangle \\ &> R^2, \end{aligned}$$

which contradicts $x \in C$. Hence $\langle tx + (1-t)x^* - p_B(x^*), p_B(x^*) - x^* \rangle \geq 0$ which implies

$$\begin{aligned} \|x - x^*\|^2 &= \|x - p_B(x^*)\|^2 + 2\langle x - p_B(x^*), p_B(x^*) - x^* \rangle + \|p_B(x^*) - x^*\|^2 \\ &\geq \|p_B(x^*) - x^*\|^2. \end{aligned}$$

In particular, we are ensured that $\frac{\|p_B(x^*) - x^*\|}{\|x - x^*\|} \leq 1$. We now split the analysis in the case where $x \in B_R(m) \cap C$ and where $x \in \overline{B_R(m)} \cap C$. In the first case, the previous computation implies the following lower-bound:

$$\begin{aligned} \langle \nabla V(x), x - x^* \rangle &\geq \alpha \|x - x^*\| (\|m - x^*\| - R) - L(R + \|x - x^*\| - \|m - x^*\|) \|x - x^*\| \\ &\geq \alpha \|x - x^*\| (\|x - x^*\| - \|x - m\| - R) \\ &\quad - L(R + \|x - x^*\| - \|x - x^*\| + \|m - x\|) \|x - x^*\| \\ &\geq \alpha \|x - x^*\|^2 - 2R(L + \alpha) \|x - x^*\|. \end{aligned}$$

Conversely, in the case where $x \in C \cap \overline{B_R(m)}$, define $t_2 = \sup\{t \in [0, 1] \mid x(t) \in B_R(m)\}$; again note that we have $t_2 = \frac{\|x(t_2) - x^*\|}{\|x - x^*\|}$. By definition, note that for any $t > t_2$, $x(t) \notin B_R(m)$ hence we get the lower-bound

$$\begin{aligned} \langle \nabla V(x), x - x^* \rangle &= \int_0^{t_1} \langle (x - x^*), \nabla^2 V(tx + (1-t)x^*)(x - x^*) \rangle \\ &\quad + \int_{t_1}^{t_2} \langle (x - x^*), \nabla^2 V(tx + (1-t)x^*)(x - x^*) \rangle \\ &\quad + \int_{t_2}^1 \langle (x - x^*), \nabla^2 V(tx + (1-t)x^*)(x - x^*) \rangle \\ &\geq \alpha \|x - x^*\|^2 \left(1 - \frac{\|x(t_2) - x^*\|}{\|x - x^*\|} + \frac{\|x(t_1) - x^*\|}{\|x - x^*\|} \right) \\ &\quad - L \|x - x^*\| (\|x(t_2) - x^*\| - \|x(t_1) - x^*\|) \\ &\geq \alpha \|x - x^*\|^2 - 2R(L + \alpha) \|x - x^*\|. \end{aligned}$$

Hence we recover for all three cases $x \notin C$, $x \in C \cap B_R(m)$ and $x \in C \cap \overline{B_R(m)}$ that

$$\langle \nabla V(x), x - x^* \rangle \geq \alpha \|x - x^*\|^2 - 2R(L + \alpha) \|x - x^*\|.$$

□

We can now prove our result.

Proof. Using Lemma 28, we have the lower bound

$$\langle \nabla V(x), x - x^* \rangle \geq \alpha \|x - x^*\|^2 - 2R(L + \alpha) \|x - x^*\|.$$

Using the Laplacian identity for $f(x) = \|x - x^*\|^2$, we get that

$$\int \langle \nabla V(x), x - x^* \rangle d\mu(x) = d,$$

which combined with the upper-bound above yields

$$\int \alpha \|x - x^*\|^2 d\mu(x) \leq 2R(L + \alpha) \int \|x - x^*\| d\mu(x) + d.$$

Denoting $\theta^2 = \mathbb{E}_{X \sim \mu}[\|X - x^*\|^2]$, it holds by Cauchy-Schwarz that

$$\theta^2 \leq \frac{2R(L + \alpha)}{\alpha} \theta + \frac{d}{\alpha}.$$

This implies that θ^2 is upper-bounded as

$$\begin{aligned} \theta^2 &\leq \frac{1}{2} \left(\frac{2R(L + \alpha)}{\alpha} + \sqrt{\frac{4R^2(L + \alpha)^2}{\alpha^2} + \frac{4d}{\alpha}} \right)^2 \\ &\leq \left(\frac{4R^2(L + \alpha)^2}{\alpha^2} + \frac{4R^2(L + \alpha)^2}{\alpha^2} + \frac{4d}{\alpha} \right) \\ &= \frac{8R^2(L + \alpha)^2}{\alpha^2} + \frac{4d}{\alpha}. \end{aligned}$$

□

B.2 Proof of Corollary 20

Proof. We start by using Proposition 17 which ensures that as long as $1 - e^{-2t} \leq \frac{\epsilon^{-t}}{2L}$, it holds that

$$\|\nabla^2 \log(p_t^V)\| \leq 2Le^t.$$

In other terms, as long as

$$e^t \leq \frac{1 + \sqrt{16L^2 + 1}}{4L},$$

it holds that

$$\|\nabla^2 \log(p_t^V)\| \leq \frac{1 + \sqrt{1 + 16L^2}}{2}.$$

Furthermore we recall that Proposition 9 yields the (uniform-in-time) lower-bound

$$\nabla^2 \log(p_t^V) \succeq -(L + 1)I_d.$$

Finally, in the case where $1 - e^{-2t} > \frac{\epsilon^{-t}}{2L}$, we have in particular that

$$\frac{e^{-2t}}{(1 - e^{-2t})^2} \leq 4L^2,$$

which combined with equation 14 yields

$$\nabla^2 \log(p_t^V) \preceq 4L^2 \left[8R^2 \left(\frac{L + 2\alpha}{\alpha} \right)^2 + \frac{4d}{\alpha} + 1 \right] I_d.$$

Hence, it holds for all t that

$$\|\nabla^2 \log(p_t^V)\| \leq 4(L + 1)^2 \left[8R^2 \left(\frac{L + 2\alpha}{\alpha} \right)^2 + \frac{4d}{\alpha} + 1 \right].$$

□

B.3 Proof of Proposition 22

We first prove the estimate on the gradient.

Proof. Recall that for all $z \in \mathbb{R}^d$, the score of the current distribution p_t reads

$$\nabla \log(p_t)(z) = \frac{-1}{(1 - e^{-2t})} \frac{\int (z - xe^{-t}) e^{-\frac{\|z - xe^{-t}\|^2}{2(1 - e^{-2t})}} e^{-V(x)} dx}{\int e^{-\frac{\|z - xe^{-t}\|^2}{2(1 - e^{-2t})}} e^{-V(x)} dx}.$$

In particular, at $z = 0$, the score can be re-written as an expectation

$$\nabla \log(p_t)(0) = \frac{e^{-t}}{1 - e^{-2t}} \mathbb{E}_{q_t}[X],$$

with $q_t \propto e^{-\frac{\|xe^{-t}\|^2}{2(1 - e^{-2t})}} e^{-V(x)}$. Note that q_t is also α -strongly log-concave outside some ball of radius R , has condition number at most κ and is such that $\nabla \log(q_t)(0) = 0$. Thus, Proposition 18 applies and we recover by Jensen inequality

$$\|\nabla \log(p_t)(0)\|^2 \leq \frac{e^{-2t}}{(1 - e^{-2t})^2} (8R^2(\kappa^2 + 1) + \frac{4d}{\alpha}).$$

We now recall the alternative formula for the score

$$\nabla \log(p_t)(z) = \frac{-e^t}{(1 - e^{-2t})} \frac{\int \nabla V((z - x)e^t) e^{-V((z - x)e^t)} e^{-\frac{\|x\|^2}{2(1 - e^{-2t})}} dx}{\int e^{-V((z - x)e^t)} e^{-\frac{\|x\|^2}{2(1 - e^{-2t})}} dx}.$$

Making the change of variable $u = (z - x)e^t$ for both the numerator and the denominator, we recover

$$\nabla \log(p_t)(z) = \frac{-e^t}{(1 - e^{-2t})} \frac{\int \nabla V(u) e^{-V(u)} e^{-\frac{\|z - e^{-t}u\|^2}{2(1 - e^{-2t})}} du}{\int e^{-V(u)} e^{-\frac{\|z - e^{-t}u\|^2}{2(1 - e^{-2t})}} du}.$$

In particular, for $z = 0$ we get

$$\begin{aligned} \|\nabla \log(p_t)(0)\|^2 &\leq e^{2t} \mathbb{E}_{q_t}[\|\nabla V(X)\|^2] \\ &\leq L^2 e^{2t} (8R^2(\kappa^2 + 1) + \frac{4d}{\alpha}). \end{aligned}$$

Finally, remark that $t \mapsto \frac{e^{-t}}{1 - e^{-2t}}$ decreases and $t \mapsto e^t$ increases. When we evaluate both functions at $t = \frac{1}{2} \log((L + 1)/L)$ we recover in both cases $\sqrt{L}\sqrt{L + 1}$ which implies that for all $t > 0$, we have

$$\|\nabla \log(p_t^V)\|^2 \leq (L + 1)^2 (8R^2(\kappa^2 + 1) + \frac{4d}{\alpha}).$$

□

Before proving the second estimate, we need this intermediary result.

Lemma 29 *Let V be an L -smooth potential that is α -strongly convex outside some ball of radius R and such that $\nabla V(0) = 0$. It holds that,*

$$V(x) - V(0) \geq \frac{\alpha}{2} \|x\|^2 - R(L + \alpha) \|x\|.$$

Proof. We write the difference $V(x) - V(0)$ in integral form:

$$\begin{aligned} V(x) - V(0) &= \int_0^1 \langle \nabla V(tx), x \rangle dt \\ &= \int_0^1 \frac{1}{t} \langle \nabla V(tx), tx \rangle dt. \end{aligned}$$

Now we use Lemma 28 with $x^* = 0$ to recover

$$\begin{aligned} V(x) - V(0) &\geq \int_0^1 \frac{1}{t} (\alpha t^2 \|x\|^2 - tR(L + \alpha)\|x\|) dt \\ &= \frac{\alpha}{2} \|x\|^2 - R(L + \alpha)\|x\|. \end{aligned}$$

We can now prove the second estimate. □

Proof. We recall that

$$\frac{d\mu(x)}{dx} = \frac{e^{-V(x)}}{\int e^{-V(x)} dx},$$

which can be re-written as

$$\frac{d\mu(x)}{dx} = \frac{e^{-(V(x)-V(0))}}{\int e^{-(V(x)-V(0))} dx}.$$

By smoothness and since we assumed $\nabla V(0) = 0$, the numerator can be lower-bounded by $e^{-L\|x\|^2}$. To upper bound the denominator, we use the previous result to recover

$$\begin{aligned} \int e^{-(V(x)-V(0))} dx &\leq \int e^{-\frac{\alpha}{2}\|x\|^2 + R(L+\alpha)\|x\|} dx \\ &= \int e^{-\frac{\alpha}{4}\|x\|^2 - \frac{\alpha}{4}\|x\|^2 + R(L+\alpha)\|x\|} dx. \end{aligned}$$

Now observe that we have for all x

$$-\frac{\alpha}{4}\|x\|^2 + R(L + \alpha)\|x\| \leq \frac{R^2(L + \alpha)^2}{\alpha},$$

which implies

$$\begin{aligned} \int e^{-(V(x)-V(0))} dx &\leq \int e^{\frac{R^2(L+\alpha)^2}{\alpha}} e^{-\frac{\alpha}{4}\|x\|^2} dx \\ &= e^{\frac{R^2(L+\alpha)^2}{\alpha}} \left(\frac{4\pi}{\alpha}\right)^{d/2}. \end{aligned}$$

Hence we finally recover

$$\frac{d\mu(x)}{dx} \geq e^{-L\|x\|^2} e^{-\frac{R^2(L+\alpha)^2}{\alpha}} \left(\frac{4\pi}{\alpha}\right)^{-d/2}.$$

□

B.4 Proof of Corollary 23

Proof. The estimates in Proposition 22 and in Corollary 20 yield

$$K_1 \lesssim L^4 \left(R^4 \kappa^4 + \frac{d^2}{\alpha^2}\right) L^2 \left(R^2 \kappa^2 + d\left(\frac{1}{\alpha} + L\right)\right) \asymp L^6 \left(R^2 \kappa^2 + \frac{d}{\alpha}\right)^3,$$

and

$$K_2 \lesssim L^2 \left(R^2 \kappa^2 + \frac{d}{\alpha}\right) + dL^2 \left(R^2 \kappa^2 + \frac{d}{\alpha}\right) \asymp dL^2 \left(R^2 \kappa^2 + \frac{d}{\alpha}\right).$$

Hence we obtain that $K_1 + K_2 \asymp L^6 \left(R^2 \kappa^2 + \frac{d}{\alpha}\right)^3$ and yields as a total complexity

$$\sum_{k=1}^N n_k \lesssim d^7 L^{d+8} \epsilon^{-2(d+3)} \left(R^2 \kappa^2 + \frac{d}{\alpha}\right)^{2(d+4)}.$$

□

C Discussion on concurrent work

In Ding et al. [18], the authors assume that there exists a *known* Gaussian distribution $\pi^{\eta, \Sigma} \sim \mathcal{N}(\eta, \Sigma)$ such that $d\mu/d\pi^{\eta, \Sigma}$ is (i) strictly bounded from above and (ii) is uniformly Lipschitz. In the simple case where μ is a Gaussian those two conditions imply that necessarily $\pi^{\eta, \Sigma} = \mu$. In the case where $\mu = 0.5\mathcal{N}(m, 1) + 0.5\mathcal{N}(-m, 1)$, condition (i) implies $\Sigma \geq 1$ and condition (ii) implies $\Sigma \leq 1$ which yields $\Sigma = 1$. The ratio thus becomes

$$\frac{d\mu}{d\pi^{\beta, \Sigma}} = e^{(\eta^2 - m^2)/2} \frac{e^{xm/2} + e^{-xm/2}}{e^{-\eta x/2}}.$$

In particular, there exists no value of η such that the ratio is uniformly Lipschitz.