

PARTITION-THEORETIC MODEL OF PRIME DISTRIBUTION

AIDAN BOTKIN, MADELINE L. DAWSEY, DAVID J. HEMMER
MATTHEW R. JUST AND ROBERT SCHNEIDER

ABSTRACT. We make an application of ideas from partition theory to a problem in multiplicative number theory. We posit a deterministic model of prime number distribution from first principles related to properties of integer partitions, that predicts the prime number theorem, and suggests new conjectures regarding predictable variations in prime gaps. The model predicts that, for $n \geq 2$,

$$p_n = 1 + 2 \sum_{k=1}^{n-1} \left\lfloor \frac{d(k)}{2} \right\rfloor + \varepsilon(n),$$

where p_k is the k th prime number, $d(k)$ is the divisor function, and $\varepsilon(k) \geq 0$ is an error that is negligible asymptotically; both the main term and error are enumerative functions in our conceptual model. We refine the error to give numerical estimates of $\pi(n)$ similar to those provided by the logarithmic integral, and much more accurate than $\text{li}(n)$ up to $n = 10,000$ where the estimate is *almost exact*. We also test a seemingly dubious prediction suggested by the model, that prime-indexed primes p_2, p_3, p_5, p_7 , etc., are more likely to be involved in twin prime pairs than are arbitrary primes; surprisingly, computations show prime-indexed primes *are* more likely to be twin primes up to at least $n = 10,000,000,000$, with the bias diminishing but persistent as n increases.

1. INTRODUCTION

1.1. Model of primes from partition first principles. The prime numbers mimic a random sequence in key ways, and many successful models of prime distribution proceed from probabilistic hypotheses [30]. In his lecture notes [17], Terence Tao writes compellingly as follows:

We do have a number of extremely convincing and well supported [random] models for the primes (and related objects) that let us predict what the answer to many prime number theory questions (both multiplicative and non-multiplicative) should be.... Indeed, the models are so effective at this task that analytic number theory is in the curious position of being able to confidently predict the answer to a large proportion of the open problems in the subject, whilst not possessing a clear way forward to rigorously confirm these answers!

In this paper, a sequel to [9], we formulate a *deterministic* model of prime number distribution based on information about the integers gleaned from facts about integer partitions.¹ In short, the comparison of two multiplicative statistics defined on partitions, the norm [27] and supernorm [9], imposes constraints on the number line unrelated to integer factorizations, which we leverage to give an estimate of the n th prime gap. Computations show the model to be reasonably accurate numerically. We justify the model from partition-theoretic first principles, and refine the error to derive an explicit prediction for the value of the n th prime number which is surprisingly good for small numbers, up to $n = 10,000$, yielding an estimate for $\pi(n)$ (the number of primes less than or equal to n) that is *almost exact* when n is small; which predicts the prime number theorem asymptotically as n increases; and which makes predictions about local behaviors of prime gaps that would be difficult to deduce without ideas about partitions. We give computational evidence for one of these predictions. Finally, we note deficiencies in the model, and suggest further refinements.

¹Early versions of this model were presented previously by some of the authors in [13] and the appendix of [8].

Here is the central claim of our model, which we formulated from first principles about partitions. For $x \in \mathbb{R}$, let $\lceil x \rceil$, $\lfloor x \rfloor$ denote the *ceiling* and *floor functions*; we define $\lfloor x \rfloor := 0$ if $x < 0$.

Partition model of prime numbers. *The n th prime $p_n \in \mathbb{P}$, $n \geq 2$, is modeled by the formula*

$$p_n = 1 + 2 \sum_{k=1}^{n-1} \left\lfloor \frac{d(k)}{2} \right\rfloor + \varepsilon(n),$$

where $d(k)$ is the divisor function and $\varepsilon(k) \geq 0$ is an explicit error term that is negligible by comparison. The case $\varepsilon(n) := 0$ for all $n \geq 1$ we refer to as *Model 1*. More nuanced consideration of the error in *Model 1* leads to a refined model we call *Model 2*, which is the case

$$\varepsilon(n) := \lfloor \pi_2(p_{n-1}) - 2\gamma n \rfloor,$$

where $\pi_2(k)$ is the number of semiprimes less than or equal to $k \geq 1$, and $\gamma = 0.5772\dots$ is the Euler-Mascheroni constant. For computations, we use an approximation to the error term,

$$\varepsilon(n) := \lfloor n \log \log n - 2\gamma n \rfloor,$$

to yield a computational model we call *Model 2**.

The reader is referred to [Table 1](#) that gives a comparison of this model's predictions for $\pi(n)$, alongside corresponding predictions of the prime number theorem.

n	$\pi(n)$	$n/\log n$	$\text{li}(n)$	Model 1	Model 2*
10	4	4.34...	6.16...	4	4
100	25	21.71...	30.12...	27	25
1000	168	144.76...	177.60...	183	169
10,000	1229	1085.73...	1246.13...	1351	1231
100,000	9592	8685.88...	9629.80...	10,601	9616
1,000,000	78,498	72,382.41...	78,627.54...	86,738	78,738

TABLE 1. Comparing estimates for $\pi(n)$

1.2. Concepts and notations. Let \mathbb{Z}^+ denote the *natural numbers*. Let \mathbb{P} denote the *prime numbers*. Let $p_i \in \mathbb{P}$ denote the i th prime number, viz. $p_1 = 2, p_2 = 3, p_3 = 5$, etc.; we define $p_0 := 1$, and we refer to the subscript $i \in \mathbb{Z}^+$ of $p_i \in \mathbb{P}$ as the *index* of the prime number. For $n \in \mathbb{Z}^+$, we write its *prime factorization* as $n = p_1^{a_1} p_2^{a_2} p_3^{a_3} \cdots p_i^{a_i} \cdots$, $a_i \in \mathbb{Z}_{\geq 0}$, where only finitely many primes p_i have nonzero *multiplicity* (number of repetitions); we omit the factor p_i from the notation if $a_i = 0$. For $n \in \mathbb{Z}^+$, let $\pi(n)$ denote the *number of primes less than or equal to n* .

Let \mathcal{P} denote the set of *integer partitions*, unordered finite multisets of natural numbers including the empty partition $\emptyset \in \mathcal{P}$ (see [\[3\]](#)). For nonempty partition $\lambda \in \mathcal{P}$, we notate $\lambda = (\lambda_1, \lambda_2, \lambda_3, \dots, \lambda_r)$, with the *parts* $\lambda_i \in \mathbb{Z}^+$ written in weakly decreasing order $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_r \geq 1$. Similarly to prime factorizations, we also write partitions in *part-multiplicity* notation $\lambda = \langle 1^{m_1} 2^{m_2} 3^{m_3} \dots i^{m_i} \dots \rangle$, omitting any part $i \geq 1$ if $m_i = 0$. For $\lambda \in \mathcal{P}$, let $|\lambda| \geq 0$ denote the *size* (sum of parts), let $\ell(\lambda) = r$ be the *length* (number of parts), and let $m_i = m_i(\lambda) \geq 0$ be the *multiplicity* of $i \in \mathbb{Z}^+$ as a part of partition λ . Let $p(n)$ be the *number of partitions of size $n \geq 0$* .

There are further important *partition statistics*, i.e., functions $f: \mathcal{P} \rightarrow \mathbb{Z}$, such as the Dyson rank, the Andrews-Garvan crank, et al., that encode combinatorial properties in their values, and often enjoy modular congruence relations and other nice behaviors [\[3\]](#). Our model of prime distribution arises from consideration of two newly-defined, *multiplicative* partition statistics.

1.3. Multiplicative partition statistics and a multiplicative theory of (additive) partitions. In [22, 23, 25] and a series of subsequent publications, the fifth author (Schneider) introduced a *multiplicative* theory of integer partitions that parallels multiplicative number theory in many respects,² and studied a variety of new, multiplicative partition statistics – many of them representing partition-theoretic analogues of well-known arithmetic functions like the Möbius function $\mu(n)$, the Euler phi function $\varphi(n)$, and other functions from multiplicative number theory [4]. In these works, the fifth author presents (and proves many cases of) the following philosophy:

Theorems in multiplicative number theory are special cases of more general theorems in partition theory. Theorems in \mathbb{Z}^+ have counterpart theorems in \mathcal{P} , and vice versa. Facts about partitions map to facts about integers, which may be difficult to deduce without appealing to partition theory.

First applications of this philosophy were chiefly connected to the theories of Dirichlet series through the study of new classes of partition zeta functions [19, 22, 26], and to q -series and quasimodular forms through work on the q -bracket of Bloch and Okounkov [23, 24]. In [20, 21], Ono, Wagner and the fifth author make a computational application to give limit formulas for arithmetic densities using partition-theoretic q -series, that are comparable to those using Dirichlet series; this is generalized in [28]. A particularly stunning, recent application of this philosophy is made by Craig, Ono and van Ittersum [7], in which the partition-theoretic coefficients of certain quasimodular forms are proved to be intrinsically connected to the prime numbers.³

In [23, 25], a *partition multiplication* operation is introduced; this comes with a concept of “subpartitions” analogous to partition divisors, and a theory of partition Dirichlet convolution.

Definition 1. For two partitions $\lambda, \gamma \in \mathcal{P}$, we define the *partition product* $\lambda \cdot \gamma \in \mathcal{P}$ (or simply $\lambda\gamma$) to be the partition obtained by concatenating the parts of λ and γ (and then reordering by size by notational convention). For instance, $(5, 3, 2) \cdot (4, 3, 1, 1) = (5, 4, 3, 3, 2, 1, 1)$.⁴

A new multiplicative partition statistic is introduced in [22] that is central to the study of partition zeta functions, a multiplicative analogue of the size $|\lambda|$ called the *norm* $N(\lambda)$ of $\lambda \in \mathcal{P}$.

Definition 2. The *norm* of an integer partition $\lambda = (\lambda_1, \lambda_2, \lambda_3, \dots, \lambda_r)$ is the product of its parts:

$$(1) \quad N(\lambda) := \lambda_1 \lambda_2 \lambda_3 \cdots \lambda_r = 1^{m_1} 2^{m_2} 3^{m_3} \cdots i^{m_i} \cdots \in \mathbb{Z}^+;$$

we define $N(\emptyset) := 1$ (it is an empty product).

Note that $N(\lambda\gamma) = N(\lambda)N(\gamma)$ for $\lambda, \gamma \in \mathcal{P}$; see [27] for more about the partition norm. Along these lines, the multiplicative and additive branches of number theory enjoy many further analogies.

1.4. Isomorphism between partitions and natural numbers. In [9], the present authors Dawsey, Just and Schneider define another multiplicative partition statistic, the *supernorm* $\widehat{N}(\lambda)$.

Definition 3. The *supernorm* of an integer partition $\lambda = (\lambda_1, \lambda_2, \lambda_3, \dots, \lambda_r)$ is the product

$$(2) \quad \widehat{N}(\lambda) := p_{\lambda_1} p_{\lambda_2} \cdots p_{\lambda_r} = 2^{m_1} 3^{m_2} 5^{m_3} \cdots p_i^{m_i} \cdots \in \mathbb{Z}^+,$$

where $p_i \in \mathbb{P}$ is the i th prime number, $i \geq 1$, and $m_j = m_j(\lambda) \geq 0$ as above; we define $\widehat{N}(\emptyset) := 1$.

Note that $\widehat{N}(\lambda\gamma) = \widehat{N}(\lambda)\widehat{N}(\gamma)$ for $\lambda, \gamma \in \mathcal{P}$. In [9], it is proved that \widehat{N} induces an isomorphism of monoids.

Theorem (Dawsey–Just–Schneider). *The supernorm map $\widehat{N}: \mathcal{P} \rightarrow \mathbb{N}$ is an isomorphism between monoid (\mathcal{P}, \cdot) with partition multiplication and monoid (\mathbb{Z}^+, \cdot) with integer multiplication.*

²Influenced by ideas of Alladi-Erdős [1], Andrews [2], Granville [11], Granville-Soundararajan [12] and Ono [18].

³Follow-up works such as [6, 10] apply similar ideas to prime powers.

⁴Equivalently, one sums the multiplicities of corresponding parts in the two partitions.

It is the central theme of [9] that *the supernorm translates facts about partitions to analogous facts about integers, with partition parts mapping to prime factors, and vice versa*. For example, in [9], bijections between integers with certain prime factorizations, and partition formulas for natural densities of subsets of prime numbers, are proved. In a follow-up note [29], the fifth author extends partition multiplication to the set \mathcal{O} of overpartitions (see e.g. [5]),⁵ and proves that an extension $\widehat{N}_{\mathcal{O}}: \mathcal{O} \rightarrow \mathbb{Q}^+$ of the supernorm to overpartitions induces a *group* isomorphism between (\mathcal{O}, \cdot) with partition multiplication, and (\mathbb{Q}^+, \cdot) with rational multiplication. In [9], J. Lagarias proves further, striking properties of the supernorm that uniquely characterize the map \widehat{N} in a lattice-theoretic context [14]. Moreover, Lagarias and Sun prove interesting asymptotic and statistical relations using sums of the reciprocal of the supernorm over different finite subsets of \mathcal{P} in [15].

1.5. Comparing the norm and supernorm. Immediately one wonders about the magnitude of the supernorm compared to the partition norm; some analysis to this effect is undertaken in [9]. Here are some easy relations. First, a partition λ with no part equal to 1 respects the inequalities

$$(3) \quad \ell(\lambda) \leq |\lambda| \leq N(\lambda) \leq \widehat{N}(\lambda).$$

A simple observation about the supernorm versus the norm that one can make from direct computation⁶ is as follows: *partitions with greater lengths or having many parts equal to 1, have supernorms a lot larger than their norms; furthermore, partitions with fewer parts and no part equal to 1, have supernorms closer to their norms*. This is compatible with the prime number theorem, although we do not use the prime number theorem to formulate our model. Recall the well-known asymptotic estimate $p_n \sim n \log n$ as n increases, which is an equivalent statement of the prime number theorem. Substituting this estimate for primes $p_i \geq 3$ in equation (2) suggests

$$(4) \quad \widehat{N}(\lambda) \approx 2^{m_1(\lambda)} \prod_{i \geq 2} (i \log i)^{m_i(\lambda)} \approx N(\lambda) \cdot 2^{m_1(\lambda)} \prod_{i \geq 2} (\log i)^{m_i(\lambda)}, \quad i \geq 2.$$

This approximation has considerable error at smaller indices $i \geq 2$. However, it does indicate the above empirical observation is compatible with known estimates of prime distribution.

Moreover, in [9] it is proved that if λ has no part equal to 1, then

$$(5) \quad N(\lambda) \leq \widehat{N}(\lambda) \leq N(\lambda)^{\frac{\log 3}{\log 2}},$$

noting $\log 3 / \log 2 = 1.5849 \dots$, so the supernorm of the partition is not far off in order of magnitude from the norm. These inequalities tell us that *partitions having fixed norm and no part equal to 1 are mapped into a relatively small interval on the number line by the supernorm*.

We observe, further, that if λ is an unrestricted partition of size $|\lambda| = n$ and p_n is the n th prime number, then the supernorm falls on the interval

$$(6) \quad p_n \leq \widehat{N}(\lambda) \leq 2^n,$$

with equality on the left-hand side when $\lambda = \langle n^1 \rangle$, and equality on the right when $\lambda = \langle 1^n \rangle$, which is proved in [9]. By similar reasoning, if λ is a partition with no part equal to 1, $|\lambda| = n$, then

$$(7) \quad p_n \leq \widehat{N}(\lambda) \leq 3^{n/2},$$

with equality on the right when n is even and $\lambda = \langle 2^{n/2} \rangle$. The above inequalities suggest *partitions of fixed size are mapped across a relatively large interval on the number line by the supernorm*, by comparison with partitions of fixed norm.

Combining the inequalities above, we see for λ a partition with no part equal to 1,

$$(8) \quad p_{|\lambda|} \leq \widehat{N}(\lambda) \leq N(\lambda)^{\frac{\log 3}{\log 2}}.$$

⁵As cited in [29], the overpartition product was suggested to the fifth author by P. Cuthbertson and H. Waldron.

⁶Thanks to Alexander Walker for computing tables of the norm and supernorm in Python for the undergraduate project ‘‘A map between binary numbers and partitions’’ mentored by Schneider and Just (Univ. of Georgia, 2020).

One further observation we have made computationally but have not proved, is that for λ a partition with no part equal to 1, the following inequality *almost always* holds:

$$(9) \quad p_{N(\lambda)} \leq \widehat{N}(\lambda) \text{ almost always (conjecture).}$$

This is compatible with the prime number theorem since $p_{ij} \sim ij \log(ij) \leq ij(\log i)(\log j) \sim p_i p_j$ as i and j increase, and is not incompatible with (8), since $|\lambda| \leq N(\lambda)$ by (3) implies that $p_{|\lambda|} \leq p_{N(\lambda)}$. Inequality (9) suggests *partitions of fixed norm equal to n almost always have supernorms lying above p_n* (similarly to fixed size above). We treat (9) as an empirical observation, as well.

From these inequalities, the central observation of this paper comes into view: *comparison of the partition norm and supernorm imposes constraints on the prime numbers*. For instance, from (8) we can deduce an upper bound for the n th prime number.

Proposition 4. *For $n \geq 2$, we have that*

$$p_n \leq n^{\frac{\log 3}{\log 2}}.$$

Proof. Take $\lambda = (n)$, $n \geq 2$; then substitute $|\lambda| = N(\lambda) = n$ on the left and right sides of (8). \square

In Section 2, we formulate a model of primes based on the above observations about partitions.

2. PARTITION MODEL OF PRIME DISTRIBUTION

2.1. Partition-theoretic model of prime gaps. While the set \mathbb{P} is highly enigmatic in number theory and known for its seemingly stochastic growth, from the perspective of the supernorm, prime numbers have somewhat regular behavior: the n th prime p_n represents the supernorm $\widehat{N}(\lambda)$ of the partition $\lambda = (n)$ with a *single part* $n \in \mathbb{Z}^+$ (recalling \emptyset maps to 1). The sequence of primes is simply the image of the sequence of natural numbers under the map \widehat{N} , while the composite integers fill in the gaps in a more complicated way.

From this perspective, the complexities of prime distribution are a manifestation of the complex proliferation of *partitions with multiple parts*, which are mapped by the supernorm to *gaps between primes*. Thus prime gaps have a combinatorial interpretation: they are the images of certain subsets of \mathcal{P} under the supernorm map, which suggests the following heuristic.

Heuristic. *Measuring prime gaps is equivalent to enumerating partitions that map into the respective prime gaps under the supernorm \widehat{N} :*

$$p_{n+1} - p_n = \# \left\{ \lambda \in \mathcal{P} : p_n \leq \widehat{N}(\lambda) < p_{n+1} \right\}.$$

Our goal based on this heuristic is to identify partitions that map into each prime gap so we can count them, at least approximately. Below, we refine the above heuristic from elementary considerations to formulate an initial, if overly simplistic, partition model of prime gaps that agrees with the facts quite well. We then address sources of error to improve the model significantly.

2.2. Simplifying assumption. We wish to simplify the problem as much as possible. The goal is to estimate the length of the interval $[p_n, p_{n+1})$ with high accuracy for each $n \geq 1$. That is, we want to enumerate partitions that we expect to map just above p_n under the supernorm. Let us summarize observations we made in Section 1.5, that may aid us in identifying such partitions:

- (1) Partitions λ with a smaller number $\ell(\lambda)$ of parts and with no part equal to 1 should have supernorms closer to their norms in magnitude, based on considerations preceding (4).
- (2) Partitions λ with fixed norm $N(\lambda) = n$ and with no part equal to 1 should have supernorms of roughly comparable magnitudes to each other, based on consideration of (5).

- (3) Partitions λ with fixed norm $N(\lambda) = n$ and with no part equal to 1 should almost always respect $p_n \leq \widehat{N}(\lambda) \leq n^{\frac{\log 3}{\log 2}}$, based on consideration of (8) and (9).

Taken together, the preceding observations produce a qualitative model in our minds: *The number line just above p_n is dominated by the images under \widehat{N} of partitions of norm n , having no part equal to 1, and possessing a small number of parts.* (We note partitions with no 1's yield odd supernorm values; for each odd value, we must also count the even number that follows.)

Furthermore, the norm- n partitions with *many* parts are mapped by \widehat{N} closer to the upper bound $n^{\log 3 / \log 2}$ and do not map into the gap $[p_n, p_{n+1})$, but rather give rise to a lengthy “tail” of leftover integers that contribute to subsequent prime gaps more diffusely, perhaps somewhat randomly. The interval $[p_n, p_{n+1})$ also includes integers arising from the “tails” of subsets of partitions with norms less than n ; we anticipate their contribution is negligible asymptotically.

To translate this qualitative vision into a computational model one can test and make predictions from, we shall assume the following extreme simplification to address the observations above.⁷

Assumption 1. *Assume odd integers in the interval $[p_n, p_{n+1})$ are the images of partitions with norm equal to $n \geq 2$, with no part equal to 1, having one or two parts, under the supernorm. Then*

$$(10) \quad p_{n+1} - p_n = 2 \cdot \#\{\lambda \in \mathcal{P} : N(\lambda) = n, m_1(\lambda) = 0, \ell(\lambda) = 1 \text{ or } 2\}.$$

Remark. *We assume images of the aforementioned partitions dominate the number line above p_n .*

Integers having two prime factors are known as *semiprimes*. We expect this simplification yields an *underestimate* of prime gaps based on the assumption all odd numbers are prime or semiprime.

Now observe that the partitions of norm $n \geq 2$, with no part equal to 1 and having one or two parts, are precisely the set consisting of the partition (n) into one part, together with partitions (d_1, d_2) where $d_1 d_2 = n$, $d_1 \geq d_2$; that is, partitions whose parts are pairs of divisors of n . There are $\lceil d(n)/2 \rceil$ such divisor pairs where $d(k)$ is the *divisor function*, setting $d(0) := 0$,⁸ including $d_1 = n, d_2 = 1$, which we associate to partition (n) . Then Assumption 1 translates to the relation

$$(11) \quad p_{n+1} - p_n = 2 \left\lceil \frac{d(n)}{2} \right\rceil.$$

We refer the reader to Table 2 to see (11) does correctly predict most of the prime gaps between 1 and 100.⁹ Computations show this near-correctness diminishes as n increases, but as we will show, equation (11) is compatible with the prime number theorem asymptotically.

2.3. Initial model. Since $p_n = 1 + \sum_{k=0}^{n-1} (p_{k+1} - p_k)$ by telescoping series, defining $p_0 := 1$ as we did above, then equation (11) leads us to a model of prime numbers.

Model 1. *The prime numbers are the sequence p_1, p_2, p_3, \dots , of positive integers defined by $p_1 = 2$ and for $n \geq 2$, by the relation*

$$p_n = 1 + 2 \sum_{k=1}^{n-1} \left\lceil \frac{d(k)}{2} \right\rceil.$$

The first ten terms of this sequence are

$$2, 3, 5, 7, 11, 13, 17, 19, 23, 27,$$

and from here our initial model does not give the correct sequence of (odd) values. However, Model 1 does almost give the correct sequence of *prime gaps* at small numbers.

⁷In [8], the second, fourth and fifth authors give more detailed discussion of this simplifying assumption.

⁸We define $d(0) := 0$ for our purposes here, as there are no positive integers d_1, d_2 , such that $d_1 d_2 = 0$.

⁹These computations were carried out by the fifth listed author (Schneider), by hand.

One can estimate $\pi(n)$ by counting the primes up to n predicted by the model. Table 2 and Table 1 display that Model 1 does capture aspects of prime distribution qualitatively: at small numbers it emulates the unpredictable prime gaps reasonably well, and at large numbers, it yields numerical estimates for $\pi(n)$ that are comparable to the prime number theorem.

In the next subsection we discuss explicit predictions of Model 1, and how we interpret them.

2.4. Predictions from Model 1. Below we show that Model 1 implies the main asymptotic term of the prime number theorem, and produces unusual predictions about local fluctuations in prime gaps including the infinitude of twin primes as a special case.

The *prime number theorem* was conjectured by Gauss and Legendre in the 1790s based on newly published tables of primes, and was known as the “prime number conjecture” for a century, until Hadamard and de la Vallée Poussin proved it at the very end of the nineteenth century [30]. The prime number theorem is usually expressed as an asymptotic estimate for $\pi(n)$, viz. the estimate

$$(12) \quad \pi(n) \sim \frac{n}{\log n}$$

as n increases, or by the asymptotically equivalent (and evidently more accurate) estimate provided by the *logarithmic integral* $\text{li}(n)$:

$$(13) \quad \pi(n) \sim \text{li}(n) := \int_2^n \frac{dt}{\log t}.$$

The prime number theorem is equivalently formulated as an asymptotic estimate for the n th prime:

$$(14) \quad p_n \sim n \log n.$$

The estimate (14) follows from (12) by noting that $\pi(p_n) = n \sim p_n / \log p_n \sim p_n / \log n$. This final formulation of the prime number theorem is suggested naturally by Model 1. The first prediction of the model is a statement of fact.

Prediction 1 (Theorem). *Model 1 predicts the prime number theorem’s estimate for the n th prime,*

$$p_n \sim n \log n \text{ as } n \rightarrow \infty.$$

Proof. It follows from Assumption 1 that

$$(15) \quad p_n = 1 + \sum_{0 \leq i \leq n-1} (p_{i+1} - p_i) \sim \sum_{1 \leq i \leq n-1} d(i) = n \log n + n(2\gamma - 1) + O(\sqrt{n})$$

where $\gamma = 0.5772\dots$ is the Euler-Mascheroni constant; the final identity is due to Dirichlet [4]. \square

Remark. *We note there is a second-order error term of magnitude $n \log \log n$ implied by the prime number theorem, arising from $n = \pi(p_n) \sim p_n / \log(p_n)$, since $\log p_n \sim \log(n \log p_n) \sim \log n + \log \log n$, that is missing from Prediction 1. We address this missing error in Section 3.*

An interesting feature of Model 1 is that it suggests *predictable local behaviors of prime gaps*.

Prediction 2. *For $n \geq 2$, Model 1 predicts the n th prime gap is larger or smaller, depending on if n has a larger or smaller number of divisors, respectively.*

Proof. This is an immediate from (11), as $d(n)$ effectively controls the prime gap in this model. \square

Prediction 2 is highly counterintuitive – it says that *the indices of primes influence the prime gaps* – and cannot be strictly accurate since this model is an underestimate, to the degree it holds true at all, and doesn’t claim to measure the entire prime gap. Seeing that Model 1 is compatible with the prime number theorem, which itself is compatible with probabilistic hypotheses [30], the authors interpret Prediction 2 as representing a probabilistic statement, i.e., *the n th prime gap has an increased probability of being larger or smaller, depending on if $d(n)$ is large or small*. This unusual prediction can be checked computationally; we test one case in Section 3 below.

From the perspective of Model 1, the twin prime conjecture is a special case of Prediction 2.

Prediction 3. *Model 1 predicts twin prime pairs occur at prime-indexed primes, i.e., at primes p_n such that $n \in \mathbb{P}$. Moreover, Model 1 predicts the set of twin primes is infinite.*

Proof. This prediction is a special case of the previous prediction, the case $d(n) = 2$ if and only if $n \in \mathbb{P}$. Since prime-indexed primes are associated to twin prime pairs in Model 1, the infinitude of twin primes follows from the infinitude of the primes. \square

We note, once again, this prediction cannot be claimed to explicitly hold since we expect Model 1 is an underestimate of most prime gaps; this model does not have the precision to definitively predict a gap of minimal size 2, ruling out possible contributions from partitions with more than two parts mapping into the gap under \widehat{N} . As with Prediction 2, the authors interpret Prediction 3 as potentially a probabilistic statement, i.e., *prime-indexed primes have an increased probability of belonging to twin prime pairs, by comparison with arbitrary prime numbers.*

This probabilistic prediction can be checked computationally. In Section 3 below, we test Prediction 3, and find to our surprise that prime-indexed primes *do* have a slightly greater probability of belonging to twin prime pairs.

2.5. Critique of Model 1. As we noted previously, while Model 1 does emulate certain features of prime distribution, we expect *a priori* that it represents an underestimate of prime gaps in general, since the model does not account for odd numbers with more than two prime factors. A more refined model might account for these contributions. Moreover, as noted above, there is a missing error of order $n \log \log n$ by comparison with prime number theorem estimates for p_n .

While the model impressed itself, so to speak, on the authors from observations about the partition norm and supernorm, some of our assumptions are based on empirical observations that are not proved, although we noted they are compatible with the prime number theorem. Moreover, we have not proved (or consistently observed) that *even one* partition with norm n should necessarily map into the n th prime gap under the supernorm; it is an assumption justified by its predictive success. Model 1 also does not suggest that non-prime odd integers should respect anything like the correct integer ordering; it is not a very good model of *composite* numbers.

In Section 3 below, we check Prediction 3 and examine the error in Model 1, and make further simplifying assumptions that lead to a second model with better computational accuracy.

3. TESTING AND REFINING THE MODEL

3.1. Computational evidence for weak version of Prediction 4. In Section 2.4 we make a probabilistic conjecture about prime-indexed primes based on Prediction 3, viz. they are more likely to belong to twin prime pairs than are arbitrary prime numbers. While being an unusual prediction, this represents a weaker version of Prediction 3 that can be tested computationally.

We checked this prediction using the high-performance computing shared facility at Michigan Technological University.¹⁰ Let us define

$$(16) \quad \text{Prob}_1(n) := \frac{\#\{\text{prime-indexed primes } p \leq n \text{ such that } p + 2 \text{ is also prime}\}}{\#\{\text{prime-indexed primes } p \leq n\}},$$

the probability that a random prime-indexed prime $\leq n$ belongs to a twin prime pair, and define

$$(17) \quad \text{Prob}_2(n) := \frac{\#\{\text{primes } p \leq n \text{ such that } p + 2 \text{ is also prime}\}}{\pi(n)},$$

the probability that a random unrestricted prime $\leq n$ is a twin prime. Then to test the probabilistic interpretation of Prediction 3, we compute the ratio

$$(18) \quad P(n) := \frac{\text{Prob}_1(n)}{\text{Prob}_2(n)} \text{ as } n \text{ increases.}$$

¹⁰These computations were made by the third author (Hemmer) with Wolfram Mathematica and SageMath CAS.

n	p_n	$p_{n+1} - p_n$ (Actual)	$p_{n+1} - p_n$ (Model 1)
1	2	1	1
2	3	2	2
3	5	2	2
4	7	4	4
5	11	2	2
6	13	4	4
7	17	2	2
8	19	4	4
9	23	6	4
10	29	2	4
11	31	6	2
12	37	4	6
13	41	2	2
14	43	4	4
15	47	6	4
16	53	6	6
17	59	2	2
18	61	6	6
19	67	4	2
20	71	2	6
21	73	6	4
22	79	4	4
23	83	6	2
24	89	8	8
25	97	4	4



TABLE 2. Comparing actual prime gaps to predictions from Model 1; we highlight entries where the prediction for the n th prime gap is off.

By probabilistic hypotheses, we generally expect the ratio $P(n)$ to be around one: prime-indexed primes and arbitrary primes have the same probability of being twin primes in a random model. Then we expect $P(n) = 1$ broadly with local oscillations around one within some margin of error, that should diminish in amplitude as $n \rightarrow \infty$. In a random model, we expect roughly as many oscillations below one as above. *Our computations contradict these expectations.*

Result of computations. *The value of $y = P(n)$ is almost always greater than $y = 1$ with minor oscillations in the value locally, except for less frequent major oscillations dipping below $y = 1$ occurring up to around $n = 100,000$, with a significant bias above $y = 1$ on this interval, as displayed in Figure 1. Figure 2 displays that there are other instances of $P(n) < 1$ between five million and nine million. Further computations¹¹ reveal that from $n = 9,000,000$ up to at least 10 billion, $P(n)$ never dips down to $y = 1$ again, while appearing to asymptotically approach $y = 1$ from above, displaying a consistent if diminishing bias toward prime-indexed primes belonging to twin prime pairs.*

Then the seemingly dubious probabilistic prediction from Model 1 that prime-indexed primes are more likely to be twin primes, is supported by our computations. These computations lend some

¹¹The authors will share our results and code with the reader upon request, and intend to post them publicly.

FIGURE 1. Plot of $P(n)$ up to $n = 170,000$.FIGURE 2. Plot of $P(n)$ up to $n = 100,000,000$.

strength to our viewing the predictions of Model 1 as being suggestive of probabilistic phenomena. Of course, as of this writing it is not known whether the set of twin primes is finite or infinite.

3.2. Refining the error term. We now turn our attention from formulating a conceptual model of prime distribution, to improving our computational model and its numerical predictions for $\pi(n)$. As is standard, we define a semiprime to be an integer having exactly two factors. Let $\pi_k(n)$ denote the number of positive integers less than or equal to $n \geq 1$ having exactly $k \geq 0$ prime factors (so $\pi_0(n) = 1$, $\pi_1(n) = \pi(n)$ for all $n \geq 1$). All the integers up to n are enumerated by

$$(19) \quad n = 1 + \pi(n) + \pi_2(n) + \pi_3(n) + \cdots + \pi_k(n) + \dots$$

with only finitely many nonzero terms, depending on n . Substituting $n \mapsto p_n$ in (19) gives

$$(20) \quad p_n = 1 + n + \pi_2(p_n) + \pi_3(p_n) + \dots$$

This provides an explicit formula for the n th prime gap in terms of values $\pi_k(p), p \in \mathbb{P}$:

$$(21) \quad p_{n+1} - p_n = 1 + [\pi_2(p_{n+1}) - \pi_2(p_n)] + [\pi_3(p_{n+1}) - \pi_3(p_n)] + \dots$$

Now, (20) and (21) are not conceptually compatible with Model 1; the equations track factorizations of even as well as odd integers in the prime gap, while Assumption 1 only has us enumerating odd semiprimes and doubling the number. But these identities can inform corrections to Model 1.

It is an estimate due to Landau [16] that for $n > 1, k \geq 1$, as $n \rightarrow \infty$ we have

$$(22) \quad \pi_k(n) \sim \pi(n) \cdot \frac{(\log \log n)^{k-1}}{(k-1)!} \sim \frac{n (\log \log n)^{k-1}}{\log n (k-1)!}.$$

It follows for $k \geq 1$,¹² that

$$(23) \quad \pi_k(p_{n+1}) - \pi_k(p_n) \sim \frac{(\log \log n)^{k-1}}{(k-1)!}.$$

For fixed $n \geq 3$, this final ratio approaches zero as $k \rightarrow \infty$. Therefore, we expect integers with $k = 3$ prime factors are the next most populous set in the n th prime gap, apart from semiprimes. To improve our model, we should model the contributions of integers with three prime factors.

3.3. Second model. One apparent generalization of Assumption 1 is that we might enumerate partitions of norm n with no part equal to 1 and having *three* parts, as well as partitions with one or two parts, to improve our model by including odd integers with three prime factors under the map \widehat{N} . But as we increase the number of parts, our guiding assumption – the supernorms of norm- n partitions with few parts should dominate the number line just above p_n – becomes weaker. Moreover, length-3 partitions (and factorizations) are nontrivial to locate computationally. Model 1 proceeded from partition-theoretic considerations that were easy to formulate explicitly; we seek improvements that avoid having to generate partitions or prime factorizations computationally.

It would be ideal to use Landau’s estimate (22) to derive an asymptotic correction term for our model as n increases. As noted above, this is problematic as Landau’s estimate enumerates both odd and even integers, and Model 1 produces even numbers simply as enumerative “doubles” of the odd semiprimes. Furthermore, even integers with a fixed number of prime factors are more numerous than their odd counterparts on any interval, since they are divisible by powers of two, the smallest prime. Some – perhaps many – of the even numbers counted by the Landau estimate (23) of $\pi_3(p_{n+1}) - \pi_3(p_n)$ are already represented as doubles to odd semiprimes in Model 1.

We posit a second simplifying assumption based on the following elementary observation: *every integer in the n th prime gap having three prime factors, is the image of a semiprime less than p_n multiplied by a prime.* We will assume the converse also holds, as a simplistic approach to modeling the contribution of integers with three factors.

Assumption 2. *Assume every semiprime less than p_n maps into the interval $[p_n, p_{n+1})$ under multiplication by some prime less than p_n , and that the integers resulting from these products are not already enumerated by Model 1. Then by (22), this yields a contribution of*

$$(24) \quad \pi_2(p_n) \sim \pi(p_n) \log \log p_n \sim n \log \log n$$

integers having exactly three prime factors in the n th prime gap.

We count these integers with three factors in our updated model, noting the assumption produces an overestimate. *Assumption 2 also fills in the missing error of order $n \log \log n$ noted previously.*

We make one further simplifying assumption that is solely computational, and as such, represents an *ad hoc* revision – yet it leads to a significantly more accurate computational estimate for $\pi(n)$

¹²Noting $\log(p_n) \sim \log n \sim \log(n+1) \sim \log(p_{n+1})$.

that is *almost exact* at small numbers. One may want to replace the sum in (15) with an integral at large numbers. Note the asymptotic estimate on the right side of (15) is equivalently expressed

$$(25) \quad p_n \sim \int_0^n \log t \, dt + 2\gamma n.$$

This integral estimate is simpler, and unaffected asymptotically, if we delete the $2\gamma n$ term. By omitting $2\gamma n$, we avoid errors connected to approximating the (possibly irrational) constant γ .

Assumption 3. *Assume that subtracting $2\gamma n$ from the model will not hurt the reasonable accuracy of the model at small numbers, and will have negligible impact on the estimate for p_n asymptotically.*

Combining Assumptions 2 and 3 above with the statement of Model 1, we posit a refined model. We add a correction term $\lfloor \pi_2(p_{n-1}) - 2\gamma n \rfloor$ (using $\lfloor x \rfloor := 0$ if $x < 0$) to the formula for p_n . Assumption 2 provides number-theoretic justification for adding the $\pi_2(p_{n-1}) \sim n \log \log n$ term, as noted above; subtracting the $2\gamma n$ term is an *ad hoc* correction for computational simplicity. We use the floor function to ensure that our outputs are still integers, and because the floor function rounds down, which may compensate somewhat for Assumption 2 being an overestimate.

We only correct the $(n-1)$ th prime gap just below p_n , to estimate p_n , and do not correct the preceding gaps, to avoid compounding any new errors that may arise from our new assumptions. In practice, our computations show that either correction significantly improves the model; correcting only the $(n-1)$ th gap yields much more accurate computational results and faster run-times.¹³

Model 2. *The prime numbers are the sequence p_1, p_2, p_3, \dots , of positive integers defined by $p_1 = 2$ and for $n \geq 2$, by the relation*

$$p_n = 1 + 2 \sum_{k=1}^{n-1} \left\lfloor \frac{d(k)}{2} \right\rfloor + \lfloor \pi_2(p_{n-1}) - 2\gamma n \rfloor.$$

Remark. *Recall that we define $\lfloor x \rfloor := 0$ if $x < 0$ for the purposes of this paper.*

Unfortunately for our team, this conceptual model turns out to be too computationally intensive to test at large numbers on our personal computers. To check its numerical predictions as n increases, we utilize Landau's estimate (22). Noting the floor function error term in Model 2 is equal to zero below $n = 18$, and grows slowly from there, by (24) we make the substitution $\pi_2(p_{n-1}) \mapsto n \log \log n$ to produce a computational version of Model 2. Although this approximation introduces a source of potential error, it yields surprisingly good estimates for $\pi(n)$.

Model 2* (computational version). *The prime numbers are the sequence p_1, p_2, p_3, \dots , of positive integers defined by $p_1 = 2$ and for $n \geq 2$, by the relation*

$$p_n = 1 + 2 \sum_{k=1}^{n-1} \left\lfloor \frac{d(k)}{2} \right\rfloor + \lfloor n \log \log n - 2\gamma n \rfloor.$$

Remark. *For further computational ease, note $2 \lfloor d(k)/2 \rfloor = d(k)$ for all $k \neq m^2, m \in \mathbb{Z}^+$. On the other hand, if k is a perfect square then $2 \lfloor d(k)/2 \rfloor = d(k) + 1$. Thus for $N \geq 1$, we have that*

$$(26) \quad 2 \sum_{k=1}^N \left\lfloor \frac{d(k)}{2} \right\rfloor = \sum_{k=1}^N d(k) + \lfloor \sqrt{N} \rfloor = \sum_{k=1}^N \left\lfloor \frac{N}{k} \right\rfloor + \lfloor \sqrt{N} \rfloor,$$

by well-known principles related to the summatory function of $d(k)$ (see e.g. [4]).

One can directly count the number of primes Model 2* predicts up to any positive integer n . We refer the reader to Table 1 to confirm that Model 2* yields quite an accurate estimate for $\pi(n)$.

¹³All relevant programming was carried out by the first listed author (Botkin), using Wolfram Mathematica CAS.

3.4. Predictions from Model 2. With the additional correction term, Model 2 gives very good estimates of prime distribution at small numbers; and the second-order corrections do not impact the model's compatibility with the prime number theorem from an asymptotic perspective. Model 2 and Model 2* are both improvements on Model 1 from a computational perspective.

However, Model 2 makes one opposite prediction from Model 1.

Prediction 4. *Model 2 predicts the set of twin primes is finite.*

Proof. The correction term $\lfloor \pi_2(p_{n-1}) - 2\gamma n \rfloor \geq 1$ for all but finitely many small values of n . \square

The correction being an increasing function of n , while it yields accurate estimates for $\pi(n)$ at small numbers, prevents prime gaps in Model 2 from getting too small.¹⁴ In fact, Model 2 predicts there should be no twin prime pairs greater than about 100.

3.5. Critique of Model 2. We record a few concerns that might be addressed in a future study.

If the reader considers Assumption 2 to be well founded, then replacing $\pi_2(p_{n-1})$ with Landau's asymptotic introduces potential inaccuracy at values of n small enough that the asymptotic does not provide a good approximation; yet predictions from Model 2* are almost exact at small numbers. These second-order corrections might be addressed more carefully.

We note in Table 1 that at $n = 1,000,000$, the Model 2* prediction for $\pi(n)$ is less accurate than that of the logarithmic integral, whereas Model 2* is much closer than $\text{li}(n)$ to the value of $\pi(n)$ in all previous rows. The estimates for $\pi(n)$ from both Model 1 and Model 2* are overestimates because we expect both models to underestimate prime gaps; while we proved the model is in line with the main term of the prime number theorem, the missing contribution of partitions (and prime factorizations) of lengths greater than three creeps up. Further corrections, either refinements of our simplifying assumptions or new ideas entirely, might produce models of higher accuracy.

Finally, we make an aesthetic critique. Model 2 and Model 2* produce sequences of integers that fail to imitate the primes in a key way. The updated models predict *arbitrary primes are sometimes even*, as the correction term $\lfloor \pi_2(p_{n-1}) - 2\gamma n \rfloor$ can be odd or even. Noting two is the only even prime, we suggest as a further refinement of Model 2, replacing $\lfloor x \rfloor$ with a variant $\lfloor x \rfloor^*$ such that $\lfloor x \rfloor^* = \lfloor x \rfloor$ if $\lfloor x \rfloor$ is even, and $\lfloor x \rfloor^* = \lfloor x \rfloor + 1$ if $\lfloor x \rfloor$ is odd. This will ensure all prime gaps are even and all prime numbers are odd, and in rounding up from odd to even values, may compensate here and there for the model's underestimate of prime gaps.

4. CONCLUDING REMARKS

The authors posit Model 1 as a conceptual model that attempts to give an explanation for aspects of prime distribution based on elementary assumptions, which we derive from observations about integer partitions. The model is compatible with major observations such as the prime number theorem and twin prime conjecture, and predicts local fluctuations in prime gaps for which the prime number theorem lacks the precision to account. We treat these predictions about fluctuations as probabilistic statements, and check one such prediction, finding a small bias toward prime-indexed primes being twin primes. By contrast, Model 2 is a computational model that introduces simple corrections to Model 1, to yield estimates of prime distribution that are surprisingly good.

Because it is compatible with observations of primes and suggestive of new phenomena that can be tested, we hope the approach we present provides a useful complement to probabilistic models.

ACKNOWLEDGEMENTS

The authors are very grateful to Krishnaswami Alladi, George E. Andrews, William Craig, Jeffrey Lagarias, Paul Pollack, Ken Ono, James Sellers, Andrew V. Sills and Alexander Walker

¹⁴The minimum size of the n th prime gap should grow roughly like $\log \log n$, under Assumption 2.

for useful comments on earlier drafts of this study, and to Ivan Ramirez Zuniga for advice about scientific modeling.

REFERENCES

- [1] K. Alladi, and P. Erdős, *On an additive arithmetic function*, Pacific Journal of Mathematics **71.2** (1977): 275-294.
- [2] G. E. Andrews, *Partition identities*, Advances in Mathematics **9.1** (1972): 10-51.
- [3] G. E. Andrews *The Theory of Partitions*, Encyclopedia of Mathematics and its Applications, vol. 2, Addison-Wesley, 1976. Reissued, Cambridge University Press, 1998.
- [4] T. M. Apostol, *Introduction to Analytic Number Theory*, Springer Science & Business Media, 1998.
- [5] S. Corteel, and J. Lovejoy, Overpartitions, *Transactions of the American Mathematical Society* **356.4** (2004): 1623-1635.
- [6] W. Craig, *Higher level q -multiple zeta values with applications to quasimodular forms and partitions*, arXiv preprint [arXiv:2409.13874](https://arxiv.org/abs/2409.13874) (2024).
- [7] W. Craig, J. W. van Ittersum and Ken Ono, *Integer partitions detect the primes*, Proceedings of the National Academy of Sciences **121.39** (2024): e2409417121.
- [8] M. L. Dawsey, M. Just, and R. Schneider, *A “supernormal” partition statistic (v1)*, arXiv preprint [arxiv:2107.14284v1](https://arxiv.org/abs/2107.14284v1) (2021).
- [9] M. L. Dawsey, M. Just, and R. Schneider, *A “supernormal” partition statistic*, The Journal of Number Theory **241** (2022): 120-141.
- [10] K. Gomez, *MacMahonesque partition functions detect sets related to primes*, arXiv preprint [arXiv:2409.14253](https://arxiv.org/abs/2409.14253) (2024).
- [11] A. Granville, *The anatomy of integers and permutations*, preprint (2008).
- [12] A. Granville, and K. Soundararajan, *it Pretentious multiplicative functions and an inequality for the zeta-function*, CRM Proceedings and Lecture Notes, Vol. 46, 2008.
- [13] M. Just, *On a map between partitions and integers* (joint work with R. Schneider), Modular Forms, Arithmetic and Women in Mathematics Conference (MAAIM), Emory University, Atlanta, Georgia, U.S.A., 19 November, 2019.
- [14] J. C. Lagarias, *Characterizing the supernorm partition statistic*, The Ramanujan Journal (2023): 1-13.
- [15] J. C. Lagarias and C. Sun, *Asymptotics of reciprocal supernorm partition statistics*, arXiv preprint [arXiv:2308.15303](https://arxiv.org/abs/2308.15303) (2023).
- [16] E. Landau, *Handbuch der Lehre von der Verteilung der Primzahlen Vol. 1*, B. G. Teubner, 1909.
- [17] T. Tao, *Probabilistic models and heuristics for the primes*, <https://terrytao.wordpress.com/2015/01/04/254a-supplement-4-probabilistic-models-and-heuristics-for-the-primes-optional/> (4 Jan., 2015), Web.
- [18] K. Ono, *The Web of Modularity: Arithmetic of the Coefficients of Modular Forms and q -series: Arithmetic of the Coefficients of Modular Forms and q -series*, No. 102, American Mathematical Soc., 2004.
- [19] K. Ono, L. Rolén, and R. Schneider, *Explorations in the theory of partition zeta functions*, in *Exploring the Riemann Zeta Function, 190 years from Riemann’s Birth*, ed. H. Montgomery, A. Nikeghbali, M. Rassias, (Springer, 2017), pp. 223–264.
- [20] K. Ono, R. Schneider, and I. Wagner, *Partition-theoretic formulas for arithmetic densities*, Analytic Number Theory, modular forms and q -series, Springer Proc. Math. Stat. **221** (2017): 611–624.
- [21] K. Ono, R. Schneider, and I. Wagner, *Partition-theoretic formulas for arithmetic densities, II*, Hardy–Ramanujan J. **43** (2020): 1–16.
- [22] R. Schneider, *Partition zeta functions*, Research in Number Theory **2** (2016), article 9, 17 pp.
- [23] R. Schneider, *Arithmetic of partitions and the q -bracket operator*, Proc. Amer. Math. Soc. **145.5** (2017): 1953–1968.
- [24] R. Schneider, *Jacobi’s triple product, mock theta functions, unimodal sequences and the q -bracket*, Int. J. Number Theory **14.07** (2018): 1961–1981.
- [25] R. Schneider, *Eulerian series, zeta functions and the arithmetic of partitions* (Advisor: Ken Ono), PhD dissertation, Emory University (2018).
- [26] R. Schneider and A. V. Sills. *Analysis and combinatorics of partition zeta functions*. Int. J. Number Theory (2020): 1-10.
- [27] R. Schneider and A. V. Sills, *The product of parts or “norm” of a partition*, Integers **20A** (2020), paper #A13, 16 pp.
- [28] R. Schneider, *Partition-theoretic Frobenius-type limit formulas*, The Ramanujan Journal (2024): 1-8.
- [29] R. Schneider, *Algebra of overpartitions*, arXiv preprint [arXiv:2410.00941](https://arxiv.org/abs/2410.00941) (2024).

- [30] G. Tenenbaum, M. M. France, and P. G. Spain, *The Prime Numbers and Their Distribution*, Vol. 6, American Mathematical Society, 2000.

DEPARTMENT OF MATHEMATICAL SCIENCES
MICHIGAN TECHNOLOGICAL UNIVERSITY
HOUGHTON, MICHIGAN 49931, U.S.A.
Email address: `abotkin@mtu.edu`

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF TEXAS AT TYLER
TYLER, TEXAS 75799, U.S.A.
Email address: `mdawsey@uttyler.edu`

DEPARTMENT OF MATHEMATICAL SCIENCES
MICHIGAN TECHNOLOGICAL UNIVERSITY
HOUGHTON, MICHIGAN 49931, U.S.A.
Email address: `djemmer@mtu.edu`

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF GEORGIA
ATHENS, GEORGIA, 30602, U.S.A.
Email address: `justmatt@uga.edu`

DEPARTMENT OF MATHEMATICAL SCIENCES
MICHIGAN TECHNOLOGICAL UNIVERSITY
HOUGHTON, MICHIGAN 49931, U.S.A.
Email address: `robertsc@mtu.edu`