Sampling the Bayesian Elastic Net

Christopher M. Hans* and Ningyi Liu

Department of Statistics, The Ohio State University, Columbus, OH, 43210, USA December 2024

Abstract

The Bayesian elastic net and its variants have become popular approaches to regression in many areas of research. The model is characterized by the prior distribution on the regression coefficients, the negative log density of which corresponds to the elastic net penalty function. While straightforward Markov chain Monte Carlo (MCMC) methods exist for sampling from the posterior distribution of the regression coefficients given the penalty parameters, full Bayesian inference—where the MCMC algorithms are expanded to integrate over uncertainty in the penalty parameters—remains a challenge. Sampling the penalty parameters (and the regression model error's variance parameter under some forms of the prior) is complicated by the presence of an intractable integral expression in the normalizing constant for the prior on the regression coefficients. Though sampling methods have been proposed that avoid the need to compute the normalizing constant, all correctly-specified methods for updating the remaining parameters that have been described in the literature involve at least one "Metropolis-within-Gibbs" update, requiring specification and tuning of proposal distributions. The computational landscape is complicated by the fact that two different forms of the Bayesian elastic net prior have been introduced in the literature, and two different representations (with and without data augmentation) of the prior suggest different MCMC algorithms for sampling the regression coefficients. We first provide a comprehensive review of the forms and representations of the Bayesian elastic net prior, discussing all combinations of these different treatments of the prior together for the first time and introducing one combination of form and representation that has yet to appear in the literature. We then introduce MCMC algorithms for full Bayesian inference for all combinations of prior form and

^{*}Corresponding author: hans@stat.osu.edu

representation. The algorithms allow for direct sampling of all parameters at low computational cost without any "Metropolis-within-Gibbs" steps, avoiding potential problems with slow convergence and mixing due to poor choice of proposal distribution. The key to the new approach is a careful transformation of the parameter space and an analysis of the resulting full conditional density functions that allows for efficient rejection sampling of the transformed parameters. We make empirical comparisons between our sampling approaches and other existing MCMC methods in the literature for a variety of potential data structures.

Key Words: elastic net; lasso; MCMC; orthant normal distribution; prior distribution; regression; regularization; rejection sampling; shrinkage

1 Introduction

The Bayesian elastic net and its variants have become popular approaches to regression in many areas of research. Li and Lin (2010) and Hans (2011) introduced the Bayesian elastic net model in the normal linear regression setting, $y = \mathbf{1}\alpha + X\beta + \varepsilon$, where y is an $n \times 1$ response vector, X is an $n \times p$ matrix of regressors, and $\varepsilon \sim N(0, \sigma^2 I_n)$. In this framework, the Bayesian elastic net is characterized by the prior on the regression coefficients,

$$\pi_c(\beta \mid \sigma^2, \lambda_1, \lambda_2) \propto \exp\left\{-\frac{1}{2\sigma^2} \left(\lambda_2 \beta^T \beta + \lambda_1 |\beta|_1\right)\right\},\tag{1}$$

where $|\beta|_1 = \sum_{j=1}^p |\beta_j|$ is the ℓ_1 -norm of β . Under the non-informative prior on the intercept parameter, $\pi(\alpha) \propto 1$, the integrated likelihood function is

$$p(y \mid X, \beta, \sigma^2) = \int p(y \mid X, \alpha, \beta, \sigma^2) p(\alpha) d\alpha = (2\pi\sigma^2)^{-(n-1)/2} n^{-1/2} e^{-\frac{1}{2\sigma^2} (y^* - X^*\beta)^T (y^* - X^*\beta)},$$
(2)

where y^* and X^* are the mean-centered response vector and column-mean-centered matrix of regressors, respectively. The posterior distribution $\pi_c(\beta \mid y, \sigma^2)$ then satisfies

$$-2\sigma^2 \log \pi_c(\beta \mid y, \sigma^2) = \text{const.} + (y^* - X^*\beta)^T (y^* - X^*\beta) + \lambda_2 \beta^T \beta + \lambda_1 |\beta|_1.$$
(3)

The non-constant component of (3) is the elastic net objective function (Zou and Hastie, 2005) with penalty parameters λ_1 and λ_2 so that, for any fixed value of σ^2 , the posterior mode of β corresponds to an elastic net estimate. We use the integrated likelihood (2) throughout and assume that y and the columns of X have been mean-centered, dropping the "*" in the notation.

The literature on the connection between Bayesian posterior modes and estimators described as solutions to penalized optimization problems is quite rich. Tibshirani (1996) made the first such connection for lasso regression, the Bayesian side of which was more fully developed by Park and Casella (2008) and Hans (2009, 2010). Bayesian connections to the adaptive lasso (Zou, 2006) have been considered by Griffin and Brown (2007, 2011), Alhamzawi *et al.* (2012), Leng *et al.* (2014), Alhamzawi and Ali (2018), Kang *et al.* (2019), and Wang *et al.* (2019), among others.

Ročková and George (2018) introduced a fully-Bayes, adaptive approach to Bayesian variable selection that combines the lasso penalty function with the ideas that underly "spike-and-slab" priors. Wang (2012) introduced a Bayesian formulation of the graphical lasso (Meinshausen and Bühlmann, 2006; Yuan and Lin, 2007; Friedman *et al.*, 2008), while Kyung *et al.* (2010) and others have studied connections to the group and fused lassos (Yuan and Lin, 2006; Tibshirani *et al.*, 2005).

Bayesian regression models with connections to the elastic net have also received extensive attention in the literature. After Zou and Hastie (2005) noted that their elastic net estimator could be viewed as the mode of a Bayesian posterior distribution, Kyung *et al.* (2010), Li and Lin (2010) and Hans (2011) sought to fully characterize the corresponding Bayesian model. Li and Lin (2010) represented the prior as a scale-mixture of normal distributions, discussed inference when σ^2 was unknown, and introduced Bayesian approaches for selecting the penalty parameters. Hans (2011) also described the scale-mixture of normals representation, introduced an additional, "direct" representation of the prior, considered full Bayesian inference on the penalty parameters, and introduced methods for Bayesian elastic net variable selection and model averaging for prediction. More recently, Lee *et al.* (2015) identified an error in Kyung *et al.* (2010)'s representation of the prior. Roy and Chakraborty (2017) corrected the error and also studied optimal selection of the penalty parameters. Wang and Wang (2023) introduced an MCMC algorithm for full Bayesian elastic net inference that was designed to avoid the need to approximate any integrals in any of the sampling steps.

Two main forms of the Bayesian elastic net prior distribution are common in the literature. Li and Lin (2010) and Hans (2011) considered the prior as parameterized in (1), where the two components of the penalty function, $\lambda_2\beta^T\beta$ and $\lambda_1|\beta_1|$, are both scaled by $2\sigma^2$. We refer to this form of the prior as the "commonly-scaled" parameterization, and we subscript prior and posterior densities under the common scaling with "c" for clarity, e.g., $\pi_c(\beta \mid \sigma^2, \lambda_1, \lambda_2)$ in (1). Kyung *et al.* (2010) and Roy and Chakraborty (2017) scale the penalty terms differentially:

$$\pi_d(\beta \mid \sigma^2, \lambda_1 \lambda_2) \propto \exp\left\{-\frac{\lambda_2}{2\sigma^2}\beta^T \beta - \frac{\lambda_1}{\sigma}|\beta|_1\right\}.$$
(4)

This version of the prior has the useful property that λ_1 and λ_2 do not depend on the units of the response variable: coupled with a scale-invariant prior on σ , an analyst would not need to adjust the fixed values of (or priors for) λ_1 and λ_2 in order to obtain the same posterior if the response variable was rescaled linearly. We refer to this form of the prior as the "differentially-scaled" parameterization and subscript corresponding prior and posterior densities with "d" to emphasize the differential scaling. We omit the subscript when the distinction between the two forms of the prior is not relevant. For fixed values of σ^2 , λ_1 , and λ_2 , the differentially-scaled prior is simply a reparameterization of the commonly-scaled prior: given the same values of σ^2 and λ_2 , one can obtain the same posterior distribution under both priors through choice of the λ_1 parameter specific to each prior. It is useful, though, to consider both versions of the prior separately because the interpretation of λ_1 is specific to the form of the scaling.

While the papers that study the Bayesian elastic net take different perspectives and have varying objectives, they have in common the theme that full Bayesian inference under the elastic net prior presents computational challenges. The $|\beta|_1$ term in the prior makes direct integration of the posterior challenging, and so one of the keys to Bayesian elastic net regression modeling is the ability to easily sample from the posterior. Several MCMC algorithms have been proposed to obtain samples from the conditional posterior of β given σ^2 , λ_1 and λ_2 . One approach—similar to the one used by Park and Casella (2008) for the Bayesian lasso—is to demarginalize the prior on β by introducing latent variables, τ^2 , that can be exploited to conduct a data augmentation Gibbs sampler. Li and Lin (2010) and Hans (2011) extend this idea to the elastic net penalty function and introduce corresponding data augmentation Gibbs samplers under the common scale prior, and Roy and Chakraborty (2017) and Wang and Wang (2023) consider the data augmentation approach under the differentially-scaled prior. As an alternative to data augmentation Gibbs sampling, Hans (2011) describes an alternative Gibbs sampler for the Bayesian elastic net that updates each β_j one at a time, conditionally on the others, without requiring the inclusion of latent variables in the sampling scheme.

The more difficult challenges to computation become apparent when we assign prior distributions to σ^2 , λ_1 , and λ_2 and wish to make inference based on the joint posterior distribution. As seen in Section 2, the normalizing constant for the Bayesian elastic net prior (1) contains the term $\Phi(-\lambda_1/(2\sigma\sqrt{\lambda_2}))^{-p}$, where $\Phi(\cdot)$ is the standard normal cumulative distribution function (cdf). The same term appears in the joint prior density for β and τ^2 under the data augmentation representation of the prior. Sampling from or integrating the joint posterior distribution therefore requires dealing with $\Phi(-\lambda_1/(2\sigma\sqrt{\lambda_1})^{-p})$, an integral expression with no closed form solution. Noting that the standard normal cdf can evaluated numerically to relatively high precision when its argument is not too close to $\pm\infty$, Hans (2011) sampled σ^2 , λ_1 , and λ_2 via random-walk "Metropolis-within-Gibbs" updates for $\log \sigma^2$, $\log \lambda_1$, and $\log \lambda_2$, with their respective full conditional distributions as the target distributions. While effective, this approach requires specifying step-size parameters for the random walks, poor choices of which can lead to slow convergence and mixing and the need to iteratively tune and re-run the MCMC algorithm. Li and Lin (2010) avoided the issue of sampling the penalty parameters by devising data-adaptive methods for selecting values for them, and attempted to sample σ^2 directly from its full conditional distribution via rejection sampling. Unfortunately, as shown in Appendix A, their rejection sampling algorithm contains an error and does not produce samples from the desired target distribution.

The computational situation is slightly improved when the differentially-scaled form of the prior (4) is used. As shown in Section 2, the awkward term in the normalizing constant is then $\Phi(-\lambda_1/\sqrt{\lambda_2})^{-p}$, which no longer depends on σ^2 . Posterior sampling of σ^2 under this form of the prior is straightforward, as demonstrated by Roy and Chakraborty (2017). Full Bayesian inference under priors on λ_1 and λ_2 , however, still must involve methods for handling the analytically intractable integral expression. Motivated by the desire to avoid numerical computation of $\Phi(\cdot)$, Wang and Wang (2023) devised a clever exchange algorithm (Murray *et al.*, 2006) that introduces p additional latent variables in such a way as to remove the term involving $\Phi(\cdot)$ from the joint posterior of the augmented parameter space. Despite avoiding computation of $\Phi(\cdot)$, the algorithm still requires one parameter be updated via the Metropolis–Hastings algorithm using a random-walk proposal, necessitating the selection of a step-size parameter for the random walk.

All of the correctly-specified MCMC approaches described above for fully-Bayes inference under the Bayesian elastic net use at least one Metropolis step. From the point of view of a practitioner, it would be better if the associated selection of random-walk step-size parameters could be avoided entirely. The computational landscape is further complicated by the fact that the two different forms (common and differential scaling) and two different representations (with and without data augmentation) of the prior suggest different approaches to MCMC for full Bayesian inference for Bayesian elastic net regression. In Section 2, we provide a comprehensive review of the forms and representations of the Bayesian elastic net prior and the existing approaches to computation. We discuss together for the first time all combinations of the different treatments of form and representation of the prior, and we introduce one combination of form and treatment that has yet to be discussed in the literature. We use this review to highlight the computational difficulties associated with full Bayesian inference. To solve the computational problems, we introduce in Section 3 new MCMC algorithms for full Bayesian inference under all combinations of prior form and representation. The algorithms allow for direct sampling of all parameters at low computational cost without using any "Metropolis-within-Gibbs" steps, avoiding potential problems with slow convergence and mixing due to poor choice of proposal distribution. The key to the new approach is a careful transformation of the parameter space and an analysis of the resulting full conditional density functions that allows for efficient rejection sampling of the transformed parameters. We make empirical comparisons in Section 4 between our new sampling approaches and other existing MCMC methods in the literature for a variety of potential data structures.

2 Existing Approaches to Model Specification and Posterior Computation

The Bayesian elastic net prior distribution can be represented directly (without data augmentation) or hierarchically (with data augmentation). We refer to these two representations of the prior as the "direct" and "DA" representations, respectively. Both forms of the prior (common and differential scaling) have a direct and DA representation. Table 1 indicates where these combinations of form and representation originally appeared in the literature. In this section we review the four combinations of representation and form, provide the corresponding posterior distributions, and discuss existing approaches to posterior computation, highlighting computational challenges.

	Direct	Data Augmentation (DA)
Common Scaling	Hans (2011)	Li and Lin (2010), Hans (2011)
Differential Scaling	*	Kyung <i>et al.</i> (2010) ^{**} , Roy and Chakraborty (2017)

Table 1: Citations for original descriptions of the four combinations of form (rows) and representation (columns) of the Bayesian elastic net prior. The direct representation of the differentially-scaled prior (*) is introduced in Section 2.1. Kyung *et al.* (2010)'s description of the DA representation of the differentially-scaled prior (**) contained an error which was corrected by Roy and Chakraborty (2017).

2.1 Direct Representation of the Prior

Hans (2011) introduced the direct characterization of the commonly-scaled elastic net prior (1). Handling the term $|\beta|_1 = \sum_{j=1}^p |\beta_j|$ by treating separately the cases $\beta_j < 0$ and $\beta_j \ge 0$, the independent priors on each β_j can be expressed as two separate, symmetric, truncated normal distributions that are weighted to have matching density at the origin. Combining the cases together,

$$\pi_{c}(\beta \mid \sigma^{2}, \lambda_{1}, \lambda_{2}) = \prod_{j=1}^{p} \left\{ \frac{1}{2} \cdot \mathbf{N}^{-} \left(\beta_{j} \mid \frac{\lambda_{1}}{2\lambda_{2}}, \frac{\sigma^{2}}{\lambda_{2}} \right) + \frac{1}{2} \cdot \mathbf{N}^{+} \left(\beta_{j} \mid -\frac{\lambda_{1}}{2\lambda_{2}}, \frac{\sigma^{2}}{\lambda_{2}} \right) \right\}$$
$$\equiv \prod_{j=1}^{p} \left\{ \frac{1}{2} \cdot \frac{\mathbf{N} \left(\beta_{j} \mid \frac{\lambda_{1}}{2\lambda_{2}}, \frac{\sigma^{2}}{\lambda_{2}} \right)}{\Phi \left(\frac{-\lambda_{1}}{2\sigma\sqrt{\lambda_{2}}} \right)} \mathbf{1}(\beta_{j} < 0) + \frac{1}{2} \cdot \frac{\mathbf{N} \left(\beta_{j} \mid -\frac{\lambda_{1}}{2\lambda_{2}}, \frac{\sigma^{2}}{\lambda_{2}} \right)}{\Phi \left(\frac{-\lambda_{1}}{2\sigma\sqrt{\lambda_{2}}} \right)} \mathbf{1}(\beta_{j} \ge 0) \right\}.$$
(5)

The notation $N^{-}(x \mid m, s^2)$ and $N^{+}(x \mid m, s^2)$ denotes the normalized density functions for, respectively, negatively and non-negatively truncated univariate normal distributions, where m and s^2 are, respectively, the mean and variance of an underlying, non-truncated, normal random variable. The corresponding density functions are:

$$\mathbf{N}^{-}(x \mid m, s^{2}) \equiv \frac{\mathbf{N}(x \mid m, s^{2})}{\Phi(-m/s)} \mathbf{1}(x < 0), \quad \mathbf{N}^{-}(x \mid m, s^{2}) \equiv \frac{\mathbf{N}(x \mid m, s^{2})}{1 - \Phi(-m/s)} \mathbf{1}(x \ge 0),$$

where $N(x \mid m, s^2) \equiv (2\pi s^2)^{-1/2} \exp\{-(x-m)^2/(2s^2)\}$ is the probability density function for a normal distribution with mean m and variance s^2 , and $\Phi(x) \equiv \int_{-\infty}^x (2\pi)^{-1/2} e^{-u^2/2} du$ is the standard normal cdf.

Isolating some of the constant terms in (5) and combining the univariate densities into a multi-

variate density, the prior can also be written as

$$\pi_c(\beta \mid \sigma^2, \lambda_1, \lambda_2) = 2^{-p} \Phi\left(\frac{-\lambda_1}{2\sigma\sqrt{\lambda_2}}\right)^{-p} \sum_{z \in \mathcal{Z}} N\left(\beta \mid -\frac{\lambda_1}{2\lambda_2}z, \frac{\sigma^2}{\lambda_2}I_p\right) \mathbf{1}(\beta \in \mathcal{O}_z), \quad (6)$$

where $N(x \mid m, S)$ is the density function for a *p*-dimensional multivariate normal distribution with mean vector *m* and covariance matrix *S*. The sum is taken over all 2^p possible *p*-vectors *z* having elements $z_j \in \{-1, 1\}$, with \mathcal{Z} being the set of all such vectors. The notation \mathcal{O}_z refers to the orthant of \mathbb{R}^p where each coordinate is restricted by *z* to be negative ($z_j = -1$) or non-negative ($z_j = 1$). The elastic net prior distribution can therefore be thought of as a collection of truncated multivariate normal distributions defined separately on the 2^p orthants of \mathbb{R}^p . The location vector for the normal distribution in each orthant depends on the orthant, but the orientations of the normal distributions are the same for all orthants. The specific values of the location and orientation parameters ensure that the prior density is continuous, but not differentiable, along the coordinate axes. Hans (2011) calls this an "orthant normal distribution" and writes the density as

$$\pi_{c}(\beta \mid \sigma^{2}, \lambda_{1}, \lambda_{2}) = \sum_{z \in \mathcal{Z}} 2^{-p} \mathbf{N}^{[z]} \left(\beta \mid -\frac{\lambda_{1}}{2\lambda_{2}} z, \frac{\sigma^{2}}{\lambda_{2}} I_{p} \right)$$
$$\equiv \sum_{z \in \mathcal{Z}} 2^{-p} \frac{\mathbf{N} \left(\beta \mid -\frac{\lambda_{1}}{2\lambda_{2}} z, \frac{\sigma^{2}}{\lambda_{2}} I_{p} \right)}{\mathbf{P} \left(z, \frac{-\lambda_{1}}{2\lambda_{2}} z, \frac{\sigma^{2}}{\lambda_{2}} I_{p} \right)} \mathbf{1}(\beta \in \mathcal{O}_{z}),$$

where $N^{[z]}(\cdot | \cdot, \cdot)$ denotes the density function for a multivariate normal distribution truncated to orthant \mathcal{O}_z , and $P(z, \cdot, \cdot)$ is the probability assigned to that orthant by the underlying normal distribution. For the commonly-scaled elastic net prior, the orthant probabilities are all equal and generate the $\Phi(-\lambda_1/(2\sigma\sqrt{\lambda_2}))^{-p}$ term in (6):

$$\begin{split} \mathbf{P}\left(z, \frac{-\lambda_1}{2\lambda_2} z, \frac{\sigma^2}{\lambda_2} I_p\right) &= \int_{\mathcal{O}_z} \mathbf{N}\left(\beta \mid \frac{-\lambda_1}{2\lambda_2} z, \frac{\sigma^2}{\lambda_2} I_p\right) d\beta \\ &= \left[\prod_{j : z_j = -1} \int_{-\infty}^0 \mathbf{N}\left(\beta_j \mid \frac{\lambda_1}{2\lambda_2}, \frac{\sigma^2}{\lambda_2}\right) d\beta_j\right] \times \left[\prod_{j : z_j = 1} \int_0^\infty \mathbf{N}\left(\beta_j \mid \frac{-\lambda_1}{2\lambda_2}, \frac{\sigma^2}{\lambda_2}\right) d\beta_j\right] \\ &= \left[\prod_{j : z_j = -1} \Phi\left(\frac{-\lambda_1}{2\sigma\sqrt{\lambda_2}}\right)\right] \times \left[\prod_{j : z_j = 1} \left(1 - \Phi\left(\frac{\lambda_1}{2\sigma\sqrt{\lambda_2}}\right)\right)\right] \\ &= \Phi\left(\frac{-\lambda_1}{2\sigma\sqrt{\lambda_2}}\right)^p. \end{split}$$

Examples of this density function when p = 1 and p = 2 are shown in Figure 1. We refer to this representation of the prior as the "direct" representation under the common scaling. It is sometimes convenient to work with the prior density by properly normalizing the expression in (1), retaining the $|\beta|_1$ term and avoiding the summation over the orthants:

$$\pi_c(\beta \mid \sigma^2, \lambda_1, \lambda_2) = 2^{-p} (2\pi)^{-p/2} (\sigma^2/\lambda_2)^{-p/2} e^{-\frac{p\lambda_1^2}{8\sigma^2\lambda_2}} \Phi\left(-\frac{\lambda_1}{2\sigma\sqrt{\lambda_2}}\right)^{-p} \exp\left\{-\frac{\lambda_2}{2\sigma^2} \beta^T \beta - \frac{\lambda_1}{2\sigma^2} |\beta|_1\right\}.$$
(7)

The direct representation of the differentially-scaled prior (4) has not been explicitly described in the literature, but it is easy to show that

$$\pi_{d}(\beta \mid \sigma^{2}, \lambda_{1}\lambda_{2}) = \prod_{j=1}^{p} \left\{ \frac{1}{2} \cdot \mathbf{N}^{-} \left(\beta_{j} \mid \frac{\sigma\lambda_{1}}{\lambda_{2}}, \frac{\sigma^{2}}{\lambda_{2}} \right) + \frac{1}{2} \cdot \mathbf{N}^{+} \left(\beta_{j} \mid -\frac{\sigma\lambda_{1}}{\lambda_{2}}, \frac{\sigma^{2}}{\lambda_{2}} \right) \right\}$$
$$= 2^{-p} \Phi \left(-\frac{\lambda_{1}}{\sqrt{\lambda_{2}}} \right)^{-p} \sum_{z \in \mathcal{Z}} \mathbf{N} \left(\beta \mid -\frac{\sigma\lambda_{1}}{\lambda_{2}} z, \frac{\sigma^{2}}{\lambda_{2}} I_{p} \right) \mathbf{1}(\beta \in \mathcal{O}_{z}).$$
(8)

The term in the normalizing constant involving $\Phi(\cdot)$ does not depend on σ^2 under this scaling of the prior. As in (7), we can express the properly normalized, differentially-scaled prior in terms of $|\beta|_1$ as:

$$\pi_d(\beta \mid \sigma^2, \lambda_1, \lambda_2) = 2^{-p} (2\pi)^{-p/2} (\sigma^2/\lambda_2)^{-p/2} e^{-\frac{p\lambda_1^2}{2\lambda_2}} \Phi\left(-\frac{\lambda_1}{\sqrt{\lambda_2}}\right)^{-p} \exp\left\{-\frac{\lambda_2}{2\sigma^2} \beta^T \beta - \frac{\lambda_1}{\sigma} |\beta|_1\right\}$$
(9)



Figure 1: Elastic net prior density $\pi_c(\beta \mid \sigma^2 = 1, \lambda_1 = 1, \lambda_2 = 1)$. Left panel: prior density for β_1 when p = 1; right panel: contours of joint prior density for β_1 and β_2 when p = 2.

2.2 Data Augmentation Representation of the Prior

Li and Lin (2010) and Hans (2011) provide an alternate characterization of the commonly-scaled elastic net prior distribution by representing it via demarginalization as a scale-mixture of normal distributions (Andrews and Mallows, 1974; West, 1987). Introducing latent scale variables, τ_j^2 , Hans (2011) defines the hierarchical model

$$\begin{split} \beta_{j} \mid \sigma^{2}, \lambda_{1}, \lambda_{2}, \tau_{j}^{2} & \stackrel{\mathrm{ind}}{\sim} & \mathrm{N}\left(0, \frac{\sigma^{2}}{\lambda_{2}}(1 - \tau_{j}^{2})\right), \\ \tau_{j}^{2} \mid \sigma^{2}, \lambda_{1}, \lambda_{2} & \stackrel{\mathrm{iid}}{\sim} & \mathrm{Inv-Gamma}_{(0,1)}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{\lambda_{1}}{2\sigma\sqrt{\lambda_{2}}}\right)^{2}\right), \end{split}$$

for j = 1, ..., p, where Inv-Gamma_(0,1)(\cdot, \cdot) is an inverse gamma distribution truncated to the interval (0, 1) with density function

$$\pi_{c}(\tau_{j}^{2} \mid \sigma^{2}, \lambda_{1}, \lambda_{2}) = \frac{1}{2\sqrt{2\pi}} \Phi\left(\frac{-\lambda_{1}}{2\sigma\sqrt{\lambda_{2}}}\right)^{-1} \left(\frac{\lambda_{1}}{2\sigma\sqrt{\lambda_{2}}}\right) (\tau_{j}^{2})^{-3/2} e^{-\frac{\lambda_{1}^{2}}{8\sigma^{2}\lambda_{2}}\tau_{j}^{-2}}, \quad 0 < \tau_{j}^{2} < 1.$$
(10)

As shown in Hans (2011), the marginal density of β_j under this model is (7), the commonly-scaled elastic net prior. Li and Lin (2010) provide a similar result, but parameterize the scale-mixture

slightly differently:

$$\begin{split} \beta_{j} \mid \sigma^{2}, \lambda_{1}, \lambda_{2}, \tau_{j}^{2} & \stackrel{\mathrm{ind}}{\sim} & \mathrm{N}\left(0, \frac{\sigma^{2}}{\lambda_{2}} \frac{\tau_{j}^{2} - 1}{\tau_{j}^{2}}\right), \\ \tau_{j}^{2} \mid \sigma^{2}, \lambda_{1}, \lambda_{2} & \stackrel{\mathrm{iid}}{\sim} & \mathrm{Gamma}_{(1,\infty)}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{\lambda_{1}}{2\sigma\sqrt{\lambda_{2}}}\right)^{2}\right), \end{split}$$

where the gamma distribution with rate parameter $\lambda_1^2/(8\sigma^2\lambda_2)$ is truncated to the interval $(1,\infty)$. Li and Lin (2010) express the normalizing constant in the density for the truncated gamma distribution in terms of the upper incomplete gamma function (DiDonato and Morris, 1986), $\Gamma_U(\alpha, x) = \int_x^\infty t^{\alpha-1}e^{-t}dt$. The relevant term in the normalizing constant of this density is $\Gamma_U\left(\frac{1}{2}, \frac{1}{2}\left(\frac{\lambda_1}{2\sigma\sqrt{\lambda_2}}\right)^2\right)$, which is equivalent to $2\sqrt{\pi}\Phi\left(\frac{-\lambda_1}{2\sigma\sqrt{\lambda_2}}\right)$ due to the identity $2\sqrt{\pi}\Phi(-x) = \Gamma_U\left(\frac{1}{2}, \frac{x^2}{2}\right)$ for $x \ge 0$. We refer to any version of this representation of the prior as the data augmentation ("DA") or scale-mixture-of-normals ("SMN") representation.

Roy and Chakraborty (2017) introduced the correct DA representation of the differentiallyscaled prior (4) via the hierarchical representation

$$\beta_j \mid \sigma^2, \lambda_1, \lambda_2, \tau_j^2 \stackrel{\text{ind}}{\sim} \mathbf{N}\left(0, \frac{\sigma^2}{\lambda_2} \left(\frac{\lambda_2 \tau_j^2}{1 + \lambda_2 \tau_j^2}\right)\right),$$
(11)

$$\tau_j^2 \mid \lambda_1, \lambda_2 \stackrel{\text{iid}}{\sim} \text{UH}\left(1, \frac{1}{2}, \frac{\lambda_1^2}{2}, \lambda_2\right),$$
 (12)

where the UH (p, r, s, λ) distribution is a limit of the compound confluent hypergeometric (CCH) distribution (Gordy, 1998). The density function for this distribution is

$$\pi_d(\tau_j^2 \mid \lambda_1, \lambda_2) = \frac{1}{2\sqrt{2\pi}} \Phi\left(-\frac{\lambda_1}{\sqrt{\lambda_2}}\right)^{-1} \lambda_1 \lambda_2^{1/2} e^{-\frac{\lambda_1^2}{2\lambda_2}} (1 + \lambda_2 \tau_j^2)^{-1/2} e^{-\frac{\lambda_1^2 \tau_j^2}{2}}, \quad \tau_j^2 > 0.$$
(13)

Table 2 summarizes the different prior scalings and data augmentation representations that have appeared in the literature.

	Prior scaling $\pi(\beta \mid \sigma^2, \lambda_1, \lambda_2) \propto$	SMN variance $\beta_j \mid \sigma^2, \lambda_2, \tau_j^2 \sim \mathbf{N}(0, \cdot)$	Mixing distribution $\pi(\tau_i^2 \mid \sigma^2, \lambda_1, \lambda_2) \propto$	Mixing distribution family
Li and Lin (2010)	$\exp\left\{-\tfrac{\lambda_2}{2\sigma^2}\beta^T\beta-\tfrac{\lambda_1}{2\sigma^2} \beta _1\right\}$	$\frac{\sigma^2}{\lambda_2} \left(\frac{\tau_j^2 - 1}{\tau_j^2} \right)$	$\left(\tau_{j}^{2}\right)^{-\frac{1}{2}}e^{-\tau_{j}^{2}\frac{\lambda_{j}^{2}}{8\sigma^{2}\lambda_{2}}}, \tau_{j}^{2} \ge 1,$	$\operatorname{Gamma}_{(1,\infty)}\left(rac{1}{2},rac{\lambda_1^2}{8\sigma^2\lambda_2} ight)$
Hans (2011)	$\exp\left\{-\tfrac{\lambda_2}{2\sigma^2}\beta^T\beta-\tfrac{\lambda_1}{2\sigma^2} \beta _1\right\}$	$\tfrac{\sigma^2}{\lambda_2} \left(1 - \tau_j^2 \right)$	$\left(\tau_{j}^{2}\right)^{-\frac{3}{2}}e^{-\tau_{j}^{-2}\frac{\lambda_{1}^{2}}{8\sigma^{2}\lambda_{2}}}, 0 \le \tau_{j}^{2} \le 1,$	Inv-Gamma _(0,1) $\left(\frac{1}{2}, \frac{\lambda_1^2}{8\sigma^2\lambda_2}\right)$
Kyung et al. (2010)	$\exp\left\{-\frac{\lambda_2}{2\sigma^2}\beta^T\beta - \frac{\lambda_1}{\sigma} \beta _1\right\}$	*	*	*
Roy and Chakraborty (2017)	$\exp\left\{-\frac{\lambda_2}{2\sigma^2}\beta^T\beta-\frac{\lambda_1}{\sigma} \beta _1\right\}$	$\frac{\sigma^2}{\lambda_2} \left(\frac{\lambda_2 \tau_j^2}{1 + \lambda_2 \tau_j^2} \right)$	$\left(1+\lambda_2\tau_j^2\right)^{-1/2}e^{-\tau_j^2\frac{\lambda_1^2}{2}}, \tau_j^2 \ge 0$	$\operatorname{UH}\left(1,rac{1}{2},rac{\lambda_{1}^{2}}{2},\lambda_{2} ight)$

Table 2: Prior scaling and data augmentation parameterization in the Bayesian elastic net literature. Double horizontal lines differentiate between approaches for scaling the ℓ_1 -norm term in the prior density. Entries in the "SMN variance" column are the variances of the normal distribution in the scalemixture-of-normals (SMN) representation of the prior; entries in the "Mixing distribution" column are the density functions for those random variances (with the distributional families named in the final column). "Gamma_(1,∞)" is a gamma distribution truncated to the interval (1, ∞), "Inv-Gamma_(0,1)" is an inverse gamma distribution truncated to the interval (0, 1), and the "UH" is distribution is a limit of the compound confluent hypergeometric (CCH) distribution (Gordy, 1998). The SMN representation described in Kyung *et al.* (2010) is inconsistent with the claimed marginal distribution of β .

2.3 Posterior Computation: Sampling β

The two main approaches for representing the prior distribution lead to two main approaches for sampling β from its posterior. Under the direct representation of the prior, Hans (2011) showed that the conditional posterior distribution of β given σ^2 , λ_1 , and λ_2 is an orthant normal distribution:

$$\pi(\beta \mid y, \sigma^2, \lambda_1, \lambda_2) = \sum_{z \in \mathcal{Z}} \omega_z \mathbf{N}^{[z]} \left(\beta \mid \mu_z, \sigma^2 R\right), \tag{14}$$

where the ω_z are non-negative, orthant-specific weights that sum to one. The parameters of the underlying normal distributions that generate this posterior have connections to Bayesian ridge regression (Jeffreys, 1961; Raiffa and Schlaifer, 1961; Hoerl and Kennard, 1970). Under both the common and differential scalings of the prior, $R = (X^T X + \lambda_2 I_p)^{-1}$ so that $\sigma^2 R$ is the same as the posterior covariance matrix for Bayesian ridge regression for a fixed λ_2 . The orthant-specific location vectors under the commonly-scaled prior are $\mu_z = \hat{\beta}_R - \frac{\lambda_1}{2}Rz$, where $\hat{\beta}_R = RX^T y$ is the ridge regression estimate of β ; under the differentially-scaled prior, the location vectors are $\mu_z =$ $\hat{\beta}_R - \sigma \lambda_1 Rz$ instead. Under both scalings, the orthant-specific weights are $\omega_z = \omega^{-1} \frac{P(z,\mu_z,\sigma^2 R)}{N(0|\mu_z,\sigma^2 R)}$, where $P(z,\mu_z,\sigma^2 R) = \int_{\mathcal{O}_z} N(u \mid \mu_z,\sigma^2 R) du$ and $\omega = \sum_{z \in \mathcal{Z}} \frac{P(z,\mu_z,\sigma^2 R)}{N(0|\mu_z,\sigma^2 R)}$. A contour plot of the joint posterior density function $\pi_c(\beta \mid y, \sigma^2, \lambda_1, \lambda_2)$ for an example data set when p = 2 is shown in the left panel of Figure 2. The posterior is Gaussian within each of the four orthants (quadrants) of \mathbb{R}^2 , and the density function is continuous (bot not differentiable) along the coordinate axes due to the $|\beta|_1$ term in the prior. This particular example illustrates a situation where the λ_1 penalty term is large enough that the posterior mode lies on one of the coordinate axes ($\beta_2 = 0$).

Sampling directly from (14) is challenging. An obvious approach is to first sample z from the discrete distribution over the 2^p orthants (each having probability ω_z) and then, conditionally on the sampled orthant, to sample from a multivariate normal distribution truncated to \mathcal{O}_z . This requires the ability to compute numerically the orthant probabilities $P(z, \mu_z, \sigma^2 R)$ and the ability to sample directly from the multivariate truncated normal distribution $N^{[z]}(\beta \mid \mu_z, \sigma^2 R)$. When p is not too large, the former can sometimes be achieved using, e.g., the pmvnorm function in the R package mvtnorm (Genz, 1992; Genz and Bretz, 2009; Genz *et al.*, 2024; R Core Team, 2024), though if λ_1 is very large, the probabilities might be too small to compute accurately. The latter, sampling directly from the truncated multivariate normal distribution, may be difficult even when p is small. When p is large, direct sampling from (14) is not practical.

Hans (2011) shows how these issues can be avoided via Gibbs sampling. The full conditional posterior distribution for β_j is a one-dimensional orthant normal distribution:

$$\pi(\beta_j \mid y, \beta_{-j}, \sigma^2, \lambda_1, \lambda_2) = (1 - \phi_j) \mathbf{N}^-(\beta_j \mid \mu_j^-, s_j^2) + \phi_j \mathbf{N}^+(\beta_j \mid \mu_j^+, s_j^2).$$
(15)

The scale parameters are $s_j^2 = \sigma^2/(x_j^T x_j + \lambda_2)$. Hans (2011) provides an interpretable expression for the location parameter for the positive component under the commonly-scaled prior (1):

$$\mu_{j}^{+} = \hat{\beta}_{R,j} + \left\{ \sum_{i \neq j} \left(\hat{\beta}_{R,i} - \beta_{i} \right) \frac{x_{i}^{T} x_{j}}{x_{j}^{T} x_{j} + \lambda_{2}} \right\} + \frac{-\lambda_{1}}{2(x_{j}^{T} x_{j} + \lambda_{2})},$$
(16)

where $\hat{R}_{R,i}$ is the *i*th component of the ridge regression estimate $\hat{\beta}_R$ for the given value of λ_2 , and $+\lambda_1$ replaces $-\lambda_1$ in the corresponding expression for μ_j^- . Under the differentially-scaled prior

(4), the trailing term for μ_j^{\pm} is $\mp \sigma \lambda_1 / (x_j^T x_j + \lambda_2)$ instead. While (16) has a familiar form—it looks like the usual formula for the conditional mean of a Bayesian ridge regression posterior with an additional penalty term involving λ_1 —a more computationally efficient expression under the commonly-scaled prior (1) is

$$\mu_j^+ = \frac{x_j^T y - (x_j^T X_{-j})\beta_{-j} - \lambda_1/2}{x_j^T x_j + \lambda_2},$$

where $(-\lambda_1/2)$ is replaced with $(+\lambda_1/2)$ in the expression for μ_j^- . Under the differentially-scaled prior (4), $\pm \lambda_1/2$ is replaced by $\pm \sigma \lambda_1$. The expression is computationally efficient because $X^T X$ and $X^T y$ can be precomputed before the start of the MCMC algorithm. Completing the description of the full conditional density, the negative and non-negative components of (15) are weighted by

$$\phi_j = \left\{ \frac{\Phi(\mu_j^+/s_j)}{\mathsf{N}(0 \mid \mu_j^+, s_j^2)} \right\} \left/ \left\{ \frac{\Phi(\mu_j^+/s_j)}{\mathsf{N}(0 \mid \mu_j^+, s_j^2)} + \frac{\Phi(-\mu_j^-/s_j)}{\mathsf{N}(0 \mid \mu_j^-, s_j^2)} \right\}.$$

An example full conditional distribution is shown in Figure 2. It is easy to sample from these distributions. The standard normal cdf, $\Phi(\cdot)$, can be computed to high numerical precision (e.g., using the pnorm function in R), and efficient algorithms exist for sampling from univariate truncated normal distributions (e.g., the rejection sampling approach of Geweke, 1991).

The DA representation of the prior suggests an alternative approach for sampling from the posterior distribution of β . Li and Lin (2010) and Hans (2011) describe a two-stage, data augmentation Gibbs sampler that samples alternately from the conditional posterior of β given the latent scale parameters, τ^2 , and then from the conditional posterior of the latent scale parameters, τ^2 , given β . Under both scalings of the priors, the full vector of regression coefficients, β , is sampled from the normal distribution N($\hat{\beta}_{R_*}, \sigma^2 R_*$), where $\hat{\beta}_{R_*} = R_* X^T y$. Under the commonly-scaled prior (1), R_* has the form $R_c = (X^T X + \lambda_2 S_{\tau}^{-1})^{-1}$, where $S_{\tau} = \text{diag}(1 - \tau_j^2)$; under the differentiallyscaled prior (4), R_* has the form $R_d = (X^T X + D_{\tau}^{-1})^{-1}$, where $D_{\tau} = \text{diag}((\tau_j^{-2} + \lambda_2)^{-1})$. Under the commonly-scaled prior (1), the latent scale parameters are updated by sampling ζ_j independently from inverse Gaussian distributions with shape parameters $\lambda_1^2/(4\lambda_2\sigma^2)$ and means $\lambda_1/(2\lambda_2|\beta_j|)$, and then transforming $\tau_j^2 = \zeta_j/(1 + \zeta_j)$. Under the differentially-scaled prior (4), the latent scale parameters are updated by sampling ζ_j independently from inverse Gaussian dis-



Figure 2: Left: contours of the joint posterior density function $\pi_c(\beta_1, \beta_2 \mid y, \sigma^2, \lambda_1, \lambda_2)$ for an example data set with p = 2. Right: corresponding full conditional density function $\pi_c(\beta_1 \mid y, \beta_2 = 0.5, \sigma^2, \lambda_1, \lambda_2)$ when $\beta_2 = 0.5$ (the dashed line in the plot on the left).

tributions with shape parameters λ_1^2 and means $\sigma \lambda_1 / |\beta_j|$, and then transforming $\tau_j^2 = 1/\zeta_j$. This sampling scheme has the advantage that the β_j are updated as a block, which may be more effective when the β_j are highly correlated in the posterior. The trade-off for the potential reduction in autocorrelation is the need to simulate an additional p latent variables.

2.4 Posterior Computation: Sampling σ^2

A common choice of prior for σ^2 in Bayesian linear regression is an inverse gamma distribution, Inv-Gamma($\nu_a/2, \nu_b/2$), or its improper limit with $\nu_a = \nu_b = 0$ so that $\pi(\sigma^2) \propto 1/\sigma^2$. Kyung *et al.* (2010) Li and Lin (2010), Hans (2011), Roy and Chakraborty (2017), and Wang and Wang (2023) all used a version of this prior in their treatments of the Bayesian elastic net. Under this prior, the form of the full conditional distribution for σ^2 depends on the form and representation for the prior on β . The full conditional has the simplest form under the differentially-scaled prior on β . As shown in Roy and Chakraborty (2017), the corresponding full conditional distribution for σ^2 under the DA representation of the differentially-scaled prior on β is

$$\sigma^2 \mid y, \beta, \tau^2, \lambda_1, \lambda_2 \sim \text{Inv-Gamma}\left(\frac{n+p-1+\nu_a}{2}, \frac{\nu_b + (y-X\beta)^T (y-X\beta) + \sum_{j=1}^p \beta_j^2 (\tau_j^{-2} + \lambda_2)}{2}\right)$$

an inverse gamma distribution that does not depend on λ_1 .

The full conditional for σ^2 under the direct representation of the differentially-scaled prior has not yet been studied in the literature, but it can be shown that

$$\frac{1}{\sigma^2} \mid y, \beta, \tau^2, \lambda_1, \lambda_2 \sim \text{MHN}\left(\frac{n+p-1+\nu_a}{2}, \frac{(y-X\beta)^T(y-X\beta)+\lambda_2\beta^T\beta+\nu_b}{2}, \lambda_1|\beta|_1\right),$$

a modified half-normal distribution (see Section 3.1 for details). Algorithms for sampling efficiently from this distribution exist (see, e.g., Sun *et al.*, 2023).

The full conditional distribution for σ^2 under the commonly-scaled prior on β is more complex. Hans (2011) worked with this full conditional under the direct representation of the prior, which has density function

$$\pi_c(\sigma^2 \mid y, \beta, \lambda_1, \lambda_2) \propto \Phi\left(-\frac{\lambda_1}{2\sigma\sqrt{\lambda_2}}\right)^{-p} (\sigma^2)^{-(n+p-1+\nu_a)/2-1} \times \exp\left\{-\frac{1}{\sigma^2} \left((y-X\beta)^T (y-X\beta) + \lambda_2\beta^T\beta + \lambda_1|\beta|_1 + p\lambda_1^2/(4\lambda_2) + \nu_b\right)/2\right\}.$$

This is a non-standard distribution that involves an analytically intractable integral expression, $\Phi(-\lambda_1/(2\sigma\sqrt{\lambda_2}))$. Noting that as long as $\lambda_1/(2\sigma\sqrt{\lambda_2})$ is not too large, $\Phi(-\lambda_1/(2\sigma\sqrt{\lambda_2}))$ (or its logarithm) can be evaluated numerically to relatively high precision, Hans (2011) used a "Metropoliswithin-Gibbs" step to update σ^2 on its log scale by sampling a proposal, $\log \sigma^{2*}$, from a normal distribution centered at the current value, $\log \sigma^2$, with a pre-specified innovation variance. The probability of accepting the proposed value (as opposed to staying at the current value) was calculated using the ratio of the full conditional for $\log \sigma^2$ evaluated at the proposed and current values. This algorithm tends to work well in practice, though poor choice of the innovation variance can result in slow convergence and mixing.

Li and Lin (2010) and Hans (2011) both work with the full conditional for σ^2 under the DA

representation of the commonly-scaled prior for β . The full conditional has density function

$$\pi_{c}(\sigma^{2} \mid y, \beta, \tau^{2}, \lambda_{1}, \lambda_{2}) \propto \Phi\left(-\frac{\lambda_{1}}{2\sigma\sqrt{\lambda_{2}}}\right)^{-p} (\sigma^{2})^{-(n+2p-1+\nu_{a})/2-1} \times \exp\left\{-\frac{1}{2\sigma^{2}}\left(\nu_{b} + (y - X\beta)^{T}(y - X\beta) + \lambda_{2}\beta^{T}S_{\tau}^{-1}\beta + \frac{\lambda_{1}^{2}}{4\lambda_{2}}\sum_{j=1}^{p}\tau_{j}^{-2}\right)\right\}$$

Hans (2011) used a random-walk Metropolis algorithm for updating $\log \sigma^2$. Li and Lin (2010) reexpressed this density in terms of the upper incomplete gamma function (DiDonato and Morris, 1986), $\Gamma_U(\alpha, x) = \int_x^\infty t^{\alpha-1} e^{-t} dt$, through the equivalence $2\sqrt{\pi} \Phi\left(\frac{-\lambda_1}{2\sigma\sqrt{\lambda_2}}\right) = \Gamma_U\left(\frac{1}{2}, \frac{1}{2}\left(\frac{\lambda_1}{2\sigma\sqrt{\lambda_2}}\right)^2\right)$, and proposed obtaining exact samples from this full conditional distribution via rejecting sampling using an inverse gamma proposal distribution. Unfortunately, the derivation of the acceptance probability for the algorithm contains an error and the resulting samples do not come from the desired target distribution. We identify the problem in detail in Appendix A.

2.5 Posterior Computation: Sampling λ_1 and λ_2

Posterior sampling of λ_1 and λ_2 is non-trivial under both scalings of the prior on β whether or not data augmentation is used. The term $\Phi(-\lambda_1/(2\sigma\sqrt{\lambda_2}))^{-p}$ (under prior (7)) or $\Phi(-\lambda_1/\sqrt{\lambda_2})^{-p}$ (under prior (9)) appears in the posterior, making direct sampling of these parameters difficult. Li and Lin (2010) and Roy and Chakraborty (2017) eschew full Bayesian inference and instead propose methods for selecting values for these hyperparameters. Hans (2011) uses separate random-walk Metropolis updates for $\log \lambda_1$ and $\log \lambda_2$ with their full conditionals as the target densities. This approach requires specification of a step-size parameter for the normal proposal, which can be difficult to select and tune. Motivated by the desire to avoid numerical computation of the standard normal cdf, $\Phi(\cdot)$, Wang and Wang (2023) devise a clever exchange algorithm (Murray *et al.*, 2006) that introduces p additional latent variables in such a way as to remove the term involving $\Phi(\cdot)$ from the joint posterior of the augmented parameter space. Despite avoiding computation of $\Phi(\cdot)$, the algorithm still requires one parameter to be updated via the Metropolis–Hastings algorithm using a random-walk proposal, necessitating the selection of a step-size parameter.

All of the correctly-specified approaches reviewed above for full Bayesian inference for the

Bayesian elastic net require the specification of at least one step-size parameter in a Metropolis– Hastings step in a Gibbs sampler. This can be challenging in practice, as appropriate scales for the step sizes are not always obvious before running the MCMC algorithm. If poor step sizes are chosen, the resulting Markov chains will mix slowly. Practitioners who make use of appropriate post-sampling MCMC diagnostics might notice this, adjust the step size, and rerun the chain (perhaps iterating this procedure several times); practitioners who simply use the original MCMC output as is will produce low-quality summaries of the posterior.

We introduce in Section 3 a new approach to posterior sampling for full Bayesian inference for the Bayesian elastic net that avoids these issues entirely. We use a simple transformation of the parameter space to (i) reduce the number of parameters whose full conditional densities have a term involving $\Phi(\cdot)$ and (ii) produce log-concave full conditional distributions that can be easily sampled via a highly-efficient rejection sampling algorithm using automatically-tuned, piece-wise exponential proposal distributions. Importantly, the approach requires no tuning on the part of the analyst.

3 Efficient Rejection Sampling for Full Bayesian Infer-

ence

Full Bayesian inference for Bayesian elastic net regression proceeds by assigning prior distributions to σ^2 , λ_1 , and λ_2 , and making inferences based on the joint posterior $\pi(\beta, \sigma^2, \lambda_1, \lambda_2 \mid y)$ and its margins. Li and Lin (2010), Hans (2011), and Roy and Chakraborty (2017) all assign to σ^2 an inverse gamma prior, $\sigma^2 \sim$ Inv-Gamma($\nu_a/2, \nu_b/2$), or the improper prior with $\pi(\sigma^2) \propto \sigma^{-2}$. Under the commonly-scaled prior for β , Hans (2011) considered the hyperprior distributions $\lambda_1 \sim$ Gamma($L, \nu_1/2$) and $\lambda_2 \sim$ Gamma($R, \nu_2/2$), where the gamma distributions are parameterized to have mean $2L/\nu_1$ and $2R/\nu_2$. When the prior is parameterized as in (1), Zou and Hastie (2005) noted that $\lambda = \lambda_1 + \lambda_2$ represents the total penalization and $\alpha = \lambda_2/(\lambda_1 + \lambda_2)$ represents the proportion of the total penalization that is attributable to the ℓ_2 -norm component. As noted in Hans (2011), when the gamma priors on λ_1 and λ_2 are independent and $\nu_1 = \nu_2 = \nu$, the induced priors on the transformed parameters are $\lambda \sim \text{Gamma}(R + L, \nu/2)$ and $\alpha \sim \text{Beta}(R, L)$, with λ and α independent. Considering the hyperpriors from these two perspectives gives the user a range of interpretations to consider when specifying L, R, ν_1 , and ν_2 . We use these prior distributions for σ^2 , λ_1 , and λ_2 unless noted otherwise.

3.1 Rejection sampling for a class of distributions

Our first approach to posterior sampling, described in Section 3.2, requires the ability to generate random variates from density functions of the form

$$f(x) \propto \Phi(-x)^{-q} x^{a-1} e^{-bx^2 - cx - d/x}, \ x > 0,$$
(18)

where $q \in \{0, 1, ...\}$. The function f is a density function under various conditions on q, a, b, c, and d. Three conditions are of special interest to us.

First, the conditions $\{q = 0, a \in \mathbb{R}, b = 0, c > 0, d > 0\}$ correspond to the family of generalized inverse Gaussian (GIG) distributions (Barndorff-Nielsen, 1977, 1978). Devroye (2014) introduced a rejection sampling algorithm for sampling GIG random variates X based on log-concavity of the density of log X. Second, the conditions $\{q = 0, a > 0, b > 0, c \in \mathbb{R}, d = 0\}$ correspond to the family of modified half normal (MHN) distributions. Sun (2020) and Sun *et al.* (2023) introduce efficient rejection sampling algorithms for obtaining samples from this class of distributions.

We can use the rejection sampling algorithms of Devroye (2014) and Sun *et al.* (2023) to obtain samples from the GIG and MHN distributions, respectively. When $a \ge 1$, the GIG and MHN distributions both have log concave densities, in which case we could also use rejection sampling techniques that exploit log concavity. We describe such an approach here that can be used to obtain samples from (18) when q = 0, $a \ge 1$, and b, c, and d are such that f(x) is integrable and log concave. The approach is strongly connected to the work of Devroye (1984, 1986), Gilks (1992), and Gilks and Wild (1992) in the sense that a piece-wise exponential hull is used to bound the target density, with proposals drawn from the corresponding piece-wise exponential distribution. The approach is not "adaptive" in the sense that the piece-wise exponential hull is not refined if a proposal is rejected, but it is "adapted" to the target density because information about the log density's mode and curvature at the mode are used to construct the proposal density. The approach was used by Hans (2009) to sample σ^2 for Bayesian lasso regression.

To construct the piece-wise exponential hull, we make use of the unique mode, x_* , of the distribution. For the GIG distribution with $a \ge 1$, the unique mode occurs at $(a-1+\sqrt{(a-1)^2+4cd})/(ac)$; for the MHN distribution with $a \ge 1$, the unique mode occurs at $(-c + \sqrt{c^2 + 8b(a-1)})/(4b)$. The piece-wise exponential hull is created by first placing a knot point at the mode, x_* . Several knot points are then placed at appropriate distances above and below the mode. To determine where to place these additional knot points, we make use of the curvature of $\log f$ at is mode, $f''(x_*) = -(a-1)/x_* - 2b - 2d/x_*^3$, by noting that if f was the density for a Gaussian distribution, $s_{x_*} = |f''(x_*)|^{-1/2}$ would be the standard deviation of the distribution and would provide a scale to inform us about where to place the knots. Using this second-order Taylor polynomial approximation to $\log f(x)$, we place one knot at $x_* + s_{x_*}/2$, and then K additional knots at $x_* + ks_{x_*}$, $k = 1, \ldots, K$. Below the mode, we place knots at $x_* - s_{x_*}/2$ and $x_* - ks_{x_*}, k = 1, \ldots, K$. Any negative knots are then removed from the set; if no knots remain below the mode, a single knot is then placed at $x_*/2$. Lines tangent to log f at the knot points are then used to construct a piecewise linear upper hull for $\log f$, with change points occurring at the intersections of the tangent lines. The piece-wise linear upper hull is exponentiated to obtain a piece-wise exponential hull for f, which can then be rescaled and used as a proposal distribution for rejection sampling. The key elements of the approach are depicted graphically in Figure 3. Practical experience suggests that a small number of knot points (K = 2 or K = 3) results in high acceptance rates with low computational overhead.

The third set of conditions under which we will need to sample from f(x) is $\{q \in \{1, 2, ...\}, a > 0, b \ge q/2, c > 0, d = 0\}$. This density function is more challenging because it contains the non-trivial term $\Phi(-x)^{-q}$. Under the more strict condition that $a \ge 1$, we can show that f(x) is log concave, and a modified version of our rejection sampling algorithm can be used to obtain samples from f(x).

Proposition 1. The function $f(x) \propto \Phi(-x)^{-q} x^{a-1} e^{-bx^2 - cx}$, x > 0, is integrable and log concave when $q \in \{1, 2, ...\}$, $a \ge 1$, $b \ge q/2$, and c > 0.



Figure 3: Illustration of the rejection sampling appraoch described in Section 3.1 for a target density f(x) with q = 0, a = 3, b = 2, c = 2, and d = 0 (a modified half normal distribution). The left panel displays the piece-wise linear hull (blue line) for $\log f(x)$ (black line), where both functions are shifted to take the value 0 at the mode, $x_* = 0.5$. The right panel displays the piece-wise exponential hull (blue line) for f(x) (black line), where both functions are scaled to take the value 1 at the mode. The black points indicate the locations of the "knot points", with "*" corresponding to the mode, x_* . The vertical red dashed lines partition the support of f according to the intersections of the lines tangent to f at the knot points and define the change points for the piece-wise exponential proposal distribution. The rejection sampler for this example has an acceptance probability of approximately 0.954.

Proof. It is clear that $\int_0^t f(x)dx < \infty$ for all $t < \infty$ because $2^q \leq \Phi(-x)^{-q} < \infty$ for all $0 \leq x \leq t < \infty$. By Feller (1968), $\Phi(-x) \geq (2\pi)^{-1/2}(x^{-1} - x^{-3})e^{-x^2/2}$, and so for large enough x, $\Phi(-x)^{-q} \leq (2\pi)^{q/2}x^{3q}e^{qx^2/2}$. For large enough t,

$$\begin{split} \int_{t}^{\infty} \Phi(-x)^{-q} x^{a-1} e^{-bx^{2}-cx} dx &= \int_{t}^{\infty} \Phi(-x)^{-q} e^{-qx^{2}/2} x^{a-1} e^{-(b-q/2)x^{2}-cx} dx \\ &\leq (2\pi)^{q/2} \int_{t}^{\infty} x^{3q+a-1} e^{-(b-q/2)x^{2}-cx} dx \\ &< \infty, \end{split}$$

and so f(x) is integrable.

Now let $Z \sim N(0, I_q)$ be a q-vector of independent, standard normal random variables, and let $|Z|_1 = \sum_{j=1}^q |Z_j|$. For x > 0, it can be shown that $\mathbb{E}\left[e^{-x|Z|_1}\right] = 2^q e^{qx^2/2} \Phi(-x)^q$. We can then write $f(x) \propto \mathbb{E}\left[e^{-x|Z|_1}\right]^{-1} x^{a-1} e^{-(b-q/2)x^2 - cx}$. The term $x^{a-1} e^{-(b-q/2)x^2 - cx}$ is log concave when $a \ge 1$, $b \ge q/2$, and c > 0, and so we need only show log convexity of $h(x) \equiv \mathbb{E}\left[e^{-x|Z|_1}\right]$.

For any $\alpha \in [0, 1]$, $x_1 > 0$, and $x_2 > 0$,

$$\begin{split} h((1-\alpha)x_1 + \alpha x_2) &= \int_{\mathbb{R}^q} e^{-((1-\alpha)x_1 + \alpha x_2)|z|_1} \mathbf{N}(z \mid 0, I_q) dz \\ &= \int_{\mathbb{R}^q} \left(e^{-x_1|z|_1} \mathbf{N}(z \mid 0, I_q) \right)^{1-\alpha} \left(e^{-x_2|z|_1} \mathbf{N}(z \mid 0, I_q) \right)^{\alpha} dz \\ &\leq \left\{ \int_{\mathbb{R}^q} \left(e^{-x_1|z|_1} \mathbf{N}(z \mid 0, I_q) \right) dz \right\}^{1-\alpha} \left\{ \int_{\mathbb{R}^q} \left(e^{-x_2|z|_1} \mathbf{N}(z \mid 0, I_q) \right) dz \right\}^{\alpha} \\ &= h(x_1)^{1-\alpha} h(x_2)^{\alpha}, \end{split}$$

where the inequality is due to Hölder's inequality. The function h(x) is therefore log convex and hence f(x) is log concave.

To implement the same rejection sampling algorithm as above, we need to be able to find the mode of $\log f(x)$, which has derivative

$$\frac{d}{dx}\log f(x) = q\frac{\phi(-x)}{\Phi(-x)} + \frac{a-1}{x} - 2bx - c.$$

The mode is no longer available as the unique, positive root of a polynomial. A mode-finding algorithm could be used to approximate x_* at some additional computational cost. Instead, we note that in order to construct an integrable, piece-wise exponential hull for f(x), we need only identify at least one knot point above the mode because the support of x is bounded below by zero; additionally identifying at least one point below the mode will help improve the quality of the piece-wise exponential approximation. We can facilitate the choice of such knot points using the following result.

Proposition 2. When $q \in \{1, 2, ...\}$, $a \ge 1$, $b \ge q/2$, and c > 0, the function $f(x) \propto \Phi(-x)^{-q}x^{a-1}e^{-bx^2-cx}$, x > 0, has a unique mode, x_* , satisfying

$$\begin{split} \frac{a-1}{c} < x_* < \frac{a-1+q}{c}, & \text{if } a \geq 1, q = 2b, \\ 0 < x_* < \frac{\sqrt{c^2 + 4q(2b-q)} - c}{2(2b-q)}, & \text{if } a = 1, q > 2b, \text{ and} \\ \frac{\sqrt{c^2 + 4(a-1)(2b-q)} - c}{2(2b-q)} < x_* < \frac{\sqrt{c^2 + 4(a-1+q)(2b-q)} - c}{2(2b-q)}, & \text{if } a > 1, q > 2b. \end{split}$$

Proof. The function f(x) has a unique mode because it is log concave, and the derivative

$$\frac{d}{dx}\log f(x) = q\frac{\phi(-x)}{\Phi(-x)} + \frac{a-1}{x} - 2bx - c$$

is decreasing in x for the same reason. By Gordon (1941), we have

$$x < \frac{\phi(-x)}{\Phi(-x)} < x + \frac{1}{x},$$

for x > 0, and so we can bound the derivative of $\log f(x)$ as

$$g_L(x) \equiv \frac{a-1}{x} - (2b-q)x - c < \frac{d}{dx}\log f(x) < \frac{a-1+q}{x} - (2b-q)x - c \equiv g_U(x).$$

When b = q/2 and a > 1, $g_L(x)$ and $g_U(x)$ are both are decreasing functions. The sign of $\frac{d}{dx} \log f(x)$ is therefore positive when x < (a-1)/c due to $g_L(x)$ and negative when x > (a-1+q)/c due to $g_U(x)$ and so, by the intermediate value theorem, $(a-1)/c < x_* < (a-1+q)/c$.

Next, when b = q/2 and a = 1, $g_L(x) = -c < 0$ for all x > 0 and so the lower bound is not useful. The upper bound, $g_U(x)$, is decreasing, and so $\frac{d}{dx} \log f(x)$ is negative when x > q/c. By the intermediate value theorem, we must have $0 < x_* < q/c$ because $\lim_{x \to 0^+} \frac{d}{dx} \log f(x) = \infty$.

Now focusing on the case b > q/2, when a = 1 we have $g_L(x) = -(2b-q)x - c < 0$ for all x > 0, and so the lower bound is not useful. The upper bound is $g_U(x) = q/x - (2b-q)x - c$, a decreasing function. The sign of $\frac{d}{dx} \log f(x)$ is therefore negative when $x > \frac{\sqrt{c^2 + 4q(2b-q)} - c}{2(2b-q)}$, and so by the same arguments as above we must have $0 < x_* < \frac{\sqrt{c^2 + 4q(2b-q)} - c}{2(2b-q)}$.

Finally, when b > q/2 and a > 1, $g_L(x) = (a - 1)/x - (2b - q)x - c$ and $g_U(x) = (a - 1 + q)/x - (2b - q)x - c$, and both are decreasing functions. The same arguments as above yields the resulting bounds on x_* .

We can therefore find one point above the mode and, when a > 1, one point below the mode that can be used to construct a piecewise linear upper hull for $\log f(x)$. With only one or two knot points which might be ill-positioned depending on the quality of the bound(s), the resulting piecewise exponential proposal distribution may result in a high rejection rate, requiring either additional well-placed knot points to start or a strategy for adapting the hull as proposals are rejected. Given the difficulty of finding a suitable set of knot points, we instead use the traditional adaptive rejection sampling algorithm Gilks and Wild (1992) to sample from f(x) when q > 0. In practice, we use the ars package (Perez Rodriguez, 2024) in R (R Core Team, 2024), supplying the ars function with inputs $\log f(x)$, $\frac{d}{dx} \log f(x)$, a lower bound of x = 0, and one (or two) initial knot points above (and below) the mode.

3.2 Rejection sampling for full Bayesian inference

As described in Section 2.5, the full conditional density functions for λ_1 and λ_2 under the differentiallyscaled prior are not available in closed form due to the $\Phi(-\lambda_1/\sqrt{\lambda_2})$ term in (8) under the direct representation or in (13) under the data augmentation representation. The same is true for σ^2 under the commonly-scaled prior due to the $\Phi(-\lambda_1/(2\sigma\sqrt{\lambda_2}))$ term in (6) and (10). As an improvement to existing MCMC methods for full Bayesian inference, we consider a transformation of the parameter space that (i) confines the awkward $\Phi(\cdot)$ term to a single full conditional distribution and (ii) results in log-concave full conditional density functions for all parameters that do not have "standard" (easy to sample from) full conditional distributions. We then exploit log-concavity to construct efficient rejection sampling algorithms for these parameters. The form of the transformation depends on whether the commonly- or differentially-scaled prior for β is used. We start with the commonly-scaled prior.

3.2.1 Sampling under the commonly-scaled prior

Under prior (7), define the transformation $(\sigma^2, \lambda_1, \lambda_2) \rightarrow (u_1 = \sigma^2, u_2 = \sqrt{\lambda_2}/\sigma, \theta = \lambda_1/(2\sigma\sqrt{\lambda_2}))$. The reparameterized prior on the regression coefficients is then

$$\pi_c(\beta \mid u_1, u_2, \theta) = 2^{-p} (2\pi)^{-p/2} u_2^p e^{-\frac{p\theta^2}{2}} \Phi(-\theta)^{-p} \exp\left\{-u_2^2 \beta^T \beta/2 - u_2 \theta |\beta|_1\right\}.$$

The awkward term involving $\Phi(\cdot)$ is now a function of only θ . Transforming the prior on σ^2 , λ_1 , and λ_2 yields

$$\pi(u_1, u_2, \theta) \propto u_1^{R+L-\nu_a/2-1} u_2^{2R+L-1} \theta^{L-1} \exp\left\{-u_1 u_2^2 \nu_2/2 - u_1 u_2 \theta \nu_1 - u_1^{-1} \nu_b/2\right\}.$$
 (19)

Combining these priors with the likelihood function yields the following full conditional posterior distributions for u_1 , u_2 , and θ :

$$u_{1} \mid y, \beta, u_{2}, \theta \sim \operatorname{GIG} \left(R + L - (\nu_{a} + n - 1)/2, u_{2}^{2}\nu_{2} + 2u_{2}\theta\nu_{1}, (y - X\beta)^{T}(y - X\beta) + \nu_{b} \right),$$

$$u_{2} \mid y, \beta, u_{1}, \theta \sim \operatorname{MHN} \left(2R + L + p, \frac{u_{1}\nu_{2} + \beta^{T}\beta}{2}, \theta(u_{1}\nu_{1} + |\beta|_{1}) \right),$$

$$\pi_{c}(\theta \mid y, \beta, u_{1}, u_{2}) \propto \Phi(-\theta)^{-p} \theta^{L-1} \exp\left\{ -p\theta^{2}/2 - \theta u_{2}(u_{1}\nu_{1} + |\beta|_{1}) \right\}.$$

We obtain samples from these distributions as follows.

The full conditional posterior distribution for u_1 is a generalized inverse Gaussian (GIG) distribution (Barndorff-Nielsen, 1977, 1978). As discussed in Section 3.1, we can use the rejection sampling method of Devroye (2014) to sample u_1 from its full conditional. The full conditional for u_1 will be log concave when $R + L - (\nu_a + n - 1)/2 \ge 1$, in which case we could also use the method described in Section 3.1 that uses information about the full conditional at its mode to sample from this distribution.

The full conditional posterior distribution for u_2 is a modified half normal (MHN) distribution and will always be log concave as we assume R > 0 and L > 0 in the prior. We can therefore use either the rejection sampling method introduced by Sun *et al.* (2023) or the rejection sampling method described in Section 3.1 that uses information about the full conditional at its mode to sample from the full conditional.

The full conditional for θ has the form of (18) with $q = p \in \{1, 2, ..., \}$, b = p/2, and c > 0. By Proposition 1, the full conditional will be log concave as long as $L \ge 1$ in the prior on λ_1 , and we can use adaptive rejection sampling by identifying at least one knot point to the right of the distribution's mode via Proposition 2. The full conditional is not log concave when 0 < L < 1, in which case other methods for sampling θ would be required. While this might be considered a limitation to our approach to sampling, we note that the rejection sampling method described below under the DA representation of the prior requires only that L > 0, and so we can always simply use the DA Gibbs sampler when 0 < L < 1.

Focusing now on the DA representation of the prior, under the commonly-scaled prior (1) and the same transformation $(\sigma^2, \lambda_1, \lambda_2) \rightarrow (u_1 = \sigma^2, u_2 = \sqrt{\lambda_2}/\sigma, \theta = \lambda_1/(2\sigma\sqrt{\lambda_2}))$, the reparameterized joint prior on β and τ is

$$\pi_{c}(\beta,\tau \mid u_{1}, u_{2},\theta) = \pi_{c}(\beta \mid \tau, u_{1}, u_{2},\theta)\pi_{c}(\tau \mid u_{1}, u_{2},\theta)$$

$$\propto \Phi(-\theta)^{-p} \theta^{p} u_{2}^{p} \left[\prod_{j=1}^{p} \tau_{j}^{-3/2} (1-\tau_{j})^{-1/2}\right] \times$$

$$\exp\left\{-\frac{u_{2}^{2}}{2} \beta^{T} S_{\tau}^{-1} \beta - \frac{\theta^{2}}{2} \sum_{j=1}^{p} \tau_{j}^{-1}\right\}.$$
(20)

The awkward term involving $\Phi(\cdot)$ is now a function of only θ . Combining the transformed prior (19) with (20) and the likelihood yields the following full conditional distributions:

$$\begin{split} u_1 \mid y, \beta, \tau, u_2, \theta &\sim & \text{GIG} \left(R + L - (\nu_a + n - 1)/2, u_2^2 \nu_2 + 2u_2 \theta \nu_1, (y - X\beta)^T (y - X\beta) + \nu_b \right), \\ u_2 \mid y, \beta, \tau, u_1, \theta &\sim & \text{MHN} \left(2R + L + p, \frac{u_1 \nu_2 + \beta^T S_{\tau}^{-1} \beta}{2}, u_1 \theta \nu_1 \right), \\ \pi_c(\theta \mid y, \beta, \tau, u_1, u_2) &\propto & \Phi \left(-\theta \right)^{-p} \theta^{p+L-1} \exp \left\{ -\frac{\theta^2}{2} \left(\sum_{j=1}^p \tau_j^{-1} \right) - \theta u_1 u_2 \nu_1 \right\}. \end{split}$$

As above, we use the method of Devroye (2014) to sample from the inverse Gaussian full conditional for u_1 . The full conditional for u_2 will always be log concave and so we use either the method of Sun *et al.* (2023) or the rejection sampling method described in Section 3.1 to update u_2 . The full conditional for θ has the form of (18) with $q = p \in \{1, 2, ..., \}$, a = p + L > 1, $b = \sum_{j=1}^{p} \tau_j^{-1}/2$, and c > 0. Because $0 < \tau_j < 1$, we have $b = \sum_{j=1}^{p} \tau_j^{-1}/2 > p/2$, and by Proposition 1 the full conditional is log concave and we can sample from this distribution using adaptive rejection sampling by identifying at least one knot point to the right of the distribution's mode via Proposition 2.

3.2.2 Sampling under the differentially-scaled prior

Under the differentially-scaled prior (9), consider the transformation $(\lambda_2, \lambda_1) \rightarrow (u_2 = \sqrt{\lambda_2}, \theta = \lambda_1/\sqrt{\lambda_2})$. The reparameterized prior on the regression coefficients is

$$\pi_d(\beta \mid \sigma^2, u_2, \theta) = 2^p (2\pi)^{-p/2} (\sigma^2)^{-p/2} u_2^p e^{-p\theta^2/2} \Phi(-\theta)^{-p} \exp\left\{-\frac{u_2^2}{2\sigma^2} \beta^T \beta - \frac{\theta u_2}{\sigma} |\beta|_1\right\},$$

the prior on σ^2 remains an inverse gamma distribution (or its improper limit), and the prior on the transformed parameters is

$$\pi(u_2,\theta) \propto u_2^{2R+L-1} \theta^{L-1} e^{-u_2^2 \nu_2/2 - u_2 \theta \nu_1/2}.$$
(21)

Combining these priors with the likelihood function yields the following full conditional posterior distributions:

$$\begin{split} & \frac{1}{\sigma^2} \mid y, \beta, u_2, \theta \quad \sim \quad \mathrm{MHN}\left(\frac{\nu_a + p + n - 1}{2}, \frac{(y - X\beta)^T (y - X\beta) + u_2^2 \beta^T \beta + \nu_b}{2}, \theta u_2 |\beta|_1\right), \\ & u_2 \mid y, \beta, \sigma^2, \theta \quad \sim \quad \mathrm{MHN}\left(2R + L + p, \frac{\beta^T \beta / \sigma^2 + \nu_2}{2}, \theta \left(|\beta|_1 / \sigma + \nu_1 / 2\right)\right), \\ & \pi_d(\theta \mid y, \beta, \sigma^2, u_2) \quad \propto \quad \Phi\left(-\theta\right)^{-p} \theta^{L-1} \exp\left\{-\theta^2 p / 2 - \theta u_2 \left(|\beta|_1 / \sigma + \nu_1 / 2\right)\right\}. \end{split}$$

The full conditionals for σ^{-2} and u_2 are log concave, and so we can either use the rejection sampling methods described in Section 3.1 or the method of Sun *et al.* (2023) to obtain samples from the full conditionals. The full conditional for θ will be log concave when $L \ge 1$, in which case we can use rejection sampling as described in Section 3.1. When 0 < L < 1, the full conditional is not log concave and we cannot use this particular method of rejection sampling. As in Section 3.2.1, when 0 < L < 1 we can instead implement a DA Gibbs sampler as described below.

Under the DA representation of the differentially-scaled prior and the same transformation

 $(\lambda_2, \lambda_1) \to (u_2 = \sqrt{\lambda_2}, \theta = \lambda_1/\sqrt{\lambda_2})$, the reparameterized joint prior on β and τ^2 is

$$\pi_{d}(\beta, \tau^{2} \mid \sigma^{2}, u_{2}, \theta) = \pi_{d}(\beta \mid \tau^{2}, \sigma^{2}, u_{2}, \theta) \pi_{d}(\tau^{2} \mid \sigma^{2}, u_{2}, \theta)$$

$$\propto (\sigma^{2})^{-p/2} \Phi (-\theta)^{-p} \theta^{p} u_{2}^{2p} \left[\prod_{j=1}^{p} (\tau_{j}^{2})^{-1/2} \right] \times$$

$$\exp \left\{ -\frac{\theta^{2}}{2} \left(p + u_{2}^{2} \sum_{j=1}^{p} \tau_{j}^{2} \right) - \frac{1}{2\sigma^{2}} \sum_{j=1}^{p} \beta_{j}^{2} \left(\tau_{j}^{-2} + u_{2}^{2} \right) \right\}. (22)$$

Combining the transformed prior (21) with (22) and the likelihood yields the following full conditional posterior distributions, which can be sampled as described above:

$$\begin{split} \sigma^{2} \mid y, \beta, \tau^{2}, u_{2}, \theta &\sim \text{Inv-Gamma}\left(\frac{p + \nu_{a} + n - 1}{2}, \frac{\nu_{b} + (y - X\beta)^{T}(y - X\beta) + \sum_{j=1}^{p} \beta_{j}^{2}(\tau_{j}^{-2} + u_{2}^{2})}{2}\right) \\ u_{2} \mid y, \beta, \tau^{2}, \sigma^{2}, \theta &\sim \text{MHN}\left(2p + 2R + L, \frac{\beta^{T}\beta/\sigma^{2} + \nu_{2} + \theta^{2}\sum_{j=1}^{p} \tau_{j}^{2}}{2}, \frac{\theta\nu_{1}}{2}\right), \\ \pi_{d}(\theta \mid y, \beta, \tau^{2}, \sigma^{2}, u_{2}) &\propto \Phi\left(-\theta\right)^{-p} \theta^{p+L-1} \exp\left\{-\frac{\theta^{2}}{2}\left(p + u_{2}^{2}\sum_{j=1}^{p} \tau_{j}^{2}\right) - \theta u_{2}\nu_{1}/2\right\}. \end{split}$$

4 Simulations

We conduct a simulation study to document the relative performance of the various approaches to MCMC for full Bayesian elastic net inference under a several scenarios for data generation. The existing, correctly-specified approaches to MCMC for full Bayesian modeling under the elastic net are the Metropolis–Hastings (MH) approach of Hans (2011), the exchange algorithm (EX) approach of Wang and Wang (2023), and the transformation and rejection sampling (RS) approach introduced in this paper. Because we are not introducing new statistical models, our comparisons focus on dynamics of the Markov chains generated by the MCMC algorithms. We use effective sample size (ESS) for model parameters as a measure of efficacy of a given algorithm for a given data set. ESS is computed using the R package mcmcse (Flegal *et al.*, 2021) based on Gong and Flegal (2015).

We consider the four simulation settings used by Zou and Hastie (2005) in their original study of the elastic net (see also Hans, 2011). In Simulation 1, n = 20 observations are simulated from the normal linear regression model $y = X\beta + \varepsilon$ with $\beta = (3, 1.5, 0, 0, 2, 0, 0, 0)^T$. The error term is generated according to $\varepsilon \sim N(0, \sigma^2 I_n)$ with $\sigma = 3$. Each row of the $n \times p$ design matrix X is generated independently from a N(0, V) distribution with covariance matrix V, where $V_{ij} = 0.5^{|i-j|}$ for $1 \le i, j \le p = 8$. Simulation 2 uses $\beta_j = 0.85, j = 1, \ldots, 8$, but is otherwise the same as Simulation 1. Simulation 3 considers a higher-dimensional setting and larger sample size, with n = 100 and p = 40. The regression coefficients are $\beta_j = 2$ for $j = 1, \ldots, 10$ and $j = 21, \ldots, 30$, and $\beta_j = 0$ for all other j. The errors are generated with $\sigma = 15$, and the regressors are generated with $V_{ij} = 0.5$ for $i \ne j$ and $V_{ii} = 1$. Simulation 4 is the same as Simulation 3, but $\beta_j = 3$ for $j = 1, \ldots, 15, \beta_j = 0$ for $j = 16, \ldots, 40$, and the regressors are generated using a block-diagonal covariance matrix V as follows. The first block $(1 \le i, j \le 5)$ has variances of 1.01 and covariances of 1, the second ($6 \le i, j \le 10$) and third ($11 \le i, j \le 15$) are the same as the first, and the final block is the 25×25 identity matrix. Covariances between all blocks are zero.

Fifty data sets y and X were generated for each simulation setting. No explicit "intercept" term was included in the generation of the response variables, but we follow the discussion in Section 1 and construct posterior distributions based on the integrated likelihood (2) obtained by marginalizing an intercept term from the regression model under a flat prior. Computationally, this means using the mean-centered y^* and X^* data in all expressions reported in this paper.

We illustrate the performance of the MH and RS algorithms under the data augmentation representation of the differentially-scaled prior for β in (11) and (12). The prior on σ^2 is the same in both cases, with $\nu_a = \nu_b = 1$, but we consider two different priors for λ_1 and λ_2 . The first prior has $L = \nu_1 = R = \nu_2 = 1$, which results in a uniform prior for $\lambda_1/(\lambda_1 + \lambda_2)$, the proportion of the total penalty allocated to the $|\beta|_1$ term in the prior. We therefore call this prior the "weak" prior, as it does not strongly favor either term in the penalty function. The second prior has L = 6, $\nu_L = 4$, R = 2, and $\nu_R = 4$. This prior puts a Beta(6, 2) prior on $\lambda_1/(\lambda_1 + \lambda_2)$ and represents a biasing of the prior in favor of stronger ℓ_1 -norm penalization. We call this prior the "strong" prior.

The RS algorithm is implemented as described in Section 3.2.2 under the data augmentation representation. The MH algorithm updates $\log \sigma^2$, $\log \lambda_1$, and $\log \lambda_2$ using random-walk Metropolis updates as described in Hans (2011). Standard deviations for the random-walk innovations, s_{σ^2} , s_{λ_1} , and s_{λ_2} , must be tuned and selected. Poor choices of these parameters can result in poorly-

mixing chains. Experimentation with the simulated data settings revealed that $s_{\sigma^2} = 1$, $s_{\lambda_1} = 1$, and $s_{\lambda_2} = 1$ resulted in generally good performance, and so we used these values in our comparisons.

We compare the efficacy of the MH and RS MCMC algorithms with the exchange algorithm (EX) MCMC sampler of Wang and Wang (2023), which requires tuning only a single Metropolis-Hastings update as part of its sampling scheme. We note that the prior distributions used by Wang and Wang (2023) are different than those used in this paper, and so the posterior distribution sampled by the EX algorithm is different than the posteriors sampled by the RS and MH algorithms (which, for a given prior strength, are identical). The prior for β under the EX algorithm setup is parameterized using a differential scaling, and the penalty parameters are transformed via $(\lambda_1, \lambda_2) \rightarrow (\lambda = \lambda_1 + \sqrt{\lambda_2}, \alpha = \lambda_1/(\lambda_1 + \sqrt{\lambda_2}))$. The prior distributions for λ and α do not contain any tunable parameters and so do not allow for the direct inclusion of prior information about the relative strengths of the two penalty perms. One step in the MCMC algorithm requires specification of s_{α} , the standard deviation of a random-walk proposal for sampling the parameter $\log(\alpha/(1-\alpha))$. Poor choices for this scale parameter can result in poorly-mixing Markov chains. We consider two values of s_{α} in our simulations. The value $s_{\alpha} = 1$ was chosen because it resulted in generally well-performing samplers across the four simulations; we refer to this algorithm as "EX". The value $s_{\alpha} = 0.1$ was chosen because in general it resulted in poor-performing samplers; we refer to this algorithm as "EX-B".

For each of the fifty simulated data sets in each of the five simulation setups, we compute the ESS for parameters β_j , σ^2 , λ_1 , and λ_2 based on 10,000 MCMC iterations of each algorithm after a burn-in of 100 iterations (with starting values chosen so that convergence to stationarity should be very quick). We also compute the ESS for λ and α (the transformed parameters under the EX algorithm) for completeness. We are comparing ESS across six different sampler/posterior setups: the new rejection sampling methods under the weak and strong prior setups (RS-W and RS-S, respectively), the random-walk Metropolis sampler under the weak and strong prior setups (MH-W and MH-S, respectively), and the exchange algorithm sampler under the good and poor step sizes (EX and EX-B, respectively).

To reduce nuisance variability when making comparisons, we treat the simulated data sets as a

blocking factor in the simulation experiment and compute, for each simulated data set, percentage improvement in ESS for MH-W, MH-S, RS-W, RS-S, and EX-B relative to the ESS for the EX algorithm, (ESS_{method} – ESS_{EX})/ESS_{EX} × 100. Figure 4 displays these percentage improvements (or reductions, if negative) for the β_j across the five simulation studies. We discuss primarily the comparison between the RS-* and EX-* methods because the MH-* methods often perform no better on average than the RS-* methods and can require extensive step-size tuning to work well. Focusing first on Simulations 1 and 2 where p = 8, we see that the distribution of percentage improvement in ESS for the RS-* methods is strongly right-skewed when the true $\beta_j \neq 0$. For some simulated data sets the RS-* methods perform worse than EX or EX-B, the reduction in ESS in those cases is small relative to the upside improvement when RS-* performs better. When the true $\beta_j = 0$, $j \in \{3, 4, 6, 7, 8\}$, the distribution of percent improvement for RSS-* is more symmetric around zero, with the EX-* and RS-* methods performing similarly. This is quantified in Table 3, where we see that the average improvement for RS-* is positive relative to EX for all regression coefficients (with larger improvements within a simulation setting when $\beta_j \neq 0$).

In contrast to Simulations 1 and 2, the higher-dimensional (p = 40) setting in Simulation 3 indicates that the EX-* methods both tend to perform better than the RS-* methods. This simulation setting has two blocks of regressors with $\beta_j = 2$ (j = 1, ..., 15 and j = 21, ..., 30), and two blocks of regressors with $\beta_j = 0$. All regressors are equally and moderately correlated via an exchangeable covariance structure. We display ESS improvement for the first two β_j in each of the four blocks. While the EX and EX-B methods perform similarly, the RS-* methods both perform worse than they did in Simulations 1 and 2 relative to EX. While the percent improvement in ESS tends to be negative for the RS-* methods in Simulation 3, we note that the reduction tends to be modest, with most values falling between 0% and -20%. This is quantified in Table 3, where we see that the average ESS percentage reduction tends to be modest (and maxes out at around 20%) under both priors across the simulated data sets.

Finally, Simulation 4 is another higher-dimensional setting (p = 40) but where the regressors are generated using a block-diagonal covariance matrix. ESS results are shown for the first two β_j in each of the four blocks; the true values are non-zero in the first three blocks and zero in the last block. There is strong correlation among the regressors within blocks, but independence across blocks. We see that RS-S tends to performs as well or slightly better than EX for the nonzero coefficients, with RS-W tending to perform slightly worse. All methods perform similarly when $\beta_j = 0$. The average improvement results in Table 3 again indicate that when the RSS-* methods perform worse, the reduction in ESS is not too large.

Figure 5 displays the same information for the other parameters. The distributions of the ESS improvements for the RS-* methods tend to be symmetric or right skewed across the simulations. While in some cases the RS-* methods show a reduction in ESS relative to EX (e.g., σ^2 in Simulations 1 and 2), in others we see a much larger improvement (e.g., λ_1 for RS-S across all simulations). The impact of the poorly-chosen step size for α in the exchange algorithm is most apparent in this figure: the effective sample sizes for α tend to be much worse for EX-B than for EX. In terms of average percent improvement, Table 3 shows that RS-S does quite well for σ^2 , λ_1 , and λ_2 parameters in Simulations 1 and 2. Despite performing reasonably well for the the regression coefficients, β_j , RS-W does not do particularly well in any of the settings for σ^2 , λ_1 , and λ_2 .

		Prior	β_1	β_2	β_3	β_4	β_5	β_6	β_7	β_8
Simulation 1	Weak	42.85	19.98	3.19	2.24	5.03	1.58	1.57	2.79	
	Strong	59.73	22.63	8.64	2.69	18.06	7.26	4.92	2.48	
Simulation 2	ion 2	Weak	7.61	15.23	25.92	9.98	10.32	8.41	15.45	11.85
	Strong	5.87	16.12	28.74	12.26	11.77	7.15	17.17	17.56	
			Prior	σ^2)	λ_1	λ_2	λ	α	
Simulation		lation 1	Weak	-14.12	-34.1	5 -28	.88 3	6.99	-60.1	
			Strong	21.79	149.8	6 1	1.9 21	3.22	49	
Simulation 2		Weak	-21.40	-43.8	6 -39	.82	2.62	-65.76		
			Strong	19.83	166.7	5 18	.39 35	3.03	38.27	
Simulation 3		Weak	-58.98	-25.8	6 -76	.51 -2	1.78	-71.12		
		liation 5	Strong	-40.95	90.4	2 -53	.17 5	3.84	-31.35	
Cim	Simu	lation 1	Weak	-56.86	-33.7	2 -69	.57 -4	7.39	-65.9	
Sint		1au011 4	Strong	-42.93	58.4	7 -39	.56 -	4.85	-31.16	
	-									

Table 3: Average percent improvement in ESS across fifty simulated data sets for the MCMC algorithm using rejection sampling (RS) versus the exchange algorithm (EX) with a well-chosen proposal standard deviation. Positive numbers in bold indicate better average performance for RS, e.g., 36.77 indicates the ESS for RS was 36.77% larger than it was for EX.



Sampler 🖨 EX-B 🛱 MH-S 🛱 MH-W 🛱 RS-S 🛱 RS-W

Figure 4: Percent improvement in ESS for fifty simulated data sets under four simulation settings for several MCMC algorithms. The exchange algorithm (EX) with a well-chosen proposal standard deviation is the baseline. ESSes for all eight β_j are shown for Simulations 1 and 2; a selection of relevant β_j are shown for Simulations 3 and 4. EX-B: exchange algorithm with a poorly-chosen step size; MH-*: random-walk Metropolis updates for σ^2 , λ_1 , and λ_2 under Strong and Weak priors; RS-*: the new rejection sampling methods under the two prior settings.



Sampler 🖨 EX-B 🖨 MH-S 🖨 MH-W 🖨 RS-S 🖨 RS-W

Figure 5: Percent improvement in ESS for fifty simulated data sets under four simulation settings for several MCMC algorithms. The exchange algorithm (EX) with a well-chosen proposal standard deviation is the baseline. The parameters $\lambda = \lambda_1 + \sqrt{\lambda_2}$ and $\alpha = \lambda_1/(\lambda_1 + \sqrt{\lambda_2})$ are a core part of the exchange algorithm sampler. EX-B: exchange algorithm with a poorly-chosen step size; MH-*: random-walk Metropolis updates for σ^2 , λ_1 , and λ_2 under Strong and Weak priors; RS-*: the new rejection sampling methods under the two prior settings.

		1 electricities of the forty p_1 with average 70 improvement in 1000 .					
		$\geq 0\%$	$\geq -5\%$	$\geq -10\%$	$\geq -20\%$		
Simulation 3	Weak Prior	0	0	15	100		
	Strong Prior	0	2.5	97.5	100		
Simulation 4	Weak Prior	10	92.5	100	100		
	Strong Prior	37.5	100	100	100		

Percentage of the forty β_i with average % improvement in ESS:

Table 4: Percentage of the number of β_j (out of 40) for which the average percent improvement (or reduction, when negative) in ESS for RS-* relative to EX is greater than or equal to various thresholds,

5 Discussion

Full Bayesian inference for the Bayesian elastic net regression model is challenging due to the $\Phi(\cdot)$ term in the normalizing constant for the prior on β that is a function of λ_1 , λ_2 , and sometimes σ^2 (depending on the parameterization of the model). All existing, correctly-specified methods for posterior sampling use at least one Metropolis–Hastings update that requires specification and tuning of a proposal distribution. We have introduced transformations for the commonly- and differentially-scaled priors (under both direct and DA representations) that result in "well known" full conditional distributions that can be easily sampled from for all but one parameter, θ . Careful analysis of the full conditional for θ reveals that rejection sampling approaches that take advantage of log-concavity of the target density function can be used to efficiently produce samples directly from the full conditional. A key to this approach is that the rejection sampling methods are automatic in the sense that no tuning is required. Access to MCMC algorithms that practitioners can run directly without having to interactively manipulate will make these statistical models more broadly impactful.

We compared our new sampling methods with the existing, exchange algorithm-based approach of Wang and Wang (2023). The exchange algorithm sampler is cleverly designed to avoid computation of $\Phi(\cdot)$ while at the same time minimizing the number of tuning parameters, and it worked quite well across the simulation settings we considered. Despite its strong performance, there are several reasons why one might prefer the new rejection sampling approaches. First, the posterior distribution in Wang and Wang (2023) from which the exchange algorithm is sampling has no user-specified parameters in the prior. While this might be useful from a particular objective

Bayes point of view where a goal is to provide users with methods that can be used reliably without much user input, having the ability to shift the prior toward stronger ℓ_1 - or ℓ_2 -norm penalization gives the user the ability to move beyond default prior settings to build a model that is a better match for their analysis. Second, while the exchange algorithm was not too sensitive to the choice of the random-walk scale parameter, s_{α} , particularly poor choices reduced the ESSes for some of the parameters. Avoiding the need to specify such a parameter is desirable.

We also note that the higher effective sample sizes for the parameters σ^2 , λ , and α under the exchange algorithm-based MCMC sampler might be due to other sampling techniques used by Wang and Wang (2023) to improve the Markov chain dynamics. Wang and Wang (2023) use a partially-collapsed (Liu, 1994; van Dyk and Park, 2008) update for σ^2 and implement a generalized Gibbs step (Liu and Sabatti, 2000; Liu, 2004) for σ^2 , λ , and α that is based on a scale transformation group to accelerate convergence of the chain. Incorporating similar techniques into the rejection samplers introduced in this paper may produce similar improvements and is an area of future work.

Finally, we note that the performance of MCMC algorithms can be very sensitive to parameterization The (re)parameterization introduced in this paper was chosen primarily to produce full conditional distributions with log-concave densities that could be sampled easily via automatic rejection sampling techniques. Other transformations might be available that yield similar computational simplicity while also further improving the Markov chain dynamics and will be the subject of future research.

A Rejection sampling σ^2 with an inverse gamma proposal

Li and Lin (2010) propose using rejection sampling to sample from the full conditional for σ^2 in their formulation of the Bayesian elastic net. They define the function $f(\sigma^2) \propto \pi(\sigma^2 | y, \beta, \tau^2, \lambda_1, \lambda_2)$ (see equation (6) in Li and Lin, 2010) to be

$$f(\sigma^2) = (\sigma^2)^{-a-1} \left\{ \Gamma_U\left(\frac{1}{2}, \frac{\lambda_1^2}{8\sigma^2\lambda_2}\right) \right\}^{-p} e^{-b/\sigma^2},$$
(23)

where a = n/2 + p and b > 0 depends on y and the other parameters in the model. We note that

the full conditional in Li and Lin (2010) is similar to but slightly different than the full conditional in (17) due to small differences in model specification and parameterization.

Li and Lin (2010) propose rejection sampling from $f(\sigma)$ using an inverse gamma proposal distribution with density function $h(\sigma^2) = b^a \Gamma(a)^{-1} (\sigma^2)^{-a-1} e^{-b/\sigma^2}$, where a and b are the same as in (23). Li and Lin (2010) describe the following algorithm for rejection sampling using this proposal distribution: "generate a candidate Z from h and a u from uniform(0,1), and then accept Z if $u \leq \Gamma(\frac{1}{2})^p b^a f(Z)/\Gamma(a)h(Z)$ or, equivalently, if $\log(u) \leq p \log(\Gamma(\frac{1}{2})) - p \log(\Gamma_U(\frac{1}{2}, \frac{\lambda_1^2}{8Z\lambda_2}))$." By the definition of the upper incomplete gamma function, we have

$$\Gamma_U\left(\frac{1}{2}, \frac{\lambda_1^2}{8Z\lambda_2}\right) = \int_{\lambda_1^2/(8Z\lambda_2)}^{\infty} t^{-1/2} e^{-t} dt$$

$$\leq \int_0^{\infty} t^{-1/2} e^{-t} dt$$

$$= \Gamma\left(\frac{1}{2}\right),$$
(24)

which means that $p \log \left(\Gamma\left(\frac{1}{2}\right)\right) - p \log \left(\Gamma_U\left(\frac{1}{2}, \frac{\lambda_1^2}{8Z\lambda_2}\right)\right) \ge 0$ with probability one. We also have that $\log(u) \le 0$ with probability one, and so the rejection sampling algorithm will accept every proposal with probability one, which can only be true if $f(\sigma^2) \propto h(\sigma^2)$, which is not the case, and the algorithm does not produce samples from $f(\sigma^2)$. The contradiction appears to arise from a reversal of the bound in (24) in one step of the construction of the algorithm in Li and Lin (2010). In fact, an inverse gamma distribution with parameters a and b that match those in (23) cannot be used as a proposal distribution for rejection sampling. The ratio of the target to the proposal densities is $f(\sigma^2)/h(\sigma^2) = b^{-a}\Gamma(a) \left\{\Gamma_U\left(\frac{1}{2}, \frac{\lambda_1^2}{8\sigma^2\lambda}\right)\right\}^{-p}$. We see that $\lim_{\sigma^2 \to 0} f(\sigma^2)/h(\sigma^2) = \infty$ and rejection sampling cannot be implemented using this proposal distribution.

References

Alhamzawi, R. and Ali, H. T. M. (2018). The Bayesian adaptive lasso regression. *Mathematical Biosciences* 303, 75–82.

Alhamzawi, R., Yu, K., and Benoit, D. F. (2012). Bayesian adaptive lasso quantile regression.

Statistical Modelling **12**, 3, 279–297.

- Andrews, D. and Mallows, C. (1974). Scale mixtures of normal distributions. *Journal of the Royal Statistical Society*, Series B **36**, 99–102.
- Barndorff-Nielsen, O. (1977). Exponentially decreasing distributions for the logarithm of particle size. *Proceedings of the Royal Society of London* **A353**, 401–419.
- Barndorff-Nielsen, O. (1978). Hyperbolic distributions and distributions on hyperbolae. *Scandinavian Journal of Statistics* **5**, 151–157.
- Devroye, L. (1984). A simple algorithm for generating random variates with a log-concave density. *Computing* **33**, 247–257.
- Devroye, L. (1986). Non-uniform Random Variate Generation. Springer, New York, 1st edn.
- Devroye, L. (2014). Random variate generation for the generalized inverse Gaussian distribution. *Statistics and Computing* **24**, 239–246.
- DiDonato, A. R. and Morris, A. H. (1986). Computation of the incomplete gamma function ratios and their inverse. *ACM Transactions on Mathematical Software (TOMS)* **12**, 4, 377–393.
- Feller, W. (1968). An Introduction to Probability Theory and Its Applications, vol. 1. Wiley, New York, 3rd edn.
- Flegal, J. M., Hughes, J., Vats, D., Dai, N., Gupta, K., and Maji, U. (2021). mcmcse: Monte Carlo Standard Errors for MCMC. Riverside, CA, and Kanpur, India. R package version 1.5-0.
- Friedman, J., Hastie, T., and Tibshirani, R. (2008). Sparse inverse covariance estimation with the graphical lasso. *Biostatistics* **9**, 3, 432–441.
- Genz, A. (1992). Numerical computation of multivariate normal probabilities. *Journal of Computational and Graphical Statistics* **1**, 141–150.
- Genz, A. and Bretz, F. (2009). *Computation of Multivariate Normal and t Probabilities*. Lecture Notes in Statistics. Springer-Verlag, Heidelberg.

- Genz, A., Bretz, F., Miwa, T., Mi, X., Leisch, F., Scheipl, F., Bornkamp, B., Maechler, M., and Hothorn, T. (2024). *mvtnorm: Multivariate Normal and t Distributions*. R package version 1.2-5.
- Geweke, J. (1991). Efficient simulation from the multivariate normal and student-t distributions subject to linear constraints. In *Computer Science and Statistics: Proceedings of the 23rd Symposium on the Interface*, 571–578. American Statistical Association, Alexandria, VA.
- Gilks, W. R. (1992). Derivative-free adaptive rejection sampling for gibbs sampling. In *Bayesian Statistics 4* Editors: J.M. Bernardo, J.O. Berger, A.P. Dawid and A.F.M. Smith, 641–649. Oxford Univ. Press.
- Gilks, W. R. and Wild, P. (1992). Adaptive rejection sampling for Gibbs sampling. *Applied Statistics* **41**, 337–348.
- Gong, L. and Flegal, J. (2015). A practical sequential stopping rule for high-dimensional Markov chain Monte Carlo. *Journal of Computational and Graphical Statistics* **25**, 684–700.
- Gordon, R. D. (1941). Values of Mills' ratio of area to bounding ordinate and of the Normal probability integral for large values of the argument. *The Annals of Mathematical Statistics* **12**, 364–366.
- Gordy, M. B. (1998). A generalization of generalized beta distributions. Tech. rep., Division of Research Statistics, Division of Monetary Affairs, Federal Reserve Board.
- Griffin, J. E. and Brown, P. J. (2007). Bayesian adaptive lassos with non-convex penalization. Working paper 07-02, CRiSM.
- Griffin, J. E. and Brown, P. J. (2011). Bayesian hyper-lassos with non-convex penalization. *Australian & New Zealand Journal of Statistics* **53**, 4, 423–442.
- Hans, C. (2009). Bayesian lasso regression. Biometrika 96, 835-845.
- Hans, C. (2010). Model uncertainty and variable selection in Bayesian lasso regression. *Statistics and Computing* **20**, 221–229.

- Hans, C. (2011). Elastic net regression modeling with the orthant normal prior. *Journal of the American Statistical Association* **106**, 1383–1393.
- Hoerl, A. and Kennard, R. (1970). Ridge regression: Biased estimation for nonorthogonal problems. *Technometrics* 12, 55–67.

Jeffreys, H. (1961). Theory of Probability. Oxford University Press, London, 3rd edn.

- Kang, K., Song, X., Hu, X. J., and Zhu, H. (2019). Bayesian adaptive group lasso with semiparametric hidden Markov models. *Statistics in Medicine* 38, 9, 1634–1650.
- Kyung, M., Gill, J., Ghosh, M., and Casella, G. (2010). Penalized regressions, standard errors, and Bayesian lassos. *Bayesian Analysis* 5, 369–412.
- Lee, K. H., Chakraborty, S., and Sun, J. (2015). Survival prediction and variable selection with simultaneous shrinkage and grouping priors. *Statistical Analysis and Data Mining: The ASA Data Science Journal* 8, 114–127.
- Leng, C., Tran, M.-N., and Nott, D. (2014). Bayesian adaptive lasso. *Annals of the Institute of Statistical Mathematics* **66**, 221–244.
- Li, Q. and Lin, N. (2010). The Bayesian elastic net. Bayesian Analysis 5, 151-170.
- Liu, J. S. (1994). The collapsed Gibbs sampler in Bayesian computations with applications to a gene regulation problem. *Journal of the American Statistical Association* **89**, 958–966.
- Liu, J. S. (2004). General conditional sampling. In *Monte Carlo Strategies in Scientific Computing*, 161–181. Springer.
- Liu, J. S. and Sabatti, C. (2000). Generalised Gibbs sampler and multigrid Monte Carlo for Bayesian computation. *Biometrika* 87, 353–369.
- Meinshausen, N. and Bühlmann, P. (2006). High-dimensional graphs and variable selection with the lasso. *The Annals of Statistics* **34**, 3, 1436–1462.

- Murray, I., Ghahramani, Z., and MacKay, D. J. C. (2006). MCMC for doubly-intractable distributions. In *Proceedings of the 22nd annual conference on uncertainty in artificial intelligence* (*UAI*), 359–366. AUAI Press.
- Park, T. and Casella, G. (2008). The Bayesian Lasso. *Journal of the American Statistical Association* **103**, 681–686.
- Perez Rodriguez, P. (2024). ars: Adaptive Rejection Sampling. R package version 0.8.
- R Core Team (2024). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Raiffa, H. and Schlaifer, R. (1961). Applied Statistical Decision Theory. Harvard University, Boston.
- Ročková, V. and George, E. I. (2018). The spike-and-slab LASSO. *Journal of the American Statistical Association* **113**, 521, 431–444.
- Roy, V. and Chakraborty, S. (2017). Selection of tuning parameters, solution paths and standard errors for Bayesian lassos. *Bayesian Analysis* **12**, 753–778.
- Sun, J. (2020). Modified-half-normal distribution and different methods to estimate average treatment effect. Ph.D. thesis, University of Louisville.
- Sun, J., Kong, M., and Pal, S. (2023). The Modified-Half-Normal distribution: Properties and an efficient sampling scheme. *Communications in Statistics Theory and Methods* **52**, 1591–1613.
- Tibshirani, R. (1996). Regression shrinkage and selection via the Lasso. *Journal of the Royal Statistical Society*, Series B **58**, 267–288.
- Tibshirani, R., Saunders, M., Rosset, S., Zhu, J., and Knight, K. (2005). Sparsity and smoothness via the fused lasso. *Journal of the Royal Statistical Society Series B: Statistical Methodology* 67, 1, 91–108.
- van Dyk, D. and Park, T. (2008). Partially collapsed Gibbs samplers: Theory and methods. *Journal of the American Statistical Association* **103**, 790–796.

- Wang, C., Li, Q., Song, X., and Dong, X. (2019). Bayesian adaptive lasso for additive hazard regression with current status data. *Statistics in Medicine* **38**, 20, 3703–3718.
- Wang, H. (2012). Bayesian graphical lasso models and efficient posterior computation. *Bayesian Analysis* 7, 4, 867–886.
- Wang, H.-B. and Wang, J. (2023). An exact sampler for fully Baysian elastic net. *Computational Statistics* **38**, 1721–1734.
- West, M. (1987). On scale mixtures of normal distributions. Biometrika 74, 646-648.
- Yuan, M. and Lin, Y. (2006). Model selection and estimation in regression with grouped variables. *Journal of the Royal Statistical Society Series B: Statistical Methodology* 68, 1, 49–67.
- Yuan, M. and Lin, Y. (2007). Model selection and estimation in the Gaussian graphical model. *Biometrika* 94, 1, 19–35.
- Zou, H. (2006). The adaptive lasso and its oracle properties. *Journal of the American Statistical Association* **101**, 476, 1418–1429.
- Zou, H. and Hastie, T. (2005). Regularization and variable selection via the elastic net. *Journal of the Royal Statistical Society*, B **67**, 301–320.