

# Robustness of quantum many-body scars in the presence of Markovian bath

Xiang-Ping Jiang,<sup>1</sup> Mingdi Xu,<sup>2</sup> Xuanpu Yang,<sup>2</sup> Yucheng Wang,<sup>3,4,5</sup> and Lei Pan<sup>2,\*</sup>

<sup>1</sup>Zhejiang Lab, Hangzhou 311121, China

<sup>2</sup>School of Physics, Nankai University, Tianjin 300071, China

<sup>3</sup>Shenzhen Institute for Quantum Science and Engineering,  
Southern University of Science and Technology, Shenzhen 518055, China

<sup>4</sup>International Quantum Academy, Shenzhen 518048, China

<sup>5</sup>Guangdong Provincial Key Laboratory of Quantum Science and Engineering,  
Southern University of Science and Technology, Shenzhen 518055, China

A generic closed quantum many-body system will inevitably tend to thermalization, whose local information encoded in the initial state eventually scrambles into the full space, known as quantum ergodicity. A paradigmatic exception in closed quantum systems for strong ergodicity breaking is known as many-body localization, where strong disorder-induced localization prevents the occurrence of thermalization. It is generally recognized that a localized quantum system would be delocalized under dissipation induced by the environment. However, this consequence recently has received challenges where an exotic dissipation-induced localization mechanism is proposed, and transitions between localized and extended phases are found. In this Letter, we promote this mechanism to systems for weak ergodicity breaking hosting quantum many-body scars (QMBS). We find that the system relaxes to a steady state dominated by QMBS, and the dissipative dynamics exhibit dynamic revivals by suitably preparing an initial state. We point out an experimental realization of the controlled dissipation with a cold atomic setup. This makes the signature of ergodicity breaking visible over dissipative dynamics and offers potential possibilities for experimentally preparing stable QMBS with associated coherent dynamics.

**Introduction.** Within the framework of statistical mechanics, generic thermodynamic closed systems relax to thermal equilibrium independent of initial conditions [1, 2]. The thermalization of closed quantum systems can be interpreted via the eigenstate thermalization hypothesis (ETH) [10–12] which states that the eigenstate itself is thermalized. However, there exist counter examples violating above paradigm such as disordered systems. A prototype is many-body localization (MBL) [3, 4] which is a consequence of the balance between many-body interactions and disorder-induced localization. The MBL emerges as a paradigm for strong ergodicity breaking [5–9] and the corresponding quench dynamics can persist to long lasting times, providing an alternative to distinguish those systems with thermalizing dynamics. However realistic systems cannot be fully isolated from the external environment which naturally elicits question that if the MBL immunizes the influence of environment on large timescales.

Intuitively, any localized system will be inevitably thermalized when it couples a thermal environment that the local information of initial state scrambles in the whole space. Previous theoretical studies and experimental measurements about dissipative dynamics of the MBL in presence of environment showed that dissipation spoils the MBL and induces ergodicity eventually with a slow relaxation dynamics [13–15]. This is physically evident that any local dephasing will drive the system into maximally mixed state (infinite temperature state) independent on choice of the system Hamiltonian despite a MBL exhibits the stretched exponential relaxation signature differ from an ergodic system. However, above

analysis is valid for cases that dissipative operators are hermitian or local (on-site) operators. Recently, several works re-investigated the fate of localization properties in disordered systems under a special type of *quasi-local* dissipation operators, and found that the system can be driven into new steady states exhibiting signatures of localization [16–21] and a dissipation induced localized-extended transition was proposed [18]. This provides a feasible route towards preparing non-thermal quantum states violating the ETH via engineered dissipation.

In the last few years, a new type of prototype violating the ETH (weak ergodicity breaking) was discovered in the Rydberg atom arrays where persistent revivals occur for some special initial states while for most initial states the system relaxes rapidly [22, 23]. The behavior of long-time coherent dynamics originates from quantum many-body scarring mechanism [24, 25] where a small set of eigenstates dubbed quantum many-body scars (OMBS) violating the ETH. Unlike strong ergodicity breaking such as the MBL where entire spectrum violates the ETH, the number of QMBS is typically a measure-zero set which is exponentially smaller than the whole Hilbert space dimension in the thermodynamic limit. This could make one expect that QMBS do not show an experimentally observable dynamic feature since they are buried in the energy spectrum. However, those systems containing QMBS as equally spaced towers in the spectrum can result in dynamical revival from special initial states, that was observed in Rydberg atom experiments. Although there have been extensive research on QMBS in closed systems [26–28], the study of QMBS in the framework of open quantum systems and

non-hermitian physics is still in its infancy [29–32].

In this Letter, we study the impact of dissipation on systems with QMBS and explore whether dissipation can drive a system to steady states exhibiting structures of QMBS instead of destroying them and then ascertain if the persistent revivals can be witnessed in corresponding dissipative dynamics for particular initial states. We demonstrate that by utilizing a set of quasi-local dissipation operators acting on a pair of neighboring sites, systems can be brought into a specific steady state consisting mainly of QMBS. We confirm the above finding through three representative models containing QMBS, i.e., the PXP model with approximate QMBS, the toy model with exact QMBS embedding within nonintegrable models [38], and the generalized  $J_1 - J_2$  model where QMBS can be analytically solved by spectrum generating algebra [39–43]. Moreover, we provide an experimental proposal to realize such quasi-local dissipation operators and observe related dissipative scarring dynamics based on current techniques about Rydberg atomic simulators.

*Dissipative PXP model.* A well-known system containing QMBS is the PXP model serving as an effective model [25] for Rydberg atoms array, which plays a significant role and the corresponding Hamiltonian in one dimension is

$$H_{\text{PXP}} = \sum_j P_j \sigma_{j+1}^x P_{j+2}, \quad (1)$$

where  $P_j = |0_j\rangle\langle 0_j| = \frac{1-\sigma_j^z}{2}$  is the projector onto  $|0_j\rangle$  state denoting no Rydberg excitation at the site  $j$  where the basis on each site consists of  $|n_j\rangle$  with Rydberg excitation number  $n_j = 0, 1$  in which case the Pauli matrix is represented as  $\sigma_j^x = |0_j\rangle\langle 1_j| + |1_j\rangle\langle 0_j|$  ( $\sigma_j^z = |1_j\rangle\langle 1_j| - |0_j\rangle\langle 0_j|$ ) represents the Pauli matrix in  $x$  ( $z$ ) direction. The PXP Hamiltonian (1) has the constraint stemming from the Rydberg blockade effect [44–47] that each spin can freely rotate only if both its neighbors are no Rydberg excitation  $|0_j\rangle$ , while it is frozen otherwise. Despite its simplicity, the PXP Hamiltonian is nonintegrable whose level statistics approaches the Wigner-Dyson distribution in thermodynamic limit. However, there exists a subset of nearly equidistant special eigenstates (QMBS) in the spectrum that forms a band as tower structure extending the whole spectrum. As shown in Fig.1(a), one can see the visible separation from the bulk of spectrum. The QMBS in PXP model are responsible to the periodic revivals on quench dynamics whose period is approximately inversely proportional to energy separation between QMBS.

The QMBS are generally fragile to perturbation [48] or disorder [49]. It is naturally expected that QMBS are also unstable against to the interaction between the system and external environments. Suppose the Rydberg

atom chain described by the PXP model is coupled to a Markovian bath, whose dynamics is described by the following Lindblad master equation [33, 34]

$$\frac{d\rho(t)}{dt} = \mathcal{L}[\rho(t)] = -i[H_S, \rho(t)] + \mathcal{D}[\rho(t)] \quad (2)$$

where  $\rho(t)$  is the density matrix of the system,  $H_S$  denotes the system Hamiltonian and  $\mathcal{L}$  is referred to as the Liouvillian superoperator. The first part in right hand side of the Lindblad equation (2) is the unitary evolution and the second part represents dissipative evolution, which is written by

$$\mathcal{D}[\rho(t)] = \sum_j \gamma_j \left( 2O_j \rho O_j^\dagger - \{O_j^\dagger O_j, \rho\} \right). \quad (3)$$

Here  $\{, \}$  denotes the anticommutator, and  $O_j$  is dissipation operator or jump operator at the site  $j$  with dissipation strength  $\gamma_j$ . As with the system Hamiltonian determining time evolution of a closed system, the dissipative dynamics of open quantum system is governed by Liouvillian superoperator. The solution is formally expressed as  $\rho(t) = e^{\mathcal{L}t} \rho_0$  and the system reaches its steady state  $\rho_{ss} = \lim_{t \rightarrow \infty} \rho(t) = \lim_{t \rightarrow \infty} e^{\mathcal{L}t} \rho_0$  for arbitrary initial state  $\rho_0 = \rho(0)$ . The steady state is closely relevant to the choice of dissipation operators  $O_j$ , and a particular concern is to design the dissipation operators such that the steady state serves as the required quantum state. The central thesis of this paper is that making QMBS as steady state by using specific dissipation operators as simple as possible.

Here we consider the following type of jump operators

$$O_j = (\sigma_j^+ + e^{i\theta} \sigma_{j+1}^+) (\sigma_j^- - e^{i\theta} \sigma_{j+1}^-), \quad (4)$$

These operators were first introduced in Refs.[35, 36] to realize topological phases as steady with engineered dissipation. The physical implementation can be based on a Bose-Hubbard model in superconducting resonators array [37], and also be proposed on optical Raman lattices [18]. Physically, above jump operator acts on a pair of nearest-neighbor sites and changes the relative phase between them but does not alter particle number. Moreover, it not only includes the local on-site dephasing term  $n_j = \sigma_j^+ \sigma_j^-$  but also the (quasi-local) hopping term  $e^{i\theta} \sigma_j^+ \sigma_{j+1}^- - e^{i\theta} \sigma_{j+1}^+ \sigma_j^-$  which is crucial to our objectives. Now we consider such dissipation operators (4) act on the PXP model, and the corresponding steady state can be obtained from the zero-mode of Liouvillian superoperator. In Fig.1(b), we plot the distribution of the steady state on eigenstates of PXP Hamiltonian, where a structure is clearly visible that the QMBS obtain larger weight than those states in the vicinity of energy. One of physical significances of QMBS in PXP model is the high overlap with product states ( $|Z_2\rangle$  and  $|Z_2'\rangle$ ),

which can be prepared in Rydberg atom experiments. Hence, to discover the intrinsic structure of QMBS, we calculate the distribution of the steady state density matrix  $\rho_{ss}$  on local basis as displayed in Fig.2(a). One can see the steady state has anomalously high overlaps with two product states  $|Z_2\rangle = |101010 \cdots 101010\rangle$  and  $|Z'_2\rangle = |010101 \cdots 010101\rangle$ .




FIG. 1. (a) The expectation value of mean magnetization  $\langle \sigma^z \rangle = \sum_j \langle \sigma_j^z \rangle / L$  in the PXP model as a function of eigenenergy  $E$ . Most of points are concentrated in the thermal bulk, but there are outermost eigenstates (indicated by crosses) as QMBS violating the ETH. (b) The distribution of steady-state density matrix on eigenstates of the PXP model as function of energy  $E$  corresponding with (a). It is clearly visible that QMBS have larger weights than thermal states in the vicinity of energy. Here the system size is chosen as  $L = 16$ , and the dissipative phase and strength is  $\theta = \pi$  and  $\gamma = 0.1$  respectively.

We further examine this feature from dynamic perspective by investigating the quantum fidelity, i.e., overlap between the density matrix at any time  $\rho(t)$  and the initial time  $\rho_0$  defined as follows [50]

$$F[\rho(t), \rho_0] = \text{Tr} \left[ \sqrt{\sqrt{\rho(t)} \rho_0 \sqrt{\rho(t)}} \right]. \quad (5)$$




FIG. 2. (a) The distribution of steady-state density matrix on local basis of the PXP model where two states ( $|Z_2\rangle$  and  $|Z'_2\rangle$ ) have markedly large values. (b) Time evolutions of the quantum fidelity for different initial product states. The system rapidly relaxes to negligible steady-state value (red dashed line) from the initial state  $|g\rangle = |000 \cdots 000\rangle$  without Rydberg excitations. In contrast, the fidelity shows periodic oscillations before reach a non-zero steady-state value, and the period is  $T = 2\pi/\Delta E$  where  $\Delta E$  is the mean energy spacing of QMBS near the middle of spectrum. The system size and parameters in dissipation are same as the Fig.1

Fig.2 (b) plots the time evolution of quantum fidelity for two different initial states which shows that the fidelity approaches zero rapidly and dynamical revival complete absence for the initial state without Rydberg excitations  $|g\rangle = |000 \cdots 000\rangle$ . In contrast, a remarkable periodic oscillation in time  $|Z_2\rangle$  with period  $T = 2\pi/\Delta E$  (see the inset of Fig.2(b)) where  $\Delta E \approx 1.33$  is the energy separation between adjacent QMBS near zero-energy. Moreover, the long-time steady value is non-zero reflect-

ing the structure in steady-state distribution on local basis as shown in Fig.2(a). It is worth pointing out that the conclusion above is far from trivial. This is because that jump operators we required can be principally constructed from non-Hermitian ones that make the steady state in the form  $\rho_{ss} = |\phi_n\rangle\langle\phi_n|$  for any eigenstates  $|\phi_n\rangle$  of the system Hamiltonian. This kind of steady state is nothing but the dark state, i.e.,  $O_j|\phi_n\rangle = 0$  for any  $j$ . However, this generally require the condition that the jump operator is non-local which is usually unrealistic. More critically, to guarantee the above dark state as an eigenstate, one has to require the priori information of the eigenstate  $|\phi_n\rangle$  which could be unfeasible. Thus, seeking out a local or quasi-local operator acting only a few neighboring sites is more realistic and desirable that is our important guideline.

*Dissipative toy model.* In addition to the PXP model studied above, our protocol can also apply to those systems with exactly solvable QMBS. To illustrate this, we use the toy model which hosts exact QMBS as a concrete example whose Hamiltonian is written as [38]

$$H_{\text{toy}} = \frac{\Omega}{2} \sum_j \sigma_j^x + \sum_j V_{j-1,j+2} P_{j,j+1}, \quad (6)$$

where  $P_{j,j+1} = (1 - \vec{\sigma}_j \cdot \vec{\sigma}_{j+1})/4$  represent the local projector on spin singlet of a neighboring pair and  $V_{ij}$  denotes an interaction operator acting on two spins  $(i, j)$ , which is generically expressed as  $V_{i,j} = \sum_{\alpha\beta} J_{ij}^{\alpha\beta} \sigma_i^\alpha \sigma_j^\beta$  with arbitrary coupling coefficients  $J_{ij}^{\alpha\beta}$  ( $\alpha\beta = x, y, z$ ). This toy model is non-integrable due to the randomness of interaction coefficients  $J_{ij}$ . We can also find that the Hamiltonian (6) does not commute with the local projector  $P_{j,j+1}$  and total magnetism  $S^x = \sum_j \sigma_j^x/2$  and thereby there is no any apparent local symmetries. However, one can verify that states in the subspace  $|\mathcal{S}_n\rangle = |S = L/2, m_x = (n - L/2 - 1)\rangle$  ( $n = 1, 2, \dots, L+1$ ) with the total number of sites  $L$ , are eigenstates whose energies are exactly equidistant  $E = \Omega m_x$ . In contrast, other states outside of the subspace are affected by interaction terms  $V_{j-1,j+2}$  which hybridize each other to form thermal eigenstates. The toy model belongs to embedding formalism [51] where the QMBS are embedded into a generic non-integrable model. We will see these exactly solvable states serve as QMBS and can also become steady states of a Liouvillian. We first numerically compute the bipartite entanglement entropy for all eigenstates of the toy model as shown in Fig.3(a) where some states with anomalously low entanglement and equally spaced energies are clearly visible while other states have large entanglement entropy. To investigate properties of QMBS under dissipation, we calculate the Liouvillian spectrum under the dissipation operators (3) at  $\theta = 0$ , where the zero-energy state (steady-state) are exactly composed of QMBS, i.e.,  $\rho_{ss}^n = |\mathcal{S}_n\rangle\langle\mathcal{S}_n|$ . A more striking

consequence is that there exists a string of states at equal spacing on the imaginary axis which are non-decay modes because real parts of them are zero. In fact, all non-decay modes originate from degenerate steady-state subspace where we can build up a series of independent non-decay modes of the Liouvillian superoperator by means of QMBS as  $|\mathcal{S}_m\rangle\langle\mathcal{S}_n|$  whose eigenvalue is  $i(E_n - E_m) = i\Omega(n - m)$ . The degenerate steady states and the associated equally spaced imaginary parts can contribute to nontrivial dissipative dynamics with perfect revival without decay as shown in the Fig.3(c). To sum up, by using non-local dissipators, a system can be driven to the steady state mainly composed of QMBS for either PXP model or the toy model with exactly solvable QMBS. Moreover, our approach can also apply to those systems where QMBS are produced by the spectrum generating algebra (see Supplementary Material).

*Experimental realization.* As we pointed out that the local or quasi-local operators are much easier to implement than those non-local ones. Here we provide an experimental scheme to concretely realize our dissipation operators (3). First, we produce two lattice systems, one of which is system lattice and another is auxiliary lattice as illustrated in Fig.4. Then coherently couples two neighbouring sites of the system to an auxiliary site between them with anti-symmetric couplings  $\Omega$  and  $-\Omega$  by spatial modulation of the Raman laser [35]. In this way, the annihilation part of dissipation operators (3) is realized. Second, after the annihilation of anti-symmetric superposition of atoms (or spins) in the system lattice, the creation process sequentially is realized by spontaneous radiation where the atom decays back to the system sites from auxiliary lattice. This creation process is symmetric since the spontaneous emission is isotropic. In this way, one can realize the jump operator of the form implement. An arbitrary phase  $\theta$  can be implemented via resonators array coupled by superconductive qubits [16, 37]. Further, we propose a quench dynamics scheme to measure the scarred nature of the steady state. Suppose the system has relaxed to the QMBS dominated steady state under the action of quasi-local dissipation operators, then we remove dissipation and thereby the system evolves under its Hamiltonian. where diagonal elements  $\rho_{nn}$  of the density matrix in eigenbasis  $\{|\phi_n\rangle\}$  remains unchanged and thereby the QMBS with associated scarred quench dynamics can be detected after removing dissipation.




FIG. 3. (a) Bipartite entanglement entropy of eigenstates for the toy model as a function of energy. Except for thermal states in the bulk which have large entropies, a group of peculiar eigenstates with low entropies which are nothing but QMBS. (b) Spectrum of the Liouvillian superoperator for the dissipative phase  $\theta = 0$  and dissipation strength  $\gamma = 0.1$ . Here we just show eigenvalues with 200 smallest and 200 largest real parts. (c) Time evolution of quantum fidelity from the initial state  $\rho_0 = |\phi_0\rangle\langle\phi_0|$  with  $|\phi_0\rangle = |\uparrow\uparrow\uparrow \dots \uparrow\uparrow\rangle$  which shows a perfect revival with period  $2\pi/\Omega$ . Here we choose the system parameter  $\Omega = 1$ , and coupling coefficients  $J_{j-1,j+2}$  are chosen by independent normal distribution. The system size is  $L = 10$  with periodic boundary condition.




FIG. 4. Schematic diagram for realizing dissipation operators (3) at  $\theta = 0$ . The upper (lower) lattice represents the auxiliary (system) lattice. A spatial modulation of the Raman laser yields a position-dependent coupling, which excites only the antisymmetric superposition of atoms sitting on neighbouring lattice sites. Particles in two adjacent sites of the system lattice are coupled to an auxiliary site in between by the spatial modulation of a Raman laser yielding opposite coupling (Rabi frequency)  $\pm\Omega$ , which corresponds to annihilation process  $(\sigma_j - \sigma_{j+1})$ . Then the creation process  $(\sigma_j^\dagger + \sigma_{j+1}^\dagger)$  is realized by isotropically spontaneous radiation where atoms decay with a rate  $\Gamma$  back to original system sites.

**Summary and outlook.** We have investigated the robustness of QMBS under the impact of dissipation. By choosing a kind of quasi-local dissipation operators, a system with QMBS can be driven to the steady state composed mainly by the QMBS. We confirm this by studying three categories of QMBS. Our results reveal that the dissipation can induce a weak ergodicity breaking steady-state, and the corresponding dissipative dynamics exhibit scarred relaxation with dynamical revival by suitably preparing an initial state. We emphasize that our protocol is generic because the dissipation operators are model-independent, which means that no prior knowledge is required. Similar to the dissipation-induced localization, the mechanism behind this protocol where the quasi-local dissipation-induced QMBS is universal. Besides the QMBS, other types of weak ergodicity breaking, such as Hilbert space fragmentation could be an inter-

esting future direction to study the dissipation-induced non-thermal steady-state, which may deepen the understanding of thermalization and ergodicity breaking in open quantum many-body systems. We also discussed the experimental realization of the quasi-local dissipation operators in cold atom setup, which could provide a routine to make the signature of ergodicity breaking visible over dissipative dynamics and offer potential possibilities for experimentally preparing stable QMBS.

## ACKNOWLEDGEMENTS

The work is supported by National Natural Science Foundation of China (Grant No. 12304290 and Grant No. 12104205). WY acknowledges support from the National Key R&D Program of China under Grant No.2022YFA1405800. LP also acknowledges support from the Fundamental Research Funds for the Central Universities.

\* [panlei@nankai.edu.cn](mailto:panlei@nankai.edu.cn)

- [1] J. Eisert, M. Friesdorf, and C. Gogolin, Quantum many-body systems out of equilibrium, *Nat. Phys.* **7**, 124 (2015).
- [2] L. D' Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics, *Adv. Phys.* **65**, 239 (2016).
- [3] R. Nandkishore and D. A. Huse, Many-body localization and thermalization in quantum statistical mechanics, *Annu. Rev. Condens. Matter Phys.* **6**, 15 (2015).
- [4] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, Colloquium: Many-body localization, thermalization, and entanglement, *Rev. Mod. Phys.* **91**, 021001 (2019).

- [5] B. L. Altshuler, Y. Gefen, A. Kamenev, and L. S. Levitov, Quasiparticle Lifetime in a Finite System: A Nonperturbative Approach, *Phys. Rev. Lett.* **78**, 2803 (1997).
- [6] D. Basko, I. Aleiner, and B. Altshuler, Metal–insulator transition in a weakly interacting many-electron system with localized single-particle states, *Ann. Phys. (Amsterdam)* **321**, 1126 (2006).
- [7] I. V. Gornyi, A. D. Mirlin, and D. G. Polyakov, Interacting Electrons in Disordered Wires: Anderson Localization and Low- $T$  Transport, *Phys. Rev. Lett.* **95**, 206603 (2005).
- [8] V. Oganesyan and D. A. Huse, Localization of interacting fermions at high temperature, *Phys. Rev. B* **75**, 155111 (2007).
- [9] A. Pal and D. A. Huse, Many-body localization phase transition, *Phys. Rev. B* **82**, 174411 (2010).
- [10] J. M. Deutsch, Quantum statistical mechanics in a closed system, *Phys. Rev. A* **43**, 2046 (1991).
- [11] M. Srednicki, Chaos and quantum thermalization, *Phys. Rev. E* **50**, 888 (1994).
- [12] M. Rigol, V. Dunjko, and M. Olshanii, *Nature (London)* **452**, 854 (2008).
- [13] E. Levi, M. Heyl, I. Lesanovsky, and J. P. Garrahan, Robustness of Many-Body Localization in the Presence of Dissipation, *Phys. Rev. Lett.* **116**, 237203 (2016).
- [14] H. P. Lüschen, P. Bordia, S. S. Hodgman, M. Schreiber, S. Sarkar, A. J. Daley, M. H. Fischer, E. Altman, I. Bloch, and U. Schneider, Signatures of Many-Body Localization in a Controlled Open Quantum System, *Phys. Rev. X* **7**, 011034 (2017).
- [15] A. R.-Abadal, J. Choi, J. Zeiher, S. Hollerith, J. Rui, I. Bloch, and C. Gross, Many-Body Delocalization in the Presence of a Quantum Bath, *Phys. Rev. X* **9**, 041014 (2019).
- [16] I. Yusipov, T. Lapyteva, S. Denisov, and M. Ivanchenko, Localization in Open Quantum Systems, *Phys. Rev. Lett.* **118**, 070402 (2017).
- [17] I. Vakulchyk, I. Yusipov, M. Ivanchenko, S. Flach, and S. Denisov, Signatures of many-body localization in steady states of open quantum systems, *Phys. Rev. B* **98**, 020202(R) (2018).
- [18] Y. Liu, Z. Wang, C. Yang, J. Jie, and Y. Wang, Dissipation-Induced Extended-Localized Transition, *Phys. Rev. Lett.* **132**, 216301 (2024).
- [19] Y. Hu, C. Yang, and Y. Wang, Can dissipation induce a transition between many-body localized and thermal states? [arXiv:2407.13655](https://arxiv.org/abs/2407.13655) (2024).
- [20] X.-P. Jiang, X. Yang, Y. Hu, and L. Pan, Dissipation induced ergodic-nonergodic transitions in finite-height mosaic Wannier-Stark lattices, [arXiv:2407.17301](https://arxiv.org/abs/2407.17301) (2024).
- [21] X. Yang, X.-P. Jiang, Z. Wei, Y. Wang, and L. Pan, Dissipation induced transition between extension and localization in the three-dimensional Anderson model, [arXiv:2409.20319](https://arxiv.org/abs/2409.20319) (2024).
- [22] H. Bernien, S. Schwartz, A. Keesling, H. Levine, A. Omran, H. Pichler, S. Choi, A. S. Zibrov, M. Endres, M. Greiner, et al., Probing many-body dynamics on a 51-atom quantum simulator, *Nature* **551**, 579 (2017).
- [23] D. Bluvstein, A. Omran, H. Levine, A. Keesling, G. Semeghini, S. Ebadi, T. T. Wang, A. A. Michailidis, N. Maskara, W. W. Ho, S. Choi, M. Serbyn, M. Greiner, V. Vuletić, and M. D. Lukin, Controlling quantum many-body dynamics in driven rydberg atom arrays, *Science* **371**, 1355 (2021).
- [24] C. J. Turner, A. A. Michailidis, D. A. Abanin, M. Serbyn, and Z. Papić, Weak ergodicity breaking from quantum many-body scars, *Nat. Phys.* **14**, 745 (2018).
- [25] C. J. Turner, A. A. Michailidis, D. A. Abanin, M. Serbyn, and Z. Papić, Quantum scarred eigenstates in a rydberg atom chain: Entanglement, breakdown of thermalization, and stability to perturbations, *Phys. Rev. B* **98**, 155134 (2018).
- [26] M. Serbyn, D. A. Abanin, and Z. Papić, Quantum many-body scars and weak breaking of ergodicity, *Nat. Phys.* **17**, 675 (2021).
- [27] S. Moudgalya, B. A. Bernevig, and N. Regnault, Quantum many-body scars and hilbert space fragmentation: A review of exact results, *Rep. Prog. Phys.* **85**, 086501 (2022).
- [28] A. Chandran, T. Iadecola, V. Khemani, and R. Moessner, Quantum many-body scars: A quasiparticle perspective, *Annu. Rev. Condens. Matter Phys.* **14**, 443 (2022).
- [29] K. Pakrouski, P. N. Pallegar, F. K. Popov, and I. R. Klebanov, Group theoretic approach to many-body scar states in fermionic lattice models, *Phys. Rev. Research* **3**, 043156 (2021).
- [30] Q. Chen, S. A. Chen, and Z. Zhu, Weak ergodicity breaking in non-hermitian many-body systems, *SciPost Phys.* **15**, 052 (2023).
- [31] H.-R. Wang, D. Yuan, S.-Y. Zhang, Z. Wang, D.-L. Deng, and L.-M. Duan, Embedding Quantum Many-Body Scars into Decoherence-Free Subspaces, *Phys. Rev. Lett.* **132**, 150401 (2024).
- [32] R. Shen, F. Qin, J.-Y. Desaulles, Z. Papić, and C. H. Lee, Enhanced Many-Body Quantum Scars from the Non-Hermitian Fock Skin Effect, *Phys. Rev. Lett.* **133**, 216601 (2024).
- [33] G. Lindblad, On the generators of quantum dynamical semigroups, *Commun. Math. Phys.* **119**, 48 (1976).
- [34] V. Gorini, A. Kossakowski, and E. C. Sudarsahan, Completely positive dynamical semigroups of N-level systems, *J. Math. Phys.* **17**, 821 (1976).
- [35] S. Diehl, A. Micheli, A. Kantian, B. Kraus, H. P. Büchler, and P. Zoller, Quantum states and phases in driven open quantum systems with cold atoms, *Nat. Phys.* **4**, 878 (2008).
- [36] B. Kraus, H. P. Büchler, S. Diehl, A. Kantian, A. Micheli, and P. Zoller, Preparation of entangled states by quantum Markov processes, *Phys. Rev. A* **78**, 042307 (2008).
- [37] D. Marcos, A. Tomadin, S. Diehl, and P. Rabl, Photon condensation in circuit quantum electrodynamics by engineered dissipation, *New J. Phys.* **14**, 055005 (2012).
- [38] W. Ho, S. Choi, H. Pichler, and M. D. Lukin, Periodic orbits, entanglement, and quantum many-body scars in constrained models: Matrix product state approach, *Phys. Rev. Lett.* **122**, 040603 (2019).
- [39] D. K. Mark and O. I. Motrunich,  $\eta$ -pairing states as true scars in an extended Hubbard model, *Phys. Rev. B* **102**, 075132 (2020).
- [40] S. Moudgalya, N. Regnault, and B. A. Bernevig,  $\eta$ -pairing in Hubbard models: From spectrum generating algebras to quantum many-body scars, *Phys. Rev. B* **102**, 085140 (2020).
- [41] J. Ren, C. Liang, and C. Fang, Quasisymmetry Groups and Many-Body Scar Dynamics, *Phys. Rev. Lett.* **126**, 120604 (2021).
- [42] L.-H. Tang, N. O’Dea, and A. Chandran, Multimagnon quantum many-body scars from tensor operators, *Phys.*

- [Rev. Research 4, 043006 \(2022\)](#).
- [43] K. Pakrouski, P. N. Pallegar, F. K. Popov, I. R. Klebanov, Many Body Scars as a Group Invariant Sector of Hilbert Space, [Phys. Rev. Lett. 125, 230602 \(2020\)](#).
  - [44] B. Olmos, M. Mller, and I. Lesanovsky, B. Olmos, M. Mller, and I. Lesanovsky, Thermalization of a strongly interacting 1d Rydberg lattice gas, [New J. Phys. 12, 013024 \(2010\)](#).
  - [45] I. Lesanovsky, B. Olmos, and J. P. Garrahan, Thermalization in a Coherently Driven Ensemble of Two-Level Systems, [Phys. Rev. Lett. 105, 100603 \(2010\)](#).
  - [46] C. Ates, J. P. Garrahan, and I. Lesanovsky, Thermalization of a Strongly Interacting Closed Spin System: From Coherent Many-Body Dynamics to a Fokker-Planck Equation, [Phys. Rev. Lett. 108, 110603 \(2012\)](#).
  - [47] S. Ji, C. Ates, J. P. Garrahan, and I. Lesanovsky, Equilibration of quantum hard rods in one dimension, [J. Stat. Mech. 2013, P02005](#).
  - [48] C.-J. Lin, A. Chandran, and O. I. Motrunich, Slow thermalization of exact quantum many-body scar states under perturbations, [Phys. Rev. Research 2, 033044 \(2020\)](#).
  - [49] I. Mondragon-Shem, M. G. Vavilov, and I. Martin, Fate of Quantum Many-Body Scars in the Presence of Disorder, [PRX Quantum 2, 030349 \(2021\)](#).
  - [50] P. Zanardi, H. T. Quan, X. Wang, and C. P. Sun, Mixed-state fidelity and quantum criticality at finite temperature, [Phys. Rev. A 75, 032109 \(2007\)](#).
  - [51] N. Shiraishi and T. Mori, Systematic construction of counterexamples to the eigenstate thermalization hypothesis, [Phys. Rev. Lett. 119, 030601 \(2017\)](#).

Supplementary Material for “Robustness of quantum many-body scars in the presence of Markovian baths”

THE QMBS AND DISSIPATIVE SCARRED DYNAMICS IN THE MODEL WITH CONSERVED QUANTITY

In this supplementary material, we apply our protocol to systems in which the analytically solvable QMBS is equally spaced and can be constructed by spectrum generating algebra. We will show that the QMBS can become exact steady states under the quasi-local dissipators introduced in the maintext (see Eq.(4)). To show this, we take the  $J_1 - J_2$  model with Dzyaloshinskii-Moriya interaction (DMI) as an example whose Hamiltonian is

$$H_{J_1 - J_2} = \sum_j (J_1 \mathbf{s}_j \cdot \mathbf{s}_{j+1} + J_2 \mathbf{s}_j \cdot \mathbf{s}_{j+2}) + H_{\text{DMI}} + h_z S_{\text{tot}}^z, \quad (\text{S1})$$

where  $H_{\text{DMI}} = \sum_j \hat{\mathbf{z}} \cdot (\mathbf{s}_j \times \mathbf{s}_{j+1})$  denotes the DMI and  $S_{\text{tot}} = \sum_{j=1}^L s_j^z$ . As discussed in Ref.[39], there are QMBS composed of equally spaced zero-momentum ( $k = 0$ ) magnon tower. Since the number of spin-up  $N_\uparrow$  and spin-down  $N_\downarrow$  is conserved, and also the dissipation operators do not change them, hence one can solve the Liouvillian superoperator and the steady-state individually in each subspace  $\{N_\uparrow, N_\downarrow\}$ . As shown in Fig.S1(a), the steady-state is the exact QMBS in each subspace and scar tower has equally spaced energy which means it might be possible to generates dissipative dynamics with periodic revival. In fact, if we prepare a ferromagnetic state  $|\rightarrow\rightarrow\rightarrow\cdots\rightarrow\rightarrow\rightarrow\rangle$  as the initial state which is an equally weighted superposition of QMBS ( $k = 0$  magnon), then the dissipative dynamics exhibits a perfect revival without decay, as shown in Fig.S1(b).




FIG. S1. (a) Distribution of the steady-state density matrix on eigenstate of the  $J_1 - J_2$  model in the eigenbasis of each subspace  $\{N_\uparrow, N_\downarrow\}$  with the dissipative phases  $\theta = 0$ . (b) The dissipative dynamics of quantum fidelity from a ferromagnetic initial state  $|\phi_0\rangle = |\rightarrow\rightarrow\rightarrow\cdots\rightarrow\rightarrow\rightarrow\rangle$ . Here the parameters are chosen as  $J_1 = 1, J_2 = -0.2, D = 1, h = 0.5$  with system size  $L = 12$ .