## On Kato's ramification filtration

Subhadip Majumder

## Abstract

For a Henselian discrete valued field K of characteristic p > 0, Kato [36] defined a ramification filtration  $\{\operatorname{fil}_n H^q(K, \mathbb{Q}_p/\mathbb{Z}_p(q-1))\}_{n\geq 0}$  on  $H^q(K, \mathbb{Q}_p/\mathbb{Z}_p(q-1))$ . One can also define a ramification filtration on  $H^q(U, \mathbb{Z}/p^m(q-1))$  using the local Kato-filtration, where U is the complement of a simple normal crossing divisor in a regular scheme X of characteristic p > 0. The main objective of this thesis is to provide a cohomological description of these filtrations using de Rham-Witt sheaves and present several applications.

To achieve our goal, we study a theory of the filtered de Rham-Witt complex of Ffinite regular schemes of characteristic p > 0 and prove several properties which are well known for the classical de Rham-Witt complex of regular schemes. As applications, we prove a refined version of Jannsen-Saito-Zhao's duality [31] over finite fields, and a similar duality for smooth projective curves over local fields. As another application, we prove a Lefschetz theorem for unramified and ramified Brauer group (with modulus) of smooth projective F-finite schemes over a field of characteristic p > 0. Further applications are given in [49] and [50].

This work is a slightly modified version of the author's Ph.D. thesis, submitted in July 2024 at TIFR, Mumbai.

## Contents

Notations       Introduction				
	1.1	de Rham-Witt Complex	11	
	1.2	The log de Rham-Witt complex	15	
		1.2.1 Definitions	15	
		1.2.2 Log structure associated to SNCD	17	
2	The	e filtered de Rham-Witt complex	25	
	2.1	Definition and properties	25	
3	Stru	acture of the filtered de Rham-Witt complex I	33	
	3.1	Filtered de Rham-Witt complex of complete local $\mathbb{F}_p$ -algebras	33	
	3.2	Cartier homomorphism I	44	
4	Stru	acture of the filtered de Rham-Witt complex II	47	
	4.1	de Rham complex of multivariate power series ring	47	
	4.2	From complete to the non-complete case	48	
	4.3	Cartier homomorphism II	55	
5	Con	nparison with Kato's filtration	63	
	5.1	Comparison with Local Kato filtration	63	
		5.1.1 Kato filtration	64	
		5.1.2 Proof of the main theorem	67	
	5.2	Comparison with Global Kato filtration	73	
		5.2.1 Global Kato filtration	73	
		5.2.2 Relation with Brauer group with modulus	80	

6	Dua	ality theorem for Hodge-Witt cohomology over finite fields	83		
	6.1	Definition of complexes and their properties	83		
	6.2	Pairings of complexes	87		
	6.3	The duality theorem over finite fields	89		
7	A d	luality theorem for 2-local fields	93		
	7.1	Kato topology	93		
	7.2	The reciprocity homomorphism	96		
	7.3	Duality for $\kappa_q^m(A n)$ and $\varpi_q^m(A n)$	97		
8	Dua	ality theorem for Hodge-Witt cohomology over local fields	101		
	8.1	Pairing of complexes	101		
	8.2	Relation with local duality	102		
		8.2.1 The maps $(\psi_{\underline{n},x}^q)^*$ vs. $\widehat{\rho}_{K,m}^{q,\underline{n}+1}$	104		
	8.3	Topologies on the cohomology groups	108		
		8.3.1 Topology of $H^i_{\text{ét}}(X, W_m \Omega^q_{X D_{\underline{n}+1}, \log})$ for $i \neq 1 \dots \dots \dots \dots$	108		
		8.3.2 Topology of $H^1_{\text{ét}}(X, W_m \Omega^2_{X D_{n+1}, \log})$	110		
		8.3.3 The Brauer-Manin pairing with modulus $\ldots \ldots \ldots \ldots \ldots$	114		
		8.3.4 Topology of $H^1_{\text{\'et}}(X, W_m \Omega^1_{X D_{n+1}, \log})$	116		
		8.3.5 The duality theorem over local fields $\ldots \ldots \ldots \ldots \ldots \ldots$	122		
9	Lefs	schetz hypersurface theorem	125		
	9.1	Some exact sequences	125		
	9.2	Weak Lefschetz for <i>p</i> -adic étale cohomology	128		
	9.3	Weak Lefschetz for $\ell$ -adic étale cohomology	132		
	9.4	Applications	133		
		9.4.1 Lefschetz for Picard groups	134		
		9.4.2 Lefschetz for Brauer groups	135		
		9.4.3 Lefschetz in complete intersection case	137		
Bibliography 141					

## Notations

We shall work over a field k of characteristic p > 0 throughout this thesis, unless otherwise stated. We let  $\mathbf{Sch}_{k}^{\text{ft}}$  denote the category of separated and finite type k-schemes. We let  $\mathbf{Sch}_{k}$  denote the category of separated Noetherian k-schemes. The product  $X \times_{\text{Spec}(k)} Y$ in  $\mathbf{Sch}_{k}$  will be written as  $X \times Y$ . We let  $X^{(q)}$  (resp.  $X_{(q)}$ ) denote the set of points on X having codimension (resp. dimension) q.

Let X be a scheme and let  $\mathcal{M}_X$  be a sheaf of monoids on X. We shall denote a log scheme  $(X, \mathcal{M}_X, \alpha: \mathcal{M}_X \to \mathcal{O}_X)$  on a scheme X by  $X_{\log}$ . If X = Spec(A), we shall also write these as  $(A, \alpha: \mathcal{M}_A \to A)$  and  $A_{\log}$ . By  $X_{\text{triv}}$ , we shall mean the (trivial) log structure given by the usual inclusion  $\alpha: \mathcal{O}_X^{\times} \to \mathcal{O}_X$ . For a scheme X, we let  $\operatorname{Sch}_X$  denote the category of schemes over X. All rings are assumed to be Noetherian commutative rings with unity. For a reduced ring A, we shall let Q(A) denote the ring of total quotients of A.

We let  $\operatorname{Sch}_{k/\operatorname{zar}}$  (resp.  $\operatorname{Sch}_{k/\operatorname{nis}}$ , resp.  $\operatorname{Sch}_{k/\acute{et}}$ ) denote the Zariski (resp. Nisnevich, resp. étale) site of  $\operatorname{Sch}_k$ . We let  $\epsilon: \operatorname{Sch}_{k/\acute{et}} \to \operatorname{Sch}_{k/\operatorname{nis}}$  denote the canonical morphism of sites. If  $\mathcal{F}$  is a sheaf on  $\operatorname{Sch}_{k/\operatorname{nis}}$ , we shall denote  $\epsilon^* \mathcal{F}$  also by  $\mathcal{F}$  as long as the usage of the étale topology is clear in a context. For  $X \in \operatorname{Sch}_k$ , we shall let  $F: X \to X$ denote the absolute Frobenius morphism. For a scheme X endowed with a topology  $\tau \in \{\operatorname{zar}, \operatorname{nis}, \acute{et}\}, X_{\tau}$  denotes the small  $\tau$ -site of X. For a closed subset  $Y \subset X$ , and a  $\tau$ -sheaf  $\mathcal{F}$ , we let  $H^*_{\tau,Y}(X, \mathcal{F})$  denote the  $\tau$ -cohomology groups of  $\mathcal{F}$  with support in Y. If  $\tau$  is fixed in a section, we shall drop it from the notations of the cohomology groups. We shall let  $D(X_{\tau})$  denote the bounded derived category of sheaves on  $X_{\tau}$ . If  $X = \operatorname{Spec} A$ , then the cohomology group  $H^i_{\tau}(X, \mathcal{F})$  may also be written as  $H^i_{\tau}(A, \mathcal{F})$ .

For any field extension k' over k, X(k') will denote the set of k'-rational points of X. Div(X) will denote the set of all Weil divisors of X. For a reduced effective Weil divisor E, Div<sub>E</sub>(X) will denote the set of all Weil divisors, whose supports are inside E.

 $\mathbb{Z}$  will denote the set of all integers and  $\mathbb{N}$  (resp.  $\mathbb{N}_0$ ) will denote the set of all positive (resp. non negative) integers. [.] (resp. [.]) will denote the greatest (resp. least) integer function. For  $\underline{n} = (n_1, \ldots, n_r) \in \mathbb{Z}^r$  and  $p \in \mathbb{N}$ , we define  $\underline{n}/p := (\lfloor n_1/p \rfloor, \ldots, \lfloor n_r/p \rfloor)$ .

We shall let  $\mathbf{Ab}$  denote the category of abelian groups. For an abelian group A, we shall write  $\operatorname{Tor}_{\mathbb{Z}}^{1}(A,\mathbb{Z}/n)$  as  ${}_{n}A$  and A/nA as A/n. For  $A, B \in \mathbf{Ab}$ , the tensor product  $A \otimes_{\mathbb{Z}} B$  will be written as  $A \otimes B$ . We shall let  $A\{p\}$  (resp.  $A\{p'\}$ ) denote the subgroup of elements of A which are annihilated by a power of (resp. an integer prime to) p. We let  $\mathbf{Ab}_{\mathbf{p}}$  be the category of p-primary torsion abelian groups. We define  $A_{\text{tor}} \coloneqq A\{p\} \cup A\{p'\}$ . We let  $\mathbf{Tab}$  denote the category of topological abelian groups with continuous homomorphisms. A locally compact Hausdorff topological group G is called "torsion-by-profinite" if there is an exact sequence

$$0 \to G_{\rm pf} \to G \to G_{\rm dt} \to 0$$

where  $G_{pf}$  is an open and profinite subgroup of G and  $G_{dt}$  is a torsion group with the quotient topology (necessarily discrete). The category of "torsion-by-profinite" will be denoted as **Pfd**.

By an *n*-local field, we mean a complete discrete valued field whose residue field is an (n-1)-local field, where the 0-local fields are defined to be finite fields. The 1-local fields will also be mentioned as local fields.

We shall write  $H^q_{\text{\acute{e}t}}(X, \mathbb{Q}/\mathbb{Z}(q-1))$  (resp.  $H^q_{\text{\acute{e}t}}(A, \mathbb{Q}/\mathbb{Z}(q-1))$ ) as  $H^q(X)$  (resp.  $H^q(A)$ ) for any  $q \ge 0$  (see [35, § 3.2, Defn. 1]).

For a Noetherian scheme X, we shall let  $\pi_1^{ab}(X)$  denote the abelianized étale fundamental group of X and consider it as a topological group with its profinite topology. We shall write  $\pi_1^{ab}(\operatorname{Spec}(F))$  interchangeably as  $G_F$ , where the latter is the abelianized absolute Galois group of a field F. We shall let  $\operatorname{Br}(X)$  denote the cohomological Brauer group of X, which is defined as the étale cohomology group  $H^2_{\acute{e}t}(X, \mathbb{G}_m)$ .

RLR is abbreviated as regular local ring. SNCD is abbreviated as simple normal crossing divisor.

## Introduction

The étale fundamental group and the Brauer group are two of the most extensively studied objects in algebraic and arithmetic geometry. The first one was introduced by Grothendieck in the 1960's in [19]. Given a connected scheme X and a geometric point  $\overline{x}$ , the étale fundamental group  $\pi_1^{\text{ét}}(X,\overline{x})$  is defined as the inverse limit of the automorphism groups  $\operatorname{Aut}_X(X_\alpha(\overline{x}))$ , where  $X_\alpha$  varies over all finite étale covers of X and  $X_\alpha(\overline{x})$  be the geometric fiber over  $\overline{x}$ . The étale fundamental group of a field k is the absolute Galois group  $\operatorname{Gal}(\overline{k}/k)$ . So the theory of étale fundamental group can be considered as a generalization of the Galois theory of fields to the case of schemes. The maximal abelian quotient of  $\pi_1^{\text{ét}}(X,\overline{x})$  is denoted by  $\pi_1^{ab}(X,\overline{x})$ .

On the other hand, the Brauer group of a field k,  $\operatorname{Br}(k)$  was introduced by R. Brauer. It is defined as the group of equivalent classes of central simple algebras over k, with addition given by the tensor product of algebras. Another important interpretation of the Brauer group of a field k is that it classifies the projective varieties over k that become isomorphic to projective space when base changed to  $\overline{k}$ , the algebraic closure of k. Such a variety is called a Severi-Brauer variety. Brauer group was generalized from field to any commutative rings by Auslander and Goldman in [5] and later to arbitrary schemes by Grothendieck, where one uses Azumaya algebras instead of central simple algebras and projective bundles which are étale locally trivial instead of projective space in the previous definition of Brauer group of a field. This group is often known as the Brauer-Azumaya group denoted by  $\operatorname{Br}_{Az}(X)$ , for a scheme X (cf. [12, Chapter 3]). On the other hand, another generalisation of the Brauer group of a field to a scheme is the cohomological Brauer group  $\operatorname{Br}(X) \coloneqq H^2_{\acute{e}t}(X, \mathbb{G}_m)$ . It is well known (cf. [12, Thm. 4.2.1]) that  $\operatorname{Br}_{Az}(X) \hookrightarrow \operatorname{Br}(X)_{tor}$ ; and it is an isomorphism if X is a quasi-projective scheme. Also,  $\operatorname{Br}(X)$  is torsion if X is a regular and integral scheme by [12, Thm. 3.5.5].

The goal of this thesis is to study the wild ramification part of  $\pi_1^{ab}(-)$ , Br(-) and more generally of the *p*-adic étale cohomology of a scheme which is a complement of a simple normal crossing divisor in a proper smooth variety of characteristic p > 0. In an attempt to do so, we study a theory of filtered de Rham-Witt complex and its role in Kato's local ramification theory [36].

## Ramification theory

Let k be a field and X be a proper smooth k scheme. The group  $\pi_1^{ab}(X)$  (or Br(X)) is often known as the unramified (abelian) étale fundamental group (resp unramified Brauer group). However, if  $U \hookrightarrow X$  is an open subscheme of X whose complement is of codimension 1 (if codimension > 1, then  $\pi_1^{ab}(-)$  and Br(-) are unchanged by purity), then to study  $\pi_1^{ab}(U)$  (resp. Br(U)), one needs to consider all the abelian étale covers (resp Brauer classes) of U with ramifications along its complement. This is where the Ramification theory comes into the picture.

Let K be a Henselian discrete valued field. If K is a local field, the ramification theory of  $G_K$  is classical ([61]). If residue field of K is not necessarily finite, then the ramification theory for  $H^1_{\text{ét}}(K, \mathbb{Q}/\mathbb{Z}) = \text{Hom}_{cts}(G_K, \mathbb{Q}/\mathbb{Z})$  was developed by Brylinski [9], Kato [36], Matsuda [52] (see also [1]). As in the classical case, these authors defined a filtration  $\{\text{fil}_n H^1_{\text{ét}}(K, \mathbb{Q}/\mathbb{Z})\}_n$ , that classifies all abelian étale covers of K whose ramification at the closed point of  $\mathcal{O}_K$  is bounded by  $n \ge 0$ . More generally, for any  $q \ge 0$ , Kato [36] also defined a ramification filtration of  $H^{q+1}(K)$  (which we write as  $\text{fil}_n^{bk} H^{q+1}(K)$ ), where, in modern language,  $H^{q+1}(K)$  is same as  $H^{q+1}(K, \mathbb{Q}/\mathbb{Z}(q))$ , the étale motivic cohomology of Suslin and Voevodsky. These filtrations are usually difficult to study and also unlike the unramified case, they don't have any cohomological interpretation in general.

Now to study global ramification theory, one can use Kato's local ramification theory to define a filtration of  $H^1_{\text{\acute{e}t}}(U, \mathbb{Q}/\mathbb{Z})$ , namely  $\{\operatorname{fil}_D H^1_{\text{\acute{e}t}}(U, \mathbb{Q}/\mathbb{Z})\}_D$  so that

$$H^{1}_{\text{\acute{e}t}}(U, \mathbb{Q}/\mathbb{Z}) = \varinjlim_{D} \operatorname{fil}_{D} H^{1}_{\text{\acute{e}t}}(U, \mathbb{Q}/\mathbb{Z}),$$

where D varies over all effective divisors whose support is contained in  $X \\ V$ . The profinite group  $\pi_1^{ab}(X, D) := (\operatorname{fil}_D H^1_{\operatorname{\acute{e}t}}(U, \mathbb{Q}/\mathbb{Z}))^{\vee}$  classifies all the abelian étale covers of U whose ramifications along its complement is bounded by D. Here  $(-)^{\vee}$  means the discrete dual. One can also define a ramification filtration for  $\operatorname{Br}(U)$  (see Definition 0.0.2) in a similar way.

In [43], Kerz and Saito constructed a two term complex (which we write as  $W_m \mathcal{F}_D^{0,\bullet}$ ) in  $\mathcal{D}^b(X_{\acute{e}t}, \mathbb{Z}/p^m)$ , such that  $\mathbb{H}^1_{\acute{e}t}(X, W_m \mathcal{F}_D^{0,\bullet}) \cong \operatorname{fil}_D H^1_{\acute{e}t}(U, \mathbb{Z}/p^m) = {}_{p^m}(\operatorname{fil}_D H^1_{\acute{e}t}(U, \mathbb{Q}/\mathbb{Z}))$ , when  $E = X \setminus U$  is a simple normal crossing divisor and X is smooth over a finite field. This allowed them to get a Lefschetz-type result for  $\pi_1^{ab}(X, D)$  and the higher dimensional ramified class field theory over finite fields in [44].

One of the objectives of this thesis is to study the ramification theory of a regular Ffinite scheme of characteristic p > 0 using Kato's filtration and give several applications.

We recall that a scheme X is called F-finite, if the absolute Frobenius map  $F: X \to X$ is a finite morphism. Let X be a regular F-finite scheme of dim N and  $E = \bigcup_{i=1}^{r} E_i$  be a simple normal crossing divisor, where  $E_i$ 's are the irreducible components. Let  $U = X \setminus E$ and  $j: U \to X$ . We fix an  $m \ge 1$ . Let  $D = \sum_{1 \le i \le r} n_i E_i$  with  $n_i \ge 0$  and  $K_i = Frac(\mathcal{O}_{X,x}^h)$  if  $\overline{\{x\}} = E_i$ . Also we set  $H_m^{q+1}(K_i) = H^{q+1}(K_i, \mathbb{Z}/p^m(q))$ . We define

$$\operatorname{fil}_{D}^{\log} H^{q+1}(U, \mathbb{Z}/p^{m}(q)) \coloneqq \operatorname{Ker}\left(H^{q+1}(U, \mathbb{Z}/p^{m}(q)) \to \bigoplus_{1 \le i \le r} \frac{H_{m}^{q+1}(K_{i})}{\operatorname{fil}_{n_{i}}^{\operatorname{bk}} H^{q+1}(K_{i}) \cap H_{m}^{q+1}(K_{i})}\right)$$

The arrow on the right hand side is induced by canonical restrictions.

The first main result of this thesis is the following theorem whose q = 0 case is the Kerz-Saito's result mentioned above :

**Theorem 0.0.1.** (Theorem 5.2.6) For any  $q \ge 0$ , there exists a two term complex  $W_m \mathcal{F}_D^{q,\bullet}$  of étale sheaves such that

$$\mathbb{H}^{1}_{\acute{e}t}(X, W_m \mathcal{F}^{q, \bullet}_D) \cong \mathrm{fil}_D H^{q+1}(U, \mathbb{Z}/p^m(q)).$$

**Definition 0.0.2.** We let  $\operatorname{Br}^{\operatorname{div}}(X|D)$  denote the subgroup of  $\operatorname{Br}(U)$  consisting of elements  $\chi$  such that for every  $x \in D^{(0)}$ , the image  $\chi_x$  of  $\chi$  under the canonical map  $\operatorname{Br}(U) \to \operatorname{Br}(K_x)$  lies in  $\operatorname{fil}_{n_x} \operatorname{Br}(K_x)$ , where  $n_x$  is multiplicity of D at x.

As an immediate corollary of Theorem 0.0.1, we have the following.

Corollary 0.0.3. (Corollary 5.2.10) We have an exact sequence

$$0 \to \operatorname{Pic}(U)/p^m \to \mathbb{H}^1_{\acute{e}t}(X, W_m \mathcal{F}_D^{1, \bullet}) \to {}_{p^m} \operatorname{Br}^{\operatorname{div}}(X|D) \to 0$$

The above result can be thought of as the ramified version of the following well-known exact sequence :

$$0 \to \operatorname{Pic}(X)/p^m \to H^1(X, W_m \Omega^1_{X, \log}) \to {}_{p^m} \operatorname{Br}(X) \to 0,$$

where  ${}_{p^m}A := \operatorname{Ker}(A \xrightarrow{p^m} A)$ . Corollary 0.0.3 will have applications to prove a Lefschetz theorem for  $\operatorname{Br}^{\operatorname{div}}(X|D)$  and also a ramified duality with modulus for curves over local fields in the later sections of this thesis.

The following theorem in local ramification theory is the key result for proving Theorem 0.0.1.

**Theorem 0.0.4.** (Theorem 5.1.16) Let K be an F-finite Henselian discrete valued field of characteristic p > 0. Then, for all  $n, q \ge 0$  there exists a two term complex  $W_m \mathcal{F}_n^{q,\bullet}$  in  $\mathcal{D}^b(K_{\acute{e}t}, \mathbb{Z}/p^m)$  such that

$$\mathbb{H}^{1}_{\acute{e}t}(K, W_m \mathcal{F}^{q, \bullet}_n) \cong {}_{p^m} \mathrm{fil}_n^{bk} H^{q+1}(K).$$

### Filtered de Rham-Witt complex

To construct the two term complex  $W_m \mathcal{F}_D^{q,\bullet}$ , we study a theory of filtered de Rham-Witt complex of X. We continue with the same notations. Also, we allow  $n_i$  in the definition of D to be any element in  $\mathbb{Z}$ .

For any  $q \ge 0$ , we first define a quasi-coherent sheaf of  $W_m \mathcal{O}_X$  modules,  $\operatorname{fil}_D W_m \Omega_U^q \subset j_* W_m \Omega_U^q$  (see Definition 2.1.6), which satisfies the following properties (Lemma 2.1.7):

1. 
$$d(\operatorname{fil}_D W_m \Omega_U^q) \subset \operatorname{fil}_D W_m \Omega_U^{q+1}$$
,

- 2.  $F(\operatorname{fil}_D W_{m+1}\Omega_U^q) \subset \operatorname{fil}_D W_m\Omega_U^q$ ,
- 3.  $V(\operatorname{fil}_D W_m \Omega_U^q) \subset \operatorname{fil}_D W_{m+1} \Omega_U^q$ ,
- 4.  $R(\operatorname{fil}_D W_{m+1}\Omega_U^q) = \operatorname{fil}_{D/p} W_m \Omega_U^q$ , where  $D/p = \sum_i \lfloor n_i/p \rfloor E_i$ ,
- 5.  $\lim_{n \to \infty} \operatorname{fil}_{nE} W_m \Omega_U^q \xrightarrow{\cong} j_* W_m \Omega_U^q.$

Also we have  $\operatorname{fil}_D W_m \Omega_U^q \subset \operatorname{fil}_{D'} W_m \Omega_U^q$ , if  $D' \geq D$  and  $\operatorname{supp}(D') \subset E$ . Moreover,  $\operatorname{fil}_D W_m \Omega_U^q \subset W_m \Omega_X^q$  if D < 0 (i.e, all its components have multiplicity < 0). We also remark that, for D = empty divisor,  $\operatorname{fil}_D W_m \Omega_U^{\bullet} = W_m \Omega_X^{\bullet}(\log E)$ , the de Rham-Witt sheaf for the canonical log structure given by E (cf. [29]). However, if U = X, then  $\operatorname{fil}_D W_m \Omega_U^{\bullet} = W_m \Omega_X^{\bullet}$ .

At this point, we refer the reader to [31] and [22], where the authors studied a similar filtered de Rham-Witt complex tailored to their respective objectives. See also [62], where a version of relative de Rham-Witt complex with modulus has been studied from the perspective of motives with modulus.

The following technical result plays an important role in studying fil<sub>D</sub> $W_m \Omega_U^q$ .

**Theorem 0.0.5.** (Theorem 4.3.17(2)) We have a short exact sequence of Zariski sheaves of  $W_m \mathcal{O}_X$ -modules

$$0 \to V^{m-1}(\operatorname{fil}_D\Omega_U^q) + dV^{m-1}(\operatorname{fil}_D\Omega_U^{q-1}) \to \operatorname{fil}_DW_m\Omega_U^q \xrightarrow{R} \operatorname{fil}_{D/p}W_{m-1}\Omega_U^q \to 0.$$

The above theorem is an analogue of [30, I.3.2] and [51, 1.16].

Recall that  $Z_1 W_m \Omega_U^q = F(W_{m+1} \Omega_U^q) = \operatorname{Ker}(F^{m-1}d : W_m \Omega_U^q \to \Omega_U^{q+1})$ . We define

$$Z_1 \operatorname{fil}_D W_m \Omega_U^q \coloneqq \operatorname{fil}_D W_m \Omega_U^q \cap j_* Z_1 W_m \Omega_U^q.$$

The following key proposition, which is analogous to [32, Lemma 4.1.3], allows one to define the two term complex  $W_m \mathcal{F}_D^{q,\bullet}$ . Recall that there is a higher Cartier operator  $C: Z_1 W_m \Omega_U^q \to W_m \Omega_U^q$ .

$$C: Z_1 \operatorname{fil}_D W_m \Omega^q_U \to \operatorname{fil}_{D/p} W_m \Omega^q_U$$

Moreover, there are isomorphisms

$$\overline{F}: \operatorname{fil}_{D/p} W_m \Omega_U^q \xrightarrow{\cong} \frac{Z_1 \operatorname{fil}_D W_m \Omega_U^q}{dV^{m-1} (\operatorname{fil}_D \Omega_U^{q-1})}; \quad \overline{C}: \frac{Z_1 \operatorname{fil}_D W_m \Omega_U^q}{dV^{m-1} (\operatorname{fil}_D \Omega_U^{q-1})} \xrightarrow{\cong} \operatorname{fil}_{D/p} W_m \Omega_U^q.$$

As a result, we have

**Definition 0.0.7.** (Definition 4.3.16) For  $D \ge -E_1 - E_2 - \cdots - E_r$ , we define

$$W_m \mathcal{F}_D^{q,\bullet} \coloneqq \left( Z_1 \operatorname{fil}_D W_m \Omega_U^q \xrightarrow{1-C} \operatorname{fil}_D W_m \Omega_U^q \right)$$

The complex  $W_m \mathcal{F}_D^{q,\bullet}$  can be considered as a ramified analogue of  $W_m \Omega_{X,\log}^q$ . Here,  $W_m \Omega_{X,\log}^q$  is the logarithmic Hodge-Witt sheaf on  $X_{\text{\acute{e}t}}$ , defined as the image of the map

$$d\log : \mathcal{K}_{q,X}^{M} \to W_{m}\Omega_{X}^{q}$$
$$d\log(u_{,}, \dots, u_{q}) = d\log[u_{1}]_{m} \wedge \dots \wedge d\log[u_{q}]_{m},$$

where  $\mathcal{K}_{q,X}^M$  be the Milnor K-theory sheaf. The following result is the analogue of the classical p - R exact sequence for  $W_m \Omega_{X,\log}^q$  (and also [31, Thm 1.1.6]).

**Theorem 0.0.8.** (Theorem 4.3.17(11)) There exists a distinguished triangle in the derived category of Zariski sheaves of abelian groups on X

$$W_1 \mathcal{F}_D^{q,\bullet} \xrightarrow{V^{m-1}} W_m \mathcal{F}_D^{q,\bullet} \xrightarrow{R} W_{m-1} \mathcal{F}_{D/p}^{q,\bullet} \xrightarrow{+} W_1 \mathcal{F}_D^{q,\bullet}[1].$$

## Relation with Jannsen-Saito-Zhao's sheaf

In this section, we continue with the additional assumption that X is smooth and finitetype over an F-finite field k.

Let's continue with the notations of the previous subsections. Moreover, we assume D > 0. Then  $D_{\text{red}} = E$ . Jannsen, Saito and Zhao in [31] have defined a subsheaf  $W_m \Omega^q_{X|D,\log} \subset W_m \Omega^q_{X,\log}$  in  $X_{\text{\acute{e}t}}$  and shown an isomorphism

$$\operatorname{dlog}: \mathcal{K}^{M}_{q,X|D} / (p^{m} \mathcal{K}^{M}_{q,X} \cap \mathcal{K}^{M}_{q,X|D}) \xrightarrow{\cong} W_{m} \Omega^{q}_{X|D,\log}$$

Here,  $\mathcal{K}_{q,X|D}^{M}$  is the relative Milnor K-theory sheaf in  $X_{\text{\acute{e}t}}$  defined in [69] as the image of the cup product

$$\operatorname{Ker}(\mathcal{O}_X^{\times} \to \mathcal{O}_E^{\times}) \otimes j_* \mathcal{K}_{q-1,U}^M \to j_* \mathcal{K}_{q,U}^M, \text{ where } j: U \hookrightarrow X.$$

This log-version of relative Milnor K-theory sheaf was also considered in [34], [35], [36], [37], [38] to establish Higher dimensional local class field theory and abelianized ramification theory. Part (1) of our next result relates Jannsen-Saito-Zhao's sheaf with our filtration in the following way :

Lemma 0.0.9. (Lemma 6.1.2) We have exact sequences

$$1. \quad 0 \to W_m \Omega^q_{X|D,\log} \to \mathrm{fil}_{-D} W_m \Omega^q_U \xrightarrow{1-\overline{F}} \frac{\mathrm{fil}_{-D} W_m \Omega^q_U}{dV^{m-1}(\mathrm{fil}_{-D} \Omega^{q-1}_U)},$$

2. 
$$0 \to j_*(W_m \Omega^q_{U,\log}) \to Z_1 W_m \Omega^q_X(\log E) \xrightarrow{1-C} W_m \Omega^q_X(\log E) \to 0.$$

The item (1) in the above lemma is already known by [31] if m = 1 or D is of the form  $p^{m-1}D'$ . Moreover, it is also right exact in the first case.

Next, we present some applications of the above results.

# First application: Ramified duality over finite fields with fixed modulus

We get the following refined version of Jannsen-Saito-Zhao's ramified duality for smooth schemes over finite fields. We write  $D_{\underline{n}} = \sum_{1 \leq i \leq r} n_i E_i$ , for  $\underline{n} = (n_1, \ldots, n_r) \in \mathbb{Z}^r$ . We say  $\underline{n} \geq \underline{0}$  if  $n_i \geq 0$  for all i.

**Theorem 0.0.10.** (Theorem 6.3.4, Corollary 6.3.6) Let X be a connected smooth projective variety over a finite field k and dim X = N. Then for all  $\underline{n} \ge \underline{0}$ , we have a perfect pairing of finite abelian groups

$$\mathbb{H}^{i}_{\acute{e}t}(X, W_m \mathcal{F}^{q, \bullet}_{\underline{D}_{\underline{n}}}) \times H^{N+1-i}_{\acute{e}t}(X, W_m \Omega^{N-q}_{X|\underline{D}_{\underline{n}+1}, \log}) \xrightarrow{\cup} \mathbb{Z}/p^m \tag{0.0.1}$$

$$H^{i}_{\acute{e}t}(X, j_{*}(W_{m}\Omega^{q}_{U, \log})) \times H^{N+1-i}_{\acute{e}t}(X, W_{m}\Omega^{N-q}_{X|E, \log}) \xrightarrow{\cup} \mathbb{Z}/p^{m}.$$

In view of Lemma 0.0.9(2), the second duality of the above theorem can be considered as the Milne's duality for the log scheme X, whose log structure is given by E (cf. [29]). Now by taking limits over <u>n</u> in the top pairing of (0.0.1), we recover the ramified duality of Jannsen-Saito-Zhao :

$$H^{i}_{\text{\acute{e}t}}(U, W_{m}\Omega^{q}_{U, \log}) \times \varprojlim_{D} H^{N+1-i}_{\text{\acute{e}t}}(X, W_{m}\Omega^{N-q}_{X|D, \log}) \to \mathbb{Z}/p^{m}.$$
(0.0.2)

### Second application: Duality for open curves over local fields

Now we proceed to our next main result which is the duality theorem for open curves over local fields of characteristic p > 0. Since there is no duality theorem for p-adic étale motivic cohomology for a smooth projective variety X over local fields in the literature, except dim X = 1 ([40, Prop 4], [47, Thm 4.7]), we consider only dim 1 case. In this case, the group  $H^i_{\acute{e}t}(X, W_m \Omega^q_{X|D_{n+1},\log})$  is not finite in general. Moreover, it might have non-discrete topology induced from the adic topology of the base field k. Considering these non trivial topologies, we have the following theorem.

**Theorem 0.0.11.** (Theorem 8.3.16) Let X be a smooth projective geometrically connected curve over a local field k of characteristic p > 0. Then for all  $\underline{n} \ge \underline{0}$ , we have perfect pairings of topological abelian groups

- $1. \qquad H^{i}_{\acute{e}t}(X, W_{m}\Omega^{2}_{X|D_{\underline{n}+1}, \log})^{\mathrm{pf}} \times \mathbb{H}^{2-i}_{\acute{e}t}(X, W_{m}\mathcal{F}^{0, \bullet}_{D_{\underline{n}}}) \to \mathbb{Z}/p^{m},$ where the group  $H^{i}_{\acute{e}t}(X, W_{m}\Omega^{2}_{X|D_{\underline{n}+1}, \log})$  is Hausdorff for all i (also profinite if  $i \neq 1$ ).
- 2.  $\begin{aligned} &H^{i}_{\acute{e}t}(X, W_{m}\Omega^{1}_{X|D_{\underline{n}+1}, \log}) \times \mathbb{H}^{2-i}_{\acute{e}t}(X, W_{m}\mathcal{F}^{1, \bullet}_{D_{\underline{n}}}) \to \mathbb{Z}/p^{m}, \ i \neq 1, \\ & \text{where } H^{i}_{\acute{e}t}(X, W_{m}\Omega^{1}_{X|D_{\underline{n}+1}, \log}) \text{ has profinite topology if } i = 0 \text{ and discrete topology if } i = 2, \end{aligned}$
- 3.  $H^{1}_{\acute{e}t}(X, W_{m}\Omega^{1}_{X|D_{n+1},\log})^{\wedge} \times \mathbb{H}^{1}_{\acute{e}t}(X, W_{m}\mathcal{F}^{1,\bullet}_{D_{\underline{n}}}) \to \mathbb{Z}/p^{m},$ where  $H^{1}_{\acute{e}t}(X, W_{m}\Omega^{1}_{X|D_{\underline{n}+1},\log})$  is an extension of a discrete group by an adic topological group and  $\mathbb{H}^{1}_{\acute{e}t}(X, W_{m}\mathcal{F}^{1,\bullet}_{D_{\underline{n}}})$  is an extension of a discrete group by a profinite group and  $(H^{1}_{\acute{e}t}(X, W_{m}\Omega^{1}_{X|D_{\underline{n}+1},\log}))^{\wedge}$  is the "discrete by profinite" completion of  $H^{1}_{\acute{e}t}(X, W_{m}\Omega^{1}_{X|D_{\underline{n}+1},\log}).$

Here, for a topological group G,  $G^{\text{pf}}$  means the topological profinite completion of G, (i.e.  $G^{\text{pf}} = \varprojlim_{\{U: \ [G:U] < \infty\}} G/U$ , where U is an open subgroup of G). Now, by taking limits over n, we have the following corollary.

**Corollary 0.0.12.** (Theorem 8.3.16(4)) Under the same hypothesis of Theorem 0.0.11, we have perfect pairing of topological abelian groups

1. 
$$\lim_{\underline{i}} H^{i}_{\acute{e}t}(X, W_{m}\Omega^{2}_{X|D_{\underline{n}+1}, \log})^{\mathrm{pf}} \times H^{2-i}_{\acute{e}t}(U, \mathbb{Z}/p^{m}) \to \mathbb{Z}/p^{m},$$

- $2. \qquad \lim_{\underline{i}} H^{i}_{\acute{e}t}(X, W_{m}\Omega^{1}_{X|D_{\underline{n}+1}, \log}) \times H^{2-i}_{\acute{e}t}(U, W_{m}\Omega^{1}_{U, \log}) \to \mathbb{Z}/p^{m}, \ i \neq 1,$
- 3.  $\lim_{\underline{h} \to \underline{h}} H^1_{\acute{e}t}(X, W_m \Omega^1_{X|D_{\underline{n}+1}, \log})^{\wedge} \times H^1_{\acute{e}t}(U, W_m \Omega^1_{U, \log}) \to \mathbb{Z}/p^m.$

We note that the groups in the pairing (1) and (2) above are zero if i = 0. As an application of the above theorem, we can deduce the *p*-primary part of the class field theory for *U*. This is proved in [48]. We also remark that the class field theory for *X* was established by Bloch [76], S. Saito [75], Yoshida [77] and the prime to *p*-part of the class field theory for *U* was established by T. Hiranouchi in [27].

## Third application: Lefschetz hypersurface theorem

Finally, we give one more application of the theory of filtered de Rham-Witt complex, namely, the Lefschetz hypersurface theorem.

**Theorem 0.0.13.** (Theorem 9.2.4) Let  $X \to \mathbb{P}_k^L$  be a smooth projective variety over an F-finite field k of dim N and D as before. Let Y be a smooth hypersurface section of X that intersects D transversally. Let  $D' = D \times_X Y$ . If Y is sufficiently ample (w.r.t D in (1) and (2)), then for all  $m \ge 1$ , the group homomorphisms

1. 
$$\mathbb{H}^{i}_{\acute{e}t}(X, W_m \mathcal{F}^{q, \bullet}_{D, U}) \to \mathbb{H}^{i}_{\acute{e}t}(Y, W_m \mathcal{F}^{q, \bullet}_{D', U'})$$

$$2. \qquad H^i_{\acute{e}t}(X, W_m \Omega^q_{X|D,\log}) \to H^i_{\acute{e}t}(Y, W_m \Omega^q_{Y|D',\log})$$

3. 
$$H^i_{\acute{e}t}(X, W_m \Omega^q_{X, \log}) \to H^i_{\acute{e}t}(Y, W_m \Omega^q_{Y, \log})$$

are isomorphisms if  $i + q \le N - 2$  and injective if i + q = N - 1.

Recall that the Lefschetz theorem for the étale fundamental groups of smooth projective schemes over a field was proven by Grothendieck (see [20, Expos'e XII, Corollaire 3.5]). Kerz-Saito [43] also proved a similar Lefschetz theorem for  $\pi_1^{ab}(X, D)$ , the ramified étale fundamental groups (with modulus) of smooth projective scheme X. To the best of our knowledge, a similar Lefschetz theorem for the Brauer groups is not known. As an application of the above theorem, we get the following Lefschetz theorem for  $\operatorname{Br}^{\operatorname{div}}(X|D)$ and  $\operatorname{Br}(X)$ .

**Theorem 0.0.14.** (Theorem 9.4.2) Let's assume  $N \ge 4$  in Theorem 0.0.13. Then, for sufficiently ample Y (w.r.t D in second case), we have

$$\operatorname{Br}(X)\{p\} \xrightarrow{\cong} \operatorname{Br}(Y)\{p\}, \quad \operatorname{Br}^{\operatorname{div}}(X|D)\{p\} \xrightarrow{\cong} \operatorname{Br}^{\operatorname{div}}(Y|D')\{p\}$$

Moreover, if k is either a finite field, local field, or algebraically closed field, then we have

$$\operatorname{Br}(X) \xrightarrow{\cong} \operatorname{Br}(Y), \quad \operatorname{Br}^{\operatorname{div}}(X|D) \xrightarrow{\cong} \operatorname{Br}^{\operatorname{div}}(Y|D').$$

As a corollary, we get the following result whose prime to p-part is already known by [12, Cor 5.5.4].

**Corollary 0.0.15.** (Corollary 9.4.9) Let k be any field and X be any smooth complete intersection in  $\mathbb{P}_k^N$  of dim d > 2. Then, we have  $\operatorname{Br}(k) \xrightarrow{\cong} \operatorname{Br}(X)$ .

## Further applications

The filtered de Rham-Witt complex studied here has further applications in understanding the log-crystalline cohomology of U, a topic that will be discussed in [50]. Additionally, the duality results established in this thesis can also be extended to curves over higher local fields. These duality results can, in turn, be used to prove class field theory for such curves, a topic that will be explored in [49].

## Organization of the thesis

The thesis contains nine chapters, which can be divided into four parts. We give an outline of the content here.

Part 1 is the Chapter 1, which recalls the preliminaries required in the later Chapters. We recall some basic properties of the de Rham-Witt complex and the logarithmic Hodge-Witt sheaves for regular schemes in Section 1.1. In Section 1.2, we discuss a few properties of the de Rham-Witt complex of Hyodo-Kato for log schemes whose log structure is given by a simple normal crossing divisor (SNCD).

Part 2 comprises Chapters 2, 3, and 4. Chapter 2 studies the basic definitions and properties of the theory of filtered de Rham-Witt complex. There we define the sheaf  $\operatorname{fil}_D W_m \Omega_U^q$ , where U is the complement of a simple normal crossing divisor E in a regular scheme X.

In Chapter 3, we study the structure of the above sheaf and prove various results when X is spectrum of a ring of the form  $S[[\pi]]$  ( $\pi$  is a variable).

In Chapter 4, we study the structure of the above sheaf for regular local rings. We also extend various results of the classical de Rham-Witt complex to the filtered setup when X is spectrum of an F-finite regular local ring, and finally to F-finite regular schemes.

Part 3 consists of Chapter 5, which establishes a cohomological description of Kato's local and global ramification filtration using the filtered de Rham-Witt complex studied

in Chapter 4. § 5.1 deals with the local Kato filtration and § 5.2 deals with the global Kato filtration.

Part 4 is composed of Chapters 6-9, which consist of applications of the concepts developed in Parts 2 and 3.

In Chapters 6-8, we prove duality theorems for logarithmic Hodge-Witt sheaves with modulus over finite fields and 1-local fields. In § 6.1, we prove a relation between Jannsen-Saito-Zhao's sheaf  $W_m \Omega_{X|D,\log}^q$  and  $\operatorname{fil}_{-D} W_m \Omega_U^q$ , define various complexes. In § 6.2, we define pairings between various complexes, which induce the pairings in the duality theorem. In § 6.3, we prove the duality theorems over finite fields and in § 8, we prove duality theorems over 1-local fields.

Finally, in Chapter 9, we prove a Lefschetz hypersurface theorem for the logarithmic Hodge-Witt cohomology with modulus. As an application, we prove a Lefschetz theorem for unramified Brauer group and ramified Brauer group (with modulus) in § 9.4.

## Chapter 1

## Preliminaries

## 1.1 de Rham-Witt Complex

In this section, we shall recall definitions of the de Rham-Witt complex and logarithmic Hodge-Witt sheaf for a regular scheme over a field of characteristic p > 0. We shall also review some basic properties of these sheaves, which are classically known in the case of smooth schemes over perfect fields. These properties were later extended to arbitrary regular schemes in [63] using the following two theorems.

**Theorem 1.1.1.** (cf. [60, Thm 2.5]) Let  $u : A \to A'$  be a regular morphism of noetherian rings. Then A' is a filtered inductive limit of smooth A algebras.

Recall that a morphism of schemes is called regular if it is flat and fibers are geometrically regular. For example the canonical map  $k[x_1, \ldots, x_n] \rightarrow k[[x_1, \ldots, x_n]]$  is regular. More generally, the map  $A \rightarrow \widehat{A}$  is regular for any excellent local ring. As a corollary of Theorem 1.1.1, we get

**Corollary 1.1.2.** Any regular local ring A of characteristic p is filtering inductive limit  $\varinjlim_{\lambda} A(\lambda)$  of finitely generated smooth  $\mathbb{F}_p$  algebras.

**Theorem 1.1.3.** (cf. [4, VII, Thm 5.7], [67, Thm 6.6]) Let  $\{X_i\}_{i\in I}$  be a filtering projective system of Noetherian schemes such that the transition maps are affine and  $\lim_{i \to i} X_i = X$  is noetherian. Let  $\{F_i\}_i$  be a compatible system of sheaves on  $\{X_{i,\tau}\}$  ( $\tau =$ zar or ét) with limit sheaf F on X. Then we have

$$H^n(X_{\tau}, F) \cong \varinjlim_i H^n(X_{i,\tau}, F_i).$$

Recall from [30], the de Rham-Witt complex  $W_m \Omega^{\bullet}_X$  over a scheme X of characteristic p is a strictly anti-commutative differential graded complex :

$$W_m \Omega^0_X = W_m \mathcal{O}_X \xrightarrow{d} W_m \Omega^1_X \xrightarrow{d} \cdots \xrightarrow{d} W_m \Omega^i_X \xrightarrow{d} \cdots$$

 $W_m \mathcal{O}_X$  is called the sheaf of Witt vectors of length m and  $W_m X = (|X|, W_m \mathcal{O}_X)$  be the corresponding Witt scheme. Let  $[-]_m : \mathcal{O}_X \to W_m \mathcal{O}_X$  be the multiplicative Teichmüllar function given by  $[x]_m = (x, 0, ..., 0)$  for any  $x \in \mathcal{O}_X$ . Let  $F : W_m \Omega_X^q \to W_{m-1} \Omega_X^q$ ,  $V : W_m \Omega_X^q \to W_{m+1} \Omega_X^q$  and  $R : W_m \Omega_X^q \to W_{m-1} \Omega_X^q$  be the Frobenius, Verschiebung and projection, respectively (cf. [30]). The homomorphisms F, V and R, satisfy the following properties.

- 1. FV = VF = p,
- 2. FdV = d, dF = pFd, pdV = Vd,
- 3.  $Fd[a]_m = [a]_m^{p-1}d[a]$ , for  $a \in \mathcal{O}_X$ ,
- 4. V(F(x)y) = xV(y), V(xdy) = VxdVy for  $x, y \in W_*\Omega_X^{\bullet}$ ,
- 5. Rd = dR, RV = VR, RF = FR.

Let  $H^q_{dR}(X)$  denote the q-th cohomology of the de Rham complex of X. Let  $C^{-1}: \Omega^q_X \to \underline{H}^q_{dR}(X)$  be the inverse Cartier homomorphism characterized by the following properties.

1.  $C^{-1}(x\omega) = x^p \omega$ , for  $x \in \mathcal{O}_X$ ,  $\omega \in \Omega_X^q$ ,

2. 
$$C^{-1}(dx_1\cdots dx_q) = (x_1\cdots x_q)^{p-1}dx_1\cdots dx_q$$
, for  $x_i \in \mathcal{O}_X$ .

We let  $Z_1\Omega_X^q = \operatorname{Ker}(d:\Omega_X^q \to \Omega_X^{q+1})$  and  $B_1\Omega_X^q = \operatorname{Image}(d:\Omega_X^{q-1} \to \Omega_X^q)$ .

**Proposition 1.1.4.** ([63, Prop 2.5]) If X is a regular scheme over  $\mathbb{F}_p$ , then  $C^{-1}$  is isomorphism.

*Proof.* If X is smooth scheme, then this is well known. To prove for general case, we may assume X is strictly Henselian local. Now the proposition follows from Theorem 1.1.1 and Theorem 1.1.3.

We write C for the composite map  $Z_1\Omega_X^q \twoheadrightarrow \underline{H}_{dR}^q(X) \xrightarrow{(C^{-1})^{-1}} \Omega_X^q$ . Assuming the  $\mathcal{O}_X$ modules  $Z_i\Omega_X^q$  and  $B_i\Omega_X^q$  are defined for some  $i \ge 1$ , we let  $Z_{i+1}\Omega_X^q$  (resp.  $B_{i+1}\Omega_X^q$ ) be the inverse image of  $Z_i\Omega_X^q$  (resp.  $B_i\Omega_X^q$ ) under the composite map  $Z_1\Omega_X^q \twoheadrightarrow H_{dR}^q(X) \xrightarrow{C} \Omega_X^q$ such that there is an induced isomorphism  $C^{-i}: \Omega_X^q \xrightarrow{\cong} Z_i\Omega_X^q/B_i\Omega_X^q$ , for  $i \ge 1$ . As a result, we get the following corollary.

#### 1.1. DE RHAM-WITT COMPLEX

Corollary 1.1.5. We have the exact sequences

$$0 \to Z_{m+1}\Omega_X^q \to Z_m\Omega_X^q \xrightarrow{dC^m} B_1\Omega_X^{q+1} \to 0, \qquad (1.1.1)$$

$$0 \to B_m \Omega_X^q \to B_{m+1} \Omega_X^q \xrightarrow{C^m} B_1 \Omega_X^q \to 0, \qquad (1.1.2)$$

for  $m \ge 1, q \ge 0$ .

*Proof.* Using the Cartier isomorphism, the proof is same as given in [51, Lem 1.14]. So we omit the details.  $\Box$ 

**Proposition 1.1.6.** (cf. [30]) Let X be a regular scheme over  $\mathbb{F}_p$ . Fix  $m \ge 0$  and  $q \ge 0$ . Then we have

1.  $0 \rightarrow V^n W_{m-n} \Omega_X^q + dV^n W_{m-n} \Omega_X^{q-1} \rightarrow W_m \Omega_X^q \xrightarrow{R^{m-n}} W_n \Omega_X^q \rightarrow 0$  is exact for  $m \ge n \ge 0$ ; 2. Ker  $(p^n : W_m \Omega_X^q \rightarrow W_m \Omega_X^q) = \text{Ker} (R^n : W_m \Omega_X^q \rightarrow W_{m-n} \Omega_X^q)$ , for  $n \ge 1$ ; 3. Ker  $(V^m : \Omega_X^q \rightarrow W_{m+1} \Omega_X^q) = B_m \Omega_X^q$ ; 4. Ker  $(V^m : \Omega_X^q \rightarrow W_{m+1} \Omega_X^q) = B_m \Omega_X^{q-1} = B_{m+1} \Omega_X^q = F^m dW_{m+1} \Omega_X^{q-1}$ ; 5. Ker  $(dV^m : \Omega_X^{q-1} \rightarrow W_{m+1} \Omega_X^q) = Z_{m+1} \Omega_X^{q-1}$ ; 6. Ker  $(dV^m : \Omega_X^{q-1} \rightarrow W_{m+1} \Omega_X^q) = Z_m \Omega_X^{q-1} = F^m W_{m+1} \Omega_X^{q-1}$ ; 7. Ker $(F^l : W_{m+l} \Omega_X^q \rightarrow W_m \Omega_X^q) = V^m (W_l \Omega_K^q)$  for all  $l \ge 1$ ; 8. Ker $(V : W_m \Omega_X^q \rightarrow W_{m+1} \Omega_X^q) = dV^{m-1} (\Omega_X^{q-1})$ .

*Proof.* All these statements are true if X is smooth. See [30, Propositions I.3.2, I.3.4] for (1) and (2), respectively. For (3)-(6), use Theorem I.3.8 of loc. cit. together with (I.3.21.1.4) and (I.3.21.1.3) of loc. cit. for the second equality of (4) and (6), respectively. For (7) and (8), use (I.3.21.1.2) and (I.3.21.1.4), respectively. For the general case, we may assume X is strictly local and reduces to the smooth case by Theorem 1.1.1 and Theorem 1.1.3.

**Definition 1.1.7.** We let  $R^q_{m-1,X} := \operatorname{Ker}\left((V^{m-1}, dV^{m-1}) : \Omega^q_X \oplus \Omega^{q-1}_X \to W_m \Omega^q_X\right)$ .

We note that  $R_{m-1,X}^q \subset B_m \Omega_X^q \oplus Z_{m-1} \Omega_X^{q-1}$  by Proposition 1.1.6(4) and (6).

**Corollary 1.1.8.** For a regular scheme X over  $\mathbb{F}_p$ , we have

$$0 \to R^q_{m-1,X} \to B_m \Omega^q_X \oplus Z_{m-1} \Omega^{q-1}_X \xrightarrow{C^{m-1}, \ dC^{m-1}} B_1 \Omega^q_X \to 0.$$

Proof. First note that  $C^{m-1}: B_m\Omega_X^q \to B_1\Omega_X^q$  (resp.  $dC^{m-1}: Z_{m-1}\Omega_X^{q-1} \to B_1\Omega_X^q$ ) is surjective by Corollary 1.1.5. Now if  $(a,b) \in \operatorname{Ker}(C^{m-1}, dC^{m-1})$ , by Proposition 1.1.6 (4) and (6), we can write  $(a,b) = (F^{m-1}da', F^{m-1}b')$ , for  $a',b' \in W_m\Omega_X^{q-1}$ , such that  $dR^{m-1}(a'+b') = 0$  (noting CF = R). Now  $V^{m-1}(a) + dV^{m-1}(b) = p^{m-1}d(a'+b') = 0$  by (2) of Proposition 1.1.6. This implies  $\operatorname{Ker}(C^{m-1}.dC^{m-1}) \subset R_{m-1,X}^q$ .

Now, for the other direction, take any  $(a, b) \in R^q_{m-1,X}$ . Then there exists  $b' \in Z_{m-1}\Omega^{q-1}_X$ such that  $(a, b) - (0, b') \in \operatorname{Ker}(C^{m-1}, dC^{m-1})$ . Since  $\operatorname{Ker}(C^{m-1}.dC^{m-1}) \subset R^q_{m-1,X}$ , we get that  $(0, b') \in R^q_{m-1,X}$ . Now by Proposition 1.1.6(5), we get  $b' \in Z_m \Omega^{q-1}_X$  and hence  $(0, b') \in \operatorname{Ker}(C^{m-1}.dC^{m-1})$  by (1.1.1). This implies  $(a, b) \in \operatorname{Ker}(C^{m-1}, dC^{m-1})$ . This finishes the proof.

Proposition 1.1.6 and the relations between F, d, V listed above immediately imply:

**Lemma 1.1.9.** The Frobenius map  $F: W_{m+1}\Omega_X^q \to W_m\Omega_X^q$  induces the homomorphism  $\overline{F}: W_m\Omega_X^q \to W_m\Omega_X^q / dV^{m-1}\Omega_X^{q-1}$ .

Now we recall the definition of logarithmic Hodge-Witt sheaf.

**Definition 1.1.10.** For a scheme X over  $\mathbb{F}_p$ , and  $m, q \ge 1$ , we define the logarithmic Hodge-Witt sheaf  $W_m \Omega^q_{X,\log,\tau}$  by  $\operatorname{Im}(\operatorname{dlog}: (\mathcal{O}_X^{\times})^{\otimes q} \to W_m \Omega^q_X)$ , where dlog is defined by

$$\operatorname{dlog}(a_1 \otimes \cdots \otimes a_q) = \operatorname{dlog}[a_1]_m \cdots \operatorname{dlog}[a_q]_m,$$

where the image is taken  $\tau = zar$ , nis or étale locally.

The above map induces a map dlog:  $\mathcal{K}_{i,X}^M/p^m \to W_m \Omega_{X,\log}^i$ ,  $\tau$ -locally, which is known to be an isomorphism when X is regular (cf. [58, Thm. 1.2, Cor. 4.2]).

The following proposition is known for smooth case and can be proved for regular case by using Theorems 1.1.1 and 1.1.3

**Proposition 1.1.11.** ([63, Prop 2.8, 2.10, 2.12]; cf. [30],[11, Lem 2,3]) Let X be a regular scheme over  $\mathbb{F}_p$ . Then we have the following exact sequences in  $X_{\acute{e}t}$ .

 $1. \quad 0 \to W_n \Omega_{X,\log}^q \xrightarrow{\underline{p}^m} W_{m+n} \Omega_{X,\log}^q \xrightarrow{R^n} W_m \Omega_{X,\log}^q \to 0.$   $2. \quad 0 \to W_m \Omega_{X,\log}^q \to W_m \Omega_X^q \xrightarrow{1-\overline{F}} W_m \Omega_X^q / dV^{m-1} \Omega_X^{q-1} \to 0.$   $3. \quad 0 \to W_m \Omega_{X,\log}^q \to Z_1 W_m \Omega_X^q \xrightarrow{1-C} W_m \Omega_X^q \to 0.$ 

## 1.2 The log de Rham-Witt complex

In this section, we shall recall the definitions and properties of log de Rham-Witt complex over a log scheme. The log structure, we shall be interested in is the one induced by a SNCD.

### **1.2.1** Definitions

We recall the definition of log structure on a scheme [39] (see also [64, Chap 2]).

**Definition 1.2.1.** Let Y be a scheme. A pre-log structure on Y is a pair  $(\mathcal{M}_Y, \alpha)$ , where  $\mathcal{M}_Y$  is a sheaf of commutative monoids on  $Y_{\text{\acute{e}t}}$  (or  $Y_{\text{zar}}$ ) and  $\alpha : \mathcal{M}_Y \to (\mathcal{O}_Y, \cdot)$  is a morphism of sheaf of monoids on Y, with respect to multiplication on  $\mathcal{O}_Y$ . The pre-log structure  $(\mathcal{M}_Y, \alpha)$  is called a log structure if  $\alpha^{-1}(\mathcal{O}_Y^{\times}) \xrightarrow{\alpha} \mathcal{O}_Y^{\times}$  is an isomorphism. A log scheme is scheme with a log structure on it.

**Definition 1.2.2.** Let  $(\mathcal{M}_Y, \alpha)$  be a pre-log structure on Y. Let  $\mathcal{M}_Y^a$  be the push-out of the following diagram in the category of sheaves of monoid on Y.

$$\mathcal{O}_Y^{\times} \stackrel{\alpha}{\leftarrow} \alpha^{-1}(\mathcal{O}_Y^{\times}) \to \mathcal{M}_Y.$$

That is,  $\mathcal{M}_Y^a = (\mathcal{O}_Y^{\times} \bigoplus \mathcal{M}_Y)/\sim$ , where, locally,  $(a,b) \sim (a',b')$  if there exists  $c_1, c_2 \in \alpha^{-1}(\mathcal{O}_Y^{\times})$  such that  $a\alpha(c_1) = a'\alpha(c_2)$  and  $bc_2 = b'c_1$ . One can check  $(\mathcal{M}_Y^a, \alpha^a)$  gives a log structure on Y, where  $\alpha^a : \mathcal{M}_Y^a \to \mathcal{O}_Y$  is given by  $\alpha^a(u,v) = u\alpha(v)$ .  $(\mathcal{M}_Y^a, \alpha^a)$  is called the log structure associated to  $(\mathcal{M}_Y, \alpha)$ .

Given a pre-log structure  $(\mathcal{M}_Y, \alpha)$  on Y, we shall denote the associated log scheme as  $(Y, \alpha : \mathcal{M}_Y \to \mathcal{O}_Y)$ , if no confusion is possible.

A morphism between two log schemes  $(Y, \alpha : \mathcal{M}_Y \to \mathcal{O}_Y)$  and  $(Y', \alpha' : \mathcal{M}'_{Y'} \to \mathcal{O}_{Y'})$ is a morphism of schemes  $f : Y \to Y'$  and a homomorphism  $h : f^{-1}(\mathcal{M}'_{Y'}) \to \mathcal{M}_Y$ , such that the following diagram commutes.

$$\begin{array}{cccc}
f^{-1}(\mathcal{M}'_{Y'}) \xrightarrow{h} & \mathcal{M}_{Y} \\
f^{-1}\alpha' \downarrow & \downarrow \alpha \\
f^{-1}(\mathcal{O}_{Y'}) \longrightarrow \mathcal{O}_{Y}.
\end{array} (1.2.1)$$

**Definition 1.2.3.** Let  $f: Y_1 \to Y_2$  be a morphism of schemes and  $(\mathcal{M}_{Y_2}, \alpha)$  be a log structure on  $Y_2$ . Then the pull back log structure  $f^*\mathcal{M}_{Y_2}$  on  $Y_1$  is defined to be the log structure associated to the pre-log structure  $f^{-1}\mathcal{M}_{Y_2} \xrightarrow{f^{-1}(\alpha)} f^{-1}\mathcal{O}_{Y_2} \to \mathcal{O}_{Y_1}$  on  $Y_1$ .

At this point, we refer the reader to [39] (also [64, Chap 2]) for a comprehensive discussion on log schemes and the definitions of various types of morphisms (e.g. integral, saturated, smooth, Cartier type etc).

Now we recall the definition of log Witt complex associated to a log structure (cf. [14, § 1], [28, § 3]). Let  $A_{\text{log}}$  denote a log ring  $(A, \alpha : M_A \to (A, \cdot))$  of characteristic P > 0 (i.e. char A = p). This induces a log structure  $(M_A, \alpha_m : M_A \to W_m(A))$  on  $W_m(A)$ , where  $\alpha_m := [\cdot]_m \circ \alpha$  and  $[\cdot]_m : A \to W_m(A)$  is the Teichmüller map.

**Definition 1.2.4.** A log differential graded ring  $(E^*, M_E)$  over a log ring  $(E^0, M_E, \alpha : M_E \to E^0)$  is a differential graded ring  $E^*$ , together with a map of monoids, dlog :  $M_E \to (E^1, +)$  such that  $d \circ dlog = 0$  and  $d\alpha(a) = \alpha(a) dlog(a)$  for all  $a \in M_E$ .

For any log ring  $A_{\text{log}}$ , a universal log differential graded ring always exists (see [28, Section 2.2, P 33]). This is called the de Rham complex of the log ring  $A_{\text{log}}$  and denoted by  $\Omega^{\star}_{A_{\text{log}}}$ .

**Definition 1.2.5.** A log Witt complex over  $A_{\log}$  consists of :

- 1. a pro-log differential graded ring  $(E^*, M_E)$  and a strict map of pro-log rings  $\lambda : (W_{\cdot}(A), M_A) \to (E^0_{\cdot}, M_E);$
- 2. a strict map of pro-log graded rings

$$F: E_m^\star \to E_{m-1}^\star,$$

such that  $\lambda F = F\lambda$ ,  $F \operatorname{dlog}_m \lambda(a) = \operatorname{dlog}_{m-1}\lambda(a)$  for all  $a \in M_A$  and  $Fd\lambda([a]_m) = \lambda([a]_{m-1})^{p-1}d\lambda([a]_{m-1})$  for all  $a \in A$ ; where  $\operatorname{dlog}_m : M_E \to E^1_{m+1}$ ;

3. and a strict map of pro graded modules over the pro graded rings  $E^*$ .

$$V: F_{\star}E_m^{\star} \to E_{m+1}^{\star}$$

such that  $V\lambda = \lambda V$ , FV = p and FdV = d, where  $F_{\star}E_m^{\star}$  indicates that  $E_m^{\star}$  is considered as  $E_{m+1}^0$ -module via  $F: E_{m+1}^0 \to E_m^0$ .

We write R for the structure map of the pro-system  $E^*$  and call it the restriction map. For any log ring  $A_{\log}$ , a universal log Witt complex always exists (cf. [28, Prop. 3.2.2]). It is called the de Rham-Witt complex of the log ring  $A_{\log}$  and denoted by  $W_{\cdot}\Omega^*_{A_{\log}}$ . Furthermore,  $W_1\Omega^{\bullet}_{A_{\log}}$  coincides with the de Rham complex  $\Omega^{\bullet}_{A_{\log}}$  of the log ring  $A_{\log}$ . One can also extend these definitions and results to any log scheme Y of characteristic p in a natural way.

#### 1.2. THE LOG DE RHAM-WITT COMPLEX

### 1.2.2 Log structure associated to SNCD

Let X be a connected regular Noetherian scheme of characteristic p > 0. We let K denote the function field of X. Let  $E \subset X$  be a simple normal crossing divisor (SNCD) with irreducible components  $E_1, \ldots, E_r$ . Recall that this means that  $E_i$  is a regular divisor on X for each i and the scheme theoretic intersection  $E_{i_1} \cap \cdots \cap E_{i_s}$   $(1 \le i_1 < \cdots < i_s \le r)$  is regular of codimension s in X. Let  $j: U \hookrightarrow X$  be the inclusion of the open complement of E.

We let  $X_{\log}$  denote the log scheme  $(X, \mathcal{M}_X, \alpha_X)$ , where  $\alpha_X \colon \mathcal{M}_X = \mathcal{O}_X \cap j_* \mathcal{O}_U^{\times} \hookrightarrow \mathcal{O}_X$  is the usual inclusion. Here the intersection is taken inside the constant sheaf given by K, the function field of X. For  $m \ge 1$ , let  $W_m(X)_{\log}$  be the log scheme structure on  $W_m(X)$ associated to the pre-log structure  $(\mathcal{M}_X, \alpha_m \colon \mathcal{M}_X \to W_m \mathcal{O}_X)$ , where  $\alpha_m \coloneqq [\cdot]_m \circ \alpha$  and  $[\cdot]_m \colon \mathcal{O}_X \to W_m \mathcal{O}_X$  is the Teichmüller map.

We let  $W_m \Omega^{\bullet}_{X_{\log}}$  denote the de Rham-Witt complex of the log scheme  $X_{\log}$ . Note that  $W_1 \Omega^{\bullet}_{X_{\log}}$  coincides with the de Rham complex  $\Omega^{\bullet}_{X_{\log}}$  of the log scheme  $X_{\log}$ . When X = Spec(A), we shall write  $W_m \Omega^q_{X_{\log}}$  as  $W_m \Omega^q_{A_{\log}}$ , for  $q \ge 0$ .

Recall that the canonical map of de Rham-Witt complexes  $W_m \Omega_X^{\bullet} \to j_*(W_m \Omega_U^{\bullet})$ is injective (because  $W_m \Omega_X^{\bullet} \to \eta_*(W_m \Omega_K^{\bullet})$  is injective, where  $\eta$ : Spec  $(K) \to X$  is the inclusion (cf. [46, Prop. 2.8])). This fact will be implicitly used throughout this thesis.

If  $X = \operatorname{Spec}(A)$ , where A is a regular local  $\mathbb{F}_p$ -algebra and  $E_i = \operatorname{Spec}(A/(x_i))$ , we let  $\pi = \prod_{i=1}^r x_i$  and let  $A_{\pi} = A[\pi^{-1}]$ . We let  $S_0 = \{0\}$  and for  $1 \leq j \leq r$ , let  $S_j$  be the collection of ordered subsets  $J = \{i_1 < \cdots < i_j\}$  of  $\{1, \ldots, r\}$ . For  $J = \{i_1 < \cdots < i_j\} \in S_j$ , we let  $W_m \Omega_A^q(\log \pi)_J = W_m \Omega_A^{q-j} \operatorname{dlog}[x_{i_1}]_m \wedge \cdots \wedge \operatorname{dlog}[x_{i_j}]_m$  as a  $W_m(A)$ -submodule of  $W_m \Omega_{A_{\pi}}^q$ . For  $J = S_0$ , we let  $W_m \Omega_A^q(\log \pi)_J = W_m \Omega_A^q$ . We let  $W_m \Omega_X^q(\log E)$  be the image  $W_m \Omega_A^q(\log \pi)$  of the canonical map  $\bigoplus_{j=0}^{\min\{q,r\}} \bigoplus_{J \in S_j} W_m \Omega_A^q(\log \pi)_J \to W_m \Omega_{A_{\pi}}^q$ . If X is not necessarily local, we let  $W_m \Omega_X^q(\log E)$  be the Zariski subsheaf of  $j_* W_m \Omega_U^q$  whose stalk at a point  $x \in X$  is  $W_m \Omega_{X_x}^q(\log E_x)$ , where  $X_x = \operatorname{Spec}(\mathcal{O}_{X,x})$  and  $E_x = X_x \times_X E$ .

Purpose of this section is to identify  $W_{\Omega_{X_{\log}}}^{\bullet}$  with  $W_{\Omega_X}^{\bullet}(\log E)$  (cf. Theorem 1.2.13).

**Lemma 1.2.6.**  $W_{\cdot}\Omega_{X}^{\bullet}(\log E)$  is a log Witt complex over  $X_{\log}$ . In particular, there is a canonical (strict) map of log Witt complexes  $\theta_{X}: W_{\cdot}\Omega_{X_{\log}}^{\bullet} \to W_{\cdot}\Omega_{X}^{\bullet}(\log E)$ .

Proof. To endow  $W_{\cdot}\Omega_{X}^{\bullet}(\log E)$  with the structure of a log Witt complex over  $X_{\log}$ , we let dlog  $\circ$   $[\cdot]_{m}: \mathcal{M}_{X} \to W_{m}\Omega_{X}^{1}(\log E)$  be the map locally defined by  $a \mapsto \operatorname{dlog}[a]_{m}$ . We take  $\lambda: (W_{\cdot}\mathcal{O}_{X}, \mathcal{M}_{X}) \to (W_{\cdot}\Omega_{X}^{0}(\log E), \mathcal{M}_{X})$  to be the identity map. Since  $j_{*}W_{\cdot}\Omega_{U}^{\bullet}$  is a Witt complex over X and  $W_{\cdot}\Omega_{X}^{\bullet}(\log E)$  is its subcomplex, we only need to verify that the latter is preserved by the Frobenius and Verchiebung operators of  $j_{*}W_{\cdot}\Omega_{U}^{\bullet}$ . Since F is multiplicative and  $F(\operatorname{dlog}[a]_m) = \operatorname{dlog}[a]_{m-1}$ , the claim is clear for F. Since  $V(w\operatorname{dlog}[a]_m) = V(wF(\operatorname{dlog}[a]_{m+1})) = V(w)\operatorname{dlog}[a]_{m+1}$ , we see that V also preserves  $W_{\cdot}\Omega^{\bullet}_X(\log E)$ .

Recall that an  $\mathbb{F}_p$ -scheme Y is called F-finite if the absolute Frobenius map  $F: Y \to Y$  is a finite morphism. In the sequel, we shall use that every essentially of finite type scheme over an F-finite field of characteristic p as well as its Henselizations and completions is F-finite (cf. [58, § 1.1]).

**Lemma 1.2.7.** Let A be a regular local F-finite  $\mathbb{F}_p$ -algebra of dim N with maximal ideal  $\mathfrak{m} = (x_1, \ldots, x_N)$  and residue field  $\mathfrak{f}$ . Then  $\Omega_A^1$  is free A-module of finite rank. Moreover,  $\{dx_1, \ldots, dx_N\}$  is a part of basis of  $\Omega_A^1$ .

*Proof.* Since A is regular and F-finite,  $\Omega_A^1$  is free (by [74, Lem 2.1.2]) and finitely generated over A (by [66, Prop. 1], taking  $k = \mathbb{F}_p$ ). Now consider the exact sequence

$$0 \to \mathfrak{m}/\mathfrak{m}^2 \xrightarrow{d \otimes 1} \Omega^1_A \otimes_A \mathfrak{f} \to \Omega^1_{\mathfrak{f}} \to 0.$$
 (1.2.2)

Here,  $d \otimes 1$  is injective because  $\mathfrak{f}$  is 0-smooth (separable) over  $\mathbb{F}_p$  (see [53, Thm 25.2, 26.9]). Also, it follows from (1.2.2) that  $\Omega_{\mathfrak{f}}^1$  has finite rank. Let  $y_1, \ldots, y_s \in A$  such that image of  $dy_1, \ldots, dy_s$  forms a basis of  $\Omega_{\mathfrak{f}}^1$ . Then by (1.2.2), we get that  $d\overline{x}_1, \ldots, d\overline{x}_N, d\overline{y}_1, \ldots, d\overline{y}_s$  forms a basis of  $\Omega_A^1 \otimes_A \mathfrak{f}$ . Since  $\Omega_A^1$  is finitely generated flat A module, by Nakayama lemma,  $dx_1, \ldots, dx_N, dy_1, \ldots, dy_s$  forms a basis of  $\Omega_A^1$ .

**Lemma 1.2.8.** Let A be as in Lemma 1.2.7. Let  $\pi = x_1 \cdots x_r$ ,  $r \leq N$ . Let  $A_{\log}$  be the log ring whose log structure on A is induced by  $\pi$ . Then the canonical maps of log differential graded A-algebras :  $\Omega^{\bullet}_{A_{\log}} \xrightarrow{\theta_A} \Omega^{\bullet}_A(\log \pi) \xleftarrow{\psi_A} \Lambda^{\bullet}_A(\Omega^1_A(\log \pi))$  are isomorphisms. Moreover, each  $\Omega^q_A(\log \pi)$  is a free A-module of finite rank.

*Proof.* We continue with the same notations of the proof of Lemma 1.2.7. We let  $\widetilde{\Omega}^1_A(\log \pi) = (\underset{1 \leq j \leq s}{\oplus} Ady_j) \oplus (\underset{1 \leq i \leq r}{\oplus} Ad\log(x_i)) \oplus (\underset{r+1 \leq i \leq N}{\oplus} Adx_i)$ . Since  $\Omega^{\bullet}_{A_{\log}}$  is a DG (differential graded) complex over A, there is a unique morphism of differential graded A-algebras  $\phi_A: \Omega^{\bullet}_A \to \Omega^{\bullet}_{A_{\log}}$ . As  $\Omega^{\bullet}_{A_{\log}}$  has a log DG complex structure, we also have a commutative diagram of monoids

$$\begin{array}{c}
M_A \xrightarrow{\text{dlog}} \Omega^1_{A_{\text{log}}} \\
\underset{\text{dlog}}{\swarrow} \downarrow_{\text{can}} \\
\Omega^1_{A_{\pi}},
\end{array} (1.2.3)$$

where  $M_A = \{ux_1^{i_1} \cdots x_r^{i_r} | u \in A^{\times} \text{ and } i_j \in \mathbb{N} \ \forall \ 1 \leq j \leq r\}.$ 

### 1.2. THE LOG DE RHAM-WITT COMPLEX

Since  $\widetilde{\Omega}_{A}^{1}$  is a free A-module, we have a well-defined A-linear map  $\theta'_{A}: \widetilde{\Omega}_{A}^{1}(\log \pi) \to \Omega_{A_{\log}}^{1}$  such that  $\theta'_{A}(ad\log(x_{i})) = ad\log(x_{i})$  for all  $a \in A$ . Hence, we get a morphism of differential graded A-algebras  $\theta'_{A}: \Lambda_{A}^{\bullet}(\widetilde{\Omega}_{A}^{1}(\log \pi)) \to \Omega_{A_{\log}}^{\bullet}$ . It is clear that there is a monoid homomorphism dlog:  $M_{A} \to \widetilde{\Omega}_{A}^{1}(\log \pi)$  which is compatible with  $\theta'_{A}$ . In particular, the latter is a morphism of log differential graded A-algebras which is identity in degree zero. We conclude from the universal property of  $\Omega_{A_{\log}}^{\bullet}$  that

$$\theta'_A: \Lambda^{\bullet}_A(\widetilde{\Omega}^1_A(\log \pi)) \xrightarrow{\cong} \Omega^{\bullet}_{A_{\log}}$$

On the other hand, as  $\widetilde{\Omega}^1_A(\log \pi)$  is a free A-module, the canonical map

$$\Lambda^{\bullet}_{A}(\widetilde{\Omega}^{1}_{A}(\log \pi)) \to \Lambda^{\bullet}_{A}(\widetilde{\Omega}^{1}_{A}(\log \pi)) \otimes_{A} A_{\pi} = \Lambda^{\bullet}_{A}(\Omega^{1}_{A_{\pi}}) = \Omega^{\bullet}_{A_{\pi}}$$

is injective and its image lies in  $\Omega^{\bullet}_{A}(\log \pi)$ . We thus get a commutative diagram

$$\Lambda^{\bullet}_{A}(\widetilde{\Omega}^{1}_{A}(\log\pi)) \xrightarrow{\widetilde{\psi}^{\bullet}_{A}} \Omega^{\bullet}_{A}(\log\pi)$$

$$(1.2.4)$$

$$\Omega^{\bullet}_{A\pi}.$$

It is clear from the definition of  $\Omega_A^{\bullet}(\log \pi)$  and Lemma 1.2.7 that  $\widetilde{\psi}_A^{\bullet}$  is surjective and hence it is an isomorphism. This implies second part of the lemma. In particular  $\widetilde{\psi}_A^1 : \widetilde{\Omega}_A^1(\log \pi) \to \Omega_A^1(\log \pi)$  is isomorphism. This forces  $\psi_A : \Lambda_A^{\bullet}(\Omega_A^1(\log \pi)) \to \Omega_A^{\bullet}(\log \pi)$ to be isomorphism. Also we have,

$$\theta_A = \widetilde{\psi}_A^{\bullet} o(\theta_A')^{-1} : \Omega_{A_{\log}}^{\bullet} \xrightarrow{\cong} \Omega_A^{\bullet}(\log \pi).$$

This achieves the first part of the lemma.

We let A be as in Lemma 1.2.8 and let  $\mathbb{F}_p^{[n]} = \mathbb{F}_p[X_1, \ldots, X_n]$ , for  $n \in \mathbb{N}$ . There are homomorphisms of log rings

$$\mathbb{F}_{p,\log} \to (\mathbb{F}_p^{[N]})_{\log} \xrightarrow{\gamma} A_{\log}, \text{ where } \gamma(X_i) = x_i \ \forall \ 1 \le i \le N.$$
(1.2.5)

The log structure of  $\mathbb{F}_p$  is trivial while that of  $\mathbb{F}_p^{[N]}$  (resp. A) is given by the simple normal crossing divisor V((X)) (resp.  $V((\pi))$ ) on  $\mathbb{A}_{\mathbb{F}_p}^N$  (resp. Spec (A)), where  $X = X_1 \cdots X_r$ . The first arrow in (1.2.5) is log smooth while the second arrow is a morphism of log rings such that the underline morphism of rings is regular. To see the latter claim, first note that A is formally **m**-smooth over  $(\mathbb{F}_p^{[N]})_{(X_1,\cdots,X_N)}$ , as  $\mathbb{F}_p$  is perfect and A is regular (cf. [65, Lem. 1.3]). It follows now from André's theorem (cf. [3]) that  $\mathbb{F}_p^{[N]} \to A$  is regular. We let  $H^i_{dR}(A_{\log})$  denote the *i*-th cohomology of the log de Rham complex  $\Omega^{\bullet}_{A_{log}}$ .

**Lemma 1.2.9.** For  $q \ge 0$ , there is an isomorphism of A-modules  $C_q^{-1}: \Omega_{A_{\log}}^q \xrightarrow{\cong} H^q_{dR}(A_{\log})$ .

Proof. Since the map  $\gamma: \mathbb{F}_p^{[N]} \to A$  is regular, Theorem 1.1.1 implies that there is a direct system  $\{A(\lambda)\}$  of smooth (in particular, finite type)  $\mathbb{F}_p^{[N]}$ -algebras such that  $A = \lim_{\lambda \to A} A(\lambda)$ . We endow each  $A(\lambda)$  with the log structure given by the pull-back of the log structure of  $(\mathbb{F}_p^{[N]})_{\log}$ . Since the log structure of A also has this property, it follows that we have a direct system of log smooth  $(\mathbb{F}_p^{[N]})_{\log}$ -algebras  $\{A(\lambda)_{\log}\}$  such that  $A_{\log} = \lim_{\lambda \to A} A(\lambda)_{\log}$  in the category of log rings. As  $\mathbb{F}_{p,\log} \to (\mathbb{F}_p^{[N]})_{\log}$  is log smooth, we get that each  $A(\lambda)_{\log}$  is log smooth over  $\mathbb{F}_{p,\log}$ .

Since the homomorphism of log rings  $\mathbb{F}_{p,\log} \to (\mathbb{F}_p^{[N]})_{\log}$  is clearly integral, and so is the homomorphism  $(\mathbb{F}_p^{[N]})_{\log} \to A(\lambda)_{\log}$  by the definition of latter's log structure ([39, Cor 4.4(i)]), it follows that  $\mathbb{F}_{p,\log} \to A(\lambda)_{\log}$  is integral. By a similar reason, this map is also saturated. It follows from Kato's criterion that each  $\mathbb{F}_{p,\log} \to A(\lambda)_{\log}$  is of Cartier type (cf. [29, § 2.12]). We conclude from loc. cit. that there is an isomorphism  $C_q^{-1}: \Omega_{A(\lambda)_{\log}}^q \xrightarrow{\cong} H^q_{dR}(A(\lambda)_{\log})$  given by  $C_q^{-1}(ad\log(b_1) \wedge \cdots \wedge d\log(b_q)) = a^p d\log(b_1) \wedge \cdots \wedge d\log(b_q)$ . Since this is clearly natural with respect to the homomorphisms of log rings, we get after passing to the limit that there is a similar inverse Cartier isomorphism for A as well. It is clear from the description that  $C_q^{-1}$  is A-linear if we endow  $\operatorname{Ker}(d)$  with the A-module structure via latter's Frobenius endomorphism.

We let  $Z_1\Omega_{A_{\log}}^q = \operatorname{Ker}(d:\Omega_{A_{\log}}^q \to \Omega_{A_{\log}}^{q+1})$  and  $B_1\Omega_{A_{\log}}^q = \operatorname{Image}(d:\Omega_{A_{\log}}^{q-1} \to \Omega_{A_{\log}}^q)$ . Taking the inverse of  $C_{\bullet}^{-1}$ , we get the following.

**Corollary 1.2.10.** For  $q \ge 0$ , there is an A-linear Cartier homomorphism  $C_q: Z_1 \Omega_{A_{\log}}^q \to \Omega_{A_{\log}}^q$  which induces an isomorphism of A-modules  $C_q: H^q_{dR}(A_{\log}) \xrightarrow{\cong} \Omega^q_{A_{\log}}$ .

Now  $Z_i \Omega^q_{A_{\log}}$  and  $B_i \Omega^q_{A_{\log}}$  can be defined like the usual case of regular schemes (cf. Section 1.1). We thus get a filtration

$$0 = B_0 \Omega_{A_{\log}}^q \subseteq B_1 \Omega_{A_{\log}}^q \subseteq \dots \subseteq \bigcup_i B_i \Omega_{A_{\log}}^q \subseteq \bigcap_i Z_i \Omega_{A_{\log}}^q \subseteq \dots \subseteq Z_1 \Omega_{A_{\log}}^q \subseteq Z_0 \Omega_{A_{\log}}^q = \Omega_{A_{\log}}^q \subseteq \Omega_{A_{\log}^q}^q \subseteq \Omega_{A_{\log}}^q \subseteq \Omega_{A_{\log}^q}^q \subseteq \Omega_{A_{\log}^q}^$$

of  $\Omega_{A_{\log}}^q$  together with an isomorphism  $C_q^{-i}: \Omega_{A_{\log}}^q \xrightarrow{\cong} Z_i \Omega_{A_{\log}}^q / B_i \Omega_{A_{\log}}^q$  for all  $i \ge 0$ . Using this isomorphism, proof of the following corollary is analogous to that of Corollary 1.1.5.

Corollary 1.2.11. We have the exact sequences

$$0 \to Z_{m+1}\Omega^q_{A_{\log}} \to Z_m\Omega^q_{A_{\log}} \xrightarrow{dC^m} B_1\Omega^{q+1}_{A_{\log}} \to 0, \qquad (1.2.6)$$

$$0 \to B_m \Omega^q_{A_{\log}} \to B_{m+1} \Omega^q_{A_{\log}} \xrightarrow{C^m} B_1 \Omega^q_{A_{\log}} \to 0, \qquad (1.2.7)$$

for  $m \ge 1, q \ge 0$ .

#### 1.2. THE LOG DE RHAM-WITT COMPLEX

**Lemma 1.2.12.** For every  $m \ge 1$ , the canonical map  $W_m \Omega^q_{A_{lor}} \to W_m \Omega^q_{A_{\pi}}$  is injective.

Proof. We shall prove the lemma by induction on m. The base case m = 1 follows from Lemma 1.2.8. For  $q, r \ge 0$  and  $m \ge 1$ , we let  $\operatorname{fil}_V^r W_m \Omega_{A_{\log}}^q = V^r (W_{m-r} \Omega_{A_{\log}}^q) + dV^r (W_{m-r} \Omega_{A_{\log}}^{q-1}) \subset W_m \Omega_{A_{\log}}^q$ . This yields the (decreasing) V-filtration on  $W_m \Omega_{A_{\log}}^q$  as a  $W_m(A)$ -module. Furthermore, we have an exact sequence (cf. [28, Lem. 3.2.4])

$$0 \to \operatorname{fil}_V^r W_m \Omega_{A_{\log}}^q \to W_m \Omega_{A_{\log}}^q \xrightarrow{R^{m-r}} W_r \Omega_{A_{\log}}^q \to 0, \qquad (1.2.8)$$

which is natural in  $A_{\log}$ . We let  $R_r^q(A_{\log}) = \text{Ker}((V^r, dV^r) : \Omega_{A_{\log}}^q \oplus \Omega_{A_{\log}}^{q-1} \to W_{r+1}\Omega_{A_{\log}}^q)$ . By comparing the above exact sequence with the analogous exact sequence for  $A_{\pi}$  and for  $m \ge 2, r = m - 1$ , and using induction on m, it suffices to show that the square

is Cartesian.

Suppose now that there is an element  $\alpha \in (\Omega_{A_{\log}}^q \oplus \Omega_{A_{\log}}^{q-1}) \cap R_{m-1}^q(A_{\pi}) \subset (\Omega_{A_{\pi}}^q \oplus \Omega_{A_{\pi}}^{q-1}).$ Then  $\alpha \in B_m \Omega_{A_{\pi}}^q \oplus Z_{m-1} \Omega_{A_{\pi}}^{q-1}$  by Corollary 1.1.8. We write  $\alpha = (\alpha_0, \alpha_1)$  with  $\alpha_0 \in B_m \Omega_{A_{\pi}}^q \cap \Omega_{A_{\log}}^q$  and  $\alpha_1 \in Z_{m-1} \Omega_{A_{\pi}}^{q-1} \cap \Omega_{A_{\log}}^{q-1}$ . We claim that  $\alpha \in B_m \Omega_{A_{\log}}^1 \oplus Z_{m-1} \Omega_{A_{\log}}^1$ .

To prove the claim, we first note that  $B_1\Omega_{A_{\log}}^q \subset \Omega_{A_{\log}}^q \hookrightarrow \Omega_{A_{\pi}}^q$  by Lemma 1.2.8, and this implies that the map  $B_1\Omega_{A_{\log}}^q \to B_1\Omega_{A_{\pi}}^q$  is injective. We next note that as  $\Omega_{A_{\log}}^{\bullet} \to \Omega_{A_{\pi}}^{\bullet}$ , our assumption already implies that  $\alpha \in Z_1\Omega_{A_{\log}}^q \oplus Z_1\Omega_{A_{\log}}^{q-1}$ . Using (1.1.1), (1.2.6) and induction on m, we conclude that  $\alpha_1 \in Z_{m-1}\Omega_{A_{\log}}^1$ . Since  $\alpha_0 \in B_m\Omega_{A_{\pi}}^q \cap \Omega_{A_{\log}}^q \subset Z_i\Omega_{A_{\pi}}^q$ for every  $i \ge 1$ , the same argument shows that  $\alpha_0 \in Z_m\Omega_{A_{\log}}^q$ . Using the commutative diagram of short exact sequences

we conclude that  $\alpha_0 \in B_m \Omega^q_{A_{\log}}$ . This proves the claim.

We now use [51, (1.15.3)] (with a limit argument as in Lemma 1.2.9) and Corollary 1.1.8, which give us a commutative diagram of short exact sequences

Since the right vertical arrow is injective, we conclude from the above claim and a diagram chase that  $\alpha \in R^q_{m-1}(A_{\log})$ . This proves that (1.2.9) is Cartesian and completes the proof of the lemma.

We can now prove the main result of this section.

**Theorem 1.2.13.** Let X be a regular F-finite  $\mathbb{F}_p$ -scheme and let  $E \subset X$  be a simple normal crossing divisor. Then the canonical strict map of log Witt complexes  $\theta_X: W_{\cdot}\Omega^{\bullet}_{X_{\log}} \to W_{\cdot}\Omega^{\bullet}(\log E)$  is an isomorphism.

*Proof.* We can assume X = Spec(A) with A as in Lemma 1.2.8. We now look at the commutative diagram of log Witt complexes over  $A_{\text{log}}$ :



The diagonal arrow on the right is injective by Lemma 1.2.12. It is clear from the definition of  $W_m \Omega^{\bullet}_A(\log \pi)$  that it lies in  $W_m \Omega^{\bullet}_{A_{\log}}$  as a log Witt subcomplex of  $W_m \Omega^{\bullet}_{A_{\pi}}$ . In particular, the inclusion  $W_m \Omega^{\bullet}_A(\log \pi) \hookrightarrow W_m \Omega^{\bullet}_{A_{\pi}}$  uniquely factors through an inclusion of log Witt complexes  $\psi_A: W_m \Omega^{\bullet}_A(\log \pi) \to W_m \Omega^{\bullet}_{A_{\log}}$ . Since  $W_m \Omega^{\bullet}_{A_{\log}}$  is the universal log Witt-complex over  $A_{\log}$ , this forces  $\psi_A$  and  $\theta_A$  to be isomorphisms and inverses of each other.

**Corollary 1.2.14.** Let X be as in Theorem 1.2.13. Then for every  $q, r, s \ge 0$  with  $r + s = m \ge 1$ , there is a short exact sequence of  $W_m \mathcal{O}_X$ -modules

$$0 \to V^r(W_s\Omega_X^q(\log E)) + dV^r(W_s\Omega_X^{q-1}(\log E)) \to W_m\Omega_X^q(\log E) \xrightarrow{R^s} W_r\Omega_X^q(\log E) \to 0.$$
(1.2.13)

*Proof.* Combine  $[28, \S 3]$  and Theorem 1.2.13.

**Corollary 1.2.15.** Let S be a regular F-finite  $\mathbb{F}_p$  algebra. Let A = S[[X]]] (where X is a variable). Then we have a canonical isomorphism of  $W_m(S)$ -modules

$$W_m \Omega^q_A \oplus W_m \Omega^{q-1}_S \operatorname{dlog}[X]_m \xrightarrow{\cong} W_m \Omega^q_A(\log X)$$

for every  $q \ge 0, m \ge 1$ 

### 1.2. THE LOG DE RHAM-WITT COMPLEX

*Proof.* Note that  $\{W_m \Omega_A^q \oplus W_m \Omega_S^{q-1} \operatorname{dlog}[X]_m\}_{m,q}$  is already a log de Rham-Witt complex over  $A_{\log}$  and there is a canonical map  $W_m \Omega_A^q \oplus W_m \Omega_S^{q-1} \operatorname{dlog}[X]_m \to W_m \Omega_A^q (\log X)$  which is compatible with the following monoid maps.

 $\alpha_m : M_A \to W_m \Omega^1_A(\log X); \qquad \alpha'_m : M_A \to W_m \Omega^1_A \oplus W_m(S) d\log[X]_m,$ 

where  $M_A = \{uX^n : u \in A^{\times}, n \in \mathbb{Z}\}$  and  $\alpha'_m(uX^n) = (\operatorname{dlog}[u]_m, n\operatorname{dlog}[X]_m)$ . Now corollary follows from Theorem 1.2.13 and the universal property of  $W_m\Omega^{\bullet}_{A_{\log}}$ .

## Chapter 2

## The filtered de Rham-Witt complex

Let X be a connected Noetherian regular F-finite  $\mathbb{F}_p$ -scheme of Krull dimension N. Let  $\operatorname{Div}(X)$  denote the group of Weil divisors on X. For any  $D = \sum_{i=1}^r n_i D_i \in \operatorname{Div}(X)$ , we let |D| denote the support of D and let  $D/p = \sum_{i=1}^r \lfloor n_i/p \rfloor D_i$ . For any  $\mathbb{F}_p$ -scheme Y and integer  $m \ge 1$ , we let  $W_m(Y) = W_m(H^0_{\operatorname{\acute{e}t}}(Y, \mathcal{O}_Y))$ . The elements of  $W_m(Y)$  will be denoted by m-tuples  $\underline{a} = (a_{m-1}, \ldots, a_0)$ . In this chapter, we shall define the filtered de Rham-Witt complex of X and prove some basic properties.

## 2.1 Definition and properties

**Definition 2.1.1.** We let  $D = \sum_{i=1}^{r} n_i D_i \in \text{Div}(X)$  and let  $j: U \hookrightarrow X$  be the inclusion of the open complement of D. For a morphism  $u: \widetilde{X} \to X$  of schemes, we let  $\widetilde{U} = U \times_X \widetilde{X}$  and  $\widetilde{D} = D \times_X \widetilde{X}$ . We let

$$\operatorname{fil}_D W_m(\widetilde{U}) = \{ \underline{a} = (a_{m-1}, \dots, a_0) \in W_m(\widetilde{U}) : a_i^{p^i} \in H^0_{\operatorname{\acute{e}t}}(\widetilde{X}, \mathcal{O}_{\widetilde{X}}(\widetilde{D})) \, \forall \, i \}.$$

We let  $\operatorname{fil}_D W_m \mathcal{O}_U$  be the presheaf on  $\operatorname{Sch}_X$  such that  $\operatorname{fil}_D W_m \mathcal{O}_U(\widetilde{X}) = \operatorname{fil}_D W_m(\widetilde{U})$ .

One easily checks that  $\operatorname{fil}_D \mathcal{O}_U = \mathcal{O}_X(D)$  and  $\operatorname{fil}_D W_m \mathcal{O}_U$  actually restricts to a sheaf on  $X_{\operatorname{zar}}$  and we have the inclusions of étale sheaves  $\operatorname{fil}_D W_m \mathcal{O}_U \subset \operatorname{fil}_{D'} W_m \mathcal{O}_W \subset h_* W_m \mathcal{O}_W$ whenever  $D \leq D'$  and  $h: W = X \setminus |D'| \hookrightarrow X$  is the inclusion. Note also that the canonical map  $j^*(\operatorname{fil}_D W_m \mathcal{O}_U) \to W_m \mathcal{O}_U$  is an isomorphism. If  $P \in X$  is a point and  $x_i$  is a generator of the ideal of  $D_i$  in  $\mathcal{O}_{X,P}$ , then one checks that  $a_i^{p^i} \in (\mathcal{O}_X(D))_P = \mathcal{O}_{X,P} \cdot \prod_j x_j^{-n_j}$  iff  $a_i^{p^i} \prod_j x_j^{n_j} \in \mathcal{O}_{X,P}$ . Hence, we obtain

$$(\operatorname{fil}_D W_m \mathcal{O}_U)_P = \{ \underline{a} \in W_m(\mathcal{O}_{X,P}[\pi^{-1}]) : a_i^{p^i} \prod_j x_j^{n_j} \in \mathcal{O}_{X,P} \ \forall \ i \},$$
(2.1.1)

where  $\pi = \prod_i x_i$ .

If X = Spec(A), where A is a regular local ring and  $\{x_1, \ldots, x_r\}$  are generators of the ideals of the irreducible components of D, we shall write  $\text{fil}_D W_m \mathcal{O}_U$  as  $\text{fil}_I W_m(A_\pi)$ or  $\text{fil}_{\underline{n}} W_m(A_\pi)$  or  $\text{fil}_I W_m(A_\pi)$ , where  $\pi = \prod_i x_i$  and  $I \subset Q(A)$  is the invertible ideal (x), where  $x = \prod_i x_i^{n_i}$  and  $\underline{n} = (n_1, \ldots, n_r) \in \mathbb{Z}^r$ . We shall write  $\mathcal{O}_X(D)$  as  $Ax^{-1}$ .

**Lemma 2.1.2.** fil<sub>D</sub> $W_m \mathcal{O}_U$  is a sheaf of abelian groups. Moreover, we have an exact sequence

$$0 \to \operatorname{fil}_D \mathcal{O}_U \xrightarrow{V^{m-1}} \operatorname{fil}_D W_m \mathcal{O}_U \xrightarrow{R} \operatorname{fil}_{D/p} W_{m-1} \mathcal{O}_U \to 0.$$
(2.1.2)

Proof. We may assume  $X = \operatorname{Spec} A$  and following the same notation as above, let D is given by the ideal (x), where  $x = \prod_i x_i^{n_i}$ . We only need to show that the map R is well-defined and surjective and  $\operatorname{fil}_D W_m \mathcal{O}_U$  is a sheaf of abelian groups. Rest of the part follows from (2.1.1). We let  $\underline{a} \in \operatorname{fil}_I W_m(A_\pi)$  and let  $I^{1/p} = (\prod_i x_i^{\lfloor n_i/p \rfloor})$ . Then  $R(\underline{a}) = \underline{b} = (b_{m-1}, \ldots, b_0)$ , where  $b_i = a_{i+1}$ . We can write (because A is UFD)  $a_{i+1} = a'_{i+1} \prod_j x_j^{t_j}$ , where  $t_j \in \mathbb{Z}$  for every j and  $a'_{i+1} \in A$  is not divisible by any  $x_j$ . By our hypothesis,  $(a_{i+1})^{p^{i+1}}x \in A$ . This implies,  $p^{i+1}t_j + n_j \ge 0$  which is equivalent to  $p^it_j + \lfloor n_j/p \rfloor \ge 0$ . So we get  $(b_i)^{p^i}x' \in A$  for  $0 \le i \le m-1$ , where  $x' = \prod_j x_j^{\lfloor n_j/p \rfloor}$ . This implies that  $\underline{b} \in \operatorname{fil}_{I^{1/p}} W_m(A_\pi)$ . Hence R is well defined. Also if  $b = (b_{m-2}, \ldots, b_0) \in \operatorname{fil}_{I^{1/p}} W_m(A_\pi)$  then by reversing the above argument we get  $(b_{m-2}, \ldots, b_0, 0) \in \operatorname{fil}_I W_m(A_\pi)$ . So R is surjective and hence we win.

Now we show  $\operatorname{fil}_D W_m \mathcal{O}_U$  is a sheaf of abelian groups by induction on m. If m = 1, then  $\operatorname{fil}_D \mathcal{O}_U = \mathcal{O}_X(D)$  is a sheaf of abelian group. If m > 1, let  $\underline{a} = (a_{m-1}, \ldots, a_0)$  and  $\underline{a'} = (a'_{m-1}, \ldots, a'_0)$  are in  $\operatorname{fil}_D W_m \mathcal{O}_U$ . Enough to show  $\underline{a} - \underline{a'} \in \operatorname{fil}_D W_m \mathcal{O}_U$ . Since R is additive, by induction and the above argument we have  $R(\underline{a} - \underline{a'}) \in \operatorname{fil}_{D/p} W_{m-1} \mathcal{O}_U$ . So, we can write  $R(\underline{a} - \underline{a'}) = (c_{m-2}, \ldots, c_0)$  satisfying the relation (2.1.1). Then the previous argument shows  $(c_{m-2}, \ldots, c_0, 0) \in \operatorname{fil}_D W_m \mathcal{O}_U$ . Now note that  $\underline{a} - \underline{a'} = (c_{m-2}, \ldots, c_0, a_0 - a'_0)$  which is clearly in  $\operatorname{fil}_D W_m \mathcal{O}_U$  (because,  $a_0 - a'_0 \in \operatorname{fil}_D \mathcal{O}_U$ ).

**Lemma 2.1.3.** For two Weil divisors D and D' with complements U and U', respectively, we have  $\operatorname{fil}_D W_m \mathcal{O}_U + \operatorname{fil}_{D'} W_m \mathcal{O}_{U'} \subset \operatorname{fil}_{D+D'} W_m \mathcal{O}_{U\cap U'}$  and  $\operatorname{fil}_D W_m \mathcal{O}_U \cdot \operatorname{fil}_{D'} W_m \mathcal{O}_{U'} \subset \operatorname{fil}_{D+D'} W_m \mathcal{O}_{U\cap U'}$  under the addition and multiplication of  $W_m \mathcal{O}_\eta$ . In particular,  $\operatorname{fil}_D W_m \mathcal{O}_U$  is a subsheaf of  $W_m \mathcal{O}_X$ -modules inside  $W_m \mathcal{O}_\eta$ .

*Proof.* Since we can check this locally, we assume that X = Spec(A), where A is a local ring. We let  $\{x_1, \ldots, x_r\}$  and  $\{y_1, \ldots, y_s\}$  be the generators of the ideals of the irreducible components of D and D', respectively. We let  $x = \prod_i x_i^{m_i}$  and  $y = \prod_j y_j^{n_j}$ , where  $D = \sum_i m_i D_i$  and  $D' = \sum_j n_j D'_j$ . Write I = (x) and I' = (y). Suppose  $\underline{a} \in \text{fil}_I W_m(A_x)$  and

#### 2.1. DEFINITION AND PROPERTIES

 $\underline{b} \in \operatorname{fil}_{I'} W_m(A_y). \text{ We can write } \underline{a} = \sum_{i=0}^{m-1} V^i([a_{m-1-i}]) \text{ and } \underline{b} = \sum_{i=0}^{m-1} V^i([b_{m-1-i}]). \text{ This yields}$  $\underline{a} \cdot \underline{b} = \sum_{i,j=0}^{m-1} V^i([a_{m-1-i}]) V^j([b_{m-1-j}]).$ 

For  $0 \le i, j \le m - 1$ , using the identity xV(y) = V(Fx.y) we have

$$V^{i}([a_{m-1-i}])V^{j}([b_{m-1-j}]) = \begin{cases} 0 & \text{if } i+j \ge m\\ V^{i+j}([a_{m-1-i}^{p^{j}}b_{m-1-j}^{p^{i}}]) & \text{if } i+j < m. \end{cases}$$
(2.1.3)

By (2.1.1), the condition  $\underline{a} \in \operatorname{fil}_{I}W_{m}(A_{x})$  and  $\underline{b} \in \operatorname{fil}_{I'}W_{m}(A_{y})$  means that  $xa_{i}^{p^{i}} \in A$ and  $yb_{i}^{p^{i}} \in A$  for each i. From this, we get for i + j < m that  $(a_{m-1-i}^{p^{j}}b_{m-1-j}^{p^{i}})^{p^{m-1-i-j}}xy = a_{m-1-i}^{p^{m-1-i}}b_{m-1-j}^{p^{m-1-j}}xy \in A$ . This implies that  $V^{i}([a_{m-1-i}])V^{j}([b_{m-1-j}]) \in \operatorname{fil}_{II'}W_{m}(A_{xy})$ . By Lemma 2.1.2 we conclude that  $\underline{a} \cdot \underline{b} \in \operatorname{fil}_{II'}W_{m}(A_{xy})$ . Also, the claim about the addition immediately follows from Lemma 2.1.2, observing that  $\operatorname{fil}_{D}W_{m}\mathcal{O}_{U}$  (and  $\operatorname{fil}_{D'}W_{m}\mathcal{O}_{U'}) \subset \operatorname{fil}_{D+D'}W_{m}\mathcal{O}_{U\cap U'}$ . This concludes the proof of the first part of the lemma. The second part follows from the first by taking D' = 0 and noting that  $\operatorname{fil}_{0}W_{m}\mathcal{O}_{X} = W_{m}\mathcal{O}_{X}$ .

Notation 2.1.4. For D as in Definition 2.1.1, let  $W_m \mathcal{O}_X(D)$  be the sheaf on  $X_{\acute{e}t}$  which is locally defined as  $W_m \mathcal{O}_X(D) = [x^{-1}]_m \cdot W_m \mathcal{O}_X \subset j_* W_m \mathcal{O}_U$ , where  $[-]_m : j_* \mathcal{O}_U \to j_* W_m \mathcal{O}_U$ is the Teichmüller homomorphism, and D is locally defined by the invertible ideal  $(x) \subset K(X)$  = the fraction field of X, where  $x = \prod_i x_i^{m_i}$ . One knows that  $W_m \mathcal{O}_X(D)$  is a sheaf of invertible  $W_m \mathcal{O}_X$ -modules. For a sheaf of  $W_m \mathcal{O}_X$ -modules  $\mathcal{M}$ , we let  $\mathcal{M}(D) = \mathcal{M} \otimes_{W_m \mathcal{O}_X} W_m \mathcal{O}_X(D)$ . If A is a local ring and D is defined by I = (x), we also write  $\mathcal{M}(D)$  as  $M(x^{-1})$  when  $\mathcal{M}$  is defined by a  $W_m(A)$ -module M.

**Lemma 2.1.5.** One has the identity  $\operatorname{fil}_{p^{m-1}D} W_m \mathcal{O}_U = W_m \mathcal{O}_X(D)$ .

*Proof.* Since we can check this locally, we assume that X = Spec(A) is as in the proof of Lemma 2.1.3. We let  $I = (x^{p^{m-1}})$ . Following our notations, we see that  $\underline{a} = (a_{m-1}, \ldots, a_0) \in \text{fil}_I W_m(A_x)$  if and only if  $a_i^{p^i} x^{p^{m-1}} \in A$  for all  $0 \le i \le m - 1$ . Since A is UFD, this is equivalent to  $a_i x^{p^{m-1-i}} \in A$  for all  $0 \le i \le m - 1$ , or equivalently  $\underline{a} \cdot [x]_m = (a_{m-1}, \ldots, a_i x^{p^{m-1-i}}, \ldots, a_0 x^{m-1}) \in W_m(A)$ . This finishes the proof.

Let us now assume that  $E \subset X$  is a simple normal crossing divisor with irreducible components  $\{E_1, \ldots, E_r\}$ . We let  $j: U \hookrightarrow X$  be the inclusion of the complement of E. Let  $\text{Div}_E(X) \cong \mathbb{Z}^r$  be the group of Weil divisors of X whose support lie in E.

**Definition 2.1.6.** For  $D = \sum_{i=1}^{r} m_i [E_i] \in \operatorname{Div}_E(X)$  and an integer  $q \ge 0$ , we let  $\operatorname{fil}_D W_m \Omega_U^q = \operatorname{fil}_D W_m \mathcal{O}_U \cdot W_m \Omega_X^q (\log E) + d(\operatorname{fil}_D W_m \mathcal{O}_U) \cdot W_m \Omega_X^{q-1} (\log E)$  considered as a subsheaf of  $j_* W_m \Omega_U^q$ . We shall write  $\operatorname{fil}_D W_1 \Omega_U^q$  as  $\operatorname{fil}_D \Omega_U^q$ .

Throughout our discussion, we shall treat  $\operatorname{fil}_D W_m \Omega_U^q$  as a subsheaf of  $j_* W_m \Omega_U^q$ . If  $X = \operatorname{Spec}(A)$ , where A is a regular local ring with maximal ideal

 $\mathfrak{m} = (x_1, \dots, x_r, y_1, \dots, y_s)$  and  $\{x_1, \dots, x_r\}$  are generators of the ideals of the irreducible components of E, then we shall write  $\operatorname{fil}_D W_m \Omega_U^q$  as  $\operatorname{fil}_I W_m \Omega_{A_\pi}^q$  or  $\operatorname{fil}_I W_m \Omega_{A_x}^q$ , where  $\pi = \prod_i x_i$  and  $I = (\prod_i x_i^{m_i})$  and  $x = \prod_i x_i^{m_i}$ .

Lemma 2.1.7. We have the following.

- 1. fil<sub>D</sub> $W_m \Omega_U^q$  is a sheaf of finite type  $W_m \mathcal{O}_X$ -modules.
- 2.  $\operatorname{fil}_D\Omega^q_U = \Omega^q_X(\log E)(D)$ . In particular,  $\operatorname{fil}_{-E}\Omega^d_U = \Omega^d_X$ , where d is the rank of  $\Omega^1_X$ .
- 3.  $\lim_{n \to \infty} \operatorname{fil}_{nE} W_m \Omega_U^q \xrightarrow{\cong} j_* W_m \Omega_U^q.$
- 4.  $d(\operatorname{fil}_D W_m \Omega_U^q) \subset \operatorname{fil}_D W_m \Omega_U^{q+1}$ .
- 5.  $F(\operatorname{fil}_D W_{m+1}\Omega_U^q) \subset \operatorname{fil}_D W_m\Omega_U^q$ .
- 6.  $V(\operatorname{fil}_D W_m \Omega_U^q) \subset \operatorname{fil}_D W_{m+1} \Omega_U^q$ .
- $7. R(\mathrm{fil}_D W_{m+1}\Omega_U^q) = \mathrm{fil}_{D/p} W_m \Omega_U^q.$
- 8.  $\operatorname{fil}_D W_m \Omega_U^q \cdot \operatorname{fil}_{D'} W_m \Omega_U^{q'} \subset \operatorname{fil}_{D+D'} W_m \Omega_U^{q+q'}$  if  $|D'| \subset E$ .
- 9.  $\operatorname{fil}_{p^{m-1}D} W_m \Omega^q_U \xrightarrow{\cong} W_m \Omega^q_X(\log E)(D).$

10.  $\operatorname{fil}_0 W_m \Omega_U^q = W_m \Omega_X^q (\log E)$  and  $\operatorname{fil}_D W_m \Omega_U^q \subset W_m \Omega_X^q$  if  $m_i < 0$  for every  $1 \le i \le r$ .

Proof. Since all claims of the lemma can be checked locally, we shall assume (whenever necessary in the proof) that  $X = \operatorname{Spec}(A)$  and  $D \subset X$  are as in Definition 2.1.6. The first part of item (1) is clear. To prove that  $\operatorname{fil}_D W_m \Omega_U^q$  is a sheaf of finite type  $W_m \mathcal{O}_X$ -modules, we can find an effective Weil divisor  $D' \in \operatorname{Div}_E(X)$  such that D'' := D + D' is effective. In particular,  $\operatorname{fil}_D W_m \Omega_U^q \subset \operatorname{fil}_{D''} W_m \Omega_U^q \subset \operatorname{fil}_{p^{m-1}D''} W_m \Omega_U^q$ . Since  $W_m \mathcal{O}_X$  is a sheaf of Noetherian rings (cf. [58, Lem. 2.9]), it suffices to show that  $\operatorname{fil}_{p^{m-1}D''} W_m \Omega_U^q$  is a sheaf of finite type  $W_m \mathcal{O}_X$ -modules. But this follows from item (9) because  $W_m \Omega_X^q(\log E)$  is a sheaf of finite type  $W_m \mathcal{O}_X$ -modules (by loc. cit.) and  $W_m \mathcal{O}_X(D'')$  is a sheaf of invertible  $W_m \mathcal{O}_X$ -modules.

When q = 0, we have  $\operatorname{fil}_D \Omega_U^q = \operatorname{fil}_D \mathcal{O}_X = \mathcal{O}_X(D) = \Omega_X^0(\log E)(D)$ . It follows that  $\operatorname{fil}_D \mathcal{O}_U \cdot \Omega_X^q(\log E) = \mathcal{O}_X(D) \cdot \Omega_X^q(\log E) = \Omega_X^q(\log E) = \Omega_X^q(\log E)(D)$  when q > 0. On the other hand,

$$d(\operatorname{fil}_{D}\mathcal{O}_{U}) \cdot \Omega_{X}^{q-1}(\log E) = d(\mathcal{O}_{X}(D)) \cdot \Omega_{X}^{q-1}(\log E)$$
  

$$\subset \mathcal{O}_{X}(D)\Omega_{X}^{1}(\log E) \cdot \Omega_{X}^{q-1}(\log E)$$
  

$$\subset \Omega_{X}^{q}(\log E)(D).$$

#### 2.1. DEFINITION AND PROPERTIES

This proves (2).

For (3), we recall that  $j_*W_m\Omega_U^q \cong j_*W_m\mathcal{O}_U \otimes_{W_m\mathcal{O}_X} W_m\Omega_X^q = j_*W_m\mathcal{O}_U \cdot W_m\Omega_X^q$ . Meanwhile, one checks directly from (2.1.1) that  $j_*W_m\mathcal{O}_U = \varinjlim_n \operatorname{fil}_{nE} W_m\mathcal{O}_U$ . This proves the desired isomorphism.

Next, we note that  $d(\operatorname{fil}_D W_m \mathcal{O}_U \cdot W_m \Omega_X^q(\log E)) \subset d(\operatorname{fil}_D W_m \mathcal{O}_U) \cdot W_m \Omega_X^q(\log E) + \operatorname{fil}_D W_m \mathcal{O}_U \cdot W_m \Omega_X^{q+1}(\log E) = \operatorname{fil}_D W_m \Omega_U^{q+1}$ . We also have

$$d(d(\operatorname{fil}_D W_m \mathcal{O}_U) \cdot W_m \Omega_X^{q-1}(\log E)) \subset d(\operatorname{fil}_D W_m \mathcal{O}_U) \cdot W_m \Omega_X^q(\log E) \subset \operatorname{fil}_D W_m \Omega_U^{q+1}.$$

This proves (4).

We now prove (5). When q = 0, the claim is clear using (2.1.1) and the fact that  $F = R \circ \overline{F}$ , where  $\overline{F}(\underline{a}) = (a_i^p)$ . We now assume q > 0 and let  $a \in \operatorname{fil}_I W_{m+1}(A_{\pi}), w \in W_{m+1}\Omega_A^q(\log \pi)$ . We then have  $F(aw) = F(a)F(w) \in \operatorname{fil}_I W_m(A_{\pi})W_m\Omega_A^q(\log \pi)$  by the q = 0 case and Theorem 1.2.13.

Suppose next that  $\underline{a} \in \operatorname{fil}_{I}W_{m+1}(A_{\pi})$  and  $w \in W_{m+1}\Omega_{A}^{q-1}(\log \pi)$ . Letting  $\underline{a} = \sum_{i=0}^{m} V^{i}([a_{m-i}])$ , we get  $Fd(\underline{a}) = Fd[a_{m}] + \sum_{i=0}^{m-1} FdV^{i+1}[a_{m-1-i}] = Fd[a_{m}] + d\underline{a}'$ , where  $\underline{a}' = (a_{m-1}, \ldots, a_{0}) \in \operatorname{fil}_{I}W_{m}(A_{\pi})$ . In particular,  $d\underline{a}' \cdot F(w) \in d(\operatorname{fil}_{I}W_{m}(A_{\pi})) \cdot W_{m}\Omega_{A}^{q-1}(\log \pi) \subset \operatorname{fil}_{I}W_{m}\Omega_{A_{\pi}}^{q}$ . We also have

$$\begin{aligned} \operatorname{fil}_{I} W_{m} \Omega_{A_{\pi}}^{q} \cdot W_{m} \Omega_{A}^{q'}(\log \pi) &\subset \operatorname{fil}_{I} W_{m}(A_{\pi}) W_{m} \Omega_{A}^{q+q'}(\log \pi) \\ &\quad + d(\operatorname{fil}_{I} W_{m}(A_{\pi})) W_{m} \Omega_{A}^{q+q'-1}(\log \pi) \\ &= \operatorname{fil}_{I} W_{m} \Omega_{A_{\pi}}^{q+q'}. \end{aligned}$$

$$(2.1.4)$$

Since  $F(d[a_m]w) = Fd[a_m] \cdot F(w)$  and  $F(w) \in W_m \Omega_A^{q-1}(\log \pi)$  by Theorem 1.2.13, it remains to show using (2.1.4) that  $Fd[a_m] \in \operatorname{fil}_I W_m \Omega_{A_\pi}^1$ .

We write  $a_m = a' \prod_i x_i^{n_i}$ , where  $n_i \in \mathbb{Z}$  for every *i* and  $a' \in A$  is not divisible by any  $x_i$ . Then the condition  $\underline{a} \in \operatorname{fil}_I W_{m+1}(A_{\pi})$  implies that  $[\prod_i x_i^{n_i}]_{m+1} \in \operatorname{fil}_I W_{m+1}(A_{\pi})$ . Letting  $\gamma = \prod_i x_i^{n_i}$ , we get

$$Fd[a_m]_{m+1} = Fd[a']_{m+1} \cdot F([\gamma]_{m+1}) + F([a_m]_{m+1}) d\log([\gamma]_m)$$

Note that  $\operatorname{dlog}[\gamma]_m = \sum_i n_i \cdot \operatorname{dlog}[x_i]_m \in W_m \Omega^1_A(\log \pi)$  (cf. Theorem 1.2.13). Hence,  $F([a_m]_{m+1})\operatorname{dlog}([\gamma]_m) \in \operatorname{fil}_I W_m \Omega^1_{A_{\pi}}$ . On the other hand, we have  $Fd[a']_{m+1} \cdot F([\gamma]_{m+1}) \in \operatorname{fil}_I W_m (A_{\pi}) \cdot W_m \Omega^1_A \subset \operatorname{fil}_I W_m \Omega^1_{A_{\pi}}$ . This proves (5).

We next show (6). As the q = 0 case is clear using (2.1.1), we shall assume q > 0. We let  $\underline{a} \in \operatorname{fil}_I W_m(A_\pi)$  and  $w \in W_m \Omega^q_A(\log \pi)$ . We can write w as the sum of elements of the form  $V^{j_0}([b_0])dV^{j_1}([b_1])\cdots dV^{j_t}([b_t])d\log[x_{i_1}]\cdots d\log[x_{i_s}]$ , where  $1 \le i_1 < \cdots < i_s \le r$ , t = q - s and  $b_i \in A$ . Letting  $\underline{b} = \underline{a}V^{j_0}[b_0]$  and  $w' = dV^{j_1}([b_1])\cdots dV^{j_t}([b_t])$ , we get

$$V(\underline{b}w') = V(\underline{b})dV^{j_1+1}[b_1]\cdots dV^{j_t+1}([b_t]) \in \mathrm{fil}_I W_{m+1}(A_{\pi}) \cdot W_{m+1}\Omega_A^t.$$

This yields

$$V(\underline{a}w) = V(\underline{b}w' \operatorname{dlog}[x_{i_1}] \cdots \operatorname{dlog}[x_{i_s}]) = V(\underline{b}w') \cdot \operatorname{dlog}[x_{i_1}] \cdots \operatorname{dlog}[x_{i_s}]$$

$$\in \operatorname{fil}_I W_{m+1}(A_{\pi}) \cdot W_{m+1}\Omega_A^t \cdot W_{m+1}\Omega_A^s(\operatorname{log}\pi)$$

$$\subset \operatorname{fil}_I W_{m+1}\Omega_A^t(\operatorname{log}\pi) \cdot W_{m+1}\Omega_A^s(\operatorname{log}\pi)$$

$$\subset \operatorname{fil}_{m+1} W_{m+1}\Omega_{A_{\pi}}^q.$$

If  $\underline{a} \in \operatorname{fil}_{I}W_{m}(A_{\pi})$  and  $w \in W_{m}\Omega_{A}^{q-1}(\log \pi)$ , then  $V(\underline{a}\underline{\cdot}w) = dV(\underline{a})\cdot V(w) \in d(\operatorname{fil}_{I}W_{m+1}(A_{\pi}))\cdot W_{m+1}\Omega_{A}^{q-1}(\log \pi) \subset \operatorname{fil}_{I}W_{m+1}\Omega_{A_{\pi}}^{q} \cdot W_{m+1}\Omega_{A}^{q-1}(\log \pi) \subset \operatorname{fil}_{I}W_{m+1}\Omega_{A_{\pi}}^{q}$ . This proves (6).

To prove (7) for q = 0 follows from Corollary 2.1.2. The q > 0 case now easily follows from q = 0 case and Corollary 1.2.14 because R is multiplicative and commutes with d.

For (8), using Lemma 2.1.3 and an elementary computation, it suffices to check that  $\operatorname{fl}_{I}W_{m}(A_{\pi}) \cdot d(\operatorname{fl}_{I'}W_{m}(A_{\pi})) \subset \operatorname{fl}_{II'}W_{m}\Omega^{1}_{A_{\pi}}$ , where  $x = \prod_{i} x_{i}^{m_{i}}$  (resp.  $y = \prod_{i} x_{i}^{n_{i}}$ ) and I = (x) (resp. I' = (y)). We let  $\underline{a} = \sum_{i=0}^{m-1} V^{i}([a_{m-1-i}]) \in \operatorname{fl}_{I}W_{m}(A_{\pi})$  and  $\underline{b} = \sum_{i=0}^{m-1} V^{i}([b_{m-1-i}]) \in \operatorname{fl}_{I'}W_{m}(A_{\pi})$ . This yields  $\underline{a} \cdot d\underline{b} = \sum_{i,j} V^{i}([a_{m-1-i}]) dV^{j}([b_{m-1-j}])$ . It suffices now to show that  $V^{i}([a_{m-1-i}]) dV^{j}([b_{m-1-j}]) \in \operatorname{fl}_{II'}W_{m}\Omega^{1}_{A_{\pi}}$  for every  $0 \leq i, j \leq m-1$ .

If we write  $b_{m-1-i} = b' \prod_l x_l^{r_l}$ , where  $b' \in A$ ,  $r_l \in \mathbb{Z}$  and no  $x_l$  divides b', then we get that  $\prod_l [x_l^{r_l}] \in \operatorname{fil}_{I'} W_m(A_\pi)$ . We write  $\gamma' = \prod_l x_l^{r_l}$ .

We first assume  $i \ge j$  and write

$$V^{i}([a_{m-1-i}])dV^{j}([b_{m-1-j}]) = V^{j}(V^{i-j}[a_{m-1-i}]d([b_{m-1-j}])).$$

It is enough to show using (6) that  $V^{i-j}[a_{m-1-i}]d([b_{m-1-j}]) \in \operatorname{fil}_{II'}W_{m-j}\Omega^1_{A_{\pi}}$ . We now compute

$$d([b_{m-1-j}])V^{i-j}([a_{m-1-i}]) = d([b'])[\gamma']V^{i-j}([a_{m-1-i}]) + [b'\gamma']V^{i-j}([a_{m-1-i}])dlog([\gamma']).$$

The terms  $[\gamma']V^{i-j}([a_{m-1-i}])$  and  $[b'\gamma']V^{i-j}([a_{m-1-i}])$  lie fil<sub>II'</sub> $W_m(A_\pi)$  by Lemma 2.1.3. Since  $d([b']) \in W_m\Omega^1_A$  and  $dlog([\gamma']) \in W_m\Omega^1_A(log\pi)$  (cf. Theorem 1.2.13), we conclude that  $d([b_{m-1-j}])V^{i-j}([a_{m-1-i}]) \in fil_{II'}W_m\Omega^1_{A_\pi}$ .

If  $i \leq j$ , then by Leibniz rule, Lemma 2.1.3 and item (4) it reduces to the case  $i \geq j$ . Hence we are done.

To prove (9), we write  $I' = I^{p^{m-1}}$  and note that  $\operatorname{fil}_{I'} W_m \Omega_{A_{\pi}}^q = \operatorname{fil}_{I'} W_m (A_{\pi}) \cdot W_m \Omega_A^q (\log \pi) + d(\operatorname{fil}_{I'} W_m (A_{\pi})) \cdot W_m \Omega_A^{q-1} (\log \pi) = W_m \Omega_A^q (\log \pi) (x^{-1}) + d([x^{-1}]_m W_m (A)) \cdot W_m \Omega_A^{q-1} (\log \pi)$ by Lemma 2.1.5. On the other hand,

$$d([x^{-1}]_m W_m(A)) \cdot W_m \Omega_A^{q-1}(\log \pi) \subset [x^{-1}]_m d(W_m(A)) \cdot W_m \Omega_A^{q-1}(\log \pi) + [x^{-1}]_m W_m(A) d\log([x]_m) \cdot W_m \Omega_A^{q-1}(\log \pi) \subset W_m \Omega_A^q (\log \pi) (x^{-1}).$$

The desired isomorphism now follows.

Finally, to prove (10), we only need to check its second part as the first part is clear. To prove its second part, it suffices to show that  $\operatorname{fl}_{-E} W_m \Omega_U^q \subset W_m \Omega_X^q$ . To show this, we let  $\underline{a} = (a_{m-1}, \ldots, a_0) \in \operatorname{fl}_I W_m(A_\pi)$  and  $w \in W_m \Omega_A^q(\log \pi)$ . Letting  $\underline{a} =$  $\sum_i V^i([a_{m-1-i}]_{m-i})$ , we have that  $a_i^{p^i} \pi^{-1} \in A$ . This implies that  $\pi \mid a_i^{p^i}$ . Equivalently,  $\pi \mid a_i$  (because A is a UFD). We write  $a_i = \pi \alpha_i$ , where  $\alpha_i \in A$ . We can write w as the sum of elements of the form

$$V^{j_0}([b_0])dV^{j_1}([b_1])\cdots dV^{j_t}([b_t])d\log([x_{i_1}]_m)\cdots d\log([x_{i_s}]_m),$$

where  $1 \le i_1 < \cdots < i_s \le r$ , t = q - s and  $b_i \in A$ . We have

$$V^{i}([a_{m-1-i}]_{m-i})dlog([x_{i_{1}}]_{m})\cdots dlog([x_{i_{s}}]_{m}) =$$
$$V^{i}([\alpha_{m-i-1}]_{m-i}[\pi]_{m-i}dlog([x_{i_{1}}]_{m-i})\cdots dlog([x_{i_{s}}]_{m-i})).$$

But this last term clearly lies in  $W_m \Omega_A^q$ . Taking the sum over  $i \in \{0, \ldots, m-1\}$ , we get that  $\underline{a} \operatorname{dlog}([x_{i_1}]_m) \cdots \operatorname{dlog}([x_{i_s}]_m) \in W_m \Omega_A^q$ . But this implies that  $\underline{a} w \in W_m \Omega_A^q$ . We have thus shown that  $\operatorname{fil}_I W_m(A_\pi) W_m \Omega_A^q(\log \pi) \subset W_m \Omega_A^q$ .

If  $\underline{a} \in \operatorname{fil}_{I}W_{m}(A_{\pi})$  and  $w \in W_{m}\Omega_{A}^{q-1}(\log \pi)$ , then  $d(\underline{a})w = d(\underline{a}w) - \underline{a}d(w)$ . We showed above that  $\underline{a}w \in W_{m}\Omega_{A}^{q-1}$  and  $\underline{a}d(w) \in W_{m}\Omega_{A}^{q}$ . It follows that  $d(\underline{a})w \in W_{m}\Omega_{A}^{q}$ . We have thus shown that  $\operatorname{fil}_{I}W_{m}(A_{\pi}) \subset W_{m}\Omega_{A}^{q}$ .

## Chapter 3

# Structure of the filtered de Rham-Witt complex I

The main purpose of this chapter is to provide a concrete description of the filtered de Rham-Witt complex defined in § 2, when X is spectrum of a ring of the form  $S[[\pi]]$  and the SNCD is given by the ideal ( $\pi$ ). Using this description, we also prove some results that are well-known in the case of de Rham-Witt complex of the regular schemes.

### 3.1 Filtered de Rham-Witt complex of complete local $\mathbb{F}_p$ -algebras

We let S be an F-finite Noetherian regular local  $\mathbb{F}_p$ -algebra. Let  $A = S[[\pi]]$  and  $K = A_{\pi}$ . For  $n \in \mathbb{Z}$  and the invertible ideal  $I = (\pi^n) \subset K$ , we shall denote fil<sub>I</sub> $W_m \Omega_K^q$  by fil<sub>n</sub> $W_m \Omega_K^q$  and write  $M(\pi^n)$  as M(n) (cf. 2.1.4) for any  $W_m(A)$ -module M. We let  $\operatorname{gr}_n W_m \Omega_K^q = \frac{\operatorname{fil}_n W_m \Omega_K^q}{\operatorname{fil}_{n-1} W_m \Omega_K^q}$ .

The goal of this section is to prove some key results about the structure of  $\operatorname{fil}_n W_m \Omega_K^q$ .

We shall use the following theorem of Geisser-Hesselholt [15, Thm. B] for proving our main results. Let  $I_p$  denote the set of positive integers prime to p.

**Theorem 3.1.1.** (*Geisser-Hesselholt*) Every element  $\omega \in W_m \Omega_A^q$  can be uniquely written as an infinite series

$$w = \sum_{i \ge 0} a_{0,i}^{(m)} [\pi]_m^i + \sum_{i \ge 0} b_{0,i}^{(m)} [\pi]_m^i d[\pi]_m + \sum_{s \ge 1 j \in I_p} \left( V^s (a_{s,j}^{(m-s)} [\pi]_{m-s}^j) + dV^s (b_{s,j}^{(m-s)} [\pi]_{m-s}^j) \right),$$

where the components  $a_{s,j}^{(n)} \in W_n \Omega_S^q$  and  $b_{s,j}^{(n)} \in W_n \Omega_S^{q-1}$ .

For  $q, s \ge 0, m \ge 1$  and  $i \in \mathbb{Z}$ , we let

$$A_{i}^{m,q}(S) = W_{m}\Omega_{S}^{q}[\pi]_{m}^{i} + W_{m}\Omega_{S}^{q}[\pi]_{m}^{i}dlog([\pi]_{m});$$

$$B_{i,s}^{m,q}(S) = V^{s}\left(W_{m-s}\Omega_{S}^{q}[\pi]_{m-s}^{i}\right) + dV^{s}\left(W_{m-s}\Omega_{S}^{q-1}[\pi]_{m-s}^{i}\right).$$

$$(S), B_{i,s}^{m,q}(S) \subset W_{m}\Omega_{K}^{q}.$$
(3.1.1)

Note that  $A_i^{m,q}(S), B_{i,s}^{m,q}(S) \subset W_m \Omega_K^q$ .

**Definition 3.1.2.** For  $q \ge 0$ ,  $m \ge 1$  and  $n \in \mathbb{Z}$ , we let

$$F_n^{m,q}(S) = \begin{cases} A_{n/p^{m-1}}^{m,q}(S) & \text{if } p^{m-1} \mid n \\ B_{n/p^{m-1-s},s}^{m,q}(S) & \text{if } s = \min\{j > 0 \mid p^{m-1-j} \mid n\}. \end{cases}$$

**Corollary 3.1.3.** For every  $m \ge 1$  and  $q \ge 0$ , there is a canonical decomposition

$$\theta_A^{m,q} \colon \prod_{n \ge 0} F_n^{m,q}(S) \xrightarrow{\cong} W_m \Omega_A^q(\log \pi).$$
(3.1.2)

*Proof.* We note from Theorem 3.1.1 that the canonical (once the uniformizer  $\pi$  is fixed) maps from  $W_m \Omega_S^q$ ,  $F_n^{m,q}(S)$  to  $W_m \Omega_A^q$  (for  $n \ge 1$ ) induce an isomorphism

$$\overline{\theta}_A^{m,q}: W_m\Omega_S^q \bigoplus \prod_{n \ge 1} F_n^{m,q}(S) \xrightarrow{\cong} W_m\Omega_A^q.$$

Then  $\overline{\theta}^{m,q}_A$  can be uniquely extended to

$$\theta_A^{m,q} : \prod_{n \ge 0} F_n^{m,q}(S) = W_m \Omega_S^{q-1} \operatorname{dlog}[\pi]_m \bigoplus W_m \Omega_S^q \bigoplus \prod_{n \ge 1} F_n^{m,q}(S)$$
$$\xrightarrow{Id \oplus \overline{\theta}_A^{m,q}} W_m \Omega_S^{q-1} \operatorname{dlog}[\pi]_m \bigoplus W_m \Omega_A^q \xrightarrow{\cong} W_m \Omega_A^q(\log \pi)$$

where the last isomorphism follows from Lemma 1.2.15.

In order to extend this decomposition to  $W_m\Omega_K^q$ , we need a relation between the groups  $\{F_n^{m,q}(S)\}$  for positive and negative values of  $n \in \mathbb{Z}$ . This can be achieved by the following lemma.

**Lemma 3.1.4.** For all  $n, l \in \mathbb{Z}, m \ge 1, q \ge 0$ , we have

$$[\pi]_m^l F_n^{m,q}(S) = F_{n+p^{m-1}l}^{m,q}(S).$$

*Proof.* We first consider the case  $n = p^{m-1}i$  with  $i \in \mathbb{Z}$ . In this case, one checks that  $F_n^{m,q}(S) = [\pi]_m^i F_0^{m,q}(S)$ . This implies that  $[\pi]_m^l F_n^{m,q}(S) = [\pi]_m^{i+l} F_0^{m,q}(S) = F_{p^{m-1}(i+l)}^{m,q}(S) = F_{p^{m-1}(i+l)}^{m,q}(S)$ .

# 3.1. FILTERED DE RHAM-WITT COMPLEX OF COMPLETE LOCAL $\mathbb{F}_p\text{-}\mathrm{ALGEBRAS}$

Suppose now that  $n = p^{m-1-s}i$  with  $0 < s \le m-1$  and  $|i| \in I_p$ . Let  $t = n + p^{m-1}l = p^{m-1-s}i + p^{m-1}l = p^{m-1-s}(i+p^sl) = p^{m-1-s}i'$ , where we let  $i' = i + p^sl \in I_p$ . We let r = m-s. We then have

$$\begin{aligned} [\pi]_{m}^{l}F_{n}^{m,q}(S) &= [\pi]_{m}^{l}(V^{s}(W_{r}\Omega_{S}^{q}[\pi]_{r}^{i}) + dV^{s}(W_{r}\Omega_{S}^{q-1}[\pi]_{r}^{i})) \\ &= V^{s}(W_{r}\Omega_{S}^{q}[\pi]_{r}^{i'}) + [\pi]_{m}^{l}dV^{s}(W_{r}\Omega_{S}^{q-1}[\pi]_{r}^{i}) \\ &\subset_{\dagger} V^{s}(W_{r}\Omega_{S}^{q}[\pi]_{r}^{i'}) + dV^{s}(W_{r}\Omega_{S}^{q-1}[\pi]_{r}^{i'}) \\ &+ V^{s}(W_{r}\Omega_{S}^{q-1}[\pi]_{r}^{i}F^{s}d[\pi]_{r}^{l}) \\ &\subset F_{t}^{m,q}(S) + V^{s}(W_{r}\Omega_{S}^{q-1}[\pi]_{r}^{i}F^{s}d[\pi]_{r}^{l}). \end{aligned}$$

To obtain  $\subset_{\dagger}$ , we applied Leibniz rule (xdy = d(xy) - ydx) on  $[\pi]_m^l dV^s (W_r \Omega_S^{q-1}[\pi]_r^i)$ . We also used the relations aV(b) = V(F(a).b) and  $F(d[t]_r) = [t]_{r-1}^{p-1}d[t]_{r-1}$ , for  $a, b \in W_r \Omega_A^q$  and  $t \in A$  in other equalities. Meanwhile, we have

$$V^{s}(W_{r}\Omega_{S}^{q-1}[\pi]_{r}^{i}F^{s}d[\pi]_{r}^{l}) \subset V^{s}(W_{r}\Omega_{S}^{q-1}d([\pi]_{r}^{i'}))$$

$$\subset V^{s}d(W_{r}\Omega_{S}^{q-1}[\pi]_{r}^{i'}) + V^{s}(d(W_{r}\Omega_{S}^{q-1})[\pi]_{r}^{i'})$$

$$\subset dV^{s}(W_{r}\Omega_{S}^{q-1}[\pi]_{r}^{i'}) + V^{s}(W_{r}\Omega_{S}^{q}[\pi]_{r}^{i'})$$

$$= F_{t}^{m,q}(S).$$

This shows that  $[\pi]_m^l F_n^{m,q}(S) \subset F_{n+p^{m-1}l}^{m,q}(S)$  for all  $l, n \in \mathbb{Z}$ . But this also implies  $[\pi]_m^{-l} F_{n+p^{m-1}l}^{m,q}(S) \subset F_n^{m,q}(S)$  and hence we are done.

Combining Corollary 3.1.3 and Lemma 3.1.4, we get

**Corollary 3.1.5.** For every  $m \ge 1$  and  $q \ge 0$ , there is a canonical decomposition

$$\theta_K^{m,q} : \left(\bigoplus_{n<0} F_n^{m,q}(S)\right) \bigoplus \left(\prod_{n\geq 0} F_n^{m,q}(S)\right) \xrightarrow{\cong} W_m \Omega_K^q.$$

*Proof.* We first claim that the canonical map

$$W_m\Omega^q_A(\log\pi) \otimes_{W_m(A)} W_m(K) \to W_m\Omega^q_K \tag{3.1.3}$$

is an isomorphism of  $W_m(K)$ -modules. Indeed, we know by [30, 0.1.5.3] that  $W_m(K) = W_m(A)[u^{-1}]$ , where the latter is the localization of  $W_m(A)$  obtained by inverting  $u = [\pi]_m$ . This implies that  $M \otimes_{W_m(A)} W_m(K) \cong M[u^{-1}]$  for  $M \in \{W_m \Omega_A^q, W_m \Omega_A^q(\log \pi)\}$ . Since  $W_m \Omega_A^q \subset W_m \Omega_A^q(\log \pi) \subset W_m \Omega_K^q$ , we get  $W_m \Omega_A^q[u^{-1}] \subset W_m \Omega_A^q(\log \pi)[u^{-1}] \subset W_m \Omega_K^q$ . The claim now follows because the composite inclusion is an isomorphism by [26, Thm. C]. By (3.1.3), we now have

$$W_{m}\Omega_{K}^{q} \cong W_{m}\Omega_{A}^{q}(\log\pi)[u^{-1}] \cong \varinjlim_{l\geq 0}[\pi]_{m}^{-l}W_{m}\Omega_{A}^{q}(\log\pi)$$

$$\cong^{1} \varinjlim_{l\geq 0}[\pi]_{m}^{-l}\prod_{n\geq 0}F_{n}^{m,q}(S) \cong \varinjlim_{l\geq 0}\prod_{n\geq 0}[\pi]_{m}^{-l}F_{n}^{m,q}(S)$$

$$\cong^{2} \varinjlim_{l\geq 0}\prod_{n\geq 0}F_{n-p^{m-1}l}^{m,q}(S) \cong \varinjlim_{l\geq 0}\prod_{n\geq -p^{m-1}l}F_{n}^{m,q}(S)$$

$$\cong \left[\varinjlim_{l\geq 0}\left(\bigoplus_{-p^{m-1}l\leq n\leq -1}F_{n}^{m,q}(S)\right)\right] \bigoplus \left[\varinjlim_{l\geq 0}\left(\prod_{n\geq 0}F_{n}^{m,q}(S)\right)\right]$$

$$\cong \left[\varinjlim_{l\geq 0}\left(\bigoplus_{-l\leq n\leq -1}F_{n}^{m,q}(S)\right)\right] \bigoplus \left(\prod_{n\geq 0}F_{n}^{m,q}(S)\right)$$

$$\cong \left(\bigoplus_{n< 0}F_{n}^{m,q}(S)\right) \bigoplus \left(\prod_{n\geq 0}F_{n}^{m,q}(S)\right),$$

where the isomorphisms  $\cong^1$  and  $\cong^2$  hold by Corollary 3.1.3 (induced by  $(\theta_A^{m,q})^{-1}$ ) and Lemma 3.1.4, respectively.

Our task now is to describe the image of  $\operatorname{fil}_n W_m \Omega_K^q$  under  $\theta_K^{m,q}$ . We shall achieve this in several steps. We begin by checking some further properties of the groups  $F_n^{m,q}(S)$ .

**Lemma 3.1.6.** For any  $n \in \mathbb{Z}, m \ge 1, q \ge 0$ , we have the following.

1. 
$$d(F_n^{m,q}(S)) \in F_n^{m,q+1}(S)$$
.  
2.  $V(F_n^{m,q}(S)) \in F_n^{m+1,q}(S)$ .  
3.  $F(F_n^{m,q}(S)) \in F_n^{m-1,q}(S)$ .  
4.  $R(F_n^{m,q}(S)) = \begin{cases} F_{n/p}^{m-1,q}(S) & \text{if } p | n \\ 0 & \text{if } p \neq n . \end{cases}$ 

*Proof.* We divide the proof into two cases: when  $p^{m-1}|n$  and when  $p^{m-1} + n$ . Case  $1: n = p^{m-1}i$ .

In this case,  $F_n^{m,q}(S) = (W_m \Omega_S^q + W_m \Omega_S^{q-1} \operatorname{dlog}([\pi]_m))[\pi]_m^i$ . Since  $d(W_m \Omega_S^q [\pi]_m^i) \subset W_m \Omega_S^{q+1}[\pi]_m^i + W_m \Omega_S^q [\pi]_m^i \operatorname{dlog}([\pi]_m) = F_n^{m,q+1}(S)$ , and,  $d(W_m \Omega_S^{q-1}[\pi]_m^i \operatorname{dlog}([\pi]_m)) \subset W_m \Omega_S^q [\pi]_m^i \operatorname{dlog}([\pi]_m) \subset F_n^{m,q+1}(S)$ , the claim (1) follows.

For (2), note that if  $i \in I_p$ , we have  $V(W_m \Omega_S^q[\pi]_m^i) \in F_n^{m+1,q}(S)$  by latter's definition. Next, we have

$$V(W_m \Omega_S^{q-1} \operatorname{dlog}([\pi]_m)[\pi]_m^i) = V(W_m \Omega_S^{q-1} d([\pi]_m^i))$$
  

$$\subset V d(W_m \Omega_S^{q-1}[\pi]_m^i) + V(W_m \Omega_S^q[\pi]_m^i)$$
  

$$\subset F_n^{m+1,q}(S).$$

If i = pi', then  $n = p^m i'$  and  $V(W_m \Omega_S^q[\pi]_m^i) \subset W_{m+1} \Omega_S^q[\pi]_{m+1}^{i'} \subset F_n^{m+1,q}(S)$ . Also, we have  $V(W_m \Omega_S^{q-1} \text{dlog}[\pi]_m[\pi]_m^i) \subset W_{m+1} \Omega_S^{q-1} \text{dlog}[\pi]_{m+1}[\pi]_{m+1}^{i'} \subset F_n^{m+1,q}(S)$ .

# 3.1. FILTERED DE RHAM-WITT COMPLEX OF COMPLETE LOCAL $\mathbb{F}_p\text{-}\mathrm{ALGEBRAS}$

For (3), write  $n = p^{m-2}i'$ , where i' = pi. We then get  $F(W_m \Omega_S^q[\pi]_m^i) \subset W_{m-1} \Omega_S^q[\pi]_{m-1}^{i'}$ which lies in  $F_n^{m-1,q}(S)$ . Similarly,

$$F(W_m\Omega_S^{q-1}\mathrm{dlog}([\pi]_m)[\pi]_m^i) \subset W_{m-1}\Omega_S^{q-1}\mathrm{dlog}([\pi]_{m-1})[\pi]_{m-1}^{i'} \subset F_n^{m-1,q}(S).$$

For (4), write  $n' = n/p = p^{m-2i}$  so that  $R(W_m \Omega_S^q[\pi]_m^i) = W_{m-1} \Omega_S^q[\pi]_{m-1}^i$ . Similarly,  $R(W_m \Omega_S^{q-1} \text{dlog}[\pi]_m[\pi]_m^i) = W_{m-1} \Omega_S^{q-1} \text{dlog}[\pi]_{m-1}[\pi]_{m-1}^i$ . Hence  $R(F_n^{m,q}(S)) = F_{n'}^{m-1,q}(S)$ .

Case 2:  $n = p^{m-1-s}i$  with  $s \ge 1$  and  $|i| \in I_p$ . In this case, we have  $F_n^{m,q}(S) = V^s(W_r\Omega_S^q[\pi]_r^i) + dV^s(W_r\Omega_S^{q-1}[\pi]_r^i)$ , where r = m - s = (m+1) - (s+1). But this implies that  $d(F_n^{m,q}(S)) = dV^s(W_r\Omega_S^q[\pi]_r^i) \subset F_n^{m,q+1}(S)$ . This proves (1).

For (2), we note that  $V(F_n^{m,q}(S)) = V^{s'}(W_r\Omega_S^q[\pi]_r^i) + VdV^s(W_r\Omega_S^{q-1}[\pi]_r^i) \subset V^{s'}(W_r\Omega_S^q[\pi]_r^i) + dV^{s'}(W_r\Omega_S^{q-1}[\pi]_r^i) = F_n^{m+1,q}(S)$  if we let s' = s + 1.

For (3), suppose first that s = 1 so that  $n = p^{m-2}i$ . Then

$$F(F_n^{m,q}(S)) = FV(W_{m-1}\Omega_S^q[\pi]_{m-1}^i) + FdV(W_{m-1}\Omega_S^{q-1}[\pi]_{m-1}^i)$$
  

$$\subset W_{m-1}\Omega_S^q[\pi]_{m-1}^i + d(W_{m-1}\Omega_S^{q-1}[\pi]_{m-1}^i)$$
  

$$\subset (W_{m-1}\Omega_S^q + W_{m-1}\Omega_S^{q-1}d\log([\pi]_{m-1}))[\pi]_{m-1}^i = F_n^{m-1,q}(S).$$

If  $s \ge 2$ , we let  $s' = s - 1 \ge 1$  so that  $n = p^{m-2-s'}i$  and r = (m-1) - s'. This yields

$$F(F_n^{m,q}(S)) = FV^s(W_r\Omega_S^q[\pi]_r^i) + FdV^s(W_r\Omega_S^{q-1}[\pi]_r^i) \subset V^{s'}(W_r\Omega_S^q[\pi]_r^i) + dV^{s'}(W_r\Omega_S^{q-1}[\pi]_r^i) = F_n^{m-1,q}(S).$$

Finally, for (4), note that

$$RV^{s}(W_{r}\Omega_{S}^{q}[\pi]_{r}^{i}) + RdV^{s}(W_{r}\Omega_{S}^{q-1}[\pi]_{r}^{i}) = V^{s}(W_{r-1}\Omega_{S}^{q}[\pi]_{r-1}^{n'/p^{m-2-s}}) + dV^{s}(W_{r-1}\Omega_{S}^{q-1}[\pi]_{r-1}^{n'/p^{m-2-s}}),$$

where n' = n/p. The latter group is in  $F_{n'}^{m-1,q}(S)$  if s < m-1 and is zero if s = m-1. This concludes the proof.

**Lemma 3.1.7.** We have  $F_n^{m,0}(S) \cdot F_{n'}^{m,q}(S) \subset F_{n+n'}^{m,q}(S)$  for  $q, n, n' \ge 0, m \ge 1$ .

Proof. First note that  $W_m(S) \cdot F_{n'}^{m,q}(S) \subset F_{n'}^{m,q}(S)$ . Indeed, if  $p^{m-1} \mid n'$  then this is obvious. If  $p^{m-1} \neq n'$  then only need to show  $W_m(S) \cdot dV^s(W_r(S)[\pi]_r^i) \subset F_{n'}^{m,q}(S)$ , where  $n' = p^{m-1-s}i$ ,  $|i| \in I_p$ . But this follows easily by Leibniz rule.

We break the proof into two cases. We shall assume  $1 \le s \le m-1$  and write r = m-s. Case 1:  $n = p^{m-1}i$ . In this case,  $F_n^{m,0}(S) = W_m(S)[\pi]_m^i$ . Hence, we get

$$F_n^{m,0}(S) \cdot F_{n'}^{m,q}(S) = W_m(S)[\pi]_m^i \cdot F_{n'}^{m,q}(S) \subset F_{n+n'}^{m,q}(S),$$

by the above observation and Lemma 3.1.4.

Case 2:  $n = p^{m-1-s}i$ , where  $|i| \in I_p$ . In this case,  $F_n^{m,0}(S) = V^s(W_r(S)[\pi]_{m-s}^i)$ . Thus, we get

$$F_{n}^{m,0}(S) \cdot F_{n'}^{m,q}(S) = V^{s}(W_{r}(S)[\pi]_{r}^{i}) \cdot F_{n'}^{m,q}(S)$$

$$\subset^{1} V^{s}([\pi]_{r}^{i}.F_{n'}^{m-s,q}(S)) \subset^{2} V^{s}(F_{n'+n}^{m-s,q}(S))$$

$$\subset^{3} F_{n'+n}^{m-s,q}(S),$$

where the equality  $\subset^1$  and  $\subset^3$  uses Lemma 3.1.6, item (3) and (2), respectively and also the above observation, while  $\subset^2$  uses Lemma 3.1.4.

**Lemma 3.1.8.**  $F_n^{m,q}(S) \hookrightarrow \operatorname{fil}_{-n} W_m \Omega_K^q$  (by  $\theta_A^{m,q}$ ) for  $n, q \ge 0, m \ge 1$ .

*Proof.* First, by Theorem 3.1.1, for all  $q, i \ge 0$ , we have

$$W_m \Omega_S^q[\pi]_m^i \hookrightarrow W_m \Omega_A^q[\pi]_m^i \text{ and } W_m \Omega_S^q[\pi]_m^i \operatorname{dlog}([\pi]_m) \hookrightarrow W_m \Omega_A^q[\pi]_m^i \operatorname{dlog}([\pi]_m).$$

Now we consider two cases.

If  $n = p^{m-1}i$  and  $i \ge 0$ , we get

$$F_n^{m,q}(S) = A_i^{m,q}(S) = (W_m \Omega_S^q + W_m \Omega_S^{q-1} \operatorname{dlog}([\pi]_m))[\pi]_m^i$$
  

$$\hookrightarrow (W_m \Omega_A^q + W_m \Omega_A^{q-1} \operatorname{dlog}([\pi]_m))[\pi]_m^i$$
  

$$= \operatorname{fil}_{-n} W_m \Omega_K^q,$$

where the last equality holds by Lemma 2.1.7(9).

If  $n = p^{m-1-s}i$  with  $1 \le s \le m-1$  and  $|i| \in I_p$ , then

$$F_n^{m,q}(S) = B_{i,s}^{m,q} = V^s(W_r\Omega_S^q[\pi]_r^i) + dV^s(W_r\Omega_S^{q-1}[\pi]_r^i)$$
  

$$\Rightarrow V^s(\operatorname{fil}_{-p^{r-1}i}W_r\Omega_K^q) + dV^s(\operatorname{fil}_{-p^{r-1}i}W_r\Omega_K^{q-1})$$
  

$$= V^s(\operatorname{fil}_{-n}W_r\Omega_K^q) + dV^s(\operatorname{fil}_{-n}W_r\Omega_K^{q-1})$$
  

$$\subset \operatorname{fil}_{-n}W_m\Omega_K^q,$$

where r = m - s and the last inclusion holds by Lemma 2.1.7(4), (6).

## 3.1. FILTERED DE RHAM-WITT COMPLEX OF COMPLETE LOCAL $\mathbb{F}_p$ -ALGEBRAS

In order to describe  $\operatorname{fil}_n W_m \Omega_K^q$  in terms of  $\{F_n^{m,q}\}$ , we need to dispose of the case q = 0 first. We do this in what follows next. One deduces from Lemma 3.1.8 for q = 0 that  $F_n^{m,0}(S) \subset \operatorname{fil}_{-n} W_m(K)$  for  $n \ge 0$ .

**Lemma 3.1.9.** For  $n \in \mathbb{Z}$ , the composite map  $\lambda_n^{m,0} \colon F_n^{m,0}(S) \hookrightarrow \operatorname{fil}_{-n} W_m(K) \twoheadrightarrow \operatorname{gr}_{-n} W_m(K)$ is an isomorphism.

*Proof.* The assertion of the lemma is obvious if m = 1, because  $F_n^{1,0}(S) = S\pi^n$  and  $\operatorname{fil}_{-n}W_1(K) = \pi^n A$ .

For m > 1, Lemma 2.1.2 implies that the sequence

$$0 \to \operatorname{fil}_{n} W_{1}(K) \xrightarrow{V^{m-1}} \operatorname{fil}_{n} W_{m}(K) \xrightarrow{R} \operatorname{fil}_{\lfloor n/p \rfloor} W_{m-1}(K) \to 0$$
(3.1.4)

is exact for every  $n \in \mathbb{Z}$ . Comparing this sequence for -n and -n-1 and noting that  $\operatorname{fil}_{-n-1}W_m(K) \subset \operatorname{fil}_{-n}W_m(K)$ , for all  $m \ge 1, n \in \mathbb{Z}$ , we get an exact sequence

$$0 \to \operatorname{gr}_{-n} W_1(K) \xrightarrow{V^{m-1}} \operatorname{gr}_{-n} W_m(K) \xrightarrow{R} \frac{\operatorname{fil}_{\lceil n/p \rceil} W_{m-1}(K)}{\operatorname{fil}_{\lceil n+1/p \rceil} W_{m-1}(K)} \to 0.$$
(3.1.5)

If we write n + 1 = pt + q with  $0 \le q < p$ , we see that  $V^{m-1}$  is an isomorphism if  $q \ne 1$ . Since s = m - 1 (cf. Definition 3.1.2) in this case, we find also that the image of  $V^{m-1}$  is  $F_n^{m,0}(S)$ .

If q = 1, we can rewrite (3.1.5) as

$$0 \to \operatorname{gr}_{-n} W_1(K) \xrightarrow{V^{m-1}} \operatorname{gr}_{-n} W_m(K) \xrightarrow{R} \operatorname{gr}_{-t} W_{m-1}(K) \to 0.$$

We now look at the diagram

It is easy to check that this diagram is commutative and the top row is exact. Now, the left and the right vertical arrows are bijective by induction. We conclude that the same is true for the middle vertical arrow too.  $\Box$ 

Before we proceed further, we compare the topologies of  $W_m \Omega_A^q(\log \pi)$  induced by various filtrations. We let  $\tau_1$  be the topology on  $W_m \Omega_A^q(\log \pi)$  induced by the filtration  $\{W_m \Omega_A^q(\log \pi)(n)\}_{n\geq 0}$  (cf. 2.1.4, § 3.1). We let  $\tau_2$  be the topology on  $W_m \Omega_A^q(\log \pi)$ induced by the filtration  $\{W_m((\pi^n))W_m \Omega_A^q(\log \pi)\}_{n\geq 0}$ . We let  $\tau_3$  be the topology on  $W_m \Omega_A^q(\log \pi)$  induced by the filtration  $\{I^n W_m \Omega_A^q(\log \pi)\}_{n\geq 0}$ , where  $I = W_m((\pi))$ . Finally, we let  $\tau_4$  be the topology on  $W_m \Omega_A^q(\log \pi)$  induced by the filtration  $\{fil_{-n} W_m \Omega_K^q\}_{n\geq 0}$ .

#### Lemma 3.1.10. The topologies defined above all agree.

*Proof.* The topologies  $\tau_1$ ,  $\tau_2$  and  $\tau_3$  will agree on  $W_m \Omega_A^q(\log \pi)$  if they do so on  $W_m(A)$ . The agreement between  $\tau_1$  and  $\tau_2$  in this special case is an elementary exercise and the the same between  $\tau_2$  and  $\tau_3$  follows from [15, Cor. 2.3]. The agreement between  $\tau_1$  and  $\tau_4$  on  $W_m \Omega_A^q(\log \pi)$  follows from Lemma 2.1.7(9).

**Corollary 3.1.11.** For  $n \ge 0$ , the map  $\theta_A^{m,q}$  of (3.1.2) induces an isomorphism

$$\theta_n^{m,0} \colon \prod_{n' \ge n} F_{n'}^{m,0}(S) \xrightarrow{\cong} \operatorname{fil}_{-n} W_m(K).$$

Proof. We first conclude from Theorem 3.1.1 (or [15, Cor 2.3]) that  $W_m(A)$  is complete with respect to its *I*-adic topology. Since *A* is *F*-finite, the ring  $W_m(A)$  is Noetherian (cf. [58, Lem. 2.9]). Furthermore, Lemma 2.1.7(1) implies that  $\operatorname{fil}_{-n}W_m(K)$  is an ideal of  $W_m(A)$ . It follows that the  $\operatorname{fil}_{-n}W_m(K)$  is *I*-adically complete, and hence, by Lemma 3.1.10, complete with respect to  $\tau_4$  topology. That is, the canonical map  $\operatorname{fil}_{-n}W_m(K) \to \varprojlim_{n'\geq 0} \frac{\operatorname{fil}_{-n-n'}W_m(K)}{\operatorname{fil}_{-n-n'}W_m(K)}$  is an isomorphism of  $W_m(A)$ -modules. Meanwhile, one deduces from Lemma 3.1.9 that the canonical map  $\bigoplus_{i=0}^{n'-1} F_{n+i}^{m,0}(S) \to \frac{\operatorname{fil}_{-n-M'}W_m(K)}{\operatorname{fil}_{-n-n'}W_m(K)}$  is an isomorphism. Passing to the limit as  $n' \to \infty$ , we get an isomorphism

$$\prod_{n' \ge n} F_{n'}^{m,0}(S) \xrightarrow{\cong} \operatorname{fil}_{-n} W_m(K),$$

which is induced by the canonical map  $F_{n'}^{m,0}(S) \to W_m(A)$  (i.e. by  $\theta_A^{m,0}$ ). Hence we achieve the proof of the lemma.

Our next task is to generalize Corollary 3.1.11 to the case when q > 0. For  $n \ge 0$ , we let  $\operatorname{fil}_{-n}^{\prime} W_m \Omega_K^q = \theta_A^{m,q} (\prod_{j\ge n} F_j^{m,q}(S))$ . By Corollary 3.1.3, this is an exhaustive descending filtration of  $W_m \Omega_A^q(\log \pi) = \operatorname{fil}_0^{\prime} W_m \Omega_K^q$ .

**Lemma 3.1.12.** For  $n, q \ge 0, m \ge 1$ ,  $\operatorname{fil}_{-n}^{\prime} W_m \Omega_K^q \subset W_m \Omega_A^q(\log \pi)$  is a  $W_m(A)$ -submodule.

Proof. We only need to show that  $\operatorname{fil}_{-n}^{\prime}W_m\Omega_K^q$  is closed under the action of  $W_m(A)$ . We let  $\omega = \sum_{j\geq n} \alpha_j \in \operatorname{fil}_{-n}^{\prime}W_m\Omega_K^q$ , where  $\alpha_j \in \theta_A^{m,q}(F_j^{m,q}(S))$ . Write  $\omega_i = \sum_{j=n}^i \alpha_j$ . Since  $W_m\Omega_A^q(\log \pi)$  is finitely generated  $W_m(A)$ -module (cf. Lemma 2.1.7(1)), it is complete with respect to *I*-adic topology. So, we can write  $\omega = \lim_{i\to\infty} \omega_i$ , where the limit is taken with respect to  $\tau_3$  topology. Now let  $\gamma = \sum_{j\geq 0} \beta_j \in W_m(A)$ , where  $\beta_j \in \theta_A^{m,0}(F_j^{m,0}(S))$ . Similarly to above, we can write  $\gamma = \lim_{i\to\infty} \gamma_i$ , where  $\gamma_i = \sum_{j=0}^i \beta_j$  and the limit is taken

# 3.1. FILTERED DE RHAM-WITT COMPLEX OF COMPLETE LOCAL $\mathbb{F}_p\text{-}\mathrm{ALGEBRAS}$

in 
$$\tau_3$$
. Then  $\lim_{i\to\infty} \gamma_i \cdot \omega_i = \gamma \cdot \omega$ . But  $\gamma_i \cdot \omega_i \subset \theta_A^{m,q}(\bigoplus_{j=n}^{2i} F_j^{m,q}(S))$  by Lemma 3.1.7. Hence  $\gamma \cdot \omega \in \sum_{j\geq n} \theta_A^{m,q}(F_j^{m,q}(S)) = \operatorname{fil}'_{-n} W_m \Omega_K^q$ .

**Lemma 3.1.13.**  $d(\operatorname{fil}_{-n}^{\prime}W_m\Omega_K^q) \subset \operatorname{fil}_{-n}^{\prime}W_m\Omega_K^{q+1}$  for every  $n, q \ge 0$  and  $m \ge 1$ .

Proof. We continue with the same notations of the proof of Lemma 3.1.12. For  $\omega \in \operatorname{fl}_{-n}^{\prime}W_m\Omega_K^q$ , we can write  $\omega = \lim_{i\to\infty}\omega_i$ , where the limit is taken with respect to  $\tau_3$ , and hence with respect to  $\tau_4$  by Lemma 3.1.10. We now let  $n' \geq 0$  be any integer. We can then find  $i \gg 0$  such that  $\omega - \omega_i \in \operatorname{fl}_{-n'}W_m\Omega_K^q$  for all  $i \gg n$ . This implies that  $d(\omega) - d(\omega_i) = d(\omega - \omega_i) \in \operatorname{fl}_{-n'}W_m\Omega_K^{q+1}$  for all  $i \gg n$  by Lemma 2.1.7(4). But this means that  $d(\omega) = \lim_{i\to\infty} d(\omega_i) = \sum_{j\geq n} d(\alpha_j)$ , where  $d(\alpha_j) \subset d(\theta_A^{m,q}(F_j^{m,q}(S))) = {}^1 \theta_A^{m,q+1}(d(F_j^{m,q}(S))) \subset {}^2 \theta_A^{m,q+1}(F_j^{m,q+1}(S))$ . Here the equality  $= {}^1$  holds because the map  $\theta_A^{m,q}$  is induced by the natural map  $F_j^{m,q}(S) \to W_m\Omega_A^q(\log\pi)$ , which commutes with d. On the other hand, the inclusion  $\subset {}^2$  holds by Lemma 3.1.6(1).

**Lemma 3.1.14.** We have  $\operatorname{fil}_{-n} W_m \Omega_K^q \subseteq \operatorname{fil}'_{-n} W_m \Omega_K^q$  for every  $n, q \ge 0$  and  $m \ge 1$ .

*Proof.* By definition of fil<sub>-n</sub> $W_m \Omega_K^q$  and Corollaries 3.1.3 and 3.1.11, we have

$$\begin{aligned} \operatorname{fil}_{-n} W_m \Omega_K^q &= \operatorname{fil}_{-n} W_m(K) W_m \Omega_A^q(\log \pi) + d(\operatorname{fil}_{-n} W_m(K)) W_m \Omega_A^{q-1}(\log \pi) \\ &= \operatorname{fil}'_{-n} W_m(K) W_m \Omega_A^q(\log \pi) + d(\operatorname{fil}'_{-n} W_m(K)) W_m \Omega_A^{q-1}(\log \pi) \\ &\subset \operatorname{fil}'_{-n} W_m(K) W_m \Omega_A^q(\log \pi) + d(\operatorname{fil}'_{-n} W_m(K) W_m \Omega_A^{q-1}(\log \pi)) \\ &+ \operatorname{fil}'_{-n} W_m(K) d(W_m \Omega_A^{q-1}(\log \pi)). \end{aligned}$$

Next, note that every element of  $\operatorname{fil}_{-n}^{\prime}W_m(K)W_m\Omega_A^q(\log \pi)$  is finite sum of elements of the form  $\gamma \cdot \omega$ , where  $\gamma \in \operatorname{fil}_{-n}^{\prime}W_m(K)$  and  $\omega \in W_m\Omega_A^q(\log \pi) = \operatorname{fil}_0^{\prime}W_m\Omega_K^q$ . Recalling the notations from the proof of Lemma 3.1.12, we write  $\omega = \lim_{i\to\infty} \omega_i$ , where  $\omega_i = \sum_{j=0}^i \alpha_j$  and  $\gamma = \lim_{i\to\infty} \gamma_i$ , where  $\gamma_i = \sum_{j=n}^i \beta_j$  (note the change in indices of  $\alpha_j$  and  $\beta_j$ ). The limits are taken in  $\tau_3$ . As in the proof of Lemma 3.1.12,  $\gamma \cdot \omega = \lim_{i\to\infty} \gamma_i \cdot \omega_i \subset \operatorname{fil}_{-n}^{\prime}W_m\Omega_K^q$ . This implies  $\operatorname{fil}_{-n}^{\prime}W_m(K)W_m\Omega_A^q(\log \pi) \subset \operatorname{fil}_{-n}^{\prime}W_m\Omega_K^q$ . Similarly,

$$\operatorname{fil}_{-n}' W_m(K) d(W_m \Omega_A^{q-1}(\log \pi)) \subset \operatorname{fil}_{-n}' W_m(K) W_m \Omega_A^q(\log \pi) \subset \operatorname{fil}_{-n}' W_m \Omega_K^q.$$

Finally,  $d(\operatorname{fil}'_{-n}W_m(K)W_m\Omega_A^{q-1}(\log \pi)) \subset d(\theta_A^{m,q-1}(\prod_{j\geq n}F_j^{m,q-1}(S))) = d(\operatorname{fil}'_{-n}W_m\Omega_K^{q-1})$ , and the latter group lies in  $\operatorname{fil}'_{-n}W_m\Omega_K^q$  by Lemma 3.1.13. This finishes the proof.  $\Box$ 

**Corollary 3.1.15.** For  $n \ge 0$ ,  $\operatorname{fil}_{-n}^{\prime} W_m \Omega_K^q$  is open (hence closed) in  $W_m \Omega_A^q(\log \pi)$  with respect to the topology  $\tau_i$  for  $1 \le i \le 4$ .

*Proof.* Combining Lemmas 3.1.12 and 3.1.14, we get  $\operatorname{fil}_{-n}^{\prime} W_m \Omega_K^q$  is  $\tau_4$ -open. Now the corollary follows from Lemma 3.1.10.

**Lemma 3.1.16.** We have  $\operatorname{fil}_{-n}^{\prime} W_m \Omega_K^q = \operatorname{fil}_{-n} W_m \Omega_K^q$  for every  $n \ge 0$ .

Proof. In view of Lemma 3.1.14, we only need to show that  $\operatorname{fil}_{-n}^{\prime}W_m\Omega_K^q \subset \operatorname{fil}_{-n}W_m\Omega_K^q$ . The latter assertion is obvious if n = 0 because  $\operatorname{fil}_{-n}W_m\Omega_K^q = W_m\Omega_A^q(\log \pi) = \operatorname{fil}_{-n}^{\prime}W_m\Omega_K^q$  by Corollary 3.1.3. We assume therefore that  $n \ge 1$ . In particular,  $\operatorname{fil}_{-n}^{\prime}W_m\Omega_K^q \subset W_m\Omega_A^q$ . We now let  $\omega \in \operatorname{fil}_{-n}^{\prime}W_m\Omega_K^q$ . Recalling the notations from the proof of Lemma 3.1.12 we have that  $\omega = \lim_{i\to\infty}\omega_i$ , where the limit is taken with respect to  $\tau_3$  (cf. [15, Proof of Thm. B, p. 488]). Meanwhile, Lemma 3.1.8 says that  $\omega_i \in \operatorname{fil}_{-n}W_m\Omega_K^q$  for each  $i \ge n$ . Since  $\operatorname{fil}_{-n}W_m\Omega_K^q$  is  $\tau_4$ -open (and hence closed) in  $W_m\Omega_A^q(\log \pi)$ , by Corollary 3.1.15; we conclude that  $\omega \in \operatorname{fil}_{-n}W_m\Omega_K^q$ .

We shall now describe  $\operatorname{fil}_n W_m \Omega_K^q$  for arbitrary  $n \in \mathbb{Z}$ . For any  $l \in \mathbb{Z}$ , we let  $t = n - p^{m-1}l$ .

**Lemma 3.1.17.** We have  $\operatorname{fil}_t W_m(K) = [\pi]_m^l \operatorname{fil}_n W_m(K)$  for every  $n \in \mathbb{Z}$ .

Proof. Let  $x = (a_{m-1}, \ldots, a_0) \in \operatorname{fil}_n W_m(K)$ , that is,  $a_i^{p^i} \pi^n \in A$  for each *i*. Then  $[\pi]_m^l x = (a_{m-1}\pi^l, a_{m-2}\pi^{pl}, \ldots, a_0\pi^{p^{m-1}l})$ . Letting  $b_i = a_i\pi^{p^{m-1-i}l}$ , we get  $(b_i)^{p^i}\pi^t = a_i^{p^i}\pi^{p^{m-1}l+t} = a_i^{p^i}\pi^n \in A$ . This implies that  $[\pi]_m^l \operatorname{fil}_n W_m(K) \subset \operatorname{fil}_t W_m(K)$  for all  $l, n \in \mathbb{Z}$ . Then, also, we have  $[\pi]_m^{-l} \operatorname{fil}_t W_m(K) \subset \operatorname{fil}_n W_m(K)$  and hence the lemma.

**Lemma 3.1.18.** We have  $\operatorname{fil}_t W_m \Omega_K^q = [\pi]_m^l \operatorname{fil}_n W_m \Omega_K^q$  for every  $n \in \mathbb{Z}$ .

*Proof.* By definition,  $\operatorname{fil}_t W_m \Omega_K^q = \operatorname{fil}_t W_m(K) W_m \Omega_A^q(\log \pi) + d(\operatorname{fil}_t W_m(K)) W_m \Omega_A^{q-1}(\log \pi)$ . By Lemma 3.1.17,  $\operatorname{fil}_t W_m(K) W_m \Omega_A^q(\log \pi) = [\pi]_m^l \operatorname{fil}_n W_m(K) W_m \Omega_A^q(\log \pi)$  and

$$d(\operatorname{fil}_{t}W_{m}(K))W_{m}\Omega_{A}^{q-1}(\log \pi) = d([\pi]_{m}^{l}\operatorname{fil}_{n}W_{m}(K))W_{m}\Omega_{A}^{q-1}(\log \pi)$$

$$\subset [\pi]_{m}^{l}d(\operatorname{fil}_{n}W_{m}(K))W_{m}\Omega_{A}^{q-1}(\log \pi)$$

$$+\operatorname{fil}_{n}W_{m}(K)[\pi]_{m}^{l}W_{m}\Omega_{A}^{q-1}(\log \pi)\operatorname{dlog}([\pi]_{m})$$

$$\subset [\pi]_{m}^{l}(d(\operatorname{fil}_{n}W_{m}(K))W_{m}\Omega_{A}^{q-1}(\log \pi)$$

$$+\operatorname{fil}_{n}W_{m}(K)W_{m}\Omega_{A}^{q-1}\operatorname{dlog}([\pi]_{m}))$$

$$= [\pi]_{m}^{l}\operatorname{fil}_{n}W_{m}\Omega_{K}^{q}.$$

This shows that  $\operatorname{fil}_t W_m \Omega_K^q \subseteq [\pi]_m^l \operatorname{fil}_n W_m \Omega_K^q$  for all  $l \in \mathbb{Z}$ . The reverse inclusion also follows like before (cf. Lemma 3.1.17).

We can now describe the image of fil<sub>•</sub> $W_m \Omega_K^q$  under the isomorphism  $\theta_K^{m,q}$  of Corollary 3.1.5 as follows.

## 3.1. FILTERED DE RHAM-WITT COMPLEX OF COMPLETE LOCAL $\mathbb{F}_p\text{-}\mathrm{ALGEBRAS}$

**Proposition 3.1.19.** For  $n \in \mathbb{Z}$ , there is a canonical decomposition

$$\theta_K^{m,q} \colon \prod_{j \ge -n} F_j^{m,q}(S) \xrightarrow{\cong} \operatorname{fil}_n W_m \Omega_K^q$$

*Proof.* For  $n \leq 0$ , this follows from Lemma 3.1.16. Suppose now  $n \geq 0$ . Choose  $l \gg 0$  such that  $t = n - p^{m-1}l < 0$ . Lemma 3.1.18 then says that  $\operatorname{fil}_n W_m \Omega_K^q = [\pi]_m^{-l} \operatorname{fil}_t W_m \Omega_K^q$ . On the other hand, we have

$$[\pi]_{m}^{-l} \operatorname{fil}_{t} W_{m} \Omega_{K}^{q} \cong [\pi]_{m}^{-l} \left( \prod_{j \ge -t} F_{j}^{m,q}(S) \right) \cong \prod_{j \ge -t} [\pi]_{m}^{-l} F_{j}^{m,q}(S) \cong \prod_{j \ge -t} F_{j-p^{m-1}l}^{m,q}(S),$$

where the first and the last isomorphisms are by Lemma 3.1.16 (induced by  $(\theta_A^{m,q})^{-1}$ ) and Lemma 3.1.4, respectively. By a change of variable  $j' = j - p^{m-1}l$ , we get

$$\prod_{j \ge -t} F_{j-p^{m-1}l}^{m,q}(S) = \prod_{j' \ge -n} F_{j'}^{m,q}(S)$$

Hence, we get an isomorphism  $\operatorname{fil}_n W_m \Omega_K^q \cong \prod_{j' \ge -n} F_{j'}^{m,q}(S)$ , whose inverse is induced by the canonical map  $F_{j'}^{m,q}(S) \to \operatorname{fil}_n W_m \Omega_K^q$ , for  $j' \ge -n$  (because  $\cong^1$  is induced by  $(\theta_A^{m,q})^{-1}$ ). We conclude that the resulting (inverse) isomorphism is  $\theta_K^{m,q}$ .

Our first application of Proposition 3.1.19 is the following.

**Proposition 3.1.20.** For every  $n \in \mathbb{Z}$ , there is a short exact sequence

$$0 \to V^{m-1}(\operatorname{fil}_n\Omega_K^q) + dV^{m-1}(\operatorname{fil}_n\Omega_K^{q-1}) \to \operatorname{fil}_nW_m\Omega_K^q \xrightarrow{R} \operatorname{fil}_{\lfloor n/p \rfloor}W_{m-1}\Omega_K^q \to 0.$$

*Proof.* Surjectivity of R is already shown in Lemma 2.1.7(7). To prove the exactness in the middle, we let  $A = V^{m-1}(\operatorname{fil}_n \Omega_K^q) + dV^{m-1}(\operatorname{fil}_n \Omega_K^{q-1})$  and  $B = \operatorname{Ker}(R)$ . From the exact sequence (cf. Proposition 1.1.6(1))

$$0 \to V^{m-1}(\Omega_K^q) + dV^{m-1}(\Omega_K^{q-1}) \to W_m \Omega_K^q \xrightarrow{R} W_{m-1}\Omega_K^q \to 0, \qquad (3.1.7)$$

and Lemma 2.1.7(4) and (6), we see that  $A \subset B = (V^{m-1}(\Omega_K^q) + dV^{m-1}(\Omega_K^{q-1})) \cap \operatorname{fil}_n W_m \Omega_K^q$ 

We now let  $x \in B$ . By Proposition 3.1.19, we can uniquely write  $x = \sum_{i \ge -n} a_i$  with  $a_i \in F_i^{m,q}(S)$ . By Corollary 3.1.5, we can also write  $x = V^{m-1}(\sum_{i \ge -l} b_i) + dV^{m-1}(\sum_{i \ge -l} c_i)$  for some  $l \gg |n|$  and  $b_i \in F_i^{1,q}(S)$ ,  $c_i \in F_i^{1,q-1}(S)$ . Hence, we get  $\sum_{i \ge -n} a_i = \sum_{i \ge -l} (V^{m-1}(b_i) + dV^{m-1}(c_i))$ . Letting  $a'_i = (V^{m-1}(b_i) + dV^{m-1}(c_i))$ , we get from Lemma 3.1.6 that  $a'_i \in F_i^{m,q}(S)$ . We conclude from Corollary 3.1.5 that  $a_i = a'_i$  for each  $i \ge -l$ . In particular,  $a'_i = 0$  for i < -n.

We thus get

$$x = \sum_{i \ge -n} V^{m-1}(b_i) + \sum_{i \ge -n} dV^{m-1}(c_i) = V^{m-1}(\sum_{i \ge -n} b_i) + dV^{m-1}(\sum_{i \ge -n} c_i)$$

Note here that V and d are continuous with respect to the topology  $\tau_2$  by virtue of Lemmas 2.1.7 and 3.1.10. Since  $b_i \in F_i^{1,q}(S)$  and  $c_i \in F_i^{1,q-1}(S)$ , Proposition 3.1.19 implies that  $\sum_{i\geq -n} b_i \in \operatorname{fil}_n \Omega_K^q$  and  $\sum_{i\geq -n} c_i \in \operatorname{fil}_n \Omega_K^{q-1}$ . Letting  $b = \sum_{i\geq -n} b_i$  and  $c = \sum_{i\geq -n} c_i$ , we get that  $x = V^{m-1}(b) + dV^{m-1}(c)$ , where  $b \in \operatorname{fil}_n \Omega_K^q$  and  $c \in \operatorname{fil}_n \Omega_K^{q-1}$ . It follows that  $x \in A$ .

### 3.2 Cartier homomorphism I

Recall from [32] (see also [22, § 10]) that for any regular  $\mathbb{F}_p$ -scheme X, there is a unique Cartier homomorphism  $C: Z_1 W_m \Omega_X^q \to W_m \Omega_X^q$  such that  $W_{m+1} \Omega_X^q \xrightarrow{R} W_m \Omega_X^q$ is the composition  $W_{m+1} \Omega_X^q \xrightarrow{F} Z_1 W_m \Omega_X^q \xrightarrow{C} W_m \Omega_X^q$  and  $Z_1 W_m \Omega_X^q := \text{Image}(F) =$  $\text{Ker}(W_m \Omega_X^q \xrightarrow{F^{m-1}d} \Omega_X^{q+1})$ . Our next goal is to explain the behavior of the Cartier homomorphism when restricted to  $\text{fl}_n W_m \Omega_K^q$ . We first need to prove the following lemma. We let  $Z_1 \text{fl}_n W_m \Omega_K^q := \text{fl}_n W_m \Omega_K^q \cap Z_1 W_m \Omega_K^q$  for  $n \in \mathbb{Z}$ .

**Lemma 3.2.1.**  $Z_1 \operatorname{fil}_n W_m \Omega_K^q = F(\operatorname{fil}_n W_{m+1} \Omega_K^q)$ . In particular the Cartier homomorphism restricts to a map

$$C: Z_1 \operatorname{fil}_n W_m \Omega_K^q \to \operatorname{fil}_{|n/p|} W_m \Omega_K^q.$$

Proof. We only need to show that  $\operatorname{fl}_n W_m \Omega_K^q \cap Z_1 W_m \Omega_K^q \subset F(\operatorname{fl}_n W_{m+1} \Omega_K^q)$ . We now let  $x \in \operatorname{fl}_n W_m \Omega_K^q \cap Z_1 W_m \Omega_K^q = \operatorname{fl}_n W_m \Omega_K^q \cap F(W_{m+1} \Omega_K^q)$ . By Corollary 3.1.5, Lemma 3.1.6 and Proposition 3.1.19, we can write  $x = \sum_{i \ge -n} a_i = \sum_{i \ge -l} a'_i = F(\sum_{i \ge -l} b_i)$  with  $a_i \in F_i^{m,q}(S)$  and  $a'_i = F(b_i) \in F_i^{m,q}(S)$  for some  $b_i \in F_i^{m+1,q}(S)$  and for some l >> |n|. By the uniqueness of the decomposition in Corollary 3.1.5, we must have  $a_i = a'_i$  for every  $i \ge -l$ . In particular,  $a'_i = 0$  for i < -n. This implies that  $x = \sum_{i \ge -n} a_i = F(\sum_{i \ge -n} b_i)$ . Since  $\sum_{i \ge -n} b_i \in \operatorname{fl}_n W_{m+1} \Omega_K^q$ , the desired claim follows.

*Remark* 3.2.2. In the next chapter, we shall also prove a Cartier isomorphism (cf. Lemma 4.3.3), which is analogous to the classical case.  $\Box$ 

The following key proposition will be used later in  $\S$  5 to study local Kato-filtration.

**Proposition 3.2.3.** Let  $n \ge 0$  and  $\omega \in Z_1 W_m \Omega_K^q$ . Then  $\omega \in Z_1 \operatorname{fil}_n W_m \Omega_K^q$  if and only if  $(1-C)(\omega) \in \operatorname{fil}_n W_m \Omega_K^q$ .

#### 3.2. CARTIER HOMOMORPHISM I

Proof. If  $\omega \in Z_1 \operatorname{fil}_n W_m \Omega_K^q$ , then  $C(\omega) = CF(\omega') = R(\omega')$  for some  $\omega' \in \operatorname{fil}_n W_{m+1} \Omega_K^q$  by Lemma 3.2.1. It follows from Lemma 3.1.6 that  $C(\omega) \in \operatorname{fil}_{\lfloor n/p \rfloor} W_m \Omega_K^q \subset \operatorname{fil}_n W_m \Omega_K^q$ . In particular, the map  $Z_1 \operatorname{fil}_n W_m \Omega_K^q \xrightarrow{1-C} \operatorname{fil}_n W_m \Omega_K^q$  is defined.

Suppose now that  $\omega \in Z_1 W_m \Omega_K^q$  is such that  $(1 - C)(\omega) \in \operatorname{fil}_n W_m \Omega_K^q$ . We can write  $\omega = F(\alpha)$  for some  $\alpha \in W_{m+1} \Omega_K^q$ . This yields  $(1 - C)(\omega) = (1 - C)F(\alpha) = F(\alpha) - R(\alpha)$ . By Corollary 3.1.5, we can write  $\alpha$  uniquely as  $\alpha = \sum_{i \ge -l} a_i$  for some  $a_i \in F_i^{m+1,q}(S)$  and  $l \gg 0$ . By Lemma 3.1.6, we get

$$F(\alpha) - R(\alpha) = \sum_{i \ge -l} F(a_i) - \sum_{i \ge -l} R(a_i) \in \operatorname{fil}_n W_m \Omega_K^q.$$
(3.2.1)

If  $l \leq n$ , then  $-n \leq -l$  and hence  $\alpha \in \operatorname{fil}_n W_{m+1}\Omega_K^q$  so that  $\omega \in Z_1 \operatorname{fil}_n W_m \Omega_K^q$ . Let l > n. Lemma 3.1.6 says that  $R(a_{-l}) \in F_{-l/p}^{m,q}(S)$  if p|l and  $R(a_{-l}) = 0$  if  $|l| \in I_p$ . Since  $F(a_{-l}) \in F_{-l}^{m,q}$  and  $\sum_{i\geq -l+1} F(a_i) - \sum_{i\geq -l} R(a_i) \in \operatorname{fil}_{l-1} W_m \Omega_K^q$ , it follows from (3.2.1) that  $F(a_{-l}) = 0$ . An induction argument says that  $F(a_i) = 0$  for all i < -n. Thus, we get  $\omega = F(\alpha) = \sum_{i\geq -n} F(a_i) \in \operatorname{fil}_n W_m \Omega_K^q$ .

## Chapter 4

# Structure of the filtered de Rham-Witt complex II

The main purpose of this chapter is to provide a description of the filtered de Rham-Witt complex of a regular local ring, based on the description given in the previous chapter. Finally, we extend various results of the de Rham-Witt complex of regular schemes to the filtered setup.

### 4.1 de Rham complex of multivariate power series ring

We let S be an F-finite Noetherian regular local  $\mathbb{F}_p$ -algebra and let  $A = S[[x_1, \ldots, x_d]]$ . We let  $1 \leq r \leq d$  and  $K = A_{\pi}$ , where  $\pi = \prod_{1 \leq i \leq r} x_i$ . For  $\underline{n} = (n_1, \ldots, n_r) \in \mathbb{Z}^r$ , and the invertible ideal  $I = (x_1^{n_1} \cdots x_r^{n_r}) \subset K$ , we shall denote  $\operatorname{fil}_I W_m \Omega_K^q$  by  $\operatorname{fil}_{\underline{n}} W_m \Omega_K^q$  and write  $M(x_1^{-n_1} \cdots x_r^{-n_r})$  as  $M(\underline{n})$  for any  $W_m(A)$ -module M (cf. Notation 2.1.4). Also  $\underline{n}/p$  will denote  $(\lfloor n_1/p \rfloor, \ldots, \lfloor n_r/p \rfloor) \in \mathbb{Z}^r$ . We let  $q \geq 0$  and let

$$F_0^{1,q}(S) = \bigoplus_{\substack{0 \le i_0, 1 \le i_1 < \dots < i_s \le r \\ r+1 \le i_{s+1} < \dots < i_{q-i_0} \le d}} \Omega_S^{i_0} \operatorname{dlog}(x_{i_1}) \cdots \operatorname{dlog}(x_{i_s}) d(x_{i_{s+1}}) \cdots d(x_{i_{q-i_0}}).$$
(4.1.1)

Note that  $F_0^{1,q}(S) \subset \Omega_K^q$ . For  $\underline{m} = (m_1, \dots, m_d) \in \mathbb{Z}^d$ , we let  $F_{\underline{m}}^{1,q}(S) = (\prod_i x_i^{m_i}) F_0^{1,q}(S)$ .

**Lemma 4.1.1.** For any  $\underline{n} = (n_1, \ldots, n_r) \in \mathbb{Z}^r$ , there is a canonical isomorphism of A-modules

$$\operatorname{fil}_{\underline{n}}\Omega_K^q \xrightarrow{\cong} \prod_{\underline{m} \in \mathbb{N}_0^d} (x_1^{-n_1} \cdots x_r^{-n_r}) F_{\underline{m}}^{1,q}(S).$$

*Proof.* From the known description of  $\Omega^q_{S[x_1,...,x_d]}$  and [58, Lem. 2.9, 2.11] for q = 0, we get that  $\Omega^q_A(\log \pi) \cong \prod_{\underline{m} \in \mathbb{N}^d_0} F^{1,q}_{\underline{m}}(S)$ . The asserted expression for  $\operatorname{fil}_{\underline{n}}\Omega^q_K$  now follows from its definition and item (2) of Lemma 2.1.7.

For  $\underline{m} = ((m_1, \ldots, m_r), (m_{r+1}, \ldots, m_d)) \in \mathbb{Z}^r \times \mathbb{N}_0^{d-r}$ , we let  $x^{\underline{m}} = \prod_{1 \leq i \leq d} x_i^{m_i}$ . We say  $\underline{m} \leq \underline{m'}$  if  $m_i \leq m'_i$  for each  $1 \leq i \leq d$ . We define  $\underline{m} < \underline{m'}$  similarly. Taking the colimit of  $\operatorname{fil}_{\underline{n}}\Omega_K^q$  as  $\underline{n} \to \infty$ , we get

**Corollary 4.1.2.** Every element  $\omega \in \Omega_K^q$  can be written uniquely as an infinite series  $\omega = \sum_{\underline{m} \in \mathbb{Z}^r \times \mathbb{N}_0^{d^{-r}}} a_{\underline{m}} x^{\underline{m}}$ , where  $a_{\underline{m}} \in F_0^{1,q}(S)$  for all  $\underline{m}$  and  $a_{\underline{m}} = 0$  if  $m_i \ll 0$  for some  $1 \le i \le r$ . Furthermore,  $\omega \in \Omega_A^q(\log \pi)$  if and only if  $a_{\underline{m}} = 0$  for every  $\underline{m}$  such that  $m_i < 0$  for some  $1 \le i \le r$  and  $\omega \in \operatorname{fil}_{\underline{n}} \Omega_K^q$  if and only if  $a_{\underline{m}} = 0$  for every  $\underline{m}$  such that  $m_i < -n_i$  for some  $1 \le i \le r$ .


Notation 4.1.3. For any  $1 \leq i \leq r$  and  $n \in \mathbb{Z}$ , we let  $E_{i,n}^{1,q}(K) = \{\omega = \sum_{\underline{m} \in \mathbb{Z}^r \times \mathbb{N}^{d-r}} a_{\underline{m}} x^{\underline{m}} : a_{\underline{m}} = 0 \text{ if } m_i < -n\}$  and  $\widetilde{E}_{i,n}^{1,q}(K) = \{\omega = \sum_{\underline{m} \in \mathbb{Z}^r \times \mathbb{N}^{d-r}} a_{\underline{m}} x^{\underline{m}} : a_{\underline{m}} = 0 \text{ if } m_i \geq -n\}$ . We let  $\operatorname{fil}_{\underline{n}}^i \Omega_K^q = \bigcap_{1 \leq i \leq i} E_{j,n_j}^{1,q}(K)$ . We then get a filtration

$$\Omega_K^q = \operatorname{fil}_{\underline{n}}^0 \Omega_K^q(K) \supset \operatorname{fil}_{\underline{n}}^1 \Omega_K^q \supset \dots \supset \operatorname{fil}_{\underline{n}}^r \Omega_K^q = \operatorname{fil}_{\underline{n}} \Omega_K^q.$$
(4.1.2)

### 4.2 From complete to the non-complete case

We let A be a regular local F-finite  $\mathbb{F}_p$ -algebra with maximal ideal  $\mathfrak{m} = (x_1, \ldots, x_d)$ . We let  $\pi = x_1 \cdots x_r$  and  $K = A_{\pi}$ , where  $1 \leq r \leq d$  and L = Q(A). We let  $\widehat{A}$  denote the  $\mathfrak{m}$ -adic completion of A with the maximal ideal  $\widehat{\mathfrak{m}}$  and let  $\widehat{K} = \widehat{A}_{\pi}$ . For  $1 \leq i \leq r$ , we let  $A_i = A_{(x_i)}$ ,  $\widehat{A}_i = \widehat{A}_{(x_i)}$  and  $\widehat{K}_i = Q(\widehat{A}_i)$ . For  $\underline{n} = (n_1, \ldots, n_r) \in \mathbb{Z}^r$  and the invertible ideal  $I = (x_1^{n_1} \cdots x_r^{n_r}) \subset K$ , we let  $\operatorname{fil}_{\underline{n}} W_m \Omega_K^q = \operatorname{fil}_I W_m \Omega_K^q$ .  $\underline{n}/p$  will denote the same as before. We also let  $\underline{tn} = (tn_1, \ldots, tn_r)$  for any  $t \in \mathbb{Z}$ . We let  $X = \operatorname{Spec}(A)$  and  $X_f = \operatorname{Spec}(A[f^{-1}])$  for any  $f \in A \setminus \{0\}$ .

**Lemma 4.2.1.** The map  $\psi_i: W_m \Omega_K^q \to W_m \Omega_{\widehat{K}_i}^q$ , induced by the inclusion  $K \hookrightarrow \widehat{K}_i$ , is injective.



*Proof.* We look at the commutative diagram

Since A is a regular local  $\mathbb{F}_p$ -algebra, the composition  $W_m\Omega_A^q \to W_m\Omega_L^q$  of all left vertical arrows with  $\gamma_1$  is injective (cf. [46, Prop. 2.8]). It follows that  $\alpha_1$  is injective. The arrow  $\alpha_2$  is injective by [58, Lem. 2.9, 2.11]. As the vertical arrows on the top floor are the localizations, it follows that  $\beta_1$  and  $\beta_2$  are injective. Since each  $\delta_i$  is an isomorphism by [26, Thm. C], it follows that  $\gamma_1$  and  $\gamma_2$  are injective. This achieves the proof.

Since A is F-finite, one knows that  $W_m(A)$  is a Noetherian local ring whose maximal ideal is  $\mathfrak{M}_m = \{\underline{a} = (a_{m-1}, \ldots, a_0) : a_{m-1} \in \mathfrak{m}\}$ . For any  $W_m(A)$ -module M, we let  $M^{\wedge}$ denote the  $\mathfrak{M}_m$ -adic (equivalently,  $W_m(\mathfrak{m})$ -adic) completion of M. By [58, Lem. 2.9, 2.11], one knows that  $W_m \Omega_A^q$  is a finite type  $W_m(A)$ -module and

$$W_m\Omega_A^q \otimes_{W_m(A)} W_m(\widehat{A}) \xrightarrow{\cong} W_m\Omega_{\widehat{A}}^q \xleftarrow{\cong} (W_m\Omega_A^q)^{\wedge}.$$
(4.2.2)

**Lemma 4.2.2.** There are canonical isomorphisms of  $W_m(\widehat{A})$ -modules

$$W_m\Omega^q_A(\log\pi) \otimes_{W_m(A)} W_m(\widehat{A}) \xrightarrow{\cong} W_m\Omega^q_{\widehat{A}}(\log\pi) \xleftarrow{\cong} W_m\Omega^q_A(\log\pi)^{\wedge}.$$
(4.2.3)

*Proof.* Since  $W_m \Omega_A^q(\log \pi)$  is a finite type  $W_m(A)$ -module, the first and the last terms in (4.2.3) are canonically isomorphic. By the definition of  $W_m \Omega_A^q(\log \pi)$  and (4.2.2), we see that the first arrow in (4.2.3) is surjective. We now look at the commutative diagram

The top horizontal arrow is injective because  $W_m(\widehat{A})$  is a flat  $W_m(A)$ -module. Furthermore,

$$W_{m}\Omega_{K}^{q} \otimes_{W_{m}(A)} W_{m}(\widehat{A}) \cong W_{m}\Omega_{A}^{q} \otimes_{W_{m}(A)} W_{m}(A_{\pi}) \otimes_{W_{m}(A)} W_{m}(\widehat{A})$$
  
$$\cong W_{m}\Omega_{\widehat{A}}^{q} \otimes_{W_{m}(\widehat{A})} (W_{m}(\widehat{A}) \otimes_{W_{m}(A)} W_{m}(K))$$
  
$$\cong W_{m}\Omega_{\widehat{A}}^{q} \otimes_{W_{m}(\widehat{A})} W_{m}(\widehat{A})_{[\pi]_{m}}$$
  
$$\cong W_{m}\Omega_{\widehat{A}}^{q} \otimes_{W_{m}(\widehat{A})} W_{m}(\widehat{K}) \cong W_{m}\Omega_{\widehat{K}}^{q}.$$

$$(4.2.5)$$

This shows that the right vertical arrow in (4.2.4) is bijective (as the composition of the isomorphisms in (4.2.5) is same as the right vertical arrow in (4.2.4)). It follows that the left vertical arrow is injective. This finishes the proof.

We shall need the following well-known commutative algebra fact to prove the next result.

**Lemma 4.2.3.** Let R be a commutative Noetherian local ring with maximal ideal  $\mathfrak{m}$ and let M be a finitely generated R-module. Assume that  $N' \subset M^{\wedge}$  is an  $\widehat{R}$ -submodule. Assume further that  $N \subset M \cap N'$  is an R-submodule which is dense in N' in the  $\mathfrak{m}$ -adic topology. Then  $N' = N^{\wedge}$  and  $N = M \cap N'$ .

*Proof.* First note that  $N' \,\subset M^{\wedge}$  is closed in the m-adic topology (cf. [53, Thms 8.13, 8.10(i)]). We see from [53, Thm. 8.1(i)] that  $N^{\wedge}$  is the closure of N in  $M^{\wedge}$ . This implies  $N^{\wedge} = N'$ . For the second claim, note that  $M/N \hookrightarrow (M/N)^{\wedge} \cong M^{\wedge}/N^{\wedge}$ , where the injectivity of the first arrow follows from the separateness of the m-adic topology (cf. Theorem 8.10(i) of loc. cit.) and the second isomorphism follows from Theorem 8.1(ii) of loc. cit. □

Lemma 4.2.4. The maps

- 1.  $d: \operatorname{fil}_{\underline{n}} W_m \Omega_K^q \to \operatorname{fil}_{\underline{n}} W_m \Omega_K^{q+1};$
- 2.  $V: \operatorname{fil}_{\underline{n}} W_m \Omega_K^q \to \operatorname{fil}_{\underline{n}} W_{m+1} \Omega_K^q;$
- 3.  $F: \operatorname{fil}_{\underline{n}} W_{m+1}\Omega_K^q \to \operatorname{fil}_{\underline{n}} W_m\Omega_K^q;$
- 4.  $R: \operatorname{fil}_{\underline{n}} W_{m+1}\Omega_K^q \to \operatorname{fil}_{\underline{n}/p} W_m\Omega_K^q$

are continuous in the  $W_m(\mathfrak{m})$ -adic topology.

Proof. The claims (3) and (4) are easy to verify. To prove (1) and (2), we note that  $\operatorname{fil}_{\underline{n}} W_m \Omega_K^q$  is a finite type  $W_m(A)$ -module by Lemma 2.1.7. We let  $J_s \subset W_m(A)$  be the ideal  $([x_1]_m^s, \ldots, [x_d]_m^s)$ . Then the  $W_m(\mathfrak{m})$ -adic topology of  $\operatorname{fil}_{\underline{n}} W_m \Omega_K^q$  coincides with the

topology given by the filtration  $\{J_s \operatorname{fil}_{\underline{n}} W_m \Omega_K^q\}$  by [15, Cor. 2.3] (see its proof). Hence, it suffices to note that given  $s \ge 1$ , one has  $d([x_i]_m^{p^m t} \operatorname{fil}_{\underline{n}} W_m \Omega_K^q) \subset J_s \operatorname{fil}_{\underline{n}} W_m \Omega_K^{q+1}$  and  $V([x_i]_m^{pt} \operatorname{fil}_{\underline{n}} W_m \Omega_K^q) \subset J_s \operatorname{fil}_{\underline{n}} W_{m+1} \Omega_K^q$  for all  $t \ge s$  and all i.

**Lemma 4.2.5.** For  $m \ge 1$ , we have the following.

- 1. The map  $F: W_{m+1}(A) \to W_m(A)$  is a finite ring homomorphism.
- 2. The  $W_m(\mathfrak{m})$ -topology of  $W_m(A)$  coincides with its  $W_{m+1}(\mathfrak{m})$ -topology via F.
- 3. For any A-submodule M of  $\Omega_K^{q-1}$  (resp.  $\Omega_K^q$ ),  $dV^{m-1}(M)$  (resp.  $V^{m-1}(M)$ ) is a  $W_{m+1}(A)$ -submodule of  $W_m \Omega_K^q$ , where  $W_{m+1}(A)$  acts on the latter via F.

*Proof.* The item (1) is well-known result of Langer-Zink (cf. [58, Thm. 2.6]). For (2), we can replace (for any m) the  $W_m(\mathfrak{m})$ -adic topology by the one given by the filtration  $\{J_s\}$  defined in the proof of Lemma 4.2.4. Our claim now follows because  $F([x_i]_{m+1}) = [x_i]_m^p$  for every i. To prove (3), we let  $a \in W_{m+1}(A)$  and  $x \in M$ . We get

$$F(a)dV^{m-1}(x) = F(a)FdV^{m}(x) = F(adV^{m}(x)) = FdV^{m}(F^{m}(a)x) - F(daV^{m}(x))$$
$$= FdV^{m}(F^{m}(a)x) - FV^{m}(F^{m}d(a)x) = FdV^{m}(F^{m}(a)x) = dV^{m-1}(F^{m}(a)x),$$

because  $FV^m = 0$ . This finishes the proof as  $F^m(a)x \in M$ . The other case is easier :  $F(a)V^{m-1}(x) = V^{m-1}(F^m(a)x)$ .

**Lemma 4.2.6.** The canonical map  $\operatorname{fil}_{\underline{n}} W_m \Omega_K^q \to \operatorname{fil}_{\underline{n}} W_m \Omega_{\widehat{K}}^q$  induces an isomorphism of  $W_m(\widehat{A})$ -modules

$$\phi: (\operatorname{fil}_{\underline{n}} W_m \Omega_K^q)^{\wedge} \xrightarrow{\cong} \operatorname{fil}_{\underline{n}} W_m \Omega_{\widehat{K}}^q.$$

Proof. Since  $\operatorname{fl}_{\underline{n}} W_m \Omega_K^q$  is a finite type  $W_m(A)$ -module, we know that the canonical map  $\operatorname{fl}_{\underline{n}} W_m \Omega_K^q \otimes_{W_m(A)} W_m(\widehat{A}) \to (\operatorname{fl}_{\underline{n}} W_m \Omega_K^q)^{\wedge}$  is an isomorphism. On the other hand, we have an injection  $\operatorname{fl}_{\underline{n}} W_m \Omega_K^q \otimes_{W_m(A)} W_m(\widehat{A}) \hookrightarrow W_m \Omega_K^q \otimes_{W_m(A)} W_m(\widehat{A}) \cong W_m \Omega_{\widehat{K}}^q$ , where the last isomorphism is by (4.2.5). It follows that  $\phi$  is injective.

Now, first consider the case q = 0. If m = 1 then the isomorphism is clear by Lemma 2.1.7(2). For m > 1, consider the commutative diagram  $W_m(\widehat{A})$  modules.

where  $W = W_m(A)$  and  $\widehat{W} = W_m(\widehat{A})$ . The top row is obtained by tensoring (2.1.2) (taking  $U = \operatorname{Spec} K, D = (\prod_{1 \le i \le r} x_i^{n_i})$ ) with  $W_m(\widehat{A})$  (over  $W_m(A)$ ) and noting that  $W_m(\widehat{A})$ is the  $W_m(\mathfrak{m})$ -adic completion of  $W_m(A)$  (cf. (4.2.2) for q = 0) and hence flat over  $W_m(A)$ . Here the module structure on fil<sub>n</sub>K is given via  $F^{m-1}$ . By Lemma 4.2.5(2), we see that the  $W_m(\mathfrak{m})$ -adic topology on fil<sub>n</sub>K (via  $F^{m-1}$ ) is same as the  $\mathfrak{m}$ -adic topology. Hence fil<sub>n</sub>K  $\otimes_{W_m(A)} W_m(\widehat{A}) = (\operatorname{fil}_n K)^{\wedge}$ , the  $\mathfrak{m}$ -adic completion of fil<sub>n</sub>K as A module. We also note that the  $W_{m-1}(\mathfrak{m})$ -topology of  $W_{m-1}(A)$  coincides with its  $W_m(\mathfrak{m})$ -topology via the surjection  $R: W_m(A) \twoheadrightarrow W_{m-1}(A)$ . It follows that for any finite type  $W_{m-1}(A)$ module M, one has isomorphisms

$$M \otimes_{W_m(A)} W_m(\widehat{A}) \cong M_m^{\wedge} \cong M_{m-1}^{\wedge} \cong M \otimes_{W_{m-1}(A)} W_{m-1}(\widehat{A}), \tag{4.2.7}$$

where  $M_{m}^{\wedge}$  (resp.  $M_{m-1}^{\wedge}$ ) is the  $W_{m}(\mathfrak{m})$  (resp.  $W_{m-1}(\mathfrak{m})$ )-adic completion of M. Thus, the left and right vertical arrows are isomorphism by m = 1 case and induction, respectively. Hence, the middle arrow is also an isomorphism. In particular, we get that  $\operatorname{fil}_{\underline{n}}W_{m}(K)$  is dense in  $\operatorname{fil}_{\underline{n}}W_{m}(\widehat{K})$ .

For q > 0 case, recall that  $\operatorname{fil}_{\underline{n}} W_m \Omega_{\widehat{K}}^q$  is the sum of  $\operatorname{fil}_{\underline{n}} W_m(\widehat{K}) W_m \Omega_{\widehat{A}}^q(\log \pi)$  and  $d(\operatorname{fil}_{\underline{n}} W_m(\widehat{K})) W_m \Omega_{\widehat{A}}^{q-1}(\log \pi)$ . Hence by Lemmas 4.2.2 and 4.2.4 and the q = 0 case we get that  $\operatorname{fil}_{\underline{n}} W_m \Omega_K^q$  is dense in  $\operatorname{fil}_{\underline{n}} W_m \Omega_{\widehat{K}}^q$ . On the other hand, the lemma is clear if  $\underline{n} = p^{m-1} \underline{n}''$  for some  $\underline{n}''$ ; by Lemma 2.1.7 (9) and Lemma 4.2.2. So we can choose some  $\underline{n}'$  such that  $p^{m-1} \underline{n}' > \underline{n}$  and so  $\operatorname{fil}_{\underline{n}} W_m \Omega_R^q \subset \operatorname{fil}_{p^{m-1} \underline{n}'} W_m \Omega_R^q$  for  $R \in \{K, \widehat{K}\}$ . Now, we apply Lemma 4.2.3 with  $M = \operatorname{fil}_{p^{m-1} \underline{n}'} W_m \Omega_K^q$  and  $N = \operatorname{fil}_{\underline{n}} W_m \Omega_K^q$  to conclude the proof.  $\Box$ 

**Corollary 4.2.7.** Consider  $F(\operatorname{fil}_{\underline{n}}W_{m+1}\Omega_{K}^{q})$  as a  $W_{m+1}(A)$ -module (via F). Then the canonical map  $F(\operatorname{fil}_{\underline{n}}W_{m+1}\Omega_{K}^{q}) \to F(\operatorname{fil}_{\underline{n}}W_{m+1}\Omega_{\widehat{K}}^{q})$  induces an isomorphism of  $W_{m+1}(\widehat{A})$ -modules (via F) :  $\phi : F(\operatorname{fil}_{\underline{n}}W_{m+1}\Omega_{K}^{q})^{\wedge} \xrightarrow{\cong} F(\operatorname{fil}_{\underline{n}}W_{m+1}\Omega_{\widehat{K}}^{q})$ .

Proof. Consider  $\operatorname{fil}_{\underline{n}}W_m\Omega_K^q$  as a finite  $W_{m+1}(A)$ -module via F (see Lemma 4.2.5(1)), so that  $F(\operatorname{fil}_{\underline{n}}W_{m+1}\Omega_K^q) \subset \operatorname{fil}_{\underline{n}}W_m\Omega_K^q$  is a  $W_{m+1}(A)$ -submodule. By Lemmas 4.2.4(3) and 4.2.6 we know that  $F(\operatorname{fil}_{\underline{n}}W_{m+1}\Omega_K^q)$  is dense in  $F(\operatorname{fil}_{\underline{n}}W_{m+1}\Omega_{\widehat{K}}^q)$  with respect to  $W_m(\widehat{\mathfrak{m}})$ -adic topology. Also by Lemmas 4.2.5(2) and 4.2.6, we have

$$\operatorname{fil}_{\underline{n}} W_m \Omega_{\widehat{K}}^q \cong \operatorname{fil}_{\underline{n}} W_m \Omega_K^q \otimes_{W_{m+1}(A)} W_{m+1}(\widehat{A}),$$

the  $W_{m+1}(\mathfrak{m})$ -adic completion (via F) of  $\operatorname{fil}_{\underline{n}} W_m \Omega_K^q$ . Now, we can apply Lemma 4.2.3 with  $M = \operatorname{fil}_{\underline{n}} W_m \Omega_K^q$  and  $N = F(\operatorname{fil}_{\underline{n}} W_{m+1} \Omega_K^q)$  (considering them as modules over  $W_{m+1}(A)$ , via F) to conclude the proof.

**Definition 4.2.8.** We let  $\psi_i: W_m \Omega_K^q \to W_m \Omega_{\widehat{K}_i}^q$  be the map induced by the inclusion  $K \hookrightarrow \widehat{K}_i$ . We let  $\operatorname{fil}_n^{"} W_m \Omega_K^q$  denote the kernel of the canonical map

$$\psi = (\psi_i) : W_m \Omega_K^q \to \bigoplus_{1 \le i \le r} \frac{W_m \Omega_{\widehat{K}_i}^q}{\operatorname{fil}_{n_i} W_m \Omega_{\widehat{K}_i}^q}.$$

It is clear that  $\operatorname{fil}_{\underline{n}} W_m \Omega_K^q \subset \operatorname{fil}_{\underline{n}}^{\prime\prime} W_m \Omega_K^q$ .

**Lemma 4.2.9.** If A is complete, then one has an exact sequence of  $W_m(A)$ -modules

$$0 \to V^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_K^q) + dV^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_K^{q-1}) \to \operatorname{fil}_{\underline{n}}''W_m\Omega_K^q \xrightarrow{R} \operatorname{fil}_{\underline{n}/p}''W_{m-1}\Omega_K^q.$$

*Proof.* We let  $T = V^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_K^q) + dV^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_K^{q-1})$ . Using the exact sequence (3.1.7), we only need to show that every element

$$\omega \in \left( V^{m-1}(\Omega_K^q) + dV^{m-1}(\Omega_K^{q-1}) \right) \bigcap \operatorname{fil}_{\underline{n}}^{\prime\prime} W_m \Omega_K^q$$

lies in T. We shall show more generally that for every  $0 \le i \le r$ , we can write  $\omega = V^{m-1}(x) + dV^{m-1}(y)$  with  $x \in \operatorname{fil}_{\underline{n}}^{i}\Omega_{K}^{q}$  and  $y \in \operatorname{fil}_{\underline{n}}^{i}\Omega_{K}^{q-1}$  (cf. Notation 4.1.3). This will finish the proof of the lemma.

Now, we can write  $\omega = V^{m-1}(x_0) + dV^{m-1}(y_0)$  for some  $x_0 \in \Omega_K^q = \operatorname{fil}_{\underline{n}}^0 \Omega_K^q$  and  $y_0 \in \Omega_K^{q-1} = \operatorname{fil}_{\underline{n}}^0 \Omega_K^{q-1}$ . Suppose next that  $i \geq 1$  and that we have found  $x \in \operatorname{fil}_{\underline{n}}^{i-1} \Omega_K^q$  and  $y \in \operatorname{fil}_{\underline{n}}^{i-1} \Omega_K^{q-1}$  such that  $\omega = V^{m-1}(x) + dV^{m-1}(y)$ . In the notations of Corollary 4.1.2, we can uniquely write  $x = \sum_{\underline{m}} a_{\underline{m}} x^{\underline{m}}$  and  $y = \sum_{\underline{m}} b_{\underline{m}} x^{\underline{m}}$  such that  $a_{\underline{m}} = 0 = b_{\underline{m}}$  for any  $\underline{m}$  such that  $m_j < -n_j$  for some j < i. We can uniquely write  $x = x_1 + x_2$  and  $y = y_1 + y_2$  such that

1.  $x_1, x_2 \in \operatorname{fil}_{\underline{n}}^{i-1}\Omega_K^q$  and  $y_1, y_2 \in \operatorname{fil}_{\underline{n}}^{i-1}\Omega_K^{q-1}$ .

2. 
$$x_1 \in \operatorname{fil}_{\underline{n}}^i \Omega_K^q$$
,  $y_1 \in \operatorname{fil}_{\underline{n}}^i \Omega_K^{q-1}$ . In particular,  $\psi_i(V^{m-1}(x_1) + dV^{m-1}(y_1)) \in \operatorname{fil}_{n_i} W_m \Omega_{\widehat{K}_i}^q$ 

3. No term of the infinite series  $x_2$  (resp.  $y_2$ ) lies in  $\operatorname{fil}_n^i \Omega_K^q$  (resp.  $\operatorname{fil}_n^i \Omega_K^{q-1}$ ).

The above conditions imply that

$$\psi_i(V^{m-1}(x_2) + dV^{m-1}(y_2)) = \psi_i(\omega - (V^{m-1}(x_1) + dV^{m-1}(y_1))) \in \operatorname{fil}_{n_i} W_m \Omega^q_{\widehat{K}_i}.$$

It follows from Lemma 3.1.6 that

$$V^{m-1}(\psi_i(x_2)) + dV^{m-1}(\psi_i(y_2)) \in \left[\prod_{j < -n_i} F_j^{m,q}(\kappa(\widehat{A}_i))\right] \cap \operatorname{fil}_{n_i} W_m \Omega_{\widehat{K}_i}^q,$$

where  $\kappa(\widehat{A}_i)$  is the residue field of  $\widehat{A}_i$ .

But the above intersection is trivial by Corollary 3.1.5 and Proposition 3.1.19. It follows that  $\psi_i(V^{m-1}(x_2) + dV^{m-1}(y_2)) = V^{m-1}(\psi_i(x_2)) + dV^{m-1}(\psi_i(y_2)) = 0$ . In particular,  $V^{m-1}(x_2) + dV^{m-1}(y_2) = 0$  by Lemma 4.2.1. We conclude that  $\omega = V^{m-1}(x_1) + dV^{m-1}(y_1)$ . We are now done by the condition (2) above and proceeding inductively.

**Proposition 4.2.10.** There is a short exact sequence of  $W_m(A)$ -modules

$$0 \to V^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_K^q) + dV^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_K^{q-1}) \to \operatorname{fil}_{\underline{n}}W_m\Omega_K^q \xrightarrow{R} \operatorname{fil}_{\underline{n}/p}W_{m-1}\Omega_K^q \to 0.$$

*Proof.* Surjectivity of R follows from Lemma 2.1.7(7). It remains therefore to show that  $\operatorname{Ker}(R) \subset V^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_K^q) + dV^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_K^{q-1})$ . First we assume that A is complete. In this case, it follows from Lemma 4.2.9 because  $\operatorname{fil}_{\underline{n}}W_m\Omega_K^q \subset \operatorname{fil}_n''W_m\Omega_K^q$ .

Suppose now that A is not complete.

We now let  $E = V^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_{K}^{q}) + dV^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_{K}^{q-1})$  and  $E' = \operatorname{Ker}(R)$ . One easily verifies that E is a  $W_{m}(A)$ -submodule of  $\operatorname{fil}_{\underline{n}}W_{m}\Omega_{K}^{q}$ . Indeed, Lemma 2.1.7 implies that  $E \subset \operatorname{fil}_{\underline{n}}W_{m}\Omega_{K}^{q}$ . Moreover, one has  $a(V^{m-1}(b) + dV^{m-1}(c)) = V^{m-1}(F(a)b - Fd(a)c) + dV^{m-1}(F(a)c)$  for any  $a \in W_{m}(A)$ ,  $b \in \operatorname{fil}_{\underline{n}}\Omega_{K}^{q}$  and  $c \in \operatorname{fil}_{\underline{n}}\Omega_{K}^{q-1}$ . Lemma 2.1.7 implies once again that  $a(V^{m-1}(b) + dV^{m-1}(c)) \in E$ . Using the complete case, it remains therefore to show that  $E = \left[V^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_{\widehat{K}}^{q}) + dV^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_{\widehat{K}}^{q-1})\right] \cap \operatorname{fil}_{\underline{n}}W_{m}\Omega_{K}^{q}$ . But this follows from Lemma 4.2.3 because E is dense in  $V^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_{\widehat{K}}^{q}) + dV^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_{\widehat{K}}^{q}) + dV^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_{\widehat{K}}^{q}) = dV^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_{\widehat{K}}^{q-1})$  by Lemmas 4.2.4 and 4.2.6.

**Corollary 4.2.11.** The inclusions  $K \hookrightarrow \widehat{K}_i$  induce an exact sequence

$$0 \to \operatorname{fil}_{\underline{n}} W_m \Omega_K^q \to W_m \Omega_K^q \xrightarrow{\psi} \bigoplus_{1 \le i \le r} \frac{W_m \Omega_{\widehat{K}_i}^q}{\operatorname{fil}_{n_i} W_m \Omega_{\widehat{K}_i}^q}$$

*Proof.* The corollary is equivalent to the claim that the inclusion  $\operatorname{fil}_{\underline{n}} W_m \Omega_K^q \subset \operatorname{fil}_{\underline{n}}^{"} W_m \Omega_K^q$  is an equality. To prove this claim, we first assume that A is complete and prove by induction on m. The case m = 1 follows directly from Corollary 4.1.2. For  $m \ge 2$ , we use the commutative diagram

Proposition 4.2.10 says that the top row in (4.2.8) is exact and Lemma 4.2.9 implies that the bottom row is exact at  $\operatorname{fil}_n^{\prime\prime} W_m \Omega_K^q$ . The left and the right vertical arrows are

#### 4.3. CARTIER HOMOMORPHISM II

bijective by induction. A diagram chase shows that the middle vertical arrow is also bijective.

For A not necessarily complete, it suffices to show that  $\operatorname{fil}_{\underline{n}} W_m \Omega_K^q = \operatorname{fil}_{\underline{n}} W_m \Omega_{\widehat{K}}^q \cap W_m \Omega_K^q$ . But this follows because  $W_m(\widehat{A})$  is a faithfully flat  $W_m(A)$ -algebra and there are isomorphisms

$$\frac{W_m\Omega_K^q}{\operatorname{fil}_{\underline{n}}W_m\Omega_K^q} \otimes_{W_m(A)} W_m(\widehat{A}) \cong \frac{W_m\Omega_K^q \otimes_{W_m(A)} W_m(\widehat{A})}{\operatorname{fil}_{\underline{n}}W_m\Omega_K^q \otimes_{W_m(A)} W_m(\widehat{A})} \cong \frac{W_m\Omega_{\widehat{K}}^q}{\operatorname{fil}_{\underline{n}}W_m\Omega_{\widehat{K}}^q}, \tag{4.2.9}$$

where the second isomorphism follows from (4.2.5) and Lemma 4.2.6.

Lemma 4.2.12. The canonical maps

$$\frac{\operatorname{fil}_{\underline{n}} W_m \Omega_K^q}{dV^{m-1}(\operatorname{fil}_{\underline{n}} \Omega_K^{q-1})} \to \frac{W_m \Omega_K^q}{dV^{m-1}(\Omega_K^{q-1})}$$
$$\frac{\operatorname{fil}_{\underline{n}} W_m \Omega_K^q}{V^{m-1}(\operatorname{fil}_{\underline{n}} \Omega_K^q)} \to \frac{W_m \Omega_K^q}{V^{m-1}(\Omega_K^q)}$$

are injective.

*Proof.* We shall prove the injectivity of the first map as the other one is completely analogous. We only need to show that every element  $\omega \in \operatorname{fl}_{\underline{n}} W_m \Omega_K^q \cap dV^{m-1}(\Omega_K^{q-1})$  lies in  $dV^{m-1}(\operatorname{fl}_{\underline{n}} \Omega_K^{q-1})$ . If A is complete, this is proven exactly as we proved Lemma 4.2.9 (with x = 0). If A is not complete, it suffices to show that

$$\operatorname{fil}_{\underline{n}} W_m \Omega_K^q \bigcap dV^{m-1}(\operatorname{fil}_{\underline{n}} \Omega_{\widehat{K}}^{q-1}) \subseteq dV^{m-1}(\operatorname{fil}_{\underline{n}} \Omega_K^{q-1}).$$

To that end, we note that  $\operatorname{fl}_{\underline{n}}W_m\Omega_{\widehat{K}}^q = (\operatorname{fl}_{\underline{n}}W_m\Omega_K^q)^{\wedge}$  by Lemma 4.2.6. In particular,  $\operatorname{fl}_{\underline{n}}\Omega_K^{q^{-1}}$  is dense in  $\operatorname{fl}_{\underline{n}}\Omega_{\widehat{K}}^{q^{-1}}$  with respect to the  $\widehat{\mathfrak{m}}$ -adic topology. It follows from Lemma 4.2.4 that  $dV^{m-1}(\operatorname{fl}_{\underline{n}}\Omega_K^{q^{-1}})$  is a dense subgroup of  $dV^{m-1}(\operatorname{fl}_{\underline{n}}\Omega_{\widehat{K}}^{q^{-1}})$  with respect to the  $W_m(\mathfrak{m})$ -adic topology. We conclude from Lemma 4.2.5 that  $\operatorname{fl}_{\underline{n}}W_m\Omega_{\widehat{K}}^q$ is the  $W_{m+1}(\mathfrak{m})$ -adic completion of the  $W_{m+1}(A)$ -module  $\operatorname{fl}_{\underline{n}}W_m\Omega_K^q$  via  $F:W_{m+1}(A) \to$  $W_m(A)$  and  $dV^{m-1}(\operatorname{fl}_{\underline{n}}\Omega_K^{q^{-1}})$  is a dense subgroup of  $dV^{m-1}(\operatorname{fl}_{\underline{n}}\Omega_{\widehat{K}}^{q^{-1}})$  with respect to the  $W_{m+1}(\mathfrak{m})$ -adic topology. By Lemma 4.2.3, it remains to show that  $dV^{m-1}(\operatorname{fl}_{\underline{n}}\Omega_K^{q^{-1}})$  is a  $W_{m+1}(A)$ -submodule of  $\operatorname{fl}_{\underline{n}}W_m\Omega_K^q$ . But this follows again from Lemma 4.2.5.

### 4.3 Cartier homomorphism II

**Definition 4.3.1.** We define  $Z_1 \operatorname{fil}_{\underline{n}} W_m \Omega_K^q \coloneqq Z_1 W_m \Omega_K^q \cap \operatorname{fil}_{\underline{n}} W_m \Omega_K^q$ , for all  $\underline{n} \in \mathbb{Z}^r$  and  $m \ge 1$ . Equivalently, we have  $Z_1 \operatorname{fil}_{\underline{n}} W_m \Omega_K^q = \operatorname{Ker}(F^{m-1}d : \operatorname{fil}_{\underline{n}} W_m \Omega_K^q \to \operatorname{fil}_{\underline{n}} \Omega_K^{q+1})$ .

**Lemma 4.3.2.** The Cartier map  $C: Z_1 W_m \Omega_K^q \to W_m \Omega_K^q$  restricts to a  $W_{m+1}(A)$ -linear map

$$C: Z_1 \operatorname{fil}_{\underline{n}} W_m \Omega_K^q \to \operatorname{fil}_{\underline{n}/p} W_m \Omega_K^q$$

such that  $C(dV^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_K^{q-1})) = 0.$ 

Proof. By Corollary 4.2.11, we only need to show that  $C \circ \psi_i(\omega) = \psi_i \circ C(\omega)$  lies in  $\operatorname{fil}_{[n_i/p]} W_m \Omega_{\widehat{K}_i}^q$  for every  $\omega \in Z_1 W_m \Omega_K^q \cap \operatorname{fil}_{\underline{n}} W_m \Omega_K^q$  and  $1 \leq i \leq r$ . But this is shown in the proof of Lemma 3.2.1.

Lemma 4.3.3. We have the following.

1. 
$$Z_1 \operatorname{fil}_{\underline{n}} W_m \Omega_K^q = F(\operatorname{fil}_{\underline{n}} W_{m+1} \Omega_K^q)$$

2.  $F: \operatorname{fil}_{\underline{n}} W_{m+1}\Omega_K^q \to \operatorname{fil}_{\underline{n}} W_m \Omega_K^q$  induces isomorphisms

$$\overline{F}: \operatorname{fil}_{\underline{n}/p} W_m \Omega_K^q \xrightarrow{\cong} \frac{Z_1 \operatorname{fil}_{\underline{n}} W_m \Omega_K^q}{dV^{m-1} (\operatorname{fil}_{\underline{n}} \Omega_K^{q-1})}; \quad \overline{C}: \frac{Z_1 \operatorname{fil}_{\underline{n}} W_m \Omega_K^q}{dV^{m-1} (\operatorname{fil}_{\underline{n}} \Omega_K^{q-1})} \xrightarrow{\cong} \operatorname{fil}_{\underline{n}/p} W_m \Omega_K^q.$$

*Proof.* We first note that the lemma is classical for  $W_{\cdot}\Omega_{K}^{\bullet}$  (cf. [30] and [32]). To prove (1) and (2), we use Proposition 4.2.10 and the identity  $FV^{m} = 0$  to get a commutative diagram

$$\begin{aligned} \operatorname{fil}_{\underline{n}} W_{m+1} \Omega_{K}^{q} & \xrightarrow{F} \operatorname{fil}_{\underline{n}} W_{m} \Omega_{K}^{q} \\ R & \downarrow \\ R & \downarrow \\ \operatorname{fil}_{\underline{n}/p} W_{m} \Omega_{K}^{q} & \xrightarrow{\overline{F}} \frac{\operatorname{fil}_{\underline{n}} W_{m} \Omega_{K}^{q}}{dV^{m-1} (\operatorname{fil}_{\underline{n}} \Omega_{K}^{q-1})}. \end{aligned}$$

$$(4.3.1)$$

Since  $dV^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_{K}^{q-1}) = FdV^{m}(\operatorname{fil}_{\underline{n}}\Omega_{K}^{q-1}) \subset F(\operatorname{fil}_{\underline{n}}W_{m+1}\Omega_{K}^{q})$ , we thus get maps

$$\operatorname{fil}_{\underline{n}/p} W_m \Omega_K^q \xrightarrow{\overline{F}} \frac{F(\operatorname{fil}_{\underline{n}} W_{m+1} \Omega_K^q)}{dV^{m-1}(\operatorname{fil}_{\underline{n}} \Omega_K^{q-1})} \hookrightarrow \frac{Z_1 \operatorname{fil}_{\underline{n}} W_m \Omega_K^q}{dV^{m-1}(\operatorname{fil}_{\underline{n}} \Omega_K^{q-1})} \xrightarrow{\overline{C}} \operatorname{fil}_{\underline{n}/p} W_m \Omega_K^q.$$
(4.3.2)

We now look at the commutative diagram

One knows classically that the composition of any two adjacent maps in the bottom row is identity. Since the vertical arrows are injective by Lemma 4.2.12, we conclude that the same holds in the top row too. This proves the lemma.  $\Box$ 

#### 4.3. CARTIER HOMOMORPHISM II

We let  $B_0 \operatorname{fll}_{\underline{n}} \Omega_K^q = 0$  and  $B_1 \operatorname{fll}_{\underline{n}} \Omega_K^q = d\operatorname{fll}_{\underline{n}} \Omega_K^{q-1}$ . We let  $Z_0 \operatorname{fll}_{\underline{n}} \Omega_K^q = \operatorname{fll}_{\underline{n}} \Omega_K^q$ . For  $i \ge 2$ , we let  $Z_i \operatorname{fll}_{\underline{n}} \Omega_K^q$  (resp.  $B_i \operatorname{fll}_{\underline{n}} \Omega_K^q$ ) be the inverse image of  $Z_{i-1} \operatorname{fll}_{\underline{n}} \Omega_K^q$  (resp.  $B_{i-1} \operatorname{fll}_{\underline{n}} \Omega_K^q$ ) under the composite map  $Z_{i-1} \operatorname{fll}_{\underline{n}} \Omega_K^q \twoheadrightarrow \frac{Z_{i-1} \operatorname{fll}_{\underline{n}} \Omega_K^q}{B_1 \operatorname{fll}_{\underline{n}} \Omega_K^q} \xrightarrow{\overline{C}} \operatorname{fll}_{\underline{n}/p} \Omega_K^q$ . The following is easily deduced from (4.3.1) and (4.3.3).

Lemma 4.3.4. 1.  $Z_i \operatorname{fil}_{\underline{n}} \Omega_K^q = F^i (\operatorname{fil}_{\underline{n}} W_{i+1} \Omega_K^q), \quad B_i \operatorname{fil}_{\underline{n}} \Omega_K^q = F^{i-1} d(\operatorname{fil}_{\underline{n}} W_i \Omega_K^{q-1}).$ 

- $\mathcal{2}. \ \ Z_i \mathrm{fil}_{\underline{n}} \Omega^q_K \supset Z_{i+1} \mathrm{fil}_{\underline{n}} \Omega^q_K \supset B_j \mathrm{fil}_{\underline{n}} \Omega^q_K \supset B_{j-1} \mathrm{fil}_{\underline{n}} \Omega^q_K \quad \forall \quad i \geq 0, \quad j \geq 1.$
- 3. The maps  $\overline{C}: \frac{Z_i \operatorname{fil}_{\underline{n}} \Omega_K^q}{B_i \operatorname{fil}_{\underline{n}} \Omega_K^q} \xrightarrow{\cong} \frac{Z_{i-1} \operatorname{fil}_{\underline{n}/p} \Omega_K^q}{B_{i-1} \operatorname{fil}_{\underline{n}/p} \Omega_K^q} \text{ and } \overline{F}: \frac{Z_{i-1} \operatorname{fil}_{\underline{n}/p} \Omega_K^q}{B_{i-1} \operatorname{fil}_{\underline{n}/p} \Omega_K^q} \xrightarrow{\cong} \frac{Z_i \operatorname{fil}_{\underline{n}} \Omega_K^q}{B_i \operatorname{fil}_{\underline{n}} \Omega_K^q} \text{ are inverses of each other.}$

**Lemma 4.3.5.** We have  $Z_i \operatorname{fil}_{\underline{n}} \Omega_K^q = Z_i \Omega_K^q \cap \operatorname{fil}_{\underline{n}} \Omega_K^q$  and  $B_i \operatorname{fil}_{\underline{n}} \Omega_K^q = B_i \Omega_K^q \cap \operatorname{fil}_{\underline{n}} \Omega_K^q$ .

*Proof.* If i = 1, the first claim follows from Definition 4.3.1 while the second claim follows from Lemma 4.2.12. To prove the first claim for  $i \ge 2$ , we look at the commutative diagram of exact sequences

The right vertical arrows is bijective by induction and we showed above that the left vertical arrow is bijective. This implies the same for the middle vertical arrow. The second claim for  $i \ge 2$  is proven similarly by replacing the  $Z_i$ -groups (resp.  $Z_{i-1}$ -groups) by the  $B_i$ -groups (resp.  $B_{i-1}$ -groups) in the above diagram.

Lemma 4.3.6. There exists a short exact sequence

$$0 \to V^{m-1}(Z_1 \operatorname{fil}_{\underline{n}}\Omega_K^q) + dV^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_K^{q-1}) \to Z_1 \operatorname{fil}_{\underline{n}}W_m \Omega_K^q \xrightarrow{R} Z_1 \operatorname{fil}_{\underline{n}/p} W_{m-1}\Omega_K^q \to 0.$$
(4.3.4)

Proof. By comparing with the classical case and using Lemma 2.1.7, one checks that (4.3.4) is a well defined complex. Moreover, Proposition 4.2.10 and Lemma 4.3.3(1) together imply that the map R is surjective. We now let  $\omega \in \operatorname{Ker}(R)$ . We then know from Proposition 4.2.10 that  $\omega = V^{m-1}(a) + dV^{m-1}(b)$  for some  $a \in \operatorname{fil}_{\underline{n}}\Omega_K^q$  and  $b \in \operatorname{fil}_{\underline{n}}\Omega_K^{q-1}$ . By Definition 4.3.1, we then get  $da = F^{m-1}d(V^{m-1}(a) + dV^{m-1}(b)) = 0$ . In other words,  $\omega \in V^{m-1}(Z_1\operatorname{fil}_{\underline{n}}\Omega_K^q) + dV^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_K^{q-1})$ .

**Definition 4.3.7.** For  $m \ge 1, n \in \mathbb{Z}^r$ , we let

$$Z_1 R_{m,\underline{n}}^q(K) = \operatorname{Ker}(Z_1 \operatorname{fil}_{\underline{n}} \Omega_K^q \oplus \operatorname{fil}_{\underline{n}} \Omega_K^{q-1} \xrightarrow{V^m, \ dV^m} Z_1 \operatorname{fil}_{\underline{n}} W_{m+1} \Omega_K^q) \text{ and} R_{m,\underline{n}}^q(K) = \operatorname{Ker}(\operatorname{fil}_{\underline{n}} \Omega_K^q \oplus \operatorname{fil}_{\underline{n}} \Omega_K^{q-1} \xrightarrow{V^m, \ dV^m} \operatorname{fil}_{\underline{n}} W_{m+1} \Omega_K^q).$$

We observe that  $Z_1 R_{m,\underline{n}}^q(K) = R_{m,\underline{n}}^q(K)$ . Indeed, any element  $(a,b) \in R_{m,\underline{n}}^q(K)$ has the property that  $a \in Z_1 \Omega_K^q \cap \operatorname{fil}_{\underline{n}} \Omega_K^q = Z_1 \operatorname{fil}_{\underline{n}} \Omega_K^q$  by Corollary 1.1.8. In particular,  $(a,b) \in Z_1 R_{m-1,\underline{n}}^q(K)$ .

**Definition 4.3.8.** For  $\underline{n} \ge (-1, \ldots, -1)$ , we let  $C_{\underline{n}, \bullet}^{m, q}(A)$  denote the 2-term complex

$$\left(Z_1 \operatorname{fil}_{\underline{n}} W_m \Omega_K^q \xrightarrow{1-C} \operatorname{fil}_{\underline{n}} W_m \Omega_K^q\right).$$

$$(4.3.5)$$

We look at the diagram (for  $\underline{n} \ge (-1, \ldots, -1)$ )

In this diagram, we let  $\phi = 1 - \widetilde{C}$ , where  $\widetilde{C}(a, b) = C(a)$ . Since  $CdV^{m-1} = 0$  by Lemma 4.3.2, this diagram is commutative. We have observed that  $Z_1R_{m-1,\underline{n}}^q(K) = R_{m-1,\underline{n}}^q(K)$ . Finally, we observe that all vertical arrows are defined because  $\operatorname{fil}_{\underline{n}/p}W_m\Omega_K^q \subset \operatorname{fil}_{\underline{n}}W_m\Omega_K^q$  under our assumption on  $\underline{n}$ . We now claim that  $\phi: R_{m-1,\underline{n}}^q(K) \to R_{m-1,\underline{n}}^q(K)$  is bijective. To show this, it is enough to check that  $\widetilde{C} = (C,0)$  is nilpotent on  $R_{m-1,\underline{n}}^q(K)$ . But this follows from the fact that  $R_{m-1,\underline{n}}^q(K) \subset B_m\Omega_K^q \oplus \Omega_K^{q-1}$  (cf. Corollary 1.1.8) and  $C^{m+1} = 0$  on  $B_m\Omega_K^q$ . Combining these claims with Lemma 4.3.6, we deduce the following.

**Corollary 4.3.9.** In the derived category of abelian groups, there exists a distinguished triangle (cf. Definition 4.3.8)

$$C^{1,q}_{\underline{n},\bullet}(A) \xrightarrow{V^{m-1}} C^{m,q}_{\underline{n},\bullet}(A) \xrightarrow{R} C^{m-1,q}_{\underline{n}/p,\bullet}(A) \xrightarrow{+} C^{1,q}_{\underline{n},\bullet}(A)[1].$$

*Proof.* We let  $F_{\underline{n},\bullet}^{1,q}(A)$  (resp.  $\widetilde{F}_{\underline{n},\bullet}^{1,q}(A)$ ) denote the 2-term complex consisting of the middle (resp. left) column of (4.3.6). Using Lemma 4.3.6, we only need to show that the composite map

$$C^{1,q}_{\underline{n},\bullet}(A) \hookrightarrow F^{1,q}_{\underline{n},\bullet}(A) \twoheadrightarrow \frac{F^{1,q}_{\underline{n},\bullet}(A)}{\widetilde{F}^{1,q}_{\underline{n},\bullet}(A)}$$

is a quasi-isomorphism. But this follows from (4.3.6) and the claims following it.  $\Box$ 

Lemma 4.3.10. We have the following.

1. Ker $(F^{m-1}d: \operatorname{fil}_{\underline{n}}W_m\Omega_K^q \to \operatorname{fil}_{\underline{n}}\Omega_K^{q+1}) = F(\operatorname{fil}_{\underline{n}}W_{m+1}\Omega_K^q).$ 

#### 4.3. CARTIER HOMOMORPHISM II

 $\begin{aligned} & \mathcal{2}. \ \operatorname{Ker}(F^{m-1}:\operatorname{fil}_{\underline{n}}W_{m}\Omega_{K}^{q} \to \operatorname{fil}_{\underline{n}}\Omega_{K}^{q}) = V(\operatorname{fil}_{\underline{n}}W_{m-1}\Omega_{K}^{q}). \\ & \mathcal{3}. \ \operatorname{Ker}(dV^{m-1}:\operatorname{fil}_{\underline{n}}\Omega_{K}^{q} \to \operatorname{fil}_{\underline{n}}W_{m}\Omega_{K}^{q+1}) = F^{m}(\operatorname{fil}_{\underline{n}}W_{m+1}\Omega_{K}^{q}). \\ & \mathcal{4}. \ \operatorname{Ker}(V:\operatorname{fil}_{\underline{n}}W_{m}\Omega_{K}^{q} \to \operatorname{fil}_{\underline{n}}W_{m+1}\Omega_{K}^{q}) = dV^{m-1}(\operatorname{fil}_{\underline{n}}\Omega_{K}^{q-1}). \\ & \mathcal{5}. \ \operatorname{Ker}(V^{m-1}:\operatorname{fil}_{\underline{n}}\Omega_{K}^{q} \to \operatorname{fil}_{\underline{n}}W_{m}\Omega_{K}^{q}) = F^{m-1}dV(\operatorname{fil}_{\underline{n}}W_{m-1}\Omega_{K}^{q-1}). \\ & \mathcal{6}. \ \operatorname{Ker}(dV^{m-1}:\operatorname{fil}_{\underline{n}}\Omega_{K}^{q} \to \operatorname{fil}_{\underline{n}}W_{m}\Omega_{K}^{q+1}/V^{m-1}\operatorname{fil}_{\underline{n}}\Omega_{K}^{q+1}) = F^{m-1}(\operatorname{fil}_{\underline{n}}W_{m}\Omega_{K}^{q}). \\ & \mathcal{7}. \ \operatorname{Ker}(V^{m-1}:\operatorname{fil}_{\underline{n}}\Omega_{K}^{q} \to \operatorname{fil}_{\underline{n}}W_{m}\Omega_{K}^{q}/dV^{m-1}\operatorname{fil}_{\underline{n}}\Omega_{K}^{q-1}) = F^{m}dV(\operatorname{fil}_{\underline{n}}W_{m}\Omega_{K}^{q-1}). \end{aligned}$ 

*Proof.* We first note that all statements are classical for  $W_{\cdot}\Omega_{K}^{\bullet}$ . Now, the item (1) follows directly from Lemma 4.3.3(1).

The item (2) is a special case of the more general claim that

$$\operatorname{Ker}(F^{l}:\operatorname{fil}_{\underline{n}}W_{m+l}\Omega_{K}^{q} \to \operatorname{fil}_{\underline{n}}W_{m}\Omega_{K}^{q}) = V^{m}(\operatorname{fil}_{\underline{n}}W_{l}\Omega_{K}^{q})$$

$$(4.3.7)$$

for all  $m \ge 0$  and  $l \ge 1$ . Since this general claim is known for  $W_{\cdot}\Omega_{K}^{\bullet}$  by Proposition 1.1.6, one gets that  $V^{m}(\operatorname{fil}_{\underline{n}}W_{l}\Omega_{K}^{q}) \subseteq \operatorname{Ker}(F^{l}:\operatorname{fil}_{\underline{n}}W_{m+l}\Omega_{K}^{q} \to \operatorname{fil}_{\underline{n}}W_{m}\Omega_{K}^{q})$ . We shall prove the reverse inclusion by induction on l. We first consider the case l = 1. It suffices to show that  $V^{m}(\Omega_{K}^{q}) \cap \operatorname{fil}_{\underline{n}}W_{m+1}\Omega_{K}^{q} = V^{m}(\operatorname{fil}_{\underline{n}}\Omega_{K}^{q})$ . But this follows from Lemma 4.2.12.

We now assume  $l \ge 2$  and  $m \ge 1$  (the m = 0 case is trivial). We let  $\omega \in \operatorname{fil}_{\underline{n}} W_{m+l} \Omega_K^q$ be such that  $F^l(\omega) = 0$ . We let  $\omega' = F(\omega)$  so that  $\omega' \in \operatorname{fil}_{\underline{n}} W_{m+l-1} \Omega_K^q$  and  $F^{l-1}(\omega') = 0$ . The induction hypothesis implies that  $F(\omega) = \omega' = V^m(x)$  for some  $x \in \operatorname{fil}_{\underline{n}} W_{l-1} \Omega_K^q$ . This implies that  $F^{l-2}d(x) = F^{m+l-2}dV^m(x) = F^{m+l-2}dF(\omega) = 0$ . We conclude from item (1) that x = F(y) from some  $y \in \operatorname{fil}_{\underline{n}} W_l \Omega_K^q$ . This yields

$$F(\omega - V^m(y)) = F(\omega) - FV^m(y) = F(\omega) - V^m F(y) = F(\omega) - V^m(x) = 0.$$

Using the l = 1 case, we get  $\omega - V^m(y) \in V^m(\operatorname{fil}_n W_l \Omega_K^q)$ . In particular,  $\omega \in V^m(\operatorname{fil}_n W_l \Omega_K^q)$ .

To prove (3), it suffices to show that

$$F^{m}(\operatorname{fil}_{\underline{n}}W_{m+1}\Omega_{K}^{q}) = F^{m}(W_{m+1}\Omega_{K}^{q}) \bigcap \operatorname{fil}_{\underline{n}}W_{m+1}\Omega_{K}^{q}$$

But this follows from Lemmas 4.3.4 and 4.3.5 because of the classical identity  $Z_m \Omega_K^q = F^m(W_{m+1}\Omega_K^q)$  (cf. Proposition 1.1.6).

The proof of (4) is identical to that of the l = 1 case of (4.3.7) using Lemma 4.2.12 and Proposition 1.1.6.

To prove (5), we can assume  $m \ge 2$ . In the latter case, we only need to show that  $\operatorname{Ker}(V^{m-1}) \subset F^{m-2}d(\operatorname{fil}_{\underline{n}}W_{m-1}\Omega_K^{q-1})$ . By Lemma 4.3.4, we can replace  $F^{m-2}d(\operatorname{fil}_{\underline{n}}W_{m-1}\Omega_K^{q-1})$ 

by  $B_{m-1} \operatorname{fil}_{\underline{n}} \Omega_K^q$ . By Proposition 1.1.6, we are thus reduced to showing that  $B_{m-1} \Omega_K^q \cap \operatorname{fil}_{\underline{n}} \Omega_K^q$ =  $B_{m-1} \operatorname{fil}_{\underline{n}} \Omega_K^q$ . But this follows from Lemma 4.3.5.

The proof of (6) and (7) are analogous to the proof of (3) and (5), respectively, using Lemma 4.2.12.  $\hfill \Box$ 

**Lemma 4.3.11.** Assuming  $\underline{n} \ge (-1, ..., -1)$ , we have

$$d(\operatorname{fil}_{\underline{n}} W_m \Omega_K^q) \subset \operatorname{Image}(Z_1 \operatorname{fil}_{\underline{n}} W_m \Omega_K^{q+1} \xrightarrow{1-C} \operatorname{fil}_{\underline{n}} W_m \Omega_K^{q+1}).$$

*Proof.* For  $\omega \in \text{fil}_{\underline{n}} W_m \Omega_K^q$ , we have  $d(\omega) = F dV(\omega) - R dV(\omega) + F dV^2 R(\omega) - R dV^2 R(\omega) + F dV^3 R^2(\omega) - R dV^3 R^2(\omega) + \dots + F dV^m R^{m-1}(\omega) - R dV^m R^{m-1}(\omega)$ , as  $R dV^m R^{m-1}(\omega) = 0$ . Hence, we have

$$d(\omega) = (F - R)((dV + dV^2R + \dots + dV^mR^{m-1})(\omega)) = (1 - C)Fd(x),$$

where we let  $x = (V + V^2 R + \dots + V^m R^{m-1})(\omega)$ . It suffices therefore to show that  $Fd(x) \in Z_1 \operatorname{fil}_{\underline{n}} W_m \Omega_K^{q+1}$ . To this end, we first note that  $F^{m-1}d(Fd(x)) = pF^m d^2(x) = 0$ , and this implies that  $Fd(x) \in Z_1 W_m \Omega_K^{q+1}$ . Secondly, Lemma 2.1.7 implies that  $Fd(x) \in \operatorname{fil}_{\underline{n}} W_m \Omega_K^{q+1}$  because  $\omega \in \operatorname{fil}_{\underline{n}} W_m \Omega_K^q$ . This achieves the proof.  $\Box$ 

Remark 4.3.12. The same proof also shows that

$$d(\operatorname{fil}_{-\underline{1}} W_m \Omega_K^q) \subset \operatorname{Image}(\operatorname{fil}_{-\underline{1}} W_{m+1} \Omega_K^{q+1} \xrightarrow{R-F} \operatorname{fil}_{-\underline{1}} W_m \Omega_K^{q+1}).$$

**Lemma 4.3.13.** The abelian group  $\frac{\operatorname{fil}_{\underline{n}}W_m\Omega_K^q}{d(\operatorname{fil}_{\underline{n}}W_m\Omega_K^{q-1})}$  is generated by the images of elements of the form  $V^i([x]_{m-i})\operatorname{dlog}([x_1]_m) \wedge \cdots \wedge \operatorname{dlog}([x_q]_m)$ , where  $[x]_{m-i} \in \operatorname{fil}_{\underline{n}}W_{m-i}(K)$ ,  $x_i \in K^{\times}$  and  $0 \leq i \leq m-1$ .

Proof. The q = 0 case of the lemma is clear and so we assume  $q \ge 1$ . We know from Lemma 2.1.7(2) that  $\operatorname{fil}_{\underline{n}}\Omega_K^q = \pi^{-\underline{n}}\Omega_A^q(\log\pi)$   $(\pi^{-\underline{n}} = x_1^{-n_1}\cdots x_r^{-n_r})$ , which is clearly generated as an abelian group by elements of the desired kind. This proves our assertion when m = 1. Using this case and induction on m, the general case follows at once by Proposition 4.2.10 and the observation that  $[x]_{m-i-1} \in \operatorname{fil}_{\underline{n}/p}W_{m-i-1}(K)$ implies  $[x]_{m-i} \in \operatorname{fil}_{\underline{n}}W_{m-i}(K)$  and  $R(V^i([x]_{m-i})w_1) = V^i([x]_{m-i-1})w_2$  if we let  $w_1 = \operatorname{dlog}([x_1]_m) \wedge \cdots \wedge \operatorname{dlog}([x_q]_m)$  and  $w_2 = \operatorname{dlog}([x_1]_{m-1}) \wedge \cdots \wedge \operatorname{dlog}([x_q]_{m-1})$ .

Recall from Proposition 1.1.6 that the multiplication map  $p: W_m \Omega_K^q \to W_m \Omega_K^q$  has a factorization  $W_m \Omega_K^q \xrightarrow{R} W_{m-1} \Omega_K^q \xrightarrow{p} W_m \Omega_K^q$ .

**Lemma 4.3.14.** Let  $\omega \in W_{m-1}\Omega_K^q$ . Then  $\omega \in \operatorname{fil}_{\underline{n}/p}W_{m-1}\Omega_K^q$  if and only if  $\underline{p}(\omega) \in \operatorname{fil}_{\underline{n}}W_m\Omega_K^q$ .

Proof. The 'only if' part is clear because  $R: \operatorname{fl}_{\underline{n}} W_m \Omega_K^q \to \operatorname{fl}_{\underline{n}/p} W_{m-1} \Omega_K^q$  is surjective. For the reverse implication, we let  $\widetilde{\omega} \in W_m \Omega_K^q$  be such that  $R(\widetilde{\omega}) = \omega$ . This yields  $VF(\widetilde{\omega}) = p\widetilde{\omega} = \underline{p}(w) \in \operatorname{fl}_{\underline{n}} W_m \Omega_K^q$ . It follows from Lemma 4.3.10(2) that  $VF(\widetilde{\omega}) = V(y')$  for some  $y' \in \operatorname{fl}_{\underline{n}} W_{m-1} \Omega_K^q$ . Proposition 1.1.6(8) now implies that  $F(\widetilde{\omega}) - y' = FdV^{m-1}(z')$  for some  $z' \in \Omega_K^{q-1}$ . That is,  $F(\widetilde{\omega} - dV^{m-1}(z')) = y' \in \operatorname{fl}_{\underline{n}} W_{m-1} \Omega_K^q$ . Lemma 4.3.10(1) in turn implies that  $F(\widetilde{\omega} - dV^{m-1}(z')) = F(y'')$  for some  $y'' \in \operatorname{fl}_{\underline{n}} W_m \Omega_K^q$ . By Proposition 1.1.6(7), we can find  $z'' \in \Omega_K^q$  such that  $\widetilde{\omega} - dV^{m-1}(z') - y'' = V^{m-1}(z'')$ . That is,  $\widetilde{\omega} - y'' = V^{m-1}(z'') + dV^{m-1}(z')$ . Equivalently,  $R(\widetilde{\omega} - y'') = 0$ . That is,  $\omega = R(\widetilde{\omega}) = R(y'') \in \operatorname{fl}_{\underline{n}/p} W_m \Omega_K^q$ . This concludes the proof.

The following result will be used later on in this thesis.

**Lemma 4.3.15.** Let  $X = \operatorname{Spec} A$  and  $X_{\pi} = D((\pi))$ . Then  $H^{i}_{\operatorname{zar}}(X_{\pi}, \mathcal{K}^{M}_{q, X_{\pi}}) = 0$  for i > 0.

*Proof.* We shall prove the lemma by induction on r = no. of components of the divisor given by  $\pi$ . Suppose first that r = 1 and let  $Y = \text{Spec}(A/(\pi))$ . For any regular scheme W and  $n \ge 1$ , we let  $G_n(W)$  denote the Gersten complex

$$\bigoplus_{w \in W^{(0)}} K_n^M(k(w)) \xrightarrow{\partial} \bigoplus_{w \in W^{(1)}} K_{n-1}^M(k(w)) \xrightarrow{\partial} \cdots$$

We then have an exact sequence

$$0 \to G_{q-1}(Y)[-1] \to G_q(X) \to G_q(X_{\pi}) \to 0.$$

Since  $H^i(G_q(X)) = H^i(G_{q-1}(Y)) = 0$  for i > 0 by [42, Prop. 10(8)]), it follows that  $H^i_{\text{zar}}(X_{\pi}, \mathcal{K}^M_{q, X_{\pi}}) = H^i(G_q(X_{\pi})) = 0$  for i > 0.

Suppose now that  $r \ge 2$  and set  $\pi_1 = x_2 \cdots x_r$  and  $A_1 = A_{\pi_1}$ . We let,  $X_1 = \text{Spec}(A_1)$ ,  $Y_1 = \text{Spec}(A_1/(x_1))$ , and  $U = X_1 \setminus Y = X_{\pi}$ . Then we have the exact sequence

$$0 \to G_{q-1}(Y_1)[-1] \to G_q(X_1) \to G_q(U) \to 0.$$

We see by induction on r that  $H^i(G_{q-1}(Y_1)) = 0 = H^i(G_q(X_1))$  for i > 0. Hence we conclude that  $H^i_{\text{zar}}(X_{\pi}, \mathcal{K}^M_{q, X_{\pi}}) = H^i(G_q(U)) = 0$  for i > 0.

**Definition 4.3.16.** Let X be a Noetherian regular F-finite  $\mathbb{F}_p$ -scheme and let  $E = \sum_i E_i$ be a simple normal crossing divisor on X with irreducible components  $E_1, \ldots, E_r$ . Let  $j: U \hookrightarrow X$  be the inclusion of the complement of E in X. Let  $\eta_i$  denote the generic point of  $E_i$  and let  $\widehat{K}_i$  be the quotient field of  $\widehat{\mathcal{O}_{X,\eta_i}}$ . Let  $j_i: \operatorname{Spec}(\mathcal{O}_{X,\eta_i}) \to X$  denote the canonical map. If  $D = \sum_i n_i E_i \in \operatorname{Div}_E(X)$  with  $n_i \ge -1$  for each i, we let  $W_m \mathcal{F}_D^{q,\bullet}$  denote the 2-term complex of Zariski sheaves

$$\left(Z_1 \operatorname{fil}_D W_m \Omega_U^q \xrightarrow{1-C} \operatorname{fil}_D W_m \Omega_U^q\right), \tag{4.3.8}$$

where  $Z_1 \operatorname{fil}_D W_m \Omega_U^q \coloneqq \operatorname{fil}_D W_m \Omega_U^q \cap j_* Z_1 W_m \Omega_U^q$ .

Combining the local results — Corollary 4.2.11, Proposition 4.2.10, Lemmas 4.3.3 and 4.3.10, Corollary 4.3.9 — together with Lemma 2.1.7(3), we obtain the following.

**Theorem 4.3.17.** Let  $D \in \text{Div}_E(X)$ . We have the following.

1. There exists an exact sequence of Zariski sheaves of  $W_m \mathcal{O}_X$ -modules

$$0 \to \operatorname{fil}_D W_m \Omega_U^q \to j_*(W_m \Omega_U^q) \to \bigoplus_{i=1}^r \frac{(j_i)_*(W_m \Omega_{\widehat{K}_i}^q)}{(j_i)_*(\operatorname{fil}_{n_i} W_m \Omega_{\widehat{K}_i}^q)}.$$

2. There is a short exact sequence of Zariski sheaves of  $W_m \mathcal{O}_X$ -modules

$$0 \to V^{m-1}(\operatorname{fil}_D\Omega_U^q) + dV^{m-1}(\operatorname{fil}_D\Omega_U^{q-1}) \to \operatorname{fil}_DW_m\Omega_U^q \xrightarrow{R} \operatorname{fil}_{D/p}W_{m-1}\Omega_U^q \to 0.$$

3. Ker
$$(F^{m-1}d: \operatorname{fil}_D W_m \Omega_U^q \to \operatorname{fil}_D \Omega_U^{q+1}) = Z_1 \operatorname{fil}_D W_m \Omega_U^q = F(\operatorname{fil}_D W_{m+1} \Omega_U^q)$$

4. There exist isomorphisms of Zariski sheaves of  $W_m \mathcal{O}_X$ -modules

$$\overline{F}: \operatorname{fil}_{D/p} W_m \Omega_U^q \xrightarrow{\cong} \frac{Z_1 \operatorname{fil}_D W_m \Omega_U^q}{dV^{m-1} (\operatorname{fil}_D \Omega_U^{q-1})}; \quad \overline{C}: \frac{Z_1 \operatorname{fil}_D W_m \Omega_U^q}{dV^{m-1} (\operatorname{fil}_D \Omega_U^{q-1})} \xrightarrow{\cong} \operatorname{fil}_{D/p} W_m \Omega_U^q.$$

5. Ker
$$(F^{m-1}: \operatorname{fil}_D W_m \Omega_U^q \to \operatorname{fil}_D \Omega_U^q) = V(\operatorname{fil}_D W_{m-1} \Omega_U^q).$$

- 6. Ker $(dV^{m-1}: \operatorname{fil}_D\Omega^q_U \to \operatorname{fil}_DW_m\Omega^{q+1}_U) = F^m(\operatorname{fil}_DW_{m+1}\Omega^q_U) = Z_m \operatorname{fil}_D\Omega^q_U.$
- 7. Ker $(V: \operatorname{fil}_D W_m \Omega_U^q \to \operatorname{fil}_D W_{m+1} \Omega_U^q) = dV^{m-1} (\operatorname{fil}_D \Omega_U^{q-1}).$

8. Ker
$$(V^m: \operatorname{fil}_D \Omega^q_U \to \operatorname{fil}_D W_{m+1} \Omega^q_U) = F^m dV(\operatorname{fil}_D W_m \Omega^{q-1}_U) = B_m \operatorname{fil}_D \Omega^q_U.$$

- 9. Ker $(dV^{m-1}: \operatorname{fil}_D\Omega_U^q \to \operatorname{fil}_DW_m\Omega_U^{q+1}/V^{m-1}\operatorname{fil}_D\Omega_U^{q+1}) = F^{m-1}(\operatorname{fil}_{\underline{n}}W_m\Omega_K^q).$
- 10. Ker $(V^{m-1}: \operatorname{fil}_D \Omega^q_U \to \operatorname{fil}_D W_m \Omega^q_U / dV^{m-1} \operatorname{fil}_D \Omega^{q-1}_U) = F^m dV(\operatorname{fil}_D W_m \Omega^{q-1}_U).$
- 11. Let  $D \ge -E$ . In the derived category of Zariski sheaves of abelian groups on X, there exists a distinguished triangle

$$W_1 \mathcal{F}_D^{q\bullet} \xrightarrow{V^{m-1}} W_m \mathcal{F}_D^{q,\bullet} \xrightarrow{R} W_{m-1} \mathcal{F}_{D/p}^{q,\bullet} \xrightarrow{+} W_1 \mathcal{F}_D^{q\bullet} [1].$$

12.  $\lim_{D \ge -E} W_m \mathcal{F}_D^{q,\bullet} = \left( j_* Z_1 W_m \Omega_U^q \xrightarrow{1-C} j_* W_m \Omega_U^q \right) = R j_* W_m \Omega_U^q.$ 

## Chapter 5

## **Comparison with Kato's filtration**

Let  $k = \mathbb{F}_p$ . For  $q \in \mathbb{Z}$  and  $n = p^m r \in \mathbb{N}$  with (p, r) = 1, we let  $\mathbb{Z}/n(q) = \mathbb{Z}/r(q) \oplus W_m \Omega_{(-,\log)}^q[-q]$  and consider it as a complex of étale sheaves on the big étale site **Et**\_k of Spec(k), where the first summand on the right is the standard étale twist (cf. [56, p. 163]). If  $r' \in \mathbb{N}$  such that  $r \mid r'$  and (p, r') = 1, we have a natural map  $\mathbb{Z}/r(q) \to \mathbb{Z}/r'(q)$ , induced by the maps  $Z/r \cong \frac{1}{r}\mathbb{Z}/\mathbb{Z} \to \frac{1}{r'}\mathbb{Z}/\mathbb{Z} \cong \mathbb{Z}/r'$  and  $\mu_r \to \mu_{r'}$  ( $\mu_r$  is the sheaf of r-th roots of unity on  $\mathbf{Et}_k$ ). On the other hand, the canonical map  $\overline{p}: W_m \Omega_{(-,\log)}^q \to W_{m+1} \Omega_{(-,\log)}^q$  induces a canonical map  $\mathbb{Z}/p^m(q) \to \mathbb{Z}/p^{m+1}(q)$ . As a result, we have canonical maps  $\mathbb{Z}/n(q) \to \mathbb{Z}/n'(q)$ , when  $n \mid n'$ . For ring R, we let  $H_m^q(R) = H_{\text{ét}}^q(\text{Spec}(R), \mathbb{Z}/p^m(q-1))$ . We let  $H^q(R) = H_{\text{ét}}^q(\text{Spec}(R), \mathbb{Q}/\mathbb{Z}(q-1)) = \lim_{n \to \infty} H_{\text{ét}}^q(R),$  with the transition maps induced by the map  $\mathbb{Z}/p^m(q) \to \mathbb{Z}/p^{m+1}(q)$ . For a scheme X, one defines the groups  $H^q(X)$  and  $H_m^q(X)$  in a similar way. The purpose of this chapter is to provide a cohomological description of Kato's ramification filtration.

### 5.1 Comparison with Local Kato filtration

We let  $K_*^M(R)$  denote the (improved) Milnor K-theory of R as defined in [42]. This coincides with the Milnor K-theory of R as defined in [33] if R is a local ring with infinite residue field (cf. [22, Lem. 2.2]). We let dlog:  $K_q^M(R)/p^m \to W_m \Omega_R^q$  denote the dlog map induced by dlog( $\{x_1, \ldots, x_q\}$ ) = dlog[ $x_1$ ]<sub> $m \land \cdots \land$  dlog[ $x_q$ ]<sub>m</sub> (cf. [30], [74, Lem. 3.2.8]) and let  $W_m \Omega_{R,\log}^q$  denote its image. This map is multiplicative with respect to the product structures of  $K_*^M(R)$  and  $W_m \Omega_R^{\bullet}$ . The map dlog:  $K_q^M(R)/p^m \to W_m \Omega_{R,\log}^q$  is bijective if R is a regular local ring (cf. [58, Thm. 5.1]).</sub>

#### 5.1.1 Kato filtration

If R is a regular local ring, the exact sequence

$$0 \to W_m \Omega^q_{(-),\log} \to Z_1 W_m \Omega^q_{(-)} \xrightarrow{1-C} W_m \Omega^q_{(-)} \to 0$$
(5.1.1)

of sheaves on  $(\operatorname{Spec} R)_{\text{ét}}$  gives an exact sequence

$$0 \to W_m \Omega^q_{R,\log} \to Z_1 W_m \Omega^q_R \xrightarrow{1-C} W_m \Omega^q_R \xrightarrow{\delta^q_m} H^{q+1}_m(R) \to 0.$$
(5.1.2)

We also have a commutative diagram (cf.  $[36, \S 1.3]$ )

$$\begin{array}{cccc}
K_q^M(R)/p^m \otimes W_m \Omega_R^{q'} & \stackrel{\cup}{\longrightarrow} W_m \Omega_R^{q+q'} \\
& & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & &$$

where the bottom row is induced by the Bloch-Kato map  $\beta_R^q: K_q^M(R) \to H_{\text{\acute{e}t}}^q(R, \mathbb{Z}/p^m(q))$ , followed by the cup product on the étale cohomology of  $W_m \Omega_{(-),\log}^{\bullet}$  on Spec (R). The top row is induced by the dlog map, followed by the wedge product. We shall denote the composite map  $W_m \Omega_R^q \xrightarrow{\delta_m^q} H_m^{q+1}(R) \to H^{q+1}(R)$  also by  $\delta_m^q$ .

We now let A be an F-finite Henselian discrete valuation ring containing k. Let  $\mathfrak{m} = (\pi)$  denote the maximal ideal of A. Let  $\mathfrak{f} = A/\mathfrak{m}$  and K = Q(A).

Lemma 5.1.1. The map  $H^1_{\acute{e}t}(K, W_m\Omega^q_{K,\log}) \rightarrow {}_{p^m}H^{q+1}(K)$  is a bijection.

*Proof.* We let X = Spec(K). Since the étale sheaf  $\mathcal{K}_{q,X}^M$  has no *p*-torsion (see [16, Thm. 8.1]), there is an exact sequence of sheaves

$$0 \to \mathcal{K}^M_{q,X}/p^m \to \varinjlim_r \mathcal{K}^M_{q,X}/p^r \xrightarrow{p^m} \varinjlim_r \mathcal{K}^M_{q,X}/p^r \to 0.$$

Considering the cohomology and observing that  $H^0_{\text{\acute{e}t}}(X, \mathcal{K}^M_{q,X}/p^r) \cong H^0_{\text{\acute{e}t}}(X, W_r \Omega^q_{X, \log}) \cong W_r \Omega^q_{K, \log} \cong K^M_q(K)/p^r$ , we get an exact sequence

$$0 \to K_q^M(K)/p^m \to K_q^M(K) \otimes \mathbb{Q}_p/\mathbb{Z}_p \xrightarrow{p^m} K_q^M(K) \otimes \mathbb{Q}_p/\mathbb{Z}_p \xrightarrow{\partial} H^1_{\text{\'et}}(X, W_r\Omega_{X, \log}^q) \to H^{q+1}(K)\{p\} \xrightarrow{p^m} H^{q+1}(K)\{p\}.$$

Since  $K_q^M(K) \otimes \mathbb{Q}_p/\mathbb{Z}_p \xrightarrow{p^m} K_q^M(K) \otimes \mathbb{Q}_p/\mathbb{Z}_p$  is surjective, the lemma follows.

We fix integers  $n, q \ge 0$  and  $m \ge 1$ . Let  $T_n^{m,q}(K)$  denote the cokernel of the map  $(1-C): Z_1 \operatorname{fil}_n W_m \Omega_K^q \to \operatorname{fil}_n W_m \Omega_K^q$ .

**Lemma 5.1.2.** Let  $X = \operatorname{Spec} A, m \ge 1, q \ge 0$ . Let  $E = V((\pi))$ ,  $U = X \setminus E$  and  $D_n = nE$ , for  $n \ge 0$ . Then, we have

1. 
$$H^i_{\acute{e}t}(X, Z_1 \operatorname{fil}_{D_n} W_m \Omega^q_U) = H^i_{\acute{e}t}(X, \operatorname{fil}_{D_n} W_m \Omega^q_U) = 0, \text{ for } i \ge 1.$$
  
2.

$$\mathbb{H}_{\acute{e}t}^{i}(X, W_m \mathcal{F}_{D_n}^q) = \begin{cases} K_q^M(K)/p^m & \text{if } i = 0\\ T_n^{m,q}(K) & \text{if } i = 1\\ 0 & \text{if } i > 1. \end{cases}$$

Proof. (1) Recall that higher cohomology of coherent sheaves on affine space is zero. Hence  $H^i_{\text{ét}}(X, Z_1 \text{fil}_{D_n} W_m \Omega_U^q) = H^i_{\text{ét}}(W_m(X), Z_1 \text{fil}_{D_n} W_m \Omega_U^q) = 0$ . Similarly, we have  $H^i_{\text{ét}}(X, \text{fil}_{D_n} W_m \Omega_U^q) = 0$ . Here,  $W_m(X)$  denotes the scheme  $(X, W_m \mathcal{O}_X)$ .

(2) We look at the following hypercohomology long exact sequence

$$0 \to \mathbb{H}^{0}_{\text{\acute{e}t}}(W_m \mathcal{F}^q_{D_n}) \to H^{0}_{\text{\acute{e}t}}(Z_1 \text{fil}_{D_n} W_m \Omega^q_U) \xrightarrow{1-C} H^{0}(\text{fil}_{D_n} W_m \Omega^q_U) \to \mathbb{H}^{1}_{\text{\acute{e}t}}(W_m \mathcal{F}^q_{D_n}) \to 0,$$
(5.1.4)

where  $\mathbb{H}^{i}_{\acute{e}t}(\mathcal{E})$  (resp.  $H^{i}_{\acute{e}t}(\mathcal{E})$ ) is the short hand notation for  $\mathbb{H}^{i}_{\acute{e}t}(X, \mathcal{E})$  (resp.  $H^{i}_{\acute{e}t}(X, \mathcal{E})$ ) and the last term of (5.1.4) is 0 because of item (1). Since  $W_m \Omega^q_A \subset \operatorname{fil}_n W_m \Omega^q_K \subset W_m \Omega^q_K$ , we have

$$K_q^M(A)/p^m \xrightarrow{\text{dlog}} \operatorname{Ker}(1-C:Z_1W_m\Omega_A^q \to W_m\Omega_A^q)$$
  

$$\subset \operatorname{Ker}(1-C:Z_1\operatorname{fil}_nW_m\Omega_K^q \to \operatorname{fil}_nW_m\Omega_K^q) = \mathbb{H}^0_{\operatorname{\acute{e}t}}(A, W_m\mathcal{F}_{D_n}^q)$$
  

$$\subset \operatorname{Ker}(1-C:Z_1W_m\Omega_K^q \to W_m\Omega_K^q) = K_q^M(K)/p^m.$$

Since A is DVR, we note that  $dlog(K_q^M(K)/p^m) \subset Z_1 W_m \Omega_K^q$  is generated by (as a group) elements of the form

- 1.  $dlog([a_1]_m) \wedge \cdots \wedge dlog([a_q]_m)$  and
- 2. dlog( $[a_1]_m$ )  $\wedge \cdots \wedge dlog([a_{q-1}]_m) \wedge dlog([\pi_x]_m)$ ,

where where  $a_i \in A^*$ . However, the elements of the form (1) and (2) also lie within  $Z_1 \text{fil}_0 W_m \Omega_K^q \subset Z_1 \text{fil}_n W_m \Omega_K^q$ , for  $n \ge 0$ . Hence, we conclude that

$$K_q^M(K)/p^m \cong \mathbb{H}^0_{\text{\'et}}(A, W_m \mathcal{F}^q_{D_n}), \text{ and}$$
$$\mathbb{H}^1_{\text{\'et}}(X, W_m \mathcal{F}^q_{D_n}) = \operatorname{Coker}(1 - C : Z_1 \operatorname{fil}_n W_m \Omega^q_K \to \operatorname{fil}_n W_m \Omega^q_K) = T_n^{m,q}(K)$$

Also, we get that  $\mathbb{H}^{i}_{\text{ét}}(X, W_m \mathcal{F}^{q}_{D_n}) = 0$  by looking at the hypercohomology long exact sequence and using item (1). This proves the lemma.

We have an exact sequence

$$0 \to W_m \Omega^q_{K,\log} \to Z_1 \operatorname{fil}_n W_m \Omega^q_K \xrightarrow{1-C} \operatorname{fil}_n W_m \Omega^q_K \xrightarrow{\partial^{m,q}_n} T^{m,q}_n(K) \to 0$$
(5.1.5)

which canonically maps to the exact sequence (5.1.2) with R = K. We let  $\overline{\delta}_n^{m,q}: T_n^{m,q}(K) \to H^{q+1}(K)$  be the induced map.

By Lemma 2.1.7(8) (for D = 0,  $D' = nV(\pi)$ ), we see that the product map on the top row of (5.1.3) (with R = K) restricts to a map

$$K_q^M(K)/p^m \otimes \operatorname{fil}_n W_m \Omega_K^{q'} \xrightarrow{\cup} \operatorname{fil}_n W_m \Omega_K^{q+q'}.$$
 (5.1.6)

It follows from Lemmas 4.3.11 and 4.3.13 that the composite map  $K_q^M(K) \otimes \operatorname{fil}_n W_m(K) \xrightarrow{\cup} \operatorname{fil}_n W_m \Omega_K^q \xrightarrow{\partial_n^{m,q}} T_n^{m,q}(K)$  is surjective. Combining this with Proposition 3.2.3 and Corollary 4.2.11 (for r = 1), we get the following.

Corollary 5.1.3. The map

$$\overline{\delta}_n^{m,q}: T_n^{m,q}(K) \to \operatorname{Image}(\operatorname{fil}_n W_m(K) \otimes K_q^M(K) \xrightarrow{\lambda_m^q} H^{q+1}(K))$$

is a bijection.

*Proof.* By Lemma 5.1.1, we only need to show the map  $T_n^{m,q}(K) \to H_m^{q+1}(K)$  is injective. If K is complete then this follows from Proposition 3.2.3. If K is not complete, then we reduce to the complete case by Corollary 4.2.11 (for r = 1).

We now recall Kato's filtration on  $H^q(K)$  from [36]. For any A-algebra R, let  $R^h$  denote the Henselization of R with respect to the ideal  $\mathfrak{m}R$ . We let  $\operatorname{Spec}(R/\mathfrak{m}R) \xrightarrow{\iota}$  $\operatorname{Spec}(R) \xleftarrow{j} \operatorname{Spec}(R[\pi^{-1}])$  be the inclusions and let

$$V^{q}(R) = \lim_{t \to l} H^{q}_{\text{ét}}(\operatorname{Spec}(R/\mathfrak{m}R), \iota^{*}Rj_{*}(\mathbb{Z}/l(q-1))) \cong H^{q}(R^{h}[\pi^{-1}]).$$
(5.1.7)

The following definition is due to Kato [36, Defn. 2.1], inspired by a filtration of  $W_m(K)$  by Brylinski [9].

**Definition 5.1.4.** We let  $\operatorname{fil}_n^{\operatorname{bk}} H^q(K)$  be the set of all elements  $\chi \in H^q(K)$  such that  $\{\chi, 1 + \pi^{n+1}T\} = 0$  in  $V^{q+1}(A[T])$  under the product map

$$H^{q}(K) \otimes K_{1}^{M}((A[T])^{h}[\pi^{-1}]) \to V^{q+1}(A[T]).$$

We let  $\operatorname{gr}_0^{\operatorname{bk}} H^q(K) = \operatorname{fil}_0^{\operatorname{bk}} H^q(K)$  and  $\operatorname{gr}_n^{\operatorname{bk}} H^q(K) = \operatorname{fil}_n^{\operatorname{bk}} H^q(K) / \operatorname{fil}_{n-1}^{\operatorname{bk}} H^q(K)$  if  $n \ge 1$ .

The following result of Kato (cf. [36, Thm. 3.2]) provides a description of  $\operatorname{fil}_n^{\operatorname{bk}} H^q(K)$  in terms of Milnor K-theory and Witt-vectors.

**Theorem 5.1.5.** One has the following.

1.  $\operatorname{fil}_n^{\operatorname{bk}} H^1(K)\{p\} = \bigcup_{m\geq 1} \delta_m^0(\operatorname{fil}_n W_m(K)).$ 

2.  $\operatorname{fl}_n^{\operatorname{bk}} H^{q+1}(K)\{p\}$  coincides with the image of

$$K_q^M(K) \otimes \operatorname{fil}_n^{\operatorname{bk}} H^1(K) \{p\} \xrightarrow{\cup} H^{q+1}(K).$$

In [43, Page 2 of the Erratum], Kerz and Saito proved the following refined version of Theorem 5.1.5(1).

**Theorem 5.1.6.** One has  $_{p^m}\operatorname{fil}_n^{\operatorname{bk}} H^1(K) = \delta_m^0(\operatorname{fil}_n W_m(K)).$ 

### 5.1.2 Proof of the main theorem

The main result of this section is the following refinement of Theorem 5.1.5.

**Theorem 5.1.7.** For  $n, q \ge 0$  and  $m \ge 1$ , the map  $\delta_m^q$ :  $\operatorname{fil}_n W_m \Omega_K^q \to H^{q+1}(K)$  induces an isomophism

$$\overline{\delta}_n^{m,q}: T_n^{m,q}(K) \xrightarrow{\cong} {}_{p^m} \operatorname{fil}_n^{\operatorname{bk}} H^{q+1}(K).$$

We shall prove this theorem in several steps. We fix  $n, q \ge 0$  and  $m \ge 1$  as before. When n > 0, we shall write  $n = p^r l$ , where  $r \ge 0$  and (p, l) = 1. We let  $S_n^{m,q}(K) = {}_{p^m} \operatorname{fil}_n^{\operatorname{bk}} H^{q+1}(K)$ . Since  $T_n^{m,q}(K)$  is a  $p^m$ -torsion group, we already deduce from Corollary 5.1.3, Theorem 5.1.5 and (5.1.3) that the map  $\overline{\delta}_n^{m,q}$  has a factorization  $\overline{\delta}_n^{m,q}: T_n^{m,q}(K) \hookrightarrow S_n^{m,q}(K)$ . In what follows, our goal is to show that this map is surjective.

We begin with the following.

Lemma 5.1.8. There is a commutative diagram

Proof. To prove the lemma, it suffices to show that VC = CV and  $V(Z_1W_m\Omega_K^q) \subset Z_1W_{m+1}\Omega_K^q$  (cf. Lemma 2.1.7(6)). For showing the first claim, it is enough to show that VCF = CVF. But this is clear because VCF = VR = RV = CFV = CVF. For the second claim, note that  $\omega \in Z_1W_m\Omega_K^q$  implies  $F^{m-1}d(\omega) = 0$ . This implies in turn that  $F^m dV(\omega) = F^{m-1}d(\omega) = 0$ . That is,  $V(\omega) \in Z_1W_{m+1}\Omega_K^q$ .

Using Lemma 5.1.8, we get a commutative diagram (cf.  $[36, \S 1.3]$ )

where the middle V is the unique map induced by the commutative diagram (5.1.8) and the right vertical arrow is the canonical inclusion  ${}_{p^m} \operatorname{fil}_n^{\operatorname{bk}} H^{q+1}(K) \rightarrow {}_{p^{m+1}} \operatorname{fil}_n^{\operatorname{bk}} H^{q+1}(K)$ . Since the canonical inclusion  $H^{q+1}_m(K) \rightarrow H^{q+1}_{m+1}(K)$  is induced by  $\overline{p} = V : W_m \Omega^q_{(-,\log)} \rightarrow W_{m+1} \Omega^q_{(-,\log)}$ , we see that the right square of (5.1.9) is commutative.

The commutative diagram

$$T_{n-1}^{m,q}(K) \to T_n^{m,q}(K)$$

$$\overline{\delta}_{n-1}^{m,q} \bigvee_{\lambda \in \mathcal{S}_n^{m,q}} \overline{\delta}_n^{m,q}$$

$$S_{n-1}^{m,q}(K) \hookrightarrow S_n^{m,q}(K)$$
(5.1.10)

implies that  $T_{n-1}^{m,q}(K) \hookrightarrow T_n^{m,q}(K)$  for every  $n \ge 1$ . We let  $\widetilde{T}_n^{m,q}(K) = \frac{T_n^{m,q}(K)}{T_{n-1}^{m,q}(K)}$  and  $\widetilde{S}_n^{m,q}(K) = \frac{S_n^{m,q}(K)}{S_{n-1}^{m,q}(K)}$  and let  $\psi_n^{m,q}: \widetilde{T}_n^{m,q}(K) \to \widetilde{S}_n^{m,q}(K)$  be the map induced by the vertical arrows in (5.1.10). Since the commutative diagram (5.1.8) is compatible with the change in values of  $n \ge 0$ , it induces maps  $V: \widetilde{T}_n^{m,q}(K) \to \widetilde{T}_n^{m+1,q}(K)$  as n varies over all positive integers. We let  $\widetilde{M}_n^{m,q}(K) = \frac{\widetilde{T}_n^{m,q}(K)}{V(\widetilde{T}_n^{m-1,q}(K))}$  and  $\widetilde{N}_n^{m,q}(K) = \frac{\widetilde{S}_n^{m,q}(K)}{\widetilde{S}_n^{m-1,q}(K)}$ . It is clear that  $\overline{\delta}_n^{m,q}$  descends to a homomorphism  $\overline{\delta}_n^{m,q}: \widetilde{M}_n^{m,q}(K) \to \widetilde{N}_n^{m,q}(K)$ .

Form now on, we shall assume n > 1, unless otherwise stated.

**Lemma 5.1.9.** The map  $V: \widetilde{T}_n^{m,q}(K) \to \widetilde{T}_n^{m+1,q}(K)$  is surjective for every  $m \ge r+1$ . *Proof.* Since  $V(adlog([x_1]_m) \land \dots \land dlog([x_q]_m) = V(a)dlog([x_1]_m) \land \dots \land dlog([x_q]_m)$ , (5.1.8) implies that the diagram

$$K_{q}^{M}(K) \otimes \operatorname{fil}_{n} W_{m}(K) \xrightarrow{\partial_{n}^{m,q}} T_{n}^{m,q}(K)$$

$$id \otimes V \downarrow \qquad \qquad \qquad \downarrow V$$

$$K_{q}^{M}(K) \otimes \operatorname{fil}_{n} W_{m+1}(K) \xrightarrow{\partial_{n}^{m+1,q}} T_{n}^{m+1,q}(K)$$
(5.1.11)

is commutative. Furthermore, the horizontal arrows are surjective by Corollary 5.1.3. It suffices therefore to show that  $\frac{\operatorname{fil}_n W_{m+1}(K)}{\operatorname{fil}_{n-1} W_{m+1}(K) + V(\operatorname{fil}_n W_m(K))} = 0$  for  $m \ge r+1$ .

To that end, we look at the commutative diagram of exact sequences

Since  $m \ge r+1$ , we see that  $\lfloor (n-1)/p^m \rfloor = \lfloor n/p^m - 1/p^m \rfloor = \lfloor n/p^m \rfloor$ . This implies that the right vertical arrow in (5.1.12) is bijective, which proves the asserted claim.

For the next three lemmas, we assume that  $1 \le m \le r$  and consider the map

$$\widetilde{\theta}_1: A \otimes \overbrace{A^{\times} \otimes \cdots \otimes A^{\times}}^{M} \to \widetilde{T}_n^{m,q}(K)$$

$$\widetilde{\theta}_1(x \otimes y_1 \otimes \cdots \otimes y_q) = \widehat{c}_n^{m,q}([x\pi^{-np^{1-m}}]_m \operatorname{dlog}([y_1]_m) \wedge \cdots \wedge \operatorname{dlog}([y_q]_m)) \mod T_{n-1}^{m,q}(K)$$

Note that this is defined because  $np^{1-m} = p^{r+1-m}l \in \mathbb{N}$  and  $[x\pi^{-np^{1-m}}]_m \in \operatorname{fil}_n W(K)$ .

**Lemma 5.1.10.**  $\widetilde{\theta}_1$  descends to a group homomorphism  $\theta_1^q: \Omega_{\mathfrak{f}}^q \to \widetilde{M}_n^{m,q}(K)$ .

Proof. For  $\{y_1, \ldots, y_q\} \in A^{\times}$ , we write  $\operatorname{dlog}([y_1]_m) \wedge \cdots \wedge \operatorname{dlog}([y_q]_m) = \operatorname{dlog}(\underline{y})$ . If  $x = a\pi$  for some  $a \in A$ , then  $\partial_n^{m,q}([x]_m \operatorname{dlog}(\underline{y})) = \partial_n^{m,q}([a\pi^{-(n-p^{m-1})p^{1-m}}]_m \operatorname{dlog}(\underline{y})) \in T_{n-1}^{m,q}(K)$ , as  $[a\pi^{-(n-p^{m-1})p^{1-m}}]_m \in \operatorname{fil}_{n-1}W_m(K)$ . Suppose next that  $y_i = 1 + a\pi$  for some  $a \in A$  and  $1 \leq i \leq q$ . By [36, Lem. 3.5], we see that as an element of  $\operatorname{fil}_n W_m \Omega_K^1$ ,  $[x]_m \operatorname{dlog}([1 + a\pi]_m)$  actually lies in  $\operatorname{fil}_{n-1} W_m \Omega_K^1$ . It follows that  $[x]_m \operatorname{dlog}(\underline{y}) \in \operatorname{fil}_{n-1} W_m \Omega_K^q$  and hence  $\partial_n^{m,q}([x]_m \operatorname{dlog}(\underline{y}))$  dies in  $\widetilde{T}_n^{m,q}(K)$ . We have thus shown that  $\widetilde{\theta}_1$  descends a group homomorphism

$$\widetilde{\theta}_1: F \otimes F^* \otimes \dots \otimes F^* \to \widetilde{T}_n^{m,q}(K) \twoheadrightarrow \widetilde{M}_n^{m,q}(K).$$
(5.1.13)

We now let  $\{x_1, \ldots, x_l\}$  and  $\{x'_1, \ldots, x'_{l'}\}$  be elements of  $A^{\times}$  such that  $\sum_i x_i - \sum_j x'_j = a\pi$  for some  $a \in A$ . Let  $y_1, \ldots, y_{q-1} \in A^{\times}$ . We claim  $\tilde{\theta}_1$  kills elements of the form

$$\left(\sum_{i} x_{i} \otimes x_{i} - \sum_{j} x_{i}' \otimes x_{i}'\right) \otimes y_{1} \otimes \cdots \otimes y_{q-1}$$

Writing  $w_0 = [\pi^{-np^{1-m}}]_m \operatorname{dlog}[y_1]_m \cdots \operatorname{dlog}[y_{q-1}]_m$ , we get the following in  $T_n^{m,q}(K)$ .

$$w_0\left(\sum_i [x_i]_m \operatorname{dlog}[x_i]_m - \sum_j [x_i']_m \operatorname{dlog}[x_j']_m\right) = w_0 d\left(\sum_i [x_i]_m - \sum_j [x_j']_m\right)$$

$$= w_0 d\left(\left[\sum_i x_i - \sum_j x_i'\right]\right) + V(\underline{b}) = w_0 d[a\pi]_m + w_0 dV(\underline{b})$$

for some  $\underline{b} = (a_{m-2}, \dots, a_1, a_0) \in W_{m-1}(A)$ .

Since  $w_0 \in \operatorname{fil}_n W_m \Omega_K^{q-1}$ , we get that  $[a\pi]_m w_0 \in \operatorname{fil}_{n-1} W_m \Omega_K^{q-1}$ , and hence  $w_0 d[a\pi]_m = [a\pi]_m w_0 \operatorname{dlog}[a\pi]_m \in \operatorname{fil}_{n-1} W_m \Omega_K^q$ . So, it dies in  $\widetilde{M}_n^{m,q}(K)$ . On the other hand,

$$w_0 dV(\underline{b}) = dV(F(w_0)\underline{b}) - V(Fd(w_0)\underline{b}) \in (1 - C)(\operatorname{fil}_n W_m \Omega_K^q) + V(\operatorname{fil}_n W_{m-1} \Omega_K^q)$$

by Lemmas 2.1.7(4),(5),(6) and 4.3.11 and hence also dies in  $\widetilde{M}_n^{m,q}(K)$ . We conclude from [6, Lem. 4.2] that  $\widetilde{\theta}_1$  descends to a homomorphism  $\theta_1^q: \Omega_{\mathfrak{f}}^q \to \widetilde{M}_n^{m,q}(K)$ .

An identical proof shows that  $\widetilde{\phi}: A \otimes A^{\times} \otimes \cdots \otimes A^{\times} \to \widetilde{T}_{n}^{r+1,q}(K)$  given by  $\widetilde{\theta}_{1}(x \otimes y_{1} \otimes \cdots \otimes y_{q}) = \partial_{n}^{m,q}([x\pi^{-l}]_{m} \operatorname{dlog}([y_{1}]_{m}) \wedge \cdots \wedge \operatorname{dlog}([y_{q}]_{m})) \mod T_{n-1}^{r+1,q}(K)$ , induces a homomorphism

$$\phi^q:\Omega^q_{\mathfrak{f}} \to \widetilde{M}_n^{r+1,q}(K). \tag{5.1.14}$$

Lemma 5.1.11.  $\theta_1^q(Z_1\Omega_f^q) = 0.$ 

*Proof.* By [36, § 3, p. 111],  $Z_1\Omega_{\mathfrak{f}}^q$  is generated by the elements of the types

- 1.  $x \operatorname{dlog}(x) \wedge \operatorname{dlog}(y_1) \wedge \cdots \wedge \operatorname{dlog}(y_{q-1})$  and
- 2.  $x^p \operatorname{dlog}(y_1) \wedge \cdots \wedge \operatorname{dlog}(y_q),$

where  $x \in \mathfrak{f}, y_j \in \mathfrak{f}^*, 1 \leq j \leq q$ .

We now let  $x, y_1, \ldots, y_q \in A^{\times}$ ,  $w = x^p \pi^{-np^{1-m}}$  and  $\operatorname{dlog}(\underline{y})_l = \operatorname{dlog}([y_1]_l) \wedge \cdots \wedge \operatorname{dlog}([y_q]_l)$ . Since m-1 < r and  $p^r | n$ , we can write  $w = (x\pi^{-(n/p)p^{1-m}})^p$ . Letting  $w' = x\pi^{-(n/p)p^{1-m}}$ , we get by Lemma 4.3.3(1) that

$$[w']_m \in \operatorname{fil}_{n/p} W_m(K) \text{ and } [w]_m \operatorname{clog}(\underline{y})_m = F([w']_{m+1} \operatorname{clog}(\underline{y})_{m+1}) \in Z_1 \operatorname{fil}_n W_m \Omega_K^q$$

In particular,  $\partial_n^{m,0} \circ (1-C)([w]_m \operatorname{dlog}(\underline{y})_m) = 0$  in  $T_n^{m,q}(K)$ . From this, we get

$$\partial_n^{m,q}([w]_m \operatorname{dlog}(\underline{y})_m) = \partial_n^{m,q} \circ C([w]_m \operatorname{dlog}(\underline{y})_m) = \partial_n^{m,q}([w']_m \operatorname{dlog}(\underline{y})_m)$$

Since the last term is in  $\operatorname{fil}_{n/p} W_m \Omega_K^q \subset \operatorname{fil}_{n-1} W_m \Omega_K^q$ , it follows that  $\partial_n^{m,q}([w]_m \operatorname{dlog}(\underline{y})_m)$  dies in  $\widetilde{M}_n^{m,q}(K)$ . This shows that the type (2) elements goes to zero under the map  $\theta_1^q$ .

Next, we let  $a = \pi^{-np^{1-m}}$  and  $w_l'' = \operatorname{dlog}([y_1]_l) \wedge \cdots \wedge \operatorname{dlog}([y_{q-1}]_l)$  so that

$$[xa]_m dlog([x]_m)w''_m = [a]_m d([x]_m)w''_m = d([xa]_mw''_m) - [x]_m d([a]_m)w''_m$$

Lemma 4.3.11 implies  $\partial_n^{m,q}([xa]_m \operatorname{dlog}([x]_m)w_m'') = \partial_n^{m,q}([x]_m d([a]_m)w_m'')$ . On the other hand,

$$[x]_m d([a]_m) w''_m = [x]_m w''_m (-n/p^{m-1}) [a]_m d\log[\pi]_m$$
  
=  $p(-n/p^m) [xa]_m w''_m d\log[\pi]_m$   
=  $(-n/p^m) VF([xa]_m w''_m d\log[\pi]_m).$ 

Since  $F([xa]_m w'' \operatorname{dlog}[\pi]_m) \in \operatorname{fil}_n W_{m-1}\Omega_K^q$ , we conclude that  $\partial_n^{m,q}([x]_m d([a]_m)w'')$  dies in  $\widetilde{M}_n^{m,q}(K)$ . Hence, so does  $\partial_n^{m,q}([xa]_m \operatorname{dlog}([x]_m)w'')$  (image of the elements of type (1)). This finishes the proof.

We next consider the map

$$\widetilde{\theta}_2 : A \otimes A^{\times} \otimes \cdots \otimes A^{\times} \to \widetilde{T}_n^{m,q}(K)$$

$$\widetilde{\theta}_2(a \otimes y_1 \otimes \cdots \otimes y_{q-1}) = \partial_n^{m,q}([x\pi^{-np^{1-m}}]_m \operatorname{dlog}([y_1]_m) \wedge \cdots \wedge \operatorname{dlog}([y_{q-1}]_m) \wedge \operatorname{dlog}([\pi]_m)),$$
  
mod  $T_{n-1}^{m,q}(K).$ 

**Lemma 5.1.12.**  $\widetilde{\theta}_2$  descends to a group homomorphism  $\theta_2^q: \frac{\Omega_{\mathfrak{f}}^{q-1}}{Z_1\Omega_{\mathfrak{f}}^{q-1}} \to \widetilde{M}_n^{m,q}(K).$ 

Proof. By Lemma 2.1.7(8), we see that the multiplication by  $\operatorname{dlog}([\pi]_m)$  preserves  $\operatorname{fil}_n W_m \Omega_K^{\bullet}$ . Since  $F^{m-1}d(\operatorname{adlog}([\pi]_m) = F^{m-1}d(\operatorname{a})\operatorname{dlog}([\pi]_m)$  and  $C(\operatorname{adlog}([\pi]_m)) = C(a)\operatorname{dlog}([\pi]_m)$  for  $a \in Z_1 W_! \Omega_K^{\bullet}$ , it follows that the map  $W_m \Omega_K^{q-1} \xrightarrow{\operatorname{dlog}([\pi]_m)} W_m \Omega_K^q$  induces maps

$$T_n^{m,q-1}(K) \xrightarrow{\operatorname{dlog}([\pi]_m)} T_n^{m,q}(K), \ \widetilde{T}_n^{m,q-1}(K) \xrightarrow{\operatorname{dlog}([\pi]_m)} \widetilde{T}_n^{m,q}(K).$$

Since  $V(adlog([\pi]_m)) = V(a)dlog([\pi]_m)$ , we get that these maps also induce

$$\widetilde{M}_n^{m,q-1}(K) \xrightarrow{\operatorname{dlog}([\pi]_m)} \widetilde{M}_n^{m,q}(K).$$

Since  $\theta_2^q = \operatorname{dlog}([\pi]_m) \circ \theta_1^{q-1} = \theta_1^q \circ \operatorname{dlog}([\pi]_m)$ , we are done by Lemmas 5.1.11.

Combining the previous three lemmas, we get a homomorphism

$$(\theta_1^q, \theta_2^q) \colon \frac{\Omega_{\mathfrak{f}}^q}{Z_1 \Omega_{\mathfrak{f}}^q} \bigoplus \frac{\Omega_{\mathfrak{f}}^{q-1}}{Z_1 \Omega_{\mathfrak{f}}^{q-1}} \to \widetilde{M}_n^{m,q}(K) \quad \text{for} \quad 1 \le m \le r.$$
(5.1.15)

Lemma 5.1.13. We have the following.

- 1. The map  $\phi^q: \Omega^q_{\mathfrak{f}} \to \widetilde{M}_n^{r+1,q}(K)$  is surjective.
- 2. For  $1 \le m \le r$ , the map  $(\theta_1^q, \theta_2^q)$  in (5.1.15) is surjective.

Proof. The map  $\frac{\operatorname{fil}_n W_m(K)}{\operatorname{fil}_{n-1} W_m(K)} \otimes K_q^M(K) \to \widetilde{T}_n^{m,q}(K)$  is surjective by Corollary 5.1.3 and  $\operatorname{fil}_n W_m(K) = [\pi^{-\alpha}]_m W_m(A)$  by Lemma 2.1.7(9), where  $\alpha = np^{1-m}$  if  $1 \leq m \leq r$  and  $\alpha = l$  if m = r + 1. Using Lemmas 4.3.11 and 4.3.13, it only remains to show that we can replace  $K_q^M(K)$  by  $K_q^M(A)$  on the left hand side of this surjection in order to prove (1). To show this, it is enough to consider the case q = 1 and to show that  $[x\pi^{-l}]_{r+1}\operatorname{dlog}([\pi]_{r+1})$  lies in the image of  $A \otimes A^{\times}$  under  $\phi^1$  if  $x \in A^{\times}$ . But  $[x\pi^{-l}]_{r+1}\operatorname{dlog}([\pi]_{r+1}) = -l^{-1}[x]_{r+1}d([\pi^{-l}]_{r+1})$ . It follows that  $\partial_n^{r+1,1}([x\pi^{-l}]_{r+1}\operatorname{dlog}([\pi]_{r+1})) = \phi^1(l^{-1}x \otimes x)$ .

To prove (2), we can repeat the above steps which reduces finally to showing that  $\partial_n^{m,1}([x\pi^{-\alpha}]_m \operatorname{dlog}([y]_m))$  lies in the image of  $(\theta_1^1, \theta_2^1)$ , where  $y \in K^{\times}$ . To show this, we write  $y = u\pi^{\beta}$  for some  $u \in A^{\times}$  and  $\beta \in \mathbb{Z}$ . This yields  $[x\pi^{-\alpha}]_m \operatorname{dlog}([y]_m) = [x\pi^{-\alpha}]_m \operatorname{dlog}([u]_m) + \beta [x\pi^{-\alpha}]_m \operatorname{dlog}([\pi]_m) = \theta_1^1(x \otimes u) + \theta_2^1(\beta x)$ . This finishes the proof.

**Lemma 5.1.14.** For  $m \ge 1$ , the map  $\overline{\delta}_n^{m,q}: \widetilde{M}_n^{m,q}(K) \to \widetilde{N}_n^{m,q}(K)$  is bijective.

*Proof.* When  $m \ge r+2$ , we are done by Lemma 5.1.9 and [36, Lem 3.6.1] as both groups are zero. Otherwise, we look at the maps

$$\Omega_{\mathfrak{f}}^{q} \xrightarrow{\phi^{q}} \widetilde{M}_{n}^{r+1,q}(K) \xrightarrow{\overline{\delta}_{n}^{r+1,q}} \widetilde{N}_{n}^{r+1,q}(K)$$

and (when  $1 \le m \le r$ )

$$\frac{\Omega_{\mathfrak{f}}^{q}}{Z_{1}\Omega_{\mathfrak{f}}^{q}} \bigoplus \frac{\Omega_{\mathfrak{f}}^{q-1}}{Z_{1}\Omega_{\mathfrak{f}}^{q-1}} \xrightarrow{\theta_{1}^{q}, \theta_{2}^{q}} \widetilde{M}_{n}^{m,q}(K) \xrightarrow{\overline{\delta}_{n}^{m,q}} \widetilde{N}_{n}^{m,q}(K).$$

Kato shows in [36, § 3] (see the last step in proof of Theorem 3.2 on p. 114) that the above two composite maps are bijective. We now apply Lemma 5.1.13 to finish the proof.  $\Box$ 

#### **Lemma 5.1.15.** For $m \ge 1$ , the square (5.1.10) is Cartesian.

Proof. Since  $T_{n-1}^{m,q}(K) \to T_n^{m,q}(K)$  is injective, as we observed following the diagram (5.1.10), it is sufficient to show that the nap  $\overline{\delta}_n^{m,q}: \widetilde{T}_n^{m,q}(K) \to \widetilde{S}_n^{m,q}(K)$  is injective. We shall prove this by induction on m. When m = 1, the map  $\widetilde{T}_n^{m,q}(K) \to \widetilde{M}_n^{m,q}(K)$  is bijective and we are done by Lemma 5.1.14. We now assume  $m \ge 2$  and look at the commutative diagram

It is clear from the definition of the groups  $S_n^{m,q}(K)$  that  $S_n^{m-1,q}(K) \cap S_{n-1}^{m,q}(K) = S_{n-1}^{m-1,q}(K)$ , and this implies that the bottom row of (5.1.16) is exact. The left vertical arrow in this diagram is injective by induction and the right vertical arrow is bijective by Lemma 5.1.14. It follows that the middle vertical arrow is injective.

We can now finally prove the following.

**Theorem 5.1.16.** For  $m \ge 1$  and  $n, q \ge 0$ , the map

$$\overline{\delta}_n^{m,q}: T_n^{m,q}(K) \to {}_{p^m} \operatorname{fil}_n^{\operatorname{bk}} H^{q+1}(K)$$

is an isomorphism, where  $T_n^{m,q}(K) = \mathbb{H}^1_{\acute{e}t}(\operatorname{Spec} A, W_m \mathcal{F}_n^{q,\bullet}).$ 

Proof. By Corollary 5.1.3, we only need to show that  $\overline{\delta}_n^{m,q}$  is surjective. For this, we let  $w \in {}_{p^m} \operatorname{fil}_n^{\operatorname{bk}} H^{q+1}(K) = S_n^{m,q}(K)$ . By Lemma 5.1.1, we see that  $w \in H^1_{\operatorname{\acute{e}t}}(K, W_m \Omega^q_{K, \log})$ . Since  $\delta^{m,q} \colon W_m \Omega^q_K \twoheadrightarrow H^1_{\operatorname{\acute{e}t}}(K, W_m \Omega^q_{K, \log})$  and by Lemma 2.1.7(3) it follows that  $\varinjlim_n T_n^{m,q}(K) = H^1_{\operatorname{\acute{e}t}}(K, W_m \Omega^q_{K, \log})$ . In particular,  $w \in T_{n'}^{m,q}(K) \cap S_n^{m,q}(K)$  for all  $n' \gg 0$ . We conclude from Lemma 5.1.15 that  $w \in T_n^{m,q}(K)$ .

## 5.2 Comparison with Global Kato filtration

Let X and  $W_m \mathcal{F}_D^{q,\bullet}$  be as in Definition 4.3.16. We also recall all the notations from there. Furthermore, assume  $D \ge 0$  (i.e.  $n_i \ge 0$ ). In this section we shall describe the group  $\mathbb{H}^1_{\text{ét}}(X, W_m \mathcal{F}_D^{q,\bullet})$  in terms of local Kato filtration studied in § 5.1. Recall the complex

$$W_m \mathcal{F}_D^{q,\bullet} = \left[ Z_1 \mathrm{fil}_D W_m \Omega_U^q \xrightarrow{1-C} \mathrm{fil}_D W_m \Omega_U^q \right].$$

We note that  $j^* W_m \mathcal{F}_D^{q,\bullet} = W_m \Omega_{U,\log}^q$ .

#### 5.2.1 Global Kato filtration

For  $x_i \in E^{(0)}$ , let  $K_i = Q(\mathcal{O}^h_{X,x_i})$  and  $j_i : H^1_{\text{ét}}(U, W_m \Omega^q_{U,\log}) \to H^1_{\text{ét}}(K_i, W_m \Omega^q_{K_i,\log})$  be the canonical restriction map. Lemma 5.1.1 implies  $H^1_{\text{\acute{e}t}}(K_i, W_m \Omega^q_{K_i,\log}) = {}_{p^m}(H^{q+1}(K_i)).$ 

We define the global Kato filtration of  $H_m^{q+1}(U) = H_{\text{ét}}^1(U, W_m \Omega_{U,\log}^q)$  as follows:

**Definition 5.2.1.** (cf. [43, Defn 2.7]) For any  $q \ge 0, m \ge 1$ , we let

$$\operatorname{fil}_{D}^{\log} H^{1}_{\operatorname{\acute{e}t}}(U, W_{m}\Omega^{q}_{U, \log}) \coloneqq ker\left(H^{1}_{\operatorname{\acute{e}t}}(U, W_{m}\Omega^{q}_{U, \log}) \xrightarrow{\oplus j_{i}} \bigoplus_{x_{i} \in E^{(0)}} \frac{H^{1}_{\operatorname{\acute{e}t}}(K_{i}, W_{m}\Omega^{q}_{K_{i}, \log})}{p^{m}(\operatorname{fil}_{n_{i}}^{\operatorname{bk}} H^{q+1}(K_{i}))}\right)$$

For q = 0, we have

$$\pi_1^{ab}(X,D)/p^m = \operatorname{Hom}_{\mathbf{Ab}}(\operatorname{fil}_D^{\log} H^1_{\operatorname{\acute{e}t}}(U,\mathbb{Z}/p^m), \mathbb{Q}/\mathbb{Z}),$$
(5.2.1)

where  $\pi_1^{ab}(X, D)$  is the (logarithmic) abelianized étale fundamental group with modulus D, defined as  $\operatorname{Hom}_{Ab}(\operatorname{fil}_D^{\log} H^1_{\operatorname{\acute{e}t}}(U, \mathbb{Q}/\mathbb{Z}), \mathbb{Q}/\mathbb{Z})$  (cf. [43, Def 2.7]). Recall from [43, Lem 3.6], that we have an isomorphism

$$\mathbb{H}^{1}_{\text{\acute{e}t}}(X, W_m \mathcal{F}_D^{q, \bullet}) \cong \operatorname{fil}_D^{\log} H^{1}_{\text{\acute{e}t}}(U, W_m \Omega_{U, \log}^q), \text{ if } q = 0.$$

Our goal in this section is to extend the above isomorphism to arbitrary  $q \ge 0$ . This will be one of the key results for proving the duality theorem for Hodge-Witt cohomology with modulus over local fields in § 8 and the Lefschetz theorem for Brauer groups in § 9.

We begin with the following lemma.

**Lemma 5.2.2.** Let k be an F-finite field of characteristic p > 0. Let  $R = k[[Y_1, Y_2]]$ ,  $\pi = Y_1^n Y_2^m, n, m \in \mathbb{Z}$  and  $K = R_{Y_1Y_2}$  be localisation of R by the element  $Y_1Y_2$ . Let's write (cf. (4.1.1))

$$F_0^{1,q} = F_0^{1,q}(R) = \Omega_k^q \oplus \Omega_k^{q-1} \mathrm{dlog} Y_1 \oplus \Omega_k^{q-1} \mathrm{dlog} Y_2 \oplus \Omega_k^{q-2} \mathrm{dlog} Y_1 \mathrm{dlog} Y_2$$

and

$$Z_1 F_0^{1,q} = Z_1 \Omega_k^q \oplus (Z_1 \Omega_k^{q-1}) \operatorname{dlog} Y_1 \oplus (Z_1 \Omega_k^{q-1}) \operatorname{dlog} Y_2 \oplus (Z_1 \Omega_k^{q-2}) \operatorname{dlog} Y_1 \operatorname{dlog} Y_2$$

Then  $Z_1 \operatorname{fil}_{(n,m)} \Omega_K^q$  has the following unique presentation.

$$Z_1 \text{fil}_{(n,m)} \Omega_K^q = \sum_{\substack{ip \ge -n \\ jp \ge -m}} Y_1^{ip} Y_2^{jp} \left( Z_1 F_0^{1,q} \right) + \sum_{\substack{i \ge -n \\ j \ge -m \\ p \nmid i \text{ or } p \nmid j}} d(Y_1^i Y_2^j F_0^{1,q-1}).$$

Moreover, one has

- 1.  $(\operatorname{fil}_{(n,m)}\Omega_K^q)_{Y_1Y_2} = \Omega_K^q$  and  $(Z_1\operatorname{fil}_{(n,m)}\Omega_K^q)_{Y_1Y_2} = Z_1\Omega_K^q$ , where we consider  $Z_1\Omega_K^q$ and  $Z_1\operatorname{fil}_{(n,m)}\Omega_K^q$  as *R*-modules via Frobenius action.
- 2. Every element  $\omega \in (\operatorname{fil}_{(n,m)}\Omega_K^q)_{Y_1}$  can be written uniquely as

$$\omega = \sum_{\substack{i \ge -N_0 \\ j \ge -m}} Y_1^i Y_2^j a_{i,j}, \text{ for some } N_0 > 0 \text{ and } a_{i,j} \in F_0^{1,q}.$$

3. Every element  $\omega \in (\operatorname{fil}_{(n,m)}\Omega_K^q)_{Y_2}$  can be written uniquely as

$$\omega = \sum_{\substack{i \ge -n \\ j \ge -M_0}} Y_1^i Y_2^j b_{i,j}, \text{ for some } M_0 > 0 \text{ and } b_{i,j} \in F_0^{1,q}.$$

#### 5.2. COMPARISON WITH GLOBAL KATO FILTRATION

4. Every element  $\omega \in (Z_1 \operatorname{fil}_{(n,m)} \Omega^q_K)_{Y_1}$  can be written uniquely as

$$\omega = \sum_{\substack{i \ge -N'_0 \\ jp \ge -m}} Y_1^{ip} Y_2^{jp} e_{i,j} + \sum_{\substack{i \ge -N'_0 \\ j \ge -m \\ p + i \text{ or } p + j}} d(Y_1^i Y_2^j f_{i,j}),$$

for some  $N'_0 > 0$  and  $e_{i,j} \in Z_1 F_0^{1,q}, f_{i,j} \in F_0^{1,q-1}$ .

5. Every element  $\omega \in (Z_1 \operatorname{fil}_{(n,m)} \Omega^q_K)_{Y_2}$  can be written uniquely as

$$\omega = \sum_{\substack{ip \ge -n \\ j \ge -M'_0}} Y_1^{ip} Y_2^{jp} e'_{i,j} + \sum_{\substack{i\ge -n \\ j \ge -M'_0 \\ p \neq i \text{ or } p \neq j}} d(Y_1^i Y_2^j f'_{i,j}),$$

for some  $M'_0 > 0$  and  $e'_{i,j} \in Z_1 F_0^{1,q}, f'_{i,j} \in F_0^{1,q-1}.$ 

*Proof.* The first part of the lemma follows from Corollary 4.1.2 and the Cartier isomorphism in Lemma 4.3.3(2) for m = 1. The rest of the part is an easy consequence of the first part and Corollary 4.1.2,

**Lemma 5.2.3.** Let R be as above.  $\mathfrak{m} = (Y_1, Y_2)$ . Let  $n \ge 0, m \ge 0$ . Consider the map

$$1 - C : Z_1 \operatorname{fil}_{(n,m)} \Omega_K^q \to \operatorname{fil}_{(n,m)} \Omega_K^q.$$

Then the induced map

$$(1-C)^*: H^2_{\mathfrak{m}}(R, Z_1 \operatorname{fil}_{(n,m)}\Omega^q_K) \to H^2_{\mathfrak{m}}(R, \operatorname{fil}_{(n,m)}\Omega^q_K)$$

is injective, where  $H^*_{\mathfrak{m}}(R, -)$  is the Zariski (eqv. étale) cohomology with support at the closed point  $V(\mathfrak{m})$  of Spec R.

*Proof.* For any R-module M, note that

$$H^2_{\mathfrak{m}}(R, M) = H^1(\operatorname{Spec} R - \{\mathfrak{m}\}, M).$$

and thus, by using čech cohomology, we have

$$H^1(\operatorname{Spec} R - \{\mathfrak{m}\}, M) = \operatorname{coker}(M_{Y_1} \oplus M_{Y_2} \xrightarrow{(+-)} M_{Y_1Y_2}).$$

Now consider the following commutative diagram.

where f, g, h are the maps induced by 1 - C.

To show  $(1 - C)^*$  is injective, we need to show that  $h(\alpha) \in \text{Image } \theta_2$  implies  $\alpha \in \text{Image } \theta_1$ , for all  $\alpha \in (Z_1 \text{fil}_{(n,m)} \Omega_K^q)_{Y_1 Y_2}$ .

Let  $\alpha \in (Z_1 \operatorname{fil}_{(n,m)} \Omega_K^q)_{Y_1 Y_2} = Z_1 \Omega_K^q$ , such that  $h(\alpha) \in \operatorname{Image} \theta_2$ . So by Lemma 5.2.2, we can write  $h(\alpha) = a - b$ , where

$$a = \sum_{\substack{i \ge -N \\ j \ge -m}} Y_1^i Y_2^j a_{i,j} \text{ and } b = \sum_{\substack{i \ge -n \\ j \ge -M}} Y_1^i Y_2^j b_{i,j}; \quad a_{i,j}, b_{i,j} \in F_0^{1,q}.$$

and  $\alpha$  as

$$\alpha = \sum_{\substack{ip \ge -N \\ jp \ge -M}} Y_1^{ip} Y_2^{jp} c_{i,j} + \sum_{\substack{i\ge -N \\ j\ge -M \\ p \nmid i \text{ or } p \nmid j}} d(Y_1^i Y_2^j d_{i,j}); \ c_{i,j} \in Z_1 F_0^{1,q} \text{ and } d_{i,j} \in F_0^{1,q-1}$$
(5.2.3)

for some N >> 0, M >> 0.

Note that  $d(Y_1^i Y_2^j d_{i,j}) = Y_1^i Y_2^j (id_{i,j} \cdot \operatorname{dlog} Y_1 + jd_{i,j} \cdot \operatorname{dlog} Y_2 + d(d_{i,j}))$ . Hence,  $d(Y_1^i Y_2^j d_{i,j})$ can be written as  $Y_1^i Y_2^j d_{i,j}''$ , where  $d_{i,j}'' = (id_{i,j} \cdot \operatorname{dlog} Y_1 + jd_{i,j} \cdot \operatorname{dlog} Y_2 + d(d_{i,j})) \in F_0^{1,q}$ . So, we can rearrange the sum in (5.2.3) and write it as

$$\alpha = \sum_{\substack{i \ge -n \\ j \ge -M}} Y_1^i Y_2^j c'_{i,j} + \sum_{\substack{j \ge -m \\ i \ge -N}} Y_1^i Y_2^j d'_{i,j} + \sum_{\substack{-N < i < -n \\ -M < j < -m}} Y_1^i Y_2^j e'_{i,j},$$
(5.2.4)

where  $c'_{i,j}, d'_{i,j}, e'_{i,j} \in F_0^{1,q}$  and each term of the above sum lies in  $Z_1\Omega_K^q$ .

Hence, we conclude that  $\sum_{\substack{i\geq -n\\j>-M}} Y_1^i Y_2^j c'_{i,j}$  (resp.  $\sum_{\substack{j\geq -m\\i>-N}} Y_1^i Y_2^j d'_{i,j}$ ) lies in  $(Z_1 \operatorname{fil}_{(n,m)}\Omega_K^q)_{Y_2}$  (resp.  $(Z_1 \operatorname{fil}_{(n,m)}\Omega_K^q)_{Y_1}$ ) (cf. Lemma 5.2.2(4),(5)). It follows that

$$\tau = \sum_{\substack{i \ge -n \\ j > -M}} Y_1^i Y_2^j c'_{i,j} + \sum_{\substack{j \ge -m \\ i > -N}} Y_1^i Y_2^j d'_{i,j} \in \text{Image } \theta_1.$$
(5.2.5)

Let's write  $c = \sum_{\substack{-N < i < -n \\ -M < j < -m}} Y_1^i Y_2^j e'_{i,j}.$ 

Claim: c = 0.

Since  $h(\alpha) \in \text{Image } \theta_2$ , it follows from (5.2.5) and from commutativity of (5.2.2) that  $h(c) = h(\alpha) - h(\tau) \in \text{Image } \theta_2$ . We shall show that this implies c = 0.

Indeed, if  $-N_0$  be the least exponent of  $Y_1$  appearing in c, then write c as

$$c = \sum_{j < -m} Y_1^{-N_0} Y_2^j e'_{-N_0,j} + \sum_{\substack{-N_0 < i < -n \\ j < -m}} Y_1^i Y_2^j e'_{i,j}.$$

Now from the expression of  $\alpha$  in (5.2.3), we note that,  $Y_1^i Y_2^j e'_{i,j}$  is either of the form  $d(Y_1^i Y_2^j \overline{e}_{i,j})$  (if p + i or p + j), or it is of the form  $Y_1^{pi_1} Y_2^{pi_2} e'_{i,j}$  (in case p divides both i, j), where  $\overline{e}_{i,j} \in F_0^{1,q-1}$ . So

$$C(Y_1^i Y_2^j e'_{i,j}) = \begin{cases} 0, & \text{if } p + i \text{ or } p + j \\ Y_1^{i/p} Y_2^{j/p} C(e'_{i,j}), & \text{otherwise.} \end{cases}$$
(5.2.6)

That is,  $C(Y_1^i Y_2^j e'_{i,j})$  is either zero or the exponent of  $Y_1$  in  $C(Y_1^i Y_2^j e'_{i,j})$  is strictly greater than *i* (because i < 0). This implies (1 - C)(c) can be written as

$$(1-C)(c) = \sum_{j < -m} Y_1^{-N_0} Y_2^j e'_{-N_0,j} + \sum_{\substack{i > -N_0 \\ j}} Y_1^i Y_2^j e''_{i,j},$$

where  $e'_{i,j}, e''_{i,j} \in F_0^{1,q}$ . Since  $(1-C)(c) \in \text{Image } \theta_2$ , it implies that every term of the above sum has the property that either the exponent of  $Y_1$  is at least -n or the exponent of  $Y_2$ is at least -m. Hence by the uniqueness of Lemma 5.2.2, we conclude

$$Y_1^{-N_0}Y_2^j e'_{-N_0,j} = 0$$
, for all  $j < -m$ .

Thus,  $-N_0$  is not the least exponent of  $Y_1$  appearing in c, which is a contradiction. Hence we get the claim.

Since the claim is proved, we get  $\alpha \in \text{Image } \theta_1$ .

Recall, for any bounded complex of abelian sheaves  $\mathcal{F}$  on  $X_{\text{\acute{e}t}}$ , we have the coniveau spectral sequence

$$E_1^{p,q} = \bigoplus_{x \in X^{(p)} \cap E} \mathbb{H}_x^{p+q}(X, \mathcal{F}) \implies \mathbb{H}_E^{p+q}(X, \mathcal{F}),$$
(5.2.7)

where  $E^{(p)}$  = set of codimension p points of E.

This spectral sequence will be used to prove the next two lemmas.

#### Lemma 5.2.4. $\mathbb{H}^{1}_{E}(X, W_{m}\mathcal{F}_{D}^{q,\bullet}) = 0.$

*Proof.* Let  $\mathcal{F}_m = W_m \mathcal{F}_D^{q,\bullet}$ , for  $D \ge 0$ .

Claim 1: For  $x \in X^{(1)} \cap E$ , we have  $\mathbb{H}^1_x(X, \mathcal{F}_m) = 0$ .

*Proof*: We let  $A_x = \mathcal{O}_{X,x}^h, K_x = Q(A_x)$  and  $\pi_x$  be a uniformizer of  $A_x$ . Since  $\mathbb{H}^1_x(X, \mathcal{F}_m) = \mathbb{H}^1_x(A_x, \mathcal{F}_m)$ , we have the following localisation exact sequence

$$\dots \to \mathbb{H}^{0}_{\text{\acute{e}t}}(A_{x}, \mathcal{F}_{m}) \xrightarrow{f} \mathbb{H}^{0}_{\text{\acute{e}t}}(K_{x}, \mathcal{F}_{m}) \to \mathbb{H}^{1}_{x}(A_{x}, \mathcal{F}_{m}) \to \mathbb{H}^{1}_{\text{\acute{e}t}}(A_{x}, \mathcal{F}_{m}) \xrightarrow{g} \mathbb{H}^{1}_{\text{\acute{e}t}}(K_{x}, \mathcal{F}_{m}).$$
(5.2.8)

By Lemma 5.1.2(2), we see that f is an isomorphism in (5.2.8). Also g is injective by Proposition 3.2.3, because  $\mathbb{H}^1_{\text{ét}}(A_x, \mathcal{F}_m) = \operatorname{Coker}(1 - C : Z_1 \operatorname{fil}_{D_x} W_m \Omega^q_{K_x} \to \operatorname{fil}_{D_x} W_m \Omega^q_{K_x})$ (by Lemma 5.1.2(2)). Hence we get  $\mathbb{H}^1_x(A_x, \mathcal{F}_m) = 0$ . This proves Claim 1.

Using Claim 1, we get  $E_{\infty}^{1,0} = 0$ . Also note that,  $E_{\infty}^{0,1} = 0$ . Hence the lemma follows.

**Lemma 5.2.5.** Let  $\mathcal{F}_m$  be as in Lemma 5.2.4. Then the canonical map  $\mathbb{H}^2_E(X, \mathcal{F}_m) \to \bigoplus_{x \in X^{(1)} \cap E} \mathbb{H}^2_x(X, \mathcal{F}_m)$  is injective.

*Proof.* We make the following claim.

Claim 2 : Let  $x \in X^{(2)} \cap E$ . Then  $\mathbb{H}^2_x(X, \mathcal{F}_m) = 0$ .

Proof: By Theorem 4.3.17(11), it is enough to prove the claim for m = 1 and for all  $D \ge 0$ . Since  $\mathbb{H}^2_x(X, \mathcal{F}_1) = \mathbb{H}^2_x(R_x, \mathcal{F}_1)$ , where  $R_x = \mathcal{O}^h_{X,x}$ ; by hypercohomology long exact sequence we have

$$0 = H_x^1(R_x, \operatorname{fil}_D\Omega_{U_x}^q) \to \mathbb{H}_x^2(R_x, \mathcal{F}_1) \to H_x^2(R_x, Z_1 \operatorname{fil}_D\Omega_{U_x}^q) \xrightarrow{(1-C)^*} H_x^2(R_x, \operatorname{fil}_D\Omega_{U_x}^q),$$

where  $U_x = \operatorname{Spec} R_x - E_x$ , where  $E_x$  is pull back of E in  $\operatorname{Spec} R_x$ .

Here  $H_x^1(R_x, \operatorname{fil}_D\Omega_{U_x}^q) = 0$  ([7, Prop 3.5.4.b]) because codimension of  $\{x\}$  in X is 2 and  $\operatorname{fil}_D\Omega_{U_x}^q$  is free  $R_x$ -module. So, to prove the claim, it is enough to show that  $(1-C)^*$  is injective.

Let  $\mathfrak{m}$  be the maximal ideal of  $R_x$ ,  $\widehat{R}_x$  be its  $\mathfrak{m}$  adic completion and  $\widehat{U}_x$  = be the inverse image of  $U_x$  to Spec  $\widehat{R}_x$ . Since  $Z_1 \operatorname{fil}_D \Omega^q_{U_x}$  and  $\operatorname{fil}_D \Omega^q_{U_x}$  are coherent sheaves on Spec  $R_x$ , and by Corollary 4.2.7 and Lemmas 4.2.6, 4.3.3 their  $\mathfrak{m}$ -adic completions are  $Z_1 \operatorname{fil}_D \Omega^q_{\widehat{U}_x}$  and  $\operatorname{fil}_D \Omega^q_{\widehat{U}_x}$ , respectively; [7, Prop 3.5.4.(d)] implies  $H^2_x(R_x, M) \cong H^2_x(\widehat{R}_x, \widehat{M})$  for  $M \in \{\operatorname{fil}_D \Omega^q_{U_x}, Z_1 \operatorname{fil}_D \Omega^q_{U_x}\}$ . Moreover, we have a commutative diagram

Hence, it is enough to show the bottom arrow is injective. By Cohen's structure theorem, we can write  $\widehat{R}_x = k[[X, Y]]$ , where k is the residue field of  $R_x$ ; D will be of the form  $V(\pi)$ , where  $\pi = X^n Y^m$ , where  $n \ge 0$  and  $m \ge 0$ . Now the claim follows from Lemma 5.2.3.

Using claim 3, we get  $E_{\infty}^{2,0} = 0$  in (5.2.7). Also,  $E_{\infty}^{0,2} = 0$ . Hence

$$\mathbb{H}^2_E(X,\mathcal{F}_m) = E^{1,1}_{\infty} \hookrightarrow E^{1,1}_1 = \bigoplus_{x \in X^{(1)} \cap E} \mathbb{H}^2_x(X,\mathcal{F}_m).$$

This finishes the proof.

Now we prove the main theorem of this chapter.

**Theorem 5.2.6.** We have a canonical isomporphism

$$\mathbb{H}^{1}_{\acute{e}t}(X, W_m \mathcal{F}^{q, \bullet}_D) \cong \mathrm{fil}^{\log}_D H^{1}_{\acute{e}t}(U, W_m \Omega^q_{U, \log}).$$

*Proof.* We have the following localisation exact sequence

$$\mathbb{H}^{1}_{E}(X, W_{m}\mathcal{F}_{D}^{q, \bullet}) \to \mathbb{H}^{1}_{\text{\acute{e}t}}(X, W_{m}\mathcal{F}_{D}^{q, \bullet}) \to H^{1}_{\text{\acute{e}t}}(U, W_{m}\Omega_{U, \log}^{q}) \to \mathbb{H}^{2}_{E}(X, W_{m}\mathcal{F}_{D}^{q, \bullet}).$$

By Lemma 5.2.4, we have  $\mathbb{H}^1_E(X, W_m \mathcal{F}^{q, \bullet}_D) = 0$ . Now we consider the following commutative diagram of localisation exact sequences

where  $A_x = \mathcal{O}_{X,x}^h, K_x = Q(A_x), x \in E^{(0)}, D_x$  is the pull back of D in Spec  $A_x$ . The vertical arrows are restriction maps induced by the morphism Spec  $A_x \to X$ . Moreover, the map (2) in the bottom row is surjective. This follows because the next term  $\mathbb{H}^2(A_x, W_m \mathcal{F}_{D_x}^{q,\bullet}) =$ 0, by looking at the hypercohomology long exact sequence for  $W_m \mathcal{F}_{D_x}^{q,\bullet}$  and noting that  $H^1_{\text{ét}}(A_x, \text{fil}_{n_x} W_m \Omega_{K_x}^q) = 0 = H^2_{\text{ét}}(A_x, Z_1 \text{fil}_{n_x} W_m \Omega_{K_x}^q)$  (cf. Lemma 5.1.2).

Also, the map (1) in the bottom row of (5.2.10) is injective by Lemmas 5.1.1, 5.1.2(2) and Corollary 5.1.3. Furthermore, we have  $\mathbb{H}^{1}_{\acute{e}t}(A_x, W_m \mathcal{F}^{q, \bullet}_{D_x}) \cong {}_{p^m} \operatorname{fil}_{n_x}^{\mathrm{bk}} H^{q+1}(K_x)$  by Theorem 5.1.7.

As a result, composite of the morphisms  $H^1_{\text{\'et}}(U, W_m \Omega^q_{U, \log}) \to \mathbb{H}^2_E(X, W_m \mathcal{F}^{q, \bullet}_D) \to \mathbb{H}^2_x(A_x, W_m \mathcal{F}^{q, \bullet}_{D_x})$  coincides with the map  $H^1_{\text{\'et}}(U, W_m \Omega^q_{U, \log}) \to \frac{H^1(K_x, W_m \Omega^q_{K_x, \log})}{p^m \operatorname{fil}_{n_x}^{\operatorname{bk}} H^{q+1}(K_x)}$ , induced by canonical restriction. Hence, we have

$$\mathbb{H}^{1}_{\text{\acute{e}t}}(X, W_{m}\mathcal{F}_{D}^{q, \bullet}) \cong \operatorname{Ker}\left(H^{1}_{\text{\acute{e}t}}(U, W_{m}\Omega_{U, \log}^{q}) \to \mathbb{H}^{2}_{E}(X, W_{m}\mathcal{F}_{D}^{q, \bullet})\right) \\
\cong \operatorname{Ker}\left(H^{1}_{\text{\acute{e}t}}(U, W_{m}\Omega_{U, \log}^{q}) \to \bigoplus_{x \in X^{(1)} \cap E} \mathbb{H}^{2}_{x}(X, W_{m}\mathcal{F}_{D}^{q, \bullet})\right) \\
= \operatorname{Ker}\left(H^{1}_{\text{\acute{e}t}}(U, W_{m}\Omega_{U, \log}^{q}) \to \bigoplus_{x \in X^{(1)} \cap E} \frac{H^{1}(K_{x}, W_{m}\Omega_{K_{x}, \log}^{q})}{p^{m}\operatorname{fil}_{n_{x}}^{\operatorname{bk}}H^{q+1}(K_{x})}\right),$$

where, the second isomorphism follows from Lemma 5.2.5. This finishes the proof of the theorem.

79

#### 5.2.2 Relation with Brauer group with modulus

For q = 1, we shall establish a relationship between the group  $\mathbb{H}^1_{\text{ét}}(X, W_m \mathcal{F}_D^{q, \bullet})$  and the group  $\mathrm{Br}^{\mathrm{div}}(X|D)$  defined in [47, Def 8.7].

**Definition 5.2.7.** We let  $\operatorname{Br}^{\operatorname{div}}(X|D)$  denote the subgroup of  $\operatorname{Br}(U)$  consisting of elements  $\chi$  such that for every  $x \in E^{(0)}$ , the image  $\chi_x$  of  $\chi$  under the canonical map  $\operatorname{Br}(U) \to \operatorname{Br}((K_x))$  lies in  $\operatorname{fil}_{n_x} \operatorname{Br}(K_x)$ . We call  $\operatorname{Br}^{\operatorname{div}}(X|\underline{0})$  to be the tame Brauer group of U, where  $\underline{0}$  is the zero divisor.

Remark 5.2.8. Our definition is slightly different from the one defined in [47]. Here we have shifted the filtration index by 1.  $\hfill \Box$ 

We note the following observations.

Lemma 5.2.9. We have

- 1.  $\operatorname{Br}^{\operatorname{div}}(X|\underline{0})$  contains the prime to p-part of  $\operatorname{Br}(U)$ .
- 2.  $\operatorname{Br}^{\operatorname{div}}(X|D) \subset \operatorname{Br}^{\operatorname{div}}(X|D')$  if  $D \leq D'$ ;  $\lim_{|D| \subset E} \operatorname{Br}^{\operatorname{div}}(X|D) = \operatorname{Br}(U)$ .
- 3.  $_{\ell^m} \operatorname{Br}^{\operatorname{div}}(X|D) = _{\ell^m} \operatorname{Br}(U) \text{ for all } D \ge 0, \ m \ge 1 \text{ and any prime } \ell \neq p.$

*Proof.* (1) follows from the fact that any prime to *p*-torsion element of Br(U) maps to  $fil_0Br(K_x)$  for all  $x \in E^{(0)}$  (see [36, Prop 6.1]). (2) easily follows from the definition. (3) follows from (1) and (2).

We conclude the chapter with the following corollary.

Corollary 5.2.10. We have an exact sequence

$$0 \to \operatorname{Pic}(U)/p^m \to \mathbb{H}^1_{\acute{e}t}(X, W_m \mathcal{F}_D^{1, \bullet}) \to {}_{p^m} \operatorname{Br}^{\operatorname{div}}(X|D) \to 0.$$
(5.2.11)

*Proof.* Recall that we have an isomorphism of étale sheaves  $dlog: \mathbb{G}_{m,U}/p^m \to W_m \Omega^1_{U,log}$ on  $U_{\acute{e}t}$ . We consider the following exact sequence of étale sheaves on  $U_{\acute{e}t}$ .

$$0 \to \mathbb{G}_{m,U} \xrightarrow{p^m} \mathbb{G}_{m,U} \xrightarrow{\mathrm{dlog}} W_m \Omega^1_{U,\log} \to 0$$

Taking cohomology and identifying  $H^1_{\text{\acute{e}t}}(U, \mathbb{G}_{m,U})$  (resp.  $H^2_{\text{\acute{e}t}}(U, \mathbb{G}_{m,U})$ ) with Pic(U) (resp. Br(U)), we get the following commutative diagram of exact sequences.

where the vertical maps are induced by canonical restrictions. Now the result follows using snake lemma, Theorem 5.2.6 and noting that  $\frac{p^m \operatorname{Br}(K_x)}{p^m \operatorname{fil}_{n_x} \operatorname{Br}(K_x)} \subset p^m \left(\frac{\operatorname{Br}(K_x)}{\operatorname{fil}_{n_x} \operatorname{Br}(K_x)}\right)$ .

## Chapter 6

# Duality theorem for Hodge-Witt cohomology over finite fields

In this chapter, we shall prove a duality theorem for the cohomology of the logarithmic Hodge-Witt sheaves with modulus over finite fields. We begin by defining the relevant complexes of sheaves, proving some of their properties and constructing their pairings over a more general field before we restrict to finite fields.

### 6.1 Definition of complexes and their properties

Let X be a Noetherian regular F-finite  $\mathbb{F}_p$ -scheme. Let d denote the rank of the locally free sheaf  $\Omega_X^1$ . Let  $E = E_1 + \dots + E_r$  be a simple normal crossing divisor on X and let  $j: U \hookrightarrow X$  be the inclusion of the complement of E. For  $\underline{n} = (n_1, \dots, n_r) \in \mathbb{Z}^r$ , we let  $D_{\underline{n}} = \sum_{i=1}^r n_i [E_i] \in \text{Div}_E(X)$ . We shall write  $\text{fil}_{D_{\underline{n}}} W_m \Omega_U^q$  as  $\text{fil}_{\underline{n}} W_m \Omega_U^q$ . We shall say that  $\underline{n} \ge \underline{n}'$  if  $n_i \ge n'_i$  for each i. We shall have analogous interpretations for the notations  $\underline{n} \le \underline{n}'$  and  $\underline{n} = \underline{n}'$ . We let  $\underline{tn} \coloneqq (tn_1, \dots, tn_r)$ ,  $\underline{t} \coloneqq (t, \dots, t) \in \mathbb{Z}^r$  for  $t \in \mathbb{Z}$ ,  $\underline{n} \pm \underline{n}' \coloneqq (n_1 \pm n'_1, \dots, n_r \pm n'_r)$  and  $\underline{n} \pm t \coloneqq \underline{n} \pm \underline{t}$ . We also let  $\underline{n}/p = (\lfloor n_1/p \rfloor, \dots, \lfloor n_r/p \rfloor)$ . We shall have analogous notations for  $\underline{n}/p^i$ , for  $i \ge 1$ .

We let  $W_m \Omega_{X,\log}^q$  denote the image of the map  $\operatorname{dlog}: \mathcal{K}_{q,X}^M \to W_m \Omega_X^q$  (cf. § 5). Letting  $\mathcal{K}_{q,X}^m = \mathcal{K}_{q,X}^M/p^m$ , this induces an isomorphism of Zariski sheaves  $\operatorname{dlog}: \mathcal{K}_{q,X}^m \xrightarrow{\cong} W_m \Omega_{X,\log}^q$ . For  $\underline{n} \geq \underline{0}$ , we let  $\mathcal{I}_{\underline{n}} = \operatorname{Ker}(\mathcal{O}_X \twoheadrightarrow \mathcal{O}_{D_{\underline{n}}})$  and let  $\mathcal{K}_{q,X|D_{\underline{n}}}^M$  be the étale (or Nisnevich) sheaf on X given by the image of the canonical map  $(1 + \mathcal{I}_{\underline{n}})^{\times} \otimes j_* \mathcal{O}_U^{\times} \otimes \cdots \otimes j_* \mathcal{O}_U^{\times} \to j_* \mathcal{K}_{q,U}^M$ . One knows that  $\mathcal{K}_{q,X|D_{\underline{n}}}^M \subset \mathcal{K}_{q,X}^M$  (cf. [31, Prop. 1.1.3]). We let  $\mathcal{K}_{q,X|D_{\underline{n}}}^m = \mathcal{K}_{q,X|D_{\underline{n}}}^M/(p^m \mathcal{K}_{q,X}^M \cap \mathcal{K}_{q,X|D_{\underline{n}}}^M)$ . Letting  $W_m \Omega_{X|D_{\underline{n}},\log}^q$  be the image of the map

#### CHAPTER 6. DUALITY THEOREM FOR HODGE-WITT COHOMOLOGY OVER 84 FINITE FIELDS

dlog:  $\mathcal{K}_{q,X|D_{\underline{n}}}^{M} \to W_{\underline{m}}\Omega_{X}^{q}$  for  $\underline{n} \geq \underline{1}$ , one gets an isomorphism dlog:  $\mathcal{K}_{q,X|D_{\underline{n}}}^{m} \xrightarrow{\cong} W_{\underline{m}}\Omega_{X|D_{\underline{n}},\log}^{q}$  (cf. [31, Thm. 1.1.5]). One also has the following.

### Lemma 6.1.1. $W_m \Omega^q_{X|D_n,\log} \subset \operatorname{fil}_{\underline{n}} W_m \Omega^q_U$ , for $\underline{n} \geq \underline{1}$ .

Proof. Using the definition of  $W_m \Omega_{X|D_n,\log}^q$  and Lemma 2.1.7(8, 10), it suffices to show that  $\operatorname{dlog}([1 + at]_m) \in \operatorname{fil}_{\underline{n}} W_m \Omega_K^1$  in the notations of § 4.2 provided A is regular local ring,  $t = \prod_{i=1}^r x_i^{n_i}$  with each  $n_i \ge 1$  and  $a \in A$ . To this end, we use [14, Lem. 1.2.3] to write  $[1 + at]_m = (ta_{m-1}, \ldots, ta_0) + [1]_m$ , where  $a_i \in A$ . This implies that  $\operatorname{dlog}([1 + at]_m) = [1 + at]_m^{-1}(d(x) + d([1]_m))$ , where  $x = (ta_{m-1}, \ldots, ta_0)$ . Since  $d([1]_m) \in W_m \Omega_{\mathbb{F}_p}^1 = 0$  and  $x \in \operatorname{fil}_{\underline{n}} W_m(K)$ , we conclude from Lemma 2.1.7(4) that  $\operatorname{dlog}([1 + at]_m) \in \operatorname{fil}_{\underline{n}} W_m \Omega_K^1$ .

We now let  $\underline{n} \ge -1$ ,  $q \ge 0$  and  $m \ge 1$ , and consider the following complexes in  $\mathcal{D}(X_{\text{ét}})$ .

$$W_m \mathcal{F}_{\underline{n}}^{q,\bullet} = \left[ Z_1 \operatorname{fil}_{\underline{n}} W_m \Omega_U^q \xrightarrow{1-C} \operatorname{fil}_{\underline{n}} W_m \Omega_U^q \right]; \tag{6.1.1}$$

$$W_m \mathcal{G}_{\underline{n}}^{q,\bullet} = \left[ \operatorname{fil}_{-\underline{n}-1} W_m \Omega_U^q \xrightarrow{1-\overline{F}} \frac{\operatorname{fil}_{-\underline{n}-1} W_m \Omega_U^q}{dV^{m-1} (\operatorname{fil}_{-\underline{n}-1} \Omega_U^{q-1})} \right]; \tag{6.1.2}$$

$$W_m \mathcal{F}^{q,\bullet} = \left[ Z_1 W_m \Omega_X^q \xrightarrow{1-C} W_m \Omega_X^q \right]; \quad W_m \mathcal{G}^{q,\bullet} = \left[ W_m \Omega_X^q \xrightarrow{1-\overline{F}} \frac{W_m \Omega_X^q}{dV^{m-1}(\Omega_X^{q-1})} \right]; \quad (6.1.3)$$

$$W_m \mathcal{H}^{\bullet} = \left[ W_m \Omega_X^d \xrightarrow{1-C} W_m \Omega_X^d \right] \cong W_m \Omega_{X,\log}^d, \quad (\text{Note } Z_1 W_m \Omega_X^d = W_m \Omega_X^d). \tag{6.1.4}$$

These complexes are defined in view of Theorem 4.3.17. The following lemma explains these complexes as objects of  $\mathcal{D}(X_{\text{ét}})$  for certain values of <u>n</u>.

Lemma 6.1.2. The sequences (cf. Lemma 6.1.1)

- $1. \quad 0 \to W_m \Omega^q_{X|D_{\underline{n}}, \log} \to \mathrm{fil}_{-\underline{n}} W_m \Omega^q_U \xrightarrow{1-\overline{F}} \frac{\mathrm{fil}_{-\underline{n}} W_m \Omega^q_U}{dV^{m-1} (\mathrm{fil}_{-\underline{n}} \Omega^{q-1}_U)};$
- $\mathcal{Q}. \quad 0 \to W_m \Omega^q_{X|D_{\underline{1}}, \log} \to \mathrm{fil}_{-\underline{1}} W_m \Omega^q_U \xrightarrow{1-\overline{F}} \xrightarrow{\mathrm{fil}_{-\underline{1}} W_m \Omega^q_U} \xrightarrow{d_U} \to 0;$
- $3. \quad 0 \to W_m \Omega^q_{X|D_{\underline{1}}, \log} \to Z_1 \operatorname{fil}_{-\underline{1}} W_m \Omega^q_U \xrightarrow{1-C} \operatorname{fil}_{-\underline{1}} W_m \Omega^q_U \to 0;$

$$4. \quad 0 \to j_*(W_m \Omega^q_{U,\log}) \to Z_1 \operatorname{fil}_{\underline{0}} W_m \Omega^q_U \xrightarrow{1-C} \operatorname{fil}_{\underline{0}} W_m \Omega^q_U \to 0;$$

5. 
$$0 \to j_*(W_m \Omega^q_{U,\log}) \to \operatorname{fil}_{\underline{0}} W_m \Omega^q_U \xrightarrow{1-\overline{F}} \frac{\operatorname{fil}_{\underline{0}} W_m \Omega^q_U}{dV^{m-1}(\operatorname{fil}_{\underline{0}} \Omega^{q-1}_U)} \to 0;$$

$$6. \quad 0 \to \Omega^q_{X|D_{\underline{n}},\log} \to Z_1 \operatorname{fil}_{-\underline{n}} \Omega^q_U \xrightarrow{1-C} \operatorname{fil}_{(-\underline{n})/p} \Omega^q_U \to 0;$$

are exact, where  $\underline{n} \ge 1$  in (1) and (6).

*Proof.* It is clear that  $(1 - \overline{F})(W_m \Omega^q_{X|D_{\underline{n}},\log}) = 0$ . Using [11, Lem. 2], (1) will therefore follow if we show that

$$\operatorname{fil}_{-\underline{n}} W_m \Omega_U^q \bigcap W_m \Omega_{X,\log}^q \subset W_m \Omega_{X|D_{\underline{n}},\log}^q \quad \text{for} \quad \underline{n} \ge \underline{0}.$$

$$(6.1.5)$$

We shall prove this inclusion by induction on m. The case m = 1 follows directly from Lemma 2.1.7(2) and [31, Thm. 1.2.1]. We now let  $m \ge 2$  and  $x \in \operatorname{fil}_{-\underline{n}} W_m \Omega_U^q \cap W_m \Omega_{X,\log}^q$ . Lemma 2.1.7(5,7) then implies that

$$R(x) \in \operatorname{fil}_{(-\underline{n})/p} W_{m-1}\Omega_U^q \bigcap W_{m-1}\Omega_{X,\log}^q \text{ and } F(x) \in \operatorname{fil}_{-\underline{n}} W_{m-1}\Omega_U^q \bigcap W_{m-1}\Omega_{X,\log}^q$$

Since we also have R(x) = F(x), it follows that  $R(x) \in \operatorname{fil}_{\underline{n}} W_{m-1} \Omega^q_U \cap W_{m-1} \Omega^q_{X,\log}$ . In particular, the induction hypothesis implies that  $R(x) = F(x) \in W_{m-1} \Omega^q_{X|D_n,\log}$ .

We now look at the commutative diagram (cf. [31, Proof of Thm. 2.3.1])

whose bottom row is exact (cf. [11, Lem. 3]). Note that the map R on the top is surjective by definition of the sheaves  $W_m \Omega_{X|D_n,\log}^q$ . Using this diagram, we can find  $y \in W_m \Omega_{X|D_n,\log}^q$  and  $z \in \Omega_{X,\log}^q$  such that  $\underline{p}^{m-1}(z) = x - y \in \operatorname{fil}_{-\underline{n}} W_m \Omega_U^q$ . We conclude from Lemma 4.3.14 that  $z \in \operatorname{fil}_{-\underline{n}'} \Omega_U^q \cap \Omega_{X,\log}^q$ , where  $\underline{n}' = -(-\underline{n}/p^{m-1})$ . In particular,  $z \in$  $\Omega_{X|D_{n'},\log}^q$  by induction. It follows that  $x - y = \underline{p}^{m-1}(z) \in W_m \Omega_{X|D_p^{m-1}\underline{n}',\log}^q \subset W_m \Omega_{X|D_n,\log}^q$ , and hence,  $x \in W_m \Omega_{X|D_n,\log}^q$ . This proves (1).

To prove (2), we only need to show that  $1 - \overline{F}$  is surjective. By Remark 4.3.12 and Lemma 4.3.13, and the fact that

$$\overline{F}(w \operatorname{dlog}([x_1]_m) \wedge \cdots \wedge \operatorname{dlog}([x_q]_m)) = \overline{F}(w) \operatorname{dlog}([x_1]_m) \wedge \cdots \wedge \operatorname{dlog}([x_q]_m)),$$

it is sufficient to find pre-images of elements of the form  $V^i([x]_{m-i})$  under the map  $\operatorname{fil}_{\underline{1}}W_m(K) \xrightarrow{1-\overline{F}} \operatorname{fil}_{\underline{1}}W_m(K)$ . We shall show that this map is in fact surjective. Using the commutative diagram of exact sequences

$$0 \to \operatorname{fil}_{-\underline{1}} W_1(K)^{V_{m-1}^{m-1}} \operatorname{fil}_{-\underline{1}} W_m(K) \xrightarrow{R} \operatorname{fil}_{-\underline{1}} W_{m-1}(K) \to 0$$

$$1-\overline{F} \downarrow \qquad 1-\overline{F} \downarrow \qquad 1-\overline{F} \downarrow \qquad 0 \to \operatorname{fil}_{-\underline{1}} W_1(K)^{V_{m-1}^{m-1}} \operatorname{fil}_{-\underline{1}} W_m(K) \xrightarrow{R} \operatorname{fil}_{-\underline{1}} W_{m-1}(K) \to 0$$

$$(6.1.7)$$

#### CHAPTER 6. DUALITY THEOREM FOR HODGE-WITT COHOMOLOGY OVER 86 FINITE FIELDS

and induction on m, it suffices to show that  $\operatorname{fil}_{-1}W_1(K) \xrightarrow{1-\overline{F}} \operatorname{fil}_{-1}W_1(K)$  is surjective when the ring A in Lemma 4.3.11 is strict Henselian. But this is elementary (cf. [31, Claim 1.2.2]).

To prove the surjectivity of 1 - C in (3), it suffices to show, using Lemma 4.3.3(2) and the commutative diagram

that  $1 - \overline{F}$  is surjective. The latter claim follows from (2). The remaining part of (3) is easily deduced from (1) and the above diagram.

The proof of (6) is identical using the commutative diagram and [31, Thm 1.2.1]

The proof of the surjectivity of 1 - C (resp.  $1 - \overline{F}$ ) in (4) (resp. (5)) is identical to that in (3) (resp. (2)). Also, one checks using an analogue of (6.1.8) for fil<sub>0</sub> $W_m \Omega_U^q$  that  $\operatorname{Ker}(1 - C) = \operatorname{Ker}(1 - \overline{F})$ . We thus have to only show that  $\operatorname{Ker}(1 - C) = j_*(W_m \Omega_{U,\log}^q)$  to finish the proofs of (4) and (5).

To that end, we look at the commutative diagram of exact sequences

The middle and the right vertical arrows are the canonical inclusion maps. It follows that  $\operatorname{Ker}(1-C) \hookrightarrow j_*(W_m \Omega^q_{U,\log})$ . To prove this inclusion is a bijection, we note that the map dlog:  $j_*(\mathcal{K}^M_{q,U}) \to j_*(W_m \Omega^q_{U,\log})$  has a factorization

Hence, it suffices to show that  $j_*(\mathcal{K}^M_{q,U}) \to j_*(W_m \Omega^q_{U,\log})$  is surjective. Since  $j_*(\mathcal{K}^m_{q,U}) \to j_*(W_m \Omega^q_{U,\log})$  is an isomorphism, we are reduced to showing that the canonical map  $j_*(\mathcal{K}^M_{q,U}) \to j_*(\mathcal{K}^m_{q,U})$  is surjective in  $X_{\text{\acute{e}t}}$ . To prove the latter, it is enough to show that the map  $j_*(\mathcal{K}^M_{q,U}) \to j_*(\mathcal{K}^m_{q,U})$  is surjective in  $X_{\text{zar}}$ .

To prove the surjectivity of  $j_*(\mathcal{K}^M_{q,U}) \to j_*(\mathcal{K}^m_{q,U})$  in  $X_{\text{zar}}$ , we can assume that X = Spec(A) and  $U = X_{\pi}$  as in § 4.2. In this case, the exact sequence

$$0 \to \mathcal{K}^{M}_{q,X_{\pi}} \xrightarrow{p^{m}} \mathcal{K}^{M}_{q,X_{\pi}} \to \mathcal{K}^{m}_{q,X_{\pi}} \to 0$$
(6.1.12)

reduces the problem to showing that  $H^1_{\text{zar}}(X_{\pi}, \mathcal{K}^M_{q, X_{\pi}}) = 0$ , which is Lemma 4.3.15.  $\Box$ 

**Corollary 6.1.3.** (cf. [31, Thm. 1.1.6]) For  $\underline{n} \ge 1$  and  $1 \le r \le m - 1$ , one has an exact sequence

$$0 \to W_r \Omega^q_{X|D_{[\underline{n}/p^{m-r}]},\log} \xrightarrow{\underline{p}^{m-r}} W_m \Omega^q_{X|D_{\underline{n}},\log} \xrightarrow{R^r} W_{m-r} \Omega^q_{X|D_{\underline{n}},\log} \to 0.$$

Proof. The surjectivity of  $R^r$  follows directly from the definition of  $W_m \Omega_{X|D_n,\log}^q$ . The remaining part of the corollary follows from its classical non-modulus version (cf. [11, Lem. 3]) and the identity  $\operatorname{fil}_{-\underline{n}} W_m \Omega_U^q \cap W_m \Omega_{X,\log}^q = W_m \Omega_{X|D_n,\log}^q$  that we showed in the proof of Lemma 6.1.2.

**Lemma 6.1.4.** One has  $W_m \mathcal{F}_0^{0,\bullet} \cong \mathbb{Z}/p^m$  and  $W_m \mathcal{G}_0^{d,\bullet} \cong W_m \Omega^d_{X,\log}$ 

*Proof.* The first isomorphism is clear from the definition. For the second isomorphism, it is enough to show (by Lemma 6.1.2(2)) that the canonical map  $W_m \Omega^d_{X|D_{\underline{1}},\log} \to W_m \Omega^d_{X,\log}$  is an isomorphism. But this follows from [45, Prop 2.2.5], because the proof works in our situation.

## 6.2 Pairings of complexes

We continue with the set up of § 6.1 and define the pairings between the complexes. We consider the first two pairings

$$\langle , \rangle_0^{0,0} : Z_1 \operatorname{fil}_{\underline{n}} W_m \Omega_U^q \times \operatorname{fil}_{-\underline{n}^{-1}} W_m \Omega_U^{d-q} \to \operatorname{fil}_{-1} W_m \Omega_U^d \subset W_m \Omega_X^d$$
(6.2.1)

and

$$\langle , \rangle_1^{1,0}: \operatorname{fil}_{\underline{n}} W_m \Omega_U^q \times \operatorname{fil}_{-\underline{n}^{-1}} W_m \Omega_U^{d-q} \to \operatorname{fil}_{-1} W_m \Omega_U^d \subset W_m \Omega_X^d \tag{6.2.2}$$

by letting  $\langle w_1, w_2 \rangle_0^{0,0} \coloneqq w_1 \wedge w_2$  and  $\langle w_1, w_2 \rangle_1^{1,0} \coloneqq w_1 \wedge w_2$ .

#### CHAPTER 6. DUALITY THEOREM FOR HODGE-WITT COHOMOLOGY OVER 88 FINITE FIELDS

We define the third pairing

$$\langle , \rangle_1^{0,1}: Z_1 \operatorname{fil}_{\underline{n}} W_m \Omega_U^q \times \frac{\operatorname{fil}_{-\underline{n}-1} W_m \Omega_U^{d-q}}{dV^{m-1} (\operatorname{fil}_{-\underline{n}-1} \Omega_U^{d-q-1})} \to \operatorname{fil}_{-1} W_m \Omega_U^d \subset W_m \Omega_X^d$$
(6.2.3)

by letting  $\langle w_1, w_2 \rangle_1^{0,1} := -C(w_1 \wedge w_2).$ 

Note that the first two pairings are defined by Lemma 2.1.7, and to show that the third pairing is also defined, we only need to prove that  $C(w_1 \wedge w_2) = 0$  if  $w_1 \in Z_1 \operatorname{fil}_{\underline{n}} W_m \Omega_U^q$  and  $w_2 \in dV^{m-1}(\operatorname{fil}_{\underline{n}-1} \Omega_U^{d-q-1})$ . To prove it, we can write  $w_1 = F(w_1')$  (cf. Theorem 4.3.17(3)) and  $w_2 = dV^{m-1}(w_2')$ , where  $w_1' \in \operatorname{fil}_{\underline{n}} W_{m+1} \Omega_U^q$  and  $w_2' \in \operatorname{fil}_{\underline{n}-1} \Omega_U^{d-q-1}$ . Then

$$C(w_1 \wedge w_2) = C(F(w_1') \wedge dV^{m-1}(w_2')) = CF(w_1' \wedge dV^m(w_2')) = R(w_1' \wedge dV^m(w_2'))$$
$$= R(w_1') \wedge dRV^m(w_2') = 0,$$

because  $RV^m = 0$ .

**Lemma 6.2.1.** For  $\underline{n} \ge \underline{0}$ , the above pairings of étale sheaves induce a pairing of the complexes

$$W_m \mathcal{F}^{q,\bullet}_{\underline{n}} \times W_m \mathcal{G}^{d-q,\bullet}_{\underline{n}} \xrightarrow{\langle , \rangle} W_m \mathcal{H}^{\bullet}$$
 (6.2.4)

such that one has a commutative diagram of pairings

in which the vertical arrows are the canonical maps.

*Proof.* To prove that (6.2.4) is defined, it is enough to show (cf.  $[55, \S 1, p. 175]$ ) that

$$(1-C)(w_1 \wedge w_2) = -C(w_1 \wedge (1-\overline{F})(w_2)) + (1-C)(w_1) \wedge w_2$$

for all  $w_1 \in Z_1 \operatorname{fil}_{\underline{n}} W_m \Omega_U^q$  and  $w_2 \in \operatorname{fil}_{\underline{-n}-1} W_m \Omega_U^{d-q}$ . But this can be easily checked using [55, Lem. 1.1(c)] and the identity  $C\overline{F} = \operatorname{id}$  (cf. [22, Lem. 7.1]). The commutativity of (6.2.5) is clear.

Using Lemma 6.1.2(1), we get the following.

**Corollary 6.2.2.** Lemma 6.2.1 remains valid if one replaces  $W_m \mathcal{G}_{\underline{n}}^{q,\bullet}$  by  $W_m \Omega_{X|D_{n+1},\log}^q$ .

Using this corollary and considering the hypercohomology, we get a commutative diagram of the cup product pairings of abelian groups (for  $\underline{n} \ge \underline{0}$ )

## 6.3 The duality theorem over finite fields

We now specialize to the situation when X is projective as well as geometrically connected over k, where k is a finite field. Other assumptions remain the same as in § 6.1. Then dim  $X = \operatorname{rank} \Omega_X^1 = d$ . We fix  $\underline{n} \geq \underline{0}$ . By [57, Cor. 1.12], there is a bijective trace map  $\operatorname{Tr}: H^{d+1}_{\text{ét}}(X, W_m \Omega^d_{X, \log}) \xrightarrow{\cong} \mathbb{Z}/p^m$ . Composing the pairings in (6.2.6) with this trace map, we get a pairing

$$\mathbb{H}^{i}_{\text{\acute{e}t}}(X, W_m \mathcal{F}^{q, \bullet}_{\underline{n}}) \times H^{d+1-i}_{\text{\acute{e}t}}(X, W_m \Omega^{d-q}_{X|D_{\underline{n}+1}, \log}) \xrightarrow{\cup} \mathbb{Z}/p^m$$
(6.3.1)

which is compatible with the pairing for the cohomology of the sheaves  $W_m \Omega^{\bullet}_{X,\log}$  obtained in [57, Thm. 1.14]. Our goal is to show that this is a perfect pairing of finite abelian groups. We begin with the following.

## **Lemma 6.3.1.** The groups $\mathbb{H}^{i}_{\acute{e}t}(X, W_m \mathcal{F}^{q, \bullet}_{\underline{n}})$ and $H^{i}_{\acute{e}t}(X, W_m \Omega^{q}_{X|D_n, \log})$ are all finite.

Proof. To prove the finiteness of  $\mathbb{H}^{i}_{\acute{e}t}(X, W_m \mathcal{F}^{q,\bullet}_{\underline{n}})$ , it suffices to show that the cohomology of  $Z_1 \operatorname{fll}_{\underline{n}} W_m \Omega^q_U$  and  $\operatorname{fll}_{\underline{n}} W_m \Omega^q_U$  are finite. But Lemma 2.1.7(1) says that these are sheaves of coherent  $W_m \mathcal{O}_X$ -modules. In particular, their Zariski (equivalently, étale) cohomology are finite type  $W_m(k)$ -modules. The desired claim now follows because  $W_m(k)$  is finite. The finiteness of  $H^i_{\acute{e}t}(X, W_1 \Omega^q_{X|D_{\underline{n}}, \log})$  follows by considering the long exact cohomology sequence associated to the sheaf exact sequence of [31, Thm. 1.2.1]. The desired finiteness for m > 1 now follows by induction using Corollary 6.1.3.

For a  $\mathbb{Z}/p^m$ -module M, we let  $M^* = \operatorname{Hom}_{\mathbb{Z}/p^m}(M, \mathbb{Z}/p^m)$ . Recall that a pairing of finite  $\mathbb{Z}/p^m$ -modules  $M \times N \to \mathbb{Z}/p^m$  is called perfect if the induced map  $M \to N^*$  is an isomorphism. Equivalently, the map  $N \to M^*$  is an isomorphism.

In the following lemma, we don't assume that k is finite and X is projective, because it will be used in other chapters.

**Lemma 6.3.2.** Let X be as in Section 6.1. Then the  $\mathcal{O}_X$  modules  $Z_1 \operatorname{fil}_{\underline{n}} \Omega_U^q$  and  $\frac{\operatorname{fil}_{\underline{n}} \Omega_U^{q+1}}{d(\operatorname{fil}_{\underline{n}} \Omega_U^q)}$  (via Frobenius) are locally free of finite rank, for all  $\underline{n} \in \mathbb{Z}^r$ .

#### CHAPTER 6. DUALITY THEOREM FOR HODGE-WITT COHOMOLOGY OVER 90 FINITE FIELDS

Proof. The sheaf of  $\mathcal{O}_X$ -modules  $\operatorname{fil}_{\underline{n}}\Omega_U^q$  is locally free of finite rank by Lemmas 2.1.7(2) and 1.2.8. To see that the same is true for  $Z_1\operatorname{fil}_{\underline{n}}\Omega_U^q$  and  $\frac{\operatorname{fil}_{\underline{n}}\Omega_U^{q+1}}{d(\operatorname{fil}_{\underline{n}}\Omega_U^q)}$ , recall first that if A is the local ring of a closed point on  $E \subset X$  with maximal ideal  $(x_1, \ldots, x_r, x_{r+1}, \ldots, x_N)$ , then  $\Omega_A^1(\log \pi)$  is a free A-module with basis  $\{\operatorname{dlog}(x_1), \ldots, \operatorname{dlog}(x_r), dx_{r+1}, \ldots, dx_N, dy_1, \ldots, dy_s\}$ , for some  $y_1, \ldots, y_s \in A$  (cf, Lemma 1.2.8). Also,  $\{x_1, \ldots, x_N, y_1, \ldots, y_s\}$ forms a differential basis (see Lemma 1.2.7) and hence a p-basis (see [66, § II, § III, Thm. 1] for the definition of p-basis and its relation with differential basis) of A over  $\mathbb{F}_p$ .

Using the above facts about the local rings of X, it is easy to check that locally (depending on  $\underline{n}$ ) we can find two finite dimensional  $\mathbb{F}_p$ -vector spaces  $V_1, V_2$  and an  $\mathbb{F}_p$ linear map  $d_0: V_1 \to V_2$  such that  $d: \Omega_X^q(\log E)(D_{\underline{n}}) \to \Omega_X^{q+1}(\log E)(D_{\underline{n}})$  is locally of the form  $(V_1 \xrightarrow{d_0} V_2) \otimes_{\mathbb{F}_p} \mathcal{O}_X^p$  (cf. [55, Lem. 1.7]). In particular, Ker(d) and Coker(d) are locally free  $\mathcal{O}_X^p$ -modules of finite rank. The ring isomorphism  $\mathcal{O}_X \xrightarrow{p} \mathcal{O}_X^p$  now implies that they are locally free  $\mathcal{O}_X$ -modules of finite rank. We can now apply Lemma 2.1.7(2) to conclude that  $Z_1 \operatorname{fil}_{\underline{n}} \Omega_U^q$  and  $\frac{\operatorname{fil}_{\underline{n}} \Omega_U^{q+1}}{d(\operatorname{fil}_{\underline{n}} \Omega_U^q)}$  are finite rank locally free sheaves of  $\mathcal{O}_X$ -modules.

**Lemma 6.3.3.** (6.3.1) is a perfect pairing of finite groups when m = 1.

*Proof.* Now, we let  $\mathcal{E}_{\underline{n}}^{q} = \frac{\operatorname{fil}_{\underline{n}}\Omega_{U}^{q}}{d(\operatorname{fil}_{\underline{n}}\Omega_{U}^{q-1})}$  and q' = d - q. To prove the lemma, we note that  $\Omega_{X|\underline{D}_{\underline{n}},\log}^{q} \cong W_{1}\mathcal{G}_{\underline{n}}^{q,\bullet}$  in  $\mathcal{D}(X_{\operatorname{\acute{e}t}})$  for every  $q, \underline{n} \ge 0$  by [31, Thm. 1.2.1]. Using the definition of the above pairings, we thus get a commutative diagram of long exact sequences

Since (6.2.2) and (6.2.3) are perfect pairings of locally free sheaves (cf. Step 2 in the proof of [31, Thm. 4.1.4]), it follows from the Grothnedick-Serre duality for the structure map  $X \to \text{Spec}(\mathbb{F}_p)$  that  $\alpha_i$  and  $\gamma_i$  in (6.3.2) are bijective maps between finite groups for all *i*. It follows that  $\beta_i$  is also an isomorphism of finite groups for all *i*. This concludes the proof.

The main result of this section is the following.

**Theorem 6.3.4.** Assume that X is a smooth projective and geometrically connected scheme of dimension d over a finite field k of characteristic p. Then (6.3.1) is a perfect pairing of finite abelian groups for all  $q, \underline{n} \ge \underline{0}$  and  $m \ge 1$ .

*Proof.* The finiteness part is already shown in Lemma 6.3.1. To prove the perfectness of the pairing, we can assume m > 1 by Lemma 6.3.3. We let i' = d + 1 - i and q' = d - q. By Theorem 4.3.17(11) and Corollary 6.1.3, we get a commutative diagram (cf. [21, Lem. 4.4] and its proof for the commutativity)

In this diagram, we have used the shorthand  $\mathbb{H}_{\acute{e}t}^*(\mathcal{E}^{\bullet})$  for the (hyper)cohomology  $\mathbb{H}_{\acute{e}t}^*(X, \mathcal{E}^{\bullet})$ and the vertical arrows are induced by the pairing (6.3.1). Since  $\lceil (\underline{n}+1)/p \rceil = \lfloor \underline{n}/p \rfloor + 1$ , it follows by induction on m that  $\alpha_i$  and  $\gamma_i$  are bijective for all i. We conclude that  $\beta_i$ is bijective for all i.

*Remark* 6.3.5. By the above theorem, we get the following isomorphism of finite groups

$$H^{d+1-i}_{\text{\acute{e}t}}(X, W_m \Omega^{d-q}_{X|D_{\underline{n}+1}, \log}) \xrightarrow{\cong} \operatorname{Hom}_{\mathbb{Z}/p^m}(\mathbb{H}^i_{\text{\acute{e}t}}(X, W_m \mathcal{F}^{q, \bullet}_{\underline{n}}), \mathbb{Z}/p^m)$$

Taking limit over <u>n</u> (note that we have canonical maps  $W_m \Omega_{X|D_{\underline{n'}+1},\log}^{d-q} \to W_m \Omega_{X|D_{\underline{n}+1},\log}^{d-q}$ and  $W_m \mathcal{F}_{\underline{n}}^{q,\bullet} \to W_m \mathcal{F}_{\underline{n'}}^{q,\bullet}$  if  $\underline{n'} \ge \underline{n}$ .), we have the following isomorphism of abelian groups.

$$\underbrace{\lim_{\underline{n}} H^{d+1-i}_{\text{ét}}(X, W_m \Omega^{d-q}_{X|D_{\underline{n}+1}, \log})}_{\mathbb{Z}/p^m} \xrightarrow{\cong} \operatorname{Hom}_{\mathbb{Z}/p^m}(\underbrace{\lim_{\underline{n}} \mathbb{H}^i_{\text{ét}}(X, W_m \mathcal{F}^{q, \bullet}_{\underline{n}}), \mathbb{Z}/p^m}) \\ \cong \operatorname{Hom}_{\mathbb{Z}/p^m}(H^i_{\text{ét}}(U, W_m \Omega^q_{U, \log}), \mathbb{Z}/p^m),$$
(6.3.4)

where the last isomorphism follows from the fact that  $\varinjlim_{\underline{n}} W_m \mathcal{F}_{\underline{n}}^{q,\bullet} \cong Rj_* W_m \Omega_{U,\log}^q$ . We can endow the left hand side of (6.3.4) with profinite topology and  $H^i_{\text{\acute{e}t}}(U, W_m \Omega_{U,\log}^q)$  with discrete topology (cf. [31, P. 1325]). Hence, (6.3.4) gives an isomorphism of profinite groups. On the other hand, Theorem 6.3.4 also shows that the map

$$\langle \alpha, - \rangle : \lim_{\stackrel{\frown}{\underline{n}}} H^{d+1-i}_{\text{\acute{e}t}}(X, W_m \Omega^{d-q}_{X|D_{\underline{n}+1}, \log}) \to \mathbb{Z}/p^m$$

is continuous, for  $\alpha \in H^i_{\text{ét}}(U, W_m \Omega^q_{U, \log})$  (cf. [31, Lem 4.1.3]). Combining all these claims, we can conclude the following perfect pairing of topological abelian groups (cf. [31, Thm. 4.1.4]).

$$H^{i}_{\text{\acute{e}t}}(U, W_{m}\Omega^{q}_{U, \log}) \times \varprojlim_{\underline{n}} H^{d+1-i}_{\text{\acute{e}t}}(X, W_{m}\Omega^{d-q}_{X|D_{\underline{n}+1}, \log}) \xrightarrow{\cup} \mathbb{Z}/p^{m}.$$

### CHAPTER 6. DUALITY THEOREM FOR HODGE-WITT COHOMOLOGY OVER 92 FINITE FIELDS

**Corollary 6.3.6.** For X as in Theorem 6.3.4, there is a perfect pairing of finite abelian groups

$$H^{i}_{\acute{e}t}(X, j_{*}(W_{m}\Omega^{q}_{U, \log})) \times H^{d+1-i}_{\acute{e}t}(X, W_{m}\Omega^{d-q}_{X|E, \log}) \xrightarrow{\cup} \mathbb{Z}/p^{m}.$$

*Proof.* Combine Lemma 6.1.2 and Theorem 6.3.4.

## Chapter 7

## A duality theorem for 2-local fields

In this chapter, we shall prove a duality theorem for 2-local fields using the class field theory developed in [35] and [37]. This will be used in § 8 later.

## 7.1 Kato topology

Since we shall use Kato's topology extensively in the next chapter, we briefly recall it here. More details can be found in  $[34, \S 7]$  (see also  $[47, \S 7.1]$ ).

Let K be a 2-local field of characteristic p > 0, i.e., a complete discrete valuation field of characteristic p > 0 whose residue field  $\mathfrak{f}$  is a local field. Recall that, a local field is a complete discrete valued field with finite residue field. Let A denote the ring of integers of K with maximal ideal  $(\pi)$ . One knows that there is a canonical isomorphism of rings  $\phi:\mathfrak{f}[[\pi]] \stackrel{\cong}{\to} A$  such that the composition  $\mathfrak{f} \to \mathfrak{f}[[\pi]] \stackrel{\phi}{\to} A \to \mathfrak{f}$  is the identity map (cf. [53, Thm 28.3, also the proof of Lem. 1]). The Kato topology on A is the unique topology such that  $\phi$  is an isomorphism of topological rings when  $\mathfrak{f}[[\pi]]$  is endowed with the product of the valuation (also called the adic) topology of  $\mathfrak{f}$ . We remark that Kato topology on A does not depends on the choice of the uniformizer  $\pi$  and  $\phi$ . The Kato topology of  $A^{\times}$  is the subspace topology induced from that of A. The Kato topology of  $K^{\times}$  is the unique topology which is compatible with the group structure of  $K^{\times}$  and for which  $A^{\times}$  is open in  $K^{\times}$ . The Kato topology of  $K_2^M(K)$  is the finest topology which is compatible with the group structures and for which the product map  $K^{\times} \times K^{\times} \to K_2^M(K)$ is continuous. This makes  $K_2^M(K)$  into a topological abelian group.

Remark 7.1.1. The canonical topological isomorphism  $\phi: \mathfrak{f}[[\pi]] \xrightarrow{\cong} A$  implies  $A^* \xrightarrow{\cong} \mathfrak{f}^* \oplus (1 + \pi.\mathfrak{f}[[\pi]])$  is also a topological isomorphism, where  $\mathfrak{f}^* \oplus (1 + \pi.\mathfrak{f}[[\pi]])$  is given direct sum topology.

Now we fix some notations.

Notation 7.1.2. If A is a Henselian discrete valuation ring with maximal ideal  $\mathfrak{m}$  and field of fraction K, we shall write  $K_n^M(A|\mathfrak{m}^r) \coloneqq H_{nis}^0(\operatorname{Spec} A, \mathcal{K}_{n,\operatorname{Spec} A|V(\mathfrak{m}^r)}^M)$  (see § 6.1) as  $\operatorname{fil}_r K_n^M(K)$ ) for  $n, r \ge 1$  and  $\operatorname{fil}_0 K_n^M(K) = K_n^M(A)$ , where  $K_n^M$  is the *n*th Milnor Ktheory  $K_n^M$ . Let  $m, q \ge 1$  and  $n \ge 0$  be three integers. If char A = p, we let  $\kappa_q^m(A|n) = \frac{\operatorname{fil}_n K_q^M(K)}{p^m K_q^M(K) \cap \operatorname{fil}_n K_q^M(K)}$  and for  $n \ge 1$ , define  $\varpi_q^m(A|n)$  so that there is an exact sequence

$$0 \to \kappa_q^m(A|n) \to \kappa_q^m(A|1) \to \varpi_q^m(A|n) \to 0.$$
(7.1.1)

It also follows that

$$\varpi_q^m(A|n) = \operatorname{Ker}\left(\frac{K_q^M(K)}{\operatorname{fil}_n K_q^M(K) + p^m K_q^M(K)} \xrightarrow{\theta} \frac{K_q^M(K)}{\operatorname{fil}_1 K_q^M(K) + p^m K_q^M(K)}\right),$$
(7.1.2)

where  $\theta$  is the natural quotient map.

For any integers  $m, r \ge 0$ , the Kato topology of  $K_2^M(K)$  induces the quotient topology on  $K_2^M(K)/m$  and the subspace topology on fil<sub>r</sub> $K_2^M(K)$ . We shall use the term Kato topology for these topologies as well. By the Kato topologies of  $\kappa_r^m(A|n)$  (resp.  $\varpi_r^m(A|n)$ ), we shall mean the subspace topology of  $\kappa_r^m(A|n)$  induced from that of  $K_2^M(K)/p^m$  (resp. the quotient topology of  $\kappa_r^m(A|1)$  via (7.1.1)). Let  $\partial_K: K_2^M(K) \twoheadrightarrow \mathfrak{f}^{\times}$ be the Tame symbol map. Via this surjective homomorphism, the Kato topology of  $K_2^M(K)$  induces the quotient topology (called the Kato topology) on  $\mathfrak{f}^{\times}$ . On the other hand, the latter is also equipped with its valuation topology.

#### **Lemma 7.1.3.** The two topologies of $\mathfrak{f}^{\times}$ coincide.

Proof. To show that the adic topology of  $\mathfrak{f}^{\times}$  is finer than the Kato topology, it suffices to show that if  $U \subset \mathfrak{f}^{\times}$  is a subgroup which is open in the Kato topology, then it contains an open subgroup in the adic topology. To that end, we look at the composite map  $T: K^{\times} \times K^{\times} \to K_2^M(K) \xrightarrow{\partial_K} \mathfrak{f}^{\times}$ . By definition, T is continuous with respect to the Kato topology of  $\mathfrak{f}^{\times}$ . It follows that the map  $T_{\pi}: K^{\times} \to \mathfrak{f}^{\times}$ , given by  $T_{\pi}(a) = \partial_K(\{a, \pi\})$  is a continuous homomorphism. In particular,  $T_{\pi}^{-1}(U)$  contains an open subgroup of  $K^{\times}$ . By definition of the Kato topology of  $K^{\times}$  and [34, § 7, Rem. 1],  $T_{\pi}^{-1}(U)$  contains a subgroup of the form  $V_0 \times V_1$ , where  $V_0 \subset \mathfrak{f}^{\times}$  is an open subgroup in the adic topology and  $V_1 \subset \prod_{i\geq 1} \mathfrak{f}^{\pi i}$ is an open subgroup in the Kato topology of  $\pi A \subset A$ . It is clear from the definition of Tthat  $T_{\pi}(V_0 \times V_1) = V_0$  which implies that  $V_0 \subset U$ .

To show that the Kato topology of  $\mathfrak{f}^{\times}$  is finer than the adic topology, it is equivalent to show that  $\partial_K$  is continuous with respect to the Kato topology on  $K_2^M(K)$  and adic topology on  $\mathfrak{f}^{\times}$ . But this is equivalent to show that the map  $T: K^* \times K^* \to \mathfrak{f}^*$  is continuous with respect to the Kato topology on  $K^* \times K^*$  and adic topology on  $\mathfrak{f}^{\times}$ . Let  $x \in \mathfrak{f}^*$  and  $xU_n$  be a basic open set containing x. Here,  $U_n = 1 + \pi_1^n \mathcal{O}_{\mathfrak{f}}$  and  $\pi_1$  is a uniformizer of  $\mathcal{O}_{\mathfrak{f}}$ . Also, let  $a, b \in K^*$  such that T(a, b) = x. We need to find two Kato open subsets A and B in  $K^*$  containing a and b, respectively, such that  $T(A \times B) \subset xU_n$ . We write  $a = \pi^{n_a}a_1$  and  $b = \pi^{n_b}b_1$ , such that  $a_1, b_1 \in A^*$  and  $n_a, n_b \in \mathbb{Z}$ . Note,  $T(a, b) = \overline{a_1}^{n_b} \cdot \overline{b_1}^{n_a} = x$ , where  $\overline{a}_1, \overline{b}_1$  are the images of  $a_1$  and  $b_1$ , respectively, in  $\mathfrak{f}^*$ . Since the map  $\tau : \mathfrak{f}^* \times \mathfrak{f}^* \to \mathfrak{f}^*$ , given by  $\tau(c, d) = c^{n_b} \cdot d^{-n_a}$  is continuous homomorphism of topological groups (with respect to adic topology), there exists n' such that  $\tau(\overline{a}_1 U_{n'} \times \overline{b}_1 U_{n'}) \subset xU_n$ . Then, for  $A = a.U_{n'}(1 + \pi.\mathfrak{f}[[\pi]])$  and  $B = b.U_{n'}(1 + \pi.\mathfrak{f}[[\pi]])$ , we have  $T(A \times B) = \tau(\overline{a}_1 U_{n'} \times \overline{b}_1 U_{n'}) \subset xU_n$ . Since A and B are Kato open in  $K^*$  (cf. Remark 7.1.1), we conclude that  $\partial_K$  is continuous.

In the sequel, the default topology of  $K_2^M(K)/n$   $(n \ge 0)$  and its subspaces (e.g.,  $K_2^M(A)/n$ ) will be the Kato topology, and the same for  $\mathfrak{f}^*/n$  will be the adic topology. If K' is a Henselian discrete valuation field with completion K, then the Kato topology of  $K'^*$  is the subspace topology induced from that of  $K^*$ . The Kato topology on  $K_2^M(K')$  is the finest topology such that the product map  $K'^* \times K'^* \to K_2^M(K')$  is continuous. It is clear that the canonical map  $K_2^M(K') \to K_2^M(K)$  is continuous.

Notation 7.1.4. Before we proceed further, we recall some topological notions. For  $G \in \mathbf{Ab}$ , we let  $G^{\vee} = \operatorname{Hom}_{\mathbf{Ab}}(G, \mathbb{Q}/\mathbb{Z})$ . For  $G \in \mathbf{Tab}$  (see page VIII for the notations  $\mathbf{Ab}, \mathbf{Tab}$ ), we let  $G^* = \operatorname{Hom}_{\mathbf{Tab}}(G, \mathbb{Q}/\mathbb{Z})$ . If G is either a profinite or a torsion locally compact Hausdorff topological abelian group, then  $G^*$  is a topological abelian group with its compact open topology and satisfies the Pontryagin duality, i.e., the evaluation map  $G \to (G^*)^*$  is an isomorphism in  $\mathbf{Tab}$  (cf. [68, § 2.9]). For  $A, B \in \mathbf{Tab}$ , we let  $\operatorname{Hom}_{\mathrm{cf}}(A, B) = (\operatorname{Hom}_{\mathbf{Tab}}(A, B))_{\mathrm{tor}}$ . We shall write  $\operatorname{Hom}_{\mathrm{cf}}(A, \mathbb{Q}/\mathbb{Z})$  as  $A^*_{\mathrm{tor}}$ . In this thesis, we shall endow  $\mathbb{Q}$  as well as the direct sums of its subgroups and subquotients (e.g., finite abelian groups) with the discrete topology.

**Lemma 7.1.5.** The map  $\partial_K : K_2^M(K) \twoheadrightarrow \mathfrak{f}^{\times}$  induces an exact sequence

$$0 \to (\mathfrak{f}^{\times})^{\star} \to K_2^M(K)^{\star} \to K_2^M(A)^{\star} \to 0.$$
(7.1.3)

Proof. Since  $K_2^M(A) = \text{Ker}(\partial_K)$ , the exactness of (7.1.3) except at  $K_2^M(A)^*$  follows immediately from Lemma 7.1.3. To prove the exactness at  $K_2^M(A)^*$ , we let  $\phi_K \colon K_2^M(K) \to K_2^M(A)$  be given by  $\phi_K(\{a,b\}) = \{a,b\} - \{\partial_K(\{a,b\}),\pi\}$ . Since  $\partial_K$ , the map  $\psi_K \colon \mathfrak{f}^* \to K_2^M(K)$ , where  $\psi_K(a) = \{a,\pi\}$  and the identity map on  $K_2^M(K)$  are group homomorphism, we see that  $\phi_K$  is a group homomorphism whose restriction to  $K_2^M(A)$  is identity.

It remains to show that  $\phi_K$  is continuous with respect to the Kato topology. Equivalently, we need to show that  $\phi_K: K_2^M(K) \to K_2^M(K)$  is continuous. As  $\partial_K$  is continuous, the latter claim follows if we show that the map  $\psi_K: \mathfrak{f}^{\times} \to K_2^M(K)$  is continuous. But this map is the composition of  $\mathfrak{f}^{\times} \hookrightarrow K^{\times} \to K_2^M(K)$ , where the first map is the canonical inclusion and the second map takes a to  $\{a, \pi\}$ . The second map is clearly continuous and the first map is continuous by [34, § 7, Rem. 1].

Remark 7.1.6. The same proof as that of Lemma 7.1.5 shows that the sequence

$$0 \to (\mathfrak{f}^*/n)^* \to (K_2^M(K)/n)^* \to (K_2^M(A)/n)^* \to 0$$
(7.1.4)

is exact for every  $n \in \mathbb{Z}$ . Also, by Remark 7.1.1, we get  $(K^{\times}/n)^{\star} \twoheadrightarrow (A^{\times}/n)^{\star} \twoheadrightarrow$  $(\operatorname{fil}_1 K^{\times}/n)^{\star}$  for every  $n \in \mathbb{Z}$ .

## 7.2 The reciprocity homomorphism

Let K be a Henselian discrete valuation field of characteristic p > 0 whose residue field  $\mathfrak{f}$  is a local field. Let A denote the ring of integers of K and  $\mathfrak{m} = (\pi)$  the maximal ideal of A. For any site  $\tau$  on Spec A (resp. Spec K), the cohomology groups  $H^i_{\tau}(\operatorname{Spec} A, \mathcal{F})$ (resp.  $H^i_{\tau}(\operatorname{Spec} K, \mathcal{F})$ ) will be written as  $H^i_{\tau}(A, \mathcal{F})$  (resp.  $H^i_{\tau}(K, \mathcal{F})$ ).

For any  $r \ge 1$ , we consider  $H^r(K)$  (see § 5 for the definition) as a discrete topological group. For  $q \le 2$ , recall from [34] that the reciprocity homomorphism

$$\rho_K^q: K_q^M(K) \to \operatorname{Hom}_{\operatorname{\mathbf{Ab}}}(H^{3-q}(K), \mathbb{Q}/\mathbb{Z})$$

is induced by the cup product pairing

$$K_q^M(K) \times H^{3-q}(K) \xrightarrow{\mathrm{NR}_K \times \mathrm{id}} H_{\mathrm{\acute{e}t}}^q(K, \mathbb{Q}/\mathbb{Z}(q)) \times H^{3-q}(K) \xrightarrow{\cup} H^3(K) \cong \mathbb{Q}/\mathbb{Z}.$$
(7.2.1)

One knows that  $\rho_K^q$  is a continuous homomorphism with respect to the Kato topology of  $K_q^M(K)$  and the profinite topology of  $(H^{3-q}(K))^*$ . Indeed, the continuity in the case of q = 2 follows from [34, Thm. 1(iii) and Thm. 2(1)], while the continuity in the case of q = 1 follows from Thm. 2(2) of loc. cit.

Notation 7.2.1. Let's write  $\operatorname{fil}_n H^q(K)$  for  $\operatorname{fil}_n^{bk} H^q(K)$  defined in Definition 5.1.4. Also, we write  $\operatorname{fil}_n H^q(K, \mathbb{Z}/p^m(q-1))$  or  $\operatorname{fil}_n H^1_{\operatorname{\acute{e}t}}(K, W_m \Omega_{K,\log}^{q-1}))$  for  $T_n^{m,q}(K)$  defined in (5.1.5). Letting  $G_K^{q,n} = \left(\frac{H^q(K)}{\operatorname{fil}_{n-1}H^q(K)}\right)^*$  and  $G_{K,m}^{q,n} = \left(\frac{H^q(K,\mathbb{Z}/p^m(q-1))}{\operatorname{fil}_{n-1}H^q(K,\mathbb{Z}/p^m(q-1))}\right)^*$ , we have the following.

**Lemma 7.2.2.** For  $n \ge 1, 1 \le q \le 2$  and  $m \ge 1$ , (7.2.1) induces continuous homomorphisms (cf. Notations 7.1.2, 7.2.1)

(1) 
$$\rho_K^{q,n}$$
: fil<sub>n</sub> $K_q^M(K) \to G_K^{3-q,n}$ ; (2)  $\rho_{K,m}^{q,n}$ :  $\kappa_2^m(A|n) \to G_{K,m}^{3-q,n}$ .

*Proof.* The existence of  $\rho_{K,m}^{q,n}$  follows from that of  $\rho_K^{q,n}$  using Theorem 5.1.16. Item (1) follows from (7.2.1) and [36, Rem. 6.6].

## 7.3 Duality for $\kappa_q^m(A|n)$ and $\varpi_q^m(A|n)$

**Lemma 7.3.1.** Let A be as in § 7.2 and  $n \ge 1, m \ge 1$ ,  $q \ge 1$  are integers, the following hold.

- 1. The canonical maps  $K_q^M(K)/p^m \to K_q^M(\widehat{K})/p^m$  and  $\kappa_q^m(A|n) \to \kappa_q^m(\widehat{A}|n)$  are injective, where  $\widehat{A}$  (resp.  $\widehat{K}$ ) is the  $\mathfrak{m}$ -adic completion of A (resp. K).
- 2. The canonical maps  $\frac{K_q^M(K)}{\operatorname{fil}_n K_q^M(K)} \to \frac{K_q^M(\widehat{K})}{\operatorname{fil}_n K_q^M(\widehat{K})}$  and  $\varpi_q^m(A|n) \to \varpi_q^m(\widehat{A}|n)$  are bijective.
- 3.  $H^q_{\acute{e}t}(R, \mathbb{Z}/p^m(q-1)) \to H^q_{\acute{e}t}(\widehat{R}, \mathbb{Z}/p^m(q-1))$  is bijective if  $R \in \{A, K\}$ .
- 4.  $\operatorname{fil}_n H^q(K, \mathbb{Z}/p^m(q-1)) \to \operatorname{fil}_n H^q(\widehat{K}, \mathbb{Z}/p^m(q-1))$  is bijective.

Proof. The first part of item (1) follows because  $W_m \Omega_K^2 \hookrightarrow W_m \Omega_{\widehat{K}}^2$ . For the second part, we note that  $\kappa_q^m(A|n) \in K_q^M(K)/p^m$  (and similarly in the case of  $\widehat{A}$ ). The first part of item (2) and the (R = K) case of item (3) follow from [38, Lem. 1] and [37, Lem. 21]. For the second part of item (2), we note that the result is true for  $\frac{K_q^M(K)}{\operatorname{fil}_n K_q^M(K) + p^m K_q^M(K)}$ (follows from the first part of (2)). Then the claim for  $\varpi_q^m(A|n)$  follows from the previous observation and (7.1.2). To prove (R = A) case of item (3) first consider the following commutative diagram obtained by the localisation sequence for the sheaf  $W_m \Omega_{(-),\log}^{q-1}$  in  $\mathcal{D}(\operatorname{Spec} B_{\text{\acute{e}t}}, \mathbb{Z}/p^m)$ , for B = A and  $\widehat{A}$ .

where x, x' are the closed points of Spec A and Spec  $\widehat{A}$ , respectively. Using (R = K) case, it is enough to show the natural map  $\gamma : H^2_x(A, W_m \Omega^{q-1}_{(-),\log}) \to H^2_{x'}(\widehat{A}, W_m \Omega^{q-1}_{(-),\log})$  is isomorphism. Since  $W_m \Omega^{q-1}_{(-),\log} \cong \left[ Z_1 W_m \Omega^{q-1}_{(-)} \xrightarrow{1-C} W_m \Omega^{q-1}_{(-)} \right]$ , we have the following

commutative diagram (letting q' = q - 1).

$$H^{1}_{x}(Z_{1}W_{m}\Omega_{A}^{q'}) \rightarrow H^{1}_{x}(W_{m}\Omega_{A}^{q'}) \rightarrow H^{2}_{x}(W_{m}\Omega_{A,\log}^{q'}) \rightarrow H^{2}_{x}(Z_{1}W_{m}\Omega_{A}^{q'}) \rightarrow H^{2}_{x}(W_{m}\Omega_{A}^{q'})$$

$$\downarrow^{\alpha_{1}} \qquad \qquad \downarrow^{\beta_{1}} \qquad \qquad \downarrow^{\gamma} \qquad \qquad \downarrow^{\alpha_{2}} \qquad \qquad \downarrow^{\beta_{2}}$$

$$H^{1}_{x'}(Z_{1}W_{m}\Omega_{\widehat{A}}^{q'}) \rightarrow H^{1}_{x'}(W_{m}\Omega_{\widehat{A}}^{q'}) \rightarrow H^{2}_{x'}(W_{m}\Omega_{\widehat{A},\log}^{q'}) \rightarrow H^{2}_{x'}(Z_{1}W_{m}\Omega_{\widehat{A}}^{q'}) \rightarrow H^{2}_{x'}(W_{m}\Omega_{\widehat{A}}^{q'}),$$

$$(7.3.2)$$

where  $H_x^1(W_m\Omega_A^{q'})$  (and similarly the other groups) is the short hand notation for  $H_x^1(A, W_m\Omega_A^{q'})$ . First, we note that the local cohomologies of  $W_m\Omega_A^{q'}$  and  $Z_1W_m\Omega_A^{q'}$  on Spec A is same as those on Spec  $W_m(A)$ . We recall that the  $\mathfrak{M}_m$ -adic (equivalently  $W_m(\mathfrak{m})$ -adic) completion of  $W_m(A)$  is  $W_m(\widehat{A})$  (cf. Lemma 4.2.2, for q = 0). Since  $Z_1W_m\Omega_A^{q'}, W_m\Omega_A^{q'}$  can be regarded as finitely generated  $W_m(A)$ -modules, and their  $\mathfrak{M}_m$ -adic completions are  $Z_1W_m\Omega_{\widehat{A}}^{q'}, W_m\Omega_{\widehat{A}}^{q'}$  (by Corollary 4.2.7, Lemmas 4.2.6, 4.3.3), respectively, we get that  $\alpha_i$  and  $\beta_i$  (i = 1, 2) are isomorphisms by [7, Prop 3.5.4.(d)]. Thus we get  $\gamma$  is isomorphism.

Proof of (4) is identical to that in (3), (R = A) case, except that one need to use  $W_m \mathcal{F}_n^{q-1,\bullet}$  complex instead of  $W_m \Omega_{(-),\log}^{q-1}$  and also Corollary 8.2.5 to obtain a diagram like (7.3.1).

**Lemma 7.3.2.** Let A and K be as in § 7.2. For  $m \ge 1, 1 \le q \le 2$  and  $n \ge 1$ , the map  $\rho_{K,m}^{q,n}$  of Lemma 7.2.2 induces a monomorphism (cf. Notation 7.1.2)

$$\widetilde{\rho}_{K,m}^{q,n}: \varpi_q^m(A|n) \hookrightarrow \left(\frac{\operatorname{fil}_{n-1}H^1_{\acute{e}t}(K, W_m\Omega_{K,\log}^{2-q})}{\operatorname{fil}_0H^1_{\acute{e}t}(K, W_m\Omega_{K,\log}^{2-q})}\right)^{\star},$$

where the dual on the right is taken with respect to the discrete topology.

*Proof.* First note that the map  $\tilde{\rho}_{K,m}^{q,n}$  is defined by the following commutative diagram.

$$0 \longrightarrow \kappa_q^m(A|n) \longrightarrow \kappa_q^m(A|1) \longrightarrow \varpi_q^m(A|n) \longrightarrow 0$$

$$\downarrow^{\rho_{K,m}^{q,n}} \qquad \qquad \downarrow^{\rho_{K,m}^{q,1}} \qquad \qquad \downarrow^{\rho_{K,m}^{q,n}}$$

$$0 \rightarrow \left(\frac{H_{\acute{e}t}^1(K, W_m \Omega_{K,\log}^{2-q})}{\operatorname{fil}_{n-1}H_{\acute{e}t}^1(K, W_m \Omega_{K,\log}^{2-q})}\right)^* \rightarrow \left(\frac{H_{\acute{e}t}^1(K, W_m \Omega_{K,\log}^{2-q})}{\operatorname{fil}_{0}H_{\acute{e}t}^1(K, W_m \Omega_{K,\log}^{2-q})}\right)^* \rightarrow \left(\frac{\operatorname{fil}_{n-1}H_{\acute{e}t}^1(K, W_m \Omega_{K,\log}^{2-q})}{\operatorname{fil}_{0}H_{\acute{e}t}^1(K, W_m \Omega_{K,\log}^{2-q})}\right)^* \rightarrow 0.$$

Now we need to show that the map  $\tilde{\rho}_{K,m}^{q,n}$  is injective. We can assume that K is complete in view of Lemma 7.3.1. We now let  $n' = \lfloor n/p \rfloor$  and claim that the following is a

### 7.3. DUALITY FOR $\kappa_q^m(A|n)$ AND $\varpi_q^m(A|n)$

99

commutative diagram of exact sequences.

The exactness of the top row follows by comparing the exact sequence of Corollary 6.1.3 (taking r = m - 1) for  $\kappa_q^m(A|n)$  and  $\kappa_q^m(A|1)$ .

Now by Theorem 4.3.17(11), we have an exact sequence

$$\cdots \to \mathbb{H}^{0}_{\mathrm{\acute{e}t}}(W_{1}\mathcal{F}^{2-q,\bullet}_{\lfloor n/p \rfloor}) \to \mathbb{H}^{1}_{\mathrm{\acute{e}t}}(W_{m-1}\mathcal{F}^{2-q,\bullet}_{n}) \xrightarrow{V} \mathbb{H}^{1}_{\mathrm{\acute{e}t}}(W_{m}\mathcal{F}^{2-q,\bullet}_{n}) \xrightarrow{R^{m-1}} \mathbb{H}^{1}_{\mathrm{\acute{e}t}}(W_{1}\mathcal{F}^{2-q,\bullet}_{\lfloor n/p \rfloor}) \to \cdots,$$

where we use the short hand notation  $\mathbb{H}^{i}_{\text{ét}}(\mathcal{E})$  for  $\mathbb{H}^{i}(A, \mathcal{E})$ . By Lemma 5.1.2(2), we have the short exact sequence

$$0 \to \mathbb{H}^{0}_{\text{\acute{e}t}}(W_{m-1}\mathcal{F}_{n}^{2-q,\bullet}) \xrightarrow{V=\overline{p}} \mathbb{H}^{0}_{\text{\acute{e}t}}(W_{m}\mathcal{F}_{n}^{2-q,\bullet}) \xrightarrow{R^{m-1}} \mathbb{H}^{0}_{\text{\acute{e}t}}(W_{1}\mathcal{F}_{\lfloor n/p \rfloor}^{2-q,\bullet}) \to 0.$$

and hence the following short exact sequence.

$$0 \to \mathbb{H}^{1}_{\text{\acute{e}t}}(W_{m-1}\mathcal{F}_{n}^{2-q,\bullet}) \xrightarrow{V} \mathbb{H}^{1}_{\text{\acute{e}t}}(W_{m}\mathcal{F}_{n}^{2-q,\bullet}) \xrightarrow{R^{m-1}} \mathbb{H}^{1}_{\text{\acute{e}t}}(W_{1}\mathcal{F}_{\lfloor n/p \rfloor}^{2-q,\bullet}) \to 0.$$

Comparing the above exact sequence with the similar one with n = 0, we get the following exact sequence.

$$0 \to \frac{\mathbb{H}^{1}_{\text{\acute{e}t}}(W_{m-1}\mathcal{F}_{n}^{2-q,\bullet})}{\mathbb{H}^{1}_{\text{\acute{e}t}}(W_{m-1}\mathcal{F}_{0}^{2-q,\bullet})} \xrightarrow{V} \frac{\mathbb{H}^{1}_{\text{\acute{e}t}}(W_{m}\mathcal{F}_{n}^{2-q,\bullet})}{\mathbb{H}^{1}_{\text{\acute{e}t}}(W_{m}\mathcal{F}_{0}^{2-q,\bullet})} \xrightarrow{R^{m-1}} \frac{\mathbb{H}^{1}_{\text{\acute{e}t}}(W_{1}\mathcal{F}_{\lfloor n/p \rfloor}^{2-q,\bullet})}{\mathbb{H}^{1}_{\text{\acute{e}t}}(W_{1}\mathcal{F}_{0}^{2-q,\bullet})} \to 0.$$

Here, we have also used the fact that the map  $\mathbb{H}^1_{\text{\acute{e}t}}(W_m \mathcal{F}_0^{2-q,\bullet}) \to \mathbb{H}^1_{\text{\acute{e}t}}(W_m \mathcal{F}_n^{2-q,\bullet})$  is injective, which follows from Corollary 5.1.3 and Lemma 5.1.2(2). Now, to show the exactness of the bottom row in (7.3.3), we take dual in the above exact sequence, use Theorem 5.1.16 and observe that  $n'-1 = \lfloor (n-1)/p \rfloor$ . Thus, the commutativity of (7.3.3) follows from the commutativity of (6.3.3). It follows from (7.3.3) and an induction argument that it suffices to show  $\tilde{\rho}_{K_1}^{q,n}$  is injective.

argument that it suffices to show  $\widetilde{\rho}_{K,1}^{q,n}$  is injective. For this, note that  $\varpi_q^1(A|n)$  and  $\left(\frac{\operatorname{fil}_{n-1}H_{\operatorname{\acute{e}t}}^1(K,\Omega_{K,\log}^{2-q})}{\operatorname{fil}_0H_{\operatorname{\acute{e}t}}^1(K,\Omega_{K,\log}^{2-q})}\right)$  have filtrations given by

$$\left\{\frac{\kappa_q^1(A|i)}{\kappa_q^1(A|n)}\right\}_{i\leq n} \text{ and } \left\{\left(\frac{\operatorname{fil}_i H^1_{\operatorname{\acute{e}t}}(K, \Omega_{K, \log}^{2-q})}{\operatorname{fil}_0 H^1_{\operatorname{\acute{e}t}}(K, \Omega_{K, \log}^{2-q})}\right)\right\}_{i\leq n-1}, \text{ respectively.}$$

It suffices to show that for all  $i \leq n-1$ , the following map induced by  $\rho_{K,1}^{q,n}$ 

$$\frac{\kappa_q^1(A|i)}{\kappa_q^1(A|i+1)} \to \left(\frac{\operatorname{fil}_i H^1_{\operatorname{\acute{e}t}}(K, \Omega_{K, \log}^{2-q})}{\operatorname{fil}_{i-1} H^1_{\operatorname{\acute{e}t}}(K, \Omega_{K, \log}^{2-q})}\right)$$

is injective. But this follows from [37, Lem 22.2 and 22.3] by identifying the above map with the map induced by the following pairings which don't have any left and right kernels.

$$\Omega_{\mathfrak{f}}^{1} \times \mathfrak{f} \to \Omega_{\mathfrak{f}}^{1} \xrightarrow{\mathrm{Tr \ o \ res}} \mathbb{F}_{p}, \quad \text{if } p \neq i; \qquad \qquad \mathfrak{f}/\mathfrak{f}^{p} \times \mathfrak{f}/\mathfrak{f}^{p} \to \Omega_{\mathfrak{f}}^{1} \xrightarrow{\mathrm{Tr \ o \ res}} \mathbb{F}_{p}, \quad \text{if } p|i$$
$$(v,w) \mapsto \mathrm{Tr}(\mathrm{res}(v.w)); \qquad \qquad (v,w) \mapsto \mathrm{Tr}(\mathrm{res}(dv.w)).$$

We conclude this chapter with a stronger claim about the dual of  $\tilde{\rho}_{K,m}^{q,n}$ , which we can easily deduce from what Kato already showed about  $(\rho_K^q)^*$ .

**Lemma 7.3.3.** For  $m \ge 1$ ,  $n \ge 0$  and q = 1, 2, the map

$$(\widehat{\rho}_{K,m}^{q,n})^{\star} : \frac{\operatorname{fil}_{n-1}H^{1}_{\acute{e}t}(K, W_{m}\Omega_{K,\log}^{2-q})}{\operatorname{fil}_{0}H^{1}_{\acute{e}t}(K, W_{m}\Omega_{K,\log}^{2-q})} \to (\varpi_{q}^{m}(A|n))^{\star}$$

is an isomorphism, where the dual on the right is taken with respect to the Kato topology. *Proof.* We can assume that K is complete. We consider the commutative diagram

where the vertical arrows are the duals of the reciprocity maps. The bottom row is exact by Remark 7.1.6. The left vertical arrow is bijective by [35, § 3, Thm. 3] and Lemma 5.1.1. It follows that the right vertical arrow is surjective. It is also injective by [36, Rem. 6.6], noting that  $\operatorname{fil}_0 H^1_{\operatorname{\acute{e}t}}(K, W_m \Omega^{2-q}_{K, \log}) = \operatorname{fil}_0^{bk} H^{3-q}(K) \cap H^1_{\operatorname{\acute{e}t}}(K, W_m \Omega^{2-q}_{K, \log})$ , which is Theorem 5.1.16. In particular,  $(\widetilde{\rho}^n_{K,m})^*$  is injective.

To prove that  $(\widetilde{\rho}_{K,m}^n)^*$  is surjective, let  $\chi \in (\varpi_2^m(A|n))^*$ . The surjectivity of  $(\rho_{K,m}^{q,1})^*$ implies that there exists an element  $\chi' \in \frac{H^1_{\text{\acute{e}t}}(K, W_m \Omega_{K,\log}^{2-q})}{\operatorname{fil}_0 H^1_{\text{\acute{e}t}}(K, W_m \Omega_{K,\log}^{2-q})}$  such that  $(\rho_{K,m}^{q,1})^*(\chi') = \chi$ . We only have to show that  $\chi' \in \frac{\operatorname{fil}_{n-1}H^1_{\text{\acute{e}t}}(K, W_m \Omega_{K,\log}^{2-q})}{\operatorname{fil}_0 H^1_{\text{\acute{e}t}}(K, W_m \Omega_{K,\log}^{2-q})}$ . But this again follows from [36, Rem. 6.6] and Theorem 5.1.16, where the latter one asserts  $\operatorname{fil}_n H^1_{\text{\acute{e}t}}(K, W_m \Omega_{K,\log}^{2-q}) = \operatorname{fil}_n^{bk} H^{3-q}(K) \cap$  $H^1_{\text{\acute{e}t}}(K, W_m \Omega_{K,\log}^{2-q})$ .

## Chapter 8

# Duality theorem for Hodge-Witt cohomology over local fields

In this chapter, we shall prove a duality theorem for the cohomology of logarithmic Hodge-Witt sheaves with modulus for curves over local fields. Due to the lack of duality results for logarithmic Hodge-Witt cohomology (without modulus) for varieties over local fields, except in dimension one (cf. [40, Prop 4], [47, Thm 4.7]), we focus solely on the case of curves. Let X be a smooth projective geometrically connected curve over a local field k of characteristic p > 0. We let  $E = \{x_1, \ldots, x_r\}, j: U \hookrightarrow X, D_n$  be as in § 6.1. Also, let  $A_x := \mathcal{O}_{X,x}^h, K_x := Q(A_x)$  and  $X_x^h = \operatorname{Spec} A_x$  for any  $x \in X_{(0)}$ .

Since the cohomology group  $H^i_{\text{ét}}(X, W_m \Omega^q_{X|D_n, \log})$  is not finite in general (unlike the case in § 6.3), we choose a different approach to prove the duality theorem in this case. We begin by recalling some pairings of complexes.

### 8.1 Pairing of complexes

By Lemma 6.2.1 and Corollary 6.2.2, we have a commutative diagram (for  $\underline{n} \ge \underline{0}$ )

For any  $x \in X_{(0)}$ , let  $\tau_x^h$ : Spec  $(k(x)) \hookrightarrow X_x^h$  and  $\tau_x$ : Spec  $(k(x)) \hookrightarrow X$  be the inclusion maps, and let  $j_x: X_x^h \to X$  be the canonical étale map. Recall (cf. [74, § 3.1]) that any pairing of complexes of sheaves  $\mathcal{A} \otimes \mathcal{B} \to \mathcal{C}$  on  $X_{\text{ét}}$  gives rise to a pairing  $\tau_x^* \mathcal{A} \otimes^L \tau_x^! \mathcal{B} \to$ 

#### CHAPTER 8. DUALITY THEOREM FOR HODGE-WITT COHOMOLOGY OVER 102 LOCAL FIELDS

 $\tau_x^! \mathcal{C}$  in  $\mathcal{D}(k(x)_{\text{ét}})$ . Applying this to (8.1.1), we get a commutative diagram of pairings

in  $\mathcal{D}(k(x)_{\text{ét}})$  which is compatible with (8.1.1). We shall write  $W_m \mathcal{F}_{\underline{n}}^{q,\bullet}$  as  $W_m \mathcal{F}_{\underline{n}}^{q}$  in the sequel.

## 8.2 Relation with local duality

For any  $\underline{n} \ge 1, q \ge 0$  and any smooth scheme X (need not be a curve) over a field, we let  $W_m \mathcal{W}_{\underline{n}}^q$  be defined by the following short exact sequence

$$0 \to W_m \Omega^q_{X|D_{\underline{n}},\log} \to W_m \Omega^q_{X|D_{\underline{1}},\log} \to W_m \mathcal{W}^q_{\underline{n}} \to 0, \qquad (8.2.1)$$

of Nisnevich (resp. étale) sheaves on X. We recall the following result without proof, as it will be used in various places throughout the thesis.

**Proposition 8.2.1.** Let  $X, D_{\underline{n}}$  be as above. For  $\nu \in \{1, \ldots, r\}$ , we let  $\delta_{\nu} = (0, \ldots, 1, \ldots, 0) \in \mathbb{N}^r$ , where 1 is in the  $\nu$ -th place. For  $\underline{n} \ge \underline{1}, q \ge 0$  and  $m \ge 1$ , we define

$$\operatorname{gr}^{\underline{n},\nu}W_m\Omega^q_{X\log} \coloneqq W_m\Omega^q_{X|D_{\underline{n}},\log}/W_m\Omega^q_{X|D_{\underline{n}+\nu},\log},$$

as Nisnevich (resp. étale) sheaf on X. Then we have

- 1.  $\operatorname{gr}^{\underline{n},\nu}W_m\Omega^q_{X_{\log}}$  is a coherent  $\mathcal{O}^{p^e}_{D_{\nu}}$ -module, for some  $e \gg 0$ .
- 2.  $W_m W_{\underline{n}}^q$  has a filtration whose successive quotients are of the form  $\operatorname{gr}_{\underline{n},\nu} W_m \Omega_{X\log}^q$ .

*Proof.* We may assume X is local. In this case, Item (1) is same as [45, Prop. 2.2.5] (see also [31, Prop. 1.1.9]). Item (2) follows from (8.2.1).

From now on, we restrict ourselves to the case of curves over local fields, assuming that X is as defined at the beginning of this chapter. In this case, we can write

$$W_m \mathcal{W}_{\underline{n}}^q = \bigoplus_{x \in E} (\tau_x)_* \circ (\tau_x)^* (W_m \mathcal{W}_{\underline{n}}^q).$$

In particular, the proper base change theorem for the torsion étale sheaves implies that

$$H^{i}_{\text{\acute{e}t}}(X, W_{m}\mathcal{W}^{q}_{\underline{n}}) = \bigoplus_{x \in E} H^{i}_{\text{\acute{e}t}}(X^{h}_{x}, W_{m}\mathcal{W}^{q}_{\underline{n}}).$$

$$(8.2.2)$$

#### 8.2. RELATION WITH LOCAL DUALITY

For  $n \ge 0$ , we let

$$W_m \mathcal{V}_{\underline{n}}^q = \left[ \frac{Z_1 \operatorname{fil}_{D_{\underline{n}}} W_m \Omega_U^q}{Z_1 W_m \Omega_X^q (\log E)} \xrightarrow{1-C} \frac{\operatorname{fil}_{D_{\underline{n}}} W_m \Omega_U^q}{W_m \Omega_X^q (\log E)} \right]$$

so that we have a term-wise short exact sequence of complexes on  $X_{\text{nis}}$  (resp.  $X_{\text{\acute{e}t}}$ ):

$$0 \to W_m \mathcal{F}^q_{\underline{0}} \to W_m \mathcal{F}^q_{\underline{n}} \to W_m \mathcal{V}^q_{\underline{n}} \to 0.$$
(8.2.3)

By [47, Prop. 4.4] (see also [40, § 3, Prop. 4]), there is a bijective trace homomorphism  $\operatorname{Tr}_X: H^2_{\operatorname{\acute{e}t}}(X, W_m \Omega^2_{X, \log}) \xrightarrow{\cong} \mathbb{Z}/p^m$ . By definition of the pairing between complexes of sheaves (cf. [55, § 1, p. 175]) the pairing on the top row of (8.1.1) is equivalent to a morphism of complexes  $W_m \Omega^{2-q}_{X|D_1, \log} \to \mathcal{H}om'(W_m \mathcal{F}^q_0, W_m \mathcal{H})$ . Composing this with the canonical map  $\mathcal{H}om'(W_m \mathcal{F}^q_0, W_m \mathcal{H}) \to \mathbf{R}\mathcal{H}om'(W_m \mathcal{F}^q_0, W_m \mathcal{H})$ , we get a morphism  $\theta: W_m \Omega^{2-q}_{X|D_1, \log} \to \mathbf{R}\mathcal{H}om'(W_m \mathcal{F}^q_0, W_m \mathcal{H})$  in  $\mathcal{D}(X_{\operatorname{\acute{e}t}})$ . Similarly, the bottom row of (8.1.1) gives rise to a morphism in the derived category  $\theta_{\underline{n}}: W_m \Omega^{2-q}_{X|D_{\underline{n}+1}, \log} \to \mathbf{R}\mathcal{H}om'(W_m \mathcal{F}^q_{\underline{n}}, W_m \mathcal{H})$ . Moreover, we have a commutative diagram

Letting  $\psi_{\underline{n}}^{2-q}$  denote the induced map between the homotopy fibers, we get a commutative diagram of distinguished triangles in  $\mathcal{D}(X_{\text{\acute{e}t}}, \mathbb{Z}/p^m)$ :

Now recall that, for  $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \mathcal{D}(X_{\text{\'et}}, \mathbb{Z}/p^m)$ , we have canonical isomorphism

 $\mathbf{R}\mathrm{Hom}^{\cdot}(\mathbb{Z}/p^{m},\mathbf{R}\mathcal{H}om^{\cdot}(\mathcal{B}^{\cdot},\mathcal{C}^{\cdot})) \xrightarrow{\cong} \mathbf{R}\mathrm{Hom}^{\cdot}(\mathcal{B}^{\cdot},\mathcal{C}^{\cdot}),$ 

(cf. [71, Prop. 4.3]) and also the canonical map

$$\mathbf{R}^{i}$$
Hom  $(\mathcal{B}, \mathcal{C}) \to$  Hom  $(\mathbf{R}^{j}$ Hom  $(\mathbb{Z}/p^{m}, \mathcal{B}), \mathbf{R}^{i+j}$ Hom  $(\mathbb{Z}/p^{m}, \mathcal{C})),$ 

for  $i, j \in \mathbb{Z}$ . So, any map  $\mathcal{A} \to \mathbf{R}\mathcal{H}om^{\cdot}(\mathcal{B}^{\cdot}, \mathcal{C}^{\cdot})$  in  $\mathcal{D}(X_{\text{\acute{e}t}}, \mathbb{Z}/p^m)$  induces a canonical map

$$H^i_{\mathrm{\acute{e}t}}(X,\mathcal{A}) = \mathbf{R}^i \operatorname{Hom}^{\cdot}(\mathbb{Z}/p^m,\mathcal{A}) \to \mathbf{R}^i \operatorname{Hom}^{\cdot}(\mathcal{B}^{\cdot},\mathcal{C}^{\cdot}) \to$$

#### CHAPTER 8. DUALITY THEOREM FOR HODGE-WITT COHOMOLOGY OVER 104 LOCAL FIELDS

$$\to \operatorname{Hom}(\mathcal{R}^{2-i}\operatorname{Hom}(\mathbb{Z}/p^m,\mathcal{B}),\mathcal{R}^2\operatorname{Hom}(\mathbb{Z}/p^m,\mathcal{C})) = \operatorname{Hom}(H^{2-i}_{\operatorname{\acute{e}t}}(X,\mathcal{B}),H^2_{\operatorname{\acute{e}t}}(X,\mathcal{C})).$$

Thus, applying  $\mathbf{R}^{i}$ Hom  $(\mathbb{Z}/p^{m}, (-))$  to (8.2.5) and using the above canonical maps, we get a commutative diagram of long exact hypercohomology sequences

where  $n \ge 0$ .

If  $x \in E$ , then (8.1.2) gives rise to a commutative diagram

which is compatible with (8.2.4).

Since  $\tau_x^*$  is exact, it is easy to check that the homotopy fibers of the left and the right vertical arrows are  $\tau_x^* W_m \mathcal{W}_n^{2^{-q}}[-1]$  and  $\mathbf{R}\mathcal{H}om(\tau_x^! W_m \mathcal{V}_n^q, \tau_x^! W_m \mathcal{H})$ , respectively. We also note that the maps  $j_x^* W_m \mathcal{W}_n^{2^{-q}} \to (\tau_x^h)_* \tau_x^* W_m \mathcal{W}_n^{2^{-q}}$  and  $(\tau_x^h)_* \tau_x^! W_m \mathcal{V}_n^q \to j_x^* W_m \mathcal{V}_n^q$ are isomorphisms on the étale (and Nisnevich) site of  $X_x^h$ . Since the canonical (i.e., forget support) map  $H_x^2(X, W_m \Omega_{X,\log}^2) \to H_{\acute{e}t}^2(X, W_m \Omega_{X,\log}^2)$  is bijective (cf. [47, Prop. 4.4]), letting  $\psi_{\underline{n},x}^{2^{-q}}$  denote the induced map between the mapping cones of the vertical arrows in (8.2.7), considering the corresponding maps between the hypercohomology and using the proper base change, we obtain a commutative diagram

$$0 \longrightarrow G_{\underline{n}}^{2-q,0}(x) \longrightarrow H^{0}(W_{m}\Omega_{X_{x}^{h}|D_{\underline{1}},\log}^{2-q}) \to H^{0}(\tau_{x}^{*}W_{m}\mathcal{W}_{\underline{n}+1}^{2-q}) \xrightarrow{\partial} G_{\underline{n}}^{2-q,1}(x) \longrightarrow \cdots$$

$$\begin{array}{c} \theta_{\underline{n}}^{*} \\ \theta_{\underline{n}}^{*} \\ 0 \to \mathbb{H}_{x}^{2}(W_{m}\mathcal{F}_{\underline{n}}^{q})^{\vee} \longrightarrow \mathbb{H}_{x}^{2}(W_{m}\mathcal{F}_{\underline{0}}^{q})^{\vee} \longrightarrow \mathbb{H}_{x}^{1}(W_{m}\mathcal{V}_{\underline{n}}^{q})^{\vee} \longrightarrow \mathbb{H}_{x}^{1}(W_{m}\mathcal{F}_{\underline{n}}^{q})^{\vee} \to \cdots \end{array}$$

$$(8.2.8)$$

of étale cohomology on  $X_x^h$  whose rows are exact and which is compatible with (8.2.6). Here, we let  $G_{\underline{n}}^{q,i}(x) = H^i_{\text{ét}}(X_x^h, W_m \Omega^q_{X_x^h|D_{n+1}, \log}).$ 

## 8.2.1 The maps $(\psi_{\underline{n},x}^q)^*$ vs. $\widehat{\rho}_{K,m}^{q,\underline{n}+1}$

Our goal now is to compute the cohomology of  $W_m \mathcal{W}_{\underline{n}}^q$  and  $W_m \mathcal{V}_{\underline{n}}^q$ , and show that  $(\psi_{\underline{n},x}^q)^*$  coincides with the reciprocity map  $\widetilde{\rho}_{K,m}^{q,n_x+1}$  of Lemma 7.3.2. We write  $D_{\underline{n}} = \sum_{x \in E} n_x[x]$ ,

where  $n_x$  is the multiplicity of  $D_{\underline{n}}$  at  $x \in E$  (i.e.  $n_x = n_i$  if  $x = x_i \in E$ ).

**Lemma 8.2.2.** Let  $q \ge 0, \underline{n} \ge \underline{1}$ . For  $x \in D_{\underline{n}}$ , we have the following.

1.  $H^i_x(X, W_m \mathcal{W}^q_{\underline{n}}) \xrightarrow{\cong} H^i_{\acute{e}t}(X^h_x, W_m \mathcal{W}^q_{\underline{n}}), \text{ for all } i \ge 0.$ 2.

$$H^0_{\acute{e}t}(X^h_x, W_m \mathcal{W}^q_{\underline{n}}) = \begin{cases} 0 & \text{if } q = 0\\ \varpi^m_q(A_x | n_x) & \text{if } q \ge 1. \end{cases}$$

3.  $H^i_{\acute{e}t}(X^h_x, W_m \mathcal{W}^q_{\underline{n}}) = 0 \text{ for } i \ge 1 \text{ and } q \ge 0.$ 

*Proof.* The items (1) and (2) are clear from the definition of  $W_m W_{\underline{n}}^q$  while (3) follows from Proposition 8.2.1 which says that  $W_m W_{\underline{n}}^q$  has a finite decreasing filtration whose graded quotients are coherent sheaves on  $D_{\underline{n}}$ .

Lemma 8.2.3. For  $x \in X^{(1)}$ , we have

$$H^1_x(X^h_x, W_m\Omega^q_{X|D_{\underline{n}}, \log}) \cong \frac{K^M_q(K_x)}{\operatorname{fil}_{n_x}K^M_q(K_x) + p^m K^M_q(K_x)}.$$

Proof. If  $x \notin D_{\underline{n}}$ , then  $H_x^1(X_x^h, W_m\Omega_{X|D_{\underline{n}},\log}^q) = H_x^1(X_x^h, W_m\Omega_{X,\log}^q) = H^0(k(x), W_m\Omega_{X,\log}^{q-1})$ =  $K_{q-1}^M(k(x))/p^m$ , where the second equality is a consequence of the Purity theorem [63, Thm 3.2] for  $W_m\Omega_{X_x^h,\log}^q$  on  $X_{x,\text{\acute{e}t}}^h$ . But this is same as the right hand side group in the lemma, because  $n_x = 0$  and so  $\operatorname{fil}_{n_x}K_q^M(K_x) = K_q^M(A_x)$ . If  $x \in D_{\underline{n}}$ , we claim  $H_{\text{\acute{e}t}}^1(X_x^h, W_m\Omega_{X_x^h|D_{n_x},\log}^q) = 0$  for all  $n_x \ge 1$ . By induction on m and Corollary 6.1.3, it is enough to prove the claim for m = 1. In this case, it is enough to show the map (cf. [31, Thm 1.2.1])

$$1 - \overline{F} : \operatorname{fil}_{-n_x} \Omega^q_{A_x} \to \operatorname{fil}_{-n_x} \Omega^q_{A_x} / d(\operatorname{fil}_{-n_x} \operatorname{fil}_{n_x} \Omega^{q-1}_{A_x}), \quad X^h_x = \operatorname{Spec} A_x;$$

is surjective. From the proof of Lemma 4.3.13, we note that the abelian group  $\operatorname{fl}_{-n_x}\Omega_{A_x}^q$ is generated by elements of the form  $\operatorname{adlog}(x_1) \wedge \cdots \wedge \operatorname{dlog}(x_q)$ , where  $a \in \operatorname{fl}_{-n_x} K_x = \pi_x^{n_x} A_x$ and  $x_i \in K_x$ , for all *i*. Here  $\pi_x$  is a uniformizer of  $A_x$ . Now, it is enough to show that  $1 - \overline{F} : \operatorname{fl}_{-n_x} K_x \to \operatorname{fl}_{-n_x} K_x$  is surjective. Let  $a \in \pi_x^{n_x} A_x$ . Since  $\overline{a}$ , the image of *a* is zero in k(x), by Hensel lemma, there exists  $y \in A_x$  such that  $y^p - y - a = 0$  in  $A_x$ . Let  $\overline{y}$  be the image of *y* in k(x). Then,  $\overline{y}^p - \overline{y} = 0 \in k(x)$ , which implies  $\overline{y} \in \mathbb{F}_p \subset A_x$ . Then letting  $y_1 = y - \overline{y}$ , we get  $y_1(y_1^{p-1} - 1) = a \in \pi_x^{n_x} A_x$ . But, note that  $y_1 \in \pi_x A_x$ , which implies  $(y_1^{p-1} - 1)$  is a unit in  $A_x$ . Hence, we conclude that  $y_1 \in \pi_x^{n_x} A_x$  and the claim follows.

Now the corollary follows from the localisation sequence for  $W_m \Omega^q_{X^h_{c}|D_{n_{\sigma}}, \log}$ .

**Lemma 8.2.4.** For  $x \in E, \underline{n} \ge \underline{0}$ , we have

CHAPTER 8. DUALITY THEOREM FOR HODGE-WITT COHOMOLOGY OVER 106 LOCAL FIELDS

*Proof.* This follows from Lemma 5.1.2 and Theorem 5.1.16.

Corollary 8.2.5. We have

$$\mathbb{H}_{x}^{i}(X_{x}^{h}, W_{m}\mathcal{F}_{\underline{n}}^{q}) = \begin{cases} \frac{H_{\acute{et}}^{1}(K_{x}, W_{m}\Omega_{K_{x}, \log}^{q})}{\operatorname{fil}_{n_{x}}H_{\acute{et}}^{1}(K_{x}, W_{m}\Omega_{K_{x}, \log}^{q})} & \text{if } i = 2\\ 0 & \text{if } i \neq 2. \end{cases}$$

*Proof.* First note that, the map  $\mathbb{H}^{i}_{\text{ét}}(X^{h}_{x}, W_{m}\mathcal{F}^{q}_{\underline{n}}) \to \mathbb{H}^{i}_{\text{ét}}(K_{x}, W_{m}\mathcal{F}^{q}_{\underline{n}})$  is injective by Theorem 5.1.16. Now the corollary follows from Lemma 8.2.4 and localisation exact sequence

$$\dots \to \mathbb{H}^{i}_{x}(X, W_{m}\mathcal{F}^{q}_{\underline{n}}) \to \mathbb{H}^{i}_{\mathrm{\acute{e}t}}(X^{h}_{x}, W_{m}\mathcal{F}^{q}_{\underline{n}}) \to \mathbb{H}^{i}_{\mathrm{\acute{e}t}}(K_{x}, W_{m}\mathcal{F}^{q}_{\underline{n}}) \to \mathbb{H}^{i+1}_{x}(X, W_{m}\mathcal{F}^{q}_{\underline{n}}) \to \cdots.$$

We also observe that  $\mathbb{H}^{i}_{\text{ét}}(K_{x}, W_{m}\mathcal{F}^{q}_{\underline{n}}) = H^{i}_{\text{ét}}(K_{x}, W_{m}\Omega^{q}_{K_{x},\log}) = 0$  for  $i \geq 2$ , because  $cd_{p}(K_{x}) \leq 1$ .

Lemma 8.2.6. We have

$$\mathbb{H}^{i}_{\acute{e}t}(X, W_{m}\mathcal{V}^{q}_{\underline{n}}) = \begin{cases} \bigoplus_{x \in E} \frac{\operatorname{fil}_{n_{x}}H^{1}_{\acute{e}t}(K_{x}, W_{m}\Omega^{q}_{K_{x}, \log})}{\operatorname{fil}_{0}H^{1}_{\acute{e}t}(K_{x}, W_{m}\Omega^{q}_{K_{x}, \log})} & \text{if } i = 1\\ 0 & \text{otherwise.} \end{cases}$$

*Proof.* Since  $\mathbb{H}^{i}_{\text{ét}}(X, W_m \mathcal{V}^q_{\underline{n}}) \cong \bigoplus_{x \in E} \mathbb{H}^{i}_{\text{ét}}(X^h_x, W_m \mathcal{V}^q_{\underline{n}})$ , we can replace X by  $X^h_x$  in order to prove the lemma. We now consider the long exact sequence

$$\dots \to \mathbb{H}^{0}_{\text{\acute{e}t}}(W_m \mathcal{F}^q_{\underline{0}}) \xrightarrow{\alpha_0} \mathbb{H}^{0}_{\text{\acute{e}t}}(W_m \mathcal{F}^q_{\underline{n}}) \to \mathbb{H}^{0}_{\text{\acute{e}t}}(W_m \mathcal{V}^q_{\underline{n}}) \to \mathbb{H}^{1}_{\text{\acute{e}t}}(W_m \mathcal{F}^q_{\underline{0}}) \xrightarrow{\alpha_1} \dots,$$
(8.2.9)

where  $\mathbb{H}^{i}_{\text{ét}}(W_m \mathcal{F}^q_{\underline{0}})$  (and similarly the other groups) is the short hand notation for  $\mathbb{H}^{i}_{\text{ét}}(X^h_x, W_m \mathcal{F}^q_{\underline{0}})$  for all i. It follows from (5.2.8) that  $\alpha_0$  is a bijection between two groups, each isomorphic to  $K^M_q(K_x)/p^m$ .

Lemma 8.2.4 and Theorem 5.1.16 imply that  $\alpha_1$  is injective. It follows that  $\mathbb{H}^0_{\text{ét}}(X^h_x, W_m \mathcal{V}^q_{\underline{n}}) = 0$ . We now apply Lemma 8.2.4 and (8.2.9) to conclude the proof.

Lemma 8.2.7. The diagram

commutes, where the horizontal arrows are canonical isomorphisms induced by Lemmas 8.2.2, 8.2.6 and (8.2.2).

*Proof.* The diagram (8.2.10) is the composition of two squares

Since (8.2.6) and (8.2.8) are compatible, it follows that the left square in (8.2.11) is commutative. It remains to show that the right square commutes. We can therefore assume  $E = \{x\}$ . We write  $\underline{n} = n$ .

Now, it follows from Lemma 8.2.2 that the map  $\kappa_q^m(A_x|1) \cong H^0_{\text{\acute{e}t}}(X_x^h, W_m \Omega_{X|D_1,\log}^q)$   $\rightarrow H^0_{\text{\acute{e}t}}(X_x^h, W_m \mathcal{W}_{n+1}^q)$  is surjective. Corollary 8.2.5 (with  $\underline{n} = 0$ ) and Lemma 8.2.6 imply that  $\mathbb{H}^2_x(X_x^h, W_m \mathcal{F}_0^{2-q})^{\vee} \rightarrow \mathbb{H}^1_x(X_x^h, \mathcal{V}_n^{2-q})^{\vee}$  is also surjective. We conclude that (8.2.8) breaks into a commutative diagram of short exact sequences

By definition of Kato's reciprocity map (cf.  $\S$  7.2), it follows from Corollary 8.2.5 that the left square in (8.2.12) is the same as the left square in the commutative diagram

$$\begin{array}{ccc} 0 \to \kappa_q^m(A_x|n+1) \to \kappa_q^m(A_x|1) \longrightarrow \varpi_q^m(A_x|n+1) \longrightarrow 0 & (8.2.13) \\ & & & & & & & \\ \rho_{K_x,m}^{q,n+1} & & & & & & \\ 0 \longrightarrow G_{K_x,m}^{3-q,n+1} \longrightarrow G_{K_x,m}^{3-q,1} \longrightarrow \left( \frac{\operatorname{fil}_n H_{\operatorname{\acute{e}t}}^1(K_x, W_m \Omega_{K_x, \log}^{2-q})}{\operatorname{fil}_0 H_{\operatorname{\acute{e}t}}^1(K_x, W_m \Omega_{K_x, \log}^{2-q})} \right)^{\vee} \to 0. \end{array}$$

From this, it immediately follows that  $(\psi_{n,x}^q)^* = \widehat{\rho}_{K_x,m}^{q,n+1}$ 

CHAPTER 8. DUALITY THEOREM FOR HODGE-WITT COHOMOLOGY OVER 108 LOCAL FIELDS

## 8.3 Topologies on the cohomology groups

Our next goal is to endow the cohomology groups  $H^i_{\text{\acute{e}t}}(X, W_m \Omega^q_{X|D_{\underline{n}+1}, \log})$  and  $\mathbb{H}^j_{\text{\acute{e}t}}(X, W_m \mathcal{F}^q_{\underline{n}})$  with the structure of topological abelian groups so that we get a perfect pairing of topological abelian groups consisting them. We fix some notations.

Notation 8.3.1. For  $\underline{n}, i, j \ge 0, 1 \le q \le 2$ , we let

$$G_{\underline{n}}^{q,i}(m) \coloneqq H^i_{\text{\'et}}(X, W_m \Omega^q_{X|D_{\underline{n}+1}, \log}) , F_{\underline{n}}^{q,j}(m) \coloneqq \mathbb{H}^j_{\text{\'et}}(X, W_m \mathcal{F}^q_{\underline{n}}).$$

Note that by Lemma 6.1.4,  $G_{\underline{0}}^{2,i}(m) = H_{\text{\acute{e}t}}^i(X, W_m \Omega_{X,\log}^2)$  and  $F_{\underline{0}}^{0,j}(m) = H_{\text{\acute{e}t}}^j(X, \mathbb{Z}/p^m)$ . It is clear that  $G_{\underline{n}}^{q,i}(m) = 0$  for  $i \ge 3$  (cf. [56, Chap. VI, Rem. 1.5]). As for i = 2, it follows from Lemmas 8.2.2, 8.2.6 and (8.2.6) that

$$G_{\underline{n}}^{q,2}(m) = G_{\underline{0}}^{q,2}(m), \ F_{\underline{n}}^{q,0}(m) = F_{\underline{0}}^{q,0}(m), \ \text{for } n \ge 0.$$
 (8.3.1)

8.3.1 Topology of  $H^i_{\acute{et}}(X, W_m \Omega^q_{X|D_{\underline{n}+1}, \log})$  for  $i \neq 1$ 

We consider four cases.

**Case 1** (i = 2, q = 2): The pairing (8.1.1) induces an isomorphism  $G_{\underline{0}}^{2,2}(m) \xrightarrow{\cong} F_{\underline{0}}^{0,0}(m)^{\vee} \cong \mathbb{Z}/p^m$  by [40, § 3, Prop. 4] (see also [47, Thm. 4.7]). We therefore endow  $G_{\underline{n}}^{2,2}(m)$  with the discrete topology via the isomorphism (8.3.1) and hence

$$G_{\underline{n}}^{2,2}(m) \times F_{\underline{n}}^{0,0}(m) \to \mathbb{Z}/p^m.$$

$$(8.3.2)$$

is a perfect pairing.

**Case 2** (i = 0, q = 2): First consider the case  $\underline{n} = 0$ . In this case, by [40, § 3, Prop. 4] (see also [47, Thm. 4.7]),  $F_{\underline{0}}^{0,2}(m)$  has discrete topology,  $G_{\underline{0}}^{2,0}(m)$  has profinite topology and the map  $G_{\underline{0}}^{2,0}(m) \to F_{\underline{0}}^{0,2}(m)^{\vee}$  is an isomorphism of profinite groups. It follows then from Lemma 7.3.2, Lemma 8.2.7, (8.2.6) and that the map  $G_{\underline{n}}^{2,0}(m) \to F_{\underline{n}}^{0,2}(m)^{\vee}$  is an isomorphism. Since  $F_{\underline{0}}^{0,2}(m)$  has discrete topology and  $F_{\underline{n}}^{0,2}(m)$  is its quotient, the latter is a discrete topological abelian group. We therefore endow  $G_{\underline{n}}^{2,0}(m)$  with the profinite topology so that

$$G_{\underline{n}}^{2,0}(m) \times F_{\underline{n}}^{0,2}(m) \to \mathbb{Z}/p^m.$$
 (8.3.3)

is a perfect pairing of topological abelian groups. We also note that  $\varinjlim_{\underline{n}} F_{\underline{n}}^{0,2}(m) = H_{\text{\acute{e}t}}^2(U, \mathbb{Z}/p^m) = 0$  (by Theorem 4.3.17(12)), because U is affine and  $cd_pU \leq 1$ . Hence

$$\lim_{\underline{m}} G_{\underline{n}}^{2,0}(m) = (\lim_{\underline{m}} F_{\underline{n}}^{0,2}(m))^{\vee} = 0.$$
(8.3.4)

**Case 3** (i = 0, q = 1): First consider the sub case  $\underline{n} = \underline{0}$ . Since  $W_m \mathcal{F}_{\underline{0}}^1 \cong j_* W_m \Omega_{U,\log}^1$  by Lemma 6.1.2(4), we consider the following exact sequences.

$$0 \to W_m \Omega^1_{X|D_{\underline{1}},\log} \to W_m \Omega^1_{X,\log} \to \bigoplus_{x \in E} (\tau_x)_* W_m \Omega^1_{\operatorname{Spec}\kappa(x),\log} \to 0,$$
(8.3.5)

$$0 \to W_m \Omega^1_{X, \log} \to j_* W_m \Omega^1_{U, \log} \xrightarrow{\operatorname{res}} \bigoplus_{x \in E} (\tau_x)_* W_m \Omega^0_{\operatorname{Spec} \kappa(x), \log} \to 0,$$

where  $\kappa(x)$  is the residue field of x and res is induced by the residue map of Milnor K-theory. This induces the following commutative diagram.

By a similar argument to that in Lemma 8.2.7, we conclude that the right vertical arrow is the reciprocity map of 1-local field  $\kappa(x)$ , which is known to be injective. The middle vertical arrow is isomorphism by [47, Thm 4.7]. This implies the left vertical arrow is also isomorphism. Since  $H^2_{\text{ét}}(X, W_m \Omega^1_{X, \log})$  is discrete (cf. [47, Prop. 3.6]),  $F^{1,2}_{\underline{0}}(m)$  being its quotient, is a discrete group. Therefore  $G^{1,0}_{\underline{0}}(m)$  is endowed with profinite topology and

$$G_{\underline{0}}^{1,0}(m) \times F_{\underline{0}}^{1,2}(m) \to \mathbb{Z}/p^m$$
 (8.3.7)

is a perfect pairing of topological abelian groups. Now by Lemma 7.3.2, Lemma 8.2.7, and (8.2.6) we get  $G_{\underline{n}}^{1,0}(m) \xrightarrow{\cong} F_{\underline{n}}^{1,2}(m)^{\vee}$  is isomorphism for  $n \ge 0$ .  $F_{\underline{n}}^{1,2}(m)$  is discrete (being quotient of  $F_0^{1,2}(m)$ ). So we endow  $G_{\underline{n}}^{1,0}(m)$  with profinite topology and hence get a perfect pairing

$$G_{\underline{n}}^{1,0}(m) \times F_{\underline{n}}^{1,2}(m) \to \mathbb{Z}/p^m$$
 (8.3.8)

for all  $n \ge 0$ .

Since  $\varinjlim_{\underline{n}} F_{\underline{n}}^{1,2}(m) = H^2_{\text{\acute{e}t}}(U, W_m \Omega^1_{U, \log}) = 0$  (by Theorem 4.3.17(12) and  $cd_p U \leq 1$ ), we

have

$$\lim_{\underline{n}} G_{\underline{n}}^{1,0}(m) = (\lim_{\underline{n}} F_{\underline{n}}^{1,2}(m))^{\vee} = 0.$$
(8.3.9)

Case 4 (i = 2, q = 1): Again first we consider the sub case <u>n</u> = 0. (8.3.5) induces a

#### CHAPTER 8. DUALITY THEOREM FOR HODGE-WITT COHOMOLOGY OVER 110 LOCAL FIELDS

commutative diagram

$$\begin{array}{cccc} 0 \to H^{0}_{\text{\acute{e}t}}(X, W_{m}\Omega^{1}_{X, \log}) \to F^{1,0}_{\underline{0}}(m) & \longrightarrow \bigoplus_{x \in E} \mathbb{Z}/p^{m} \longrightarrow H^{1}_{\text{\acute{e}t}}(X, W_{m}\Omega^{1}_{X, \log}) \\ & 1 & 1 & 2 & & & & & & \\ 0 \to H^{2}_{\text{\acute{e}t}}(X, W_{m}\Omega^{1}_{X, \log})^{\vee} \to G^{1,2}_{\underline{0}}(m)^{\vee} \to \bigoplus_{x \in E} H^{1}_{\text{\acute{e}t}}(\kappa(x), W_{m}\Omega^{1}_{\kappa(x), \log})^{\vee} \to H^{1}_{\text{\acute{e}t}}(X, W_{m}\Omega^{1}_{X, \log})^{\vee}, \\ \end{array}$$

$$\begin{array}{c} (8.3.10) \\ \end{array}$$

where (1) is isomorphism and (1') is injection by [47, Prop. 3.6, Thm 4.7]. (3) is the dual of the isomorphism  $H^1_{\text{\acute{e}t}}(\kappa(x), W_m \Omega^1_{\kappa(x), \log}) \xrightarrow{inv_x} \mathbb{Z}/p^m$  (invariant map of class field theory of  $\kappa(x)$ ) and so is isomorphism. This implies (2) is isomorphism. By loc.cit,  $H^0_{\text{\acute{e}t}}(X, W_m \Omega^1_{X, \log})$  is profinite. Thus we endow discrete topology on  $G^{1,2}_{\underline{0}}(m)$  so that  $F^{1,0}_{\underline{0}}(m) \xrightarrow{\cong} G^{1,2}_{\underline{0}}(m)^{\vee}$  is an isomorphism of profinite group. Now by (8.3.1), we know  $G^{1,2}_{\underline{n}}(m) = G^{1,2}_{\underline{0}}(m)$  and  $F^{1,0}_{\underline{n}}(m) = F^{1,0}_{\underline{0}}(m) = H^0_{\text{\acute{e}t}}(U, W_m \Omega^1_{U, \log})$ . Hence we have perfect pairings

$$G_{\underline{n}}^{1,2}(m) \times F_{\underline{n}}^{1,0}(m) \to \mathbb{Z}/p^m, \text{ for all } n \ge 0,$$

$$\lim G_{\underline{n}}^{1,2}(m) \times H_{\underline{n}}^0(U, W_n, \Omega_{\underline{n}}^1, \dots) \to \mathbb{Z}/p^m$$
(8.3.11)

$$\varprojlim_{\underline{n}} G^{1,2}_{\underline{n}}(m) \times H^0_{\text{\'et}}(U, W_m \Omega^1_{U, \log}) \to \mathbb{Z}/p^m$$

# 8.3.2 Topology of $H^1_{\acute{e}t}(X, W_m \Omega^2_{X|D_{n+1}, \log})$


Now consider the case (i = 1, q = 2). To define topology on  $G_{\underline{n}}^{2,1}(m)$  we shall use a recipe.

Let G and G' be two abelian groups such that G' is a torsion group. Let  $G' \times G \to \mathbb{Q}/\mathbb{Z}$  be a pairing which has no left kernel. Let  $\beta: G' \to G^{\vee}$  be the induced injective homomorphism. We let G be endowed with the weakest topology with respect to which the map  $\beta(f): G \to \mathbb{Q}/\mathbb{Z}$  is continuous for every  $f \in G'$  (called the weak topology induced by G'). This makes G a topological abelian group for which the identity element has a fundamental system of open neighborhoods of the form  $\operatorname{Ker}(\beta(f_1)) \cap \cdots \cap \operatorname{Ker}(\beta(f_r))$  with  $f_1, \ldots, f_r \in G'$ . It is clear that  $\beta$  has a factorization  $\beta: G' \to G^*$ , where the dual on the right is taken with respect to the weak topology.

#### **Lemma 8.3.2.** The map $\beta: G' \to G^*$ is a bijection.

*Proof.* Let  $f \in G^*$ . We can find  $f_1, \ldots, f_r \in G'$  such that the continuous homomorphism

 $f: G \to \mathbb{Q}/\mathbb{Z}$  has the property that  $H := \bigcap_{i=1}^{r} \operatorname{Ker}(\beta(f_i)) \subset \operatorname{Ker}(f)$ . We now look at



Since  $\mathbb{Q}/\mathbb{Z}$  is injective, there exists  $\tilde{f}$  such that the above diagram is commutative. Since G' is a torsion group, we see that each summand of the the group on the bottom is a finite cyclic group. It follows that we can replace  $\mathbb{Q}/\mathbb{Z}$  by a finite cyclic group in (8.3.12). Using the standard description of homomorphisms between finite cyclic groups, we can find integers  $n_1, \ldots, n_r$  such that  $f = \tilde{f} \circ \Delta = \sum_{i=1}^r n_i \beta(f_i) = \beta(\sum_{i=1}^r n_i f_i) \in G'$ .  $\Box$ 

For a topological abelian group G whose topology is  $\tau$ , we let  $G^{\text{pf}}$  denote the profinite completion of G with respect to  $\tau$ . That is,  $G^{\text{pf}}$  is the inverse limit  $\lim_{t \to U} G/U$  with the inverse limit topology, where U runs through the  $\tau$ -open subgroups of finite index in G. It is clear that  $G^{\text{pf}}$  is a profinite abelian group and the profinite completion map  $G \to G^{\text{pf}}$  is continuous with dense image. We note the following elementary lemma.

**Lemma 8.3.3.** If G is a topological abelian group, then the map  $(G^{\text{pf}})^* \to G^*$  induces a topological isomorphism  $(G^{\text{pf}})^* \to (G^*)_{\text{tor}}$  for the discrete topology of  $(G^*)_{\text{tor}}$ . In particular if G is torsion topological abelian group, then  $(G^{\text{pf}})^* \to G^*$  is isomorphism.

Proof. Since the profinite completion map  $G \to G^{\text{pf}}$  is continuous with dense image, it follows that the canonical map  $(G^{\text{pf}})^* \to G^*$  is injective and its image lies in  $(G^*)_{\text{tor}}$  (as any continuous map  $f: G^{\text{pf}} \to \mathbb{Q}/\mathbb{Z}$  has finite order). On the other hand, any continuous map  $f; G \to \mathbb{Q}/\mathbb{Z}$  of finite order factors through G/Ker f and hence through  $G^{\text{pf}}$ , because Ker f is an open subgroup of finite index in G. This proves the lemma.

Let G be a torsion topological abelian group and let  $G^*$  be endowed with the discrete topology. Then the evaluation map  $\operatorname{ev}_G: G \to (G^*)^{\vee}$  is continuous with respect to the profinite topology of the target group. This implies that  $\operatorname{ev}_G$  has a canonical factorization  $G \to G^{\operatorname{pf}} \xrightarrow{\operatorname{ev}_G^{\operatorname{pf}}} (G^*)^{\vee}$  with  $\operatorname{ev}_G^{\operatorname{pf}}$  continuous.

**Corollary 8.3.4.** The map  $ev_G^{pf}: G^{pf} \to (G^*)^{\vee}$  is a topological isomorphism.

*Proof.* Follows from Lemma 8.3.3 using the Pontryagin duality.

111

#### CHAPTER 8. DUALITY THEOREM FOR HODGE-WITT COHOMOLOGY OVER 112 LOCAL FIELDS

To apply the above recipe, we use the pairing  $F_{\underline{n}}^{0,1}(m) \times G_{\underline{n}}^{2,1}(m) \to \mathbb{Z}/p^m$  (cf. (8.1.1)) to endow  $G_{\underline{n}}^{2,1}(m)$  with the structure of a topological abelian group via the weak topology induced by  $F_{\underline{n}}^{0,1}(m)$ . To check that the hypothesis of Lemma 8.3.2 is satisfied, we look at the commutative diagram

obtained by dualizing (8.2.6). Lemmas 8.2.2 and 8.2.6 imply that the rows of this diagram are exact. In combination with Lemma 7.3.3, Lemma 8.2.7 and [40, Prop. 3.4] (see also [47, Thm. 4.7]), these results also imply that  $(\theta_{\underline{n}}^{\star})^{\vee}$  is a monomorphism. This proves that the desired hypothesis is satisfied.

We conclude from Lemma 8.3.2 that with respect to the weak topology of  $G_{\underline{n}}^{2,1}(m)$ , the map  $(\theta_{\underline{n}}^{*})^{\vee}: F_{\underline{n}}^{0,1}(m) \to G_{\underline{n}}^{2,1}(m)^{*}$  is a bijection.

Remark 8.3.5. Via the pairing  $F_{\underline{0}}^{0,1}(m) \times G_{\underline{0}}^{2,1}(m) \to \mathbb{Z}/p^m$ , we can endow  $G_{\underline{0}}^{2,1}(m)$  with the weak topology induced by  $F_{\underline{0}}^1(m)$ . Using the Pontryagin duality between the profinite topology of  $G_{\underline{0}}^{2,1}(m)$  and the discrete topology of  $F_{\underline{0}}^1(m)$  (cf. [47, Thm 4.7]), it is easy to check that  $\{\operatorname{Ker}(\beta(f))|f \in F_{\underline{0}}^{0,1}(m)\}$  forms a sub-base for the profinite topology of  $G_{\underline{0}}^{2,1}(m)$ . It follows that the weak topology of  $G_{\underline{0}}^{2,1}(m)$  coincides with the profinite topology. Furthermore,  $G_{\underline{n}}^{2,1}(m) \to G_{\underline{0}}^{2,1}(m)$  is continuous.

By combining the isomorphism  $(\theta_{\underline{n}}^*)^{\vee}: F_{\underline{n}}^{0,1}(m) \to G_{\underline{n}}^{2,1}(m)^*$ , Corollary 8.3.4 and the Pontryagin duality, we get the following case of the duality.

**Corollary 8.3.6.** For  $n \ge 0$ , there is a perfect pairing of topological abelian groups

$$(G^{2,1}_{\underline{n}}(m))^{\mathrm{pf}} \times F^{0,1}_{\underline{n}}(m) \to \mathbb{Z}/p^m.$$

Hence, we have an isomprphism  $(G_{\underline{n}}^{2,1}(m))^{\text{pf}} \cong \pi_1^{ab}(X, D_n)/p^m$ .

*Proof.* First part of the corollary is clear from the previous discussion. The second part follows from the first part, (5.2.1) and Theorem 5.2.6.

**Case 5** (i = 1, q = 2): We have a perfect pairing of topological abelian groups

$$(G_{\underline{n}}^{2,1}(m))^{\mathrm{pf}} \times F_{\underline{n}}^{0,1}(m) \to \mathbb{Z}/p^m, \qquad (8.3.14)$$

where  $(G_{\underline{n}}^{2,1}(m))^{\text{pf}}$  has profinite topology and  $F_{\underline{n}}^{0,1}(m)$  has discrete topology.

Even though we don't know much about the topology of  $G_{\underline{n}}^{2,1}(m)$ , the following lemma ensures that the topology is separated. This property is crucial while proving Class field theory in modulus setup.

**Lemma 8.3.7.** The profinite completion map  $G_{\underline{n}}^{2,1}(m) \to (G_{\underline{n}}^{2,1}(m))^{\text{pf}}$  is injective, and the induced subspace topology of  $G_{\underline{n}}^{2,1}(m)$  coincides with its weak topology. In particular, the latter topology is Hausdorff.

*Proof.* By Corollary 8.3.6, the injectivity claim is equivalent to that the map  $\theta_{\underline{n}}^*$  in the following (cf. (8.2.6)) is injective.

But this follows because  $(\psi_{\underline{n}}^2)^*$  is injective by Lemma 7.3.2 and Lemma 8.2.7 while the maps  $\theta^*$  are bijective by [40, § 3, Prop. 4] (see also [47, Thm. 4.7]). The claim, that the weak topology of  $G_{\underline{n}}^{2,1}(m)$  coincides with the subspace topology induced from the profinite topology of  $(G_{\underline{n}}^{2,1}(m))^{\text{pf}}$ , follows from Lemma 8.3.2 whose proof shows that every open subgroup of  $G_{\underline{n}}^{2,1}(m)$  has finite index.

**Lemma 8.3.8.** For  $\underline{n} \geq \underline{n}'$ , the canonical map  $G_{\underline{n}}^{2,1}(m) \to G_{\underline{n}'}^{2,1}(m)$  is surjective continuous and the induced map  $(G_{\underline{n}}^{2,1}(m))^{\text{pf}} \to (G_{\underline{n}'}^{2,1}(m))^{\text{pf}}$  is a topological quotient.

*Proof.* We consider the exact sequence in  $X_{\text{ét}}$ 

$$0 \to W_m \Omega^2_{X|D_{\underline{n}},\log} \to W_m \Omega^2_{X|D_{\underline{n}'},\log} \to W_m \Omega^2_{X|D_{\underline{n}'},\log} / W_m \Omega^2_{X|D_{\underline{n}},\log} \to 0.$$

Since  $\mathcal{A} = W_m \Omega^2_{X|D_{\underline{n}'},\log}/W_m \Omega^2_{X|D_{\underline{n}},\log}$  is supported on E, by Proposition 8.2.1, we have a filtration of  $\mathcal{A}$  whose successive quotients are coherent sheaves on E. Hence  $H^1_{\text{ét}}(X, \mathcal{A}) = 0$ . This implies the map  $G^{2,1}_{\underline{n}'}(m) \to G^{2,1}_{\underline{n}'}(m)$  is surjective. Now we look at the commutative diagram

$$G_{\underline{n}}^{2,1}(m) \longrightarrow G_{\underline{n}'}^{2,1}(m) \tag{8.3.16}$$

$$(G_{\underline{n}}^{2,1}(m))^{\mathrm{pf}} \longrightarrow (G_{\underline{n}'}^{2,1}(m))^{\mathrm{pf}}.$$

By Theorem 5.2.6, we see that the transition map  $F_{\underline{n}'}^{0,1}(m) \to F_{\underline{n}}^{0,1}(m)$  is injective. We conclude from Corollary 8.3.6 that the bottom horizontal arrow in (8.3.16) is a surjective continuous homomorphism between two profinite groups. It must therefore be a topological quotient map. Finally, the claim, that the horizontal arrow in the top row of (8.3.16) is continuous, follows from Lemma 8.3.7 which asserts that each term of this row is a subspace of the corresponding term in the bottom row.

#### CHAPTER 8. DUALITY THEOREM FOR HODGE-WITT COHOMOLOGY OVER 114 LOCAL FIELDS

Corollary 8.3.9. We have a perfect pairing of topological groups

$$\lim_{\underline{\leftarrow} \underline{n}} (G_{\underline{n}}^{2,1}(m))^{\mathrm{pf}} \times H^1_{\acute{e}t}(U, \mathbb{Z}/p^m) \to \mathbb{Z}/p^m.$$

*Proof.* By Theorem 4.3.17(12), we have  $\varinjlim_{\underline{n}} F_{\underline{n}}^{0,1}(m) = H^1_{\text{\'et}}(U, \mathbb{Z}/p^m)$ . We now combine Corollary 8.3.6, [22, Lem. 7.2] and [47, Lem. 2.6] to conclude the proof.

#### 8.3.3 The Brauer-Manin pairing with modulus

In this section we recall the Brauer-Manin pairing for modulus studied in [47]. This will be necessary to endow  $G_n^{1,1}$  with a topology.

Let  $\operatorname{Pic}(X)$  be the group of isomorphism classes of line bundles on X. For a fixed effective divisor D, the relative Picard group  $\operatorname{Pic}(X|D)$  (usually written as  $\operatorname{Pic}(X,D)$ in the literature) is the set of isomorphism classes of pairs  $(\mathcal{L}, u: \mathcal{O}_D \xrightarrow{\cong} \mathcal{L}|_D)$ , where  $\mathcal{L}$  is an invertible sheaf on X and  $\mathcal{L}|_D \coloneqq \mathcal{L} \otimes_{\mathcal{O}_X} \mathcal{O}_D$ . One says that  $(\mathcal{L}, u) \cong (\mathcal{L}', u')$  if there is an isomorphism  $w: \mathcal{L} \xrightarrow{\cong} \mathcal{L}'$  such that  $u' = w|_D \circ u$ .  $\operatorname{Pic}(X|D)$  is an abelian group under the tensor product of invertible sheaves and their trivializations along D whose identity element is  $(\mathcal{O}_X, \operatorname{id})$ . There is a canonical homomorphism  $\operatorname{Pic}(X|D) \to \operatorname{Pic}(X)$  which is surjective. By [73, Lem 2.1] we have  $\operatorname{Pic}(X|D) = H^1_{\operatorname{\acute{e}t}}(X, \mathcal{K}^M_{1,X|D}) = H^1_{\operatorname{Zar}}(X, \mathcal{K}^M_{1,X|D})$ .

Let  $\operatorname{Pic}^{0}(X)$  be the subgroup of degree zero divisor classes in  $\operatorname{Pic}(X)$  and  $\operatorname{Pic}(X)$ be the Picard scheme of X.  $\operatorname{Pic}(X)(k)$  has a structure of totally disconnected locally compact Hausdorff topological group, whose topology is induced by the adic (a.k.a the valuation) topology of k (cf. [47, § 6.3]). We call the topology to be adic topology. Recall that the connected component  $\operatorname{Pic}^{0}(X)$  of  $\operatorname{Pic}(X)$  is a connected abelian variety over k. It follows from [12, Thm 10.5.1] that the group of k rational points  $\operatorname{Pic}^{0}(X)(k)$  is compact as adic topological group. Hence,  $\operatorname{Pic}^{0}(X)(k)$  is a profinite (because profinite = compact + totally disconnected + Hausdorff) topological abelian group. This topology also induces an adic topology on  $\operatorname{Pic}(X)$  (note,  $\operatorname{Pic}(X) = \operatorname{Pic}(X)(k)$  if  $X(k) \neq \phi$ ) such that the connected component  $\operatorname{Pic}^{0}(X)$  is a profinite topological group.

Since we have an exact sequence :  $\mathbb{Z}^r \to \operatorname{Pic}(X) \to \operatorname{Pic}(U) \to 0$ , we endow  $\operatorname{Pic}(U)$ with the quotient topology. Also we have an exact sequence of topological group :  $0 \to \operatorname{Pic}^0(X) \to \operatorname{Pic}(X) \to \mathbb{Z} \to 0$ . As a result we also conclude that the quotient topology on  $\operatorname{Pic}(X)/p^m$  and  $\operatorname{Pic}(U)/p^m$  are profinite for all  $m \ge 1$ .

By [47, Thm 5.6], there exists a group scheme  $\operatorname{Pic}(X|D_{\underline{n}})$  ( $\underline{n} \geq \underline{1}$ ) over k such that  $\operatorname{Pic}(X|D_{\underline{n}}) = \operatorname{Pic}(X|D_{\underline{n}})(k)$ . Hence  $\operatorname{Pic}(X|D_{\underline{n}})$  has a structure of adic topological group whose topology is induced from k, which is locally compact and Hausdorff.

Let's write  $\operatorname{Br}(X|D_n)$  for the Brauer group with modulus  $D_n$  defined in Definition 5.2.7 (cf. [47, Defn 8.9]). We endow  $\operatorname{Br}(X)$  and  $\operatorname{Br}(X|D_n)$  with discrete topology. By [70, § 9], [47, Prop 9.4, 11.2] there exist continuous pairings (called the Brauer-Manin pairings)

$$\operatorname{Br}(X) \times \operatorname{Pic}(X) \to \mathbb{Q}/\mathbb{Z}, \quad \operatorname{Br}(X|D_{\underline{n}}) \times \operatorname{Pic}(X|D_{\underline{n}+1}) \to \mathbb{Q}/\mathbb{Z}; \quad \underline{n} \ge \underline{0},$$
(8.3.17)

compatible with the maps  $Br(X) \hookrightarrow Br(X|D_{\underline{n}})$  and  $Pic(X|D_{\underline{n}+1}) \twoheadrightarrow Pic(X)$ . Moreover, by [70, Thm 9.2], [47, Lem 12.2(1)], we have isomorphisms

$$\operatorname{Br}(X) \cong (\operatorname{Pic}(X))^*, \quad \operatorname{Br}(X|D_{\underline{n}}) \cong (\operatorname{Pic}(X|D_{\underline{n}+1}))^*.$$
(8.3.18)

Since  $\operatorname{Br}(X|D_{\underline{n}})$  is torsion group, it follows that  $(\operatorname{Pic}(X|D_{\underline{n}+1}))^* = (\operatorname{Pic}(X|D_{\underline{n}+1}))^*_{\operatorname{tor}}$ . As a result, we get  $(\operatorname{Pic}(X|D_{\underline{n}+1})^{\operatorname{pf}})^* = (\operatorname{Pic}(X|D_{\underline{n}+1}))^*$  by Lemma 8.3.3. Taking  $p^m$  torsion, we get  $(\operatorname{Pic}(X|D_{\underline{n}+1})^{\operatorname{pf}}/p^m)^* = (\operatorname{Pic}(X|D_{\underline{n}+1})/p^m)^*$ . Note that the quotient topology on  $\operatorname{Pic}(X|D_{\underline{n}+1})^{\operatorname{pf}}/p^m$  is profinite. Hence, by Pontryagin duality and Corollary 8.3.4, we have

$$\operatorname{Pic}(X|D_{\underline{n}+1})^{\operatorname{pf}}/p^{m} \cong ((\operatorname{Pic}(X|D_{\underline{n}+1})/p^{m})^{*})^{\vee} \cong (\operatorname{Pic}(X|D_{\underline{n}+1})/p^{m})^{\operatorname{pf}}.$$
(8.3.19)

Now (8.3.18) induces the following isomorphisms.

(1) 
$$_{p^m}\operatorname{Br}(X) \cong (\operatorname{Pic}(X)/p^m)^*$$
, (2)  $_{p^m}\operatorname{Br}(X|D_{\underline{n}}) \cong (\operatorname{Pic}(X|D_{\underline{n}+1})/p^m)^*$ , (8.3.20)

(3) 
$$\operatorname{Pic}(X)/p^m \cong (p^m \operatorname{Br}(X))^*$$
, (4)  $\operatorname{Pic}(X|D_{\underline{n}+1})^{\operatorname{pf}}/p^m \cong (p^m \operatorname{Br}(X|D_{\underline{n}}))^*$ ,

where the last two isomorphisms of (8.3.20) are obtained by taking Pontryagin dual of the first two isomorphisms, (8.3.19) and also noting  $\text{Pic}(X)/p^m$  is profinite. Moreover, the isomorphisms in bottom row are topological isomorphisms.

We end this section by recalling the following commutative diagram of short exact sequences of topological groups (see the proof of [47, Cor 9.6]).

where  $\alpha_{p^m}$  and  $\gamma_{p^m}$  are induced by (8.3.17) and  $\beta_{p^m}$  is induced from bottom row of (6.2.6).

#### CHAPTER 8. DUALITY THEOREM FOR HODGE-WITT COHOMOLOGY OVER 116 LOCAL FIELDS

## 8.3.4 Topology of $H^1_{\acute{e}t}(X, W_m \Omega^1_{X|D_{n+1}, \log})$

Finally, we consider the case (i = 1, q = 1). First we prove a general proposition.

**Proposition 8.3.10.** Let Y be an N-dimensional smooth scheme finite type over an F-finite field k. Let  $D_{\underline{n}}$  be as in § 6.1. For all  $\underline{n} \ge \underline{1}, d \ge N, q \ge 0, m \ge 1$ , we have

$$H^{d}_{\acute{e}t}(Y, \mathcal{K}^{M}_{q,Y|\underline{D_{n}}}/p^{m}) \xrightarrow{\cong} H^{d}_{\acute{e}t}(Y, W_{m}\Omega^{q}_{Y|\underline{D_{n}}, \log}).$$

*Proof.* The map is induced by the canonical map  $d\log : \mathcal{K}^M_{q,Y|D_n}/p^m \to W_m \Omega^q_{Y|D_n,\log}$  which is an isomorphism of Nisnevich (hence also étale) sheaves if  $\underline{n} = \underline{1}$  by [45, Thm 2.2.4]. Since  $\mathcal{K}^M_{q,Y|D_n}$  is *p*-torsion free, the exact sequence

$$0 \to \mathcal{K}^{M}_{q,Y|D_{\underline{n}}} \to \mathcal{K}^{M}_{q,Y|D_{\underline{1}}} \to \frac{\mathcal{K}^{M}_{q,Y|D_{\underline{1}}}}{\mathcal{K}^{M}_{q,Y|D_{\underline{n}}}} \to 0$$

induces another exact sequence of the form

$$0 \to {}_{p^m} \left( \frac{\mathcal{K}^M_{q,Y|D_{\underline{1}}}}{\mathcal{K}^M_{q,Y|D_{\underline{n}}}} \right) \to \mathcal{K}^M_{q,Y|D_{\underline{n}}} / p^m \xrightarrow{\phi} \mathcal{K}^M_{q,Y|D_{\underline{1}}} / p^m$$
(8.3.22)

for every  $\underline{n} \geq \underline{1}$ . It is enough to show that  $H^d_{\text{\acute{e}t}}(Y, {}_{p^m}\left(\frac{\mathcal{K}^M_{q,Y|D_{\underline{1}}}}{\mathcal{K}^M_{q,Y|D_{\underline{n}}}}\right)) = 0$  for  $d \geq N$  and Image  $\phi \cong W_m \Omega^q_{Y|D_{\underline{n}},\log}$ , for all  $m \geq 1$ . First we claim that, for  $m \gg 0$ , we have an isomorphism of Nisnevich (resp. étale) sheaves

$$\operatorname{dlog}: \frac{\mathcal{K}_{q,Y|D_{\underline{1}}}^{M}}{\mathcal{K}_{q,Y|D_{\underline{n}}}^{M}} \xrightarrow{\cong} \frac{W_{m}\Omega_{Y|D_{\underline{1}},\log}^{q}}{W_{m}\Omega_{Y|D_{\underline{n}},\log}^{q}}.$$
(8.3.23)

To prove the claim, note that the bijective map  $dlog: \mathcal{K}^M_{q,Y|D_{\underline{1}}}/p^m \to W_m \Omega^q_{Y|D_{\underline{1}},log}$  and the surjective map  $dlog: \mathcal{K}^M_{q,Y|D_{\underline{n}}} \to W_m \Omega^q_{Y|D_{\underline{n}},log}$  together imply that

$$\operatorname{dlog}: \frac{\mathcal{K}_{q,Y|D_{\underline{n}}}^{M} + p^{m}\mathcal{K}_{q,Y|D_{\underline{1}}}^{M}}{p^{m}\mathcal{K}_{q,Y|D_{\underline{1}}}^{M}} \xrightarrow{\cong} W_{m}\Omega_{Y|D_{\underline{n}},\log}^{q}$$

$$(8.3.24)$$

is an isomorphism of Nisnevich (resp. étale) sheaves for all  $m \ge 1$ . Furthermore, Image  $\phi \cong W_m \Omega_{Y|D_{\underline{n}},\log}^q$ . Now, (8.3.23) easily follows once we observe that  $p^m \mathcal{K}_{q,Y|D_{\underline{1}}}^M \subset \mathcal{K}_{q,Y|D_{\underline{n}}}^M$  for m >> 0. Next, we note that the sheaf  $\frac{W_m \Omega_{Y|D_{\underline{1}},\log}^q}{W_m \Omega_{Y|D_{\underline{n}},\log}^q}$  which is supported on E, has a filtration whose successive quotients are coherent sheaves on Y (cf. Proposition 8.2.1(1)). Hence  $H_{\text{\'et}}^d(Y, \frac{\mathcal{K}_{q,Y|D_{\underline{1}}}^M}{\mathcal{K}_{q,Y|D_{\underline{n}}}^M}) \cong H_{\text{\'et}}^d(Y, \frac{W_m \Omega_{Y|D_{\underline{1}},\log}^q}{W_m \Omega_{Y|D_{\underline{n}},\log}^q}) = 0$  for  $d \ge N$ . As the right

hand side of (8.3.23) is  $p^m$ -torsion, we equivalently get  $H^d_{\text{\acute{e}t}}(Y, p^m\left(\frac{\mathcal{K}^M_{q,Y|D_1}}{\mathcal{K}^M_{q,Y|D_n}}\right)) = 0$  for  $d \ge N, m >> 0.$  Hence,  $H^d_{\text{\acute{e}t}}(Y, \mathcal{K}^M_{q, Y|D_n}/p^m) \xrightarrow{\cong} H^d_{\text{\acute{e}t}}(Y, W_m \Omega^q_{Y|D_n, \log})$  for m >> 0.

Now we prove by decreasing induction on m that  $H^d_{\acute{e}t}(Y, {}_{p^m}\left(\frac{\mathcal{K}^M_{q,Y|D_1}}{\mathcal{K}^M_{q,Y|D_n}}\right)) = 0$ , for all  $m \ge 1$ . Fix some *m* for which we have  $p^m \left(\frac{\mathcal{K}_{q,Y|D_{\underline{1}}}^M}{\mathcal{K}_{q,Y|D_{\underline{n}}}^M}\right) = \frac{\mathcal{K}_{q,Y|D_{\underline{1}}}^M}{\mathcal{K}_{q,Y|D_{\underline{n}}}^M} = \frac{W_m \Omega_{Y|D_{\underline{1}},\log}^q}{W_m \Omega_{Y|D_{\underline{n}},\log}^q}$ . Now consider the following exact sequence (for 1 < i < m).

$$0 \rightarrow {}_{p^{i}}W_{m}\Omega^{q}_{Y|D_{\underline{n}},\log} \xrightarrow{\phi_{1}} {}_{p^{i}}W_{m}\Omega^{q}_{Y|D_{\underline{1}},\log} \rightarrow {}_{p^{i}}\left(\frac{W_{m}\Omega^{q}_{Y|D_{\underline{1}},\log}}{W_{m}\Omega^{q}_{Y|D_{\underline{n}},\log}}\right) \rightarrow$$
$$\rightarrow W_{m}\Omega^{q}_{Y|D_{\underline{n}},\log}/p^{i} \xrightarrow{\phi_{2}} W_{m}\Omega^{q}_{Y|D_{\underline{1}},\log}/p$$

Enough to show  $H^d_{\text{\'et}}(Y, \operatorname{Coker} \phi_1) = 0 = H^d_{\text{\'et}}(Y, \operatorname{Ker} \phi_2)$  for all  $d \ge N$ . To prove this, we make the following claims.

**Claim 1:**  $_{p^i}W_m\Omega^q_{Y|D_{\underline{n}},\log} \cong W_i\Omega^q_{Y|D_{\underline{\lceil n/p^{m-i}\rceil},\log}}, \text{ for } i, m \ge 1, q \ge 0 \text{ and } \underline{n} \ge \underline{1}.$ Proof: Since  $W_m\Omega^q_{Y|D_{\underline{n}},\log} \subset W_m\Omega^q_Y, \text{ and } \operatorname{Ker}(p^i : W_m\Omega^q_Y \to W_m\Omega^q_Y) = \operatorname{Ker}(R^i : W_m\Omega^q_Y \to W_{m-i}\Omega^q_Y), \text{ we conclude that } \operatorname{Ker}(p^i : W_m\Omega^q_{Y|D_{\underline{n}},\log} \to W_m\Omega^q_{Y|D_{\underline{n}},\log}) = \operatorname{Ker}(R^i : W_m\Omega^q_Y)$  $W_m \Omega^q_{Y|D_n,\log} \to W_{m-i} \Omega^q_{Y|D_n,\log}$ ). But, by Corollary 6.1.3, we have an isomorphism

$$\overline{p}^{m-i}: W_i\Omega^q_{Y|D_{[\underline{n}/p^{m-i}]}, \log} \xrightarrow{\cong} \operatorname{Ker}(R^i: W_m\Omega^q_{Y|D_{\underline{n}}, \log} \to W_{m-i}\Omega^q_{Y|D_{\underline{n}}, \log}).$$

This proves claim 1.

From this claim, we obtain Coker  $\phi_1 \cong \frac{W_i \Omega_{Y|D_{\underline{1}},\log}^q}{W_i \Omega_{Y|D_{\underline{n}},\log}^q}$  is supported on *E*. Therefore, by Proposition 8.2.1, we have  $H^d_{\text{ét}}(Y, \text{Coker } \phi_1) = 0$  for  $d \ge N$ .

Claim 2: Ker  $\phi_2 \cong W_{m-i}\Omega^q_{Y|D_{\lceil n/p^i\rceil},\log}/W_{m-i}\Omega^q_{Y|D_{\underline{n}},\log}.$ 

Proof: To see this, first we note that (as  $p = \overline{p}R$ )

$$p^{i}(W_{m}\Omega^{q}_{Y|D_{\underline{n}},\log}) = \overline{p}^{i}R^{i}(W_{m}\Omega^{q}_{Y|D_{\underline{n}},\log}) = \overline{p}^{i}(W_{m-i}\Omega^{q}_{Y|D_{\underline{n}},\log}).$$

Now let  $\alpha \in W_m \Omega^q_{Y|D_n,\log}$  be any local section such that  $\alpha = p^i \beta$ , where  $\beta \in W_m \Omega^q_{Y|D_1,\log}$  be a local section. Since  $R^i(\beta) \in W_{m-i}\Omega^q_{Y,\log}$  and  $\overline{p}^i R^i(\beta) = \alpha \in \mathrm{fil}_{D-\underline{n}} W_m \Omega^q_U$ , by repeatedly applying Lemma 4.3.14, we get  $R^i\beta \in W_{m-i}\Omega^q_{Y,\log} \cap \operatorname{fil}_{D_{-\lceil \underline{n}/p^i\rceil}}W_{m-i}\Omega^q_U = W_{m-i}\Omega^q_{Y\mid D_{\lceil n/p^i\rceil},\log}$ where the last equality follows from Lemma 6.1.2(1). So we get

$$\alpha = \overline{p}^i R^i \beta \in \overline{p}^i (W_{m-i} \Omega^q_{Y|D_{[\underline{n}/p^i]}, \log}).$$

#### CHAPTER 8. DUALITY THEOREM FOR HODGE-WITT COHOMOLOGY OVER 118 LOCAL FIELDS

This implies Ker  $\phi_2 \subset \frac{\overline{p}^i(W_{m-i}\Omega^q_{Y|D_{[\underline{n}/p^i]}}, \log)}{\overline{p}^i(W_{m-i}\Omega^q_{Y|D_{\underline{n}}, \log})}$ . Also we have  $\frac{\overline{p}^i(W_{m-i}\Omega^q_{Y|D_{[\underline{n}/p^i]}}, \log)}{\overline{p}^i(W_{m-i}\Omega^q_{Y|D_{\underline{n}}, \log})} \subset \text{Ker } \phi_2$ , because  $[\underline{n}/p^i] \geq \underline{1}$  and so  $\overline{p}^i(W_{m-i}\Omega^q_{Y|D_{[\underline{n}/p^i]}}, \log) \subset \overline{p}^i(W_{m-i}\Omega^q_{Y|D_{\underline{1}}, \log}) = p^i(W_m\Omega^q_{Y|D_{\underline{1}}, \log})$ . Now the claim 2 follows from the isomorphism

$$\frac{W_{m-i}\Omega^q_{Y|D_{[\underline{n}/p^i]},\log}}{W_{m-i}\Omega^q_{Y|D_{\underline{n}},\log}} \xrightarrow{\overline{p}^i} \frac{\overline{p}^i(W_{m-i}\Omega^q_{Y|D_{[\underline{n}/p^i]},\log})}{\overline{p}^i(W_{m-i}\Omega^q_{Y|D_{\underline{n}},\log})}.$$

By Proposition 8.2.1, Ker  $\phi_2$  has a filtration whose successive quotients are coherent sheaves supported on E. Hence  $H^d_{\text{ét}}(Y, \text{Ker } \phi_2) = 0$  for all  $d \ge N$ . This proves the proposition.

**Corollary 8.3.11.** For  $m \ge 1, \underline{n} \ge \underline{0}$ , we have an exact sequence

$$0 \to \operatorname{Pic}(X|D_{\underline{n}+1})/p^{m} \to H^{1}_{\acute{e}t}(X, W_{m}\Omega^{1}_{X|D_{\underline{n}+1}, \log}) \to {}_{p^{m}}H^{2}_{\acute{e}t}(X, \mathcal{K}^{M}_{1, X|D_{\underline{n}+1}}) \to 0.$$
(8.3.25)

*Proof.* Note that  $\operatorname{Pic}(X|D_{\underline{n}}) \cong H^1_{\operatorname{\acute{e}t}}(X, \mathcal{K}^M_{1,X|D_{\underline{n}}})$ . Now the corollary immediately follows from Proposition 8.3.10, using the exact sequence  $0 \to \mathcal{K}^M_{1,X|D_{\underline{n}+1}} \xrightarrow{p^m} \mathcal{K}^M_{1,X|D_{\underline{n}+1}} \to \mathcal{K}^M_{1,X|D_{\underline{n}+1}}/p^m \to 0$  in  $X_{\operatorname{\acute{e}t}}$ .

**Case 6** (i = 1, q = 1): We endow  $G_{\underline{n}}^{1,1}(m)$  with the structure of weakest topological abelian group such that  $\operatorname{Pic}(X|D_{\underline{n}+1})/p^m$  becomes an open subgroup of  $G_{\underline{n}}^{1,1}(m)$ in (8.3.25). As a result we see the induced quotient topology on  ${}_{p^m}H^2_{\operatorname{\acute{e}t}}(X, \mathcal{K}_{1,X|D_{\underline{n}+1}}^M)$ in (8.3.25)) is discrete (cf. [47, Lem 12.5]) and (8.3.25) becomes an exact sequence of topological abelian groups. We let

$$(G_{\underline{n}}^{1,1}(m))^{\wedge} \coloneqq H^{1}_{\text{\'et}}(X, W_{m}\Omega^{1}_{X|D_{\underline{n}+1}, \log})^{\wedge} = \varprojlim_{V} H^{1}_{\text{\'et}}(X, W_{m}\Omega^{1}_{X|D_{\underline{n}+1}, \log})/V,$$

where V varies over all open subgroups of finite index in  $\operatorname{Pic}(X|D_{\underline{n}+1})/p^m$ . Here we also claim that

$$H^2_{\text{\acute{e}t}}(X, \mathcal{K}^M_{1,X|D_{\underline{n}}}) \cong H^2_{\text{\acute{e}t}}(X, \mathcal{K}^M_{1,X|D_{\underline{n}'}})$$

$$(8.3.26)$$

for  $\underline{n} \geq \underline{n}' \geq \underline{1}$ . To see this, we consider the exact sequence  $0 \to \mathcal{K}_{1,X|D_{\underline{n}}}^{M} \to \mathcal{K}_{1,X|D_{\underline{n}}'}^{M} \to \mathcal{K}_{1,X|D_{\underline{n}}'}^{M} \to 0$  in  $X_{\text{\acute{e}t}}$ . By a similar argument to that in Proposition 8.3.10, we see that  $\frac{\mathcal{K}_{1,X|D_{\underline{n}}}^{M}}{\mathcal{K}_{1,X|D_{\underline{n}}}^{M}} \cong \frac{W_m \Omega_{X|D_{\underline{n}'},\log}^1}{W_m \Omega_{X|D_{\underline{n}},\log}^1}$ , for some m >> 0. Now, by Proposition 8.2.1, we get  $H_{\text{\acute{e}t}}^i(X, \frac{W_m \Omega_{X|D_{\underline{n}'},\log}^1}{W_m \Omega_{X|D_{\underline{n}},\log}^1}) = 0$  for i = 1, 2 and hence the claim.

A locally compact Hausdorff topological group G is called "torsion-by-profinite" if there is an exact sequence

$$0 \to G_{\rm pf} \to G \to G_{\rm dt} \to 0$$

where  $G_{pf}$  is an open and profinite subgroup of G and  $G_{dt}$  is a torsion group with the quotient topology (necessarily discrete). Note that, the Pontryagin dual  $G^* =$  $\operatorname{Hom}_{\mathbf{Tab}}(G,\mathbb{T})$  is again a "torsion-by-profinite" group. Moreover, if G is torsion, then  $G^* = \operatorname{Hom}_{\mathbf{Tab}}(G,\mathbb{Q}/\mathbb{Z})$ . We also recall that "torsion-by-profinite" groups satisfy Pontryagin duality, namely, the evaluation map  $G \to (G^*)^*$  is topological isomorphism (cf. [68, § 2.9]). The category of "torsion-by-profinite" will be denoted as **Pfd**. Note that **Pfd** contains both profinite and discrete groups.

We have the following corollary:

**Corollary 8.3.12.** For all  $\underline{n} \ge 0$ , we have the following exact sequences in Pfd.

$$0 \to \operatorname{Pic}(X|D_{\underline{n}+1})^{\operatorname{pf}}/p^{m} \to (G_{\underline{n}}^{1,1}(m))^{\wedge} \to {}_{p^{m}}H^{2}_{\acute{e}t}(X,\mathcal{K}^{M}_{1,X|D_{\underline{1}}}) \to 0, \qquad (8.3.27)$$

$$0 \to \left( {}_{p^m} H^2_{\acute{e}t}(X, \mathcal{K}^M_{1, X|D_{\underline{1}}}) \right)^* \to \left( G^{1,1}_{\underline{n}}(m) \right)^* \to \left( \operatorname{Pic}(X|D_{\underline{n}+1})^{\operatorname{pf}}/p^m \right)^* \to 0.$$
(8.3.28)

*Proof.* Note that, (8.3.25) and (8.3.26) induce an exact sequence of topological abelian groups

$$0 \to (\operatorname{Pic}(X|D_{\underline{n}+1})/p^m)/V \to H^1_{\operatorname{\acute{e}t}}(X, W_m \Omega^1_{X|D_{\underline{n}+1}, \log})/V \to {}_{p^m} H^2_{\operatorname{\acute{e}t}}(X, \mathcal{K}^M_{1, X|D_{\underline{1}}}) \to 0$$

for any open subgroup V of finite index in  $\operatorname{Pic}(X|D_{\underline{n}+1})/p^m$ . Now (8.3.27) is obtained by taking inverse limit over V and noting that  $\operatorname{Pic}(X|D_{\underline{n}+1})^{\operatorname{pf}}/p^m \cong (\operatorname{Pic}(X|D_{\underline{n}+1})/p^m)^{\operatorname{pf}}$ , which is (8.3.19).

Now by taking Pontryagin dual in (8.3.27), we get an exact sequence in Pfd

$$0 \to ({}_{p^m} H^2_{\text{\'et}}(X, \mathcal{K}^M_{1, X|D_{\underline{1}}}))^* \to ((G^{1,1}_{\underline{n}}(m))^{\wedge})^* \to (\operatorname{Pic}(X|D_{\underline{n}+1})^{\operatorname{pf}}/p^m)^* \to 0.$$

We also note that  $(\operatorname{Pic}(X|D_{\underline{n}+1})/p^m)^* = ((\operatorname{Pic}(X|D_{\underline{n}+1})/p^m)^{\mathrm{pf}})^*$  (by Lemma 8.3.3). Since  $\operatorname{Pic}(X|D_{\underline{n}+1})/p^m$  is an open subgroup of  $G_{\underline{n}}^{1,1}(m)$ , the exact sequence (8.3.28) is also obtained by taking topological dual in (8.3.25). Compairing this exact sequence with the above one, we get  $((G_{\underline{n}}^{1,1}(m))^*)^* = (G_{\underline{n}}^{1,1}(m))^*$ . Hence, (8.3.28) is also topologically exact.

By Corollary 5.2.10, we have an exact sequence

$$0 \to \operatorname{Pic}(U)/p^m \to F_{\underline{n}}^{1,1}(m) \to {}_{p^m}\operatorname{Br}(X|D_{\underline{n}}) \to 0, \qquad (8.3.29)$$

#### CHAPTER 8. DUALITY THEOREM FOR HODGE-WITT COHOMOLOGY OVER 120 LOCAL FIELDS

where we endow  $F_{\underline{n}}^{1,1}(m)$  with the structure of weakest topological group such that  $\operatorname{Pic}(U)/p^m$  becomes an open subgroup in (8.3.29). Hence (8.3.29) becomes a short exact sequence in **Pfd** (because  $\operatorname{Pic}(U)/p^m$  is a profinite group). Taking Pontryagin dual, we have the following exact sequence in **Pfd**.

$$0 \to (_{p^m} \operatorname{Br}(X|D_{\underline{n}}))^* \to (F_{\underline{n}}^{1,1}(m))^* \to (\operatorname{Pic}(U)/p^m)^* \to 0$$
(8.3.30)

Now recall from [47, Lem 12.5] that there is a pairing

$$\operatorname{Pic}(U) \times H^{2}_{\operatorname{\acute{e}t}}(X, \mathcal{K}^{M}_{1, X|D_{1}}) \to \mathbb{Q}/\mathbb{Z}$$

$$(8.3.31)$$

compatible with the left pairing of (8.3.17) and induces an isomorphism

$$H^2_{\text{\'et}}(X, \mathcal{K}^M_{1, X|D_{\underline{1}}}) \cong (\operatorname{Pic}(U))^*$$

As a result, we get the following isomorphisms of topological abelian group.

$$(1) {}_{p^m} H^2_{\text{\'et}}(X, \mathcal{K}^M_{1, X|D_1}) \xrightarrow{\cong} (\operatorname{Pic}(U)/p^m)^{\star} \quad (2) \operatorname{Pic}(U)/p^m \xrightarrow{\cong} ({}_{p^m} H^2_{\text{\'et}}(X, \mathcal{K}^M_{1, X|D_1}))^{\star},$$

$$(8.3.32)$$

where the right isomorphism is obtained from the left one by applying Pontryagin duality and noting that  $\operatorname{Pic}(U)/p^m$  is profinite.


We now consider the following diagram of exact sequences.

$$0 \longrightarrow \operatorname{Pic}(U)/p^{m} \longrightarrow F_{\underline{n}}^{1,1}(m) \longrightarrow {}_{p^{m}}\operatorname{Br}(X|D_{\underline{n}}) \longrightarrow 0 \qquad (8.3.33)$$

$$\delta_{p^{m}} \downarrow \cong \beta'_{p^{m}} \downarrow \qquad \qquad \downarrow \alpha'_{p^{m}}$$

$$0 \rightarrow ({}_{p^{m}}H_{\operatorname{\acute{e}t}}^{2}(X, \mathcal{K}_{1,X|D_{\underline{1}}}^{M}))^{\vee} \rightarrow (G_{\underline{n}}^{1,1}(m))^{\vee} \rightarrow (\operatorname{Pic}(X|D_{\underline{n}+1})/p^{m})^{\vee} \rightarrow 0,$$

where  $\delta_{p^m}$ ,  $\beta'_{p^m}$ ,  $\alpha'_{p^m}$  are the maps induced from (8.3.31), (6.2.6) and (8.3.17), respectively. Moreover,  $\delta_{p^m}$  is isomorphism by (8.3.32)(2), because  ${}_{p^m}H^2_{\text{ét}}(X, \mathcal{K}^M_{1,X|D_1})$  is discrete. We claim that the above diagram is commutative. For this consider the diagram.



#### 8.3. TOPOLOGIES ON THE COHOMOLOGY GROUPS

where the front face and the back face are the left square of (8.3.33) and (8.3.21), respectively after replacing  $(-)^*$  by  $(-)^{\vee}$  in the latter square. Note that the top floor and bottom floor are commutative by naturality of cohomology, while the right sided face is commutative by compatibility of the pairing (6.2.6). The left face is commutative by the construction of  $\delta_{p^m}$  (see [47, Lem 12.5]). Since the back face is commutative (by (8.3.21)), we conclude the front is also commutative.

Now the right square of (8.3.33) commutes because of the construction of the Brauer-Manin pairing in [47, Prop 9.4] and the following commutative diagram (cf. [47, Prop 4.4]).

where  $G_{\underline{n},x}^{1,1} = H_x^1(X_x^h, W_m \Omega_{X|D_{\underline{n}+1},\log}^1) = K_x^{\times}/(\operatorname{fil}_{n_x+1}K_x^{\times} + p^m K_x^{\times})$  (resp.  $\mathbb{Z}/p^m$ ) and  $F_{\underline{n},x}^{1,1} = \mathbb{H}_{\operatorname{\acute{e}t}}^1(X_x^h, W_m \mathcal{F}_{\underline{n}}^1) = {}_{p^m}\operatorname{fil}_{n_x}\operatorname{Br}(K_x)$ (resp.  ${}_{p^m}\operatorname{Br}(A_x) = \mathbb{Z}/p^m$ ), if  $x \in D_{\underline{n}}$  (resp.  $x \notin D_{\underline{n}}$ ) by Lemma 8.2.3 and Theorem 5.1.16, respectively. Here the top pairing is induced by the local class field theory for Brauer group (cf. [35, § 3.4]). Now we prove the main result of this section.

**Proposition 8.3.13.** The pairing  $G_{\underline{n}}^{1,1}(m) \times F_{\underline{n}}^{1,1}(m) \to \mathbb{Z}/p^m$  induces isomorphisms  $F_{\underline{n}}^{1,1}(m) \xrightarrow{\cong} (G_{\underline{n}}^{1,1}(m))^*, \quad (G_{\underline{n}}^{1,1}(m))^{\wedge} \xrightarrow{\cong} (F_{\underline{n}}^{1,1}(m))^*.$ 

*Proof.* By (8.3.32)(2), (8.3.20)(2) and the exact sequence (8.3.28), we get that the map  $\beta'_{p^m}$  in (8.3.33) is injective and its image lies in  $(G_{\underline{n}}^{1,1}(m))^*$ . Moreover, this is surjective onto  $(G_{\underline{n}}^{1,1}(m))^*$ . This follows by replacing the bottom row of (8.3.33) by (8.3.28) and applying (8.3.20) and (8.3.32) once again.

For the second one, note that  $((G_{\underline{n}}^{1,1}(m))^{\wedge})^* = (G_{\underline{n}}^{1,1}(m))^*$  (see the proof of Corollary 8.3.12). Then  $F_{\underline{n}}^{1,1}(m) \xrightarrow{\cong} ((G_{\underline{n}}^{1,1}(m))^{\wedge})^*$  is a topological isomorphism in **Pfd** (because  $\delta_{p^m}$  is topological isomorphism). Now, by taking Pontryagin dual we get the isomorphism  $(G_{\underline{n}}^{1,1}(m))^{\wedge} \xrightarrow{\cong} (F_{\underline{n}}^{1,1}(m))^*$  in **Pfd**.

Remark 8.3.14. We call  $G_{\underline{n}}^{1,1}(m)^{\wedge}$  to be the "torsion-by-profinite" completion of  $G_{\underline{n}}^{1,1}(m)$ .

Corollary 8.3.15. We have a perfect pairing of topological abelian groups

$$\lim_{\underline{n}} (G_{\underline{n}}^{1,1}(m))^{\wedge} \times H^{1}_{\acute{e}t}(U, W_{m}\Omega^{1}_{U,\log}) \to \mathbb{Z}/p^{m}$$

#### CHAPTER 8. DUALITY THEOREM FOR HODGE-WITT COHOMOLOGY OVER 122 LOCAL FIELDS

Proof. We note that  $\lim_{\underline{\to}\underline{n}} F_{\underline{n}}^{1,1}(m) = H_{\acute{e}t}^1(U, W_m \Omega_{U,\log}^1)$  by Theorem 4.3.17(12). Also, note that the map  $\operatorname{Br}(X|D_{\underline{n}}) \to \operatorname{Br}(X|D_{\underline{n}'})$  is injective if  $\underline{n'} \geq \underline{n}$ . Hence, by (8.3.20), the transition map  $\operatorname{Pic}(X|D_{\underline{n'}+1})^{\mathrm{pf}}/p^m \to \operatorname{Pic}(X|D_{\underline{n}+1})^{\mathrm{pf}}/p^m$  is topological quotient map. Also, by (8.3.27) (resp. (8.3.29)), the transition map  $(G_{\underline{n'}}^{1,1}(m))^{\wedge} \to (G_{\underline{n}}^{1,1}(m))^{\wedge}$  (resp.  $F_{\underline{n}}^{1,1}(m) \to F_{\underline{n'}}^{1,1}(m))$  is topological quotient (resp. topological injection). We endow  $\lim_{\underline{\leftarrow}\underline{n}} (G_{\underline{n}}^{1,1}(m))^{\wedge}$  (resp.  $H_{\acute{e}t}^1(U, W_m \Omega_{U,\log}^1)$ ) with inverse limit (resp. direct limit) topology. As a results we have the following exact sequences in **Pfd**.

$$0 \to \varprojlim_{\underline{n}} \operatorname{Pic}(X|D_{\underline{n}+1})^{\operatorname{pf}}/p^m \to \varprojlim_{\underline{n}}(G_{\underline{n}}^{1,1}(m))^{\wedge} \to {}_{p^m}H^2_{\operatorname{\acute{e}t}}(X, \mathcal{K}^M_{1,X|D_{\underline{1}}}) \to 0,$$
$$0 \to \operatorname{Pic}(U)/p^m \to H^1_{\operatorname{\acute{e}t}}(U, W_m\Omega^1_{U, \log}) \to {}_{p^m}\operatorname{Br}(U) \to 0.$$

Moreover, note that the canonical map  $G = \lim_{\underline{\leftarrow} \underline{n}} (G_{\underline{n}}^{1,1}(m))^{\wedge} \to (G_{\underline{n}}^{1,1}(m))^{\wedge}$  is surjective tive (because the transition map  $\operatorname{Pic}(X|D_{\underline{n}'+1})^{\operatorname{pf}}/p^m \to \operatorname{Pic}(X|D_{\underline{n}+1})^{\operatorname{pf}}/p^m$  is surjective if  $\underline{n'} \geq \underline{n}$ ). By [47, Lem 2.6] we conclude that  $\lim_{\underline{\leftarrow} \underline{n}} (G_{\underline{n}}^{1,1}(m))^* \to G^*$  is isomorphism (note  $(G_{\underline{n}}^{1,1}(m))^* = ((G_{\underline{n}}^{1,1}(m))^{\wedge})^*)$ . Now, by Proposition 8.3.13, we have topological isomorphism  $H^1_{\operatorname{\acute{e}t}}(U, W_m \Omega^1_{U, \log}) \xrightarrow{\cong} G^*$  in **Pfd**. Hence by Pontryagin duality for "torsionby-profinite" groups, we get the result.

#### 8.3.5 The duality theorem over local fields

Finally, we summarise all the duality we proved the in previous subsections.

**Theorem 8.3.16.** Let  $X, U, D_{\underline{n}}$  be as in the beginning of § 8. Then for all  $i \ge 0$  and  $\underline{n} \ge \underline{0}$ , we have perfect pairings of topological abelian groups

1.  $H^{i}_{\acute{e}t}(X, W_{m}\Omega^{j}_{X|D_{\underline{n}+1}, \log})^{\mathrm{pf}} \times \mathbb{H}^{2-i}_{\acute{e}t}(X, W_{m}\mathcal{F}^{2-j, \bullet}_{D_{\underline{n}}}) \to \mathbb{Z}/p^{m}, \ (i, j) \neq (1, 1), (2, 1);$ where  $H^{i}_{\acute{e}t}(X, W_{m}\Omega^{j}_{X|D_{n+1}, \log})$  is already profinite if  $i \neq 1$ .

2. 
$$H^{2}_{\acute{e}t}(X, W_{m}\Omega^{1}_{X|D_{n+1}, \log}) \times \mathbb{H}^{0}_{\acute{e}t}(X, W_{m}\mathcal{F}^{1, \bullet}_{D_{n}}) \to \mathbb{Z}/p^{m},$$
  
where  $H^{2}_{\acute{e}t}(X, W_{m}\Omega^{1}_{X|D_{n+1}, \log})$  has discrete topology.

3.  $\begin{aligned} H^{1}_{\acute{e}t}(X, W_{m}\Omega^{1}_{X|D_{\underline{n}+1}, \log})^{\wedge} \times \mathbb{H}^{1}_{\acute{e}t}(X, W_{m}\mathcal{F}^{1, \bullet}_{D_{\underline{n}}}) \to \mathbb{Z}/p^{m}, \\ where \ H^{1}_{\acute{e}t}(X, W_{m}\Omega^{1}_{X|D_{\underline{n}+1}, \log}) \text{ is an extension of discrete group by an adic topological} \\ group \ and \ \mathbb{H}^{1}_{\acute{e}t}(X, W_{m}\mathcal{F}^{1, \bullet}_{D_{\underline{n}}}) \text{ is in } \mathbf{Pfd} \ and \ (H^{1}_{\acute{e}t}(X, W_{m}\Omega^{1}_{X|D_{\underline{n}+1}, \log}))^{\wedge} \text{ is the "torsion} \\ by \ profinite" \ completion \ of \ H^{1}_{\acute{e}t}(X, W_{m}\Omega^{1}_{X|D_{\underline{n}+1}, \log}) \ (cf. \ Remark \ 8.3.14). \end{aligned}$ 

#### 8.3. TOPOLOGIES ON THE COHOMOLOGY GROUPS

4.  $\lim_{\stackrel{\frown}{\leftarrow}} H^{i}_{\acute{e}t}(X, W_{m}\Omega^{j}_{X|D_{\underline{n}+1}, \log})^{\wedge} \times H^{2-i}_{\acute{e}t}(U, W_{m}\Omega^{2-j}_{U, \log}) \to \mathbb{Z}/p^{m},$ where  $(-)^{\wedge}$  is the "torsion-by-profinite" completion for i = j = 1 and the profinite completion, otherwise. (Note, these groups are all zero if i = 0.)

*Proof.* Combine (8.3.2) (8.3.3), (8.3.8), (8.3.11), Corollary 8.3.6, Proposition 8.3.13. For the item (4), combine (8.3.4), (8.3.9), Corollaries 8.3.9, 8.3.15.

# Chapter 9

# Lefschetz hypersurface theorem

Lefschetz hypersurface (a.k.a the Weak Lefschetz) theorem in étale cohomology (cf. [56, Thm VI.7.1], [41]) plays an important role for studying  $\ell$ -adic étale cohomology of smooth schemes over a field. In this chapter, we shall consider the *p*-adic analogue of these theorems.

Let  $X, E, U, D = D_{\underline{n}}$  be as in § 6.1. Moreover, we also assume X is smooth projective over an F-finite base field k and we fix an embedding  $X \hookrightarrow \mathbb{P}_k^L$ . Let dim X = N. Let Y be a general smooth hypersurface section such that  $E' = Y \times_X E$  is also a simple normal crossing divisor. Let  $D' \coloneqq D \times_X Y$  and  $i \colon Y \hookrightarrow X$  denote the closed embedding. Also, let  $j' \colon U' \hookrightarrow Y$  be the inclusion of the complement of E' in Y. Also, write V for the complement of  $E \cup Y$  in X.

Main purpose of this chapter is to prove a Lefschetz theorem for the cohomology groups  $H^i_{\text{ét}}(X, W_m \Omega^q_{X, \log}), \mathbb{H}^i_{\text{ét}}(X, W_m \mathcal{F}^{q, \bullet}_D)$  and  $H^i_{\text{ét}}(X, W_m \Omega^q_{X|D, \log})$ .

### 9.1 Some exact sequences

Let  $X_{\log}$  and  $Y_{\log}$  be the log schemes whose log structures on X and Y are given by the sheaves of monoids

$$\mathcal{M}_X = \mathcal{O}_X \cap j_* \mathcal{O}_U^*$$
 and  $\mathcal{M}'_Y = \mathcal{O}_Y \cap j'_* \mathcal{O}_{U'}^*$ , respectively.

We identify  $\Omega^1_{X_{\log}}$  (resp.  $\Omega^1_{Y_{\log}}$ ) with  $\Omega^1_X(\log E)$  (resp.  $\Omega^1_Y(\log E')$ ) by Lemma 1.2.8.

**Lemma 9.1.1.** For all  $q \ge 0$ , we have the following exact sequence.

$$0 \to \mathcal{I}/\mathcal{I}^2 \otimes_{\mathcal{O}_Y} \Omega_Y^{q-1}(\log E') \xrightarrow{\overline{d}} i^* \Omega_X^q(\log E) \to \Omega_Y^q(\log E') \to 0, \tag{9.1.1}$$

where  $\mathcal{I}$  is the ideal sheaf  $\mathcal{O}_X(-Y)$ .

*Proof.* Note that  $i: Y_{\log} \to X_{\log}$  is a strict closed immersion of log schemes. This means,  $Y \to X$  is a closed immersion and  $i^* \mathcal{M}_X \xrightarrow{\cong} \mathcal{M}'_Y$  (cf. [59, p.275]). Hence by [59, Prop IV.2.3.2], we have an exact sequence

$$\mathcal{I}/\mathcal{I}^2 \xrightarrow{\overline{d}} i^* \Omega^1_X(\log E) \to \Omega^1_Y(\log E') \to 0.$$
(9.1.2)

Since X is F-finite and smooth over k, we have an exact sequence of finite type locally free sheaves :

$$0 \to f^*\Omega^1_k \to \Omega^1_X \to \Omega^1_{X/k} \to 0,$$

by [53, Thm 25.1], where  $f: X \to \operatorname{Spec} k$ . So we conclude rank  $\Omega^1_X(\log E) = \operatorname{rank} \Omega^1_X = N + \operatorname{rank} \Omega^1_k$ , where the first equality follows from the description of  $\Omega^1_A(\log \pi)$  in Lemma 1.2.8.

Similarly, we also get rank  $\Omega_Y^1(\log E') = N - 1 + \operatorname{rank} \Omega_k^1$ . Since  $\mathcal{I}/\mathcal{I}^2$  is locally free of rank one, by counting rank we get the map  $\overline{d}$  in (9.1.2) is injective. Now the lemma follows by taking higher exterior power in (9.1.2) and [24, Ex II 5.16(d)].

We refer the reader to Definition 2.1.6 for the definition of  $\operatorname{fil}_D \Omega_U^q$ .

**Corollary 9.1.2.** For any  $D \in Div_E(X)$ , we have the following exact sequences.

$$0 \to \operatorname{fil}_{D-Y}\Omega_V^q \to \operatorname{fil}_D\Omega_U^q \to i_*\operatorname{fil}_{D'}\Omega_{U'}^q \to 0, \qquad (9.1.3)$$

$$0 \to \operatorname{fil}_{-Y}\Omega^q_{V_1} \to \Omega^q_X \to i_*\Omega^q_Y \to 0, \qquad (9.1.4)$$

where  $V_1 \coloneqq X - Y$  and  $D' \coloneqq D \times_X Y$ .

*Proof.* Note that, the second exact sequence follows from (9.1.3), considering trivial log structure on X and Y (i.e. putting E to be zero divisor in (9.1.3)).

For the first one, note that (9.1.1) gives the following exact sequence of  $\mathcal{O}_X$  modules.

$$0 \to \mathcal{O}_X(-Y) \cdot \Omega^q_X(\log E) + d(\mathcal{O}_X(-Y)) \cdot \Omega^{q-1}_X(\log E) \to \Omega^q_X(\log E) \to i_*\Omega^q_Y(\log E') \to 0.$$
(9.1.5)

Now note,

$$\mathcal{O}_X(-Y) \cdot \Omega_X^q(\log E) + d(\mathcal{O}_X(-Y)) \cdot \Omega_X^{q-1}(\log E) = \Omega_X^q(\log(E+Y))(-Y)$$
  
= fil\_-Y \Over \Over U\_V^q. (9.1.6)

Thus, we obtain the following exact sequence.

$$0 \to \operatorname{fil}_{-Y}\Omega^q_V \to \Omega^q_X(\log E) \to i_*\Omega^q_Y(\log E') \to 0.$$
(9.1.7)

Now tensoring by  $\mathcal{O}_X(D)$  over  $\mathcal{O}_X$  and using Lemma 2.1.7(2), we get the result.

**Corollary 9.1.3.** For any  $D \in \text{Div}_E(X)$  and  $q \ge 0$ , we have the following exact sequences.

$$0 \to Z_1 \operatorname{fil}_{D-Y} \Omega^q_V \to Z_1 \operatorname{fil}_D \Omega^q_U \xrightarrow{\theta_D} i_* Z_1 \operatorname{fil}_{D'} \Omega^q_{U'} \to 0.$$
(9.1.8)

$$0 \to Z_1 \operatorname{fil}_{-Y} \Omega^q_{V_1} \to Z_1 \Omega^q_X \xrightarrow{\theta} i_* Z_1 \Omega^q_Y \to 0.$$
(9.1.9)

*Proof.* By Corollary 9.1.2, we have the following diagram.

Since  $Z_1 \operatorname{fil}_D \Omega_U^q = \operatorname{Ker} (d : \operatorname{fil}_D \Omega_U^q \to \operatorname{fil}_D \Omega_U^{q+1})$ , this gives the following left exact sequence.

$$0 \to Z_1 \operatorname{fil}_{D-Y} \Omega^q_V \to Z_1 \operatorname{fil}_D \Omega^q_U \xrightarrow{\theta_D} i_* Z_1 \operatorname{fil}_{D'} \Omega^q_{U'},$$

where, we note that  $i_*Z_1 \operatorname{fil}_{D'} \Omega^q_{U'} = \ker i_*d$ , because  $i_*$  is (left) exact. To show the right exactness of (9.1.8), we consider the following diagram.

$$\begin{aligned} \operatorname{fil}_{D} W_{2} \Omega_{U}^{q} & \xrightarrow{\theta_{D}} i_{*} \operatorname{fil}_{D'} W_{2} \Omega_{U'}^{q} \\ F & \downarrow & i_{*} F \\ Z_{1} \operatorname{fil}_{D} \Omega_{U}^{q} & \xrightarrow{\theta_{D}} i_{*} Z_{1} \operatorname{fil}_{D'} \Omega_{U'}^{q}. \end{aligned}$$

Note that  $i_*F$  are surjective by Lemma 4.3.3(1) and exactness of  $i_*$ . Also, the top  $\theta_D$  is surjective because locally the map is  $\operatorname{fl}_{\underline{n}}W_2\Omega^q_{(R_\pi)} \twoheadrightarrow \operatorname{fl}_{\underline{n}}W_2\Omega^q_{((R/t)_{\overline{n}})}$ , where R is a RLR and  $t, \pi, \overline{\pi}$  and  $\underline{n}$  corresponds to Y, E, E' and D, respectively. This map is surjective by definition of  $\operatorname{fl}_{\underline{n}}W_2\Omega^q_{(R_\pi)}$  (cf. Definition 2.1.6) and noting that  $\operatorname{fl}_{\underline{n}}W_2(R_\pi) \twoheadrightarrow \operatorname{fl}_{\underline{n}}W_2(R_\pi)$ . Hence, we conclude the bottom  $\theta_D$  is also surjective.

The proof of (9.1.9) is similar and hence we skip the proof.

Now recall the complexes

$$W_m \mathcal{F}_{D,U}^{q,\bullet} = \left[ Z_1 \mathrm{fil}_D W_m \Omega_U^q \xrightarrow{1-C} \mathrm{fil}_D W_m \Omega_U^q \right], W_m \mathcal{F}_X^{q,\bullet} = \left[ Z_1 W_m \Omega_X^q \xrightarrow{1-C} W_m \Omega_X^q \right],$$

where  $D \ge 0$  in  $\operatorname{Div}_E(X)$ . Here, we use the notation  $W_m \mathcal{F}_{D,U}^{q,\bullet}$  instead of  $W_m \mathcal{F}_D^{q,\bullet}$ , to keep track of the log structure corresponding to the open subscheme U. Let

$$\mathcal{N}_{D,U}^{q,\bullet} \coloneqq \left[ Z_1 \operatorname{fil}_{-D} \Omega_U^q \xrightarrow{1-C} \operatorname{fil}_{(-D)/p} \Omega_U^q \right], \quad \text{for } D \ge E.$$

By Lemma 6.1.2(6), we know  $\mathcal{N}_{D,U}^{q,\bullet}$  is quasi-isomorphic to  $\Omega_{X|D,\log}^q$ .

Then Corollaries 9.1.2 and 9.1.3 give :

**Corollary 9.1.4.** We have the following short exact sequences of complexes in  $\mathcal{D}(X_{\acute{e}t})$ .

$$0 \to W_1 \mathcal{F}_{D-Y,V}^{q,\bullet} \to W_1 \mathcal{F}_{D,U}^{q,\bullet} \to i_* W_1 \mathcal{F}_{D',U'}^{q,\bullet} \to 0$$
(9.1.11)

$$0 \to W_1 \mathcal{F}^{q,\bullet}_{-Y,V_1} \to W_1 \mathcal{F}^{q,\bullet}_X \to i_* W_1 \mathcal{F}^{q,\bullet}_Y \to 0.$$
(9.1.12)

$$0 \to \mathcal{N}_{D-Y,V}^{q,\bullet} \to \mathcal{N}_{D,U}^{q,\bullet} \to i_* \mathcal{N}_{D',U'}^{q,\bullet} \to 0.$$
(9.1.13)

In particular, we have exact triangles in  $\mathcal{D}(X_{\acute{e}t})$ .

$$W_1 \mathcal{F}^{q,\bullet}_{-Y,V_1} \to \Omega^q_{X,\log} \to i_* \Omega^q_{Y,\log}, \tag{9.1.14}$$

$$\mathcal{N}_{D-Y,V}^{q,\bullet} \to \Omega_{X|D,\log}^q \to i_* \Omega_{Y|D',\log}^q.$$
(9.1.15)

## 9.2 Weak Lefschetz for *p*-adic étale cohomology

Recall that we have morphisms of sheaves

$$W_m \mathcal{F}_{D,U}^{q,\bullet} \to i_* W_m \mathcal{F}_{D',U'}^{q,\bullet}, \ W_m \Omega_{X|D,\log}^q \to i_* W_m \Omega_{Y|D',\log}^q, \ W_m \Omega_{X,\log}^q \to i_* W_m \Omega_{Y,\log}^q.$$

These induce the following group homomorphisms.

$$\begin{split} & \mathbb{H}^{i}_{\text{\acute{e}t}}(X, W_{m}\mathcal{F}^{q, \bullet}_{D, U}) \to \mathbb{H}^{i}_{\text{\acute{e}t}}(X, i_{*}W_{m}\mathcal{F}^{q, \bullet}_{D', U'}) =^{*} \mathbb{H}^{i}_{\text{\acute{e}t}}(Y, W_{m}\mathcal{F}^{q, \bullet}_{D', U'}) \\ & H^{i}_{\text{\acute{e}t}}(X, W_{m}\Omega^{q}_{X|D, \log}) \to H^{i}_{\text{\acute{e}t}}(X, i_{*}W_{m}\Omega^{q}_{Y|D', \log}) =^{*} H^{i}_{\text{\acute{e}t}}(Y, W_{m}\Omega^{q}_{Y|D', \log}), \\ & H^{i}_{\text{\acute{e}t}}(X, W_{m}\Omega^{q}_{X, \log}) \to H^{i}_{\text{\acute{e}t}}(X, i_{*}W_{m}\Omega^{q}_{Y, \log}) =^{*} H^{i}_{\text{\acute{e}t}}(Y, W_{m}\Omega^{q}_{Y, \log}), \end{split}$$

where the equalities "=\*" hold because of exactness of  $i_*$ .

**Lemma 9.2.1.** Fix a divisor  $D \in \text{Div}_E(X)$ . Then there exists a smooth hypersurface section Y such that Y intersects E transversally. Moreover, for all  $n \ge 1$  and

- 1. for all  $i, q \ge 0$  such that  $i + q \le N 1$ , we have
  - (a)  $H^i(X, \Omega^q_X(\log E)(D nY)) = 0,$
  - (b)  $H^{i}(X, \Omega^{q}_{X}(-nY)) = 0;$
- 2. for all  $i, q \ge 0$  such that  $i + q \le N 2$ , we have
  - (a)  $H^i(X, i_*\Omega^q_V(\log E')(D nY)) = 0$
  - (b)  $H^{i}(X, i_{*}\Omega^{q}_{Y}(-nY)) = 0;$

*Proof.* Let  $\mathcal{O}_X(1)$  be the ample line bundle on X corresponding to the embedding  $X \hookrightarrow \mathbb{P}_k^L$ . For  $D \in \text{Div}_E(X)$ , there exists  $n_0 > 0$  such that

$$H^{i}(X, \Omega^{q}_{X}(\log E)(D-n)) = H^{i}(X, \Omega^{q}_{X}(-n)) = 0, \text{ for all } n \ge n_{0}, i+q \le N-1.$$
(9.2.1)

Using Serre's vanishing theorem, this is a consequence of the Grothendieck-Serre duality and the fact the  $\Omega_X^q(\log E)$ ,  $\Omega_X^q$  are locally free sheaves of finite rank (see [18, Lem 5.1]). We also note that we obtain the identities of (9.2.1) for all i, q such that  $i \leq N - 1$ . But, for the proof, we only need the condition  $i + q \leq N - 1$ .

By Bertini theorem [17, Thm 3.9, 4.5], there exists  $H \in H^0(\mathbb{P}^L_k, \mathcal{O}(n'))$  for all  $n' >> n_0$ such that  $Y = H \cap X$  and  $Y_i = H \cap E_i$  are smooth for all  $i \leq r$ . In particular, Y intersects E transversally. This Y together with (9.2.1) prove the item (1) of the lemma.

Now we set  $A_n^q \coloneqq \Omega_V^q(\log E')(D - nY)$  and  $B_n^q \coloneqq \Omega_V^q(-nY)$ , where  $n \ge 1, q \ge 0$ .

Claim:  $H^i(Y, A_n^q) = H^i(Y, B_n^q) = 0$  for all  $i + q \le N - 2$  and  $n \ge 1$ .

Proof: We show this only for  $A_n^q$ , because the other case is analogous. We proceed by induction on  $q \ge 0$ .

If q = 0 then  $A_n^q = \mathcal{O}_Y(D - nY)$ . In this case the claim is clear from the exact sequence

$$0 \to \mathcal{O}_X(D - (n+1)Y) \to \mathcal{O}_X(D - nY) \to i_*\mathcal{O}_Y(D - nY) \to 0,$$

together with (9.2.1).

Now for  $q \ge 1$ , we consider the following two exact sequences.

$$0 \to \Omega_Y^{q-1}(\log E')(D - (n+1)Y) \xrightarrow{d} i^* \Omega_X^q(\log E)(D - nY) \to \Omega_Y^q(\log E')(D - nY) \to 0,$$

$$(9.2.2)$$

$$0 \to \Omega_X^q(\log E)(D - (n+1)Y) \to \Omega_X^q(\log E)(D - nY) \to i_* i^* \Omega_X^q(\log E)(D - nY) \to 0,$$

where, the first one is obtained from Lemma 9.1.1. The second exact sequence together with (9.2.1) imply  $H^i(Y, i^*\Omega_X^q(\log E)(D-nY)) = 0$  for all  $i+q \leq N-2$ . Also, by induction hypothesis,  $H^i(Y, A_{n+1}^{q-1}) = 0$  for all  $i+q \leq N-1$ . Hence, from the top exact sequence of (9.2.2) we conclude  $H^i(Y, A_n^q) = 0$  for all  $i+q \leq N-2$ .

**Definition 9.2.2.** A smooth hypersurface section Y is called sufficiently ample with respect to X (resp. sufficiently ample with respect to (X, D)) if the condition (1.b) (resp. (1.a) for every D' such that  $|D'| \leq |D|$ ) of Lemma 9.2.1 is satisfied.

*Remark* 9.2.3. By virtue of Lemma 9.2.1, a sufficiently ample smooth hypersurface section Y w.r.t X (resp. (X, D)) always exists. Moreover, we note in the proof of the lemma that condition (2.a) (resp. (2.b)) follows from (1.a) (resp. (1.b)).

Now we prove the main theorem of this chapter. The following theorem also extends [43, Thm 3.4].

**Theorem 9.2.4.** Let X, E be as in the beginning of § 9 and  $D \ge 0$  in  $\text{Div}_E(X)$ . Let Y be a smooth hypersurface section of X and  $D' = D \times_X Y$ . If Y is sufficiently ample w.r.t X(resp. (X, D), (X, -D)), then for all m the group homomorphisms

$$H^{i}_{\acute{e}t}(X, W_{m}\Omega^{q}_{X, \log}) \to H^{i}_{\acute{e}t}(Y, W_{m}\Omega^{q}_{Y, \log}), \qquad (9.2.3)$$

$$\mathbb{H}^{i}_{\acute{e}t}(X, W_m \mathcal{F}^{q, \bullet}_{D, U}) \to \mathbb{H}^{i}_{\acute{e}t}(Y, W_m \mathcal{F}^{q, \bullet}_{D', U'}), \qquad (9.2.4)$$

and 
$$H^i_{\acute{e}t}(X, W_m \Omega^q_{X|D,\log}) \to H^i_{\acute{e}t}(Y, W_m \Omega^q_{Y|D',\log})$$
 (9.2.5)

are isomorphisms if  $i + q \le N - 2$  and injective if i + q = N - 1.

*Proof.* We shall proceed by induction on m. By Proposition 1.1.11(1), Theorem 4.3.17(11) and Corollary 6.1.3, it is enough to show, for all  $j \ge 0$ , the maps

$$H^{i}(X, \Omega^{q}_{X, \log}) \to H^{i}(Y, \Omega^{q}_{Y, \log}), \qquad (9.2.6)$$
$$\mathbb{H}^{i}_{\text{\acute{e}t}}(X, W_{1}\mathcal{F}^{q, \bullet}_{D/p^{j}, U}) \to \mathbb{H}^{i}_{\text{\acute{e}t}}(Y, W_{1}\mathcal{F}^{q, \bullet}_{D'/p^{j}, V}), \\H^{i}_{\text{\acute{e}t}}(X, \Omega^{q}_{X|[D/p^{j}], \log}) \to H^{i}_{\text{\acute{e}t}}(Y, \Omega^{q}_{Y|[D'/p^{j}], \log}), \qquad (9.2.6)$$

are bijective (resp. injective) if  $i + q \le N - 2$  (resp. i + q = N - 1).

Since  $D \ge 0$  in each of the above cases, we note that  $D/p^j = D/p^{j+1}$  and  $\lfloor D/p^j \rfloor = \lfloor D/p^{j+1} \rfloor$  for j >> 0.

In view of Corollary 9.1.4, it is enough to show, for all  $i + q \le N - 1$  and  $0 \le \overline{D} \le D$ ,

$$\mathbb{H}^{i}_{\text{ét}}(X, W_{1}\mathcal{F}^{q, \bullet}_{-Y, V_{1}}) = \mathbb{H}^{i}_{\text{ét}}(X, W_{1}\mathcal{F}^{q, \bullet}_{\overline{D}-Y, V}) = \mathbb{H}^{i}_{\text{ét}}(X, \mathcal{N}^{q, \bullet}_{\overline{D}-Y, U}) = 0.$$
(9.2.7)

We shall show the left and the middle groups are zero. Proof for the right one is analogous to that of middle one. For any  $q \ge 1$ , there exists a residue homomorphism (see the proof of [8, Lem 6.2])

$$Res^q : \Omega^q_X(\log(E+Y)) \to i_*\Omega^{q-1}_Y(\log E')$$

which is characterized by

#### 9.2. WEAK LEFSCHETZ FOR *p*-ADIC ÉTALE COHOMOLOGY

- $Res^{q}(\omega) = 0$ , if  $\omega \in \Omega^{q}_{X}(\log E)$  and
- $\operatorname{Res}^{q}(\omega \wedge \operatorname{dlog}(f)) = \overline{\omega}$ , if  $\omega \in \Omega_X^{q-1}(\log E)$  is a lift of  $\overline{\omega} \in \Omega_Y^{q-1}(\log E')$ ,

where f is a local equation of Y. Similarly, we have a residue homomorphism

$$Res^q: \Omega^q_X(\log Y) \to i_*\Omega^{q-1}_Y$$

So we have exact sequences

$$0 \to \Omega_X^q(\log E) \to \Omega_X^q(\log(E+Y)) \xrightarrow{Res^q} i_*\Omega_Y^{q-1}(\log E') \to 0, \qquad (9.2.8)$$

$$0 \to \Omega_X^q \to \Omega_X^q(\log Y) \xrightarrow{Res^q} i_* \Omega_Y^{q-1} \to 0.$$
(9.2.9)

Now applying  $-\otimes_{\mathcal{O}_X} \mathcal{O}(\overline{D} - Y)$  in (9.2.8) and  $-\otimes_{\mathcal{O}_X} \mathcal{O}(-Y)$  in (9.2.9), we get

$$0 \to \Omega_X^q (\log E)(\overline{D} - Y) \to \Omega_X^q (\log(E + Y))(\overline{D} - Y) \xrightarrow{Res^q} i_* \Omega_Y^{q-1} (\log E')(\overline{D} - Y) \to 0.$$
(9.2.10)

$$0 \to \Omega_X^q(-Y) \to \Omega_X^q(\log Y)(-Y) \xrightarrow{Res^q} i_*\Omega_Y^{q-1}(-Y) \to 0.$$
(9.2.11)

Using the exact sequences (9.2.10), (9.2.11) and hypothesis on Y, we get

$$H^{i}(X, \Omega^{q}_{X}(\log(E+Y))(\overline{D}-Y)) = 0 = H^{i}(X, \Omega^{q}_{X}(\log Y)(-Y)), \qquad (9.2.12)$$

for all  $i \in [0, N - q - 1]$  and  $0 \le \overline{D} \le D$ .

Finally, to prove (9.2.7), enough to show for all  $i \in [0, N - q - 1]$  and  $0 \le \overline{D} \le D$ ,

$$H^{i}(X, Z_{1}\Omega_{X}^{q}(\log(E+Y))(\overline{D}-Y)) = 0 = H^{i}(X, Z_{1}\Omega_{X}^{q}(\log Y)(-Y)).$$

For q = 0, this is clear by the hypothesis on Y and noting that  $Z_1 \mathcal{O}_X(\overline{D} - Y) \cong \mathcal{O}_X(\overline{D}/p - Y)$  (see 4.3.3(2)).

For  $q \ge 1$ , consider the exact sequence (cf. Lemma 4.3.4)

$$0 \to Z_1 \Omega_X^{q-1} (\log(E+Y))(\overline{D}-Y) \to \Omega_X^{q-1} (\log(E+Y))(\overline{D}-Y) \xrightarrow{d} Z_1 \Omega_X^q (\log(E+Y))(\overline{D}-Y) \xrightarrow{C} \Omega_X^q (\log(E+Y))(\overline{D}/p-Y) \to 0.$$

Now, by induction on q and (9.2.12), the claim follows and hence the theorem.

131

### 9.3 Weak Lefschetz for $\ell$ -adic étale cohomology

Before we proceed further, we review some Weak Lefschetz theorems for  $\ell$ -adic étale cohomology. Let  $\ell$  be any prime number different from p.

**Theorem 9.3.1.** Let k be either finite or 1-local or separably closed field of characteristic  $p \ge 0$ . Let X, E, E', U, U' and Y be as in the beginning of § 9. Then, for all  $j \in \mathbb{Z}$  we have the following.

- 1. The map  $H^i_{\acute{e}t}(X, \mathbb{Z}/\ell^m(j)) \to H^i_{\acute{e}t}(Y, \mathbb{Z}/\ell^m(j))$  is isomorphism if  $i \leq N-2$  and injective if i = N-1.
- 2. The map  $H^i_{\acute{e}t}(U, \mathbb{Z}/\ell^m(j)) \to H^i_{\acute{e}t}(U', \mathbb{Z}/\ell^m(j))$  is isomorphism if  $i \leq N-2$  and injective if i = N-1.
- 3. The map  $H^i_E(X, \mathbb{Z}/\ell^m(j)) \to H^i_{E'}(Y, \mathbb{Z}/\ell^m(j))$  is isomorphism if  $i \leq N-1$  and injective if i = N.

*Proof.* First we prove item (1). Let's write  $\Lambda = \mathbb{Z}/\ell^m$ . Recall there is a perfect pairing of finite abelian groups (see [31, Introduction], [56, Chapter VI,§11], [23, Thm 9.9])

$$H^{i}_{c,\text{\acute{e}t}}(X \smallsetminus Y, \Lambda(j)) \times H^{2N+e-i}_{\acute{e}t}(X \smallsetminus Y, \Lambda(N+e'-j)) \to H^{2N+e}_{c,\acute{e}t}(X \smallsetminus Y, \Lambda(N+e')) \cong \Lambda,$$
(9.3.1)

where e' = 1 if k is an 1-local field and 0 otherwise; e is the cohomological dimension of k. Note, e = 0 if k is separably closed, e = 1 if k is a finite field and e = 2 if k is an 1-local field.

Now, for the smooth hypersurface section Y in X, we look at the exact sequence

$$\cdots \to H^i_{c,\text{\acute{e}t}}(X \smallsetminus Y, \Lambda(j)) \to H^i_{\text{\acute{e}t}}(X, \Lambda(j)) \to H^i_{\text{\acute{e}t}}(Y, \Lambda(j)) \to H^{i+1}_{c,\text{\acute{e}t}}(X \smallsetminus Y, \Lambda(j)) \to \cdots.$$

By (9.3.1),  $H^i_{c,\text{\acute{e}t}}(X \smallsetminus Y, \Lambda(j)) \xrightarrow{\cong} (H^{2N+e-i}_{\text{\acute{e}t}}(X \smallsetminus Y, \Lambda(N+e'-j)))^{\vee}$ . Since,  $X \smallsetminus Y$  is affine, its  $\ell$ -cohomological dimension is  $\leq N + e$  ([56, VI, Thm 7.2]). Hence,  $H^i_{c,\text{\acute{e}t}}(X \smallsetminus Y, \Lambda(j)) = (H^{2N+e-i}_{\text{\acute{e}t}}(X \smallsetminus Y, \Lambda(N-j)))^{\vee} = 0$  if i < N. This proves item (1).

Now we shall prove item (2) and (3) simultaneously using induction on the number of components of E. We write  $E = \bigcup_{t=1}^{r} E_t$ , where r is the number of irreducible components of E. Since Y intersects E transversally (by hypothesis on Y), we have  $E' = \bigcup_{t=1}^{r} E'_t$ , where  $E'_t = E_t \cap Y$ . Now, we look at the following commutative diagram obtained by the localisation sequence.

where  $H^i_{\text{ét}}(-)$  (resp.  $H^i_*(-)$ ) is the short hand notation for  $H^i_{\text{ét}}(-, \Lambda(j))$  (resp.  $H^i_*(-, \Lambda(j))$ ) and all the vertical arrows are the canonical restriction maps.

To start the induction, let r = 1 (i.e, E and E' are irreducible). Then, we claim that item (2) and (3) are true. Indeed, by purity [56, VI, Thm 5.1], we have  $H_{\text{\acute{e}t}}^{i-2}(E, \Lambda(j-1)) \xrightarrow{\cong} H_E^i(X, \Lambda(j))$  and a similar isomorphism for E' and Y. These isomorphisms are compatible with the map  $\phi_i$  (see [10, A.2] and [54, Ch 10]). Since dim E = N - 1, by item (1), we get that  $\phi_i$  is bijective if  $i \leq N - 1$  and injective if i = N. A diagram chase in (9.3.2) shows that  $\psi_i$  is bijective if  $i \leq N - 2$  and injective if i = N - 1. This proves the claim.

Now if r > 1, we consider the following commutative diagram.

where  $\overline{E} = \bigcup_{t=2}^{r} E_t$  and  $\overline{E}' = \overline{E} \cap Y$ . We note that  $E \smallsetminus \overline{E}$  (resp.  $E' \smallsetminus \overline{E}'$ ) is a smooth divisor in  $X \smallsetminus \overline{E}$  (resp.  $Y \smallsetminus \overline{E}'$ ). So, using purity theorem again, we can canonically identify the map  $\gamma_i$  with the map  $H^{i-2}_{\text{\acute{e}t}}(E \smallsetminus \overline{E}, \Lambda(j-1)) \to H^{i-2}_{\text{\acute{e}t}}(E' \smallsetminus \overline{E}', \Lambda(j-1))$ .

Since  $E \setminus \overline{E} = E_1 \setminus (E_1 \cap \overline{E})$  and  $E_1 \cap \overline{E}$  is a simple normal crossing divisor in  $E_1$ with r-1 irreducible components, our induction hypothesis implies that item (2) of the theorem is valid if we take  $X = E_1$  and  $U = E_1 \setminus \overline{E}$ . That is,  $\gamma_i$  is bijective if  $i-2 \leq N-1-2 = N-3$  (i.e,  $i \leq N-1$ ) and injective if i = N. Also, by induction hypothesis, item (3) is valid by taking  $\overline{E}$  instead of E. Hence  $\alpha_i$  is bijective if  $i \leq N-1$ and injective if i = N. An easy diagram chase in (9.3.3) shows that  $\beta_i$  is bijective if  $i \leq N-1$  and injective if i = N. That is, item (3) is true when number of irreducible components of E is r. Moreover, note that  $\beta_i = \phi_i$ . Using item (1), a diagram chase in (9.3.2) shows that item (2) is valid if number of irreducible components of E is r. Hence, by induction, we have proved item (2) and (3) simultaneously.

## 9.4 Applications

In this section, we shall prove a Lefschetz theorem for Brauer groups, as a application of Theorem 9.2.4.

#### 9.4.1 Lefschetz for Picard groups

We need a version of Noether-Lefschetz theorem for Picard groups over an arbitrary field, which we are going to prove now. Before we proceed further, we let  $Ab_p$  be the category of *p*-primary torsion abelian groups.

**Proposition 9.4.1.** Let k be any field. Let X be a smooth geometrically connected projective variety over k such that dim  $X \ge 4$ . Let Y be a smooth hypersurface section which satisfies q = 0 case of the conditions (1.b) and (2.b) of Lemma 9.2.1 (cf. Definition 9.2.2). Then the canonical map

$$\operatorname{Pic}(X) \xrightarrow{\tau} \operatorname{Pic}(Y)$$

is injective and Coker  $\tau \in \mathbf{Ab}_{\mathbf{p}}$  (where p = char k).

*Proof.* Case 1: k is algebraically closed. In this case  $\tau$  is an isomorphism by [25, Thm IV.3.1]. One notes that the conditions (i) and (ii) of loc. cit. Thm IV.3.1 are always valid for any smooth hypersurface section Y of dim  $\geq 2$  (see Thm IV.1.5 of loc. cit. for the Leff condition). On the other hand, the condition (iii) of loc. cit. Thm IV.3.1 is also satisfied because of the q = 0 case of the condition (2.b) of Lemma 9.2.1.

**Case 2:** k is separably closed but not algebraically closed. Let  $\overline{k}$  be an algebraic closure of k. Also, let  $\overline{X} = X \times_{\text{Spec} k} \text{Spec} \overline{k}$  and  $\overline{Y} = Y \times_{\text{Spec} k} \text{Spec} \overline{k}$ . By flat base change theorem [24, Thm III.9.3], we see that  $(\overline{X}, \overline{Y})$  also satisfies the q = 0 case of the conditions (1.b) and (2.b) of the Lemma 9.2.1. Moreover, because X is geometrically connected (equivalently,  $H^0(X, \mathcal{O}_X) = k$ ), the condition (1.b) of Lemma 9.2.1 implies  $H^0(Y, \mathcal{O}_Y) = k$ , by looking at the exact sequence

$$0 = H^0(X, \mathcal{O}_X(-Y)) \to H^0(X, \mathcal{O}_X) \xrightarrow{\cong} H^0(Y, \mathcal{O}_Y) \to H^1(X, \mathcal{O}_X(-Y)) = 0.$$

In particular  $\overline{Y}$  is connected. Now, case 1 implies  $\operatorname{Pic}(\overline{X}) \xrightarrow{\overline{\tau}} \operatorname{Pic}(\overline{Y})$  is an isomorphism. We note that the maps  $\operatorname{Pic}(X) \to \operatorname{Pic}(\overline{X})$  and  $\operatorname{Pic}(Y) \to \operatorname{Pic}(\overline{Y})$  are injective (see [72]) and their cokernels are in  $\operatorname{Ab}_{\mathbf{p}}$ , if char k = p > 0. To see the latter fact, we consider the exact sequence :

$$0 \to \mathcal{O}_X^{\times} \to \mathcal{O}_{X_{k'}}^{\times} \to \mathcal{O}_{X_{k'}}^{\times} / \mathcal{O}_X^{\times} \to 0,$$

which gives the exact sequence :  $\operatorname{Pic}(X) \xrightarrow{\theta_{k'}} \operatorname{Pic}(X_{k'}) \to H^1_{\operatorname{\acute{e}t}}(X, \mathcal{O}_{X_{k'}}^{\times}/\mathcal{O}_X^{\times})$ , where k' is any finite inseparable extension of k. Since  $\mathcal{O}_{X_{k'}}^{\times}/\mathcal{O}_X^{\times}$  is annihilated by  $p^M$  for some M > 0, we get that  $H^1_{\operatorname{\acute{e}t}}(X, \mathcal{O}_{X_{k'}}^{\times}/\mathcal{O}_X^{\times})$  and hence Coker  $\theta_{k'}$  is  $p^M$ -torsion. Now, by taking

#### 9.4. APPLICATIONS

direct limit over k', we get that the cokernel of  $\theta$  and  $\theta'$  in the following diagram are in  $Ab_p$ . Then, the following diagram

$$0 \to \operatorname{Pic}(X) \xrightarrow{\theta} \operatorname{Pic}(\overline{X}) \to \operatorname{Coker} \theta \to 0$$

$$\downarrow^{\tau} \cong \downarrow^{\overline{\tau}} \qquad \downarrow^{\alpha}$$

$$0 \to \operatorname{Pic}(Y) \xrightarrow{\theta'} \operatorname{Pic}(\overline{Y}) \to \operatorname{Coker} \theta' \to 0$$

$$(9.4.1)$$

implies that there is an exact sequence

$$0 \to \operatorname{Pic}(X) \xrightarrow{\tau} \operatorname{Pic}(Y) \to \operatorname{Ker}\alpha \to 0.$$
(9.4.2)

In particular, Coker  $\tau \in \mathbf{Ab}_{\mathbf{p}}$ . This proves our assertion in Case 2.

**Case 3:** k is arbitrary. Let  $k^s$  be a separable closure of k and  $X^s = X \times_{\text{Spec } k} \text{Spec } k^s$ ,  $Y^s = Y \times_{\text{Spec } k} \text{Spec } k^s$ . By a similar argument given in case 2, we see that  $X^s$  and  $Y^s$ are geometrically connected and the pair  $(X^s, Y^s)$  satisfies q = 0 case of the conditions (1.b) and (2.b) of Lemma 9.2.1. In particular, by case 2, we have an exact sequence of  $G = \text{Gal}(k^s/k)$ -modules

$$0 \to \operatorname{Pic}(X^s) \xrightarrow{\tau^s} \operatorname{Pic}(Y^s) \to \operatorname{Coker} \tau^s \to 0,$$

where Coker  $\tau^s \in \mathbf{Ab}_{\mathbf{p}}$ . As a result, we get an exact sequence

$$0 \to \operatorname{Pic}(X^s)^G \xrightarrow{(\tau^s)^G} \operatorname{Pic}(Y^s)^G \to (\operatorname{Coker} \tau^s)^G.$$

Thus  $\operatorname{Coker}((\tau^s)^G) \in \operatorname{Ab}_p$ . Now by [12, Prop 5.4.2], we have a commutative diagram of exact sequences :

This implies  $\tau$  is injective and Coker  $\tau \in \mathbf{Ab}_{\mathbf{p}}$ . This proves the proposition.

#### 9.4.2 Lefschetz for Brauer groups

Now we prove the following Lefschetz theorem for Br(X) and  $Br^{div}(X|D)$ .

**Theorem 9.4.2.** Let k be either a finite field or 1-local field or algebraically closed field such that char k = p > 0. Let X be a smooth geometrically connected projective variety over k such that dim  $X \ge 4$ . Then, for a sufficiently ample (cf. Definition 9.2.2) hypersurface section Y (w.r.t X or (X, D)), we have

- 1.  $\operatorname{Br}(X) \xrightarrow{\cong} \operatorname{Br}(Y)$ ,
- 2.  $\operatorname{Br}^{\operatorname{div}}(X|D) \xrightarrow{\cong} \operatorname{Br}^{\operatorname{div}}(Y|D')$ , where D and D' are as in Theorem 9.2.4.

*Proof.* For the item (1), we consider the following diagram

By Theorem 9.2.4, the middle vertical arrow is isomorphism for all  $m \ge 1$ . Then, taking direct limit over m, the above diagram induces an exact sequence

$$0 \to A \otimes_{\mathbb{Z}} \mathbb{Q}_p / \mathbb{Z}_p \to \operatorname{Br}(X) \{p\} \to \operatorname{Br}(Y) \{p\} \to 0,$$

where  $A := \operatorname{Coker}(\operatorname{Pic}(X) \to \operatorname{Pic}(Y))$ . On the other hand, by Proposition 9.4.1 and Remark 9.2.3, we know that A is a torsion group. Hence  $A \otimes_{\mathbb{Z}} \mathbb{Q}_p / \mathbb{Z}_p = 0$  and  $\operatorname{Br}(X) \{p\} \xrightarrow{\cong} \operatorname{Br}(Y) \{p\}$ .

On the other hand, let  $\ell$  be a prime different from p. Then, using Theorem 9.3.1 instead of Theorem 9.2.4, a similar argument as above shows  $\operatorname{Br}(X)\{\ell\} \xrightarrow{\cong} \operatorname{Br}(Y)\{\ell\}$ . Thus, we get  $\operatorname{Br}(X) \xrightarrow{\cong} \operatorname{Br}(Y)$  (because  $\operatorname{Br}(-)$  is torsion for regular scheme).

Now we consider item (2). Recall, U and U' are the complements of E and E' in X and Y, respectively (see § 9). Using the exact sequence (see Corollary 5.2.10)

$$0 \to \operatorname{Pic}(U)/p^m \to \mathbb{H}^1_{\operatorname{\acute{e}t}}(X, W_m \mathcal{F}_D^{1, \bullet}) \to {}_{p^m} \operatorname{Br}^{\operatorname{div}}(X|D) \to 0$$

and Theorem 9.2.4, the proof of  $\operatorname{Br}^{\operatorname{div}}(X|D)\{p\} \xrightarrow{\cong} \operatorname{Br}^{\operatorname{div}}(Y|D')\{p\}$  is analogous to that of item (1). Here one also needs to use that  $(\operatorname{Coker}(\operatorname{Pic}(U) \to \operatorname{Pic}(U'))) \otimes_{\mathbb{Z}} \mathbb{Q}_p/\mathbb{Z}_p = 0$ . But this follows from the above projective case because  $\operatorname{Pic}(Y) \twoheadrightarrow \operatorname{Pic}(U')$ .

Now we consider the prime to p case. Note that  $\operatorname{Br}^{\operatorname{div}}(X|D)\{\ell\} = \operatorname{Br}(U)\{\ell\}$  and  $\operatorname{Br}^{\operatorname{div}}(Y|D')\{\ell\} = \operatorname{Br}(U')\{\ell\}$ , where  $\ell$  is a prime different from p, by Lemma 5.2.9. In this case, the theorem follows by a similar argument using Theorem 9.3.1.

Remark 9.4.3. The proof of Theorem 9.4.2 also shows that the isomorphisms  $\operatorname{Br}(X)\{p\} \xrightarrow{\cong} \operatorname{Br}(Y)\{p\}$  and  $\operatorname{Br}^{\operatorname{div}}(X|D)\{p\} \xrightarrow{\cong} \operatorname{Br}^{\operatorname{div}}(Y|D')\{p\}$  hold if we take any *F*-finite field *k* in the above theorem.

#### 9.4.3 Lefschetz in complete intersection case

In this section we shall prove a Lefschetz theorem for smooth complete intersection varieties. We recall some notations. For a field k,  $k^s$  will be its separable closure and  $\overline{k}$  will be an algebraic closure of k containing  $k^s$ . For a k-scheme X, we shall write  $X^s = X \times_{\text{Spec } k} \text{Spec } k^s$  and  $\overline{X} = X \times_{\text{Spec } k} \text{Spec } \overline{k}$ .

**Definition 9.4.4.** (cf. [2, Def 5.5]) Let X be a smooth k-scheme of dim N and  $\mathcal{L}$  be an ample line bundle on X. Then  $(X, \mathcal{L})$  is called a Kodaira pair if  $H^i(X, \Omega^q_{X/k} \otimes_{\mathcal{O}_X} \mathcal{L}^{-n}) = 0$  for all  $i + q \leq N - 1$  and  $q \geq 0, n \geq 1$ .

Here are few examples.

**Examples :** (1)  $(\mathbb{P}_k^N, \mathcal{O}(n))$  is a Kodaira pair for n > 0.

(2) If X is smooth projective over a characteristic 0 field k with an ample line bundle  $\mathcal{L}$ , then  $(X, \mathcal{L})$  is a Kodaira pair by [13, Cor 2.11].

**Lemma 9.4.5.** Let X be smooth over an F-finite field k (of char p > 0) with an ample line bundle  $\mathcal{L}$ . If  $(X, \mathcal{L})$  is a Kodaira pair then  $H^i(X, \Omega_X^q \otimes_{\mathcal{O}_X} \mathcal{L}^{-n}) = 0$  for all  $i+q \le N-1$ and  $q \ge 0, n \ge 1$ .

*Proof.* Since k is F-finite, and X is smooth over k, we have a short exact sequence of finite type locally free sheaves

$$0 \to f^*\Omega^1_k \to \Omega^1_X \to \Omega^1_{X/k} \to 0,$$

where  $f: X \to \operatorname{Spec} k$ . Then by [24, Ex II.5.16(d)], there is a decreasing filtration  $\{F^t\}_{0 \le t \le q}$  of  $\Omega^q_X \otimes_{\mathcal{O}_X} \mathcal{L}^{-n}$  such that  $F^t/F^{t+1} \cong \Lambda^t(f^*\Omega^1_k) \otimes_{\mathcal{O}_X} \Omega^{q-t}_{X/k} \otimes_{\mathcal{O}_X} \mathcal{L}^{-n} \cong \Omega^t_k \otimes_k (\Omega^{q-t}_{X/k} \otimes_{\mathcal{O}_X} \mathcal{L}^{-n})$ . Now the hypothesis on  $(X, \mathcal{L})$  implies  $H^i(X, F^t/F^{t+1}) = 0$  for all  $i + q \le N - 1$  and  $t \le q$ . Hence we get the lemma.

*Remark* 9.4.6. The above lemma implies that the condition (1.b) (and hence (2.b)) of Lemma 9.2.1 holds if  $(X, \mathcal{O}_X(Y))$  is Kodaira pair.

**Lemma 9.4.7.** Let  $X \hookrightarrow \mathbb{P}^N_k$  be any smooth complete intersection subvariety. Let Y be any hypersurface section of X. Then  $(X, \mathcal{O}_X(Y))$  is a Kodaira pair.

Proof. This follows from [2, Cor 5.8] in the following way. Since  $(\mathbb{P}_k^N, \mathcal{O}(1))$  is a Kodaira pair, loc.cit. implies  $(X, \mathcal{O}_X(1))$  is also a Kodaira pair. In particular,  $(X, \mathcal{O}_X(n))$  is also Kodaira pair for all  $n \ge 1$ . But  $\mathcal{O}_X(Y) \sim \mathcal{O}_X(n)$  for some n > 0 and we get the lemma.

**Proposition 9.4.8.** Let Y be any smooth complete intersection in  $\mathbb{P}_k^N$ . Let dim Y = d. Then we have the canonical isomorphism  $H^i_{\acute{e}t}(\mathbb{P}_k^N, W_m \Omega^q_{\mathbb{P}_k^N, \log}) \xrightarrow{\cong} H^i_{\acute{e}t}(Y, W_m \Omega^q_{Y, \log})$  for all  $i, q \ge 0$  such that  $i + q \le d - 1$ .

Proof. Since Y is smooth and complete intersection, we can write  $Y = Y_1 \cap H$ , where  $Y_1$  is smooth complete intersection in  $\mathbb{P}^N_k$  of dim d+1 and H is a smooth hypersurface section. We have seen in Theorem 9.2.4 that the condition (1.b) of Lemma 9.2.1 (taking  $X = Y_1$ ) implies the isomorphism  $H^i_{\text{ét}}(Y_1, W_m \Omega^q_{Y_1, \log}) \xrightarrow{\cong} H^i_{\text{ét}}(Y, W_m \Omega^q_{Y, \log})$ , for all  $i \leq d - q - 1$ . Hence, to prove the isomorphism, it is enough to show that  $(Y_1, \mathcal{O}_{Y_1}(H))$  is a Kodaira pair (by Remark 9.4.6). But this follows from Lemma 9.4.7. Thus we have proven the isomorphism  $H^i_{\text{ét}}(Y_1, W_m \Omega^q_{Y_1, \log}) \xrightarrow{\cong} H^i_{\text{ét}}(Y, W_m \Omega^q_{Y, \log})$ , for all  $i \leq d - q - 1$ . Now, inductively, repeating the same argument for  $Y_1$  we can prove the proposition.

The following corollary is a generalization of [12, Cor. 5.5.4], where the result is proven for prime to p-part of the Brauer Groups.

**Corollary 9.4.9.** Let k be any field. If Y is a smooth complete intersection in  $\mathbb{P}_k^N$  of dim  $d \ge 3$ , then  $\operatorname{Br}(k) \cong \operatorname{Br}(Y)$ .

*Proof.* By [12, Cor 5.5.4], one knows  $Br(k) \cong Br_1(Y)$ , where  $Br_1(Y) = Ker(Br(Y) \rightarrow Br(Y^s))$ . Also it was proven in loc. cit. that  $Br(Y^s)\{\ell\} = 0$  for all  $\ell$  coprime to p. So, it is enough to show  $Br(Y^s)\{p\} = 0$ .

We claim that  $\operatorname{Br}(Y^s) \hookrightarrow \operatorname{Br}(\overline{Y})$ . By [12, Thm 5.2.5], it is enough to show  $Y^s$  is geometrically connected and  $H^1(Y^s, \mathcal{O}_{Y^s}) = 0$ . This can be seen by using decreasing induction on dim  $Y^s$ . Note that  $Y^s$  is a smooth complete intersection in  $\mathbb{P}_{k^s}^N$  of dim dand hence is geometrically connected. If d = N (i.e,  $Y^s = \mathbb{P}_{k^s}^N$ ), then the vanishing of  $H^1(Y^s, \mathcal{O}_{Y^s})$  is clear. If d < N, we can write  $Y^s = Y_1 \cap H$  as in Proposition 9.4.8, where  $Y_1$  is smooth complete intersection of dim d + 1 in  $\mathbb{P}_{k^s}^N$  and H is a smooth hypersurface section. Then

$$0 \to \mathcal{O}_{Y_1}(-H) \to \mathcal{O}_{Y_1} \to \mathcal{O}_Y \to 0$$

induces an exact sequence

$$H^1(Y_1, \mathcal{O}_{Y_1}) \to H^1(Y, \mathcal{O}_Y) \to H^2(Y_1, \mathcal{O}_{Y_1}(-H)).$$

By induction hypothesis,  $H^1(Y_1, \mathcal{O}_{Y_1}) = 0$ . Also, note that  $H^2(Y_1, \mathcal{O}_{Y_1}(-H)) = 0$ , because  $(Y_1, \mathcal{O}_{Y_1}(H))$  is a Kodaira pair (by Lemma 9.4.7) and dim  $Y_1 = d + 1 \ge 4$ . Hence the claim is proved.

Now, to prove the corollary, it is enough to show  $\operatorname{Br}(\overline{Y})\{p\} = 0$ . For that we consider the diagram (9.4.3), replacing X and Y by  $\mathbb{P}_{\overline{k}}^{N}$  and  $\overline{Y}$ , respectively. By [20,

XII, Cor 3.7], we have  $\operatorname{Pic}(\mathbb{P}_{\overline{k}}^{N}) \xrightarrow{\cong} \operatorname{Pic}(\overline{Y})$ . Hence, using Proposition 9.4.8, we get  $\operatorname{Br}(\mathbb{P}_{\overline{k}}^{N})\{p\} \cong \operatorname{Br}(\overline{Y})\{p\}$ . But  $\operatorname{Br}(\mathbb{P}_{\overline{k}}^{N}) = \operatorname{Br}(\overline{k}) = 0$  by [12, Thm 6.1.3]. Hence the claim is proved.

# Bibliography

- A. Abbes, T. Saito, Ramification of local fields with imperfect residue fields, Amer. J. Math., **124**, (2002), no. 5, 879–920.
- [2] B. Antieau, B. Bhatt and A. Mathew, Counterexamples to Hochschild-Kostant-Rosenberg in characteristic p, Forum of Mathematics, Sigma (2021), 9, 1–26. 137
- [3] M. André, Localisation de la lissité formelle, Manuscripta Math, 13, (1974), 297-307. 19
- [4] M. Artin, A. Grothendieck and J.-L. Verdier, *Théorie des topos et cohomologie étale des schémas*, Lecture Notes in Math. 269, 270. 11
- [5] M. Auslander, O. Goldman, The Brauer group of a commutative ring, Trans. Amer. Math. Soc., vol. 97, no. 3, 1960, pp. 367–409.
- [6] S. Bloch, K. Kato, *p*-adic étale cohomology, Publ. Math. IHES, 63, (1986), 107–152.
   70
- [7] W. Bruns, J. Herzog, Cohen-Macaulay Rings, Cambridge University Press, Cambridge, Revised Edition 1998. 78, 98
- [8] F. Binda, S. Saito, Relative cycles with moduli and regulator maps, J. of the Inst. of Math. Jussieu 18 (2019), no. 6, 1233–1293. 130
- [9] J.-P. Brylinski, Théorie du corps de classes de Kato et revetments abéliens de surfaces, Ann. Inst. Fourier (Grenoble), 33, (1983), 23-38. 2, 66
- [10] D.-C. Cisinski, F. Déglise, Étale motives, Compos. Math., 152, (2016), 556–666.
   133
- [11] J.-L. Colliot-Thélène, J.-J. Sansuc, C. Soulé, Torsion dans le groupe de Chow de codimension deux Duke Math. J., 50, (1983), 763–801. 14, 85, 87

- [12] J.-L. Colliot-Thélène, A. Skorobogatov, The Brauer-Grothendieck group, Ergebnisse der Mathematik und ihrer Grenzgebiete 3, Folge. A Series of Modern Surveys in Mathematics, 71, Springer, Berlin, 2021. 1, 9, 114, 135, 138, 139
- [13] P. Deligne and L. Illusie, 'Relèvements modulo p<sup>2</sup> et décomposition du complexe de de Rham', Invent. Math. 89(2) (1987), 247–270 137
- T. Geisser, L. Hesselholt, The de Rham-Witt complex and p-adic vanishing cycles, J. Amer. Math. Soc., 19, (2006), 1–36. 16, 84
- [15] T. Geisser, L. Hesselholt, On the K-theory of complete regular local F<sub>p</sub>-algebras, Topology, 45, (2006), 475–493. 33, 40, 42, 51
- [16] T. Geisser, M. Levine, The K-theory of fields in characteristic p, Invent. Math., 139, (2000), 459–493.
- [17] M. Ghosh, A. Krishna, Bertini theorems revisited J. London Math. Soc., 108, (2023), 1163-1192. 129
- [18] M. Ghosh, A. Krishna, Zero-cycles on normal projective varieties. Journal of the Institute of Mathematics of Jussieu; 22(5), (2023), 2241-2295. 129
- [19] A. Grothendieck, SGA1 (1960-61), Lecture Notes in Mathematics-Vol. 224, SpringerVerlag, New York, 1971.
- [20] A. Grothendieck, et al. Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2). Sém. géométrie algébrique du Bois-Marie, 1962. Augmenté d'un exposé de Michèle Raynaud. Documents Math'ematiques 4, Soc. Math. France, 2005./ 8, 139
- [21] R. Gupta, A. Krishna, Duality theorem and applications to Brauer groups over finite fields, (2024). 91
- [22] R. Gupta, A. Krishna, Ramified class field theory and duality over finite fields, arXiv:2104.03029v1 [math.AG], (2021). 4, 44, 63, 88, 114
- [23] R. Gupta, A. Krishna, J. Rathore, Tame class field theory over local fields, arXiv:2209.02953, [math.AG], (2022). 132
- [24] R. Hartshorne, Algebraic Geometry, Graduate Text in Mathematics, 52, (1997), Springer-Verlag. 126, 134, 137

- [25] R. Hartshorne, Ample subvarieties of algebraic varieties. Berlin, Springer, 1970. 134
- [26] L. Hesselholt, The big de Rham-Witt complex, Acta Math., 214, (2015), 135–207.
   35, 49
- [27] T. Hiranouchi, Class field theory for open curves over local fields, J. Théor. Nombres Bordeaux, 30, no 2, (2018), 501–524. 8
- [28] L. Hesselholt, I. Madsen, On the K-theory of local fields, Ann. Math., 158, (2003), 1-113. 16, 21, 22
- [29] O. Hyodo, K. Kato, Semi-stable reduction and crystalline cohomology with logarithmic poles, Astérisque, 223, (1994), 221–268. 4, 6, 20
- [30] L. Illusie, Complexe de de Rham-Witt et cohomologie cristalline, Ann. Sci. Ec. Norm. Supér., (4) 12, (1979), 501–661. 4, 12, 13, 14, 35, 56, 63
- [31] U. Jannsen, S. Saito, Y. Zhao, Duality for relative logarithmic de Rham-Witt sheaves and wildly ramified class field theory over finite fields, Comp. Math., 154, (2008), 1306–1331. i, 4, 5, 6, 83, 84, 85, 86, 87, 89, 90, 91, 102, 105, 132
- [32] K. Kato, Duality theorems for the p-primary étale cohomology I, Algebraic and Topological Theories (Kinosaki, 1984), Kinokuniya-shoten, (1985), 127–148. 4, 44, 56
- [33] K. Kato, Milnor K-theory and Chow group of zero cycles, Applications of algebraic K-theory to algebraic geometry and number theory, Contemp. Math., 55, Amer. Math. Soc, Providence, RI, (1986), 241–255.
- [34] K. Kato, A generalization of local class field theory by using K-groups. I, J. Fac. Sci., Univ. Tokyo, 27, (1980), 303–376. 6, 93, 94, 96
- [35] K. Kato, A generalization of local class field theory by using K-groups. II, J. Fac. Sci., Univ. Tokyo, Sect. I A 27, (1980), no. 3, 603–683. vi, 6, 93, 100, 121
- [36] K. Kato, Swan conductors for characters of degree one in the imperfect residue field case, in Algebraic K-Theory and Algebraic Number Theory (Honolulu, Hawaii, 1987), Contemp. Math., 83, Amer. Math. Soc., Providence, (1989), 101–131. i, 2, 6, 64, 66, 67, 68, 69, 70, 72, 80, 97, 100

- [37] K. Kato, Existence theorem for higher local fields, In: Invitation to higher local fields, Geometry & Topology Monographs, 3, (2000), 165–195. 6, 93, 97, 100
- [38] K. Kato, Residue Homomorphims in Milnor K-theory Advanced Studies in Pure Mathematics, Galois Groups and their Representations, 2, (1983), 153–172. 6, 97
- [39] K. Kato, Logarithmic structures of Fontaine-Illusie, in Algebraic Analysis, Geometry, and Number Theory, J.-I. Igusa ed., 1988, Johns Hopkins University, pp. 191–224. 15, 16, 20
- [40] K. Kato, S. Saito, Unramified class field theory of arithmetic surfaces, Ann. of Math., 118, No. 2, (1983), 241–275. 7, 101, 103, 108, 112, 113
- [41] N. Katz, Applications of the weak Lefschetz theorem. Appendix to: B. Poonen and J.F. Voloch, Random Diophantine equations, Arithmetic of higherdimensional algebraic varieties, Progress in Math. 226 Birkhauser, 2004, pp. 175-184. 125
- [42] M. Kerz, Milnor K-theory of local rings with finite residue fields, J. Alg. Geom., 19, (2010), 173–191. 61, 63
- [43] M. Kerz, S. Saito, Lefschetz theorem for abelian fundamental group with modulus, Algebra & Number Theory, 8, (2014), 689–701. Correction to the article Lefschetz theorem for abelian fundamental group with modulus, Algebra & Number Theory, 16, (2022), 2001-2003. 2, 8, 67, 73, 74, 130
- [44] M. Kerz, S. Saito, Chow group of 0-cycles with modulus and higher-dimensional class field theory, Duke Math. J., 165, (2016), 2811–2897. 2
- [45] M. Kerz, Y. Zhao, Higher ideles and class field theory, Nagoya Math. J., 236, (2019), 214–250. 87, 102, 116
- [46] A. Krishna, J. Park, De Rham-Witt sheaves via algebraic cycles, Comp. Math., 157, (2021), 2089–2132. 17, 49
- [47] A. Krishna, J. Rathore, S. Sadhukhan, Duality for cohomology of split tori on curves over local fields, arXiv:2302.06069, [math.AG], (2023).
  7, 80, 93, 101, 103, 104, 108, 109, 110, 112, 113, 114, 115, 118, 120, 121, 122
- [48] A. Krishna, S. Majumder, Class field theory for curves over local fields, arXiv:2307.15416, [math.AG], (2023). 8

- [49] A. Krishna, S. Majumder, Class field theory for curves over higher local fields, In preparation, (2025). i, 9
- [50] A. Krishna, S. Majumder, Some applications of the filtered de Rham-Witt complex, In preparation, (2025). i, 9
- [51] P. Lorenzon, Logarithmic Hodge-Witt Forms and Hyodo-Kato Cohomology, J. Algebra, 249, (2002), 247–265. 4, 13, 21
- [52] S. Matsuda, On the Swan conductors in positive characteristic, Amer. J. Math., 119, (1997), 705–739.
- [53] H. Matsumura, Commutative ring theory, Cambridge studies in advanced mathematics, 8, Cambridge University Press, 1986. 18, 50, 93, 126
- [54] C. Mazza, V. Voevodsky, C. Weibel, Lecture notes on motivic cohomology, Clay Mathematics Monographs, 2, American Mathematical Society, Providence, (2006) 133
- [55] J. Milne, Duality in the flat cohomology of a surface, Ann. Sci. Ec. Norm. Supér.,
  (4), 9, (1976), 171–201. 88, 90, 103
- [56] J. Milne, Etale Cohomology, Princeton Math. Ser., 33, Princeton University Press, 1980. 63, 108, 125, 132, 133
- [57] J. Milne, Values of zeta functions of varieties over finite fields, Amer. J. Math., 108, (1986), 297–360.
- [58] M. Morrow, K-theory and logarithmic Hodge-Witt sheaves of formal schemes in characteristic p, Ann. Sci. Éc. Norm. Supér., (4), 52, no. 6, (2019), 1537–1601. 14, 18, 28, 40, 48, 49, 51, 63
- [59] A. Ogus, Lectures on Logarithmic Algebraic Geometry, Cambridge studies in advanced mathematics, 178, Cambridge University Press, 2018. 126
- [60] D. Popescu, General Neron Desingularization and approximation, Nagoya Math. J., 104, (1986), 85–115. 11
- [61] J.-P. Serre, Local fields, Graduate Texts in Mathematics, 67, Springer, (1979). 2
- [62] F. Ren, K. Rülling, Duality for Hodge-Witt cohomology with modulus, arXiv:2403.18763, [math.AG], (2024). 4

- [63] A. Shiho, On logarithmic Hodge-Witt cohomology of regular schemes, J. Math. Sci. Univ. Tokyo, 14, (2007), 567–635. 11, 12, 14, 105
- [64] A. Shiho, Crystalline fundamental groups. I. Isocrystals on log crystalline site and log convergent site, J. Math. Sci. Univ. Tokyo, 7, (2000), no.4, 509–656. 15, 16
- [65] H. Tanimoto, Smoothness of Noetherian rings, Nagoya Math. J., 95, (1984), 163– 179.
- [66] A. Tyc, Differential basis, p-basis and smoothness in characteristic p > 0, Proc. AMS, 103, (1988), 389–394. 18, 90
- [67] Panin, I., The equi-characteristic case of the Gersten conjecture, Proc. Steklov Inst. of Math. 241, (2003), 154–163 11
- [68] L. Ribes, P. Zalesskii, Profinite Groups, 2nd, Ergebnisse der Mathematik und iher Grenzgibiete, 40, Springer, (2010). 95, 119
- [69] K. Rülling, S. Saito, Higher Chow groups with modulus and relative Milnor Ktheory, Trans. Amer. Math. Soc., 370, (2018), 987–1043.
- [70] S. Saito, Arithmetic on two dimensional local rings, Invent. Math., 85, (1986), 379–414. 115
- [71] J. Gamst, K. Hoechsmann, Products in Sheaf-Cohomology, Tohôku Math. J., 22, (1970), 143-162. 103
- [72] Stacks project authors, The Stacks project, Lemma 0CC5(2024) 134
- [73] A. Suslin, V. Voevodsky, Singular homology of abstract algebraic varieties, Invent. Math., 123, (1996), 61–94. 114
- [74] Y. Zhao, Duality for relative logarithmic de Rham-Witt sheaves on semistable schemes over  $\mathbb{F}_q[[t]]$ , Doc. Math., **23**, (2018), 1925–1967. **18**, 63, 101
- [75] S. Saito, Class field theory for curves over local fields, J. Number Theory 21, (1985), no. 1, p. 44–80. 8
- [76] S. Bloch, Algebraic K-theory and class field theory for arithmetic surfaces, Ann. Math., 114, (1981), 229–265.
- [77] T. Yoshida, Finiteness theorems in the class field theory of varieties over local fields, J. Number Theory 101, (2003), no. 1, p. 138–150.