ASYMPTOTICS FOR THE MAGNETIC DIRICHLET-TO-NEUMANN EIGENVALUES IN GENERAL DOMAINS

BERNARD HELFFER, AYMAN KACHMAR, AND FRANÇOIS NICOLEAU

ABSTRACT. Inspired by a paper by T. Chakradhar, K. Gittins, G. Habib and N. Peyerimhoff, we analyze their conjecture that the ground state energy of the magnetic Dirichlet-to-Neumann operator tends to infinity as the magnetic field tends to infinity. More precisely, we prove refined conjectures for general two dimensional domains, based on the analysis in the case of the half-plane and the disk by two of us (B.H. and F.N.). We also extend our analysis to the three dimensional case, and explore a connection with the eigenvalue asymptotics of the magnetic Robin Laplacian.

Keywords: Magnetic Dirichlet to Neumann operator, Dirichlet and Robin eigenvalues, Eigenvalue asymptotics.

2020 Mathematics Subject Classification. Primary 58J50, Secondary 35P20.

1. INTRODUCTION

1.1. The magnetic Dirichlet to Neumann operator. This paper is a continuation of [17] where the ground state energy of the Dirichlet to Neumann operator in the case with a constant magnetic field in the unit disk $D(0,1) \subset \mathbb{R}^2$ was studied. Here, we extend this problem in the case of general bounded domains in \mathbb{R}^n (n = 2, 3).

Let Ω be a bounded connected subset of \mathbb{R}^n , with smooth boundary $\partial\Omega$ consisting of a finite number of connected components; in short we say that Ω is a regular domain of \mathbb{R}^n . For any $u \in \mathcal{D}'(\Omega)$, the magnetic Schrödinger operator on Ω is defined as

(1.1)
$$H_A u = (D - A)^2 u = -\Delta u - 2i A \cdot \nabla u + (|A|^2 - i \operatorname{div} A)u,$$

where $D = -i\nabla$, $-\Delta$ is the usual positive Laplace operator on \mathbb{R}^n and $A = \sum_{j=1}^n A_j dx_j$ is the

1-form magnetic potential. We often identify the 1-form magnetic potential A with the vector field $\overrightarrow{A} = (A_1, ..., A_n)$, and we assume that $\overrightarrow{A} \in C^{\infty}(\overline{\Omega}; \mathbb{R}^n)$. The magnetic field is given by the 2-form B = dA. We will also use the notation $\overrightarrow{H} = \operatorname{curl} \overrightarrow{A}$.

Since zero does not belong to the spectrum of the Dirichlet realization of H_A , the boundary value problem

(1.2)
$$\begin{cases} H_A \ u = 0 & \text{in } \Omega, \\ u_{|\partial\Omega} = f \in H^{1/2}(\partial\Omega), \end{cases}$$

Date: January 15, 2025.

has a unique solution $u \in H^1(\Omega)$, that we call the magnetic harmonic extension of f. The Dirichlet to Neumann map, (in what follows D-to-N map), is defined by

(1.3)
$$\begin{array}{cccc} \Lambda_A : H^{1/2}(\partial\Omega) &\longmapsto & H^{-1/2}(\partial\Omega) \\ f &\longmapsto & (\partial_{\nu} u + i \langle A, \vec{\nu} \rangle \ u)_{|\partial\Omega} \,, \end{array}$$

where $\vec{\nu}$ is the outward normal unit vector field on $\partial\Omega$. More precisely, we define the D-to-N map using the equivalent weak formulation :

(1.4)
$$\langle \Lambda_A f, g \rangle_{H^{-1/2}(\partial\Omega) \times H^{1/2}(\partial\Omega)} = \int_{\Omega} \langle (-i\nabla - A)u, (-i\nabla - A)v \rangle \, dx \,,$$

for any $g \in H^{1/2}(\partial\Omega)$ and $f \in H^{1/2}(\partial\Omega)$ such that u is the unique solution of (1.2) and v is any element of $H^1(\Omega)$ so that $v_{|\partial\Omega} = g$. Clearly, the D-to-N map is a positive operator.

We recall that since Ω is assumed to be bounded the spectrum of the D-to-N operator is discrete and is given by an increasing sequence of eigenvalues

$$(1.5) 0 \le \mu_1 \le \mu_2 \le \dots \le \mu_n \le \dots$$

which tends to $+\infty$.

Due to the identity in (1.4), the D-to-N lowest eigenvalue $\mu_1 := \lambda_1^{\text{DN}}(A, \Omega)$ can be expressed in the variational form as

(1.6)
$$\lambda_1^{\mathrm{DN}}(A,\Omega) = \inf_{u \in C^{\infty}(\overline{\Omega}), \|u\|_{\partial\Omega} = 1} \|(-i\nabla - A)u\|_{\Omega}^2,$$

and the eigenvalues $\mu_j := \lambda_j^{\text{DN}}(A, \Omega)$ can be expressed by the min-max principle,

(1.7)
$$\lambda_{j}^{\mathrm{DN}}(A,\Omega) = \inf_{\substack{M \subset C^{\infty}(\overline{\Omega}) \\ \dim(M) = j}} \left(\max_{u \in M, \|u\|_{\partial\Omega} = 1} \|(-i\nabla - A)u\|_{\Omega}^{2} \right),$$

where $\|\cdot\|_{\Omega}$ and $\|\cdot\|_{\partial\Omega}$ denote the L^2 -norms in $L^2(\Omega; \mathbb{C})$ and $L^2(\partial\Omega; \mathbb{C})$ respectively. For j = 1, we simply write $\lambda^{\text{DN}}(A, \Omega)$ instead of $\lambda_1^{\text{DN}}(A, \Omega)$.

1.2. **Planar domains.** In the two dimensional case, we prove accurate asymptotics for the lowest eigenvalue of the magnetic D-to-N operator.

Our first result concerns the constant magnetic field.

Theorem 1.1. Let Ω be a regular domain in \mathbb{R}^2 and A be a magnetic potential with constant magnetic field with norm 1. Then the ground state energy of the D-to-N map Λ_{bA} satisfies

(1.8)
$$\lim_{b \to +\infty} b^{-1/2} \lambda^{\mathrm{DN}}(bA, \Omega) = \hat{\alpha} := \frac{\alpha}{\sqrt{2}},$$

where $-\alpha$ is the unique negative zero of the parabolic cylindrical function $D_{1/2}(z)$.

This theorem was conjectured in [17]. We recall that, for $\nu \in \mathbb{R}$, the parabolic cylinder function

 $D_{\nu}(z)$ is the (normalized) solution of the differential equation

(1.9)
$$w'' + \left(\nu + \frac{1}{2} - \frac{z^2}{4}\right) w = 0,$$

which tends to 0 as $z \to +\infty$. More precisely, $D_{\nu}(z)$ has the following asymptotic expansion

(1.10)
$$D_{\nu}(z) = e^{-\frac{z^2}{4}} z^{\nu} \left(1 + \mathcal{O}\left(\frac{1}{z^2}\right)\right), \ z \to +\infty.$$

We refer to ([17], Section 2) for more details on the parabolic cylinder functions. At last, the positive real α appearing in Theorem 1.1 is approximately equal to

(1.11)
$$\alpha = 0.7649508673...$$

One can actually get a two-terms asymptotics where the second term takes account of the curvature of the boundary. The following result is a generalization of [17, Theorem 1.1], in the case of the disk.

Theorem 1.2. Let Ω be a regular domain in \mathbb{R}^2 and A be a magnetic potential with constant magnetic field with norm 1. Then the ground state energy of the D-to-N map Λ_{bA}^{DN} satisfies

$$\lambda^{\mathrm{DN}}(bA,\Omega) = \hat{\alpha}b^{\frac{1}{2}} - \frac{\hat{\alpha}^2 + 1}{3} \max_{x \in \partial\Omega} \kappa_x + o(1),$$

where κ_x denotes the curvature at x.

We actually prove that the asymptotics in Theorem 1.2 holds for the j'th eigenvalue, for every fixed $j \geq 2$, and when the magnetic field is only supposed to be constant in a neighborhood of $\partial\Omega$. If Ω is simply connected, the following inequality holds [31]

$$\max_{x\in\partial\Omega}\kappa_x\geq\sqrt{\pi/|\Omega|}\,,$$

and we obtain as corollary of Theorem 1.2:

Corollary 1.3. Let Ω be a regular domain in \mathbb{R}^2 and A be a magnetic potential with constant magnetic field with norm 1. Suppose that Ω is simply connected and \mathcal{B} is a disk with the same area as Ω . Then there exists $b_1(\Omega) > 0$ such that, for all $b \ge b_1(\Omega)$,

$$\lambda^{\mathrm{DN}}(bA,\Omega) \leq \lambda^{\mathrm{DN}}(bA,\mathcal{B}).$$

The inequality in Corollary 1.3 is reminiscent of an inequality for the magnetic Laplacian [8], and it would be interesting to investigate whether it holds for all b > 0. For the magnetic Laplacian, there is progress in the study of this question [5, 20, 21]. The geometric isoperimetric inequality also yields that $\lambda^{\text{DN}}(bA, \Omega) \leq \lambda^{\text{DN}}(bA, \mathcal{B}_*)$, where \mathcal{B}_* is a disk with the same perimeter as Ω .

If the magnetic field is variable, not vanishing in $\overline{\Omega}$ and constant along the boundary one could expect a more general result in the spirit of the one of N. Raymond [32] devoted to the ground state energy of the Neumann magnetic Laplacian. The second term will also involve the normal derivative of the magnetic field to the boundary.

We will also consider the case of variable magnetic field in 2D and in 3D in the same spirit as for the analysis of the Neumann problem appearing in surface superconductivity [26, 13, 32, 33, 14].

We prove in particular the following theorem (we refer to [15, 14, 27] for the Neumann problem).

Theorem 1.4. Let Ω be a regular domain in \mathbb{R}^2 , A be a magnetic potential with non vanishing magnetic field B(x) in $\overline{\Omega}$, then the ground state energy of the D-to-N map Λ_{bA} satisfies

(1.12)
$$\lambda^{\text{DN}}(bA,\Omega) = \hat{\alpha} (\inf_{x \in \partial\Omega} |B(x)|)^{\frac{1}{2}} b^{\frac{1}{2}} + o(b^{\frac{1}{2}}).$$

Remark 1.5. We will prove a more general result valid for a larger class of magnetic fields and for the low-lying eigenvalues. Actually, Theorem 1.4 is still true if the magnetic field B does not vanish on $\partial\Omega$ and if the set $\mathcal{Z}(B) := \{x \in \Omega : B(x) = 0\}$ consists of a finite number of smooth curves such that $|\nabla B| > 0$ on $\mathcal{Z}(B)$. See Assumption 4.1 and Example 1 for other conditions. Interestingly, only the values of the magnetic field on the boundary contributes to the main term in the asymptotics for the D-to-N operator. In fact, unlike the Neumann magnetic Laplacian, there is no contribution involving $\inf_{x \in \Omega} |B(x)|$.

1.3. Three dimensional case. We have a similar result for variable magnetic fields in 3D which is in correspondence with known results obtained in the analysis of the ground state energy of the Neumann realization of the magnetic Laplacian (see [27, 14, 33]):

Theorem 1.6. Let Ω be a regular bounded domain in \mathbb{R}^3 , A be a magnetic potential with non vanishing magnetic field B(x) in $\overline{\Omega}$, then the ground state energy of the D-to-N map Λ_{bA}^{DN} satisfies

(1.13)
$$\lim_{b \to +\infty} b^{-1/2} \lambda^{\mathrm{DN}}(bA, \Omega) = \inf_{x \in \partial\Omega} \left(\lambda^{\mathrm{DN}}(\vartheta(x)) |B(x)|^{\frac{1}{2}} \right),$$

where, for $x \in \partial \Omega$,

• $\vartheta(x)$ is defined by

$$\langle \vec{H}(x) | \vec{\nu} \rangle = -|B(x)| \sin \vartheta(x)$$
.

- $\vec{H}(x)$ is the magnetic vector field associated with B(x) considered as a 2-form by the Hodge-map.
- $\vec{\nu}$ is the exterior normal at $x \in \partial \Omega$.
- $\lambda^{\text{DN}}(\vartheta)$ is the ground state energy (see (6.3)) relative to the half space when the magnetic field is constant.

There are two important consequences of Theorem 1.6:

• When B is constant with magnitude 1, it follows that

$$\lim_{b \to +\infty} b^{-1/2} \lambda^{\mathrm{DN}}(bA, \Omega) = \inf_{x \in \partial\Omega} \lambda^{\mathrm{DN}}(\vartheta(x)),$$

which is consistent with Theorem 1.1 for 2D domains.

• More generally, if we know only that |B(x)| is constant, as for the helical magnetic field $B(x) = (\cos(\tau x_3), \sin(\tau x_3), 0)$ encountered in liquid crystals [29, 11], then

$$\lim_{b \to +\infty} b^{-1/2} \lambda^{\mathrm{DN}}(bA, \Omega) = |B|^{\frac{1}{2}} \inf_{x \in \partial \Omega} \lambda^{\mathrm{DN}}(\vartheta(x)).$$

Notice, that if $\partial\Omega$ has a component homeomorphic to the sphere \mathbb{S}^2 , then the hairy ball Theorem applied to the tangential part of the magnetic field at the boundary implies that there exists a point $x \in \partial\Omega$ such that $\vartheta(x) = 0$ and we deduce

$$\inf_{x \in \partial \Omega} \lambda^{\mathrm{DN}}(\vartheta(x)) = \hat{\alpha} \,.$$

1.4. The magnetic Robin Laplacian. We can get information about $\lambda^{\text{DN}}(bA, \Omega)$ by comparing with the lowest eigenvalue of a Robin problem

(1.15)
$$\mu(\hat{\lambda}) = \mu(bA, \hat{\lambda}, \Omega) = \inf_{u \neq 0} \frac{\|(-i\nabla - bA)u\|_{\Omega}^2 - \lambda \|u\|_{\partial\Omega}^2}{\|u\|_{\Omega}^2}$$

In two dimensions and under constant magnetic field, two-term and three-term asymptotics for $\mu(\hat{\lambda})$ are available [18, 7] in the regime where $b \to +\infty$ and

$$\hat{\lambda} = \hat{\alpha}b^{1/2} + o(b^{1/2}).$$

If we choose

$$\hat{\lambda}^* = \hat{\alpha} b^{1/2} + o(b^{1/2})$$
 such that $\mu(\hat{\lambda}^*) \le 0$

and

$$\hat{\lambda}_* = \hat{\alpha} b^{1/2} + o(b^{1/2})$$
 such that $\mu(\hat{\lambda}_*) \ge 0$,

then, we get using the characterization in (1.6),

$$\hat{\lambda}_* \le \lambda^{\mathrm{DN}}(bA, \Omega) \le \hat{\lambda}^*.$$

We will use this approach in Section 5 to analyze the splitting of the low-lying eigenvalues under a constant magnetic field and we obtain

Theorem 1.7. Suppose that Ω is a regular domain in \mathbb{R}^2 such that the curvature of the boundary has a unique non-degenerate maximum. Let A be a vector field on Ω with constant magnetic field curl A = 1. Then there is a constant $K_*(\Omega) > 0$ such that

$$\lambda_2(bA,\Omega) - \lambda_1(bA,\Omega) = K_*(\Omega)b^{-1/4} + o(b^{-1/4}) \quad as \ b \to +\infty.$$

The Robin problem is not analyzed under non-constant magnetic field in two dimensions. For the three dimensional case, the existing results in [12] only cover the regime $|\hat{\lambda}| = o(b^{1/2})$, whereas the relevant regime for our setting is $|\hat{\lambda}| \propto b^{1/2}$.

1.5. Weak field limit. Our final result concerns the limit as $b \to 0$ in simply connected domains. Assuming that $\Omega \subset \mathbb{R}^n$ is simply connected, n = 2, 3, there is a unique vector field $A_{\Omega} \in H^1(\Omega; \mathbb{R}^n)$ such that [9, Prop. D.1.1]

$$\operatorname{curl} A_{\Omega} = 1 \text{ and } \operatorname{div} A_{\Omega} = 0 \text{ on } \Omega, \quad \vec{\nu} \cdot A_{\Omega} = 0 \text{ on } \partial\Omega.$$

Theorem 1.8. Let Ω be a regular domain in \mathbb{R}^n and A be a vector field generating a constant magnetic field curl A = 1. Then, the lowest eigenvalue of the D-t-N operator in Ω satisfies

$$\lambda^{\mathrm{DN}}(bA,\Omega) = \frac{b^2}{|\partial\Omega|} \int_{\Omega} |A_{\Omega}|^2 dx + o(b^2) \quad \text{as } b \to 0$$

Compared with the magnetic Laplacian, the coefficient of b^2 is the average of $|A_{\Omega}|^2$ over Ω [9, Proposition 1.5.2].

By [8, Proposition 3.1] and the geometric isoperimetric inequality, we obtain as corollary of Theorem 1.8:

Corollary 1.9. Let Ω be a regular domain in \mathbb{R}^2 and A be a magnetic potential with constant magnetic field with norm 1. Suppose that Ω is simply connected and \mathcal{B} is a disk with the same area as Ω . Then there exists $b_0(\Omega) > 0$ such that, $|b| \leq b_0(\Omega)$,

$$\lambda^{\mathrm{DN}}(bA,\Omega) \le \lambda^{\mathrm{DN}}(bA,\mathcal{B}).$$

Remark 1.10. When Ω is the disk of radius R, the magnetic potential $A_{\Omega}(x, y) = \frac{1}{2}(-y, x)$ and the lowest eigenvalue of the D-t-N operator is explicitly given by (see [16, Remark 6.1] and [17, Remark 5.10]),

$$\lambda^{\rm DN}(bA, D(0, R)) = \frac{bR}{2} \frac{I_0'(\frac{bR^2}{4})}{I_0(\frac{bR^2}{4})},$$

where

(1.16)
$$I_0(z) = \sum_{k=0}^{+\infty} \frac{z^{2k}}{2^{2k} (k!)^2}$$

is the modified Bessel function of the first kind of order 0. So, in this case, we can give a more accurate asymptotics:

(1.17)
$$\lambda^{\text{DN}}(bA, D(0, R)) = \frac{R^3 b^2}{16} + O(b^4).$$

1.6. **Organization.** In Section 2, we collect preliminaries to be used throughout the paper. In Sections 3 and 4, we prove upper and lower bounds on the low-lying eigenvalues in the two dimensional case, which yield Theorems 1.1, 1.2 and 1.4. For generic 2D domains, we study the splitting of the low-lying eigenvalues in Section 5. The proof of Theorem 1.6 in 3D domains occupies Section 6. We prove Theorem 1.8. in Section 7. Finally, there are two appendices on gauge transformations and a reference operator in the half-space, respectively.

2. Preliminaries

2.1. The half-plane. In the half-plane $\mathbb{R}^2_+ = \{(t,x) : t > 0\}$, for $b \in \mathbb{R}$, we introduce

(2.1)
$$E(b) = \inf_{u \in C^{\infty}(\overline{\mathbb{R}^{2}_{+}}), \|u\|_{\partial \mathbb{R}^{2}_{+}} = 1} \|(-i\nabla - bA_{\tau})u\|_{\mathbb{R}^{2}_{+}}^{2},$$

where $A_{\tau}(t,x) = (0,t)$, for $(t,x) \in \mathbb{R}^2_+$. Notice that A_{τ} is tangent to the boundary of \mathbb{R}^2_+ , and it generates a constant unit magnetic field with norm 1.

The sign of b is irrelevant by the invariance under the unitary transformation of complex conjugation, since

$$(2.2) E(b) = E(-b)$$

By scaling (see [17, Section 3]), E(b) and $\hat{\alpha}$ can be expressed as¹

(2.3)
$$E(b) = b^{1/2}E(1), \quad \hat{\alpha} = E(1).$$

2.2. Harmonic oscillator with Robin condition. The constant $\hat{\alpha}$ is also related to the harmonic oscillator on the half-axis

$$-\frac{d^2}{dt^2} + (t-\xi)^2 \quad \text{on } \mathbb{R}_+,$$

with Robin boundary condition at t = 0, (i.e with the boundary condition $u'(0) = \gamma u(0)$). The lowest eigenvalue of this operator was studied in [25, 18], and the other eigenvalues are more recently studied in [7].

We denote by $\|\cdot\|_2$ the L^2 -norm on \mathbb{R}_+ , and for b > 0, $\gamma, \xi \in \mathbb{R}^2$, we introduce the lowest eigenvalue of the Robin magnetic harmonic oscillator:

$$\mu(\gamma,\xi;b) = \inf_{\|f\|_2=1} \left(\int_0^{+\infty} \left(|f'(t)|^2 + (bt-\xi)^2 |f(t)|^2 \right) dt + \gamma |f(0)|^2 \right)$$

If we minimize over $\xi \in \mathbb{R}$, we set

$$\Theta(\gamma; b) = \inf_{\xi \in \mathbb{R}} \mu(\gamma, \xi; b),$$

and we have by scaling

(2.4)
$$\Theta(\gamma; b) = b\Theta(b^{-1/2}\gamma; 1), \quad \mu(\gamma, \xi; b) = b\mu(b^{-1/2}\gamma, b^{-1/2}\xi; 1)$$

This shows that it suffices to consider the case b = 1, and we thus introduce

(2.5)
$$\Theta(\gamma) := \Theta(\gamma; 1), \quad \mu(\gamma, \xi) := \mu(\gamma; \xi, 1).$$

The Neumann case $\gamma = 0$ corresponds to the de Gennes model, and

$$\Theta_0 = \Theta(0)$$

is the so-called de Gennes constant. It is known that approximatively $\Theta_0 \approx 0.590106$. As function of $\xi \in]-\infty, +\infty[$, the eigenvalue $\mu(\gamma, \xi)$ decreases from $+\infty$ until it reaches a unique minimum attained at

(2.6)
$$\xi(\gamma) = \sqrt{\Theta(\gamma) + \gamma^2},$$

then $\mu(\gamma, \xi)$ increases to 1. The function $\mathbb{R} \ni \gamma \to \Theta(\gamma) \in (-\infty, 1)$ is smooth and increasing [18, Theorem II.1 and Proposition II.5], and it has a unique zero $\gamma_0 < 0$, (see Figure 1):

(2.7)
$$\Theta(\gamma_0) = 0$$

Furthermore, the derivative of $\Theta(\cdot)$ is given in [18, Proposition II.5] as

(2.8)
$$\Theta'(\gamma) = |u_{\gamma}(0)|^2$$

where u_{γ} is a eigenfunction corresponding to the eigenvalue $\Theta(\gamma)$, normalized in $L^2(\mathbb{R}_+)$.

¹In [17], the authors consider m(b) = E(2b) and $\alpha = m(1)$.

We can also express the relation between $\xi(\gamma)$ and $\Theta(\gamma)$ in terms of the parabolic cylindrical functions as in [17, Eqs. (2.41)-(2.42)]. In fact, one can prove that $\xi(\gamma)$ satisfies the implicit equation:

$$\sqrt{2} \quad D'_{\frac{\Theta(\gamma)-1}{2}}(-\sqrt{2} \ \xi(\gamma)) = \gamma \ D_{\frac{\Theta(\gamma)-1}{2}}(-\sqrt{2} \ \xi(\gamma)).$$

Under the constraint (2.6), or equivalently using the relation ([28], p. 327),

(2.9)
$$D'_{\nu}(z) - \frac{z}{2}D_{\nu}(z) + D_{\nu+1}(z) = 0,$$

we see that $\xi(\gamma)$ is also solution of

(2.10)
$$-\sqrt{2}D_{\frac{\Theta(\gamma)+1}{2}}(-\sqrt{2}\xi(\gamma)) = (\gamma + \xi(\gamma)) \ D_{\frac{\Theta(\gamma)-1}{2}}(-\sqrt{2}\xi(\gamma)).$$

In light of (2.10), we have the following:

- If we take $\gamma = 0$, we recover [17, Eqs (2.40)-(2.41)].
- If we take $\gamma = \gamma_0$, knowing that $\xi(\gamma_0) = -\gamma_0$ and $\Theta(\gamma_0) = 0$, we derive obviously $D_{\frac{1}{2}}(\sqrt{2}\gamma_0) = 0$ in (2.10). Since $\hat{\alpha} = \alpha/\sqrt{2}$ and $-\alpha$ is the unique negative zero of $D_{\frac{1}{2}}$, we finally get that

$$\hat{\alpha} = -\gamma_0 \,.$$

FIGURE 1. Graph of the function $\Theta(\gamma)$.

2.3. D-to-N on the half-axis.

We can also derive the relation $\hat{\alpha} = -\gamma_0$, where γ_0 is the unique zero of $\Theta(\cdot)$, directly from the following characterization of $\hat{\alpha} = \alpha/\sqrt{2}$ (see [17, Eq. (6.7)])

(2.12)
$$\hat{\alpha} = \inf_{\substack{f(0) \neq 0\\\xi \in \mathbb{R}}} \frac{\int_0^{+\infty} \left(|f'(t)|^2 + (t-\xi)^2 |f(t)|^2 \right) dt}{|f(0)|^2}$$

This point of view is helpful to prove the following.

Proposition 2.1. There exists a function f_* in the Schwartz space $\mathcal{S}(\overline{\mathbb{R}_+})$ such that

(i) $f_* > 0$ on \mathbb{R}_+ and $f_*(0) = 1$. (ii)

$$\int_{\mathbb{R}_+} \left(|f'_*(t)|^2 + (t - \hat{\alpha})^2 |f_*(t)|^2 \right) dt = \hat{\alpha}.$$

 $(2.11) \qquad \qquad \hat{\alpha} =$

(iii)

$$\int_{\mathbb{R}_{+}} (t - \hat{\alpha}) |f_{*}(t)|^{2} dt = 0, \quad \int_{\mathbb{R}_{+}} (t - \hat{\alpha})^{2} |f_{*}(t)|^{2} dt = \frac{\hat{\alpha}}{4}, \quad \int_{\mathbb{R}_{+}} (t - \hat{\alpha})^{3} |f_{*}(t)|^{3} dt = \frac{1}{6} (1 - 2\hat{\alpha}^{2}).$$
(iv)

$$\int_{\mathbb{R}_+} f'_*(t) f_*(t) dt = -\frac{1}{2} \,, \quad \int_{\mathbb{R}_+} t |f'_*(t)|^2 dt = \frac{1}{3} + \frac{\hat{\alpha}^2}{12} \,.$$

Proof. Knowing that $\hat{\alpha} = -\gamma_0$ and $\Theta(\gamma_0) = 0$, we infer from (2.6) that $\xi(\gamma_0) = \hat{\alpha}$ and $\mu(-\hat{\alpha}, \hat{\alpha}) = 0$. Choose a positive and normalized (in $L^2(\mathbb{R}_+)$) ground state φ_* of $\mu(-\hat{\alpha}, \hat{\alpha})$ and put $f_* = (1/|\varphi_*(0)|)\varphi_*$. Then f_* satisfies

$$-f_*'' + (t - \hat{\alpha})^2 f_* = 0 \text{ on } \mathbb{R}_+, \quad f_*'(0) = -\hat{\alpha}, \quad f_*(0) = 1.$$

We get (ii) and $\int_{\mathbb{R}_+} f'_*(t) f_*(t) dt = -\frac{1}{2}$ by integration by parts. For the identities in (iii), we reproduce the calculations in [3]. We notice that for

$$v = 2pf'_* - p'f_*, \ p(t) = (t - \hat{\alpha})^k \text{ and } k \in \{0, 1, 2\},\$$

we have

$$\left(-\partial_t^2 + (t-\hat{\alpha})^2\right)v = \left(-4(t-\hat{\alpha})^2p' - 4(t-\hat{\alpha})p\right)f_*,$$

and we use integration by parts to write

$$\int_{\mathbb{R}_+} \left(-4(t-\hat{\alpha})^2 p' - 4(t-\hat{\alpha})p \right) |f_*(t)|^2 dt = \left(v'(0) + \hat{\alpha}v(0) \right) f_*(0).$$

The formulas of (iii) follow by considering successively k = 0, 1, 2. Finally, the last identity in (iv) is obtained by integration by parts

$$\int_{\mathbb{R}_+} t |f'_*(t)|^2 dt = -\int_{\mathbb{R}_+} f_*(t) f'_*(t) dt - \int_{\mathbb{R}_+} t f_*(t) f''_*(t) dt$$

and by using $f_*''(t) = (t - \hat{\alpha})^2 f_*(t)$ and the identities in (iii).

Remark 2.2. Although not needed in the proof, notice that we have

$$f_*(t) = \frac{D_{-1/2}(\sqrt{2}t - \alpha)}{D_{-1/2}(-\alpha)}$$

Note that the parabolic cylinder function $D_{-\frac{1}{2}}(z)$ is also related with the usual Hankel function: $D_{-\frac{1}{2}}(z) = \sqrt{\frac{z}{2\pi}} K_{\frac{1}{4}}(\frac{z^2}{4})$. Using Mathematica, we get numerically

$$\int_0^{+\infty} f_*(t)^2 dt \approx 0.6861814388 \,.$$

2.4. A useful identity. Recall that $W^{1,\infty}(\Omega;\mathbb{R})$ is the space of $L^{\infty}(\Omega)$ real-valued functions f such that $\nabla f \in L^{\infty}(\Omega)$.

In various proofs, we will use the now standard identity given by the following proposition.

Proposition 2.3. Suppose that $A \in H^1(\Omega; \mathbb{R}^2)$. If $u \in H^1(\Omega)$ and $w \in W^{1,\infty}(\Omega, \mathbb{R})$, then

$$\operatorname{\mathsf{Re}}\int_{\Omega}(-i\nabla - A)u \cdot \overline{(-i\nabla - A)(w^2u)}\,dx = \int_{\Omega}|(-i\nabla - A)(wu)|^2\,dx - \int_{\Omega}|\nabla w|^2|u|^2\,dx$$

Proof. This is a direct consequence of the two identities

$$(-i\nabla - A)(w^2u) = w(-i\nabla - A)(wu) - iwu\nabla w, \quad w(-i\nabla - A)u = (-i\nabla - A)(wu) + iu\nabla w.$$

2.5. Gauge invariance. If $U \subset \Omega$ is an open set and $\phi : U \to \mathbb{R}$ is in $H^1(U)$, then for any function $u \in H^1(U)$, we have the following identities on U,

$$|(-i\nabla - bA)u|^2 = |(-i\nabla - bA')v|^2, \quad |u|^2 = |v|^2, \quad \operatorname{curl} A = \operatorname{curl} A',$$

where $v = u e^{-ib\phi}$ and $A' = A - \nabla \phi$. This amounts to a local gauge transformation.

If U is simply connected and we know that the vector potentials A and A' have the same curl on U, then we can find a function ϕ defined on U such that $A' = A - \nabla \phi$.

We will use local gauge transformations to transform a given vector potential to a more convenient one. For instance, if B = dA is not constant, we describe below how we can locally approximate A by a vector potential with constant magnetic field, up to a local gauge transformation.

Proposition 2.4. Suppose that $\partial\Omega$ is C^1 , $A \in H^1(\Omega; \mathbb{R}^2)$ and $B = \operatorname{curl} A$ is C^1 on $\overline{\Omega}$. There exists a constant C > 0 such that, for every $p \in \overline{\Omega}$, there exists a function $\phi_p : U_p \to \mathbb{R}$ such that

$$|A(x) - \nabla \phi_p - B(p)A_0(x-p)| \le C |x-p|^2 \quad (x \in U_p)$$

where $A_0(x) = \frac{1}{2}(-x_2, x_1)$, and $U_p = \overline{\Omega} \cap \mathcal{B}(p, \delta)$ for some $\delta > 0$.

A stronger version of Proposition 2.4 is given in [26, Lemma 3.1], when the point p is in Ω .

Proof of Proposition 2.4. We can extend A to a compactly supported vector field in $H^1(\mathbb{R}^2; \mathbb{R}^2)$, and we obtain an extension of $B = \operatorname{curl} A$ to all of \mathbb{R}^2 as well. In a disk $\mathcal{B}(p, \delta)$ centered at p, consider the Coulomb gauge

$$A'(x) = 2\left(\int_0^1 B(p + s(x - p))sds\right)A_0(x - p).$$

Noticing that $\operatorname{curl} A = \operatorname{curl} A'$, we can write $A = A' - \nabla \phi$ in $\mathcal{B}(p, \delta)$. Since B is C^1 on $\overline{\Omega}$, we get $A'(x) = B(p)A_0(x-p) + \mathcal{O}(|x-p|^2), x \in U_p$.

2.6. Parallel coordinates in two dimensions. In the course of the proofs, we will often deal with functions supported in a neighborhood of a boundary point of Ω . In such cases, it is convenient to carry out the computations in *parallel coordinates* that we introduce below.

Pick a connected component Γ of the boundary of Ω , and denote by L its length. By means of the arc-length parametrization, we can identify Γ and $\mathbb{R}/L\mathbb{Z}$, where $s \in \mathbb{R}/L\mathbb{Z}$ is the curvilinear coordinate of a point $x \in \Gamma$. We choose $t_0 > 0$ sufficiently small such that

$$\Omega_0 := \{ x \in \Omega : \operatorname{dist}(x, \Gamma) < t_0 \}$$

is diffeomorphic to $(0, t_0) \times (\mathbb{R}/L\mathbb{Z})$. More precisely, we introduce the diffeomorphism

(2.13)
$$\Phi_0: (0, t_0) \times (\mathbb{R}/L\mathbb{Z}) \to \Omega_0.$$

such that, for $x = \Phi_0(t, s) \in \Omega_0$, $t = \text{dist}(x, \Gamma)$ and s is the curvilinear coordinate of p(x), the the orthogonal projection of x on Γ . Thus, t denotes the normal distance to Γ , and s measures the tangential distance along Γ .

Note that, if Ω is simply connected, the boundary consists of a single connected component, $\Gamma = \partial \Omega$.

3. Upper bounds in two dimensional domains

3.1. Non vanishing magnetic field.

Proposition 3.1. Let Ω be a regular domain in \mathbb{R}^2 and A be a vector potential with a magnetic field $B = \operatorname{curl} A$ that vanishes nowhere on $\partial \Omega$. Suppose that B is C^1 on a neighborhood of $\partial \Omega$. Then, for every fixed $j \geq 1$, the j'th eigenvalue value of the D-to-N map Λ_{bA} satisfies

$$\lambda_j^{\mathrm{DN}}(bA,\Omega) \leq \left(\inf_{x\in\partial\Omega} |B(x)|\right)^{\frac{1}{2}} \hat{\alpha} \, b^{\frac{1}{2}} + \mathcal{O}(b^{1/3}) \quad , \quad b \to +\infty \, .$$

Proof. Since the D-to-N operators with vector potentials A and -A are unitarily equivalent, it suffices to consider B > 0.

Step 1. The test function.

Choose a point $p \in \partial \Omega$ such that $B(p) = \min_{x \in \partial \Omega} B(x)$. By Proposition 2.4, we can assume that, modulo a (local) gauge transformation, that A satisfies,

$$A(x) = B(p)A_0(x-p) + \mathcal{O}(|x-p|^2).$$

Let Γ be the connected component of $\partial\Omega$ that contains p. Working in parallel coordinates $(t,s) \in (0,t_0) \times (\mathbb{R}/L\mathbb{Z})$, centered at the point p, we introduce the functions

$$v(t,s) = u(t,s) \cdot e^{-ib\varphi(s,t)}, \quad u(t,s) = b^{\rho/2}\chi_1(b^{\rho}s) \cdot \chi(b^{\rho}t) \cdot f_*(B(p)^{1/2}b^{1/2}t) \cdot e^{-i(B(p)b)^{1/2}\hat{\alpha}s},$$

where $\chi_1 \in C_c^{\infty}(-1, 1)$ is normalized in $L^2(\mathbb{R})$, χ is a smooth cut-off function, equal to 1 in a neighborhood of 0, $\rho \in (0, \frac{1}{2})$, and f_* is the function introduced in Proposition 2.1. The function φ is real-valued and amounts to a (local) gauge transformation in the (t, s) coordinates (its choice will be explained below).

With Φ_0 the diffeomorphism introduced in (2.13), $v \circ \Phi_0$ defines a function in $H^1(\Omega)$, which will be the test function with which we will work. Since χ_1 is nomalized in $L^2(\mathbb{R})$, we get that the restriction of $v \circ \Phi_0$ to $\partial\Omega$ is normalized in $L^2(\partial\Omega)$. Put $A^{\text{lin}}(x) := B(p)A_0(x-p)$. We choose the function φ such that (see Appendix A):

(3.1)
$$\|(-i\nabla - bA^{\ln})v \circ \Phi_0\|_{\Omega}^2 = \int_{\mathbb{R}^2_+} \Big(|\partial_t u|^2 + (1 - tk(s))^{-2} |(-i\partial_s + B(p)b(t - \frac{1}{2}t^2k(s))u|^2 \Big) (1 - tk(s))dtds,$$

where k(s) is the curvature at the point of curvilinear coordinate s.

Step 2. Some estimates.

Our choice of the function u yields

$$\int_{\mathbb{R}^2_+} |u(t,s)|^2 dt ds \le \left(B(p)b \right)^{-1/2} \int_{\mathbb{R}_+} |f_*(\tau)|^2 d\tau.$$

With $\epsilon \in (0, 1)$, we have by the Cauchy-Schwarz inequality,

$$\int_{\mathbb{R}^2_+} \left| \left(-i\partial_s + B(p)b(t - \frac{1}{2}t^2k(s) \right) u \right|^2 dt ds \le (1 + \epsilon) \int_{\mathbb{R}^2_+} \left| (-i\partial_s + B(p)bt) u \right|^2 dt ds + \mathcal{O}(\epsilon^{-1}b^{\frac{3}{2} - 4\rho}).$$

Moreover, a routine calculation yields

$$\int_{\mathbb{R}^2_+} \left| (-i\partial_s + B(p)bt) u \right|^2 dt ds \le (1+\epsilon) \left(B(p)b \right)^{1/2} \int_{\mathbb{R}_+} (\tau - \hat{\alpha})^2 |f_*(\tau)|^2 d\tau + \mathcal{O}(\epsilon^{-1}b^{-\frac{1}{2}+2\rho}).$$

Using that f_* is a Schwartz function, we have²

$$\int_{\mathbb{R}^2_+} |\partial_t u|^2 dt ds = \left(B(p)b \right)^{1/2} \int_{\mathbb{R}^2_+} |f'_*(\tau)|^2 d\tau + \mathcal{O}(b^{-\infty}).$$

Returning back to (3.1), we bound 1 - tk(s) from above by $1 + \mathcal{O}(b^{-\rho})$. Using ii. in Proposition 2.1, and choosing $\epsilon = b^{-\rho}$ and $\rho = 1/3$, we get

$$\|(-i\nabla - bA^{\operatorname{lin}})v \circ \Phi_0\|_{\Omega}^2 \le (B(p)b)^{1/2}\hat{\alpha} + \mathcal{O}(b^{1/3}).$$

Step 3. Finishing the proof for the lowest eigenvalue.

Using the Cauchy-Schwarz inequality, one has

$$\begin{aligned} \|(-i\nabla - bA)v \circ \Phi_0\|_{\Omega}^2 &\leq (1 + b^{-1/6}) \|(-i\nabla - bA^{\ln})v \circ \Phi_0\|_{\Omega}^2 + \mathcal{O}(b^{1/6}\|b(A - A^{\ln})v \circ \Phi_0\|_{\Omega}^2) \\ &\leq (B(p)b)^{1/2}\hat{\alpha} + \mathcal{O}(b^{1/3}). \end{aligned}$$

Since $v \circ \Phi_0^{-1}$ is normalized in $L^2(\partial \Omega)$, and B(p) is the minimum of B on the boundary, this finishes the proof of the proposition for j = 1, thanks to the characterization of $\lambda_1^{\text{DN}}(bA, \Omega)$ in (1.6).

Step 4. Finishing the proof for the *j*'th eigenvalue, j > 1.

²We write $F = \mathcal{O}(b^{-\infty})$ if, for any $n \in \mathbb{N}$, $F = \mathcal{O}(b^{-n})$ as $b \to +\infty$.

Consider $\chi_1, \dots, \chi_j \in C_c^{\infty}(-1, 1)$ that constitute an orthonormal set in $L^2(\mathbb{R})$ and such that their supports are pairwise disjoint. We slightly modify the test function by introducing, for every $j \in \mathbb{N}$,

$$v_j(t,s) = u_j(t,s) \cdot e^{-ib\varphi(s,t)}, \quad u_j(t,s) = b^{\rho/2}\chi_j(b^\rho s) \cdot \chi(b^\rho t) \cdot f_*(B(p)^{1/2}b^{1/2}t) \cdot e^{-i(B(p)b)^{1/2}\hat{\alpha}s}.$$

Then, with $\rho = \frac{1}{3}$, and $q \in \{1, \dots, j\}$,

$$\|(-i\nabla - bA)v_q \circ \Phi_0\|_{\Omega}^2 \le \left(B(p)b\right)^{1/2}\hat{\alpha} + \mathcal{O}(b^{1/3}).$$

and for $q' \neq q$,

$$\langle (-i\nabla - bA)v_q \circ \Phi_0, (-i\nabla - bA)v_{q'} \circ \Phi_0 \rangle_{\Omega} = 0$$

Let $M_j = \text{Span}(v_1 \circ \Phi_0, \dots, v_j \circ \Phi_0)$ and $M_j^{\partial\Omega}$ be the space of its restriction to $\partial\Omega$. We observe that the $v_p(0, \cdot)$ $(p = 1, \dots, j)$ form an orthonormal basis of $M_j^{\partial\Omega}$. Hence, dim $(M_j) = j$, and we conclude by using the variational formulation in (1.7),

$$\lambda_j(bA,\Omega) \le \max_{g \in M_j} \frac{\|(-i\nabla - bA)g\|_{\Omega}^2}{\|g\|_{\partial\Omega}^2} \le (B(p)b)^{1/2}\hat{\alpha} + \mathcal{O}(b^{1/3}).$$

3.2. Improvement in the constant magnetic field.

Proposition 3.2. Let Ω be a regular domain in \mathbb{R}^2 and A be a vector potential with magnetic field $B = \operatorname{curl} A$. Suppose that B = 1 on a neighborhood of $\partial \Omega$. Then, for every fixed $j \ge 1$, the *j*'th eigenvalue value of the D-to-N map Λ_{bA} satisfies

$$\lambda_j^{\mathrm{DN}}(bA,\Omega) \le \hat{\alpha} \, b^{\frac{1}{2}} - \frac{\hat{\alpha}^2 + 1}{3} \max_{x \in \partial \Omega} k(x) + \mathcal{O}(b^{-1/6}) \quad (b \to +\infty),$$

where k is the curvature of $\partial \Omega$.

Proof.

Step 1. The test function.

The test function has a similar structure to the one constructed in Proposition 3.1, but since the magnetic field is constant on a neighborhood of $\partial\Omega$, we can carry out the computations to sub-leading terms.

Choose a point $p \in \partial \Omega$ such that $k(p) = \max_{x \in \partial \Omega} k(x)$, and let Γ be the connected component of $\partial \Omega$ that contains p. Let Φ_0 be the coordinate transformation as introduced in (2.13), but we center it at p, i.e. $\Phi_0^{-1}(p) = (0, 0)$.

Consider $\chi_1, \chi_2, \dots \in C_c^{\infty}(-1, 1)$ that constitute an orthonormal set in $L^2(\mathbb{R})$. For every $j \in \mathbb{N}$, the test function has the form $v_j \circ \Phi_0^{-1}$, with v_j defined as

$$v_j(t,s) = u_j(t,s) \cdot e^{-ib\varphi(s,t)}, \quad u_j(t,s) = b^{\rho/2}\chi_j(b^{\rho}s) \cdot \chi(b^{\rho}t) \cdot f_*(b^{1/2}t) \cdot e^{-ib^{1/2}\hat{\alpha}s}.$$

Here, χ is a cut-off function, equal to 1 in a neighborhood of 0, $\rho \in (0, \frac{1}{2})$, and f_* is the function introduced in Proposition 2.1. Moreover, we can suppose that the functions χ_1, \dots, χ_j have pairwise disjoint supports,

The function φ is defined in Appendix A. We can use

(3.2)
$$\|(-i\nabla - bA)v_j \circ \Phi_0\|_{\Omega}^2 = \int_{\mathbb{R}^2_+} \left(|\partial_t u_j|^2 + (1 - tk(s))^{-2} \left| \left(-i\partial_s + b(t - \frac{1}{2}t^2k(s))u_j \right|^2 \right) (1 - tk(s)) dt ds.$$

Restricting the functions $v_j \circ \Phi_0$ to $\partial \Omega$, we obtain an orthonormal set in $L^2(\partial \Omega)$.

Step 2. Some estimates.

Consider $u \in \{u_j : j \in \mathbb{N}\}$. Since f_* is a Schwartz function, the function u satisfies, for $m \ge 0$,

(3.3)
$$\int_{\mathbb{R}^{2}_{+}} t^{m} |u|^{2} dt ds = b^{-\frac{m+1}{2}} \int_{\mathbb{R}^{+}} \tau^{m} |f_{*}(\tau)|^{2} d\tau + \mathcal{O}(b^{-\infty}),$$
$$\int_{\mathbb{R}^{2}_{+}} t^{m} |\partial_{t} u|^{2} dt ds = b^{-\frac{m-1}{2}} \int_{\mathbb{R}^{+}} \tau^{m} |f_{*}'(\tau)|^{2} d\tau + \mathcal{O}(b^{-\infty}).$$

In the support of u, we have $k(s) = k(0) + \mathcal{O}(b^{-\rho})$. Hence,

(3.4)
$$\int_{\mathbb{R}^{2}_{+}} t^{m} |u|^{2} k(s) dt ds = b^{-\frac{m+1}{2}} k(0) \int_{\mathbb{R}_{+}} \tau^{m} |f_{*}(\tau)|^{2} d\tau + \mathcal{O}\left(b^{-\frac{m+1}{2}-\rho}\right),$$
$$\int_{\mathbb{R}^{2}_{+}} t^{m} |\partial_{t} u|^{2} k(s) dt ds = b^{-\frac{m-1}{2}} k(0) \int_{\mathbb{R}_{+}} \tau^{m} |f_{*}'(\tau)|^{2} d\tau + \mathcal{O}\left(b^{-\frac{m-1}{2}-\rho}\right)$$

In particular, we have

(3.5)
$$\int_{\mathbb{R}^2_+} |\partial_t u|^2 (1 - tk(s)) dt ds = b^{1/2} \int_{\mathbb{R}_+} |f'_*(\tau)|^2 d\tau - k(0) \int_{\mathbb{R}_+} \tau |f'_*(\tau)|^2 d\tau + \mathcal{O}(b^{-\rho}).$$

Step 3. More estimates.

F

Since the functions f_* and χ are real-valued, it is straightforward to verify that

(3.6)
$$\left| \left(-i\partial_s + b(t - \frac{1}{2}t^2k(s))u \right|^2 = F + G, \right.$$

where

$$= b \left| \left(b^{1/2} (t - \frac{1}{2} t^2 k(s)) - \hat{\alpha} \right) u \right|^2, \quad G = b^{3\rho} |\partial_s \chi(b^\rho s)|^2 |\chi(b^\rho t) f_*(b^{1/2} t)|^2.$$

Writing $(1 - tk(s))^{-1} = \mathcal{O}(1)$ in the support of u, and doing the change of variables $(\sigma, \tau) = (b^{\rho}s, b^{1/2}t)$, we get

(3.7)
$$\int_{\mathbb{R}^2_+} (1 - tk(s))^{-1} G \, dt ds = \mathcal{O}(b^{2\rho - \frac{1}{2}}).$$

To get an accurate estimate of the integral of $(1 - tk(s))^{-1}F$, we write

$$(1 - tk(s))^{-1} = 1 + tk(s) + \mathcal{O}(t^2)$$
 and $k(s) = k(0) + \mathcal{O}(b^{-\rho})$

in the support of u, and we expand the square to get

$$(1 - tk(s))^{-1}F = (1 + tk(0))F + \mathcal{O}((b^{-\rho}t + t^2)F),$$

$$F = b[(b^{1/2}t - \hat{\alpha})^2 - b^{1/2}t^2k(s)(b^{1/2}t - \hat{\alpha}) + \mathcal{O}(bt^4)]|u|^2$$

Doing a routine calculation, we obtain

(3.8)
$$\int_{\mathbb{R}^{2}_{+}} (1 - tk(s))^{-1} F \, dt ds = b^{1/2} \int_{\mathbb{R}_{+}} (\tau - \hat{\alpha})^{2} |f_{*}(\tau)|^{2} d\tau - k(0) \int_{\mathbb{R}_{+}} \left[\tau^{2} (\tau - \hat{\alpha}) - \tau (\tau - \hat{\alpha})^{2} \right] |f_{*}(\tau)|^{2} d\tau + \mathcal{O}(b^{-\rho}).$$

Returning to (3.6), we infer from (3.7) and (3.8),

(3.9)
$$\int_{\mathbb{R}^{2}_{+}} (1 - tk(s))^{-1} \left| \left(-i\partial_{s} - b(t - \frac{1}{2}t^{2}k(s)) u \right|^{2} dt ds = b^{1/2} \int_{\mathbb{R}^{+}} (\tau - \hat{\alpha})^{2} |f_{*}(\tau)|^{2} d\tau - k(0) \int_{\mathbb{R}^{+}} \left[\tau^{2}(\tau - \hat{\alpha}) - \tau(\tau - \hat{\alpha})^{2} \right] |f_{*}(\tau)|^{2} d\tau + \mathcal{O}(b^{-\rho}) + \mathcal{O}(b^{2\rho - \frac{1}{2}}).$$

Step 4. Finishing the proof.

We introduce the following constant

$$C_* = -\int_{\mathbb{R}_+} \tau |f'_*(\tau)|^2 d\tau - \int_{\mathbb{R}_+} \left[\tau^2 (\tau - \hat{\alpha}) - \tau (\tau - \hat{\alpha})^2 \right] |f_*(\tau)|^2 d\tau,$$

which can also be expressed as^3

$$C_* = -\int_{\mathbb{R}_+} \tau |f'_*(\tau)|^2 d\tau - \hat{\alpha} \int_{\mathbb{R}_+} (\tau - \hat{\alpha})^2 |f_*(\tau)|^2 d\tau.$$

Then, we collect (3.5) and (3.9) and choose ρ such that $-\rho = 2\rho - \frac{1}{2}$, i.e. $\rho = 1/6$. Eventually, for $v \in \{v_j : j \in \mathbb{N}\}$, we infer from (3.2),

$$\|(-i\nabla - bA)v \circ \Phi_0\|_{\Omega}^2 = \hat{\alpha}b^{1/2} + C_*k(0) + \mathcal{O}(b^{-1/6}),$$

where we used (ii) in Proposition 2.1.

The space $M_j = \text{Span}(v_n \circ \Phi_0^{-1}: 1 \le n \le j)$ has dimension j and its restriction to $\partial\Omega$ has an orthonormal basis consisting of the functions $v_p(0, \cdot)$ $(p = 1, \dots, j)$. Thus, we conclude by (1.7) that

$$\lambda_j^{\text{DN}}(bA, \Omega) \le \hat{\alpha} b^{1/2} + C_* k(0) + \mathcal{O}(b^{-1/6}).$$

To finish the proof, we recall that $k(0) = \max_{x \in \partial \Omega} k(x)$, and we use (iii) and (iv) in Proposition 2.1 to deduce that

$$C_* = -\frac{\hat{\alpha}^2 + 1}{3} \,.$$

³This form is similar to the one in the displayed equation appearing after Eq. (11.27) in [13].

4. LOWER BOUNDS IN TWO DIMENSIONS

4.1. Non vanishing magnetic fields.

Assumption 4.1 (Admissible magnetic fields). Let Ω be a regular domain in \mathbb{R}^2 and $A: \Omega \to \mathbb{R}^2$ be a vector potential with a magnetic field $B = \operatorname{curl} A$ such that

$$\liminf_{b\to+\infty} \left(b^{-\zeta} \inf_{\|u\|_{\Omega}=1} \|(-i\nabla - bA)u\|_{\Omega}^2 \right) > 0,$$

for some $\zeta > 1/2$.

Example 1 (Magnetic fields satisfying Assumption 4.1).

- (i) If B is C^1 on $\overline{\Omega}$ and |B| > 0 everywhere on $\overline{\Omega}$, then by [13], Assumption 4.1 holds with $\zeta = 1$.
- (ii) If B is a non-vanishing step function and the discontinuity set consists of a finite number of smooth curves in Ω , then Assumption 4.1 holds with $\zeta = 1$ (see [1, 2]).
- (iii) If B is C^1 on $\overline{\Omega}$, |B| > 0 on $\partial\Omega$, and the set $\mathcal{Z}(B) = \{x \in \Omega : B(x) = 0\}$ consists of a finite number of smooth curves such that $|\nabla B| > 0$ on $\mathcal{Z}(B)$, then by [30, Theorem 4], Assumption 4.1 holds with $\zeta = 2/3$.

Proposition 4.2. Let Ω be a regular domain in \mathbb{R}^2 and A be a vector potential with a magnetic field $B = \operatorname{curl} A$ that satisfies Assumption 4.1 and does not vanish on $\partial\Omega$. Suppose that B is C^1 on a neighborhood of $\partial\Omega$. Then, there is $\delta > 0$ such that, the ground state energy of the D-to-N map Λ_{bA} satisfies

$$\lambda_1^{\mathrm{DN}}(bA,\Omega) \ge \left(\inf_{x \in \partial \Omega} |B(x)|\right)^{\frac{1}{2}} b^{\frac{1}{2}} \hat{\alpha} + \mathcal{O}(b^{\frac{1}{2}-\delta})) \quad , \quad b \to +\infty \,.$$

Proof. We choose $t_0 > 0$ such that B is C^1 and does not vanish on

$$\mathcal{N} := \{ x \in \overline{\Omega} : \operatorname{dist}(x, \partial \Omega) < t_0 \}.$$

Let $u \in C^{\infty}(\overline{\Omega})$. In the sequel, all estimates will be uniform with respect to u, and with respect to b in a neighborhood of $+\infty$.

As a consequence of Assumption 4.1, there exist positive constants Θ_1, b_0 and $\zeta > \frac{1}{2}$ such that

(4.1)
$$\|(-i\nabla - bA)u\|_{\Omega}^2 \ge \Theta_1 b^{\zeta} \|u\|_{\Omega}^2 \quad \text{for all } b \ge b_0,$$

and by [13],

(4.2)
$$\|(-i\nabla - bA)u\|_{\Omega}^2 \ge \Theta_1 b \|u\|_{\Omega}^2 \quad \text{if supp } u \subset \mathcal{N}.$$

We now proceed to the proof of Proposition 4.2, which we split into several steps.

Step 1:

Consider a constant $\rho \in (\frac{1}{4}, \zeta)$ and a partition of unity

$$\chi_1^2 + \chi_2^2 = 1 \quad \text{on } \Omega,$$

$$t(x, \partial \Omega) < b^{-\rho} \quad \text{supp } \chi_1 \subset \{\text{dist}(x, \partial \Omega) < 0\}$$

where $\chi_1 = 1$ on $\{ \operatorname{dist}(x, \partial \Omega) < b^{-\rho} \}$, $\operatorname{supp} \chi_1 \subset \{ \operatorname{dist}(x, \partial \Omega) < 2b^{-\rho} \}$ and

$$|\nabla \chi_1| + |\nabla \chi_2| = \mathcal{O}(b^{\rho}).$$

Then, with $\epsilon = b^{-\delta}$ and $\delta > 0$, we write

$$\begin{aligned} \|(-i\nabla - bA)u\|_{\Omega}^{2} \\ &= (1-\epsilon)\|(-i\nabla - bA)u\|_{\Omega}^{2} + \epsilon\|(-i\nabla - bA)u\|_{\Omega}^{2} \\ &= (1-\epsilon)\sum_{j=1}^{2}\|(-i\nabla - bA)\chi_{j}u\|_{\Omega}^{2} + \epsilon\|(-i\nabla - bA)u\|_{\Omega}^{2} + \mathcal{O}(b^{2\rho}\|u\|_{\Omega}^{2}) \\ &\geq (1-\epsilon)\sum_{j=1}^{2}\|(-i\nabla - bA)\chi_{j}u\|_{\Omega}^{2} + (\epsilon\Theta_{1}b^{\zeta} + \mathcal{O}(b^{2\rho}))\|u\|_{\Omega}^{2}, \end{aligned}$$

where we have used (4.1), and Proposition 2.3 with $w = \chi_j$. For instance, assuming that

$$(4.3) 2\rho - \zeta + \delta < 0,$$

we obtain that, for b sufficiently large,

(4.4)
$$\|(-i\nabla - bA)u\|_{\Omega}^{2} \ge (1-\epsilon)\|(-i\nabla - bA)\chi_{1}u\|_{\Omega}^{2} + \frac{\epsilon\Theta_{1}b^{\zeta}}{2}\|u\|_{\Omega}^{2}$$

Step 2:

For the sake of simplicity, we suppose that Ω is simply connected, to ensure that

$$\Omega_0 := \{ x \in \Omega : \operatorname{dist}(x, \partial \Omega) < t_0 \}$$

is diffeomorphic to $(0, t_0) \times (\mathbb{R}/L\mathbb{Z})$, and use the parallel coordinates defined by the transformation Φ_0 in (2.13). The proof can be easily adjusted to cover the case the non-simply connected case where $\partial\Omega$ consists of a finite number of connected components, by treating doing the computations on each connected component.

We introduce a partition of unity of Ω_0 ,

$$\sum_{j} g_j^2 = 1, \quad \sum_{j} |\nabla g_j|^2 = \mathcal{O}(b^{2\rho}),$$

such that

supp
$$g_j \subset \Phi_0((0, t_0) \times (s_j - b^{-\rho}, s_j + b^{-\rho}).$$

With $v = \chi_1 u$ supported in $\{|B| > 0\}$, we use (4.2) to write

$$\|(-i\nabla - bA)v\|_{\Omega}^2 \ge (1-\epsilon)\|(-i\nabla - bA)v\|_{\Omega}^2 + \epsilon\Theta_1 b\|v\|_{\Omega}^2.$$

Consequently, we have, with $v_j = g_j v$,

(4.5)
$$\|(-i\nabla - bA)v\|_{\Omega}^{2} \ge (1-\epsilon)\sum_{j} \|(-i\nabla - bA)v_{j}\|_{\Omega}^{2} + \epsilon\Theta_{1}b\|v\|_{\Omega}^{2} + \mathcal{O}(b^{2\rho}\|v\|_{\Omega}^{2}).$$

For all j, pick $p_j \in \partial \Omega \cap \operatorname{supp} g_j$ and put $A_j^{\text{lin}}(x) = B(p_j)A_0(x-p_j)$. In a neighborhood of p_j , we apply a gauge transformation

$$(v_j, A) \mapsto (v'_j = v e^{-ib\phi_j}, A'_j = \nabla \phi_j + A_j^{\text{lin}})$$

as indicated in Proposition 2.4. For instance, we have

$$|A - A'_j| \le C b^{-4\rho}$$
 on $\operatorname{supp} v_j$.

To lighten the notation, we skip the ' when referring to the new configuration (v'_j, A'_j) . We have by the Cauchy-Schwarz inequality,

$$\|(-i\nabla - bA)v_j\|_{\Omega}^2 \ge (1-\epsilon)\|(-i\nabla - bA_j^{\ln})v_j\|_{\Omega}^2 - C\epsilon^{-1}b^{2-4\rho}\|v_j\|^2.$$

Inserting this into (4.5) and arguing as in (4.4), we get

(4.6)
$$\|(-i\nabla - bA)v\|_{\Omega}^{2} = (1 - 2\epsilon) \sum_{j} \|(-i\nabla - bA_{j}^{\mathrm{lin}})v_{j}\|_{\Omega}^{2} + \frac{\epsilon\Theta_{1}b}{2} \|v\|_{\Omega}^{2},$$

provided that

(4.7)
$$1 - 4\rho + 2\delta < 0.$$

Step 3.

Now we write a lower bound for $\|(-i\nabla - bA_j^{\text{lin}})v_j\|_{\Omega}^2$. Notice that $\operatorname{curl} A_j^{\text{lin}} = B(p_j)$ is constant. As recalled in Appendix A,

$$\begin{aligned} \|v_j\|_{\Omega}^2 &= \int_{\mathbb{R}^2_+} |\tilde{v}_j|^2 (1 - tk(s)) dt ds, \\ |(-i\nabla - bA_j^{\text{lin}})v_j\|_{\Omega}^2 &= \int_{\mathbb{R}^2_+} \left(|\partial_t \tilde{v}_j|^2 + (1 - tk(s))^{-2} |(-i\partial_s^2 + B(p_j)b(t - \frac{1}{2}t^2k(s)))\tilde{v}_j|^2 \right) (1 - tk(s)) dt ds, \end{aligned}$$

where

$$\tilde{v}_j(t,s) = e^{-ib\varphi_j(t,s)}v \circ \Phi_0(t,s)$$

is obtained after expressing v_j in the (t,s) coordinates and after performing a gauge transformation.

Recall that the support of v_j is contained in $\{\operatorname{dist}(x,\partial\Omega) \leq 2b^{-\rho}\} \cap \operatorname{supp} g_j$. Consequently, on the support of \tilde{v}_j ,

$$(1 - tk(s)) = 1 + \mathcal{O}(b^{-\rho}), \quad (1 - tk(s))^{-2} = 1 + \mathcal{O}(b^{-\rho}),$$

and a routine application of Hölder's inequality yields,

$$|(-i\partial_s + b(t - \frac{1}{2}t^2k(s)))\tilde{v}_j|^2 \ge (1 - \epsilon)|(-i\partial_s + B(p_j)bt)\tilde{v}_j|^2 + \mathcal{O}(\epsilon^{-1}b^{2-4\rho}|\tilde{v}_j|^2).$$

Collecting the previous estimates, we get

$$\|(-i\nabla - bA_j^{\ln})v_j\|_{\Omega}^2 \ge (1 - \epsilon + \mathcal{O}(b^{-\rho}))\|(-i\nabla - B(p_j)bA_0)\tilde{v}_j\|_{\mathbb{R}^2_+}^2 + \mathcal{O}(\epsilon^{-1}b^{2-4\rho}\|v_j\|_{\Omega}^2).$$

where we notice that v_j is supported in $\overline{\mathbb{R}^2_+}$. Inserting this into (4.6) and using (2.3), we obtain

$$\|(-i\nabla - bA)\chi_1 u\|_{\Omega}^2 \ge (1 + \mathcal{O}(\epsilon) + \mathcal{O}(b^{-\rho})) \Big(\inf_j |B(p_j)|\Big)^{1/2} b^{1/2} \hat{\alpha} \sum_{j=1}^2 \|\tilde{v}_j\|_{\partial\mathbb{R}^2_+}^2 + \frac{\epsilon\Theta_1 b}{4} \|v\|_{\Omega}^2.$$

Step 4.

To finish the proof, we observe that

$$\sum_{j} \|\tilde{v}_{j}\|_{\partial \mathbb{R}^{2}_{+}}^{2} = \int_{\partial \Omega} \left(\sum_{j} g_{j}^{2} \right) |u|^{2} ds(x) = \|u\|_{\partial \Omega}^{2},$$

and we choose (ρ, δ) such that

$$\frac{1}{4} + \frac{\delta}{2} < \rho < \frac{\zeta - \delta}{2} \text{ and } 0 < \delta < \frac{1}{2} \left(\zeta - \frac{1}{2} \right),$$

to ensure that the conditions in (4.3) and (4.7) are respected.

We insert (4.8) into (4.4) and we recall that $\epsilon = b^{-\delta}$. Eventually, if we choose $\rho > \delta$ we obtain

$$\|(-i\nabla - bA)u\|_{\Omega}^{2} \ge \left(1 + \mathcal{O}(b^{-\delta})\right) \left(\inf_{x \in \partial\Omega} |B(x)|\right) b^{1/2} \hat{\alpha} \|u\|_{\partial\Omega}^{2}.$$

Using the variational formulation of $\lambda_1^{\text{DN}}(bA,\Omega)$ achieves the proof.

4.2. Concentration of the magnetic harmonic extension. In the case of a magnetic field which is constant on a neighborhood of $\partial\Omega$, we would like to get a more accurate lower bound for $\lambda^{\text{DN}}(bA,\Omega)$ that matches with the upper bound in Proposition 3.2. As an intermediate step, we need some information on the concentration of the magnetic harmonic extension of ground states.

For all b > 0, let f_b be a normalized eigenfunction (in $L^2(\partial \Omega)$) of the D-to-N eigenvalue $\lambda^{\text{DN}}(bA, \Omega)$. Let us denote by u_b its magnetic harmonic extension to Ω , that is

(4.9)
$$(-i\nabla - bA)^2 u_b = 0 \quad \text{on } \Omega, \quad u_b|_{\partial\Omega} = f_b.$$

The weak formulation of the D-to-N map yields

(4.10)
$$\forall v \in H^1(\Omega), \quad \langle (-i\nabla - bA)u_b, (-i\nabla - bA)v \rangle_{\Omega} - \lambda^{\mathrm{DN}}(bA, \Omega) \langle f_b, v |_{\partial\Omega} \rangle_{\partial\Omega} = 0,$$

where $\langle \cdot, \cdot \rangle_{\Omega}$ and $\langle \cdot, \cdot \rangle_{\partial\Omega}$ denote the inner product in $L^2(\Omega)$ and $L^2(\partial\Omega)$, respectively.

The next proposition states that u_b decays exponentially away from the boundary.

Proposition 4.3. Let Ω be a regular domain in \mathbb{R}^2 and A be a vector potential with a magnetic field $B = \operatorname{curl} A$ that does not vanish on $\overline{\Omega}$. Suppose that B is C^1 on $\overline{\Omega}$ and that B = 1 on a neighborhood of $\partial\Omega$. Let

$$0 < \delta < m(B,\Omega) := \inf_{x \in \overline{\Omega}} |B(x)|.$$

Then, there exists $C_{\delta}, b_{\delta} > 0$ such that, for all $b \ge b_{\delta}$, the magnetic harmonic extension u_b of f_b satisfies

$$\int_{\Omega} |u_b|^2 \exp\left(\delta b^{1/2} t(x)\right) dx \le C_{\delta} \int_{\Omega} |u_b|^2 dx,$$
$$\int_{\Omega} |(-i\nabla - bA)u_b|^2 \exp\left(\delta b^{1/2} t(x)\right) dx \le C_{\delta} b \int_{\Omega} |u_b|^2 dx,$$

where $t(x) = \operatorname{dist}(x, \partial \Omega)$.

Proof. The proof relies on the method of Agmon estimates, with due adjustments to fit the D-to-N operator.

Step 1. Link with a magnetic Robin Laplacian.

To lighten the notation, we will write $\lambda(b)$ for $\lambda^{\text{DN}}(bA, \Omega)$. Knowing from the previous sections that $\lambda(b) = \hat{\alpha}b^{1/2} + o(b^{1/2})$, we have the following lower bounds [19, Theorem 1.1 (2)], for all $\psi \in H^1(\Omega)$ with support in $\{B = 1\}$,

(4.11)
$$\|(-i\nabla - bA)\psi\|_{\Omega}^2 - \lambda(b)\|\psi\|_{\partial\Omega}^2 \ge \left(\Theta(-\hat{\alpha})b + o(b)\right)\|\psi\|_{\Omega}^2,$$

and

(4.12)
$$\|(-i\nabla - bA)\psi\|_{\Omega}^2 - 2\lambda(b)\|\psi\|_{\partial\Omega}^2 \ge \left(\Theta(-2\hat{\alpha})b + o(b)\right)\|\psi\|_{\Omega}^2,$$

where $\Theta(-\hat{\alpha}) = 0$ and $\Theta(-2\hat{\alpha}) < 0$, by the considerations in Subsection 2.2. Essentially, the lower bounds in (4.11) and (4.11) result from

$$\|(-i\nabla - bA)\psi\|_{\Omega}^{2} - \gamma \|\psi\|_{\partial\Omega}^{2} \ge \left(\Theta(\gamma, b) + o(b)\right) \|\psi\|_{\Omega}^{2},$$

the scaling relation in (2.4), and the continuity of $\Theta(\cdot)$.

Step 2.

Using (4.10) with
$$v = w^2 u_b$$
 and $w = \exp(\delta b^{1/2} t(x))$, we get by Proposition 2.3,
(4.13) $0 = \|(-i\nabla - bA)wu_b\|_{\Omega}^2 - \lambda(b)\|u_b\|_{\partial\Omega}^2 - \delta^2 b\|wu_b\|_{\Omega}^2$.

Consider R > 1 to be chosen sufficiently large, and a partition of unity

where
$$\chi_1 = 1$$
 on $\{\operatorname{dist}(x, \partial \Omega) < Rb^{-1/2}\}$, $\operatorname{supp} \chi_1 \subset \{\operatorname{dist}(x, \partial \Omega) < 2Rb^{-1/2}\}$ and
 $|\nabla \chi_1| + |\nabla \chi_2| = \mathcal{O}(R^{-1}b^{1/2}).$

We then have

$$\|(-i\nabla - bA)(wu_b)\|_{\Omega}^2 = \sum_{j=1}^2 \|(-i\nabla - bA)(\chi_j wu_b)\|_{\Omega}^2 + \mathcal{O}(R^{-2}b)\|wu_b\|_{\Omega}^2,$$

 $v_1^2 + v_2^2 = 1$ on Ω

with

$$\|(-i\nabla - bA)(\chi_2 w u_b)\|_{\Omega}^2 \ge m(B, \Omega) b \|\chi_2 w u_b\|_{\Omega}^2,$$

which follows from [9, Lemma 1.4.1], since $\chi_2 w u_b$ is compactly supported and B does not vanish in Ω

Eventually, by using (4.11) with $\psi = \chi_1 w u_b$, and noticing that $w|_{\partial\Omega} = 1$, we get from (4.13),

$$0 \ge o(b) \|\chi_1 w u_b\|_{\Omega}^2 + m(B,\Omega) b \|\chi_2 w u_b\|_{\Omega}^2 - \left(\delta^2 b + \mathcal{O}(R^{-2}b)\right) \|w u_b\|_{\Omega}^2.$$

On the support of χ_1 , we know that $w = \mathcal{O}(1)$, so we have

$$\left(m(B,\Omega) - \delta^2 + \mathcal{O}(R^{-2})\right) b \|\chi_1 w u_b\|_{\Omega}^2 \le C b \|u_b\|_{\Omega}^2,$$

where the constant C depends on δ and R.

Step 3.

To conclude, since $0 < \delta < m(B, \Omega) \leq 1$, we choose R sufficiently large so that

$$m(B,\Omega) - \delta^2 + \mathcal{O}(R^{-2}) \ge \frac{1}{2}(m(B,\Omega) - \delta^2)$$

and obtain the estimate

$$\|wu_b\|_{\Omega}^2 = \mathcal{O}(\|u_b\|_{\Omega}^2).$$

Returning to (4.13) and writing

$$\delta^2 b \|wu_b\|_{\Omega}^2 = \frac{1}{2} \|(-i\nabla - bA)(wu_b)\|_{\Omega}^2 + \frac{1}{2} \|(-i\nabla - bA)(wu_b)\|_{\Omega}^2 - \lambda(b) \|u_b\|_{\partial\Omega}^2,$$

we get from (4.12),

$$\delta^2 b \|wu_b\|_{\Omega}^2 \ge \frac{1}{2} \|(-i\nabla - bA)(wu_b)\|_{\Omega}^2 + \left(\frac{1}{2}\Theta(-2\hat{\alpha})b + o(b)\right)\|wu_b\|_{\Omega}^2,$$

which eventually yields

$$\|(-i\nabla - bA)(wu_b)\|_{\Omega}^2 = \mathcal{O}(b\|wu_b\|_{\Omega}^2)$$

			1
			1
L			1
-	-	-	

Remark 4.4. The decay estimates in Proposition 4.3 continue to hold if u_b is the magnetic harmonic extension of a $f_b \in H^{1/2}(\partial\Omega)$ with the D-to-N energy $\langle \Lambda_{bA}f_b, f_b \rangle \leq Cb^{1/2}$, where $C \in \mathbb{R}_+$.

Corollary 4.5. Given an integer $m \ge 0$, there exists $C_m, b_m > 0$ such that, for all $b \ge b_m$,

$$\int_{\Omega} |u_b|^2 (t(x))^m dx \le C_m \, b^{-\frac{m+1}{2}},$$
$$\int_{\Omega} |(-i\nabla - bA)u_b|^2 (t(x))^m dx \le C_m \, b^{-\frac{m-1}{2}}$$

Proof. Knowing that $||u_b||_{\partial\Omega} = 1$ and $\lambda^{\text{DN}}(bA, \Omega) \leq \alpha b^{1/2} + o(b^{1/2})$, (1.6) yields

$$\Theta_1 b \| u_b \|_{\Omega}^2 = \mathcal{O}(b^{1/2}).$$

The result in the corollary follows from Proposition 4.3 since $z^m \leq m! e^z$ for $z \geq 0$.

4.3. Two terms asymptotics. The main idea, following what has been done in Surface Superconductivity is to use the result in [17] for disks, the radius being locally chosen as the inverse of the curvature when it is positive. The starting point is in the case of the disk \mathcal{B}_R of radius R

(4.14)
$$\lambda^{\text{DN}}(bA, \mathcal{B}_R) = \hat{\alpha}b^{1/2} - \frac{\hat{\alpha}^2 + 1}{3}R^{-1} + \mathcal{O}(b^{-1/2}).$$

The analysis in [17] can be applied to $\mathcal{B}_R^{\text{ext}}$, the exterior of the disk \mathcal{B}_R , and we get

(4.15)
$$\lambda^{\text{DN}}(bA, \mathcal{B}_R^{\text{ext}}) = \hat{\alpha}b^{1/2} + \frac{\hat{\alpha}^2 + 1}{3}R^{-1} + \mathcal{O}(b^{-1/2}).$$

Hence we can also consider boundary points with negative curvature. We will give another proof of (4.15) below, which relies on a known result for a model with a Robin boundary condition.

To cover later all the cases with one notation we introduce for $R \in \mathbb{R}$

$$\lambda^{\mathrm{DN}}(b,R) = \begin{cases} \lambda^{\mathrm{DN}}(bA,\mathcal{B}_R) & \text{if } R > 0\\ \hat{\alpha}b^{1/2} & \text{if } R = 0\\ \lambda^{\mathrm{DN}}(bA,\mathcal{B}_{-R}^{\mathrm{ext}}) & \text{if } R < 0 \,, \end{cases}$$

and observe that

(4.16)
$$\lambda^{\text{DN}}(b,R) = \hat{\alpha}b^{1/2} - \frac{\hat{\alpha}^2 + 1}{3}R^{-1} + \mathcal{O}(b^{-1/2}).$$

We will prove the following proposition.

Proposition 4.6. Let Ω be a regular domain in \mathbb{R}^2 and A be a vector potential with a magnetic field $B = \operatorname{curl} A$ that does not vanish on $\overline{\Omega}$. Suppose that B is C^1 on $\overline{\Omega}$ and that B = 1 on a neighborhood of $\partial\Omega$. Then, the ground state energy of the D-to-N map Λ_{bA} satisfies

$$\lambda^{\mathrm{DN}}(bA,\Omega) \ge \hat{\alpha} \, b^{\frac{1}{2}} - \frac{\hat{\alpha}^2 + 1}{3} \max_{x \in \partial \Omega} k(x) + \mathcal{O}(b^{-1/6}) \quad , \quad b \to +\infty \,,$$

where k is the curvature of $\partial \Omega$.

4.3.1. Warmup. Consider $R_* > 0$, $\rho \in (0, \frac{1}{2})$ and a smooth complex-valued function \tilde{u} on $(0, t_0) \times (\mathbb{R}/L\mathbb{Z})$ such that

$$\operatorname{supp} \tilde{u} \subset [s_1, s_2] \times (0, b^{-\rho}),$$

where $|s_2 - s_1| < 2\pi R_* < L$, $b \ge b_0$ and $R_* > b^{-\rho}$. We introduce the energy

$$q_*(\tilde{u}) = \int_{\mathbb{R}/L\mathbb{Z}} \int_0^{t_0} \left(|\partial_t \tilde{u}|^2 + (1 - R_*^{-1}t)^{-2} |(-i\partial_s + b(t - \frac{1}{2}R_*^{-1}t^2)\tilde{u}| \right)^2 (1 - R_*^{-1}t) dt ds.$$

Using (4.14) and reverting to polar coordinates with the change of variables $r = R_* - t$, $\theta = s - s_1$, we get

(4.17)
$$q_*(\tilde{u}) \ge \lambda^{\mathrm{DN}}(b, R_*) \int_{\mathbb{R}/L\mathbb{Z}} |\tilde{u}(0, s)|^2 ds.$$

A similar analysis applies if $R_* \leq 0$, by using (4.16), and we find that (4.17) continues to hold in this case too.

4.3.2. A perturbed model. To deal with the exterior of a disk, or more general exterior/interior domains, we consider an approximate model with a constant curvature β .

Consider $\beta \in \mathbb{R}$, $\rho \in (\frac{1}{4}, \frac{1}{2})$, and the following energy

$$q_{\beta}^{\text{app}}(u) = \int_{\mathbb{R}} \int_{0}^{b^{-\rho}} \left(|\partial_{t}u|^{2} + (1+2t\beta) | (-i\partial_{s} + b(t-\frac{1}{2}t^{2}\beta)) u|^{2} \right) (1-t\beta) dt ds,$$

= $C^{\infty}(\mathbb{R} \times (0, b^{-\rho}))$

where $u \in C_c^{\infty}(\mathbb{R} \times (0, b^{-\rho}))$.

Lemma 4.7. There exist constants $C, b_0 > 0$, such that, for $b \ge b_0$ and $u \in C_c^{\infty}(\mathbb{R} \times (0, b^{-\rho}))$, we have

$$q_{\beta}^{\mathrm{app}}(u) \ge \left(b^{1/2}\hat{\alpha} - \frac{\hat{\alpha}^2 + 1}{3}\beta - Cb^{-\frac{1}{2}}\right) \int_{\mathbb{R}} |u(0,s)|^2 ds.$$

Moreover, the constants C, b_0 can be chosen independently of β when it varies in a bounded interval.

Proof.

Step 1.

Let us introduce

$$\lambda = \inf_{\substack{u \in C_c^\infty(\mathbb{R} \times (0, b^{-\rho})) \\ u|_{\{0\} \times \mathbb{R}} \neq 0}} \frac{q_\beta^{\mathrm{app}}(u)}{\int_{\mathbb{R}} |u(0, s)|^2 ds}.$$

Arguing as in Propositions 3.1 and Proposition 4.2, we can show that

$$\lambda = \hat{\alpha}b^{1/2} + o(b^{1/2}).$$

Step 2.

For $d \in \mathbb{R}$, let

$$\hat{\lambda}(d) = \hat{\alpha}b^{1/2} + d.$$

We introduce the ground state energy

$$G(\hat{\lambda}(d)) = \inf_{\substack{u \in C_c^{\infty}(\mathbb{R} \times (0, b^{-\rho}))\\ u \neq 0}} \frac{q_{\beta}^{\mathrm{app}}(u) - \hat{\lambda}(d) \int_{\mathbb{R}} |u(0, s)|^2 ds}{\int_{\mathbb{R}} \int_0^{b^{-\rho}} |u|^2 (1 - \beta t) dt ds}$$

With $\gamma = -b^{-1/2}\hat{\lambda}(d)$ and $\Theta(\gamma)$ as in (2.4), we know from [18, Lemma V.9]

$$G(\hat{\lambda}(d)) = \left(\Theta(\gamma)b - \beta C_1(\gamma)b^{1/2} + \mathcal{O}(b^{-\rho + \frac{1}{4}})\right) \int_{\mathbb{R}} \int_0^{b^{-\rho}} |u|^2 (1 - \beta t) dt ds,$$

where

$$C_1(\gamma) = -\left\langle \left((\tau - \xi(\gamma))^3 + \partial_\tau \right) f_\gamma, f_\gamma \right\rangle_{L^2(\mathbb{R}_+)},$$

and f_{γ} is a normalized ground state of $\Theta(\gamma)$.

This constant is calculated in [7, Lemma B.4],

(4.18)
$$C_1(\gamma) = \frac{1}{3} (1 - \gamma \xi(\gamma)) |u_{\gamma}(0)|^2.$$

Knowing that $\gamma = -\hat{\alpha} - db^{-1/2}$ and that $\Theta(\cdot)$, $\xi(\cdot)$, and u are smooth, we get by Taylor's expansion at $\hat{\alpha}$

$$\Theta(\gamma) = -d\Theta'(-\hat{\alpha})b^{-1/2} + \mathcal{O}(d^2b^{-1}),$$

$$\xi(\gamma) = \hat{\alpha} - \frac{d\Theta'(-\hat{\alpha})}{2\hat{\alpha}}b^{-1/2} + \mathcal{O}(d^2b^{-1}),$$

$$C_1(\gamma) = \frac{1}{3}(1+\hat{\alpha}^2)|u_{-\hat{\alpha}}(0)|^2 + \mathcal{O}(db^{-1/2})$$

We can choose $\zeta > 0$,

$$d = d_* = -\frac{\beta}{3}(1 + \hat{\alpha}^2) \frac{|u_{-\hat{\alpha}}(0)|^2}{\Theta'(-\hat{\alpha})} - \zeta b^{-1/2},$$

and $\hat{\lambda}_* = \hat{\lambda}(d_*)$ such that

$$G^{\operatorname{app}}(\hat{\lambda}_*) := \Theta(\gamma)b - \beta C_1(\gamma)b^{1/2} + \mathcal{O}(b^{-\rho + \frac{1}{4}}) \ge 0.$$

Step 3.

Returning to λ introduced in Step 1, and by writing

$$q^{\rm app}(u) = q^{\rm app}(u) - \hat{\lambda}_* \int_{\mathbb{R}} |u(0,s)|^2 ds + \hat{\lambda}_* \int_{\mathbb{R}} |u(0,s)|^2 ds$$

we get

$$q^{\text{app}}(u) \ge G^{\text{app}}(\hat{\lambda}_{*}) \int_{0}^{b^{-\rho}} |u|^{2} (1 - \beta t) dt ds + \hat{\lambda}_{*} \int_{\mathbb{R}} |u(0,s)|^{2} ds \ge \hat{\lambda}_{*} \int_{\mathbb{R}} |u(0,s)|^{2} ds,$$

and consequently,

$$\lambda \ge \hat{\lambda}_* = \hat{\alpha} b^{1/2} + d_*.$$

To finish the proof, notice that, by [18, Proposition II.5],

$$\Theta'(-\hat{\alpha}) = |u_{-\hat{\alpha}}(0)|^2.$$

Hence

$$d_* = -\frac{\beta}{3}(1 + \hat{\alpha}^2) - \zeta b^{-1/2} \,.$$

4.3.3. Reduction to a tubular domain. To simplify the presentation, suppose that Ω is simply connected. Let f_b be a normalized eigenfunction of the D-to-N operator, and denote by $u = u_b$ the magnetic harmonic extension of f_b . Consider a partition of unity on Ω , $\chi_1^2 + \chi_2^2 = 1$, with supp $\chi_1 \subset \{t(x) < b^{-\rho}\}$, where $t(x) = \text{dist}(x, \partial\Omega)$. Then, thanks to Proposition 4.3 and Corollary 4.5,

(4.19)
$$\lambda^{\text{DN}}(bA,\Omega) = \|(-i\nabla - bA)u\|_{\Omega}^{2} \\ = \|(-i\nabla - bA)\chi_{1}u\|_{\Omega}^{2} + \mathcal{O}(b^{-\infty}).$$

The support of $\chi_1 u$ is contained in $\Omega_0 = \{t(x) < t_0\}$, which is diffeomorphic to $\mathcal{S} := (0, t_0) \times (\mathbb{R}/L\mathbb{Z})$.

4.3.4. Reduction to constant curvature. We work with the parallel coordinates $(t, s) = \Phi_0^{-1}(x)$ and argue as in Step 2 in the proof of Proposition 4.2. In fact, we introduce a partition of unity of $\mathbb{R}/L\mathbb{Z}$

$$\sum_{j} g_j^2 = 1, \quad \sum_{j} |\nabla g_j| = \mathcal{O}(b^{\delta}), \quad \operatorname{supp} g_j \subset [s_j - b^{-\delta}, s_j + b^{-\delta}],$$

where $0 < \delta < \frac{1}{2}$ is to be chosen later on. We write

(4.20)
$$\|(-i\nabla - bA)\chi_1 u\|_{\Omega}^2 = \sum_j q(\tilde{u}_j) + \mathcal{O}(b^{2\delta} \|u\|_{\Omega}^2),$$

where

$$q(\tilde{u}_j) = \int_{\mathcal{S}} \left(|\partial_t \tilde{u}_j|^2 + (1 - tk(s))^{-2} | (-i\partial_s - b \, a(t,s)) \tilde{u}_j|^2 \right) (1 - tk(s)) ds dt,$$

and

$$a(s,t) = -t + \frac{1}{2}t^2k(s), \quad \tilde{u}_j = g_j(s)e^{i\varphi_j(s,t)}(\chi_1 u \circ \Phi_0^{-1})(s,t).$$

Writing

$$a_j(s,t) = -t + \frac{1}{2}R_j^{-1}t^2, \quad R_j^{-1} = \begin{cases} k_j & \text{if } k_j \neq 0\\ 0 & \text{if } k_j = 0 \end{cases}, \quad k_j = k(s_j),$$

we have

$$\begin{split} a(s,t) &= a_j(s,t) + \mathcal{O}(b^{-\delta}t^2), \\ 1 - tk(s) &= 1 - R_j^{-1} + \mathcal{O}(b^{-\delta}t), \\ (1 - tk(s))^{-2} &= (1 - R_j^{-1}t)^{-2} + \mathcal{O}(b^{-\delta}t), \end{split}$$

and

$$q(\tilde{u}_j) = q_j(\tilde{u}_j) + \mathcal{O}(E_{j,1}) + \mathcal{O}(E_{j,2}) + \mathcal{O}(E_{j,3})$$

where

$$\begin{split} q_{j}(\tilde{u}_{j}) &= \int_{\mathcal{S}} \Big(|\partial_{t}\tilde{u}|^{2} + (1 - R_{j}^{-1}t)^{-2} |(-i\partial_{s} + b(t - \frac{1}{2}R_{j}^{-1}t^{2})\tilde{u}|^{2} \Big) (1 - R_{j}^{-1}t) dt ds \\ E_{j,1} &= b^{-\delta} \int \Big(|\partial_{t}\tilde{u}_{j}|^{2} + |(-i\partial_{s} - b\,a)\tilde{u}_{j}|^{2} \Big) t\, dt ds, \\ E_{j,2} &= b^{2-2\delta} \int |\tilde{u}_{j}|^{2} t^{4} dt ds, \\ E_{j,3} &= b^{1-\delta} ||t^{2}\tilde{u}_{j}||_{2} ||(-i\partial_{s} - b\,a)\tilde{u}_{j}||_{2}. \end{split}$$

Using Corollary 4.5, we write

$$\sum_{j} E_{j,1} = \mathcal{O}(b^{-\delta}), \quad \sum_{j} E_{j,2} = \mathcal{O}(b^{-\frac{1}{2}-2\delta}), \quad \sum_{j} E_{j,3} = \mathcal{O}(b^{-\delta}),$$

and by using (4.17), we get

$$q(\tilde{u}_j) \ge \lambda^{\mathrm{DN}}(b, R_j) \int_{\mathbb{R}/L\mathbb{Z}} |\tilde{u}_j(0, s)|^2 ds + \mathcal{O}(b^{-\delta}) \quad \text{if } k_j > 0,$$

and a similar estimate when $k_j \leq 0$. Inserting this into (4.20) and noticing that,

$$\sum_{j} \int_{\mathbb{R}/L\mathbb{Z}} |\tilde{u}_{j}(0,s)|^{2} ds = \int_{\partial\Omega} |u|^{2} ds = 1,$$
$$b^{2\delta} ||u||_{\Omega}^{2} = \mathcal{O}(b^{-\frac{1}{2}+2\delta}) \text{ by Corollary 4.5,}$$

we get

$$\|(-i\nabla - bA)\chi_1 u\|^2 \ge \inf_j \lambda^{\mathrm{DN}}(b, R_j) + \mathcal{O}(b^{-\delta}) + \mathcal{O}(b^{2\delta - \frac{1}{2}}).$$

By invoking (4.16) and using that

$$R_j^{-1} = k(s_j) \le \max_{x \in \partial \Omega} k_x,$$

we get

$$\inf_{j} \lambda^{\mathrm{DN}}(b, R_j) \ge \hat{\alpha} b^{\frac{1}{2}} - \frac{\hat{\alpha}^2 + 1}{3} \max_{x \in \partial \Omega} k_x + \mathcal{O}(b^{-1/2}).$$

Optimizing over δ , we choose $\delta = \frac{1}{6}$, and we get from (4.19)

$$\lambda^{\mathrm{DN}}(bA,\Omega) \ge \hat{\alpha}b^{\frac{1}{2}} - \frac{\hat{\alpha}^2 + 1}{3} \max_{x \in \partial\Omega} k_x + \mathcal{O}(b^{-\frac{1}{6}}).$$

5. Splitting of eigenvalues in 2D

Assumption 5.1. Suppose that Ω is a regular domain such that the curvature k has a unique maximum of curvilinear coordinate 0, and that $k_2 := -k''(0) > 0$.

Theorem 5.2. Suppose that Assumption 5.1 holds. Let A be a vector field on Ω with constant magnetic field curl A = 1. Then, for every $j \in \mathbb{N}$, the j'th eigenvalue of the D-to-N map satisfies

$$\lambda_j(bA,\Omega) = \hat{\alpha}b^{\frac{1}{2}} - \frac{\hat{\alpha}^2 + 1}{3} \max_{x \in \partial\Omega} k(x) + (2j-1)c_*b^{-1/4} + o(b^{-1/4}) \quad as \ b \to +\infty,$$

where $c_* = c_*(\Omega) > 0$ is a constant.

5.1. Robin eigenvalues. We will establish a link between the D-to-N eigenvalues, and the eigenvalues of the magnetic Laplacian with Robin boundary condition. For $j \in \mathbb{N}$ and $\gamma \in \mathbb{R}$, we introduce the eigenvalues

$$\mu_{j}^{\mathrm{R}}(\gamma, b) = \min_{\dim(M)=j} \left(\max_{\|u\|_{\Omega}=1} \|(-i\nabla - bA)u\|_{\Omega}^{2} + b^{1/2}\gamma \|u\|_{\partial\Omega}^{2} \right).$$

Proposition 5.3. For all $j \in \mathbb{N}$ and b > 0, the equation

$$\mu_i^{\rm R}(\gamma, b) = 0$$

has a unique solution $\gamma_i(b)$, and $\gamma_i(b) < 0$. Furthermore,

$$\lambda_1^{\rm DN}(bA,\Omega) = -b^{1/2}\gamma_1(b),$$

and if for all $j \leq N$, the eigenvalue $\mu_j(\gamma_j(b), b)$ is simple, then

$$\lambda_j^{\rm DN}(bA,\Omega) = -b^{1/2}\gamma_j(b) \quad for \ 2 \le j \le N.$$

Proof. Suppose that j and b are fixed. The function $\gamma \mapsto \mu_j(\gamma, b)$ is increasing and we know that $\lambda_j(0, b) > 0$. By the min-max principle,

$$\mu_j^{\mathrm{R}}(\gamma, b) \le 2\mu_j^{\mathrm{R}}(\gamma, 0) + 2b^2 \|A\|_{\infty}^2,$$

and we have the spectral asymptotics [6]

$$\mu_j^{\rm R}(\gamma, 0) = -\gamma^2 + o(\gamma^2) \quad (\gamma \to -\infty).$$

This proves that $\gamma \mapsto \mu_j^{\mathrm{R}}(\gamma, b)$ has a unique zero $\gamma_j(b)$, and $\gamma_j(b) < 0$. Furthermore, by monotonicity, we get $\gamma_{j+1}(b) \leq \gamma_j(b)$.

Let ψ_j be an eigenfunction of $\mu_j(\gamma_j(b), b)$, $g_j = \psi_j|_{\partial\Omega}$ is an eigenfunction of the D-to-N operator with eigenvalue $-b^{1/2}\gamma_j(b)$. Hence, $(-b^{1/2}\gamma_j(b))$ is a non-decreasing sequence of eigenvalues of the D-to-N operator Λ_{bA} .

Consider an orthonormal basis $\{f_j : j \in \mathbb{N}\}$ in $L^2(\partial\Omega)$ such that every f_j is an eigenfunction of the D-to-N operator, corresponding to the j'th eigenvalue $\lambda_j^{\text{DN}}(bA, \Omega)$. If u_j is the magnetic harmonic extension of f_j , then it is a zero mode for the magnetic Laplacian $(-i\nabla - bA)^2$ with Robin boundary condition

$$\vec{\nu} \cdot (-i\nabla - bA)u_j = \lambda_j^{\mathrm{DN}}(bA, \Omega)u_j \quad \text{on } \partial\Omega.$$

Since $\{u_j : j \in \mathbb{N}\}$ is linearly independent, we get that $\lambda_j^{\mathrm{DN}}(bA, \Omega) = -b^{1/2}\gamma_j(b)$, for all $j \leq N$, provided that the eigenvalues $\mu_1^{\mathrm{R}}(\gamma_1(b), b), \cdots, \mu_N^{\mathrm{R}}(\gamma_N(b), b)$ are simple. \Box

5.2. **Proof of Theorem 5.2.** Suppose that $j \in \mathbb{N}$ and $\gamma \in \mathbb{R}$ are fixed. If Ω satisfies Assumption 5.1, and if the magnetic field $B = \operatorname{curl} A$ is equal to 1, it follows from [7] that

(5.1)
$$\mu_j^{\rm R}(\gamma, b) = b\Theta(\gamma) - b^{1/2}C_1(\gamma) \max_{x \in \partial\Omega} k(x) + (2j-1)b^{1/4}C_2(\gamma) + o(b^{1/4}) \quad (b \to +\infty),$$

locally uniformly with respect to γ . The coefficients in (5.1) are given as follows:

- $\Theta(\gamma)$ is introduced in (2.5);
- $C_1(\gamma)$ is introduced in (4.18);
- $C_2(\gamma) = \frac{1}{2}\sqrt{k_2C_1(\gamma)\partial_{\xi}^2\mu(\gamma,\xi(\gamma))}, \ \mu(\gamma,\xi)$ is introduced in (2.5), and $\xi(\gamma)$ is introduced in (2.6).

Thanks to Propositions 3.2, 4.6 and 5.3, the unique zero of $\mu_j^{\rm R}(\gamma, b) = 0$ satisfies

$$\gamma_j(b) = -b^{-1/2}\lambda_j^{\rm DN}(bA,\Omega) = -\hat{\alpha} + \frac{\hat{\alpha}^2 + 1}{3}k_*b^{-1/2} + o(b^{-1/2}),$$

where $k_* = \max_{x \in \partial \Omega} k(x)$. Furthermore,

$$\Theta(-\hat{\alpha}) = 0, \quad C_1(-\hat{\alpha}) = \frac{\hat{\alpha}^2 + 1}{3} \Theta'(-\hat{\alpha})$$

We write by Taylor's formula at $-\hat{\alpha}$,

$$\Theta(\gamma_j(b)) = (\gamma_j(b) + \hat{\alpha})\Theta'(-\hat{\alpha}) + \mathcal{O}(b^{-1}),$$

$$C_1(\gamma_j(b)) = C_1(-\hat{\alpha}) + \mathcal{O}(b^{-1/2}) = \frac{\hat{\alpha}^2 + 1}{3}\Theta'(-\hat{\alpha}) + \mathcal{O}(b^{-1/2}),$$

$$C_2(\gamma_j(b)) = C_2(-\hat{\alpha}) + o(1) \qquad (b \to +\infty).$$

Inserting these formulas into (5.1) and noting that $\mu_j^{\rm R}(\gamma_j(b), b) = 0$, we get

$$\gamma_j(b) = -\hat{\alpha} + \frac{\hat{\alpha}^2 + 1}{3}k_*b^{-1/2} - (2j-1)\frac{C_2(-\hat{\alpha})}{\Theta'(-\hat{\alpha})}b^{-3/4} + o(b^{-3/4}).$$

Thanks to Proposition 5.3, we finish the proof of Theorem 5.2, where the constant c_* is

$$c_* = \frac{C_2(-\hat{\alpha})}{\Theta'(-\hat{\alpha})}.$$

BERNARD HELFFER, AYMAN KACHMAR, AND FRANÇOIS NICOLEAU

6. The case of dimension 3

6.1. Introduction. In the unbounded case, the definition of the magnetic D-to-N map is not quite as simple as in the case of bounded domains. For compactly supported magnetic fields, the D-to-N operator in the half-space \mathbb{R}^3_+ was well defined in ([24], Appendix B) using the Lax-Phillips method. For such compactly supported magnetic fields, the solvability of the direct problem in an infinite slab Σ was also studied in [23]. We recall that an infinite slab is defined as

(6.1)
$$\Sigma = \{ x = (x', x_n) \in \mathbb{R}^n : x' = (x_1, \dots, x_{n-1}) \in \mathbb{R}^{n-1}, \ 0 < x_n < L \}, \ n \ge 3$$

At last, for non-compactly supported electromagnetic fields, the D-to-N map on an unbounded open set $\Omega \subset \mathbb{R}^3$ corresponding to a closed waveguide was studied in [22]. Here by closed waveguide, we mean that there exists a C^2 bounded open simply connected set $\omega \subset \mathbb{R}^2$ such that $\Omega \subset \omega \times \mathbb{R}$.

On the other hand, we do not really need to introduce the D-to-N operator in \mathbb{R}^3_+ but only use the corresponding ground state energy given by the variational approach and this does not involve explicitly a D-to-N operator. So we choose to avoid to refer to this operator in the proofs and will come back to this question which is interesting in itself in the last subsection.

6.2. The case of the half-space. Following what was done in Surface Superconductivity (see Lu-Pan [27] and Helffer-Morame [14]), we have to look at the non-homogeneous Dirichlet problem in $\mathbb{R}^3_+ = \{(x_1, x_2, x_3) \in \mathbb{R}^3, x_1 > 0\}$ for the family, parametrized by the angle $(\frac{\pi}{2} - \vartheta)$ of the magnetic field \vec{H} (considered as a vector in \mathbb{R}^3) with the normal vector (1, 0, 0) at a point of $\partial \mathbb{R}^3_+$.

As in ([17], Section 3), after scaling, we can always assume that the magnetic field b = 1. Thus, we consider the Dirichlet realization of the magnetic Laplacian in \mathbb{R}^3_+ :

(6.2)
$$H^{\text{Dir}}(\vartheta) := D_{x_1}^2 + D_{x_2}^2 + (D_{x_3} + \cos \vartheta x_1 - \sin \vartheta x_2)^2 \quad , \ \vartheta \in [0, \frac{\pi}{2}] \,,$$

and consider (we implicitly assume the condition that the denominator is not zero)

(6.3)
$$\lambda^{DN}(\vartheta) := \inf_{\phi \in C_0^{\infty}(\overline{\mathbb{R}^3_+})} \frac{||\nabla_{A_{\vartheta}}\phi||^2}{\int |\phi(0, x_2, x_3)|^2 dx_2 dx_3},$$

where

$$A = A_{\vartheta} = (0, 0, -\cos\vartheta x_1 + \sin\vartheta x_2).$$

The main result of this subsection is the following :

Proposition 6.1. One has :

(6.4)
$$\inf_{\vartheta \in [0, \frac{\pi}{2}]} \lambda^{DN}(\vartheta) = \hat{\alpha} = \alpha / \sqrt{2} \,,$$

and this infimum is uniquely realized at $\vartheta = 0$.

Remark 6.2. It follows from Proposition 6.1 that $\lambda^{DN}(\vartheta)$ is minimal when the magnetic field \vec{H} is parallel to the hyperplane $x_1 = 0$.

To prove this proposition, we first make a partial Fourier transform in the x_3 -variable and we can reduce the computation to a (ϑ, τ) -family of operators on $\mathbb{R}^2_+ = \{x_1 > 0\}$:

(6.5)
$$H^{\mathrm{Dir}}(\vartheta,\tau) := D_{x_1}^2 + D_{x_2}^2 + (\tau + \cos\vartheta x_1 - \sin\vartheta x_2)^2 , \ \tau \in \mathbb{R}.$$

We now introduce

$$\lambda^{\mathrm{DN}}(\vartheta,\tau) := \inf_{\phi \in C_0^{\infty}(\overline{\mathbb{R}^2_+})} \frac{||\nabla_{\check{A}}\phi||^2}{\int |\phi(0,x_2)|^2 dx_2} \,,$$

where

$$\dot{A} = \dot{A}_{\vartheta,\tau} = (0, 0, \tau + \cos \vartheta x_1 - \sin \vartheta x_2)$$

Our first result is

Lemma 6.3.

$$\lambda^{\mathrm{DN}}(\vartheta) := \inf_{\tau \in \mathbb{R}} \lambda^{\mathrm{DN}}(\vartheta, \tau) \,.$$

Proof. For the upper-bound, we can use sequences of the form $\chi_n(t)e^{it\tau}\theta(x_1,x_2)$. For the lower bound, we use the partial Fourier transform.

The next point is to observe that:

Lemma 6.4. If
$$\vartheta \in (0, \frac{\pi}{2}]$$
,

(6.6)
$$\lambda^{\mathrm{DN}}(\vartheta,\tau) = \lambda^{DN}(\vartheta,0)$$

Proof. This is evident through a translation in the tangential x_2 variable.

We will also need

Lemma 6.6. If $\vartheta = 0$,

Lemma 6.5. If $\vartheta \in (0, \frac{\pi}{2})$, there exists $\phi_{\vartheta} \in H^1_{\tilde{A}}(\mathbb{R}^2_+)$ such that

$$\int \phi_{\vartheta}(0, x_2)^2 dx_2 = 1$$

and

$$\lambda^{\mathrm{DN}}(\vartheta, 0) = ||\nabla_{\tilde{A}} \phi_{\vartheta}||^2$$

where

$$\tilde{A} = \check{A}_{\vartheta,0}$$
.

Proof. This is a consequence of the compact injection of $H^1_{\tilde{A}}(\mathbb{R}^2_+)$ in $L^2(\mathbb{R}^2_+)$.

Hence, the parameter τ is relevant only in the case when $\vartheta = 0$, and this is the object of the following lemma :

(6.7)
$$\lambda^{\rm DN}(0,\tau) = \hat{\alpha} \,.$$

Proof. In this case, the Laplacian becomes

(6.8)
$$H(0,\tau) = D_{x_1}^2 + D_{x_2}^2 + (\tau + x_1)^2 .$$

After a Fourier transform in the x_2 -variable, we get a new family of magnetic Laplacians :

(6.9)
$$\tilde{H}(\xi_2,\tau) = D_{x_1}^2 + \xi_2^2 + (\tau + x_1)^2 \quad , \ \xi_2 \in \mathbb{R} \, .$$

We now have to analyze the family of the associated energies $\lambda^{DN}(\xi_2, \tau)$ depending of two parameters (ξ_2, τ) . Using the variational characterization of the ground state, it is clear that the infimum is obtained for $\xi_2 = 0$, and this latest case was analyzed in ([17], Proposition 3.1). Thus, the proof is complete.

Now, let us study the case $\vartheta = \frac{\pi}{2}$. One gets :

Lemma 6.7. If $\vartheta = \frac{\pi}{2}$,

(6.10)
$$\lambda^{\mathrm{DN}}(\frac{\pi}{2},\tau) = 1.$$

Proof. By Lemma 6.4, we can always assume that $\tau = 0$. So, in this case, we have :

(6.11)
$$H\left(\frac{\pi}{2},0\right) = D_{x_1}^2 + D_{x_2}^2 + x_2^2.$$

Thanks to a decomposition using the Hermite functions basis in the x_2 variable, we have to look at the family of Hamiltonians in \mathbb{R}^+ :

(6.12)
$$\hat{H}(k) = D_{x_1}^2 + (2k+1) , \ k \in \mathbb{N}.$$

If $\lambda(k)$ denotes the lowest eigenvalue of D-to-N map associated with $\hat{H}(k)$, one easily gets $\lambda(k) = \sqrt{2k+1}$. Thus,

(6.13)
$$\lambda^{\text{DN}}\left(\frac{\pi}{2},\tau\right) = \lambda^{DN}\left(\frac{\pi}{2},0\right) = \inf_{k \ge 0} \sqrt{2k+1} = 1.$$

End of the proof of Proposition 6.1:

Let us set $\Omega = \{(x_1, x_2) \in \mathbb{R}^2, x_1 > 0\}$. In the following, we shall show that for any $\vartheta \in (0, \frac{\pi}{2})$, there exists a suitable $g(\vartheta) > \hat{\alpha}$ such that, for any $u \in C^{\infty}(\overline{\Omega})$,

(6.14)
$$\int_{\Omega} |\nabla u|^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 dx_2 \ge g(\vartheta) \int_{\partial \Omega} |u(0, x_2)|^2 dx_2.$$

To this end, we follow a similar strategy to ([14], Subsection 3.4). We introduce an interpolation parameter $\rho \in [0, 1]$, and we write the integrand in the (LHS) of (6.14) as

$$\begin{aligned} |\nabla u|^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 &= \rho^2 \left(|\partial_{x_1} u|^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 \right) \\ &+ (1 - \rho^2) |\partial_{x_1} u|^2 + |\partial_{x_2} u|^2 + (1 - \rho^2) (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 \,. \end{aligned}$$

First, thanks to ([17], Proposition 3.1), one immediately gets :

(6.15)
$$\int_0^{+\infty} \left(|\partial_{x_1} u|^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 \right) dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u(0, x_2)^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 dx_1 \ge \hat{\alpha} \sqrt{\cos \vartheta} |u|^2 + \hat{\alpha} \sqrt{\cos \vartheta}$$

Integrating with respect to the x_2 variable, we obtain :

$$(6.16) \quad \rho^2 \int_{\Omega} \left(|\partial_{x_1} u|^2 + (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 \right) \, dx_1 dx_2 \geq \rho^2 \hat{\alpha} \sqrt{\cos \vartheta} \, \int_{\partial \Omega} |u(0, x_2)|^2 \, dx_2 \, .$$

Secondly, we observe that :

(6.17)
$$\int_{\mathbb{R}} \left(|\partial_{x_2} u|^2 + (1-\rho^2)(x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 \right) dx_2 \geq \sqrt{1-\rho^2} \sin \vartheta \int_{\mathbb{R}} |u|^2 dx_2.$$

This last inequality is a consequence of the lower bound for the harmonic oscillator, (in the x_2 variable). On one hand, integrating (6.17) with respect to the x_1 variable, one easily gets :

$$\int_{\Omega} \left((1-\rho^2) |\partial_{x_1} u|^2 + |\partial_{x_2} u|^2 + (1-\rho^2) (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 \right) dx_1 dx_2$$

$$\geq (1-\rho^2) \left(\int_{\Omega} |\partial_{x_1} u|^2 + (1-\rho^2)^{-\frac{1}{2}} \sin \vartheta |u|^2 dx_1 dx_2 \right).$$

On the other hand, using the following lower bound for the D-to-N map associated with the Hamiltonian $-\partial_{x_1}^2 + (1-\rho^2)^{-\frac{1}{2}}\sin\vartheta$ on the interval $(0, +\infty)$,

(6.18)
$$\int_{0}^{+\infty} |\partial_{x_{1}}u|^{2} + (1-\rho^{2})^{-\frac{1}{2}}\sin\vartheta \ |u|^{2} \ dx_{1} \ge (1-\rho^{2})^{-\frac{1}{4}}\sqrt{\sin\vartheta} \ |u(0,x_{2})|^{2},$$

and integrating again over the variable x_2 , we finally get

$$\int_{\Omega} \left((1-\rho^2) |\partial_{x_1} u|^2 + |\partial_{x_2} u|^2 + (1-\rho^2) (x_1 \cos \vartheta - x_2 \sin \vartheta)^2 |u|^2 \right) \, dx_1 dx_2 \\ \geq (1-\rho^2)^{\frac{3}{4}} \sqrt{\sin \vartheta} \int_{\partial \Omega} |u(0,x_2)|^2 \, dx_2 \, .$$

As a conclusion, we have obtained :

(6.19)
$$\lambda^{\mathrm{DN}}(\vartheta) \ge \left(\rho^2 \hat{\alpha} \sqrt{\cos \vartheta} + (1-\rho^2)^{\frac{3}{4}} \sqrt{\sin \vartheta}\right)$$

In particular, choosing $\rho = \cos \vartheta$, we get

(6.20)
$$\lambda^{\mathrm{DN}}(\vartheta) \ge g(\vartheta) := \hat{\alpha}(\cos\vartheta)^{\frac{5}{2}} + (\sin\vartheta)^2 > \hat{\alpha} , \ \forall \vartheta \in (0, \frac{\pi}{2}].$$

For the last inequality, we can observe that with $X = \cos \vartheta \in [0, 1]$, we have always:

$$\hat{\alpha}X^{5/2} + 1 - X^2 > \hat{\alpha}$$
.

This concludes the proof of Proposition 6.1.

6.3. Lower bounds in general domains. We have

Proposition 6.8. Let Ω be a regular bounded domain in \mathbb{R}^3 and A be a magnetic potential with constant magnetic field with norm 1, then the ground state energy of the D-to-N map Λ_{bA}^{DN} satisfies

(6.21)
$$\liminf_{b \to +\infty} b^{-1/2} \lambda^{\mathrm{DN}}(bA, \Omega) \ge \hat{\alpha}$$

More generally, we have

Proposition 6.9. Let Ω be a regular bounded domain in \mathbb{R}^3 , A be a magnetic potential with non vanishing C^{∞} magnetic field in $\overline{\Omega}$, then, with the notation of Theorem 1.6, the ground state energy of the D-to-N map Λ_{bA}^{DN} satisfies

(6.22)
$$\liminf_{b \to +\infty} b^{-1/2} \lambda^{\mathrm{DN}}(bA, \Omega) \ge \inf_{x \in \partial\Omega} \left(\lambda^{DN}(\vartheta(x)) |B(x)|^{\frac{1}{2}} \right).$$

(see also [27] or [14] and (B.1) in Appendix B)

Proof. We follow the proof of Proposition 1.4 given in the case of dimension 2. For step 1, we have to replace the 2D lower bound of the Neumann problem by the (3D)-statement proven by Lu-Pan [27]

(6.23)
$$\liminf_{b \to +\infty} b^{-1} \lambda^{\operatorname{Ne}}(bA, \Omega) \ge \min\left(\inf_{x \in \overline{\Omega}} |B(x)|, \inf_{x \in \partial\Omega} \sigma(\vartheta(x))|B(x)|\right),$$

or a more accurate version with remainder.

For step 2, the equivalent of Proposition 2.4 is given in Lemma 5.4 from [27].

Finally, we can implement the constant magnetic field results obtained in the previous subsection.

6.4. Upper bounds in general domains. As observed in [33] the proof for the Neumann problem is only sketched in [26]. On the other hand, we only state the following version, with the notation of Proposition 6.9

Proposition 6.10. Let Ω be a regular bounded domain in \mathbb{R}^3 , A be a magnetic potential with non vanishing C^{∞} magnetic field in $\overline{\Omega}$, then the ground state energy of the D-to-N map Λ_{bA}^{DN} satisfies

(6.24)
$$\limsup_{b \to +\infty} b^{-1/2} \lambda^{\mathrm{DN}}(bA, \Omega) \le \inf_{x \in \partial\Omega} \left(\lambda^{DN}(\vartheta(x)) |B(x)|^{\frac{1}{2}} \right).$$

Proof. We distinguish three cases depending on the value of $\vartheta(x)$ where $\inf_{x \in \partial \Omega} \left(\lambda^{DN}(\vartheta(x)) |B(x)|^{\frac{1}{2}} \right)$ is attained.

Case 1. We assume that there exists $p \in \partial \Omega$ such that $\vartheta(p) \in (0, \frac{\pi}{2})$ and

(6.25)
$$\lambda^{\mathrm{DN}}(\vartheta(p))|B(p)|^{\frac{1}{2}} = \inf_{x \in \partial\Omega} \left(\lambda^{\mathrm{DN}}(\vartheta(x))|B(x)|^{\frac{1}{2}}\right).$$

In this case, after using Lemma 3.4 in [27] (which extends proposition 2.4 to the (3D)-case), we can take in the new system of coordinates centered at p such that Ω is locally defined by $\{x_1 > 0\}$, the quasimode

$$u(x_1, x_2, x_3) = b^{\frac{\nu}{2}} \chi(b^{\rho} x_3) \chi(b^{\rho} x_1) \chi(b^{\rho} x_2) \phi_{\vartheta(p)}((bB(p)^{1/2} x_1, (bB(p))^{1/2} x_2))$$

where ϕ_{ϑ} is defined in Lemma 6.5 and $\rho \in (0, \frac{1}{2})$. **Case 2**. We assume that $\vartheta(p) = \frac{\pi}{2}$ in (6.24). We take as quasi-mode

$$u(x_1, x_2, x_3) = b^{\frac{\rho}{2}} \chi(b^{\rho} x_3) \chi(b^{\rho} x_1) \chi(b^{\rho} x_2) (bB(p))^{1/2} \exp\left(-\frac{1}{2} (bB(p)) x_2^2\right) \exp\left(-(bB(p))^{1/2} x_1\right) + \frac{1}{2} \exp\left(-\frac{1}{2} (bB(p)) x_2^2\right) \exp\left(-\frac{1}{2} (bB(p)) x_$$

Case 3. We assume that $\vartheta(p) = 0$ in (6.24).

Notice that this is always the case when the magnetic field is constant. We are essentially like in the (2D) case and take

$$u(x_1, x_2, x_3) = b^{\rho} \chi(b^{\rho} x_2) \chi(b^{\rho} x_3) \cdot \chi(b^{\rho} x_1) \cdot f_* \big((bB(p))^{1/2} x_1 \big) \cdot e^{-i(bB(p)x_1)^{1/2} \hat{\alpha} x_3}.$$

6.5. On the D-to-N operator relative to \mathbb{R}^3_+ . Let us denote $H^1_A(\mathbb{R}^3_+) = \{ u \in L^2(\mathbb{R}^3_+), D_{x_1}u \in L^2(\mathbb{R}^3_+), D_{x_2}u \in L^2(\mathbb{R}^3_+), (D_{x_3} + \cos \vartheta x_1 - \sin \vartheta x_2)u \in L^2(\mathbb{R}^3_+) \} \}.$ Since $H^1_A(\mathbb{R}^3_+) \subset H^1_{\text{loc}}(\overline{\mathbb{R}^3_+})$, we can define the trace space

$$\check{H}^{\frac{1}{2}}(\mathbb{R}^2) = \{ u \in L^2_{loc}(\mathbb{R}^2), \, \exists \tilde{u} \in H^1_A(\mathbb{R}^3_+) \text{ s.t. } \tilde{u}_{x_1=0} = u \}$$

To define the D-to-N operator, we can now start "formally" from the weak form given in (1.4) with $\Omega = \mathbb{R}^3_+$ and $A = A_\vartheta$:

(6.26)
$$\langle \Lambda_A f, g \rangle_{\check{H}^{-1/2}(\partial\Omega) \times \check{H}^{1/2}(\partial\Omega)} = \int_{\Omega} \langle (-i\nabla - A)u, (-i\nabla - A)v \rangle \, dx \, ,$$

for any $g \in \check{H}^{1/2}(\partial \Omega)$ and $f \in \check{H}^{1/2}(\partial \Omega)$ such that u is the unique solution see (6.2) of

$$H^{\mathrm{Dir}}(\vartheta)u = 0, \ u_{\partial\Omega} = f,$$

and v is any element of $H^1(\Omega)$ so that $v_{|\partial\Omega} = g$.

Remark 6.11. It would be interesting to verify various technical details in order to associate a self-adjoint realization to this weak definition of the D-t-N operator in 3D but this is not needed for the results presented in this paper.

7. WEAK MAGNETIC FIELD.

Proof of Theorem 1.8.

Step 1.

Since Ω is simply connected, $\lambda^{\text{DN}}(bA,\Omega) = \lambda^{\text{DN}}(bA_{\Omega},\Omega)$. We write $\hat{\lambda} = \lambda^{\text{DN}}(bA_{\Omega},\Omega)$. Using f = 1 as trial state in (1.6), we get

$$0 \le \hat{\lambda} \le \frac{b^2}{|\partial \Omega|} \int_{\Omega} |A_{\Omega}|^2 dx.$$

Step 2.

We introduce $E_1(\hat{\lambda}, b) = \mu(bA_{\Omega}, \Omega, \hat{\lambda})$ as in (1.15) and we observe that $E_1(\hat{\lambda}, b) = 0$ (see Proposition 5.3 with $\gamma = b^{-1/2}\hat{\lambda}$). The second eigenvalue satisfies

$$\liminf_{b \to 0^+} E_2(\hat{\lambda}, b) > 0,$$

which follows from the min-max principle and the following lower bound

$$\||(-i\nabla - bA_{\Omega})f\|_{\Omega}^{2} - \hat{\lambda}\|f\|_{\partial\Omega}^{2} \ge \left(\frac{1}{2}\|\nabla f\|_{\Omega}^{2} - 2\hat{\lambda}\|f\|_{\partial\Omega}^{2}\right) - 2b^{2}\|A_{\Omega}f\|_{\Omega}^{2}.$$

Step 3.

Let $f_{\hat{\lambda}}$ be the positive and normalized (in $L^2(\Omega)$) ground state of $E_1(\hat{\lambda}, 0)$. By [10, Theorem 2.1],

$$E_1(\hat{\lambda}, 0) = -\frac{|\partial \Omega|}{|\Omega|}\hat{\lambda} + o(\hat{\lambda}) \quad (\hat{\lambda} \to 0).$$

Writing

$$E_1(\hat{\lambda}, 0) = \|\nabla f_{\hat{\lambda}}\|_{\Omega}^2 - \hat{\lambda} \|f_{\hat{\lambda}}\|_{\partial\Omega}^2 \ge \frac{1}{2} \|\nabla f_{\hat{\lambda}}\|_{\Omega}^2 + \frac{1}{2} E_1(2\hat{\lambda}, 0),$$

we deduce that

$$\frac{1}{2} \|\nabla f_{\hat{\lambda}}\|_{\Omega}^2 \le E_1(\hat{\lambda}, 0) - \frac{1}{2} E_1(2\hat{\lambda}, 0) = o(\hat{\lambda}),$$

and by the Poincaré inequality

$$\|f_{\hat{\lambda}} - \langle f_{\hat{\lambda}} \rangle\|_{\Omega}^2 = o(\hat{\lambda}),$$

where $\langle f_{\hat{\lambda}} \rangle$ is the average of $f_{\hat{\lambda}}$ over Ω . The normalization of $f_{\hat{\lambda}}$ in $L^2(\Omega)$ then yields

$$\langle f_{\hat{\lambda}} \rangle = \frac{1}{|\Omega|^{1/2}} + o(\hat{\lambda}).$$

Step 4.

Since div $A_{\Omega} = 0$, it follows from (1.1) that

$$H_{bA_{\Omega}}f_{\hat{\lambda}} = E_1(\hat{\lambda}, 0)f_{\hat{\lambda}} + b^2|A_{\Omega}|^2f_{\hat{\lambda}} - 2ibA_{\Omega}\cdot\nabla f_{\hat{\lambda}}.$$

Moreover, since $\vec{\nu}\cdot A_\Omega=0,\,f_{\hat{\lambda}}$ satisfies the boundary condition

$$\partial_{\nu} f_{\hat{\lambda}} + i\vec{\nu} \cdot A_{\Omega} f_{\hat{\lambda}} = \hat{\lambda} f_{\hat{\lambda}} \text{ on } \partial\Omega$$

By Step 3, we have

$$\|2ibA_{\Omega}\cdot\nabla f_{\hat{\lambda}}\|_{\Omega} \le b\|\nabla f_{\hat{\lambda}}\|_{\Omega} = o(b\hat{\lambda}^{1/2}).$$

We introduce the real-valued function $g_{\hat{\lambda}}$ as the solution of

$$-\Delta g_{\hat{\lambda}} - C(\hat{\lambda})f_{\hat{\lambda}} + |A_{\Omega}|^2 f_{\hat{\lambda}} = 0 \text{ on } \Omega, \quad \partial_{\nu}g_{\hat{\lambda}} = \hat{\lambda}g_{\hat{\lambda}} \text{ on } \partial\Omega,$$

where

$$C(\hat{\lambda}) = \int_{\Omega} |A_{\Omega}|^2 f_{\hat{\lambda}}^2 dx = \frac{1}{|\Omega|} \int_{\Omega} |A_{\Omega}|^2 dx + o(\hat{\lambda}).$$

With $u = f_{\hat{\lambda}} + b^2 g_{\hat{\lambda}}$, we have

$$H_{bA_{\Omega}}u = \left(E_1(\hat{\lambda}, 0) + b^2 C(\hat{\lambda})\right)f_{\hat{\lambda}} - 2ibA_{\Omega} \cdot \nabla f_{\hat{\lambda}} + b^2 \left(b^2 |A_{\Omega}|^2 - 2iA_{\Omega} \cdot \nabla\right)g_{\hat{\lambda}}.$$

By Step 1, this yields

$$\left\| H_{bA_{\Omega}}u - \left(E_1(\hat{\lambda}, 0) + \frac{b^2}{|\Omega|} \|A_{\Omega}\|_{\Omega}^2 \right) u \right\|_{\Omega} = o(b^2) \quad (b \to 0^+),$$

and by Step 2, the spectral theorem yields

$$E_1(\hat{\lambda}, b) = E_1(\hat{\lambda}, 0) + \frac{b^2}{|\Omega|} ||A_\Omega||_{\Omega}^2 + o(b^2)$$

34

Step 5.

To finish the proof, we use that $E_1(\hat{\lambda}, b) = 0$ and that $\hat{\lambda} = O(b^2)$, along with the asymptotics of $E_1(\hat{\lambda}, 0)$ in Step 2.

Acknowledgments. A.K. is partially supported by CUHK-SZ grant no. UDF01003322 and UF02003322. F.N is supported by the French National Research Project GDR Dynqua. The authors would like to thank Vladimir Lotoreichik for the helpful comments.

Appendix A. On the choice of gauge

Considering A as a 1-form is helpful in passing from the Cartesian to parallel coordinates,

$$A_1dx_1 + A_2dx_2 = \tilde{A}_1dt + \tilde{A}_2ds,$$

where

$$\partial_t \tilde{A}_2 - \partial_s \tilde{A}_1 = -(1 - tk(s))B,$$

and $B = \operatorname{curl} A$ is the magnetic field in Cartesian coordinates.

Suppose that B is constant. Choosing a simply connected set $V \subset \mathbb{R}/L\mathbb{Z}$, we define a function $\varphi : \mathbb{R}/L\mathbb{Z} \times V$ such that

$$\partial_t \varphi = \tilde{A}_1, \quad \partial_s \varphi - B(t - \frac{1}{2}t^2k(s)) = \tilde{A}_2(s,t) \quad \text{on } V.$$

Such a function is given by

$$\varphi(t,s) = \int_0^t \tilde{A}_1(\tau,s)d\tau + g(s),$$

where $g: \mathbb{R}/L\mathbb{Z} \to \mathbb{R}$ satisfies $g'(s) = \tilde{A}_2(0, s)$ on V, and g(s) = 0 outside a neighborhood of V; this last condition ensures the periodicity of g.

That this is relevant is apparent from

$$\begin{split} \|(-i\nabla - bA)u\|_{\Phi_0^{-1}(V)}^2 &= \int_V \left(|(-i\partial_t - b\tilde{A}_1)\tilde{u}|^2 + (1 - tk(s))^{-2} |(-i\partial_s - b\tilde{A}_2)\tilde{u}|^2 \right) (1 - tk(s)) dt ds \\ &= \int_V \left(|\partial_t (e^{-ib\varphi}\tilde{u})|^2 + (1 - tk(s))^{-2} |(-i\partial_s + bB(t - \frac{1}{2}t^2k(s)))e^{-2ib\varphi}\tilde{u}|^2 \right) (1 - tk(s)) dt ds \end{split}$$

APPENDIX B. A REMINDER FOR MODELS IN HALF-SPACES.

We refer to [27] and [14] for the proof of the results which are recalled here as presented in [15]. If N is a unit vector in \mathbb{R}^3 , we now consider the Neumann realization in $\Omega := \{x \in \mathbb{R}^3 \mid x \cdot N > 0\}$. After a rotation, we can assume in the proofs that N = (1, 0, 0), so Ω is $\mathbb{R}^3_+ := \{x_1 > 0\}$. After scaling, we can assume that h = 1 and |H| = 1. Here \vec{H} is the magnetic vector field curl \vec{A}

associated with the magnetic field B considered as a 2-form.

After some rotation in the (x_2, x_3) variables, we can assume that the new magnetic field is $(\beta_1, \beta_2, 0)$ and we are reduced to the problem of analyzing :

$$P(\beta_1, \beta_2) := D_{x_1}^2 + D_{x_2}^2 + (D_{x_3} + \beta_2 x_1 - \beta_1 x_2)^2 ,$$

in $\{x_1 > 0\}$, where :

$$\beta_1^2 + \beta_2^2 = 1 \; .$$

We introduce :

$$\beta_2 = \cos \vartheta , \ \beta_1 = \sin \vartheta ,$$

and we observe that, if N is the external normal to $x_1 = 0$, we have :

(B.1)
$$\langle \vec{H} | N \rangle = -\sin \vartheta$$

By partial Fourier transform, we arrive to :

(B.2)
$$L(\vartheta, \tau) = D_{x_1}^2 + D_{x_2}^2 + (\tau + \cos \vartheta \, x_1 - \sin \vartheta \, x_2)^2 ,$$

in $x_1 > 0$ and with Neumann condition on $x_1 = 0$. The bottom of the spectrum of $L(\vartheta, \tau)$ is given by :

(B.3)
$$\sigma(\vartheta) := \inf \operatorname{sp}_{\operatorname{ess}} \left(L(\vartheta, D_t) \right) = \inf_{\tau} \left(\inf \operatorname{sp}_{\operatorname{ess}}(L(\vartheta, \tau)) \right) \,.$$

Proposition B.1. .

The bottom of the spectrum of the Neumann realization of $H_{A,\Omega}^{\text{Ne}}$ in $\Omega := \{x \in \mathbb{R}^3 \mid x \cdot N > 0\}$ is :

(B.4) $\inf \, \operatorname{sp}_{\operatorname{ess}} H^{\operatorname{Ne}}_{A,\Omega} = \sigma(\vartheta) \, b \, h \; ,$

where $\vartheta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ is defined by (B.1).

By symmetry considerations, we observe also that :

(B.5)
$$\sigma(\vartheta) = \sigma(-\vartheta) = \sigma(\pi - \vartheta)$$

It is consequently enough to look at the restriction to $[0, \frac{\pi}{2}]$.

B.1. **Properties of** $\vartheta \mapsto \sigma(\vartheta)$. Let us now list the main properties of the function $\vartheta \mapsto \sigma(\vartheta)$ on $[0, \frac{\pi}{2}]$. Most of them are established in [27] but see also [14].

(1) σ is continuous on $[0, \frac{\pi}{2}]$.

(2)

$$(B.6) \sigma(0) = \Theta_0 < 1 .$$

(3)

(B.7)
$$\sigma(\frac{\pi}{2}) = 1$$

(4)

(B.8)
$$\sigma(\vartheta) \ge \Theta_0(\cos\vartheta)^2 + (\sin\vartheta)^2 .$$

- (5) If $\vartheta \in]0, \frac{\pi}{2}[$, the spectrum of $L(\vartheta, \tau)$ is independent of τ and its essential spectrum is contained in $[1, +\infty]$.
- (6) For $\vartheta \in]0, \frac{\pi}{2}[, \sigma(\vartheta)]$ is an isolated eigenvalue of $L(\vartheta, \tau)$, with multiplicity one.
- (7) The function σ is strictly increasing on $[0, \frac{\pi}{2}]$.

An immediate consequence of this analysis is

36

Proposition B.2.

When b = |H| is fixed the bottom of the spectrum of $H_{A,\Omega}^{\text{Ne}}$ in $\Omega := \{x \cdot N > 0\}$ is minimal when $\vartheta = 0$ that is, according to (B.1), when the magnetic field vector satisfies $\vec{H} \cdot N = 0$.

References

- W. Assaad. The breakdown of superconductivity in the presence of magnetic steps. Commun. Contemp. Math. 23 (2021), Paper No. 2050005, 53 pp.
- W. Assaad and A. Kachmar. Lowest energy band function for magnetic steps. J. Spectr. Theory 12 (2022), 813-833.
- [3] A. Bernoff and P. Sternberg. Onset of superconductivity in decreasing fields for general domains. J. Math. Phys. 39 (1998), 1272–1284.
- [4] T. Chakradhar, K. Gittins, G. Habib, and N. Peyerimhoff. A note on the magnetic Steklov operator on functions. arXiv:2410.07462 v1 and v2 (2024).
- [5] B. Colbois, C. Léna, L. Provenzano, A. Savo. A reverse Faber-Krahn inequality for the magnetic Laplacian. J. Math. Pures Appl. 192 (2024), article no. 103632.
- [6] D. Daners and J.B. Kennedy. On the asymptotic behaviour of the eigenvalues of a Robin problem. Differential Integral Equations 23 (2010), 659–669.
- [7] R. Fahs, L. Le Treust, N. Raymond, and S. Vũ Ngọc. Boundary states of the Robin magnetic Laplacian. Doc. Math. 29 (2024), 1157–1200.
- [8] S. Fournais and B. Helffer. Inequalities for the lowest magnetic Neumann eigenvalue. Lett. Math. Phys. 109 (2019), 1683-1700.
- [9] S. Fournais and B. Helffer. Spectral Methods in Surface Superconductivity. Progress in Nonlinear Differential Equations and Their Applications, Vol. 77, Birkhäuser, (2010).
- [10] T. Giorgi and R. Smits. Eigenvalue estimates and critical temperature in zero fields for enhanced surface superconductivity. Z. Angew. Math. Phys. 58 (2007), 224-245.
- B. Helffer and A. Kachmar. Helical magnetic fields and semi-classical asymptotics of the lowest eigenvalue. Doc. Math. 28 (2023), 857–901.
- [12] B. Helffer, A. Kachmar and N. Raymond. Magnetic confinement for the 3D Robin Laplacian. Pure Appl. Funct. Anal. 7 (2022), 601–639
- [13] B. Helffer and A. Morame. Magnetic bottles in connection with superconductivity. J. Funct. Anal. 185 (2001), 604–680.
- [14] B. Helffer and A. Morame. Magnetic bottles for the Neumann problem: the case of dimension 3. Proc. Indian Acad. Sci. 112 (2002), 71–84.
- [15] B. Helffer and A. Morame. Magnetic bottles for the Neumann problem: curvature effects in the case of dimension 3 (general case) Ann. Scient. Éc. Norm. Sup., 4e série, 37 (2004), 105–170.
- [16] B. Helffer and F. Nicoleau. Trace formulas for the magnetic Dirichlet to Neumann operator. ArXiv:2407.08671, to appear in Asymptotic Analysis.
- [17] B. Helffer and F. Nicoleau. On the magnetic Dirichlet to Neumann operator on the disk- strong diamagnetism and strong magnetic field limit- ArXiv:2411.15522 v2, (2024).
- [18] A. Kachmar. On the ground state energy for a magnetic Schrödinger operator and the effect of the de Gennes boundary condition. J. Math. Phys. 47 (2006), art. no. 072106.
- [19] A. Kachmar. Diamagnetism versus Robin condition and concentration of ground states. Asymptot. Anal. 98 (2016), no. 4, 341–375.
- [20] A. Kachmar, V. Lotoreichik. On the isoperimetric inequality for the magnetic Robin Laplacian with negative boundary parameter. J. Geom. Anal. 32 (2022), 182; Correction 32 (2022), 237.
- [21] A. Kachmar, V. Lotoreichik. A geometric bound on the lowest magnetic Neumann eigenvalue via the torsion function. SIAM J. Math. Anal. 56 (2024), 5723–5745.
- [22] Y. Kian. Determination of non-compactly supported electromagnetic potentials in unbounded closed waveguide. Revista Matemàtica Iberoamericana, 36 (2020), 671-710.

- [23] K. Krupchyk, M. Lassas, G. Uhlmann. Inverse problems with partial data for a magnetic Schrödinger operator in an Infinite slab or bounded domain. Comm. Math. Phys., 327 (2014), 993-1009.
- [24] X. Li. Inverse boundary value problems with partial data in unbounded domains. Inverse Problems, 28 (2012), 085003.
- [25] K. Lu and X. Pan. Gauge invariant eigenvalue problems in \mathbb{R}^2 and in \mathbb{R}^2_+ . Trans. Amer. Math. Soc. 352 (2000), 1247–276.
- [26] K. Lu and X. Pan. Eigenvalue problems of Ginzburg-Landau operator in bounded domains. Journal of Mathematical Physics 4 (1999), 2647–2667.
- [27] K. Lu and X. Pan. Surface nucleation of superconductivity in 3-dimensions. J. of Diff. Equations 168 (2000), 386–452.
- [28] W. Magnus, F. Oberhettinger and R.P. Soni. Formulas and theorems for the special functions of mathematical physics, 3rd enlarged ed, Grundlehren der Mathematischen Wissenschaften, Volume 52, Springer, (1965).
- [29] X.B. Pan. An eigenvalue variation problem of magnetic Schrödinger operator in three dimensions. Discrete Contin. Dyn. Syst. 24 (2009), 933–978.
- [30] X.B. Pan and K.H. Kwek. Schrödinger operators with non-degenerately vanishing magnetic fields in bounded domains. Trans. Amer. Math. Soc. 354 (2002), 4201–4227.
- [31] K. Pankrashkin. An inequality for the maximum curvature through a geometric flow. Arch. Math. 105 (2015), 297–300.
- [32] N. Raymond. Sharp asymptotics for the Neumann Laplacian with variable magnetic field: case of dimension 2. Ann. Henri Poincaré 10 (2009), 95–122.
- [33] N. Raymond. On the semiclassical (3D)- Neumann Laplacian with variable magnetic field. Asymptotic Analysis 68 (2010), 1–40.

(B. Helffer) LABORATOIRE DE MATHÉMATIQUES JEAN LERAY, CNRS, NANTES UNIVERSITÉ, 44000 NANTES, FRANCE.

Email address: Bernard.Helffer@univ-nantes.fr

(A. Kachmar) School of Science and Engineering, The Chinese University of Hong Kong Shenzhen, Guangdong, 518172, P.R. China.

Email address: akachmar@cuhk.edu.cn

(F. Nicoleau) LABORATOIRE DE MATHÉMATIQUES JEAN LERAY, CNRS, NANTES UNIVERSITÉ, 44000 NANTES, FRANCE.

Email address: francois.nicoleau@univ-nantes.fr