arXiv:2501.01008v1 [eess.SP] 2 Jan 2025

CONFINED ORTHOGONAL MATCHING PURSUIT FOR SPARSE RANDOM COMBINATORIAL MATRICES*

XINWEI ZHAO[†], JINMING WEN[‡], HONGQI YANG[§], AND XIAO MA[¶]

Abstract. Orthogonal matching pursuit (OMP) is a commonly used greedy algorithm for recovering sparse signals from compressed measurements. In this paper, we introduce a variant of the OMP algorithm to reduce the complexity of reconstructing a class of K-sparse signals $\boldsymbol{x} \in \mathbb{R}^n$ from measurements $\boldsymbol{y} = \boldsymbol{A}\boldsymbol{x}$, where $\boldsymbol{A} \in \{0,1\}^{m \times n}$ is a sparse random combinatorial matrix with $d \ (d \leq m/2)$ ones per column. The proposed algorithm, referred to as the confined OMP algorithm, utilizes the properties of \boldsymbol{x} and \boldsymbol{A} to remove much of the redundancy in the dictionary (also referred to as \boldsymbol{A}) and thus fewer column indices of \boldsymbol{A} need to be identified. To this end, we first define a confined set Γ with $|\Gamma| \leq n$ and then prove that the support of \boldsymbol{x} is a subset of Γ with probability 1 if the distributions of non-zero components of \boldsymbol{x} satisfy a certain condition. During the process of the confined OMP algorithm, the possibly chosen column indices are strictly confined into the confined set Γ . We further develop lower bounds on the probability of exact recovery of \boldsymbol{x} using OMP algorithm and confined OMP algorithm with K iterations, respectively. The obtained theoretical results of confined OMP algorithm can be used to optimize the column degree d of \boldsymbol{A} . Finally, experimental results show that the confined OMP algorithm is more efficient in reconstructing a class of sparse signals compared to the OMP algorithm.

Key words. Compressed sensing, exact recovery probability, orthogonal matching pursuit, sparse signal recovery, sparse random combinatorial matrices

1. Introduction. Compressed sensing (CS) as a novel sampling theory [6, 12] has attracted much attention in recent twenty years. In CS, it is common to encounter the following linear model

$$(1.1) y = Ax,$$

where $A \in \mathbb{R}^{m \times n}$ is a measurement matrix with $m < n, x \in \mathbb{R}^n$ is an unknown *K*-sparse signal (i.e., x has at most *K* non-zero elements) and $y \in \mathbb{R}^m$ is a known measurement vector. The CS recovery algorithm aims to recover x from (1.1) by solving the following minimization problem

(1.2)
$$\min \|\boldsymbol{x}\|_0 \quad \text{s.t.} \; \boldsymbol{y} = \boldsymbol{A}\boldsymbol{x}$$

where $\|\boldsymbol{x}\|_0 \stackrel{\triangle}{=} |\{i : x_i \neq 0\}|$ and x_i is the *i*-th element of \boldsymbol{x} . Unfortunately, as shown in [18], the problem (1.2) is NP-hard in general. Two methods are commonly used for solving this problem. One focuses on the reconstruction of sparse signals by considering a convex relaxation of (1.2), such as solving the l_1 -minimization problem [5, 36, 7, 15]. There are also algorithms solving (1.2) directly, such as greedy algorithms [35, 30, 8, 29, 32, 25, 37, 24, 39] and thresholding algorithms [4, 17, 33, 11, 2].

Funding: This work was supported by the National Key R&D Program of China (No. 2021YFA1000500). (Corresponding author: Xiao Ma.)

[†]School of Computer Science and Engineering, Sun Yat-sen University, Guangzhou, 510006, China (zhaoxw9@mail2.sysu.edu.cn).

[‡]Department of Information Science and Technology, Jinan University, Guangzhou 510632, China (jinming.wen@mail.mcgill.ca).

[§]Guangdong Province Key Laboratory of Computational Science, School of Computer Science and Engineering, Sun Yat-sen University, Guangzhou 510006, China (mcsyhq@mail.sysu.edu.cn).

[¶]School of Computer Science and Engineering, Sun Yat-sen University, Guangzhou, 510006, China (maxiao@mail.sysu.edu.cn).

Algorithm 1.1 Orthogonal matching pursuit algorithm

Input: $\boldsymbol{y} \in \mathbb{R}^{m}, \boldsymbol{A} \in \mathbb{R}^{m \times n}$, and K. Initialize: $k = 0, \boldsymbol{r}^{(0)} = \boldsymbol{y}$, and $\Lambda^{(0)} = \emptyset$. while $\boldsymbol{\cdot} \boldsymbol{r}^{(k)} \neq 0$ and $k \leq K$ are met' do k = k + 1, $t^{(k)} = \arg \max |\boldsymbol{A}_{i}^{T} \boldsymbol{r}^{(k-1)}|$, (Identification) $\Lambda^{(k)} = \Lambda^{(k-1)} \cup \{t^{(k)}\}$, (Augmentation) $\hat{\boldsymbol{x}}_{\Lambda^{(k)}}^{(k)} = \arg \min_{\boldsymbol{x} \in \mathbb{R}^{k}} \|\boldsymbol{y} - \boldsymbol{A}_{\Lambda^{(k)}} \boldsymbol{x}\|_{2}$, (Estimation) $\boldsymbol{r}^{(k)} = \boldsymbol{y} - \boldsymbol{A}_{\Lambda^{(k)}} \hat{\boldsymbol{x}}_{\Lambda^{(k)}}^{(k)}$. (Residual update) end while Output: $\hat{\boldsymbol{x}}^{(k)}$.

Among greedy algorithms, the OMP algorithm [35] is one of the most commonly used algorithms. As a greedy algorithm, the OMP algorithm identifies the support (index set of non-zero elements) of the sparse signal \boldsymbol{x} in an iterative manner and thus iteratively performs local optimal updates. Specifically, the process of the OMP algorithm at each iteration can be summarized into four steps [37] (see Algorithm 1.1 for details):

- Identification: select the column of A maximally correlated with the residual $r^{(k-1)}$.
- Augmentation: add the index of the chosen column into the estimated support set $\Lambda^{(k)}$.
- Estimation: estimate the values of elements whose indices are in the estimated support set $\Lambda^{(k)}$ by solving a least squares problem.
- Residual update: eliminate the vestige of columns in $\Lambda^{(k)}$ from the measurement vector \boldsymbol{y} , resulting in a new residual used for the next iteration.

Among these, the computational complexity of the OMP algorithm is mainly dominated by the identification step and the estimation step. In order to enhance the computational efficiency and recovery performance of the OMP algorithm, there have been some studies on the modified OMP algorithm, mainly focusing on the identification step. For example, the generalized OMP algorithm [37] (a.k.a. orthogonal multi-matching pursuit algorithm [25]) allows multiple indices maximally correlated with the residual are chosen at each iteration so that fewer number of iterations are required. The existing methods seek to efficiently select the "true" column indices from the redundant dictionary (measurement matrix A). This raises a question: Is it possible to reduce the redundancy of the dictionary A?

Before answering this question, we introduce some useful tools to characterize the performance of the recovery algorithm. In [6], Candès and Tao introduced the concept of restricted isometry property (RIP) and showed that if A satisfies RIP with relatively small restricted isometry constant (RIC) δ_{2K} , any K-sparse signal can be exactly recovered by solving a l_1 -minimization problem. In particular, it has been proved in [41] that $\delta_{K+1} < 1/\sqrt{K+1}$ is sufficient for the OMP algorithm to recover any K-sparse signal from (1.1) in K iterations. The mutual coherence, denoted as μ_m , is also an important parameter for A, which indicates the maximum absolute correlation between normalized columns of A. It has been showed in [34] that any K-sparse signal can be exactly recovered by the OMP algorithm if $K < (\mu_m^{-1} + 1)/2$. In [35], the authors developed a lower bound on the probability that any K-sparse signal can be exactly recovered from (1.1) by using the OMP algorithm in K iterations, where the matrix \mathbf{A} in (1.1) is a random Gaussian matrix. In [40], this lower bound is further tightened with the aid of prior information of \mathbf{x} . Unfortunately, the lower bound techniques proposed in [35] and [40] are only suitable for the case of Gaussian matrices.

In CS, the construction of measurement matrices is also one of the main concerns. In general, the random measurement matrices can be classified into dense and sparse matrices. It has been verified that many dense matrices, such as Gaussian matrices and Fourier matrices, satisfy the RIP with overwhelming probability [1] and have provably good recovery performance. On the other hand, sparse random matrices also attract much attention [3, 22, 23, 10, 26, 28] since the sparsity can enable the computation of the matrix-product to be remarkably efficient and save the storage space in practice [19]. In particular, sparse binary-valued measurement matrices are commonly used in some applications, including group testing [20], DNA Microarrays [31], and single-pixel imaging [14]. Furthermore, many studies [10, 26, 27] showed that the sparse binary-valued measurement matrices are as "good" as the dense ones both in theory and in practice.

Assume that the measurement matrix $\mathbf{A} \in \{0, 1\}^{m \times n}$ is a sparse combinatorial matrix with independent columns, where each column is chosen uniformly among the vectors with d ($d \leq m/2$) ones. Now we answer the previous question. Yes, much redundancy of \mathbf{A} can be removed if \mathbf{x} is a signal defined in Definition 2.5. Actually, this observation can be traced back to the work of Khajehnejad et al. in [23]. They showed that the redundancy of a sparse matrix constructed by the expander theory can be eliminated if non-zero elements of \mathbf{x} are non-negative. Our work is more general and takes the observation in [23] as a special case. Specifically, the sparse signal we consider is not limited to be non-negative, but a class of signals defined in Definition 2.5, including the Gaussian sparse signal. Furthermore, the considered measurement matrix \mathbf{A} in this paper is more general than that constructed by the expander theory in [23]. The contributions of this paper are summarized as follows.

- We first define the confined set Γ with $|\Gamma| \leq n$ and prove that the support of \boldsymbol{x} defined in Definition 2.5 is a subset of Γ with probability 1. To theoretically clarify the effectiveness of removing the redundancy of \boldsymbol{A} , we present the expectations of the sparsity of \boldsymbol{y} and the size of Γ .
- We propose a variant of OMP algorithm, referred to as the confined OMP algorithm, by introducing the confined set Γ into the identification step. The possibly chosen column indices are strictly confined into the confined set Γ . We further analyze the complexities of OMP algorithm and confined OMP algorithm. The analysis results show that the identification efficiency of confined OMP algorithm is at least $\frac{nKd-K}{|\Gamma|Kd-K+n}$ times that of OMP algorithm. Furthermore, the experimental results show that the confined OMP algorithm achieves a large reduction in complexity if $K \ll m$.
- We develop a lower bound on the probability of exact recovery of any K-sparse signal x using OMP algorithm over a sparse random combinatorial matrix. As far as we know, our work is the first to develop the recovery lower bound of OMP algorithm over a sparse random combinatorial matrix. We further develop a lower bound on the probability of exact recovery of x defined in Definition 2.5 using confined OMP algorithm over a sparse random combinatorial matrix.

The paper is organized as follows. We define the confined set Γ and present the

proposed algorithm in section 2. In section 3, the expectations of the sparsity of \boldsymbol{y} and the size of Γ are investigated, and then the recovery performance bounds of OMP algorithm and confined OMP algorithm are provided. Experimental results are presented in section 4 and section 5 concludes the paper.

Notation: we use boldface lowercase letters to denote column vectors and boldface uppercase letters to denote matrices. The support of \boldsymbol{x} is denoted by Ω and the complement of Ω is $\Omega^c = [n] \setminus \Omega = \{i : i \in [n], i \notin \Omega\}$, where [n] represents the set $\{1, 2, \dots, n\}$. Let x_i and \boldsymbol{A}_j be the *i*-th element of \boldsymbol{x} and the *j*-th column of \boldsymbol{A} , respectively. We use $A_{i,j}$ to denote the element of \boldsymbol{A} located at the *i*-th row and the *j*-th column. We denote by \boldsymbol{x}_{Λ} the sub-vector of \boldsymbol{x} that contains the entries of \boldsymbol{x} indexed by the set Λ , and \boldsymbol{A}_{Λ} the sub-matrix of \boldsymbol{A} that contains the columns of \boldsymbol{A} indexed by the set Λ . We use \boldsymbol{A}^T to denote the transport of \boldsymbol{A} and $\mathbf{E}[X]$ to represent the expectation of X.

2. Proposed algorithm. This section will first introduce the inherent properties between random combinatorial matrices and sparse signals, and then present the details of the proposed algorithm.

2.1. The confined set. For a sufficiently small number $\epsilon > 0$, let $\mathcal{E} = \{i : |y_i| \le \epsilon\}$ where y_i is the *i*-th element of y. The definition of the confined set is as follows.

DEFINITION 2.1 (The confined set). The confined set Γ is specified by defining its complement Γ^c as $\Gamma^c = \bigcup_{i \in \mathcal{E}} \Gamma^c_i$ where $\Gamma^c_i = \{j : A_{i,j} = 1, |y_i| \leq \epsilon\}$. That is, $\Gamma = [n] \setminus \Gamma^c$.

The following Theorem gives the probability of $\{\Omega \subseteq \Gamma\}$.

THEOREM 2.2 (The probability of $\{\Omega \subseteq \Gamma\}$). Suppose that in (1.1), $A \in \{0,1\}^{m \times n}$ is a random combinatorial matrix with independent columns, where each column is chosen uniformly among the vectors with d ones. Furthermore, the K non-zero components of x are independent and identically distributed (i.i.d.), with the same cumulative distribution function (CDF) $F_X(x) = \mathbb{P}\{X \leq x\}$. Then, the probability of $\{\Omega \subseteq \Gamma\}$ is given by

(2.1)
$$\mathbb{P}\{\Omega \subseteq \Gamma\} = \left(1 - \sum_{\ell=1}^{K} \binom{K}{\ell} \left(F_X^{*\ell}(\epsilon) - F_X^{*\ell}(-\epsilon)\right)\right)^{|\mathcal{E}|}$$

where $F_X^{*\ell}(x) = (\underbrace{F_X * F_X * \cdots * F_X}_{\ell \text{ times}})(x)$ and the asterisk * denotes the convolution

operation.

Proof. See Appendix A.

With Theorem 2.2, we have the following two Corollaries.

COROLLARY 2.3. If the probability density function (PDF) $f_X(x)$ of non-zero components of x is a continuous function, then the probability $\mathbb{P}\{\Omega \subseteq \Gamma\} \to 1$ as $\epsilon \to 0$.

Proof. Obviously, $F_X^{*\ell}(x)$ is a continuous function provided that $f_X(x)$ is a continuous function. According to Theorem 2.2, we have $\lim_{\epsilon \to 0} \mathbb{P}\{\Omega \subseteq \Gamma\} = 1$.

COROLLARY 2.4. If the values of non-zero components of x share the same polarity, then the probability $\mathbb{P}\{\Omega \subseteq \Gamma\} \to 1 \text{ as } \epsilon \to 0.$

Algorithm 2.1 Confined orthogonal matching pursuit algorithm

Input: $\boldsymbol{y} \in \mathbb{R}^{m}$, $\boldsymbol{A} \in \{0, 1\}^{m \times n}$, ϵ , and K. Initialize: k = 0, $\boldsymbol{r}^{(0)} = \boldsymbol{y}$, and $\Lambda^{(0)} = \emptyset$. Preprocessing: $\mathcal{E} = \{i : |y_i| \le \epsilon\}$ and $\Gamma = [n] \setminus \bigcup_{i \in \mathcal{E}} \Gamma_i^c$; if $|\Gamma| = K$ then $\hat{\boldsymbol{x}}_{\Gamma} = \underset{\boldsymbol{x} \in \mathbb{R}^{|\Gamma|}}{\operatorname{arg\,min}} \|\boldsymbol{y} - \boldsymbol{A}_{\Gamma}\boldsymbol{x}\|_2$, return $\hat{\boldsymbol{x}}$. end if while ' $\boldsymbol{r}^{(k)} \neq 0$ and $k \le K$ are met' do k = k + 1, $t^{(k)} = \underset{i \in \Gamma}{\operatorname{arg\,max}} |\boldsymbol{A}_i^T \boldsymbol{r}^{(k-1)}|$, (Identification) $\hat{\boldsymbol{\lambda}}^{(k)}_{\Lambda^{(k)}} = \underset{\boldsymbol{x} \in \mathbb{R}^k}{\operatorname{arg\,min}} \|\boldsymbol{y} - \boldsymbol{A}_{\Lambda^{(k)}}\boldsymbol{x}\|_2$, (Estimation) $\hat{\boldsymbol{x}}^{(k)}_{\Lambda^{(k)}} = \underset{\boldsymbol{x} \in \mathbb{R}^k}{\operatorname{arg\,min}} \|\boldsymbol{y} - \boldsymbol{A}_{\Lambda^{(k)}}\boldsymbol{x}\|_2$, (Estimation) $\boldsymbol{r}^{(k)} = \boldsymbol{y} - \boldsymbol{A}_{\Lambda^{(k)}} \hat{\boldsymbol{x}}^{(k)}_{\Lambda^{(k)}}$. (Residual update) end while return $\hat{\boldsymbol{x}}^{(K)}$.

The Corollaries indicate that by setting $\epsilon \to 0$, both Gaussian sparse signals and nonnegative sparse signals can ensure $\mathbb{P}\{\Omega \subseteq \Gamma\} = 1$. We call these signals as *confined signals* in this paper. The definition of the confined signal is given as follows.

DEFINITION 2.5 (The confined signal). The confined signal is a K-sparse signal whose non-zero components are i.i.d. and, for $\ell = 1, 2, \dots, K$, their CDFs satisfy $F_X^{*\ell}(\epsilon) - F_X^{*\ell}(-\epsilon) \to 0$ as $\epsilon \to 0$.

2.2. Confined OMP algorithm. The confined OMP is a modified OMP algorithm by introducing a confined set Γ . The key feature of the confined OMP algorithm is to introduce a confined set Γ into the identification step such that the possibly chosen column indices are strictly confined into the confined set Γ . In the following, we assume that the signal x is defined in Definition 2.5 and has exactly K non-zero elements. The details of the confined OMP algorithm are summarized in Algorithm 2.1. The necessary explanations of the confined OMP algorithm are as follows.

- (**Preprocessing**) For a sufficiently small number ϵ , obtain the set $\mathcal{E} = \{i : |y_i| \leq \epsilon\}$ first. Then, obtain the confined set $\Gamma = [n] \setminus \bigcup_{i \in \mathcal{E}} \Gamma_i^c$, where $\Gamma_i^c = \{j : A_{i,j} = 1, |y_i| \leq \epsilon\}$. With $\Omega \subseteq \Gamma$, we have $\Omega = \Gamma$ if they have the same size. Thus, the identification is already done without the subsequent iterations.
- (Identification) If $|\Gamma| > K$, the iterative processing is executed. In each iteration, correlations between columns whose indices are in Γ and the residual are compared. The column index corresponding to the maximal correlation is chosen as the new element of the estimated support set $\Lambda^{(k)}$.

We discuss the complexity of OMP algorithm and confined OMP algorithm as follows. It is known that the matrix-vector product can be divided into two steps: multiplication and addition. For a binary-valued matrix, the matrix-vector product only involves addition, no multiplication is required. If non-zero elements of \boldsymbol{x} are real numbers, the addition requires at most n(d-1) float point operations (flops). Furthermore, the selection of the maximum inner-product value in the identification requires n-1 flops. Thus, the identification of OMP algorithm requires at most TABLE 1

 $Comparison \ of \ the \ complexity \ of \ the \ OMP \ algorithm \ and \ the \ confined \ OMP \ algorithm \ in \ the \ identification \ step.$

Algorithm	OMP	confined OMP	
Condition	null	$ \Gamma = K$	$ \Gamma > K$
Flops	Knd - K	n	$K \Gamma d - K + n$

Kn(d-1) + K(n-1) flops.

Thanks to $|\Gamma| \leq n$, the confined OMP algorithm requires fewer inner products in the identification step compared to the conventional OMP algorithm. Specifically, no inner product is required if $|\Gamma| = K$. Otherwise, the identification requires at most $K|\Gamma|(d-1)+K(|\Gamma|-1)$ flops, where $K|\Gamma|(d-1)$ flops are required for the matrix-vector product operations and $K(|\Gamma|-1)$ flops for the selections of the maximum innerproduct values. Furthermore, the preprocessing step involves additional operations. Specifically, there are extra n flops to obtain \mathcal{E} and the complexity of obtaining Γ is negligible. Consider identification and preprocessing together, these steps requires nflops if $|\Gamma| = K$. Otherwise, at most $K|\Gamma|(d-1) + K(|\Gamma|-1) + n$ flops are required. Table 1 summarizes the complexity of OMP algorithm and confined OMP algorithm in the identification step. The improvement of the proposed algorithm over OMP algorithm is around Kd times in terms of identification efficiency if $|\Gamma| = K$, and $\frac{Knd-K}{K|\Gamma|d-K+n}$ times otherwise.

It can be foreseen that the confined OMP algorithm achieves a large reduction in complexity if $|\Gamma| \ll n$. Especially in case where $|\Gamma| = K$, the confined OMP algorithm eliminates the need for the identification. Even if $|\Gamma| = n$, the computational complexity of OMP algorithm and confined OMP algorithm are comparable, since the extra complexity introduced by the preprocessing step is negligible.

3. Analysis of the proposed algorithm. In this section, we first investigate the expectations of the sparsity of y and the size of the confined set Γ . Then, we study the lower bounds on the exact recovery probability of OMP algorithm and confined OMP algorithm in the case of using sparse random combinatorial matrices. Note that the lower bound on the exact recovery probability of OMP algorithm is valid for any K-sparse signal, whereas that of confined OMP algorithm is only valid for the signal defined in Definition 2.5.

3.1. The sparsity of \boldsymbol{y} . Studying the sparsity of \boldsymbol{y} helps to derive the subsequent Lemmas and Theorems. For $k \in \{1, 2, \dots, K\}$, let $\nu^{(k)} = n - |\mathcal{E}|$ be the number of "non-zero" elements of $\boldsymbol{y} = \boldsymbol{A}\boldsymbol{x}$, where the signal \boldsymbol{x} is defined in Definition 2.5 and has k non-zero elements. Obviously, the value of $\nu^{(k)}$ ranges from d to kd. The following Lemma gives the probability of $\nu^{(k)}$ for $k = 1, 2, \dots, K$.

LEMMA 3.1. For any integer K, it holds that $\mathbb{P}\{\nu^{(1)} = d\} = 1$ and

(3.1)
$$\mathbb{P}\{\nu^{(k)} = v\} = \sum_{z=\max\{v-d,d\}}^{\min\{v,(k-1)d\}} \frac{\binom{z}{(v-z)}\binom{m-z}{d-v+z}}{\binom{m}{d}} \mathbb{P}\{\nu^{(k-1)} = z\}$$

for $k = 2, 3, \dots, K$ and $v = d, d + 1, \dots, kd$, where $\mathbb{P}\{\nu^{(k-1)} = z\}$ can be calculated by (3.1) recursively.

Proof. See Appendix B.



FIG. 1. Empirical means and expectations of $\nu^{(K)}$ for different column degree d.

Once the probability $\mathbb{P}\{\nu^{(K)} = v\}$ is obtained, one can easily obtain the expectation of $\nu^{(K)}$.

THEOREM 3.2 (The expectation of $\nu^{(K)}$). Suppose that in (1.1), $\mathbf{A} \in \{0, 1\}^{m \times n}$ is a random combinatorial matrix with d ones per column and the signal \mathbf{x} is defined in Definition 2.5. Then,

(3.2)
$$\mathbf{E}[\nu^{(K)}] = \sum_{v=d}^{Kd} v \cdot \mathbb{P}\{\nu^{(K)} = v\},$$

where $\mathbb{P}\{\nu^{(K)} = v\}$ is given in Lemma 3.1.

Figure 1 shows empirical means and expectations of $\nu^{(K)}$ for different column degree d, where $\mathbf{A} \in \{0, 1\}^{100 \times 256}$ and \mathbf{x} is a Gaussian sparse signal with exactly K non-zero elements. It can be seen that the empirical results match well with their expectations.

3.2. The size of Γ . To clarify the effectiveness of removing the redundancy of \boldsymbol{A} , it is necessary to consider the size of Γ theoretically. Furthermore, the size of Γ is crucial to the complexity and recovery performance analysis of the proposed algorithm. The following Theorem will give $\mathbf{E}[|\Gamma|]$.

THEOREM 3.3 (The expectation of $|\Gamma|$). Suppose that in (1.1), $A \in \{0,1\}^{m \times n}$ is a random combinatorial matrix with d ones per column and the signal x is defined in Definition 2.5. Then,

(3.3)
$$\mathbf{E}[|\Gamma|] = K + (n-K) \cdot \sum_{\nu=d}^{Kd} \frac{\binom{\nu}{d}}{\binom{m}{d}} \mathbb{P}\{\nu^{(K)} = \nu\},$$

where $\mathbb{P}\{\nu^{(K)} = v\}$ is given in Lemma 3.1.

Proof. See Appendix C.

Figure 2 shows empirical means and expectations of $|\Gamma|$ for different column degree d, where $A \in \{0, 1\}^{100 \times 256}$ and x is a Gaussian sparse signal with exactly K non-zero elements. It can be seen that the empirical results match well with their expectations. We also observe that $\mathbf{E}[|\Gamma|] \approx K$ when K is relatively small. These observations confirm the effectiveness of removing the redundancy of A.



FIG. 2. Empirical means and expectations of $|\Gamma|$ for different column degree d.

3.3. Recovery performance analysis of OMP. In this subsection, we provide a lower bound on the probability that the OMP algorithm exactly recovers any K-sparse signal \boldsymbol{x} in K iterations for $\boldsymbol{A} \in \{0, 1\}^{m \times n}$.

We first define the coherence between two columns of \boldsymbol{A} as

(3.4)
$$\mu_{i,j} = \frac{|\mathbf{A}_i^T \mathbf{A}_j|}{\|\mathbf{A}_i\|_2 \|\mathbf{A}_j\|_2}$$

where $i, j \in [n]$ and $i \neq j$. Different from the random Gaussian matrices [40], the values of $\mu_{i,j}$ are discrete for the binary-valued matrix **A**. That is, we have $\mu_{i,j} \in \{0, 1/d, 2/d, \dots, 1\}$. We further use g(s) for $s \in \{0, 1/d, 2/d, \dots, 1\}$ to denote the distribution function of $\mu_{i,j}$:

(3.5)
$$g(s) = \mathbb{P}\left\{\mu_{i,j} = s\right\} = \frac{\binom{d}{sd}\binom{m-d}{d-sd}}{\binom{m}{d}},$$

where $\{\mu_{i,j} = s\}$ means the event that the column A_i shares exactly sd non-zero positions with the column A_j .

We introduce several distribution functions used in Theorem 3.4. Let

$$\mathcal{P} = \left\{ \sum_{i=1}^{K-1} s_i : s_i \in \{0, 1/d, 2/d, \cdots, 1\} \text{ for } i = 1, 2, \cdots, K-1 \right\}$$

and

$$\mathcal{Q} = \left\{ \sum_{i=1}^{K} s_i^2 : s_i \in \{0, 1/d, 2/d, \cdots, 1\} \text{ for } i = 1, 2, \cdots, K \right\}.$$

Due to the randomness of s_i , we have a distribution function h(z) over \mathcal{Q} , given by

$$h(z) = \sum_{s_1^2 + \dots + s_K^2 = z} g(s_1)g(s_2) \cdots g(s_K).$$

Similarly, we have a distribution function $\phi(z)$ over \mathcal{P} , given by

$$\phi(z) = \sum_{s_1 + \dots + s_{K-1} = z} g(s_1)g(s_2) \cdots g(s_{K-1}).$$

Then, the CDF associated with $\phi(z)$ is given by

$$\Phi(z) = \sum_{z' \le z} \phi(z').$$

Let $r_i = \sum_{j=1}^{K-1} s_j$ for $i = 1, 2, \dots, K$ and r_m be the maximum variable of r_i . Then, by the order statistics [9], the distribution function of r_m is given by

$$\phi_m(z) = (\Phi(z))^K - (\Phi(z) - \phi(z))^K,$$

where $z \in \mathcal{P}$.

THEOREM 3.4 (The lower bound on the exact recovery probability of OMP). Suppose that in (1.1), **A** is a random combinatorial matrix with independent columns, where the degree of each column is d satisfying $(1+\varepsilon) \log m \le d \le m/2$ for a constant $\varepsilon > 0$. The signal **x** is any K-sparse signal. Define $\mathbb{S}_{omp} = \{OMP \text{ exactly recovers } \mathbf{x} \text{ with } K \text{ iterations} \}$. Then, for a sufficiently large m, it holds that

(3.6)
$$\mathbb{P}\{\mathbb{S}_{omp}\} \ge \left(\sum_{p \in \mathcal{P}} \sum_{q \in \mathcal{Q}} \mathbf{1}(p, q) \phi_m(p) h(q)\right)^K - o(1),$$

where the indicator function is

$$\mathbf{1}(p,q) = \begin{cases} 1 & \text{if } \frac{\sqrt{\max\{0,1-p\}}}{\sqrt{K}} > \sqrt{q}, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. See Appendix D.

3.4. Recovery performance analysis of confined OMP. In this subsection, we will present a lower bound on the probability that the confined OMP algorithm exactly recovers the sparse signal \boldsymbol{x} for $\boldsymbol{A} \in \{0,1\}^{m \times n}$, where \boldsymbol{x} is defined in Definition 2.5.

THEOREM 3.5 (The lower bound on the exact recovery probability of confined OMP). Suppose that in (1.1), \mathbf{A} is a random combinatorial matrix with independent columns, where the degree of each column is d satisfying $(1 + \varepsilon) \log m \le d \le m/2$ for a constant $\varepsilon > 0$. The sparse signal \mathbf{x} is defined in Definition 2.5 and has exactly K non-zero elements. Define $\mathbb{S}_{\text{comp}} = \{\text{confined OMP exactly recovers } \mathbf{x} \text{ with } K \text{ iterations} \}$. Then, for a sufficiently large m, it holds that

(3.7)
$$\mathbb{P}\{\mathbb{S}_{\text{comp}}\} \ge \left(1 - \sum_{\nu=d}^{Kd} \frac{\binom{\nu}{d}}{\binom{m}{d}} \mathbb{P}\{\nu^{(K)} = \nu\}\right)^{n-K} - o(1),$$

where $\mathbb{P}\{\nu^{(k)} = v\}$ is given in Lemma 3.1.

Proof. See Appendix E.

4. Experimental results. This section presents the experimental results that demonstrate the advantage of the confined OMP algorithm in terms of recovery performance and complexity. Furthermore, we use the lower bound on the probability $\mathbb{P}\{\mathbb{S}_{comp}\}$ to optimize the column degree d. The experimental results were obtained



FIG. 3. Recovery performance for K-sparse (a) Gaussian signals and (b) flat signals versus the sparsity K.



FIG. 4. (a) Average number of inner-product operations in the identification step and (b) average CPU times of different greedy algorithms for Gaussian signals with different K.

by MATLAB R2023a on a desktop computer with Intel(R) Core(TM) i7-11700 CPU @ 2.50 GHz. In the following simulations, we generate a K-sparse signal whose support is chosen at random. In addition, we consider two types of K-sparse signals: (a) Gaussian signals and (b) flat signals. The non-zero elements of the Gaussian signal are independently and randomly drawn from a standard Gaussian distribution. Furthermore, the support of the flat signal is randomly chosen and the nonzero elements are set to 1.

4.1. Recovery performances for different sparsity K. We first show the efficiency and recovery performance of the proposed confined OMP algorithm for different sparsity K. In Figure 3 and Figure 4, each column of the measurement matrix $A \in \{0,1\}^{100 \times 256}$ is chosen independently and uniformly among the vectors with 10 ones.

In Figure 3, we can see that the confined OMP algorithm achieves performance gains for both Gaussian and flat signals. In particular, the performance gain for the flat signal is appreciable. It is reasonable because when the sparsity K is relative small, the size of Γ is not as large, which enables the confined OMP algorithm to screen out more interference.



FIG. 5. Recovery performance for 4-sparse (a) Gaussian signals and (b) flat signals versus measurement m.



FIG. 6. (a) Average number of inner-product operations in the identification step and (b) average CPU times of different greedy algorithms for Gaussian signals with different measurements m.

Figure 4 shows the average inner-product operations in the identification step and average CPU times of different greedy algorithms for Gaussian signals with different K. From Figure 4(a), we observe that the number of inner-product operations of OMP algorithm increases linearly with the increase of K. The reason is that OMP algorithm requires Kn inner-product operations in the identification step to recover a K-sparse signal. For the confined OMP algorithm, however, the identification is eliminated if $|\Gamma| = K$. Otherwise, $K|\Gamma|$ inner-product operations are required to recover a K-sparse signal. As shown in Figure 2, the expectation of $|\Gamma|$ is much smaller than n for a relatively small K. Thus, in Figure 4(a), the confined OMP algorithm achieves a significant reduction in number of inner-product operations for a relatively small K. The average CPU time shown in Figure 4(b) relates to the number of inner-product operations. Their trends of curves shown in Figure 4(a) and (b) are almost the same. In particular, the confined OMP algorithm achieves a reduction of about 85% in average CPU time to recover a 6-sparse Gaussian signal when compared to that of OMP algorithm. Furthermore, we also present the average CPU time of the batch OMP algorithm proposed in [32] for comparison. It can be seen that the

complexity reduction of the confined OMP algorithm is much greater than that of the batch OMP algorithm for a relatively small K.

Remark: The batch OMP algorithm is an efficient implementation of the OMP algorithm. Their recovery performances are the same. Thus, we omit the performance comparison between the batch OMP algorithm and the proposed algorithm. More details about the batch OMP algorithm can be found in [32].

4.2. Recovery performances for different measurements m. We further show the efficiency and recovery performance of the confined OMP algorithm for different measurements m. In Figure 5 and Figure 6, each column of the measurement matrix $\mathbf{A} \in \{0, 1\}^{m \times 256}$ is chosen independently and uniformly among the vectors with 10 ones. The sparsity K is set to 4.

In Figure 5, the recovery performances of confined OMP algorithm are better than those of OMP algorithm, but performance gains become smaller as m increases. However, we observe in Figure 6 that the number of inner-product operations for the confined OMP algorithm tends to 0 as m increases, resulting in a reduction of about 86% in average CPU time. These observations indicate that the complexity of the confined OMP algorithm is mainly contributed by the complexity of solving least-squares problems when $m \geq 100$.

4.3. Lower bounds. The lower bounds on the probability of \mathbb{S}_{omp} and \mathbb{S}_{comp} can be obtained by (3.6) and (3.7), respectively. In experimental tests, the column degree d of each chosen \boldsymbol{A} satisfies $\log m < d \leq m/2$ for a sufficiently large m. In this case, the probability $\mathbb{P}\left\{\det\left(\boldsymbol{A}_{\Lambda^{(k)}}^T\boldsymbol{A}_{\Lambda^{(k)}}\right)\neq 0\right\}$ is approximately equal to 1. Thus, the effect of the second term for both (3.6) and (3.7) is negligible.

The lower bounds on $\mathbb{P}{S_{omp}}$ and $\mathbb{P}{S_{comp}}$ are shown in Figure 3 and Figure 5. It can be seen that the lower bound on $\mathbb{P}{S_{comp}}$ is much tighter than that on $\mathbb{P}{S_{omp}}$. However, for both OMP algorithm and confined OMP algorithm, there exists a non-negligible gap between theory and practice. The reasons are as follows.

- We first consider the lower bound on the probability $\mathbb{P}\{\mathbb{S}_{omp}\}\)$. The scalings of $\|A_{\Omega}^{T}A_{\Omega}x'_{\Lambda^{(k)}}\|_{\infty}$ and $\|A_{\Omega^{c}}^{T}A_{\Omega}x'_{\Lambda^{(k)}}\|_{\infty}$ may lead to the gap between the two sides of (D.4). Furthermore, the bound on λ_{min} derived in Corollary D.4 is loose for a large sparsity K, which causes that the bound on $\mathbb{P}\{\mathbb{S}_{omp}\}$ also being loose.
- We further consider the lower bound on the probability $\mathbb{P}\{S_{\text{comp}}\}$. This bound given in (3.7) is actually the exact recovery probability conditioned on the event $\{|\Gamma| = K\}$. Thus, the gap between the analytical curve and experimental curve is equal to the exact recovery probability conditioned on the event $\{|\Gamma| > K\}$. This conditional probability derived in Appendix F is loose for the same reasons as in the case of bound for the OMP algorithm.

Remark: In [35] and [40], lower bounds on the exact recovery probability of OMP algorithm over Gaussian matrices also are loose. However, these theoretical results may provide a guide to determine whether the greedy algorithms are appropriate for reconstruction of sparse signals [35]. Otherwise, another sparse recovery algorithm is considered instead.

4.4. Optimization of column degree d. As shown in (3.7), the lower bound on $\mathbb{P}\{\mathbb{S}_{\text{comp}}\}$ is related to the column degree d of A. We expect to optimize the recovery performance of confined OMP algorithm by optimizing the lower bound on $\mathbb{P}\{\mathbb{S}_{\text{comp}}\}$. In other words, the optimized target is to increase the probability $\mathbb{P}\{|\Gamma| = K\}$. In the following experimental simulations, the column degree d of a measurement matrix



FIG. 7. Empirical and theoretical recovery performances of confined OMP as a function of column degree d.

 $A \in \{0, 1\}^{100 \times 256}$ satisfies $d > \log m$ such that the probability that the least squares have a unique solution is approximately equal to 1. The experimental and theoretical simulations are based on flat signals with K = 10 and K = 5, respectively. It can be seen from Figure 7 that the trend of theoretical curve is almost the same with that of experimental curve. Both of them achieve the best recovery performance when d = 12.

5. Conclusion and future research problem. This paper proposed a variant of OMP algorithm, referred to as confined OMP algorithm, to recover a class of sparse signals. We proved that the support of \boldsymbol{x} is contained in the confined set Γ if the signal \boldsymbol{x} is defined in Definition 2.5. We further presented the expectation of $|\Gamma|$ to show that much redundancy of \boldsymbol{A} can be removed, resulting in a improvement of the proposed algorithm in terms of identification efficiency. We also developed lower bounds on the probability $\mathbb{P}\{\mathbb{S}_{omp}\}$ and $\mathbb{P}\{\mathbb{S}_{comp}\}$ over sparse random combinatorial matrices. Finally, experimental results showed that confined OMP algorithm achieves a recovery performance gain and a reduction in complexity compared with OMP algorithm.

There are several future research problems arising from the confined OMP algorithm:

• As shown in [13], $K < spark(\mathbf{A})/2$ is a sufficient and necessary condition for the perfect recovery of a K-sparse signal with (1.2), where $spark(\mathbf{A})$ is the spark of \mathbf{A} and defined as $spark(\mathbf{A}) := \min\{\|\mathbf{x}\|_0 : \mathbf{A}\mathbf{x} = \mathbf{0}, \mathbf{x} \neq \mathbf{0}\}$. It is advisable to directly perform $\hat{\mathbf{x}}_{\Gamma} = \arg\min_{\mathbf{x}\in\mathbb{R}^{|\Gamma|}} \|\mathbf{y} - \mathbf{A}_{\Gamma}\mathbf{x}\|_2$ if $|\Gamma| < spark(\mathbf{A})$. In

this case, only one least squares needs to be performed. However, the main challenge is to determine the spark of A. Fortunately, Theorem D.1 implies that, for a sufficiently large m, the spark of a sparse random combinatorial matrix is equal to m+1 with probability 1-o(1). Our future work will focus on addressing this problem.

• In many applications, the sparsity K is unknown in practice. Fortunately, as shown in (3.2), the expectation of $\nu^{(K)}$ is related to the sparsity K if the signal is defined in Definition 2.5. The sparsity K is predictable according to the sparsity of y. It may be interesting to design a blind recovery algorithm that can achieve the same recovery performance as a non-blind recovery algorithm with a moderate increase in complexity.

XINWEI ZHAO, JINMING WEN, HONGQI YANG, AND XIAO MA

• As shown in Corollary D.4, the minimum eigenvalue λ_{min} of $\mathbf{A}_{\Omega}^{T} \mathbf{A}_{\Omega}$ ranges from $1 - r_m$ to $1 + r_m$, where $r_m \in \{0, 1/d, \cdots, (K-1)d\}$. Since $\mathbf{A}_{\Omega}^{T} \mathbf{A}_{\Omega}$ is considered as a positive definite matrix, we have $\lambda_{min} > 0$. As K increases, the lower bound on λ_{min} becomes less tight, resulting in a loose lower bound on $\mathbb{P}\{S_{omp}\}$. It may be challenging to give a tight bound on λ_{min} or σ_{min} in the future.

Appendix A. The proof of Theorem 2.2. Assume that the sparse signal x has exactly K non-zero elements. The following Lemma is useful for the proof of Theorem 2.2.

LEMMA A.1. Suppose that the K non-zero components of \mathbf{x} in (1.1), denoted without loss of generality by X_1, X_2, \dots, X_K , are i.i.d., with the same CDF $F_X(x)$ and PDF $f_X(x)$. Given $|y_i| \leq \epsilon$, for $\ell = 1, 2, \dots, K$ and $i \in \mathcal{E}$, the probability $\mathbb{P}\{|\Gamma_i^c \cap \Omega| = \ell\}$ is given by

(A.1)
$$\mathbb{P}\{|\Gamma_i^c \cap \Omega| = \ell\} = \binom{K}{\ell} \left(F_X^{*\ell}(\epsilon) - F_X^{*\ell}(-\epsilon)\right).$$

Proof. The event $\{|\Gamma_i^c \cap \Omega| = \ell\}$ is equivalent to the event that the sum of ℓ (out of K) non-zero components is less than or equal to ϵ . Without loss of generality, we assume that $Y = X_1 + X_2 + \cdots + X_{\ell}$. Consider the simple case that $\ell = 2$, the CDF of $Y = X_1 + X_2$ is calculated as

$$F_Y(y) = \mathbb{P}\{Y \le y\} = \mathbb{P}\{X_1 + X_2 \le y\}$$
$$= (F_X * F_X)(y) = \int_{-\infty}^{+\infty} f_X(\tau) F_X(y - \tau) d\tau$$

Similarly, extending Y to the sum of multiple components, its CDF $F_Y(y)$ is given by

$$F_Y(y) = \underbrace{(F_X * F_X * \dots * F_X)(y)}_{\ell \text{ times}}(y) = F_X^{*\ell}(y).$$

Once the CDF $F_Y(y)$ is obtained, one can calculate the probability of the event $\{|\Gamma_i^c \cap \Omega| = \ell\}$:

$$\mathbb{P}\{|\Gamma_i^c \cap \Omega| = \ell\} = \binom{K}{\ell} \left(F_Y(\epsilon) - F_Y(-\epsilon)\right) = \binom{K}{\ell} \left(F_X^{*\ell}(\epsilon) - F_X^{*\ell}(-\epsilon)\right).$$

In the following, we present the proof of Theorem 2.2.

Proof. The support of x is a subset of Γ if and only if the support of x is disjoint from the set Γ^c . In other words, the event $\{\Omega \subseteq \Gamma\}$ is equivalent to the event $\{|\Gamma^c \cap \Omega| = 0\}$. Further, the event $\{|\Gamma^c \cap \Omega| = 0\}$ is equivalent to the event that $\{|\Gamma_i^c \cap \Omega| = 0\}$ for all $i \in \mathcal{E}$. Thus, we have

$$\begin{split} \mathbb{P}\{\Omega \subseteq \Gamma\} &= \mathbb{P}\left\{\left| \bigcup_{i \in \mathcal{E}} \Gamma_i^c \cap \Omega \right| = 0 \right\} \\ &= \mathbb{P}\left\{ \left| \bigcup_{i \in \mathcal{E}} \Gamma_i^c \cap \Omega \right| = 0 \right\} \\ &= \prod_{i \in \mathcal{E}} \mathbb{P}\{\left|\Gamma_i^c \cap \Omega\right| = 0\} \\ &= \prod_{i \in \mathcal{E}} \left(1 - \mathbb{P}\{\left|\Gamma_i^c \cap \Omega\right| \neq 0\}\right) \\ &= \left(1 - \sum_{\ell=1}^K \binom{K}{\ell} \left(F_X^{*\ell}(\epsilon) - F_X^{*\ell}(-\epsilon)\right)\right)^{|\mathcal{E}|}, \end{split}$$
(A.2a)

where (A.2a) follows from Lemma A.1.

Appendix B. The proof of Lemma 3.1.

Proof. It is known that \boldsymbol{y} is a linear combination of K columns of \boldsymbol{A} . For K = 1, the value of $\nu^{(1)}$ is always equal to d since the degree of each column of \boldsymbol{A} is d. As a result, we have

$$\mathbb{P}\{\nu^{(1)} = d\} = 1.$$

For $k = 2, 3, \dots, K$, the value of $\nu^{(k)}$ ranges from d to kd. Here, the value of $\nu^{(k)}$ is equivalent to the number of non-zero elements for a vector resulting from the element-wise OR operation applied to k columns of A. It is hard to calculate the probability of $\nu^{(k)}$ directly since it is related to k columns of A. Fortunately, it can be modeled as a Markov model. Specifically, by the law of total probability, the probability $\mathbb{P}(\nu^{(k)})$ is

(B.1)
$$\mathbb{P}(\nu^{(k)}) = \sum_{\nu^{(k-1)}} \mathbb{P}(\nu^{(k)} | \nu^{(k-1)}) \mathbb{P}(\nu^{(k-1)}).$$

The transition probability $\mathbb{P}(\nu^{(k)}|\nu^{(k-1)})$ only depends on $\nu^{(k-1)}$ and $\nu^{(k)}$. This is equivalent to a linear combination of a vector with $\nu^{(k-1)}$ non-zero elements and any column of \mathbf{A} . If $\nu^{(k-1)} < \max\{\nu^{(k)} - d, d\}$ and $\nu^{(k-1)} > \min\{\nu^{(k)}, (k-1)d\}$, the transition probability $\mathbb{P}(\nu^{(k)}|\nu^{(k-1)})$ is obviously 0. Otherwise, the transition probability is

(B.2)
$$\mathbb{P}(\nu^{(k)}|\nu^{(k-1)}) = \frac{\binom{\nu^{(k-1)}}{\nu^{(k)}-\nu^{(k-1)}}\binom{m-\nu^{(k-1)}}{d-\nu^{(k)}+\nu^{(k-1)}}}{\binom{m}{d}}$$

where $\nu^{(k)} \in \{d, d+1, \cdots, kd\}$ and $\nu^{(k-1)} \in \{\max\{\nu^{(k)} - d, d\}, \max\{\nu^{(k)} - d, d\} + 1, \cdots, \min\{\nu^{(k)}, (k-1)d\}\}$. Here, (B.2) represents the probability that any column of \boldsymbol{A} shares exactly $\nu^{(k)} - \nu^{(k-1)}$ non-zero positions with a vector having $\nu^{(k-1)}$ non-zero elements.

The probability $\mathbb{P}(\nu^{(k-1)})$ can be calculated by (B.1) recursively. As a consequence, we get (3.1).

Appendix C. The proof of Theorem 3.3.

Proof. As shown in Theorem 2.2, the support of \boldsymbol{x} is a subset of the confined set Γ with probability 1. Thus, there are at least K out of n columns whose indices are in Γ . That is, we have $\mathbf{E}[|\Gamma|] \geq K$.

Assume that $|\Gamma| > K$ and there exists a column A_j for $j \in \Gamma \setminus \Omega$. Given $\nu^{(K)} = v$, the probability of the event $\{j \in \Gamma \setminus \Omega | \nu^{(K)} = v\}$ is given by

(C.1)
$$\mathbb{P}\left\{j\in\Gamma\backslash\Omega\middle|\nu^{(K)}=\upsilon\right\}=\frac{\binom{\upsilon}{d}}{\binom{m}{d}}.$$

Here, (C.1) indicates the probability that the column A_j shares exactly d non-zero positions with y having v non-zero elements. If this is not the case, then $j \notin \Gamma$ since the *i*-th element of A_j must be 0 for $i \in \mathcal{E}$.

Following from Lemma 3.1, the value of v ranges from d to Kd. Thus, by the law of total probability, we have

(C.2)
$$\mathbb{P}\left\{j\in\Gamma\backslash\Omega\right\} = \sum_{v=d}^{Kd} \mathbb{P}\left\{j\in\Gamma\backslash\Omega\middle|\nu^{(K)}=v\right\} \mathbb{P}\left\{\nu^{(K)}=v\right\}$$
$$= \sum_{v=d}^{Kd} \frac{\binom{v}{d}}{\binom{m}{d}} \mathbb{P}\left\{\nu^{(K)}=v\right\},$$

where $\mathbb{P}\left\{\nu^{(K)}=\upsilon\right\}$ is given in Lemma 3.1.

In summary, there are K columns whose indices are in $\Omega \subseteq \Gamma$. For the remaining n - K column indices, each of them belongs to $\Gamma \setminus \Omega$ with probability given in (C.2). As a result, we get $\mathbf{E}[|\Gamma|]$ shown in (3.3).

Appendix D. The proof of Theorem 3.4. There are two Corollaries used in the proof of Theorem 3.4. These two Corollaries are obtained from two different Theorems, which will be proved below respectively.

THEOREM D.1 (Theorem 1.2 in [16]). Fix $\varepsilon > 0$, and let d = d(m) be any function of m satisfying $\min\{d, m - d\} \ge (1 + \varepsilon) \log m$. Then, for an $m \times m$ random combinatorial matrix \mathbf{Q} with independent rows, where each row is chosen uniformly among the vectors with d ones, we have

$$\lim_{m \to \infty} \mathbb{P}\{\boldsymbol{Q} \text{ is singular}\} \to 0.$$

COROLLARY D.2. Fix $\varepsilon > 0$, and let d = d(m) be any function of m satisfying $(1+\varepsilon)\log m \leq d \leq m/2$. Then, for an $m \times K$ random combinatorial matrix A_{Ω} with d ones per column and a sufficiently large m, we have

(D.1)
$$\mathbb{P}\left\{\det\left(\boldsymbol{A}_{\Omega}^{T}\boldsymbol{A}_{\Omega}\right)\neq0\right\}=1-o(1),$$

where $det(\cdot)$ represents the determinant of argument.

Proof. According to Theorem D.1, we know that the $m \times m$ random combinatorial matrix \boldsymbol{Q} is nonsingular with probability 1 - o(1) for a sufficiently large m. In other words, m columns of \boldsymbol{Q}^T are linearly independent with probability 1 - o(1) for a sufficiently large m. For the $m \times K$ random combinatorial matrix \boldsymbol{A}_{Ω} with a sufficiently large m, K columns of \boldsymbol{A}_{Ω} are obviously linearly independent with probability 1 - o(1) for 1 - o(1). Furthermore, it is known that the rank of $\boldsymbol{A}_{\Omega}^T \boldsymbol{A}_{\Omega}$ is equal to that of \boldsymbol{A}_{Ω} . Thus, we get (D.1).

Remark: In general, the sparsity K is much smaller than m. Thus, as the increase of m, the probability $\mathbb{P}\{\det\left(\boldsymbol{A}_{\Omega}^{T}\boldsymbol{A}_{\Omega}\right)\neq0\}$ converges to 1 more rapidly than the probability $\mathbb{P}\{\det(\boldsymbol{Q})\neq0\}$ shown in Theorem D.1.

Let M be an $m \times m$ matrix and $r_i = \sum_{j=1, j \neq i}^m |M_{i,j}|$ be the sum of the absolute values of the non-diagonal elements in the *i*-th row of M for $i = 1, 2, \dots, m$. Define a circle centered at $M_{i,i}$ with a radius of r_i as a Gershgorin disc $D(M_{i,i}, r_i)$.

THEOREM D.3 (Gershgorin's circle Theorem [21]). Every eigenvalues of M lies in one of Gershgorin discs $D(M_{i,i}, r_i)$.

COROLLARY D.4. Let $\boldsymbol{G} = \boldsymbol{A}_{\Omega}^{T} \boldsymbol{A}_{\Omega}$. The minimum eigenvalue λ_{min} of \boldsymbol{G} lies in the Gershgorin disc $D(1, r_m)$, where $r_m = \max_{i \in \Omega} \sum_{j \in \Omega, j \neq i} \mu_{i,j}$.

Proof. Without loss of generality, we assume that $\Omega = \{1, 2, \dots, |\Omega|\}$. The symmetric matrix G is given by

$$\boldsymbol{G} = \boldsymbol{A}_{\Omega}^{T} \boldsymbol{A}_{\Omega} = \begin{pmatrix} 1 & \mu_{1,2} & \mu_{1,3} & \cdots & \mu_{1,|\Omega|} \\ \mu_{2,1} & 1 & \mu_{2,3} & \cdots & \mu_{2,|\Omega|} \\ \mu_{3,1} & \mu_{3,2} & 1 & \cdots & \mu_{3,|\Omega|} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mu_{|\Omega|,1} & \mu_{|\Omega|,2} & \mu_{|\Omega|,3} & \cdots & 1 \end{pmatrix}$$

It can be seen that the centers of all Gershgorin discs are the same. Thus, all eigenvalues of G lie in the largest Gershgorin disc, i.e., $D(1, r_m)$.

Assume that each column of A is normalized. The proof of Theorem 3.4 is presented as follows.

Proof. Define \mathbb{D} as the event $\left\{ \det \left(\boldsymbol{A}_{\Lambda^{(K)}}^T \boldsymbol{A}_{\Lambda^{(K)}} \right) \neq 0 \right\}$. To ensure the *K*-sparse signal \boldsymbol{x} can be exactly recovered by the OMP algorithm in *K* iterations, the probability $\mathbb{P}\{\mathbb{S}_{\text{omp}}\}$ is equivalent to

(D.2)
$$\mathbb{P}\{\mathbb{S}_{omp}\} = \prod_{k=1}^{K} \mathbb{P}\left\{ \|\boldsymbol{A}_{\Omega}^{T} \boldsymbol{r}^{(k-1)}\|_{\infty} > \|\boldsymbol{A}_{\Omega^{c}}^{T} \boldsymbol{r}^{(k-1)}\|_{\infty} \Big| \mathbb{D}\right\} \mathbb{P}\left\{\mathbb{D}\right\}.$$

During the identification step of OMP algorithm, the algorithm picks a "true" column from \mathbf{A}_{Ω} whenever $\|\mathbf{A}_{\Omega}^{T}\mathbf{r}^{(k-1)}\|_{\infty} > \|\mathbf{A}_{\Omega}^{T}\mathbf{r}^{(k-1)}\|_{\infty}$. These chosen columns determine which elements of \mathbf{x} are non-zero. The values of these elements of \mathbf{x} are determined by solving a least-squares problem, which has a unique solution if $\mathbf{A}_{\Lambda^{(K)}}^{T}\mathbf{A}_{\Lambda^{(K)}}$ has full rank, i.e., det $\left(\mathbf{A}_{\Lambda^{(K)}}^{T}\mathbf{A}_{\Lambda^{(K)}}\right) \neq 0$. Based on Corollary D.2, the probability $\mathbb{P}\left\{\mathbb{D}\right\} = 1 - o(1)$ if the column degree d satisfies $(1 + \varepsilon) \log m \leq d \leq m/2$ for a constant $\varepsilon > 0$ and a sufficiently large m. Thus, we have

(D.3)
$$\mathbb{P}\{\mathbb{S}_{\text{omp}}\} \ge \prod_{k=1}^{K} \mathbb{P}\left\{ \|\boldsymbol{A}_{\Omega}^{T}\boldsymbol{r}^{(k-1)}\|_{\infty} > \|\boldsymbol{A}_{\Omega^{c}}^{T}\boldsymbol{r}^{(k-1)}\|_{\infty} \Big| \mathbb{D}\right\} - o(1).$$

In (D.3), the first term is based on the assumption that the OMP algorithm has picked k "true" columns from A_{Ω} at the (k+1)-th iteration. Thus, the residual $r^{(k-1)}$ can be rewritten as [38]

$$\boldsymbol{r}^{(k-1)} = \boldsymbol{y} - \boldsymbol{A}_{\Lambda^{(k)}}^T \hat{\boldsymbol{x}}_{\Lambda^{(k)}},$$

where the estimated support set $\Lambda^{(k)} \subseteq \Omega$. Since $\boldsymbol{y} = \boldsymbol{A}_{\Omega} \boldsymbol{x}_{\Omega}$ and $\boldsymbol{A}_{\Lambda^{(k)}}$ is a submatrix of A_{Ω} , the residual $r^{(k-1)}$ also can be expressed as a linear combination of columns of A_{Ω} [38], resulting in

$$\boldsymbol{r}^{(k)} = \boldsymbol{A}_{\Omega} \boldsymbol{x}'_{\Lambda^{(k)}} = \boldsymbol{A} \boldsymbol{x}',$$

where the support of x' is contained in the support of x. Thus, the first term of (D.3) can be interpreted as

(D.4)
$$\prod_{k=1}^{K} \mathbb{P}\left\{ \|\boldsymbol{A}_{\Omega}^{T}\boldsymbol{A}_{\Omega}\boldsymbol{x}_{\Lambda^{(k)}}^{\prime}\|_{\infty} > \|\boldsymbol{A}_{\Omega^{c}}^{T}\boldsymbol{A}_{\Omega}\boldsymbol{x}_{\Lambda^{(k)}}^{\prime}\|_{\infty} \middle| \mathbb{D} \right\}.$$

We expect to eliminate $x'_{\Lambda^{(k)}}$ in (D.4) since the distribution of $x'_{\Lambda^{(k)}}$ is unknown. We denote by σ_{min} the minimum singular value of $\mathbf{A}_{\Omega}^{T}\mathbf{A}_{\Omega}$ and λ_{min} the minimum eigenvalue of $\mathbf{A}_{\Omega}^{T}\mathbf{A}_{\Omega}$. Conditioned on the event \mathbb{D} , $\mathbf{A}_{\Omega}^{T}\mathbf{A}_{\Omega}$ is actually a positive definite matrix. In the following, We consider the scaling of $\|\boldsymbol{A}_{\Omega}^{T}\boldsymbol{A}_{\Omega}\boldsymbol{x}'_{\Lambda^{(k)}}\|_{\infty}$.

(D.5a)
$$\|\boldsymbol{A}_{\Omega}^{T}\boldsymbol{A}_{\Omega}\boldsymbol{x}_{\Lambda^{(k)}}^{\prime}\|_{\infty} \geq \frac{1}{\sqrt{K}}\|\boldsymbol{A}_{\Omega}^{T}\boldsymbol{A}_{\Omega}\boldsymbol{x}_{\Lambda^{(k)}}^{\prime}\|_{2}$$

(D.5b)
$$\geq \frac{\sigma_{min}}{\sqrt{K}} \| \boldsymbol{x}_{\Lambda^{(k)}}' \|_2$$

(D.5c)
$$= \frac{\sqrt{\lambda_{min}}}{\sqrt{K}} \| \boldsymbol{x}'_{\Lambda^{(k)}} \|_2$$

(D.5d)
$$\geq \frac{\sqrt{\max\{0, 1 - r_m\}}}{\sqrt{K}} \| \boldsymbol{x}'_{\Lambda^{(k)}} \|_2$$

Here, (D.5b) is based on the basic property of singular value [21]. Since $A_{\Omega}^{T} A_{\Omega}$ is a positive definite matrix, in (D.5c), we have $\sigma_{min} = \sqrt{\lambda_{min}}$ for $\lambda_{min} > 0$. According positive definite matrix, in (D.6c), we have $\sigma_{min} = \sqrt{\lambda_{min}}$ for $\lambda_{min} > 0$. Recording to Corollary D.4, all eigenvalues of $\boldsymbol{A}_{\Omega}^{T} \boldsymbol{A}_{\Omega}$ lie in a Gershgorin circle centered at 1 with a radius of $r_m = \max_{i \in \Omega} \sum_{j \in \Omega, j \neq i} \mu_{i,j}$. With $1 - r_m \leq \lambda_{min} \leq 1 + r_m$ and $\lambda_{min} > 0$, the value of λ_{min} is lower bounded by $\max\{0, 1 - r_m\}$ in (D.5d). Let $\psi = \arg\max_{i \in \Omega} |\boldsymbol{A}_i^T \boldsymbol{r}^{(k)}|$. We further consider the scaling of $\boldsymbol{A}_{\Omega^c}^T \boldsymbol{A}_{\Omega}$.

(D.6a)
$$\|\boldsymbol{A}_{\Omega^{c}}^{T}\boldsymbol{A}_{\Omega}\boldsymbol{x}_{\Lambda^{(k)}}^{\prime}\|_{\infty} = |\boldsymbol{A}_{\psi}^{T}\boldsymbol{A}_{\Omega}\boldsymbol{x}_{\Lambda^{(k)}}^{\prime}|$$

$$(D.6b) = \|\boldsymbol{A}_{\psi}^{\star} \boldsymbol{A}_{\Omega} \boldsymbol{x}_{\Lambda^{(k)}}^{\star}\|_{2}$$

(D.6c)
$$\leq \|\boldsymbol{A}_{\psi}^{t}\boldsymbol{A}_{\Omega}\|_{2}\|\boldsymbol{x}_{\Lambda^{(k)}}^{\prime}\|_{2}$$

(D.6d)
$$= \sqrt{\sum_{i \in \Omega} |\boldsymbol{A}_{\psi}^{T} \boldsymbol{A}_{i}|^{2} \|\boldsymbol{x}_{\Lambda^{(k)}}^{\prime}\|_{2}}$$

(D.6e)
$$= \sqrt{\sum_{i \in \Omega} \mu_{\psi,i}^2} \| \boldsymbol{x}_{\Lambda^{(k)}} \|_2$$

Here, $|\mathbf{A}_{\psi}^{T} \mathbf{A}_{\Omega} \mathbf{x}_{\Lambda^{(k)}}'|$ is a non-zero value, which can be expressed as $\|\mathbf{A}_{\psi}^{T} \mathbf{A}_{\Omega} \mathbf{x}_{\Lambda^{(k)}}'\|_{2}$ in (D.6b). Furthermore, (D.6c) is based on the Cauchy–Schwarz inequality.

With the scalings of $\|\boldsymbol{A}_{\Omega}^{T}\boldsymbol{A}_{\Omega}\boldsymbol{x}_{\Lambda^{(k)}}^{'}\|_{\infty}$ and $\|\boldsymbol{A}_{\Omega^{c}}^{T}\boldsymbol{A}_{\Omega}\boldsymbol{x}_{\Lambda^{(k)}}^{'}\|_{\infty}$, (D.4) is lower bound

by

$$\begin{split} \prod_{k=1}^{K} \mathbb{P}\left\{ \|\boldsymbol{A}_{\Omega}^{T}\boldsymbol{A}_{\Omega}\boldsymbol{x}_{\Lambda^{(k)}}'\|_{\infty} > \|\boldsymbol{A}_{\Omega^{c}}^{T}\boldsymbol{A}_{\Omega}\boldsymbol{x}_{\Lambda^{(k)}}'\|_{\infty} \Big| \mathbb{D} \right\} \\ & \geq \prod_{k=1}^{K} \mathbb{P}\left\{ \frac{\sqrt{\max\{0, 1-r_{m}\}}}{\sqrt{K}} \|\boldsymbol{x}_{\Lambda^{(k)}}'\|_{2} > \sqrt{\sum_{i\in\Omega} \mu_{\psi,i}^{2}} \|\boldsymbol{x}_{\Lambda^{(k)}}'\|_{2} \Big| \mathbb{D} \right\} \\ (\text{D.7a}) & = \mathbb{P}\left\{ \frac{\sqrt{\max\{0, 1-r_{m}\}}}{\sqrt{K}} > \sqrt{\sum_{i\in\Omega} \mu_{\psi,i}^{2}} \Big| \mathbb{D} \right\}^{K}, \end{split}$$

where (D.7a) is independent of k.

Once the distributions of $\sum_{i\in\Omega} \mu_{\psi,i}^2$ and r_m are obtained, one can calculate (D.7a) easily. We first consider the distribution of $\sum_{i\in\Omega} \mu_{\psi,i}^2$. Since columns of \boldsymbol{A} are independent, the distribution of $\mu_{\psi,i}$ for $i \in \Omega$ is given in (3.5). We denote by h(z) the distribution function of $\sum_{i\in\Omega} \mu_{\psi,i}^2$. Let

$$Q = \left\{ \sum_{i=1}^{K} s_i^2 : s_i \in \{0, 1/d, 2/d, \cdots, 1\} \text{ for } i = 1, 2, \cdots, K \right\}.$$

Then, the distribution function h(z) over \mathcal{Q} is given by

$$h(z) = \sum_{s_1^2 + \dots + s_K^2 = z} g(s_1)g(s_2)\cdots g(s_K).$$

We further consider the distribution of $r_m = \max_{i \in \Omega} \sum_{j \in \Omega, j \neq i} \mu_{i,j}$. Without loss of generality, we assume that $\Omega = \{1, 2, \dots, |\Omega|\}$. Let $r_i = \sum_{j=1, j \neq i}^{|\Omega|} \mu_{i,j}$ for $i = 1, 2, \dots, |\Omega|$ and $\phi_m(z)$ be the distribution function of r_m . There are two steps to calculate the distribution of r_m . First, we need to obtain the distribution of r_i , and then use the order statistics [9] to obtain the distribution of the maximum value of $r_1, r_2, \dots, r_{|\Omega|}$. We first consider the distribution of r_i . The value of r_i is the sum of K-1 variables $\mu_{i,j}$ whose the distribution function is given in (3.5). Let

$$\mathcal{P} = \left\{ \sum_{i=1}^{K-1} s_i : s_i \in \{0, 1/d, 2/d, \cdots, 1\} \text{ for } i = 1, 2, \cdots, K-1 \right\}$$

and $\phi(z)$ be the distribution function of r_i . Then, the distribution function $\phi(z)$ over \mathcal{P} is given by

$$\phi(z) = \sum_{s_1 + \dots + s_{K-1} = z} g(s_1)g(s_2) \cdots g(s_{K-1})$$

In the next step, we need to use the order statistics to obtain the distribution of the maximum value of r_i . Prior to this, we have to get the CDF of $\phi(z)$, which is given by

$$\Phi(z) = \sum_{z' \le z} \phi(z').$$

With the distribution $\phi(z)$ and its CDF $\Phi(z)$, the distribution of r_m over \mathcal{P} is

$$\phi_m(z) = (\Phi(z))^K - (\Phi(z) - \phi(z))^K.$$

So far, we have obtained the distributions of $\sum_{i \in \Omega} \mu_{\psi,i}^2$ and r_m . By the law of total probability, the base of (D.7a) can be calculated as

$$\begin{split} \mathbb{P}\left\{\frac{\sqrt{\max\{0,1-r_m\}}}{\sqrt{K}} > \sqrt{\sum_{i\in\Omega}\mu_{\psi,i}^2} \left| \mathbb{D} \right\} \\ (\mathrm{D.8a}) &= \sum_{p\in\mathcal{P}}\sum_{q\in\mathcal{Q}} \mathbb{P}\left\{\frac{\sqrt{\max\{0,1-p\}}}{\sqrt{K}} > \sqrt{q} \left| r_m = p, \sum_{i\in\Omega}\mu_{\psi,i}^2 = q, \mathbb{D} \right\} \right. \\ &\times \mathbb{P}\left\{r_m = p \left| \mathbb{D} \right\} \mathbb{P}\left\{\sum_{i\in\Omega}\mu_{\psi,i}^2 = q \left| \mathbb{D} \right\} \right. \\ (\mathrm{D.8b}) &= \sum_{p\in\mathcal{P}}\sum_{q\in\mathcal{Q}} \mathbb{P}\left\{\frac{\sqrt{\max\{0,1-p\}}}{\sqrt{K}} > \sqrt{q} \left| r_m = p, \sum_{i\in\Omega}\mu_{\psi,i}^2 = q, \mathbb{D} \right\} \phi_m(p)h(q) \right. \end{split}$$

In (D.8a), $\{r_m = p\}$ and $\left\{\sum_{i \in \Omega} \mu_{\psi,i}^2 = q\right\}$ are independent and both of them are independent of the event \mathbb{D} . Furthermore, the first factor in (D.8b) can be represented by an indicator function, which is given by

$$\mathbf{1}(p,q) = \begin{cases} 1 & \text{if } \frac{\sqrt{\max\{0,1-p\}}}{\sqrt{K}} > \sqrt{q}, \\ 0 & \text{otherwise.} \end{cases}$$

As a result, (D.4) is lower bounded by (D.9)

$$\prod_{k=1}^{K} \mathbb{P}\left\{ \|\boldsymbol{A}_{\Omega}^{T}\boldsymbol{A}_{\Omega}\boldsymbol{x}_{\Lambda^{(k)}}^{\prime}\|_{\infty} > \|\boldsymbol{A}_{\Omega^{c}}^{T}\boldsymbol{A}_{\Omega}\boldsymbol{x}_{\Lambda^{(k)}}^{\prime}\|_{\infty} \Big| \mathbb{D}\right\} \geq \left(\sum_{p \in \mathcal{P}} \sum_{q \in \mathcal{Q}} \mathbf{1}(p,q)\phi_{m}(p)h(q)\right)^{K}.$$

Combined with (D.3) and (D.9), we get (3.6).

Appendix E. The proof of Theorem 3.5.

Proof. We first define \mathbb{D} as the event $\left\{ \det \left(\boldsymbol{A}_{\Lambda^{(K)}}^T \boldsymbol{A}_{\Lambda^{(K)}} \right) \neq 0 \right\}$. By the law of total probability, we have

$$\mathbb{P}\{\mathbb{S}_{\text{comp}}\} = \mathbb{P}\{\mathbb{S}_{\text{comp}} | |\Gamma| = K\} \mathbb{P}\{|\Gamma| = K\} + \mathbb{P}\{\mathbb{S}_{\text{comp}} | |\Gamma| > K\} \mathbb{P}\{|\Gamma| > K\}$$

(E.1a)
$$= \mathbb{P}\left\{\mathbb{D}\right\} \mathbb{P}\left\{|\Gamma| = K\right\} + \mathbb{P}\left\{\mathbb{S}_{\text{comp}} \left||\Gamma| > K\right\} \mathbb{P}\left\{|\Gamma| > K\right\}$$

(E.1b)
$$\geq \mathbb{P} \{ \mathbb{D} \} \mathbb{P} \{ |\Gamma| = K \}$$

As shown in Algorithm 2.1, the identification is already done if $|\Gamma| = K$. Thus, the probability $\mathbb{P} \{ \mathbb{S}_{\text{comp}} | |\Gamma| = K \}$ is reduced to the probability that the least squares has a unique solution, i.e., $\mathbb{P} \{ \mathbb{D} \}$. In (E.1a), the derivation of the lower bound on $\mathbb{P} \{ \mathbb{S}_{\text{comp}} | |\Gamma| > K \}$ is similar to that of OMP algorithm. The details are given in Appendix F. However, our experimental results reveal that the first term of (E.1a) is dominant. For simplicity, we only consider the contribution of the first term of (E.1a), resulting in (E.1b).

With Corollary D.2, the probability $\mathbb{P} \{\mathbb{D}\} = 1 - o(1)$ for a sufficiently large *m*. Thus, we have

$$\mathbb{P}\{\mathbb{S}_{\text{comp}}\} \ge \mathbb{P}\{|\Gamma| = K\} - o(1).$$

21

Given any $j \in \Omega^c$, the probability $\mathbb{P}\{|\Gamma| = K\}$ is equivalent to the probability $\mathbb{P}\{j \notin I\}$ Γ ^{$|\Omega^c|}, which is given by</sup>$

$$\mathbb{P}\{|\Gamma| = K\} = \mathbb{P}\{j \notin \Gamma\}^{|\Omega^c|}$$
$$= (1 - \mathbb{P}\{j \in \Gamma \setminus \Omega\})^{n-K}$$
$$= \left(1 - \sum_{v=d}^{Kd} \frac{\binom{v}{d}}{\binom{m}{d}} \mathbb{P}\{\nu^{(K)} = v\}\right)^{n-K},$$

where $\mathbb{P}\{\nu^{(k)} = v\}$ is given in Lemma 3.1 and $\mathbb{P}\{j \in \Gamma \setminus \Omega\}$ is obtained in (C.2). Thus, we get (3.7).

Appendix F. The lower bound on $\mathbb{P} \{ \mathbb{S}_{comp} | |\Gamma| > K \}$. During the identification step of confined OMP algorithm, the choice of column index is confined into the set Γ . We use q'(s) to denote the distribution of coherence between two columns whose indices are in Γ . The distribution q'(s) is different from the distribution q(s)given in (3.5), which is given by

(F.1)
$$g_1(s) = \mathbb{P}\left\{\mu_{i,j} = s\right\} = \sum_{\upsilon=d}^{Kd} \frac{\binom{d}{sd}\binom{\upsilon-d}{d-sd}}{\binom{\upsilon}{d}} \mathbb{P}\left\{\nu^{(K)} = \upsilon\right\} \quad \text{for } i, j \in \Gamma,$$

where $\mathbb{P}\{\nu^{(k)} = v\}$ is given in Lemma 3.1 and $s \in \{0, 1/d, \dots, 1\}$.

The following steps are similar to those of OMP algorithm. We denote by h'(z)the distribution function of $\sum_{i \in \Omega} \mu_{\psi',i}^2$ where $\psi' = \underset{i \in \Gamma \setminus \Omega}{\operatorname{arg\,max}} |\mathbf{A}_i^T \mathbf{r}^{(k)}|$. The distribution

function h'(z) over \mathcal{Q} is given by

$$h'(z) = \sum_{s_1^2 + \dots + s_K^2 = z} g'(s_1)g'(s_2) \cdots g'(s_K).$$

We use $\phi'(z)$ to denote the distribution function of $r_i = \sum_{j \in \Omega, j \neq i} \mu_{i,j}$ for $i \in \Omega$. Then, we have

$$\phi'(z) = \sum_{s_1 + \dots + s_{K-1} = z} g'(s_1) g'(s_2) \cdots g'(s_{K-1}),$$

where $z \in \mathcal{P}$. We further get the CDF of $\phi'(z)$, which is given by

$$\Phi'(z) = \sum_{z' \le z} \phi'(z').$$

With the distribution $\phi'(z)$ and its CDF $\Phi'(z)$, the distribution of r_m is given by

$$\phi'_{m}(z) = (\Phi'(z))^{K} - (\Phi'(z) - \phi'(z))^{K}$$

As a result, the probability $\mathbb{P}\left\{\mathbb{S}_{\text{comp}} | |\Gamma| > K\right\}$ is lower bounded by

(F.2)
$$\mathbb{P}\left\{\mathbb{S}_{\text{comp}}\Big||\Gamma| > K\right\} \ge \left(\sum_{p \in \mathcal{P}} \sum_{q \in \mathcal{Q}} \mathbf{1}(p,q) \phi'_m(p) h'(q)\right)^K - o(1),$$

where the indicator function is

$$\mathbf{1}(p,q) = \begin{cases} 1 & \text{if } \frac{\sqrt{\max\{0,1-p\}}}{\sqrt{K}} > \sqrt{q}, \\ 0 & \text{otherwise.} \end{cases}$$

REFERENCES

- R. BARANIUK, M. DAVENPORT, R. DEVORE, AND M. WAKIN, A simple proof of the restricted isometry property for random matrices, Constr. Approx., 28 (2008), pp. 253–263, https:// doi.org/https://doi.org/10.1007/s00365-007-9003-x.
- A. BECK AND M. TEBOULLE, A fast iterative shrinkage-thresholding algorithm for linear inverse problems, SIAM J. Imag. Sci., 2 (2009), pp. 183–202, https://doi.org/10.1137/080716542.
- [3] R. BERINDE AND P. INDYK, Sparse recovery using sparse random matrices, MIT-CSAIL Tech. Rep., (2008).
- T. BLUMENSATH AND M. E. DAVIES, Iterative hard thresholding for compressed sensing, Appl. Comput. Harmon. Anal., 27 (2009), pp. 265–274, https://doi.org/https://doi.org/10.1016/ j.acha.2009.04.002.
- [5] E. CANDÈS, J. ROMBERG, AND T. TAO, Robust uncertainty principles: exact signal reconstruction from highly incomplete frequency information, IEEE Trans. Inf. Theory, 52 (2006), pp. 489–509, https://doi.org/10.1109/TIT.2005.862083.
- [6] E. CANDÈS AND T. TAO, Decoding by linear programming, IEEE Trans. Inf. Theory, 51 (2005), pp. 4203–4215, https://doi.org/10.1109/TIT.2005.858979.
- S. S. CHEN, D. L. DONOHO, AND M. A. SAUNDERS, Atomic decomposition by basis pursuit, SIAM Rev., 43 (2001), pp. 129–159, https://doi.org/10.1137/S003614450037906X.
- W. DAI AND O. MILENKOVIC, Subspace pursuit for compressive sensing signal reconstruction, IEEE Trans. Inf. Theory, 55 (2009), pp. 2230–2249, https://doi.org/10.1109/TIT.2009. 2016006.
- [9] H. A. DAVID AND H. N. NAGARAJA, Order statistics, John Wiley & Sons, New York, USA, 2004.
- [10] A. G. DIMAKIS, R. SMARANDACHE, AND P. O. VONTOBEL, Ldpc codes for compressed sensing, IEEE Trans. Inf. Theory, 58 (2012), pp. 3093–3114, https://doi.org/10.1109/TIT.2011. 2181819.
- [11] D. DONOHO, De-noising by soft-thresholding, IEEE Trans. Inf. Theory, 41 (1995), pp. 613–627, https://doi.org/10.1109/18.382009.
- D. DONOHO, Compressed sensing, IEEE Trans. Inf. Theory, 52 (2006), pp. 1289–1306, https:// doi.org/10.1109/TIT.2006.871582.
- [13] D. L. DONOHO AND M. ELAD, Optimally sparse representation in general (nonorthogonal) dictionaries via l¹ minimization, Proc. Natl. Acad. Sci. U.S.A., 100 (2003), pp. 2197– 2202, https://doi.org/10.1073/pnas.0437847100.
- [14] M. F. DUARTE, M. A. DAVENPORT, D. TAKHAR, J. N. LASKA, T. SUN, K. F. KELLY, AND R. G. BARANIUK, Single-pixel imaging via compressive sampling, IEEE Signal Process Mag., 25 (2008), pp. 83–91, https://doi.org/10.1109/MSP.2007.914730.
- [15] B. EFRON, T. HASTIE, I. JOHNSTONE, AND R. TIBSHIRANI, Least angle regression, Ann. Statist., 32 (2004), pp. 407 – 499, https://doi.org/10.1214/009053604000000067.
- [16] A. FERBER, M. KWAN, AND L. SAUERMANN, Singularity of sparse random matrices: simple proofs, Comb. Probab. Comput., 31 (2022), p. 21–28, https://doi.org/10.1017/ S0963548321000146.
- [17] S. FOUCART, Hard thresholding pursuit: An algorithm for compressive sensing, SIAM J. Numer. Anal., 49 (2011), pp. 2543–2563, https://doi.org/10.1137/100806278.
- [18] S. FOUCART AND H. RAUHUT, A mathematical introduction to compressive sensing, Birkhäuser, Basel, Switzerland, 1th ed., 2013.
- [19] A. GILBERT AND P. INDYK, Sparse recovery using sparse matrices, Proc. IEEE, 98 (2010), pp. 937–947, https://doi.org/10.1109/JPROC.2010.2045092.
- [20] A. C. GILBERT, M. A. IWEN, AND M. J. STRAUSS, Group testing and sparse signal recovery, in 2008 42nd Asilomar Conference on Signals, Systems and Computers, 2008, pp. 1059–1063, https://doi.org/10.1109/ACSSC.2008.5074574.
- [21] R. A. HORN AND C. R. JOHNSON, Matrix analysis, Cambridge Univ. Press, Cambridge, U.K., 2012.
- [22] S. JAFARPOUR, W. XU, B. HASSIBI, AND R. CALDERBANK, Efficient and robust compressed sensing using optimized expander graphs, IEEE Trans. Inf. Theory, 55 (2009), pp. 4299– 4308, https://doi.org/10.1109/TIT.2009.2025528.
- [23] M. A. KHAJEHNEJAD, A. G. DIMAKIS, W. XU, AND B. HASSIBI, Sparse recovery of nonnegative signals with minimal expansion, IEEE Trans. Signal Process., 59 (2011), pp. 196–208, https://doi.org/10.1109/TSP.2010.2082536.
- [24] S. KWON, J. WANG, AND B. SHIM, Multipath matching pursuit, IEEE Trans. Inf. Theory, 60 (2014), pp. 2986–3001, https://doi.org/10.1109/TIT.2014.2310482.
- [25] E. LIU AND V. N. TEMLYAKOV, The orthogonal super greedy algorithm and applications in

compressed sensing, IEEE Trans. Inf. Theory, 58 (2012), pp. 2040–2047, https://doi.org/10.1109/TIT.2011.2177632.

- [26] X.-J. LIU, S.-T. XIA, AND F.-W. FU, Reconstruction guarantee analysis of basis pursuit for binary measurement matrices in compressed sensing, IEEE Trans. Inf. Theory, 63 (2017), pp. 2922–2932, https://doi.org/10.1109/TIT.2017.2677965.
- [27] M. LOTFI AND M. VIDYASAGAR, Compressed sensing using binary matrices of nearly optimal dimensions, IEEE Trans. Signal Process., 68 (2020), pp. 3008–3021, https://doi.org/10. 1109/TSP.2020.2990154.
- [28] W. LU, T. DAI, AND S.-T. XIA, Binary matrices for compressed sensing, IEEE Trans. Signal Process., 66 (2018), pp. 77–85, https://doi.org/10.1109/TSP.2017.2757915.
- [29] D. NEEDELL AND J. TROPP, CoSaMP: Iterative signal recovery from incomplete and inaccurate samples, Appl. Comput. Harmon. Anal., 26 (2009), pp. 301–321, https://doi.org/https:// doi.org/10.1016/j.acha.2008.07.002.
- [30] D. NEEDELL AND R. VERSHYNIN, Signal recovery from incomplete and inaccurate measurements via regularized orthogonal matching pursuit, IEEE J. Sel. Top. Signal Process., 4 (2010), pp. 310–316, https://doi.org/10.1109/JSTSP.2010.2042412.
- [31] F. PARVARESH, H. VIKALO, S. MISRA, AND B. HASSIBI, Recovering sparse signals using sparse measurement matrices in compressed dna microarrays, IEEE J. Sel. Top. Signal Process, 2 (2008), pp. 275–285, https://doi.org/10.1109/JSTSP.2008.924384.
- [32] R. RUBINSTEIN, M. ZIBULEVSKY, AND M. ELAD, Efficient implementation of the k-svd algorithm using batch orthogonal matching pursuit, Technical Report Computer Science Department, Technion, 40 (2008), pp. 1–15.
- [33] J. TANNER AND K. WEI, Normalized iterative hard thresholding for matrix completion, SIAM J. Sci. Comput., 35 (2013), pp. S104–S125, https://doi.org/10.1137/120876459.
- [34] J. TROPP, Greed is good: algorithmic results for sparse approximation, IEEE Trans. Inf. Theory, 50 (2004), pp. 2231–2242, https://doi.org/10.1109/TIT.2004.834793.
- [35] J. A. TROPP AND A. C. GILBERT, Signal recovery from random measurements via orthogonal matching pursuit, IEEE Trans. Inf. Theory, 53 (2007), pp. 4655–4666, https://doi.org/10. 1109/TIT.2007.909108.
- [36] M. J. WAINWRIGHT, Sharp thresholds for high-dimensional and noisy sparsity recovery using l₁ -constrained quadratic programming (Lasso), IEEE Trans. Inf. Theory, 55 (2009), pp. 2183– 2202, https://doi.org/10.1109/TIT.2009.2016018.
- [37] J. WANG, S. KWON, AND B. SHIM, Generalized orthogonal matching pursuit, IEEE Trans. Signal Process., 60 (2012), pp. 6202–6216, https://doi.org/10.1109/TSP.2012.2218810.
- [38] J. WANG AND B. SHIM, On the recovery limit of sparse signals using orthogonal matching pursuit, IEEE Trans. Signal Process., 60 (2012), pp. 4973–4976.
- [39] J. WEN AND H. LI, Binary sparse signal recovery with binary matching pursuit*, Inverse Probl., 37 (2021), p. 065014, https://doi.org/10.1088/1361-6420/abf903.
- [40] J. WEN, R. ZHANG, AND W. YU, Signal-dependent performance analysis of orthogonal matching pursuit for exact sparse recovery, IEEE Trans. Signal Process., 68 (2020), pp. 5031–5046.
- [41] J. WEN, Z. ZHOU, J. WANG, X. TANG, AND Q. MO, A sharp condition for exact support recovery with orthogonal matching pursuit, IEEE Trans. Signal Process., 65 (2017), pp. 1370–1382, https://doi.org/10.1109/TSP.2016.2634550.