arXiv:2501.01112v1 [math.CO] 2 Jan 2025 [arXiv:2501.01112v1 \[math.CO\] 2 Jan 2025](http://arxiv.org/abs/2501.01112v1)

CONNECTED IDEALS OF CHORDAL GRAPHS

KANOY KUMAR DAS, AMIT ROY, AND KAMALESH SAHA

ABSTRACT. For $t \geq 2$, the t-independence complex of a graph G is the collection of all $A \subseteq V(G)$ such that each connected component of the induced subgraph $G[A]$ has at most $t-1$ vertices. The Stanley-Reisner ideal $I_t(G)$ of the t-independence complex of G, called t-connected ideal, is generated by monomials in a polynomial ring R corresponding to all $A \subseteq V(G)$ of size t such that $G[A]$ is connected. This class of ideals is a natural generalization of the edge ideals of graphs. In this paper, we investigate the t-connected ideals of chordal graphs. In particular, we prove that for a chordal graph G and for all t

 $reg(R/I_t(G)) = (t-1)\nu_t(G)$ and $pd(R/I_t(G)) = bight(I_t(G)),$

where $\nu_t(G)$ denotes the induced matching number of the corresponding hypergraph of $I_t(G)$, and reg, pd and bight stand for the regularity, projective dimension, and big height, respectively. As a consequence of the above results, we completely characterize when the t-connected ideal of a chordal graph has a linear resolution as well as when it satisfies the Cohen-Macaulay property. The above formulas and their consequences can be seen as a nice generalization of the classical results corresponding to the edge ideals of chordal graphs.

1. INTRODUCTION

An integral focus of research in the area of commutative algebra is the study of monomial ideals, particularly square-free monomial ideals, due to their strong connections with combinatorics and topology. A general objective in this area is to express or translate algebraic properties of a certain class of ideals in terms of the combinatorial or topological properties of the associated objects. There are several ways to associate a combinatorial object to a square-free monomial ideal; the most popular among them are the following two: (a) using the Stanley-Reisner correspondence to associate an abstract simplicial complex, and (b) associating a simple hypergraph (or clutter). Both these identifications have certain advantages and are frequently used to study square-free monomial ideals. Additionally, many algebraic properties of an ideal in a polynomial ring depend on the characteristics of the base field. However, if these come from the structure of the associated combinatorial object, then they are independent of the choice of the base field.

The graded minimal free resolution of a graded module gives insight into its structure and measures its complexity. Determining the minimal free resolution of a graded module is computationally a challenging task. So, researchers try to get some estimation of the minimal free resolution via two important invariants: (i) the Castelnuovo-Mumford regularity (or simply, regularity) that measures the width of a minimal free resolution, and (ii) the projective dimension, which gives the length of a minimal free resolution. These two invariants have been extensively investigated for several classes of monomial ideals, more notably in the case of edge ideals of graphs. Indeed, a celebrated theorem of Fröberg [\[10\]](#page-13-0) gave an algebraic interpretation of chordal graphs in terms of the linearity of the minimal free resolution of edge ideals. More precisely, the edge ideal $I(G)$ of a graph G has a linear resolution if and only if the complement of G is chordal. On the other hand, when G is chordal, a precise combinatorial formula for the regularity and projective dimension of $I(G)$ are well-known.

To extend the study to square-free monomial ideals, various generalizations of edge ideals, such as path ideals, clique ideals, etc., have been introduced (see [\[4,](#page-13-1) [17\]](#page-13-2)). Note that the edge ideal of a graph G can be realized as the Stanley-Reisner ideal of a simplicial complex obtained from G , often referred to as the independence complex of G . In the literature, there is a notion of higher

²⁰²⁰ *Mathematics Subject Classification.* 05E40, 13F55, 05C05.

Key words and phrases. connected ideals, chordal graphs, regularity, projective dimension, linear resolution, Cohen-Macaulay property.

independence complexes of a graph G, generalizing the independence complex of G. For $t \geq 2$, the t-independence complex of a graph G, denoted by $\text{Ind}_t(G)$, is the collection of all $A \subseteq V(G)$ such that each connected component of $G[A]$ has at most $t - 1$ vertices. Recently, Stanley-Reisner ideals of these complexes have been considered in [\[1,](#page-12-0) [2,](#page-12-1) [7\]](#page-13-3). In this article, we broaden this study by considering the class of chordal graphs. It turns out that the monomial generators of the Stanley-Reisner ideal of the t -independence complex of a graph G correspond to the connected subgraphs of size t in G. This ideal is called the t-connected ideal of G, and we denote it by $I_t(G)$.

There are several motivations for studying higher independence complexes or connected ideals of graphs, some of which are listed below.

- (1) The *t*-independence complex of a graph appeared in the work of Szabó and Tardos [\[23\]](#page-13-4), where they introduced and discussed generalizations of the problem of independent transversal in graphs.
- (2) In [\[16\]](#page-13-5), Meshulam related the homology groups of $\text{Ind}_2(G)$ with the domination number of G. Recently, Deshpande-Shukla-Singh [\[8\]](#page-13-6) extends Meshulam's result by relating the homology groups of $Ind_t(G)$ with the distance t-domination number of G. It is important to note that the distance t -domination number is a well-known invariant in graph theory (see [\[25\]](#page-13-7) and the references therein).
- (3) In [\[8\]](#page-13-6), the authors have also shown that Ind_t(G) of a chordal graph are either contractible or homotopy equivalent to a wedge of spheres. Note that, for $t = 2$, $\text{Ind}_2(G)$ of a chordal graph G is sequentially Cohen-Macaulay, whereas for each $t \geq 3$, the complexes $\mathrm{Ind}_t(G)$ may not be sequentially Cohen-Macaulay (see [\[1,](#page-12-0) Proposition 4.3]), which is quite surprising.
- (4) In [\[19\]](#page-13-8), Paolini and Salvetti established a connection between the twisted cohomology of the classical braid groups and the cohomology of higher independence complexes related to the corresponding Coxeter graphs (see also [\[21\]](#page-13-9)).
- (5) Also, the notion of t -independent set has been explored from a purely graph-theoretic point of view. Specifically, it is related to the idea of clustered graph coloring (see [\[22,](#page-13-10) [26\]](#page-13-11)).
- (6) The class of t-connected ideals of graphs is a natural generalization of edge ideals as $I_2(G)$ = $I(G)$. In this regard, one should note that the t-path ideals of graphs are also a generalization of edge ideals, which are relatively well-studied, and for $t \leq 3$, t-path ideals coincide with t-connected ideals.
- (7) The authors in [\[7\]](#page-13-3) showed that a partial analogue of Fröberg's theorem naturally extends to all t-connected ideals.

It is well-known that any square-free monomial ideal can be seen as an edge ideal of a simple hypergraph. Let H be a t-uniform hypergraph, and $I(H)$ denote its edge ideal in a polynomial ring R. Then the regularity (respectively, projective dimension) of $R/I(H)$ is bounded below by $(t - 1)\nu(\mathcal{H})$ (respectively, bight($I(\mathcal{H})$)), where $\nu(\mathcal{H})$ denote the induced matching number of H. These bounds are attained for various classes of simple graphs (i.e., 2-uniform hypergraphs), including the chordal one (see $[3]$). In the case of general t -uniform hypergraphs, only a few classes are known for which $reg(R/I(\mathcal{H}))$ attains the lower bound.

Let $\mathcal{H}_t(G)$ be a t-uniform hypergraph induced from a simple graph G such that $\mathcal{H}_2(G) = G$, i.e., $I(\mathcal{H}_t(G))$ can be viewed as a generalization of $I(G)$; for instance, path ideals, connected ideals, and clique ideals of graphs, etc. One of the natural questions in this context is to ask for which classes of such hypergraphs $\mathcal{H}_t(G)$ the well-known results corresponding to $I(G)$ carry forward to higher t. Note that among different classes of simple graphs, chordal graphs have garnered special attention due to their connections with various branches of mathematics and computer science, as well as the fact that several algebraic invariants of their edge ideals can be expressed in terms of combinatorial invariants of the underlying graphs. Thus, it is worthwhile to first explore the above question in the context of chordal graphs.

In this paper, we investigate the *t*-connected ideal $I_t(G)$ corresponding to a chordal graph G. Specifically, we are interested to know whether the regularity and the projective dimension of such ideals can be expressed in terms of the combinatorial invariants of the associated hypergraphs, as mentioned above. The first main theorem along this direction is the following:

Theorem [3.6.](#page-7-0) *Let* G *be a chordal graph. Then for any* $t \geq 2$ *,*

$$
reg(R/I_t(G)) = (t-1)\nu_t(G),
$$

where $\nu_t(G)$ *denotes the induced matching number of the hypergraph corresponding to* $I_t(G)$ *.*

As a corollary of the above theorem, we characterize when the t-connected ideal of a chordal graph has a linear resolution as follows:

Corollary [3.7.](#page-8-0) Let G be a chordal graph and $t \geq 2$ be an integer. Then $I_t(G)$ has a linear resolution if *and only if* G *is* t -gap-free (i.e., $\nu_t(G) = 1$).

The above corollary extends [\[2,](#page-12-1) Theorem 5.1], where they proved the same result for trees, a subclass of chordal graphs. Next, we establish the following combinatorial formula for the projective dimension of $R/I_t(G)$:

Theorem [4.5.](#page-9-0) *Let* G *be a chordal graph. Then for all* $t \geq 2$, $pd(R/I_t(G)) = bight(I_t(G))$ *.*

As an application of the above theorem, we combinatorially characterize all Cohen-Macaulay tconnected ideals of chordal graphs, which ensures that the Cohen-Macaulay property of such ideals does not depend on the characteristic of the base field.

Corollary [4.8.](#page-11-0) *Let* G *be a chordal graph and* $t \geq 2$ *be an integer. Then* $I_t(G)$ *is Cohen-Macaulay if and only if* $I_t(G)$ *is unmixed.*

The above result generalizes a famous theorem of Herzog-Hibi-Zheng [\[14\]](#page-13-13), where they classified all Cohen-Macaulay chordal graphs.

In the spirit of Theorem [3.6](#page-7-0) and [4.5,](#page-9-0) one can try to obtain similar formulas for the regularity and projective dimension in the case of path ideals and clique ideals of graphs. We remark that an extensive amount of work is available in the literature on the t-path ideals of graphs (see [\[6,](#page-13-14) [12\]](#page-13-15) and the references therein). Meanwhile, recently in [\[6\]](#page-13-14), we were able to show that the regularity and projective dimension formulas of edge ideals of chordal graphs in terms of the induced matching number and big height do not extend to t-path ideals for $t \geq 4$, even for the class of trees. Now, if one considers the t-clique ideal of a tree, then it is easy to see that the ideal is a zero ideal for $t \geq 3$. Regarding clique ideals of chordal graphs, we show that the above-mentioned formula of regularity can not be extended to higher t (Example [5.2\)](#page-12-2).

The paper is structured in the following way. In Section [2,](#page-2-0) we recall some standard notions and results of combinatorics and commutative algebra. In Section [3,](#page-5-0) we establish the regularity formula of t-connected ideals of chordal graphs and characterize when such ideals have linear resolutions. In Section [4,](#page-8-1) we derive the formula for the projective dimension of t-connected ideals of chordal graphs and characterize when such ideals are Cohen-Macaulay. We make some concluding remarks in Section [5.](#page-11-1)

2. PRELIMINARIES

In this section, we recall some preliminary notions from combinatorics and commutative algebra, which are used throughout the rest of the paper.

2.1. **Graph Theory and Combinatorics:** A graph G is a pair $(V(G), E(G))$, where $V(G)$ is called the vertex set of G and $E(G)$, a collection of subsets of $V(G)$ of size 2, is known as the edge set of G. We now recall some useful notations related to a graph G that will be needed in the later sections.

(1) If $x_1, \ldots, x_m \in V(G)$, then $G \setminus \{x_1, \ldots, x_m\}$ denotes the graph with the vertex set $V(G) \setminus$ ${x_1, \ldots, x_m}$ and the edge set $\{u, v\} \in E(G) \mid x_i \notin \{u, v\}$ for each $i \in [m]$. If $m = 1$, then $G \setminus \{x_1\}$ is simply denoted by $G \setminus x_1$.

- (2) For $C \subseteq V(G)$, the set of *neighbors* of C, denoted by $N_G(C)$, is the set $\{w \in V(G) \setminus C \mid$ $\{w, x\} \in E(G)$ for some $x \in C$. The set of *closed neighbors* of C is the set $N_G(C) \cup C$ and is denoted by $N_G[C]$. If $C = \{a\}$ for some $a \in V(G)$, then we simply denote the sets $N_G(\{a\})$ and N_G [{a}] as $N_G(a)$ and $N_G[a]$, respectively.
- (3) Let $W \subseteq V(G)$. Then the *induced subgraph of G on* W, denoted by $G[W]$, is the graph on the vertex set W with the edge set $\{e \in E(G) \mid e \subseteq W\}$. Note that $G[W] = G \setminus (V(G) \setminus W)$.
- (4) Let $t~\geq~2$ be an integer, and $\mathcal{U}~=~\{C_1,\ldots,C_r~:~|C_i|~=~t, G[C_i]$ is connected, $C_i \, \cap \,$ $C_j = \emptyset$ for all $1 \leq i \leq j \leq r$. We say that $\mathcal U$ is a *t*-connected induced matching of G if $E(G[\cup_{i=1}^{r} C_i]) = \cup_{i=1}^{r} E(G[C_i]).$ The *t*-connected induced matching number of G, denoted by $\nu_t(G)$, is given by $\nu_t(G) = \max\{|\mathcal{U}| : \mathcal{U}$ is a t-connected induced matching of $G\}$. If $t = 2$, then $\nu_2(G)$ is called the *induced matching number* of G and is also denoted in the literature as $\nu(G)$. We say G is *t*-*gap*-free whenever $\nu_t(G) = 1$.

Various classes of simple graphs:

- (1) A *path* graph P_n of length n is a graph with the vertex set $\{x_1, \ldots, x_{n+1}\}$, and the edge set $\{\{x_i, x_{i+1}\} \mid 1 \leq i \leq n\}$. A cycle C_n of length n is a graph with the vertex set $\{x_1, \ldots, x_n\}$, and the edge set $\{\{x_1, x_n\}, \{x_i, x_{i+1}\} \mid 1 \leq i \leq n-1\}.$
- (2) For a positive integer m, a *complete graph* K_m is a graph on m vertices such that there is an edge between any two distinct vertices.
- (3) A graph G is called *chordal* if it contains an induced cycle C_n of length at most 3. If G is a chordal graph, then G contains at least one vertex x such that $N_G(x)$ is a complete graph (see [\[9\]](#page-13-16)). Such a vertex is called a *simplicial vertex* of G. Note that any induced subgraph of a chordal graph is again a chordal graph.

2.2. The **t-connected ideal:** Let G be a graph with $V(G) = \{x_1, \ldots, x_n\}$, and R denotes the polynomial ring $\mathbb{K}[x_1,\ldots,x_n]$, where $\mathbb K$ is a field. The square-free monomial ideal

$$
I_t(G) = \left\langle \mathbf{x}_C := \prod_{x_i \in C} x_i \mid C \subseteq V(G), |C| = t, \text{ and } G[C] \text{ is connected} \right\rangle
$$

in the polynomial ring R is called the t*-connected ideal* of G.

2.3. Connected ideal as edge ideal of a hypergraph: A *hypergraph* H is a pair $(V(H), E(H))$, where $E(\mathcal{H})\subseteq 2^{V(\mathcal{H})},$ and for any two $\mathcal{E}_1,\mathcal{E}_2\in E(\mathcal{H}),$ $\mathcal{E}_1\not\subset\mathcal{E}_2.$ The sets $V(\mathcal{H})$ and $E(\mathcal{H})$ are called the vertex set and edge set of H, respectively. For a fixed positive integer m, if $|\mathcal{E}| = m$ for each $\mathcal{E} \in E(\mathcal{H})$, then we say that H is an *m*-uniform hypergraph. Note that if H is a 2-uniform hypergraph, then H is just a graph. As in the case of graphs, if $A \subseteq V(H)$, then $H \setminus A$ denotes the hypergraph with the vertex set $V(\mathcal{H}) \setminus A$, and the edge set $\{\mathcal{E} \in E(\mathcal{H}) \mid \mathcal{E} \cap A = \emptyset\}$. Similarly, for any $A \subseteq V(H)$, the hypergraph $H \setminus (V(H) \setminus A)$ is called the *induced subhypergraph* of H on the vertex set A. For $x \in V(\mathcal{H})$, we simply write $\mathcal{H} \setminus x$ to denote the hypergraph $\mathcal{H} \setminus \{x\}$. A subset $U \subseteq V(H)$ is called a *vertex cover* of H if for any edge $\mathcal{E} \in E(H)$ one has $\mathcal{E} \cap U \neq \emptyset$. A *minimal vertex cover* of H is a vertex cover that is minimal with respect to inclusion.

Let H be a hypergraph on the vertex set $\{x_1, \ldots, x_n\}$ and let $R = \mathbb{K}[x_1, \ldots, x_n]$. Corresponding to each $\mathcal{E} \in E(\mathcal{H})$, one can assign the monomial $\mathbf{x}_{\mathcal{E}} = \prod_{x_j \in \mathcal{E}} x_j$ in R . Then the ideal $\langle \mathbf{x}_{\mathcal{E}} |$ $\mathcal{E} \in E(\mathcal{H})$ is called the *edge ideal* of H, and is denoted by $I(\mathcal{H})$. Let $I \subseteq R$ be a square-free monomial ideal with the unique minimal monomial generating set $\mathcal{G}(I)$. Then I can be viewed as an edge ideal of a hypergraph \mathcal{H}_I , where $V(\mathcal{H}_I)=\{x_1,\ldots,x_n\}$ and $E(\mathcal{H}_I)=\{\{x_{i_1},\ldots,x_{i_r}\}\mid$ $x_{i_1}\cdots x_{i_r}\,\in\mathcal{G}(I)\}.$ In other words, we have $I\,=\,I(\mathcal{H}_I).$ It is well-known in the literature that the minimal prime ideals of I (equivalently, the associated primes of I since I is a radical ideal) are exactly the ideals generated by the minimal vertex covers of \mathcal{H}_I . Consequently, the *height* of I (resp., the *big height* of I), denoted by $ht(I)$ (resp., bight(I)), is the minimum (resp., maximum) cardinality of a minimal vertex cover of \mathcal{H}_I .

Let G be a graph on the vertex set $\{x_1, \ldots, x_n\}$. Consider the ideal $I_t(G)$ in the polynomial ring R. Since $I_t(G)$ is a square-free monomial ideal, from the previous discussion, we can associate a hypergraph, say $\mathcal{H}_t(G)$, on the vertex set $\{x_1, \ldots, x_n\}$ such that $I_t(G) = I(\mathcal{H}_t(G))$. More precisely,

- $V(\mathcal{H}_t(G)) = V(G),$
- $\bullet \: E(\mathcal{H}_t(G)) = \{\{x_{i_1},\ldots,x_{i_t}\} \subseteq V(G) \mid G[\{x_{i_1},\ldots,x_{i_t}\}] \text{ is connected }\}.$

2.4. **Some algebraic invariants:** Let *I* be a graded ideal in the polynomial ring $R = \mathbb{K}[x_1, \ldots, x_n].$ Then, a graded minimal free resolution of R/I is an exact sequence

$$
\mathcal{F} : 0 \to F_r \xrightarrow{\partial_r} \cdots \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} F_0 \xrightarrow{\partial_0} R/I \to 0,
$$

where F_0 = R, F_i = $\oplus_{j\in\mathbb{N}}R(-j)^{\beta_{i,j}(R/I)}$ for each $i\,\geq\,1,$ ∂_0 is the natural quotient map, and $R(-j)$ is the polynomial ring R with its grading twisted by j. The numbers $\beta_{i,j}(R/I)$ are uniquely determined, and called the i th N-graded *Betti numbers* of R/I in degree j. The *Castelnuovo-Mumford regularity* (or simply called the *regularity*) of R/I , denoted by $reg(R/I)$, is the number $max\{j - i \mid$ $\beta_{i,j}(R/I) \neq 0$. The invariant $\max\{i \mid \beta_{i,j}(R/I) \neq 0\}$ is called the *projective dimension* of R/I , and is denoted by $pd(R/I)$. Let I be a graded ideal generated in a single degree r. Then, we say that I has a *r*-linear resolution (or simply, a linear resolution) if $\text{reg}(R/I) = r - 1$.

The following are some well-known results regarding regularity and projective dimension, which we are going to use in the subsequent sections.

 ${\tt Lemma~2.1.}$ $[13]$ Let I_1 \subseteq R_1 $=$ $\mathbb{K}[x_1,\ldots,x_n]$ and I_2 \subseteq R_2 $=$ $\mathbb{K}[y_1,\ldots,y_m]$ be two graded ideals. *Consider the ideal* $I = I_1R + I_2R \subseteq R = \mathbb{K}[x_1,\ldots,x_n,y_1,\ldots,y_m]$ *. Then*

- (i) reg $(R/I) = \text{reg}(R_1/I_1) + \text{reg}(R_2/I_2)$ *,*
- *(ii)* $pd(R/I) = pd(R_1/I_1) + pd(R_2/I_2)$.

Lemma 2.2. [\[5,](#page-13-18) Lemma 2.10, Lemma 5.1] *Let* $I \subseteq R$ *be a square-free monomial ideal and let* x_i *be a variable appearing in some generator of* I*. Then*

 (i) reg $(R/I) \leq \max\{ \text{reg}(R/(I : x_i)) + 1, \text{reg}(R/(I, x_i)) \}.$ Moreover, $reg(R/I) \in \{ reg(R/(I : x_i)) + 1, reg(R/(I, x_i)) \}.$ (*ii*) $pd(R/I) \leq max{pd(R/(I : x_i)), pd(R/(I, x_i))}.$

Lemma 2.3. (cf. [\[20,](#page-13-19) Chapter 18]) *Let* J *and* K *be two graded ideals of* R*. Then*

 (i) reg($R/(J + K)$) < max{reg(R/J), reg(R/K), reg($R/(J \cap K)$) – 1}, *(ii)* $pd(R/(J + K)) \leq max{pd(R/J), pd(R/K), pd(R/(J \cap K)) + 1}.$

2.5. **Bounds on regularity and projective dimension:** Let H be a hypergraph. A *matching* in H is a collection of pairwise disjoint edges of H. More precisely, a subset $\mathcal{D} \subseteq E(\mathcal{H})$ is called a matching of H if for any two distinct edges $\mathcal{E}_1, \mathcal{E}_2 \in \mathcal{D}$, one has $\mathcal{E}_1 \cap \mathcal{E}_2 = \emptyset$. An *induced matching* is a matching D in H such that the edge set of the induced subhypergraph of H on the vertices of D is precisely the set D . The following lower bound on the regularity in terms of the induced matching is well-known.

Lemma 2.4. [\[11,](#page-13-20) Theorem 4.2] *Let* H *be a hypergraph, and* D *an induced matching of* H*. Then*

$$
reg(R/I(\mathcal{H})) \geq \sum_{\mathcal{E} \in \mathcal{D}}(|\mathcal{E}| - 1).
$$

Let us define $\nu(\mathcal{H}) = \max\{|\mathcal{D}| : \mathcal{D}$ is an induced matching of $\mathcal{H}\}$, and call this to be the *induced matching number* of the hypergraph H. Then for a simple graph $G, \nu(G)$ gives a crude lower bound of reg($R/I(G)$). For our purpose, given a simple graph G , we call an induced matching of the hypergraph $\mathcal{H}_t(G)$ to be a t-connected induced matching of G. Thus, we have $\nu(\mathcal{H}_t(G)) = \nu_t(G)$, where $\nu_t(G)$ is defined as in Section [2.1.](#page-2-1) Consequently, in our case, we have the following lower bound for the regularity of t -connected ideals of G .

Lemma 2.5. *Let* G *be a finite simple graph. Then* $reg(R/I_t(G)) \ge (t-1)\nu_t(G)$ *.*

Using the Alexander dual of square-free monomial ideals (see [\[13\]](#page-13-17)) and Terai's formula [\[24,](#page-13-21) Theorem 2.1], one can get an analogous bound for the projective dimension of $R/I_t(G)$ in terms of bight($I_t(G)$) as follows:

Lemma 2.6. *Let* G *be a finite simple graph. Then* $\text{pd}(R/I_t(G)) \geq \text{bight}(I_t(G)).$

Note: Let s be the maximum cardinality of the set of vertices in a connected component of a graph G. Then $I_t(G) = \langle 0 \rangle$ for all $t > s$. Thus, it is enough to focus on non-zero t-connected ideals, and sometimes we will assume this without mentioning it specifically.

3. CASTELNUOVO-MUMFORD REGULARITY AND LINEARITY

In this section, we compute the regularity of t -connected ideals of chordal graphs in terms of the t-connected induced matching number. As a consequence, we characterize when such an ideal has a linear resolution. Let us start with the following easy observation.

Proposition 3.1. *Let* $I \subseteq R$ *be a monomial ideal, and* $x_1, \ldots, x_r \in R$ *are some indeterminates. Then* $(I : x_r) + \langle x_1, \ldots, x_{r-1} \rangle = ((I + \langle x_1, \ldots, x_{r-1} \rangle) : x_r).$

Proposition 3.2. *Let* G *be a connected graph and* $x \in V(G)$ *be a simplicial vertex.* If $A \subseteq V(G)$ *such that* $x \in A$, $|A| = t \geq 2$, and $G[A]$ *is connected, then* $G[A \setminus \{x\}]$ *is also connected.*

Proof. If possible, let us assume that $G[A \setminus \{x\}]$ is a disconnected graph. Let H_1 and H_2 be any two connected components of $G[A \setminus \{x\}]$. Since $G[A]$ is connected, there are $y_1 \in V(H_1)$ and $y_2 \in$ $V(H_2)$ such that $\{x, y_1\}, \{x, y_2\} \in E(G[A])$. Then $\{y_1, y_2\} \in E(G[A])$, as x is a simplicial vertex. This is a contradiction to the fact the H_1 and H_2 are connected components in $G[A \setminus \{x\}]$.

Lemma 3.3. *Let* G *be a graph, and* $C \subseteq V(G)$ *such that* $|C| = t - 1$ *, and* $w \in N_G(C)$ *. Then* $\nu_t(G \setminus (N_G[C] \cup N_G[w])) \leq \nu_t(G) - 1.$

Proof. Let U be a t-connected induced matching of $G \setminus (N_G[C] \cup N_G[w])$ such that $|U| = \nu_t(G \setminus E)$ $(N_G[C] \cup N_G[w])$). Then the inequality follows from the fact that $U \cup \{C \cup \{w\}\}\$ is a t-connected induced matching of G .

The next lemma plays a crucial role in establishing the main results of this article. To describe the lemma precisely, we first introduce some notations as follows.

Notations: Let x be a vertex of the graph G, and $t \geq 2$ be an integer. Define the set $\mathcal{A}_x := \{C \subseteq$ $V(G):|C|=t-1, x \in C, G[C]$ is connected}. Without loss of generality, let $\mathcal{A}_x = \{C_1, \ldots, C_k\}.$ Then for $1 \leq i \leq k$, we define

 $\mathcal{B}_{C_i} := \{ w \in N_G(C_i) \mid C_i \cup \{w\} \neq C_j \cup \{w'\} \text{ for any } w' \in N_G(C_j), \text{ where } 1 \leq j \leq i-1 \}.$ By construction, $\mathcal{B}_{C_1} \neq \emptyset$ when $I_t(G) \neq \langle 0 \rangle$.

Lemma 3.4. Let G be a chordal graph and $x \in V(G)$ be a simplicial vertex. Let $\mathcal{A}_x = \{C_1, \ldots, C_k\}$. *For* $1 \leq i \leq k$ *, define*

$$
J_i := \mathbf{x}_{C_i} \langle w \mid w \in \mathcal{B}_{C_i} \rangle,
$$

$$
K_i := I(\mathcal{H}_t(G) \setminus (\bigcup_{j=1}^i C_j)).
$$

If $\mathcal{B}_{C_i}\neq\emptyset$, then

\n- (1)
$$
J_i + K_i = I(\mathcal{H}_t(G) \setminus (\bigcup_{j=1}^{i-1} C_j)),
$$
\n- (2) $J_i \cap K_i = \mathbf{x}_{C_i} L_i$, where for any $w \in \mathcal{B}_{C_i}$, we have $(L_i : w) = M_i + N_i + Q_i$, where $M_i = \langle v \mid v \in N_G(C_i) \setminus \{w\} \rangle$, $N_i = \langle v \mid v \in N_G(w) \setminus N_G[C_i] \rangle$, $Q_i = \langle \mathbf{x}_C \mid C \subseteq V(G), |C| = t, G[C]$ is connected, $C \cap (N_G[C_i] \cup N_G[w]) = \emptyset$.
\n

Proof. (1) Follows immediately from the construction of the ideals J_i and K_i .

(2) Fix any $1 \leq i \leq k$ and any $w \in \mathcal{B}_{C_i}$. We first show that $M_i + N_i + Q_i \subseteq (L_i : w)$. Let $v \in N_G(\dot{C}_i) \setminus \{w\}$. Then we can write $wv\mathbf{x}_{C_i} = \text{lcm}(w\mathbf{x}_{C_i}, wv\mathbf{x}_{C_i\setminus\{x\}})$. Since $w, v \in N_G(C_i)$, we see that $G[C_i \cup \{w, v\}]$ is a connected graph. Hence by Proposition [3.2,](#page-5-1) $G[(C_i \setminus \{x\}) \cup \{w, v\}]$ also connected. Also, note that $C_j \nsubseteq (C_i \setminus \{x\}) \cup \{w, v\}$ for any $1 \leq j \leq i$, as $x \in C_j$ for all such j. Hence we get $wv_{C_i\setminus\{x\}} \in K_i$. Moreover, from our choice of $w \in \mathcal{B}_{C_i}$, it is easy to see that $w\mathbf{x}_{C_i}\in J_i.$ Thus, we have $wv\mathbf{x}_{C_i}\in J_i\cap K_i,$ and therefore, $v\in (L_i:w).$ Now, let $v\in N_G(w)\setminus N_G[C_i].$ Then again, we can write $wv\mathbf{x}_{C_i}=\operatorname{lcm}(w\mathbf{x}_{C_i},wv\mathbf{x}_{C_i\setminus\{x\}}).$ By almost similar arguments as above, we obtain $w\mathbf{x}_{C_i}\in J_i$ and $wv\mathbf{x}_{C_i\setminus\{x\}}\in K_i$, and thus, $\widetilde{w\mathbf{x}_{C_i}}\in J_i\cap K_i$, which gives $v\in (L_i:w).$ Finally, let $\mathbf{x}_C\in Q_i,$ where $C\subseteq V(G)$ such that $|C|=t,$ $G[C]$ is connected, and $C \cap (N_G[C_i] \cup N_G[w]) = \emptyset.$ Then we write

$$
w\mathbf{x}_{C_i}\mathbf{x}_C = \text{lcm}(w\mathbf{x}_{C_i},\mathbf{x}_C),
$$

where $w\mathbf{x}_{C_i} \in J_i$ and $\mathbf{x}_C \in K_i$. Thus, $w\mathbf{x}_{C_i}\mathbf{x}_C \in J_i \cap K_i$ and hence $\mathbf{x}_C \in (L_i:w)$.

We now proceed to show that $(L_i:w) \subseteq M_i + N_i + Q_i$. Let $A \subseteq V(G)$ be such that $\mathbf{x}_A \in \mathcal{G}(K_i)$. We consider the following two cases:

Case-I. Let $A \cap N_G[C_i] = \emptyset$. Then $w \notin A$ and hence $\text{lcm}(w\mathbf{x}_{C_i}, \mathbf{x}_A) = w\mathbf{x}_{C_i}\mathbf{x}_A$. Now if $A \cap$ $N_G[w]\neq \emptyset$, then $A\cap N_G[w]\subseteq N_G(w)\setminus N_G[C_i],$ since $w\notin A.$ Also if $A\cap N_G[w]=\emptyset,$ then clearly $A\cap (N_G[C_i]\cup N_G[w])=\emptyset.$ Thus, in any case, we have

$$
(\operatorname{lcm}(w\mathbf{x}_{C_i}, \mathbf{x}_A):w\mathbf{x}_{C_i}) = \mathbf{x}_A \in N_i + Q_i.
$$

Case-II. Let $A \cap N_G[C_i] \neq \emptyset$. Note that, since $C_i \nsubseteq A, A \nsubseteq C_i,$ and $G[C_i \cup A]$ is connected, there exists some $a \in A \setminus C_i$ and $b \in C_i \setminus A$ such that $\{a, b\} \in E(G)$. Thus $A \cap N_G(C_i) \neq \emptyset$.

First, consider the case when $A \cap N_G(C_i) = \{w\}$. Recall that $|C_i| = t - 1$, and $|A| = t$. Since $C_i \nsubseteq A,$ there exists some $v \in A \setminus C_i$ such that $v \neq w$ and $v \notin N_G[C_i].$ Since $G[A]$ is connected, there exists a shortest path $v = y_0, y_1, \ldots, y_r = w$ such that $y_i \in A$ for all $0 \le i \le r$. Note that $y_{r-1}\,\in\, N_G(w),$ and $v\notin\, N_G[C_i].$ So if $y_{r-1}\,\in\, C_i,$ then there exists $1\,\leq\, i\,\leq\, r-2$ such that $y_i\in A\cap N_G(C_i),$ which is a contradiction to the fact that $A\cap N_G(C_i)=\{w\}.$ Thus $y_{r-1}\notin N_G[C_i]$ and hence $y_{r-1}\in N_G(w)\setminus N_G[C_i].$ Therefore, in this case $(\mathrm{lcm}(w\mathbf{x}_{C_i},\mathbf{x}_A):w\mathbf{x}_{C_i})\in N_i.$ Finally, assume that there exists $w' \in A \cap N_G(C_i)$, where $w' \neq w$. Then $(\text{lcm}(w\mathbf{x}_{C_i}, \mathbf{x}_A): w\mathbf{x}_{C_i}) \in$ $M_i.$ Therefore, $(L_i:w)=M_i+N_i+Q_i$.

In the following example, we illustrate some of the notations used in Lemma [3.4](#page-5-2) with the aid of Figure [1.](#page-6-0)

FIGURE 1. A chordal graph G.

Example 3.5. Let G be the chordal graph as in Figure [1](#page-6-0) and $x₅$ a simplicial vertex in G. Consider the 4-connected ideal $I(\mathcal{H}_4(G))$ inside the polynomial ring $R = \mathbb{K}[x_1, \ldots, x_{14}]$. In this case observe that $\mathcal{A}_{x_5} = \{C_1, C_2, \ldots, C_9\}$, where $C_1 = \{x_3, x_4, x_5\}$, $C_2 = \{x_3, x_5, x_6\}$, $C_3 =$ ${x_4, x_5, x_6}, C_4 = {x_2, x_4, x_5}, C_5 = {x_1, x_4, x_5}, C_6 = {x_2, x_3, x_5}, C_7 = {x_1, x_3, x_5}, C_8 =$ ${x_5, x_6, x_7}$, and $C_9 = {x_5, x_6, x_8}$. It is easy to see that $\mathcal{B}_{C_1} = {x_1, x_2, x_6}$ and thus $J_1 =$ $\langle x_1x_3x_4x_5, x_2x_3x_4x_5, x_3x_4x_5x_6\rangle$. Note that, $x_4\notin \mathcal{B}_{C_2}$ since $\{x_4\}\cup C_2=\{x_6\}\cup C_1$. Consequently, $\mathcal{B}_{C_2} = \{x_1, x_2, x_7, x_8\}$ and thus $J_2 = \langle x_1x_3x_5x_6, x_2x_3x_5x_6, x_3x_5x_6x_7, x_3x_5x_6x_8\rangle$. Similarly, one can determine the sets \mathcal{B}_{C_i} and the ideals $J_{C_i},$ for $3\leq i\leq 9.$

We are now ready to prove the main theorem of this section.

Theorem 3.6. *Let* G *be a chordal graph. Then for any* $t \geq 2$ *,*

$$
reg(R/I_t(G)) = (t-1)\nu_t(G).
$$

Proof. In view of Lemma [2.4,](#page-4-0) it is enough to prove that $reg(R/I_t(G)) \leq (t-1)\nu_t(G)$. We prove this using induction on $|V(G)|.$ First and foremost, if $|V(G)|\leq t,$ then either $I_t(G)=\langle \prod_{x\in V(G)} x\rangle$ or $I_t(G) = \langle 0 \rangle$. In either case, it is easy to see that reg $(R/I_t(G)) = (t - 1)\nu_t(G)$. Therefore, we may assume that $|V(G)| \geq t + 1$. Also, we can assume that $\nu_t(G) \geq 1$. Moreover, we will write $I_t(G) = I(\mathcal{H}_t(G))$, where $\mathcal{H}_t(G)$ is the hypergraph corresponding to the t-connected ideal of G. Now, let $x \in V(G)$ be a simplicial vertex of G, and $A_x = \{C_1, \ldots, C_k\}$. Following Lemma [3.4,](#page-5-2) whenever $\mathcal{B}_{C_i}\neq\emptyset$ for some $i\in\{1,\ldots,k\},$ we denote

$$
J_i = \mathbf{x}_{C_i} \langle w \mid w \in \mathcal{B}_{C_i} \rangle,
$$

\n
$$
K_i = I(\mathcal{H}_t(G) \setminus (\cup_{j=1}^i C_j)),
$$

\n
$$
J_i \cap K_i = \mathbf{x}_{C_i} L_i.
$$

Claim: $reg(R/L_i) \leq (t-1)\nu_t(G) - (t-2)$ for each $1 \leq i \leq k$. *Proof of the claim.* Let $\mathcal{B}_{C_i} = \{w_1, \ldots, w_s\}$. Then by Lemma [3.4,](#page-5-2) $L_i + \langle w_1, \ldots, w_s \rangle = \langle w_1, \ldots, w_s \rangle$. Thus,

$$
reg(R/(L_i + \langle w_1, \ldots, w_s \rangle)) = 0 \le (t - 1)\nu_t(G) - (t - 2).
$$

Now by Lemma [3.4,](#page-5-2)

$$
((L_i + \langle w_1, \dots, w_{s-1} \rangle) : w_s) = (L_i : w_s) + \langle w_1, \dots, w_{s-1} \rangle
$$

\n
$$
= \langle v \mid v \in N_G(C_i) \setminus \{w_s\} \rangle + \langle v \mid v \in N_G(w_s) \setminus N_G[C_i] \rangle
$$

\n
$$
+ \langle \mathbf{x}_C \mid C \subseteq V(G), |C| = t, G[C] \text{ is connected },
$$

\n
$$
C \cap (N_G[C_i] \cup N_G[w_s]) = \emptyset \rangle
$$

\n
$$
= \langle v \mid v \in N_G(C_i) \setminus \{w_s\} \rangle + \langle v \mid v \in N_G(w_s) \setminus N_G[C_i] \rangle
$$

\n
$$
+ I_t(G \setminus (N_G[C_i] \cup N_G[w_s])).
$$

Then

$$
reg(R/((L_i + \langle w_1, \dots, w_{s-1} \rangle) : w_s)) = reg(R/I_t(G \setminus (N_G[C_i] \cup N_G[w_s])))
$$

$$
\leq (t-1)\nu_t(G \setminus (N_G[C_i] \cup N_G[w_s]))
$$

$$
\leq (t-1)(\nu_t(G) - 1),
$$

where the first inequality is by the induction hypothesis and the second inequality follows from Lemma [3.3.](#page-5-3) Hence, by Lemma [2.2,](#page-4-1) we have $reg(R/(L_i + \langle w_1, \ldots, w_{s-1} \rangle) \le (t-1)\nu_t(G) - (t-2)$. Now for each $2 \le j \le s - 1$, similarly using Lemma [3.4,](#page-5-2) we have

$$
((L_i + \langle w_1, \dots, w_{s-j})): w_{s-j+1}) = (L_i : w_{s-j+1}) + \langle w_1, \dots, w_{s-j} \rangle
$$

\n
$$
= \langle v \mid v \in N_G(C_i) \setminus \{w_{s-j+1}\}\rangle + \langle v \mid v \in N_G(w_{s-j+1}) \setminus N_G[C_i] \rangle
$$

\n
$$
+ \langle \mathbf{x}_C \mid C \subseteq V(G), |C| = t, G[C] \text{ is connected},
$$

\n
$$
C \cap (N_G[C_i] \cup N_G[w_{s-j+1}]) = \emptyset \rangle
$$

\n
$$
= \langle v \mid v \in N_G(C_i) \setminus \{w_{s-j+1}\}\rangle + \langle v \mid v \in N_G(w_{s-j+1}) \setminus N_G[C_i] \rangle
$$

\n
$$
+ I_t(G \setminus (N_G[C_i] \cup N_G[w_{s-j+1}])).
$$

Thus for each $2 \leq j \leq s - 1$, we get

reg
$$
(R/((L_i + \langle w_1, \ldots, w_{s-j})) : w_{s-j+1})) = \text{reg}(R/I_t(G \setminus (N_G[C_i] \cup N_G[w_{s-j+1}])))
$$

 $\leq (t-1)\nu_t(G \setminus (N_G[C_i] \cup N_G[w_{s-j+1}]))$
 $\leq (t-1)(\nu_t(G) - 1),$

Therefore, repeatedly applying Lemma [2.2,](#page-4-1) we obtain $reg(R/L_i) \leq (t-1)(\nu_t(G)-1) - 1 =$ $(t-1)\nu_t(G) - (t-2)$. This completes the proof of the claim.

Now, consider the ideal $K_k = I(\mathcal{H}_t(G)\backslash \bigcup_{j=1}^k C_j) = I(\mathcal{H}_t(G\backslash x)) = I_t(G\backslash x).$ By the induction hypothesis, reg $(R/K_k) \le (t-1)\nu_t(G \setminus x) \le (t-1)\nu_t(G)$. Also, reg $(R/J_k) = t-1 \le (t-1)\nu_t(G)$. Moreover, $reg(R/(J_k \cap K_k)) = reg(R/\mathbf{x}_{C_i}L_k) \leq (t-1)+(t-1)\nu_t(G)-(t-2) = (t-1)\nu_t(G)+1.$ Hence, by Lemma [2.3,](#page-4-2) ${\rm reg}(R/(J_k+K_k))\leq (t-1)\nu_t(G).$ Note that $J_k+K_k=I(\mathcal{H}\setminus \bigcup_{j=1}^{k-1} C_j)=0$ $K_{k-1}.$ We now write $I(\mathcal{H}\setminus \bigcup_{j=1}^{k-2} C_j)=J_{k-1}+K_{k-1}$ and continue the above process. Note that if for some $1 \le i \le k$, $\mathcal{B}_{C_i} = \emptyset$, then $K_i = K_{i-1}$. Hence, after a finite number of steps, we obtain

$$
reg(R/I_t(G)) = reg(R/I(\mathcal{H}_t(G))) = reg(R/(J_1 + K_1) \le (t - 1)\nu_t(G),
$$

and this completes the proof.

As an application of the above theorem, we get a complete classification of chordal graphs whose t-connected ideals have linear resolutions as follows.

Corollary 3.7. *Let* G *be a chordal graph and* $t \geq 2$ *be an integer. Then* $I_t(G)$ *has a linear resolution if and only if G is* t -gap-free (i.e., $\nu_t(G) = 1$).

The above result can be considered as an extension of [\[2,](#page-12-1) Theorem 5.1], where they have shown that for a tree T and for $t \geq 2$, $I_t(T)$ has a linear resolution if and only if T is t-gap-free.

Example 3.8. Let us consider the graph G as shown in Figure [1.](#page-6-0) Then one can deduce that

$$
\nu_t(G) = \begin{cases}\n4 & \text{for } t = 2, \\
3 & \text{for } t = 3, \\
2 & \text{for } t = 4, 5, 6, \\
1 & \text{for } t = 7, ..., 14, \\
0 & \text{for } t > 14.\n\end{cases}
$$

Therefore, using Theorem [3.6,](#page-7-0) we can derive $reg(R/I_t(G))$ for all $t \geq 2$. Note that $I_t(G) = \langle 0 \rangle$ for all $t > 14$. If $I_t(G) \neq \langle 0 \rangle$, then due to Corollary [3.7,](#page-8-0) $I_t(G)$ has a linear resolution if and only if $t = 7, \ldots, 14.$

4. PROJECTIVE DIMENSION AND COHEN-MACAULAY PROPERTY

In this section, we compute the projective dimension of the t-connected ideal of a chordal graph in terms of the big height of the corresponding ideal. As a corollary, we classify when a t -connected ideal of a chordal graph is Cohen-Macaulay.

Proposition 4.1. *Let* G *be a disjoint union of two graphs* G_1 *and* G_2 *, i.e.,* $G = G_1 \sqcup G_2$ *. Then* $bight(I_t(G)) = bight(I_t(G_1)) + bight(I_t(G_2)).$

Proof. Since $G = G_1 \sqcup G_2$, we have $\mathcal{G}(I_t(G)) = \mathcal{G}(I_t(G_1)) \sqcup \mathcal{G}(I_t(G_2))$. Thus, p is a minimal prime ideal of $I_t(G)$ if and only if $\mathfrak{p} = \mathfrak{p}_1 + \mathfrak{p}_2$, where \mathfrak{p}_1 and \mathfrak{p}_2 are minimal prime ideals of $I_t(G_1)$ and $I_t(G_2)$, respectively. Hence, the result follows.

Proposition 4.2. Let H be an induced subgraph of a graph G. Then $\text{bight}(I_t(H)) \leq \text{bight}(I_t(G))$ *.*

Proof. Since H is an induced subgraph of G, any t-connected set in H is also a t-connected set in G . Thus, we have $I_t(H) \subseteq I_t(G)$. Let p be an associated prime of $I_t(H)$ such that $\text{ht}(\mathfrak{p}) = \text{bight}(I_t(H))$. Now, let us consider the prime ideal $\mathfrak{q} = \mathfrak{p} + \langle V(G) \setminus V(H) \rangle$. Note that q is a prime ideal containing $I_t(G)$. Then there exists a minimal prime ideal \mathfrak{p}' of $I_t(G)$ such that $\mathfrak{p}' \subseteq \mathfrak{q}$. It is easy to observe from the structure of $\frak q$ that $\frak p\subseteq\frak p'$ as $\frak p$ is a minimal prime ideal containing $I_t(H).$ Therefore, $\text{bight}(I_t(H)) \leq \text{ht}(\mathfrak{p}') \leq \text{bight}(I_t(G)).$

Proposition 4.3. *Let* $C \subseteq V(G)$ *be such that* $|C| = t - 1$ *with* $t \geq 2$ *and* $G[C]$ *is connected. Let J denote the ideal* $\langle wx_C | w \in N_G(C) \rangle$ *. Then* $\text{pd}(R/J) \leq \text{bight}(I_t(G))$ *.*

Proof. It is easy to see that $pd(R/J) = pd(R/(N_G(C)))$, and the Koszul complex is the minimal free resolution of $R/N_G(C)$. Thus, $pd(R/J) = |N_G(C)|$. To show bight $(I_t(G)) \geq |N_G(C)|$, note that $\mathfrak{p} = \langle V(G) \setminus C \rangle$ is a prime ideal containing $I_t(G)$. Then there exists a minimal prime ideal q of $I_t(G)$ such that $\mathfrak{q} \subseteq \mathfrak{p}$. Since $J \subseteq I_t(G) \subseteq \mathfrak{q}$ and $C \cap \mathcal{G}(\mathfrak{q}) = \emptyset$, we must have $\langle N_G(C) \rangle \subseteq \mathfrak{q}$. Hence, bight $(I_t(G)) \geq ht(q) \geq |N_G(C)|$ as desired.

The following lemma on the big height of $I_t(G)$ is important in the proof of the main theorem of this section.

Lemma 4.4. *Let* x *be a simplicial vertex of a graph* G and $C \subseteq V(G)$ *be such that* $x \in C$, $|C| = t - 1$ *for some integer* $t \geq 2$ *, and* $G[C]$ *is connected. Then for each* $y \in N_G(C)$ *, we have*

 $bight(I_t(G)) \geq |N_G(C)| + |N_G(y) \setminus N_G[C]| + bight(I_t(G \setminus (N_G[C] \cup N_G[y])).$

Proof. First consider the case when $N_G(y) \subseteq N_G[C]$. In this case, if p is a minimal prime ideal of $I_t(G \setminus (N_G[C] \cup N_G[y]))$ such that $\text{ht}(\mathfrak{p}) = \text{bight}(I_t(G \setminus (N_G[C] \cup N_G[y]))$, then one can verify that $\mathfrak{q} = \mathfrak{p} + \langle N_G(C) \rangle$ is a prime ideal containing $I_t(G)$ as $|C| = t-1$. Now, if \mathfrak{q}' is a minimal prime ideal of $I_t(G)$ with $\mathfrak{q}'\subseteq \mathfrak{q}$, then we must have $\mathfrak{p}\subseteq \mathfrak{q}'$. Also, note that $\langle N_G(C)\rangle\subseteq \mathfrak{q}'$ as $y'\mathbf{x}_C\in I_t(G)$ for all $y' \in N_G(C)$. Thus, we have $\mathfrak{q}' = \mathfrak{q}$, and consequently,

$$
bight(I_t(G)) \ge ht(\mathfrak{q}') = |N_G(C)| + ht(\mathfrak{p})
$$

= |N_G(C)| + |N_G(y) \setminus N_G[C]| + bight(I_t(G \setminus (N_G[C] \cup N_G[y])),

where $|N_G(y)\setminus N_G[C]|=0$ in this case. Now, suppose $N_G(y)\nsubseteq N_G[C]$. Similar to the first case, if p is a minimal prime ideal of $I_t(G\setminus (N_G[C\cup N_G[y]))$ such that ht(p) = bight($I_t(G\setminus (N_G[C\cup N_G[y]))$), then

$$
\mathfrak{q} = \mathfrak{p} + \langle x \rangle + \langle N_G(C) \setminus \{y\} \rangle + \langle N_G(y) \setminus N_G[C] \rangle
$$

is a prime ideal containing $I_t(G)$. Let $\mathfrak q'$ be a minimal prime ideal of $I_t(G)$ such that $\mathfrak q' \subseteq \mathfrak q$. Then, it is easy to see that $\frak{p}\subseteq\frak{q}'$. Note that for each $y'\in N_G(C)$ with $y'\neq y$, we have $yy'\mathbf{x}_{C\setminus\{x\}}\in I_t(G)$ by Proposition [3.2.](#page-5-1) Hence, $\langle N_G(C) \setminus \{y\} \rangle \subseteq \mathfrak{q}'$ as $((C \setminus \{x\}) \cup \{y\}) \cap \mathcal{G}(\mathfrak{q}') = \emptyset$. Again, we have $x \in \mathfrak{q}'$ as $y\mathbf{x}_C \in I_t(G)$. Observe that for each $w \in N_G(y) \setminus N_G[C]$, $G[C \cup \{y,w\}]$ is connected, and thus, $G[C \setminus \{x\} \cup \{y,w\}]$ is connected by Proposition [3.2.](#page-5-1) In particular, $yw\mathbf{x}_{C\setminus\{x\}} \in I_t(G)$ for each $w\in N_G(y)\setminus N_G[C]$, which implies $\langle N_G(y)\setminus N_G[C]\rangle\subseteq \mathfrak{q}'$ as $((C\setminus\{x\})\cup\{\hat{y\}})\cap\mathcal{G}(\mathfrak{q}')=\emptyset.$ Hence, $q' = q$, and consequently,

$$
bight(I_t(G)) \geq ht(\mathfrak{q}') = |N_G(C)| + |N_G(y) \setminus N_G[C]| + bight(I_t(G \setminus (N_G[C] \cup N_G[y])).
$$

This completes the proof.

We now proceed to prove the main theorem of this section.

Theorem 4.5. Let G be a chordal graph. Then for all $t \geq 2$, $pd(R/I_t(G)) = bight(I_t(G))$.

Proof. Due to Lemma [2.6,](#page-5-4) it is enough to show that $\text{pd}(R/I_t(G)) \leq \text{bight}(I_t(G))$. We proceed by induction on $|V(G)|$. The statement is trivially true for $|V(G)| < t$. If $|V(G)| = t$, then either $I_t(G) = \langle 0 \rangle$ or $I_t(G) = \langle \mathbf{x}_{V(G)} \rangle$ depending on whether G is connected or not. In both cases, it is easy to see that $\text{pd}(R/I_t(G)) = \text{bight}(I_t(G))$. Therefore, we may assume that $|V(G)| > t$. Note that, in case G is disconnected, and G_1, \ldots, G_r are all connected components of G such that $|V(G_i)| < t$ for all $i \in [r]$, then again $I_t(G) = \langle 0 \rangle$, and thus $\text{pd}(R/I_t(G)) = \text{bight}(I_t(G))$. Therefore, we may further assume that G has at least one connected component with at least t many vertices. Now, let us consider the following two cases:

Case-I. For each simplicial vertex x of G, if $C \in \mathcal{A}_x$, then $V(G) \neq N_G[C]$. Note that since G has at least one connected component with at least t many vertices, we can find a simplicial vertex x of G and some $C \in \mathcal{A}_x$ such that $C \subsetneq N_G[C]$. Based on this observation, we consider two subcases:

$$
\Box
$$

Subcase-I(A). There exists a simplicial vertex x of G and $C \in \mathcal{A}_x$ such that $G[N_G[C]]$ forms a connected component of G . Then we have

$$
I_t(G) = I_t(G[N_G[C]]) + I_t(G \setminus N_G[C]).
$$

From our assumption in Case-I, it follows that $1 \leq |N_G[C]| < |V(G)|$ and $1 \leq |V(G \setminus N_G[C])| <$ $|V(G)|$. Therefore, by the induction hypothesis, Lemma [2.1](#page-4-3) and Proposition [4.1,](#page-8-2) we have

$$
pd(R/I_t(G)) = pd(R/I_t(G[N_G[C]])) + pd(R/I_t(G \setminus N_G[C]))
$$

\n
$$
\leq bight(I_t(G[N_G[C]])) + bight(I_t(G \setminus N_G[C]))
$$

\n
$$
= bight(I_t(G)).
$$

Subcase-I(B). For each simplicial vertex x of G and each $C \in \mathcal{A}_x$, $G[N_G|C]$ does not form a connected component of G. In particular, $C \subsetneq N_G[C]$ for any such C. Recall that $I_t(G) = I(\mathcal{H}_t(G))$. Now, fix a simplicial vertex x of G. Let $A_x = \{C_1, \ldots, C_k\}$. For $1 \le i \le k$, we define

$$
J_i = \langle \mathbf{x}_{C_i} w \mid w \in \mathcal{B}_{C_i} \rangle \text{ and } K_i = I(\mathcal{H}_t(G) \setminus (\bigcup_{j=1}^i C_j)).
$$

Note that by construction, $\mathcal{B}_{C_1}\neq\emptyset$ as $C_1\subsetneq N_G[C_1].$ Fix some $i\in\{1,\ldots,k\}$ such that $\mathcal{B}_{C_i}\neq\phi.$ Then we are in the situation of Lemma [3.4,](#page-5-2) and thus, $J_i+K_i=I(\mathcal{H}_t(G)\setminus (\bigcup_{j=1}^{i-1} C_j)).$ Now, we proceed to prove the following claim:

Claim 1: $pd(R/J_i \cap K_i) \leq bight(I_t(G)) - 1$.

Proof of the Claim 1. We have $J_i \cap K_i = \mathbf{x}_{C_i} L_i$, where L_i is generated by the monomials $\frac{\text{lcm}(m,m')}{\mathbf{x}_{C_i}}$, where $m \in J_i$ and $m' \in K_i$. Thus, it is enough to prove that $\text{pd}(R/L_i) \leq \text{bight}(I_t(G)) - 1$. Let $\mathcal{B}_{C_i} = \{w_1, \ldots, w_s\}$. Then by Lemma [3.4,](#page-5-2) $L_i + \langle w_1, \ldots, w_s \rangle = \langle w_1, \ldots, w_s \rangle$, and hence, ${\rm pd}(R/L_i+\langle w_1, \cdots, w_s\rangle)=s.$ Note that ${\cal B}_{C_i}\,\subseteq\, N_G(C_i).$ Moreover, since $G[N_G[C_i]]$ does not form a connected component of $G,$ there exists some $y\in N_G(C_i)$ such that $N_G(y)\setminus N_G[C_i]\neq\emptyset.$ In this case, by Lemma [4.4,](#page-9-1) we have $\mathrm{bight}(I_t(G))\geq |N_G(C_i)|+|N_G(y)\setminus N_G[C_i]|\geq s+1.$ Thus,

$$
\mathrm{pd}(R/L_i + \langle w_1, \cdots, w_s \rangle) \le \mathrm{bight}(I_t(G)) - 1. \tag{1}
$$

Now by Lemma [3.4,](#page-5-2) $(L_i:w_j)=\langle N_G(C_i)\setminus\{w_j\}\rangle+\langle N_G(w_j)\setminus N_G[C_i]\rangle+I_t(G\setminus(N_G[C_i]\cup N_G(w_j)))$ for each $j \in [s]$. Hence, using Lemma [4.4](#page-9-1) and the induction hypothesis, we obtain

$$
\mathrm{pd}(R/(L_i:w_j))\leq \mathrm{bight}(I_t(G))-1,\tag{2}
$$

for each $j \in [s].$ Observe that $((L_i + \langle w_1, \ldots, w_{s-1}\rangle) : w_s) = (L_i : w_s)$ by Proposition [3.1](#page-5-5) and Lemma [3.4.](#page-5-2) Thus, using Equation [\(2\)](#page-10-0), we get $pd(R/((L_i+\langle w_1, \ldots, w_{s-1}\rangle) : w_s)) \leq \text{bight}(I_t(G))$ 1. Consequently, by Lemma [2.2](#page-4-1) and the Equation [\(1\)](#page-10-1), we obtain $\text{pd}(R/L_i + \langle w_1, \ldots, w_{s-1} \rangle \leq$ bight($I_t(G)$)−1. In view of Proposition [3.1](#page-5-5) and Lemma [3.4,](#page-5-2) we again have $((L_i+\langle w_1, \ldots, w_{s-2}\rangle)$: $w_{s-1}) = (L_i : w_{s-1}).$ Thus, proceeding similarly as before and using Lemma [2.2](#page-4-1) repeatedly, we finally obtain $\text{pd}(R/L_i) \leq \text{bight}(I_t(G)) - 1$, and this completes the proof of the above claim.

It is easy to observe that $I_t(G\setminus x)=I(\mathcal{H}_t(G)\setminus \cup_{j=1}^k C_j)=K_k.$ Since $G\setminus x$ is an induced sub-graph of G, by Proposition [4.2](#page-8-3) and the induction hypothesis, we have $pd(R/K_k) \leq bid(t_t(G))$. Moreover, ${\rm pd}(R/J_k)={\rm pd}(R/\langle\mathcal{B}_{C_k}\rangle)=|\mathcal{B}_{C_k}|\leq |N_G(C_k)|$, and proceeding as in the proof of Proposition [4.3,](#page-8-4) we have $\text{pd}(R/J_k) \leq \text{bight}(I_t(G))$. Thus, using the Claim 1 and by Lemma [2.3,](#page-4-2) we obtain ${\rm pd}(R/J_k+K_k)\leq {\rm bight}(I_t(G)),$ where $J_k+K_k=I(\mathcal{H}_t(G)\setminus\cup_{j=1}^{k-1} C_j)=K_{k-1}$ by Lemma [3.4.](#page-5-2) Next, using Lemma [3.4,](#page-5-2) we write $K_{k-2} = J_{k-1} + K_{k-1}$ and continue the above process. Note that $\mathcal{B}_{C_1}\neq\emptyset$, and if for some $i\in\{2,\ldots,k\},$ $\mathcal{B}_{C_i}=\emptyset$, then $K_i=K_{i-1}.$ Thus using Lemma [3.4](#page-5-2) and Lemma [2.3](#page-4-2) repeatedly, we get $\text{pd}(R/J_i + K_i) \leq \text{bight}(I_t(G))$ for each $i \in [k]$. In particular, $\text{pd}(R/I_t(G)) = \text{pd}(R/J_1 + K_1) \leq \text{bight}(I_t(G))$, as desired.

Case-II. There exists a simplicial vertex x of G, and some $C_1 \in \mathcal{A}_x$ such that $V(G) = N_G[C_1]$. In particular, G is a connected graph. As before, since G has at least one connected component with at least t many vertices, we may as well assume that $C_1 \subsetneq N_G[C_1]$. Without loss of generality, let

 $\mathcal{A}_x = \{C_1, \ldots, C_k\}$, where $N_G[C_i] = V(G)$ for $1 \leq i \leq l$, and $N_G[C_i] \subsetneq V(G)$ for $l+1 \leq i \leq k$. As before, define $J_i = \langle \mathbf{x}_{C_i} w \mid w \in \mathcal{B}_{C_i} \rangle$ and $K_i = I(\mathcal{H}_t(G) \setminus (\bigcup_{j=1}^i C_j))$ for $i \in [k].$ Fix some $i\in [k]$ such that $\mathcal{B}_{C_i}\neq\emptyset.$ First, we consider the case when $l+1\leq i\leq k.$ In this case, since G is connected and $N_G[C_i]\subsetneq V(G),$ there exists some $y\in N_G(C_i)$ such that $N_G(y)\setminus N_G[C_i]\neq\emptyset.$ In particular, $G[N_{G}[C_{i}]]$ does not form a connected component of $G.$ Thus proceeding as in Subcase-I(B), we obtain $\text{pd}(R/J_i \cap K_i) \leq \text{bight}(I_t(G)) - 1$. Now consider the case when $i \in [l]$. Let $\mathcal{B}_{C_i} = \{w_1, \ldots, w_r\}$ and $N_G(C_i) = \{w_1, \ldots, w_r, w_{r+1}, \ldots, w_s\}$ for some $s \geq r$. Then $J_i =$ $\mathbf{x}_{C_i}\langle w_1, \ldots, w_r \rangle.$

Claim 2: For each $i \in [l]$, if $\mathcal{B}_{C_i} \neq \emptyset$, then $\text{pd}(R/J_i \cap K_i) \leq \text{bight}(I_t(G)) - 1$.

Proof of the Claim 2. We first aim to show that

$$
J_i \cap K_i = \mathbf{x}_{C_i} \langle w_m w_n \mid 1 \leq m < n \leq r \rangle + \mathbf{x}_{C_i} \langle w_m w_n \mid m \in [r], r+1 \leq n \leq s \rangle. \tag{3}
$$

Indeed, by Proposition [3.2,](#page-5-1) for each $1 \leq m \leq n \leq r$, $G[(C_i \setminus \{x\}) \cup \{w_m, w_n\}]$ is a connected subgraph of G. Thus $\mathbf{x}_{C_i}w_mw_n = \text{lcm}(w_m\mathbf{x}_{C_i},w_mw_n\mathbf{x}_{C_i\setminus\{x\}})$, where $w_m\mathbf{x}_{C_i}\in J_i$, and $w_mw_n\mathbf{x}_{C_i\setminus\{x\}}\in K_i.$ Similarly, if $m\in [r]$ and $r+1\leq n\leq s,$ then by Proposition [3.2,](#page-5-1) $G[(C_i\setminus$ ${x}$) ∪ ${w_m, w_n}$] is also connected. Thus, if M denotes the right-hand side of Equation [\(3\)](#page-11-2), then $M\subseteq J_i\cap K_i.$ Conversely, if $A\subseteq V(G)$ such that $C_i\nsubseteq A, |A|\,=\,t,$ and $G[A]$ is connected, then $\{w_m, w_n\} \subseteq A$ for some $w_m, w_n \in N_G(C_i)$. Thus $J_i \cap K_i \subseteq M$ and consequently, $\text{pd}(R/J_i \cap K_i) = \text{pd}(R/I(G'))$, where $I(G')$ is the edge ideal of the graph G' with $V(G') = N_G(C_i)$ and $E(G') = \{\{w_m, w_n\}, \{w_p, w_q\} \mid 1 \leq m < n \leq r, p \in [r], r+1 \leq q \leq s\}.$ It is easy to see that the complement of G' is a disconnected graph, and thus, using [\[15,](#page-13-22) Theorem 4.2.6] we have $\text{pd}(R/J_i \cap K_i) = |N_G(C_i)|-1$. This completes the proof of Claim 2 since $\text{bight}(I_t(G)) \geq |N_G(C_i)|$, by Proposition [4.3.](#page-8-4)

Thus, for each $i\in[k]$ we observe that if $\mathcal{B}_{C_i}\neq\emptyset$, then ${\rm pd}(R/J_i\cap K_i)\leq {\rm bight}(I_t(G))-1.$ Hence, we are in the same situation as in Subcase-I(B). Proceeding as before, we see that $\text{pd}(R/J_i + K_i) \leq$ bight($I_t(G)$) for each $i \in [k]$, and in particular, $\text{pd}(R/I_t(G)) = \text{pd}(R/J_1 + K_1) \leq \text{bight}(I_t(G)).$ This completes the proof of the Theorem.

Example 4.6. Let G be the graph as in Figure [1.](#page-6-0) Observe that $\{x_4, x_5, x_6, x_7, x_8, x_{12}, x_{13}, x_{14}\}$, is a vertex cover of $\mathcal{H}_4(G)$ with maximum possible cardinality. Thus by Theorem [4.5,](#page-9-0) $pd(R/I_4(G)) =$ 8.

Remark 4.7. In this context, one should note that if $I(\mathcal{H})$ is a sequentially Cohen-Macaulay edge ideal of a hypergraph H, then $pd(R/I(H)) = bight(I(H))$ [\[18,](#page-13-23) Corollary 3.33]. Also, it is wellknown that if G is chordal, then $I(G)$ is sequentially Cohen-Macaulay. Now, due to these facts and Theorem [4.5,](#page-9-0) one can ask whether, for a chordal graph G and $t \geq 3$, $I_t(G)$ is sequentially Cohen-Macaulay or not. However, the answer to this question is negative (see [\[1,](#page-12-0) Proposition 4.3]). Moreover, a natural question arises from this discussion is the following: if $I(G)$ is sequentially Cohen-Macaulay, then do we have $pd(R/I_t(G)) = bight(I_t(G))$ for all $t \geq 3$? Although this happens in the case of chordal graphs, this question has a negative answer in general. For example, if we consider the cycle C_5 of length 5, then $I(C_5)$ is sequentially Cohen-Macaulay but $pd(R/I₃(C₅)) = 3 > 2 = bight(I₃(C₅)).$

Next, as a corollary of Theorem [4.5,](#page-9-0) we generalize a famous result of Herzog-Hibi-Zheng [\[14\]](#page-13-13), where they combinatorially classified all Cohen-Macaulay chordal graphs.

Corollary 4.8. *Let* G *be a chordal graph and* $t \geq 2$ *be an integer. Then* $I_t(G)$ *is Cohen-Macaulay if and only if* $I_t(G)$ *is unmixed.*

Proof. The proof follows from Theorem [4.5](#page-9-0) and the Auslander-Buchsbaum formula. □

5. CONCLUDING REMARKS

In this section, by *hypergraphs* $H_t(G)$ *induced from a graph* G, we mean a class of *t*-uniform hypergraphs for which $H_2(G) = G$. In other words, the edge ideal $I(\mathcal{H}_t(G))$ can be viewed as a

higher degree generalization of $I(G)$. In this article, we have shown in Theorem [3.6](#page-7-0) and [4.5](#page-9-0) that if $I(H_t(G))$ corresponds to the t-connected ideal of a chordal graph G, then reg($R/I(H_t(G))$) = $(t-1)\nu(\mathcal{H}_t(G))$ and $\text{pd}(R/I(\mathcal{H}_t(G))) = \text{bight}(I(\mathcal{H}_t(G)))$. Also, it follows from [\[7,](#page-13-3) Theorem 3.12] that if the complement of G is chordal, then $I(\mathcal{H}_t(G))$ has a linear resolution. In view of this, the following question arises naturally, which nicely extends the edge ideals to a higher degree from the perspective of chordal graphs.

Question 5.1. *What type of t-uniform hypergraphs* $\mathcal{H}_t(G)$ *induced from a graph* G *satisfy the following three conditions simultaneously for all* $t \geq 2$ *:*

- *(i)* $reg(R/I(\mathcal{H}_t(G))) = (t-1)\nu(\mathcal{H}_t(G))$ *when G is chordal*,
- *(ii)* $\text{pd}(R/I(\mathcal{H}_t(G))) = \text{bight}(I(\mathcal{H}_t(G)))$ *when G is chordal,*
- *(iii)* $I(\mathcal{H}_t(G))$ *has a linear resolution when the complement of G is chordal.*

First, one may think of answering the above question for the existing classes of edge ideals of t-uniform hypergraphs induced from a graph, such as the t -path ideals and the t -clique ideals of graphs. Note that the t-path ideals fail to satisfy conditions (i) and (ii) of the above question (see [\[6,](#page-13-14) Theorem 5.3, 5.8]). However, to the best of our knowledge, it is still not known whether t -path ideals satisfy the condition (iii) or not.

Next, let us consider the t-clique ideal of a graph [\[17,](#page-13-2) Definition 3.1]. Then condition (iii) of the above question holds true [\[17,](#page-13-2) Corollary 3.4]. We do not know about the condition (ii). However, condition (i) is not true for t-cliques ideals, which follows from the following example.

Example 5.2. Let $G_{t,r} = G_1 \cup \cdots \cup G_{r+1}$ be a graph with $G_i \simeq K_t$ for all $i \in \{1, \ldots, r+1\}$ and there is a vertex $x \in V(G_{t,r})$ such that $V(G_i) \cap V(G_j) = \{x\}$ for all distinct i and j. Let $\mathcal{H}_t(G_{t,r})$ be the corresponding hypergraph of the t-connected ideal of $G_{t,r}$. Then the t-connected ideal of $G_{t,r}$ is $I(\mathcal{H}_t(\hat{G}_{t,r})) = X\setminus \mathbf{x}_{V(G_i)\setminus \{x\}} \mid 1 \leq i \leq r+1$). Then one can easily obtain that $reg(R/I(\mathcal{H}_t(G_{t,r}))) = (t-2)(r+1) + 1$, whereas the induced matching number of $\mathcal{H}_t(G_{t,r})$ is 1. Therefore, we have $reg(R/I(\mathcal{H}_t(G_{t,r}))) - (t-1)\nu(\mathcal{H}_t(G_{t,r})) = (t-2)r$. In other words, the regularity can be arbitrarily larger than the general lower bound for any given $t > 2$.

Moving on, in the case of edge ideals of graphs, there are several classes of graphs other than the chordal one for which $reg(R/I(G)) = \nu(G)$ (see [\[3,](#page-13-12) Theorem 14]) and $pd(R/I(G)) = bight(I(G))$ (for example, sequentially Cohen-Macaulay edge ideals) hold. In this article, we have extended the above formulas in Theorem [3.6](#page-7-0) and [4.5](#page-9-0) for the t-connected ideals of chordal graphs. Thus, the following question naturally arises in this context.

Question 5.3. *Find those classes of graph* G *for which* reg($R/I_t(G)$) = (t-1) $\nu_t(G)$ and $\text{pd}(R/I_t(G))$ = bight($I_t(G)$) for all $t \geq 2$, where $I_t(G)$ denotes the *t*-connected ideal of G.

Acknowledgements. The first and the second authors are supported by Postdoctoral Fellowships at Chennai Mathematical Institute. The third author would like to thank the National Board for Higher Mathematics (India) for the financial support through the NBHM Postdoctoral Fellowship. All the authors are partially supported by a grant from the Infosys Foundation.

Data availability statement. Data sharing is not applicable to this article as no new data were created or analyzed in this study.

Conflict of interest. The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

- [1] F. M. Abdelmalek, P. Deshpande, S. Goyal, A. Roy, and A. Singh. Chordal graphs, higher independence and vertex decomposable complexes. *Internat. J. Algebra Comput.*, 33(3):481–498, 2023. [2,](#page-1-0) [12](#page-11-3)
- [2] H. Ananthnarayan, O. Javadekar, and A. Maithani. Linear quotients of connected ideals of graphs. *arXiv:2401.01046*, 2024. [2,](#page-1-0) [3,](#page-2-2) [9](#page-8-5)
- [3] A. Banerjee, S. K. Beyarslan, and H. Huy Tài. Regularity of edge ideals and their powers. In *Advances in algebra*, volume 277 of *Springer Proc. Math. Stat.*, pages 17–52. Springer, Cham, 2019. [2,](#page-1-0) [13](#page-12-3)
- [4] A. Conca and E. De Negri. M-sequences, graph ideals, and ladder ideals of linear type. *J. Algebra*, 211(2):599–624, 1999. [1](#page-0-0)
- [5] H. Dao, C. Huneke, and J. Schweig. Bounds on the regularity and projective dimension of ideals associated to graphs. *J. Algebraic Combin.*, 38(1):37–55, 2013. [5](#page-4-4)
- [6] K. K. Das, A. Roy, and K. Saha. On the path ideals of chordal graphs. *arXiv:2405.15897*, 2024. [3,](#page-2-2) [13](#page-12-3)
- [7] P. Deshpande, A. Roy, A. Singh, and A. Van Tuyl. Fröberg's theorem, vertex splittability and higher independence complexes. *To appear in the Journal of Commutative Algebra, arXiv:*2311.02430, 2023. [2,](#page-1-0) [13](#page-12-3)
- [8] P. Deshpande, S. Shukla, and A. Singh. Distance r-domination number and r-independence complexes of graphs. *European J. Combin.*, 102:Paper No. 103508, 14, 2022. [2](#page-1-0)
- [9] G. A. Dirac. On rigid circuit graphs. *Abh. Math. Sem. Univ. Hamburg*, 25:71–76, 1961. [4](#page-3-0)
- [10] R. Fröberg. On Stanley-Reisner rings. In *Topics in algebra, Part 2 (Warsaw, 1988)*, volume 26, Part 2 of *Banach Center Publ.*, pages 57–70. PWN, Warsaw, 1990. [1](#page-0-0)
- [11] H. T. Hà. Regularity of squarefree monomial ideals. In *Connections between algebra, combinatorics, and geometry*, volume 76 of *Springer Proc. Math. Stat.*, pages 251–276. Springer, New York, 2014. [5](#page-4-4)
- [12] N. T. Hang and T. Vu. Projective dimension and regularity of 3-path ideals of unicyclic graphs. *arXiv:2402.16166*, 2024. [3](#page-2-2)
- [13] J. Herzog and T. Hibi. *Monomial ideals*, volume 260 of *Graduate Texts in Mathematics*. Springer-Verlag London, Ltd., London, 2011. [5,](#page-4-4) [6](#page-5-6)
- [14] J. Herzog, T. Hibi, and X. Zheng. Cohen-Macaulay chordal graphs. *J. Combin. Theory Ser. A*, 113(5):911–916, 2006. [3,](#page-2-2) [12](#page-11-3)
- [15] S. Jacques. Betti numbers of graph ideals. *arXiv:*0410107, 2004. [12](#page-11-3)
- [16] R. Meshulam. Domination numbers and homology. *J. Combin. Theory Ser. A*, 102(2):321–330, 2003. [2](#page-1-0)
- [17] S. Moradi. t-clique ideal and t-independence ideal of a graph. *Communications in Algebra*, 46(8):3377–3387, 2018. [1,](#page-0-0) [13](#page-12-3)
- [18] S. Morey and R. H. Villarreal. Edge ideals: algebraic and combinatorial properties. In *Progress in commutative algebra 1*, pages 85–126. de Gruyter, Berlin, 2012. [12](#page-11-3)
- [19] G. Paolini and M. Salvetti. Weighted sheaves and homology of Artin groups. *Algebr. Geom. Topol.*, 18(7):3943–4000, 2018. [2](#page-1-0)
- [20] I. Peeva. *Graded syzygies*, volume 14 of *Algebra and Applications*. Springer-Verlag London, Ltd., London, 2011. [5](#page-4-4)
- [21] M. Salvetti. Some combinatorial constructions and relations with Artin groups. In *Combinatorial methods in topology and algebra*, volume 12 of *Springer INdAM Ser.*, pages 121–126. Springer, Cham, 2015. [2](#page-1-0)
- [22] E. Sampathkumar. Generalizations of independence and chromatic numbers of a graph. *Discrete Math.*, 115(1- 3):245–251, 1993. [2](#page-1-0)
- [23] T. Szabó and G. Tardos. Extremal problems for transversals in graphs with bounded degree. *Combinatorica*, 26(3):333–351, 2006. [2](#page-1-0)
- [24] N. Terai. Alexander duality theorem and Stanley-Reisner rings. Number 1078, pages 174–184. 1999. Free resolutions of coordinate rings of projective varieties and related topics (Japanese) (Kyoto, 1998). [6](#page-5-6)
- [25] F. Tian and J.-M. Xu. A note on distance domination numbers of graphs. *Australas. J. Combin.*, 43:181–190, 2009. [2](#page-1-0)
- [26] D. R. Wood. Defective and clustered graph colouring. *Electron. J. Combin.*, DS23:71, 2018. [2](#page-1-0)

CHENNAI MATHEMATICAL INSTITUTE, INDIA *Email address*: kanoydas@cmi.ac.in; kanoydas0296@gmail.com

CHENNAI MATHEMATICAL INSTITUTE, INDIA *Email address*: amitiisermohali493@gmail.com

CHENNAI MATHEMATICAL INSTITUTE, INDIA *Email address*: ksaha@cmi.ac.in; kamalesh.saha44@gmail.com