Congruence properties of Lehmer-Euler numbers

Takao Komatsu

Faculty of Education Nagasaki University Nagasaki 852-8521 Japan komatsu@nagasaki-u.ac.jp

Guo-Dong Liu

Department of Mathematics and Statistics Huizhou University Huizhou, Guangdong 516007 China lgd@hzu.edu.cn

Abstract

Certain generalization of Euler numbers was defined in 1935 by Lehmer using cubic roots of unity, as a natural generalization of Bernoulli and Euler numbers. In this paper, Lehmer's generalized Euler numbers are studied to give certain congruence properties together with recurrence and explicit formulas of the numbers. We also show a new polynomial sequence and its properties. Some identities including Euler and central factorial numbers are obtained.

Keywords: Lehmer numbers, Euler numbers, Bernoulli numbers, congruence property, determinant expression, recurrence formula, central factorial numbers

1 Introduction

In 1935, D. H. Lehmer [13] introduced and investigated generalized Euler numbers W_n , which are defined as

$$\sum_{n=0}^{\infty} W_n \frac{t^n}{n!} = \frac{3}{e^t + e^{\omega t} + e^{\omega^2 t}} = \left(\sum_{l=0}^{\infty} \frac{t^{3l}}{(3l)!}\right)^{-1},$$

where $\omega = \frac{-1+\sqrt{-3}}{2}$ and $\omega^2 = \bar{\omega} = \frac{-1-\sqrt{-3}}{2}$ are the cubic roots of unity. Note that $W_n = 0$ if and only if $3 \nmid n$, and $(-1)^n W_{3n} > 0$. They can be considered as analogues of Euler numbers E_n defined as

$$\sum_{n=0}^{\infty} E_n \frac{t^n}{n!} = \frac{2}{e^t + e^{-t}} = \frac{1}{\cosh t} = \left(\sum_{l=0}^{\infty} \frac{t^{2l}}{(2l)!}\right)^{-1}$$

,

and its subsequence of the absolute values of its non-zero elements $\{\mathcal{W}_{3n}\}_{n\geq 0}$, where $\mathcal{W}_n := (-1)^n W_n$, is recorded in [23, A002115]:

 $1, 1, 19, 1513, 315523, 136085041, 105261234643, 132705221399353, 254604707462013571, 705927677520644167681, 2716778010767155313771539, \ldots$

We shall call W_n the (nth) Lehmer-Euler number in this paper for short.

In [1] several properties on determinants and representations of the Lehmer-Euler numbers are studied. Some generalizations are shown in [10]. In this paper, we show several congruence properties together with recurrence and explicit formulas of the numbers.

2 Basic properties of W_n

In this section, we remember some fundamental properties of Lehmer-Euler numbers W_n from [1, 10]. They are used to show congruence properties in the next main section.

Theorem 1. (i) (Recurrence formula) We have $W_0 = 1$ and

$$W_{3n} = -\sum_{k=0}^{n-1} {3n \choose 3k} W_{3k} \quad (n \ge 1).$$

(ii) (Explicit formula) For $n \ge 1$, we have

$$W_{3n} = (3n)! \sum_{k=1}^{n} (-1)^k \sum_{\substack{i_1 + \dots + i_k = n \\ i_1, \dots, i_k \ge 1}} \frac{1}{(3i_1)! \cdots (3i_k)!}.$$

(iii) (Determinant expression) For $n \ge 1$, we have

$$W_{3n} = (-1)^n (3n)! \begin{vmatrix} \frac{1}{3!} & 1 & 0 \\ \frac{1}{6!} & \frac{1}{3!} \\ \vdots & \vdots & \ddots & 1 & 0 \\ \frac{1}{(3n-3)!} & \frac{1}{(3n-6)!} & \cdots & \frac{1}{3!} & 1 \\ \frac{1}{(3n)!} & \frac{1}{(3n-3)!} & \cdots & \frac{1}{6!} & \frac{1}{3!} \end{vmatrix}.$$

By using Trudi's formula ([2, 21, 26]) we can obtain another explicit expression and inversion relation of W_n .

Theorem 2. For $n \ge 1$, we have

$$W_{3n} = (3n)! \sum_{t_1+2t_2+\dots+nt_n=n} \binom{t_1+\dots+t_n}{t_1,\dots,t_n} \prod_{l=1}^n \left(\frac{-1}{(3l)!}\right)^{t_l}.$$

By using the inversion formula (see, e.g. [11]), we obtain the following. **Proposition 1.** For $n \ge 1$, we have

$$\frac{(-1)^n}{(3n)!} = \begin{vmatrix} \frac{W_3}{3!} & 1 & 0 \\ \frac{W_6}{6!} & \frac{W_3}{3!} \\ \vdots & \vdots & \ddots & 1 & 0 \\ \frac{W_{3n-3}}{(3n-3)!} & \frac{W_{3n-6}}{(3n-6)!} & \cdots & \frac{W_3}{3!} & 1 \\ \frac{W_{3n}}{(3n)!} & \frac{W_{3n-3}}{(3n-3)!} & \cdots & \frac{W_6}{6!} & \frac{W_3}{3!} \end{vmatrix} .$$

3 Congruence properties

In this section, we give some congruence properties of W_n .

Theorem 3. For any non-negative integer n, we have

$$W_{3n} \in \mathbb{Z}$$
 and $W_{3n} \equiv (-1)^n \pmod{9}$.

Proof. By Theorem 1 (i), for n > 0, we have

$$W_{3n} = -\sum_{k=0}^{n-1} \binom{3n}{3k} W_{3k}.$$

Since $\binom{3n}{3k}$ is an integer and $W_0 = 1$, W_{3n} is an integer for all nonnegative integer n. By

$$\sum_{k=0}^{n} \binom{3n}{3k} x^{3k} = \frac{1}{3} \sum_{j=0}^{2} (1 + \omega^{j} x)^{3n},$$

we get

$$\sum_{k=0}^{n} \binom{3n}{3k} (-1)^{k} = \frac{1}{3} \sum_{j=0}^{2} (1-\omega^{j})^{3n} = 3^{n-1} \left((-\sqrt{-3})^{n} + (\sqrt{-3})^{n} \right).$$
(1)

Hence, if n is odd positive, then

$$\sum_{k=0}^{n} \binom{3n}{3k} (-1)^k = 0$$

and if n = 2n' is even positive, then

$$\sum_{k=0}^{n} \binom{3n}{3k} (-1)^{k} = 2(-1)^{n'} 3^{3n'-1} \equiv 0 \pmod{9}.$$

By induction with Theorem 1 (i), for n > 0, we have

$$W_{3n} \equiv (-1)^n - \sum_{k=0}^n \binom{3n}{3k} (-1)^k \equiv (-1)^n \pmod{9}.$$

From Theorem 3, for $m \ge 0$, we have

$$W_{6m} \equiv 1 \text{ and } W_{6m+3} \equiv -1 \pmod{3^2}$$
.

For a prime p, the following Lucas's Theorem ([16], see also [15]) is well-known.

Lemma 1 (Lucas's Theorem). If p is a prime and m and n are non-negative integers with base p representations $m = m_r p^r + \cdots + m_1 p + m_0$ and $n = n_r p^r + \cdots + n_1 p + n_0$, respectively, then

$$\binom{m}{n} = \binom{m_r p^r + \dots + m_1 p + m_0}{n_r p^r + \dots + n_1 p + n_0} \equiv \prod_{i=0}^r \binom{m_i}{n_i} \pmod{p}.$$

By using Lemma 1, we have the following result.

Theorem 4. For $n \ge 0$, we have

$$W_{9n} \equiv (-1)^n$$
, $W_{9n+3} = (-1)^{n-1}$, $W_{9n+6} \equiv (-1)^{n-1} 8 \pmod{3^3}$.

In particular, for $m \ge 0$,

$$\begin{split} W_{18m} &\equiv 1, \quad W_{18m+3} \equiv -1, \quad W_{18m+6} \equiv -8, \\ W_{18m+9} &\equiv -1, \quad W_{18m+12} \equiv 1, \quad W_{18m+15} \equiv 8 \pmod{3^3}. \end{split}$$

Proof. When n = 0, we know that $W_0 = 1$, $W_3 = -1$ and $W_6 = 19 \equiv -8 \pmod{27}$. By Lucas's Theorem, we have for $\ell = 0, 1, 2$

$$\binom{9n}{9k+3\ell} \equiv \binom{3n}{3k+\ell} \pmod{3}.$$

In fact, these congruences hold for $\pmod{3^l}$ $(\forall l \ge 1)$ if n is enough large. Since

$$\sum_{k=0}^{n-1} \binom{3n}{3k+1} x^{3k+1} = \frac{1}{3} \sum_{j=0}^{2} \omega^{2j} (1+\omega^j)^{3n} ,$$

by setting x = -1 we have

$$\begin{split} \sum_{k=0}^{n-1} \binom{3n}{3k+1} (-1)^{k-1} &= \frac{1}{3} \sum_{j=0}^{2} \omega^{2j} (1+\omega^j)^{3n} \\ &= \frac{1}{3} \left(\omega^2 (-3\sqrt{-3})^n + \omega(3\sqrt{-3})^n \right) \\ &= 3^{n-1} \left(\omega^2 (-\sqrt{-3})^n + \omega(\sqrt{-3})^n \right). \end{split}$$

Hence,

$$\sum_{k=0}^{n-1} \binom{3n}{3k+1} (-1)^{k-1} = \begin{cases} (-1)^{\frac{n+1}{2}} 3^{\frac{3n-1}{2}} & \text{when } n \text{ is odd;} \\ (-1)^{\frac{n}{2}} 3^{\frac{3n}{2}-1} & \text{when } n \text{ is even.} \end{cases}$$

Since

$$\sum_{k=0}^{n-1} \binom{3n}{3k+2} x^{3k+2} = \frac{1}{3} \sum_{j=0}^{2} \omega^{j} (1+\omega^{j})^{3n} ,$$

we have

$$\sum_{k=0}^{n-1} \binom{3n}{3k+2} (-1)^{k-1} 8 = -\frac{8}{3} \sum_{j=0}^{2} \omega^{j} (1+\omega^{j})^{3n}$$
$$= -8 \cdot 3^{n-1} \left(\omega (-\sqrt{-3})^{n} + \omega^{2} (\sqrt{-3})^{n} \right).$$

Hence,

$$\sum_{k=0}^{n-1} \binom{3n}{3k+2} (-1)^{k-1} 8 = \begin{cases} (-1)^{\frac{n+1}{2}} 8 \cdot 3^{\frac{3n-1}{2}} & \text{when } n \text{ is odd;} \\ (-1)^{\frac{n}{2}} 8 \cdot 3^{\frac{3n}{2}-1} & \text{when } n \text{ is even.} \end{cases}$$

By Theorem 1 (i) together with the proof of Theorem 3, for $n \ge 3$, we have

$$W_{9n} = -\sum_{k=0}^{3n-1} {9n \choose 3k} W_{3k}$$

= $-\sum_{k=0}^{n-1} {9n \choose 9k} W_{9k} - \sum_{k=0}^{n-1} {9n \choose 9k+3} W_{9k+3} - \sum_{k=0}^{n-1} {9n \choose 9k+6} W_{9k+6}$
= $(-1)^n - 0 - 0 = (-1)^n \pmod{3^3}.$

When n = 1, we have $W_9 = -1513 \equiv -1 \pmod{3^3}$. When n = 2, we have $W_{18} = 105261234643 \equiv 1 \pmod{3^3}$. In fact, for $n \ge n_0$

$$W_{9n} \equiv \begin{cases} (-1)^n \pmod{3^{\frac{3n_0-1}{2}}} & \text{if } n_0 \text{ is odd;} \\ (-1)^n \pmod{3^{\frac{3n_0}{2}-1}} & \text{if } n_0 \text{ is even.} \end{cases}$$

Since

$$\sum_{k=0}^{n} \binom{3n+1}{3k} x^{3k} = \frac{1}{3} \sum_{j=0}^{2} (1+\omega^{j} x)^{3n+1},$$

for x = -1, we have

$$\sum_{k=0}^{n} \binom{3n+1}{3k} (-1)^{k} = \frac{1}{3} \sum_{j=0}^{2} (1-\omega^{j})^{3n+1}$$
$$= \frac{1}{3} \left((-3\sqrt{-3})^{n} (1-\omega) + (3\sqrt{-3})^{n} (1-\omega^{2}) \right).$$

Hence,

$$\sum_{k=0}^{n} \binom{3n+1}{3k} (-1)^{k} = \begin{cases} (-1)^{\frac{n+1}{2}} 3^{\frac{3n-1}{2}} & n \text{ is odd;} \\ (-1)^{\frac{n}{2}} 3^{\frac{3}{2}n} & n \text{ is even.} \end{cases}$$

Since

$$\sum_{k=0}^{n} \binom{3n+1}{3k+1} x^{3k+1} = \frac{1}{3} \sum_{j=0}^{2} \omega^{2j} (1+\omega^{j} x)^{3n+1},$$

we have

$$\begin{split} \sum_{k=0}^{n} \binom{3n+1}{3k+1} (-1)^{k-1} &= \frac{1}{3} \sum_{j=0}^{2} \omega^{2j} (1-\omega^{j})^{3n+1} \\ &= \frac{1}{3} \left(\omega^{2} (-3\sqrt{-3})^{n} (1-\omega) + \omega (3\sqrt{-3})^{n} (1-\omega^{2}) \right) \\ &= \frac{1}{3} \left((\omega^{2} - 1) (-3\sqrt{-3})^{n} + (\omega - 1) (3\sqrt{-3})^{n} \right). \end{split}$$

Hence,

$$\sum_{k=0}^{n} \binom{3n+1}{3k+1} (-1)^{k-1} = \begin{cases} (-1)^{\frac{n-1}{2}} 3^{\frac{3n-1}{2}} & n \text{ is odd;} \\ (-1)^{\frac{n}{2}+1} 3^{\frac{3}{2}n} & n \text{ is even.} \end{cases}$$

Since

$$\sum_{k=0}^{n-1} \binom{3n+1}{3k+2} x^{3k+2} = \frac{1}{3} \sum_{j=0}^{2} \omega^{j} (1+\omega^{j} x)^{3n+1},$$

we have

$$\sum_{k=0}^{n-1} \binom{3n+1}{3k+2} (-1)^{k-1} 8 = -\frac{8}{3} \sum_{j=0}^{2} \omega^{j} (1-\omega^{j})^{3n+1}$$
$$= -\frac{8}{3} \left(\omega (-3\sqrt{-3})^{n} (1-\omega) + \omega^{2} (3\sqrt{-3})^{n} (1-\omega^{2}) \right)$$
$$= -\frac{8}{3} \left((\omega - \omega^{2}) (-3\sqrt{-3})^{n} + (\omega^{2} - \omega) (3\sqrt{-3})^{n} \right).$$

Hence,

$$\sum_{k=0}^{n-1} \binom{3n+1}{3k+2} (-1)^k = \begin{cases} (-1)^{\frac{n+1}{2}} 16 \cdot 3^{\frac{3n-1}{2}} & n \text{ is odd;} \\ 0 & n \text{ is even.} \end{cases}$$

By Theorem 1 (i), for $n \ge 2$, we have

$$\begin{split} W_{9n+3} &= -\sum_{k=0}^{3n+1} \binom{9n+3}{3k} W_{3k} \\ &= -\sum_{k=0}^{n} \binom{9n+3}{9k} W_{9k} - \sum_{k=0}^{n-1} \binom{9n+3}{9k+3} W_{9k+3} - \sum_{k=0}^{n-1} \binom{9n+3}{9k+6} W_{9k+6} \\ &\equiv -0 + (-1)^{n-1} - 0 = (-1)^{n-1} \pmod{3^3}. \end{split}$$

When n = 1, we have $W_{12} = 315523 \equiv 1 \pmod{3^3}$. In fact, for $n \ge n_0$

$$W_{9n+3} \equiv \begin{cases} (-1)^{n-1} \pmod{3^{\frac{3n_0+1}{2}}} & \text{if } n_0 \text{ is odd;} \\ (-1)^{n-1} \pmod{3^{\frac{3n_0}{2}}} & \text{if } n_0 \text{ is even.} \end{cases}$$

Since

$$\sum_{k=0}^{n} \binom{3n+2}{3k} x^{3k} = \frac{1}{3} \sum_{j=0}^{2} (1+\omega^{j} x)^{3n+2},$$

for x = -1, we have

$$\begin{split} \sum_{k=0}^{n} \binom{3n+2}{3k} (-1)^{k} &= \frac{1}{3} \sum_{j=0}^{2} (1-\omega^{j})^{3n+2} \\ &= \frac{1}{3} \left((-3\sqrt{-3})^{n} (1-\omega)^{2} + (3\sqrt{-3})^{n} (1-\omega^{2})^{2} \right) \\ &= - \left(\omega (-3\sqrt{-3})^{n} + \omega^{2} (3\sqrt{-3})^{n} \right). \end{split}$$

Hence,

$$\sum_{k=0}^{n} \binom{3n+2}{3k} (-1)^{k} = \begin{cases} (-1)^{\frac{n+1}{2}} 3^{\frac{3n+1}{2}} & n \text{ is odd;} \\ (-1)^{\frac{n}{2}} 3^{\frac{3}{2}n} & n \text{ is even.} \end{cases}$$

Since

$$\sum_{k=0}^{n} \binom{3n+2}{3k+1} x^{3k+1} = \frac{1}{3} \sum_{j=0}^{2} \omega^{2j} (1+\omega^{j}x)^{3n+2},$$

we have

$$\sum_{k=0}^{n} \binom{3n+2}{3k+1} (-1)^{k-1} = \frac{1}{3} \sum_{j=0}^{2} \omega^{2j} (1-\omega^{j})^{3n+2}$$
$$= \frac{1}{3} \left(\omega^{2} (-3\sqrt{-3})^{n} (1-\omega)^{2} + \omega (3\sqrt{-3})^{n} (1-\omega^{2})^{2} \right)$$
$$= -\left((-3\sqrt{-3})^{n} + (3\sqrt{-3})^{n} \right).$$

Hence,

$$\sum_{k=0}^{n} \binom{3n+2}{3k+1} (-1)^{k-1} = \begin{cases} 0 & n \text{ is odd;} \\ (-1)^{\frac{n}{2}-1} 2 \cdot 3^{\frac{3}{2}n} & n \text{ is even.} \end{cases}$$

Since

$$\sum_{k=0}^{n} \binom{3n+2}{3k+2} x^{3k+2} = \frac{1}{3} \sum_{j=0}^{2} \omega^{j} (1+\omega^{j}x)^{3n+2},$$

we have

$$\begin{split} \sum_{k=0}^{n} \binom{3n+2}{3k+2} (-1)^{k-1} &= -\frac{8}{3} \sum_{j=0}^{2} \omega^{j} (1-\omega^{j})^{3n+2} \\ &= -\frac{8}{3} \left(\omega^{2} (-3\sqrt{-3})^{n} (1-\omega)^{2} + \omega (3\sqrt{-3})^{n} (1-\omega^{2})^{2} \right) \\ &= 8 \left(\omega^{2} (-3\sqrt{-3})^{n} + \omega (3\sqrt{-3})^{n} \right). \end{split}$$

Hence,

$$\sum_{k=0}^{n} \binom{3n+2}{3k+2} (-1)^{k-1} 8 = \begin{cases} (-1)^{\frac{n+1}{2}} 8 \cdot 3^{\frac{3n+1}{2}} & n \text{ is odd;} \\ (-1)^{\frac{n}{2}} 8 \cdot 3^{\frac{3}{2}n} & n \text{ is even.} \end{cases}$$

By Theorem 1 (i), for $n \ge 2$, we have

$$W_{9n+6} = -\sum_{k=0}^{3n+1} \binom{9n+6}{3k} W_{3k}$$

= $-\sum_{k=0}^{n} \binom{9n+6}{9k} W_{9k} - \sum_{k=0}^{n} \binom{9n+6}{9k+3} W_{9k+3} - \sum_{k=0}^{n-1} \binom{9n+6}{9k+6} W_{9k+6}$
= $-0 - 0 + (-1)^{n-1} 8 = (-1)^{n-1} 8 \pmod{3^3}.$

When n = 1, we have $W_{15} = -136085041 \equiv 8 \pmod{3^3}$.

$$W_{9n+6} \equiv -1 - 24 - 3 \equiv -1 \pmod{3^3}$$
.

In fact, for $n \ge n_0$

$$W_{9n+6} \equiv \begin{cases} (-1)^{n-1}8 \pmod{3^{\frac{3n_0+1}{2}}} & \text{if } n_0 \text{ is odd;} \\ (-1)^{n-1}8 \pmod{3^{\frac{3n_0}{2}}} & \text{if } n_0 \text{ is even.} \end{cases}$$

-		

3.1 Lehmer-Euler numbers modulo powers of three

A classical result on Euler numbers modulo power of two due to Stern [24] asserts that

$$E_{2n} \equiv E_{2m} \pmod{2^k} \iff 2n \equiv 2m \pmod{2^k}.$$
 (2)

In 1875 Stern [24] gave a brief sketch of a proof of (2), then Frobenius [5] amplified Stern's sketch in 1910. In 1979 Ernvall [4] said that he could

not understand Frobenius' proof and gave his own proof using the umbral calculus. In 2000 an induction proof of (2) was given by Wagstaff [27]. Sun [25] obtained an explicit congruence for Euler numbers modulo powers of two to give a new proof of (2). Liu [14] gave a still new proof of (2) by using the Euler numbers of order 2 and the central factorial numbers.

In the case of the Lemer-Euler numbers, the following is predicted.

 $W_{3n} \equiv W_{3m} \pmod{3^k} \iff 3n \equiv 3m \pmod{3^k}.$

But unfortunately this does not hold. Together with Theorems 3 and 4, indeed, we have the following.

Proposition 2. We have

 $\{ W_{3n} \pmod{3^2} \}_{n=0}^{\infty} = 1, \overline{8, 1} \pmod{2}, \\ \{ W_{3n} \pmod{3^3} \}_{n=0}^{\infty} = 1, \overline{26, 19, 26, \overline{1, 8, 1}} \pmod{6}, \\ \{ W_{3n} \pmod{3^4} \}_{n=0}^{\infty} = 1, \overline{26, 19, 26, \overline{1, 8, 1}} \pmod{6}, \\ \{ W_{3n} \pmod{3^4} \}_{n=0}^{\infty} = 1, \overline{242, 19, 80}, \overline{1, 62, 55, 53, 19, 53, 55, 62, 1} \pmod{2 \cdot 3^2}, \\ \{ W_{3n} \pmod{3^5} \}_{n=0}^{\infty} = 1, \overline{242, 19, 188, 109, 62, 28, 107, 19, 80, 82, 224, 217, 53}, \\ \overline{181, 53, 217, 224, 82, 80, 19, 107, 28, 62, 109, 188, 19, 242}, \\ \overline{1, 224, 55, 134, 181, 215, 136, 224, 163, 161, 19, 26, 190, 62, 190, 26, 19, 161}, \\ \overline{163, 224, 136, 215, 181, 134, 55, 224, 1} \pmod{2 \cdot 3^3}.$

Remark. For $k \geq 6$, the palindromic property does not hold for $\{W_{3n} \pmod{3^k}\}_{n=0}^{\infty}$.

Based on the above results, we predict that the following can be generally stated.

Conjecture 1. If $3n \equiv 3m \pmod{2 \cdot 3^k}$, then $W_{3n} \equiv W_{3m} \pmod{3^{k+1}}$.

Proof of Proposition 2. Concerning the case of $W_{3n} \pmod{3^4}$, we can show

$$W_{27n} \equiv (-1)^n, \quad W_{27n+3} \equiv (-1)^{n-1},$$

$$W_{27n+6} \equiv W_{27n+24} \equiv (-1)^n 19, \quad W_{27n+15} \equiv (-1)^{n-1} 19,$$

$$W_{27n+9} \equiv W_{27n+21} \equiv (-1)^n 26,$$

$$W_{27n+12} \equiv W_{27n+18} \equiv (-1)^n 28 \pmod{3^4}.$$

For example, similarly to the proof of Theorem 4, the first identity can be shown by Theorem 1 (i) and the facts of the following relations.

$$\begin{split} \sum_{k=0}^{n} \binom{27n}{27k} (-1)^{k} &\equiv \begin{cases} 0 & \text{if } n = 1 \text{ or } n \geq 3 \\ -18 & \text{if } n = 2, \end{cases} \\ \sum_{k=0}^{n-1} \binom{27n}{27k+3} (-1)^{k-1} &\equiv \begin{cases} -9 & \text{if } n = 1 \\ 0 & \text{if } n \geq 2, \end{cases} \\ \sum_{k=0}^{n} \binom{27n}{27k+6} (-1)^{k} 19 &\equiv \begin{cases} 36 & \text{if } n = 1 \\ 0 & \text{if } n \geq 2, \end{cases} \\ \sum_{k=0}^{n-1} \binom{27n}{27k+9} (-1)^{k} 26 &\equiv \begin{cases} -3 & \text{if } n = 1 \\ 9 & \text{if } n = 2 \\ 0 & \text{if } n \geq 3, \end{cases} \\ \sum_{k=0}^{n} \binom{27n}{27k+12} (-1)^{k} 28 &\equiv \begin{cases} 45 & \text{if } n = 1 \\ 0 & \text{if } n \geq 2, \end{cases} \\ \sum_{k=0}^{n-1} \binom{27n}{27k+12} (-1)^{k-1} 19 &\equiv \begin{cases} 36 & \text{if } n = 1 \\ 0 & \text{if } n \geq 2, \end{cases} \\ \sum_{k=0}^{n-1} \binom{27n}{27k+15} (-1)^{k-1} 19 &\equiv \begin{cases} 36 & \text{if } n = 1 \\ 0 & \text{if } n \geq 2, \end{cases} \\ \sum_{k=0}^{n-1} \binom{27n}{27k+15} (-1)^{k} 28 &\equiv \begin{cases} 3 & \text{if } n = 1 \\ 9 & \text{if } n \geq 2, \end{cases} \\ 0 & \text{if } n \geq 3, \end{cases} \\ \sum_{k=0}^{n-1} \binom{27n}{27k+21} (-1)^{k} 26 &\equiv \begin{cases} -36 & \text{if } n = 1 \\ 0 & \text{if } n \geq 2, \end{cases} \\ \sum_{k=0}^{n-1} \binom{27n}{27k+21} (-1)^{k} 26 &\equiv \begin{cases} -36 & \text{if } n = 1 \\ 0 & \text{if } n \geq 2, \end{cases} \\ \sum_{k=0}^{n-1} \binom{27n}{27k+21} (-1)^{k} 26 &\equiv \begin{cases} -36 & \text{if } n = 1 \\ 0 & \text{if } n \geq 2, \end{cases} \\ \sum_{k=0}^{n-1} \binom{27n}{27k+21} (-1)^{k} 26 &\equiv \begin{cases} -36 & \text{if } n = 1 \\ 0 & \text{if } n \geq 2, \end{cases} \\ \sum_{k=0}^{n-1} \binom{27n}{27k+24} (-1)^{k} 19 &\equiv \begin{cases} 9 & \text{if } n = 1 \\ 0 & \text{if } n \geq 2, \end{cases} \\ (\text{mod } 3^{4}) . \end{cases} \end{split}$$

Other identities are similarly shown and their proofs are omitted.

4 Incomplete Lehmer-Euler numbers

The various incomplete numbers have been introduced and studied (see, e.g., [8, 9, 11]). In order to generalize the numbers W_n , we shall introduce two kinds of incomplete Lehner-Euler numbers. For $m \geq 1$, define the

incomplete Lehmer-Euler numbers $W_{n,\leq m}$ and $W_{n,\geq m}$ by

$$\sum_{n=0}^{\infty} W_{n,\leq m} \frac{t^n}{n!} = \frac{1}{1 + \sum_{l=1}^{m} \frac{t^{3l}}{(3l)!}}$$
(3)

and

$$\sum_{n=0}^{\infty} W_{n,\geq m} \frac{t^n}{n!} = \frac{1}{1 + \sum_{l=m}^{\infty} \frac{t^{3l}}{(3l)!}},$$
(4)

respectively. When $m \to \infty$, $W_n = W_{n, \leq \infty}$, and when m = 1, $W_n = W_{n, \geq 1}$. Hence, both incomplete numbers are reduced to the original Lehmer-Euler numbers. Moreover, note that $W_{n, \leq m} = W_{n, \geq m} = 0$ if and only if $3 \nmid n$.

These incomplete Lehmer-Euler numbers satisfy the following recurrence and explicit formulas and determinant expressions. The proposition can be proven similar to Theorem 1, thus we omit the proof.

Proposition 3. (i) *(Recurrence formulas) We have the following.*

(a)

$$W_{3n,\le m} = -\sum_{k=\max\{n-m,0\}}^{n-1} \binom{3n}{3k} W_{3k,\le m} \quad (n\ge 1)$$

with $W_{0,\leq m} = 1$. (b)

$$W_{3n,\geq m} = -\sum_{k=0}^{n-m} \binom{3n}{3k} W_{3k,\geq m} \quad (n\geq m)$$

with $W_{0,\geq m} = 1$ and $W_{3,\geq m} = \cdots = W_{3m-3,\geq m} = 0$.

(ii) (Explicit formulas) For n, m ≥ 1, we have the following:
(a)

$$W_{3n,\leq m} = (3n)! \sum_{k=1}^{n} (-1)^k \sum_{\substack{i_1 + \dots + i_k = n \\ 1 \leq i_1, \dots, i_k \leq m}} \frac{1}{(3i_1)! \cdots (3i_k)!}$$

(b)

$$W_{3n,\geq m} = (3n)! \sum_{k=1}^{n} (-1)^k \sum_{\substack{i_1 + \dots + i_k = n \\ i_1, \dots, i_k \geq m}} \frac{1}{(3i_1)! \cdots (3i_k)!}.$$

(iii) (Determinant expressions) For n ≥ m ≥ 1, we have the following:
(a)

$$W_{3n,\leq m} = (-1)^{n} (3n)! \begin{vmatrix} \frac{1}{3!} & 1 & 0 \\ \vdots & \ddots & \ddots \\ \frac{1}{(3m)!} & & \ddots \\ 0 & \ddots & & \ddots \\ 0 & \ddots & & 1 \\ 0 & \frac{1}{(3m)!} & \cdots & \frac{1}{3!} \\ 0 & \frac{1}{(3m)!} & \cdots & \frac{1}{3!} \end{vmatrix}$$

•

(b)

$$W_{3n,\geq m} = (-1)^n (3n)! \begin{vmatrix} 0 & 1 & 0 \\ \vdots & \ddots & \ddots \\ 0 & & \ddots \\ \frac{1}{(3m)!} & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & 1 \\ \frac{1}{(3n)!} & \cdots & \frac{1}{(3m)!} & 0 & \cdots & 0 \\ & & & & & & \\ \hline n-m+1 & & & & & \\ \hline n-m+1 & & & & & \\ \hline \end{matrix} \right|.$$

The other explicit formulas, due to Trudi's formula, are also obtained for the incomplete cases, and the proof is similar to Theorem 2, thus we omit the proof.

Proposition 4. For $n \ge m \ge 1$, we have

$$W_{3n,\leq m} = (3n)! \sum_{t_1+2t_2+\dots+mt_m=n} {t_1+\dots+t_m \choose t_1,\dots,t_m} \prod_{l=1}^m \left(\frac{-1}{(3l)!}\right)^{t_l}$$

and

$$W_{3n,\geq m} = (3n)! \sum_{mt_m + (m+1)t_{m+1} + \dots + nt_n = n} \binom{t_m + t_{m+1} + \dots + t_n}{t_m, t_{m+1}, \dots, t_n} \prod_{l=m}^n \left(\frac{-1}{(3l)!}\right)^{t_l}.$$

5 Higher order Lehmer-Euler numbers

For positive integers r and α , the generalized Lehmer-Euler numbers of order α , which is denoted by $W_{r,n}^{(\alpha)}$ is defined by

$$\sum_{n=0}^{\infty} W_{r,n}^{(\alpha)} \frac{t^n}{n!} = \left(\frac{r}{\sum_{j=0}^{r-1} e^{\zeta^j t}}\right)^{\alpha} = \left(\sum_{l=0}^{\infty} \frac{t^{rl}}{(rl)!}\right)^{-\alpha},$$
(5)

where $\zeta = \zeta_r$ is the primitive *r*-th root of unity. Hence, when r = 3 and $\alpha = 1$, $W_n = W_{1,n}^{(1)}$ is the original Lehmer-Euler number. Since

$$(1+w)^{-\alpha} = \sum_{l=0}^{\infty} {\alpha+l-1 \choose l} (-w)^l \quad (|w|<1),$$

putting

$$1 + w = \frac{1}{r} \sum_{j=0}^{r-1} e^{\zeta^{j} t},$$

we have

$$\begin{split} W_{r,n}^{(\alpha)} &= \left. \frac{d^{n}}{dt^{n}} \left(\frac{r}{\sum_{j=0}^{r-1} e^{\zeta^{j}t}} \right)^{\alpha} \right|_{t=0} \\ &= \sum_{l=0}^{n} \left(\alpha + l - 1 \\ l \right) \left. \frac{d^{n}}{dt^{n}} \left(1 - \frac{1}{r} \sum_{j=0}^{r-1} e^{\zeta^{j}t} \right)^{l} \right|_{t=0} \\ &= \sum_{l=0}^{n} \left(\alpha + l - 1 \\ l \right) \left. \frac{d^{n}}{dt^{n}} \sum_{k=0}^{l} \left(l \\ k \right) \left(-\frac{1}{r} \sum_{j=0}^{r-1} e^{\zeta^{j}t} \right)^{k} \right|_{t=0} \\ &= \sum_{l=0}^{n} \left(\alpha + l - 1 \\ l \right) \sum_{k=0}^{l} \left(-\frac{1}{r} \right)^{k} \left(l \\ k \right) \\ &\times \left. \frac{d^{n}}{dt^{n}} \sum_{\substack{i_{1}+i_{2}+\dots+i_{r}=k}{i_{1},i_{2},\dots,i_{r}\geq 0}} \frac{k!}{i_{1}!i_{2}!\dots i_{r}} \prod_{j=1}^{r} e^{\zeta^{j-1}i_{j}t} \right|_{t=0} \\ &= \sum_{k=0}^{n} \left(-\frac{1}{r} \right)^{k} \sum_{\substack{i_{1}+i_{2}+\dots+i_{r}=k}{i_{1},i_{2},\dots,i_{r}\geq 0}} \frac{k!}{i_{1}!i_{2}!\dots i_{r}!} \left(\sum_{j=1}^{r} \zeta^{j-1}i_{j} \right)^{n} \end{split}$$

$$\times {\binom{\alpha+l-1}{l}\binom{l}{k}}$$

$$= \sum_{k=0}^{n} \left(-\frac{1}{r}\right)^{k} {\binom{\alpha+k-1}{k}} \sum_{\substack{i_{1}+i_{2}+\dots+i_{r}=k\\i_{1},i_{2},\dots,i_{r}\geq 0}} \frac{k!}{i_{1}!i_{2}!\dots i_{r}!} \left(\sum_{j=1}^{r} \zeta^{j-1}i_{j}\right)^{n}$$

$$\times \sum_{l=0}^{n-k} {\binom{\alpha+k+l-1}{l}}$$

$$= \sum_{k=0}^{n} \left(-\frac{1}{r}\right)^{k} {\binom{\alpha+k-1}{k}} {\binom{\alpha+n}{n-k}} \sum_{\substack{i_{1}+i_{2}+\dots+i_{r}=k\\i_{1},i_{2},\dots,i_{r}\geq 0}} \frac{k!}{i_{1}!i_{2}!\dots i_{r}!} \left(\sum_{j=1}^{r} \zeta^{j-1}i_{j}\right)^{n}.$$

When $\alpha = 1$, the generalized Lehmer-Euler numbers $W_{r,n}$ are reduced as

$$W_{r,n} = \sum_{k=0}^{n} \left(-\frac{1}{r}\right)^{k} \binom{n+1}{n-k} \sum_{\substack{i_{1}+i_{2}+\cdots+i_{r}=k\\i_{1},i_{2},\cdots,i_{r}\geq 0}} \frac{k!}{i_{1}!i_{2}!\cdots i_{r}!} \left(\sum_{j=1}^{r} \zeta^{j-1}i_{j}\right)^{n}.$$

If r = 3, we have an expression of the Lehmer-Euler numbers of higher-order:

$$W_n = \sum_{k=0}^n \left(-\frac{1}{3}\right)^k \binom{\alpha+n}{n-k} \binom{\alpha+k-1}{k} \sum_{\substack{i_1+i_2+i_3=k\\i_1,i_2,i_3\ge 0}} \frac{k!}{i_1!i_2!i_3!} \left(i_1+i_2\omega+i_3\omega^2\right)^n.$$

If r = 2, we have an expression of the Euler numbers of higher-order ([17]):

$$E_n^{(\alpha)} = \sum_{k=0}^n \left(-\frac{1}{2}\right)^k \binom{\alpha+n}{n-k} \binom{\alpha+k-1}{k} \sum_{j=0}^k \binom{k}{j} (k-2j)^n \,. \tag{6}$$

6 A new polynomial sequence and its properties

For a complex number x and a non-negative integer k, the polynomial sequence $\Delta(x, k)$ is defined by

$$\Delta(x,k+1) = (x+1)(2x+1)\Delta(x+1,k) - x^2\Delta(x,k) \quad (k \ge 0)$$
 (7)

with $\Delta(x,0) = 1$. By using the recurrence relation in (7), we have

$$\Delta(x,1) = x^2 + 3x + 1 \,,$$

$$\begin{split} \Delta(x,2) &= x^4 + 10x^3 + 25x^2 + 20x + 5, \\ \Delta(x,3) &= x^6 + 21x^5 + 140x^4 + 385x^3 + 490x^2 + 287x + 61. \end{split}$$

The central factorial numbers t(n, k) of the first kind (also known as the Stirling numbers of the first kind with level 2 [18]) are given by the formula

$$x\left(x+\frac{n}{2}-1\right)\left(x+\frac{n}{2}-2\right)\cdots\left(x-\frac{n}{2}+1\right) = \sum_{k=0}^{\infty} t(n,k)x^{k}$$
(8)

and its exponential generating function is given by

$$\left(2\log\left(\frac{x}{2} + \sqrt{\frac{x^2}{4} + 1}\right)\right)^k = k! \sum_{n=k}^{\infty} t(n,k) \frac{x^n}{n!}.$$
(9)

(see, e.g., [22]). It follows from (8) and (9) that

$$t(n,k) = t(n-2,k-2) - \frac{(n-2)^2}{4}t(n-2,k)$$
(10)

with t(n,0) = 0 (n > 0), t(n,n) = 1 $(n \ge 0)$ and t(n,k) = 0 (n + k is odd, k > n or k < 0.

The central factorial numbers T(n, k) of the second kind (also known as the Stirling numbers of the second kind with level 2 [19]) are given by the formula

$$x^{n} = \sum_{k=0}^{\infty} T(n,k) x\left(x + \frac{k}{2} - 1\right) \left(x + \frac{k}{2} - 2\right) \cdots \left(x - \frac{k}{2} + 1\right)$$
(11)

and its exponential generating function is given by

$$(e^{\frac{x}{2}} - e^{-\frac{x}{2}})^k = k! \sum_{n=k}^{\infty} T(n,k) \frac{x^n}{n!}.$$
 (12)

(see, e.g., [22]). It follows from (11) and (12) that

$$T(n,k) = T(n-2,k-2) + \frac{k^2}{4}T(n-2,k)$$
(13)

with T(n,0) = 0 (n > 0), T(n,n) = 1 $(n \ge 0)$ and T(n,k) = 0 (n+k is odd, k > n or k < 0).

In this section, we show the identity between Euler numbers and the central factorial numbers of the second kind in terms of the polynomial sequence $\Delta(x, k)$ in (7).

Theorem 5. For non-negative integers n and k, we have

$$E_{2n+2k} = \sum_{j=0}^{n} \frac{(-1)^{j-k}(2j)!\Delta(j,k)}{2^j} T(2n,2j)$$

Remark. When k = 0, the identity in Theorem 5 is reduced to

$$E_{2n} = \sum_{j=0}^{n} \frac{(-1)^j (2j)!}{2^j} T(2n, 2j) \,. \tag{14}$$

(see [20]). When n = 0, it is reduced to

$$E_{2k} = (-1)^k \Delta(0,k) \,. \tag{15}$$

The identity involving the central factorial numbers of the first kind and the polynomial sequence $\Delta(x, k)$ can be given as follows.

Theorem 6. For a non-negative integer n, we have

$$\sum_{j=0}^{n} (-4)^{n-j} t(2n+1,2j+1)\Delta(x,j) = (x+1)(x+2)\cdots(x+2n).$$

Remark. When x = 0, by (15), it is reduced to

$$\sum_{j=0}^{n} 4^{n-j} t(2n+1,2j+1) E_{2j} = (-1)^n (2n)!.$$

Proof of Theorem 5. We shall prove by induction. When k = 0, the identity is valid because of (14) in [20]. Suppose that the identity is valid for some k. By (7) and (13), we have

$$E_{2n+2(k+1)} = E_{2(n+1)+2k}$$

= $\sum_{j=0}^{n+1} \frac{(-1)^{j-k}(2j)!\Delta(j,k)}{2^j} T(2n+2,2j)$
= $\sum_{j=0}^{n+1} \frac{(-1)^{j-k}(2j)!\Delta(j,k)}{2^j} (T(2n,2j-2)+j^2T(2n,2j))$

$$\begin{split} &= \sum_{j=0}^{n} \frac{(-1)^{j-k+1}(2j+2)!\Delta(j+1,k)}{2^{j+1}} T(2n,2j) \\ &\quad + \sum_{j=0}^{n} \frac{(-1)^{j-k}(2j)!j^2\Delta(j,k)}{2^j} T(2n,2j) \\ &= \sum_{j=0}^{n} \frac{(-1)^{j-k+1}(2j)!}{2^j} \left((j+1)(2j+1)\Delta(j+1,k) - j^2\Delta(j,k) \right) T(2n,2j) \\ &= \sum_{j=0}^{n} \frac{(-1)^{j-(k+1)}(2j)!\Delta(j,k+1)}{2^j} T(2n,2j) \,. \end{split}$$

Hence, the identity is valid for k + 1.

Proof of Theorem 6. When n = 0, by $t(1,1)\Delta(x,0) = 1$, the identity is valid. Suppose that the identity is valid for some n. By (7) and (10), we have

$$\begin{split} &\sum_{j=0}^{n+1} (-4)^{n-j+1} t(2n+3,2j+1)\Delta(x,j) \\ &= \sum_{j=0}^{n+1} (-4)^{n-j+1} \left(t(2n+1,2j-1) - \frac{(2n+1)^2}{4} t(2n+1,2j+1) \right) \Delta(x,j) \\ &= \sum_{j=0}^{n+1} (-4)^{n-j+1} t(2n+1,2j-1)\Delta(x,j) \\ &\quad + \sum_{j=0}^n (-4)^{n-j} (2n+1)^2 t(2n+1,2j+1)\Delta(x,j) \\ &= \sum_{j=0}^n (-4)^{n-j} t(2n+1,2j+1)\Delta(x,j+1) \\ &\quad + (2n+1)^2 (x+1) (x+2) \cdots (x+2n) \\ &= \sum_{j=0}^n (-4)^{n-j} t(2n+1,2j+1) \left((x+1) (2x+1)\Delta(x+1,j) - x^2 \Delta(x,j) \right) \\ &\quad + (2n+1)^2 (x+1) (x+2) \cdots (x+2n) \\ &= (x+1) (2x+1) (x+2) \cdots (x+2n) + (2n+1)^2 (x+1) (x+2) \cdots (x+2n) \\ &= (x+1) (2x+1) (x+2) \cdots (x+2n) + (2n+1)^2 (x+1) (x+2) \cdots (x+2n) \\ &= (x+1) (x+2) \cdots (x+2n+1) (x+2n+2) . \end{split}$$

Hence, the identity is valid for k + 1.

References

- R. Barman and T. Komatsu, Lehmer's generalized Euler numbers in hypergeometric functions, J. Korean Math. Soc. 56 (2019), 485–505.
- [2] F. Brioschi, Sulle funzioni Bernoulliane ed Euleriane, Annali di Mat.,
 i. (1858), 260–263; Opere Mat., i. pp. 343–347.
- [3] P. J. Cameron, Some sequences of integers, Discrete Math. 75 (1989), 89–102.
- [4] R. Ernvall, Generalized Bernoulli numbers, generalized irregular primes, and class number, Ann. Univ. Turku Ser.A I 178 (1979), 1– 72.
- [5] F. G. Frobenius, Über die Bernoullischen Zahlen und die Eulerschen Polynome, in: Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin, 1910, pp. 809–847; also in: Gesammelte Abhandlungen III, Springer-Verlag, 1968, pp. 440–478.
- [6] J. W. L. Glaisher, Expressions for Laplace's coefficients, Bernoullian and Eulerian numbers etc. as determinants, Messenger (2) 6 (1875), 49-63.
- T. Komatsu, Complementary Euler numbers, Period. Math. Hungar. 75 (2017), 302–314.
- [8] T. Komatsu, Incomplete poly-Cauchy numbers, Monatsh. Math. 180 (2016), No. 2, 271–288.
- [9] T. Komatsu, K. Liptai and I. Mező, Incomplete poly-Bernoulli numbers associated with incomplete Stirling numbers, Publ. Math. Debr. 88 (2016), No. 3-4, 357-368.
- [10] T. Komatsu and R. K. Pandey, On hypergeometric Cauchy numbers of higher grade, AIMS Mathematics 6 (2021), Issue 7, 6630–6646.
- T. Komatsu and J. L. Ramirez, Some determinants involving incomplete Fubini numbers, An. Ştiinţ. Univ. "Ovidius" Constanţa Ser. Mat. 26 (2018), no.3, 143–170. DOI: 10.2478/auom-2018-0038

- T. Komatsu and H. Zhu, *Hypergeometric Euler numbers*, AIMS Math. 5 (2020), 1284–1303.
- [13] D. H. Lehmer, Lacunary recurrence formulas for the numbers of Bernoulli and Euler, Ann. of Math. (2), 36 (1935), 637–649.
- [14] G. Liu, On congruences of Euler numbers modulo powers of two, J. Number Theory 128 (2008), 3063–3071.
- [15] A. D. Loveless, A congruence for products of binomial coefficients modulo a composite, Integers 7 (2007), #A44.
- [16] E. Lucas, Théorie des Fonctions Numériques Simplement Périodiques, Amer. J. Math. 1 (1878), 184–196, 197–240, 289–321.
- [17] Q.-M. Luo, An explicit formula for the Euler numbers of higher order, Tamkang J. Math. 36 (2005), No. 4, 315–317.
- [18] T. Komatsu, J. L. Ramírez and D. Villamizar, A combinatorial approach to the Stirling numbers of the first kind with higher level, Studia Sci. Math. Hungar. 58 (2021), no.3, 293–307.
- [19] T. Komatsu, J. L. Ramírez and D. Villamizar, A combinatorial approach to the generalized central factorial numbers, Mediterr. J. Math. 18 (2021), no.5, Paper No. 192, 14 pp.
- [20] G.-D. Liu and W.-P. Zhang, Applications of an explicit formula for the generalized Euler numbers, Acta Math. Sin. (Eng. Ser.) 24 (2008), no.2, 343–352.
- [21] T. Muir, The theory of determinants in the historical order of development, Four volumes, Dover Publications, New York, 1960.
- [22] J. Riordan, Combinatorial identities, John Wiley & Sons, Inc., New York-London-Sydney, 1968.
- [23] N. J. A. Sloane, *The on-line encyclopedia of integer sequences*, available at oeis.org.
- [24] M. A. Stern, Zur Theorie der Eulerschen Zahlen, J. Reine Angew. Math. 79 (1875), 67–98.
- [25] Z.-W. Sun, On Euler numbers modulo powers of two, J. Number Theory 115(2) (2005), 371–380.

- [26] N. Trudi, Intorno ad alcune formole di sviluppo, Rendic. dell' Accad. Napoli (1862), 135–143.
- [27] S. S. Wagstaff Jr., Prime divisors of the Bernoulli and Euler numbers, in: Number Theory for the Millennium, III, Urbana, IL, 2000, A K Peters, Natick, MA, 2002, pp. 357–374.