

DISPROOF OF CLAIMS ABOUT PURE CUBIC FIELDS

SIHAM AOUISSI AND DANIEL C. MAYER

ABSTRACT. Let $\Gamma = \mathbb{Q}(\sqrt[3]{n})$ be a pure cubic field with normal closure $k = \mathbb{Q}(\sqrt[3]{n}, \zeta)$, where $n > 1$ denotes a cube free integer, and ζ is a primitive cube root of unity. Suppose k possesses an elementary bicyclic 3-class group $\text{Cl}_3(k)$, and the conductor of $k/\mathbb{Q}(\zeta)$ has the shape $f \in \{pq_1q_2, 3pq, 9pq\}$ where $p \equiv 1 \pmod{9}$ and $q, q_1, q_2 \equiv 2, 5 \pmod{9}$ are primes. It is disproved that there are only two possible capitulation types $\varkappa(k)$, either type a.1, (0000), or type a.2, (1000), as claimed by Ismaili and El Mesaoudi. Evidence is provided, theoretically and experimentally, of two further types, b.10, (0320), and d.23, (1320).

1. THREE ERRONEOUS CLAIMS

To exclude any misunderstanding from the very beginning, we postpone the introduction, and we clearly point out that Ismaili and El Mesaoudi state three erroneous theorems, Théorème 2 in section 3 on page 53, Théorème 3 in section 4 on page 56, and Théorème 4 in section 5 on page 57, in their paper [13]. Only for Théorème 2, an explicit proof is established in [13, pp. 53–55], which uses an incomplete version of a group theoretic theorem, Satz 14.17 on page 371, by Huppert [9], and consequently arrives at the erroneous claim. For Théorème 3, Ismaili and El Mesaoudi refer to the proof of Théorème 2, immediately after Théorème 3. For Théorème 4, they also refer to the proof of Théorème 2, immediately before Proposition 8 and Théorème 4. All three theorems concern the capitulation kernels, $\varkappa(k) = (\ker(T_i : \text{Cl}_3(k) \rightarrow \text{Cl}_3(K_i)))_{1 \leq i \leq 4}$, of the 3-class group $\text{Cl}_3(k) \simeq (\mathbb{Z}/3\mathbb{Z}) \times (\mathbb{Z}/3\mathbb{Z})$ of the normal closure $k = \mathbb{Q}(\sqrt[3]{n}, \zeta)$ of a pure cubic field $\Gamma = \mathbb{Q}(\sqrt[3]{n})$ in the four unramified cyclic cubic relative extensions K_1, \dots, K_4 of k . The conductors of $k/\mathbb{Q}(\zeta)$ are $f \in \{pq_1q_2, 3pq, 9pq\}$ with primes $p \equiv 1 \pmod{9}$ and $q, q_1, q_2 \equiv 2, 5 \pmod{9}$. All three theorems erroneously exclude two capitulation types $\varkappa(k) \in \{(1320), (0320)\}$, based on the incomplete use of Huppert's theorem, and state that only two capitulation types $\varkappa(k) \in \{(1000), (0000)\}$ are possible. In the particular situation where the auxiliary pure cubic field $\mathbb{Q}(\sqrt[3]{p})$ possesses a differential principal factorization (DPF) of type α [1, Thm. 2.1, p. 254], the capitulation is even reduced to the single type $\varkappa(k) = (0000)$.

Date: Tuesday, 20 February 2024.

2010 Mathematics Subject Classification. 11R16, 11R20, 11R29, 11R37, 20D15.


Key words and phrases. Pure cubic number fields, normal closure, relative conductor, multipliers, cubic residue symbols, differential principal factors, unramified cyclic cubic extensions, capitulation, 3-class field tower, finite 3-groups, elementary bicyclic commutator quotient, maximal subgroups, abelian quotient invariants, kernels and targets of Artin transfers.

Research of second author supported by the Austrian Science Fund (FWF): projects J0497-PHY, P26008-N25, and by the Research Executive Agency of the European Union (EUREA).

2. INTRODUCTION

In his Doctoral Thesis of 1992 [10], Ismaili determined the possible capitulation types of the normal closures of certain pure cubic fields. In cooperation with El Mesaoudi [13], the possible capitulation types were narrowed down in 2005. We show theoretically that some group theoretic arguments, used in the elimination process, were not justified, and the purging went too far. Our criticism is underpinned by experimental counter examples.

Let $n > 1$ be a cube free integer, and $\Gamma = \mathbb{Q}(\sqrt[3]{n})$ be the pure cubic field with radicand n , i.e., a non-Galois field of degree three. Denote by ζ a primitive cube root of unity, and let $k_0 = \mathbb{Q}(\zeta)$ be the third cyclotomic field. Then $k = \Gamma \cdot k_0$ is the normal closure of Γ , an absolutely dihedral field of degree six. In the last chapter of his Ph.D. Thesis [10, § 3.4, pp. 44–59], [3], Ismaili considers normal closures k of pure cubic fields Γ subject to the following conditions (see Figure 1):



- (1) the class number h_Γ of Γ is exactly divisible by 3 and the 3-class group of k is elementary bicyclic, $\text{Cl}_3(k) \simeq (\mathbb{Z}/3\mathbb{Z}) \times (\mathbb{Z}/3\mathbb{Z})$,
- (2) the relative genus field $(k/k_0)^*$ coincides with the Hilbert 3-class field $F_3^1(k)$ of k ,
- (3) the conductor of k/k_0 is of the shape $f \in \{3pq, 9pq, pq_1q_2\}$ where $p \equiv 1 \pmod{9}$ and $q, q_1, q_2 \equiv 2, 5 \pmod{9}$ are prime numbers.

Ismaili calls normal closures k satisfying (1) and (2) of *type III* [10, Dfn. 3.1, p. 31], and determines the class field theoretic 3-extensions, which are essential for the capitulation

problem, in the following way, if (3) is satisfied additionally. (We do not pay attention to the conductors $f = 9p$ in Théorème 2, section 3, page 53 of [13], and $f = pq$ with $q \equiv 8 \pmod{9}$ in Théorème 3, section 4, page 56, and we focus on Théorème 4, section 5, page 57.) If $p \equiv 1 \pmod{3}$ is a prime number, then p splits into a product $p = \pi_1\pi_2$ of two prime elements π_1 and π_2 in k_0 . If such a prime p divides the radicand n , then p ramifies in k and there exist prime ideals $\mathfrak{P}_1, \mathfrak{P}_2$ of k such that

$$(2.1) \quad \pi_1\mathcal{O}_k = \mathfrak{P}_1^3, \quad \pi_2\mathcal{O}_k = \mathfrak{P}_2^3, \quad \text{and} \quad p\mathcal{O}_k = \mathfrak{P}_1^3\mathfrak{P}_2^3.$$

Theorem 1. *The structure of all unramified abelian 3-extensions of k , viewed as Kummer extensions, is given by adjoining cube roots of the prime elements π_1, π_2 lying over the prime divisor $p \equiv 1 \pmod{9}$ of the conductor f : the relative genus field is bicyclic bicubic,*

$$(2.2) \quad (k/k_0)^* = F_3^1(k) = k(\sqrt[3]{\pi_1}, \sqrt[3]{\pi_2}),$$

and the four unramified cyclic cubic extensions of k are given by [10, Thm. 3.18, p. 58]

$$(2.3) \quad K_1 = k(\sqrt[3]{\pi_1\pi_2}), \quad K_2 = k(\sqrt[3]{\pi_2}), \quad K_3 = k(\sqrt[3]{\pi_1}), \quad \text{and} \quad K_4 = k(\sqrt[3]{\pi_1\pi_2^2}).$$

If the automorphism group $\text{Gal}(k/\Gamma) = \langle \tau \rangle$ is generated by τ , then $\pi_2 = \pi_1^\tau$, the extensions $K_2 = K_3^\tau$ are conjugate non-Galois, and the extensions K_1 and K_4 are Galois over \mathbb{Q} . The associated norm class groups $H_i = \text{Norm}_{K_i/k}\text{Cl}_3(K_i)$, $1 \leq i \leq 4$, are generated by the classes $[\mathfrak{P}_j]$ of prime ideals $\mathfrak{P}_1, \mathfrak{P}_2 = \mathfrak{P}_1^\tau$ lying over the prime elements π_1, π_2 [10, p. 51]:

$$(2.4) \quad H_1 = \langle [\mathfrak{P}_1\mathfrak{P}_2] \rangle, \quad H_2 = \langle [\mathfrak{P}_1] \rangle, \quad H_3 = \langle [\mathfrak{P}_2] \rangle, \quad \text{and} \quad H_4 = \langle [\mathfrak{P}_1\mathfrak{P}_2^2] \rangle.$$

In order to present Ismaili's results [10] on the capitulation of 3-classes of k in K_1, \dots, K_4 , i.e. the capitulation type $\varkappa(k)$ of k , some background must be developed in Section 3.

3. CAPITULATION KERNELS AND ABELIAN TYPE INVARIANTS

Since Scholz and Taussky [24, pp. 34–38] only considered *partial capitulation* with cyclic kernels of order 3, we use the capitulation types of Nebelung [22], which are summarized in [17, Tbl. 6–7, pp. 492–493]. The latter also admit *total capitulation* with bicyclic kernel of order 9. Let \mathcal{P}_K be the principal ideals and \mathcal{O}_K the integers of a number field K .

Definition 1. Let k be an algebraic number field with elementary bicyclic 3-class group $\text{Cl}_3(k) \simeq (\mathbb{Z}/3\mathbb{Z}) \times (\mathbb{Z}/3\mathbb{Z})$. Denote by K_1, \dots, K_4 the four unramified cyclic cubic extensions of k , and by $T_i : \text{Cl}_3(k) \rightarrow \text{Cl}_3(K_i)$, $\mathfrak{a}\mathcal{P}_k \mapsto (\mathfrak{a}\mathcal{O}_{K_i})\mathcal{P}_{K_i}$, the *class extension homomorphisms* (briefly *transfers*) from k to K_i . Then, the (unramified) *capitulation type* (or *transfer kernel type*, TKT) of k [18, Dfn. 1.1] is defined by $\varkappa(k) := (\varkappa_1, \dots, \varkappa_4)$ with

$$(3.1) \quad \varkappa_i := \begin{cases} j & \text{if } \ker(T_i) = \text{Norm}_{K_j/k}\text{Cl}_3(K_j), \text{ where } 1 \leq i, j \leq 4 \\ 0 & \text{if } \ker(T_i) = \text{Cl}_3(k). \end{cases}$$

Here, $\text{Cl}_3(k)$ is the 3-class group and \mathfrak{a} an ideal of k . According to Hilbert's Theorem 94, a transfer T_i cannot be injective, that is, its kernel $\ker(T_i)$ cannot be trivial.

Definition 2. Capitulation types $\varkappa = (\varkappa_1, \dots, \varkappa_4)$ and $\lambda = (\lambda_1, \dots, \lambda_4)$ are *partially ordered* by the declaration $\varkappa \leq \lambda : \iff [\varkappa_i \leq \lambda_i \text{ component wise for all } 1 \leq i \leq 4]$, where, for each $i = 1, \dots, 4$, we put $\varkappa_i \leq \lambda_i : \iff$ either $\varkappa_i = \lambda_i$ or $\lambda_i = 0$.

The unique maximal capitulation type is a.1, (0000), but any capitulation type without zero components is minimal, since it cannot shrink further, by Hilbert's Theorem 94. In the sequel, we are only concerned with four special capitulation types [17, pp. 492–493], subject to the following partial order. Note that the types (0320) and (1000) are incomparable.

$$(3.2) \quad \begin{array}{ccccc} & & & & (0000), \text{ a.1, maximal} \\ & & & \nearrow & \uparrow \\ & \text{b.10,} & (0320) & & (1000), \text{ a.2} \\ & & \uparrow & \nearrow & \\ & \text{d.23, minimal} & (1320) & & \end{array}$$

Definition 3. Let $\varkappa = (\varkappa_1, \dots, \varkappa_4)$ be a capitulation type. A *fixed point* of \varkappa is a component $\varkappa_i = i$, for some $1 \leq i \leq 4$. A *transposition* of \varkappa is a pair of distinct components such that $\varkappa_i = j$ and $\varkappa_j = i$, for some $1 \leq i \neq j \leq 4$.

The types a.2, (1000), and d.23, (1320), possess the fixed point 1. The types b.10, (0320), and d.23, (1320), contain a transposition ($2 \mapsto 3, 3 \mapsto 2$).

Now we are in the position to state Ismaili's results on the capitulation over normal closures k of pure cubic fields Γ of type III [10, § 3.4 (iii), Thm. 3.18, p. 58].

Theorem 2. *If k is of type III, there are four possibilities for the capitulation type,*

$$(3.3) \quad \varkappa(k) \in \{(1320), (0320), (1000), (0000)\},$$

because, for exponents $1 \leq a, b \leq 2$, the power product of prime ideals

$$(3.4) \quad \mathfrak{P}_1^a \mathfrak{P}_2^b \mathcal{O}_{K_i} = \sqrt[3]{\pi_1^a \pi_2^b} \mathcal{O}_{K_i} \text{ becomes principal for } i = \begin{cases} 1 & \text{if } a = b = 1, \\ 2 & \text{if } a = 0, b = 1, \\ 3 & \text{if } a = 1, b = 0, \\ 4 & \text{if } a = 1, b = 2, \end{cases}$$

that is, the transfer kernels contain at least the classes

$$(3.5) \quad [\mathfrak{P}_1 \mathfrak{P}_2] \in \ker(T_1), \quad [\mathfrak{P}_2] \in \ker(T_2), \quad [\mathfrak{P}_1] \in \ker(T_3), \quad \text{and} \quad [\mathfrak{P}_1 \mathfrak{P}_2^2] \in \ker(T_4),$$

and since the prime ideal \mathfrak{p} lying over p in Γ certainly becomes principal in the Hilbert 3-class field $F_3^1(\Gamma)$, the product $\mathfrak{P}_1 \mathfrak{P}_2 = \mathfrak{p} \mathcal{O}_k$ becomes principal in $K_4 = k \cdot F_3^1(\Gamma)$, whence $\ker(T_4) = \langle [\mathfrak{P}_1 \mathfrak{P}_2^2], [\mathfrak{P}_1 \mathfrak{P}_2] \rangle = \text{Cl}_3(k)$. Therefore, the minimal possible capitulation type is $\varkappa(k) = (1320)$, containing a fixed point and a transposition, and since K_2 and K_3 are conjugate, their transfer kernels may expand simultaneously only, that is, $\varkappa(k) = (1320)$ becomes $\varkappa(k) = (1000)$. However, if the fixed point expands, then two other capitulation types, $\varkappa(k) = (0320)$ and $\varkappa(k) = (0000)$, arise.

In the following corollary, which describes the *Galois structure* of the relative 3-genus field $(k/k_0)^*$, we use the notation in angle brackets $\langle \text{order, identifier} \rangle$ of the SmallGroups library [4] for finite groups of small order.

Corollary 1. *If k is of type III, the absolute Galois group of the Hilbert 3-class field of k is $G = \text{Gal}(F_3^1(k)/\mathbb{Q}) \simeq \langle 54, 13 \rangle$. Since the unramified cyclic cubic relative extensions K_2 and K_3 of k are absolutely non-Galois conjugate, their splitting field is the relative 3-genus field $(k/k_0)^* = F_3^1(k)$. In the setting of Theorem 2, the other two unramified cyclic cubic relative extensions K_1 and K_4 of k are absolutely Galois with non-abelian groups $\text{Gal}(K_1/\mathbb{Q}) \simeq \langle 18, 4 \rangle$ and $\text{Gal}(K_4/\mathbb{Q}) \simeq \langle 18, 3 \rangle$. The maximal abelian normal subgroup of G is the relative 3-genus group $\text{Gal}((k/k_0)^*/k_0) \simeq \langle 27, 5 \rangle$ which is elementary tricyclic.*

Proof. Among the 15 groups of order 54, $G = \text{Gal}((k/k_0)^*/\mathbb{Q}) \simeq \langle 54, 13 \rangle$ is the unique group with 4 triplets of conjugate subgroups $\langle 18, 3 \rangle$ of order 18 and a single self-conjugate subgroup $\langle 18, 4 \rangle$ of order 18. According to the Galois correspondence, these subgroups are associated with 4 triplets of conjugate pure cubic subfields, and a single cyclic cubic subfield with conductor p , of the relative 3-genus field $(k/k_0)^*$. For the Dedekind species II, $f = pq_1q_2$, the pure cubic subfields are one with conductor p , one with conductor $q_1 \cdot q_2$, and two with conductor $p \cdot q_1 \cdot q_2$. For the Dedekind species IB, $f = 3pq$, they are one with conductor p , one with conductor $3 \cdot q$, and two with conductor $3 \cdot p \cdot q$. For the Dedekind species IA, $f = 9pq$, they are one with conductor p , one with conductor $3 \cdot q$, and two with conductor $9 \cdot p \cdot q$. The group $G = \langle 54, 13 \rangle$ possesses a unique subgroup of order 27, and this subgroup is abelian, more precisely elementary tricyclic $\langle 27, 5 \rangle$. The extensions K_2 and K_3 are associated with a pair of conjugate subgroups $\langle 3, 1 \rangle$ of order 3, and the extensions K_1 and K_4 are associated with two self-conjugate subgroups $\langle 3, 1 \rangle$ of order 3. However, the quotients $\text{Gal}(K_1/\mathbb{Q}) \simeq \text{Gal}((k/k_0)^*/\mathbb{Q})/\text{Gal}((k/k_0)^*/K_1) \simeq \langle 18, 4 \rangle$ and $\text{Gal}(K_4/\mathbb{Q}) \simeq \text{Gal}((k/k_0)^*/\mathbb{Q})/\text{Gal}((k/k_0)^*/K_4) \simeq \langle 18, 3 \rangle$ are different. \square

While Ismaili's Theorems 1 and 2 are perfectly correct, the joint paper [13, Thm. 4, p. 57] of Ismaili and El Mesaoudi uses erroneous group theoretic arguments to narrow down the four types $\varkappa(k)$ to only two. In fact, the authors try to eliminate the two capitulation types b.10, (0320), and d.23, (1320), which contain a transposition $(2 \mapsto 3, 3 \mapsto 2)$.

This transposition can be characterized by the structure of the 3-class groups $\text{Cl}_3(K_i)$ of the corresponding extensions with $i \in \{2, 3\}$.

Definition 4. Let k be an algebraic number field with elementary bicyclic 3-class group $\text{Cl}_3(k) \simeq (\mathbb{Z}/3\mathbb{Z}) \times (\mathbb{Z}/3\mathbb{Z})$. Denote by K_1, \dots, K_4 the four unramified cyclic cubic extensions of k . Then, the *abelian type invariants* (ATI) of the four 3-class groups $\alpha(k) := (\text{Cl}_3(K_i))_{1 \leq i \leq 4}$ are called the *transfer target type* (TTT) of k [18, Dfn. 1.1]. Together with the transfer kernel type (TKT) they form the *Artin pattern* of k [21, Dfn. 4.3, p. 27],

$$(3.6) \quad \text{AP}(k) := [\alpha(k), \varkappa(k)] = [(\text{Cl}_3(K_i))_{i=1}^4, (\ker(T_i))_{i=1}^4].$$

Frequently, we shall write the ATI in logarithmic form, e.g. $(111) \triangleq (3, 3, 3)$, $(21) \triangleq (9, 3)$.

In 2009, we have determined the *abelian quotient invariants* (AQI) of the four maximal subgroups M_1, \dots, M_4 , that is, the ATI of the commutator quotients M_i/M_i' , $1 \leq i \leq 4$,

of all finite 3-groups G with elementary bicyclic G/G' [19]. In particular, we proved the following remarkable theorem, using absolute identifiers of the SmallGroups database [4].

Theorem 3. *The group $B := \langle 243, 3 \rangle$ has nilpotency class $\text{cl}(B) = 3$, coclass $\text{cc}(B) = 2$, soluble length $\text{sl}(B) = 2$, nuclear rank $\nu(B) = 2$, relation rank $d_2(B) = 4$, and Artin pattern*

$$(3.7) \quad \text{AP}(B) = [(21, 111, 111, 21), (0320)].$$

- (1) *Any finite 3-group G with $G/G' \simeq (3, 3)$ and one of the TKTs b.10, (0320), d.19, (2320), d.23, (1320), or d.25, (4320), is a descendant of B .*
- (2) *Any finite 3-group G with $G/G' \simeq (3, 3)$ and coclass $\text{cc}(G) \geq 3$ is descendant of B .*
- (3) *Any descendant of B shares with B the stable part of the Artin pattern, $[\alpha_0, \varkappa_0]$ with $\alpha_0 = (\dots, 111, 111, \dots)$ and $\varkappa_0 = (\dots 32 \dots)$, in particular, the conspicuous transposition $(2 \mapsto 3, 3 \mapsto 2)$.*

Proof. The bifurcation group B with nuclear rank $\nu(B) = 2$ is determined uniquely by its identifier in the SmallGroups database [4]. Its AQI were determined in [19, § 4.3, Tbl. 4.3, p. 434]. B is the smallest group with TKT b.10, (0320). The types d.19, (2320), d.23, (1320), and d.25, (4320), arise from type b.10, (0320), by shrinking of the *polarization* (first component). The AQI of the infinitely many descendants of B were determined in [19, § 4.4, Tbl. 4.5, p. 438] and [19, § 4.5, Tbl. 4.7, p. 441] for coclass $\text{cc}(G) = 2$, and in [19, § 4.6, Thm. 4.5, pp. 444–445] for the bifurcation to coclass $\text{cc}(G) \geq 3$. The statement concerning coclass $\text{cc}(G) \geq 3$ is due to Nebelung [22], who discovered the crucial bifurcation from coclass 2 to coclass 3. Since the components (32) of the TKT cannot shrink, according to Hilbert's Theorem 94, they remain stable for any descendant of $\langle 243, 3 \rangle$. Accordingly, the corresponding components (111, 111) of the TTT (AQI) cannot expand and form the *stabilization* of the Artin pattern. \square

4. LITTLE AND BIG TWO-STAGE TOWERS

Definition 5. Let k be an algebraic number field with elementary bicyclic 3-class group $\text{Cl}_3(k) \simeq (\mathbb{Z}/3\mathbb{Z}) \times (\mathbb{Z}/3\mathbb{Z})$. Denote by K_1, \dots, K_4 the four unramified cyclic cubic extensions of k . Then, the automorphism groups $\mathfrak{G}_i := \text{Gal}(\mathbb{F}_3^1(K_i)/k)$, $1 \leq i \leq 4$, of the first Hilbert 3-class fields of K_i are called the groups of the four *little two-stage towers* of k , and the automorphism group $G := \text{Gal}(\mathbb{F}_3^2(k)/k)$ of the second Hilbert 3-class field of k is called the group of the *big two-stage tower* of k .

Theorem 4. *Let the assumptions in Definition 5 be satisfied. Generally, the connection between groups of little and big two-stage towers is given by*

$$(4.1) \quad \mathfrak{G}_i \simeq G/M'_i, \quad \mathfrak{A}_i \simeq \text{Cl}_3(K_i) \simeq M_i/M'_i$$

where M_1, \dots, M_4 denote the four maximal subgroups of G , and $\mathfrak{A}_i := \text{Gal}(\mathbb{F}_3^1(K_i)/K_i)$ is the distinguished abelian maximal subgroup of \mathfrak{G}_i , for $1 \leq i \leq 4$. The commutator quotients $G/G' \simeq \mathfrak{G}_i/\mathfrak{G}'_i \simeq \text{Cl}_3(k) \simeq (\mathbb{Z}/3\mathbb{Z}) \times (\mathbb{Z}/3\mathbb{Z})$ are isomorphic, here elementary bicyclic.

In particular, if K_i has 3-class group $\text{Cl}_3(K_i) \simeq (\mathbb{Z}/3\mathbb{Z}) \times (\mathbb{Z}/3\mathbb{Z}) \times (\mathbb{Z}/3\mathbb{Z})$, the elementary tricyclic 3-group, for some $i = 1, \dots, 4$, then the little two-stage tower has the group with SmallGroups identifier [4]

$$(4.2) \quad \mathfrak{G}_i \simeq \langle 81, 7 \rangle \simeq \text{Syl}_3(A_9).$$

Proof. For the proof of the general statements, we refer to [19, § 2, Prop. 2.1, pp. 417–419]. When the unramified cyclic cubic extension K_i/k possesses an elementary tricyclic 3-class group $\text{Cl}_3(K_i) \simeq (3, 3, 3)$, then $[\mathbb{F}_3^1(K_i) : K_i] = 3^3 = 27$ is the relative degree of its Hilbert 3-class field, and the group \mathfrak{G}_i of the little two-stage tower is of maximal class and order $\#\mathfrak{G}_i = [\mathbb{F}_3^1(K_i) : K_i] \cdot [K_i : k] = 3^3 \cdot 3 = 81$, according to [8, Satz 4, Korollar, p. 9], because $\text{Cl}_3(k) \simeq (3, 3)$. Since the distinguished abelian maximal subgroup $\mathfrak{A}_i = \text{Gal}(\mathbb{F}_3^1(K_i)/K_i) \simeq \text{Cl}_3(K_i)$ of \mathfrak{G}_i is elementary tricyclic, \mathfrak{G}_i is uniquely determined as $\langle 81, 7 \rangle \simeq \text{Syl}_3(A_9)$, the Sylow 3-subgroup of the alternating group A_9 of degree 9, according to [19, § 4.1, Thm. 4.1, p. 427]. \square

5. THEORETICAL DISPROOF OF THE CLAIMS

We state the incorrect hypothesis [13, Thm. 4, p. 57] in a form which is more detailed than in the original paper. However, we exclude that $q_i \equiv 8 \pmod{9}$, for $i = 1$ or $i = 2$.

Hypothesis 1. Let $k = \mathbb{Q}(\sqrt[3]{n}, \zeta)$ be the normal closure of a pure cubic number field $\Gamma = \mathbb{Q}(\sqrt[3]{n})$. Suppose the cubefree radicand n is a positive integer with one of the following decompositions in prime numbers $p \equiv 1 \pmod{9}$ and $q, q_1, q_2 \equiv 2, 5 \pmod{9}$:

$$(5.1) \quad \begin{aligned} n &= p^e q_1^{e_1} q_2^{e_2} \equiv \pm 1 \pmod{9} \text{ with integers } e, e_1, e_2 \in \{1, 2\} \text{ (species II),} \\ n &= p^{e_1} q^{e_2} \not\equiv \pm 1 \pmod{9} \text{ with integers } e_1, e_2 \in \{1, 2\} \text{ (species IB),} \\ n &= 3^e p^{e_1} q^{e_2} \text{ with integers } e, e_1, e_2 \in \{1, 2\} \text{ (species IA).} \end{aligned}$$

Assume the 3-class group of k is elementary bicyclic, $\text{Cl}_3(k) \simeq (\mathbb{Z}/3\mathbb{Z}) \times (\mathbb{Z}/3\mathbb{Z})$, the 3-class group of Γ is elementary cyclic, $\text{Cl}_3(\Gamma) \simeq \mathbb{Z}/3\mathbb{Z}$, and the relative 3-genus field $(k/k_0)^*$ coincides with the Hilbert 3-class field $\mathbb{F}_3^1(k)$, that is, k is of Ismaili's type III, as drawn in Figure 1. Then there are **only two** possibilities for the capitulation type of k ,

$$(5.2) \quad \varkappa(k) \in \{(1000), (0000)\},$$

in the four unramified cyclic cubic relative extensions K_1, \dots, K_4 of k . If the index of the subgroup U_0 generated by all subfield units in the group $U(\Gamma_p)$ of the pure cubic auxiliary field $\Gamma_p = \mathbb{Q}(\sqrt[3]{p})$ is equal to $(U(\Gamma_p) : U_0) = 1$, that is, Γ_p is of differential principal factorization type α , then **only the single** capitulation type $\varkappa(k) = (0000)$ is possible.

Warning 1. Ismaili and El Mesaoudi conducted the proof of [13, § 5, Thm. 4, p. 57] simultaneously with the proofs of [13, § 4, Thm. 3, p. 56] and [13, § 3, Thm. 2, p. 53] in [13, § 3, pp. 53–55], making use of a theorem of Huppert which is cited incompletely as [13, § 2, Prop. 4, p. 51]. Although we only found explicit numerical counter examples to [13, § 5, Thm. 4, p. 57] in Section 6, it must be pointed out that none of the above mentioned Theorems 2,3,4 in [13] was proved reliably, and violations of Theorems 2,3 could still exist.

Now we explain how the erroneous Hypothesis 1 can be disproved theoretically. For this purpose, we first state the correct form of the group theoretic result which was used by Ismaili and El Mesaoudi to derive the incorrect Hypothesis 1.

Let p be a prime number and G be a finite p -group with lower central series $\gamma_2 G = [G, G]$ and $\gamma_i G = [\gamma_{i-1} G, G]$ for $i \geq 3$. Denote by $\gamma_1 G$ the *two-step centralizer* of G such that $\gamma_1 G / \gamma_4 G$ is the centralizer of $\gamma_2 G / \gamma_4 G$ in $G / \gamma_4 G$. The correct form of [13, § 2, Prop. 4, p. 51] is the following theorem by Huppert [9, Satz 14.17, p. 371].

Theorem 5. *If G is a 3-group of maximal nilpotency class **with order** $\#G = 3^n$, $n > 4$, **that is**, $\#G \geq 243$, then G is metabelian, i.e., $G'' = 1$, and $\gamma_1 G$ is a metacyclic group with nilpotency class $\text{cl}(\gamma_1 G) \leq 2$.*

Proof. This theorem is [9, Satz 14.17, p. 371]. Its proof makes use of [9, Satz 14.16, p. 370], where statements about a p -group G of maximal nilpotency class **with order** $\#G = p^n$, $n > p + 1$, are proved, **that is**, $n > 3 + 1 = 4$, **if** $p = 3$. \square

Now we state and prove the correct theorem which must replace the erroneous Hypothesis 1. We prefer to express the ramification in terms of the conductor f rather than the radicand n with cumbersome and useless exponents e, e_1, e_2 .

Theorem 6. *Let $k = \mathbb{Q}(\sqrt[3]{n}, \zeta)$ be the normal closure of a pure cubic number field $\Gamma = \mathbb{Q}(\sqrt[3]{n})$. Suppose the conductor f of k/k_0 is a positive integer with one of the following decompositions in prime numbers $p \equiv 1 \pmod{9}$ and $q, q_1, q_2 \equiv 2, 5 \pmod{9}$:*

$$(5.3) \quad \begin{aligned} f &= pq_1q_2 \text{ (species II),} \\ f &= 3pq \text{ (species IB),} \\ f &= 9pq \text{ (species IA).} \end{aligned}$$

Assume the 3-class group of k is elementary bicyclic, $\text{Cl}_3(k) \simeq (\mathbb{Z}/3\mathbb{Z}) \times (\mathbb{Z}/3\mathbb{Z})$, the 3-class group of Γ is elementary cyclic, $\text{Cl}_3(\Gamma) \simeq \mathbb{Z}/3\mathbb{Z}$, and the relative 3-genus field $(k/k_0)^$ coincides with the Hilbert 3-class field $F_3^1(k)$, that is, k is of Ismaili's type III, as drawn in Figure 1. Then, up to equivalence, there are **four** possibilities for the capitulation type of k in the four unramified cyclic cubic relative extensions K_1, \dots, K_4 of k , either*

$$(5.4) \quad \varkappa(k) \in \{(1000), (0000)\},$$

if the unit norm index is $(U(k) : \text{Norm}_{K_i/k} U(K_i)) = 3$, for $i \in \{2, 3\}$, or

$$(5.5) \quad \varkappa(k) \in \{(1320), (0320)\},$$

if the unit norm index is $(U(k) : \text{Norm}_{K_i/k} U(K_i)) = 1$, for $i \in \{2, 3\}$.

*If the index of the subgroup U_0 generated by all subfield units in the group $U(\Gamma_p)$ of the pure cubic auxiliary field $\Gamma_p = \mathbb{Q}(\sqrt[3]{p})$ is equal to $(U(\Gamma_p) : U_0) = 1$, that is, Γ_p is of DPF-type α , then **only two** capitulation types are possible, depending on the unit norm index, either $\varkappa(k) = (0000)$, if $(U(k) : \text{Norm}_{K_i/k} U(K_i)) = 3$, for $i = 2, 3$, or $\varkappa(k) = (0320)$, if $(U(k) : \text{Norm}_{K_i/k} U(K_i)) = 1$, for $i = 2, 3$.*

Proof. (of Theorem 6 by disproof of Hypothesis 1.) Ismaili and El Mesaoudi try to discourage the capitulation types b.10, (0320), and d.23, (1320), by proving that in Formula (3.5) not only $[\mathfrak{P}_2] \in \ker(T_2)$ but also $[\mathfrak{P}_1] \in \ker(T_2)$ [13, § 3, Proof of Thm. 2, item (1), p. 53], and thus $\ker(T_3) \simeq \ker(T_2) = \langle [\mathfrak{P}_1], [\mathfrak{P}_2] \rangle = \text{Cl}_3(k)$. They use contraposition, trying to derive a contradiction from the assumption that $[\mathfrak{P}_1] \notin \ker(T_2)$ [13, § 3, Proof of Thm. 2, item (1), p. 54], which implies $\text{Cl}_3(K_3) \simeq \text{Cl}_3(K_2) \simeq (3, 3, 3)$, according to Theorem 3 and Formula (3.7), and consequently $\mathfrak{G}_2 \simeq \mathfrak{G}_3 \simeq \langle 81, 7 \rangle \simeq \text{Syl}_3(A_9)$ with $\#\mathfrak{G}_2 = \#\mathfrak{G}_3 = 81$, according to Theorem 4, Formula (4.2).

Then, Ismaili and El Mesaoudi **apply the incomplete theorem of Huppert** [13, § 2, Prop. 4, p. 51], without the crucial condition $\#G \geq 243$, to the groups \mathfrak{G}_2 and \mathfrak{G}_3 of the little two-stage towers [13, § 3, Proof of Thm. 2, item (1)(iv), p. 55], and conclude that the two-step centralizers $\mathfrak{A}_2 = \gamma_1 \mathfrak{G}_2$ and $\mathfrak{A}_3 = \gamma_1 \mathfrak{G}_3$ must be metacyclic, in contradiction to $\mathfrak{A}_2 \simeq \mathfrak{A}_3 \simeq (3, 3, 3)$. However, the correct theorem of Huppert cannot be applied to a group of order $3^4 = 81$, and thus **no contradiction arises** when the capitulation kernel $\ker(T_2)$ is cyclic of order 3, generated by $[\mathfrak{P}_2]$, i.e., $\varkappa(k) = (\varkappa_1 320)$ with $\varkappa_1 \in \{0, 1\}$.

Finally, the well-known Galois-cohomological theorem on the Herbrand quotient of the *unramified* cyclic relative extension K_i/k of *odd* prime degree $[K_i : k] = 3$, $\#\ker(T_i) = \#H^1(\langle \sigma_i \rangle, U(K_i)) = [K_i : k] \cdot \#\hat{H}^0(\langle \sigma_i \rangle, U(K_i)) = 3 \cdot (U(k) : \text{Norm}_{K_i/k} U(K_i))$, for $1 \leq i \leq 4$, where $\langle \sigma_i \rangle = \text{Gal}(K_i/k) \simeq \mathbb{Z}/3\mathbb{Z}$, admits the distinction between total and partial capitulation in K_2 and K_3 :

$\varkappa(k) \in \{(1000), (0000)\}$ if and only if $(U(k) : \text{Norm}_{K_i/k} U(K_i)) = 3$, for $i = 2, 3$, and
 $\varkappa(k) \in \{(1320), (0320)\}$ if and only if $(U(k) : \text{Norm}_{K_i/k} U(K_i)) = 1$, for $i = 2, 3$. \square

6. EXPERIMENTAL DISPROOF OF THE CLAIMS

In this concluding section, we underpin our theoretical disproof of the claims by Ismaili and El Mesaoudi [13, Thm. 4, p. 57] by extensive numerical counter examples under the GRH, compiled with the aid of class field theoretic routines by Fieker [7], which are implemented in the computer algebra system Magma [5], [6], [14]. Since the normal closures $k = \mathbb{Q}(\sqrt[3]{n}, \zeta)$ of pure cubic fields $\Gamma = \mathbb{Q}(\sqrt[3]{n})$ are constructed as 3-ring class fields modulo 3-admissible conductors f , they are automatically produced as *multiplets* (k_1, \dots, k_m) of pairwise non-isomorphic fields sharing the common conductor f with multiplicities $m = m(f)$ determined by the formulas in [16, Thm. 2.1, p. 833] or [20, Cor. 3.2, p. 2219]. In Table 1–3, we show that Théorème 4 is violated by conductors $f = pq_1q_2$ of Dedekind species II with primes $p \equiv 1 \pmod{9}$ and various combinations of $q_1, q_2 \equiv 2, 5 \pmod{9}$. Columns begin with prime factors p, q_1, q_2 and conductor f , continue with normalized radicand n_i , Artin pattern consisting of capitulation type (CT) \varkappa_i and log abelian type invariants α_i , and a minimal differential principal factor (DPF) (norm of an ambiguous principal ideal), split into m rows, according to the multiplicity $m = m(f)$ of the conductor, and end with the (DPF-)type of the pure cubic auxiliary field $\mathbb{Q}(\sqrt[3]{p})$, which decides whether the \varkappa_i may be of CT d.23 (for type γ) or not (for type α). The radicand $n_i = a \cdot b^2$ is called *normalized*, if $1 \leq b < a$ are square free coprime integers.

6.1. Counterexamples of Dedekind species II. First, we list lots of counterexamples of Dedekind species II with conductors of the shape $3 \nmid f = pq_1q_2$, giving rise to *doublets*, in Table 1, 2, and 3.

Only in the mixed case $q_1 \equiv 2 \pmod{9}$, $q_2 \equiv 5 \pmod{9}$ of Table 3 with most numerous hits, there occurs a single instance of capitulation type (CT) d.23, $\varkappa \sim (1320)$, because the *pure cubic auxiliary field* $\mathbb{Q}(\sqrt[3]{p})$ is of DPF type γ . The primes $p \equiv 1 \pmod{9}$ with type γ are collected in the sequence A363717 of the On-line Encyclopedia of Integer Sequences (OEIS) [23].

TABLE 1. Doublets, $m = 2$, with $f = pq_1q_2$, $q_1 \equiv 2 \pmod{9}$, $q_2 \equiv 2 \pmod{9}$

p	q_1	q_2	f	n_i	CT	\varkappa_i	α_i	DPF	Type
19	2	11	418	$836 = 2^2 \cdot 11 \cdot 19$	b.10	4001	111, 21, 21, 111	11	α
				$4598 = 2 \cdot 11^2 \cdot 19$	b.10	4001	111, 21, 21, 111	11	
19	2	83	3154	$119852 = 2^2 \cdot 19^2 \cdot 83$	b.10	3010	111, 21, 111, 21	$38 = 2 \cdot 19$	α
				$6308 = 2^2 \cdot 19 \cdot 83$	b.10	0043	21, 21, 111, 111	$76 = 2^2 \cdot 19$	
37	2	11	814	$8954 = 2 \cdot 11^2 \cdot 37$	b.10	3010	111, 21, 111, 21	11	α
				$1628 = 2^2 \cdot 11 \cdot 37$	b.10	0043	21, 21, 111, 111	11	
37	2	29	2146	$4292 = 2^2 \cdot 29 \cdot 37$	b.10	0320	21, 111, 111, 21	29	α
				$62234 = 2 \cdot 29^2 \cdot 37$	b.10	3010	111, 21, 111, 21	29	
37	2	47	3478	$6956 = 2^2 \cdot 37 \cdot 47$	b.10	3010	111, 21, 111, 21	47	α
				$163466 = 2 \cdot 37 \cdot 47^2$	b.10	3010	111, 21, 111, 21	47	
109	2	11	2398	$26378 = 2 \cdot 11^2 \cdot 109$	b.10	0402	21, 111, 21, 111	2	α
				$4796 = 2^2 \cdot 11 \cdot 109$	b.10	0320	21, 111, 111, 21	2	
127	2	11	2794	$5588 = 2^2 \cdot 11 \cdot 127$	b.10	0320	21, 111, 111, 21	2	α
				$30734 = 2 \cdot 11^2 \cdot 127$	b.10	3010	111, 21, 111, 21	2	
199	2	11	4378	$8756 = 2^2 \cdot 11 \cdot 199$	b.10	0402	32, 111, 22, 111	11	α
				$48158 = 2 \cdot 11^2 \cdot 199$	b.10	0320	32, 111, 111, 22	11	

TABLE 2. Doublets, $m = 2$, with $f = pq_1q_2$, $q_1 \equiv 5 \pmod{9}$, $q_2 \equiv 5 \pmod{9}$

p	q_1	q_2	f	n_i	CT	\varkappa_i	α_i	DPF	Type
37	23	5	4255	$97865 = 5^2 \cdot 23^2 \cdot 37$	b.10	3010	111, 21, 111, 21	23	α
				$21275 = 5^2 \cdot 23 \cdot 37$	b.10	3010	111, 21, 111, 21	23	

The conditions for the 3-class groups of k and Γ and the *class number formula* enforce that all fields k are of DPF-type β with *non-radical absolute* principal factors. The DPF in the tables is always a *cubic residue* modulo the prime p .

TABLE 3. Doublets, $m = 2$, with $f = pq_1q_2$, $q_1 \equiv 2 \pmod{9}$, $q_2 \equiv 5 \pmod{9}$

p	q_1	q_2	f	n_i	CT	\varkappa_i	α_i	DPF	Type
19	2	113	4294	$81586 = 2 \cdot 19^2 \cdot 113$	b.10	0043	21, 21, 111, 111	$76 = 2^2 \cdot 19$	α
				$4294 = 2 \cdot 19 \cdot 113$	b.10	3010	111, 21, 111, 21	$38 = 2 \cdot 19$	
19	11	5	1045	$1045 = 5 \cdot 11 \cdot 19$	b.10	3010	111, 21, 111, 21	11	α
				$19855 = 5 \cdot 11 \cdot 19^2$	b.10	4001	111, 21, 21, 111	11	
19	11	23	4807	$4807 = 11 \cdot 19 \cdot 23$	b.10	4001	111, 21, 21, 111	11	α
				$91333 = 11 \cdot 19^2 \cdot 23$	b.10	3010	111, 21, 111, 21	11	
37	2	23	1702	$1702 = 2 \cdot 23 \cdot 37$	b.10	3010	111, 21, 111, 21	23	α
				$62974 = 2 \cdot 23 \cdot 37^2$	b.10	4001	111, 21, 21, 111	23	
37	11	5	2035	$2035 = 5 \cdot 11 \cdot 37$	b.10	4001	111, 21, 21, 111	11	α
				$75295 = 5 \cdot 11 \cdot 37^2$	b.10	3010	111, 21, 111, 21	11	
37	29	5	5365	$5365 = 5 \cdot 29 \cdot 37$	b.10	0043	21, 21, 111, 111	29	α
				$198505 = 5 \cdot 29 \cdot 37^2$	b.10	3010	111, 21, 111, 21	29	
109	2	5	1090	$10900 = 2^2 \cdot 5^2 \cdot 109$	b.10	0320	21, 111, 111, 21	2	α
				$1090 = 2 \cdot 5 \cdot 109$	b.10	0043	21, 21, 111, 111	2	
163	2	5	1630	$1630 = 2 \cdot 5 \cdot 163$	b.10	4001	111, 21, 21, 111	5	α
				$16300 = 2^2 \cdot 5^2 \cdot 163$	b.10	0402	21, 111, 21, 111	5	
181	2	5	1810	$1810 = 2 \cdot 5 \cdot 181$	b.10	3010	111, 21, 111, 21	5	α
				$18100 = 2^2 \cdot 5^2 \cdot 181$	b.10	0043	21, 21, 111, 111	5	
199	2	5	1990	$1990 = 2 \cdot 5 \cdot 199$	b.10	4001	111, 32, 22, 111	5	α
				$19900 = 2^2 \cdot 5^2 \cdot 199$	b.10	4001	111, 32, 22, 111	5	
307	2	5	3070	$3070 = 2 \cdot 5 \cdot 307$	b.10	0043	21, 21, 111, 111	2	α
				$30700 = 2^2 \cdot 5^2 \cdot 307$	b.10	4001	111, 21, 21, 111	2	
379	2	5	3790	$3790 = 2 \cdot 5 \cdot 379$	b.10	4001	111, 21, 21, 111	5	α
				$37900 = 2^2 \cdot 5^2 \cdot 379$	b.10	4001	111, 21, 21, 111	5	
397	2	5	3970	$39700 = 2^2 \cdot 5^2 \cdot 397$	b.10	0043	21, 21, 111, 111	2	α
				$3970 = 2 \cdot 5 \cdot 397$	b.10	0320	21, 111, 111, 21	2	
433	2	5	4330	$4330 = 2 \cdot 5 \cdot 433$	b.10	0320	21, 111, 111, 21	2	α
				$43300 = 2^2 \cdot 5^2 \cdot 433$	b.10	0320	21, 111, 111, 21	2	
487	2	5	4870	$48700 = 2^2 \cdot 5^2 \cdot 487$	b.10	0043	22, 32, 111, 111	5	α
				$4870 = 2 \cdot 5 \cdot 487$	b.10	3010	111, 32, 111, 22	5	
523	2	5	5230	$5230 = 2 \cdot 5 \cdot 523$	b.10	3010	111, 32, 111, 21	$2^2 \cdot 5 = 20$	α
				$52300 = 2^2 \cdot 5^2 \cdot 523$	b.10	3010	111, 32, 111, 21	$2^2 \cdot 5 = 20$	
541	2	5	5410	$5410 = 2 \cdot 5 \cdot 541$	d.23	0243	21, 33, 111, 111	$2^2 \cdot 5 = 20$	γ
				$54100 = 2^2 \cdot 5^2 \cdot 541$	d.23	3210	111, 33, 111, 21	$2^2 \cdot 5 = 20$	

6.2. **Counterexamples of Dedekind species IB.** Second, we provide some counterexamples of Dedekind species IB with conductors of the shape $3 \parallel f = 3pq$, giving rise to *doublets*, in Table 4 and 5.

TABLE 4. Doublets, $m = 2$, with $f = 3pq$, $q \equiv 2 \pmod{9}$

p	q	f	n_i	CT	\varkappa_i	α_i	DPF	Type
73	2	438	$292 = 2^2 \cdot 73$	b.10	4001	111, 21, 21, 111	3	α
			$146 = 2 \cdot 73$	b.10	0043	21, 21, 111, 111	3	
73	11	2409	$8833 = 11^2 \cdot 73$	b.10	0043	21, 21, 111, 111	3	α
			$803 = 11 \cdot 73$	b.10	3010	111, 21, 111, 21	3	
199	2	1194	$398 = 2 \cdot 199$	b.10	0402	32, 111, 21, 111	$12 = 2^2 \cdot 3$	α
			$796 = 2^2 \cdot 199$	b.10	0043	21, 32, 111, 111	$12 = 2^2 \cdot 3$	
271	2	1626	$542 = 2 \cdot 271$	b.10	0402	32, 111, 22, 111	3	α
			$1084 = 2^2 \cdot 271$	b.10	0402	32, 111, 22, 111	3	
487	2	2922	$1948 = 2^2 \cdot 487$	b.10	0043	21, 32, 111, 111	$12 = 2^2 \cdot 3$	α
			$974 = 2 \cdot 487$	b.10	0043	21, 32, 111, 111	$12 = 2^2 \cdot 3$	
523	2	3138	$1046 = 2 \cdot 523$	b.10	3010	111, 21, 111, 21	3	α
			$2092 = 2^2 \cdot 523$	b.10	0043	21, 21, 111, 111	3	
541	2	3246	$2164 = 2^2 \cdot 541$	d.23	0243	21, 33, 111, 111	$6 = 3 \cdot 2$	γ
			$1082 = 2 \cdot 541$	d.23	3210	111, 33, 111, 21	$6 = 3 \cdot 2$	
577	2	3462	$1154 = 2 \cdot 577$	b.10	3010	111, 21, 111, 21	3	α
			$2308 = 2^2 \cdot 577$	b.10	0043	21, 21, 111, 111	3	
613	2	3678	$2452 = 2^2 \cdot 613$	b.10	4001	111, 21, 21, 111	3	α
			$1226 = 2 \cdot 613$	b.10	4001	111, 21, 21, 111	3	

TABLE 5. Doublets, $m = 2$, with $f = 3pq$, $q \equiv 5 \pmod{9}$

p	q	f	n_i	CT	\varkappa_i	α_i	DPF	Type
73	5	1095	$365 = 5 \cdot 73$	b.10	0043	21, 21, 111, 111	3	α
			$1825 = 5^2 \cdot 73$	b.10	4001	111, 21, 21, 111	3	
541	5	8115	$13525 = 5^2 \cdot 541$	d.23	0243	21, 33, 111, 111	$15 = 3 \cdot 5$	γ
			$2705 = 5 \cdot 541$	d.23	3210	111, 33, 111, 21	$15 = 3 \cdot 5$	

6.3. Counterexamples of Dedekind species IA. Next, we list lots of counterexamples of Dedekind species IA with conductors of the shape $9 \mid f = 9pq$, giving rise to *quartets*, in Table 6 and 7. A special remark is due to the conductor $f = 3582$: Here, two members of the quartet have irregular 3-class groups of rank 3, namely $(9, 9, 3)$ and $(9, 3, 3)$, the former with radicand $2388 = 2^2 \cdot 3 \cdot 199$ and DPF-type α , the latter with radicand $3582 = 2 \cdot 3^2 \cdot 199$, DPF-type β , and DPF $12 = 2^2 \cdot 3$.

TABLE 6. Quartets, $m = 4$, with $f = 9pq$, $q \equiv 2 \pmod{9}$

p	q	f	n_i	CT	\varkappa_i	α_i	DPF	Type
73	2	1314	$1314 = 2 \cdot 3^2 \cdot 73$	b.10	0320	21, 111, 111, 21	3	α
			$876 = 2^2 \cdot 3 \cdot 73$	b.10	3010	111, 21, 111, 21	3	
			$2628 = 2^2 \cdot 3^2 \cdot 73$	b.10	0402	21, 111, 21, 111	3	
			$438 = 2 \cdot 3 \cdot 73$	b.10	4001	111, 21, 21, 111	3	
109	2	1962	$1308 = 2^2 \cdot 3 \cdot 109$	b.10	4001	111, 21, 21, 111	2	α
			$1962 = 2 \cdot 3^2 \cdot 109$	b.10	0043	21, 21, 111, 111	2	
			$654 = 2 \cdot 3 \cdot 109$	b.10	0043	21, 21, 111, 111	2	
			$3924 = 2^2 \cdot 3^2 \cdot 109$	b.10	3010	111, 21, 111, 21	2	
127	2	2286	$1524 = 2^2 \cdot 3 \cdot 127$	b.10	0320	21, 111, 111, 21	2	α
			$2286 = 2 \cdot 3^2 \cdot 127$	b.10	0320	21, 111, 111, 21	2	
			$762 = 2 \cdot 3 \cdot 127$	b.10	0043	21, 21, 111, 111	2	
			$4572 = 2^2 \cdot 3^2 \cdot 127$	b.10	0043	21, 21, 111, 111	2	
199	2	3582	$1194 = 2 \cdot 3 \cdot 199$	b.10	3010	111, 32, 111, 21	$12 = 2^2 \cdot 3$	α
			$7164 = 2^2 \cdot 3^2 \cdot 199$	b.10	0402	32, 111, 21, 111	$12 = 2^2 \cdot 3$	
271	2	4878	$3252 = 2^2 \cdot 3 \cdot 271$	b.10	3010	111, 32, 111, 22	3	α
			$4878 = 2 \cdot 3^2 \cdot 271$	b.10	4001	111, 32, 22, 111	3	
			$1662 = 2 \cdot 3 \cdot 271$	b.10	0043	22, 32, 111, 111	3	
			$9756 = 2^2 \cdot 3^2 \cdot 271$	b.10	4001	111, 32, 22, 111	3	
19	11	1881	$1881 = 3^2 \cdot 11 \cdot 19$	b.10	4001	111, 21, 21, 111	11	α
			$6897 = 3 \cdot 11^2 \cdot 19$	b.10	3010	111, 21, 111, 21	11	
			$11913 = 3 \cdot 11 \cdot 19^2$	b.10	0043	21, 21, 111, 111	11	
			$627 = 3 \cdot 11 \cdot 19$	b.10	3010	111, 21, 111, 21	11	
37	11	3663	$13431 = 3 \cdot 11^2 \cdot 37$	b.10	4001	111, 21, 21, 111	11	α
			$3663 = 3^2 \cdot 11 \cdot 37$	b.10	4001	111, 21, 21, 111	11	
			$1221 = 3 \cdot 11 \cdot 37$	b.10	3010	111, 21, 111, 21	11	
			$40293 = 3^2 \cdot 11^2 \cdot 37$	b.10	3010	111, 21, 111, 21	11	

6.4. Prototypes of counterexamples. Finally, we present the *prototypes* of counterexamples with minimal conductors f . When several non-isomorphic fields share a common conductor they are collected in a *multiplet* (doublet or quartet), according to [16, Thm. 2.1, p. 833, Exm. 1, p. 840], [20, Cor. 3.2, p. 2219], [1, Thm. 2.2, p. 255]. For the prime divisor $p \equiv 1 \pmod{9}$ of the conductor f , the *differential principal factorization type*, [1, Thm. 2.1, p. 254], [2], of $\mathbb{Q}(\sqrt[3]{p})$, necessarily different from type β , is checked. Since $p = 541$ is the smallest prime with type γ , the type is α , for all primes $p < 541$, which prohibits a fixed point capitulation at the first component, by [13, § 3, Proof of Thm. 2, item (3), p. 55].

We go further than Ismaili and El Mesaoudi by determining the group $G = \text{Gal}(F_3^\infty(k)/k)$ of the maximal unramified pro-3-extension, i.e. the Hilbert 3-class field tower, of k , which

TABLE 7. Quartets, $m = 4$, with $f = 9pq$, $q \equiv 5 \pmod{9}$

p	q	f	n_i	CT	\varkappa_i	α_i	DPF	Type
73	5	3285	$16425 = 3^2 \cdot 5^2 \cdot 73$	b.10	0043	21, 21, 111, 111	3	α
			$1095 = 3 \cdot 5 \cdot 73$	b.10	4001	111, 21, 21, 111	3	
			$3285 = 3^2 \cdot 5 \cdot 73$	b.10	4001	111, 21, 21, 111	3	
			$5475 = 3 \cdot 5^2 \cdot 73$	b.10	4001	111, 21, 21, 111	3	
127	5	5715	$1905 = 3 \cdot 5 \cdot 127$	b.10	0043	21, 21, 111, 111	5	α
			$28575 = 3^2 \cdot 5^2 \cdot 127$	b.10	0402	21, 111, 21, 111	5	
			$9525 = 3 \cdot 5^2 \cdot 127$	b.10	0320	21, 111, 111, 21	5	
			$5715 = 3^2 \cdot 5 \cdot 127$	b.10	0402	21, 111, 21, 111	5	

is always metabelian $F_3^\infty(k) = F_3^2(k)$ in our numerical examples. Except mentioned differently, the TKT is always b.10, $\varkappa \sim (0320)$, with associated ATI $\alpha \sim [(21)(1^3)(1^3)(21)]$, which uniquely identifies $G \simeq \langle 243, 3 \rangle$ with admissible relation rank $d_2(G) = 4$ [21, Thm. 5.1, p. 28]. Since k has 3-class rank $\varrho = 2$, signature $(r_1, r_2) = (0, 3)$, torsion free Dirichlet unit rank $r = r_1 + r_2 - 1 = 2$, and contains the primitive cube roots of unity, indicated by $\theta = 1$, the bounds for the relation rank of G are given by

$$(6.1) \quad 2 = \varrho \leq d_2(G) \leq \varrho + r + \theta = 2 + 2 + 1 = 5,$$

according to Shafarevich [25]. The differential principal factorization type of $k = \mathbb{Q}(\zeta, \sqrt[3]{n})$ is always DPF-type β .

Example 1. Let $f = 3pq$ with primes $p \equiv 1 \pmod{9}$ and $q \equiv 2, 5 \pmod{9}$.

- For $q \equiv 2 \pmod{9}$, we found two prototypes (see Table 4):
 $f = 3 \cdot 73 \cdot 2 = 438$, a doublet with radicands $n \in \{2^2 \cdot 73, 2 \cdot 73\}$.
 $f = 3 \cdot 541 \cdot 2 = 3246$, a doublet with radicands $n \in \{2^2 \cdot 541, 2 \cdot 541\}$. This is the unique instance which admits a fixed point capitulation, since $\mathbb{Q}(\sqrt[3]{541})$ is of type γ , and indeed both members of the doublet share the TKT d.23, $\varkappa \sim (1320)$, with associated ATI $\alpha \sim [(33)(1^3)(1^3)(21)]$ and thus $G \simeq \langle 6561, 1990 \rangle$ with $d_2(G) = 4$.
- For $q \equiv 5 \pmod{9}$, there also occurred two prototypes (see Table 5):
 $f = 3 \cdot 73 \cdot 5 = 1095$, a doublet with radicands $n \in \{5 \cdot 73, 5^2 \cdot 73\}$.
 $f = 3 \cdot 541 \cdot 5 = 8115$, a doublet with radicands $n \in \{5 \cdot 541, 5^2 \cdot 541\}$. Since $\mathbb{Q}(\sqrt[3]{541})$ is of type γ , a fixed point is admissible in the capitulation type, and both members of the doublet share the TKT d.23, $\varkappa \sim (1320)$, with associated ATI $\alpha \sim [(33)(1^3)(1^3)(21)]$, again leading to $G \simeq \langle 6561, 1990 \rangle$ with $d_2(G) = 4$ [15].

Example 2. Let $f = 9pq$ with primes $p \equiv 1 \pmod{9}$ and $q \equiv 2, 5 \pmod{9}$. For both prototypes, 3 is cubic residue modulo 73, and the principal factors are 3 and 9, for each component of the quartets. Again, the TKT is $\varkappa \sim (0320)$, and the (logarithmic) abelian quotient invariants are $\alpha \sim [(21)(1^3)(1^3)(21)]$, for each component of the quartets.

- For $q \equiv 2 \pmod{9}$, we have (see Table 6):
 $f = 9 \cdot 73 \cdot 2 = 1314$, quartet with radicands $n \in \{2 \cdot 3 \cdot 73, 2^2 \cdot 3 \cdot 73, 2^2 \cdot 3^2 \cdot 73, 2 \cdot 3^2 \cdot 73\}$.

- For $q \equiv 5 \pmod{9}$, we got (see Table 7):
 $f = 9 \cdot 73 \cdot 5 = 3285$, quartet with radicands $n \in \{3^2 \cdot 5^2 \cdot 73, 3 \cdot 5 \cdot 73, 3^2 \cdot 5 \cdot 73, 3 \cdot 5^2 \cdot 73\}$.

Example 3. Let $f = pq_1q_2$ with primes $p \equiv 1 \pmod{9}$ and $q_i \equiv 2, 5 \pmod{9}$ for $i = 1, 2$. Again, the capitulation type is $\varkappa \sim (0320)$, and the (logarithmic) abelian quotient invariants are $\alpha \sim [(21)(1^3)(1^3)(21)]$, for each component of the doublets.

- For $q_1 \equiv q_2 \equiv 2 \pmod{9}$, there occurred (see Table 1):
 $f = 19 \cdot 2 \cdot 11 = 418$, a doublet with radicands $n \in \{2^2 \cdot 11 \cdot 19, 2 \cdot 11 \cdot 19\}$. Since 11 is cubic residue modulo 19, the principal factors are 11 and $2^2 \cdot 19$, respectively 11 and $2 \cdot 19$.
- For the mixed case $q_1 \equiv 2 \pmod{9}$, $q_2 \equiv 5 \pmod{9}$, we found (see Table 3):
 $f = 19 \cdot 5 \cdot 11 = 1045$, a doublet with radicands $n \in \{5 \cdot 11 \cdot 19, 5^2 \cdot 11 \cdot 19\}$. Since 11 is cubic residue modulo 19, the principal factors are 11 and $5 \cdot 19$, respectively 11 and 11^2 .
- For $q_1 \equiv q_2 \equiv 5 \pmod{9}$, there occurred (see Table 2):
 $f = 37 \cdot 5 \cdot 23 = 4255$, a doublet with radicands $n \in \{5 \cdot 23 \cdot 37, 5^2 \cdot 23 \cdot 37\}$. Since 23 is cubic residue modulo 37, the principal factors are 23 and $5 \cdot 37$, respectively 23 and 23^2 .

Although it is not a counter example, we illuminate the impact of the prime $p = 541$ once more, with a conductor intentionally skipped from our systematic investigations. 541 is the leading term of the sequence A363717 in the On-line Encyclopedia of Integer Sequences (OEIS) [23]. The sequence consists of the prime radicands $p \equiv 1 \pmod{9}$ of pure cubic number fields $\mathbb{Q}(\sqrt[3]{p})$ whose normal closure possesses the differential principal factorization (DPF) type γ [1, Thm. 2.1, p. 254].

Example 4. Let $f = 9p$ with a prime $p \equiv 1 \pmod{9}$. Except mentioned differently, the TKT is usually a.1, $\varkappa \sim (0000)$.

- For $f = 9 \cdot 19 = 171$, we have a doublet, each component with ATI $\alpha \sim [(1)(1)(1)(1)]$, which uniquely identifies the abelian root $G \simeq \langle 9, 2 \rangle$ with $d_2(G) = 3$ [4].
- For $f = 9 \cdot 199 = 1791$, we found a doublet, each component with ATI $\alpha \sim [(21)(11)(11)(11)]$, which uniquely identifies mainline $G \simeq \langle 81, 9 \rangle$ with $d_2(G) = 4$.
- For $f = 9 \cdot 541 = 4869$, however, there occurred a doublet, each component with TKT $\varkappa \sim (1000)$, having a fixed point, and ATI $\alpha \sim [(33)(11)(11)(11)]$, which uniquely identifies $G \simeq \langle 2187, 387 \rangle$ with $d_2(G) = 3$ in depth 1 [4].

However, eventually we point out that a pure cubic auxiliary field $\mathbb{Q}(\sqrt[3]{p})$ of DPF-type γ is **not a warranty** for *partial* capitulation in the first component (it is only necessary but not sufficient for the capitulation types a.2, $\varkappa \sim (1000)$ and d.23, $\varkappa \sim (1320)$).

Example 5. Let $f = pq$ with primes $p \equiv 1 \pmod{9}$ and $q \equiv 8 \pmod{9}$.

- For $p = 1279$ and $q = 17$, the conductor $f = 21743$ gives rise to a doublet of Dedekind species II, whose members have *total* capitulation in the first component with Artin pattern $\alpha = [32, 11, 11, 11]$, $\varkappa = (0000)$, although 1279 belongs to the sequence A363717.

7. DATA AVAILABILITY STATEMENT

Experimental results communicated in this article and the source code of Magma program scripts used for the computations may be requested from the second author by email.

8. ACKNOWLEDGEMENTS

The second author acknowledges that his research was supported by the Austrian Science Fund (FWF): projects J0497-PHY, P26008-N25, and by the Research Executive Agency of the European Union (EUREA): project Horizon Europe 2021–2027.

REFERENCES

- [1] S. Aouissi, D. C. Mayer, M. C. Ismaili, M. Talbi, and A. Azizi, 3-rank of ambiguous class groups of cubic Kummer extensions, *Period. Math. Hungar.* **81** (2020), 250–274.
- [2] S. Aouissi, A. Azizi, M. C. Ismaili, D. C. Mayer and M. Talbi, Principal factors and lattice minima in cubic fields, *Kyushu J. Math.* **76** (2022), 101–118.
- [3] M. Ayadi, A. Azizi, and M. C. Ismaili, *The capitulation problem for certain number fields*, Advanced Studies in Pure Mathematics **30** (2001), Class Field Theory - Its Centenary and Prospect, pp. 467–482.
- [4] H. U. Besche, B. Eick, and E. A. O’Brien, *The SmallGroups Library — a Library of Groups of Small Order*, 2005, an accepted and refereed GAP package, available also in MAGMA.
- [5] W. Bosma, J. Cannon, and C. Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), 235–265.
- [6] W. Bosma, J. J. Cannon, C. Fieker, and A. Steels (eds.), *Handbook of Magma functions*, Edition 2.28, Sydney, 2024.
- [7] C. Fieker, *Computing class fields via the Artin map*, Math. Comp. **70** (2001), no. 235, 1293–1303.
- [8] F.-P. Heider und B. Schmithals, *Zur Kapitulation der Idealklassen in unverzweigten primzyklischen Erweiterungen*, J. Reine Angew. Math. **336** (1982), 1–25.
- [9] B. Huppert, *Endliche Gruppen, I*, Springer-Verlag, Berlin, 1967.
- [10] M. C. Ismaili, *Sur la capitulation des 3-classes d’idéaux de la clôture normale d’un corps cubique pur*, Thèse de doctorat, Univ. Laval, Québec, 1992.
- [11] M. C. Ismaili and R. El Mesaoudi, *Sur la divisibilité exacte par 3 du nombre de classes de certains corps cubiques purs*, *Ann. Sci Math. Québec* **25** (2001), no. 2, 153–177.
- [12] M. C. Ismaili and R. El Mesaoudi, *Corps cubiques purs dont le nombre de classes est exactement divisible par 3*, *Ann. Sci Math. Québec* **28** (2004), no. 1–2, 103–112.
- [13] M. C. Ismaili and R. El Mesaoudi, *Sur la capitulation des 3-classes d’idéaux de la clôture normale de certains corps cubiques purs*, *Ann. Sci Math. Québec* **29** (2005), no. 1, 49–72.
- [14] Magma Developer Group, *Magma Computational Algebra System*, Version 2.28-16, Sydney, 2024, available from <http://magma.maths.usyd.edu.au>.
- [15] Magma Developer Group, *Magma, Data for groups of order 3⁸*, `data3to8.tar.gz`, Sydney, 2012, available from <http://magma.maths.usyd.edu.au>.
- [16] D. C. Mayer, *Multiplicities of dihedral discriminants*, Math. Comp. **58** (1992), no. 198, 831–847 and S55–S58.
- [17] D. C. Mayer, *Transfers of metabelian p-groups*, Monatsh. Math. **166** (2012), no. 3–4, 467–495.
- [18] D. C. Mayer, *The distribution of second p-class groups on coclass graphs*, J. Théor. Nombres Bordeaux **25** (2013), no. 2, 401–456.
- [19] D. C. Mayer, *Principalization algorithm via class group structure*, J. Théor. Nombres Bordeaux **26** (2014), no. 2, 415–464.

- [20] D. C. Mayer, *Quadratic p -ring spaces for counting dihedral fields*, Int. J. Number Theory **10** (2014), no. 8, 2205–2242.
- [21] D. C. Mayer, *New number fields with known p -class tower*, Tatra Mt. Math. Pub. **64** (2015), 21–57, DOI 10.1515/tmmp-2015-0040, Special Issue on Number Theory and Cryptology ‘15.
- [22] B. Nebelung, *Klassifikation metabelscher 3-Gruppen mit Faktorkommutatorgruppe vom Typ (3, 3) und Anwendung auf das Kapitulationsproblem*, Inauguraldissertation, Universität zu Köln, 1989.
- [23] OEIS Foundation Inc. (N. J. A. Sloane), *The On-Line Encyclopedia of Integer Sequences* (OEIS), 2024, Published electronically at <https://oeis.org/>.
- [24] A. Scholz und O. Taussky, *Die Hauptideale der kubischen Klassenkörper imaginär quadratischer Zahlkörper: ihre rechnerische Bestimmung und ihr Einfluß auf den Klassenkörperturm*, J. Reine Angew. Math. **171** (1934), 19–41.
- [25] I. R. Shafarevich, *Extensions with prescribed ramification points* (Russian), Publ. Math., Inst. Hautes Études Sci. **18** (1964), 71–95. (English transl. by J. W. S. Cassels in Amer. Math. Soc. Transl., II. Ser., **59** (1966), 128–149.)

ALGEBRAIC THEORIES AND APPLICATIONS RESEARCH TEAM, ECOLE NORMALE SUPÉRIEURE (ENS)
OF MOULAY ISMAIL UNIVERSITY (UMI), B.P. 3104, TOULAL, MEKNÉS, MOROCCO
Email address: s.aouissi@umi.ac.ma

NAGLERGASSE 53, 8010 GRAZ, AUSTRIA
Email address: quantum.algebra@icloud.com
URL: <http://www.algebra.at>