# THE MORDELL-TORNHEIM ZETA FUNCTION: KRONECKER LIMIT TYPE FORMULA, SERIES EVALUATIONS AND APPLICATIONS

#### SUMUKHA SATHYANARAYANA AND N. GURU SHARAN

ABSTRACT. In this paper, we establish Kronecker limit type formulas for the Mordell-Tornheim zeta function  $\Theta(r,s,t,x)$  as a function of the second as well as the third arguments. As an application of these formulas, we obtain results of Herglotz, Ramanujan, Guinand, Zagier and Vlasenko-Zagier as corollaries. We show that the Mordell-Tornheim zeta function lies centrally between many modular relations in the literature, thus providing the means to view them under one umbrella. We also give series evaluations of  $\Theta(r,s,t,x)$  in terms of Herglotz-Zagier function, Vlasenko-Zagier function and their derivatives. Using our new perspective of modular relations, we obtain a new infinite family of results called mixed functional equations.

## Contents

1. Introduction	1
1.1. Modular relations in the literature	2
2. Kronecker Limit type formula for $\Theta(r, s, t, x)$ in the Third variable	4
3. Applications in modular relations	10
3.1. Vlasenko and Zagier's functional equation	11
3.2. Guinand's functional equations	14
3.3. Ramanujan's functional equation	15
4. Kronecker limit type formula for $\Theta(r, s, t, x)$ in the Second variable	16
4.1. Mixed functional equations	20
4.2. Kronecker limit type formula	20
5. Modular relations: The Mordell-Tornheim zeta perspective	23
6. Concluding Remarks	23
7. Acknowledgment	23
References	24

#### 1. Introduction

Let

$$\mathscr{D} := \left\{ (r, s, t) \in \mathbb{C}^3 \mid \operatorname{Re}(r+t) > 1, \operatorname{Re}(s+t) > 1, \operatorname{Re}(r+s+t) > 2 \right\}. \tag{1.1}$$

In the region  $\mathcal{D}$ , the Mordell-Tornheim zeta function is given by the absolutely convergent series

$$\zeta_{\text{MT}}(r, s, t) := \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{n^r m^s (n+m)^t}.$$
(1.2)

The function  $\zeta_{\text{MT}}(r, s, t)$  can be meromorphically continued to the entire  $\mathbb{C}^3$  space, and its singular hyperplanes are given by  $r+t=1-\ell$ ,  $s+t=1-\ell$ , for any  $\ell \in \mathbb{N} \cup \{0\}$  and by r+s+t=2. Matsumoto [14] has given some details on the hyperplanes. The behavior of  $\zeta_{\text{MT}}(r, s, t)$  around its singularities is largely unexplored. In a recent work with Dixit [5], the authors considered its generalization

$$\Theta(r, s, t, x) := \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{n^r m^s (n + mx)^t},$$
(1.3)

 $<sup>2020\</sup> Mathematics\ Subject\ Classification.\ Primary\ 11M32,\ 30D30;\ Secondary\ 39B32.$ 

Keywords and phrases. Mordell-Tornheim zeta function, Kronecker limit formula, modular relation, series evaluation. Abbreviated title: Kronecker limit type formulas and their applications.

for x > 0 and  $(r, s, t) \in \mathcal{D}$  as defined in (1.1). Clearly,  $\Theta(r, s, t, 1) = \zeta_{\text{MT}}(r, s, t)$ . One can easily observe that  $\Theta(r, s, t, x)$  satisfies the following functional equations for x > 0 and  $(r, s, t) \in \mathcal{D}$ :

$$\Theta(r, s, t, x) = \Theta(r - 1, s, t + 1, x) + x \Theta(r, s - 1, t + 1, x). \tag{1.4}$$

$$\Theta(r, s, t, x) = x^{-t}\Theta(s, r, t, \frac{1}{x}). \tag{1.5}$$

The meromorphic continuation of  $\Theta(r, s, t, x)$  can be done similarly as in the case of  $\zeta_{\text{MT}}(r, s, t)$  [14, Theorem 1]. Recently, it has been shown that [5, Theorem 1], as  $t \to 0$ ,

$$\Theta(1, 1, t, x) = \frac{2}{t^2} + \frac{2\gamma - \log(x)}{t} + \gamma^2 - \gamma \log(x) - \frac{\pi^2}{6} + O(|t|), \tag{1.6}$$

where  $\gamma$  is Euler's constant. Results of this kind did not exist in the literature prior to this. Moreover, one can also derive the higher Laurent coefficients of  $\Theta(1, 1, t, x)$  as a finite sum of Arakawa and Kaneko constants [2, Corollary 5]

$$c_k := \frac{1}{k!} \int_0^\infty \left\{ \log^k(y) - \log^y \left( \frac{y}{1 - e^{-y}} \right) \right\} \left( \frac{1}{e^y - 1} - \frac{1}{y} \right) dy, \tag{1.7}$$

and integrals of the form

$$L_k^*(x) := \sum_{j=0}^k a_{k+1,j} \int_0^\infty \frac{\log^j(y)}{y} \log\left(\frac{1 - e^{-xy}}{1 - e^{-y}}\right) \log\left(\frac{1 - e^{-y}}{y}\right) dt \tag{1.8}$$

for  $k \in \mathbb{N}$ . For finer details, see the proofs of Theorem 3.4 and Proposition 3.7 in [5]. Recently, Matsumoto, Onodera and Sahoo, have derived a general asymptotic result for the Mordell-Tornheim multiple-zeta function [15].

Consider the real binary quadratic form  $Q(x,y) = ax^2 + bxy + cy^2$  with discriminant  $D = b^2 - 4ac$ . For a > 0, the associated zeta function is defined by

$$\zeta_Q(s) = \sum_{(m,n)\neq(0,0)} Q(m,n)^{-s}$$

where the sum runs over all the non-origin coordinates of the lattice  $\mathbb{Z} \times \mathbb{Z}$ .  $\zeta_Q(s)$  converges absolutely for Re(s) > 1. For D < 0, Kronecker proved that, around s = 1,

$$|D|^{\frac{s}{2}}\zeta_{Q}(s) = \frac{2\pi}{s-1} + 2\pi \left(2\gamma - \log 4 + H\left(\frac{-b+\sqrt{D}}{2a}\right)\right) + O(|s-1|),\tag{1.9}$$

where  $H(z) = -\log(y|\eta(z)|^4)$ , with  $\eta(z)$  being the Dedekind eta function. The result (1.9) is famously known as the Kronecker limit formula. Siegel [18] connected the Kronecker limit formula with the theory of *L*-functions. Over the century, several prominent mathematicians such as Hecke [10], Herglotz [11], Ramachandra [16], Stark [21] and Zagier [25] gave important contributions to this field. Note that (1.6) is a Kronecker limit type formula for the function  $\Theta(1,1,t,x)$  in the variable t, around t=0.

In Section 2, we give Kronecker limit type formulas for  $\Theta(r, s, t, x)$  in t, i.e., the third variable, in Theorem 2.3. For a fixed  $r, s \in \mathbb{C}$ , only possibilities of poles for  $\Theta(r, s, t, x)$  in t are at  $t = 1 - r - \ell$ ,  $t = 1 - s - \ell$ , for  $\ell \in \mathbb{N} \cup \{0\}$  and t = 2 - r - s. However, we show that all of these points need not have a pole necessarily. As a direct consequence of our Kronecker limit type formula, the singular part of the Laurent series of  $\Theta(r, s, t, x)$  in t is established. Interestingly, we show that the order of pole too varies by cases.  $\Theta(r, s, t, x)$  can have double, single or no pole in t, depending on the choices of t and t. This method cannot be used to obtain the Kronecker limit type formula for the  $\Theta(r, s, t, x)$  in the second variable.

1.1. Modular relations in the literature. Let Re(x) > 0. A transformation of the form F(x) = F(1/x) is called called a modular relation. Equivalently, we can also write it as form F(-1/z) = F(z), where  $z \in \mathbb{H}$ , the upper half complex plane.

Ramanujan [3] had recorded the following interesting Modular relation, on page 220 of Lost Notebook: For x > 0, let

$$\phi(x) := \psi(x) + \frac{1}{2x} - \log(x),$$

where  $\psi(x) = \frac{\Gamma'(x)}{\Gamma(x)}$ , is the logarithmic derivative of the Gamma function  $\Gamma(x)$ . Then,

$$\sqrt{x} \left\{ \frac{\gamma - \log(2\pi x)}{2x} + \sum_{n=1}^{\infty} \phi(nx) \right\} = \sqrt{\frac{1}{x}} \left\{ \frac{x(\gamma - \log\left(\frac{2\pi}{x}\right))}{2} + \sum_{n=1}^{\infty} \phi\left(\frac{n}{x}\right) \right\}$$

$$= -\frac{1}{\pi^{3/2}} \int_0^\infty \left| \Xi\left(\frac{1}{2}y\right) \Gamma\left(\frac{-1+iy}{4}\right) \right|^2 \frac{\cos\left(\frac{1}{2}y\log x\right)}{1+y^2} \, dy, \tag{1.10}$$

where

$$\Xi(y) = \xi \left(\frac{1}{2} + iy\right),$$
  
$$\xi(s) = \frac{1}{2}s(s-1)\pi^{-\frac{s}{2}}\Gamma\left(\frac{s}{2}\right)\zeta(s),$$

are the Riemann's functions. Dixit [4, Theorem 4.1] obtained a generalization of the above result in the variable z, with the Hurwitz zeta function  $\zeta(z, x)$  in the place of  $\psi(x)$  in the first equality, the modular relation.

Guinand [8] gave modular relations which are analogous to the first equality in (1.10) involving derivatives of digamma function  $\psi(x)$ . For  $z \in \mathbb{N}$  and z > 2, he showed that,

$$x^{\frac{z}{2}} \sum_{j=1}^{\infty} \psi^{(z-1)}(1+jx) = \left(\frac{1}{x}\right)^{\frac{z}{2}} \sum_{j=1}^{\infty} \psi^{(z-1)}\left(1+\frac{j}{x}\right). \tag{1.11}$$

Also, analogously, for the first derivative, he proved the identity

$$x\sum_{j=1}^{\infty} \left(\psi'(1+jx) - \frac{1}{jx}\right) - \frac{1}{2}\log(x) = \frac{1}{x}\sum_{j=1}^{\infty} \left(\psi'\left(1+\frac{j}{x}\right) - \frac{1}{\frac{j}{x}}\right) - \frac{1}{2}\log\left(\frac{1}{x}\right)$$
(1.12)

Another identity which is closely related to (1.10) is the two term functional equation of the Herglotz-Zagier function F(x). The function F(x) is defined as

$$F(x) := \sum_{n=1}^{\infty} \frac{\psi(nx) - \log(nx)}{n} \qquad (x \in \mathbb{C} \setminus (-\infty, 0]).$$
 (1.13)

This function was studied for the first time by Herglotz [11]. Later, Zagier studied this in connection with Kronecker limit formula for real quadratic fields [25]. From the asymptotic expansion of  $\psi(x)$  as  $x \to \infty$ , we can see that that the series in (1.13) converges absolutely for any  $x \in \mathbb{C} \setminus (-\infty, 0]$ . Zagier [25, Equations (7.4), (7.8)] proved the following two- and three-term functional equations for F(x):

$$F(x) + F\left(\frac{1}{x}\right) = 2F(1) + \frac{1}{2}\log^2(x) - \frac{\pi^2}{6x}(x-1)^2,$$
(1.14)

$$F(x) - F(x+1) - F\left(\frac{x}{x+1}\right) = -F(1) + \text{Li}_2\left(\frac{1}{1+x}\right),$$
 (1.15)

with [25, Equation (7.12)]

$$F(1) = -\frac{1}{2}\gamma^2 - \frac{\pi^2}{12} - \gamma_1,\tag{1.16}$$

where  $\gamma_1$  is the first Stieltjes constant. Observe that the two-term functional equation (1.14) of Herglotz-Zagier function resembles the first equality of Ramanujan's modular transformation (1.10). For any integer  $r \geq 2$ , Vlasenko and Zagier [24] defined the higher Herglotz function  $F_r(x)$ , by the following absolutely convergent series:

$$F_r(x) := \sum_{n=1}^{\infty} \frac{\psi(nx)}{n^r} \qquad (x \in \mathbb{C} \setminus (-\infty, 0]).$$
 (1.17)

They further showed that  $F_r(x)$  satisfies two- and three-term functional equations analogous [24, Proposition 4] to (1.14) and (1.15), given by,

$$F_r(x) + (-x)^{r-1} F_r\left(\frac{1}{x}\right) = \zeta(r+1) \left( (-x)^r - \frac{1}{x} \right) - \sum_{\ell=1}^r \zeta(\ell) \zeta(r-\ell+1) (-x)^{\ell-1}, \tag{1.18}$$

$$F_r(x) - F_r(x+1) + (-x)^{r-1} F_r\left(\frac{x+1}{x}\right) = \zeta(r+1) \left(\frac{(-x)^r}{x+1} - \frac{1}{x}\right) - \sum_{\ell=1}^r \zeta(r-\ell+1,\ell)(-x)^{\ell-1},\tag{1.19}$$

where the double zeta function  $\zeta(s_1, s_2)$  is defined [24, Section 1.1] by,

$$\zeta(s_1, s_2) = \sum_{n=1}^{\infty} \sum_{m=n}^{\infty} \frac{1}{m^{s_1} n^{s_2}}, \quad \text{(for } s_1 \ge 2 \text{ and } s_2 \ge 1).$$
 (1.20)

In this paper, we discover a new way to obtain these modular relations using the generalized Mordell-Tornheim zeta function  $\Theta(r, s, t, x)$ , and its properties (1.4) and (1.5). This facilitates us to develop a unified method to obtain the aforementioned two-term functional equations. The modular results discovered by Guinand, Ramanujan, Herglotz, Zagier and Vlasenko-Zagier are thus inherently connected in this way. Moreover, this approach shows that Mordell-Tornheim zeta function lies centrally between these modular relations. See Sections 3 and 5 for details.

In the Section 4, we give the series evaluations of  $\Theta(r, s, t, x)$  in terms of zeta values,  $\psi(x)$  and its derivatives, in Theorem 4.2. We further use Theorem 4.2 to get *mixed functional equations* in (4.14) as a corollary, which have not appeared in the literature. We call them mixed functional equations since they contain terms of the form both

$$\sum_{m=1}^{\infty} \frac{\psi^{(a)}(mx+1)}{m^b} \quad \text{and} \quad \sum_{m=1}^{\infty} \frac{\gamma + \psi(mx+1)}{m^c},$$


for a, b, and  $c \in \mathbb{N}$ , which appear in the Guinand's functional equation and the Vlasenko-Zagier's functional equation respectively. Of the infinite family of new identities we can obtain, one such new is, as explained in Remark 1,

$$-\sum_{m=1}^{\infty} \frac{\psi'(mx+1)}{m^2} + \sum_{m=1}^{\infty} \frac{\psi'(\frac{m}{x}+1)}{m^2} + \frac{2}{x} \sum_{m=1}^{\infty} \frac{\gamma + \psi(mx+1)}{m^3} = \zeta^2(2).$$
 (1.21)

Also as a corollary of Theorem 4.2, we derive a Kronecker limit type formula, as stated in Theorem (4.4), for  $\Theta(r, s, t, x)$  in the second variable.

# 2. Kronecker Limit type formula for $\Theta(r,s,t,x)$ in the Third variable

We first prove the Proposition 2.1, and its corollary 2.2, before using them to prove the main results of the Section-the Kronecker Limit type formula in the *third variable*, stated below in three Theorems 2.3, 2.4 and 2.5. Theorems 2.3 and 2.4 are subdivided into four cases each, whose casewise classification can be better understood by the flowcharts in Figure 1 and 2 respectively. Theorems 2.3 and 2.4 pertains to singularities of the type  $r+t=1-\ell$  for  $\ell \in \mathbb{N} \cup \{0\}$  while Theorem 2.5 pertains to singularities of the type r+s+t=2.






FIGURE 1. The cases for  $r \in \mathbb{Z} \setminus \mathbb{N}$ .

FIGURE 2. The cases for  $r \in \mathbb{N}$ .

**Proposition 2.1.** Let  $r, s, t \in \mathbb{C}$  such that  $r \notin \mathbb{N}$ , and

 $Re(t) > \max\{0, 1 - Re(r), 1 - Re(s), 2 - Re(r) - Re(s)\}$ . For any  $M \in \mathbb{N} \cup \{0\}$  and x > 0, we have

$$\frac{1}{\Gamma(t)} \int_0^\infty y^{t-1} \mathrm{Li}_s(e^{-xy}) \left( \mathrm{Li}_r(e^{-y}) - \Gamma(1-r) y^{r-1} - \sum_{k=0}^M (-1)^k \zeta(r-k) \frac{y^k}{k!} \right) dy$$

$$=\Theta(r,s,t,x) - \frac{\Gamma(1-r)\Gamma(t+r-1)}{x^{t+r-1}\Gamma(t)}\zeta(t+s+r-1) - \sum_{k=0}^{M} \frac{(-1)^k}{k!} \frac{\zeta(r-k)\Gamma(t+k)}{x^{t+k}\Gamma(t)}\zeta(t+k+s). \tag{2.1}$$

*Proof.* For Re(t), x, n > 0, the definition of the Gamma function  $\Gamma(t)$  upon a change of variable gives,

$$\int_0^\infty y^{t-1} e^{-(x+n)y} \, dy = (x+n)^{-t} \Gamma(t).$$

Since Re(r+t) > 1, divide both sides of the equation by  $n^r$  and sum over  $n \in \mathbb{N}$  and interchange the order of summation and integration to get,

$$\int_0^\infty y^{t-1} e^{-xy} \operatorname{Li}_r(e^{-y}) \, dy = \Gamma(t) \sum_{n=1}^\infty \frac{1}{n^r (x+n)^t},\tag{2.2}$$

where  $\operatorname{Li}_r(z) := \sum_{u=1}^{\infty} \frac{z^u}{u^r}$  is the polylogarithm function. For  $r \notin \mathbb{N}$ , around y = 0, as stated in [6, Equation 8, p.29], we have,

$$\operatorname{Li}_{r}(e^{-y}) = \Gamma(1-r)y^{r-1} + \sum_{k=0}^{\infty} (-1)^{k} \zeta(r-k) \frac{y^{k}}{k!} = O(y^{\max(0,\operatorname{Re}(r)-1)}). \tag{2.3}$$

The bound in (2.3) along with  $Re(t) > \max\{0, 1 - Re(r)\}$  ensures the absolute convergence, which in turn allows the interchange using [22, Theorem 2.1]. Using (2.2) and the definition of  $\Gamma(t)$  twice, we see that,

$$\int_{0}^{\infty} y^{t-1} e^{-xy} \left( \operatorname{Li}_{r}(e^{-y}) - \Gamma(1-r)y^{r-1} - \sum_{k=0}^{M} (-1)^{k} \zeta(r-k) \frac{y^{k}}{k!} \right) dy$$

$$= \Gamma(t) \sum_{n=1}^{\infty} \frac{1}{n^{r}(x+n)^{t}} - \frac{\Gamma(1-r)\Gamma(t+r-1)}{x^{t+r-1}} - \sum_{k=0}^{M} \frac{(-1)^{k}}{k!} \frac{\zeta(r-k)\Gamma(t+k)}{x^{t+k}}.$$
(2.4)

Replace x by mx in (2.4), divide both the sides of resulting equation by  $m^s\Gamma(t)$ , sum over  $m \in \mathbb{N}$  and then interchange the order of summation and integration to get,

$$\frac{1}{\Gamma(t)} \int_{0}^{\infty} y^{t-1} \operatorname{Li}_{s}(e^{-xy}) \left( \operatorname{Li}_{r}(e^{-y}) - \Gamma(1-r)y^{r-1} - \sum_{k=0}^{M} (-1)^{k} \zeta(r-k) \frac{y^{k}}{k!} \right) dy$$

$$= \Theta(r, s, t, x) - \frac{\Gamma(1-r)\Gamma(t+r-1)}{x^{t+r-1}\Gamma(t)} \zeta(t+s+r-1) - \sum_{k=0}^{M} \frac{(-1)^{k}}{k!} \frac{\zeta(r-k)\Gamma(t+k)}{x^{t+k}\Gamma(t)} \zeta(t+k+s). \tag{2.5}$$

We justify the interchange of the order of summation and integration as follows: From (2.3), we have,

$$\operatorname{Li}_{r}(e^{-y}) - \Gamma(1-r)y^{r-1} - \sum_{k=0}^{M} (-1)^{k} \zeta(r-k) \frac{y^{k}}{k!} = O(y^{M+1}), \quad \text{as} \quad y \to 0^{+}.$$
 (2.6)

Similarly,

$$\operatorname{Li}_{s}(e^{-y}) = \begin{cases} O(1), & \text{if } \operatorname{Re}(s) \ge 1, \\ O(y^{\operatorname{Re}(s)-1}), & \text{if } \operatorname{Re}(s) < 1, \end{cases} \quad \text{as} \quad y \to 0^{+}.$$
 (2.7)

Since  $Re(t) > max\{0, 1 - Re(s)\}\$ , from (2.6) and (2.7) we get

$$y^{t-1} \operatorname{Li}_{s}(e^{-xy}) \left( \operatorname{Li}_{r}(e^{-y}) - \Gamma(1-r) y^{r-1} - \sum_{k=0}^{M} (-1)^{k} \zeta(r-k) \frac{y^{k}}{k!} \right) = O\left(y^{M+\epsilon}\right), \tag{2.8}$$

for any  $\epsilon > 0$ , as  $y \to 0^+$ . Also,  $\text{Li}_s(e^{-y}) = O(y^{-K})$ , for any K > 1 as  $y \to \infty$ . Therefore,

$$y^{t-1}\operatorname{Li}_{s}(e^{-xy})\left(\operatorname{Li}_{r}(e^{-y}) - \Gamma(1-r)y^{r-1} - \sum_{k=0}^{M}(-1)^{k}\zeta(r-k)\frac{y^{k}}{k!}\right) = O\left(y^{\max\{\operatorname{Re}(t) + M - 1 - K, \operatorname{Re}(t) + r - 2 - K\}}\right), \quad (2.9)$$

for any K > 1, as  $y \to \infty$ . Using [22, Theorem 2.1] with the facts in (2.8) and (2.9), we justify the interchange in (2.5) to complete the proof.

We now give the following result, analogous to Proposition 2.1 for  $r \in \mathbb{N}$  as a Corollary.

**Corollary 2.2.** Let  $r \in \mathbb{N}$  and  $s, t \in \mathbb{C}$  such that  $\text{Re}(t) > \max\{0, 1 - \text{Re}(r), 1 - \text{Re}(s), 2 - \text{Re}(r) - \text{Re}(s)\}$ . For any  $M \in \mathbb{N} \cup \{0\}$  with  $M \ge r - 1$ , and x > 0 we have,

$$\frac{1}{\Gamma(t)} \int_{0}^{\infty} y^{t-1} \operatorname{Li}_{s}(e^{-xy}) \left( \operatorname{Li}_{r}(e^{-y}) - (-1)^{r-1} (H_{r-1} - \log(y)) \frac{y^{r-1}}{(r-1)!} - \sum_{\substack{k=0\\k\neq r-1}}^{M} (-1)^{k} \zeta(r-k) \frac{y^{k}}{k!} \right) dy$$

$$= \Theta(r, s, t, x) - \sum_{\substack{k=0\\k\neq r-1}}^{M} \frac{(-1)^{k}}{k!} \frac{\zeta(r-k)\Gamma(t+k)}{x^{t+k}\Gamma(t)} \zeta(t+k+s)$$

$$+ \frac{(-1)^{r} \Gamma(t+r-1)}{x^{t+r-1}(r-1)! \Gamma(t)} \left( \zeta(t+s+r-1) (H_{r-1} - \psi(t+r-1) + \log(x)) - \zeta'(t+s+r-1) \right). \tag{2.10}$$

*Proof.* Let  $a \in \mathbb{N}$ . Using the Laurent series expansions of  $\Gamma(1-r)$  and  $\zeta(r-a+1)$  around r=a and simplifying, we get,

$$\lim_{r \to a} \left( \operatorname{Li}_r(e^{-y}) - \Gamma(1-r)y^{r-1} - \sum_{k=0}^{M} (-1)^k \zeta(r-k) \frac{y^k}{k!} \right)$$

$$= \operatorname{Li}_a(e^{-y}) + (-1)^a (H_{a-1} - \log(y)) \frac{y^{a-1}}{(a-1)!} - \sum_{\substack{k=0\\k \neq a-1}}^{M} (-1)^k \zeta(a-k) \frac{y^k}{k!}$$
(2.11)

Take limit as  $r \to a$  on both sides of (2.1). Invoke [22, Theorem 2.2] with the bounds given in (2.8) and (2.9) to see the integral on the left-hand side is analytic in r, in the strip a - 1 < Re(r) < a + 1 with a removable singularity at a. Hence, interchanging the order of limit and integration, we get,

$$\frac{1}{\Gamma(t)} \int_{0}^{\infty} y^{t-1} \operatorname{Li}_{s}(e^{-xy}) \left( \operatorname{Li}_{a}(e^{-y}) + (-1)^{a} (H_{a-1} - \log(y)) \frac{y^{a-1}}{(a-1)!} - \sum_{\substack{k=0\\k\neq a-1}}^{M} (-1)^{k} \zeta(a-k) \frac{y^{k}}{k!} \right) dy$$

$$= \Theta(a, s, t, x) - \sum_{\substack{k=0\\k\neq a-1}}^{M} \frac{(-1)^{k}}{k!} \frac{\zeta(a-k)\Gamma(t+k)}{x^{t+k}\Gamma(t)} \zeta(s+t+k)$$

$$- \lim_{r \to a} \left( \frac{\Gamma(1-r)\Gamma(t+r-1)}{x^{t+r-1}\Gamma(t)} \zeta(s+t+r-1) - \frac{(-1)^{a}\zeta(r-a+1)\Gamma(t+a-1)}{x^{t+a-1}(a-1)!\Gamma(t)} \zeta(s+t+a-1) \right).$$

To evaluate the limit in the above equation, use the Laurent series expansions of the respective functions around r = a. Finally, replace a by r in the resultant to get (2.10).

We are now equipped to prove the Kronecker Limit type formula in third variable, around  $t = 1 - r - \ell$ .

**Theorem 2.3.** Let  $r \in \mathbb{Z} \setminus \mathbb{N}$  and  $\ell \in \mathbb{N} \cup \{0\}$  be arbitrary such that  $\ell \geq 1 - r$ . Then, the following hold. Case I: For a fixed  $s \in \mathbb{Z}$  such that  $s > r + \ell$  and  $s \neq \ell + 1$ , or  $s \in \mathbb{C} \setminus \mathbb{Z}$ , we have, around  $t = 1 - r - \ell$ ,

$$\Theta(r, s, t, x) = (-1)^{r-1} \frac{(r+\ell-1)!}{\ell!} x^{\ell} \Gamma(1-r) \zeta(s-\ell)$$

$$+ \sum_{k=0}^{r+\ell-1} \frac{(r+\ell-1)! \zeta(r-k)}{x^{1-r-\ell+k} k! (r+\ell-k-1)!} \zeta(-r+s-\ell+k+1) + O(|t-(1-r-\ell)|)$$
(2.12)

Case II: For a fixed  $s \in \mathbb{Z}$  such that  $s > r + \ell$  and  $s = \ell + 1$ , we have, around  $t = 1 - r - \ell$ ,

$$\Theta(r, s, t, x) = (-1)^{r-1} (-r)! x^{\ell} \frac{(r+\ell-1)!}{\ell! (t-(1-r-\ell))} + (-1)^{r-1} (-r)! x^{\ell} \frac{(r+\ell-1)!}{\ell!} (\gamma - \log(x) + H_{\ell} - H_{r+\ell-1}) 
+ \sum_{k=0}^{r+\ell-1} {r+\ell-1 \choose k} \frac{\zeta(r-k)}{x^{k-r-\ell+1}} \zeta(k-r+2) + O(|t-(1-r-\ell)|).$$
(2.13)

**Case III:** For a fixed  $s \in \mathbb{Z} \setminus \mathbb{N}$  such that  $s \leq r + \ell$ , we have, around  $t = 1 - r - \ell$ ,

$$\Theta(r,s,t,x) = (-1)^{r-1} x^{\ell} \frac{(-r)!(r+\ell-1)!}{\ell!} \zeta(s-\ell) + \sum_{k=0}^{r+\ell-1} \frac{\zeta(r-k)(r+\ell-1)!}{x^{k-r-\ell+1} k! (r+\ell-k-1)!} \zeta(s+1-r-\ell+k)$$

$$+ (-1)^{s-1} x^{s-1} \frac{(r+\ell-1)!(-s)!}{(r+\ell-s)!} \zeta(s-\ell) + O(|t-(1-r-\ell)|). \tag{2.14}$$

Case IV: For a fixed  $s \in \mathbb{N}$  such that  $s \leq r + \ell$ , we have, around  $t = 1 - r - \ell$ ,

$$\Theta(r, s, t, x) = x^{s-1} \frac{(r+\ell-1)!}{(r+\ell-s)!(s-1)!} \frac{\zeta(s-\ell)}{(t-(1-r-\ell))} 
+ x^{s-1} \frac{(r+\ell-1)!}{(r+\ell-s)!(s-1)!} (\gamma - H_{r+\ell-1} + H_{s-1} - \log(x)) \zeta(s-\ell) - (-1)^r x^{\ell} \frac{(-r)!(r+\ell-1)!}{\ell!} \zeta(s-\ell) 
+ \sum_{\substack{k=0\\k\neq r+\ell-s}}^{r+\ell-1} \frac{(r+\ell-1)!\zeta(r-k)}{(r+\ell-k-1)!x^{k-r-\ell+1}k!} \zeta(s-r-\ell+k+1) + O(|t-(1-r-\ell)|)$$
(2.15)

*Proof.* Let  $r \in \mathbb{Z} \setminus \mathbb{N}$  and  $\ell \in \mathbb{N} \cup \{0\}$  such that  $\ell \geq 1 - r$  be fixed and  $s \in \mathbb{C}$ . Then, for  $t > -r - \ell$ , (2.6) and (2.7) gives,

$$y^{t-1} \operatorname{Li}_{s}(e^{-xy}) \left( \operatorname{Li}_{r}(e^{-y}) - \Gamma(1-r) y^{r-1} - \sum_{k=0}^{r+\ell+\lfloor |\operatorname{Re}(s)|\rfloor+1} (-1)^{k} \zeta(r-k) \frac{y^{k}}{k!} \right) = O\left(y^{\epsilon-1}\right), \tag{2.16}$$

for some  $\epsilon > 0$ . For  $M = r + \ell + \lfloor |\operatorname{Re}(s)| \rfloor + 1$ , from (2.9) and (2.16), the integral on the left-hand side of (2.1) is now defined and analytic in the extended region  $\operatorname{Re}(t) > -r - \ell$  by using [22, Theorem 2.2]. Hence, due to the pole of  $\Gamma(t)$  at  $t = 1 - r - \ell$ , we see

$$\lim_{t \to 1 - r - \ell} \left( \frac{1}{\Gamma(t)} \int_0^\infty y^{t-1} \mathrm{Li}_s(e^{-xy}) \left( \mathrm{Li}_r(e^{-y}) - \Gamma(1 - r) y^{r-1} - \sum_{k=0}^{r + \ell + \lfloor |\mathrm{Re}(s)| \rfloor + 1} (-1)^k \zeta(r - k) \frac{y^k}{k!} \right) dy \right) = 0.$$

Hence, from (2.1)

$$\Theta(r, s, t, x) = \frac{\Gamma(1 - r)\Gamma(t + r - 1)}{x^{t + r - 1}\Gamma(t)} \zeta(r + s + t - 1) 
+ \sum_{k=0}^{r + \ell + \lfloor |\operatorname{Re}(s)| \rfloor + 1} \frac{(-1)^k \zeta(r - k)\Gamma(t + k)}{x^{t + k}k!\Gamma(t)} \zeta(s + t + k) + O(|t - (1 - r - \ell)|).$$
(2.17)

To get the required result for  $\Theta(r,s,t,x)$ , we analyze the behaviour of the right-hand side of (2.17) around  $t=1-r-\ell$ . Since  $\Gamma(t)$  has a pole at  $t=1-r-\ell$ , only the terms with a pole in the numerator survive as  $t\to 1-r-\ell$ . We handle the remaining calculations in cases, as mentioned in statement of this Theorem.

Case I: Note that,  $\Gamma(t+k)$  has a simple pole only for  $k \leq r+\ell-1$ . For any k,  $\zeta(s+t+k)$  has no pole since  $s > r+\ell$  or  $s \notin \mathbb{Z}$ .  $\Gamma(t+r-1)$  has a simple pole.  $\zeta(r+s+t-1)$  has no pole since  $s \neq \ell+1$ . Use the functional equation  $\Gamma(t+1) = t\Gamma(t)$  repeatedly to get,

$$\frac{\Gamma(t+k)}{\Gamma(t)} = t(t+1)\cdots(t+k-1).$$

Taking limit as  $t \to 1 - r - \ell$  on both sides of the above equation, for  $0 \le k \le r + \ell - 1$  we get,

$$\lim_{t \to 1 - r - \ell} \left( \frac{\Gamma(t+k)}{\Gamma(t)} \right) = (1 - r - \ell)(2 - r - \ell) \cdots (k - r - \ell) = (-1)^k \frac{(r+\ell-1)!}{(r+\ell-k-1)!}.$$
 (2.18)

Also, since  $r \in \mathbb{Z} \setminus \mathbb{N}$ ,

$$\frac{\Gamma(t+r-1)}{\Gamma(t)} = \frac{\Gamma(t+r-1)}{(t-1)(t-2)\cdots(t+r-1)\Gamma(t+r-1)} = \frac{1}{(t-1)(t-2)\cdots(t+r-1)}$$

Hence, we can see,

$$\lim_{t \to 1 - r - \ell} \left( \frac{\Gamma(t + r - 1)}{\Gamma(t)} \right) = (-1)^{r - 1} \frac{(r + \ell - 1)!}{\ell!}.$$
 (2.19)

Use (2.18) and (2.19) in (2.17) to get the required result (2.12).

Case II: Note that,  $\Gamma(t+k)$  has a simple pole only for  $k \leq r + \ell - 1$ . For any  $k, \zeta(s+t+k)$  has no pole since

 $s > r + \ell$ .  $\Gamma(t + r - 1)$  has a simple pole.  $\zeta(r + s + t - 1)$  has a simple pole since  $s = \ell + 1$ . Also, observe that, for any  $k, r \in \mathbb{N}$  such that  $k \le r - 1$ , around t = 1 - r, from Laurent series expansions of the  $\Gamma(z)$ , we have

$$\frac{\Gamma(t+r-1)}{\Gamma(t)} = (-1)^{r-1} \frac{(r+\ell-1)!}{\ell!} - (-1)^r \frac{(r+\ell-1)!}{\ell!} (H_\ell - H_{r+\ell-1})(t - (1-r-\ell)) + O(|t-(1-r-\ell)|^2). \tag{2.20}$$

Also, the Laurent series of  $x^{r+t-1}$  with additional term around  $r+t=1-\ell$  is as follows,

$$\frac{1}{x^{t+r-1}} = x^{\ell} - x^{\ell} \log(x)(t - (1 - r - \ell)) + O(|t - (1 - r - \ell)|^2). \tag{2.21}$$

The well-known Laurent series expansion of  $\zeta(z)$ ,

$$\zeta(t+r+\ell) = \frac{1}{(t-(1-r-\ell))} + \gamma - \gamma_1(t-(1-r-\ell)) + O(|t-(1-r-\ell)|^2). \tag{2.22}$$

Combining the equations (2.18), (2.20), (2.21) and (2.22) in (2.17), we get the required result (2.13).

Case III: Note that,  $\Gamma(t+k)$  has a simple pole only for  $k \leq r+\ell-1$ . Observe that  $\zeta(s+t+k)$  has a simple pole only for  $k=r+\ell-s$ , since  $s \leq r+\ell$ . Also note that  $r+\ell-s>r+\ell-1$  since  $s \in \mathbb{Z} \setminus \mathbb{N}$ .  $\Gamma(t+r-1)$  has a simple pole.  $\zeta(r+s+t-1)$  has no pole since  $s \leq r+\ell$ . Hence, from (2.17), we get the required result (2.14).

Case IV: Note that,  $\Gamma(t+k)$  has a simple pole only for  $k \leq r+\ell-1$ . Observe that  $\zeta(s+t+k)$  has a simple pole only for  $k=r+\ell-s$ , since  $s\leq r+\ell$ . Also note that  $r+\ell-s\leq r+\ell-1$  since  $s\in\mathbb{N}$ .  $\Gamma(t+r-1)$  has a simple pole.  $\zeta(r+s+t-1)$  has no pole since  $s\leq r+\ell$ . Hence, from (2.17), we get the required result (2.15).

**Theorem 2.4.** Let  $r \in \mathbb{N}$  and  $\ell \in \mathbb{N} \cup \{0\}$  be arbitrary such that  $\ell \geq 1 - r$ . Then, the following hold. Case I: For a fixed  $s \in \mathbb{Z}$  such that  $s > r + \ell$ , or  $s \in \mathbb{C} \setminus \mathbb{Z}$ , we have,

$$\Theta(r, s, t, x) = {r + \ell - 1 \choose \ell} x^{\ell} \frac{\zeta(s - \ell)}{(t - (1 - r - \ell))} + {r + \ell - 1 \choose \ell} x^{\ell} \zeta(s - \ell) (\gamma + H_{r-1} - H_{r+\ell-1}) 
+ \sum_{\substack{k=0 \\ k \neq r-1}}^{r+\ell-1} {r + \ell - 1 \choose k} x^{r+\ell-1-k} \zeta(r - k) \zeta(1 + k - \ell - r + s) + O(|t - (1 - r - \ell)|).$$
(2.23)

**Case II:** For a fixed  $s \in \mathbb{Z} \setminus \mathbb{N}$  such that  $s \leq r + \ell$ , we have,

$$\Theta(r, s, t, x) = {r + \ell - 1 \choose \ell} x^{\ell} \frac{\zeta(s - \ell)}{(t - (1 - r - \ell))} + {r + \ell - 1 \choose \ell} x^{\ell} \zeta(s - \ell) (\gamma + H_{r-1} - H_{r+\ell-1}) 
+ \sum_{\substack{k=0 \\ k \neq r-1}}^{r+\ell-1} {r + \ell - 1 \choose k} x^{r+\ell-1-k} \zeta(r - k) \zeta(1 + k - \ell - r + s) 
+ (-1)^{s-1} \frac{(r + \ell - 1)!(-s)!}{(r + \ell - s)!} x^{s-1} \zeta(s - \ell) + O(|t - (1 - r - \ell)|).$$
(2.24)

**Case III:** For a fixed  $s \in \mathbb{N}$  such that  $s \leq r + \ell$  and  $s \neq \ell + 1$ , we have,

$$\Theta(r, s, t, x) = {r + \ell - 1 \choose \ell} x^{\ell} \frac{\zeta(s - \ell)}{(t - (1 - r - \ell))} + {r + \ell - 1 \choose \ell} x^{\ell} \zeta(s - \ell) (\gamma + H_{r-1} - H_{r+\ell-1}) 
+ {r + \ell - 1 \choose s - 1} x^{s-1} \frac{\zeta(s - \ell)}{(t - (1 - r - \ell))} + {r + \ell - 1 \choose s - 1} x^{s-1} \zeta(s - \ell) (\gamma + H_{s-1} - H_{r+\ell-1} - \log(x)) 
+ \sum_{\substack{k=0 \\ k \neq r-1 \\ k \neq r+\ell-s}}^{r+\ell-1} {r + \ell - 1 \choose k} x^{r+\ell-1-k} \zeta(r - k) \zeta(1 + k - \ell - r + s) + O(|t - (1 - r - \ell)|).$$
(2.25)

**Case IV:** For a fixed  $s \in \mathbb{N}$  such that  $(s \le r + \ell \text{ and})$   $s = \ell + 1$ , we have

$$\Theta(r, s, t, x) = \binom{r + \ell - 1}{\ell} \frac{2x^{\ell}}{(t - (1 - r - \ell))^{2}} + \binom{r + \ell - 1}{\ell} \frac{x^{\ell} (2\gamma + H_{\ell} + H_{r-1} - 2H_{r+\ell-1} - \log(x))}{(t - (1 - r - \ell))}$$

$$+\frac{1}{3} {r+\ell-1 \choose \ell} x^{\ell} \left(-\pi^{2} + 3(\gamma + H_{\ell} - H_{r+\ell-1} - \log(x))(\gamma + H_{r-1} - H_{r+\ell-1}) + 3\psi'(r+\ell)\right) + \sum_{\substack{k=0 \\ k \neq r-1}}^{r+\ell-1} {r+\ell-1 \choose k} x^{r+\ell-1-k} \zeta(r-k) \zeta(k-r+2) + O(|t-(1-r-\ell)|).$$
(2.26)

*Proof.* Let  $s \in \mathbb{C}$ . Then, for  $t > -r - \ell$ , (2.11) and (2.16) gives,

$$\operatorname{Li}_{r}(e^{-y}) - (-1)^{r-1} (H_{r-1} - \log(y)) \frac{y^{r-1}}{(r-1)!} - \sum_{\substack{k=0\\k \neq r-1}}^{r+\ell + \lfloor |\operatorname{Re}(s)| \rfloor + 1} (-1)^{k} \zeta(r-k) \frac{y^{k}}{k!} = O(y^{\epsilon-1})$$
 (2.27)

for some  $\epsilon > 0$ . As in the previous theorem, for  $M = r + \ell + \lfloor |\text{Re}(s)| \rfloor + 1$ , from (2.9) and (2.27), the integral on the left-hand side of (2.10) is now defined and analytic in the extended region  $\text{Re}(t) > -r - \ell$  by using [22, Theorem 2.2]. Hence, due to the pole of  $\Gamma(t)$  at  $t = 1 - r - \ell$ , we see

$$\lim_{t \to 1-r-\ell} \left( \frac{1}{\Gamma(t)} \int_0^\infty y^{t-1} \mathrm{Li}_s(e^{-xy}) \left( \mathrm{Li}_r(e^{-y}) - (-1)^{r-1} (H_{r-1} - \log(y)) \frac{y^{r-1}}{(r-1)!} - \sum_{\substack{k=0\\k \neq r-1}}^{r+\ell+\lfloor |\mathrm{Re}(s)| \rfloor + 1} (-1)^k \zeta(r-k) \frac{y^k}{k!} \right) dy \right) = 0.$$

Hence, from (2.10),

$$\Theta(r, s, t, x) = \frac{(-1)^{r-1}\Gamma(t+r-1)}{x^{t+r-1}(r-1)!\Gamma(t)} \Big( \zeta(t+s+r-1) \left( H_{r-1} - \psi(t+r-1) + \log(x) \right) - \zeta'(t+s+r-1) \Big) + \sum_{\substack{k=0\\k\neq r-1}}^{r+\ell+\lfloor|\operatorname{Re}(s)|\rfloor+1} \frac{(-1)^k}{k!} \frac{\zeta(r-k)\Gamma(t+k)}{x^{t+k}\Gamma(t)} \zeta(t+k+s) + O(t-(1-r-\ell)).$$
(2.28)

To get the required Kronecker limit formula for  $\Theta(r, s, t, x)$ , we analyze the behaviour of the right-hand side of (2.17) around  $t = 1 - r - \ell$ . Since  $\Gamma(t)$  has a pole at  $t = 1 - r - \ell$ , only the terms with a pole in the numerator survive, as  $t \to 1 - r - \ell$ . We handle the remaining calculations in cases, as mentioned in statement of this Theorem. Case I: Note that,  $\Gamma(t+k)$  has a simple pole only for  $k \le r + \ell - 1$ . Observe that  $\zeta(s+t+k)$  and  $\zeta(r+s+t-1)$  have no poles since  $s > r + \ell$  or  $s \notin \mathbb{Z}$ .  $\Gamma(t+r-1)$  has a simple pole.  $\psi(t+r-1)$  has a simple pole. From (2.28), using (2.18) and (2.20), we get the required result (2.23).

Case II: Note that,  $\Gamma(t+k)$  has a simple pole only for  $k \leq r+\ell-1$ . Observe that  $\zeta(s+t+k)$  has a simple pole for  $k=r+\ell-s$ , moreover  $r+\ell-s>r+\ell-1$ .  $\zeta(r+s+t-1)$  has no pole since  $s\neq \ell+1$ .  $\Gamma(t+r-1)$  has a simple pole.  $\psi(t+r-1)$  has a simple pole. Also observe that the term corresponding to  $k=r+\ell-s$  appears due to the pole of  $\zeta(t+k+s)$  and not  $\Gamma(t+k)$ . From (2.28), we get the required result (2.24).

Case III: Note that,  $\Gamma(t+k)$  has a simple pole only for  $k \leq r+\ell-1$ . Observe that  $\zeta(s+t+k)$  has a simple pole for  $k=r+\ell-s$ , moreover  $r+\ell-s \leq r+\ell-1$ .  $\zeta(r+s+t-1)$  has no pole since  $s \neq \ell+1$ .  $\Gamma(t+r-1)$  has a simple pole.  $\psi(t+r-1)$  has a simple pole. As in the previous case, the term corresponding to  $k=r+\ell-s$  appears due to the pole of  $\zeta(t+k+s)$  and not  $\Gamma(t+k)$ . From (2.28), we get the required result (2.25).

Case IV: Note that,  $\Gamma(t+k)$  has a simple pole only for  $k \leq r+\ell-1$ . Observe that  $\zeta(s+t+k)$  has no pole since  $s=\ell+1$  and  $k\neq r-1$ .  $\zeta(r+s+t-1)$  and  $\zeta'(r+s+t-1)$  have a simple and a double pole respectively since  $s=\ell+1$ .  $\Gamma(t+r-1)$  has a simple pole.  $\psi(t+r-1)$  has a simple pole. Also, use the following power series, around  $t=1-r-\ell$ ,

$$\frac{\Gamma(t+r-1)}{\Gamma(t)} = (-1)^{r-1} \frac{(r+\ell-1)!}{\ell!} - (-1)^r \frac{(r+\ell-1)!}{\ell!} (\psi(\ell+1) - \psi(r+\ell))(t - (1-r-\ell)) 
- (-1)^r \frac{(r+\ell-1)!}{2\ell!} ((\psi(\ell+1) - \psi(r+\ell))^2 - \psi'(\ell+1) + \psi'(r+\ell)) (t - (1-r-\ell))^2 + O(|(t-(1-r-\ell))|^3),$$
(2.29)

$$\frac{1}{x^{t+r-1}} = x^{\ell} - x^{\ell} \log(x)(t - (1-r-\ell)) + \frac{1}{2}x^{\ell} \log^2(x)(t - (1-r-\ell))^2 + O(|(t - (1-r-\ell))|^3). \tag{2.30}$$

Use (2.29) and (2.30) in (2.28) to get the required result (2.26).

Next, we are interested to get Kronecker limit type formula for  $\Theta(r, s, t, x)$  in the third variable t, around t = 2 - r - s. We can obtain such a result for fixed  $r, s \in \mathbb{C}$  such that  $r + s \in \mathbb{N}$  and  $r + s \geq 2$  in the following way:

- $r, s \in \mathbb{N}$ : It reduces to case IV of Theorem 2.4 upon taking  $s = \ell + 1$ .
- $s \in \mathbb{N}$  and  $r \in \mathbb{Z} \setminus \mathbb{N}$ : It reduces to case II of Theorem 2.3 upon taking  $s = \ell + 1$ .
- $r \in \mathbb{N}$  and  $s \in \mathbb{Z} \setminus \mathbb{N}$ : It reduces to the previous case,  $s \in \mathbb{N}$  and  $r \in \mathbb{Z} \setminus \mathbb{N}$ , after using (1.5).

Hence, for a fixed  $r, s \in \mathbb{C} \setminus \mathbb{Z}$ , we state the formula as the following theorem.

**Theorem 2.5.** Let  $r, s \in \mathbb{C} \setminus \mathbb{Z}$  such that  $r + s \in \mathbb{N}$  and  $r + s \geq 2$ . Around t = 2 - r - s, we have

$$\Theta(r, s, t, x) = (-1)^{r+s} x^{s-1} (r+s-2)! \Gamma(1-r) \Gamma(1-s)$$

$$+ \sum_{k=0}^{r+s-2} {r+s-2 \choose k} x^{r+s-2-k} \zeta(r-k) \zeta(k-r+2) + O(|t-(2-r-s)|).$$
(2.31)

*Proof.* Then, for Re(t) > 1 - r - s, (2.6) and (2.7) gives,

$$y^{t-1} \operatorname{Li}_{s}(e^{-xy}) \left( \operatorname{Li}_{r}(e^{-y}) - \Gamma(1-r) y^{r-1} - \sum_{k=0}^{r+s+\lfloor |\operatorname{Re}(s)| \rfloor} (-1)^{k} \zeta(r-k) \frac{y^{k}}{k!} \right) = O\left(y^{\epsilon-1}\right), \tag{2.32}$$

for some  $\epsilon > 0$ . For  $M = r + s + \lfloor |\text{Re}(s)| \rfloor$ , from (2.9) and (2.32), the integral on the left-hand side of (2.1) is now defined and analytic in the extended region Re(t) > 1 - r - s by using [22, Theorem 2.2]. Hence, due to the pole of  $\Gamma(t)$  at t = 2 - r - s, we see

$$\lim_{t \to 2-r-s} \left( \frac{1}{\Gamma(t)} \int_0^\infty y^{t-1} \mathrm{Li}_s(e^{-xy}) \left( \mathrm{Li}_r(e^{-y}) - \Gamma(1-r) y^{r-1} - \sum_{k=0}^{r+s+\lfloor |\mathrm{Re}(s)| \rfloor} (-1)^k \zeta(r-k) \frac{y^k}{k!} \right) dy \right) = 0.$$

Hence, from (2.1)

$$\Theta(r, s, t, x) = \frac{\Gamma(1 - r)\Gamma(t + r - 1)}{x^{t + r - 1}\Gamma(t)} \zeta(r + s + t - 1) 
+ \sum_{k=0}^{r + s + \lfloor |\operatorname{Re}(s)| \rfloor} \frac{(-1)^k \zeta(r - k)\Gamma(t + k)}{x^{t + k}k!\Gamma(t)} \zeta(s + t + k) + O(|t - (2 - r - s)|).$$
(2.33)

Since  $\Gamma(t)$  has a pole at t=2-r-s, only the terms with a pole in the numerator survive, as  $t\to 2-r-s$ . We list the poles of numerator at t=2-r-s: Note that,  $\Gamma(t+k)$  has a simple pole only for  $k\le r+s-2$ . For any k,  $\zeta(s+t+k)$  has no pole since  $r\notin\mathbb{Z}$ .  $\Gamma(t+r-1)$  has no pole since  $s\notin\mathbb{Z}$ .  $\Gamma(t+s+t-1)$  has a simple pole. Hence, use the laurent series expansions of the functions on the right-hand side of (2.33) to get the required result (2.31).

#### 3. Applications in modular relations

We start this Section by proving a recursive formula, Theorem 3.1, for the generalized Mordell-Tornheim zeta function  $\Theta(r, s, t, x)$ . The motivation for this result originated from a recursive result for the Mordell-Tornheim zeta function  $\zeta_{\text{MT}}(r, s, t)$  obtained by Huard, Williams and Zhang [12, Equation (1.6)]. We use the specific case of Theorem 3.1, with s = r = n, as stated in Corollary 3.2, to then obtain the two-term functional equation given by Vlasenko-Zagier (1.18) as a further implication.

**Theorem 3.1.** For any  $r, s, t \in \mathbb{R}$  such that r + s + t > 2, r + t > 1 and s + t > 1 and for any  $n \in \mathbb{N}$ , we have,

$$\Theta(r, s, t, x) = \sum_{\ell=0}^{n} \binom{n}{\ell} x^{\ell} \Theta(r - n + \ell, s - \ell, t + n, x).$$

$$(3.1)$$

*Proof.* For n = 0, it is vacuously true. For n = 1, we can easily see that,

$$\Theta(r, s, t, x) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{n^{r-1} m^s (n + mx)^{t+1}} + x \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{n^r m^{s-1} (n + mx)^{t+1}} 
= \Theta(r - 1, s, t + 1, x) + x \Theta(r, s - 1, t + 1, x).$$
(3.2)

Hence, the statement holds for n = 1. We now assume the identity is true for all j < n and then prove that the statement holds for j = n. Since the statement holds for j = n - 1, we have,

$$\Theta(r, s, t, x) = \sum_{\ell=0}^{n-1} {n-1 \choose \ell} x^{\ell} \Theta(r - (n-1) + \ell, s - \ell, t + n - 1, x).$$

Use (3.2) inside the summation to get,

$$\Theta(r, s, t, x) = \sum_{\ell=0}^{n-1} \binom{n-1}{\ell} x^{\ell} \left( \Theta(r-n+\ell, s-\ell, t+n, x) + x \Theta(r-(n-1)+\ell, s-\ell-1, t+n, x) \right)$$

$$= \sum_{\ell=0}^{n-1} \binom{n-1}{\ell} x^{\ell} \Theta(r-n+\ell, s-\ell, t+n, x) + x \sum_{\ell=0}^{n-1} \binom{n-1}{\ell} x^{\ell} \Theta(r-(n-1)+\ell, s-\ell-1, t+n, x)$$

Replace the variable of summation  $\ell$  by  $\ell-1$  in the second term to get,

$$\begin{split} \Theta(r, s, t, x) &= \sum_{\ell=0}^{n-1} \binom{n-1}{\ell} x^{\ell} \Theta(r - n + \ell, s - \ell, t + n, x) + \sum_{\ell=1}^{n} \binom{n-1}{\ell-1} x^{\ell} \Theta(r - n + \ell, s - \ell, t + n, x) \\ &= \Theta(r - \ell, s, t + n, x) + \sum_{\ell=1}^{n-1} \left( \binom{n-1}{\ell} + \binom{n-1}{\ell-1} \right) x^{\ell} \Theta(r - n + \ell, s - \ell, t + n, x) + x^{n} \Theta(r, s - n, t + n, x) \\ &= \Theta(r - \ell, s, t + n, x) + \sum_{\ell=1}^{n-1} \binom{n}{\ell} x^{\ell} \Theta(r - n + \ell, s - \ell, t + n, x) + x^{n} \Theta(r, s - n, t + n, x) \\ &= \sum_{\ell=0}^{n} \binom{n}{\ell} x^{\ell} \Theta(r - n + \ell, s - \ell, t + n, x), \end{split}$$

where we have used the property  $\binom{n-1}{\ell} + \binom{n-1}{\ell-1} = \binom{n}{\ell}$  of the Binomial coefficients in the second last step. Observe that  $\Theta$  by definition is well defined at the arguments in all of the above steps. Hence, by the Principle of Mathematical Induction, (3.1) is true for all  $n \in \mathbb{N}$ .

**Corollary 3.2.** For  $r, t \in \mathbb{R}$  such that r + t > 1 and 2r + t > 2, and for any x > 0.

$$\Theta(r, r, t, x) = \sum_{\ell=0}^{r} \binom{r}{\ell} x^{\ell} \Theta(\ell, r - \ell, t + r, x).$$

*Proof.* Put s = r and n = r in (3.1) to get this result.

We now obtain the functional equations proved by Vlasenko and Zagier [24, Equation 11] as an application of the above Corollary 3.2.

## 3.1. Vlasenko and Zagier's functional equation.

Corollary 3.3. Functional equation in (1.18) given by Vlasenko-Zagier holds.

*Proof.* Let  $r \in \mathbb{N}$ ,  $r \neq 1$  be arbitrary and t > 1 - r. Let x > 0. From Corollary 3.2, we have,

$$\Theta(r, r, t, x) = \sum_{\ell=0}^{r} {r \choose \ell} x^{\ell} \Theta(\ell, r - \ell, t + r, x).$$

Separate out the first and the last term from the sum to get,

$$\Theta(r, r, t, x) = \Theta(0, r, t + r, x) + \sum_{\ell=1}^{r-1} \binom{r}{\ell} x^{\ell} \Theta(\ell, r - \ell, t + r, x) + x^{r} \Theta(r, 0, t + r, x).$$

Use inversion (1.5) for the third term on the right-hand side to get,

$$\Theta(r, r, t, x) = \Theta(0, r, t + r, x) + S(r, t) + x^{-t}\Theta(0, r, t + r, \frac{1}{x}),$$
(3.3)

where S(r,t) is defined by

$$S(r,t) := \sum_{\ell=1}^{r-1} \binom{r}{\ell} x^{\ell} \Theta(\ell, r - \ell, t + r, x).$$

We now simplify S(r,t) as follows:

$$S(r,t) = \sum_{\ell=1}^{r-1} \left( \sum_{i=1}^{\ell} (-1)^{\ell-i} {r \choose i} - \sum_{j=1}^{\ell-1} (-1)^{\ell-j} {r \choose j} \right) x^{\ell} \Theta(\ell, r - \ell, t + r, x)$$

$$= \sum_{\ell=1}^{r-1} \sum_{i=1}^{\ell} (-1)^{\ell-i} {r \choose i} x^{\ell} \Theta(\ell, r - \ell, t + r, x) - \sum_{\ell=1}^{r-1} \sum_{j=1}^{\ell-1} (-1)^{\ell-j} {r \choose j} x^{\ell} \Theta(\ell, r - \ell, t + r, x).$$

Replace  $\ell$  by  $\ell+1$  in the second term of the right-hand side to get,

$$S(r,t) = \sum_{\ell=1}^{r-1} \sum_{i=1}^{\ell} (-1)^{\ell-i} \binom{r}{i} x^{\ell} \Theta(\ell, r-\ell, t+r, x) + \sum_{\ell=0}^{r-2} \sum_{j=1}^{\ell} (-1)^{\ell-j} \binom{r}{j} x^{\ell+1} \Theta(\ell+1, r-\ell-1, t+r, x).$$

Since  $\ell = 0$  makes the summand zero, hence the second term can be rewritten as,

$$S(r,t) = \sum_{\ell=1}^{r-1} \sum_{i=1}^{\ell} (-1)^{\ell-i} \binom{r}{i} x^{\ell} \Theta(\ell,r-\ell,t+r,x) + \sum_{\ell=1}^{r-2} \sum_{j=1}^{\ell} (-1)^{\ell-j} \binom{k}{j} x^{\ell+1} \Theta(\ell+1,r-\ell-1,t+r,x).$$

Separate the  $\ell = r - 1$  term from the first term on the right-hand side to get,

$$S(r,t) = x^{r-1}\Theta(r-1,1,t+r,x) \sum_{i=1}^{r-1} (-1)^{r-i+1} \binom{r}{i}$$

$$+ \sum_{\ell=1}^{r-2} \sum_{i=1}^{\ell} (-1)^{\ell-i} \binom{r}{i} x^{\ell} \Theta(\ell,r-\ell,t+r,x) + \sum_{\ell=1}^{r-2} \sum_{j=1}^{\ell} (-1)^{\ell-j} \binom{k}{j} x^{\ell+1} \Theta(\ell+1,r-\ell-1,t+r,x)$$

$$= x^{r-1}\Theta(r-1,1,t+r,x) \sum_{i=1}^{r-1} (-1)^{r-i+1} \binom{r}{i}$$

$$+ \sum_{\ell=1}^{r-2} \sum_{i=1}^{\ell} (-1)^{\ell-i} \binom{r}{i} x^{\ell} \left( \Theta(\ell,r-\ell,t+r,x) + x\Theta(\ell+1,r-\ell-1,t+r,x) \right)$$

$$= x^{r-1}\Theta(r-1,1,t+r,x) \sum_{j=1}^{r-1} (-1)^{r-i+1} \binom{r}{j} + \sum_{j=1}^{r-2} \sum_{j=1}^{\ell} (-1)^{\ell-i} \binom{r}{j} x^{\ell} \Theta(\ell+1,r-\ell,t+r-1,x), \quad (3.4)$$

where we have used (1.4) in the last step. We note the following fact obtained by the Binomial theorem,

$$\sum_{i=1}^{r-1} (-1)^{r-i+1} \binom{r}{i} \stackrel{i \to r-i}{=} -\sum_{i=1}^{r-1} (-1)^i \binom{r}{i} = 1 + (-1)^r = \begin{cases} 0, & \text{if r is odd,} \\ 2, & \text{if r is even.} \end{cases}$$

Also, we know that

$$\sum_{i=1}^{\ell} (-1)^{\ell-i} \binom{r}{i} = (-1)^{\ell} \sum_{i=1}^{\ell} (-1)^{i} \binom{r}{i} = (-1)^{\ell} \left(-1 + \sum_{i=0}^{\ell} (-1)^{i} \binom{r}{i}\right) = (-1)^{\ell+1} + \binom{r-1}{\ell},$$

where we have used that  $\sum_{i=0}^{\ell} (-1)^i \binom{r}{i} = (-1)^{\ell} \binom{r-1}{\ell}$ . Hence, using the two facts mentioned above, (3.4) becomes,

$$S(r,t) = (1 + (-1)^r)x^{r-1}\Theta(r-1,1,t+r,x) + \sum_{\ell=1}^{r-2} \left( (-1)^{\ell+1} + \binom{r-1}{\ell} \right) x^{\ell}\Theta(\ell+1,r-\ell,t+r-1,x).$$
 (3.5)

Take the limit  $t \to 1 - r$  in (3.5) to get,

$$\lim_{t \to 1-r} S(r,t) = (1+(-1)^r)x^{r-1} \lim_{t \to 1-r} \Theta(r-1,1,t+r,x) + \sum_{\ell=1}^{r-2} \left( (-1)^{\ell+1} + \binom{r-1}{\ell} \right) x^{\ell} \zeta(\ell+1)\zeta(r-\ell), \quad (3.6)$$

since  $\Theta(a, b, 0, x) = \zeta(a)\zeta(b)$  for any a, b > 1. The limit on right-hand side is dealt as follows,

$$\lim_{t\rightarrow 1-r}\Theta(r-1,1,t+r,x)=\lim_{t\rightarrow 1-r}x^{-t-r}\Theta(1,r-1,t+r,\tfrac{1}{x})$$

$$= \lim_{t \to 1-r} \left( x^{-t-r} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{nm^{r-1}(n + \frac{m}{x})^{t+r}} \right)$$

$$= \frac{1}{x} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{nm^{r-1}(n + \frac{m}{x})}$$

$$= \frac{1}{x} \sum_{m=1}^{\infty} \frac{1}{m^{r-1}} \sum_{n=1}^{\infty} \frac{1}{n(n + \frac{m}{x})}$$

$$= \frac{1}{x} \sum_{m=1}^{\infty} \frac{1}{m^{r-1}} \sum_{n=1}^{\infty} \left( \frac{\frac{x}{m}}{n} - \frac{\frac{x}{m}}{n + \frac{m}{x}} \right)$$

$$= \sum_{m=1}^{\infty} \frac{1}{m^{r}} \left( \gamma + \psi(\frac{m}{x} + 1) \right)$$

$$= \gamma \zeta(r) + \sum_{m=1}^{\infty} \frac{\psi(\frac{m}{x} + 1)}{m^{r}}$$
(3.7)

Hence, from (3.6) and (3.7), we get,

$$\lim_{t \to 1-r} S(r,t) = (1+(-1)^r)x^{r-1} \left( \sum_{m=1}^{\infty} \frac{\psi(\frac{m}{x}+1)}{m^r} + \gamma \zeta(r) \right) + \sum_{\ell=1}^{r-2} \left( (-1)^{\ell+1} + \binom{r-1}{\ell} \right) x^{\ell} \zeta(\ell+1) \zeta(r-\ell). \quad (3.8)$$

We also need the following evaluation,

$$\Theta(0, r, t + r, x) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{m^r (n + mx)^{t+r}}$$

$$= \sum_{m=1}^{\infty} \frac{1}{m^r} \sum_{n=1}^{\infty} \frac{1}{(n + mx)^{t+r}}$$

$$= \sum_{m=1}^{\infty} \frac{\zeta(t + r, mx + 1)}{m^r}$$

$$= \sum_{m=1}^{\infty} \frac{\zeta(t + r, mx + 1) - \frac{1}{t - 1 + r}}{m^r} + \frac{\zeta(r)}{t - 1 + r}$$
(3.9)

Hence, we can also see that,

$$x^{-t}\Theta(0,r,t+r,\frac{1}{x}) = x^{-t} \left( \sum_{m=1}^{\infty} \frac{\zeta(t+r,\frac{m}{x}+1) - \frac{1}{t-1+r}}{m^r} + \frac{\zeta(r)}{t-1+r} \right).$$
 (3.10)

Substitute (3.9) and (3.10) in (3.3) to get.

$$\Theta(r, r, t, x) - \frac{\zeta(r)(1 + x^{-t})}{t - 1 + r} = \sum_{m=1}^{\infty} \frac{\zeta(t + r, mx + 1) - \frac{1}{t - 1 + r}}{m^r} + S(r, t) + x^{-t} \sum_{m=1}^{\infty} \frac{\zeta(t + r, \frac{m}{x} + 1) - \frac{1}{t - 1 + r}}{m^r}.$$

Take the limit of both sides of the above equation as  $t \to 1-r$ , and use (3.8) to get

$$\lim_{t \to 1-r} \left( \Theta(r, r, t, x) - \frac{\zeta(r)(1 + x^{-t})}{t - 1 + r} \right)$$

$$= -\sum_{m=1}^{\infty} \frac{\psi(mx+1)}{m^r} + \lim_{t \to 1-r} (S(r, t)) - x^{r-1} \sum_{m=1}^{\infty} \frac{\psi(\frac{m}{x} + 1)}{m^r}$$

$$= -\sum_{m=1}^{\infty} \frac{\psi(mx+1)}{m^r} + (1 + (-1)^r) x^{r-1} \left( \sum_{m=1}^{\infty} \frac{\psi(\frac{m}{x} + 1)}{m^r} + \gamma \zeta(r) \right)$$

$$+\sum_{\ell=1}^{r-2} \left( (-1)^{\ell+1} + {r-1 \choose \ell} \right) x^{\ell} \zeta(\ell+1) \zeta(r-\ell) - x^{r-1} \sum_{m=1}^{\infty} \frac{\psi(\frac{m}{x}+1)}{m^r}$$

$$= -\sum_{m=1}^{\infty} \frac{\psi(mx+1)}{m^r} + (-1)^r x^{r-1} \sum_{m=1}^{\infty} \frac{\psi(\frac{m}{x}+1)}{m^r} + (1+(-1)^r) x^{r-1} \gamma \zeta(r)$$

$$+\sum_{\ell=1}^{r-2} \left( (-1)^{\ell+1} + {r-1 \choose \ell} \right) x^{\ell} \zeta(\ell+1) \zeta(r-\ell). \tag{3.11}$$

Put s = r and  $\ell = 0$  in case 3 of Theorem 2.4 to get,

$$\Theta(r, r, t, x) = \frac{\zeta(r) (1 + x^{r-1})}{(t - 1 + r)} + x^{r-1} \zeta(r) (\gamma - \log(x))$$
$$+ \gamma \zeta(r) + x^{r-1} \sum_{k=1}^{r-2} {r-1 \choose k} \frac{1}{x^k} \zeta(r - k) \zeta(k+1) + O(|t - 1 + r|).$$

Upon adding and subtracting suitable terms, we get,

$$\lim_{t \to 1-r} \left( \Theta(r, r, t, x) - \frac{\zeta(r)(1+x^{-t})}{t-1+r} \right) \\
= \lim_{t \to 1-r} \left( \frac{\zeta(r) \left( x^{r-1} - x^{-t} \right)}{t-1+r} \right) + x^{r-1} \zeta(r) (\gamma - \log(x)) + \gamma \zeta(r) + x^{r-1} \sum_{k=1}^{r-2} {r-1 \choose k} \frac{1}{x^k} \zeta(r-k) \zeta(k+1) \\
= x^{r-1} \zeta(r) \log(x) + x^{r-1} \zeta(r) (\gamma - \log(x)) + \gamma \zeta(r) + x^{r-1} \sum_{k=1}^{r-2} {r-1 \choose k} \frac{1}{x^k} \zeta(r-k) \zeta(k+1) \\
= x^{r-1} \zeta(r) \gamma + \zeta(r) \gamma + x^{r-1} \sum_{k=1}^{r-2} {r-1 \choose k} \frac{1}{x^k} \zeta(r-k) \zeta(k+1) \\
= x^{r-1} \zeta(r) \gamma + \zeta(r) \gamma + \sum_{k=1}^{r-2} {r-1 \choose k} x^k \zeta(r-k) \zeta(k+1), \tag{3.12}$$

where we have replaced k by r-1-k in the last step. Hence, from (3.11) and (3.12), we see,

$$-\sum_{m=1}^{\infty} \frac{\psi(mx+1)}{m^r} + (-1)^r x^{r-1} \sum_{m=1}^{\infty} \frac{\psi(\frac{m}{x}+1)}{m^r} + (1+(-1)^r) x^{r-1} \gamma \zeta(r) + \sum_{\ell=1}^{r-2} \left( (-1)^{\ell+1} + \binom{r-1}{\ell} \right) x^{\ell} \zeta(\ell+1) \zeta(r-\ell) = x^{r-1} \zeta(r) \gamma + \zeta(r) \gamma + \sum_{k=1}^{r-2} \binom{r-1}{k} x^k \zeta(r-k) \zeta(k+1).$$

On cancellation and simplification, we get, with the convention that  $\zeta(1) = \gamma$  as taken by Vlasenko-Zagier,

$$\sum_{m=1}^{\infty} \frac{\psi(mx+1)}{m^r} + (-x)^{r-1} \sum_{m=1}^{\infty} \frac{\psi(\frac{m}{x}+1)}{m^r} = (-1)^r x^{r-1} \gamma \zeta(r) - \sum_{\ell=1}^{r-2} (-1)^{\ell} x^{\ell} \zeta(\ell+1) \zeta(r-\ell) - \zeta(r) \gamma.$$

$$= -\sum_{\ell=0}^{r-1} (-x)^{\ell} \zeta(\ell+1) \zeta(r-\ell).$$

Use the functional equation  $\psi(y+1) = \psi(y) + y^{-1}$  in both the summands on the left-hand side of the above equation and simplify to get (1.18).

## 3.2. Guinand's functional equations.

Corollary 3.4. Functional equations in (1.11) and (1.12) given by Guinand hold.

*Proof.* We use the inversion formula (1.5) with r = s = 0 and  $t \in \mathbb{R}$  such that t > 2,

$$\Theta(0,0,t,x) = x^{-t}\Theta(0,0,t,\frac{1}{x}). \tag{3.13}$$

From [1, Equation 6.4.10, p.260], for  $j \in \mathbb{N}$  and  $\psi^{(j)}$  being the  $j^{th}$  derivative of  $\psi$ , we have,

$$\psi^{(j)}(z) = (-1)^{j+1} j! \sum_{\ell=0}^{\infty} \frac{1}{(\ell+z)^{j+1}} = (-1)^{j+1} j! \zeta(j+1,z), \tag{3.14}$$

where  $\zeta(s,a)$  is the Hurwitz zeta function. Then, from (3.13), we have

$$\sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{(n+mx)^t} = x^{-t} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{(n+\frac{m}{x})^t}.$$
 (3.15)

Replace n by n-1 in both the sums on left-hand and right-hand side of the equation and then use (3.14) to see that, for  $t \in \mathbb{N}$  such that  $t \geq 3$ ,

$$(-1)^{t}(t-1)! \sum_{m=1}^{\infty} \psi^{(t-1)}(1+mx) = (-1)^{t}(t-1)! x^{-t} \sum_{m=1}^{\infty} \psi^{(t-1)}\left(1+\frac{m}{x}\right).$$

Cancel  $(-1)^t(t-1)!$  on both sides and then multiply both sides by  $x^{\frac{t}{2}}$  to get (1.11). We now prove the second equation of Guinand (1.12). From the definition of the Hurwitz zeta function,  $\zeta(s,a) = \sum_{n=0}^{\infty} (n+a)^{-s}$ , and (3.15), we can see,

$$\sum_{m=1}^{\infty} \zeta(t, mx + 1) = x^{-t} \sum_{m=1}^{\infty} \zeta(t, \frac{m}{x} + 1).$$
(3.16)

From [4, Equation 4.3], we have,

$$\zeta(z,x) = \frac{x^{1-z}}{z-1} + O(x^{-z}).$$

Adding and subtracting suitable terms from the summands on both left-hand side and right-hand side, (3.16) becomes,

$$\sum_{m=1}^{\infty} \left( \zeta(t, mx+1) + \frac{(mx)^{1-t}}{t-1} - \frac{(mx)^{1-t}}{t-1} \right) = x^{-t} \sum_{m=1}^{\infty} \left( \zeta(t, \frac{m}{x}+1) + \frac{\left(\frac{m}{x}\right)^{1-t}}{t-1} - \frac{\left(\frac{m}{x}\right)^{1-t}}{t-1} \right).$$

Simplify to get

$$\sum_{m=1}^{\infty} \left( \zeta(t, mx+1) - \frac{(mx)^{1-t}}{t-1} \right) + \frac{x^{1-t}\zeta(t-1)}{t-1} = x^{-t} \sum_{m=1}^{\infty} \left( \zeta(t, \frac{m}{x}+1) - \frac{\left(\frac{m}{x}\right)^{1-t}}{t-1} \right) + \frac{\zeta(t-1)}{x(t-1)}. \tag{3.17}$$

See the following limits hold, where we use (3.14) for the first one

$$\lim_{t \to 2} \left( \zeta(t, y+1) - \frac{y^{1-t}}{t-1} \right) = \zeta(2, y+1) - \frac{1}{y} = \psi'(1+y) - \frac{1}{y}, \tag{3.18}$$

$$\lim_{t \to 2} \left( \left( x^{1-t} - \frac{1}{x} \right) \frac{\zeta(t-1)}{t-1} \right) = -\frac{\log(x)}{x} = -2\frac{\log(x)}{2x} = -\frac{\log(x)}{2x} + \frac{\log(\frac{1}{x})}{2x}. \tag{3.19}$$

Tend  $t \to 2$  in (3.17), use (3.18) and (3.19) and then multiply both the left-hand side and right-hand side of the resultant by x to get (1.12).

#### 3.3. Ramanujan's functional equation.

Corollary 3.5. The first equality in the functional equation given by Ramanujan in (1.10) holds.

*Proof.* From (3.16) we have,

$$\sum_{m=1}^{\infty} \zeta(t, mx + 1) = x^{-t} \sum_{m=1}^{\infty} \zeta(t, \frac{m}{x} + 1).$$

From [4, Equation 4.3], we have,

$$\zeta(z,x) = \frac{x^{1-z}}{z-1} + \frac{1}{2}x^{-z} + O(x^{-z-1}).$$

Adding and subtracting suitable terms from the summands on both left-hand side and right-hand side, we deduce,

$$\sum_{m=1}^{\infty} \left( \zeta(t, mx + 1) + \frac{(mx)^{1-t}}{t - 1} + \frac{1}{2} (mx)^{-t} - \frac{(mx)^{1-t}}{t - 1} - \frac{1}{2} (mx)^{-t} \right)$$

$$= x^{-t} \sum_{m=1}^{\infty} \left( \zeta(t, \frac{m}{x} + 1) + \frac{\left(\frac{m}{x}\right)^{1-t}}{t - 1} + \frac{1}{2} \left(\frac{m}{x}\right)^{-t} - \frac{\left(\frac{m}{x}\right)^{1-t}}{t - 1} - \frac{1}{2} \left(\frac{m}{x}\right)^{-t} \right). \tag{3.20}$$

Then, on using  $\zeta(z, a+1) = \zeta(z, a) - a^{-z}$ , (3.20) becomes,

$$\begin{split} &\sum_{m=1}^{\infty} \left( \zeta(t, mx) + \frac{(mx)^{1-t}}{t-1} + \frac{1}{2} (mx)^{-t} - \frac{(mx)^{1-t}}{t-1} - \frac{1}{2} (mx)^{-t} - (mx)^{-t} \right) \\ &= x^{-t} \sum_{m=1}^{\infty} \left( \zeta(t, \frac{m}{x}) + \frac{\left(\frac{m}{x}\right)^{1-t}}{t-1} + \frac{1}{2} \left(\frac{m}{x}\right)^{-t} - \frac{\left(\frac{m}{x}\right)^{1-t}}{t-1} - \frac{1}{2} \left(\frac{m}{x}\right)^{-t} - \left(\frac{m}{x}\right)^{-t} \right). \end{split}$$

Let us denote  $\varphi(t,x)$  as follows:

$$\varphi(t,x) := \zeta(t,x) - \frac{x^{1-t}}{t-1} - \frac{1}{2}x^{-t}.$$

Then, we see.

$$\left(\sum_{m=1}^{\infty}\varphi(t,mx)+\frac{x^{1-t}}{t-1}\zeta(t-1)-\frac{1}{2}x^{-t}\zeta(t)\right)=x^{-t}\left(\sum_{m=1}^{\infty}\varphi(t,\frac{m}{x})+\frac{\left(\frac{1}{x}\right)^{1-t}}{t-1}\zeta(t-1)-\frac{1}{2}\left(\frac{1}{x}\right)^{-t}\zeta(t)\right).$$

Take the second term on the left-hand side to the right-hand side and vice versa, to get,

$$\left(\sum_{m=1}^{\infty} \varphi(t, mx) - \frac{\frac{1}{x}}{t-1} \zeta(t-1) - \frac{1}{2} x^{-t} \zeta(t)\right) = x^{-t} \left(\sum_{m=1}^{\infty} \varphi(t, \frac{m}{x}) - \frac{x}{t-1} \zeta(t-1) - \frac{1}{2} \left(\frac{1}{x}\right)^{-t} \zeta(t)\right).$$

Multiply both sides by  $x^{\frac{t}{2}}$  to get the first equality of [4, Theorem 4.1]. As shown in [4, Corollary 4.2], letting  $t \to 1$ , we get the first equality of the Ramanujan's transformation formula (1.10).

# 4. Kronecker limit type formula for $\Theta(r, s, t, x)$ in the Second variable

The method used in Section 2 cannot be used to find Kronecker limit type formula for  $\Theta(r, s, t, x)$  in the second variable s. Hence, we come with an alternative method to obtain the same, which involves the series evaluations of  $\Theta(r, s, t, x)$  in terms of Herglotz-Zagier function (1.13) and higher Herglotz functions (1.17). The evaluation is obtained by the partial fraction method, which is very effective in the theory of multiple zeta functions. For instance, Gangl, Kaneko and Zagier [7] used the partial fraction, for integers  $i, j \geq 2$ ,

$$\frac{1}{m^i n^j} = \sum_{a+b=i+j} \left( \binom{a-1}{i-1} \frac{1}{(m+n)^a n^b} + \binom{a-1}{j-1} \frac{1}{(m+n)^a n^b} \right),$$

to prove the Euler decomposition formula.

$$\zeta(i)\zeta(j) = \sum_{a+b=i+j} \left( \binom{a-1}{i-1} \zeta(a,b) + \binom{a-1}{j-1} \zeta(a,b) \right),$$

where  $\zeta(s_1, s_2)$  is the double zeta function as defined in (1.20). Guo and Xie [9] used the partial fraction method to obtain natural shuffle algebra structure for the integral representations of the multiple-zeta values. We prove the partial fraction in the following Lemma 4.1 before using it to obtain the series evaluation.

**Lemma 4.1.** Fix  $r, t \in \mathbb{N} \cup \{0\}$  such that  $r + t \geq 1$ . Then, for n, y > 0, the following partial fraction holds,

$$\frac{1}{n^r(n+y)^t} = (-1)^r \sum_{j=0}^{t-1} {j+r-1 \choose j} \frac{1}{y^{j+r}(n+y)^{t-j}} + \sum_{i=0}^{r-1} {i+t-1 \choose i} \frac{(-1)^i}{n^{r-i}y^{t+i}}.$$
 (4.1)

with the convention that  $\binom{-1}{0} = 1$ ,  $\binom{c}{-1} = 0$  for any  $c \in \mathbb{Z} \setminus \{-1\}$  with an exception  $\binom{-1}{-1} = 1$ .

*Proof.* Start with the expression on the right-hand side of (4.1) and take least common denominator to get,

$$(-1)^{r} \sum_{j=0}^{t-1} {j+r-1 \choose j} \frac{y^{t-j}(n+y)^{j}}{y^{r+t}(n+y)^{t}} + \sum_{i=0}^{r-1} {i+t-1 \choose i} \frac{(-1)^{i}n^{i}y^{r-i}}{n^{r}y^{r+t}}$$

$$= (-1)^{r} \sum_{j=0}^{t-1} {j+r-1 \choose j} \frac{n^{r}y^{t-j}(n+y)^{j}}{n^{r}y^{r+t}(n+y)^{t}} + \sum_{i=0}^{r-1} {i+t-1 \choose i} \frac{(-1)^{i}n^{i}y^{r-i}(n+y)^{t}}{n^{r}y^{r+t}(n+y)^{t}}$$

$$= \frac{A(n,y,r,t)}{n^{r}y^{r+t}(n+y)^{t}}, \tag{4.2}$$

where

$$A(n,y,r,t) := (-1)^r n^r \sum_{j=0}^{t-1} \binom{j+r-1}{j} y^{t-j} (n+y)^j + (n+y)^t \sum_{i=0}^{r-1} \binom{i+t-1}{i} (-1)^i n^i y^{r-i}.$$

Clearly, for a fixed  $r, t \in \mathbb{N}$ , A(n, y, r, t) is a polynomial of two variables n, y.

$$A(n,y,r,t) = (-1)^{r} n^{r} \sum_{j=0}^{t-1} {j+r-1 \choose j} y^{t-j} \sum_{c=0}^{j} {j \choose c} n^{c} y^{j-c} + \sum_{d=0}^{t} {t \choose d} n^{d} y^{t-d} \sum_{i=0}^{r-1} {i+t-1 \choose i} (-1)^{i} n^{i} y^{r-i}$$

$$= (-1)^{r} \sum_{j=0}^{t-1} \sum_{c=0}^{j} {j+r-1 \choose j} {j \choose c} n^{r+c} y^{t-c} + \sum_{d=0}^{t} \sum_{i=0}^{r-1} {i+t-1 \choose i} {t \choose d} (-1)^{i} n^{d+i} y^{t-d+r-i}$$

$$= (-1)^{r} \sum_{c=0}^{t-1} n^{r+c} \sum_{j=c}^{t-1} {j+r-1 \choose j} {j \choose c} y^{t-c} + \sum_{d=0}^{t} \sum_{i=0}^{r-1} {i+t-1 \choose i} {t \choose d} (-1)^{i} n^{d+i} y^{t-d+r-i}. \tag{4.3}$$

A(n,y,r,t) can be seen as a polynomial in n with coefficients as polynomials in y. Clearly, the highest degree of n possible in A(n,y,r,t) is r+t-1, where  $r,t\in\mathbb{N}$  are fixed. We now show explicitly evaluate the coefficients of  $n^a$  for  $1\leq a\leq r+t-1$  in two cases.

**Case 1:** For  $1 \le a \le r - 1$ .

Observe that the first term on the right-hand side of (4.3) does not contribute to the coefficient of  $n^a$  since least power of n possible is r. From the second term, the only way to get  $n^a$  is when  $d \le a$  and i = a - d. Hence we get,

Coefficient of 
$$n^a$$
 in  $A(n, y, r, t) = \sum_{d=0}^a \binom{a-d+t-1}{a-d} \binom{t}{d} (-1)^{a-d} y^{t+r-a}$ 

$$= (-1)^a y^{t+r-a} \sum_{d=0}^a \binom{a-d+t-1}{a-d} \binom{t}{d} (-1)^d$$

$$\stackrel{d \to a-d}{=} y^{t+r-a} \sum_{d=0}^a \binom{d+t-1}{d} \binom{t}{a-d} (-1)^d$$

$$= y^{t+r-a} \sum_{d=0}^a \frac{(d+t-1)!}{d!(t-1)!} \cdot \frac{t!}{(a-d)!(d+t-a)!} \cdot (-1)^d$$

$$= \sum_{d=0}^a \frac{(d+t-1)!}{d!(a-d)!(d+t-a)!} (-1)^d$$

$$= \frac{ty^{t+r-a}}{a!} \sum_{d=0}^a (-1)^d \binom{a}{d} p(d), \tag{4.4}$$

where  $p(d) = \frac{(d+t-1)!}{(d+t-a)!} = (d+t-1)(d+t-2)\cdots(d+t-a+1)$  is a polynomial in d of degree a-1. As explained in [19, Section 3] and [17, Corollary 2], the sum on the right-hand side of (4.4) is zero, since the polynomial p(d) has a degree smaller than a. Hence, the coefficient of  $n^a$  in A(n, y, r, t) is zero for  $1 \le a \le r - 1$ .

**Case 2:** For  $r \le a \le r + t - 1$ .

Observe that the term c = a - r in the first term of the right-hand side of (4.3) alone contributes to the coefficient of  $n^a$ . The condition  $a \le r + t - 1$  ensures that  $a - r \le t - 1$ . From the second term of the right-hand side of (4.3),

for any i, d = a - i alone contributes to the coefficient of  $n^a$ . Since  $r \le a \le r + t - 1$ , we have  $d = a - i \le t$  for any i. Since  $i \le r - 1$ , we always have  $d \ge 0$ . Hence we get,

Coefficient of 
$$n^a$$
 in  $A(n, y, r, t) = y^{t+r-a} \left( A_1(n, y, r, t) + A_2(n, y, r, t) \right)$ . (4.5)

where,

$$A_1(n, y, r, t) := (-1)^r \sum_{j=a-r}^{t-1} {j+r-1 \choose j} {j \choose a-r},$$
(4.6)

$$A_2(n, y, r, t) := \sum_{i=0}^{r-1} {i + t - 1 \choose i} {t \choose a - i} (-1)^i.$$

$$(4.7)$$

We now evaluate both  $A_1(n, y, r, t)$  and  $A_2(n, y, r, t)$  explicitly. Firstly, after expanding the binomials in  $A_1(n, y, r, t)$ , we get,

$$A_{1}(n,y,r,t) = \frac{(-1)^{r}}{(r-1)!(a-r)!} \sum_{j=a-r}^{t-1} \frac{(j+r-1)!}{(j+r-a)!}$$

$$j \to i + a - r \frac{(-1)^{r}}{(r-1)!(a-r)!} \sum_{i=0}^{r+t-a-1} \frac{(i+a-1)!}{i!}$$

$$= \frac{(-1)^{r}(a-1)!}{(r-1)!(a-r)!} \sum_{i=0}^{r+t-a-1} \binom{i+a-1}{i}$$

$$= \frac{(-1)^{r}(a-1)!}{(r-1)!(a-r)!} \binom{r+t-1}{r+t-a-1}$$

$$= \frac{(-1)^{r}}{a} \times \frac{(r+t-1)!}{(r-1)!(t-1)!} \times \frac{(t-1)!}{(a-r)!(r+t-a-1)!}$$

$$= \frac{(-1)^{r}(r+t-a)}{a} \binom{r+t-1}{t} \binom{t}{a-r}. \tag{4.8}$$

Next, we evaluate  $A_2(n, y, r, t)$ . Since  $\binom{t}{a-i} = 0$  for i < a - t, it is enough to consider the sum between  $a - t \le i \le r - 1$ , and on expanding the binomials, we get,

$$A_{2}(n, y, r, t) = t \sum_{i=a-t}^{r-1} \frac{(-1)^{i}}{i!(a-i)!} \times \frac{(i+t-1)!}{(i+t-a)!}$$

$$i \to j + a - t t \sum_{j=0}^{r+t-a-1} \frac{(-1)^{j+a-t}}{(j+a-t)!(t-j)!} \times \frac{(j+a-1)!}{j!}$$

$$= \frac{(-1)^{a+t}}{(t-1)!} \sum_{j=0}^{r+t-a-1} (-1)^{j} {t \choose j} \frac{(j+a-1)!}{(j+a-t)!}$$

$$= \frac{(-1)^{a+t}}{(t-1)!} \times \frac{(-1)^{a+r+t}(a-r-t)}{at(r-1)!} {t \choose r+t-a} (r+t-1)!$$

$$= -\frac{(-1)^{r}(r+t-a)}{a} {t \choose t} \frac{t}{(a-r)}, \tag{4.9}$$

where we have used the fact  $\binom{c}{d} = \binom{c}{c-d}$  in the last step and the following identity obtained from *Mathematica 11* in the second last step,

$$\sum_{j=0}^{r+t-a-1} (-1)^j \binom{t}{j} \frac{(j+a-1)!}{(j+a-t)!} = \frac{(-1)^{a+r+t}(a-r-t)}{at(r-1)!} \binom{t}{r+t-a} (r+t-1)!.$$

On substituting (4.8) and (4.9) in (4.5), we can see that the coefficient of  $n^a$  is zero for  $r \le a \le r + t - 1$ . Thus, both the cases combined, we have shown that the coefficient of  $n^a$  is zero for all  $1 \le a \le r + t - 1$ , and thus,

A(n, y, r, t) is independent of n, and hence,

$$A(n, y, r, t) = A(0, y, r, t) = y^{r+t}$$

Put this in (4.2) to get the required result.

We now use Lemma 4.1 to prove the following theorem for the series evaluations of  $\Theta(r, s, t, x)$ .

**Theorem 4.2.** For  $r, t \in \mathbb{N} \cup \{0\}$  such that  $r + t \ge 1$ , and  $s \in \mathbb{C}$  such that Re(s) > 2 - r - t, we have,

$$\Theta(r,s,t,x) = \sum_{i=0}^{r-2} \frac{(-1)^i}{x^{t+i}} \binom{i+t-1}{i} \zeta(s+t+i) \zeta(r-i) - \frac{(-1)^r}{x^{r+t-1}} \binom{r+t-2}{t-1} \sum_{m=1}^{\infty} \frac{\gamma + \psi(mx+1)}{m^{r+s+t-1}} + (-1)^r \sum_{i=0}^{t-2} \binom{j+r-1}{j} \frac{(-1)^{t-j}}{(t-j-1)!} \frac{1}{x^{j+r}} \sum_{m=1}^{\infty} \frac{\psi^{(t-j-1)}(mx+1)}{m^{r+s+j}}, \tag{4.10}$$

with the convention that  $\binom{-1}{0} = 1$ ,  $\binom{c}{-1} = 0$  for any  $c \in \mathbb{Z} \setminus \{-1\}$  with an exception  $\binom{-1}{-1} = 1$ .

*Proof.* From the definition (1.3), we can see

$$\Theta(r, s, t, x) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{n^r m^s (n + mx)^t} = \sum_{m=1}^{\infty} \frac{1}{m^s} \sum_{n=1}^{\infty} \frac{1}{n^r (n + mx)^t}.$$
(4.11)

Use Lemma 4.1 with y = mx in (4.11) to get,

$$\begin{split} \Theta(r,s,t,x) &= \sum_{m=1}^{\infty} \frac{1}{m^s} \sum_{n=1}^{\infty} \left( (-1)^r \sum_{j=0}^{t-1} \binom{j+r-1}{j} \frac{1}{(mx)^{j+r}(n+mx)^{t-j}} + \sum_{i=0}^{r-1} \binom{i+t-1}{i} \frac{(-1)^i}{n^{r-i}(mx)^{t+i}} \right) \\ &= \sum_{m=1}^{\infty} \frac{1}{m^s} \sum_{n=1}^{\infty} \left( (-1)^r \sum_{j=0}^{t-2} \binom{j+r-1}{j} \frac{1}{(mx)^{j+r}(n+mx)^{t-j}} + \sum_{i=0}^{r-2} \binom{i+t-1}{i} \frac{(-1)^i}{n^{r-i}(mx)^{t+i}} \right) \\ &+ \sum_{m=1}^{\infty} \frac{1}{m^s} \sum_{n=1}^{\infty} \left( (-1)^r \binom{r+t-2}{t-1} \frac{1}{(mx)^{r+t-1}(n+mx)} + (-1)^{r-1} \binom{r+t-2}{r-1} \frac{1}{n(mx)^{r+t-1}} \right) \\ &= \sum_{m=1}^{\infty} \frac{1}{m^s} \sum_{n=1}^{\infty} \left( (-1)^r \sum_{j=0}^{t-2} \binom{j+r-1}{j} \frac{1}{(mx)^{j+r}(n+mx)^{t-j}} + \sum_{i=0}^{r-2} \binom{i+t-1}{i} \frac{(-1)^i}{n^{r-i}(mx)^{t+i}} \right) \\ &- (-1)^r \binom{r+t-2}{t-1} \frac{1}{x^{r+t-1}} \sum_{j=0}^{\infty} \frac{1}{m^{r+s+t-1}} \sum_{j=0}^{\infty} \left( \frac{1}{n} - \frac{1}{n+mx} \right), \end{split}$$

where we have used  $\binom{r+t-2}{r-1} = \binom{r+t-2}{t-1}$ . From the series definition of the digamma function  $\psi(x)$ , we have,

$$\Theta(r,s,t,x) = \sum_{m=1}^{\infty} \frac{1}{m^s} \sum_{n=1}^{\infty} \left( (-1)^r \sum_{j=0}^{t-2} {j+r-1 \choose j} \frac{1}{(mx)^{j+r} (n+mx)^{t-j}} + \sum_{i=0}^{r-2} {i+t-1 \choose i} \frac{(-1)^i}{n^{r-i} (mx)^{t+i}} \right) - (-1)^r {r+t-2 \choose t-1} \frac{1}{x^{r+t-1}} \sum_{m=1}^{\infty} \frac{\gamma + \psi(mx+1)}{m^{r+s+t-1}}.$$

From (3.14), we have, for  $j \in \mathbb{N}, j \geq 2$ 

$$\frac{(-1)^j}{(j-1)!}\psi^{(j-1)}(z+1) = \sum_{\ell=1}^{\infty} \frac{1}{(\ell+z)^j}.$$
(4.12)

Hence, we can simplify  $\Theta(r, s, t, x)$  as

$$\Theta(r,s,t,x) = (-1)^r \sum_{j=0}^{t-2} {j+r-1 \choose j} \frac{1}{x^{j+r}} \sum_{m=1}^{\infty} \frac{1}{m^{r+s+j}} \sum_{n=1}^{\infty} \frac{1}{(n+mx)^{t-j}} + \sum_{i=0}^{r-2} \frac{(-1)^i}{x^{t+i}} {i+t-1 \choose i} \sum_{m=1}^{\infty} \frac{1}{m^{s+t+i}} \sum_{n=1}^{\infty} \frac{1}{n^{r-i}} - \frac{(-1)^r}{x^{r+t-1}} {r+t-2 \choose t-1} \sum_{m=1}^{\infty} \frac{\gamma + \psi(mx+1)}{m^{r+s+t-1}}.$$

Use (4.12) to get,

$$\begin{split} \Theta(r,s,t,x) &= (-1)^r \sum_{j=0}^{t-2} \binom{j+r-1}{j} \frac{(-1)^{t-j}}{(t-j-1)!} \frac{1}{x^{j+r}} \sum_{m=1}^{\infty} \frac{\psi^{(t-j-1)}(mx+1)}{m^{r+s+j}} \\ &+ \sum_{i=0}^{r-2} \frac{(-1)^i}{x^{t+i}} \binom{i+t-1}{i} \zeta(s+t+i) \zeta(r-i) - \frac{(-1)^r}{x^{r+t-1}} \binom{r+t-2}{t-1} \sum_{m=1}^{\infty} \frac{\gamma + \psi(mx+1)}{m^{r+s+t-1}}. \end{split}$$

This is the required result (4.10).

4.1. **Mixed functional equations.** The two-term functional equation given by Guinand (1.11) has higher derivatives of  $\psi(x)$  in the summand while the one given by Vlasenko-Zagier (1.18) has the higher power of m in the denominator. We call the functional equation obtained in (4.14), the mixed functional equations since they contain both, a series containing higher derivatives of  $\psi(x)$  and containing higher powers of m in the denominator each.

Remark 1. The identity (1.21) holds true.

Proof. Use (1.4) twice to get,

$$\Theta(2,2,t,x) = \Theta(0,2,t+2,x) + 2x\Theta(1,1,t+2,x) + x^2\Theta(2,0,t+2,x). \tag{4.13}$$

Use (1.5) for the third term on the right-hand side of (4.13) and tend  $t \to 0$  to get,

$$\zeta^{2}(2) = \Theta(0, 2, 2, x) + 2x\Theta(1, 1, 2, x) + \Theta(0, 2, 2, \frac{1}{x}).$$

Then, use Theorem 4.2 thrice and simplify to get (1.21).

Such identitites are new in literature. We now give a family of mixed functional equations in the following theorem as a direct corollary of Theorem 4.2.

**Theorem 4.3.** Let us define  $\mathcal{F}(x)$  as follows,

$$\mathscr{F}(x) := (-1)^r \sum_{j=0}^{t-2} \binom{j+r-1}{j} \frac{(-1)^{t-j}}{(t-j-1)!} \frac{1}{x^{j+r}} \sum_{m=1}^{\infty} \frac{\psi^{(t-j-1)}(mx+1)}{m^{2r+j}}$$

$$+ \sum_{i=0}^{r-2} \frac{(-1)^i}{x^{t+i}} \binom{i+t-1}{i} \zeta(r+t+i) \zeta(r-i) - \frac{(-1)^r}{x^{r+t-1}} \binom{r+t-2}{t-1} \sum_{m=1}^{\infty} \frac{\gamma + \psi(mx+1)}{m^{2r+t-1}}.$$

Then, we have,

$$\mathscr{F}(x) = x^{-t}\mathscr{F}\left(\frac{1}{x}\right). \tag{4.14}$$

*Proof.* Put s = r in (4.10) and substitute it in  $\Theta(r, r, t, x) = x^{-t}\Theta(r, r, t, \frac{1}{x})$ .

# 4.2. Kronecker limit type formula.

**Theorem 4.4. Case 1:** Fix an  $r \in \mathbb{N}$ , r > 2.

Then, for any  $t \in \mathbb{N} \cup \{0\}$ , s+t=1 is a singular point of  $\Theta(r,s,t,x)$ , with the following polar singularity structure in s, around s=1-t.

$$\Theta(r,s,t,x) = \frac{x^{-t}\zeta(r)}{(s-(1-t))} + \left(\gamma\zeta(r)x^{-t} + (-1)^r \sum_{j=0}^{t-2} {j+r-1 \choose j} \frac{(-1)^{t-j}}{(t-j-1)!} \frac{1}{x^{j+r}} \sum_{m=1}^{\infty} \frac{\psi^{(t-j-1)}(mx+1)}{m^{r+j-t+1}} + \sum_{i=1}^{r-2} \frac{(-1)^i}{x^{t+i}} {i+t-1 \choose i} \zeta(i+1)\zeta(r-i) - \frac{(-1)^r}{x^{r+t-1}} {r+t-2 \choose t-1} \sum_{m=1}^{\infty} \frac{\gamma+\psi(mx+1)}{m^r} + O_x(|s-(1-t)|). \quad (4.15)$$

Case 2: r = 1

Then for any  $t \in \mathbb{N}$ , s + t = 1 is a singular point of  $\Theta(1, s, t, x)$ , with the following polar singularity structure in s, around s = 1 - t.

$$\Theta(1, s, t, x) = \frac{x^{-t}}{(s - (1 - t))^2} + \frac{x^{-t}}{(s - (1 - t))} (\gamma + \log(x) + H_{t-1}) + \left(\frac{1}{x^t} (\gamma(\gamma + \log(x) + H_{t-1}) + \gamma_1)\right)$$

$$-\sum_{j=0}^{t-2} \frac{(-1)^{t-j} x^{-j-1}}{(t-j-1)!} \sum_{m=1}^{\infty} \frac{\psi^{(t-j-1)}(mx) + (-1)^{t-j} (t-j-2)! (mx)^{-(t-j-1)}}{m^{j-t+2}} + \frac{t\zeta(2)}{x^{t+1}} + \frac{1}{x^t} \sum_{m=1}^{\infty} \frac{\psi(mx) - \log(mx)}{m} + O_x(|s-(1-t)|).$$

$$(4.16)$$

*Proof.* Let us start with the case 1. From (4.10), we have,

$$\Theta(r,s,t,x) - \frac{1}{x^t}\zeta(s+t)\zeta(r) = (-1)^r \sum_{j=0}^{t-2} \binom{j+r-1}{j} \frac{(-1)^{t-j}}{(t-j-1)!} \frac{1}{x^{j+r}} \sum_{m=1}^{\infty} \frac{\psi^{(t-j-1)}(mx+1)}{m^{r+s+j}} + \sum_{i=1}^{r-2} \frac{(-1)^i}{x^{t+i}} \binom{i+t-1}{i} \zeta(s+t+i)\zeta(r-i) - \frac{(-1)^r}{x^{r+t-1}} \binom{r+t-2}{t-1} \sum_{m=1}^{\infty} \frac{\gamma + \psi(mx+1)}{m^{r+s+t-1}}.$$

Since we know, around s = 1 - t,

$$\zeta(s+t) = \frac{1}{s - (1-t)} + \gamma + O(|s - (1-t)|).$$

Hence

$$\lim_{s \to 1-t} \left( \Theta(r, s, t, x) - \frac{1}{x^t} \zeta(s+t) \zeta(r) \right) = (-1)^r \sum_{j=0}^{t-2} \binom{j+r-1}{j} \frac{(-1)^{t-j}}{(t-j-1)!} \frac{1}{x^{j+r}} \sum_{m=1}^{\infty} \frac{\psi^{(t-j-1)}(mx+1)}{m^{r+j-t+1}} + \sum_{i=1}^{r-2} \frac{(-1)^i}{x^{t+i}} \binom{i+t-1}{i} \zeta(i+1) \zeta(r-i) - \frac{(-1)^r}{x^{r+t-1}} \binom{r+t-2}{t-1} \sum_{m=1}^{\infty} \frac{\gamma + \psi(mx+1)}{m^r}.$$

Simplify to get, around s = 1 - t

$$\Theta(r,s,t,x) = \frac{x^{-t}\zeta(r)}{(s-1+t)} + \left(\gamma\zeta(r)x^{-t} + (-1)^r \sum_{j=0}^{t-2} \binom{j+r-1}{j} \frac{(-1)^{t-j}}{(t-j-1)!} \frac{1}{x^{j+r}} \sum_{m=1}^{\infty} \frac{\psi^{(t-j-1)}(mx+1)}{m^{r+j-t+1}} + \sum_{i=1}^{r-2} \frac{(-1)^i}{x^{t+i}} \binom{i+t-1}{i} \zeta(i+1)\zeta(r-i) - \frac{(-1)^r}{x^{r+t-1}} \binom{r+t-2}{t-1} \sum_{m=1}^{\infty} \frac{\gamma+\psi(mx+1)}{m^r} + O(|s-1+t|).$$

For  $t-j-1 \ge 1$ ,  $\psi^{(t-j-1)}(y) = O(y^{-(t-j-1)})$  as  $y \to \infty$ . Hence, we can see that, as  $x \to \infty$ ,

$$\frac{\psi^{(t-j-1)}(mx+1)}{m^{r+j-t+1}} = O_x\left(\frac{1}{m^r}\right).$$

Since  $r \ge 2$ , the series involving the derivatives of  $\psi(x)$  is absolutely convergent. This is the required result (4.15) for the first case.

For the case 2, put r = 1 in Theorem 4.2 to get,

$$\Theta(1, s, t, x) = -\sum_{j=0}^{t-2} \frac{(-1)^{t-j}}{(t-j-1)!} \frac{1}{x^{j+1}} \sum_{m=1}^{\infty} \frac{\psi^{(t-j-1)}(mx+1)}{m^{s+j+1}} + \frac{1}{x^t} \sum_{m=1}^{\infty} \frac{\gamma + \psi(mx+1)}{m^{s+t}}.$$
 (4.17)

We simplify the terms on the right-hand side of (4.17) as follows:

$$\sum_{m=1}^{\infty} \frac{\gamma + \psi(mx+1)}{m^{s+t}} = \gamma \zeta(s+t) + \sum_{m=1}^{\infty} \frac{\psi(mx+1)}{m^{s+t}}$$

$$= \gamma \zeta(s+t) + \frac{1}{x} \zeta(s+t+1) + \sum_{m=1}^{\infty} \frac{\psi(mx)}{m^{s+t}}$$

$$= \gamma \zeta(s+t) + \frac{1}{x} \zeta(s+t+1) + \sum_{m=1}^{\infty} \frac{\psi(mx) - \log(mx)}{m^{s+t}} + \sum_{m=1}^{\infty} \frac{\log(mx)}{m^{s+t}}$$

$$= (\gamma + \log(x))\zeta(s+t) + \frac{1}{x} \zeta(s+t+1) - \zeta'(s+t) + \sum_{m=1}^{\infty} \frac{\psi(mx) - \log(mx)}{m^{s+t}}. \tag{4.18}$$

Simplification of the other term is as follows: Since  $t-j-1 \ge 1$ , we have  $\psi^{(t-j-1)}(y+1) = \psi^{(t-j-1)}(y) - (-1)^{t-j}(t-j-1)!y^{j-t}$ . Using it we can see,

$$\sum_{m=1}^{\infty} \frac{\psi^{(t-j-1)}(mx+1)}{m^{s+j+1}} = -\frac{(-1)^{t-j}(t-j-1)!}{x^{t-j}} \zeta(s+t+1) + \sum_{m=1}^{\infty} \frac{\psi^{(t-j-1)}(mx)}{m^{s+j+1}}$$

$$= -\frac{(-1)^{t-j}(t-j-1)!}{x^{t-j}} \zeta(s+t+1) - \frac{(-1)^{t-j}(t-j-2)!}{x^{t-j-1}} \zeta(s+t)$$

$$+ \sum_{m=1}^{\infty} \frac{\psi^{(t-j-1)}(mx) + (-1)^{t-j}(t-j-2)!(mx)^{-(t-j-1)}}{m^{s+j+1}}.$$
(4.19)

Substitute (4.18) and (4.19) in (4.17) to get

$$\begin{split} \Theta(1,s,t,x) &- \frac{1}{x^t} \Biggl( \gamma + \log(x) + \sum_{j=0}^{t-2} \frac{1}{(t-j-1)} \Biggr) \zeta(s+t) + \frac{1}{x^t} \zeta'(s+t) \\ &= \frac{t}{x^{t+1}} \zeta(s+t+1) + \frac{1}{x^t} \sum_{m=1}^{\infty} \frac{\psi(mx) - \log(mx)}{m^{s+t}} \\ &- \sum_{j=0}^{t-2} \frac{(-1)^{t-j}}{(t-j-1)!} \frac{1}{x^{j+1}} \sum_{m=1}^{\infty} \frac{\psi^{(t-j-1)}(mx) + (-1)^{t-j}(t-j-2)!(mx)^{-(t-j-1)}}{m^{s+j+1}}. \end{split}$$

Taking limit as  $s \to 1 - t$ , we get.

$$\lim_{s \to 1-t} \left( \Theta(1, s, t, x) - \frac{1}{x^t} \left( \gamma + \log(x) + H_{t-1} \right) \zeta(s+t) + \frac{1}{x^t} \zeta'(s+t) \right)$$

$$= \frac{t}{x^{t+1}} \zeta(2) + \frac{1}{x^t} \sum_{m=1}^{\infty} \frac{\psi(mx) - \log(mx)}{m}$$

$$- \sum_{j=0}^{t-2} \frac{(-1)^{t-j}}{(t-j-1)!} \frac{1}{x^{j+1}} \sum_{m=1}^{\infty} \frac{\psi^{(t-j-1)}(mx) + (-1)^{t-j} (t-j-2)! (mx)^{-(t-j-1)}}{m^{j-t+2}}.$$

where  $H_n$  is the  $n^{th}$  Harmonic number, with the convention  $H_0 = 0$ . Also, see that, as  $y \to \infty$ ,

$$\psi^{(t-j-1)}(y) + (-1)^{t-j}(t-j-2)! \frac{1}{y^{t-j-1}} = O(y^{-(t-j)}),$$

which ensures the convergence of the series involving the derivatives of  $\psi(x)$ . Use the Laurent expansions of  $\zeta(s+t)$  and  $\zeta'(s+t)$  around s=1-t to get the required result (4.16).

## Corollary 4.5.

$$\Theta(1, s, 1, x) = \frac{1}{xs^2} + \frac{\gamma + \log(x)}{xs} + \frac{1}{x}(\gamma(\gamma + \log(x)) + \gamma_1) + \frac{\zeta(2)}{x^2} + \frac{1}{x} \sum_{n=1}^{\infty} \frac{\psi(mx) - \log(mx)}{m}.$$

*Proof.* Put r = t = 1 in Theorem 4.4 and tend  $s \to 0$  to get,

$$\lim_{s \to 0} \left( \Theta(1, s, 1, x) - \frac{1}{x} \left( \gamma + \log(x) \right) \zeta(s+1) + \frac{1}{x} \zeta'(s+1) \right) = \frac{\zeta(2)}{x^2} + \frac{1}{x} \sum_{m=1}^{\infty} \frac{\psi(mx) - \log(mx)}{m}.$$

Use the Laurent series expansions of  $\zeta(s+1)$  and  $\zeta'(s+1)$  around s=0 to get the result.

Remark 2. From [25, Equation 7.12],

$$\sum_{m=1}^{\infty} \frac{\psi(m) - \log(m)}{m} = -\frac{\gamma^2}{2} - \frac{\pi^2}{12} - \gamma_1,$$

and putting x = 1 in Corollary 4.5, we can see that around s = 0,

$$\Theta(1, s, 1, 1) = \frac{1}{s^2} + \frac{\gamma}{s} + \frac{6\gamma^2 + \pi^2}{12} + O(|s|). \tag{4.20}$$

#### 5. Modular relations: The Mordell-Tornheim zeta perspective

We list few of the corollaries in the Table 3 below. We start with the necessary arguments for  $\Theta$  in the first column. After using (1.4) the number of times as in column 3, we do the operation on t as in column 2, to obtain the result mentioned in column 4.

For example,  $\Theta(1,1,t,x)$  can give mixed functional equations, an instance of Vlasenko-Zagier's functional equation (1.18), based on the steps mentioned in columns 2 and 3. Observe that  $\Theta(0,0,t,x)$  is the centroid between Ramanujan's functional equation (1.10) and Guinand's functional equations (1.11) and (1.12), differing by the limiting value of t alone, bringing the three results under a single umbrella. Look at [5, Corollary 3.5] to see how Zagier's result (1.14) can be obtained.

Θ	Operation on $t$	No. of splits	Result obtained
$\Theta(0,0,t,x)$	$t \in \mathbb{N}, t \geq 3$	0	Guinand (1.11)
	$t \to 2$	0	Guinand (1.12)
	$t \to 1$	0	Ramanujan (1.10)
$\Theta(1,1,t,x)$	$t \in \mathbb{N}, t \ge 2$	0	Mixed functional equations, Theorem 4.3
	$t \to 1$	0	Vlasenko-Zagier (1.18) with $r = 2$
	$t \to 0$	1	Zagier (1.14)
	$t \in \mathbb{N}, t \ge 2$	0	Mixed functional equations, Theorem 4.3
	$t \to 1$	0	Vlasenko-Zagier (1.18) with $r = 4$
$\Theta(2,2,t,x)$	$t \to 0$	1	Vlasenko-Zagier (1.18) with $r = 3$
		2	Mixed functional equations, Remark 1
	$t \rightarrow -1$	2	Vlasenko-Zagier (1.18) with $r = 2$

FIGURE 3. The table shows the connections between the Mordell-Tornheim zeta function and the various modular relations in literature.

#### 6. Concluding Remarks

We conclude the paper with the following remarks:

- (1) In this paper, we have obtained the Kronecker limit formula for  $\Theta(r, s, t, x)$  in the variable t in Theorem 2.3 and Theorem 2.4 for  $\ell \geq 1-r$  and Theorem 2.5 for  $r+s \in \mathbb{N}, r+s \geq 2$ . We can extend Theorem 2.3 and Theorem 2.4 for all  $\ell \in \mathbb{N} \cup \{0\}$  and Theorem 2.5 for all  $r, s \in \mathbb{C}$ . However, the closed form expression for the constant term is a challenging task, which is in terms of Arakawa-Kaneko constants and the integral mentioned in (1.7) and (1.8) respectively.
- (2) We have the description of polar nature of  $\Theta(r, s, t, x)$  as a function of the *third variable* t as well as the *second variable* s (hence the *first variable* r) as a function of one variable. However, it will be very interesting to see the nature of series expansion as a function of three variables r, s and t simultaneously, in the sense of several complex variables.
- (3) As discussed in the Section 5, the Mordell-Tornheim zeta function gives a new perspective of the two-term functional equations in the literature. This interplay between  $\Theta(r,s,t,x)$  and the modular relations thus widens the scope for the study on  $\Theta(r,s,t,x)$  and its generalizations. For example, having higher order terms of  $\Theta(r,s,t,x)$  in Theorem 2.3, will give us two-term explicit functional equations Ishibashi and Higher Ishibashi functions, showed in [5, Theorem 3.4].
- (4) We have obtained two term functional equations as an corollary of Theorem (3.1). Three-term analogue of (3.1) will be an extremely desirable result. That is because, such an identity will enable us to derive three-term functional equations such as (1.15) and (1.19).
- (5) The result (4.20) could have interesting applications.

## 7. Acknowledgment

The authors would like to extend gratitude to Prof. Atul Dixit for his insights and continuous support. The first author was partially supported by the grant MIS/IITGN/R&D/MATH/AD/2324/058 of IIT Gandhinagar.

#### References

- [1] M. Abramowitz and I. A. Stegun, Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, 9th printing, New York: Dover, p. 260, 1972.
- [2] T. Arakawa and M. Kaneko, Multiple zeta values, poly-Bernoulli numbers, and related zeta functions, Nagoya Math. J. 153 (1999), 189–209.
- [3] B. C. Berndt and A. Dixit, A transformation formula involving the Gamma and Riemann zeta functions in Ramanujan's Lost Notebook, The Legacy of Alladi Ramakrishnan in the Mathematical Sciences, K. Alladi, J. Klauder, C. R. Rao, Eds, Springer, New York, 2010, pp. 199–210.
- [4] A. Dixit, Analogues of a transformation formula of Ramanujan, Int. J. Number Theory 7(5) (2011), 1151–1172.
- [5] A. Dixit, S. Sathyanarayana, and N. G. Sharan, Mordell-Tornheim zeta functions and functional equations for Herglotz-Zagier type functions, arXiv preprint (arXiv:2405.07934v1), 2024.
- [6] A. Erdelyi, W. Magnus, F. Oberhettinger and F. G. Tricomi, *Higher transcendental functions* Vol. I, Based, in part, on notes left by Harry Bateman. McGraw-Hill Book Co., Inc., New York-Toronto-London, 1953, 302 pp.
- [7] H. Gangl, M. Kaneko and D. Zagier, Double zeta values and modular forms, automorphic forms and zeta functions, Proceedings of the conference in memory of Tsuneo Arakawa, World Scientific, Hackensack (2006), 71–106.
- [8] A. P. Guinand, A note on the logarithmic derivative of the Gamma function, Edinburgh Math. Notes 38 (1952), 1-4.
- [9] L. Guo and B. Xie, The shuffle relation of fractions from multiple zeta values, Ramanujan J., 25 (2011), 307-317.
- [10] E. Hecke, Über die Kroneckersche Grenzformell für reelle quadratische Körper und die Klassenzhal relative-Abelscher Körper, Verhandl. d. Naturforschenden Gesell. i. Basel 28 (1917), 363—372.
- [11] G. Herglotz, Über die Kroneckersche Grenzformel für reele, quadratische Körper I, Ber. Ver handl. Sachsischen Akad. Wiss. Leipzig 75 (1923), 3–14.
- [12] J. G. Huard, K. S. Williams and N. Y. Zhang, On Tornheim's double series, Acta Arith. 75 (1996), 105-117.
- [13] L. Kronecker, Werke IV, Chelsea Publishing Co. (1929) p. 221–225.
- [14] K. Matsumoto, On analytic continuation of various multiple zeta-functions, Surveys in Number Theory: Papers from the Millennial Conference on Number Theory. AK Peters: CRC Press, 2002.
- [15] K. Matsumoto, K. Onodera and D. K. Sahoo, Mordell-Tornheim multiple zeta-functions, their integral analogues, and relations among multiple polylogarithms, arXiv preprint (arXiv:2409.19980v2), 2024.
- [16] K. Ramachandra, Some applications of Kronecker's limit formulas, Ann. of Math., 80 (1964), no. 1, 104–148.
- [17] S. M. Ruiz, An algebraic identity leading to Wilson's theorem, Math. Gaz., 80 (1996), no. 489, 579—582.
- [18] C. L. Siegel, Bernoullische Polynome und quadratische Zahlkörper, Collected Papers IV, p. 9-40, 1968.
- [19] M. Z. Spivey, Combinatorial sums and finite differences, Discrete Math., 307 (2007), no. 24, 3130-3146.
- [20] R. P. Stanley, Enumerative Combinatorics, Vol. 1, 2nd ed., Cambridge Stud. Adv. Math. 49, Cambridge University Press, 2012.
- [21] H. M. Stark, Values of L-functions at s = 1. I. L-functions for quadratic forms, Adv. Math. 7(3) (1971), 301—343.
- [22] N. M. Temme, Special functions: An introduction to the classical functions of mathematical physics, Wiley-Interscience Publication, New York, 1996.
- [23] E. C. Titchmarsh, The Theory of the Riemann Zeta-function, Clarendon Press, Oxford, 1986.
- [24] M. Vlasenko and D. Zagier, Higher Kronecker "limit" formulas for real quadratic fields, J. Reine Angew. Math. 679 (2013), 23-64.
- [25] D. Zagier, A Kronecker limit formula for real quadratic fields, Math. Ann. 213 (1975), 153-184.

Department of Mathematics, Central University of Karnataka, Kadaganchi, Kalaburagi, Karnataka-585367, India. *Email address*: sumukhas@cuk.ac.in, neerugarsumukha@gmail.com

Department of Mathematics, Indian Institute of Technology Gandhinagar, Palaj, Gandhinagar, Gujarat-382355, India.

Email address: gurusharan.n@iitgn.ac.in, sharanguru5@gmail.com