

HOLONOMICITY FROM A HEEGAARD-FLOER PERSPECTIVE

BENJAMIN COOPER AND ROBERT DEYESO

ABSTRACT. We construct S^r -colored knot Floer homologies and prove that they satisfy categorified recurrence relations. The associated Euler characteristic implies q -holonomicity of the corresponding sequence of colored Alexander polynomials, in analogy with the AJ conjecture for colored Jones polynomials.

CONTENTS

1. Introduction	1
2. Holonomicity of colored Alexander polynomials	4
3. Overview of Heegaard-Floer theory	9
4. Colored knot Floer homology	18
5. Weyl algebra and homological holonomicity	28
6. Holonomicity for $S^r \widehat{CFK}(K)$	35
7. Notation	38
References	39

1. INTRODUCTION

Given a knot K in S^3 and a simple Lie algebra \mathfrak{g} , the Reshetikhin-Turaev construction produces a q -series valued invariant $\langle K \rangle_V$ of K for each finite dimensional representation V of the corresponding quantum group $U_q(\mathfrak{g})$. A priori, the collection $\{\langle K \rangle_V\}_{V \in \text{Rep}(\mathfrak{g})}$ appears to contain an infinite amount of information, but the existence of recurrence relations among, or the q -holonomicity of, these knot invariants tells us that there is a great deal of redundancy. In exchange for an infinite sequence

$$f : \mathbb{Z}_{\geq 0} \rightarrow \mathbb{Z}[q^{-1}][[q]]$$

Date: 6th January, 2025.

we can make do with a finite collection of initial conditions and a recurrence relation of the form

$$\sum_{i=n}^{n+m} a_i(q^n, q) f(i) = 0 \quad \text{for} \quad a_i(u, v) \in R[u, v], \quad (1.1)$$

where $R := \mathbb{Z}[q, q^{-1}]$. Garoufalidis and Le proved that for each knot K such a recurrence relation always exists among \mathfrak{g} invariants (for $\mathfrak{g} \neq G_2$) [GL05, Thm. 6]. Unfortunately, this approach does not readily lead to an interpretation of the recurrence relation. In the simplest non-trivial case, the irreducible representations $V_n \cong S^n(V_1)$ of \mathfrak{sl}_2 ($\dim V_n = n + 1$) determine a sequence

$$f(n) = \langle K \rangle_{V_n}$$

called the colored Jones polynomials of K . The AJ conjecture, posed by Garoufalidis [Gar08, Gar04], states that the recurrence relation satisfied by this sequence is a non-commutative deformation of the A -polynomial. The commutative A -polynomial is characterized by vanishing on the space of characters which extend from the boundary torus $S^1 \times S^1 \cong \partial(S^3 \setminus \nu K)$ into the 3-manifold $S^3 \setminus \nu K$, where νK is a regular neighborhood of the knot K , see [CCG⁺94]. In foundational work, Frohman, Gelca and Lofaro [FGL02, FG00] studied similar statements. Gukov studied this problem using insights from physics [Guk05]. The conjecture is known to be true for some families of knots, see [L06].

A different story which also stems from the Reshetikhin-Turaev invariants involves Khovanov homology and its generalizations, knot homology theories, which categorify the Reshetikhin-Turaev invariants, see [KR08, Web17]. For a knot K in S^3 , and a representation V of a simple Lie algebra \mathfrak{g} , there are chain complexes of graded vector spaces $[[K]]_V$, the homology of which are invariants of K . The generating function associated to the homology is the Poincaré polynomial

$$P(K)_V(t, q) = \sum_{i, j \in \mathbb{Z}} t^i q^j \dim H^{i, j}([[K]]_V), \quad \text{satisfying} \quad P(K)_V|_{t=-1} = \langle K \rangle_V. \quad (1.2)$$

Its value at $t = -1$ recovers the Reshetikhin-Turaev invariants as the graded Euler characteristic of knot homology. The situation here is sometimes likened to the relationship between ordinary homology and Euler characteristic, while both are useful tools, the historical precedence of the latter has been overshadowed by the great worlds which have been revealed to us by the former.

If one considers super Lie algebras $\mathfrak{g} = \mathfrak{gl}(1|1)$ then the Alexander polynomial $\Delta_K(q)$ becomes an example of the Reshetikhin-Turaev construction outlined above [Vir06, Sar15]. Heegaard-Floer homology was used to define knot Floer homology $\widehat{HFK}(K)$ by Ozsváth, Szabó and Rasmussen [OS04a, Ras03a]. In analogy with Eqn.

(1.2) above, the Euler characteristic of this knot homology theory is the Alexander polynomial.

The purpose of this paper is to examine the question of q -holonomicity for these knot homology theories. The HOMFLY homology unites the \mathfrak{sl}_n knot homology theories [Ras15, Cau17] and, conjecturally, this includes the knot Floer homology as well [DGR06, BPLHW22]. Fuji, Gukov and Sułkowski [FGSk13] found examples of recurrence relations among the Poincaré polynomials $P(K)_{S^n}(q, t)$ of the HOMFLY homology theories. The colored HOMFLY polynomials $P(K)_{S^n}(q, -1)$ are known to be q -holonomic [GLL18]. When placed within the framework of [DGR06], this work suggests the existence of some recurrence relations or *homological q -holonomicity* among S^n -colored HOMFLY knot homologies; a lifting of those relations. It is therefore natural to formulate the conjecture below.

Conjecture 1.1. *There exists a notion of homological q -holonomicity for sequences of chain complexes of (filtered) graded vector spaces. For each knot K in S^3 , the sequence of S^n -colored HOMFLY homologies $\{\llbracket K \rrbracket_{S^n}\}_{n \geq 1}$ is homologically q -holonomic.*

Here we pursue a simplified version of this conjecture. Since a spectral sequence from HOMFLY homology to knot Floer homology ought to imply that homological q -holonomicity holds for knot Floer homology as well. We restrict our attention to this setting, where a robust theory has been developed and can be leveraged in the context of our problem.

Theorem 1.2. *There exists a notion of homological q -holonomicity for sequences of (filtered) chain complexes of graded vector spaces. For each knot K in S^3 , the S^n -colored knot Floer homologies $\{S^r \widehat{CFK}(K)\}_{n \geq 1}$ is homologically q -holonomic.*

Our paper is structured as follows. In §2 we show that the S^n -colored Alexander polynomials are q -holonomic in the sense of Eqn. (1.1). In §4, using immersed curve technology [HRW22], we construct the S^n -colored knot Floer homologies $S^r \widehat{HFK}(K)$ using Rozansky’s infinite braid argument [Roz14] (which the spectral sequence from the S^n -colored HOMFLY homology [Hog18, Thm. 2.3] would naturally abut). The Euler characteristics of $S^r \widehat{HFK}(K)$ correspond to limits of Alexander polynomials $\Delta_K^r(q)$ from §2. We introduce a notion of homological q -holonomicity in §5 which would apply to the conjecture above and discuss a few simple examples. In §6 we prove that the S^n -colored knot Floer homologies are homologically q -holonomic.

Acknowledgements. The first author would like to thank M. Stošić and O. Yacobi for conversations. J. Hanselman for addressing some questions about his work.

2. HOLONOMICITY OF COLORED ALEXANDER POLYNOMIALS

In §2.1 we construct the S^n -colored Alexander polynomials and in §2.2 we show that this sequence is q -holonomic.

2.1. S^n -colored Alexander polynomials. For a knot K in S^3 , the Alexander polynomial $\Delta_K(q) \in \mathbb{Z}[q, q^{-1}]$ [Lic97]. A priori, $\Delta_K(q)$ is only defined up to multiplication by $q^{\pm 1}$, but it can be expressed uniquely when the equation

$$\Delta_K(q) = \Delta_K(q^{-1})$$

is required to hold. In this paper, $\Delta_K(q)$ will be used to denote this *symmetrized form* of the Alexander polynomial.

A different normalization, $\Delta_K^1(q)$ which will be more important for us in what follows, is determined by requiring the lowest power of q to be q^0 . More precisely, if $\deg(\Delta_K)$ is the largest monomial power of q occurring in the symmetrized form of $\Delta_K(q)$ then the polynomial

$$\Delta_K^1(q) := q^{\deg(\Delta_K)} \Delta_K(q) \in \mathbb{Z}[q] \quad (2.1)$$

is the *positive form*. The S^r -colored Alexander polynomials are given by certain limits of positive forms of Alexander polynomials described in this section.

Definition 2.1. Given a knot K in S^3 and $\nu K \subset S^3$ a regular neighborhood. The (r, s) -cable $K_{r,s}$ of K is the (r, s) -torus link in the boundary torus $\partial \nu K$, winding r -times around the meridian and s -times around the longitude that is determined by the Seifert framing.

The Alexander polynomial $\Delta_{K_{r,s}}(q)$ of the (r, s) -cable $K_{r,s}$ can be expressed in terms of the Alexander polynomial $\Delta_K(q)$ of the knot K and the Alexander polynomial of the torus knot $\Delta_{T_{r,s}}(q)$ respectively [Lic97].

$$\begin{aligned} \Delta_{K_{r,s}}(q) &= \Delta_K(q^r) \Delta_{T_{r,s}}(q) \\ &= \Delta_K(q^r) \cdot \frac{q^{rs} - 1}{q^r - 1} \cdot \frac{q - 1}{q^s - 1} \end{aligned} \quad (2.2)$$

By writing the rational functions as geometric series as in Eqn. (2.4), $\Delta_{K_{r,s}}^1(q) = q^{\deg(\Delta_{K_{r,s}})} \Delta_{K_{r,s}}(q)$ becomes an element of the power series ring

$$\mathbb{Z}[[q]] := \varprojlim \mathbb{Z}[q]/(q^n). \quad (2.3)$$

The proposition below shows that the sequence of Alexander polynomials obtained by increasing the number of longitudinal full twists converges in the topology of the ring $\mathbb{Z}[[q]]$.

Proposition 2.2. *There is a limit*

$$\lim_{n \rightarrow \infty} \Delta_{K_r, rn+1}^1(q) = \Delta_K^1(q^r) \frac{q-1}{q^r-1}$$

where the rational function represents the power series

$$\frac{q-1}{q^r-1} := (1-q) \sum_{i=0}^{\infty} q^{ir} \quad (2.4)$$

Proof. For a sequence $\{f_n\}_{n \geq 0}$ of power series to converge to f in the q -adic topology, or $f_n \rightarrow f$ as $n \rightarrow \infty$, it suffices to show that for each $n \in \mathbb{Z}_{\geq 0}$, there exists $N \in \mathbb{Z}_{\geq 0}$ such that

$$f \equiv f_i \pmod{q^n} \quad \text{for all } i > N.$$

When $s > r$, the Alexander polynomial of the (r, s) -torus $\Delta_{T_{r,s}}(q)$ satisfies

$$\frac{q-1}{q^r-1} \frac{q^{rs}-1}{q^s-1} = \frac{q-1}{q^r-1} \cdot 1 + \frac{q-1}{q^r-1} \cdot q^s + \cdots + \frac{q-1}{q^r-1} \cdot q^{(r-1)s} \quad (2.5)$$

Using Eqn. (2.2) and (2.5), it follows that

$$\begin{aligned} \lim_{n \rightarrow \infty} q^{\deg(\Delta_{K_r, rn+1})} \Delta_{K_r, rn+1}(q) &= q^{r \deg(\Delta_K)} \Delta_K(q^r) \lim_{n \rightarrow \infty} \left(\frac{q-1}{q^r-1} + q^{rn+1} (*) \right) \\ &= q^{r \deg(\Delta_K)} \Delta_K(q^r) \left(\frac{q-1}{q^r-1} + 0 \right) \end{aligned}$$

where $(*)$ corresponds to the higher order terms on the right-hand side of in Eqn. (2.5). \square

Definition 2.3. The S^r -colored Alexander polynomial of a knot K is the limit of Alexander polynomials of the $(r, rn+1)$ -cables of K as $n \rightarrow \infty$.

$$\bar{\Delta}_K^r(q) := q^{r \deg(\Delta_K)} \Delta_K(q^r) \frac{q-1}{q^r-1}$$

When K is the unknot U , the Alexander polynomial $\Delta_U(q) = 1$. So, the S^r -colored Alexander polynomial of the unknot is the series below.

$$\bar{\Delta}_U^r(q) = \frac{q-1}{q^r-1} = \sum_{i=0}^{\infty} (q^{ir} - q^{ir+1}) \quad (2.6)$$

The reduced S^r -colored Alexander polynomial of a knot K is the quotient

$$\Delta_K^r(q) := \bar{\Delta}_K^r(q) / \bar{\Delta}_U^r(q). \quad (2.7)$$

Notice that $\Delta_K^1(q)$ agrees with $\bar{\Delta}_K^1(q)$ and Eqn. (2.1) above.

2.2. Recurrence relation for S^r -colored Alexander polynomials. In the remainder of this section we will show directly that the sequence of r -colored Alexander polynomials satisfy a recurrence relation of the form Eqn. (1.1) from the introduction. The relative simplicity of the expressions $\Delta_K^r(q)$ and $\bar{\Delta}_K^r(q)$ from Def. 2.3 allow us to be explicit. This construction will be refined in §6.

Definition 2.4. (Seq) Let $R := \mathbb{Z}[q, q^{-1}]$. If T is an R -algebra then there is a natural T -module $\text{Seq}(T) := T^{\mathbb{Z}_{\geq 0}}$ which consists of sequences of elements of T which are parameterized by $r \in \mathbb{Z}_{\geq 0}$. Let $\text{Seq} := \text{Seq}(R[[q]])$ be sequences of power series which are bounded in negative powers of q .

For each knot K , using Def. 2.3, there are sequences

$$\Delta_K, \bar{\Delta}_K : \mathbb{Z}_{\geq 0} \rightarrow R[[q]] \in \text{Seq} \quad \text{given by} \quad (2.8)$$

$$\Delta_K(r) := \Delta_K^r(q) \quad \text{and} \quad \bar{\Delta}_K(r) := \bar{\Delta}_K^r(q) \quad (2.9)$$

of reduced and unreduced S^r -colored Alexander polynomials respectively. We will find recurrence relations of the form

$$\sum_{i=n}^{n+m} a_i(q^n, q) \Delta_K(i) = 0 \quad \text{for} \quad a_i(u, v) \in R[u, v] \quad (2.10)$$

This will be done in two steps. First in §2.2 we find such a recurrence relation A_K for the sequence of reduced polynomials Δ_K . Then in §2.2 this relation is combined with the recurrence relation Eq. (2.6) for the unknot $\Delta_U(q)$ to produce a recurrence relation for the unreduced sequence $\bar{\Delta}_K$.

Before we start, a bit of material about the relationship between recurrence relations and the Weyl algebra is reviewed.

The Weyl algebra. It is sometimes useful to think of recurrence relations in terms of operators. There are two operators L, M acting on sequences $f \in \text{Seq}$ by

$$(Lf)(n) := f(n+1) \quad \text{and} \quad (Mf)(n) := q^n f(n) \quad (2.11)$$

These operators satisfy the relation $qLM = ML$. The *Weyl algebra* is defined by this data

$$\mathcal{A} := \mathbb{Z}[q, q^{-1}] \langle L, M \rangle / \langle qLM - ML \rangle$$

and by virtue of the assignments in Eqn. (2.11), sequences Seq are an \mathcal{A} -module. For a given sequence $f \in \text{Seq}$, the recurrence relation in Eqn. (2.10) is equivalent to the relation

$$Af = 0$$

when $A \in \mathcal{A}$ is the element $A = \sum_{i=n}^{n+m} L^i a_i(M, q)$, $a_i \in \mathbb{Z}[q, q^{-1}] \langle M \rangle$ for $i = n, \dots, n+m$. The set of recurrence relations satisfied by f can be identified with

$$\text{Ann}(f) = \{A \in \mathcal{A} : Af = 0\} \subseteq \mathcal{A}.$$

D-operations. Let $X \in \mathbb{Z}[q, q^{-1}]$ be a q -Laurent polynomial. Then there is an operator $D_X = 1 - XL \in \mathcal{A}$ which acts on sequences. In long form, if $f \in \text{Seq}$ is a sequence then

$$(D_X f)(n) := f(n) - Xf(n+1)$$

is a new sequence which is defined in terms of f and multiplication by X .

Proposition 2.5. *Suppose that $f, g \in \text{Seq}$ are sequences, $X, Y \in \mathbb{Z}[q, q^{-1}]$ are Laurent polynomials and $C \in \mathbb{Z}$ is an integer. Then the operators D_X satisfy the properties below.*

- (1) $\mathbb{Z}[q, q^{-1}]$ -linearity: $D_X(Yf) = YD_X(f)$ and $D_X(f+g) = D_X(f) + D_X(g)$
- (2) Derivation: $D_X(fg) = D_X(f)g + (X(Lf))D_1(g)$
- (3) Commutation: $D_X D_Y = D_Y D_X$
- (4) Annihilation: $D_X(q^{Cn}) = q^{Cn}(1 - Xq^C)$. In particular, if $X = q^{-C}$ then $D_X(q^{Cn}) = 0$.

Proof. All of the properties are verified directly from the definition of D_X .

- (1) For $\mathbb{Z}[q, q^{-1}]$ -linearity,

$$\begin{aligned} D_X(Yf)(n) &= Yf(n) - YXf(n+1) \\ &= Y(f(n) - Xf(n+1)) \\ &= YD_X(f)(n) \end{aligned}$$

$$\begin{aligned} D_X(f+g)(n) &= (f+g)(n) - X(f+g)(n+1) \\ &= f(n) + g(n) - X(f(n+1) + g(n+1)) \\ &= (f(n) - Xf(n+1)) + (g(n) - Xg(n+1)) \\ &= D_X(f)(n) + D_X(g)(n) \end{aligned}$$

- (2) Derivation,

$$\begin{aligned} D_X(fg)(n) &= f(n)g(n) - Xf(n+1)g(n+1) \\ &= (f(n) - Xf(n+1))g(n) + Xf(n+1)(g(n) - g(n+1)) \end{aligned}$$

- (3) For commutation,

$$\begin{aligned} (D_X D_Y f)(n) &= (D_Y f)(n) - X(D_Y f)(n+1) \\ &= (f(n) - Yf(n+1)) - X(f(n+1) - Yf(n+2)) \\ &= f(n) - (X+Y)f(n+1) + XYf(n+2) \end{aligned}$$

which is symmetric in X and Y , so $(D_X D_Y f)(n) = (D_Y D_X f)(n)$.

- (4) Annihilation: $D_X(q^{Cn})(n) = q^{Cn} - Xq^{C(n+1)} = q^{Cn}(1 - Xq^C)$

□

The lemma below is a consequence of the properties from the Prop. 2.5 above.

Lemma 2.6. *Given a knot K in S^3 , if the Alexander polynomial of K is written as*

$$\Delta_K^1(q) = \sum_{i=1}^N a_i q^{n_i} \quad \text{where} \quad a_i \in \mathbb{Z} \setminus \{0\}, n_i \text{ distinct}$$

then the operator $A_K := \prod_i D_{q^{-n_i}}$ annihilates the sequence of reduced Alexander polynomials $\Delta_K(r) := \Delta_K^r(q) = \Delta_K^1(q^r)$

Proof. By induction on the number of non-zero terms N . If $N = 1$ then $\Delta_K(q) = a_1 q^{n_1}$ and $D_{q^{-n_1}}$ annihilates the associated sequence, by Prop. 2.5 (4). Suppose $\Delta_K(q) = f(q) + a q^n$ where $f(q) := \sum_{i=1}^{N-1} a_i q^{n_i}$, with $a_i \neq 0$ for $i = 1, \dots, N-1$. By induction, the operator $D' := D_{q^{-n_1}} D_{q^{-n_2}} \cdots D_{q^{-n_{N-1}}}$ annihilates the sequence $r \mapsto f(q^r)$. So

$$\begin{aligned} (A_K \Delta_K)(r) &= ((D' D_{q^{-n}}) \Delta_K)(r) \\ &= D'(D_{q^{-n}}(f(q^r) + a q^{r n})) \\ &= D' D_{q^{-n}} f(q^r) + D'(a D_{q^{-n}}(q^{r n})) \end{aligned} \quad (2.5 \text{ (1)})$$

$$= D' D_{q^{-n}} f(q^r) + D'(a \cdot 0) \quad (2.5 \text{ (4)})$$

$$= D' D_{q^{-n}} f(q^r)$$

$$= D_{q^{-n}} D' f(q^r) \quad (2.5 \text{ (3)})$$

$$= D_{q^{-n}}(0)$$

$$= 0$$

□

Example 2.7. The operator $D_1 D_{q^{-1}} D_{q^{-2}}$ annihilates the sequence $\Delta_{3_1}(q^r) = 1 - q^r + q^{2r}$ of reduced S^r -colored Alexander polynomials from Eqn. (2.7) associated to the right-handed trefoil $K = 3_1$.

Incorporating the unknot U . Recall from Eqn. (2.6) that the r -colored Alexander polynomial of the unknot U can be identified with the power series expansion of the rational function

$$\bar{\Delta}_U^r(q) = \frac{q-1}{q^r-1} = (1-q) \sum_{i=0}^{\infty} q^{ri}$$

Lemma 2.8. *If $\bar{\Delta}_U(r) := \bar{\Delta}_U^r(q)$ defines the sequence $\bar{\Delta}_U \in \text{Seq}$ then $(M-1)(L-1)\bar{\Delta}_U = 0$*

Proof. The unknot sequence is $\bar{\Delta}_U(r) = (q-1)/(q^r-1)$. So

$$(q^r-1)\bar{\Delta}_U(r) - (q^{r-1}-1)\bar{\Delta}_U(r-1) = 0$$

which can be rewritten as $(M-1)(L-1)\bar{\Delta}_U = 0$.

□

Now for each knot K in S^3 , we would like a recurrence relation for the sequence of unreduced S^r -colored Alexander polynomials from Def. 2.3 and Eqn. (2.8).

$$\bar{\Delta}_K(r) := \bar{\Delta}_K^r(q) = \Delta_K(r)\bar{\Delta}_U(r)$$

Lemmas 2.6 and 2.8 gave recurrence relations, A_K and $(M-1)(L-1)$, for the sequences $\Delta_K(r)$ and $\bar{\Delta}_U(r)$ respectively. The existence of a recurrence for $\bar{\Delta}_K(r)$ now follows from the general theory, see [GL16, Thm. 5.2 (b)] or [Sab93, Cor. 2.1.6]. However, we are interested in explicit formulas so a direct calculation is included below.

Theorem 2.9. *The element $\bar{A}_K := (M-1)A_K \in \mathcal{A}$ annihilates the sequence $\bar{\Delta}_K \in \text{Seq}$ of unreduced S^r -colored Alexander polynomials defined by $\bar{\Delta}_K(r) := \bar{\Delta}_K^r(q)$.*

Proof. Define the coefficients a_j by the relation

$$A_K \Delta_K(r) = \sum_{j=r}^{r+d} a_j \Delta_K(j)$$

Lemma 2.6 shows $A_K \Delta_K = 0$, so the sum $\sum_{j=r}^{r+d} a_j \Delta_K(j) = 0$ is zero. Multiplying through by $\bar{\Delta}_U(r+d)$ shows that $\sum_{j=r}^{r+d} a_j \bar{\Delta}_U(r+d) \Delta_K(j) = 0$. Using the recurrence $\bar{\Delta}_U(r+d) = (q^{n+j} - 1)/(q^{n+d} - 1) \bar{\Delta}_U(r+j)$ from Lemma 2.8 for each term this sum gives $\sum_{j=r}^{r+d} a_j \frac{q^{n+j}-1}{q^{n+d}-1} \bar{\Delta}_U(r+j) \Delta_K(j) = 0$. Multiplying both sides of this sum by $q^{n+d} - 1$ gives us the relation $\sum_{j=r}^{r+d} a_j (q^{n+j} - 1) \bar{\Delta}_U(r+j) \Delta_K(j) = 0$. Simplifying further,

$$\begin{aligned} 0 &= \sum_{j=r}^{r+d} a_j (q^{n+j} - 1) \Delta_K(r+j) \bar{\Delta}_U(r+j) = (M-1) \sum_{j=r}^{r+d} a_j \Delta_K(r+j) \bar{\Delta}_U(r+j) \\ &= (M-1) A_K \Delta_K \bar{\Delta}_U \\ &= \bar{A}_K \bar{\Delta}_K \end{aligned}$$

□

3. OVERVIEW OF HEEGAARD-FLOER THEORY

This section contains a review of the Heegaard Floer materials which are needed for Section 4. In particular, we recall the bordered Heegaard Floer theory, its relation to the immersed curves interpretation of $\widehat{CFD}(S^3 \setminus \nu K)$ and how to compute $\widehat{HFK}(K)$ with its bigrading. The last subsection describes the cabling algorithm for immersed curves.

Heegaard Floer homology is a suite of closed 3-manifold invariants which was introduced by Ozsváth and Szabó [OS04c, OS04b]. For a closed 3-manifold Y and a spin^c structure \mathfrak{s} , there are graded \mathbb{F}_2 -vector space valued invariants $HF^\infty(Y, \mathfrak{s})$, $HF^\pm(Y, \mathfrak{s})$,

and $\widehat{HF}(Y, \mathfrak{s})$ of the pair (Y, \mathfrak{s}) . While these invariants are specified by combinatorial data, called w -pointed Heegaard diagrams, they measure subtle topological information about the manifold Y using pseudo-holomorphic curves.

To a knot K in a 3-manifold Y , one can associate a (z, w) -doubly pointed Heegaard diagram where the pair of marked points and handle decomposition specify K within Y . The extra z marked point induces a filtration on the Heegaard Floer chain complex associated to the w -pointed diagram. When $Y = S^3$, this results in the bi-graded knot Floer homology $\widehat{HFK}(K)$, [OS04a] and [Ras03b], which categorifies the Alexander polynomial in the sense that

$$\Delta_K(q) = \sum_{i,j} (-1)^i q^j \dim \widehat{HFK}^{i,j}(K).$$

The knot Floer invariants $\widehat{HFK}(K)$ are difficult to compute, see [Hom20]. However, Hanselman, Rasmussen and Watson [HRW24] interpreted the bordered Floer invariants [LOT18] identifying $\widehat{HFK}(K)$ with a chain complex of morphisms between immersed curves in the Fukaya category of a surface. It is this simpler setting which facilitates our study in Section 4. The remainder of this section contains a review of these ideas and their essential properties.

3.1. The curve invariant. Bordered Heegaard Floer homology, introduced by Lipshitz, Ozsváth, and Thurston [LOT18], is a relative version of Heegaard Floer homology. The bordered theory associates to a compact connected surface with one boundary component, Σ , a dg algebra \mathcal{A} . If M is a compact 3-manifold with boundary suitably¹ parameterized by a map $\phi : \Sigma \rightarrow M$ then there are two bordered invariants related to (M, ϕ) : a Type *D* structure $\widehat{CFD}(M, \phi)$, that is a left dg module over \mathcal{A} , and a Type *A* structure $\widehat{CFA}(M, \phi)$, that is a right \mathcal{A}_∞ -module over \mathcal{A} . When Y is a closed 3-manifold which is obtained by gluing 3-manifolds M and N along their boundaries using an orientation reversing homeomorphism $h : (\partial N, w_2) \rightarrow (\partial M, w_1)$

$$Y = M \sqcup_h N \Rightarrow \widehat{HF}(Y) \cong H_*(\widehat{CFA}(M, \phi) \boxtimes \widehat{CFD}(N, \bar{\phi})) \quad (3.1)$$

there is a product \boxtimes of these modules which recovers the Heegaard-Floer invariant of Y in the sense of Eqn. (3.1), see [LOT18, §9].

For 3-manifold with torus boundary, a parameterization ϕ is equivalent to choosing a pair of curves (α, β) of ∂M and fixing a basepoint $w \in \partial Y$. For knot complements

¹The theory is defined in terms of manifolds with fixed handle decompositions; we are glossing over these kinds of details here.

$M = S^3 \setminus \nu K$, the parameterization ϕ will always be described by the Seifert-framed meridian-longitude basis $\{\mu, \lambda\}$.

Hanselman, Rasmussen and Watson [HRW24, HRW22], also [HKK17, §4.4], show that Type D structures $\widehat{CFD}(M, \phi)$ are equivalent to immersed curves γ_M (elsewhere denoted by $\widehat{HF}(M)$ or $\widehat{\Gamma}(M)$) in the surface $T_M = \partial M \setminus w$, decorated by local systems and defined up to regular homotopy. For a knot complement $M := S^3 \setminus \nu K$, set $\gamma_K := \gamma_M$.

Remark 3.1. (Local systems) Knot complements $M = S^3 \setminus \nu K$ are of *loop type* [HW23b], as a consequence the associated immersed curves γ_K always have trivial local systems.

Remark 3.2. (Grading arrows) If the invariant γ_K has multiple components then they are connected by pairs of edges called grading arrows [HRW22, Definition 28]. While domains involving grading arrows do not contribute to the differential, they are considered when determining Maslov grading differences, as in Prop. 4.7.

The collection of immersed curves γ_M in T_M is almost always lifted to a curve $\bar{\gamma}_M$ in the covering space \bar{T}_M described next.

Definition 3.3. The covering space \bar{T}_M of $T_M = \partial M \setminus \{w\}$ is the cover whose fundamental group is the kernel of the composition

$$\pi_1(T_M) \rightarrow \pi_1(\partial M) \rightarrow H_1(\partial M) \rightarrow H_1(M).$$

When $H_1(M) \cong \mathbb{Z}$, this cover \bar{T}_M is homeomorphic to an infinite cylinder with z lifting to infinitely-many marked points at heights of $\frac{1}{2} + \mathbb{Z}$. The meridian μ lifts to a line $\bar{\mu}$ which passes through these lifted marked points (See Figure 1). In this paper, the infinite cylinder will always be pictured as a quotient of a horizontal infinite rectangle, heights are lower to the left and higher to the right.

A curve $\bar{\gamma}_0$ which “wraps around the cylinder” is a homologically essential simple closed curve, when the curve is oriented or the two quotient lines are ordered then the points of intersection between $\bar{\gamma}_0$ and other curves inherit an order.

Notation 3.4. For a 3-manifold M with torus boundary, a lift of the immersed curve invariant γ_M to the covering space \bar{T}_M typically depends on a spin^c structure of M and so must be denoted by $\bar{\gamma}_{(M, \mathfrak{s})}$ (or $\widehat{HF}(M, \mathfrak{s})$). However, there is a unique spin^c structure \mathfrak{s}_0 on knot complements $M_K := S^3 \setminus \nu K$ because $H^2(M_K) \cong H_1(M_K, \partial M_K) = 0$. So we will suppress \mathfrak{s}_0 from the notation $\bar{\gamma}_K$. We also adopt the notations $T_K := T_{M_K}$ and $\bar{T}_K := \bar{T}_{M_K}$ unless otherwise stated.

This curve valued knot invariant $\bar{\gamma}_K$ in \bar{T}_K has a few important structural properties which we list below.




FIGURE 1. The invariants for the unknot U and right-handed $T(2,3)$, lifted to \bar{T}_M . Integral heights between the marked points are indicated underneath.

Remark 3.5. (Structural properties)

- (1) The invariant $\bar{\gamma}_K$ contains an *essential component* $\bar{\gamma}_0$ that wraps around the cylinder \bar{T}_K . All of the other components of $\bar{\gamma}_K$ are closed, homologically inessential and lie within a neighborhood of the meridional curve $\bar{\mu}$ [HRW22, Corollary 64].
- (2) The invariant γ_K in T_K is invariant under hyperelliptic involution of T_K and so its lift $\bar{\gamma}_K$ to \bar{T}_K is unchanged under rotation by π [HRW22, Theorem 7]. As a consequence, we always take the lift of $\bar{\gamma}_K$ to be centered at height 0.

Remark 3.6. (Knot invariants)

- (1) The Seifert genus $g(K)$ is the largest height in \bar{T}_K for which any component of $\bar{\gamma}_K$ intersects $\bar{\mu}$ [HRW22, Proposition 48].
- (2) The invariant $\tau(K)$ is the height in \bar{T}_K at which $\bar{\gamma}_0$ first intersects $\bar{\mu}$ after “wrapping around the cylinder.”

The pairing theorem for bordered Heegaard-Floer invariants from Eqn. (3.1) admits an immersed curves interpretation too. The immersed curves γ_M or $\bar{\gamma}_M$ associated to Type D structures are naturally viewed as objects in the Fukaya category of T_M or \bar{T}_M respectively. In this category, there is a chain complex $(CF(\gamma, \eta), d)$ of morphisms called *intersection Floer theory*. As an \mathbb{F}_2 -vector space, $CF(\gamma, \eta)$ is generated by the transverse points of intersection $\gamma \pitchfork \eta$ between the two curves. The differential d counts bigons such as those in Figure 2, see [Abo08, §2] or [HRW24, §4.1]. The theorem below restates the pairing theorem from Eqn. (3.1) in these terms.

Theorem 3.7 ([HRW22, Theorem 2]). *Suppose that a closed 3-manifold Y is homeomorphic to a quotient $Y \cong M \sqcup_h N$, where the M and N are compact, oriented 3-manifolds with torus boundary and $h : (\partial N, w_2) \rightarrow (\partial M, w_1)$ is an orientation reversing homeomorphism. Then there is an isomorphism between the Heegaard-Floer*

homology of Y and the intersection Floer homology in T_M .

$$\widehat{HF}(M \sqcup_h N) \cong H_*(CF(\gamma_M, h(\gamma_N))).$$

This isomorphism respects relative gradings and $spin^c$ structures.

3.2. Knot Floer Homology. Knot Floer homology $\widehat{HFK}(K)$ for a knot $K \subset S^3$ is computed from a Heegaard diagram with two basepoints, w and z [OS04a, §3.1]. Such a diagram is constructed by upgrading a w -pointed Heegaard diagram for S^3 to a double (w, z) -pointed Heegaard diagram. After adding the extra z -basepoint, the Maslov grading, sensitive to bigons covering the w -point, is retained and a second *Alexander grading* is introduced which counts bigons covering the new z -point, see [Hom20].

The pairing Theorem 3.7 above allows us to introduce knot Floer homology $\widehat{HFK}(K)$ in Theorem 3.9 below. We want to glue the knot complement $M = S^3 \setminus \nu K$, which gives us the curve $\bar{\gamma}_K$, with the ∞ -framed solid torus, which determines the meridian curve $\bar{\mu}$. Without any modifications to Thm. 3.7, this pairing produces a chain complex which computes only the w -pointed homology $\widehat{HF}(S^3)$ of the gluing.

In order to recover knot Floer homology $\widehat{HFK}(K)$ and its Alexander filtration, the covering space \bar{T}_K is modified by replacing the lifts of the marked point w at heights $(\frac{2n+1}{2}, 0)$ with small disks D_n , (with radius small enough not to intersect the curve $\bar{\gamma}_K$). Within each disk D_n are lifts of marked points z and w at heights $(\frac{2n+1}{2}, -\epsilon)$ and $(\frac{2n+1}{2}, \epsilon)$, respectively. The meridian $\bar{\mu}$ is now the curve that goes between consecutive pairs of marked points.

To reconstruct the knot Floer homology, begin by setting

$$C^-(\bar{\gamma}_M, \bar{\mu}) := \mathbb{F}_2[U] \otimes_{\mathbb{F}_2} CF(\bar{\gamma}_M, \bar{\mu}).$$

This is generated over $\mathbb{F}[U]$ by the intersection points $\bar{\gamma}_M \cap \bar{\mu}$ between $\bar{\gamma}_M$ and $\bar{\mu}$. The additional z -basepoints added in the previous paragraph allow us to count bigons in new ways. If B is a bigon then denote by $n_w(B)$ and $n_z(B)$ the number of times B covers w and z respectively. Let $N_i^w(x, y) \in \mathbb{F}_2$ be the number of bigons B from x to y for which $n_w(B) = i$. The differential

$$d : C^-(\bar{\gamma}_M, \bar{\mu}) \rightarrow C^-(\bar{\gamma}_M, \bar{\mu}) \quad \text{is} \quad d(x) := \sum_{i=0}^{\infty} \sum_y U^i N_i^w(x, y) \cdot y.$$

Without modification, there is only a homotopy equivalence [HRW22, Warning §4.2] between associated graded objects

$$gr C^-(\bar{\gamma}_M, \bar{\mu}) \simeq gr CFK^-(K)$$

with respect to the Alexander filtration discussed below. The issue is that $C^-(\bar{\gamma}_M, \bar{\mu})$ is not necessarily a chain complex because of the existence of teardrop or fishtail

shaped regions. To obtain chain complex $C^-(\bar{\gamma}_M, \bar{\mu})$ which is equivalent to $CFK^-(K)$ one must deform the construction using a bounding chain that counts these teardrop-shaped regions [Han23b, Def. 3.12].

The hat-version of the C^- -construction is obtained by setting $U = 0$.

$$(\hat{C}(K), \partial^v) := (C^-(\bar{\gamma}_K, \bar{\mu}), d) \otimes_{\mathbb{F}_2[U]} \mathbb{F}_2 \quad (3.2)$$

This is always a chain complex because it agrees with the $U = 0$ specialization of the deformed C^- mentioned above². The homology of the chain complex $\hat{C}(K)$ computes knot Floer homology $\widehat{HFK}(K)$ in the sense of Thm 3.9 below. Before stating the theorem, we recall the filtration, the Alexander grading and the Maslov grading.

The *Alexander grading* A , or the q -grading, is computed using the intersection numbers n_z and n_w of bigons with the z and w -basepoints. In more detail, after choosing the symmetric presentation in Remark 3.5 (2), there is a function A on generators $\bar{\gamma}_K \frown \bar{\mu}$ of $C^-(\bar{\gamma}_M, \bar{\mu})$. If B is a bigon from x to y then

$$A(x) - A(y) = n_z(B) - n_w(B) \quad \text{and} \quad A(U \cdot x) = A(x) - 1.$$

There is a *filtration* $F^n C^-(\bar{\gamma}_M, \bar{\mu}) := \{x : A(x) \leq n\}$; the associated graded $gr C^-(\bar{\gamma}_M, \bar{\mu})$ is the chain complex which is obtained by considering only bigons B with $n_z(B) = 0$.

The *Maslov grading* M is the t -grading or homological grading. Suppose that x and y are two points of intersection between $\bar{\gamma}_K$ and $\bar{\mu}$, and B is a bigon from x to y . So the boundary of B consists of two piecewise smooth paths one from x to y in $\bar{\gamma}_K$ and the other from y to x in $\bar{\mu}$. The relative Maslov gradings can be computed using Hanselman's equation,

$$M(y) - M(x) = 2(\text{Wind}_w(B) + \text{Wght}(B) - \text{Rot}(B)). \quad (3.3)$$

see [Han23a]. In this formula, $\text{Wind}_w(B)$ is the net winding number of B around enclosed w basepoints. The other two terms involve grading arrows, see Rmk. 3.2 or [Han23a, §2.1], $\text{Wght}(B)$ is the sum of weights (counted with sign) of all grading arrows traversed by B and $\text{Rot}(B)$ is $\frac{1}{2\pi}$ times the total counterclockwise rotation along the smooth sections of B . Alternatively,

$$\text{Rot}(B) = \frac{1}{2\pi}(2\pi - a\frac{\pi}{2} - c\pi),$$

where a and c are the numbers of corners and cusps (from grading arrows) traversed respectively.

²This is because when $U = 0$, the differential of the deformed C^- -complex defined by Hanselman no longer counts teardrop-shaped regions when they cover a basepoint (as they cover a w -marked point) and any teardrop-shaped regions which do not cover a basepoint can be removed by homotopy invariance. So the contribution of the bounding chain to the differential at $U = 0$ is zero.



FIGURE 2. Bigons from x to y and from y to x , see Ex. 3.8

Example 3.8. (Maslov gradings) Figure 2 shows two important bigons for us. Each black circle is understood to have z and w marked points to their left and right sides, respectively. For both bigons, $\text{Wght}(B) = 0$ since there are no grading arrows traversed by the bigons. The bigon on the left satisfies $\text{Rot}(B) = \frac{1}{2}$ and bigon on the right satisfies $\text{Rot}(B) = -\frac{1}{2}$; the sign appears because they occur on different sides of the lateral axis. For the bigon on the left, there are no w -points so Eqn. (3.3) gives $M(y) - M(x) = 1$. For the bigon on the right, there are $(r-1)$ -many w -points and we have $M(y) - M(x) = 2r - 3$.

Once the chain complex $\hat{C}(K)$ has been given the structure above, the theorem below can be stated more precisely.

Theorem 3.9. ([HRW22, Thm. 52]) *The filtered complex $(\hat{C}(K), \partial^v)$ is homotopy equivalent to the filtered complex $\widehat{CFK}(K)$ associated to a knot by [OS04a] and [Ras03b]. In particular, the associated graded complex computes the knot Floer homology*

$$\widehat{HFK}(K) \cong H_*(gr\hat{C}(K))$$

3.3. Vertical Simplification. In this section we introduce vertically simplified bases. This will give us a reasonable toolkit for understanding the structure of $\hat{C}(K)$ from Eqn. (3.2) and its relationship to the curve $\bar{\gamma}_K$.

Definition 3.10. (Vertically simplified basis) A basis $\{x_i\}$ for a chain complex (C, d) is *vertically-simplified* when for each basis element x_i , exactly one of the following conditions holds:

- (1) $dx_i = x_j$ for some x_j ,
- (2) x_i is in the kernel but not the image of d ,
- (3) $x_i = dx_j$ for a unique x_j .

Remark 3.11. (1) Vertical simplification is equivalent to choosing an isomorphism of the form

$$C \cong (\oplus_i \Lambda(x_i)) \oplus D,$$

where d vanishes on D and $d(x_i) = 1_i$.

- (2) A bicomplex (C, d, δ) can be vertically-simplified with respect to d or with respect to δ . In general, it is unknown whether all of the bicomplexes appearing

in Heegaard Floer theory can be simplified with respect to both differentials simultaneously.

We now fix some notations related to the vertical simplification of $\hat{C}(K)$.

Notation 3.12. (1) For each knot K , the chain complex $\hat{C}(K)$ admits a vertically-simplified basis. Since $H_*(\widehat{CFK}(K), \partial^v) \cong \widehat{HF}(S^3) \cong \mathbb{F}_2\langle \chi_0 \rangle$, the subspace D above is 1-dimensional. There are isomorphisms

$$\hat{C}(K) \cong \mathbb{F}_2\langle \chi_0 \rangle \oplus \bigoplus_{i \in I} \Lambda(x_i) \quad (3.4)$$

where $\Lambda(x_i) = \mathbb{F}_2\langle 1_i, x_i \rangle$ and $\partial^v(x_i) = 1_i$.

- (2) The set I is ordered so that $j \geq i$ implies $A(x_j) \leq A(x_i)$.
- (3) The *distinguished generator* χ_0 generates $\widehat{HF}(S^3)$ under the identification from Theorem 3.9.
- (4) There is a *bottom generator* b which has the least Maslov grading among generators y which satisfy $A(y) = -g(K)$.

Remark 3.13. A vertically-simplified basis ensures that no two generators are connected by a differential that does not reduce the Alexander grading, such a basis is called *reduced*. Equivalently, there are no bigons in the differentials of $\hat{C}(K)$ or $C^-(\bar{\gamma}_K, \bar{\mu})$ that do not cross marked points. Note that, homotopy invariance of $\bar{\gamma}_K$ allows us to assume each curve segment between generators is length-minimizing. This also produces a reduced chain complex.

If a chain complex $CFK^-(K)$ which has been simultaneously horizontally and vertically simplified then the algorithm below tells us how to reconstruct the immersed curve $\bar{\gamma}_K$. This proposition is the immersed curves reformulation of [LOT18, Theorem 11.31].

Proposition 3.14. ([HRW22, Proposition 47]) *Given a basis for $CFK^-(K)$ which is both horizontally-simplified and vertically-simplified, the curve invariant $\bar{\gamma}_K$ is obtained in \bar{T}_K by:*

- (1) *For each basis element x of $CFK^-(K)$, place a short verticle segment at height $A(x)$.*
- (2) *If $CFK^-(K)$ contains a vertical arrow from x to y , then connect the top endpoints of their corresponding short segments by an arc.*
- (3) *If $CFK^-(K)$ contains a horizontal arrow from x to y , then connect the bottom endpoints of their corresponding short segments by an arc.*
- (4) *There will remain a unique segment with an unattached top endpoint and a unique segment with an unattached bottom endpoint. Attach the top endpoint to the top edge of \bar{T}_M at height 0 and attach the remaining bottom endpoint to the bottom edge of \bar{T}_M at height 0.*

As noted in Rmk. 3.11 (2), we cannot assume that such a simultaneously simplified basis exists. However, by ignoring step (3) and the latter half of step (4), we can obtain the partial information about $\bar{\gamma}_K$ discussed below. The next corollary will play a role in Lemma 4.4.

Corollary 3.15. *If K is a knot in S^3 and $\hat{C}(K) \cong \mathbb{F}_2\langle\chi_0\rangle \oplus \bigoplus_{i \in I} \Lambda(x_i)$ a vertically-simplified basis, with elements ordered as in Notation 3.12, then there is a half-completed curve $\frac{1}{2}\bar{\gamma}_K$ in \bar{T}_K which satisfies*

- (1) *Each generator y corresponds to a short vertical segment at height given by its Alexander grading $A(y)$ in \bar{T}_K .*
- (2) *A horizontal curve segment connects each pair x_i and 1_i along their top endpoints corresponding to the length n_i arrow in ∂^v .*
- (3) *The vertical segment corresponding to the distinguished generator χ_0 connects to the top edge of \bar{T}_K .*
- (4) *The vertical segment corresponding to the bottom generator b is a vertical segment of lowest height in \bar{T}_K .*

3.4. Cabling immersed curves. The Alexander polynomial transforms according to Eqn. (2.2) under the cabling operation described by Def. 2.1. In [HW23a], Hanselman and Watson discovered how the (p, q) -cable operation transforms the immersed curves description of Heegaard-Floer homology described above. If the curve $\bar{\gamma}_K$ is placed within a minimal box as in Fig. 3 then the $(3, 4)$ -cable $K_{3,4}$ of a knot K leads to the immersed curve $\bar{\gamma}_{K_{3,4}}$ in Figure 4.

Theorem 3.16 ([HW23a]). *Suppose that $\bar{\gamma}_K$ in \bar{T}_K is the immersed curve associated to a knot K in S^3 . Then the procedure below produces the immersed curve $\bar{\gamma}_{K_{p,q}}$ associated to the (p, q) -cable of K ³.*

- (1) *Create p parallel copies of \bar{T}_K . Place a copy of $\bar{\gamma}_K$ in each \bar{T}_K , so that they are scaled horizontally by a factor of p and the next lower copy of the multicurve is q units lower (in A -grading or height) than the previous copy*
- (2) *Connect the loose ends of the successive copies of the curve*
- (3) *Observe that the marked points (or disks) admit an isotopy which moves them uniformly vertically downward until they appear in order in the bottom copy of \bar{T}_K . Using this isotopy to carry the multicurve from steps (1) and (2) produces the immersed curve $\bar{\gamma}_{K_{p,q}}$.*

Notation 3.17. Step (3) allows us to view the resulting multicurve in the cover $\bar{T}_{K_{p,q}}$, but it also makes a mess, so in what follows it will be convenient to skip this step. We instead use a single zig-zag curve $\bar{\mu}$ through all unisotoped lifts of

³Compared to [HW23a], all of our diagrams have been rotated by $-\pi/2$.




FIGURE 3. A cabling pattern. A minimal box is $2g(K)$ wide.

the marked point as in Figure 4. In particular, when considering the $(r, rn + 1)$ -cable $K_{r, rn+1}$ of K , by a slight abuse of notation, we will use $\bar{\gamma}_{K_{r, rn+1}}$ to denote the immersed curve resulting from steps (1) and (2) of the above construction and \bar{T}^r to denote the ambient surface.

Notation 3.18. (\circ -dot) Moreover, when illustrating the curves associated to cablings, it is often useful to simplify the pictures by combining \bullet components above the first row into a single \circ -component, see Fig. 5 or Fig. 9.




FIGURE 4. The $T(3, 4)$ cabling applied to a pattern, see Fig. 3.

4. COLORED KNOT FLOER HOMOLOGY

Theorem 4.6 constructs a categorification of the S^r -colored Alexander polynomial which was introduced in §2.1. Corollary 3.15 contains a structure theorem for this filtered chain complex which will be used in Section 6.

Definition 4.1. (subdivision) Recall that by vertically-simplifying, every chain complex C over a field such as \mathbb{F}_2 can be decomposed into a sum

$$C \cong (\oplus_i \Lambda(x_i)) \oplus D,$$

where the differential of C vanishes on D and for each $\Lambda(x_i)$ -term, $d(x_i) := 1_i$. If C is bigraded then the *length* of a $\Lambda(x_i)$ -term is the Alexander or q -degree of d . With this in mind, we can introduce a chain complex $subdiv(C)$ which fixes D and replaces each $\Lambda(x_i)$ summand of length $A(d_i) = A(x_i) - A(1_i)$ by a direct sum of $A(d_i)$ terms of the form $\Lambda(x_{i,j})$ for $1 \leq j \leq n_i$ each of length 1. Alternatively, subdivision can be computed from the assignments below.

- (1) $subdiv(E \oplus F) := subdiv(E) \oplus subdiv(F)$
- (2) $subdiv(\mathbb{F}_2) := \mathbb{F}_2$
- (3) $subdiv((\Lambda(x_i), d)) := \bigoplus_{j=1}^{A(d)} (\Lambda(x_{i,j}), d_j)$ where $A(d_j) = 1$

If $(f_{x_i}, f_{1_i}) : \Lambda(x_i) \rightarrow \Lambda(x'_i)$ are the components of a chain map f then the assignment $subdiv(f_{x_i}, f_{1_i}) := (f_{x_i}, f_{1_i}, f_{x_i}, f_{1_i}, \dots, f_{x_i}, f_{1_i})$ extends $subdiv$ to a functor. This statement won't be used here, so further discussion is omitted.

The algebraic notion introduced above arises from an observation about the cabling procedure which is illustrated in Fig. 5 below.




FIGURE 5. Subdividing a bigon when $A(d) = 2$. For \circ -dot notation, see 3.18.

Example 4.2. Let $\Delta_m := \Lambda(\xi)$ where $d(\xi) = 1$ and $|\xi|_A = m$ be the 2-term chain complex of Alexander degree m . Then the chain complex $D_m := subdiv(\Delta_m)$ consists of m , A -degree 1, 2-term complexes:

$$D_m = \bigoplus_{i=1}^m \Lambda(\xi_i).$$

Moreover, the relation $D_m \cong D_{m-1} \oplus \Lambda(\xi_m)$ determines embeddings $\iota_m : D_{m-1} \hookrightarrow D_m$ and the limit of the sequence

$$D_1 \xrightarrow{\iota_1} D_2 \xrightarrow{\iota_2} D_3 \rightarrow \dots$$

can be identified with the unknot homology $S^r \widehat{CFK}(U)$ in Example 4.8 (up to grading considerations which will depend upon r).




FIGURE 6. A schematic for Lemma 4.4. The head consists of contributions from the lowest box. The tail begins on the non-compact curve at χ_0 and continues until reaching the penultimate box.

Definition 4.3. The 1-colored knot Floer complex is defined to be the usual knot Floer complex, $S^1\widehat{CFK}(K) := \widehat{C}(K)$, from Eqn. (3.2). Much like r -colored knot Floer complexes introduced later, the 1-colored complex has a “head-tail” decomposition

$$S^1\widehat{CFK}(K) \cong Hd(K) \oplus Tl(K)$$

compatible with the vertical-simplification Notation 3.12, in the sense that

$$Hd(K) := \bigoplus_{i \in I} \Lambda(x_i) \quad \text{and} \quad Tl(K) := \mathbb{F}_2 \langle \chi_0 \rangle$$

Lemma 4.4. Let g be the genus of the knot K and $K_{r, rn+1} := C_{r, rn+1}(K)$. When $r > 1$ and $n > 2g - 1$ the chain complex $\widehat{C}(K_{r, rn+1})$ admits a “head-tail” decomposition, an ungraded isomorphism of the form:

$$\tau_A^{\leq rn+1} \widehat{C}(K_{r, rn+1}) \cong subdiv(Hd(K)) \oplus subdiv(\Delta_m)$$

where $m = n - 2\tau(K) - 1$, $\tau_A^{\leq n} C := \{x \in C : A(x) \leq n\}$, $Hd(K)$ appears in Def. 4.3 and Δ_m in Ex. 4.2.

Proof. The basic setting is as follows. The curve invariant $\bar{\gamma}_{K_{r, rn+1}}$ in \overline{T}_M^r associated to $(r, rn+1)$ -cable $C_{r, rn+1}(K)$ of K is obtained from the curve invariant $\bar{\gamma}_K$ of K using the algorithm in Theorem 3.16. By Theorem 3.9,

$$\widehat{CFK}(C_{r, rn+1}(K)) \simeq \widehat{C}(K_{r, rn+1}) = CF(\bar{\gamma}_{K_{r, rn+1}}, \bar{\mu}) \otimes_{\mathbb{F}_2[u]} \mathbb{F}_2.$$

implies that the knot Floer complex is the intersection Floer complex. In keeping with Rmk. 3.13, we choose a length-minimizing representative of $\bar{\gamma}_K$ and a length-minimizing representative of the corresponding cabling invariant $\bar{\gamma}_{K_{r, rn+1}}$. With these choices, the generators of $\widehat{C}(K_{r, rn+1})$ form a reduced basis with respect to the vertical differential ∂^v . (Note by assuming a minimal representative of $\bar{\gamma}_{K_{r, rn+1}}$ the boxes in

the discussion of the cabling algorithm may not be so box-like, but the initial choice of a curve in a box suffices to identify and partition generators in the reduced complex.)

The key structural observation about the cabling algorithm (Thm. 3.16) is that when the $(r, rn + 1)$ -cabling parameter $n > 2g - 1$ is sufficiently large, two things are true:

- (1) the two copies of $\bar{\gamma}_K$ in the penultimate and final rows occupy disjoint heights.
- (2) the arc that connects these two copies of $\bar{\gamma}_K$ is horizontal and its length is proportionate to n

This observation is illustrated in Fig. 6. The head is determined by the copy of $\bar{\gamma}_K$ in the bottom row. The tail formed by the intersection points on the curve between the generator χ_0 and the start of the second copy of $\bar{\gamma}_K$.

Head identification: Let $\bar{\gamma}_{rK}$ be the “ r -scaled copy” of $\bar{\gamma}_K$ contained within the box in the bottom row of the cabling construction. Since $\bar{\gamma}_K$ is bound within the region of Alexander length $2g$, the curve $\bar{\gamma}_{rK}$ is bound inside the lowest $(2g - 1)r + 1$ terms. Set

$$\hat{C}_r(K) := H_*(CF(\bar{\gamma}_{rK}, \bar{\mu})).$$

Since the boxes are topologically identical we need only analyze how changing $\bar{\mu}$ to the zig-zag curve (as in Fig. 4) changes the chain complex. We assume that the chain complexes $\hat{C}(K)$ and $\hat{C}_r(K)$ have been vertically simplified and follow the conventions of §3.3. Following Cor. 3.15 (3), the vertical generator χ_0 is the distinguished vertical segment which connects to the box in the penultimate row. The remaining vertical segments correspond to terms $\Lambda(x_i)$ with $d(x_i) = 1_i$ each of which corresponds to a bigon of length n_i . To each such term there is a bigon B appearing as the left-hand side of Fig. 5. Associated to each such region is a subdivided region appearing as the right-hand side of Fig. 5.

Let us explain in detail how a bigon is subdivided. Suppose generators x_i and 1_i from $\hat{C}(K)$ are connected by a length n_i bigon. All of the bigons are always $\bar{\gamma}_{rK}$ on the right and $\bar{\mu}^r$ on the left. In $\hat{C}(K_{r, rn+1})$, the arc connecting the top endpoints of their associated curve segments makes $2(n_i - 1)$ additional intersections with $\bar{\mu}^r$. Introduce new generators $x_{i,j}$ and $1_{i,j}$. Set $x_{i,n_i} := x_i$ and $1_{i,1} := 1_i$ so that

- (1) the bigon B from $x_{i,j}$ to $1_{i,j}$, covers a single z -marked point ($n_z(B) = 1$) for $1 \leq j \leq n_i$,
- (2) the bigon B from $x_{i,j}$ to $1_{i,j+1}$, covers $r - 1$ w -marked points ($n_w(B) = r - 1$) for $1 \leq j \leq n_i - 1$,

with \bar{T}^r understood to contain pairs of z and w marked points (as in the discussion following Theorem 3.7). Labeled in this way, we collect these generators according to

increasing Alexander grading by pairs admitting a bigon covering a single z marked point:

$$1_{i,1} \leftarrow x_{i,1}, 1_{i,2} \leftarrow x_{i,2}, \dots, 1_{i,j} \leftarrow x_{i,j}, \dots, 1_{i,n_i} \leftarrow x_{i,n_i}$$

Each pair $1_{i,j} \leftarrow x_{i,j}$ is connected by a bigon pictured on the left in Figure 2. This collection agrees with $\text{subdiv}(\Lambda(x_i))$ from Def. 4.1.

The above only specifies an embedding of $\text{subdiv}(Hd(K)) \hookrightarrow \tau_A^{\leq rn+1} \hat{C}(K_{r,rn+1})$. There can be no additional generators besides those accounting for the tail below because, after performing step (3) of the cabling algorithm, the pairs of curves $(\bar{\mu}^r, \bar{\gamma}_{rK})$ and $(\bar{\mu}, \bar{\gamma}_K)$ can be assumed to be equal except for the subdivision occurring due to the adding of basepoints.

Tail identification: The remaining generators are due to the arc connecting the distinguished generator χ_0 to the copy of $\bar{\gamma}_{rK}$ in the penultimate row, the arc begins outside of the lowest box in Fig. 6 and continues until it reaches the start of the second highest box in Fig. 6. The q -grading of the distinguished generator is given by $A(\chi_0) = r\tau(K)$, after this the arc forms pairs of intersections identical to those in the discussion of the head above until it reaches the height of the left edge of the bounding box in the the penultimate row. The particulars of how the arc connect to the second copy of $\bar{\gamma}_K$ do not change our estimate. This collection of generators introduced by the arc can be identified with the dilation $\text{subdiv}(\Delta_m)$ where $m = n - 2\tau(K) - 1$. □

In order to identify the limit of $(r, rn + 1)$ -cablings of K as $n \rightarrow \infty$ in Theorem 4.6, we next recall Rozansky's convergence criteria.

Suppose that \mathcal{A} is an additive category and $K^+(\mathcal{A})$ is the homotopy category of chain complexes over \mathcal{A} which are unbounded in positive degree. If $\iota : A \rightarrow B$ is a map then the *isomorphism order* is defined to be

$$|\iota|_{\cong} := \max\{n : \tau^{\leq n}(\iota) \text{ is an isomorphism}\},$$

where $\tau^{\leq n} : K^+(\mathcal{A}) \rightarrow K^{\leq n}(\mathcal{A})$ is truncation above degree n .

Lemma 4.5. ([Roz14, Theorem 3.4]) *Suppose that \mathcal{A} is an additive category and $K^+(\mathcal{A})$ is the associated homotopy category of chain complexes. Then a directed system has a limit if and only if the isomorphism orders diverge.*

$$\varinjlim (C_1 \xrightarrow{\iota_1} C_2 \xrightarrow{\iota_2} C_3 \xrightarrow{\iota_3} \dots) \quad \Leftrightarrow \quad \lim_{i \rightarrow \infty} |\iota_i|_{\cong} = \infty$$

Recall the element $b \in \hat{C}(K)$ from Notation 3.12 (4) is the element of lowest Alexander grading and least Maslov grading. For any knot K , the chain complex $|b|^{-1} \widehat{CFK}(K) = t^{-M(b)} q^{-A(b)} \hat{C}(K)$ has been shifted so that this generator lies in degree $(0, 0)$. The corollary uses the lemma to construct chain complexes $S^r \widehat{CFK}(K)$.

Corollary 4.6. *For each knot K in S^3 , there is a limit of filtered chain complexes*

$$S^r \widehat{CFK}(K) := \lim_{n \rightarrow \infty} |b_{r,n}|^{-1} \hat{C}(K_{r, rn+1})$$

where $K_{r, rn+1} = C_{r, rn+1}(K)$ is the $(r, rn+1)$ -cable of K and $b_{r,n} \in \hat{C}(K_{r, rn+1})$ is the bottom generator. The homology of this chain complex is an invariant of the knot K .

Proof. By virtue of Lemma 4.4, there are ungraded isomorphisms

$$\tau_A^{\leq rn+1} \hat{C}(K_{r, rn+1}) \cong \text{subdiv}(Hd(K)) \oplus \text{subdiv}(\Delta_m)$$

Setting $C_{r,n} := \tau_A^{\leq rn+1} |b_{r,n}|^{-1} \hat{C}(K_{r, rn+1})$. Under these identifications we can define inclusion maps $\iota_n : C_{r,n} \hookrightarrow C_{r, n+1}$. The maps ι_n are identity on $\text{subdiv}(Hd(K))$ and agree with the inclusions ι_n appearing in Ex. 4.2 above on tail terms. The maps ι_n can be upgraded to graded maps because the curves and associated grading data are identical in the regions under consideration. The isomorphism order diverges because the reduced complexes have the structure Eqn. (3.4) and the maps ι_n are identity maps. We set

$$S^r \widehat{CFK}(K) := \lim_{n \rightarrow \infty} C_{r,n}$$

The homology of $S^r \widehat{CFK}(K)$ is an invariant of K since $\bar{\gamma}_K$ and the absolute Maslov and Alexander gradings of the generators $b_{r,n}$ involved in the grading shifts above are invariants of K . Additionally, the graded Euler characteristic of S^r -colored knot Floer homology matches the S^r -colored Alexander polynomial of K since the Euler characteristic commutes with limits (matching Definition 2.3). \square

Proposition 4.7. *There is an (ungraded) isomorphism of vector spaces*

$$\psi^r : S^r \widehat{CFK}(K) \rightarrow S^{r+1} \widehat{CFK}(K).$$

The (t, q) -degrees of ψ^r are affine linear functions of r .

Proof. If we forget about gradings then there are canonical isomorphisms $\psi^r : S^r \widehat{CFK}(K) \rightarrow S^{r+1} \widehat{CFK}(K)$ by virtue of the r -independence of the expression from Cor. 4.6:

$$S^r \widehat{CFK}(K) = \text{subdiv}(Hd(K)) \oplus \lim_{n \rightarrow \infty} \text{subdiv}(\Delta_{n-2\tau-1}).$$

Notice that it suffices to show affine linearity of ψ^r on generators 1_i since the others are related to an even-index generator by a bigon containing a z marked point which introduces a grading difference that is independent of r . We will show affine linearity of ψ^r in Alexander and Maslov gradings separately.

Alexander gradings. For a generator y , we write $A(y)$ for the Alexander grading and $A'(y)$ for the Alexander grading in the shifted complex from Cor. 4.6. We will first investigate grading changes associated to χ_0 and then turn to the other




FIGURE 7. The right-handed trefoil is $\bar{\gamma}_K$ placed into a box with distinguished generators. Each \bullet contains a z ϵ -above and a w ϵ -below.




FIGURE 8. The limit of the right-handed trefoil cabling process applied to a pattern Fig. 3 after removing extraneous bigons. In the first figure the generators satisfy $A(\chi_0) - A(b) = 2$, but here this difference $A'(\chi_0) - A'(b) = 2 * 3$ is scaled by a factor of $r = 3$.

generators. The Alexander grading of χ_0 is straightforward to determine by looking at the r -scaled, shifted copy of $\bar{\gamma}_K$ in the final row of the cabling algorithm. The generator b , which we recall has least Maslov grading among the generators with minimal Alexander grading, sits at height 0. Being a generator with least Alexander grading, its associated curve segment necessarily turns to the right, or upward in height, to intersect $\bar{\mu}^r$ between the two bottom rows, as otherwise there would be a generator with a smaller Alexander grading. Likewise, the curve segment associated to χ_0 also turns upward to begin the tail between the final two rows. This means that the difference in their heights $A'(\chi_0) - A'(b)$ is divisible by r . (see Figure 8), and if h measures heights in $\hat{C}(K)$ we see that

$$A'(\chi_0) - A'(b) = r(A(\chi_0) - A(b)). \quad (4.1)$$

We know that the difference $A(\chi_0) - A(b)$ is precisely $\tau(K) + g(K)$ since the Alexander grading of x is $\tau(K)$ by definition, and the Alexander grading is shifted before r -scaling by $-A(b) = g(K)$. Thus, the shifted Alexander grading of χ_0 in $S^r \widehat{CFK}(K)$ is

$$A'(\chi_0) = r(\tau(K) + g(K)). \quad (4.2)$$

Finally, any other generator 1_i coming from a curve segment that turns upward, (the complex is reduced), has its Alexander grading differ from that of χ_0 as a multiple of r for the same reasons.

Maslov gradings. Since the generator b can belong to any component of the curve $\bar{\gamma}_K$, determining Maslov gradings of generators relative to that of b may involve traversing a grading arrow which introduces a non-zero $2\text{Wght}(B)$ term in Hanselman's Eqn. (3.3). Fortunately, the weight m of any such grading arrow is independent of r ⁴, since a grading arrow belonging to the bottom row does not cross over any marked points when constructing the cabled curve $\bar{\gamma}_{K,r,n+1}$.

We know from the general algorithm to build $\bar{\gamma}_K$ from $\widehat{CFK}^-(K)$ (not necessarily using a horizontally- and vertically-simplified basis) from Prop. 3.14 that a sequence of immersed bigons connects any two generators in $|b_{r,n}|^{-1}\hat{C}(K_{r,n+1})$. Let $*_i B_i$ be the concatenation of such a sequence of bigons which connects the bottom generator b to the distinguished generator χ_0 (of the $\bar{\gamma}_K$ in the lowest row). Using Eqn. (3.3), we have that

$$M(\chi_0) - M(b) = 2(\text{Wind}_w + \text{Wght} - \text{Rot})(*_i B_i) \quad (4.3)$$

$$= 2\text{Wind}_w(*_i B_i) + 2m - 2\text{Rot}(*_i B_i). \quad (4.4)$$

In this formula, m is the weight of the grading arrow traversed by the first bigon B_1 to go from b to a generator on the distinguished component $\bar{\gamma}_0 \subset \bar{\gamma}_K$ which “wraps around the cylinder.” If b already lies on $\bar{\gamma}_0$ then $m = 0$ because no grading arrow is required.

Observe that the next term, $\text{Rot}(*_i B_i)$, does not depend on r , since the bigons between intersections in $\bar{\gamma}_{K_{r+1,(r+1)n+1}} \curvearrowright \bar{\mu}^{r+1}$ can be assumed to belong to the two bottom rows and are unchanged under a slight homotopy of $\bar{\mu}$ to allow for an extra row above.

However, the $\text{Wind}_w(*_i B_i)$ -term will change because of the extra row of marked points. Observe that any bigon B_i between generators of the two bottom rows contains either collections of the $(r-1)$ -many w marked points from the rows above or collections of single w marked points from the bottom row. This implies that

⁴Specifically, we can always take grading arrows to lie to the right of $\bar{\mu}$, since no marked points cross the arrows as we move to $\bar{\gamma}_{rK}$, none of its features change.

$\text{Wind}_w(B_i)$ is a linear combination of $(r - 1)$ and 1 , so that we may sum these contributions and upgrade Equation 4.3 to be

$$M(x) - M(b) = 2\text{Wind}_w(*_i B_i) + 2m - 2\text{Rot}(*_i B_i) \quad (4.5)$$

$$= 2\alpha(r - 1) + 2\beta + 2m - 2\text{Rot}(*_i B_i) \quad (4.6)$$

$$= 2\alpha(r - 1) + \delta, \quad (4.7)$$

with δ capturing the r -invariant contribution to the grading difference. An affine linear function of $r - 1$.

Since there is a sequence of bigons connecting the distinguished generator χ_0 to any generator y in the image of the vertical differential (such as 1_i), the same reasoning about the location and shape of bigons, as well as the weight of a potential grading arrow, applies to show that

$$M(y) - M(\chi_0) = 2\kappa(r - 1) + \nu. \quad (4.8)$$

(Note that an affine function of $(r - 1)$ is an affine function of r .) \square

Example 4.8. The illustration below shows two immersed curves: the stable curve $[U]_r$ associated to the unknot is drawn in red and the lift of the meridian μ is the blue curve which zigzags between the two different types of base points (see Not. 3.18). The points of intersection between the two curves constitute generators for the chain complex $S^r \widehat{CFK}(U)$, see Fig. 9.




FIGURE 9. The stable curve associated to the unknot.

The Poincaré polynomial of the S^r -colored unknot $S^r \widehat{CFK}(U)$ is

$$P(U)_{S^r}(t, q) = \frac{1 + tq}{1 - t^{2(r-1)}q}.$$

As a vector space

$$\begin{aligned} S^r \widehat{CFK}(U) &\cong \mathbb{F}_2[u] \otimes \Lambda(\xi) \\ &\cong \mathbb{F}_2 \langle u^n \xi^k : n \in \mathbb{Z}_{\geq 0}, k \in \{0, 1\} \rangle. \end{aligned}$$

The grading is given by $|u| = t^{2(r-1)}q^r$ and $|\xi| = t^1q^1$ so that $|u^n \xi^k| = |u|^n |\xi|^k$. There is a differential $d : S^r \widehat{CFK}(U) \rightarrow S^r \widehat{CFK}(U)$ which is determined by the assignments

$$d(u) := 0 \quad \text{and} \quad d(\xi) := 1$$

after extending by the Leibniz rule. The filtration is given by

$$F^\ell(S^r \widehat{CFK}(U)) := \{x : |x|_q \leq \ell\}.$$

Notice that, since the differential d decreases q -degree by one, the differential of the associated graded chain complex $S^r \widehat{CFK}(U) = gr(S^r \widehat{CFK}(U))$ is identically zero. This is a general phenomenon.

Example 4.9. In a similar way, there is a filtered chain complex $S^r \widehat{CFK}(3_1)$ associated to the right-handed trefoil, see Fig. 8. The stable curve is pictured in here.




FIGURE 10. Stable curve associated to the right-handed trefoil

The Poincaré polynomial of the right-handed trefoil is given by two terms associated to the intersection points near the “head” of the curve, together with a “tail” which can be identified with a shifted copy of the S^r -colored unknot found in Example 4.8.

$$P(3_1)_{S^r}(t, q) = 1 + tq + t^{4r-2}q^{4r}P(U)_{S^r}$$

As a vector space

$$\begin{aligned} S^r \widehat{CFK}(3_1) &:= \Lambda(c) \oplus t^{4r-2}q^{4r} S^r \widehat{CFK}(U) \\ &\cong \Lambda(c) \oplus t^{4r-2}q^{4r} (\mathbb{F}_2[u] \otimes \Lambda(\xi)) \end{aligned}$$

where $\Lambda(c) := \mathbb{F}_2\langle 1_c, c \rangle$ and $\Lambda(\xi) := \mathbb{F}_2\langle \xi, 1 \rangle$ and the bidegrees are given by $|c| = t^1q^1$, $|u| = t^{4r-2}q^{4r}t^{2(r-1)}q = t^{6r-1}q^{4r+1}$ and $|\xi| = t^{4r-2}q^{4r}t^1q^1 = t^{4r-1}q^{4r+1}$. The differential $d : S^r \widehat{CFK}(3_1) \rightarrow S^r \widehat{CFK}(3_1)$ is determined by the assignments $d(c) := 1_c$, $d(u) := 0$ and $d(\xi) := 1$.

As for the unknot, the filtration is defined by q -degree and the differential vanishes on the associated graded complex.

The proposition below summarizes the information contained in this section. The filtered graded chain complexes we constructed in this section are similar in form to the right-handed trefoil above Ex. 4.9 but with some extra $\Lambda(a)$ -type terms. The contents of this section which are used by §6 are summarized by the proposition below.

Proposition 4.10. *For each knot K , the filtered chain complex $S^r \widehat{CFK}(K)$ admits a canonical decomposition. In the category $H^0(F^*Ch_{\mathbb{Z}})$ there are isomorphisms*

$$S^r \widehat{CFK}(K) \cong (\Lambda(a_{1,r}) \oplus \Lambda(a_{2,r}) \oplus \cdots \oplus \Lambda(a_{n,r})) \oplus \theta^r S^r \widehat{CFK}(U). \quad (4.9)$$

where θ is an overall (t, q) -degree shift.

The head of the decomposition consists of a finite sum $\bigoplus_i \Lambda(a_{i,r})$ of 2-dimensional complexes,

$$\Lambda(a_{i,r}) \cong \mathbb{F}_2 \langle 1_{i,r}, a_{i,r} \rangle \quad (4.10)$$

The differential is defined to be $d(a_{i,r}) := 1_{i,r}$ with grading equal to $|a_{i,r}| = tq|1_{i,r}|$. Moreover, for each $1 \leq i \leq n$, the grading $|1_{i,r}| = t^{mr+b}q^{nr+c}$ is an affine linear function in r .

On the other hand, the tail of the decomposition is a shifted copy of the unknot complex $\theta^r \widehat{S^r \text{CFK}}(U)$ from Ex. 4.8,

$$\widehat{S^r \text{CFK}}(U) \cong \mathbb{F}_2[u] \oplus \Lambda(\xi)$$

the grading is determined by the assignments $|u| = t^{2(r-1)}q^r$ and $|\xi| = tq$ in the sense that $|u^n \xi^k| = |u|^n |\xi|^k$ and the differential is fixed by setting $d(u) := 0$ and $d(\xi) := 1$.

Remark 4.11. The stable curve in the infinite annulus isn't an object in the same Fukaya category as the curves associated to the cablings. We do not address here the question of how to describe the category in which this curve lives.

5. WEYL ALGEBRA AND HOMOLOGICAL HOLONOMICITY

5.1. Some algebraic definitions. If \mathcal{A} is an abelian category then there is an additive category $F^* \mathcal{A}$ of *filtered objects* in \mathcal{A} . An object is a sequence $E := \{F^n E\}_{n \in \mathbb{Z}}$ with inclusions $F^{n+1} E \subseteq F^n E$. Maps $f : E \rightarrow E'$ are maps in \mathcal{A} which commute with inclusions. Similarly, if \mathcal{A} is an additive category then there is an additive category $\mathcal{A}_{\mathbb{Z}}$ of *graded objects* in \mathcal{A} . An object of $\mathcal{A}_{\mathbb{Z}}$ is a sequence $\{E^n\}_{n \in \mathbb{Z}}$ of objects in \mathcal{A} and maps $f : E \rightarrow E'$ are collections $\{f^n : E^n \rightarrow E'^n\}_{n \in \mathbb{Z}}$ of maps in \mathcal{A} . There is a functor called *associated graded*

$$gr : F^* \mathcal{A} \rightarrow \mathcal{A}_{\mathbb{Z}}$$

which is defined on objects by $gr(\{F^n E\}_{n \in \mathbb{Z}}) := F^m E / F^{m+1} E$.

For an additive category \mathcal{A} , there is a category $Ch(\mathcal{A})$ of chain complexes over \mathcal{A} .

Definition 5.1. ($Ch(\mathcal{A})$) A *chain complex* $E \in Ob(Ch(\mathcal{A}))$ consists of pair $(\{E^k\}_{k \in \mathbb{Z}}, d_E^k : E^k \rightarrow E^{k+1})$ where $E^k \in Ob(\mathcal{A})$ and $d_E^{k+1} d_E^k = 0$ for all $k \in \mathbb{Z}$.

A map $f \in Hom^k(E, E')$ of degree k is a collection $f = \{f^i : E^i \rightarrow E'^{k+i}\}_{i \in \mathbb{Z}}$ of maps in \mathcal{A} . For any such map, there is a map $\delta^k f \in Hom^{k+1}(E, E')$ with components

$$(\delta^k f)^i := d_{E'}^{k+i} f^i + (-1)^{i+1} f^{i+1} d_E^i$$

In this way, the set of maps $Hom(E, E') = (\{Hom^k(E, E')\}_{k \in \mathbb{Z}}, d^k : Hom^k(E, E') \rightarrow Hom^{k+1}(E, E'))$ is a chain complex.

The composition satisfies the Leibniz rule with respect to the differential making $Ch(\mathcal{A})$ a category which is enriched over itself. More generally, a category which is enriched over chain complexes is called a *differential graded (dg) category*.

A dg category \mathcal{C} has a number of auxilliary structures stemming from the differential. A map f is a *cycle* if $\delta f = 0$. Two maps, f and g are *homotopic* or $f \simeq g$ when there is a map H such that $\delta H = f - g$. There is a category $Z^0(\mathcal{C})$ consisting of degree zero cycles and the *homotopy category* $H^0(\mathcal{C}) := Z^0(\mathcal{C})/\simeq$ consisting of cycles modulo boundaries. Two objects $E, F \in Ob(\mathcal{C})$ are *homotopy equivalent* $E \simeq F$ when they are isomorphic $E \cong F$ in the homotopy category $H^0(\mathcal{C})$. An object $E \in Ob(\mathcal{C})$ is *contractible* when $E \simeq 0$.

5.2. Action of Weyl algebra. The dg category \mathcal{Seq} introduced below will be our replacement for the collection of sequences Seq .

Definition 5.2. ($\mathcal{Seq}(\mathcal{A})$) Suppose \mathcal{A} is a dg category. Then the dg category of sequences is defined by $\mathcal{Seq}(\mathcal{A}) := \prod_{i \in \mathbb{Z}_{\geq 0}} \mathcal{A}$. In more detail, the objects are given by sequences $E = (E_i)_{i \in \mathbb{Z}_{\geq 0}}$ of objects $E_i \in Ob(\mathcal{A})$. A map of degree k from $E = (E_i)_{i \in \mathbb{Z}_{\geq 0}}$ to $E' = (E'_i)_{i \in \mathbb{Z}_{\geq 0}}$ is given by an element of the product

$$Hom_{\mathcal{Seq}(\mathcal{A})}^k(E, E') = \prod_{i \in \mathbb{Z}_{\geq 0}} Hom_{\mathcal{A}}^k(E_i, E'_i).$$

If $f : E \rightarrow E'$ then we write $f = (f_i)_{i \in \mathbb{Z}_{\geq 0}}$, or just $f = (f_i)$, for the sequence of maps which determine f . Given two maps $f : E \rightarrow E'$ and $g : E' \rightarrow E''$ then the composition and the differential are defined componentwise

$$(g \circ f) = (g_i \circ f_i)_{i \in \mathbb{Z}_{\geq 0}} \quad \text{and} \quad \delta(f) := (d_{E_i} f_i)_{i \in \mathbb{Z}_{\geq 0}} \quad (5.1)$$

The identity map is $1_E = (1_{E_i})_{i \in \mathbb{Z}_{\geq 0}}$.

For a dg category \mathcal{A} , the Grothendieck group $K_0(\mathcal{A})$ is simply an abelian group. When a category \mathcal{A} has additional structure, this structure is reflected in $K_0(\mathcal{A})$. For instance, if \mathcal{A} supports a \mathbb{Z} -action then the Grothendieck group $K_0(\mathcal{A})$ is a $\mathbb{Z}[q, q^{-1}]$ -module. If $\mathcal{A} = (\mathcal{A}, \otimes, 1)$ is a monoidal category then $K_0(\mathcal{A})$ is a ring. More details and a discussion of the ideas surrounding the statement of the proposition below can be found in the reference [KMS09].

Proposition 5.3. *Suppose that \mathcal{A} is a dg category which supports a \mathbb{Z} -action. Then there are dg functors*

$$\mathcal{M}, \mathcal{L} : \mathcal{Seq}(\mathcal{A}) \rightarrow \mathcal{Seq}(\mathcal{A})$$

and a natural isomorphism

$$R : \mathcal{M} \circ \mathcal{L} \xrightarrow{\sim} t^2 q \circ \mathcal{L} \circ \mathcal{M}. \quad (5.2)$$

which lifts the action of the Weyl algebra on the right-hand side of the map

$$\Phi : K_0(\mathcal{Seq}(\mathcal{A})) \rightarrow K_0(\mathcal{A})^{\mathbb{Z}_{\geq 0}} \quad \text{given by} \quad \Phi((E_i)_{i \in \mathbb{Z}_{\geq 0}}) := ([E_i])_{i \in \mathbb{Z}_{\geq 0}}$$

Proof. In order to introduce the map Φ , one uses $K_0(\mathcal{A} \times \mathcal{B}) \cong K_0(\mathcal{A}) \times K_0(\mathcal{B})$, see [Wei13, Ex. 2.1.4, p. 75].

Recall from §2.2 that there are maps $M : \text{Seq} \rightarrow \text{Seq}$ and $L : \text{Seq} \rightarrow \text{Seq}$ which are defined on functions f by Eqn. (2.11). In the proof of the statement above we will construct functors \mathcal{M} and \mathcal{L} which have the same effect as M and L on objects. Then we will find the isomorphism R .

The functor $\mathcal{M} : \text{Seq}(\mathcal{A}) \rightarrow \text{Seq}(\mathcal{A})$ is given by $\mathcal{M} := \prod_{i \in \mathbb{Z}_{\geq 0}} t^{2i-2} q^i$. In more detail, \mathcal{M} is defined on objects $E = (E_i)_{i \in \mathbb{Z}_{\geq 0}}$ by

$$\mathcal{M}(E) := ((\mathcal{M}E)_i)_{i \in \mathbb{Z}_{\geq 0}} \quad \text{where} \quad (\mathcal{M}E)_i := t^{2i-2} q^i E_i.$$

If $f : E \rightarrow E'$ is a map $f = (f_i)_{i \in \mathbb{Z}_{\geq 0}}$ of degree k then there is a map $\mathcal{M}(f) : \mathcal{M}(E) \rightarrow \mathcal{M}(E')$ of degree k the components of which are $(\mathcal{M}f)_i := t^{2i-2} q^i f_i$. These assignments determine a dg functor by virtue of universal properties and being a componentwise dg functor. Alternatively, a short proof of the three properties below is included.

- (i) $\mathcal{M}(g \circ f) = \mathcal{M}(g) \circ \mathcal{M}(f)$
- (ii) $\mathcal{M}(1_E) = 1_{\mathcal{M}(E)}$
- (iii) $d\mathcal{M}(f) = \mathcal{M}(df)$

In each case, we use the functoriality of the assignment $f_i \mapsto t^{2i-2} q^i f_i$. For (i), $\mathcal{M}(g \circ f)_i = t^{2i-2} q^i (g_i \circ f_i) = (t^{2i-2} q^i g_i) \circ (t^{2i-2} q^i f_i) = \mathcal{M}(g)_i \circ \mathcal{M}(f)_i$. For (ii), $\mathcal{M}(1_E)_i = t^{2i-2} q^i 1_{E_i} = 1_{t^{2i-2} q^i E_i} = 1_{\mathcal{M}(E)_i}$. For (iii), $d\mathcal{M}(f)_i = d(t^{2i-2} q^i f_i) = \mathcal{M}(df)_i$.

On the other hand, the functor $\mathcal{L} : \text{Seq}(\mathcal{A}) \rightarrow \text{Seq}(\mathcal{A})$ is given by shifting objects and maps by one. On objects $E = (E_i)_{i \in \mathbb{Z}_{\geq 0}}$ by

$$\mathcal{L}(E) := ((\mathcal{L}E)_i)_{i \in \mathbb{Z}_{\geq 0}} \quad \text{where} \quad (\mathcal{L}E)_i := E_{i+1}$$

and if $f = (f_i)_{i \in \mathbb{Z}_{\geq 0}}$ is a map then $\mathcal{L}(f)_i := f_{i+1}$. For properties (i), (ii) and (iii) above. For (i): $\mathcal{L}(g \circ f)_i = (g \circ f)_{i+1} = g_{i+1} \circ f_{i+1} = \mathcal{L}(g)_i \circ \mathcal{L}(f)_i$. For (ii): $\mathcal{L}(1_E)_i = 1_{E_{i+1}} = 1_{\mathcal{L}(E)_i}$. For (iii): $d\mathcal{L}(f)_i = df_{i+1} = \mathcal{L}(df)_i$.

Finally, for the map R in Eqn. (5.2). Given $f : E \rightarrow E'$, $f = (f_i)_{i \in \mathbb{Z}}$, the left-hand side is $(\mathcal{M} \circ \mathcal{L})(f)_i = \mathcal{M}(f_{i+1}) = t^{2i} q^{i+1} f_{i+1}$. On the right-hand side, we compute $(\mathcal{L} \circ \mathcal{M})(f)_i = t^{2i-2} q^i f_{i+1}$. The two sides differ by $t^2 q$. \square

Remark 5.4. In some cases it might be more natural to use $\mathcal{M} := \prod_i t^{2i} q^i$ or in Ex. 5.9 below, to use a two independent functors $\mathcal{M}_{t^2} := \prod_i t^{2(i-2)}$ and $\mathcal{M}_q := \prod_i q^i$.

Let us now specialize our discussion to the dg category of $\mathcal{A} = F^*Ch_{\mathbb{Z}}$ (filtered) graded chain complexes. This category will be denoted by

$$\text{Seq} := \text{Seq}(F^*Ch_{\mathbb{Z}}). \tag{5.3}$$

Shifting the internal grading, together with the filtration degree, up by 1 determines a functor $q : \mathcal{Seq} \rightarrow \mathcal{Seq}$ which generates a \mathbb{Z} -action $\mathbb{Z} \cong \langle q \rangle$. Shifting the homological degree in $F^*Ch_{\mathbb{Z}}$ yields a second automorphism $t : \mathcal{Seq} \rightarrow \mathcal{Seq}$ which commutes with the functors \mathcal{M} and \mathcal{L} . In this way, the category \mathcal{Seq} is a module over the t -extended Weyl algebra

$$\mathcal{A}^+ := \mathcal{A} \otimes_{\mathbb{Z}} \mathbb{Z}[t, t^{-1}] \quad (5.4)$$

The homotopy category $H^0(\mathcal{Seq})$ is triangulated, so that the category \mathcal{Seq} is considered to be pre-triangulated [BK90].

The next definition consists of three parts. From a very general sort of Whitehead tower to the definition of holonomicity which we use in §6.

Definition 5.5. (Holonomicity) An object E of a pre-triangulated category \mathcal{C} can be assembled from a subcategory $\mathcal{D} \subset \mathcal{C}$ if there is a length n sequence of distinguished triangles in $H^0(\mathcal{C})$ of the form

$$\begin{array}{ccccccc} X_1 & \rightarrow & X_0 & \rightarrow & Y_0 & \rightarrow & X_1[1] \\ X_2 & \rightarrow & X_1 & \rightarrow & Y_1 & \rightarrow & X_2[1] \\ & & & & \vdots & & \\ X_{n+1} & \rightarrow & X_n & \rightarrow & Y_n & \rightarrow & X_{n+1}[1] \end{array}$$

with $X_0 \cong E$, $X_{n+1} \cong 0$ and $\{Y_i\}_{i=0}^n \subset Ob(\mathcal{D})$. Such a sequence of distinguished triangles is *non-trivial* if $Y_0 \not\cong 0$. If an algebra B acts on a pre-triangulated category \mathcal{C} then let $\langle B \cdot E \rangle \subset \mathcal{C}$ be the smallest thick pre-triangulated subcategory containing objects of the form $F(E)$ for $F \in B$. An object $E \in Ob(\mathcal{Seq})$ is *holonomic* if $E \cong 0$ or $E \not\cong 0$ and it can be assembled in a non-trivial way from the smallest thick pre-triangulated category $\langle \mathcal{A}^+ \cdot E \rangle$ containing orbit of the Weyl action on E .

Under the identification given by Φ from Prop. 5.3 this condition becomes to q -holonomicity.

Remark 5.6. Our focus here is principally holonomicity as it is seen in relation to classical quantum topological ideas, but in this “categorified setting” it might also be interesting to investigate holonomicity properties of maps assigned to cobordisms.

5.3. Examples of holonomicity. This section is concluded with a collection of examples related to the definition of holonomicity from Def. 5.5 above.

Example 5.7. (Constant sequence) If $C \in Ob(F^*Ch_{\mathbb{Z}})$ then there is an object $\overline{C} \in Ob(\mathcal{Seq})$ given by $\overline{C}_n := C$ for all $n \geq 0$. There is a map $f : \overline{C} \rightarrow \mathcal{L}\overline{C}$ given by $f = (1_C : \overline{C}_n \rightarrow \overline{C}_{n+1})$. There is one distinguished triangle

$$X_1 \rightarrow X_0 \xrightarrow{f} Y_0 \rightarrow X_1[1]$$

with $X_1 \cong 0$, $X_0 = \overline{C}$ and $Y_0 = \mathcal{L}\overline{C}$.

Example 5.8. (Summing Sequence) For each complex $C \in Ob(F^*Ch_{\mathbb{Z}})$, there is an object $C^{\oplus} \in Ob(\mathcal{S}eq)$ where $C_n^{\oplus} := \bigoplus_{i=0}^n C$ for $n \geq 0$. There is a map $i : C^{\oplus} \rightarrow \mathcal{L}(C^{\oplus})$, $i = (i_n)_{n \geq 0}$ where $i_n : \bigoplus_{j=0}^n C \rightarrow (\bigoplus_{j=0}^n C) \oplus C$ is the inclusion of the first n -terms. Then the cone $C(i)_n$ is homotopy equivalent to a single copy of C . So the cone on f is isomorphic to the constant sequence, $C(f) \cong \bar{C}$ in $H^0(\mathcal{S}eq)$. Combining this observation with the construction for the constant sequence in Example 5.7 gives the following collection of distinguished triangles.

$$\begin{array}{ccccccc} X_1 & \rightarrow & X_0 & \xrightarrow{i} & Y_0 & \rightarrow & X_1[1] \\ X_2 & \rightarrow & X_1 & \xrightarrow{f} & Y_1 & \rightarrow & X_2[1] \end{array}$$

where $(X_0, X_1, X_2) = (C^{\oplus}, \bar{C}, 0)$ and $(Y_0, Y_1) = (\mathcal{L}(C^{\oplus}), \mathcal{L}(X_1))$.

On the other hand, unwinding the sequence of extensions above gives the diagram below.

$$\begin{array}{ccc} C^{\oplus} & \xrightarrow{i} & \mathcal{L}(C^{\oplus}) \\ f \downarrow & & \downarrow \mathcal{L}(f) \\ \mathcal{L}(C^{\oplus}) & \xrightarrow{\mathcal{L}(i)} & \mathcal{L}\mathcal{L}(C^{\oplus}) \end{array}$$

Holonomicity could also be formulated in terms of contractible twisted complexes of this sort.

The next example was explored by the first author and O. Yacobi circa 2012. This is the original motivation for Def. 5.5.

Example 5.9. (Pascal's Triangle) The Grassmannian $Gr(d, n)$ of d -planes in \mathbb{C}^n is stratified by Schubert varieties $Gr(d, n) = \cup_I X(I)$ [Bri05] where $I := (1 \leq i_1 < i_2 < \dots < i_d \leq n)$ and $\dim X(I) = 2 \sum_{k=1}^d (i_k - k)$. If $H(d, n) := H_*(Gr(d, n); \mathbb{Z})$ then there is an identification $H(d, n) = \bigoplus_I \mathbb{Z} \langle [X(I)] \rangle$. The Poincaré polynomial of the homology can be identified with the q -binomial coefficients as follows,

$$P(H(d, n))(t, q^{\frac{1}{2}}) = \bigoplus_k q^k \dim H_{2k}(Gr(d, n)) = \begin{bmatrix} n \\ d \end{bmatrix} = \frac{[n]!}{[d]![n-d]!}$$

where $[n] = (q^n - q^{-n})/(q - q^{-1})$ and $[n]! = [n][n-1] \cdots [1]$. This is a q -analogue of the binomial coefficient in the sense that the limit $q \rightarrow 1$ gives $n!/(d!(n-d)!)$.

There are degree zero maps $\alpha : q^{2(n-d)}H(d-1, n-1) \rightarrow H(d, n)$ and $\beta : H(d, n) \rightarrow H(d, n-1)$ which are defined by

$$\begin{aligned} \alpha([X(i_1, \dots, i_{d-1})]) &:= [X(i_1, i_2, \dots, i_{d-1}, n)] \quad \text{and} \\ \beta([X(i_1, \dots, i_d)]) &:= \begin{cases} 0 & \text{for } i_d = n \\ [X(i_1, i_2, \dots, i_d)] & \text{for } i_d < n \end{cases} \end{aligned}$$

It follows from the definitions above that there is a short exact sequence,

$$0 \rightarrow q^{2(n-d)}H(d-1, n-1) \xrightarrow{\alpha} H(d, n) \xrightarrow{\beta} H(d, n-1) \rightarrow 0.$$

This short exact sequence which determines Pascal's triangle also produces lots of combinatorial examples in our setting.

Example 5.10. (Unknot) Setting $S^r(U) := S^r \widehat{CFK}(U)$. From Ex. 4.8, the unknot homology can be identified with

$$\begin{aligned} S^r(U) &\cong \mathbb{F}_2[u] \otimes \Lambda(\xi) \\ &\cong \mathbb{F}_2 \langle u^n \xi^k : n \in \mathbb{Z}_{\geq 0}, k \in \{0, 1\} \rangle. \end{aligned}$$

The grading is given by $|u| = t^{2(r-1)}q^r$ and $|\xi| = t^1q^1$ so that $|u^n \xi^k| = |u|^n |\xi|^k$. The differential is determined by setting $d(u) := 0$ and $d(\xi) := 1$.

To build a recurrence relation for the sequence $(S^r(U))_{r \in \mathbb{Z}_{\geq 0}} \in Seq$, we begin by observing that there are maps

$$h_r : S^r(U) \rightarrow |u|S^r(U) \quad \text{where} \quad h_r(u^n \xi^k) := u^{n-1} \xi^k$$

the degree of h_r is $|h_r| = |u||u^{n-1} \xi^k| - |u^n \xi^k| = 0$, and h_r is a chain map which respects the q -filtration because it decreases q -degree: $h_r : F^\ell(S^r(U)) \rightarrow F^{\ell-r}(S^r(U)) \subset F^\ell(S^r(U))$ where $F^\ell(S^r(U)) := \{x : |x|_q \leq \ell\}$. The cone $C(h_r)$ is homotopy equivalent to the 2-dimensional subcomplex $\Lambda(\xi) := \mathbb{F}_2 \langle 1, \xi \rangle$. So there is a map $h : S^r(U) \rightarrow \mathcal{M}S^r(U)$ given by $h := (h_r)_{r \geq 0}$ such that

$$C(h) \cong \overline{\Lambda(\xi)} \quad \text{in} \quad Seq$$

is isomorphic to the constant sequence for $\Lambda(\xi)$ in Seq . The recurrence relation is now completed by following the Example 5.7 above, there is a map $f : C(h) \rightarrow \mathcal{L}(C(h))$ so that $C(f) \cong 0$ in Seq .

The diagram below illustrates our result.

$$\begin{array}{ccc} S^r(U) & \xrightarrow{h} & \mathcal{M}S^r(U) \\ \downarrow f & & \downarrow f \\ \mathcal{L}S^r(U) & \xrightarrow{\mathcal{L}(h)} & \mathcal{L}\mathcal{M}S^r(U) \end{array}$$

This diagram categorifies Lemma 2.8.

The next example is more representative of the general case in §6.

Example 5.11. (Right-handed trefoil) Set $S^r(3_1) := S^r \widehat{CFK}(3_1)$. From Ex. 4.9

$$\begin{aligned} S^r(3_1) &:= \Lambda(c) \oplus t^{4r-2} q^{4r} S^r \widehat{CFK}(U) \\ &\cong \Lambda(c) \oplus t^{4r-2} q^{4r} (\mathbb{F}_2[u] \otimes \Lambda(\xi)) \end{aligned}$$

This time the bidegrees are given by $|c| = t^1 q^1$, $|u| = t^{4r-2} q^{4r} t^{2(r-1)} q = t^{6r-1} q^{4r+1}$ and $|\xi| = t^{4r-2} q^{4r} t^1 q^1 = t^{4r-1} q^{4r+1}$. The differential is given by $d(c) := 1_c$, $d(u) := 0$ and $d(\xi) := 1$.

As with the unknot in Ex. 4.8 above, we define a map $f = (f_r)$, with components $f_r : S^r(3_1) \rightarrow |u|S^r(3_1)$ for which $f_r(c) := 0$, $f_r(1_c) := 0$ and $f_r(u^n \xi^k) := u^{n-1} \xi^k$, but now the neglected term $\Lambda(c)$ doubles in the sense that

$$C(f_r) \simeq \Lambda(c) \oplus |u|\Lambda(c) \oplus \Lambda(\xi)$$

Next there is a map $g := (g_r)$, the components $g_r : C(f_r) \rightarrow C(f_{r+1})$ are given by $g_r(c) := c$, $g_r(1_c) := 1_c$ and $g_r(x) := 0$ for $x \notin \Lambda(c)$. The cone is

$$C(g_r) \simeq [|u_r|\Lambda(c_r) \oplus \Lambda(\xi_r)] \oplus [|u_{r+1}|\Lambda(c_{r+1}) \oplus \Lambda(\xi_{r+1})]$$

where we include subscripts on the generators to emphasize their different graded degrees. There is a map $h = (h_r)$, where $h_r : C(g_r) \rightarrow \alpha C(g_{r+1})$ with $\alpha := |u_r|/|u_{r+1}| = t^{6r-1} q^{4r+1} / (t^{6(r+1)-1} q^{4(r+1)+1}) = t^{-6} q^{-4}$ is independent of r , h_r identifies the terms indicated below

$$|u_r|\Lambda(c_r) \xrightarrow{\sim} \alpha |u_{r+1}|\Lambda(c_{r+1}) \quad \text{and} \quad |u_{r+1}|\Lambda(c_{r+1}) \xrightarrow{\sim} \alpha |u_{r+2}|\Lambda(c_{r+2})$$


and h_r is zero on other terms. So that the cone $C(h_r)$ can be identified with four terms

$$C(h_r) \simeq \Lambda(\xi_r) \oplus \Lambda(\xi_{r+1}) \oplus \alpha \Lambda(\xi_{r+1}) \oplus \alpha \Lambda(\xi_{r+2})$$

If $\beta := |\xi_r|/|\xi_{r+1}| = t^{-4} q^{-2}$ then there is a degree zero isomorphism $i_r : C(h_r) \xrightarrow{\sim} \beta C(h_{r+1})$ which identifies all of the terms below pairwise,

$$i_r : \begin{bmatrix} \Lambda(\xi_r) \\ \Lambda(\xi_{r+1}) \\ \alpha \Lambda(\xi_{r+1}) \\ \alpha \Lambda(\xi_{r+2}) \end{bmatrix} \xrightarrow{\sim} \beta \begin{bmatrix} \Lambda(\xi_{r+1}) \\ \Lambda(\xi_{r+2}) \\ \alpha \Lambda(\xi_{r+2}) \\ \alpha \Lambda(\xi_{r+3}) \end{bmatrix}$$

To recap, we have a sequence of maps upon which we take cones. First $f : S^r(3_1) \rightarrow |u|S^r(3_1)$, $g : C(f) \rightarrow \mathcal{L}C(f)$, $h : C(g) \rightarrow \alpha \mathcal{L}C(g)$ and $i : C(h) \rightarrow \beta \mathcal{L}C(h)$. Finally, i is a degree zero isomorphism, $C(i) \cong 0$ in $H^0(\text{Seq})$. Here is the diagram,



The diagram in Ex. 4.9 is a kind of Koszul resolution. In §6 we will show that for each knot K there is a Koszul resolution of the sort pictured above which categorifies the Alexander A-polynomial \bar{A}_K from Thm. 2.9.

6. HOLONOMICITY FOR $S^r \widehat{CFK}(K)$

In this section we will construct a homological q -holonomic relation for each sequence $S := (S^r \widehat{CFK}(K))_{r \geq 0}$ for each knot K in S^3 . The arguments here generalize Ex. 4.8 and Ex. 4.9.

Recall Proposition 4.10, the decomposition is analogous to the observation that there is a bijection between the terms summed to produce the colored Alexander polynomials from Def. 2.3. Allowing r to vary shows that the ungraded chain complex remains constant. The maps ℓ_r introduced below make this idea more precise.

Corollary 6.1. *There are isomorphisms of ungraded chain complexes $\ell_r : S^r \widehat{CFK}(K) \rightarrow S^{r+1} \widehat{CFK}(K)$*

Proof. Set $\ell_r := (\sum_i r_i) + t$ in

$$\begin{array}{ccc}
 S^r \widehat{CFK}(K) & \xrightarrow{\sim} & (\oplus_i \Lambda(a_i^r)) \oplus \theta^r S^r \widehat{CFK}(U) \\
 \ell_r \downarrow & & \downarrow (\sum_i r_i) + t \\
 S^{r+1} \widehat{CFK}(K) & \xrightarrow{\sim} & (\oplus_i \Lambda(a_i^r)) \oplus \theta^{r+1} S^{r+1} \widehat{CFK}(U)
 \end{array}$$

The maps are determined by the following assignments, for $r_i : \Lambda(a_{i,r}) \rightarrow \Lambda(a_{i,r+1})$ set $r_i(1_{i,r}) := 1_{i,r+1}$ and $r_i(a_{i,r}) := a_{i,r+1}$, and for $t : \theta^r S^r \widehat{CFK}(U) \rightarrow \theta^{r+1} S^{r+1} \widehat{CFK}(U)$ set $t(u) := u$ and $t(\xi) := \xi$. \square

Let us now prove the reduced case (in analogy with Lemma 2.6), the head of the sequence in Prop. 4.10 can be systematically reduced with cones $C(\mathfrak{D}_i)$. The graded Euler characteristic of these cones agrees with the operators D_X from §2.2. The main result of this section, Thm. 6.6 is proven by reducing the general case, the sequence determined by Eq. (4.9), to the reduced case.

Definition 6.2. A sequence $S := (S^r)_{r \geq 0} \in \mathcal{Seq}$ will be called a *reduced sequence* when each term can be written as $S^r \cong \bigoplus_{i \in I_r} \Lambda(a_{i,r})$ in $H^0(\mathcal{Seq})$, with $|I_r| < \infty$ and the maps identifying adjacent terms

$$\ell_r(a_{i,r}) := a_{i,r+1} \quad \text{and} \quad \ell_r(1_{i,r}) := 1_{i,r+1} \quad (6.1)$$

are isomorphisms $\ell_r : S^r \xrightarrow{\sim} S^{r+1}$ of chain complexes.

Proposition 6.3. *For any such reduced sequence S , there are equivalence relations \equiv on the index sets I_r . For $i, j \in I_r$,*

$$i \equiv j \quad \text{when} \quad \frac{|\ell_r(a_{i,r})|}{|a_{i,r}|} = \frac{|\ell_r(a_{j,r})|}{|a_{j,r}|}$$

So for a reduced sequence S , we set $[I_r] := I_r / \equiv$, $|S_r| := |[I_r]|$ and $[\Lambda(a_{i,r})] := \bigoplus_{j \equiv i} \Lambda(a_{j,r})$ for the sum over corresponding congruence classes. It follows that each term S^r in the sequence S admits a factorization

$$S^r \cong \sum_{i \in [I_r]} [\Lambda(a_{i,r})]$$

into congruence classes of summands. The maps ℓ_r respect this decomposition in the sense that $\ell_r([\Lambda(a_{i,r})]) = [\Lambda(\ell_r(a_{i,r}))]$. In this sense, there is a map $\ell_r : I_r \rightarrow I_{r+1}$ given by $\ell_r(i) := i$.

Definition 6.4. A reduced sequence is *regular* when $i \equiv j$ if and only if $\ell_r(i) \equiv \ell_r(j)$.

Lemma 6.5. *A regular reduced sequence S is holonomic in the sense of Def. 5.5.*

Proof. By induction on the number of congruence classes $|S|$. If $|S| = 0$ then the sequence $S = \bar{0}$ is the constant sequence. Suppose that $|S| = n$ and pick $i \in [I_0]$ then the maps ℓ_r restrict to degree zero isomorphisms $f_{i,r} := \ell_r|_{[\Lambda(a_{i,r})]}$

$$f_{i,r} : [\Lambda(a_{i,r})] \rightarrow \alpha_i [\Lambda(\ell_r(a_{i,r}))] \quad \text{where} \quad \alpha_i := \frac{|\ell_r(a_{i,r})|}{|a_{i,r}|}$$

So there are degree zero chain maps $\mathfrak{D}_{i,r} : S^r \rightarrow \alpha_i \mathcal{L} S^r$

$$\mathfrak{D}_{i,r}(x) := \begin{cases} f_{i,r}(x) & x \in [\Lambda(a_{i,r})] \\ 0 & x \notin [\Lambda(a_{i,r})] \end{cases}$$

There is a map $\mathfrak{D}_i := (\mathfrak{D}_{i,r})_{r \geq 0}$ formed by combining the maps $\mathfrak{D}_{i,r}$. The cone $S_{new} := C(\mathfrak{D}_i)$ is homotopy equivalent to a sequence which excludes the i th component of S and doubles the others

$$C(\mathfrak{D}_{i,r}) \cong S_{\setminus i}^r \oplus \alpha_i S_{\setminus i}^r \quad \text{where} \quad S_{\setminus i}^r := \bigoplus_{j \in [I_r] \setminus \{i\}} [\Lambda(a_{j,r})]. \quad (6.2)$$

The proof is concluded by showing that the sequence $C(\mathfrak{D}_i)$ is a regular reduced sequence with $|C(\mathfrak{D}_i)| = |S| - 1$. This will occur in three steps. First we prove that the sequence is reduced. Second that the sequence is regular. Third we show that the number of congruence classes has decreased by one. The point is that the set of congruence classes of $S_{\setminus i}^r$ and $\alpha_i S_{\setminus i}^r$ are the same because α_i is just a degree shift.

(i) *The sequence $C(\mathfrak{D}_i)$ is reduced:* Eqn. (6.2) shows that each term

$$C(\mathfrak{D}_i)_r \cong \bigoplus_{i \neq j} (\Lambda(a_{j,i}) \oplus \alpha_i \Lambda(a_{j,i})) = \bigoplus_{j \in J_r} \Lambda(b_{j,i})$$

where $J_r := A_r \sqcup B_r$ where $A_r := I_r \setminus [i]$ and $B_r := I_r \setminus [i]$ and an element $j \in I \setminus [i]$ corresponds to either $\Lambda(b_{j,r}) := \Lambda(a_{j,r})$ when $j \in A_r$ and $\Lambda(b_{j,r}) := \alpha_i \Lambda(a_{j,r})$ when $j \in B_r$.

The maps $m_r : \bigoplus_{j \in J_r} \Lambda(b_{j,r}) \rightarrow \bigoplus_{j \in J_{r+1}} \Lambda(b_{j,r+1})$ determined by the assignments in Eqn. (6.1) above determine isomorphisms between adjacent terms in the new sequence because regularity of the sequence S shows ℓ_r respects the equivalence relation \equiv and induces isomorphisms between components. The maps $m_r = \ell_r|_{S^r / [\Lambda(a_{i,r})]} + \alpha_i \ell_r|_{S^r / [\Lambda(a_{i,r})]}$ are a block diagonal matrix with components given by the restriction of ℓ_r to the complement of the i th component and an α_i -shifted copy of this restriction.

(ii) *The sequence $C(\mathfrak{D}_i)$ is regular:* Since $(\alpha_i |\ell_r(a_{j,r})|) / (\alpha_i |a_{j,r}|) = |\ell_r(a_{j,r})| / |a_{j,r}|$ the fold map $\varphi : J_r \rightarrow I_r \setminus [i]$ is 2-to-1 and induces a bijection $J_r / \equiv \xrightarrow{\sim} (I_r \setminus [i]) / \equiv$. Since S^r is regular, $j, k \in I_r \setminus [i]$ implies $j \equiv k \Leftrightarrow \ell_r(j) \equiv \ell_r(k)$. Since $m_r = \ell_r$ on equivalence classes under the identification φ , $C(\mathfrak{D}_r)$ is regular as well.

(iii) *The number of congruence classes has decreased:* Observe that $|C(\mathfrak{D}_r)| = |J_r / \equiv| =_{\varphi} |(I_r / \equiv) \setminus [i]| = |I_r / \equiv| - 1$.

□

Theorem 6.6. *If K is a knot then the sequence $S^r := (S^r \widehat{CFK}(K))_{r \geq 0} \in Seq$ is holonomic.*

Proof. Recall from Prop. 4.10 that

$$S^r \widehat{CFK}(K) \cong Hd \oplus Tl$$

where $Hd := (\Lambda(a_{1,r}) \oplus \Lambda(a_{2,r}) \oplus \cdots \oplus \Lambda(a_{n,r}))$ and $Tl := \theta^r S^r \widehat{CFK}(U)$.

There is a map $f_r : S^r \widehat{CFK}(K) \rightarrow |u| S^r \widehat{CFK}(K)$ which takes $f|_{Hd} = 0$ and on the tail $f(u^n \xi^k) := u^{n-1} \xi^k$. As in Ex. 4.9, $C(f_r) \simeq T \oplus |u| T \oplus \Lambda(\xi)$. The maps f_r

combine to form a map $f : S^r \widehat{CFK}(K) \rightarrow mS^r \widehat{CFK}(K)$. We claim that the sequence of cones $C(f) \in Seq$ is regular and reduced.

Set $I_r := \{1, \dots, n, |u|1, \dots, |u|n, \xi\}$. It follows from Prop. 4.10 that

$$C(f_r) \cong Hd \oplus |u|Hd \oplus \Lambda(\xi) = \sum_{i \in I_r} \Lambda(c_{i,r})$$

where

$$\Lambda(c_{i,r}) := \begin{cases} \Lambda(a_{i,r}) & i \in \{1, \dots, n\} \\ |u|\Lambda(a_{i,r}) & i \in \{|u|1, \dots, |u|n\} \\ \Lambda(\xi) & i \in \{\xi\} \end{cases}$$

this is the same complex for each $r \geq 1$, so the maps $\ell_r : C(f_r) \rightarrow C(f_{r+1})$ given by the assignments Eqn. (6.1) are isomorphisms. It follows that the sequence $C(f)$ is reduced.

The sequence $C(f)$ is regular because, for any $i \in I_r$, the bidegrees of $|1_{i,r}|$ are fixed affine linear functions, which implies that the quotients $|\ell_r(1_{i,r})|/|1_{i,r}|$ are constant functions of r .

The result now follows from Lemma 6.5. \square

Remark 6.7. The theorem shows that holonomicity corresponds to a Koszul-type complex which is determined by the immersed curve representing the module $\widehat{CFD}(K)$.

7. NOTATION

\equiv	Prop. 6.3	$\bar{\Delta}_K^r(q)$	unreduced r -colored polynomial, Def. 2.3
\circ	Notation 3.18	$\bar{\Delta}_U^r(q)$	r -colored unknot, Eqn. (2.6)
$ x _t$	homological degree	$\Delta_K^r(q)$	reduced r -colored polynomial, Eqn. (2.7)
$ x _q$	graded degree	$a_{i,r}$	Prop. 4.10
$ x _{t,q}$	$ x _{t,q} = (x _t, x _q)$	\mathcal{A}	Weyl algebra §2.2
$ x $	multiplicative bidegree, if $ x _{t,q} = (n, m)$ then $ x = t^n q^m$	\mathcal{A}^+	Eqn. (5.4)
$1_{i,r}$	Prop. 4.10	A_K	A -polynomial, Lem. 2.6
3_1	right-handed trefoil	\bar{A}_K	unreduced A -polynomial, Thm. 2.9
θ	Prop. 4.10	$\mathcal{A}_{\mathbb{Z}}$	category of graded object, §5.1
Δ_n	Ex. 4.2	b	bottom generator, Not. 3.12
$\Delta_K(q)$	Alexander's balanced Laurent polynomial	χ_0	distinguished generator, Not. 3.12
$\Delta_K^1(q)$	positive form of Alexander polynomial	\bar{C}	Ex. 5.7
		$\hat{C}(K), C^-(K)$	Eqn. (3.2)

$CF(\gamma, \eta)$	Thm. 3.7	q^m	monomial or graded shift functor
$Ch(\mathcal{A})$	category of chain complexes in \mathcal{A} , Def. 5.1	$q^m = \cdot\{-m\}$	
D_X	difference operator, §2.2	Seq	Def. 2.4
\mathfrak{D}	Lem. 6.5	Seq(T)	Def. 2.4
$F^*\mathcal{A}$	category of filtered objects, §5.1	Seq	$Seq = Seq(F^*Ch_{\mathbb{Z}})$, Eqn. (5.3)
$\gamma_K, \bar{\gamma}_K$	§3.1	$Seq(\mathcal{A})$	sequences in the dg category \mathcal{A}
$\frac{1}{2}\bar{\gamma}_K$	Cor. 3.15	$\tau(K)$	Tau invariant Rem. 3.6
$g(K)$	Seifert genus Rem. 3.6	t^n	monomial or homological shift functor $t^n = \cdot[-n]$
$\bar{\gamma}_0$	Rmk. 3.5 (1)	$t^n q^m$	monomial or shift functor
H^0	homotopy category, §5.1	T_K, \bar{T}_K	§3.1
Hd	Def. 4.3	Tl	Def. 4.3
\widehat{HFK}	$H_*(\widehat{CFK}(K))$, knot Floer homology (balanced grading)	\widehat{CFD}	Eqn. (3.1)
$K_{r,s}$	Def. 2.1	$S^r \widehat{CFK}(K)$	S^r -colored knot Floer associated graded chain complex
ℓ_r	Cor. 6.1	$S^r \widehat{CFK}(K)$	S^r -colored knot Floer filtered graded chain complex
\mathcal{L}	Prop. 5.3	$S^r \widehat{HFK}$	$H_*(S^r \widehat{CFK}(K))$
m	Prop. 5.3	U	the unknot
νK	regular neighborhood of knot	$\mathbb{Z}_{\geq 0}$	$\mathbb{Z}_{\geq 0} := \{0, 1, 2, \dots\} \subset \mathbb{Z}$

REFERENCES

- [Abo08] Mohammed Abouzaid, *On the Fukaya categories of higher genus surfaces*, Adv. Math. **217** (2008), no. 3, 1192–1235. MR 2383898
- [BK90] A. I. Bondal and M. M. Kapranov, *Framed triangulated categories*, Mat. Sb. **181** (1990), no. 5, 669–683. MR 1055981
- [BPLHW22] Anna Beliakova, Krzysztof Putyra, Robert Louis-Hadrien, and Emmanuel Wagner, *A proof of dunfield-gukov-rasmussen conjecture*, arXiv:2210.00878 (2022).
- [Bri05] Michel Brion, *Lectures on the geometry of flag varieties*, Topics in cohomological studies of algebraic varieties, Trends Math., Birkhäuser, Basel, 2005, pp. 33–85. MR 2143072
- [Cau17] Sabin Cautis, *Remarks on coloured triply graded link invariants*, Algebr. Geom. Topol. **17** (2017), no. 6, 3811–3836. MR 3709661
- [CCG⁺94] D. Cooper, M. Culler, H. Gillet, D. D. Long, and P. B. Shalen, *Plane curves associated to character varieties of 3-manifolds*, Invent. Math. **118** (1994), no. 1, 47–84. MR 1288467
- [DGR06] Nathan M. Dunfield, Sergei Gukov, and Jacob Rasmussen, *The superpolynomial for knot homologies*, Experiment. Math. **15** (2006), no. 2, 129–159. MR 2253002
- [FG00] Charles Frohman and Răzvan Gelca, *Skein modules and the noncommutative torus*, Trans. Amer. Math. Soc. **352** (2000), no. 10, 4877–4888. MR 1675190
- [FGL02] Charles Frohman, Răzvan Gelca, and Walter Lofaro, *The A-polynomial from the noncommutative viewpoint*, Trans. Amer. Math. Soc. **354** (2002), no. 2, 735–747. MR 1862565

- [FGSk13] Hiroyuki Fuji, Sergei Gukov, and Piotr Sułkowski, *Super- A -polynomial for knots and BPS states*, Nuclear Phys. B **867** (2013), no. 2, 506–546. MR 2992793
- [Gar04] Stavros Garoufalidis, *On the characteristic and deformation varieties of a knot*, Proceedings of the Casson Fest, Geom. Topol. Monogr., vol. 7, Geom. Topol. Publ., Coventry, 2004, pp. 291–309. MR 2172488
- [Gar08] ———, *Difference and differential equations for the colored Jones function*, J. Knot Theory Ramifications **17** (2008), no. 4, 495–510. MR 2414452
- [GL05] Stavros Garoufalidis and Thang T. Q. Lê, *The colored Jones function is q -holonomic*, Geom. Topol. **9** (2005), 1253–1293. MR 2174266
- [GL16] ———, *A survey of q -holonomic functions*, Enseign. Math. **62** (2016), no. 3-4, 501–525. MR 3692896
- [GLL18] Stavros Garoufalidis, Aaron D. Lauda, and Thang T. Q. Lê, *The colored HOMFLYPT function is q -holonomic*, Duke Math. J. **167** (2018), no. 3, 397–447. MR 3761103
- [Guk05] Sergei Gukov, *Three-dimensional quantum gravity, Chern-Simons theory, and the A -polynomial*, Comm. Math. Phys. **255** (2005), no. 3, 577–627. MR 2134725
- [Han23a] Jonathan Hanselman, *Heegaard Floer homology and cosmetic surgeries in S^3* , J. Eur. Math. Soc. (JEMS) **25** (2023), no. 5, 1627–1669. MR 4592856
- [Han23b] ———, *Knot Floer homology as immersed curves*, <https://arxiv.org/abs/2305.16271> (2023).
- [HKK17] F. Haiden, L. Katzarkov, and M. Kontsevich, *Flat surfaces and stability structures*, Publ. Math. Inst. Hautes Études Sci. **126** (2017), 247–318. MR 3735868
- [Hog18] Matthew Hogancamp, *Categorified Young symmetrizers and stable homology of torus links*, Geom. Topol. **22** (2018), no. 5, 2943–3002. MR 3811775
- [Hom20] Jennifer Hom, *Lecture notes on heegaard floer homology*, <https://arxiv.org/abs/2008.01836> (2020).
- [HRW22] Jonathan Hanselman, Jacob Rasmussen, and Liam Watson, *Heegaard Floer homology for manifolds with torus boundary: properties and examples*, Proc. Lond. Math. Soc. (3) **125** (2022), no. 4, 879–967. MR 4500201
- [HRW24] ———, *Bordered Floer homology for manifolds with torus boundary via immersed curves*, J. Amer. Math. Soc. **37** (2024), no. 2, 391–498. MR 4695506
- [HW23a] Jonathan Hanselman and Liam Watson, *Cabling in terms of immersed curves*, Geom. Topol. **27** (2023), no. 3, 925–952. MR 4599309
- [HW23b] ———, *A calculus for bordered Floer homology*, Geom. Topol. **27** (2023), no. 3, 823–924. MR 4599308
- [KMS09] Mikhail Khovanov, Volodymyr Mazorchuk, and Catharina Stroppel, *A brief review of abelian categorifications*, Theory Appl. Categ. **22** (2009), No. 19, 479–508. MR 2559652
- [KR08] Mikhail Khovanov and Lev Rozansky, *Matrix factorizations and link homology*, Fund. Math. **199** (2008), no. 1, 1–91. MR 2391017
- [Lô6] Thang T. Q. Lê, *The colored Jones polynomial and the A -polynomial of knots*, Adv. Math. **207** (2006), no. 2, 782–804. MR 2271986
- [Lic97] W. B. Raymond Lickorish, *An introduction to knot theory*, Graduate Texts in Mathematics, vol. 175, Springer-Verlag, New York, 1997. MR 1472978
- [LOT18] Robert Lipshitz, Peter S. Ozsvath, and Dylan P. Thurston, *Bordered Heegaard Floer homology*, Mem. Amer. Math. Soc. **254** (2018), no. 1216, viii+279. MR 3827056
- [OS04a] Peter Ozsváth and Zoltán Szabó, *Holomorphic disks and knot invariants*, Adv. Math. **186** (2004), no. 1, 58–116. MR 2065507

- [OS04b] Peter Ozsváth and Zoltán Szabó, *Holomorphic disks and three-manifold invariants: properties and applications*, Ann. of Math. (2) **159** (2004), no. 3, 1159–1245. MR 2113020
- [OS04c] ———, *Holomorphic disks and topological invariants for closed three-manifolds*, Ann. of Math. (2) **159** (2004), no. 3, 1027–1158. MR 2113019
- [Ras03a] Jacob Rasmussen, *Floer homology and knot complements*, PhD thesis, Harvard University (2003).
- [Ras03b] Jacob Andrew Rasmussen, *Floer homology and knot complements*, ProQuest LLC, Ann Arbor, MI, 2003, Thesis (Ph.D.)—Harvard University. MR 2704683
- [Ras15] Jacob Rasmussen, *Some differentials on Khovanov-Rozansky homology*, Geom. Topol. **19** (2015), no. 6, 3031–3104. MR 3447099
- [Roz14] Lev Rozansky, *An infinite torus braid yields a categorified Jones-Wenzl projector*, Fund. Math. **225** (2014), no. 1, 305–326. MR 3205575
- [Sab93] Claude Sabbah, *Systèmes holonomes d'équations aux q -différences*, D -modules and microlocal geometry (Lisbon, 1990), de Gruyter, Berlin, 1993, pp. 125–147. MR 1206016
- [Sar15] Antonio Sartori, *The Alexander polynomial as quantum invariant of links*, Ark. Mat. **53** (2015), no. 1, 177–202. MR 3319619
- [Vir06] O. Ya. Viro, *Quantum relatives of the Alexander polynomial*, Algebra i Analiz **18** (2006), no. 3, 63–157. MR 2255851
- [Web17] Ben Webster, *Knot invariants and higher representation theory*, Mem. Amer. Math. Soc. **250** (2017), no. 1191, v+141. MR 3709726
- [Wei13] Charles A. Weibel, *The K -book*, Graduate Studies in Mathematics, vol. 145, American Mathematical Society, Providence, RI, 2013, An introduction to algebraic K -theory. MR 3076731

UNIVERSITY OF IOWA, DEPARTMENT OF MATHEMATICS, 14 MACLEAN HALL, IOWA CITY, IA 52242-1419 USA

Email address: ben-cooper@uiowa.edu

Email address: robert-deyeso@uiowa.edu