

Gravitational Wave Mountains: *current-carrying domain walls*

Anish Ghoshal^{1,*} and Yu Hamada^{2,3,†}

¹*Institute of Theoretical Physics, Faculty of Physics,
University of Warsaw, ul. Pasteura 5, 02-093 Warsaw, Poland*

²*Deutsches Elektronen-Synchrotron DESY, Notkestr. 85, 22607 Hamburg, Germany*

³*Research and Education Center for Natural Sciences, Keio University,
4-1-1 Hiyoshi, Yokohama, Kanagawa 223-8521, Japan*

Domain wall (DW) networks may have formed in the early universe following the spontaneous breaking of a discrete symmetry. Notably, several particle physics models predict the existence of current-carrying DWs, which can capture and store particles as zero modes on it. In this study, we demonstrate that gravitational waves (GWs) generated by current-carrying DWs with fermionic zero modes exhibit a novel feature: an additional peak in the GW spectrum resembling mountains, arising from metastable topological remnants, which we term “spherons.” This distinct signature could be detectable in upcoming GW observatories such as LISA and ET. The results suggest that DW networks in beyond Standard Model scenarios could emit GW signals that are significantly stronger and with greater detectability than previously expected.

Introduction— Domain walls (DWs) are typical topological defects that are formed due to cosmological phase transitions in the early universe when for instance two (nearly) degenerate vacua are present. Just after the formation of the DW network, they evolve in what is known as the scaling regime. During this time the correlation length of the network is approximately same as that of the Hubble horizon size $L \sim t$ [1–3]. The fraction of the total universe energy budget stored in the DWs increases linearly with time $\rho_{\text{DW}}/\rho_{\text{tot}} \propto t$, which can then easily dominate the total energy density at later stage leading to inconsistency with current cosmological observations; this is known as the domain-wall problem. One may avoid this scenario if there is an energy bias V_{bias} between the different vacua. Such a bias provides pressure between vacua, which drives the DW network to collapse.¹ As reviewed below, the collapsing DW network radiates gravitational wave (GW) with significant amplitude, which is expected to be observed by GW experiments. The DW evolution in the early universe has been widely studied numerically [17–25] as well as analytically [1–3, 26–31].

Generically topological defects can have rich internal structure which has the ability to carry some sort of charge without dissipation. In the case of cosmic strings [32], a current carried on the strings can prevent the string loop from collapsing, leading to a stable rotating loop (called vorton [33]). These loops might reach equilibrium configurations due to balance between

the string tension and the centrifugal force [33–35] whose classical and quantum stability has been discussed [36–45]. Besides the vortons, the cosmological and astrophysical impacts of current-carrying strings have been also discussed [46–59].

Similarly, a DW can carry current [60–64], whose internal degrees of freedom are classified into two cases: fermionic current carrier arising when sign of a fermion mass differs on both sides of the DW [60–62] and bosonic carrier arising when some $U(1)$ symmetry is broken only inside the DW [32, 63]. If DWs are current-carrying, one may expect that such a current stabilizes a closed DW by balance between the DW tension and the centrifugal force analogously to vortons. This stabilized object can be long-lived in the universe, and might have some cosmological impact in addition to standard DW networks as we see below.

BSM Model involving DW with currents— The concept of the current-carrying DW is quite common. One simple beyond Standard Model (BSM) example for the fermionic carriers is DWs in two-Higgs doublet model (2HDM) [65–71], where the Higgs potential consisting of two Higgs doublets H_1, H_2 has a \mathbb{Z}_2 symmetry $H_1 \rightarrow H_1, H_2 \rightarrow -H_2$ with a tiny bias term proportional to $H_1^\dagger H_2 + \text{h.c.}$. This \mathbb{Z}_2 symmetry is spontaneously broken in the vacuum, giving rise to a DW. Depending on the type of the Yukawa couplings [72], the SM fermions feel flipped mass terms when getting across the DWs,² and hence carry the SM gauge (and baryon/lepton number) current.

¹ It could happen that after some time t_{ann} is elapsed, the vacuum energy difference V_{bias} between the two degenerate vacua counterbalances the pressure due to the domain wall surface tension σ . This may lead DWs towards each other and annihilate before they can dominate the universe at a time t_{dom} [4–7]. Closed DWs at this phase shrink and under specific conditions, may enter within their Schwarzschild radius and form PBHs [8–14], a process known in the literature as “catastrogenesis” [15, 16].

² Depending on the parameters, this model can have DWs that break $U(1)$ electromagnetic symmetry by the condensation of the charged bosonic fields [67–70]. In such a case, the bosonic particles also can play roles of the charge/current carriers on the DWs. In this work, however, we do not consider those cases but specify ourselves to fermionic carriers.

Although the 2HDM is quite simple, the tension of the DW cannot be beyond about 10^6 GeV^3 since it is related to the electroweak scale. This can be made more general by adding a real SM-singlet scalar S to 2HDM [73–75] whose VEV is quite general, and imposing a \mathbb{Z}_2 symmetry $H_1 \rightarrow H_1, H_2 \rightarrow -H_2, S \rightarrow -S$.³ Again, depending on the couplings to the SM fermions, the DW can contain charge/current carriers. The DW tension is dominantly controlled by the VEV of S .

Also one can find a current-carrying DW in $SO(10)$ GUT as shown in Ref. [77]. In this case, one can not introduce explicit breaking terms for the discrete symmetry since it is a subgroup of the $SO(10)$ gauge symmetry, resulting in that the decay mechanism of the DWs must rely on nucleation of cosmic string loops [71, 78–83]. Nevertheless, we specify ourselves to cases with bias terms instead of the nucleation throughout this work.

Domain Walls and GW — It is known that after the production of the DWs, they form a scaling network [1–3], in which the number of the DWs remains about $\mathcal{O}(1)$ per Hubble patch. As such a DW network causes a serious cosmological problem known as DW domination, it should decay by some mechanism. One possible way is to introduce in the Lagrangian a tiny bias term breaking the discrete symmetry slightly, leading to pressure difference ΔV between different vacua (domains) separated by the DWs. The DW network collapses when the pressure is comparable to the DW surface energy at $t \simeq t_{\text{ann}}$, satisfying

$$\Delta V d_H(t_{\text{ann}})^3 \simeq \sigma d_H(t_{\text{ann}})^2 \quad \therefore t_{\text{ann}} \simeq \frac{\sigma}{\Delta V}, \quad (1)$$

where d_H is the Hubble length and σ is the DW tension. As this must occur before the DW domination era, which starts at $t_{\text{dom}}^{-1} \simeq G\sigma$, we have a necessary condition

$$\Delta V > G\sigma^2. \quad (2)$$

with G the Newton constant.

The DW network radiates GW when collapsing. One may use for a rough estimate the quadrupole formula of the GW emission rate, from which one can get the radiation rate \dot{E}_{GW} as

$$\dot{E}_{\text{GW}} \simeq -G\sigma^2 R(t)^2, \quad (3)$$

where $R(t)$ is the typical radius of the shrinking DW. Here we have used that typical oscillation frequency of the DW is given as $1/R(t)$. This formula holds for point-like object observed at infinity. Nevertheless, this gives a nice approximation of the GW amplitude in most cases [18].

³ DW solutions in this model with a different \mathbb{Z}_2 symmetry is also possible [76].

The radiated GW from DWs might be observed as stochastic GW background at present universe. It is convenient to consider the GW energy spectrum defined as

$$\Omega_{\text{GW}} \equiv \frac{1}{\rho_c} \frac{d\rho_{\text{GW}}(f)}{d \log f} \quad (4)$$

where ρ_{GW} and ρ_c are the GW energy density and critical energy density, respectively.

Since the typical length scale of the collapsing network is given by the Hubble size, $R(t_{\text{ann}}) \sim t_{\text{ann}}$, the emitted GW spectrum has a peak around $1/t_{\text{ann}}$. Away from the peak, it is well approximated by power-law tails as f^3 and f^{-1} in IR and UV regimes, respectively [18, 22]. Here the IR one is deduced from the causality argument. After the emission, the GW spectrum is red-shifted and observed today as

$$\Omega_{\text{GW},0}(f) = \Omega_{\text{GW,max}} \begin{cases} \frac{f^3}{f_{\text{peak}}^3} & f \ll f_{\text{peak}} \\ \frac{f_{\text{peak}}}{f} & f \gg f_{\text{peak}} \end{cases} \quad (5)$$

with

$$\Omega_{\text{GW,max}} = 2 \times 10^{-3} (G\sigma t_{\text{ann}})^2 \left(\frac{g_{*0}}{g_*(t_{\text{ann}})} \right)^{\frac{1}{3}} \quad (6)$$

$$f_{\text{peak}} = \frac{1}{t_{\text{ann}}} \frac{a(t_{\text{ann}})}{a(t_0)}. \quad (7)$$

Notice that some results that deviate from the UV spectral index -1 have been reported in the literature. For instance, it is estimated to be -1.7 in Ref. [23] and -1.5 in Ref. [25]. While they do not affect the GW peak amplitude significantly, the GW signal in the UV regime may be slightly suppressed.

Current-carrying Domain Walls— When DWs couple to other particles, the DWs may gain internal degrees of freedom by capturing them. Let us consider a DW consisting of a real scalar ϕ coupled to a single Dirac fermion ψ with a Yukawa coupling y . The simplest and self-contained setup is given by the following Lagrangian:

$$\mathcal{L} = \frac{1}{2}(\partial_\mu \phi)^2 - \lambda(\phi^2 - v_\phi^2)^2 + \bar{\psi} [i\cancel{\partial} - y\phi] \psi, \quad (8)$$

where the potential term forces ϕ to take vacuum expectation value v_ϕ , leading to the DW configuration. We here solve the Dirac equation for ψ in the presence of the DW configuration ϕ_{DW} located at $x = 0$:

$$\phi_{\text{DW}} = v_\phi \tanh\left(\frac{m_\phi x}{2}\right) \quad (9)$$

with m_ϕ being the mass of ϕ particle. Ignoring the y, z and t directions reduces to the equation for the x dependence,

$$i\gamma^1 \partial_x \psi = y\phi_{\text{DW}} \psi. \quad (10)$$

Since $i\gamma^1$ is hermitian, it is always possible to label the solution ψ in terms of two eigenstates corresponding to $i\gamma^1 = \pm 1$. For $i\gamma^1 = -1$ space, one gets a solution

$$\psi(x) \propto \varphi \exp\left(-\int_0^x dx' \phi_{\text{DW}}(x')\right), \quad (11)$$

with φ a spinor satisfying $i\gamma^1\varphi = -\varphi$, from which one can see that this solution is localized at $x = 0$ and decays as $|x| \rightarrow \infty$. The solutions dependent on y, z and t are easily obtained by performing the Lorentz boost in the y - or z -directions on ψ . Note that these solutions are massless modes, namely, they behave as massless particles propagating only on the DW (y and z directions). If ψ has a charge, which can be either a gauge (e.g., electric) or global (e.g., baryon number) charge, the trapped mode induces current and charge, leading to the current-carrying DW.

When the DW forms a closed sphere with a radius R , the fermion solution is localized around the sphere surface and the degrees of freedom in the direction along the sphere is labeled by the orbital angular momentum j ($= 1/2, 3/2, \dots$) [84–86]. The z component of the angular momentum j_z can be taken as $j_z = -j, -j+1, \dots, j$, which means that there are $2j+1$ degenerated states for the fixed j . A state with the angular momentum j has an energy $E \simeq (j + \frac{1}{2})/R$ as long as $E \ll m_\psi$.

During the time evolution of the DW network, the network continuously produces closed spheres made of the DW by reconnection. However, it has been observed in Ref. [25] that most closed DWs are produced during the network collapsing, $t \simeq t_{\text{ann}}$. Therefore we focus on the latter case, however the former ones may also give additional contributions to our analysis given below.

After produced, the closed DWs shrink due to the DW tension and may capture particles in the bulk. This is a crucial mechanism for the DW to get current/charge. We here introduce a yield Y for the would-be trapped particles ψ as

$$Y \equiv \frac{n_\psi}{s} \quad (12)$$

with n_ψ and s being the number density of ψ and the entropy density in the universe, respectively. The total charge captured by the single DW sphere with the radius $R(t)$ is roughly estimated as

$$Q(R) \simeq \frac{4\pi Y s}{3} (R_0^3 - R(t)^3), \quad (13)$$

where R_0 is the radius of the produced closed sphere, i.e., R_0 is around the network size $R_0 \sim t_{\text{ann}}$.⁴ One should

note that the total current J of the captured particles is negligible due to the cancellation among the trapped particles. Here we assumed the capture rate of the particles to be $\mathcal{O}(1)$ and the trapped particles ψ to be maximally asymmetric, namely, without antiparticles to avoid pair annihilation on the DW. This point will be discussed below.

Notably the captured particles contributes to the DW sphere energy M_{DW} . Denoting the highest energy level of the trapped particles as j_{max} , (i.e., the highest energy is $\sim j_{\text{max}}/R$), the total occupation number ($\sim Q$) is

$$\sum_{j=\frac{1}{2}, \frac{3}{2}, \dots}^{j_{\text{max}}} (2j+1) \sim j_{\text{max}}^2, \quad \therefore j_{\text{max}} \sim \sqrt{Q}. \quad (14)$$

The summation of the energy over j leads to

$$\sum_j \sum_{j_z=-j}^j E(j) \sim \frac{j_{\text{max}}^3}{R} \sim \frac{Q^{\frac{3}{2}}}{R}, \quad (15)$$

resulting in

$$M_{\text{DW}} \sim 4\pi\sigma R(t)^2 + \frac{Q^{\frac{3}{2}}}{R(t)} + \frac{4\pi}{3} R(t)^3 \Delta V \quad (16)$$

$$\equiv 4\pi\sigma_{\text{eff}} R(t)^2, \quad (17)$$

where we have defined the effective tension of the DW:

$$\sigma_{\text{eff}} \equiv \sigma + \frac{Q^{\frac{3}{2}}}{4\pi R(t)^3} + \frac{1}{3} R(t) \Delta V. \quad (18)$$

Note that the Q -dependence is different from that for vorton energy ($\propto Q^2$) [33].

In order to have sufficient charge/current on DW, we have two conditions. The first condition: m_ψ should not be much smaller than the kinetic energy of ψ in the bulk. This comes from the fact that the typical scale of the trapping potential that ψ feels is m_ψ . Thus ψ cannot be trapped on the DW but be transmitted or reflected when kinetic energy of the injecting particle is larger than m_ψ . Due to this condition, one finds that ψ cannot be relativistic in the thermal bath since relativistic particles have kinetic energy T ($> m_\psi$). Given this, in order to have the enough number density (or Y) of ψ , they should be decoupled from the thermal equilibrium like dark matter. Especially one possible example is asymmetric dark matter, in which the dark matter abundance in the bulk is dominated by the asymmetric part, leading to the natural suppression of pair annihilation of the trapped charge on the DW. Even without such an asymmetry in the bulk, it is possible to consider the DW with a significant CP violation, which captures particles and antiparticles with different probabilities, resulting in asymmetry only on the DW. See, e.g., Ref. [70] for CP-violating DW in 2HDM. When there is no asymmetry on the DW, still one could have enough charges due to statistical fluctuation.

⁴ It is not trivial whether the network of the current-carrying DW exhibits the scaling behavior. Although we might have deviation from the scaling, we assume such effects not to be significant for our results below. See also Discussion and conclusions.

The second condition: the capture process must be energetically favored compared to transmission or reflection. This is because otherwise the capture process is not favored to happen, leading to termination of the capture process. When the DW sphere radius varies from R to $R - \Delta R$, this condition is given as

$$\left| -\frac{Q^{\frac{3}{2}}}{R^2} + \frac{\partial M_{\text{DW}}}{\partial Q} \frac{\partial Q}{\partial R} \right| \Delta R \leq 4\pi n_\psi m_\psi R^2 \Delta R \quad (19)$$

where the L.H.S. indicates the increase of the energy due to the capture while the R.H.S. is the energy of the free particles in the case that the capture does not happen. For R to be close to R_0 , the two terms in lhs vanish, resulting in that the inequality is satisfied. This means that the capture is significant at least just after the DW starts to shrink, and this stage gives a dominant amount of the captured particles. This allows us to assume the $\mathcal{O}(1)$ capture rate as the first study.

Subsequent GW from metastable object – As stated above, the shrinking DWs can gain significant amount of charge when they shrink. This charge is enough to prevent them from shrinking due to repulsion caused by the charge, which corresponds to the second term in (16). Such an object is similar to vortons [33, 87], which are made of charge/current-carrying string loop. The stabilized radius R_{sph} is determined by the condition

$$\left. \frac{d}{dR} M_{\text{DW}} \right|_{R=R_{\text{sph}}} = 0. \quad (20)$$

At $R = R_{\text{sph}}$, the DW tension and the repulsive force of the charge are balanced, forming a metastable spherical object,⁵ which we dub *spheron*. As stated above, the spherons are assumed to be produced during the network collapse, $t = t_{\text{ann}}$ or equivalently $T = T_{\text{ann}} (\simeq$

$\sqrt{M_{\text{pl}}/t_{\text{ann}}}$). Then one may get

$$R_{\text{sph}} \sim \frac{\sqrt{Q}}{\sigma^{\frac{1}{3}}} \sim \frac{R_0 \sqrt{Y M_{\text{pl}} T_{\text{ann}}}}{\sigma^{\frac{1}{3}}}, \quad (21)$$

where we have assumed $R_0 \gg R_{\text{sph}}$ and ignored the bias pressure ΔV in M_{DW} .

We here provide a necessary condition for the trapped mode not to escape into the bulk. Using the expression of $Q(R_{\text{sph}})$ (13) and R_{sph} (21), the highest energy of the trapped particle is given as

$$E_{\text{max}} \sim \frac{j_{\text{max}}}{R_{\text{sph}}} \sim (8\pi\sigma)^{\frac{1}{3}}, \quad (22)$$

which must be smaller than the bulk mass m_ψ to prevent from escaping into the bulk. Notice that 2HDM does not contain sufficiently heavy fermions and requires some extension in the matter sector.

While spherons are classically stable after stabilized, they can decay through either quantum decay of charge-carrier particles like the case of vortons [44, 45, 52] or fission [88] by non-perturbative tunneling like Q -balls [89].⁶ One should be able to calculate this lifetime t_{dec} in principle once the model is fixed. Nevertheless, since it is highly model-dependent and beyond the scope of this work, we keep it as a free parameter and focus on the cosmological impacts. In the both decay processes, spherons lose their charge and start to shrink again at $t = t_{\text{dec}}$, and hence again oscillate asymmetrically with typical frequency $1/R_{\text{sph}}$. This leads to secondary GW radiation and its rate is estimated by simply replacing the size of DW in Eq. (3) with the size of the shrinking spheron $\dot{E}_{\text{GW}} \simeq -G\sigma_{\text{eff}}^2 R_{\text{sph}}^2$, where the spheron radius R_{sph} is given by (21).

The GW spectrum radiated from the spherons are calculated in a similar way to those of conventional DW networks. However, one should note that this is radiated at $t = t_{\text{dec}}$, which results in the different peak frequency. Thus we get the GW spectrum as

⁵ They would not be spherically symmetric if they had significant current. Nevertheless, as stated above, they cannot get significant current from the bulk particles. Thus the spherically symmetric configuration is energetically favored.

⁶ Indeed, the spheron cannot be stable against fission because

$\partial^2 M_{\text{DW}}(R_{\text{sph}})/\partial Q^2 > 0$ due to the term proportional to ΔV . Thus quantum tunneling allows it to decay into smaller spherons whose total charge is equal to the initial one. On the other hand, this fission is forbidden in classical processes as it requires a large deformation from the spherical shape for an intermediate configuration, requiring large energy cost.

$$\Omega_{\text{GW,sph}} \simeq \frac{8\pi\tau}{3H_0^2} (G\sigma_{\text{eff}}(R_{\text{sph}})R_{\text{sph}})^2 \frac{1}{t_{\text{ann}}^3} \left(\frac{a(t_{\text{ann}})}{a(t_{\text{dec}})}\right)^3 \left(\frac{a(t_{\text{dec}})}{a(t_0)}\right)^4 \left(\frac{g_{*0}}{g_*(t_{\text{dec}})}\right)^{\frac{1}{3}} \times \begin{cases} \left(\frac{f}{f_{\text{peak,sph}}}\right)^3 & f \ll f_{\text{peak,sph}} \\ \frac{f_{\text{peak,sph}}}{f} & f \gg f_{\text{peak,sph}} \end{cases} \quad (23)$$

with $f_{\text{peak,sph}} \equiv R_{\text{sph}}^{-1} a(t_{\text{dec}})/a(t_0)$, where τ is the time duration of the radiation process and we have assumed that the spherons remain stable until the sudden decay at $t = t_{\text{dec}}$, giving the number density of the spheron $t_{\text{ann}}^{-3} (a(t_{\text{ann}})/a(t_{\text{dec}}))^3$. Here we simply took the same spectral index as those for the conventional DW net-

work [18, 22]. Since the spheron radius typically decreases as $R_{\text{sph}} \propto \exp(-t/t_{\text{dec}})$ during the decay, τ may be given as $\tau \sim t_{\text{dec}}$ while the radiated GW energy must not exceed the whole energy of the spheron, resulting in $\tau \equiv \min [t_{\text{dec}}, \sigma_{\text{eff}} R_{\text{sph}}^2 / |\dot{E}_{\text{GW}}|]$.

Using Eq. (21), one may rewrite the peak frequency and its maximum value as

$$\Omega_{\text{GW,sph}}|_{\text{max}} \simeq 1 \times 10^{-7} \left(\frac{Y}{10^{-10}}\right) \left(\frac{\sigma}{10^{18} \text{ GeV}^3}\right)^{\frac{4}{3}} \left(\frac{100 \text{ MeV}}{T_{\text{dec}}}\right)^3 \min \left[1, \frac{T_{\text{dec}}^2 M_{\text{Pl}}}{\sigma}\right], \quad (24)$$

$$f_{\text{peak,sph}} \simeq 3 \times 10^{-6} \text{ Hz} \left(\frac{10^{-10}}{Y}\right)^{\frac{1}{2}} \left(\frac{\Delta V}{10^{-2} \text{ GeV}^4}\right)^{\frac{3}{4}} \left(\frac{10^{18} \text{ GeV}^3}{\sigma}\right)^{\frac{5}{12}} \left(\frac{100 \text{ MeV}}{T_{\text{dec}}}\right), \quad (25)$$

from which one can see that the parameter dependence is much different from the DW without current/charge.

Fig. 1 shows the GW spectrum obtained from the superposition of those of DW network and spherons with several benchmark cases. We here assume a step-function-like transition between IR and UV regime for simplicity. The left and right peaks of thick solid lines correspond to the DW network and spherons, respectively. Thin solid lines indicate power-law integrated sensitivity curves of future GW observatories: the Square Kilometer Array (SKA) [90, 91], Gaia and THEIA [92], LISA [93, 94], μ ARES [95], DECIGO [96], AEDGE [97], BBO [98, 99] Einstein Telescope (ET) [100, 101], and Cosmic Explorer (CE) [102, 103]. We take the threshold signal-to-noise ratio (SNR) to be $\text{SNR} = 1$. A black dotted curve is obtained by considering LISA and ET operations together, see Ref. [104] for details. We also show the constraints on ΔN_{eff} from PLANCK 2018 limits [105], as well as future reaches of cosmic microwave background (CMB) experiments like CMB-HD [106, 107]. The red shaded region (aLV) is excluded by the constraint from advanced LIGO-VIRGO [108]. One can see that the GW signal from the spherons is even larger than the conventional one depending on the parameters. Furthermore, it improves the detectability of GW in higher-frequency

region.

Fig. 2 shows parameter space of σ and ΔV in which SNR exceeds 10. Solid and dashed contours indicate that GW signals from spherons and conventional DW network are detected by future GW experiments with $\text{SNR} = 10$, respectively. The gray bottom-right region is excluded due to the DW domination (Eq. (2)). On the other hand, in the purple region, the DW network annihilation time t_{ann} , at which spherons are formed, is later than the spheron lifetime so that the spheron cannot be formed. See Supplementary Materials for the details of the SNR calculation.

When the spherons are sufficiently long-lived, their energy dominates the total energy density of the universe, resulting in early matter domination. The decay of the spherons injects entropy into the thermal bath and dilutes GW radiated by the DW network [109]. We checked that this effect is not significant in the parameter space presented above.

Discussion and conclusions— Domain Wall (DW) networks could have formed in the early universe after the spontaneous breaking of a discrete symmetry. In presence of a vacuum energy difference ΔV lifting the degeneracy between the different vacua, DWs are driven towards annihilating each other. In order to have viable




FIG. 1: GW spectrum from DW network and spheron. Thick solid lines indicate the GW spectrum that is superposition of those from the conventional DW network (left peaks) and from the spherons (right peaks). Thin solid lines are power-law integrated sensitivity curves for future GW experiments. A dashed black line is a broken power-law integrated sensitivity curve obtained by combining LISA and ET sensitivity.

cosmology, the DW network must annihilate before occupying a significant energy fraction of the universe.

We showed that GW arising due to current-carrying DW gives a novel shape in the form of an extra peak (see Fig. 1) from metastable topological remnant, which we dub as spheron. The extra contribution leads to the GW spectrum with such features detectable in upcoming GW experiments like LISA and ET. We identified the GW sensitivity reaches of the parameter space of current-carrying DW (see Fig. 2). One can find that they have nice complementarity, namely, GW from spherons can provide significant detectability in parameter space in which that from conventional DW network cannot be detected. Furthermore, the parameter dependence (especially ΔV dependence) is quite non-trivial compared to that from DW networks. This is because the position of the secondary peak (right one) depends on ΔV . Thus there is an optimized ΔV leading to the maximum SNR for each GW experiment. Nonetheless, higher frequency GW with larger ΔV can be detected in ultrahigh frequency GW detectors in the future, see Ref.[110] for a review.

We also investigate the DW network interpretation of




FIG. 2: **Left panel:** Parameter space where the two GW signals will be observed with SNR > 10 (right side of the contours) for different GW experiments. We separate the calculation into the signals from the spherons (solid contours) and the conventional DW network (DWN) (dashed lines). The gray and purple bottom-right regions are excluded due to the DW domination and the fact that the spheron has too short lifetime to be formed, respectively. The points with black and white stars indicate benchmark points corresponding to the blue lines in the upper and lower panels in Fig. 1, respectively.

stochastic GW background recently reported in PTA collaborations [111–114], and predict that if such a signal in PTA arises due to current-carrying DW network, the additional new peak corresponding to the spherons will be detectable in LISA for $T_{\text{dec}} \simeq 10 \text{ MeV}$ and $10^{-14} \lesssim Y \lesssim 10^{-10}$, see sec. in Supplementary Materials for the details.

We discussed above simple BSM examples involving 2HDM(+singlet extension) in which current-carrying DWs maybe formed. While our analysis is model-independent, our prescription for the GW spectrum and the parameter space that be tested can be easily translated into BSM microphysics. In addition, the current-carrying DW may feel more friction from bulk particles than usual [26, 115–118], as the charge capture process causes the momentum transfer between the particle and DW. This friction can make a shrinking closed DW non-relativistic and hence helps to form spherons. Nevertheless it is also non-trivial how this affects the evolution of the DW network before collapsing. Indeed, the friction tends to prevent the network from reaching the scaling regime while the network reconnection would release the

trapped particles to the bulk. A further detailed analysis is necessary to address this point. In summary, Gravitational Wave Mountains from current-carrying DW are an interesting target for planned GW searches. Ultimately, our results call for an independent confirmation based on numerical lattice simulations, which will be done elsewhere.

Acknowledgements— The authors thank Simone Blasi, David Dunsy, Ricardo Z. Ferreira, Yuta Hamada, Thomas Konstandin, Kohta Miura, Kyohei Mukaida, Wakutaka Nakano, Sabir Ramazanov, Mohamed Younes Sassi, and Tanmay Vachaspati for useful discussions. This work is supported by the Deutsche Forschungsgemeinschaft under Germany's Excellence Strategy - EXC 2121 Quantum Universe - 390833306.

SUPPLEMENTARY MATERIALS

Signal-to-noise ratio (SNR)

We now wish to present the calculation of the SNR. A similar method to what we describe in this section, up to some small updates, has also been used in [119]. For details regarding the SNR estimations see e.g. [120–124]. Interferometers measure displacements in terms of a so called dimensionless strain-noise $h_{\text{GW}}(f)$ that is related to the GW amplitude and can be converted into the corresponding energy density [92]

$$h^2 \Omega_{\text{exp}}(f) = \frac{2\pi^2 f^2}{3H_0^2} h_{\text{GW}}(f)^2 h^2, \quad (26)$$

with $H_0 = h \times 100$ (km/s)/Mpc being the Hubble rate today. We compute the SNR for a given or projected experimental noise sensitivity curves $\Omega_{\text{exp}}(f)h^2$ in order to assess the detection probability of the GW background via the following prescription [122, 125]

$$\text{SNR} \equiv \sqrt{2t_{\text{obs}} \int_{f_{\text{min}}}^{f_{\text{max}}} df \left(\frac{h^2 \Omega_{\text{GW}}(f)}{h^2 \Omega_{\text{exp}}(f)} \right)^2}, \quad (27)$$

where $h = 0.7$ and the observation period t_{obs} is taken to be 20 years for Gaia, THEIA and SKA, and 4 years for the others.

One may consider the effective dark radiation bounds during Big Bang Nucleosynthesis (BBN) and CMB decoupling. In particular, the energy density of the stochastic GW background needs to be smaller than the limit on dark radiation which is depicted in ΔN_{eff} from BBN and CMB observations. This is because the gravitons behave as massless relativistic degrees of freedom. Any change of the number of effective relativistic degrees of freedom (N_{eff}) at the time of recombination is usually set by the relation [126]

$$\int_{f_{\text{min}}}^{\infty} \frac{df}{f} h^2 \Omega_{\text{GW}}(f) \leq 5.6 \times 10^{-6} \Delta N_{\text{eff}}. \quad (28)$$




FIG. 3: Plot of SNR for GW from conventional DW network (dashed) and DW network+spherons (solid). The horizontal dotted line indicates SNR = 1.

While the lower limit for the integration is $f_{\text{min}} \simeq 10^{-10}$ Hz for BBN and $f_{\text{min}} \simeq 10^{-18}$ Hz for the CMB bounds, in practice, when e.g. several GW spectra are depicted simultaneously, as a first-order estimate, one uses the approximation to ignore the frequency dependence and to set bounds just on the energy density of the peak for a given GW spectrum; this is shown as

$$h^2 \Omega_{\text{GW}}^{\text{Peak}} \leq 5.6 \times 10^{-6} \Delta N_{\text{eff}}. \quad (29)$$

We consider the constraints on ΔN_{eff} from BBN and the PLANCK 2018 limits [105], as well as future reaches of CMB experiments such as CMB-S4 [127, 128] and CMB-HD [106, 107].

In order to calculate SNR or show sensitivity curves in Fig. 1, we need information of noise h_{GW} or Ω_{exp} . Let us summarize the references from which we read off the information of noise.

- Gaia and THEIA: noise energy density Ω_{exp} from Ref. [92]
- LISA: noise energy density Ω_{exp} from Ref. [93]
- μ ARES: noise energy density Ω_{exp} from Ref. [95]
- DECIGO: strain-noise h_{GW} from Ref. [96]
- AEDGE: strain-noise h_{GW} from Ref. [97]
- BBO: strain-noise h_{GW} from Refs. [98, 99]
- ET: strain-noise h_{GW} from Ref. [101]
- CE: strain-noise h_{GW} available on <https://dcc.cosmicexplorer.org/CE-T2000017/public>
- SKA 20 years: power-law integrated curve directly taken from Ref. [129] with an appropriate factor to compensate the difference of the threshold SNR.




FIG. 4: GW spectrum from DW network and spheron. We took the parameters σ and ΔV such that the GW spectrum from the network fits recent PTA signals (NANOGrav: gray, EPTA: red). By fixing $T_{\text{dec}} = 10 \text{ MeV}$, the additional peak from spherons can lie within LISA sensitivity range (purple) for $10^{-14} \lesssim Y \lesssim 10^{-10}$.

Fig. 3 shows plots of SNR for several future GW experiments to compare conventional DW network (dashed) and DW network+spherons (solid). The horizontal dotted line indicates SNR = 1. Clearly one can see enhancement of SNR thanks to the spherons.

DW interpretation of PTA signal

We here investigate the DW network interpretation of stochastic GW background recently reported in PTA collaborations [111–114]. In order for the GW spectrum from DW network to fit the PTA signals, we should take appropriate σ and ΔV . If these DW are current-carrying, we may have an additional peak from spherons as stated in the main text, so that we should be able to predict the position and height of the additional peak. In particular, one finds the peak within the sensitivity range of LISA when one takes $T_{\text{dec}} = 10 \text{ MeV}$ and $10^{-14} \lesssim Y \lesssim 10^{-10}$, see Fig. 4. (For $Y \lesssim 10^{-14}$, it is difficult to distinguish the additional peak.) The gray and red bars show the “violin plots” for NANOGrav with 15-years data [111] and European Pulsar Timing Array [112], respectively. The purple shaded region is the power-law integrated sensitivity for LISA, calculated in the same way as Fig. 1.

* anish.ghoshal@fuw.edu.pl

† yu.hamada@desy.de

[1] W. H. Press, B. S. Ryden and D. N. Spergel, *Dynamical Evolution of Domain Walls in an Expanding Universe*, *Astrophys. J.* **347** (1989) 590.

- [2] M. Hindmarsh, *Analytic scaling solutions for cosmic domain walls*, *Phys. Rev. Lett.* **77** (1996) 4495 [hep-ph/9605332].
- [3] P. P. Avelino, C. J. A. P. Martins and J. C. R. E. Oliveira, *One-scale model for domain wall network evolution*, *Phys. Rev. D* **72** (2005) 083506 [hep-ph/0507272].
- [4] T. W. B. Kibble, *Topology of Cosmic Domains and Strings*, *J. Phys. A* **9** (1976) 1387.
- [5] A. Vilenkin, *Gravitational Field of Vacuum Domain Walls and Strings*, *Phys. Rev. D* **23** (1981) 852.
- [6] P. Sikivie, *Of Axions, Domain Walls and the Early Universe*, *Phys. Rev. Lett.* **48** (1982) 1156.
- [7] G. B. Gelmini, M. Gleiser and E. W. Kolb, *Cosmology of Biased Discrete Symmetry Breaking*, *Phys. Rev. D* **39** (1989) 1558.
- [8] F. Ferrer, E. Masso, G. Panico, O. Pujolas and F. Rompineve, *Primordial Black Holes from the QCD axion*, *Phys. Rev. Lett.* **122** (2019) 101301 [1807.01707].
- [9] H. Deng, J. Garriga and A. Vilenkin, *Primordial black hole and wormhole formation by domain walls*, *JCAP* **04** (2017) 050 [1612.03753].
- [10] H. Deng and A. Vilenkin, *Primordial black hole formation by vacuum bubbles*, *JCAP* **12** (2017) 044 [1710.02865].
- [11] H. Deng, *Primordial black hole formation by vacuum bubbles. Part II*, *JCAP* **09** (2020) 023 [2006.11907].
- [12] Y. Gouttenoire and E. Vitagliano, *Primordial black holes and wormholes from domain wall networks*, *Phys. Rev. D* **109** (2024) 123507 [2311.07670].
- [13] R. Z. Ferreira, A. Notari, O. Pujolàs and F. Rompineve, *Collapsing domain wall networks: impact on pulsar timing arrays and primordial black holes*, *JCAP* **06** (2024) 020 [2401.14331].
- [14] D. I. Dunskey and M. Kongsore, *Primordial black holes from axion domain wall collapse*, *JHEP* **06** (2024) 198 [2402.03426].
- [15] G. B. Gelmini, A. Simpson and E. Vitagliano, *Catastrogenesis: DM, GWs, and PBHs from ALP string-wall networks*, *JCAP* **02** (2023) 031 [2207.07126].
- [16] G. B. Gelmini, J. Hyman, A. Simpson and E. Vitagliano, *Primordial black hole dark matter from catastrogenesis with unstable pseudo-Goldstone bosons*, *JCAP* **06** (2023) 055 [2303.14107].
- [17] T. Hiramatsu, M. Kawasaki, K. Saikawa and T. Sekiguchi, *Axion cosmology with long-lived domain walls*, *JCAP* **01** (2013) 001 [1207.3166].
- [18] T. Hiramatsu, M. Kawasaki and K. Saikawa, *On the estimation of gravitational wave spectrum from cosmic domain walls*, *JCAP* **02** (2014) 031 [1309.5001].
- [19] K. Saikawa, *A review of gravitational waves from cosmic domain walls*, *Universe* **3** (2017) 40 [1703.02576].
- [20] T. Krajewski, J. H. Kwapisz, Z. Lalak and M. Lewicki, *Stability of domain walls in models with asymmetric potentials*, *Phys. Rev. D* **104** (2021) 123522 [2103.03225].
- [21] C.-F. Chang and Y. Cui, *Dynamics of Long-lived Axion Domain Walls and Its Cosmological Implications*, **2309.15920**.
- [22] N. Kitajima, J. Lee, K. Murai, F. Takahashi and W. Yin, *Gravitational waves from domain wall*

- collapse, and application to nanohertz signals with QCD-coupled axions, *Phys. Lett. B* **851** (2024) 138586 [2306.17146].
- [23] R. Z. Ferreira, S. Gasparotto, T. Hiramatsu, I. Obata and O. Pujolas, *Axionic defects in the CMB: birefringence and gravitational waves*, *JCAP* **05** (2024) 066 [2312.14104].
- [24] N. Kitajima, J. Lee, F. Takahashi and W. Yin, *Stability of domain walls with inflationary fluctuations under potential bias, and gravitational wave signatures*, **2311.14590**.
- [25] I. Dankovsky, E. Babichev, D. Gorbunov, S. Ramazanov and A. Vikman, *Revisiting evolution of domain walls and their gravitational radiation with CosmoLattice*, *JCAP* **09** (2024) 047 [2406.17053].
- [26] K. Nakayama, F. Takahashi and N. Yokozaki, *Gravitational waves from domain walls and their implications*, *Phys. Lett. B* **770** (2017) 500 [1612.08327].
- [27] C. J. A. P. Martins, I. Y. Rybak, A. Avgoustidis and E. P. S. Shellard, *Extending the velocity-dependent one-scale model for domain walls*, *Phys. Rev. D* **93** (2016) 043534 [1602.01322].
- [28] C. J. A. P. Martins, I. Y. Rybak, A. Avgoustidis and E. P. S. Shellard, *Stretching and Kibble scaling regimes for Hubble-damped defect networks*, *Phys. Rev. D* **94** (2016) 116017 [1612.08863]. [Erratum: Phys.Rev.D 95, 039902 (2017)].
- [29] P. P. Avelino, *Comparing parametric and non-parametric velocity-dependent one-scale models for domain wall evolution*, *JCAP* **04** (2020) 012 [2001.06318].
- [30] O. Pujolas and G. Zahariade, *Domain wall annihilation: A QFT perspective*, *Phys. Rev. D* **107** (2023) 123527 [2212.11204].
- [31] D. Grüber, L. Sousa and P. P. Avelino, *Stochastic gravitational wave background generated by domain wall networks*, *Phys. Rev. D* **110** (2024) 023505 [2403.09816].
- [32] E. Witten, *Superconducting Strings*, *Nucl. Phys. B* **249** (1985) 557.
- [33] R. L. Davis and E. P. S. Shellard, *COSMIC VORTONS*, *Nucl. Phys. B* **323** (1989) 209.
- [34] B. Carter, *Cosmological relic distribution of conducting string loops*, *Annals of the New York Academy of Sciences* **647** (1991) 758.
- [35] J. M. Alimi, A. Blanchard, A. Bouquet, F. Martin de Volnay and J. Tran Thanh Van, eds., *Particle astrophysics: The Early universe and cosmic structures. Proceedings, 25th Rencontres de Moriond, Les Arcs, France, March 4-11, 1990*, (Gif-Sur-Yvette), Ed. Frontieres, 1990.
- [36] B. Carter and X. Martin, *Dynamic instability criterion for circular (Vorton) string loops*, *Annals Phys.* **227** (1993) 151 [hep-th/0306111].
- [37] X. Martin and P. Peter, *Dynamical stability of Witten rings*, *Phys. Rev. D* **51** (1995) 4092 [hep-ph/9405220].
- [38] E. Radu and M. S. Volkov, *Existence of stationary, non-radiating ring solitons in field theory: knots and vortons*, *Phys. Rept.* **468** (2008) 101 [0804.1357].
- [39] R. A. Battye and P. M. Sutcliffe, *Vorton construction and dynamics*, *Nucl. Phys. B* **814** (2009) 180 [0812.3239].
- [40] J. Garaud, E. Radu and M. S. Volkov, *Stable Cosmic Vortons*, *Phys. Rev. Lett.* **111** (2013) 171602 [1303.3044].
- [41] R. A. Battye, S. J. Cotterill and J. A. Pearson, *A detailed study of the stability of vortons*, *JHEP* **04** (2022) 005 [2112.08066].
- [42] R. A. Battye and S. J. Cotterill, *Stable Cosmic Vortons in Bosonic Field Theory*, *Phys. Rev. Lett.* **127** (2021) 241601 [2111.07822].
- [43] M. Ibe, S. Kobayashi, Y. Nakayama and S. Shirai, *On Stability of Fermionic Superconducting Current in Cosmic String*, *JHEP* **05** (2021) 217 [2102.05412].
- [44] Y. Abe, Y. Hamada, K. Saji and K. Yoshioka, *Quantum current dissipation in superconducting strings and vortons*, *JHEP* **02** (2023) 004 [2209.03223].
- [45] K. Harigaya, X. Niu, W. Xue and F. Yang, *Stability of Superconducting Strings*, **2412.12259**.
- [46] J. P. Ostriker, A. C. Thompson and E. Witten, *Cosmological Effects of Superconducting Strings*, *Phys. Lett. B* **180** (1986) 231.
- [47] M. F. Oliveira, A. Avgoustidis and C. J. A. P. Martins, *Cosmic string evolution with a conserved charge*, *Phys. Rev. D* **85** (2012) 083515 [1201.5064].
- [48] B. Hartmann, F. Michel and P. Peter, *Excited cosmic strings with superconducting currents*, *Phys. Rev. D* **96** (2017) 123531 [1710.00738].
- [49] R. Brandenberger, B. Cyr and R. Shi, *Constraints on Superconducting Cosmic Strings from the Global 21-cm Signal before Reionization*, *JCAP* **09** (2019) 009 [1902.08282].
- [50] B. Imtiaz, R. Shi and Y.-F. Cai, *Updated constraints on superconducting cosmic strings from the astronomy of fast radio bursts*, *Eur. Phys. J. C* **80** (2020) 500 [2001.11149].
- [51] C. J. A. P. Martins, P. Peter, I. Y. Rybak and E. P. S. Shellard, *Generalized velocity-dependent one-scale model for current-carrying strings*, *Phys. Rev. D* **103** (2021) 043538 [2011.09700].
- [52] H. Fukuda, A. V. Manohar, H. Murayama and O. Telem, *Axion strings are superconducting*, *JHEP* **06** (2021) 052 [2010.02763].
- [53] Y. Abe, Y. Hamada and K. Yoshioka, *Electroweak axion string and superconductivity*, *JHEP* **06** (2021) 172 [2010.02834].
- [54] P. Agrawal, A. Hook, J. Huang and G. Marques-Tavares, *Axion string signatures: a cosmological plasma collider*, *JHEP* **01** (2022) 103 [2010.15848].
- [55] R. Thériault, J. T. Mirocha and R. Brandenberger, *Global 21cm absorption signal from superconducting cosmic strings*, *JCAP* **10** (2021) 046 [2105.01166].
- [56] C. J. A. P. Martins, P. Peter, I. Y. Rybak and E. P. S. Shellard, *Charge-velocity-dependent one-scale linear model*, *Phys. Rev. D* **104** (2021) 103506 [2108.03147].
- [57] B. Cyr, H. Jiao and R. Brandenberger, *Massive black holes at high redshifts from superconducting cosmic strings*, *Mon. Not. Roy. Astron. Soc.* **517** (2022) 2221 [2202.01799].
- [58] I. Y. Rybak and L. Sousa, *Emission of gravitational waves by superconducting cosmic strings*, *JCAP* **11** (2022) 024 [2209.01068].
- [59] P. Auclair, S. Blasi, V. Brdar and K. Schmitz, *Gravitational waves from current-carrying cosmic strings*, *JCAP* **04** (2023) 009 [2207.03510].

- [60] R. Jackiw and C. Rebbi, *Solitons with Fermion Number 1/2*, *Phys. Rev. D* **13** (1976) 3398.
- [61] V. A. Rubakov and M. E. Shaposhnikov, *Do We Live Inside a Domain Wall?*, *Phys. Lett. B* **125** (1983) 136.
- [62] D. B. Kaplan, *A Method for simulating chiral fermions on the lattice*, *Phys. Lett. B* **288** (1992) 342 [[hep-lat/9206013](#)].
- [63] P. Peter, *Surface current carrying domain walls*, *J. Phys. A* **29** (1996) 5125 [[hep-ph/9503408](#)].
- [64] T. Vachaspati, *Kinks and Domain Walls : An Introduction to Classical and Quantum Solitons*. Oxford University Press, 2007, [10.1017/9781009290456](#).
- [65] R. A. Battye, G. D. Brawn and A. Pilaftsis, *Vacuum Topology of the Two Higgs Doublet Model*, *JHEP* **08** (2011) 020 [[1106.3482](#)].
- [66] M. Eto, M. Kurachi and M. Nitta, *Constraints on two Higgs doublet models from domain walls*, *Phys. Lett. B* **785** (2018) 447 [[1803.04662](#)].
- [67] R. A. Battye, A. Pilaftsis and D. G. Viatic, *Domain wall constraints on two-Higgs-doublet models with Z_2 symmetry*, *Phys. Rev. D* **102** (2020) 123536 [[2010.09840](#)].
- [68] R. A. Battye, A. Pilaftsis and D. G. Viatic, *Simulations of Domain Walls in Two Higgs Doublet Models*, *JHEP* **01** (2021) 105 [[2006.13273](#)].
- [69] K. H. Law and A. Pilaftsis, *Charged and CP-violating kink solutions in the two-Higgs-doublet model*, *Phys. Rev. D* **105** (2022) 056007 [[2110.12550](#)].
- [70] M. Y. Sassi and G. Moortgat-Pick, *Domain walls in the Two-Higgs-Doublet Model and their charge and CP-violating interactions with Standard Model fermions*, *JHEP* **04** (2024) 101 [[2309.12398](#)].
- [71] B. Fu, A. Ghoshal, S. F. King and M. H. Rahat, *Type-I two-Higgs-doublet model and gravitational waves from domain walls bounded by strings*, *JHEP* **08** (2024) 237 [[2404.16931](#)].
- [72] G. C. Branco, P. M. Ferreira, L. Lavoura, M. N. Rebelo, M. Sher and J. P. Silva, *Theory and phenomenology of two-Higgs-doublet models*, *Phys. Rept.* **516** (2012) 1 [[1106.0034](#)].
- [73] C.-Y. Chen, M. Freid and M. Sher, *Next-to-minimal two Higgs doublet model*, *Phys. Rev. D* **89** (2014) 075009 [[1312.3949](#)].
- [74] M. Muhlleitner, M. O. P. Sampaio, R. Santos and J. Wittbrodt, *The $N2HDM$ under Theoretical and Experimental Scrutiny*, *JHEP* **03** (2017) 094 [[1612.01309](#)].
- [75] M. Muhlleitner, M. O. P. Sampaio, R. Santos and J. Wittbrodt, *Phenomenological Comparison of Models with Extended Higgs Sectors*, *JHEP* **08** (2017) 132 [[1703.07750](#)].
- [76] M. Y. Sassi and G. Moortgat-Pick, *Electroweak Symmetry Restoration in the $N2HDM$ via Domain Walls*, [2407.14468](#).
- [77] G. Lazarides and Q. Shafi, *SUPERCONDUCTING MEMBRANES*, *Phys. Lett. B* **159** (1985) 261.
- [78] T. W. B. Kibble, G. Lazarides and Q. Shafi, *Walls Bounded by Strings*, *Phys. Rev. D* **26** (1982) 435.
- [79] A. Vilenkin and A. E. Everett, *Cosmic Strings and Domain Walls in Models with Goldstone and PseudoGoldstone Bosons*, *Phys. Rev. Lett.* **48** (1982) 1867.
- [80] A. E. Everett and A. Vilenkin, *Left-right Symmetric Theories and Vacuum Domain Walls and Strings*, *Nucl. Phys. B* **207** (1982) 43.
- [81] J. Preskill and A. Vilenkin, *Decay of metastable topological defects*, *Phys. Rev. D* **47** (1993) 2324 [[hep-ph/9209210](#)].
- [82] D. I. Dunsky, A. Ghoshal, H. Murayama, Y. Sakakihara and G. White, *GUTs, hybrid topological defects, and gravitational waves*, *Phys. Rev. D* **106** (2022) 075030 [[2111.08750](#)].
- [83] M. Eto, Y. Hamada and M. Nitta, *Composite topological solitons consisting of domain walls, strings, and monopoles in $O(N)$ models*, *JHEP* **08** (2023) 150 [[2304.14143](#)].
- [84] S. Aoki and H. Fukaya, *Curved domain-wall fermions*, *PTEP* **2022** (2022) 063B04 [[2203.03782](#)].
- [85] S. Aoki and H. Fukaya, *Curved domain-wall fermion and its anomaly inflow*, *PTEP* **2023** (2023) 033B05 [[2212.11583](#)].
- [86] S. Aoki, H. Fukaya, N. Kan, M. Koshino and Y. Matsuki, *Magnetic monopole becomes dyon in topological insulators*, *Phys. Rev. B* **108** (2023) 155104 [[2304.13954](#)].
- [87] C. J. A. P. Martins and E. P. S. Shellard, *Vorton formation*, *Phys. Rev. D* **57** (1998) 7155 [[hep-ph/9804378](#)].
- [88] T. D. Lee and Y. Pang, *Nontopological solitons*, *Phys. Rept.* **221** (1992) 251.
- [89] S. R. Coleman, *Q-balls*, *Nucl. Phys. B* **262** (1985) 263. [Addendum: *Nucl.Phys.B* 269, 744 (1986)].
- [90] G. Janssen et al., *Gravitational wave astronomy with the SKA*, *PoS AASKA14* (2015) 037 [[1501.00127](#)].
- [91] A. Weltman et al., *Fundamental physics with the Square Kilometre Array*, *Publ. Astron. Soc. Austral.* **37** (2020) e002 [[1810.02680](#)].
- [92] J. Garcia-Bellido, H. Murayama and G. White, *Exploring the early Universe with Gaia and Theia*, *JCAP* **12** (2021) 023 [[2104.04778](#)].
- [93] LISA COSMOLOGY WORKING GROUP COLLABORATION, P. Auclair et al., *Cosmology with the Laser Interferometer Space Antenna*, *Living Rev. Rel.* **26** (2023) 5 [[2204.05434](#)].
- [94] R. Flauger, N. Karnesis, G. Nardini, M. Pieroni, A. Ricciardone and J. Torrado, *Improved reconstruction of a stochastic gravitational wave background with LISA*, *JCAP* **01** (2021) 059 [[2009.11845](#)].
- [95] A. Sesana et al., *Unveiling the gravitational universe at μ -Hz frequencies*, *Exper. Astron.* **51** (2021) 1333 [[1908.11391](#)].
- [96] S. Kawamura et al., *Current status of space gravitational wave antenna DECIGO and B-DECIGO*, *PTEP* **2021** (2021) 05A105 [[2006.13545](#)].
- [97] AEDGE COLLABORATION, Y. A. El-Neaj et al., *AEDGE: Atomic Experiment for Dark Matter and Gravity Exploration in Space*, *EPJ Quant. Technol.* **7** (2020) 6 [[1908.00802](#)].
- [98] V. Corbin and N. J. Cornish, *Detecting the cosmic gravitational wave background with the big bang observer*, *Class. Quant. Grav.* **23** (2006) 2435 [[gr-qc/0512039](#)].
- [99] G. M. Harry, P. Fritschel, D. A. Shaddock, W. Folkner and E. S. Phinney, *Laser interferometry for the big bang observer*, *Class. Quant. Grav.* **23** (2006) 4887.

- [Erratum: Class.Quant.Grav. 23, 7361 (2006)].
- [100] M. Punturo et al., *The Einstein Telescope: A third-generation gravitational wave observatory*, *Class. Quant. Grav.* **27** (2010) 194002.
- [101] S. Hild et al., *Sensitivity Studies for Third-Generation Gravitational Wave Observatories*, *Class. Quant. Grav.* **28** (2011) 094013 [1012.0908].
- [102] LIGO SCIENTIFIC Collaboration, B. P. Abbott et al., *Exploring the Sensitivity of Next Generation Gravitational Wave Detectors*, *Class. Quant. Grav.* **34** (2017) 044001 [1607.08697].
- [103] D. Reitze et al., *Cosmic Explorer: The U.S. Contribution to Gravitational-Wave Astronomy beyond LIGO*, *Bull. Am. Astron. Soc.* **51** (2019) 035 [1907.04833].
- [104] A. Marriotti-Best, D. Chowdhury, A. Ghoshal and G. Tasinato, *Exploring cosmological gravitational wave backgrounds through the synergy of LISA and ET*, **2409.02886**.
- [105] PLANCK Collaboration, N. Aghanim et al., *Planck 2018 results. VI. Cosmological parameters*, *Astron. Astrophys.* **641** (2020) A6 [1807.06209]. [Erratum: *Astron. Astrophys.* 652, C4 (2021)].
- [106] N. Sehgal et al., *CMB-HD: An Ultra-Deep, High-Resolution Millimeter-Wave Survey Over Half the Sky*, **1906.10134**.
- [107] CMB-HD Collaboration, S. Aiola et al., *Snowmass2021 CMB-HD White Paper*, **2203.05728**.
- [108] KAGRA, VIRGO, LIGO SCIENTIFIC Collaboration, R. Abbott et al., *Upper limits on the isotropic gravitational-wave background from Advanced LIGO and Advanced Virgo's third observing run*, *Phys. Rev. D* **104** (2021) 022004 [2101.12130].
- [109] Y. Gouttenoire, *Primordial black holes from conformal Higgs*, *Phys. Lett. B* **855** (2024) 138800 [2311.13640].
- [110] N. Aggarwal et al., *Challenges and opportunities of gravitational-wave searches at MHz to GHz frequencies*, *Living Rev. Rel.* **24** (2021) 4 [2011.12414].
- [111] NANOGrav Collaboration, G. Agazie et al., *The NANOGrav 15 yr Data Set: Evidence for a Gravitational-wave Background*, *Astrophys. J. Lett.* **951** (2023) L8 [2306.16213].
- [112] EPTA, INPTA: Collaboration, J. Antoniadis et al., *The second data release from the European Pulsar Timing Array - III. Search for gravitational wave signals*, *Astron. Astrophys.* **678** (2023) A50 [2306.16214].
- [113] H. Xu et al., *Searching for the Nano-Hertz Stochastic Gravitational Wave Background with the Chinese Pulsar Timing Array Data Release I*, *Res. Astron. Astrophys.* **23** (2023) 075024 [2306.16216].
- [114] D. J. Reardon et al., *Search for an Isotropic Gravitational-wave Background with the Parkes Pulsar Timing Array*, *Astrophys. J. Lett.* **951** (2023) L6 [2306.16215].
- [115] M. C. Huang and P. Sikivie, *The Structure of Axionic Domain Walls*, *Phys. Rev. D* **32** (1985) 1560.
- [116] A. Vilenkin and E. P. S. Shellard, *Cosmic Strings and Other Topological Defects*. Cambridge University Press, 7, 2000.
- [117] S. Blasi, A. Mariotti, A. Rase, A. Sevrin and K. Turbang, *Friction on ALP domain walls and gravitational waves*, *JCAP* **04** (2023) 008 [2210.14246].
- [118] S. Hassan, G. R. Kane, J. March-Russell and G. Obied, *Chern-Simons Induced Thermal Friction on Axion Domain Walls*, **2410.19906**.
- [119] I. Baldes, Y. Gouttenoire, F. Sala and G. Servant, *Supercool composite Dark Matter beyond 100 TeV*, *JHEP* **07** (2022) 084 [2110.13926].
- [120] B. Allen and J. D. Romano, *Detecting a stochastic background of gravitational radiation: Signal processing strategies and sensitivities*, *Phys. Rev. D* **59** (1999) 102001 [gr-qc/9710117].
- [121] H. Kudoh, A. Taruya, T. Hiramatsu and Y. Himemoto, *Detecting a gravitational-wave background with next-generation space interferometers*, *Phys. Rev. D* **73** (2006) 064006 [gr-qc/0511145].
- [122] E. Thrane and J. D. Romano, *Sensitivity curves for searches for gravitational-wave backgrounds*, *Phys. Rev. D* **88** (2013) 124032 [1310.5300].
- [123] C. Caprini, D. G. Figueroa, R. Flauger, G. Nardini, M. Peloso, M. Pieroni, A. Ricciardone and G. Tasinato, *Reconstructing the spectral shape of a stochastic gravitational wave background with LISA*, *JCAP* **11** (2019) 017 [1906.09244].
- [124] D. Brzemiński, A. Hook and G. Marques-Tavares, *Precision early universe cosmology from stochastic gravitational waves*, *JHEP* **11** (2022) 061 [2203.13842].
- [125] C. Caprini et al., *Science with the space-based interferometer eLISA. II: Gravitational waves from cosmological phase transitions*, *JCAP* **04** (2016) 001 [1512.06239].
- [126] M. Maggiore, *Gravitational wave experiments and early universe cosmology*, *Phys. Rept.* **331** (2000) 283 [gr-qc/9909001].
- [127] CMB-S4 Collaboration, K. Abazajian et al., *CMB-S4: Forecasting Constraints on Primordial Gravitational Waves*, *Astrophys. J.* **926** (2022) 54 [2008.12619].
- [128] CMB-S4 Collaboration, K. Abazajian et al., *Snowmass 2021 CMB-S4 White Paper*, **2203.08024**.
- [129] M. Breitbach, J. Kopp, E. Madge, T. Opferkuch and P. Schwaller, *Dark, Cold, and Noisy: Constraining Secluded Hidden Sectors with Gravitational Waves*, *JCAP* **07** (2019) 007 [1811.11175].