pyBoLaNO: A Python symbolic package for normal ordering involving bosonic ladder operators

Hendry M. Lim^{a,b,*}, Donny Dwiputra^{c,a}, M. Shoufie Ukhtary^a, Ahmad R. T. Nugraha^{a,d}

^aResearch Center for Quantum Physics, National Research and Innovation Agency (BRIN), South Tangerang 15314, Indonesia ^bDepartment of Physics, Faculty of Mathematics and Natural Sciences, Universitas Indonesia, Depok 16424, Indonesia ^cAsia Pacific Center for Theoretical Physics, Pohang 37673, Korea

^dDepartment of Engineering Physics, Telkom University, Bandung 40257, Indonesia

Abstract

We present pyBoLaNO, a Python symbolic package based on SymPy to quickly normal-order (Wick-order) any polynomial in bosonic ladder operators. By extension, this package offers the normal ordering of commutators of any two polynomials in bosonic ladder operators and the evaluation of the normal-ordered expectation value evolution in the Lindblad master equation framework for open quantum systems. The package also supports multipartite descriptions and multiprocessing. We describe the package's workflow, show examples of use, and discuss its computational performance. All codes and examples are available on our GitHub repository.

to the *n*th energy level E_n is obtained by applying \hat{b}^{\dagger} to the vacuum state $|0\rangle$ a total of *n* times, $|n\rangle = \mathcal{N}\hat{b}^{\dagger n} |0\rangle$, where \mathcal{N} is a normalization constant [1, 2]. Taking into account quantum statistics, the ladder operators are defined differently for bosons and fermions—usually denoted \hat{a} and \hat{a}^{\dagger} in the latter case. Furthermore, the eigenvectors of the bosonic annihilation operator are quantum states that most closely resemble the classical harmonic oscillator, that is, the coherent states [3]. The formulation is

E-mail address: hendry01@ui.ac.id

far-reaching in quantum mechanics. The ladder operators form the basis for quantum field theory, in which a particle is considered as the excitation of the underlying quantum field [2]. They can also be found in the theoretical description of physical systems in atomic, molecular, and optical physics [3] that extends beyond simple harmonic oscilla-

PROGRAM SUMMARY Program Title: pyBoLaNO Developer's repository link: https://github.com/hendry24/py BoLaNO Licensing provisions: MIT License Programming language: Python Nature of problem: Normal (Wick) ordering involving bosonic ladder operators. Solution method: Blasiak's formulae for the normal ordering of an arbitrary monomial in bosonic ladder operators. Symbolic programming is fully provided by SymPy. **1. Introduction** Ladder operators arise in the study of the quantum tion/lowering operator \hat{b} and its Hermitian conjugate, the reation/raising operator \hat{b}^{\dagger} . They allow for the algebraic tratment of the problem, giving a simple relation between the system's eigenstates. The eigenstate $|n\rangle$ corresponding to the nth energy level E_n is obtained by applying \hat{b}^{\dagger} lob, the vacuum state $|0\rangle$ a total of n times, $|n\rangle = N\hat{b}^{\dagger n}|0\rangle$, As the system description becomes more complex, the algebraic manipulation of the dynamical equations be-As such, a calculator that does the algebra is desirable. The SymPy package [4] provides a symbolic computation framework in the Python programming language. At the time of writing this work, SymPy is on release ver. 1.13.3, which supports the ladder operators via the sympy.physics.secondquant submodule. However, the support is limited to arithmetic operations. The function wicks for normal (Wick) ordering is not implemented for bosonic ladder operators. In practice, we often desire to evaluate the normal ordering of operators involving bosonic ladder operators. An example of such cases, which motivates us for this work, is the evolution of the expectation values for open quantum systems described by the Lindblad master equation [5, 6]. If analytical expressions are not the concern, several packages are available to solve the problem numerically, e.g., QuTiP [7, 8] and QuantumOptics. jl [9]. Otherwise, to our knowledge [10-24], there is yet a symbolic computational package that facilitates the solution to the problem.

> In this work, we develop the Python package pyBoLaNO that offers fast symbolic normal ordering of expressions involving bosonic ladder operators, which ex-

^{*}Corresponding author.

Preprint submitted to Computer Physics Communications

tends to fast normal ordering of commutators and expectation value evolution from the Lindblad master equation. The package is fully built on and compatible with SymPy. It supports multipartite descriptions and multiprocessing for each additive term in the input(s).

The remainder of this paper is structured as follows. In Section 2, we elaborate on the theoretical basis for the features presented in this package. Section 3 describes the package's functionalities. The package usage is shown through some selected examples in Section 4. The computational performance of the package is discussed in Section 5. Finally, Section 6 concludes this paper.

2. Theoretical Considerations

Here, we present a brief overview of the formulations of the ladder operators available in quantum mechanics textbooks. The readers interested in only the essentials of the package may skip to Section 2.1.

We consider the classical Hamiltonian of the simple harmonic oscillator under Hooke's law [1]:

$$H = \frac{p^2}{2m} + \frac{m\omega_0^2 x^2}{2},$$
 (1)

where m is the oscillator's mass and ω_0 is its natural angular frequency. The quantization of this Hamiltonian is achieved by replacing the position x and momentum p with the corresponding Hilbert space operators. We obtain

$$\hat{H} = \frac{\hat{p}^2}{2m} + \frac{m\omega_0^2 \hat{x}^2}{2}.$$
 (2)

The time-independent Schröddinger equation (TISE) is given by

$$\hat{H}\psi = E\psi, \tag{3}$$

where ψ is the system's wave function and E is its energy. In the algebraic treatment of the problem (the other being the analytic method involving Hermite polynomials), we define the operators

$$\hat{b} = \frac{(m\omega_0 \hat{x} + i\hat{p})}{\sqrt{2\hbar m\omega_0}},\tag{4a}$$

$$\hat{b}^{\dagger} = \frac{(m\omega_0 \hat{x} - i\hat{p})}{\sqrt{2\hbar m\omega_0}},\tag{4b}$$

satisfying the commutation relations

$$\left[\hat{b},\hat{b}^{\dagger}\right] = 1,\tag{5}$$

where $\left[\hat{A}, \hat{B}\right] = \hat{A}\hat{B} - \hat{B}\hat{A}$. The Hamiltonian becomes

$$\hat{H} = \hbar\omega_0 \left(\hat{b}^{\dagger}\hat{b} + \frac{1}{2}\right). \tag{6}$$

Considering the time-independent Schrödinger equation (TISE),

$$\hat{H} \left| \psi \right\rangle = E \left| \psi \right\rangle,$$
 (7)

where ψ is the wave function and E is the energy, it can be shown that

$$\hat{H}\left(\hat{b}^{\dagger}|\psi\rangle\right) = \left(E + \hbar\omega_{0}\right)\left(\hat{b}^{\dagger}|\psi\rangle\right),\tag{8a}$$

$$\hat{H}\left(\hat{b}\left|\psi\right\rangle\right) = \left(E - \hbar\omega_{0}\right)\left(\hat{b}\left|\psi\right\rangle\right),\tag{8b}$$

meaning that operating on the system $|\psi\rangle$ with \hat{b}^{\dagger} raises its energy by a quantum of $\hbar\omega_0$, while operating with \hat{b} lowers its energy by $\hbar\omega_0$. This is akin to moving up and down the rung of a ladder, hence the name "ladder operators". By definition, the lowest energy state is called the vacuum state $|0\rangle$, which satisfies $\hat{H}(\hat{b}|0\rangle) = 0$. The *n*th eigenstate is given by

$$|n\rangle = \frac{1}{\sqrt{n!}}\hat{b}^{\dagger n} |0\rangle.$$
(9)

The formulation of ladder operators is not limited to a mechanical oscillator. To be more general, it can be formulated from any system described by a Hamiltonian resembling Eq. (2) where the *canonical* position q and momentum p take the roles of the real position and momentum considered above (see, for example Chap. 2 of Ref. [3] for the treatment of electromagnetic waves). In some cases, they are called the "raising" and "lowering" operators. In some others, they are called the "annihilation" and "creation" operators. In this paper, we shall use the term "ladder operators" to refer to both operators, while we use "annihilation operator" and "creation operator" to refer to \hat{b} and \hat{b}^{\dagger} , respectively.

The formulation can also be applied to an ensemble of many simple harmonic oscillators. In this case, we have pairs of ladder operators $(\hat{b}_j, \hat{b}_j^{\dagger})$, each associated with a single uncoupled oscillator in the ensemble indexed by j. They satisfy the commutation relations

$$\left[\hat{b}_j, \hat{b}_k\right] = \left[\hat{b}_j^{\dagger}, \hat{b}_k^{\dagger}\right] = 0, \qquad (10a)$$

$$\left[\hat{b}_{j},\hat{b}_{k}^{\dagger}\right] = \delta_{jk},\tag{10b}$$

where δ_{jk} is the Kronecker delta.

The indices also appear in the occupation number representation (see, for example, Chap. 3 of Ref. [2]). Given $|n_0, n_1, n_2, \ldots\rangle$, where n_0 denotes the number of particles in the quantum state ψ_0 , n_1 is the number of particles in the state ψ_1 , etc. [2], to decrease and increase the number of particles, we use the ladder operators with the corresponding index: \hat{b}_j and \hat{b}_j^{\dagger} , respectively. We consider $|0, 0\rangle$ and try to increase each count by one. This can be done in two ways:

$$\hat{b}_1^{\dagger} \hat{b}_0^{\dagger} \left| 0, 0 \right\rangle \propto \left| 1, 1 \right\rangle \tag{11}$$

or

$$\hat{b}_{0}^{\dagger}\hat{b}_{1}^{\dagger} |0,0\rangle \propto |1,1\rangle.$$
 (12)

Since we end up in the same state, we must have

$$\hat{b}_0^{\dagger} \hat{b}_1^{\dagger} = \lambda \hat{b}_1^{\dagger} \hat{b}_0^{\dagger} \tag{13}$$

where λ is a scalar. By quantum statistics, in our threedimensional world we have $\lambda = \pm 1$. We can similarly consider other processes, such as adding one particle in the state ψ_0 and removing one in the state ψ_1 . In general, the $\lambda = 1$ case corresponds to bosons and gives the commutation relations shown in Eqs. (10a) and (10b). Meanwhile, the $\lambda = -1$ case corresponds to fermions and gives the (anti)commutation relations

$$\{\hat{a}_j, \hat{a}_k\} = \left\{\hat{a}_j^{\dagger}, \hat{a}_k^{\dagger}\right\} = 0,$$
 (14a)

$$\left\{\hat{a}_j, \hat{a}_k^\dagger\right\} = \delta_{jk},\tag{14b}$$

where we have used a instead of b for fermionic ladder operators. In the following, we focus only on the bosonic ladder operators and drop the adjective "bosonic" for brevity.

2.1. Normal ordering and Blasiak's formulae

Normal ordering (or Wick ordering) means using commutation relations to rewrite the given expression so that all creation operators are written to the left of all annihilation operators. This is important in quantum field theory to avoid divergent integrals and convenient for calculations via Wick's theorem [2]. In quantum optics, it is convenient for calculations via the optical equivalence theorem [3]. The normal ordering of an operator \hat{A} may be denoted by : \hat{A} :, for example,

$$:\hat{b}_{j}\hat{b}_{j}^{\dagger}\hat{b}_{j}:=\hat{b}_{j}+\hat{b}_{j}^{\dagger}\hat{b}_{j}^{2}.$$
(15)

A useful result is given by Blasiak [25, 26] for the normal ordering of a monomial of ladder operators,

$$\hat{X} = \hat{b}^{\dagger r_M} \hat{b}^{s_M} \dots \hat{b}^{\dagger r_2} \hat{b}^{s_2} \hat{b}^{\dagger r_1} \hat{b}^{s_1}.$$
 (16)

Let $\mathbf{r} = (r_1, r_2, ..., r_M)$ and $\mathbf{s} = (s_1, s_2, ..., s_M)$. Also, let

$$d_l = \sum_{m=1}^{l} (r_m - s_m)$$
(17)

be the lth excess of creation operators in the monomial. Given the generalized Stirling numbers,

$$S_{\boldsymbol{r},\boldsymbol{s}}(k) = \frac{1}{k!} \sum_{j=0}^{k} \binom{k}{j} (-1)^{k-j} \prod_{m=1}^{M} (d_{m-1}+j)_{s_m}, \quad (18)$$

where $\binom{a}{b} = \frac{a!}{b!(a-b)!}$ is the binomial coefficient and $(m)_n = \frac{m!}{(m-n)!}$ is the falling factorial, we can obtain the normal-ordered expression for \hat{X} as

$$:\hat{X}:=\begin{cases} \hat{b}^{\dagger d_{M}} \sum_{\substack{k=s_{1} \\ r_{1}+r_{2}+\dots+r_{M} \\ \sum_{k=r_{M}} S_{\overline{s},\overline{r}}(k)\hat{b}^{\dagger k}\hat{b}^{k}, \quad d_{M} \ge 0\\ \sum_{k=r_{M}} S_{\overline{s},\overline{r}}(k)\hat{b}^{\dagger k}\hat{b}^{k} \hat{b}^{-d_{M}}, \quad d_{M} < 0 \end{cases}$$
(19)

where $\overline{r} = (r_M, \ldots, r_2, r_1)$ and $\overline{s} = (s_M, \ldots, s_2, s_1)$.

2.2. The Lindblad master equation

A widely used formalism to describe open quantum systems—quantum systems whose interactions with their environments are practically intractable—is the Lindblad master equation [5, 6]. For a system described by the density matrix ρ , the evolution of the system is given by

$$\frac{\mathrm{d}\rho}{\mathrm{d}t} = -\frac{i}{\hbar} \left[\hat{H}, \rho \right] + \sum_{j} \gamma_{j} \mathcal{D} \left(\hat{O}_{j}, \hat{P}_{j} \right) \left[\rho \right].$$
(20)

The Hamiltonian \hat{H} describes the closed system dynamics, while the Lindblad dissipators

$$\mathcal{D}\left(\hat{O}_{j},\hat{P}_{j}\right)\left[\rho\right] = \hat{O}_{j}\rho\hat{P}_{j}^{\dagger} - \frac{1}{2}\left\{\hat{P}_{j}^{\dagger}\hat{O}_{j},\rho\right\}$$
(21)

describe the open system dynamics. Each dissipator is defined by the operators \hat{O}_j, \hat{P}_j , which describe the open system process. The multiplying scalar $\gamma_j \geq 0$ can be interpreted as the process rate.

Theoretical treatment of open quantum systems often deals with calculating the evolution of expectation values for some physical quantities. Let a quantity A be represented by the operator \hat{A} . Given the density matrix ρ , the expectation value of A is given by tr $(\rho \hat{A})$. Multiplying Eq. (20) by \hat{A} (from both sides of the equation), we have

$$\frac{\mathrm{d}\langle A\rangle}{\mathrm{d}t} = -\frac{i}{\hbar} \mathrm{tr}\left(\left[\hat{H},\rho\right]\hat{A}\right) + \sum_{j} \gamma_{j} \mathrm{tr}\left(\mathcal{D}\left(\hat{O}_{j},\hat{P}_{j}\right)[\rho]\hat{A}\right)$$
(22)

We call the trace containing \hat{H} the "Hamiltonian trace", and those containing \mathcal{D} the "dissipator traces". For arbitrary Hamiltonian and dissipator operators, they are given by

$$\operatorname{tr}\left(\left[\hat{H},\rho\right]\hat{A}\right) = \left\langle\left[\hat{A},\hat{H}\right]\right\rangle$$
(23)

and

$$\operatorname{tr}\left(\mathcal{D}\left(\hat{O}_{j},\hat{P}_{j}\right)\left[\rho\right]\hat{A}\right) = \frac{1}{2}\left\langle \left[\hat{P}_{j}^{\dagger},\hat{A}\right]\hat{O}_{j}\right\rangle + \frac{1}{2}\left\langle \hat{P}_{j}^{\dagger}\left[\hat{A},\hat{O}_{j}\right]\right\rangle.$$
 (24)

3. Package Description

The core workflow of pyBoLaNO is shown in Fig. 1. We start with a brief overview of the SymPy objects relevant to our package. Then, we describe the workflow of the package's main functions: ops, normal_ordering / NO, NO_commutator, and LME_expval_evo; as well as the multiprocessing configuration dictionary: mp_config.



Figure 1: Core functionalities of pyBoLaNO. The main functions are shown in blue, while the core utility functions are colored black. Meanwhile, the purple-colored mp_config is a Python variable. Red connectors show symbolic workflows, while green connectors show visual workflows. Dashed rectangles indicate functionalities that are not accessible by the user.

3.1. Relevant objects in SymPy

All algebraic expressions in SymPy are based on the Expr or AtomicExpr class. The former is used for expressions with arguments (composite expressions), such as Symbol. The latter is used for expressions without arguments (atomic expressions), such as Number. The tuple of arguments of an expression object can be accessed from its args attribute. For AtomicExpr, an empty tuple is returned.

Each mathematical expression returns the corresponding operation class: Add for summation, Mul for multiplication, Pow for exponentiation, and Commutator for commutator, to name a few. These are all composite objects and may have each other as arguments. Add and Mul may have any number of arguments. Meanwhile, the first argument of Pow is the base and the second argument is the exponent. The two arguments of Commutator are the entries for each slot in the bracket. For example, given that x=Symbol("x"), then x**2+2*x is constructed as Add(Pow(Symbol("x"), Integer(2)),Mul(Integer(2), Symbol("x"))). This representation can be accessed using the srepr function. We can also check whether a given expression is contained using the has method.

SymPy has some rules to represent algebraic expressions, specifically for ordering arguments. As a consequence, Commutator(A,B) (where A and B are placeholders) may be shown as Mul(Integer(-1),Commutator(B,A))

instead. There are also rules to simplify the representation. For example, A+B+C is represented by Add(A,B,C) instead of Add(Add(A,B),C). For Commutator, SymPy automatically takes any non-commuting scalar out of the bracket and distributes the commutator when at least one of the arguments is an Add . pyBoLaNO makes the assumption that the input Commutator is represented as a linear combination of commutators.

The ladder operators are given by AnnihilateBoson and CreateBoson. One can be converted to the other using Dagger, which takes the Hermitian conjugate of its input. These, along with Commutator, are not objects imported by import sympy. Instead, the user has to run import sympy.physics.secondquant. pyBoLaNO recognizes the ladder operators and the commutator by these classes. SymPy has a class for normal ordering, namely NO, but it is not recognized by our package. Table 1 specifies SymPy objects that behave well in pyBoLaNO.

3.2. Constructing the ladder operators

Instead of calling AnnihilateBoson(k) and CreateBoson(k) from sympy.physics.secondquant, where k is the subscript, we recommend using pyBoLaNO 's ops:

ops(k)

that can be used to construct the annihilation and creation operators. The subscript k is optional and can be

SymPy object	Description
Number	Atomic expression for numbers (Integer, Float, Rational). The args attribute is an empty tuple.
Symbol	Symbol expression. Useful for variables. The args attribute is a tuple containing the string input to the object's constructor.
AnnihilateBoson	Bosonic annihilation operator. The LATEX output is \hat{b}_j . Here j is a Symbol or Integer input to the class constructor. Inputting Integer other than 0 will raise an error in the printing process. This is likely a bug. The args attribute is a tuple containing the subscript.
CreateBoson	Bosonic creation operator. The LATEX output is \hat{b}_j^{\dagger} . Similar to AnnihilateBoson.
Commutator	Takes the commutator between its two inputs. The args attribute is a tuple containing both its arguments.
Add	The sum object. The args attribute is a tuple of its summands.
Mul	The product object. The args attribute is a tuple of its factors.
Pow	The exponentiation object. The args attribute is a tuple (b,e) of its base and exponent.

Table 1: SymPy objects compatible with $\tt pyBoLaNO$. SymPy is ver. 1.13.3 .

Symbol or any Python object convertible to a string. This function preprocesses k into Symbol, which is the most compatible with AnnihilateBoson and CreateBoson. In the current version of SymPy, it also avoids the printing error that occurs when nonzero integers are used as subscripts (which is likely a bug).

3.3. Normal ordering

The normal ordering of a polynomial **q** in ladder operators is done by the **normal_ordering** function, with syntax.

normal_ordering(q)

We also offer a shorthand:

NO(q)

Blasiak's formulae, given by Eq. (19), are implemented in the _NO_Blasiak function, which evaluates the normal ordering of the given monomial in the ladder operators. Extra algorithms are executed to handle multipartite descriptions. The general workflow is described below.

- 1. Make a list of summands of q.
- 2. Make each summand of **q** into a list of factors, separated by the ladder operator subscripts. Scalars are put in a separate group.
- 3. Normal-order each factor using _NO_Blasiak .
- Multiply the sums obtained by normal-ordering each factor to give the almost-normal-ordered summand of q.
- 5. Sum all the almost-normal-ordered summands of **q** to give the almost-normal-ordered **q**.
- 6. Due to how SymPy handles noncommuting objects, the Mul of factors obtained in step 4 are yet to be normal-ordered. Instead, each term goes like $\dots \hat{b}_{j}^{\dagger a} \hat{b}_{j}^{b} \hat{b}_{k\neq j}^{\dagger c} \hat{b}_{k\neq j}^{d} \dots$ Do the final sorting as follows for each summand:
 - (a) Collect scalars into one list.
 - (b) Collect creation operators as dictionary entries whose keys are their subscripts.
 - (c) Collect annihilation operators similarly.
 - (d) Sort both dictionaries by key, then return a list of their values.
 - (e) Multiply all items in each list to give a Mul of scalars, creation operators, and annihilation operators.
 - (f) Multiply the three resulting Mul objects to give the fully normal-ordered summand.

Sum all summands and return the resulting Add.

3.4. Normal-ordered commutators

The normal ordering of a commutator between two polynomials **A** and **B** in ladder operators can be evaluated using the **do_commutator** function. The syntax is given by

do_commutator(A, B)

This function simply calls normal_ordering(A*B-B*A).

3.5. Expectation value evolution in the Lindblad master equation framework

Equation (22) can be evaluated by calling LME_expval_evo. The syntax is

LME_expval_evo(H, D, A, hbar_is_one)

Here, **H** is the Hamiltonian of the system, written as a SymPy expression. Meanwhile, **D** is a list that specifies the dynamics of the open system. Each element in **D** is a list **[gamma_j, 0_j, P_j]** containing the process rate γ_j and the operators \hat{O}_j , \hat{P}_j defining the dissipator. If $\hat{O}_j = \hat{P}_j$, the user may omit the third element from the list. If no dissipation is present, the user may input an empty list, **D**=[]. The operator corresponding to the quantity whose expectation value evolution is calculated is input to **A**. Finally, **hbar_is_one** is a Boolean to set $\hbar = 1$. This option is True by default. We note that Symbol("hbar") is used instead of **sympy.physics.quantum.hbar** to ensure that it behaves well with the package functionalities. The following happens when this function is called.

- 1. Call Hamiltonian_trace(H, A) to calculate the Hamiltonian trace with Eq. (23) in normal-ordered form. Add the result to the output.
- 2. For each entry in D, call dissipator_trace(O_j,P_j,A) to calculate the dissipator trace given by Eq. (24) in normal-ordered form. Multiply the result by the corresponding process rate gamma_j and add to the output.
- 3. Return the output Add.

The output is generally a sum that contains _expval objects. The _expval class wraps the output operators inside bra-kets. The class inherits SymPy's Symbol class and is used only for result visualization.

3.6. Mutiprocessing configurations

Multiprocessing is used in **normal_ordering** to handle each summand of the input. The user may control the multiprocessing behavior by modifying the **mp_config** variable after importing the package. The syntax is

```
mp_config["enable"] = True
mp_config["num_cpus"] = os.cpu_count()
mp_config["min_num_args"] = 2
```

where the values assigned above are the default values of the package. Here, "enable" specifies whether multiprocessing is enabled, "num_cpus" specifies the number of CPU threads to use, and "min_num_args" specifies the minimum number of summands mentioned above (arguments of the Add object). Any integer less than 2 will cause the package to set mp_config["min_num_args"]=2.

4. Some Usecases of the Package

The examples provided in this section are available in the package tutorial on our GitHub repository, which uses Jupyter Notebook to render the IAT_EX output. The following snippets show an exemplary setup of the package.

```
import pybolano as bl
bl.mpconfig["num_cpus"] = 4
```

import sympy as sm
latex = sm.latex # to print as LaTeX code.

b, bd = bl.ops()
b_1, bd_1 = bl.ops(1)
b_2, bd_2 = bl.ops(sm.Symbol("2"))

print(latex(
b_1
))

Out: b_{1}

4.1. normal_ordering

print(latex(
bl.normal_ordering(b * bd * b)
))

Out: b_{} + {b^\dagger_{}} b_{}^{2}

Multipartite input:

```
print(latex(
bl.normal_ordering(b_2 * b_1 * bd_2**2 * bd_1)
))
Out: 2 {b^\dagger_{1}} {b^\dagger_{2}} b_{1} +

→ {b^\dagger_{1}} {b^\dagger_{2}}^{2} b_{1}

→ b_{2} + 2 {b^\dagger_{2}} +

→ {b^\dagger_{2}}^{2} b_{2}
```

Polynomial input:

Input with symbols:

x = sm.Symbol("x")

print(latex(
bl.normal_ordering(x * b_1 * x**2 * bd_1**2)
))

```
Out: 2 x^{3} {b^dagger_{1}} + x^{3}

\rightarrow {b^dagger_{1}}^{2} b_{1}
```

4.2. NO_commutator

print(latex(
bl.NO_commutator(A, B)
))

Out: - b_{}

Multipartite input:

 $A = bd_1*bd_2$ $B = b_1*b_2$

print(latex(
bl.NO_commutator(A, B)
))

Out: -1 - {b^\dagger_{1}} b_{1} - \rightarrow {b^\dagger_{2}} b_{2}

Polynomial input:

```
A = b_1 + 2*b_2**2
B = bd_1**3 + 2*bd_2*b_2
```

print(latex(
bl.NO_commutator(A, B)
))

Out: 8 b_{2}^{2} + 3 {b^dagger_{1}}^{2}

Input with symbols:

x = sm.Symbol("x")
A = x*b_1
B = x**(0.5)*bd_1*b

print(latex(
bl.NO_commutator(A, B)
))

Out: x^{1.5} b_{}

4.3. LME_expval_evo

Now we consider some evolution equations in the literature as a way to validate this package, highlighting its potential usage for describing various quantum systems. The following examples also illustrate problems where it is desirable to compute the evolution of the expectation values of some observables for which pyBoLaNO may be useful.

4.3.1. The quantum simple harmonic oscillator

It is well-known that the evolution of the expected value of the annihilation operator \hat{b} in a simple harmonic Hamiltonian $\hat{H} = \hbar \omega_0 \left(\hat{b}^{\dagger} \hat{b} + 1/2 \right)$ is given by

$$\frac{\mathrm{d}\left\langle\hat{b}\right\rangle}{\mathrm{d}t} = -i\hbar\omega_0\left\langle\hat{b}\right\rangle \tag{25}$$

Here is how it is obtained with pyBoLaNO :

```
hbar, omega_0 = sm.symbols(r"hbar omega_0")
b, bd = bl.ops()
H = hbar*omega_0*bd*b
D = []
A = b
print(latex(
bl.LME_expval_evo(H, D, A, hbar_is_one=False)
))
Out: \frac{d}{d t} {\left\langle b_{}
```

Additionally, we can show that the energy is conserved, i.e. that

$$\frac{\mathrm{d}\left\langle \hat{b}^{\dagger}\hat{b}\right\rangle }{\mathrm{d}t} = 0 \tag{26}$$

Assuming the variables from the previous code block, we have

A = bd*b

print(latex(
bl.LME_expval_evo(H, D, A)
))

```
Out: frac{d}{d t} {\left[ \frac{1}{d t} \right]} 
\rightarrow {b^dagger_{} b_{} = 0
```

4.3.2. The quantum Rayleigh oscillator

A limit cycle can be defined as a stable oscillation to which the system is always attracted [27]. This is a consequence of nonlinearity and can be observed in nonlinear oscillators. The Rayleigh oscillator is one such exemplary model. We consider the evolution of the expected phase point $\langle \hat{b} \rangle$ for a quantum Rayleigh oscillator, a nonlinear quantum oscillator exhibiting a quantum limit cycle. From Ref. [10], the equation of motion is specified by (with $\hbar = 1$ and $\hat{b} = [\hat{x} + i\hat{p}]/2$, cf. Eqs. [14–16] of Ref. [10]):

$$\hat{H} = \omega_0 \hat{b}^{\dagger} \hat{b} + i \frac{\mu}{12} \left(\hat{b}^{\dagger} \hat{b}^3 - \hat{b}^{\dagger 3} \hat{b} \right) + i \frac{\mu}{24} \left(\hat{b}^4 - \hat{b}^{\dagger 4} \right) - i \frac{\mu \left(q_0^2 - 1 \right)}{4} \left(\hat{b}^2 - \hat{b}^{\dagger 2} \right)$$
(27)

and

....

$$\gamma_1 = \mu \left(q_0^2 - 1 \right), \quad \hat{O}_1 = \hat{P}_1 = \hat{b}^{\dagger}$$
 (28a)

$$\gamma_2 = \frac{3\mu}{4}, \quad \hat{O}_2 = \hat{P}_2 = \hat{b}^2$$
 (28b)

$$\gamma_3 = \mu, \quad \hat{O}_3 = \hat{P}_3 = \hat{b}^{\dagger}\hat{b} - \frac{b^{\dagger 2}}{2}$$
 (28c)

where μ is the nonlinearity parameter and q_0 is one of the Rayleigh parameters. The evolution of $\langle \hat{b} \rangle$ is governed by (cf. Eqs. [13, 17–25] of Ref. [10])

$$\frac{\mathrm{d}\left\langle\hat{b}\right\rangle}{\mathrm{d}t} = -i\omega_{0}\left\langle\hat{b}\right\rangle + \frac{\mu}{2}\left(q_{0}^{2}-1\right)\left[\left\langle\hat{b}\right\rangle + \left\langle\hat{b}^{\dagger}\right\rangle\right] \\ -\frac{\mu}{6}\left[\left\langle\hat{b}^{3}\right\rangle + \left\langle\hat{b}^{\dagger3}\right\rangle\right] - \frac{\mu}{2}\left[\left\langle\hat{b}^{\dagger}\hat{b}^{2}\right\rangle + \left\langle\hat{b}^{\dagger2}\hat{b}\right\rangle\right]$$
(29)

Here is how Eq. (29) can be obtained with pyBoLaNO :

```
omega_0, mu, q_0 = sm.symbols(r"omega_0 mu
→ q_0")
b, bd = bl.ops()
H = omega_0*bd*b \
    + sm.I*mu/12 * (bd*b**3 - bd**3*b) \
    + sm.I*mu/24 * (b**4-bd**4) \
    - sm.I*mu*(q_0**2-1)/4 * (b**2-bd**2)
D = [[mu*(q_0**2-1), bd],
    [3*mu/4, b**2],
    [mu, bd*b-bd**2/2]]
A = b
print(latex(
bl.LME_expval_evo(H, D, A)
))
```

```
Out: \frac{d}{d t} {\left\langle b_{}
     \left|\right| = \left| \frac{q_{0}^{2}}{2} \right|
\hookrightarrow
     {\left\langle b_{} \right\rangle}}{2} +
\hookrightarrow
     frac{mu q_{0}^{2} {\left}agle}
\hookrightarrow
     {b^\dagger_{}} \right\rangle}}{2} -
\hookrightarrow
     \frac{\mu {\left\langle b_{}
\hookrightarrow
     \right\rangle}}{2} - \frac{\mu
\hookrightarrow
     {\left\langle b_{}^{3} \right\rangle}}{6}
\hookrightarrow
     - \frac{\mu {\left\langle {b^\dagger_{}}
\hookrightarrow
     \right\rangle}}{2} - \frac{\mu
\hookrightarrow
     {\left\langle {b^{dagger_{}} b_{}^{2}
\hookrightarrow
     \right\rangle}}{2} - \frac{\mu
\hookrightarrow
    {\left\langle {b^{dagger_{}}^{2} b_{}
\hookrightarrow
    \right\rangle}}{2} - \frac{\mu
\hookrightarrow
     {\left\langle {b^\dagger_{}}^{3}
\hookrightarrow
     \right\rangle}}{6} - i \omega_{0}
\hookrightarrow
     {\left\langle b_{} \right\rangle}}
```

4.3.3. A bipartite quantum battery with quadratic driving

A quantum battery is a quantum system that can store energy and whose energy can be harvested for useful work [12]. Recently, Downing and Ukhtary [12] proposed a setup for a bipartite quantum battery. The system contains a charger and a holder coupled to each other, with a short quadratic pulse driving the charger. The equation of motion is specified by (with $\hbar = 1$, cf. Eqs. [1–6] of Ref. [12])

$$\hat{H} = \omega_c \hat{b}_c^{\dagger} \hat{b}_c + \omega_h \hat{b}_h^{\dagger} \hat{b}_h + g \left(\hat{b}_c^{\dagger} \hat{b}_h + \hat{b}_h^{\dagger} \hat{b}_c \right) + \frac{\Omega}{2} \delta(t) \left(\hat{b}_c^{\dagger 2} + \hat{b}_c^2 \right)$$
(30)

and

$$\gamma_1 = \gamma, \quad \hat{O}_1 = \hat{O}_1 = \hat{b}_c \tag{31}$$

where g is the coupling strength, Ω is the pulse strength, and $\delta(t)$ is the Dirac delta function. The quantities $\left\langle \hat{b}_{c}^{\dagger}\hat{b}_{c}\right\rangle$ and $\left\langle \hat{b}_{h}^{\dagger}\hat{b}_{h}\right\rangle$ are proportional to the energy of the charger and the holder, respectively. After t = 0, the driving is off, and they evolve according to (cf. Eq. [12] of Ref. [12])

$$\frac{\mathrm{d}\left\langle \hat{b}_{c}^{\dagger}\hat{b}_{c}\right\rangle}{\mathrm{d}t} = -\gamma\left\langle \hat{b}_{c}^{\dagger}\hat{b}_{c}\right\rangle - ig\left[\left\langle \hat{b}_{c}^{\dagger}\hat{b}_{h}\right\rangle - \left\langle \hat{b}_{h}^{\dagger}\hat{b}_{c}\right\rangle\right] \quad (32a)$$

$$\frac{\mathrm{d}\left\langle b_{h}^{\dagger}b_{h}\right\rangle}{\mathrm{d}t} = ig\left[\left\langle \hat{b}_{c}^{\dagger}\hat{b}_{h}\right\rangle - \left\langle \hat{b}_{h}^{\dagger}\hat{b}_{c}\right\rangle\right] \tag{32b}$$

Here are the same equations obtained using our package:

omega_c, omega_h, g, gamma = \
 sm.symbols(r"omega_c omega_h g gamma")
b_c, bd_c = bl.ops("c")
b_h, bd_h = bl.ops("h")

```
H = omega_c * bd_c*b_c \setminus
     + omega_h * bd_h*b_h \
     + g*(bd_c*b_h + bd_h*b_c)
D = [[gamma, b_c]]
A = bd_c*b_c
print(latex(
bl.LME_expval_evo(H,D,A)
))
Out: \frac{d}{d t} {\left\langle
    {b^\dagger_{\mathtt{\text{c}}}}
\hookrightarrow
     b_{\mathtt{\text{c}}} \right\rangle} = - i
\hookrightarrow
     g {\left\langle
\hookrightarrow
     {b^\dagger_{\mathtt{\text{c}}}}
\hookrightarrow
     b_{\mathtt{\text{h}}} \right\rangle} + i g
\hookrightarrow
     {\left\langle
\hookrightarrow
     b^{dagger_{\mathrm{h}}}
\hookrightarrow
     b_{\mathtt{\text{c}}} \right\rangle} -
\hookrightarrow
     \gamma {\left\langle
\hookrightarrow
     {b^\dagger_{\mathtt{\text{c}}}}
\hookrightarrow
```

Assuming the variables from the previous block,

 $A = bd_h*b_h$ print(latex(bl.LME_expval_evo(H,D,A))) Out: \frac{d}{d t} {\left\langle {b^\dagger_{\mathtt{\text{h}}}} \hookrightarrow b_{\mathtt{\text{h}}} \right\rangle} = i g \hookrightarrow {\left\langle \hookrightarrow {b^\dagger_{\mathtt{\text{c}}}} \hookrightarrow b_{\mathtt{\text{h}}} \right\rangle} - i g \hookrightarrow {\left\langle \hookrightarrow {b^\dagger_{\mathtt{\text{h}}}} \hookrightarrow b_{\mathtt{\text{c}}} \right\rangle} \hookrightarrow

4.3.4. A PT-symmetric trimer of harmonic oscillators

Bender and Boettcher [28] show that a Hamiltonian does not need to be Hermitian to have a real, nonnegative spectrum. The parity-time (\mathcal{PT}) symmetry can be viewed as a generalization to the hermicity condition—this validates the non-Hermitian quantum mechanics formulation. In non-Hermitian quantum mechanics, exceptional points (EPs) mark the crossover between the unbroken and broken \mathcal{PT} phase. Exceptional points are useful. For example, at its EPs, a system may gain enhanced sensitivity, making it desirable for sensing applications [29]. As another example, it is observed that the fluorescence rates of certain single-photon sources reach their quantum limit upon crossing an EP [30].

Downing and Saroka [16] formulate a simple model of short oligomer chains of harmonic oscillators with a \mathcal{PT} symmetric Hamiltonian in the Lindblad master equation framework, showing the emergence of EPs. We consider a trimer system whose dynamics is specified by ($\hbar = 1$, cf. Eqs. [4, 5a, 5b, 10] of Ref. [16])

$$\hat{H} = \left(\omega_0 + i\frac{\kappa}{2}\right)\hat{b}_1^{\dagger}\hat{b}_1 + \omega_0\hat{b}_2^{\dagger}\hat{b}_2 + \left(\omega_0 - i\frac{\kappa}{2}\right)\hat{b}_3^{\dagger}\hat{b}_3 + g\left(\hat{b}_1^{\dagger}\hat{b}_2 + \hat{b}_2^{\dagger}\hat{b}_3 + \text{h.c.}\right)$$
(33)

and

$$\gamma_1 = \gamma_1, \quad \dot{O}_1 = \dot{P}_1 = \dot{b}_1 \tag{34a}$$

$$\gamma_2 = \gamma_2, \quad \hat{O}_2 = \hat{P}_2 = \hat{b}_2 \tag{34b}$$

$$\gamma_3 = \gamma_3, \quad O_3 = P_3 = b_3 \tag{34c}$$

$$\gamma_4 = p_1, \quad O_4 = P_4 = b_1'$$
 (34d)
 $\gamma_4 = r_1, \quad O_4 = P_4 = b_1'$ (34d)

$$\gamma_5 = p_2, \quad O_5 = P_5 = b_2' \tag{34e}$$

$$\gamma_6 = p_3, \quad O_6 = P_6 = b_3^{\dagger}$$
 (34f)

where ω_0 is the natural frequency of all oscillators, g is the coupling strength, and κ specifies both the gain rate of oscillator 1 and loss rate of oscillator 3. Meanwhile, γ_k and p_k , k = 1, 2, 3, specify the gain and loss rates from incoherent processes for the oscillator k. The non-Hermitian parts of \hat{H} are consequences of these processes. The evolution of $\left\langle \hat{b}_k^{\dagger} \hat{b}_k \right\rangle$ are given by (cf. Eqs.[9, 14–19] of Ref. [16])

$$\frac{\mathrm{d}\left\langle\hat{b}_{1}^{\dagger}\hat{b}_{1}\right\rangle}{\mathrm{d}t} = p_{1} - \left(\gamma_{1} - p_{1}\right)\left\langle\hat{b}_{1}^{\dagger}\hat{b}_{1}\right\rangle \qquad (35a)$$
$$- ig\left\langle\hat{b}_{1}^{\dagger}\hat{b}_{2}\right\rangle + ig\left\langle\hat{b}_{2}^{\dagger}\hat{b}_{1}\right\rangle \qquad (35b)$$
$$\frac{\mathrm{d}\left\langle\hat{b}_{2}^{\dagger}\hat{b}_{2}\right\rangle}{\mathrm{d}t} = p_{2} - \left(\gamma_{2} - p_{2}\right)\left\langle\hat{b}_{2}^{\dagger}\hat{b}_{2}\right\rangle \qquad (35b)$$

$$+ ig \left\langle \hat{b}_{1}^{\dagger} \hat{b}_{2} \right\rangle - ig \left\langle \hat{b}_{2}^{\dagger} \hat{b}_{3} \right\rangle$$
$$- ig \left\langle \hat{b}_{2}^{\dagger} \hat{b}_{1} \right\rangle + ig \left\langle \hat{b}_{3}^{\dagger} \hat{b}_{2} \right\rangle$$
$$\frac{\mathrm{d} \left\langle \hat{b}_{3}^{\dagger} \hat{b}_{3} \right\rangle}{\mathrm{d}t} = p_{3} - (\gamma_{3} - p_{3}) \left\langle \hat{b}_{3}^{\dagger} \hat{b}_{3} \right\rangle$$
$$+ ig \left\langle \hat{b}_{2}^{\dagger} \hat{b}_{3} \right\rangle - ig \left\langle \hat{b}_{3}^{\dagger} \hat{b}_{2} \right\rangle$$
(35c)

Here are the equations obtained using pyBoLaNO :

```
b_2, bd_2 = bl.ops(2)
b_3, bd_3 = bl.ops(3)
H = (\text{omega}_0 + \text{sm.I}*\text{kappa}/2)*bd_1*b_1 \setminus
    + omega_0*bd_2*b_2 \
    + (omega_0 - sm.I*kappa/2)*bd_3*b_3 \
    + g*(bd_1*b_2+bd_2*b_1 + bd_2*b_3 +
     \rightarrow bd_3*b_2)
D = [[gamma_1, b_1]],
      [gamma_2, b_2],
      [gamma_3, b_3],
      [p_1, bd_1],
      [p_2, bd_2],
      [p_3, bd_3]]
A = bd_1*b_1
print(latex(
bl.LME_expval_evo(H, D, A)
))
Out: \frac{d}{d t} {\left\langle
    {b^\dagger_{1}} b_{1} \right\rangle} = - i
    g {\left\langle {b^\dagger_{1}} b_{2}
\hookrightarrow
    \right\rangle} + i g {\left\langle
\hookrightarrow
    b^{dagger_{2}} b_{1} \quad -
\hookrightarrow
    \gamma_{1} {\left\langle {b^\dagger_{1}}
\hookrightarrow
    b_{1} \right\rangle} + p_{1} {\left\langle
    b^{dagger_{1}} b_{1} \ + \
\hookrightarrow
    p_{1}
\hookrightarrow
```

Assuming the variables from the previous block,

```
A = bd_2*b_2
print(latex(
bl.LME_expval_evo(H, D, A)
))
Out: \frac{d}{d t} {\left\langle
     {b^\dagger_{2}} b_{2} \right\rangle} = i g
\hookrightarrow
     {\left\langle {b^{dagger_{1}} b_{2}
 \hookrightarrow
     \right\rangle} - i g {\left\langle
 \hookrightarrow
     b^{dagger_{2}} b_{1} \quad right - i g
\hookrightarrow
     {\left\langle {b^\dagger_{2}} b_{3}
\hookrightarrow
     \right\rangle} + i g {\left\langle
 \hookrightarrow
     {b^\dagger_{3}} b_{2} \right\rangle} -
 \hookrightarrow
     \gamma_{2} {\left\langle {b^\dagger_{2}}
\hookrightarrow
     b_{2} \right\rangle} + p_{2} {\left\langle
 \hookrightarrow
     {b^\dagger_{2}} b_{2} \right\rangle} +
 \hookrightarrow
```

```
→ p_{2}
```

and

```
A = bd_3*b_3
print(latex(
bl.LME_expval_evo(H, D, A)
))
Out: \frac{d}{d t} {\left\langle
     {b^\dagger_{3}} b_{3} \right\rangle} = i g
\hookrightarrow
     {\left\langle {b^\dagger_{2}} b_{3}
\hookrightarrow
     \right\rangle} - i g {\left\langle
\hookrightarrow
     b^{dagger_{3}} b_{2} \
\hookrightarrow
     \gamma_{3} {\left\langle {b^\dagger_{3}}}
\hookrightarrow
    b_{3} \right\rangle} + p_{3} {\left\langle
\hookrightarrow
     {b^\dagger_{3}} b_{3} \right\rangle} +
\hookrightarrow
    p_{3}
\hookrightarrow
```

4.3.5. A pair of nonreciprocal driven-dissipative quantum resonators

So far, we have considered the cases where $\hat{O}_j = \hat{P}_j$. For the last example, we will discuss a system in which this is not the case.

The notion of reciprocity, i.e., when the interaction between two objects is identical if they are swapped, is encountered in various topics in physics, such as Newton's third law of motion, electromagnetic reciprocity [31], and scattering reciprocity [32]. The lack of reciprocity (in other words: nonreciprocity) can lead to interesting phenomena and practical advantages [17]. A physical model that gives rise to nonreciprocity in open quantum systems is given by Downing and Sturges [17]. The model consists of a pair of driven-dissipative quantum resonators, where the asymmetry arises from the relative phase difference between the coherent and incoherent couplings. The dynamics is described within the Lindblad master equation framework, with ($\hbar = 1$, cf. Eqs. [1, 2] of Ref. [17])

$$\hat{H} = \Delta \sum_{k=1,2} \hat{b}_k^{\dagger} \hat{b}_k + \Omega \left(\hat{b}_1 + \hat{b}_1^{\dagger} \right) + g \left(e^{i\theta} \hat{b}_1^{\dagger} \hat{b}_2 + e^{-i\theta} \hat{b}_2^{\dagger} \hat{b}_1 \right)$$
(36)

and

$$\gamma_1 = \gamma, \quad \hat{O}_1 = \hat{P}_1 = \hat{b}_1 \tag{37a}$$

$$\gamma_2 = \gamma, \quad O_2 = P_2 = b_2 \tag{37b}$$

$$\gamma_3 = \Gamma e^{i\phi}, \quad O_3 = b_2, \quad P_3 = b_1 \tag{37c}$$

$$\gamma_4 = \Gamma e^{-i\phi}, \quad O_4 = b_1, \quad P_4 = b_2$$
 (37d)

in the rotating reference frame of the laser driving oscillator 1. Here, Δ is the detuning of the oscillators with respect to the laser, Ω is the driving strength, g is the (coherent) coupling strength, and θ is the coupling phase. The third and fourth dissipators describe the incoherent coupling between the two oscillators sharing the same bath, whose rate is taken to be complex with amplitude $0 \leq \Gamma \leq \gamma$ and phase ϕ . The quantities $\langle \hat{b}_1 \rangle$ and $\langle \hat{b}_2 \rangle$ evolve according to (cf. Eqs. [3, 4] of Ref. [17])

$$\frac{\mathrm{d}\left\langle\hat{b}_{1}\right\rangle}{\mathrm{d}t} = -i\left(\Delta - i\frac{\gamma}{2}\right)\left\langle\hat{b}_{1}\right\rangle - \left(ige^{i\theta} + \frac{\Gamma e^{i\phi}}{2}\right)\left\langle\hat{b}_{2}\right\rangle - i\Omega$$
(38a)
$$\frac{\mathrm{d}\left\langle\hat{b}_{2}\right\rangle}{\mathrm{d}t} = -\left(ige^{-i\theta} + \frac{\Gamma e^{-i\phi}}{2}\right)\left\langle\hat{b}_{1}\right\rangle - i\left(\Delta - i\frac{\gamma}{2}\right)\left\langle\hat{b}_{2}\right\rangle$$
(38b)

Here are Eqs. (38a) and (38b) obtained using the package:

```
Delta, Omega, g, theta = sm.symbols("Delta
\hookrightarrow Omega g theta")
gamma, Gamma, phi = sm.symbols("gamma Gamma
→ phi")
b_1, bd_1 = bl.ops(1)
b_2, bd_2 = bl.ops(2)
H = Delta*(bd_1*b_1 + bd_2*b_2) \setminus
    + Omega*(b_1+bd_1) \setminus
    + g*(sm.E**(sm.I*theta)*bd_1*b_2 +
     \rightarrow sm.E**(-sm.I*theta)*bd_2*b_1)
D = [[gamma, b_1]],
      [gamma, b_2],
      [Gamma*sm.E**(sm.I*phi), b_2, b_1],
      [Gamma*sm.E**(-sm.I*phi), b_1, b_2]]
A = b_1
print(latex(
bl.LME_expval_evo(H, D, A)
))
Out: \frac{d}{d t} {\left\langle b_{1}}
   \right\rangle} = - i \Delta {\left\langle
\hookrightarrow
\rightarrow b_{1} \right\rangle} - \frac{\Gamma
   {\left\langle b_{2} \right\rangle} e^{i
\hookrightarrow
    \phi}}{2} - i \Omega - i g {\left\langle
\hookrightarrow
    b_{2} \right\rangle} e^{i \theta} -
\hookrightarrow
    \frac{\gamma {\left\langle b_{1}}
\hookrightarrow
```

```
\rightarrow \right\rangle}}{2}
```

Assuming variables from the previous block,

A = b_2
print(latex(
bl.LME_expval_evo(H, D, A)
))

```
Out: \frac{d}{d t} {\left\langle b_{2}

→ \right\rangle} = - i \Delta {\left\langle

→ b_{2} \right\rangle} - \frac{\Gamma

→ {\left\langle b_{1} \right\rangle} e^{-i

→ \phi}}{2} - i g {\left\langle b_{1}}

→ \right\rangle} e^{-i \theta} -

→ \frac{\gamma {\left\langle b_{2}}

→ \right\rangle}}{2}
```

5. Performance

Since it is implemented for each summand in the input expression, the speedup gained by using multiprocessing is approximately linear (the evaluation of each summand may take different durations). Meanwhile, the evaluation of a single monomial is optimal, since explicit formulae are used.

6. Conclusion

We have developed a symbolic package pyBoLaNO for the normal ordering of polynomials in bosonic ladder operators utilizing Blasiak's formulae. By extension, it allows the user to obtain the normal-ordered equivalent of a commutator between two polynomials in bosonic ladder operators, as well as the normal-ordered expectation value evolution equation for a system described in the Lindblad master equation framework. We have described the package workflow in detail and provided the syntaxes for the core functionalities. We have exhibited some examples of use by taking recent results from the literature which also serve to validate the package. We have briefly discussed the computational cost of the normal ordering algorithm. This package is aimed at quantum physics theorists who desire fast, error-free normal ordering for bosonic ladder algebra, in particular when dealing with expectation value evolution in the Lindblad master equation framework. We welcome suggestions and constructive criticism to improve this package through its public repository.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

The package's source code and instructions for installation are publicly available at https://github.com/hendry 24/pyBoLaNO. The code used for Section 4 is compiled into a Jupyter Notebook available at https://github.com /hendry24/pyBoLaNO/blob/main/tutorial.ipynb.

Acknowledgments

H.M.L. is supported by a research assistantship from the BRIN Directorate for Talent Management. The authors acknowledge Quasi Lab and Mahameru BRIN for their mini-cluster and HPC facilities.

References

- D. J. Griffiths, D. F. Schroeter, Introduction to quantum mechanics, 3rd Edition, Cambridge University Press, 2018.
- [2] T. Lancaster, S. Blundell, Quantum field theory for the gifted amateur, 1st Edition, Oxford Univ. Press, 2014.
- [3] C. C. Gerry, P. L. Knight, Introductory quantum optics, Cambridge University Press, 2005.
- [4] A. Meurer, C. P. Smith, M. Paprocki, O. Čertík, S. B. Kirpichev, M. Rocklin, A. Kumar, S. Ivanov, J. K. Moore, S. Singh, T. Rathnayake, S. Vig, B. E. Granger, R. P. Muller, F. Bonazzi, H. Gupta, S. Vats, F. Johansson, F. Pedregosa, M. J. Curry, A. R. Terrel, Š. Roučka, A. Saboo, I. Fernando, S. Kulal, R. Cimrman, A. Scopatz, SymPy: symbolic computing in Python, PeerJ Comput. Sci. 3 (2017) e103.
- [5] M. Schlosshauer, Decoherence and the quantum-to-classical transition, Springer, 2007.
- [6] H.-P. Breuer, F. Petruccione, The theory of open quantum systems, Oxford University Press, 2002.
- [7] J. Johansson, P. Nation, F. Nori, QuTiP: An open-source Python framework for the dynamics of open quantum systems, Comput. Phys. Commun. 183 (8) (2012) 1760.
- [8] J. Johansson, P. Nation, F. Nori, QuTiP 2: A Python framework for the dynamics of open quantum systems, Comput. Phys. Commun. 184 (4) (2013) 1234.
- [9] S. Krämer, D. Plankensteiner, L. Ostermann, H. Ritsch, QuantumOptics.jl: A Julia framework for simulating open quantum systems, Comput. Phys. Commun. 227 (2018) 109.
- [10] A. Chia, L. C. Kwek, C. Noh, Relaxation oscillations and frequency entrainment in quantum mechanics, Phys. Rev. E 102 (4) (2020) 042213.
- [11] Y. Shen, W.-K. Mok, C. Noh, A. Q. Liu, L.-C. Kwek, W. Fan, A. Chia, Quantum synchronization effects induced by strong nonlinearities, Phys. Rev. A 107 (5) (2023) 053713.
- [12] C. A. Downing, M. S. Ukhtary, Hyperbolic enhancement of a quantum battery, Phys. Rev. A 109 (5) (2024) 052206.
- [13] D. O. Krimer, M. Zens, S. Rotter, Critical phenomena and nonlinear dynamics in a spin ensemble strongly coupled to a cavity. I. semiclassical approach, Phys. Rev. A 100 (1) (2019) 013855.
- M. Zens, D. O. Krimer, S. Rotter, Critical phenomena and nonlinear dynamics in a spin ensemble strongly coupled to a cavity. II. semiclassical-to-quantum boundary, Phys. Rev. A 100 (1) (2019) 013856.
- [15] B. Ahmadi, P. Mazurek, P. Horodecki, S. Barzanjeh, Nonreciprocal quantum batteries, Phys. Rev, Lett. 132 (21) (2024) 210402.
- [16] C. A. Downing, V. A. Saroka, Exceptional points in oligomer chains, Commun. Phys. 4 (1) (2021) 254.
- [17] C. A. Downing, T. J. Sturges, Directionality between drivendissipative resonators, Europhys. Lett. 140 (3) (2022) 35001.
- [18] C. A. Downing, A. Vidiella-Barranco, Parametrically driving a quantum oscillator into exceptionality, Sci. Rep. 13 (1) (2023) 11004.
- [19] L. Ben Arosh, M. C. Cross, R. Lifshitz, Quantum limit cycles and the rayleigh and van der pol oscillators, Phys. Rev. Res. 3 (2021) 013130.
- [20] E. Amitai, M. Koppenhöfer, N. Lörch, C. Bruder, Quantum effects in amplitude death of coupled anharmonic self-oscillators, Phys. Rev. E 97 (2018) 052203.
- [21] F. Minganti, A. Miranowicz, R. W. Chhajlany, F. Nori, Quantum exceptional points of non-hermitian hamiltonians and liouvillians: The effects of quantum jumps, Phys. Rev. A 100 (6) (2019) 062131.
- [22] G. Chimczak, A. Kowalewska-Kudłaszyk, E. Lange, K. Bartkiewicz, J. Peřina, The effect of thermal photons on exceptional points in coupled resonators, Sci. Rep. 13 (1) (2023) 5859.
- [23] D. Farina, G. M. Andolina, A. Mari, M. Polini, V. Giovannetti, Charger-mediated energy transfer for quantum batteries: An open-system approach, Phys. Rev. B 99 (3) (2019) 035421.
- [24] G.-Q. Zhang, Z. Chen, W. Xiong, C.-H. Lam, J. Q. You, Paritysymmetry-breaking quantum phase transition via parametric

drive in a cavity magnonic system, Phys. Rev. B 104 (6) (2021) 064423.

- [25] P. Blasiak, Combinatorics of boson normal ordering and some applications (2005). arXiv:quant-ph/0507206.
- [26] M. A. Méndez, P. Blasiak, K. A. Penson, Combinatorial approach to generalized bell and stirling numbers and boson normal ordering problem, J. Math. Phys. 46 (8) (2005) 083511.
- [27] A. Pikovsky, M. Rosenblum, J. Kurths, Synchronization: a universal concept in nonlinear sciences, Cambridge University Press, 2001.
- [28] C. M. Bender, S. Boettcher, Real spectra in non-hermitian hamiltonians having *PT* symmetry, Phys. Rev. Lett. 80 (24) (1998) 5243.
- [29] J. Wiersig, Prospects and fundamental limits in exceptional point-based sensing, Nat. Commun. 11 (1) (2020) 2454.
- [30] W. Zhou, J. Liu, J. Zhu, D. Gromyko, C. Qiu, L. Wu, Exceptional points unveiling quantum limit of fluorescence rates in non-hermitian plexcitonic single-photon sources, APL Quantum 1 (1) (2024) 016110.
- [31] C. Caloz, A. Alù, S. Tretyakov, D. Sounas, K. Achouri, Z.-L. Deck-Léger, Electromagnetic nonreciprocity, Phys. Rev. Appl. 10 (4) (2018) 047001.
- [32] L. Deák, T. Fülöp, Reciprocity in quantum, electromagnetic and other wave scattering, Ann. of Phys. 327 (4) (2012) 1050.