

Reentrant topological phases and spin density wave induced by 1D moiré potentials

Guo-Qing Zhang,¹ Ling-Zhi Tang,^{2,*} L. F. Quezada,^{1,3} Shi-Hai Dong,^{1,3,†} and Dan-Wei Zhang^{4,5,‡}

¹*Research Center for Quantum Physics, Huzhou University, Huzhou 313000, P. R. China*

²*Quantum Science Center of Guangdong-Hong Kong-Macao Greater Bay Area (Guangdong), Shenzhen 518045, China*

³*Laboratorio de Ciencias de la Información Cuántica, Centro de Investigación en Computación, Instituto Politécnico Nacional, UPALM, 07700, Ciudad de México, México*

⁴*Key Laboratory of Atomic and Subatomic Structure and Quantum Control (Ministry of Education), Guangdong Basic Research Center of Excellence for Structure and Fundamental Interactions of Matter, South China Normal University, Guangzhou 510006, China*

⁵*Guangdong Provincial Key Laboratory of Quantum Engineering and Quantum Materials, School of Physics, South China Normal University, Guangzhou 510006, China*

(Dated: January 6, 2025)

Exotic topological and correlated phases induced by moiré potentials have recently been explored in 2D twisted bilayer graphene systems. Here we investigate spin-1/2 interacting fermionic atoms in the 1D moiré optical superlattice formed by the superposition of two commensurate potentials. We reveal the reentrant topological phases and periodic-moiré-spin density wave (PM-SDW) of the ground states in both non-interacting and interacting cases. The reentrant topological phases are driven by the moiré potential with the sequent trivial-topological-trivial-topological-trivial transition. The critical exponent of topological transitions and the localization properties of the single-particle eigenstates are studied. The PM-SDW order of the many-body ground state is contributed from the moiré potential, and is enhanced by the on-site interaction but suppressed by the nearest-neighbor interaction. Our results enrich the topological physics with multiple transitions and spin-density orders in 1D moiré systems, which might be experimentally realized with ultracold atoms in tunable optical lattices.

INTRODUCTION

Topological phases characterized by unique electronic properties have become an important research field in condensed-matter physics [1–6] and artificial systems [7–17]. Topological phases are usually protected by certain symmetries and are robust against local perturbations. The correspondence of the topological invariants of the insulating bulk states and the in-gap edge states characterizes the novelty of topological phases. Interactions can enrich topological phases, such as the fractional quantum Hall effects [18] and topological Mott insulators [19–27]. These topological properties have potential applications in fault-tolerant quantum computation [28–30] and spintronics devices [31–33]. Meanwhile, quantum simulations of topological matters have achieved great progress, such as the realizations of the 1D topological insulator model [34–36], 2D Harper-Hofstadter [37] and Haldane models [38], and for the observations of chiral edge states [39, 40] with ultracold atoms in optical lattices.

On the other hand, 2D moiré systems formed by stacking two monolayer materials vertically and a tunable twist angle have recently revealed exotic physics, such as flat bands [41–47], moiré excitons [48, 49], interlayer ferromagnetism [50], and correlated topological states [51–56]. Moiré superlattices that emerge from the

misaligned placement of two periodic lattices exhibit a new lattice periodicity, which can be finite or infinite, corresponding to the commensurate or incommensurate superlattices, respectively. This provides a tunable way to engineer the band structures in moiré materials. The band width in moiré systems can be made extremely narrow, which enable to stabilize strongly-correlated superconductivity and topological insulators [57–61]. By reducing 2D twisted graphene to 1D incommensurate carbon nanotubes, the moiré potentials can emerge and alter the band structure and characteristics of the system [62, 63]. Alternatively, the moiré analogs can be achieved by finite incommensurate 1D lattice potentials [64, 65] or finite coupled resonators with different modulation lengths [66]. It has been revealed that the incommensurability can enable quasi-fractal charge-density waves (CDWs) in 1D narrow-band moiré systems [65]. However, it remain unclear whether the 1D commensurate moiré potentials can induce topological phases and spin density orders.

In this work, we address this question by exploring a 1D spin-1/2 fermionic optical lattice with a commensurate moiré potential and interatomic interactions. In the non-interacting limit, the system Hamiltonian under the periodic boundary condition (PBC) is analytically solved in the momentum space with reduced moiré Brillouin zones (MBZs). Under the open boundary condition (OBC), physical characteristics are solved by the exact diagonalization (ED) method. The topological characteristics of this 1D system are the winding number defined under the PBC [67] and its generalization to the real space under the OBC [68]. We find that the trivial

* tanglingzhi@quantumsc.cn

† dongsh2@yahoo.com

‡ danweizhang@m.scnu.edu.cn

band insulator can be driven to a nontrivial insulator by the moiré superlattice potential for more than once. This reentrant topological phase has the trivial-topological-trivial-topological-trivial multiple transition, which is revealed from the winding numbers, energy gap and the edge states. For the topological transitions, we derive the scale invariance of the winding number with a critical exponent. By adopting the band flatness and the fractal dimension (FD), we study the localization of single-particle eigenstates induced by the moiré potential. In the interacting case, the moiré enlarged periodicity makes the system's Hilbert space far from the flexible of ED method. Thus, we use the density matrix renormalization group (DMRG) method to investigate the density wave orders and topology of the many-body ground state. We unveil the occurrence of periodic-moiré-spin density wave (PM-SDW) whenever the moiré potential is activated, in both non-interacting and interacting cases. The PM-SDW order is enhanced by the on-site interaction but suppressed by the nearest-neighbor interaction. In the presence of strong nearest-neighbor interaction, the CDW order with the moiré periodicity can be exhibited. Moreover, the multiple topological transitions are preserved in the interacting case. The reentrant topological phase is characterized by the many-body Berry phase and nontrivial edge excitations, and even hosts the SDW order, which corresponds the topological PM-SDW.

RESULTS

Model Hamiltonian

We start by considering a 1D Raman optical lattice loaded with ultracold fermionic atoms [35, 36]. Two internal states of atoms are used to encode a spin-1/2 degree of freedom, and proper Raman laser beams can be used to engineer atomic hopping terms with effective spin-orbit couplings and Zeeman potentials. The tight-binding Hamiltonian of the system reads

$$\begin{aligned}
\hat{H} &= \hat{H}_0 + \hat{H}_{\text{int}} \\
\hat{H}_0 &= -t \sum_j (\hat{c}_{j,\uparrow}^\dagger \hat{c}_{j+1,\uparrow} - \hat{c}_{j,\downarrow}^\dagger \hat{c}_{j+1,\downarrow} + \text{H.c.}) \\
&\quad + t_s \sum_j (\hat{c}_{j,\uparrow}^\dagger \hat{c}_{j+1,\downarrow} - \hat{c}_{j,\downarrow}^\dagger \hat{c}_{j+1,\uparrow} + \text{H.c.}) \\
&\quad + \sum_j (m_z + m_{o,j}) (\hat{n}_{j,\uparrow} - \hat{n}_{j,\downarrow}), \\
\hat{H}_{\text{int}} &= U \sum_j \hat{n}_{j,\uparrow} \hat{n}_{j,\downarrow} + V \sum_{j,\sigma} \hat{n}_{j,\sigma} \hat{n}_{j+1,\sigma}.
\end{aligned} \tag{1}$$

Here $\hat{c}_{j,\sigma}^\dagger$ creates a fermionic atom with spin- σ ($\sigma = \uparrow, \downarrow$) on lattice site j , and $\hat{n}_{j,\sigma} = \hat{c}_{j,\sigma}^\dagger \hat{c}_{j,\sigma}$ is the particle number operator. The parameters t and t_s denote the spin-dependent and spin-flip hopping strengths,

respectively. The former term generates an effective phase and momentum kick to the atoms while preserving their spin, whereas the latter term can flip their spin. The Zeeman potential contains two parts, the uniform part with strength m_z and the spatially modulated part $m_{o,j}$. We consider the modulation with a moiré potential

$$m_{o,j} = m_o [\cos(2\pi j/a_1) + \cos(2\pi j/a_2)], \tag{2}$$

which is formed by the superposition of two periodic potentials of period a_α ($\alpha = 1, 2$). Here, we focus on the commensurate moiré potential by choosing $a_1 = 3$ and $a_2 = 7$. Thus, the lattice system has a period of $a_{12} = a_1 a_2 = 21$ with each supercell consisting of 21 sites. In addition to the on-site atomic repulsion of strength U , we include the nearest-neighbour repulsion of strength V in the interaction Hamiltonian \hat{H}_{int} in Eq. (1). We assume the lattice length to be $L = a_{12} A = 21A$ with A supercells and focus on the system at (near) half filling with the particle number $N_f = L$. We set $t = 1$ as the energy unit hereafter.

In the absence of interactions ($U = V = 0$) and moiré potential ($m_o = 0$), the single-particle Hamiltonian \hat{H}_0 reduces to the chiral AIII-class model with a topological insulator when $|m_z| < 2t$ [35], which has been realized with ultracold fermions in the 1D Raman optical lattice [36]. The chiral symmetry is given by $\hat{C} \hat{H}_0 \hat{C}^{-1} = -\hat{H}_0$, where $\hat{C} = \mathbf{I}_L \otimes \hat{\sigma}_x$ is the chiral symmetry operator with \mathbf{I}_L a L -rank identity matrix and $\hat{\sigma}_{x,y,z}$ Pauli matrices. In the presence of the moiré potential, the chirality of \hat{H}_0 presences, the corresponding momentum Hamiltonian can be derived. By taking the Fourier transformation under the PBC, we can transform the moiré Zeeman terms to the momentum space as

$$\sum_j \cos\left(\frac{2\pi j}{a_\alpha}\right) \hat{n}_{j,\sigma} = \frac{1}{2} \sum_k \left(\hat{c}_{k,\sigma}^\dagger \hat{c}_{k+\frac{2\pi}{a_\alpha},\sigma} + \text{H.c.} \right). \tag{3}$$

By constructing the Bloch basis $|BK\rangle = [\hat{c}_{k+0\pi/a_{12}\uparrow} \hat{c}_{k+2\pi/a_{12}\uparrow} \cdots \hat{c}_{k+0\pi/a_{12}\downarrow} \hat{c}_{k+2\pi/a_{12}\downarrow} \cdots]^\text{T}$ in the reduced MBZ $k \in [0, 2\pi/a_{12}]$, we obtain the Bloch Hamiltonian of \hat{H}_0 as

$$\begin{aligned}
\hat{H}_B(k) &= \sum_{i=1}^{a_{12}} [(m_z - 2t \cos k) \hat{\sigma}_z \otimes \mathbf{D}_i - 2t_s \sin k \hat{\sigma}_y \otimes \mathbf{D}_i] \\
&\quad + \frac{m_o}{2} \sum_{i=1}^{a_{12}} (\mathbf{I}_2 \otimes \mathbf{E}_i + \mathbf{I}_2 \otimes \mathbf{F}_i + \text{H.c.}),
\end{aligned} \tag{4}$$

where $\mathbf{D}_i = \text{diag}(\cdots, 0, 1_i, 0, \cdots)$ is a a_{12} -rank diagonal matrix with i -th diagonal element equal to 1, and \mathbf{E}_i (\mathbf{F}_i) is a $a_{12} \times a_{12}$ square matrix with all elements equal to 0 except $\mathbf{E}_i(i, i + a_2) = 1$ [$\mathbf{F}_i(i, i + a_1) = 1$]. Under the moiré potential, the single-particle energy spectrum generally separates to $2a_{12}$ subbands.




FIG. 1. (Color online) (a) Real-space winding number ν and (b) inverse of the zero-mode localization length Λ^{-1} as functions of m_o and m_z . Red dashed curve in (a) denotes the topological phase boundary revealed by the momentum-space winding number ν_k . Real-space winding number ν and energy gap ΔE as functions of (c) m_o for $m_z = 2.4$ and (d) m_z for $m_o = 2.8$ under the OBC. (e) Energy spectrum with respect to m_o under the OBC. The zero-energy modes in the topological regions are highlighted in red. (f) Density distributions of the two zero-energy edge modes for $m_z = 2.4$ and $m_o = 1.7$. Other parameters are $t_s = 0.95$, and $A = 32$.

Reentrant topological phases

The topological nature of the chiral-symmetric single-particle Hamiltonian can be revealed by the winding number. In the OBC, the real-space winding number of \hat{H} is given by [68]

$$\nu = \frac{1}{L} \text{Tr}'(\hat{C}\hat{P}[\hat{P}, \hat{X}]), \quad (5)$$

where the projector $\hat{P} = \sum_{l=1}^L (|\psi_l\rangle\langle\psi_l| - \hat{C}|\psi_l\rangle\langle\psi_l|\hat{C}^{-1})$ sums over the lowest half single-particle wave functions $|\psi_l\rangle$ (for the half-filling case), $\hat{C} = \mathbf{I}_L \otimes \hat{\sigma}_x$ is the real-space chiral operator, \hat{X} is the coordinate operator with $X_{j\sigma, j'\sigma'} = \delta_{jj'}\delta_{\sigma\sigma'}$, and Tr' denotes the trace per volume over the center internal $L' = L/2$ matrix elements which can avoid the boundary effect. For consistency, we also compute the winding number under the PBC ν_k from the Bloch Hamiltonian (4), which can be written as the following off-diagonal form

$$\hat{H}_B(k) = \begin{pmatrix} 0 & q(k) \\ q^\dagger(k) & 0 \end{pmatrix}, \quad (6)$$

with $q(k)$ a $a_{12} \times a_{12}$ matrix. The 1D winding number is then obtained by integral over the MBZ [67]

$$\nu_k = \frac{1}{2\pi i} \int_0^{2\pi/a_{12}} dk \text{Tr} [q^{-1} \partial_k q]. \quad (7)$$

The chiral symmetry ensures ν_k being an integer, which counts the number of times the momentum space Hamiltonian encircles the original point [69]. In general, it is not possible to obtain an analytical expression for ν_k in our model, so numerical integration is employed. Figure 1 (a) shows the topological phase diagram characterized by the real-space winding number ν . The topological phase boundary is consistent with that obtained by the momentum-space winding number ν_k , plotted as the red dashed line.

Due to the gap-closing nature at topological transition points, the localization length Λ of the zero-energy mode under the OBC diverges with $\Lambda^{-1} \rightarrow 0$ [68]. To calculate the localization length of the zero-energy mode, we can rewrite the Hamiltonian as

$$\hat{H}_0 = \begin{pmatrix} M_1 & T & 0 & 0 & \dots & 0 & 0 & 0 \\ T^\dagger & M_2 & T & 0 & \dots & 0 & 0 & 0 \\ 0 & T^\dagger & M_3 & T & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & T^\dagger & M_{L-1} & T \\ 0 & 0 & 0 & 0 & \dots & 0 & T^\dagger & M_L \end{pmatrix}, \quad (8)$$

where $M_j = m_{o,j} \hat{\sigma}_z$ and $T = it_s \hat{\sigma}_y - t \hat{\sigma}_z$. The zero-energy eigenstate $\psi = \{\psi_1, \psi_2 \dots \psi_{L-1}, \psi_L\}^T$ with $\psi_j = \{\psi_{j,\uparrow}, \psi_{j,\downarrow}\}$ can be obtained by solving $\hat{H}_0 |\psi\rangle = 0$ as

$$\psi_j = \begin{cases} -T^{-1} M_1 \psi_1 & j = 2; \\ -T^{-1} (T^\dagger \psi_{j-2} + M_{j-1} \psi_{j-1}), & j > 2. \end{cases} \quad (9)$$

By setting $\psi_1 = \{1, -1\}^T$ as the eigenstate of $\hat{\sigma}_x$, one can obtain ψ_L through Eq. (9). Hence, we obtain

$$\Lambda^{-1} = \left| \frac{1}{L} \min \left\{ \ln \left| \frac{\psi_{L,\uparrow}}{\psi_{1,\uparrow}} \right|, \ln \left| \frac{\psi_{L,\downarrow}}{\psi_{1,\downarrow}} \right| \right\} \right|. \quad (10)$$

The numerical result of Λ^{-1} is shown in Fig. 1 (b). The divergence of the localization length indicates the topological phase boundary, which is consistent with that revealed by the winding numbers shown in Fig. 1 (a).

Similar to the disorder-induced TAI [27, 70–72], the commensurate moiré potential can drive a topological phase from a trivial insulator. More interestingly, the finger-shaped elongations of the topological region in Fig. 1 (a) host reentrant topological transitions induced by the moiré potential. In Fig. 1 (c), we highlight the reentrant topological transitions by plotting ν and the single-particle energy gap $\Delta E = E_{L+1} - E_L$ under the OBC as functions of m_o with fixed $m_z = 2.4$. One can see the sequent trivial-topological-trivial-topological-trivial transition driven by the moiré potential strength, with gap closes occurring at each transition point. Such




FIG. 2. (Color online) (a) $\partial\nu/\partial m_o$ as a function of m_o with $m_z = 2.4$ and various system sizes. The finite-size critical point at $m_o = m_o^{(L)}$ is given by the peak of each curve. (b) Finite-size scaling of the distance from $m_o^{(L)}$ to the ideal transition point $m_{oc} = 1.223$. (c) $\partial\nu/\partial m_z$ as a function of m_z with $m_o = 2.8$ and various system sizes. (d) Finite-size scaling of the distance from $m_z^{(L)}$ to $m_{zc} = 1.678$. Other parameter is $t_s = 0.95$.

a reentrant topological transition is absent without the moiré potential. However, it can be induced by the uniform potential m_z with finite and proper values of m_o , such as the topological-trivial-topological-trivial transition shown in Fig. 1 (d). The finite and vanishing values of ΔE in Figs. 1 (c,d) correspond to the absence and presence of two zero-energy edge modes in the trivial and topological phases under the OBC, respectively. To be more clearly, we show the energy spectrum with respect to the moiré potential strength under the OBC in Fig. 1 (e). As the energy gap closes in topological regions, there emerges two exponentially localized states near two ends of the 1D lattice with zero energy, which are shown in Fig. 1 (f). These localized edge states will disappear when the energy gap opens. Note that the reentrant topological phase can be induced by other moiré potentials with different choices of a_1 and a_2 , and more reentrant transitions can be realized with proper superlattice and Hamiltonian parameters.

We further perform the scale invariance analysis of the real-space winding number in finite-size systems. Note that the finite-size scaling of the fidelity susceptibility and quantum entanglement has been used to explore the critical behaviors in quantum phase transitions [73–76]. Figures 2 (a) and 2 (c) show the numerical results of $\partial\nu/\partial m_o$ and $\partial\nu/\partial m_z$ as functions of m_o and m_z for various system sizes with fixed $m_z = 2.4$ and $m_o = 2.8$, respectively. With the increase of A ($L = 21A$), the peaks of the curves of $\partial\nu/\partial m_{o,z}$ at $m_{o,z}^{(L)}$ approach to the ideal topological transition points at $m_o = m_{oc} = 1.223$ and $m_z = m_{zc} = 1.678$ obtained from ν_k . The distances




FIG. 3. (Color online) The energy spectrum in momentum space for (a) $m_o = 0.1$, (b) $m_o = 1$ and (c) $m_o = 10$ under PBC. (d) Inverse flatness of s -th energy band f_s^{-1} for $s = 1, 10$ as a function of m_o . Other parameters are $t_s = 0.95$ and $m_z = 2.4$.

from $m_{o,z}^{(L)}$ to $m_{oc,zc}$ as a function of the system size L are shown in Figs. 2 (b) and 2 (d), respectively. We find that they are well fitted by the same power-law scaling form [77–79]

$$|m_o^{(L)} - m_{oc}| \propto L^\mu, |m_z^{(L)} - m_{oz}| \propto L^\mu. \quad (11)$$

By fitting the curves using Eq. (11) in the logarithm form, we obtain the critical exponents $\mu = 0.9897$ in Fig. 2 (b) and $\mu = 1.0057$ in Fig. 2 (d), respectively. This indicates that these reentrant topological transitions belong to the same universal class with $\mu \approx 1$, but different from the that of the disorder-driven topological transitions with $\mu \approx 2$ [80].

Localization properties

We proceed to study the effect of the commensurate moiré potential on the localization property of the single-particle eigenstates. Figures 3 (a-c) show the energy bands of $\hat{H}_B(k)$ in the momentum space for $m_o = 0.1, 1, 10$ with fixed $m_z = 2.4$, respectively. The results indicate a flattening of the energy bands as m_o increases. We can introduce a dimensionless parameter to characterize the flatness of s -th band, which is defined as

$$f_s = \Delta_s / W_s. \quad (12)$$

Here $\Delta_s = \max\{E_s(k) - E_{s-1}(k), E_{s+1}(k) - E_s(k)\}$ denotes the sub-band gap and $W_s = \max\|E_s(k) - E_s(k')\|_{k,k'}$ is the bandwidth. When flat band emerges, the bandwidth W_s vanishes while sub-band gap Δ_k keeps finite, and f_s will tend to infinity. For a better visibility,




FIG. 4. (Color online) (a) Real-space FD and (b) momentum-space FD of eigenstates versus m_o with $A = 32$. (c) Density distributions for typical extended, critical, and localized states for $A = 20$. (d) Finite-size scaling of the real-space FD for the three states with $L = 21A$ and $A = 64, 96, 128, 160$. Other parameters are $t_s = 0.95$ and $m_z = 2.4$.

we plot the inverse of the flatness f_s^{-1} as a function of m_o for $s = 1, 10$ sub-bands in Fig. 3 (d). It is clear that the inverse of the flatness tends to zero ($f_s \rightarrow \infty$), and the energy bands become nearly flat under moderate moiré potential, which lead to the emergence of compact localized states, similar to those in other moiré systems [42, 81, 82].

We adopt the FD to reveal the localization properties of eigenstates. For the l -th eigenstate, the real-space FD is defined as

$$\Gamma^{(l)} = - \lim_{L \rightarrow \infty} \ln(IPR^{(l)}) / \ln L, \quad (13)$$

where $IPR^{(l)} = \sum_{j,\sigma} |\psi_{j,\sigma}^{(l)}|^4$ is the real-space inverse participation ratio (IPR), and $\psi_{j,\sigma}^{(l)}$ is the probability amplitude of the l -th eigenstate. The momentum-space FD is then given as $\Gamma_k^{(l)} = - \lim_{L \rightarrow \infty} \ln(IPR_k^{(l)}) / \ln L$, where the momentum $IPR_k^{(l)}$ is obtained by applying a Fourier transformation to the real-space eigenstates. There are extended, localized and critical states in general localization systems. For an extended eigenstate, the wave function is delocalized in the real space with $\Gamma^{(l)} \sim 1$, but localized in the dual-momentum space with $\Gamma_k^{(l)} \sim 0$. A localized eigenstate is opposite and characterized by $\Gamma^{(l)} \sim 0$ and $\Gamma_k^{(l)} \sim 1$. In contrast, a critical state is delocalized in both the real and the momentum spaces, and thus has finite FDs $0 < \Gamma^{(l)} < 1$ and $0 < \Gamma_k^{(l)} < 1$.

The numerical results of $\Gamma^{(l)}$ and $\Gamma_k^{(l)}$ of all eigenstates as functions of m_o for $m_z = 2.4$ are shown in Figs. 4 (a) and 4(b), respectively. As m_o increases from 0 to a moderate value $m_o \sim 1$, a considerable part of eigenstates




FIG. 5. (Color online) (a,c) The real-space spin density fluctuation δS_j plotted for non-interacting case with $U = V = 0$ (a), and interacting case with $U = V = 1$ (c). Solid lines for $m_o = 0$ and dashed curves for $m_o = 0.01$. Inset plots are the charge density fluctuation δn_j for $m_o = 0.01$ where no CDW occurs. (b,d) The momentum-space spin density fluctuation δS_k for the non-interacting and interacting cases in (a,c), respectively. (e) The maximum local spin density S_k^{\max} as functions of m_o for non-interacting (black circles) and interacting (blue squares) cases. (f) The scaling of maximum local spin density versus system size $L = 21A$ with $A = 2, 4, 8, 16$ for the non-interacting (solid lines) and interacting (dashed lines) cases. Other parameters are $t_s = 0.95$ and $m_z = 2.4$.

cross over to the delocalized critical states from the extended states. For moderate m_o , the extended, critical, and localized states coexists in the system. For sufficiently large m_o , only critical and localized states are exhibited. The density distributions $\langle \hat{n}_j \rangle = \langle \hat{n}_{j,\uparrow} \rangle + \langle \hat{n}_{j,\downarrow} \rangle$ of three distinct eigenstates are shown in Fig. 4 (c). One can observe that the delocalized critical states in the commensurate lattice exhibit translation invariance with respect to the supercells, which reflect the moiré pattern of the superlattice. In Fig. 4 (d), we perform the finite-size scaling of the real-space FD and confirm the three distinct localization states.

Periodic-moiré-spin density waves

We first report the emergent of PM-SDW order in the Mott insulating phase driven by the moiré potential m_o ,

and then discuss the influence of the on-site interaction U and nearest-neighbor interaction V . To characterize the PM-SDW order, we calculate the spin density fluctuation

$$\delta S_j = S_j - \bar{S} \quad (14)$$

in the real space. Here $S_j = \langle \hat{n}_{j,\uparrow} \rangle - \langle \hat{n}_{j,\downarrow} \rangle$ is spin density on site j and \bar{S} is the spatially averaged spin density. The local spin density in the momentum space is given by $S_k = \sum_j S_j \cos(jk + \phi_0)$ [83], where $k = 2\pi i A/L$ ($i = 1, 2, \dots, a_{12}$) is the wave vector and ϕ_0 denotes an arbitrary phase. The spin densities may subsequently become modulated by the moiré superlattice, revealing a PM-SDW period that is commensurate with the supercell of $a_{12} = 21$ sites. Thus, one can extract these properties from the peaks of the density distribution by transforming δS_j to the momentum space δS_k . The density distribution of the PM-SDW state shows a period of a_{12} sites in the real space, which corresponds to $a_{12} - 1$ typical peaks in the momentum space, as the peak at $k = 0$ is removed by subtracting the average spin density \bar{S} in Eq. (14).

In Figs. 5 (a,b), we show the spin density fluctuation for $m_o = 0, 0.01$ in the real and momentum spaces, respectively. The periodicity of the SDW emerges when m_o changes from 0 to 0.01. These are $a_{12} - 1 = 20$ peaks in the spin density fluctuation S_k , which clearly reveal the PM-SDW nature of the many-body ground state induced by the moiré potential. The charge density fluctuation $\delta n_j = \langle \hat{n}_j \rangle - \bar{n}$ is also plotted as an inset figure, which indicates the absence of the CDW in this non-interacting case. Similar results for the interacting case with $U = V = 1$ are shown in Figs. 5 (c,d). Although the DMRG sweeps convergent to our error goal, two peaks near the magnitude of 10^{-10} are indistinguishable in the background due to the approximate same magnitude numerical errors in DMRG cutoffs. The maximum local spin density S_k^{\max} extracted from the highest peak, is plotted as functions of m_o in Fig. 5 (e). These numerical results indicate that the emergence of the PM-SDW order for both non-interacting and interacting cases is due to the moiré superlattice potential. Moreover, a very small moiré potential strength is sufficient to drive the PM-SDW order. We also consider the dependence of the PM-SDW on the system size and reveal the scaling of the maximum local spin density S_k^{\max} in Fig. 5 (f). For the ordinary Mott insulator without SDW order, when $m_o = 0$ the local spin densities keep vanishing for all system sizes. For finite m_o , the maximum local spin density per site is non-vanishing and independent on the system size, which indicate that the SDW order is preserved in the thermodynamic limit. In cold-atom systems, the SDW order could be experimentally detected using the spin-polarized scanning tunneling microscopy [84].

We further investigate the effects of the on-site and nearest-neighbor interactions on the PM-SDW. For this propose, we fix the moiré potential strength at $m_o = 2$, where the many-body ground state exhibits the PM-SDW order even in the non-interacting limit. In the




FIG. 6. (Color online) (a) Peak values of δS_k in region $[0, \pi)$ plotted in descending order for several U s with $V = 0$. (b) Distributions of δS_j in the two centered supercells. (c) Peak values of δS_k plotted in descending order for several V s with $U = 0$. (d) δn_j and δS_j for $V = 20$. (e, f) Distributions of δn_j and δS_j for $V = 2$ (e) and $V = 20$ (f). Other parameters are $t_s = 0.95$, $m_z = 2.4$ and $m_o = 2$.

presence of an on-site interaction U , particles with spin imbalances have lower energy and thus the SDW order is enhanced by increasing U . We observe the positive effect of on-site interaction on the PM-SDW from the local spin densities in the momentum space δS_k and its real-space counterpart δS_j , as shown in Figs. 6 (a) and 6 (b), respectively. Note that all peak values of δS_k are two-fold degenerate due to the reflection symmetry with respect to $k = \pi$. In Fig. 6 (a), we consider the ten peaks in region $k \in [0, \pi)$ and depict their values in descending order for several U s. Peak values of δS_k keep rising as U increases, which demonstrates the enhancement of the PM-SDW order. The corresponding real-space fluctuation δS_j in the two centered supercells is plotted in Fig. 6 (b), which shows the same moiré periodicity and increased amplitudes of the SDWs as U increases.

Strong nearest-neighbor interaction V may suppress the SDW and induce the CDW, since the spin imbalance has lower interaction energy in this case. In Fig. 6 (c), 20 peak values of δS_k are plotted in the descending order for $V = 0, 1, 2, 20$. The PM-SDWs are almost unchanged for small to moderate V s, but are inhibited for large V . We focus on the strong nearest-neighbor interaction case with $V = 20$, and depict δn_j and δS_j in the middle two supercells in Fig. 6 (d). The CDW with a period of two sites emerges in this case, while the SDW

almost vanishes. For a better comparison, we present the momentum-space counterparts δn_k and δS_k for $V = 2$ and $V = 20$ in Figs. 6 (e,f), respectively. For $V = 2$ in Fig. 6 (e), the PM-SDW dominates and a minor PM-CDW coexists with very small magnitudes. For $V = 20$ in Fig. 6 (f), the PM-SDW is strongly suppressed, and the CDW dominates with the average charge density $\bar{n} = 0$ and period of two sites. Here the CDW order shows the dominated wave number at $k = \pi$. The moiré periodicity can still be revealed from the 20 peaks for PM-CDW or PM-SDW, which are three orders of magnitude smaller than that for the CDW order at $k = \pi$.

Interacting topological phases

In the interacting case, the topology of the ground state at half filling can be characterized by the many-body Berry phase under the twisted, which is given by PBC [85–87]

$$\gamma = \frac{1}{\pi} \oint d\theta \langle \Psi^g(\theta) | i\partial_\theta | \Psi^g(\theta) \rangle \text{ mod } 2 \quad (15)$$

in units of π . Here $|\Psi^g(\theta)\rangle$ is the many-body ground state of L fermions with the twist phase θ . The quantized Berry phase is $\gamma = 1$ for the topological phase, while $\gamma = 0$ for the trivial phase. In Figs. 7 (a,b), we show the numerical results of γ in the m_o - V and m_o - U planes for $m_z = 2.4$, respectively. The reentrant topological phase is preserved for certain U s or V s, before the on-site or nearest-neighbor interactions are dominated. It's worth emphasizing that the reentrant topological phase is in the parameter region of the PM-SDW, as discussed previously. Thus, the topological PM-SDW can be exhibited for interacting fermions in the moiré superlattice.

In the interacting topological phase under the OBC, the ground states near half filling are two-fold degenerate with zero-energy excitations localized near two edge of the lattice. We numerically compute the excitation gap Δ_g under the OBC, which is defined as the energy gap between the first excited state and the ground state. The excitation gap and the corresponding Berry phase as functions of m_o for $U = 0.1$ are plotted in Fig. 7 (c). The gap closing and reopening behavior is consistent with that of the many-body Berry phase. To show the edge excitations in the topological phase, we compute the density distribution

$$\delta N_f(j) = \langle \Psi_{N_f+1}^g | \hat{n}_j | \Psi_{N_f+1}^g \rangle - \langle \Psi_{N_f}^g | \hat{n}_j | \Psi_{N_f}^g \rangle, \quad (16)$$

which is defined as the difference of the ground-state distributions between $N_f + 1$ -filling and N_f -filling with $N_f = L$ or $L - 1$. This corresponds to adding or removing one quasiparticle to the half-filling ground state. In the topological phase, the added (removed) quasiparticle tends to localize near one end of the 1D lattice, as shown in Fig. 7 (d).




FIG. 7. (Color online) (a,b) Many-body Berry phase γ in units of π plotted in the m_o - V plane with $U = 0$ (a) and in the m_o - U plane with $V = 0$ (b). (c) Excitation gap Δ_g under the OBC and γ as functions of m_o for $U = 0.1$. (d) Distribution of two zero-energy excitation modes near half filling δN_f for $U = 0.1$ and $m_o = 1.7$. Other parameters are $t_s = 0.95$, $m_z = 2.4$, $A = 2$ for γ , and $A = 8$ for others.

CONCLUSIONS

To summarize, we have demonstrated that the 1D moiré potential can induce trivial-topological-trivial-topological-trivial multiple transitions in both single-particle and many-body regions in a spin-1/2 fermionic optical superlattice. We have uncovered the topological phases with zero-energy edge modes or excitations and nontrivial topological numbers. The scaling exponents of topological transitions for both moiré and uniform Zeeman potentials have been revealed and agree with each others. We have also investigated the localization property with the moiré-induced nearly flat bands and delocalized critical states. The topology and localization in our moiré superlattices are different from those in disorder-induced TAIs. Then, we have unveiled the PM-SDW orders of the many-body ground state, which are instantly induced by turning on the moiré potential. The on-site interaction can enhance the PM-SDW, while a sufficient nearest-neighbor interaction suppresses the SDW and induces the CDW. Finally, we have generalized our findings to the interacting region by means of the DMRG method. The reentrant topological phase persists for a finite interaction strength, after which two topological regions either merge or vanish, as the nearest-neighbor and on-site interactions enhance and destroy the topology, respectively. Although we focus a typical moiré potential in this work, our findings can be extended to other commensurate moiré potentials with different supercells, such as $\{a_1, a_2\} = \{2, 5\}, \{5, 7\}, \{3, 11\}$, etc. Our proposed model provides a feasible setup to experimentally observe exotic reentrant topological phases, since the required optical superlattices and spin-

flip (spin-dependent) hoppings are already realized in cold-atom experiments. It would also be interesting to further explore reentrant topological phases and SDW orders in 2D or incommensurate moiré systems.

METHODS

The topological and localization properties of the system in the single-particle case are directly obtained from the real-space Hamiltonians \hat{H} in Eq. (1) and the momentum-space Hamiltonian $H_B(k)$ in Eq. (4). Both Hamiltonians are easy to construct, and the real-space and momentum-space physical quantities are numerically calculated by the ED method of these matrices using MATLAB version R2023b. The physical quantities of half-filling many-body ground states in the noninteracting limit are obtained from the ED method by summing over the lowest half single-particle physical quantities. As a supercell has 21 sites, the ED method is far beyond the availability with more than one supercell in the many-body interacting case. The many-body ground states at half-filling in the presence of the on-site and nearest-neighbor interactions are simulated by the DMRG method with matrix-product state representation. We use the itensor library [88] version 3.1.10 in practice. In the simulation of the many-

body Berry phase, the system size $L = 21A$ with $A = 2$ is considered, the bond dimension of the maximum matrix product states is set to 200, and 24 DMRG sweeps or 10^{-6} relative energy error goal are taken. These criteria are sufficient in our simulations as the topological invariant is robust against disorders and finite-site effect. In the calculation of other quantities, we increase the supercell number to $A = 8, 16$ ($L = 168, 336$ with $N_a = L$ spin-1/2 fermions), with up to 400 bond states and 36 DMRG sweeps or 10^{-8} relative energy error goal to achieve convergent results.

DATA AVAILABILITY

The data that used to create Figures (1-7) this paper are available from the corresponding authors upon reasonable request.

CODE AVAILABILITY

The codes are mainly based on mature commercial software and long-lasting open-source library, and are not confused to reproduce. The computer codes used in this paper are available from the corresponding authors upon reasonable request.

-
- [1] Hasan, M. Z. & Kane, C. L. Colloquium: Topological insulators. *Rev. Mod. Phys.* **82**, 3045–3067 (2010).
- [2] Qi, X.-L. & Zhang, S.-C. Topological insulators and superconductors. *Rev. Mod. Phys.* **83**, 1057–1110 (2011).
- [3] Culcer, D., Keser, A. C., Li, Y. & Tkachov, G. Transport in two-dimensional topological materials: recent developments in experiment and theory. *2D Mater.* **7**, 022007 (2020).
- [4] Bansil, A., Lin, H. & Das, T. Colloquium: Topological band theory. *Rev. Mod. Phys.* **88**, 021004 (2016).
- [5] Chiu, C.-K., Teo, J. C. Y., Schnyder, A. P. & Ryu, S. Classification of topological quantum matter with symmetries. *Rev. Mod. Phys.* **88**, 035005 (2016).
- [6] Armitage, N. P., Mele, E. J. & Vishwanath, A. Weyl and dirac semimetals in three-dimensional solids. *Rev. Mod. Phys.* **90**, 015001 (2018).
- [7] Lu, L., Joannopoulos, J. D. & Soljačić, M. Topological photonics. *Nat. photon.* **8**, 821–829 (2014).
- [8] Ozawa, T. et al. Topological photonics. *Rev. Mod. Phys.* **91**, 015006 (2019).
- [9] Zhang, D.-W., Zhu, Y.-Q., Zhao, Y. X., Yan, H. & Zhu, S.-L. Topological quantum matter with cold atoms. *Adv. Phys.* **67**, 253–402 (2018).
- [10] Coen, S. et al. Nonlinear topological symmetry protection in a dissipative system. *Nat. Commun.* **15**, 1398 (2024).
- [11] Schroer, M. D. et al. Measuring a topological transition in an artificial spin-1/2 system. *Phys. Rev. Lett.* **113**, 050402 (2014).
- [12] Roushan, P. et al. Observation of topological transitions in interacting quantum circuits. *Nature* **515**, 241–244 (2014).
- [13] Tan, X. et al. Topological maxwell metal bands in a superconducting qutrit. *Phys. Rev. Lett.* **120**, 130503 (2018).
- [14] Tan, X. et al. Experimental measurement of the quantum metric tensor and related topological phase transition with a superconducting qubit. *Phys. Rev. Lett.* **122**, 210401 (2019).
- [15] Wei, M. et al. Quantum fluctuation of the quantum geometric tensor and its manifestation as intrinsic hall signatures in time-reversal invariant systems. *Phys. Rev. Lett.* **130**, 036202 (2023).
- [16] Li, X. et al. Mapping the topology-localization phase diagram with quasiperiodic disorder using a programmable superconducting simulator. *Phys. Rev. Res.* **6**, 1042038 (2024).
- [17] Zhang, D.-W. et al. Quantum simulation of exotic pt-invariant topological nodal loop bands with ultracold atoms in an optical lattice. *Phys. Rev. A* **93**, 043617 (2016).
- [18] Tsui, D. C., Stormer, H. L. & Gossard, A. C. Two-dimensional magnetotransport in the extreme quantum limit. *Phys. Rev. Lett.* **48**, 1559–1562 (1982).
- [19] Raghu, S., Qi, X.-L., Honerkamp, C. & Zhang, S.-C. Topological mott insulators. *Phys. Rev. Lett.* **100**, 156401 (2008).
- [20] Dauphin, A., Müller, M. & Martin-Delgado, M. A. Rydberg-atom quantum simulation and chern-number

- characterization of a topological mott insulator. *Phys. Rev. A* **86**, 053618 (2012).
- [21] Kuno, Y., Shimizu, K. & Ichinose, I. Various topological mott insulators and topological bulk charge pumping in strongly-interacting boson system in one-dimensional superlattice. *New J. Phys.* **19**, 123025 (2017).
- [22] Grusdt, F., Höning, M. & Fleischhauer, M. Topological edge states in the one-dimensional superlattice bose-hubbard model. *Phys. Rev. Lett.* **110**, 260405 (2013).
- [23] Xu, Z. & Chen, S. Topological mott insulators of ultracold atomic mixtures induced by interactions in one-dimensional optical superlattices. *Phys. Rev. B* **88**, 045110 (2013).
- [24] Chen, Y.-L., Zhang, G.-Q., Zhang, D.-W. & Zhu, S.-L. Simulating bosonic chern insulators in one-dimensional optical superlattices. *Phys. Rev. A* **101**, 013627 (2020).
- [25] Zhang, D.-W. et al. Skin superfluid, topological mott insulators, and asymmetric dynamics in an interacting non-hermitian aubry-andré-harper model. *Phys. Rev. B* **101**, 235150 (2020).
- [26] Wang, Y.-X. & Qi, D.-X. Spontaneous symmetry breaking of an interacting chern insulator on a topological square lattice. *Phys. Rev. B* **99**, 075204 (2019).
- [27] Zhang, G.-Q., Tang, L.-Z., Zhang, L.-F., Zhang, D.-W. & Zhu, S.-L. Connecting topological anderson and mott insulators in disordered interacting fermionic systems. *Phys. Rev. B* **104**, L161118 (2021).
- [28] Nayak, C., Simon, S. H., Stern, A., Freedman, M. & Das Sarma, S. Non-abelian anyons and topological quantum computation. *Rev. Mod. Phys.* **80**, 1083–1159 (2008).
- [29] Poulsen Nautrup, H., Friis, N. & Briegel, H. J. Fault-tolerant interface between quantum memories and quantum processors. *Nat. Commun.* **8**, 1321 (2017).
- [30] He, M., Sun, H. & He, Q. L. Topological insulator: Spintronics and quantum computations. *Front. Phys.* **14**, 43401 (2019).
- [31] Breunig, O. & Ando, Y. Opportunities in topological insulator devices. *Nat. Rev. Phys.* **4**, 184–193 (2022).
- [32] Ou, Y. et al. Zrte2/crte2: an epitaxial van der waals platform for spintronics. *Nat. Commun.* **13**, 2972 (2022).
- [33] Sahu, P. et al. Room temperature spin-to-charge conversion in amorphous topological insulating gd-alloyed Bi_xSe_{1-x}/CoFeB bilayers. *ACS Appl. Mater. Interfaces* **15**, 38592–38602 (2023).
- [34] Su, W. P., Schrieffer, J. R. & Heeger, A. J. Solitons in polyacetylene. *Phys. Rev. Lett.* **42**, 1698–1701 (1979).
- [35] Liu, X.-J., Liu, Z.-X. & Cheng, M. Manipulating topological edge spins in a one-dimensional optical lattice. *Phys. Rev. Lett.* **110**, 076401 (2013).
- [36] Song, B. et al. Observation of symmetry-protected topological band with ultracold fermions. *Sci. Adv.* **4**, eaao4748 (2018).
- [37] Harper, P. G. Single band motion of conduction electrons in a uniform magnetic field. *Proc. Phys. Soc. Sect. A* **68**, 874 (1955).
- [38] Haldane, F. D. M. Model for a quantum hall effect without landau levels: Condensed-matter realization of the "parity anomaly". *Phys. Rev. Lett.* **61**, 2015–2018 (1988).
- [39] Stuhl, B. K., Lu, H.-L., Aycock, L. M., Genkina, D. & Spielman, I. B. Visualizing edge states with an atomic bose gas in the quantum hall regime. *Science* **349**, 1514–1518 (2015).
- [40] Mancini, M. et al. Observation of chiral edge states with neutral fermions in synthetic hall ribbons. *Science* **349**, 1510–1513 (2015).
- [41] Cao, Y. et al. Correlated insulator behaviour at half-filling in magic-angle graphene superlattices. *Nature* **556**, 80–84 (2018).
- [42] Zhang, Y.-H., Mao, D., Cao, Y., Jarillo-Herrero, P. & Senthil, T. Nearly flat chern bands in moiré superlattices. *Phys. Rev. B* **99**, 075127 (2019).
- [43] Balents, L., Dean, C. R., Efetov, D. K. & Young, A. F. Superconductivity and strong correlations in moiré flat bands. *Nat. Phys.* **16**, 725–733 (2020).
- [44] Haddadi, F., Wu, Q., Kruchkov, A. J. & Yazyev, O. V. Moiré flat bands in twisted double bilayer graphene. *Nano Lett.* **20**, 2410–2415 (2020).
- [45] Hu, J. et al. Correlated flat bands and quantum spin liquid state in a cluster mott insulator. *Commun. Phys.* **6**, 172 (2023).
- [46] Li, Y. et al. 1d electronic flat bands in untwisted moiré superlattices. *Adv. Mater.* **35**, 2300572 (2023).
- [47] Crépel, V., Regnault, N. & Queiroz, R. Chiral limit and origin of topological flat bands in twisted transition metal dichalcogenide homobilayers. *Commun. Phys.* **7**, 146 (2024).
- [48] Zhang, L. et al. Twist-angle dependence of moiré excitons in WS₂/MoSe₂ heterobilayers. *Nat. Commun.* **11**, 5888 (2020).
- [49] Guo, H., Zhang, X. & Lu, G. Shedding light on moiré excitons: A first-principles perspective. *Sci. Adv.* **6**, eabc5638 (2020).
- [50] Chen, W. et al. Direct observation of van der waals stacking-dependent interlayer magnetism. *Science* **366**, 983–987 (2019).
- [51] Dean, C. R. et al. Hofstadter's butterfly and the fractal quantum hall effect in moiré superlattices. *Nature* **497**, 598–602 (2013).
- [52] Chittari, B. L., Chen, G., Zhang, Y., Wang, F. & Jung, J. Gate-tunable topological flat bands in trilayer graphene boron-nitride moiré superlattices. *Phys. Rev. Lett.* **122**, 016401 (2019).
- [53] Zhang, Y.-H. & Senthil, T. Bridging hubbard model physics and quantum hall physics in trilayer graphene/h – BN moiré superlattice. *Phys. Rev. B* **99**, 205150 (2019).
- [54] Li, H., Kumar, U., Sun, K. & Lin, S.-Z. Spontaneous fractional chern insulators in transition metal dichalcogenide moiré superlattices. *Phys. Rev. Res.* **3**, L032070 (2021).
- [55] Serlin, M. et al. Intrinsic quantized anomalous hall effect in a moiré heterostructure. *Science* **367**, 900–903 (2020).
- [56] Park, Y., Kim, Y., Chittari, B. L. & Jung, J. Topological flat bands in rhombohedral tetralayer and multilayer graphene on hexagonal boron nitride moiré superlattices. *Phys. Rev. B* **108**, 155406 (2023).
- [57] Cao, Y. et al. Unconventional superconductivity in magic-angle graphene superlattices. *Nature* **556**, 43–50 (2018).
- [58] Xie, Y.-M. & Law, K. T. Orbital fulde-ferrell pairing state in moiré ising superconductors. *Phys. Rev. Lett.* **131**, 016001 (2023).
- [59] Nuckolls, K. P. et al. Strongly correlated chern insulators in magic-angle twisted bilayer graphene. *Nature* **588**, 610–615 (2020).

- [60] Breið, C. N. & Andersen, B. M. Chern insulator phases and spontaneous spin and valley order in a moiré lattice model for magic-angle twisted bilayer graphene. *Phys. Rev. B* **107**, 165114 (2023).
- [61] Wang, C. *et al.* Fractional chern insulator in twisted bilayer moiré. *Phys. Rev. Lett.* **132**, 036501 (2024).
- [62] Koshino, M., Moon, P. & Son, Y.-W. Incommensurate double-walled carbon nanotubes as one-dimensional moiré crystals. *Phys. Rev. B* **91**, 035405 (2015).
- [63] Zhao, S. *et al.* Observation of drastic electronic-structure change in a one-dimensional moiré superlattice. *Phys. Rev. Lett.* **124**, 106101 (2020).
- [64] Vu, D. & Das Sarma, S. Moiré versus mott: Incommensuration and interaction in one-dimensional bichromatic lattices. *Phys. Rev. Lett.* **126**, 036803 (2021).
- [65] Gonçalves, M., Amorim, B., Riche, F., Castro, E. V. & Ribeiro, P. Incommensurability enabled quasi-fractal order in 1d narrow-band moiré systems. *Nat. Phys.* (2024).
- [66] Yu, D. *et al.* Moiré lattice in one-dimensional synthetic frequency dimension. *Phys. Rev. Lett.* **130**, 143801 (2023).
- [67] Schnyder, A. P., Ryu, S. & Ludwig, A. W. W. Lattice model of a three-dimensional topological singlet superconductor with time-reversal symmetry. *Phys. Rev. Lett.* **102**, 196804 (2009).
- [68] Mondragon-Shem, I., Hughes, T. L., Song, J. & Prodan, E. Topological criticality in the chiral-symmetric class at strong disorder. *Phys. Rev. Lett.* **113**, 046802 (2014).
- [69] Yin, C., Jiang, H., Li, L., Lü, R. & Chen, S. Geometrical meaning of winding number and its characterization of topological phases in one-dimensional chiral non-hermitian systems. *Phys. Rev. A* **97**, 052115 (2018).
- [70] Li, J., Chu, R.-L., Jain, J. K. & Shen, S.-Q. Topological anderson insulator. *Phys. Rev. Lett.* **102**, 136806 (2009).
- [71] Groth, C. W., Wimmer, M., Akhmerov, A. R., Tworzydło, J. & Beenakker, C. W. J. Theory of the topological anderson insulator. *Phys. Rev. Lett.* **103**, 196805 (2009).
- [72] Tang, L.-Z., Liu, S.-N., Zhang, G.-Q. & Zhang, D.-W. Topological anderson insulators with different bulk states in quasiperiodic chains. *Phys. Rev. A* **105**, 063327 (2022).
- [73] Gu, S.-J., Kwok, H.-M., Ning, W.-Q. & Lin, H.-Q. Fidelity susceptibility, scaling, and universality in quantum critical phenomena. *Phys. Rev. B* **77**, 245109 (2008).
- [74] Albuquerque, A. F., Alet, F., Sire, C. & Capponi, S. Quantum critical scaling of fidelity susceptibility. *Phys. Rev. B* **81**, 064418 (2010).
- [75] Osterloh, A., Amico, L., Falci, G. & Fazio, R. Scaling of entanglement close to a quantum phase transition. *Nature* **416**, 608–610 (2002).
- [76] Zhang, G.-Q. & Xu, J.-B. Quantum coherence of an xy spin chain with dzyaloshinskii-moriya interaction and quantum phase transition. *J. Phys. A* **50**, 265303 (2017).
- [77] Chen, W. Scaling theory of topological phase transitions. *J. Phys. Condens. Matter* **28**, 055601 (2016).
- [78] Chen, W., Legner, M., Rüegg, A. & Sigrist, M. Correlation length, universality classes, and scaling laws associated with topological phase transitions. *Phys. Rev. B* **95**, 075116 (2017).
- [79] Yu, X.-J. & Li, W.-L. Fidelity susceptibility at the lifshitz transition between the noninteracting topologically distinct quantum critical points. *Phys. Rev. B* **110**, 045119 (2024).
- [80] Assunçao, B. D., Ferreira, G. J. & Lewenkopf, C. H. Phase transitions and scale invariance in topological anderson insulators. *Phys. Rev. B* **109**, L201102 (2024).
- [81] Ghorashi, S. A. A. *et al.* Topological and stacked flat bands in bilayer graphene with a superlattice potential. *Phys. Rev. Lett.* **130**, 196201 (2023).
- [82] Chen, Y., Huang, J., Jiang, K. & Hu, J. Decoding flat bands from compact localized states. *Sci. Bull.* **68**, 3165–3171 (2023).
- [83] Kundu, S. & Sénéchal, D. Spin density wave order in interacting type-i and type-ii weyl semimetals. *Phys. Rev. B* **103**, 085136 (2021).
- [84] Wiesendanger, R. Spin mapping at the nanoscale and atomic scale. *Rev. Mod. Phys.* **81**, 1495–1550 (2009).
- [85] Xiao, D., Chang, M.-C. & Niu, Q. Berry phase effects on electronic properties. *Rev. Mod. Phys.* **82**, 1959–2007 (2010).
- [86] Guo, H. & Shen, S.-Q. Topological phase in a one-dimensional interacting fermion system. *Phys. Rev. B* **84**, 195107 (2011).
- [87] Jünemann, J. *et al.* Exploring interacting topological insulators with ultracold atoms: The synthetic creutz-hubbard model. *Phys. Rev. X* **7**, 031057 (2017).
- [88] Fishman, M., White, S. R. & Stoudenmire, E. M. The ITensor Software Library for Tensor Network Calculations. *SciPost Phys. Codebases* 4 (2022).

ACKNOWLEDGMENTS

This work was supported by the National Key Research and Development Program of China (Grant No. 2022YFA1405304), the National Natural Science Foundation of China (Grants No. 12174126 and No. 12104166), the Guangdong Basic and Applied Basic Research Foundation (Grant No. 2024B1515020018), the Science and Technology Program of Guangzhou (Grant No. 2024A04J3004), and the Open Fund of Key Laboratory of Atomic and Subatomic Structure and Quantum Control (Ministry of Education).

AUTHOR CONTRIBUTIONS

G.Q.Z. performed the many-body calculations and wrote the first draft, L.Z.T. conceived and performed the single-particle part, L.F.Q. analyzed the data, D.W.Z. conceived the initial idea, S.H.D. and D.W.Z supervised the project. All authors discussed the results and revised the manuscript.

COMPETING INTERESTS

The authors declare no competing interests.