

# THE PELL SEQUENCE AND CYCLOTOMIC MATRICES INVOLVING SQUARES OVER FINITE FIELDS

HAI-LIANG WU, LI-YUAN WANG AND HE-XIA NI\*

ABSTRACT. In this paper, by some arithmetic properties of the Pell sequence and some  $p$ -adic tools, we study certain cyclotomic matrices involving squares over finite fields. For example, let  $1 = s_1, s_2, \dots, s_{(q-1)/2}$  be all the nonzero squares over  $\mathbb{F}_q$ , where  $q = p^f$  is an odd prime power with  $q \geq 7$ . We prove that the matrix

$$B_q((q-3)/2) = \left[ (s_i + s_j)^{(q-3)/2} \right]_{2 \leq i, j \leq (q-1)/2}$$

is a singular matrix whenever  $f \geq 2$ . Also, for the case  $q = p$ , we show that

$$\det B_p((p-3)/2) = 0 \Leftrightarrow Q_p \equiv 2 \pmod{p^2\mathbb{Z}},$$

where  $Q_p$  is the  $p$ -th term of the Pell sequence.

## CONTENTS

1. Introduction	2
1.1. Notation	2
1.2. Background and Motivation	2
1.3. The Pell sequence	5
1.4. The first result: case $m \in \{n-2, n-1\}$	5
1.5. The second result: case $m = n$	6
1.6. The third result: a variant of Carlitz's cyclotomic matrices	7
2. Proofs of Theorem 1.1 and its corollary	7
3. Preparations for the proof of Theorem 1.2	13
3.1. A lemma on almost circulant matrices	13
3.2. Some $p$ -adic preparations	14
4. Proof of Theorem 1.2	16
5. Proof of Theorem 1.3	20
References	24

---

*Key words and phrases.* the Pell sequence, cyclotomic matrices, character sums over finite fields.

2020 *Mathematics Subject Classification.* Primary 11L05, 15A15; Secondary 11R18, 12E20.

This research was supported by the Natural Science Foundation of China (Grant Nos. 12101321, 12201291 and 12371004).

\*Corresponding author.

## 1. INTRODUCTION

1.1. **Notation.** In this paper, the symbol  $p$  always denotes an odd prime. We let  $\mathbb{Q}_p$  be the  $p$ -adic number field and let  $\mathbb{Z}_p$  be the ring of all  $p$ -adic integers. As usual, we use  $\mathbb{C}_p$  to denote the completion of the algebraic closure of  $\mathbb{Q}_p$ . In addition, we let

$$\text{ord}_p : \mathbb{C}_p \rightarrow \mathbb{R}$$

be the  $p$ -adic order function over  $\mathbb{C}_p$ .

Let  $q = p^f$  be an odd prime power with  $f \in \mathbb{Z}^+$ . The symbol  $\mathbb{F}_q$  denotes the finite field with  $q$  elements and  $\mathbb{F}_q^\times = \mathbb{F}_q \setminus \{0\}$  is the multiplicative cyclic group of all nonzero elements over  $\mathbb{F}_q$ . The group of all multiplicative characters of  $\mathbb{F}_q$  is denoted by  $\widehat{\mathbb{F}_q^\times}$ , and the trivial character is written by  $\varepsilon$ . Throughout this paper, for any multiplicative character  $\psi : \mathbb{F}_q^\times \rightarrow \mathbb{C}$  (or  $\mathbb{C}_p$ ), we extend  $\psi$  to  $\mathbb{F}_q$  by letting  $\psi(0) = 0$ .

Let  $\zeta_p$  be a primitive  $p$ -th root of unity over  $\mathbb{C}$  (or  $\mathbb{C}_p$ ). Then for any  $\psi \in \widehat{\mathbb{F}_q^\times}$ , the Gauss sum of  $\psi$  is defined by

$$G_q(\psi) = \sum_{x \in \mathbb{F}_q} \psi(x) \zeta_p^{\text{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(x)},$$

where  $\text{Tr}_{\mathbb{F}_q/\mathbb{F}_p} : \mathbb{F}_q \rightarrow \mathbb{F}_p$  is the trace map. Also, for any  $\psi, \chi \in \widehat{\mathbb{F}_q^\times}$ , the Jacobi sum of  $\psi$  and  $\chi$  is defined by

$$J_q(\psi, \chi) = \sum_{x \in \mathbb{F}_q} \psi(x) \chi(1 - x).$$

For any positive integer  $a$  and nonnegative integer  $b$ , the symbol  $\binom{a}{b}$  is the binomial coefficient, that is,

$$\binom{a}{b} = \frac{a(a-1) \cdots (a-b+1)}{b!}.$$

Finally, for any square matrix  $M$  over a field,  $\det M$  denotes the determinant of  $M$ .

1.2. **Background and Motivation.** The investigations of cyclotomic matrices involving elements over finite fields were initiated by Lehmer [7] and Carlitz [1]. For instance, for any nontrivial multiplicative character  $\psi$  modulo  $p$ , Carlitz [1] first studied the cyclotomic matrices

$$(1.1) \quad C_p^-(\psi) := [\psi(j-i)]_{1 \leq i, j \leq p-1},$$

and

$$(1.2) \quad C_p^+(\psi) := [\psi(j+i)]_{1 \leq i, j \leq p-1}.$$

Carlitz [1, Theorem 4 and Theorem 5] proved that

$$\det C_p^-(\psi) = (-1)^{(p-1)/m} \cdot \frac{G_p(\psi)^{p-1}}{p},$$

and

$$\det C_p^+(\psi) = \begin{cases} \frac{1}{p} (-1)^{\frac{p-1}{2m}} G_p(\psi)^{p-1} & \text{if } m \equiv 1 \pmod{2}, \\ \frac{1}{p} (-1)^{\frac{p-1}{m}} \delta(\psi)^{p-1} G_p(\psi)^{p-1} & \text{if } m \equiv 0 \pmod{2}, \end{cases}$$

where  $m = \min\{k \in \mathbb{Z}^+ : \psi^k = \varepsilon\}$  is the order of  $\psi$  and

$$\delta(\psi) = \begin{cases} 1 & \text{if } \psi(-1) = 1, \\ -\mathbf{i} & \text{if } \psi(-1) = -1. \end{cases}$$

Along Carlitz's work, Chapman [2] further studied some variants of the matrix  $C_p^+(\psi)$ . In fact, let  $\left(\frac{\cdot}{p}\right)$  denotes the Legendre symbol, i.e., the unique quadratic multiplicative character of  $\mathbb{F}_p$ . Chapman considered the matrices

$$C_p^{(0)} := \left[ \left( \frac{i+j}{p} \right) \right]_{0 \leq i, j \leq (p-1)/2},$$

and

$$C_p^{(1)} := \left[ \left( \frac{i+j}{p} \right) \right]_{1 \leq i, j \leq (p-1)/2}.$$

Although Chapman only changed the sizes of the matrices, the calculations of  $\det C_p^{(0)}$  and  $\det C_p^{(1)}$  became extremely complicated. In fact, for  $p \equiv 1 \pmod{4}$ , let  $\epsilon_p > 1$  and  $h_p$  denote the fundamental unit and class number of  $\mathbb{Q}(\sqrt{p})$  respectively, and write

$$\epsilon_p^{h_p} = a_p + b_p \sqrt{p}$$

with  $a_p, b_p \in \mathbb{Q}$ . Then Chapman [2] showed that

$$\det C_p^{(0)} = \begin{cases} (-1)^{(p+3)/4} 2^{(p-1)/2} a_p & \text{if } p \equiv 1 \pmod{4}, \\ -2^{(p-1)/2} & \text{if } p \equiv 3 \pmod{4}, \end{cases}$$

and that

$$\det C_p^{(1)} = \begin{cases} (-1)^{(p-1)/4} 2^{(p-1)/2} b_p & \text{if } p \equiv 1 \pmod{4}, \\ 0 & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

Chapman also investigated some variants of  $C_p^-(\psi)$ , and the most well-known variant among them is Chapman's "evil determinant":

$$\det C_p^- := \det \left[ \left( \frac{j-i}{p} \right) \right]_{1 \leq i, j \leq (p+1)/2}.$$

Let

$$\epsilon_p^{(2+(-1)^{(p+3)/4})h_p} = a'_p + b'_p \sqrt{p}$$

with  $a'_p, b'_p \in \mathbb{Q}$ . Then Chapman conjectured that

$$\det C_p^- = \begin{cases} -a'_p & \text{if } p \equiv 1 \pmod{4}, \\ 1 & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

Vsemirnov [15, 16] later confirmed this tough conjecture by using ingenious and sophisticated matrix decomposition.

Sun [12] studied the variants of the above matrices from another perspective. For example, Sun [12] consider the matrix

$$(1.3) \quad S_p := \left[ \left( \frac{i^2 + j^2}{p} \right) \right]_{1 \leq i, j \leq (p-1)/2}$$

which involves the squares over  $\mathbb{F}_p$ . Sun [12, Theorem 1.2] proved that  $-\det S_p \bmod p\mathbb{Z}$  is a nonzero square over  $\mathbb{Z}/p\mathbb{Z} = \mathbb{F}_p$ . Moreover, Sun conjectured that  $-\det S_p$  is indeed a square of some integer whenever  $p \equiv 3 \pmod{4}$ . This conjecture was later confirmed by Alekseyev and Krachun. For the case  $p \equiv 1 \pmod{4}$ , if we write  $p = a^2 + 4b^2$  with  $a, b \in \mathbb{Z}$  and  $a \equiv 1 \pmod{4}$ , then Cohen, Sun and Vsemirnov conjectured that  $S_p/a$  is also a square of certain integer. This conjecture was later confirmed by the first author [17].

Recently, for any positive integer  $m$ , Sun [13] considered the matrix

$$(1.4) \quad S_p(m) := [(i^2 + j^2)^m]_{1 \leq i, j \leq (p-1)/2},$$

and obtained several interesting results concerning  $\det S_p(m)$ . For example, when  $p \equiv 3 \pmod{4}$ , Sun [13, Theorem 1.2(ii)] proved that

$$\det S_p(p-3) = \det \left[ \frac{1}{(i^2 + j^2)^2} \right]_{1 \leq i, j \leq (p-1)/2} \equiv \frac{1}{4} \prod_{r=1}^{(p-3)/4} \left( r + \frac{1}{4} \right)^2 \pmod{p\mathbb{Z}_p}.$$

After introducing the above relevant research results, we now state our research motivations. Inspired by Sun's matrix  $S_p(m)$  defined by Eq. (1.4), in this paper, we mainly consider a variant of  $S_p(m)$ . In fact, let  $q = 2n + 1$  and let

$$\{s_1 = 1, s_2, \dots, s_n\} = \{x^2 : x \in \mathbb{F}_q^\times\}.$$

Then we define the matrix

$$(1.5) \quad B_q(m) := [(s_i + s_j)^m]_{2 \leq i, j \leq n},$$

which concerns all nonzero squares over  $\mathbb{F}_q$ .

Before studying the properties of  $B_q(m)$ , let's first have a brief discussion. Let

$$h(t) = a_{m-1}t^{m-1} + \dots + a_1t + a_0$$

be a polynomial over a commutative ring with  $\deg(h(t)) \leq m-1$ . Then it is known that (see [6, Lemma 9])

$$(1.6) \quad \det [h(x_i + y_j)]_{1 \leq i, j \leq m} = a_{m-1}^l \cdot \prod_{r=0}^{l-1} \binom{l-1}{r} \cdot \prod_{1 \leq i < j \leq m} (x_i - x_j)(y_j - y_i).$$

By this we immediately see that  $B_q(m)$  is singular whenever  $m \leq n-3$ . Hence it is meaningful to consider the cases of  $n-2 \leq m \leq q-1$ . In this paper, we focus on the cases of  $m = n-2, n-1, n$ .

**1.3. The Pell sequence.** Before the statements of our main results, we introduce the Pell sequence. The Pell sequence is an infinite sequence of integers that comprise the denominators of the closest rational approximations to the square root of 2. Specifically, the Pell sequence  $\{P_i\}_{i=0}^{\infty}$  is defined by

$$P_0 = 0, P_1 = 1, P_{i+1} = 2P_i + P_{i-1},$$

and its companion sequence  $\{Q_i\}_{i=0}^{\infty}$  is defined by

$$Q_0 = Q_1 = 2, Q_{i+1} = 2Q_i + Q_{i-1}.$$

For any integer  $i \geq 0$ , it is also known that

$$P_i = \frac{1}{2\sqrt{2}} \left( (1 + \sqrt{2})^i - (1 - \sqrt{2})^i \right) = \sum_{k=0}^{\lfloor (i-1)/2 \rfloor} \binom{i}{2k+1} 2^k,$$

and

$$Q_i = \left( (1 + \sqrt{2})^i + (1 - \sqrt{2})^i \right).$$

By this, it is easy to see that

$$P_p \equiv 2^{(p-1)/2} \equiv \left( \frac{2}{p} \right) \pmod{p\mathbb{Z}},$$

and that

$$Q_p \equiv 2 \pmod{p\mathbb{Z}}.$$

**1.4. The first result: case  $m \in \{n-2, n-1\}$ .** Now we state our first result.

**Theorem 1.1.** *Let  $q = p^f = 2n + 1 \geq 7$  be an odd prime power with  $p$  prime and  $f \in \mathbb{Z}^+$ . Then the following results hold.*

(i)  $B_q(n-1)$  is a singular matrix whenever  $f \geq 2$ . Moreover, if  $f = 1$ , then

$$\det B_p(n-1) = \frac{2 \cdot ((n-1)!)^n}{(0!1! \cdots (n-1)!)^2} \cdot a_p \in \mathbb{F}_p,$$

where

$$a_p = \frac{2 - Q_p}{p} \pmod{p\mathbb{Z}} \in \mathbb{Z}/p\mathbb{Z} = \mathbb{F}_p.$$

(ii)  $B_q(n-2)$  is singular if and only if the integer  $f \geq 2$ . When  $f = 1$ , we have

$$\det B_p(n-2) = \left( -\frac{1}{2} \right)^{n-2} \cdot \frac{((n-2)!)^{n-1}}{(0!1! \cdots (n-2)!)^2} \in \mathbb{F}_p^\times.$$

By Theorem 1.1, we obtain the following result.

**Corollary 1.1.** *Let  $p = 2n + 1 \geq 7$  be an odd prime. Then the following results hold.*

(i)  $B_p(n-1)$  is singular if and only if

$$Q_p \equiv 2 \pmod{p^2\mathbb{Z}}.$$

Moreover, if  $p \equiv 1 \pmod{4}$ , then

$$\left( \frac{\det B_p(n-1)}{p} \right) = \left( \frac{2a_p}{p} \right).$$

If  $p \equiv 3 \pmod{4}$ , then

$$\left( \frac{\det B_p(n-1)}{p} \right) = (-1)^{\frac{h(-p)-1}{2}} \left( \frac{a_p}{p} \right),$$

where  $h(-p)$  is the class number of the imaginary quadratic field  $\mathbb{Q}(\sqrt{-p})$ .

(ii) Suppose  $p \equiv 1 \pmod{4}$ . Then

$$\left( \frac{\det B_p(n-2)}{p} \right) = \left( \frac{6}{p} \right).$$

Suppose  $p \equiv 3 \pmod{4}$ . Then

$$\left( \frac{\det B_p(n-2)}{p} \right) = \left( \frac{-2}{p} \right).$$

*Remark 1.1.* With the help of a computer, we verify that

$$\{p : p \text{ is a prime with } 7 \leq p \leq 10^5 \text{ and } Q_p \equiv 2 \pmod{p^2\mathbb{Z}}\} = \{13, 31\}.$$

Motivated by this, we pose the following conjecture.

**Conjecture 1.1.** *Let  $p \geq 7$  be an odd prime. Then  $B_p(n-1)$  is singular if and only if  $p \in \{13, 31\}$ .*

1.5. **The second result: case  $m = n$ .**

**Theorem 1.2.** *Let  $q = p^f = 2n + 1 \geq 7$  be an odd prime power with  $p$  prime and  $f \in \mathbb{Z}^+$ . Then  $B_q(n)$  is singular whenever  $f \geq 2$ . Moreover, if  $f = 1$ , then*

$$\det B_p(n) = (-1)^n \left( \frac{1}{2} \right)^{n-2} \cdot \frac{(n!)^{n+1}}{(0!1! \cdots n!)^2} \cdot b_p \in \mathbb{F}_p,$$

where

$$b_p = \frac{2p + \binom{2}{p} - P_p}{2p} \pmod{p\mathbb{Z}} \in \mathbb{Z}/p\mathbb{Z} = \mathbb{F}_p.$$

Similar to Corollary 1.1, we have the following result.

**Corollary 1.2.** *Let  $p = 2n + 1 \geq 7$  be a prime. Then*

$$\left( \frac{\det B_p(n)}{p} \right) = \left( \frac{-2b_p}{p} \right),$$

and hence  $B_p(n)$  is singular if and only if

$$P_p \equiv 2p + \left( \frac{2}{p} \right) \pmod{p^2\mathbb{Z}}.$$

**1.6. The third result: a variant of Carlitz's cyclotomic matrices.** Using essentially the same method appeared in the proof of Theorem 1.2, we can obtain some satisfactory results on certain variants of Carlitz's matrix  $C_p^-(\psi)$  and  $C_p^+(\psi)$  defined by Eq. (1.1) and Eq. (1.2) respectively.

In fact, let  $\mathbb{F}_q^\times = \{x_1 = 1, x_2, \dots, x_{q-1}\}$ . For any nontrivial character  $\psi \in \widehat{\mathbb{F}_q^\times}$ , we define

$$D_q^-(\psi) := [\psi(x_j - x_i)]_{2 \leq i, j \leq q-1},$$

and

$$D_q^+(\psi) := [\psi(x_j + x_i)]_{2 \leq i, j \leq q-1}.$$

We have the following result.

**Theorem 1.3.** *Let  $q = p^f \geq 3$  be an odd prime power. Then for any nontrivial character  $\psi \in \widehat{\mathbb{F}_q^\times}$ , we have*

$$\det D_q^-(\psi) = -\frac{(1 + \psi(-1))}{q^2} G_q(\psi)^{q-1},$$

and

$$\det D_q^+(\psi) = \frac{(-1)^{(q+1)/2} \cdot \psi(-1)}{q^2} \left(2 - \overline{\psi(2)}\right) G_q(\psi)^{q-1}.$$

Let  $\psi = \phi$  be the unique quadratic character of  $\mathbb{F}_q$ , i.e.,

$$\phi(x) = \begin{cases} 1 & \text{if } x \in \{s_1, s_2, \dots, s_{(q-1)/2}\}, \\ 0 & \text{if } x = 0, \\ -1 & \text{otherwise.} \end{cases}$$

Then  $D_p^-(\phi)$  is an integer matrices. Note that (see [3, Corollary 3.7.6])

$$G_q(\phi) = \begin{cases} (-1)^{f-1} \sqrt{q} & \text{if } p \equiv 1 \pmod{4}, \\ (-1)^{f-1} \mathbf{i}^f \sqrt{q} & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

By this and Theorem 1.3, we can obtain the following corollary directly.

**Corollary 1.3.** *Let  $q = p^f \geq 3$  be an odd prime power. Then  $D_q^-(\phi)$  is singular if and only if  $q \equiv 3 \pmod{4}$ . For  $q \equiv 1 \pmod{4}$ , we have*

$$\det D_q^-(\phi) = -2q^{(q-5)/2}.$$

## 2. PROOFS OF THEOREM 1.1 AND ITS COROLLARY

Recall that  $n = (q-1)/2$  and  $s_1 = 1, s_2, \dots, s_n$  are exactly all the non-zero squares over  $\mathbb{F}_q$ . Also, for any integers  $a, b$  with  $a \leq b$ , we use the symbol  $[a, b]$  to denote the set  $\{a, a+1, \dots, b\}$ .

We begin with the following result.

**Lemma 2.1.** *Let  $q = p^f = 2n+1 \geq 5$  be an odd prime power with  $p$  prime and  $f \in \mathbb{Z}^+$ . Then*

$$(-1)^{\frac{(n-1)(n-2)}{2}} \prod_{2 \leq i < j \leq n} (s_j - s_i)^2 = \left(-\frac{1}{2}\right)^{n-2} \in \mathbb{F}_p.$$

*Proof.* Let

$$F(t) = \frac{t^n - 1}{t - 1} = t^{n-1} + t^{n-2} + \cdots + t + 1$$

be a polynomial over  $\mathbb{F}_q$ . Note that  $x$  is a nonzero square over  $\mathbb{F}_q$  if and only if  $x^n = 1$ . Hence it is easy to verify that

$$(2.1) \quad F(t) = \frac{t^n - 1}{t - 1} = t^{n-1} + t^{n-2} + \cdots + t + 1 = \prod_{2 \leq j \leq n} (t - s_j).$$

By Eq. (2.1) we obtain that  $(-1)^{n-1} \prod_{2 \leq j \leq n} s_j$  is equal to the constant term of  $F(t)$ , i.e.,

$$(2.2) \quad \prod_{2 \leq j \leq n} s_j = (-1)^{n-1}.$$

Also, clearly  $(-1)^{n-1} \prod_{2 \leq j \leq n} (s_j - 1)$  is equal to the constant term of  $F(t + 1)$ , that is,

$$(2.3) \quad \prod_{2 \leq j \leq n} (s_j - 1) = (-1)^{n-1} \cdot n.$$

Next we consider the product

$$(-1)^{\frac{(n-1)(n-2)}{2}} \prod_{2 \leq i < j \leq n} (s_j - s_i)^2.$$

Let  $F'(t)$  be the formal derivative of  $F(t)$ . Then one can verify that

$$(2.4) \quad \begin{aligned} (-1)^{\frac{(n-1)(n-2)}{2}} \prod_{2 \leq i < j \leq n} (s_j - s_i)^2 &= \prod_{2 \leq i \neq j \leq n} (s_j - s_i) \\ &= \prod_{2 \leq j \leq n} \prod_{i \in [2, n] \setminus \{j\}} (s_j - s_i) \\ &= \prod_{2 \leq j \leq n} F'(s_j) \end{aligned}$$

As  $(t - 1)F(t) = t^n - 1$ , we have

$$F(t) + (t - 1)F'(t) = nt^{n-1},$$

and hence

$$F'(s_j) = \frac{ns_j^{n-1}}{s_j - 1} = \frac{n}{s_j(s_j - 1)}$$

for any  $j \in [2, n]$ . By the above results,

$$(2.5) \quad (-1)^{\frac{(n-1)(n-2)}{2}} \prod_{2 \leq i < j \leq n} (s_j - s_i)^2 = \prod_{2 \leq j \leq n} \frac{n}{s_j(s_j - 1)}.$$



Combining Eq. (2.5) with Eq. (2.2) and Eq. (2.3), we obtain

$$(-1)^{\frac{(n-1)(n-2)}{2}} \prod_{2 \leq i < j \leq n} (s_j - s_i)^2 = n^{n-2} = \left(-\frac{1}{2}\right)^{n-2} \in \mathbb{F}_p.$$

This completes the proof.  $\square$

Before the statement of next lemma, we introduce the following notations. Let  $l$  be a positive integer and let  $t_1, t_2, \dots, t_l$  be variables. Then for any  $k \in [1, l]$ , the  $k$ -th elementary symmetric polynomial of  $t_1, t_2, \dots, t_l$  is defined by

$$\sigma_k(t_1, t_2, \dots, t_l) = \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq l} \prod_{1 \leq j \leq k} t_{i_j}.$$

In addition, we let

$$\sigma_0(t_1, t_2, \dots, t_l) = 1.$$

In 2022, Grinberg, Sun and Zhao [4, Theorem 3.1] obtained the following result.

**Lemma 2.2.** *Let  $l$  be a positive integer. Then*

$$(2.6) \quad \det \left[ (x_i + y_j)^l \right]_{1 \leq i, j \leq l} = (-1)^{\frac{l(l-1)}{2}} \cdot \prod_{1 \leq i < j \leq l} (x_j - x_i)(y_j - y_i) \cdot \sum_{k=0}^l U_k,$$

where

$$U_k = \sigma_k(x_1, \dots, x_l) \sigma_{l-k}(y_1, \dots, y_l) \cdot \prod_{r \in [0, l] \setminus \{k\}} \binom{l}{r}.$$

Recall that the Pell sequence  $\{P_i\}_{i=0}^{\infty}$  is defined by

$$P_0 = 0, P_1 = 1, P_{i+1} = 2P_i + P_{i-1},$$

and its companion  $\{Q_i\}_{i=0}^{\infty}$  is defined by

$$Q_0 = Q_1 = 2, Q_{i+1} = 2Q_i + Q_{i-1}.$$

We now need the following results, which were obtained by Z.-H. Sun [9, Theorem 4.1], Z.-W. Sun [10, Final Remark] and Z.-W. Sun [11, Remark 3.1] respectively.

**Lemma 2.3.** *Let  $p \geq 7$  be a prime. Then the following congruences hold.*

$$(2.7) \quad \sum_{k=1}^{p-1} \frac{2^k}{k} \equiv \frac{2-2^p}{p} - \frac{7}{12} p^2 B_{p-3} \pmod{p^3 \mathbb{Z}_p},$$

where  $B_{p-3}$  is the  $(p-3)$ -th Bernoulli number.

$$(2.8) \quad \sum_{k=1}^{\frac{p-1}{2}} \frac{1}{k \cdot 2^k} \equiv -2^{\frac{p+1}{2}} \cdot \frac{P_p - 2^{\frac{p-1}{2}}}{p} \pmod{p \mathbb{Z}_p}.$$

$$(2.9) \quad 4 \binom{2}{p} P_p \equiv 2 + Q_p \pmod{p^2\mathbb{Z}}.$$

Now we are in a position to prove Theorem 1.1.

**Proof of Theorem 1.1.** (i) By Lemma 2.2 and Lemma 2.1 we obtain

$$(2.10) \quad \det B_q(n-1) = (-1)^{\frac{(n-1)(n-2)}{2}} \prod_{2 \leq i < j \leq n} (s_j - s_i)^2 \cdot \sum_{k=0}^{n-1} W_k = \left(-\frac{1}{2}\right)^{n-2} \cdot \sum_{k=0}^{n-1} W_k,$$

where

$$(2.11) \quad W_k = \sigma_k(s_2, \dots, s_n) \sigma_{n-1-k}(s_2, \dots, s_n) \cdot \prod_{r \in [0, n-1] \setminus \{k\}} \binom{n-1}{r}$$

for any  $k \in [0, n-1]$ .

We next consider  $W_k$ . By Eq. (2.1) we see that

$$F(t) = t^{n-1} + t^{n-2} + \dots + t + 1 = \prod_{2 \leq j \leq n} (t - s_j) = \sum_{k=0}^{n-1} (-1)^k \sigma_k(s_2, \dots, s_n) t^{n-k},$$

and hence for any  $k \in [0, n-1]$  we have

$$(2.12) \quad \sigma_k(s_2, \dots, s_n) = (-1)^k.$$

By Eq. (2.11) and Eq. (2.12) we obtain

$$(2.13) \quad W_k = (-1)^{n-1} \prod_{r \in [0, n-1] \setminus \{k\}} \binom{n-1}{r}.$$

Suppose first that  $f \geq 2$ . As  $q = p^f \geq 9$ , we have

$$n-1 = \frac{q-3}{2} = \frac{p-3}{2} \cdot 1 + \frac{p-1}{2} \cdot p + \dots + \frac{p-1}{2} \cdot p^{f-1} > \frac{p+1}{2}.$$

By the Lucas congruence, for  $j \in \{(p-1)/2, (p+1)/2\}$ , we have

$$\binom{n-1}{j} \equiv \binom{(p-3)/2}{j} \binom{(p-1)/2}{0} \dots \binom{(p-1)/2}{0} \equiv 0 \pmod{p\mathbb{Z}}.$$

Thus,

$$\left| \left\{ r \in [0, n-1] : \binom{n-1}{r} \equiv 0 \pmod{p\mathbb{Z}} \right\} \right| \geq 2.$$

This, together with Eq. (2.13), implies that  $W_k = 0$  (as an element of  $\mathbb{F}_q$ ) if  $f \geq 2$ . Hence by Eq. (2.10) we see that  $B_q(n-1)$  is singular whenever  $f \geq 2$ .

Suppose now  $f = 1$ . Then for any  $r \in [0, n-1]$  we have

$$\binom{n-1}{r} \not\equiv 0 \pmod{p\mathbb{Z}}.$$

Hence by Eq. (2.13) we obtain

$$\begin{aligned}
 \sum_{k=0}^{n-1} W_k &= (-1)^{n-1} \left( \sum_{r=0}^{n-1} \binom{n-1}{r}^{-1} \right) \cdot \prod_{r \in [0, n-1]} \binom{n-1}{r} \\
 (2.14) \qquad &= (-1)^{n-1} \cdot \frac{n}{2^n} \left( \sum_{k=1}^n \frac{2^k}{k} \right) \cdot \frac{((n-1)!)^n}{(0!1! \cdots (n-1)!)^2},
 \end{aligned}$$

where the last equality follows from Sury's identity [14]

$$\sum_{r=0}^{n-1} \binom{n-1}{r}^{-1} = \frac{n}{2^n} \sum_{k=1}^n \frac{2^k}{k}$$

and

$$\prod_{r \in [0, n-1]} \binom{n-1}{r} = \frac{((n-1)!)^n}{(0!1! \cdots (n-1)!)^2}.$$

We next consider

$$\sum_{k=1}^n \frac{2^k}{k} \pmod{p\mathbb{Z}_p}.$$

One can verify that

$$\begin{aligned}
 \sum_{k=1}^{p-1} \frac{2^k}{k} &= \sum_{k=1}^n \frac{2^k}{k} + \sum_{k=1}^n \frac{2^{p-k}}{p-k} \\
 &\equiv \sum_{k=1}^n \frac{2^k}{k} - 2 \sum_{k=1}^n \frac{1}{k \cdot 2^k} \pmod{p\mathbb{Z}_p}.
 \end{aligned}$$

Combining this with Eq. (2.7) and Eq. (2.8), we obtain

$$\begin{aligned}
 \sum_{k=1}^n \frac{2^k}{k} &\equiv \frac{2-2^p}{p} - 2^{\frac{p+3}{2}} \frac{P_p - 2^{\frac{p-1}{2}}}{p} \\
 &\equiv \frac{2-2^p}{p} - 4 \left( \frac{2}{p} \right) \frac{P_p - 2^{\frac{p-1}{2}}}{p} \pmod{p\mathbb{Z}_p}.
 \end{aligned}$$

By this and Eq. (2.9), we obtain

$$(2.15) \qquad \sum_{k=1}^n \frac{2^k}{k} \equiv \frac{-Q_p - 2^p + \left(\frac{2}{p}\right) 2^{\frac{p+3}{2}}}{p} \equiv \frac{2 - Q_p}{p} \pmod{p\mathbb{Z}_p},$$

where the last congruence follows from

$$2 - \left( -2^p + \left( \frac{2}{p} \right) 2^{\frac{p+3}{2}} \right) = 2 \left( 2^{\frac{p-1}{2}} - \left( \frac{2}{p} \right) \right)^2 \equiv 0 \pmod{p^2\mathbb{Z}}.$$

Now combining Eq. (2.15) with Eq. (2.14) and Eq. (2.10), one can verify that

$$\det B_p(n-1) = \frac{2 \cdot ((n-1)!)^n}{(0!1! \cdots (n-1)!)^2} \cdot a_p \in \mathbb{F}_p,$$

where

$$a_p = \frac{2 - Q_p}{p} \bmod p\mathbb{Z} \in \mathbb{Z}/p\mathbb{Z} = \mathbb{F}_p.$$

This completes the proof of (i).

(ii) Let  $h(t) = t^{n-2}$ . Then by Eq. (1.6) and Lemma 2.1 we obtain

$$\begin{aligned} \det B_q(n-2) &= \prod_{r=0}^{n-2} \binom{n-2}{r} \cdot \prod_{2 \leq i < j \leq n} -(s_j - s_i)^2 \\ &= \prod_{r=0}^{n-2} \binom{n-2}{r} \cdot (-1)^{\frac{(n-1)(n-2)}{2}} \prod_{2 \leq i < j \leq n} (s_j - s_i)^2 \\ (2.16) \quad &= \left(-\frac{1}{2}\right)^{n-2} \cdot \prod_{r=0}^{n-2} \binom{n-2}{r}. \end{aligned}$$

Suppose first that  $f \geq 2$ . Noting that  $q \geq 9$ , we obtain

$$n-2 = \frac{q-5}{2} = \frac{p-5}{2} \cdot 1 + \frac{p-1}{2} \cdot p + \cdots + \frac{p-1}{2} \cdot p^{f-1} > \frac{p-1}{2}.$$

By the Lucas congruence again, we clearly have

$$\binom{n-2}{(p-1)/2} \equiv \binom{(p-5)/2}{(p-1)/2} \binom{(p-1)/2}{0} \cdots \binom{(p-1)/2}{0} \equiv 0 \pmod{p\mathbb{Z}}.$$

By this and Eq. (2.16), we see that  $\det B_q(n-2) = 0$  whenever  $f \geq 2$ .

Suppose now  $f = 1$ . Then  $\binom{n-2}{r} \not\equiv 0 \pmod{p\mathbb{Z}}$  for any  $r \in [0, n-2]$ . Hence by Eq. (2.16) again we obtain

$$\det B_p(n-2) = \left(-\frac{1}{2}\right)^{n-2} \cdot \frac{((n-2)!)^{n-1}}{(0!1! \cdots (n-2)!)^2} \in \mathbb{F}_p^\times.$$

In view of the above, we have completed the proof. □

We next prove Corollary 1.1.

**Proof of Corollary 1.1.** (i) If  $p \equiv 1 \pmod{4}$ , then  $n$  is even. Thus,

$$\left(\frac{\det B_p(n-1)}{p}\right) = \left(\frac{2a_p}{p}\right) \left(\frac{(n-1)!}{p}\right)^n = \left(\frac{2a_p}{p}\right).$$

Suppose now  $p \equiv 3 \pmod{4}$  and  $p > 3$ . Then  $n$  is odd and hence

$$\left(\frac{\det B_p(n-1)}{p}\right) = \left(\frac{2a_p}{p}\right) \left(\frac{(n-1)!}{p}\right) = \left(\frac{-a_p}{p}\right) \left(\frac{n!}{p}\right) = (-1)^{\frac{h(-p)-1}{2}} \left(\frac{a_p}{p}\right).$$

The last equality follows from the Mordell congruence [8]

$$n! = \frac{p-1}{2}! \equiv (-1)^{\frac{h(-p)+1}{2}} \pmod{p\mathbb{Z}},$$

where  $h(-p)$  is the class number of the imaginary quadratic field  $\mathbb{Q}(\sqrt{-p})$ . This completes the proof of (i).

(ii) Suppose  $p \equiv 1 \pmod{4}$ . Then by the Wilson congruence, one can verify that

$$-1 \equiv (p-1)! \equiv (-1)^n \cdot (n!)^2 \equiv (n!)^2 \pmod{p\mathbb{Z}}.$$

This implies  $n! = \sqrt{-1}$  over  $\mathbb{F}_p$ . Thus,

$$\left(\frac{n!}{p}\right) \equiv (n!)^n \equiv (-1)^{n/2} \equiv \left(\frac{2}{p}\right) \pmod{p\mathbb{Z}},$$

that is,

$$\left(\frac{n!}{p}\right) = \left(\frac{2}{p}\right).$$

Now by this and Theorem 1.1(ii), and noting that  $n$  is even in this case, we obtain

$$\left(\frac{\det B_p(n-2)}{p}\right) = \left(\frac{(n-2)!}{p}\right) = \left(\frac{n!}{p}\right) \left(\frac{(n-1)n}{p}\right) = \left(\frac{6}{p}\right),$$

where the last equality follows from  $n(n-1) = 3/4$  over  $\mathbb{F}_p$ .

Suppose now  $p \equiv 3 \pmod{4}$ . Then  $n$  is odd. By Theorem 1.1(ii) we clearly have

$$\left(\frac{\det B_p(n-2)}{p}\right) = \left(\frac{-2}{p}\right).$$

In view of the above, we have completed the proof.  $\square$

### 3. PREPARATIONS FOR THE PROOF OF THEOREM 1.2

**3.1. A lemma on almost circulant matrices.** Let  $n \geq 2$  be an integer and let  $\mathbf{v} = (a_0, a_1, \dots, a_{n-1}) \in \mathbb{C}^n$ . The circulant matrix of  $\mathbf{v}$  is an  $n \times n$  matrix defined by

$$C_n(\mathbf{v}) = [a_{j-i}]_{0 \leq i, j \leq n-1},$$

where  $a_s = a_t$  whenever  $s \equiv t \pmod{n}$ , that is,

$$C_n(\mathbf{v}) = \begin{bmatrix} a_0 & a_1 & \cdots & a_{n-2} & a_{n-1} \\ a_{n-1} & a_0 & \cdots & a_{n-3} & a_{n-2} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ a_2 & a_3 & \cdots & a_0 & a_1 \\ a_1 & a_2 & \cdots & a_{n-1} & a_0 \end{bmatrix}.$$

Recently, the first author and the second author [18, Theorem 4.1] defined the almost circulant matrix  $W_n(\mathbf{v})$  of  $\mathbf{v}$  by

$$W_n(\mathbf{v}) = [a_{j-i}]_{1 \leq i, j \leq n-1},$$

and obtained the following result.

**Lemma 3.1.** *Let  $\lambda_0, \lambda_1, \dots, \lambda_{n-1}$  be all the eigenvalues of  $C_n(\mathbf{v})$ . Then*

$$\det W_n(\mathbf{v}) = \frac{1}{n} \sum_{l=0}^{n-1} \prod_{k \in [0, n-1] \setminus \{l\}} \lambda_k.$$

**3.2. Some  $p$ -adic preparations.** Let  $\pi \in \mathbb{C}_p$  with  $\pi^{p-1} + p = 0$  and let  $\zeta_\pi \in \mathbb{C}_p$  be a primitive  $p$ -th root of unity with  $\zeta_\pi \equiv 1 + \pi \pmod{\pi^2}$ .

Recall that  $q = p^f$ . Let  $\zeta_{q-1} \in \mathbb{C}_p$  be a primitive  $(q-1)$ -th root of unity. Then it is known that  $\mathbb{Q}_p(\zeta_{q-1})/\mathbb{Q}_p$  is an unramified extension and  $[\mathbb{Q}_p(\zeta_{q-1}) : \mathbb{Q}_p] = f$ . Hence

$$\mathbb{Z}_p[\zeta_{q-1}]/\mathfrak{p} \cong \mathbb{F}_q,$$

where  $\mathfrak{p} = p\mathbb{Z}_p[\zeta_{q-1}]$  is a prime ideal of  $\mathbb{Z}[\zeta_{q-1}]$ . From now on, we identify  $\mathbb{F}_q$  with  $\mathbb{Z}_p[\zeta_{q-1}]/\mathfrak{p}$ . The Teichmüller character  $\omega_q : \mathbb{F}_q \rightarrow \mathbb{C}_p$  is a multiplicative character of  $\mathbb{F}_q$  defined by

$$(3.1) \quad \omega_q(x \bmod \mathfrak{p}) \equiv x \pmod{\mathfrak{p}}$$

for any  $x \in \mathbb{Z}_p[\zeta_{q-1}]$ . Also, it is easy to verify that  $\omega_q$  is a generator of  $\widehat{\mathbb{F}_q^\times}$ . For any integer  $r \in [0, q-2]$ , letting

$$r = r_0 \cdot 1 + r_1 \cdot p + \dots + r_{f-1} \cdot p^{f-1}$$

be the decomposition of  $r$  in the base  $p$ , we define

$$(3.2) \quad s(r) = \sum_{i=0}^{f-1} r_i.$$

For  $s(r)$ , we have the following known result (see [3, Lemma 3.6.7]).

**Lemma 3.2.** *For any  $r \in [0, q-2]$  we have*

$$s(r) = (p-1) \sum_{i=0}^{f-1} \left\{ \frac{rp^i}{q-1} \right\},$$

where  $\{x\}$  denotes the fractional part of a real number  $x$ .

Recall that  $n = (q-1)/2 = (p^f - 1)/2$ . Using Lemma 3.2, we obtain the following result.

**Lemma 3.3.** *Suppose  $r \in [0, n-1]$ . Then*

$$s(n) + s(n+r) > s(r).$$

*Proof.* For any  $i \in [0, f-1]$ , we set

$$x_i(r) := \left\{ \frac{rp^i}{q-1} \right\}.$$

Then it is easy to verify that

$$(3.3) \quad \frac{1}{2} + \left\{ \frac{1}{2} + x_i(r) \right\} = \begin{cases} x_i(r) + 1 & \text{if } 0 \leq x_i(r) < 1/2, \\ x_i(r) & \text{if } 1/2 \leq x_i(r) < 1. \end{cases}$$

As  $0 \leq r \leq n-1$ , we have  $x_0(r) < 1/2$ . Hence by Eq. (3.3) we obtain

$$\frac{1}{2} + \left\{ \frac{1}{2} + x_0(r) \right\} > x_0(r).$$

By this, Lemma 3.2 and Eq. (3.3), one can verify that

$$\begin{aligned} s(n) + s(n+r) &= (p-1) \sum_{i=0}^{f-1} \left( \frac{1}{2} + \left\{ \frac{(n+r)p^i}{q-1} \right\} \right) \\ &= (p-1) \sum_{i=0}^{f-1} \left( \frac{1}{2} + \left\{ \frac{p^i}{2} + \frac{rp^i}{q-1} \right\} \right) \\ &= (p-1) \sum_{i=0}^{f-1} \left( \frac{1}{2} + \left\{ \frac{p^i-1}{2} + \frac{1}{2} + \frac{rp^i}{q-1} \right\} \right) \\ &= (p-1) \sum_{i=0}^{f-1} \left( \frac{1}{2} + \left\{ \frac{1}{2} + x_i(r) \right\} \right) \\ &> (p-1) \sum_{i=0}^{f-1} x_i(r) \\ &= s(r). \end{aligned}$$

This completes the proof.  $\square$

Recall that  $p$  is an odd prime. We next introduce the  $p$ -adic Gamma function  $\Gamma_p$ . For any  $n \in \mathbb{Z}^+$ , we define

$$\Gamma_p(n) = (-1)^n \prod_{0 < k < n, p \nmid k} k \in \mathbb{Z}_p^\times.$$

Since  $\mathbb{Z}^+$  is dense in  $\mathbb{Z}_p$  and  $\mathbb{Z}_p^\times$  is a closed multiplicative group, the  $p$ -adic Gamma function  $\Gamma_p : \mathbb{Z}_p \rightarrow \mathbb{Z}_p^\times$  is defined by

$$\Gamma_p(x) = \lim_{i \rightarrow \infty} \Gamma_p(x_i),$$

where  $\{x_i\}_{i=1}^\infty$  is a sequence of positive integers  $x_i$  with  $\lim_{i \rightarrow \infty} x_i = x$ .

In this paper, we need following result on  $\Gamma_p$ .

**Lemma 3.4.** *Suppose that  $p \geq 5$  is a prime. Let  $n \in \mathbb{Z}^+$ . Then for any  $x, y \in \mathbb{Z}_p$  we have*

$$x \equiv y \pmod{p^n \mathbb{Z}_p} \Rightarrow \Gamma_p(x) \equiv \Gamma_p(y) \pmod{p^n \mathbb{Z}_p}.$$

We conclude this section with the following result, which is known as the Gross-Koblitz formula [5, Theorem 1.7].

**Lemma 3.5.** *For any  $r \in [0, q-2]$ , let the Gauss sum*

$$G_q(\omega_q^{-r}) = \sum_{x \in \mathbb{F}_q} \omega_q^{-r}(x) \zeta_\pi^{\text{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(x)}.$$

Then

$$G_q(\omega_q^{-r}) = -\pi^{s(r)} \cdot \prod_{i=0}^{f-1} \Gamma_p \left( \left\{ \frac{rp^i}{q-1} \right\} \right),$$

where  $\Gamma_p$  is the  $p$ -adic Gamma function and  $\{x\}$  is the fractional part of a real number  $x$ .

#### 4. PROOF OF THEOREM 1.2

Let notations be as in Section 3. In this section, we fix a generator  $g$  of the cyclic group  $\mathbb{F}_q^\times$ . Recall that  $n = (q-1)/2$ . For any  $i \in [0, n-1]$ , we let

$$a_i = \omega_q^{-n} (1 + g^{2i}),$$

and let

$$\mathbf{v} = (a_0, a_1, \dots, a_{n-1}).$$

We begin with the following result.

**Lemma 4.1.** *For any  $r \in [0, n-1]$ , let*

$$(4.1) \quad \lambda_r := \frac{(-1)^r}{2} J_q(\omega_q^{-n}, \omega_q^{-r}) + \frac{(-1)^{n+r}}{2} J_q(\omega_q^{-n}, \omega_q^{-(n+r)}).$$

Then  $\lambda_0 = -1, \lambda_1, \dots, \lambda_{n-1}$  are exactly all the eigenvalues of the circulant matrix  $C_n(\mathbf{v})$ .

*Proof.* For any  $r \in [0, n-1]$ , one can verify that

$$\begin{aligned} & \sum_{0 \leq j \leq n-1} \omega_q^{-n} (1 + g^{2j-2i}) \omega_q^{-r} (g^{2j}) \\ &= \sum_{0 \leq j \leq n-1} \omega_q^{-n} (1 + g^{2j-2i}) \omega_q^{-r} (g^{2j-2i}) \omega_q^{-r} (g^{2i}) \\ &= \sum_{0 \leq j \leq n-1} \omega_q^{-n} (1 + g^{2j}) \omega_q^{-r} (g^{2j}) \omega_q^{-r} (g^{2i}). \end{aligned}$$

This implies that for any  $r \in [0, n-1]$ ,

$$C_n(\mathbf{v}) \mathbf{u}_r = y_r \mathbf{u}_r,$$

where the column vector

$$\mathbf{u}_r = (\omega_q^{-r}(g^0), \omega_q^{-r}(g^2), \dots, \omega_q^{-r}(g^{2(n-1)}))^T,$$

and

$$y_r = \sum_{0 \leq j \leq n-1} \omega_q^{-n} (1 + g^{2j}) \omega_q^{-r} (g^{2j}).$$

As  $\mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{n-1}$  are clearly linearly independent over  $\mathbb{C}$ , the numbers  $y_0, y_1, \dots, y_{n-1}$  are all the eigenvalues of  $C_n(\mathbf{v})$ . Note that

$$\frac{1}{2} (\varepsilon(x) + \omega_q^{-n}(x)) = \begin{cases} 1 & \text{if } x \in \{s_1, s_2, \dots, s_n\}, \\ 0 & \text{otherwise.} \end{cases}$$



Thus, for each  $r \in [0, n-1]$ , we have

$$\begin{aligned}
 y_r &= \sum_{0 \leq j \leq n-1} \omega_q^{-n} (1 + g^{2j}) \omega_q^{-r} (g^{2j}) \\
 &= \sum_{1 \leq j \leq n} \omega_q^{-n} (1 + s_j) \omega_q^{-r} (s_j) \\
 &= \frac{1}{2} \sum_{x \in \mathbb{F}_q} (\varepsilon(x) + \omega_q^{-n}(x)) \omega_q^{-n} (1 + x) \omega_q^{-r} (x) \\
 &= \frac{(-1)^r}{2} \sum_{x \in \mathbb{F}_q} \omega_q^{-n} (1 + x) \omega_q^{-r} (-x) + \frac{(-1)^{n+r}}{2} \sum_{x \in \mathbb{F}_q} \omega_q^{-n} (1 + x) \omega_q^{-(n+r)} (-x) \\
 &= \lambda_r.
 \end{aligned}$$

For  $\lambda_0$ , one can verify that

$$\begin{aligned}
 \lambda_0 &= \frac{1}{2} J_q (\omega_q^{-n}, \varepsilon) + \frac{(-1)^n}{2} J_q (\omega_q^{-n}, \omega_q^{-n}) \\
 &= \frac{1}{2} J_q (\omega_q^{-n}, \varepsilon) + \frac{(-1)^n}{2} J_q (\omega_q^{-n}, \omega_q^n) \\
 &= -\frac{1}{2} \omega_q(1) + \frac{(-1)^n}{2} (-\omega_q^{-n}(-1)) \\
 &= -1.
 \end{aligned}$$

In view of the above, we have completed the proof.  $\square$

Now we are in a position to prove Theorem 1.2.

**Proof of Theorem 1.2.** Recall that  $g$  is a generator of  $\mathbb{F}_q^\times$ . Then it is clear that

$$\det B_q(n) = \det \left[ \left( 1 + \frac{s_j}{s_i} \right)^n \right]_{2 \leq i, j \leq n} = \det [(1 + g^{2j-2i})^n]_{1 \leq i, j \leq n-1}.$$

By the definition (3.1) of the Teichmüller character  $\omega_q$  of  $\mathbb{F}_q$ , we see that

$$(4.2) \quad \det B_q(n) = \det [\omega_q (1 + g^{2j-2i})]_{1 \leq i, j \leq n-1} \pmod{\mathfrak{p}} = \det W_n(\mathbf{v}) \pmod{\mathfrak{p}},$$

where

$$\mathbf{v} = (\omega_q^{-n} (1 + g^0), \omega_q^{-n} (1 + g^2), \dots, \omega_q^{-n} (1 + g^{2(n-1)})),$$

and  $\mathfrak{p} = p\mathbb{Z}_p[\zeta_{q-1}]$  is a prime ideal. Hence we focus on the determinant of the almost circulant matrix  $W_n(\mathbf{v})$ . By Lemma 4.1 and Lemma 3.1 we have

$$(4.3) \quad \det B_q(n) = \det W_n(\mathbf{v}) \pmod{\mathfrak{p}} = \frac{1}{n} \sum_{l=0}^{n-1} \prod_{r \in [0, n-1] \setminus \{l\}} \lambda_r \pmod{\mathfrak{p}},$$

where  $\lambda_r$  is defined by Eq. (4.1). We next consider  $\lambda_r \pmod{\mathfrak{p}}$ . Suppose  $r \in [1, n-1]$ . Then

$$J_q(\omega_q^{-n}, \omega_q^{-(n+r)}) = \frac{G_q(\omega_q^{-n})G_q(\omega_q^{-(n+r)})}{G_q(\omega_q^{-r})}.$$

By Lemma 3.5 and Lemma 3.3, we obtain

$$\text{ord}_p(J_q(\omega_q^{-n}, \omega_q^{-(n+r)})) = \frac{1}{p-1}(s(n) + s(n+r) - s(r)) > 0.$$

This implies that

$$J_q(\omega_q^{-n}, \omega_q^{-(n+r)}) \equiv 0 \pmod{\mathfrak{p}}$$

for any  $r \in [1, n-1]$ . Hence by Eq. (4.1) we have

$$(4.4) \quad \lambda_r \equiv \frac{(-1)^r}{2} J_q(\omega_q^{-n}, \omega_q^{-r}) \pmod{\mathfrak{p}}$$

whenever  $r \in [1, n-1]$ .

**Case I:**  $f \geq 2$ .

Suppose  $r \in [1, n-1]$ . Then

$$J_q(\omega_q^{-n}, \omega_q^{-r}) = \frac{G_q(\omega_q^{-n})G_q(\omega_q^{-r})}{G_q(\omega_q^{-(n+r)})}.$$

By Lemma 3.5 we obtain

$$(4.5) \quad \text{ord}_p(J_q(\omega_q^{-n}, \omega_q^{-r})) = \frac{1}{p-1}(s(n) + s(r) - s(n+r)).$$

As  $q = p^f \geq 9$ , we have  $n-1 \geq (p+3)/2$ . By this, for  $r \in \{(p+1)/2, (p+3)/2\} \subseteq [1, n-1]$ , we have

$$s(n) + s(r) > s(n+r).$$

This, together with Eq (4.5), implies that

$$|\{1 \leq r \leq n-1 : J_q(\omega_q^{-n}, \omega_q^{-r}) \equiv 0 \pmod{\mathfrak{p}}\}| = |\{1 \leq r \leq n-1 : \lambda_r \equiv 0 \pmod{\mathfrak{p}}\}| \geq 2.$$

By this, Eq. (4.4) and Eq. (4.3), we see that  $B_q(n)$  is singular if  $f \geq 2$ .

**Case II:**  $f = 1$ .

Note first that in this case,  $\mathbb{Q}_p(\zeta_{q-1}) = \mathbb{Q}_p$  and  $\mathfrak{p} = p\mathbb{Z}_p$ . By Eq. (4.5), for any  $r \in [1, n-1]$ , we see that

$$\text{ord}_p(J_p(\omega_p^{-n}, \omega_p^{-r})) = \frac{1}{p-1}(s(n) + s(r) - s(n+r)) = \frac{1}{p-1}(n+r - (n+r)) = 0.$$

Thus, for any  $r \in [1, n-1]$ ,

$$\lambda_r \equiv \frac{(-1)^r}{2} J_q(\omega_p^{-n}, \omega_p^{-r}) \not\equiv 0 \pmod{p\mathbb{Z}_p},$$

that is,  $\lambda_r$  is a  $p$ -adic unit. By this and Eq. (4.3), we obtain

$$(4.6) \quad \det B_q(n) = -2\lambda_0\lambda_1 \cdots \lambda_{n-1} \left( \frac{1}{\lambda_0} + \frac{1}{\lambda_1} + \cdots + \frac{1}{\lambda_{n-1}} \right) \pmod{p\mathbb{Z}_p}.$$

We next consider  $\lambda_r \pmod{p\mathbb{Z}_p}$ . By Lemma 3.5 and Lemma 3.4. Suppose  $r \in [1, n-1]$ . Then

$$\begin{aligned} \lambda_r &\equiv \frac{(-1)^r}{2} J_q(\omega_p^{-n}, \omega_p^{-r}) \\ &\equiv \frac{(-1)^r}{2} \frac{G_p(\omega_p^{-n})G_p(\omega_p^{-r})}{G_p(\omega_p^{-(n+r)})} \\ &\equiv \frac{(-1)^{r+1}}{2} \cdot \frac{\Gamma_p\left(\frac{n}{p-1}\right)\Gamma_p\left(\frac{r}{p-1}\right)}{\Gamma_p\left(\frac{n+r}{p-1}\right)} \\ &\equiv \frac{(-1)^{r+1}}{2} \cdot \frac{\Gamma_p(p-n)\Gamma_p(p-r)}{\Gamma_p(p-n-r)} \\ &\equiv \frac{(-1)^r}{2} \frac{(p-n-1)! \cdot (p-r-1)!}{(p-1-n-r)!} \\ &\equiv \frac{(-1)^{r+1}}{2} \frac{(n+r)!}{n! \cdot r!} \\ &\equiv \frac{(-1)^{r+1}}{2} \binom{n+r}{r} \\ &\equiv -\frac{1}{2} \binom{n}{r} \pmod{p\mathbb{Z}_p}. \end{aligned}$$

The last congruence follows from

$$(-1)^r \cdot \binom{n+r}{r} \equiv \binom{n}{r} \pmod{p\mathbb{Z}}$$

for any  $r \in [1, n-1]$ .

By the above results and Sury's identity [14], we obtain

$$\begin{aligned} &-2\lambda_0\lambda_1 \cdots \lambda_{n-1} \left( \frac{1}{\lambda_0} + \frac{1}{\lambda_1} + \cdots + \frac{1}{\lambda_{n-1}} \right) \\ &\equiv (-1)^{n-1} \left( \frac{1}{2} \right)^{n-2} \cdot \prod_{r=0}^n \binom{n}{r} \cdot \left( 1 - 2 \sum_{r=0}^n \binom{n}{r}^{-1} \right) \\ &\equiv (-1)^{n-1} \left( \frac{1}{2} \right)^{n-2} \cdot \frac{(n!)^{n+1}}{(0!1! \cdots n!)^2} \cdot \left( 1 - \frac{n+1}{2^n} \sum_{k=1}^{n+1} \frac{2^k}{k} \right) \end{aligned}$$

$$\begin{aligned}
&\equiv (-1)^{n-1} \left(\frac{1}{2}\right)^{n-2} \cdot \frac{(n!)^{n+1}}{(0!1! \cdots n!)^2} \cdot \left(1 - \frac{1}{2^{n+1}} \sum_{k=1}^{n+1} \frac{2^k}{k}\right) \\
&\equiv (-1)^{n-1} \left(\frac{1}{2}\right)^{n-2} \cdot \frac{(n!)^{n+1}}{(0!1! \cdots n!)^2} \cdot \left(-1 - \frac{1}{2^{n+1}} \sum_{k=1}^n \frac{2^k}{k}\right) \pmod{p\mathbb{Z}_p}.
\end{aligned}$$

By this and using Eq. (2.15) and Eq. (2.9), we finally obtain

$$\begin{aligned}
&-2\lambda_0\lambda_1 \cdots \lambda_{n-1} \left(\frac{1}{\lambda_0} + \frac{1}{\lambda_1} + \cdots + \frac{1}{\lambda_{n-1}}\right) \\
&\equiv (-1)^n \left(\frac{1}{2}\right)^{n-2} \cdot \frac{(n!)^{n+1}}{(0!1! \cdots n!)^2} \cdot \frac{2p + \binom{2}{p} - P_p}{2p} \pmod{p\mathbb{Z}_p}.
\end{aligned}$$

In view of the above, we have completed the proof.  $\square$

## 5. PROOF OF THEOREM 1.3

We begin with the following lemma.

**Lemma 5.1.** *Let  $\chi_q$  be a generator of  $\widehat{\mathbb{F}_q^\times}$ . Then for any nontrivial character  $\psi \in \widehat{\mathbb{F}_q^\times}$ , the following results hold.*

$$(5.1) \quad A_q := \prod_{r=0}^{q-2} J_q(\psi, \chi_q^r) = \frac{G_q(\psi)^{q-1}}{q}.$$

$$(5.2) \quad S_q = \sum_{r=0}^{q-2} \frac{1}{J_q(\psi, \chi_q^r)} = \frac{1-q}{q} (1 + \psi(-1)).$$

$$(5.3) \quad T_q := \sum_{r=0}^{q-2} \frac{(-1)^r}{J_q(\psi, \chi_q^r)} = \frac{1-q}{q} (2 - \overline{\psi(2)}).$$

*Proof.* Since  $\chi_q$  is a generator of  $\widehat{\mathbb{F}_q^\times}$  and  $\psi \neq \varepsilon$ , there is an integer  $j \in [1, q-2]$  such that  $\chi_q^{-j} = \psi$ . One can verify that

$$\begin{aligned}
\prod_{r=0}^{q-2} J_q(\psi, \chi_q^r) &= \prod_{r=0}^{q-2} J_q(\chi_q^{-j}, \chi_q^r) \\
&= J_q(\chi_q^{-j}, \chi_q^j) \cdot \prod_{r \in [0, q-2] \setminus \{j\}} J_q(\chi_q^{-j}, \chi_q^r) \\
&= (-1)^{j+1} \cdot \prod_{r \in [0, q-2] \setminus \{j\}} \frac{G_q(\chi_q^{-j}) G_q(\chi_q^r)}{G_q(\chi_q^{-j+r})}
\end{aligned}$$

$$\begin{aligned}
 &= (-1)^{j+1} \cdot \frac{G_q(\varepsilon)}{G_q(\chi^{-j})G_q(\chi^j)} \cdot \prod_{r \in [0, q-2]} \frac{G_q(\chi_q^{-j})G_q(\chi_q^r)}{G_q(\chi_q^{-j+r})} \\
 &= \frac{G_q(\chi_q^{-j})^{q-1}}{q} \cdot \prod_{r \in [0, q-2]} \frac{G_q(\chi_q^r)}{G_q(\chi_q^{-j+r})} \\
 &= \frac{G_q(\psi)^{q-1}}{q}.
 \end{aligned}$$

This completes the proof of Eq. (5.1).

Now we prove Eq. (5.2). As  $|J_q(\chi_q^{-j}, \chi_q^r)| = \sqrt{q}$  whenever  $r \in [1, q-2] \setminus \{j\}$ , one can verify that

$$\begin{aligned}
 \sum_{r=0}^{q-2} \frac{1}{J_q(\psi, \chi_q^r)} &= \sum_{r=0}^{q-2} \frac{1}{J_q(\chi_q^{-j}, \chi_q^r)} \\
 &= \frac{1}{J_q(\chi_q^{-j}, \varepsilon)} + \frac{1}{J_q(\chi_q^{-j}, \chi_q^j)} + \frac{1}{q} \sum_{r \in [1, q-2] \setminus \{j\}} \frac{q}{J_q(\chi_q^{-j}, \chi_q^r)} \\
 &= -(1 + (-1)^j) + \frac{1}{q} \sum_{r \in [1, q-2] \setminus \{j\}} \overline{J_q(\chi_q^{-j}, \chi_q^r)} \\
 &= -(1 + \psi(-1)) + \frac{1}{q} \sum_{r \in [0, q-2]} \overline{J_q(\chi_q^{-j}, \chi_q^r)} - \frac{1}{q} \overline{J_q(\psi, \varepsilon)} - \frac{1}{q} \overline{J_q(\psi, \psi^{-1})} \\
 &= \frac{1-q}{q} (1 + \psi(-1)),
 \end{aligned}$$

where the last equality follows from

$$\sum_{r \in [0, q-2]} J_q(\chi_q^{-j}, \chi_q^r) = 0.$$

This completes the proof of Eq. (5.2)

Finally, we prove Eq. (5.3). We first claim that

$$(5.4) \quad \sum_{r \in [0, (q-3)/2]} J_q(\psi, \chi_q^{2r}) = \frac{(q-1)\psi(2)}{2}.$$

In fact, it is easy to verify that

$$\begin{aligned}
 \sum_{r \in [0, (q-3)/2]} J_q(\psi, \chi_q^{2r}) &= \sum_{r \in [0, (q-3)/2]} \sum_{x \in \mathbb{F}_q} \psi(1-x) \chi_q^{2r}(x) \\
 &= \sum_{x \in \mathbb{F}_q} \psi(1-x) \sum_{r \in [0, q-2]} \chi_q^{2r}(x) \\
 &= \frac{q-1}{2} \psi(2),
 \end{aligned}$$

where the last equality follows from

$$\sum_{r \in [0, q-2]} \chi_q^{2r}(x) = \begin{cases} (q-1)/2 & \text{if } x = \pm 1, \\ 0 & \text{otherwise.} \end{cases}$$

By Eq. (5.4) we obtain

$$\begin{aligned} S_q + T_q &= 2 \sum_{r \in [0, (q-3)/2]} \frac{1}{J_q(\psi, \chi_q^{2r})} \\ &= \frac{2}{J_q(\psi, \varepsilon)} + \frac{1 + \psi(-1)}{J_q(\psi, \psi^{-1})} + \frac{2}{q} \sum_{\substack{r \in [1, (q-3)/2] \\ \chi_q^{2r} \neq \psi^{-1}}} \frac{q}{J_q(\psi, \chi_q^{2r})} \\ &= -(3 + \psi(-1)) + \frac{2}{q} \sum_{\substack{r \in [1, (q-3)/2] \\ \chi_q^{2r} \neq \psi^{-1}}} \overline{J_q(\psi, \chi_q^{2r})} \\ &= -(3 + \psi(-1)) + \frac{2}{q} \sum_{r \in [0, (q-3)/2]} \overline{J_q(\psi, \chi_q^{2r})} - \frac{2}{q} \overline{J_q(\psi, \varepsilon)} - \frac{1 + \psi(-1)}{q} \overline{J_q(\psi, \psi^{-1})} \\ &= \frac{1-q}{q} \left( 3 + \psi(-1) - \overline{\psi(2)} \right). \end{aligned}$$

Combining this with Eq. (5.2), we have

$$T_q = (S_q + T_q) - S_q = \frac{1-q}{q} (2 - \overline{\psi(2)}).$$

In view of the above, we have completed the proof.  $\square$

From now on, we fix a generator  $g$  of  $\mathbb{F}_q^\times$ . For any nontrivial character  $\psi \in \widehat{\mathbb{F}_q^\times}$ , define the circulant matrices

$$(5.5) \quad M_q(\psi) := [\psi(g^{j-i} - 1)]_{0 \leq i, j \leq q-2},$$

and

$$(5.6) \quad N_q(\psi) := [\psi(g^{j-i} + 1)]_{0 \leq i, j \leq q-2}.$$

We need the following lemma.

**Lemma 5.2.** *Let  $\chi_q$  be a generator of  $\widehat{\mathbb{F}_q^\times}$ . Then the numbers*

$$\alpha_r = \psi(-1) J_q(\psi, \chi_q^r) \quad (r = 0, 1, \dots, q-2)$$

*are exactly all the eigenvalues of  $M_q(\psi)$ . Also, the numbers*

$$\beta_r = (-1)^r J_q(\psi, \chi_q^r) \quad (r = 0, 1, \dots, q-2)$$

*are exactly all the eigenvalues of  $N_q(\psi)$ .*

*Proof.* For any  $r \in [0, q-2]$ , it is easy to see that

$$\begin{aligned}
 \sum_{j=0}^{q-2} \psi(g^{j-i} - 1) \chi_q^r(g^j) &= \sum_{j=0}^{q-2} \psi(g^{j-i} - 1) \chi_q^r(g^{j-i}) \chi_q^r(g^i) \\
 &= \sum_{j=0}^{q-2} \psi(g^j - 1) \chi_q^r(g^j) \chi_q^r(g^i) \\
 &= \sum_{x \in \mathbb{F}_q} \psi(x - 1) \chi_q^r(x) \chi_q^r(g^i) \\
 &= \psi(-1) J_q(\psi, \chi_q^r) \chi_q^r(g^i).
 \end{aligned}$$

This implies that

$$M_q(\psi) \boldsymbol{\xi}_r = \alpha_r \boldsymbol{\xi}_r,$$

where

$$\boldsymbol{\xi}_r = (\chi_q^r(g^0), \chi_q^r(g^1), \dots, \chi_q^r(g^{q-2}))^T.$$

As these  $\boldsymbol{\xi}_r$  are linearly independent over  $\mathbb{C}$ , the numbers  $\alpha_0, \alpha_1, \dots, \alpha_{q-2}$  are all the eigenvalues of  $M_q(\psi)$ . Using essentially the same method, one can easily obtain that all the eigenvalues of  $N_q(\psi)$  are  $\beta_0, \beta_1, \dots, \beta_{q-2}$ .

In view of the above, we have completed the proof.  $\square$

Now we are in a position to prove our last theorem.

**Proof of Theorem 1.3.** (i) It is clear that

$$G(t) = \frac{t^{q-1} - 1}{t - 1} = 1 + t + \dots + t^{q-2} = \prod_{i=2}^{q-1} (t - x_i)$$

over  $\mathbb{F}_q$ . This implies

$$(5.7) \quad \prod_{i=2}^{q-2} x_i = -1.$$

Recall that  $g$  is a generator of  $\mathbb{F}_q^\times$ . By Eq. (5.7) we obtain

$$(5.8) \quad \det D_q^-(\psi) = \prod_{i=2}^{q-1} \psi(x_i) \cdot \det \left[ \psi \left( \frac{x_j}{x_i} - 1 \right) \right]_{2 \leq i, j \leq q-1} = \psi(-1) \cdot \det [\psi(g^{j-i} - 1)]_{1 \leq i, j \leq q-2}.$$

Note that  $[\psi(g^{j-i} - 1)]_{1 \leq i, j \leq q-2}$  is an almost circulant matrix. By Lemma 3.1 and Lemma 5.2, we obtain

$$\begin{aligned}
 \det [\psi(g^{j-i} - 1)]_{1 \leq i, j \leq q-2} &= \frac{1}{q-1} \cdot \alpha_0 \alpha_1 \cdots \alpha_{q-2} \cdot \left( \frac{1}{\alpha_0} + \frac{1}{\alpha_1} + \dots + \frac{1}{\alpha_{q-2}} \right) \\
 &= \frac{\psi(-1)}{q-1} \cdot \prod_{r=0}^{q-2} J_q(\psi, \chi_q^r) \cdot \left( \sum_{r=0}^{q-2} \frac{1}{J_q(\psi, \chi_q^r)} \right)
 \end{aligned}$$

$$= -\frac{(1 + \psi(-1))}{q^2} G_q(\psi)^{q-1}.$$

Combining this with Eq. (5.8), we obtain

$$\det D_q^-(\psi) = -\frac{(1 + \psi(-1))}{q^2} G_q(\psi)^{q-1}.$$

This completes the proof of (i).

(ii) Similar to Eq. (5.7), we have

$$(5.9) \quad \det D_q^+(\psi) = \psi(-1) \cdot \det [\psi(g^{j-i} + 1)]_{1 \leq i, j \leq q-2}.$$

Since  $[\psi(g^{j-i} + 1)]_{1 \leq i, j \leq q-2}$  is an almost circulant matrix, by Lemma 3.1 and Lemma 5.2, one can verify that

$$\begin{aligned} \det [\psi(g^{j-i} + 1)]_{1 \leq i, j \leq q-2} &= \frac{1}{q-1} \cdot \beta_0 \beta_1 \cdots \beta_{q-2} \cdot \left( \frac{1}{\beta_0} + \frac{1}{\beta_1} + \cdots + \frac{1}{\beta_{q-2}} \right) \\ &= \frac{(-1)^{(q-1)/2}}{q-1} \cdot \prod_{r=0}^{q-2} J_q(\psi, \chi_q^r) \cdot \left( \sum_{r=0}^{q-2} \frac{(-1)^r}{J_q(\psi, \chi_q^r)} \right) \\ &= \frac{(-1)^{(q+1)/2}}{q^2} \left( 2 - \overline{\psi(2)} \right) G_q(\psi)^{q-1}. \end{aligned}$$

By this and Eq. (5.9), we finally obtain

$$\det D_q^+(\psi) = \frac{(-1)^{(q+1)/2} \cdot \psi(-1)}{q^2} \left( 2 - \overline{\psi(2)} \right) G_q(\psi)^{q-1}.$$

In view of the above, we have completed the proof.  $\square$

**Acknowledgments** The authors would like to thank Prof. Hao Pan for his helpful comments and steadfast encouragement.

This research was supported by the Natural Science Foundation of China (Grant Nos. 12101321, 12201291 and 12371004).

## REFERENCES

- [1] L. Carlitz, Some cyclotomic matrices, *Acta Arith.* 5 (1959), 293–308.
- [2] R. Chapman, Determinants of Legendre symbol matrices, *Acta Arith.* 115 (2004), 231–244.
- [3] H. Cohen, *Number Theory, Vol. I. Tools and Diophantine Equations*, Springer, 2007.
- [4] D. Grinberg, Z.-W. Sun and L. Zhao, Proof of three conjectures on determinants related to quadratic residues, *Linear Multilinear Algebra* 70 (2022), no. 19, 3734–3746.
- [5] B. Gross and N. Koblitz, Gauss sums and the  $p$ -adic  $\Gamma$ -function, *Ann. Math* 109 (1979), 569–581.
- [6] C. Krattenthaler, Advanced determinant calculus: a complement, *Linear Algebra Appl.* 411 (2005), 68–166.
- [7] D.H. Lehmer, On certain character matrices, *Pac. J. Math.* 6 (1956), 491–499.
- [8] L.J. Mordell, The congruence  $((p-1)/2)! \equiv \pm 1 \pmod{p}$ , *Am. Math. Mon.* 68 (1961), 145–146.
- [9] Z.-H. Sun, Congruences involving Bernoulli and Euler numbers, *J. Number Theory* 128 (2008), no. 2, 280–312.



- [10] Z.-W. Sun, A congruence for primes, Proc. Amer. Math. Soc. 123 (1995), no. 5, 1341–1346.
- [11] Z.-W. Sun, On the sum  $\sum_{k \equiv r \pmod{m}} \binom{n}{k}$  and related congruences, Israel J. Math. 128 (2002), 135–156.
- [12] Z.-W. Sun, On some determinants with Legendre symbol entries, Finite Fields Appl. 56 (2019), 285–307.
- [13] Z.-W. Sun, Some determinants involving quadratic residues modulo primes, preprint, arXiv:2401.14301.
- [14] B. Sury, Sum of reciprocals of the binomial coefficients, Europ. J. Combinatorics, 14 (1993), 351–353.
- [15] M. Vsemirnov, On the evaluation of R. Chapman’s “evil determinant”, Linear Algebra Appl. 436 (2012), 4101–4106.
- [16] M. Vsemirnov, On R. Chapman’s “evil determinant”: case  $p \equiv 1 \pmod{4}$ , Acta Arith. 159 (2013), 331–344.
- [17] H.-L. Wu, Determinants concerning Legendre symbols, C. R. Math. Acad. Sci. Paris 359 (2021), 651–655.
- [18] H.-L. Wu and L.-Y. Wang, The Gross-Koblitz formula and almost circulant matrices related to Jacobi sums, preprint, arXiv:2409.13307.

(HAI-LIANG WU) SCHOOL OF SCIENCE, NANJING UNIVERSITY OF POSTS AND TELECOMMUNICATIONS,  
NANJING 210023, PEOPLE’S REPUBLIC OF CHINA

*Email address:* whl.math@smail.nju.edu.cn

(LI-YUAN WANG) SCHOOL OF PHYSICAL AND MATHEMATICAL SCIENCES, NANJING TECH UNIVERSITY,  
NANJING 211816, PEOPLE’S REPUBLIC OF CHINA

*Email address:* wly@smail.nju.edu.cn

(HE-XIA NI) DEPARTMENT OF APPLIED MATHEMATICS, NANJING AUDIT UNIVERSITY, NANJING 211815,  
PEOPLE’S REPUBLIC OF CHINA

*Email address:* nihexia@yeah.net