

COMBINATORIAL CALABI FLOW FOR IDEAL CIRCLE PATTERN

SHENGYU LI AND ZHIGANG WANG*

ABSTRACT. We study the combinatorial Calabi flow for ideal circle patterns in both hyperbolic and Euclidean background geometry. We prove that the flow exists for all time and converges exponentially fast to an ideal circle pattern metric on surfaces with prescribed attainable curvatures. As a consequence, we provide an algorithm to find the desired ideal circle patterns.

CONTENTS

1. Introduction	1
1.1. Ideal circle pattern	1
1.2. Thurston's construction and curvature map	3
1.3. Main result	4
2. Preliminaries	5
3. Combinatorial Calabi flow in hyperbolic background geometry	6
3.1. Hyperbolic background geometry	6
3.2. Long time existence	8
3.3. Convergence	9
4. Combinatorial Calabi flow in Euclidean background geometry	10
4.1. Euclidean background geometry	10
4.2. Long time existence and convergence	12
References	13

1. INTRODUCTION

1.1. Ideal circle pattern. Circle patterns was introduced by Thurston [20] to study the geometry and topology of 3-manifolds. Thurston [21] once proposed a conjecture that infinite hexagonal circle packings converge to classical Riemann

Date: January 6, 2025.

2020 Mathematics Subject Classification. Primary 52C26.

Key words and phrases. Ideal circle pattern, combinatorial Calabi flow, combinatorial Ricci flow.

*Corresponding author.

mapping, which was proved by Rodin-Sullivan [17]. Since then, circle patterns have been studied intensively in [15, 18, 19, 24].

Assume that S is a compact oriented closed surface with a triangulation \mathcal{T} and a constant Riemannian metric μ . A circle pattern \mathcal{P} on (S, μ) is a collection of oriented circles. A circle pattern \mathcal{P} is \mathcal{T} -type if there exists a geodesic triangulation $\mathcal{T}(\mu)$ of (S, μ) with the following properties: (i) $\mathcal{T}(\mu)$ is isotopic to \mathcal{T} ; (ii) the vertices of $\mathcal{T}(\mu)$ coincide with the centers of circles in \mathcal{P} .

Let V, E, F denote the sets of vertices, edges and faces of \mathcal{T} . Connecting the centers of two adjacent circles C_{v_i} and C_{v_j} of circle pattern $\mathcal{P} = \{C_v : v \in V\}$ gives an edge $e = [v_i, v_j] \in E$. For every $e \in E$, there exists an exterior intersection angle $\Theta(e) \in (0, \pi)$. For each $v \in V$, let \mathbb{D}_v (resp. D_v) denote the open (resp. closed) disk bounded by C_v . Each connected component of the set $S \setminus (\cup_{v \in V} \mathbb{D}_v)$ is called an interstice.

Definition 1. A \mathcal{T} -type circle pattern \mathcal{P} on (S, μ) is called ideal if it satisfies the following properties:

- (i) there exists a one-to-one correspondence between the interstices of \mathcal{P} and the 2-cells of \mathcal{T} ;
- (ii) every interstice of \mathcal{P} consists of a point (see, e.g., Figure 1).




FIGURE 1. An ideal circle pattern

Let $\Theta : E \rightarrow (0, \pi)$ be an angle function. A natural question asks whether there exists a \mathcal{T} -type ideal circle pattern with a given exterior intersection function Θ . Thurston [20, Chap.13] observed the relation between circle patterns and hyperbolic polyhedra. Let \mathbb{B}^3 denote the hyperbolic 3-space. Given a convex hyperbolic polyhedron \mathcal{Q} in \mathbb{B}^3 , the plane of each polyhedron face will cut a circle on the sphere $\partial\mathbb{B}^3$. All circles form a circle pattern with dual combinatorial type of \mathcal{Q} . The exterior intersection angles between adjacent circles are equal to the dihedral angles of corresponding faces of \mathcal{Q} . A convex hyperbolic polyhedron in \mathbb{B}^3 is called ideal if all its vertices are on the sphere $\partial\mathbb{B}^3$. In this way, an ideal circle pattern corresponds to a unique ideal polyhedron. Consider the following condition:

(C1) For any distinct edges e_1, e_2, \dots, e_m forming the boundary of a 2-cell of \mathcal{T} ,

$$\sum_{i=1}^m \Theta(e_i) = (m-2)\pi.$$

Let g denote the genus of surface S . Under the dual conditions of (C1), Rivin [16] proved there exists a unique ideal polyhedron with the dihedral angles given by Θ for the case $g = 0$. Equivalently, he showed that there exists a unique ideal circle pattern with given exterior intersection angles on surfaces. Bobenko-Springborn [2] further generalized this result to the case $g \geq 1$.

1.2. Thurston's construction and curvature map. Thurston's construction is presented as follows. Let S be a surface of genus $g > 1$ (resp. $g = 1$) with a triangulation \mathcal{T} . Assume $\Theta(e) \in (0, \pi)$ for each $e \in E$. Let $V = \{v_1, v_2, \dots, v_N\}$. A radius vector $r = (r_1, r_2, \dots, r_N) \in \mathbb{R}_+^N$ assigns each vertex $v_i \in V$ a positive radius r_i . For each edge $[v_i, v_j] \in E$, we assign a length

$$l_{ij} = \cosh^{-1} \left(\cosh r_i \cosh r_j + \sinh r_i \sinh r_j \cos \Theta([v_i, v_j]) \right)$$

in hyperbolic background geometry and

$$l_{ij} = \sqrt{r_i^2 + r_j^2 + 2r_i r_j \cos \Theta([v_i, v_j])}$$

in Euclidean background geometry. It is easy to check that there exists a unique geodesic triangle with lengths l_{ij} , r_i and r_j . Gluing all these hyperbolic (resp. Euclidean) triangles along the common edges gives a hyperbolic (resp. Euclidean) ideal circle pattern metric μ on the surface S with possible cone singularities at vertices of \mathcal{T} .

To describe the singularities on (S, μ) , the discrete curvature is defined to be

$$K_i = 2\pi - \sigma(v_i),$$

where $\sigma(v_i)$ is the sum of inner angles at v_i for all triangles incident to v_i . This yields the following curvature map

$$\begin{aligned} Th : \quad \mathbb{R}_+^N &\longrightarrow \mathbb{R}^N \\ (r_1, r_2, \dots, r_N) &\longmapsto (K_1, K_2, \dots, K_N). \end{aligned}$$

Then K_1, K_2, \dots, K_N are smooth functions of r . Bobenko-Springborn [2] gave complete characterizations of the images of the curvature map in both hyperbolic and Euclidean background geometry.

Theorem A (Bobenko-Springborn [2]). *Assume that Θ is an angle function satisfying the condition (C1).*

- (i) *In hyperbolic background geometry, Th is injective. Moreover, the image of Th consists of vectors (K_1, K_2, \dots, K_N) satisfying*

$$K_i < 2\pi \quad (\forall i = 1, 2, \dots, N), \tag{1.1}$$

and

$$\sum_{v_i \in A} K_i > 2\pi|A| - 2 \sum_{e \in E, \partial e \cap A \neq \emptyset} \Theta(e). \quad (1.2)$$

for any non-empty subset A of V .

- (ii) In Euclidean background geometry, Th is injective up to scaling. Moreover, the image of Th consists of vectors (K_1, K_2, \dots, K_N) satisfying and

$$\sum_{v_i \in A} K_i \geq 2\pi|A| - 2 \sum_{e \in E, \partial e \cap A \neq \emptyset} \Theta(e). \quad (1.3)$$

for any non-empty subset A of V , where the equality holds if and only if $A = V$.

1.3. Main result. Chow-Luo [3] introduced the combinatorial Ricci flow on closed surfaces and proved the existence and convergence of the flow. This yields an algorithm finding circle patterns. Ge [7] and Ge-Hua [8] introduced the combinatorial Calabi flow for circle patterns with non-obtuse exterior intersection angles in Euclidean and hyperbolic background geometry, respectively. Ge-Hua-Zhou [10] used the combinatorial Ricci flow for ideal circle pattern, providing a computational method to search for ideal circle pattern. We refer the reader to [9, 12, 13, 14, 23] for other developments concerning combinatorial flows. In this paper, we explore the combinatorial Calabi flow introduced by Ge [6] to find the desired ideal circle pattern metrics.

A curvature vector $\overline{K} = (k_1, k_2, \dots, k_N) \in \mathbb{R}^N$ is said to be hyperbolic (resp. Euclidean) **attainable** if it satisfies (1.1) and (1.2) (resp. (1.3)). In hyperbolic background geometry, let us consider the following flow

$$\frac{dr_i}{dt} = \sum_{j=1}^N (k_j - K_j) \frac{\partial K_j}{\partial r_i} \sinh^2 r_i \quad (i = 1, 2, \dots, N), \quad (1.4)$$

where $r(0) \in \mathbb{R}_+^N$ is an initial radius vector. In Euclidean background geometry, consider the flow

$$\frac{dr_i}{dt} = \sum_{j=1}^N (k_j - K_j) \frac{\partial K_j}{\partial r_i} r_i^2 \quad (i = 1, 2, \dots, N), \quad (1.5)$$

where $r(0) \in \mathbb{R}_+^N$ is an initial radius vector. Using Lyapunov functions, we get the following result.

Theorem 1. *Assume that Θ is an angle function satisfying (C1) and \overline{K} is hyperbolic (resp. Euclidean) attainable. Then the flow (1.4) (resp. (1.5)) exists for all time and converges exponentially fast to a radius vector which produces a hyperbolic (resp. Euclidean) ideal circle pattern metric on surfaces whose discrete curvatures are assigned by \overline{K} .*

The paper is organized as follows. In Section 2, we introduce several properties of circle pattern. In Section 3, we prove Theorems 2 and 3. We give the proofs of Theorems 4 and 1 in Section 4.

2. PRELIMINARIES

In this section, we present several results regarding circle pattern.

Lemma 1 ([10], Lemma 1.1). *Given $\Theta_k \in (0, \pi)$ and two positive numbers r_i, r_j , there exists a configuration of two intersection circles in both Euclidean and hyperbolic geometries, unique up to isometries, having radii r_i, r_j and meeting exterior intersection angle Θ_k .*




FIGURE 2. A two-circle configuration

Let $\Delta_{v_i v_j v_k}$ be the triangle by connecting the centers v_i, v_j and one of the intersection points of two circles (see Figure 2). Let ϑ_η be the inner angle of $\Delta_{v_i v_j v_k}$ at the centers v_η for $\eta \in \{i, j\}$. The following result is due to Ge-Hua-Zhou [10].

Lemma 2 ([10], Lemma 1.2). *Let $\Theta_k \in (0, \pi)$ be fixed.*

(i) *In hyperbolic background geometry,*

$$\frac{\partial \vartheta_i}{\partial r_i} < 0, \quad \sinh r_j \frac{\partial \vartheta_i}{\partial r_j} = \sinh r_i \frac{\partial \vartheta_j}{\partial r_i} > 0.$$

(ii) *In Euclidean background geometry,*

$$\frac{\partial \vartheta_i}{\partial r_i} < 0, \quad r_j \frac{\partial \vartheta_i}{\partial r_j} = r_i \frac{\partial \vartheta_j}{\partial r_i} > 0.$$

Lemma 3 ([10], Lemma 1.5). *In hyperbolic background geometry, for $\epsilon > 0$, there exists a positive number M such that $\vartheta_i < \epsilon$ whenever $r_i > M$.*

Recall that \mathcal{T} is a triangulation of surface S with vertices $V = \{v_1, v_2, \dots, v_N\}$. For a radius vector $r = (r_1, r_2, \dots, r_N)$, let $u_i = \ln \tanh(r_i/2)$ (resp. $u_i = \ln r_i$) in hyperbolic (resp. Euclidean) background geometry for $i = 1, 2, \dots, N$. Then the

discrete curvatures K_1, K_2, \dots, K_N are smooth functions of $u = (u_1, u_2, \dots, u_N)$. This gives the curvature map Th in terms of u :

$$\begin{aligned} Th : \quad \mathbb{R}_-^N \text{ (resp. } \mathbb{R}^N) &\longrightarrow \mathbb{R}^N \\ (u_1, u_2, \dots, u_N) &\longmapsto (K_1, K_2, \dots, K_N). \end{aligned}$$

The following result is obtained by Ge-Hua-Zhou [10].

Lemma 4 ([10], Lemmas 2.1 and 3.1). *Let Th be the curvature map.*

- (i) *In hyperbolic background geometry, the Jacobian matrix L of Th in terms of u is symmetric and positive definite.*
- (ii) *In Euclidean background geometry, the Jacobian matrix L of Th in terms of u is symmetric and positive definite when restricted in*

$$\Upsilon = \left\{ u \in \mathbb{R}^N : \sum_{i=1}^N u_i = d \right\} \quad (d \text{ is a given real number}),$$

where the null space of L is $\{k(1, 1, \dots, 1) : k \in \mathbb{R}\}$.

Recall that K_i is the discrete curvature at v_i for $i = 1, 2, \dots, N$. We write $j \sim i$ if the vertices v_i and v_j are adjacent. Using geometry arguments, Ge-Hua ([8, Lemmas 2.2 and 2.3]) obtained an initial version of the following estimate. Applying Glickenstein-Thomas's result (see [11, Proposition 9]), Li-Luo-Xu [13] also provided an independent proof of the following lemma.

Lemma 5. *In hyperbolic background geometry, for any constant $\rho > 1$, there exists a constant $M > 0$ such that*

$$\rho \sum_{j \sim i} \frac{\partial K_j}{\partial u_i} + (\rho - 1) \frac{\partial K_i}{\partial u_i} > 0$$

when $r_i > M$.

3. COMBINATORIAL CALABI FLOW IN HYPERBOLIC BACKGROUND GEOMETRY

3.1. Hyperbolic background geometry. Recall that \mathcal{T} is a triangulation of surface S with vertices $V = \{v_1, v_2, \dots, v_N\}$. For $i = 1, 2, \dots, N$, let

$$r = (r_1, r_2, \dots, r_N)$$

be a radius vector and $u_i = \ln \tanh(r_i/2)$. Assume that $\bar{K} = (k_1, k_2, \dots, k_N)$ is a hyperbolic attainable vector. By changing variables, the combinatorial Calabi flow (1.4) is equivalent to the flow

$$\frac{du_i}{dt} = \sum_{j=1}^N (k_j - K_j) \frac{\partial K_j}{\partial u_i} \quad (i = 1, 2, \dots, N), \quad (3.1)$$

where $u(0) \in \mathbb{R}_-^N$ is an initial vector.

Following Colin de Verdière [5], Chow-Luo [3] and Ge-Hua-Zhou [10], we consider the 1-form

$$\alpha = \sum_{i=1}^N (K_i - k_i) du_i.$$

It is easy to verify that α is a closed 1-form, which implies that

$$\Psi(u) = \int_{u(0)}^u \sum_{i=1}^N (K_i - k_i) du_i$$

for $u = (u_1, u_2, \dots, u_N) \in \mathbb{R}_+^N$ is well-defined. Since the Hessian matrix of Ψ is equal to the positive definite Jacobian matrix L mentioned in Lemma 4, Ψ is strictly convex. Meanwhile, Theorem A implies that there exists a unique point $r^* \in \mathbb{R}_+^N$ such that $K_i(r^*) = k_i$ for $i = 1, 2, \dots, N$. Let $u^* \in \mathbb{R}_+^N$ be the vector corresponding to r^* satisfying $K_i(u^*) = k_i$. A direct computation gives

$$\left. \frac{\partial \Psi}{\partial u_i} \right|_{u=u^*} = K_i(u^*) - k_i = 0.$$

Hence u^* is a critical point of Ψ . Motivated by Lyapunov theory [1, 4, 22], we introduce the following function

$$\Lambda(u) = \Psi(u) - \Psi(u^*).$$

Lemma 6. *In hyperbolic background geometry, we have*

$$\lim_{\|u\| \rightarrow +\infty} \Lambda(u) = +\infty.$$

To prove Lemma 6, we need the following lemma.

Lemma 7 ([9], Lemma 2.8). *Let f be a smooth strictly convex function defined in a convex set Ω with a critical point $p \in \Omega$. Then the following properties hold:*

- (i) p is the unique global minimum point of f .
- (ii) If Ω is unbounded, then

$$\lim_{\|x\| \rightarrow +\infty} f(x) = +\infty.$$

Proof of Lemma 6. By the above discussion, Ψ is strictly convex in \mathbb{R}_+^N and u^* is a critical point. It follows from Lemma 7 that

$$\lim_{\|u\| \rightarrow +\infty} \Psi(u) = +\infty.$$

Hence $\Lambda(u) \rightarrow +\infty$. □

3.2. Long time existence. In what follows, we prove the existence part of Theorem 1.

Theorem 2. *The flow (3.1) exists for all time.*

Proof. Since each K_i is a smooth function of u , (K_1, K_2, \dots, K_N) is locally Lipschitz continuous. By ODE theory [22], the flow (3.1) has a unique solution $u(t)$ in $[0, \epsilon)$ for some $\epsilon > 0$. Hence $u(t)$ exists in a maximal time interval $[0, T_0)$ with $0 < T_0 \leq +\infty$. Assume that T_0 is finite. Then for at least one $i \in \{1, 2, \dots, N\}$, there exists $t_m \rightarrow T_0$ such that

$$r_i(t_m) \rightarrow 0 \text{ or } r_i(t_m) \rightarrow +\infty.$$

For the first case, we have $u_i(t_m) \rightarrow -\infty$. Lemma 6 implies

$$\Lambda(u(t_m)) \rightarrow +\infty.$$

Since L is positive definite, a calculation yields

$$\begin{aligned} \frac{d\Lambda}{dt} &= \sum_{i=1}^N \frac{\partial \Psi}{\partial u_i} \frac{du_i}{dt} \\ &= - \sum_{i=1}^N \sum_{j=1}^N (K_i - k_i) \frac{\partial K_j}{\partial u_i} (K_j - k_j) \\ &= -(K_1 - k_1, \dots, K_N - k_N) L (K_1 - k_1, \dots, K_N - k_N)^T \\ &\leq 0. \end{aligned}$$

It follows that

$$\Lambda(u(t_m)) \leq \Lambda(u(0)),$$

which leads to a contradiction.

We now consider the second case. Let k_{\max} and k_{\min} be the maximum and minimum values among $\{k_1, k_2, \dots, k_N\}$, respectively. By Lemma 3, there exists $\eta \in (0, 1)$ and $M_1 > 0$ for any $r_i > M_1$ such that

$$K_i > 2\pi\eta > k_{\max}. \quad (3.2)$$

Meanwhile, Lemma 2 implies

$$\frac{\partial K_j}{\partial u_i} < 0 \text{ and } \frac{\partial K_i}{\partial u_i} > 0. \quad (3.3)$$

A direct computation yields

$$\begin{aligned} \frac{du_i}{dt} &= \sum_{j=1}^N (k_j - K_j) \frac{\partial K_j}{\partial u_i} \\ &= \sum_{j \sim i} (k_j - K_j) \frac{\partial K_j}{\partial u_i} + (k_i - K_i) \frac{\partial K_i}{\partial u_i} \\ &< - \left[(2\pi - k_{\min}) \sum_{j \sim i} \frac{\partial K_j}{\partial u_i} + (2\pi\eta - k_{\max}) \frac{\partial K_i}{\partial u_i} \right]. \end{aligned}$$

Note that

$$\frac{2\pi - k_{\min}}{2\pi\eta - k_{\max}} - 1 > 0.$$

It follows from Lemma 5 that there exists $M_2 > 0$ such that for $r_i > M_2$,

$$(2\pi - k_{\min}) \sum_{j \sim i} \frac{\partial K_j}{\partial u_i} + (2\pi\eta - k_{\max}) \frac{\partial K_i}{\partial u_i} > 0.$$

Let $M = \max\{M_1, M_2\}$. Then $\frac{du_i}{dt} < 0$, which implies

$$\frac{dr_i}{dt} < 0.$$

Since $r_i(t_m) \rightarrow +\infty$, we can choose a sufficiently large n such that $r_i(t_n) > M$. Let

$$a = \inf_s \{s < t_n : r_i(t) > M, \forall t \in [s, t_n]\}.$$

Then we can verify that

$$r_i(a) = M.$$

On the other hand, we have $\frac{dr_i(t)}{dt} < 0$ for $t \in [a, t_n]$. Therefore, we see that

$$r_i(a) > r_i(t_n) > M,$$

which contradicts to $r_i(a) = M$.

By the above discussions, $r_i(t)$ is uniformly bounded for $i = 1, 2, \dots, N$. Hence the flow (3.1) exists for all time and $u(t)$ stays in a compact set of \mathbb{R}_+^N . \square

3.3. Convergence. Using an energy function, we prove the convergence of the flow.

Theorem 3. *Suppose that $\bar{K} = (k_1, k_2, \dots, k_N)$ is hyperbolic attainable. The flow (3.1) converges exponentially fast to a radius vector which produces a hyperbolic ideal circle pattern metric on surfaces with discrete curvatures k_1, k_2, \dots, k_N .*

Proof. Define

$$C(u) = \sum_{i=1}^N (K_i - k_i)^2$$

as an energy function. Since $u(t)$ stays in a compact set of \mathbb{R}_+^N and L depends on u continuously, there exists $\lambda_0 > 0$ such that

$$\begin{aligned} C'(u) &= 2 \sum_{i=1}^N \sum_{j=1}^N (K_j - k_j) \frac{\partial K_j}{\partial u_i} \frac{du_i}{dt} \\ &= -2(K_1 - k_1, \dots, K_N - k_N) L^2 (K_1 - k_1, \dots, K_N - k_N)^T \\ &\leq -2\lambda_0 \sum_{i=1}^N (K_i - k_i)^2 \\ &= -2\lambda_0 C(u). \end{aligned}$$

It follows that

$$\sum_{i=1}^N (K_i - k_i)^2 = C(u) \leq C(u(0)) e^{-2\lambda_0 t}.$$

Therefore, we get

$$|K_i - k_i| \leq \sqrt{C(u(0))} e^{-\lambda_0 t}$$

for $i = 1, 2, \dots, N$. Note that $\left| \frac{\partial K_j}{\partial u_i} \right|$ is uniformly bounded. Then we have

$$|u_i - u_i^*| = \left| \sum_{j=1}^N \int_{\infty}^t (K_j - k_j) \frac{\partial K_j}{\partial u_i} dt \right| \leq \lambda e^{-\lambda_0 t}$$

for some positive number λ . As $t \rightarrow +\infty$, we derive that u converges exponentially fast to u^* . We thus finish the proof. \square

4. COMBINATORIAL CALABI FLOW IN EUCLIDEAN BACKGROUND GEOMETRY

4.1. Euclidean background geometry. Let $u_i = \ln r_i$ for $i = 1, 2, \dots, N$ and $u = (u_1, u_2, \dots, u_N)$. Suppose that $\bar{K} = (k_1, k_2, \dots, k_N)$ is a Euclidean attainable vector. Then the flow (1.5) transforms into the equivalent flow

$$\frac{du_i}{dt} = \sum_{j=1}^N (k_j - K_j) \frac{\partial K_j}{\partial u_i} \quad (i = 1, 2, \dots, N), \quad (4.1)$$

where $u(0)$ is an initial vector in \mathbb{R}^N .

A direct calculation gives

$$\sum_{i=1}^N \frac{du_i}{dt} = -(1, 1, \dots, 1)L(K_1 - k_1, \dots, K_N - k_N)^T,$$

where L is the Jacobian matrix mentioned in Lemma 4. Moreover, the fact that the matrix L has a null space $\{k(1, 1, \dots, 1) : k \in \mathbb{R}\}$ implies that

$$\sum_{i=1}^N \frac{du_i}{dt} = 0.$$

Hence $\sum_{i=1}^N u_i$ remains constant. It means that $u \in \Upsilon$, where

$$\Upsilon = \left\{ u \in \mathbb{R}^N : \sum_{i=1}^N u_i = \sum_{i=1}^N u_i(0) \right\}.$$

Let

$$\Psi(u) = \int_{u(0)}^u \sum_{i=1}^N (K_i - k_i) du_i$$

be a function defined in the set Υ . A direct calculation shows that the Hessian matrix of Ψ is equal to the Jacobian matrix L . Because u is restricted in Υ , it follows from Lemma 4 that L is positive definite. Similar to the discussion in hyperbolic background geometry, Ψ is well defined and strictly convex.

By Theorem A, we know that there exists a radius vector $r^* \in \mathbb{R}_+^N$ such that $K_i(r^*) = k_i$ for $i = 1, 2, \dots, N$. Note that

$$Th(\mathbb{R}_+^N) = Th(\Upsilon).$$

Since L is positive definite, there exists a unique vector $u^* \in \Upsilon$ corresponding to r^* such that

$$K_i(u^*) = k_i$$

for $i = 1, 2, \dots, N$. Moreover,

$$\left. \frac{\partial \Psi}{\partial u_i} \right|_{u=u^*} = K_i(u^*) - k_i = 0.$$

Hence u^* is a unique critical point of Ψ . We now consider the function

$$\Lambda(u) = \Psi(u) - \Psi(u^*).$$

By a similar argument to the proof of Lemma 6, we obtain the following result.

Lemma 8. *In Euclidean background geometry, we have*

$$\lim_{\|u\| \rightarrow +\infty} \Lambda(u) = +\infty.$$

4.2. Long time existence and convergence. Now we prove the following result.

Theorem 4. *Suppose that $\bar{K} = (k_1, k_2, \dots, k_N)$ is Euclidean attainable. The flow (4.1) exists for all time and converges exponentially fast to a radius vector that produces a Euclidean ideal circle pattern metric on surfaces with discrete curvatures k_1, k_2, \dots, k_N .*

Proof. By a similar argument to the hyperbolic background geometry, we prove that the flow (1.5) has a unique solution $u(t)$ in a maximal time interval $[0, T_0)$ with $0 < T_0 \leq +\infty$. We claim that $T_0 = +\infty$. Otherwise, assume that T_0 is finite. Then for at least one $i \in \{1, 2, \dots, N\}$, there exists a sequence $\{t_m\}$ with limit T_0 such that

$$r_i(t_m) \rightarrow 0 \text{ or } r_i(t_m) \rightarrow +\infty.$$

It shows that

$$u_i(t_m) \rightarrow -\infty \text{ or } u_i(t_m) \rightarrow +\infty.$$

No matter which case occurs, it follows from Lemma 8 that $\Psi(u(t_m)) \rightarrow +\infty$, which implies

$$\Lambda(u(t_m)) \rightarrow +\infty.$$

Nevertheless, a direct computation gives

$$\begin{aligned} \frac{d\Lambda}{dt} &= \sum_{i=1}^N \frac{\partial \Psi}{\partial u_i} \frac{du_i}{dt} \\ &= - \sum_{i=1}^N \sum_{j=1}^N (K_j - k_j) \frac{\partial K_j}{\partial u_i} (K_i - k_i) \\ &= -(K_1 - k_1, \dots, K_N - k_N) L (K_1 - k_1, \dots, K_N - k_N)^T \\ &\leq 0. \end{aligned}$$

As a result, we have

$$\Lambda(u(t_m)) \leq \Lambda(u(0)),$$

which leads to a contradiction. Hence the flow (4.1) never touches the boundary of $\Upsilon \subset \mathbb{R}^N$ in any finite time interval. Namely, the flow exists for all time and $u(t)$ stays in a compact set of Υ .

The proof of exponential convergence of the flow (4.1) is similar to the argument of the flow (3.1). \square

Finally, we give the proof of Theorem 1.

Proof of Theorem 1. Since the flow (1.4) (resp. (1.5)) is equivalent to the flow (3.1) (resp. (4.1)), the conclusion of Theorem 1 is an immediate consequence of Theorems 2, 3 and 4. \square

Acknowledgements. The present investigation was supported by the *Natural Science Foundation of Hunan Province* under Grant no. 2022JJ30185 of the P. R. China. The authors thank Prof. Ze Zhou for his stimulating discussions and Prof. Yueping Jiang for his invaluable help.

Conflicts of interest. The authors declare that they have no conflict of interest.

Data availability statement. Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

Statement of independent research outcomes. It has come to the authors' attention that Prof. Xiaoxiao Zhang at Beijing Wuzi University has independently arrived at similar conclusions, coinciding with the authors' work.

REFERENCES

- [1] E. A. Barbařhin, N. N. Krasovskii, *On the existence of Lyapunov functions in the case of asymptotic stability in the large*, (Russian) Akad. Nauk SSSR. Prikl. Mat. Meh. **18** (1954), 345–350.
- [2] A. I. Bobenko, B. A. Springborn, *Variational principles for circle patterns and Koebe's theorem*, Trans. Amer. Math. Soc. **356** (2004), 659–689.
- [3] B. Chow, F. Luo, *Combinatorial Ricci flows on surfaces*, J. Differential Geom. **63** (2003), 97–129.
- [4] F. H. Clarke, Y. S. Ledyaev, R. J. Stern, *Asymptotic stability and smooth Lyapunov functions*, J. Differential Equations **149** (1998), 69–114.
- [5] Y. Colin de Verdière, *Un principe variationnel pour les empilements de cercles*, (French) Invent. Math. **104** (1991), 655–669.
- [6] H. Ge, *Combinatorial methods and geometric equations*, Thesis (Ph.D.), Peking University, Beijing, 2012, 144 pp.
- [7] H. Ge, *Combinatorial Calabi flows on surfaces*, Trans. Amer. Math. Soc. **370** (2018), 1377–1391.
- [8] H. Ge, B. Hua, *On combinatorial Calabi flow with hyperbolic circle patterns*, Adv. Math. **333** (2018), 523–538.
- [9] H. Ge, B. Hua, Z. Zhou, *Circle patterns on surfaces of finite topological type*, Amer. J. Math. **143** (2021), 1397–1430.
- [10] H. Ge, B. Hua, Z. Zhou, *Combinatorial Ricci flows for ideal circle patterns*, Adv. Math. **383** (2021), Paper No. 107698, 26 pp.
- [11] D. Glickenstein, J. Thomas, *Duality structures and discrete conformal variations of piecewise constant curvature surfaces*, Adv. Math. **320** (2017), 250–278.
- [12] G. Hu, Z. Lei, Y. Qi, P. Zhou, *Combinatorial p -th Calabi flows for total geodesic curvatures in hyperbolic background geometry*, J. Geom. Anal. **35** (2025), Paper No. 18.
- [13] S. Li, Q. Luo, Y. Xu, *Combinatorial Calabi flow on surfaces of finite topological type*, Proc. Amer. Math. Soc. **152** (2024), 4035–4047.
- [14] A. Lin, X. Zhang, *Combinatorial p -th Calabi flows on surfaces*, Adv. Math. **346** (2019), 1067–1090.

- [15] J. Liu, Z. Zhou, *How many cages midscribe an egg*, Invent. Math. **203** (2016), 655–673.
- [16] I. Rivin, *A characterization of ideal polyhedra in hyperbolic 3-space*, Ann. of Math. (2) **143** (1996), 51–70.
- [17] B. Rodin, D. Sullivan, *The convergence of circle packings to the Riemann mapping*, J. Differential Geom. **26** (1987), 349–360.
- [18] O. Schramm, *How to cage an egg*, Invent. Math. **107** (1992), 543–560.
- [19] K. Stephenson, *Introduction to circle packing. The theory of discrete analytic functions*, Cambridge University Press, Cambridge, 2005.
- [20] W. Thurston, *Geometry and topology of 3-manifolds*, Princeton lecture notes, 1976.
- [21] W. Thurston, *The finite Riemann mapping theorem*. Invited talk, an international symposium at Purdue University on the occasion of the proof of the Bieberbach Conjecture, 1985.
- [22] W. Walter, *Ordinary Differential Equations*, Springer, New York, 1998.
- [23] X. Xu, C. Zheng, *Combinatorial curvature flows for generalized circle packings on surfaces with boundary*, Int. Math. Res. Not. IMRN **2023** (2023), 17704–17728.
- [24] Z. Zhou, *Circle patterns with obtuse exterior intersection angles*, arXiv:1703.01768.

SHENGYU LI

SCHOOL OF MATHEMATICS AND STATISTICS, HUNAN FIRST NORMAL UNIVERSITY, CHANGSHA 410205, HUNAN, P. R. CHINA.

Email address: `lishengyu@hnu.edu.cn`

ZHIGANG WANG

SCHOOL OF MATHEMATICS AND STATISTICS, HUNAN FIRST NORMAL UNIVERSITY, CHANGSHA 410205, HUNAN, P. R. CHINA.

Email address: `wangmath@163.com`