Sets preserved by a large subgroup of the special linear group

Le Quang Hung^{*} Thang Pham[†] Kaloyan Slavov[‡]

January 6, 2025

Abstract

Let *E* be a subset of the the affine plane over a finite field \mathbb{F}_q . We bound the size of the subgroup of $\mathrm{SL}_2(\mathbb{F}_q)$ that preserves *E*. As a consequence, we show that if *E* has size $\ll q^{\alpha}$ and is preserved by $\gg q^{\beta}$ elements of $\mathrm{SL}_2(\mathbb{F}_q)$ with $\beta \geq 3\alpha/2$, then *E* is contained in a line. This result is sharp in general, and will be proved by using a point-line incidence bound in \mathbb{F}_q^3 .

1 Introduction

Let \mathbb{F}_q be a finite field of order q, let E be a subset of \mathbb{F}_q^2 , and let R be a subset of $\mathrm{SL}_2(\mathbb{F}_q)$. In a recent work [2] on packing sets over finite fields, Hegyvári, Hung, Iosevich, and Pham give lower bounds for the size of

$$R(E) := \bigcup_{\theta \in R} \theta(E) = \{\theta(x) \colon x \in E, \theta \in R\}$$

in terms of |E|, |R|, q (and other quantities determined by R and E). However, in some cases, only the trivial bound $|R(E)| \ge |E|$ is available (see Proposition 2.1, Theorem 2.2, and Theorem 2.3 in [2]).

In the present paper, we ask for a description of extremal configurations, that is, sets $E \subset \mathbb{F}_q^2$ which "stay packed" under $\mathrm{SL}_2(\mathbb{F}_q)$. More precisely, our goal is to study sets $E \subset \mathbb{F}_q^2$ such that R(E) = E for a large subset R of $\mathrm{SL}_2(\mathbb{F}_q)$.

Throughout the paper, $X \ll Y$ means $X \leq CY$ for a constant C; $X \gg Y$ means $X \geq CY$ for a constant C, and $X \approx Y$ means $X \ll Y$ and $Y \ll X$.

For a subset E of \mathbb{F}_q^2 , let

$$R_E = \{ \theta \in \mathrm{SL}_2(\mathbb{F}_q) \mid \theta(E) = E \}.$$

Example 1. If E is a line, then |E| = q and $|R_E| = q^2 - q \approx q^2$.

Note that for any $E \subset \mathbb{F}_q^2$, denoting the complement $\mathbb{F}_q^2 - E$ by E^c , we have $R_E = R_{E^c}$. So Example 1 also shows that when E is the complement of a line, we have $|R_E| \approx q^2$.

Example 2. For positive divisor c of q-1, there exists a subgroup $S \subset \mathbb{F}_q^*$ with |S| = (q-1)/c.

^{*}Hanoi University of Science and Technology, Hanoi. Email: hung.lequang@hust.edu.vn

[†]University of Science, Vietnam National University, Hanoi. Email: thangpham.math@vnu.edu.vn

[‡]Department of Mathematics, ETH Zürich. Email: kaloyan.slavov@math.ethz.ch

Let

$$E = \left\{ (0, y) \in \mathbb{F}_q^2 \colon y \in S \right\}.$$

Then

$$R_E = \left\{ \begin{pmatrix} a & 0 \\ d & a^{-1} \end{pmatrix} \in \mathrm{SL}_2(\mathbb{F}_q) \colon a \in S, d \in \mathbb{F}_q \right\}.$$

Here again $|E| = |S| = (q-1)/c \approx q$ and $|R| = q|S| = q(q-1)/c \approx q^2$.

Motivated by these examples, we ask the following

Question 1.1. Let E be a subset of \mathbb{F}_q^2 that is preserved under "many" elements in $\mathrm{SL}_2(\mathbb{F}_q)$. Does it follow (under an assumption on |E|) that E is contained in a line?

We answer in the affirmative.

Theorem 1.2. Given $c_1 > 0$, there exists a constant $c_2 > 0$ with the following property. Let $\alpha, \beta \geq 0$ be such that $\beta \geq 3\alpha/2$. Let E be a subset of \mathbb{F}_q^2 with $|E| \leq c_1 q^{\alpha}$. Suppose the set R_E of matrices θ in $\mathrm{SL}_2(\mathbb{F}_q)$ such that $\theta(E) = E$ satisfies $|R_E| \geq c_2 q^{\beta}$. Then E is contained in a line.

Note that the assumption involves merely counting data for |E| and $|R_E|$ while the conclusion is a geometric structure on E. In the next corollary, we spell out some illustrative special cases.

Corollary 1.3. Let E be a subset of \mathbb{F}_q^2 . Let R_E denote the set of matrices θ in $\mathrm{SL}_2(\mathbb{F}_q)$ such that $\theta(E) = E$.

- (i) If $|E| \ll q$ and $|R_E| \gg q^{3/2}$, then E is contained in a line;
- (ii) If $|E| \ll q^{2/3}$ and $|R_E| \gg q$, then E is contained in a line;
- (iii) If $|E| \ll q^{4/3}$ and $|R_E| \gg q^2$, then E is contained in a line.

The key ingredient in the proof of Theorem 1.2 is the next result, which is of independent interest.

Theorem 1.4. Let $E \subset \mathbb{F}_q^2$. Let R_E be the set of matrices θ in $\mathrm{SL}_2(\mathbb{F}_q)$ such that $\theta(E) = E$. Suppose at least 2 lines through the origin intersect $E - \{(0,0)\}$. Then $|R_E| \ll |E|^{3/2}$.

The exponent 3/2 in Theorem 1.4 is sharp, as the example below illustrates.

Example 3. For p a prime and $q = p^2$, let $E = \mathbb{F}_p^2 \subset \mathbb{F}_q^2$. Then $R_E \supset \mathrm{SL}_2(\mathbb{F}_p)$. So |E| = q and $|R_E| \ge p^3 - p \approx q^{3/2} = |E|^{3/2}$.

In fact, Example 3 shows that Theorem 1.2 is sharp when $\alpha = 1$ in the following sense: in the statement of Corollary 1.3(i), if one keeps the assumption $|E| \ll q$ but replaces $|R_E| \gg q^{3/2}$ by $|R_E| \gg q^{3/2-\varepsilon}$ with any $\varepsilon > 0$, the statement would be false.

The example below shows that Theorem 1.2 is in fact sharp in the sense described above for all α of the form $\alpha = 1/s$ or $\alpha = 2/s$ for a positive integer s.

Example 4. Let α be of the form $\alpha = 1/s$ or $\alpha = 2/s$ for a positive integer s. Then there exist sufficiently large q and subsets $E \subset \mathbb{F}_q^2$ of size $|E| \approx q^{\alpha}$ such that $|R_E| \approx q^{3\alpha/2}$ and E is not contained in a line. Indeed, pick any positive integer r', set $r = (2/\alpha)r'$, and consider $E = \mathbb{F}_{p^{r'}}^2 \subset \mathbb{F}_q^2$, where $q = p^r$. Then $|E| = p^{2r'} = q^{\alpha}$ while R_E contains $\mathrm{SL}_2(\mathbb{F}_{p^{r'}})$, so $|R_E| \geq p^{3r'} - p^{r'} \approx q^{3\alpha/2}$.

Remark 1.5. Given $E \subset \mathbb{F}_q^2$, one can obtain an upper bound for R_E as follows. Let $P = E \times E \subseteq \mathbb{F}_q^2 \times \mathbb{F}_q^2$. Then, we know from [2] that

$$\left| I(P, R_E) - \frac{|P| |R_E|}{q^2} \right| \ll q\sqrt{|R_E| |P|} + |R_E|.$$

If $1 \ll |E|$ and $1 \ll |E^c|$, using the fact that $I(P, R_E) = |R_E||E|$, one has $|R_E| \ll q^2$ (suppose first $|E| < q^2/2$; then apply the same for the complement E^c). However, the bound from Theorem 1.4 is better as long as $|E| \ll q^{4/3}$.

It would be interesting to study analogues of Theorem 1.2 in higher dimensions.

Based on Example 4, we conjecture that Theorem 1.2 is sharp for each specific $\alpha \in [0, 4/3]$.

In light of results on the packing set problems over the reals (see, for example, [3] and references therein) Theorem 1.2 suggests that it is reasonable to state the following conjecture.

Conjecture 1.6. Let E be a compact set in \mathbb{R}^2 with $\dim_H(E) = \alpha$. Assume that there exists a set R of matrices θ in $\mathrm{SL}_2(\mathbb{R})$ such that $\theta(E) = E$ for each $\theta \in R$, with $\dim_H(R) > 3\alpha/2$. Then the set E is contained in a line. Here \dim_H means the Hausdorff dimension.

2 Proof of Theorem 1.2

Let P be a set of points and L be a set of lines in \mathbb{F}_q^3 . We denote the number of incidences between P and L by I(P,L). The proof of Theorem 1.2 makes use of an incidence bound I(P,L). A modification of the proof of Theorem 8.6 in [5] gives the following result.

Lemma 2.1. Let P be a set of points in \mathbb{F}_p^3 and L be a set of lines in \mathbb{F}_p^3 . Assume that any plane contains at most M lines from L, then we have

$$I(P,L) \le c \left(|P|^{1/2} |L|^{3/4} M^{1/4} + |P| + |L| \right),$$

with an absolute constant c.

The lemma below will allow us to apply the point–line incidence bound of Lemma 2.1.

Lemma 2.2. Let A be a subset of $\mathbb{F}_q^2 \times \mathbb{F}_q^2$ such that for $(m_1, m_2) \in A$ with $m_1 = (u_1, v_1), m_2 = (u_2, v_2)$ we have $u_1, u_2 \neq 0$ and $(v_1, v_2) \neq (0, 0)$. Assume the image of A under the first projection to \mathbb{F}_q^2 does not contain two distinct points on the same line through the origin. Then the map $f: \operatorname{SL}_2(\mathbb{F}_q) \to \mathbb{F}_q^3, \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto (a, d, c)$ satisfies the following properties.

(i) For any $(m_1, m_2) \in A$, the set

$$\ell_{m_1 \to m_2} := f\left(\{\theta \in \mathrm{SL}_2(\mathbb{F}_q) \colon \theta m_1 = m_2\}\right)$$

is a line in \mathbb{F}^3_q ;

- (ii) for $(m_1, m_2) \neq (m'_1, m'_2)$, we have $\ell_{m_1 \to m_2} \neq \ell_{m'_1 \to m'_2}$;
- (iii) f is injective on the set $SL_2(\mathbb{F}_q) \setminus \{\theta : \theta(e_1) = e_1\}$, here e_1 is the line defined by y = 0.

Proof. Let $(m_1, m_2) \in A$. Write $m_1 = (u_1, v_1)$ and $m_2 = (u_2, v_2)$. The set of all $\theta = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in$ SL₂ (\mathbb{F}_q) such that $\theta m_1 = m_2$ is described as the solution set of the following system of three equations

$$\begin{cases} au_1 + bv_1 = u_2, \\ cu_1 + dv_1 = v_2, \\ ad - bc = 1. \end{cases}$$

Solving this system, we have

$$\begin{bmatrix} a \\ d \\ c \end{bmatrix} = \begin{bmatrix} \frac{u_2}{u_1} \\ \frac{u_1}{u_2} \\ \frac{v_2}{u_1} - \frac{v_1}{u_2} \end{bmatrix} + b \begin{bmatrix} -\frac{v_1}{u_1} \\ \frac{v_2}{u_2} \\ -\frac{v_1v_2}{u_1u_2} \end{bmatrix},$$

for $b \in \mathbb{F}_q$. Now one checks that f satisfies the required properties.

Lemma 2.3. Let $E \subset \mathbb{F}_q^2$. Suppose there are exactly 2 lines through the origin that intersect $E - \{(0,0)\}$. Then $|R_E| \leq |E|$.

Proof. Since $SL_2(\mathbb{F}_q)$ acts doubly transitive on the set of lines through the origin, we can assume without loss of generality that the two lines are the two coordinate axes e_1 and e_2 . Denote by E_1 and E_2 the intersections of $E - \{(0,0)\}$ with e_1 and e_2 , respectively. Fix a point m = (x,0) on E_1 .

A matrix $\theta \in \mathrm{SL}_2(\mathbb{F}_q)$ that sends $e_1 \mapsto e_1$ and $e_2 \mapsto e_2$ has the form $\theta = \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix}$. Fixing the image θm on E_1 uniquely determines a, hence also d. Thus, the number of θ 's that preserve E_1 and E_2 is at most $|E_1|$.

Next, a matrix $\theta \in \mathrm{SL}_2(\mathbb{F}_q)$ that swaps e_1 and e_2 has the form $\theta = \begin{pmatrix} 0 & b \\ c & 0 \end{pmatrix}$. Fixing the image θm on E_2 uniquely determines c, hence also b. Thus, the number of θ 's that swap E_1 and E_2 is at most $|E_2|$.

Altogether, $|R_E| \le |E_1| + |E_2| = |E|$.

Lemma 2.4. Let \mathcal{L} be a set of $m \geq 3$ lines through the origin in \mathbb{F}_q^2 . Then the number of matrices θ in $\mathrm{SL}_2(\mathbb{F}_q)$ such that $\theta(\mathcal{L}) = \mathcal{L}$ is $\leq 2 \cdot m^3(m-1)^2$.

Proof. As in the previous lemma, we can assume that \mathcal{L} contains the two coordinate lines e_1 and e_2 . Consider a matrix $\theta = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ in $\mathrm{SL}_2(\mathbb{F}_q)$ that acts on \mathcal{L} . Deciding which element in \mathcal{L} goes under θ to e_1 amounts to fixing the ratio c/d in $\mathbb{F}_q \cup \{\infty\}$. Similarly, deciding which element in \mathcal{L} goes under θ to e_2 amounts to fixing the ratio a/b. Deciding where θ sends e_1 and e_2 determines the ratios a/c and b/d.

Therefore, unless a = d = 0 or b = c = 0, the ratio a/d is also determined by the choices we have made; then $\theta = \lambda \theta_0$ for a fixed matrix θ_0 and a scalar $\lambda \in \mathbb{F}_q$. The condition det $(\theta) = 1$ now gives

at most two possibilities for λ . If a = d = 0 or b = c = 0, we decide where θ sends an element in $\mathcal{L} - \{e_1, e_2\}$; this will determine a/d, and we finish as in the previous case.

Proposition 2.5. Let $E \subset \mathbb{F}_q^2$. Let R_E be the set of matrices θ in $SL_2(\mathbb{F}_p)$ such that $\theta(E) = E$. Let m_0 be the number of lines passing through the origin that intersect $E - \{(0,0)\}$ at exactly m_1 points (for a fixed $m_1 \geq 1$). With the constant c taken from Lemma 2.1, assume that $m_0 > 8 + 4c$. Then

$$|R_E| \le 16c^2 (m_0 m_1)^{3/2}.$$

In particular, $|R_E| \le 16c^2 |E|^{3/2}$.

Proof. Replace E by $E - \{(0,0)\}$ to assume without loss of generality that $(0,0) \notin E$. Let $\ell_1, \ldots, \ell_{m_0}$ be all the lines through the origin such that $|\ell_j \cap E| = m_1$. Let $E_j = \ell_j \cap E$ for $j = 1, \ldots, m_0$.

Let S be the set of matrices $\theta \in SL_2(\mathbb{F}_q)$ such that $\theta(E_j) \in \{E_1, \ldots, E_{m_0}\}, \forall j = 1, \ldots, m_0$. Then $R_E \subset S$. We will estimate |S|.

For $j = 1, ..., m_0$, fix a point $a_j \in E_j$. Let $B = \{a_1, ..., a_{m_0}\} - (e_1 \cup e_2)$ and $C = \left(\bigcup_{j=1}^{m_0} E_j\right) - (e_1 \cup e_2)$.

Let

$$\Omega = \{ (u_1, u_2, \theta) \in B \times C \times S \mid \theta u_1 = u_2 \}$$

Let $\theta \in S$. For any $u_1 \in B$ such that $\theta u_1 \notin e_1 \cup e_1$, we have $(u_1, \theta u_1, \theta) \in \Omega$. Therefore

$$|\Omega| \ge (m_0 - 4)|S|.$$
(1)

Set $S_1 = \{\theta \in S \mid \theta e_1 \neq e_1\}$ and $S_2 = \{\theta \in S \mid \theta e_1 = e_1\}$. For a fixed $(u_1, u_2) \in B \times C$, there exists at most one $\theta = \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} \in S_2$ such that $\theta u_1 = u_2$. Therefore $|\Omega \cap (B \times C \times S_2)| \leq |B||C| \leq m_0^2 m_1$.

We now estimate $|\Omega \cap (B \times C \times S_1)|$. Apply Lemma 2.2 with $A = B \times C$. Let

$$L = \{ \ell_{u_1 \to u_2} \mid u_1 \in B, u_2 \in C \};$$

this is a set of lines in \mathbb{F}_q^3 with $|L| \leq m_0^2 m_1$. Notice that $|\Omega \cap (B \times C \times S_1)|$ equals the number of incidences between the lines in L and the points in $f(S_1)$.

We will see below that any plane in \mathbb{F}_q^3 contains at most $2m_0$ lines from L. Once this is established, Lemma 2.1 implies

$$I(L, f(S_1)) \le c \left(|L|^{3/4} |S_1|^{1/2} (2m_0)^{1/4} + |L| + |f(S_1)| \right),$$

yielding the upper bound

$$|\Omega| \le 2c \left(m_0^{7/4} m_1^{3/4} |S|^{1/2} + m_0^2 m_1 + |S| \right).$$
(2)

Comparing the lower and upper bounds (1) and (2) for $|\Omega|$ under $m_0 > 8 + 4c$, we obtain $|S| \le 16c^2m_0^{3/2}m_1^{3/2}$, completing the proof.

We now turn to verifying the technical condition that any plane in \mathbb{F}_q^3 contains at most $2m_0$ lines from L.

First, we observe that $\ell_{u\to v}$ and $\ell_{u\to w}$ (elements of L) are not in the same plane, for any v, w not in the same line passing through the origin. Let $u = (x_0, y_0), v = (x_1, y_1), w = (x_2, y_2)$. Then, by the proof of Lemma 2.2, we have

$$\ell_{u \to v} = \left\{ \left(\frac{x_1}{x_0}, \frac{x_0}{x_1}, \frac{y_1}{x_0} - \frac{y_0}{x_1} \right) + t \left(-\frac{y_0}{x_0}, \frac{y_1}{x_1}, -\frac{y_0 y_1}{x_0 x_1} \right) : t \in \mathbb{F}_q \right\},\$$

and

$$\ell_{u \to w} = \left\{ \left(\frac{x_2}{x_0}, \frac{x_0}{x_2}, \frac{y_2}{x_0} - \frac{y_0}{x_2} \right) + t \left(-\frac{y_0}{x_0}, \frac{y_2}{x_2}, -\frac{y_0 y_2}{x_0 x_2} \right) : t \in \mathbb{F}_q \right\}.$$

Since v and w lie in two distinct lines passing through the origin, we have $\frac{y_1}{x_1} \neq \frac{y_2}{x_2}$. Therefore, the line $\ell_{u \to v}$ is not parallel to the line $\ell_{u \to w}$. Moreover, since there exists no $\theta \in \text{SL}_2(\mathbb{F}_q)$ such that $\theta(u) = v, \theta(u) = w$, we have $\ell_{u \to v} \cap \ell_{u \to w} = \emptyset$. Therefore, $\ell_{u \to v}$ and $\ell_{u \to w}$ are neither parallel nor intersect, which means that $\ell_{u \to v}, \ell_{u \to w}$ are not in the same plane.

Moreover, we will show that for any $u \in B$, any 3 distinct lines $\ell_{u \to v_1}, \ell_{u \to v_2}, \ell_{u \to v_3}$ in L are not in the same plane for any v_1, v_2, v_3 in a line passing through the origin. We use the parametric formulas for the $\ell_{u \to v_i}$ given in the proof of Lemma 2.1. Note that these 3 lines are parallel to each other. Take a point Z on one of them other than its pinpoint. A simple determinant computation shows that Z and the three pinpoints do not lie in the same plane.

Now, let W be a plane in \mathbb{F}_q^3 . For any $a \in \{a_1, \ldots, a_{m_0}\} - e_1 \cup e_2$, the plane W can contain at most 2 lines of the form $\ell_{a \to u_2}$. Therefore W contains at most $2m_0$ lines from L.

Proof of Theorem 1.4. We denote the set of lines passing through the origin and intersecting the set $E - \{(0,0)\}$ by \mathcal{L} . If $|\mathcal{L}| = 2$, the statement follows from Lemma 2.3, so assume $|\mathcal{L}| \geq 3$. We partition \mathcal{L} into subsets $\mathcal{L} = \bigcup_i \mathcal{L}_i$, where each set \mathcal{L}_i is a union of $m_0^{(i)}$ lines and each contains exactly $m_1^{(i)}$ points from E. We now fall into the following cases:

Case 1: If there exists *i* such that $|\mathcal{L}_i| \ge 8+4c$, then the theorem follows directly from Proposition 2.5.

Case 2: If there exists *i* such that $3 \leq |\mathcal{L}_i| \leq 8 + 4c$, then the theorem follows directly from Lemma 2.4.

Case 3: If for each *i*, the cardinality of \mathcal{L}_i is either 1 or 2. Since $|\mathcal{L}| \ge 3$, we can take a union of 2 or 3 of the \mathcal{L}_i 's that consists of 3 or 4 lines and is preserved by R_E . Then Lemma 2.4 gives the upper bound $|R_E| \le 2 \times 4^5$.

Putting these cases together, the proof is complete.

Proof of Theorem 1.2. Apply Theorem 1.4: let C be such that whenever E is a subset of \mathbb{F}_q^2 such that $E - \{(0,0)\}$ intersects at least 2 lines through the origin, we have $|R_E| \leq C|E|^{3/2}$. Pick $c_2 > C.c_1^{3/2}$. Let E be a subset of \mathbb{F}_q^2 with $|E| \leq c_1 q^{\alpha}$. Suppose $|R_E| \geq c_2 q^{\beta}$, where $\beta \geq 3\alpha/2$.

Suppose that $E - \{(0,0)\}$ intersects at least 2 lines through the origin. Then

$$c_2 q^{\beta} \le |R_E| \le C|E|^{3/2} \le C(c_1 q^{\alpha})^{3/2} = C c_1^{3/2} q^{3\alpha/2}$$

which is a contradiction.

3 Some discussions

3.1 Other incidence bounds imply weaker results

One might ask if we can get stronger results when we replace Lemma 2.1 by other point-line incidence bounds in the literature. In this section, we discuss this question. We first recall the following three incidence bounds.

Theorem 3.1 ([1, 6]). Let P be a set of points and L be a set of lines in \mathbb{F}_q^3 . Then the number of incidences between P and L satisfies

$$\left| I(P,L) - \frac{|P||L|}{q^2} \right| \ll q\sqrt{|P||L|}.$$

Theorem 3.2 ([4]). Let P be a set of n distinct points and L be a set of m distinct lines in \mathbb{F}_q^3 . Assume that no plane contains more than $c\sqrt{m}$ lines from L for some constant c. Then we have

$$I(P,L) \ll |L||P|^{2/5} + |P|^{6/5}$$

By a generic projection, one can show that the number of incidences between P and L is at most the number of incidences between a point set and a line set in two dimensions over an arbitrary fields, which is quite weak in general. Therefore, the main result in [9] implies the following.

Theorem 3.3. Let P be a set of points and L be a set of lines in \mathbb{F}_p^3 . Assume that $|P|^{7/8} < |L| < |P|^{8/7}$ and $|P|^{-2}|L|^{13} \ll p^{15}$, then we have

$$I(P,L) \ll |P|^{11/15} |L|^{11/15}.$$

If we just plug these incidence bounds in the proof of Theorem 1.2, without checking conditions on the sets and the underlying fields, one has

- Theorem 3.1 implies $|S| \le q^2 m_1$.
- Theorem 3.2 implies $|S| \le (m_0 m_1)^{5/3}$.
- Theorem 3.3 implies $|S| \le m_0^{7/4} m_1^{11/4}$.

All of these bounds are not better than that of Theorem 1.2.

3.2 Acting on a set of lines over prime fields

For a set \mathcal{L} of lines through the origin in \mathbb{F}_p^2 , let

$$R_{\mathcal{L}} = \{ \theta \in \mathrm{SL}_2(\mathbb{F}_p) \mid \theta(\mathcal{L}) = \mathcal{L} \}.$$

The following theorem offers an upper bound for $|R_{\mathcal{L}}|$.

Theorem 3.4. Let \mathcal{L} be a set of m lines passing through the origin in \mathbb{F}_p^2 with 1 < m < p+1. Let $R_{\mathcal{L}}$ be the set of matrices θ in $\mathrm{SL}_2(\mathbb{F}_p)$ such that $\theta(\mathcal{L}) = \mathcal{L}$. Then $|R_{\mathcal{L}}| \ll mp$.

For a subset E of \mathbb{F}_p^2 , let \mathcal{L}_E be the set of lines passing through the origin one needs to cover the whole set E. Then it is clear that $|R_E| \leq |R_{\mathcal{L}_E}|$. Thus, this theorem implies $|R_E| \ll p|\mathcal{L}_E|$, which is better than the bound in Theorem 1.4 in the regime when $|E| \gg p^{2/3} |\mathcal{L}_E|^{2/3}$.

While a point-line incidence bound due to Mockenhaupt and Tao in [5] plays a crucial role in the proof of Theorem 1.4, the following incidence bound with Möbius hyperbolae given by Rudnev and Weeler in [8] is required in our proof of Theorem 3.4.

Theorem 3.5 ([8]). For p is a prime, let \mathcal{L} be set of lines $\ell \neq \{(x, y) \in \mathbb{F}_p^2 : x = 0\}$ in \mathbb{F}_p^2 passing through the origin, and R be the set of matrices θ in $\mathrm{SL}_2(\mathbb{F}_p)$ such that $\theta(\ell) \in \mathcal{L}, \forall \ell \in \mathcal{L}$. Then, if $|R| > |\mathcal{L}|$, we have

$$I(\mathcal{L} \times \mathcal{L}, R) \ll \frac{|R||\mathcal{L}|^2}{p} + |\mathcal{L}|^{\frac{1}{2}}|R| + \min\left(|\mathcal{L}|^{\frac{7}{5}}|R|^{\frac{4}{5}}, p^{\frac{1}{3}}|\mathcal{L}|^{\frac{4}{3}}|R|^{\frac{2}{3}}\right)$$

where $I(\mathcal{L} \times \mathcal{L}, R) = |\{((\ell, \ell'), \theta) : \theta \in R, \ell, \ell' \in \mathcal{L}, \theta(\ell) = \ell'\}|.$

Proof of Theorem 3.4. If m = 2, we have $|R_{\mathcal{L}}| \leq 2(p-1)$ by the proof of Lemma 2.3. If $3 \leq m \leq C$ for a constant C, we apply Lemma 2.4. For $m \gg 1$, we argue as follows.

Without loss of generality, under m , we can assume that the*y* $-axis is not in <math>\mathcal{L}$. If $|R_{\mathcal{L}}| < |\mathcal{L}| = m$, it is clear that

$$|R_{\mathcal{L}}| \ll mp.$$

If $|R_{\mathcal{L}}| > |\mathcal{L}|$, then following Theorem 3.5, we have

$$I(\mathcal{L} \times \mathcal{L}, R_{\mathcal{L}}) \ll \frac{m^2 |R_{\mathcal{L}}|}{p} + |m|^{1/2} |R_{\mathcal{L}}| + \min\left(m^{\frac{7}{5}} |R_{\mathcal{L}}|^{\frac{4}{5}}, p^{\frac{1}{3}} m^{\frac{4}{3}} |R_{\mathcal{L}}|^{\frac{2}{3}}\right).$$

On the other hand, since $\theta(\ell) \in \mathcal{L}$ for all $\theta \in R_{\mathcal{L}}$ and $\ell \in \mathcal{L}$, we have

$$I(\mathcal{L} \times \mathcal{L}, R_{\mathcal{L}}) = |\mathcal{L}||R_{\mathcal{L}}| = m|R_{\mathcal{L}}|.$$

Therefore,

$$m|R_{\mathcal{L}}| \ll \frac{m^2|R_{\mathcal{L}}|}{p} + |m|^{1/2}|R_{\mathcal{L}}| + \min\left(m^{\frac{7}{5}}|R_{\mathcal{L}}|^{\frac{4}{5}}, p^{\frac{1}{3}}m^{\frac{4}{3}}|R_{\mathcal{L}}|^{\frac{2}{3}}\right).$$

Since $m \gg 1$, by a direct computation, one has

$$|R_{\mathcal{L}}| \ll mp. \qquad \Box$$

We can give an analogue of Theorem 1.2.

Theorem 3.6. Given $c_1 > 0$, there exists a constant $c_2 > 0$ with the following property. Let $\gamma \ge 0$. Let E be a subset of \mathbb{F}_p^2 such that the number m of lines through the origin that intersect E is at most c_1p^{γ} . Suppose $|R_E| \ge c_2p^{1+\gamma}$. Then E is contained in a line.

Remark 3.7. It should not be difficult to extend Theorem 3.5 to arbitrary finite fields with the method in [8]. However, we will again obtain a result in terms of the parameter m, which is the number of lines passing through the origin that one would need to cover the whole set E. This is a weakness compared to Theorem 1.2.

4 Acknowledgements

The authors would like to thank the Vietnam Institute for Advanced Study in Mathematics (VIASM) for the hospitality and the excellent working conditions, where part of this work was done. Thang Pham was partially supported by ERC Advanced Grant no. 882971, "GeoScape", and by the Erdős Center.

References

- M. Bennett, A. Iosevich, and J. Pakianathan, Three-point configurations determined by subsets of via the Elekes-Sharir Paradigm, Combinatorica, 34(6) (2014), 689–706.
- [2] N. Hegyvári, L. Q. Hung, A. Iosevich, and T. Pham, Packing sets under finite groups via algebraic incidence structures, arXiv:2411.05377 (2024).
- [3] A. Iosevich, P. Mattila, E. Palsson, M. Q. Pham, T. Pham, S. Senger, and C. Y. Shen, Packing sets in euclidean space by affine transformations, arXiv:2405.03087 (2024).
- [4] J. Kollár, Szemerédi-Trotter-type theorems in dimension 3, Advances in Mathematics, 271 (2015), 30–61.
- [5] G. Mockenhaupt and T. Tao, Restriction and Kakeya phenomena for finite fields, Duke Mathematical Journal, 121 (1) (2004), 35–74.
- [6] N. D. Phuong, T. Pham, and L. A. Vinh, *Incidences between planes over finite fields*, Proceedings of the American Mathematical Society, 147(5) (2019), 2185–2196.
- [7] T. Pham, Triangles with one fixed side-length, a Furstenberg-type problem, and incidences in finite vector spaces, Forum Mathematicum, in press, 2024.
- [8] M. Rudnev, and J. Weeler, On incidence bounds with Möbius hyperbolae in positive characteristic, Finite Fields and Their Applications, **78** (2022).
- [9] S. Stevens and F. De Zeeuw, An improved point-line incidence bound over arbitrary fields, Bulletin of the London Mathematical Society, 49(5) (2017), 842–858.