

# Beyond Non-Degeneracy: Revisiting Certainty Equivalent Heuristic for Online Linear Programming

Yilun Chen<sup>†</sup> and Wenjia Wang<sup>‡</sup>

<sup>†</sup>School of Data Science, the Chinese University of Hong Kong, Shenzhen (CUHK Shenzhen)

<sup>‡</sup>Data Science and Analytics Thrust, Information Hub, Hong Kong University of Science and Technology (Guangzhou)

chenyilun@cuhk.edu.cn, wenjiawang@hkust-gz.edu.cn

January 6, 2025

## Abstract

The Certainty Equivalent (CE) heuristic is a widely-used algorithm for various dynamic resource allocation problems in OR and OM. Despite its popularity, existing theoretical guarantees of CE are limited to settings satisfying restrictive fluid regularity conditions, particularly, the non-degeneracy conditions, under the widely held belief that the violation of such conditions leads to performance deterioration and necessitates algorithmic innovation beyond CE.

In this work, we conduct a refined performance analysis of CE within the general framework of online linear programming. We show that CE achieves uniformly near-optimal regret (up to a polylogarithmic factor in  $T$ ) under only mild assumptions on the underlying distribution, without relying on any fluid regularity conditions. Our result implies that, contrary to prior belief, CE effectively beats the curse of degeneracy for a wide range of problem instances with continuous conditional reward distributions, highlighting the distinction of the problem's structure between discrete and non-discrete settings. Our explicit regret bound interpolates between the mild  $(\log T)^2$  regime and the worst-case  $\sqrt{T}$  regime with a parameter  $\beta$  quantifying the minimal rate of probability accumulation of the conditional reward distributions, generalizing prior findings in the multisecretary setting.

To achieve these results, we develop novel algorithmic analytical techniques. Drawing tools from the empirical processes theory, we establish strong concentration analysis of the solutions to random linear programs, leading to improved regret analysis under significantly relaxed assumptions. These techniques may find potential applications in broader online decision-making contexts.

*Subject classifications:* Certainty Equivalent Heuristic, Regret Analysis, Degeneracy, Online Linear Programming, Network Revenue Management, Dynamic Resource Allocation, Multisecretary

# 1 Introduction

A variety of important OR and OM problems involve allocating finite, non-replenishable resources to sequentially arriving, random requests, in order to maximize cumulative rewards. Typical examples include the multisecretary problem (Kleinberg, 2005; Arlotto and Gurvich, 2019; Besbes et al., 2024), network revenue management (NRM) (Gallego and Van Ryzin, 1994; Talluri and Van Ryzin, 2006; Jasin and Kumar, 2012), dynamic bidding in repeated auctions (Balseiro et al., 2015) and order fulfillment (Jasin and Sinha, 2015), among others. Despite their diverse application background, these problems share common characteristics and can be treated through a unified modeling framework, typically termed *online linear programming* (OLP) as is noticed and explored in a recent line of research (Vera and Banerjee, 2021; Li and Ye, 2022; Bray, 2024).

The workhorse algorithm for OLP is the **certainty equivalent heuristic** (CE). Initially proposed in the broader context of stochastic control (cf. Bertsekas (2012)), CE relies on the idea of replacing all random variables with their average values and repeatedly solving the resulting static and deterministic problem (usually referred to as the fluid problem) at each decision epoch to facilitate its online decision-making. This heuristic algorithm comes with computational tractability, offering an advantage over the optimal, yet computationally burdensome dynamic programming (DP) approach. CE and its variants have been extensively researched in various contexts of OLP (or more generally, dynamic optimization with resource constraints), sometimes under alternative names such as the frequent re-solving heuristic or the online greedy (Lueker, 1998; Jasin and Kumar, 2012; Li and Ye, 2022; Bray, 2024; Balseiro, Besbes and Pizarro, 2023). A central focus in this body of work is to understand CE’s performance, typically measured by the regret, which captures the asymptotic loss in reward relative to the optimal DP (or other performance benchmarks such as the hindsight optimum or the fluid optimum), as the system scales.

Impressive regret guarantees for CE—either independent of  $T$  or growing logarithmically in  $T$  (where  $T$  is the length of the time horizon)—have been established under certain fluid regularity conditions, most notably, the *non-degeneracy conditions* (cf. Assumptions 4.1 and 4.2). Despite their prevalence, these conditions are unsatisfying for several reasons, as recently noted in the literature (Jiang et al., 2022a). First, they are not imposed directly on the problem’s primitives, in particular, the underlying distribution of the resource consumption and reward of an individual request and the initial inventory of resources. Rather, they concern the stability of the optimal solution to the fluid problem, making them difficult to intuit and verify. Second, degeneracy is likely to occur in practice (Bumpensanti and Wang, 2020), violating these conditions and thus limiting

the applicability of performance guarantees that depend on them.

Existing efforts to relax restrictive fluid regularity conditions are mostly limited to the *discrete* environment where arriving requests belong to finitely many types. In such settings, a clear understanding of the relationship between problem structure and achievable algorithmic performance has been established: (i) degeneracy causes CE to fail, and (ii) provably uniformly low regret is attainable by novel algorithmic innovations—mostly variants of the standard CE (see literature review). However, in practice, especially with the increasing availability of data and advances in prescriptive analytics, many OLP instances involve an overwhelmingly large variety of request types, making continuous distributions a more suitable modeling choice. Unfortunately, algorithmic analysis beyond fluid regular instances are substantially more challenging in non-discrete settings. Despite existing efforts such as Besbes et al. (2024) and Jiang et al. (2022a), there lacks systematic technical tools in general. Consequently, whether and when any algorithm, or even the standard CE can attain uniform low regret guarantees in these settings, and which structural features of the problem fundamentally drive regret accumulation, remain largely unknown.

The above naturally motivates the following questions. (i) *Are there weaker, more intuitive conditions under which CE achieves low regret?* (ii) *Which structures of the OLP problem determine the performance of CE (or other algorithms)? Are fluid regularity conditions essential?*

In this work, we answer question (i) in the affirmative. Our analysis reveals that, surprisingly, CE “beats” the curse of degeneracy in non-discrete settings in a broad sense, contrary to the understanding established in the discrete setting, and shedding light on question (ii). Next we highlight our contributions in detail.

**Regret guarantees.** We identify two broad classes of request distributions under which CE achieves uniformly near-optimal (up to a polylogarithmic factor in  $T$ ) hindsight regret for any initial resource inventory, without relying on fluid regularity conditions (cf. Theorem 3.1 and Proposition 3.2). These distribution classes encompass rich structures, including both well-explored OLP instances in the literature such as the multisecretary problems (cf. Example 2.1) and less understood but practically relevant ones such as the generalized linear models (cf. Example 2.3), among others. They essentially generalize the class of “gap-free” reward distributions (Bray, 2024), which have been shown as necessary for CE to achieve  $o(\sqrt{T})$  regret in the multisecretary setting (Besbes et al., 2024). In particular, Our first class allows for arbitrary resource consumption distributions but requires all the conditional reward CDFs to be supported on an interval starting from zero. The second class relaxes this condition, requiring it to hold only locally, provided that the distribution

is well-behaving.

The explicit regret scaling,  $\tilde{\Theta}\left(T^{\frac{1}{2}-\frac{1}{2(1+\beta)}}(\log T)^{\frac{2+\beta}{2+2\beta}+\mathbb{1}\{\beta=0\}}\right)$ , interpolates between the mild  $(\log T)^2$  regime and the worst-case  $\sqrt{T}$  regime with a parameter  $\beta \in [0, \infty)$ . This parameter generalizes the critical regret driver identified in the multisecretary setting (Besbes et al., 2024), quantifying the minimal probability mass accumulation of the conditional reward CDFs. Specifically,  $\beta = 0$  corresponds to uniform-like conditional reward distributions, yielding the mild regret bound of  $\mathcal{O}((\log T)^2)$ . As  $\beta \rightarrow \infty$ , the conditional reward CDFs converge towards distributions with gaps on their supports, and the achievable regret scaling worsens, approaching  $\sqrt{T}$ . Both the polylogarithmic regret bound and the interpolation between different  $o(\sqrt{T})$  regret regimes driven by the parameter  $\beta$  are novel to the literature. They establish a direct connection between the properties of the underlying request distribution and the best achievable regret scaling of CE in the general OLP setting, providing a definitive affirmative answer to question (i).

**Algorithmic insights.** Our results reveal distinct patterns of the performance of CE in non-discrete settings. Most notably, we find that CE effectively “beats” the curse of degeneracy, achieving uniformly low (and near-optimal) regret even when standard non-degeneracy conditions are violated (cf. Corollary 4.2) for a wide range of non-discrete request distributions. This contrasts sharply with earlier observations, e.g. Bumpensanti and Wang (2020) in the discrete setting, where CE provably incurs  $\Theta(\sqrt{T})$  regret whenever degeneracy occurs, regardless of the underlying distribution.

We thoroughly discuss and explain this phenomenon, attributing it to differing relationships—in discrete and non-discrete settings—between *dual uniqueness*, namely, the uniqueness of the optimal solution to the dual fluid problem, and standard non-degeneracy notions, where we identify dual uniqueness as the critical determinant of whether CE suffers from performance deterioration (cf. Proposition 4.6). In discrete settings, we prove an equivalence between dual uniqueness and other notions of non-degeneracy (cf. Lemma 4.3), an implication of the piecewise linearity of the dual fluid objective. In contrast, in non-discrete settings, the dual fluid objective is smooth when the underlying distribution belongs to the class that we identify, under which dual uniqueness always holds yet non-degeneracy conditions may fail, breaking the equivalence between them (cf. Lemma 4.4). These analyses highlight an intriguing shift in the role of standard non-degeneracy notions: while necessary to prevent CE performance deterioration in discrete settings, they prove to be overly restrictive in non-discrete settings. Overall, our results imply that the standard CE is effective for a broader range of instances than previously understood.

**Novel methodology.** We develop novel regret analysis techniques that bypass the need for fluid regularity conditions to derive improved ( $o(\sqrt{T})$ ) regret bounds in non-discrete settings. In prior literature, fluid regularity conditions are often imposed as sufficient conditions to enable strong martingale-type arguments, which requires that the set of binding resource constraints remains identical throughout the algorithm’s execution. We overcome this restrictive requirement by carefully decomposing the hindsight regret of CE (Lemma 5.1) and performing a concentration analysis of the solution to the per-step sample average approximation (SAA) of the fluid dual problem at each time  $t$  (Lemma 5.2). The concentration analysis is the key technical innovation that allows us to establish improved regret guarantees. Using the peeling device from empirical processes theory, we partition the solution space and apply union bounds with carefully controlled entropy numbers to establish robust worst-case concentration bounds. These bounds hold even when the set of binding constraints changes over time, enabling regret guarantee under significantly relaxed assumptions. The worst-case analysis leaves us with an unavoidable additional  $\log T$  in the regret bounds, which is a mild price to pay to get rid of the fluid regularity conditions. We believe the concentration analyses are of independent interest and the techniques developed may have broader applicability, particularly since such concentration properties are relevant to the performance of CE-type algorithms in various online stochastic optimization settings with continuous underlying distributions.

## 1.1 Related Literature

**CE and Fluid-Regular OLP.** In the revenue management literature, the (quantity-based) NRM problem has been formulated as an discrete OLP instance and received considerable attention (Talluri and Van Ryzin, 2006). Earlier work (Gallego and Van Ryzin, 1994) established  $\mathcal{O}(\sqrt{T})$  regret guarantee under static heuristic algorithms. Reiman and Wang (2008) obtained an improved  $o(\sqrt{T})$  regret bound by introducing re-solving in their algorithm design. Jasin and Kumar (2012) analyzed the frequent re-solving heuristic (a primal version of CE) and established an  $\mathcal{O}(1)$  regret under the non-degeneracy condition of the fluid LP. Bumpensanti and Wang (2020), among others, further confirmed that non-degeneracy is necessary for CE to achieve this optimal (fluid) regret. In non-discrete settings, where requests are modeled with continuous distributions, Lueker (1998) initiated the study of OLP with a single resource constraint and proved an  $\mathcal{O}(\log T)$  regret bound for CE (referred to as the **online greedy**). This bound was shown to be tight in the multisecretary problem by Bray (2024). Recent works, including Li and Ye (2022), Bray (2024),

and Balseiro, Besbes and Pizarro (2023) extended the  $\mathcal{O}(\log T)$  regret guarantee of CE to general multi-resource settings, but their analyses critically rely on restrictive fluid regularity conditions, which we aim to remove in this work.

**Uniform Loss in OLP.** To handle the challenges posed by fluid degeneracy on CE, a recent stream of papers have proposed new algorithms in the discrete setting and achieved  $\mathcal{O}(1)$  hindsight regret uniformly across degenerate and non-degenerate instances (Arlotto and Gurvich, 2019; Bumpensanti and Wang, 2020; Vera and Banerjee, 2021). The removal of fluid regularity conditions in non-discrete settings, however, is substantially more challenging. Existing attempts (Besbes et al., 2024; Jiang et al., 2022a) are limited to structured settings. Besbes et al. (2024) focused on the multisecretary problem and introduced an algorithmic adjustment (termed CwG) to the standard CE that attains near optimal regret guarantees for general reward distributions with gaps on their supports. They also propose a simulation-based algorithm (RAMS) in the general OLP settings, which provably achieves the regret guarantee (up to a simulation error) of any other algorithm. Jiang et al. (2022a) studied a “semi-discrete” setting with finitely many resource consumption types, achieving an  $\mathcal{O}((\log T)^2)$  regret guarantee under the **boundary-attracted algorithm**, an adjusted CE heuristic. Notably, they also provide an  $\mathcal{O}(\log T)$  regret of the standard CE in the same setting, contingent on an additional uniform second-order growth condition, for which we provide further discussion in Appendix E. Overall, the (semi)discreteness of the underlying distribution seems critical to the analysis in all the aforementioned work. Our work complements this literature with a fresh perspective that for a wide class of non-discrete request distributions, the standard CE already achieves uniformly near-optimal regret scaling, effectively “beating” the curse of degeneracy. From a methodological point of view, our work contributes technical tools for algorithmic analysis in non-discrete settings, potentially paving the way for further generalization of this literature to incorporate more complicated non-discrete cases.

**Degeneracy in Stochastic Control.** In the broader context of stochastic dynamic control, the impact of fluid degeneracy/instability on algorithm performance has been widely noted, for example, in stochastic network optimization (Huang and Neely, 2009), centralized dynamic matching (Kerimov et al., 2024; Gupta, 2024; Wei et al., 2023) and network revenue management with reusable resources (Xie et al., 2024; Balseiro, Ma and Zhang, 2023). While these settings differ from those considered in this work, they bear notable similarities, especially in the role of the fluid relaxation in algorithm design and performance analysis. Particularly, Gupta (2024); Xie et al. (2024); Chen et al. (2024) highlight the critical role of dual uniqueness in guaranteeing good algorithm performance in their respective settings. These results are primarily in the discrete setting,

similar to Bumpensanti and Wang (2020). The perspective provided in this work complements and extends this understanding of the significance of dual uniqueness by offering non-discrete evidence.

**Online Stochastic Knapsack.** There is also a rich relevant literature in theoretical computer science studying the problem of online stochastic knapsack. Kleinberg (2005) considers a multisecretary problem in the random input model and derive an (asymptotic) competitive ratio of  $1 - \mathcal{O}\left(\sqrt{\frac{1}{k}}\right)$ , where  $k$  is the fixed budget (analogous to  $\mathbf{b}$  in this paper). Subsequent work Devanur and Hayes (2009); Feldman et al. (2010); Molinaro and Ravi (2014); Agrawal et al. (2014) extend Kleinberg (2005) to the general setting of online knapsack/online packing and achieve progressively improving competitive ratios. Hajiaghayi et al. (2007); Alaei (2014); Chawla et al. (2023); Jiang et al. (2022b) study the  $k$ -unit prophet problem where the inputs are independent (but not identical) random variables. Competitive ratios of  $1 - \tilde{\Theta}\left(\sqrt{\frac{1}{k}}\right)$  for large  $k$  as well as concrete constants for small  $k$  were shown. The setting of this work is different from the aforementioned line of works: (i) the performance metric we use in this paper is the additive regret rather than the competitive ratio, and (ii) the focus is on understanding the performance of CE in the *i.i.d.* input model for regular distributions, rather than competing against the worst case. The main result of this work can be essentially translated to an  $\left(1 - \mathcal{O}\left(\frac{(\log T)^2}{T}\right)\right)$ -competitive ratio for distributions with nice structures and the initial inventory of resources scaling linearly in  $T$ , for a general online stochastic knapsack problem with *i.i.d.* inputs.

## 1.2 Organization

The paper is organized as follows. We formally set up the problem in Section 2. The main results are stated in Section 3. We then discuss the implication of our results on fluid degeneracy in Section 4. The proof sketch of our main results is provided in Section 5. The conclusions follow in Section 6.

# 2 Problem Formulation

## 2.1 Model

There are  $m$  resources with an initial capacity of  $\mathbf{b} = (b_1, b_2, \dots, b_m)^\top \in \mathbb{R}_{\geq 0}^m$ . At each time period  $t = 1, \dots, T$ , a demand request  $(\mathbf{a}_t, r_t)$  arrives, assumed to be drawn *i.i.d.* from a distribution with joint CDF  $F(\cdot)$ , where  $F, \mathbf{b}$  and  $T$  are known *a priori*. Upon the arrival of the request and the revelation of  $(\mathbf{a}_t, r_t)$ , a decision maker (DM) needs to immediately and irrevocably decide whether or not to accept it. The ACCEPT decision results in the consumption of  $\mathbf{a}_t = (a_{1t}, \dots, a_{mt})^\top$  of each

resource and an earned reward of  $r_t$ . The REJECT decision imposes no change on the resources and garners zero reward. The ACCEPT decision is feasible if and only if the remaining capacity of resource  $i$  is at least  $a_i$ , for  $i = 1, \dots, m$ . The goal of the DM is to maximize the expected total reward collected through the  $T$  periods subject to the resource capacity constraint.

A dynamic allocation policy  $\pi$  specifies, for each demand sequence realization  $\mathcal{I} \triangleq \{(\mathbf{a}_t, r_t)\}_{j=1}^T$ , a (possibly random) sequence of binary decisions  $\{x_t^\pi\}_{t=1}^T$  with  $x_t^\pi = 1(0)$  corresponding to the accept (reject) decision.  $\pi$  is *non-anticipatory* if each  $x_t^\pi$  is independent of the future request realization  $\mathcal{I}_t \triangleq \{(\mathbf{a}_j, r_j)\}_{j=t+1}^T$  and depends only on the current history  $\mathcal{H}_t \triangleq \{(\mathbf{a}_j, r_j)\}_{j=1}^t$ .  $\pi$  is *feasible* if *w.p.1* the capacity constraint is satisfied throughout, i.e.  $\sum_{t=1}^T a_{it}x_t^\pi \leq b_i, i = 1, \dots, m$ . A policy is admissible if it is both non-anticipatory and feasible. Let  $\Pi$  denote the set of admissible policies. The DM seeks to maximize  $\mathbb{E} \left[ \sum_{t=1}^T r_t x_t^\pi \right]$ , the expected total reward collected under an admissible policy  $\pi \in \Pi$ .

**Performance Metric.** The performance of an admissible policy  $\pi$  is typically measured by its revenue loss, namely  $\sup_{\tau \in \Pi} \mathbb{E} \left[ \sum_{t=1}^T r_t x_t^\tau \right] - \mathbb{E} \left[ \sum_{t=1}^T r_t x_t^\pi \right]$ , where  $\sup_{\tau \in \Pi} \mathbb{E} \left[ \sum_{t=1}^T r_t x_t^\tau \right]$  is the optimal dynamic programming (DP) value. However, the so-defined revenue loss is often not a viable performance metric due to the lacking of tractability of the optimal DP value. A common practice is to instead consider upper bounds of the revenue loss, derived by replacing the DP value by the optimal values of certain tractable relaxations. Two most popular such relaxations are the hindsight relaxation and the fluid relaxation.

**The hindsight relaxation.** Under the hindsight relaxation, the non-anticipatory constraint is removed, and the resource allocation problem becomes a multi-knapsack Linear Program (LP) (for a given demand sequence realization  $\mathcal{I}$ ):

$$\begin{aligned} \max \quad & \sum_{t=1}^T r_t x_t \\ \text{s.t.} \quad & \sum_{t=1}^T a_{it} x_t \leq b_i, \quad i = 1, \dots, m, \\ & x_t \in [0, 1], \quad t = 1, \dots, T. \end{aligned} \tag{1}$$

Taking expectation of (1) (over  $\mathcal{I}$ ) yields the optimal hindsight value, denoted by  $V_{\mathbf{b}, T}^{\text{hind}}$ .

**The fluid relaxation.** The fluid relaxation further assumes that  $T$  is prohibitively large, to the point that all randomness in (1) is averaged out (with a proper scaling on the order of  $\frac{1}{T}$ ),



which yields

$$\begin{aligned} & \sup_x \mathbb{E}_{(\mathbf{a}, r) \sim F} [rx(\mathbf{a}, r)] \\ & \text{s.t. } \mathbb{E}_{(\mathbf{a}, r) \sim F} [\mathbf{a}x(\mathbf{a}, r)] \leq \mathbf{d}, \quad x(\cdot, \cdot) \in [0, 1], \end{aligned} \tag{2}$$

where  $\mathbf{d} \triangleq \frac{\mathbf{b}}{T}$  is the normalized inventory of resources. We denote the optimal value of the above fluid program by  $V_{\mathbf{d}}^{\text{fluid}}$ .

It is fairly straightforward to see that  $V_{\mathbf{d}}^{\text{fluid}}T \geq V_{\mathbf{b}, T}^{\text{hind}}$  both are upper bounds on the optimal DP value, for  $T \geq 1$ , for which we omit the proof. We denote by  $\text{REG}_{\mathbf{b}, T}^{\text{hind}}(\pi) \triangleq V_{\mathbf{b}, T}^{\text{hind}} - \mathbb{E} \left[ \sum_{t=1}^T r_t x_t^\pi \right]$  the *hindsight regret* incurred by policy  $\pi$ , and  $\text{REG}_{\mathbf{b}, T}^{\text{fluid}}(\pi) \triangleq V_{\mathbf{d}}^{\text{fluid}}T - \mathbb{E} \left[ \sum_{t=1}^T r_t x_t^\pi \right]$  the *fluid regret* incurred by policy  $\pi$ . Throughout this paper, the hindsight regret<sup>1</sup> will be our main performance metric. We thereby denote  $\text{REG}_{\mathbf{b}, T}(\pi) \triangleq \text{REG}_{\mathbf{b}, T}^{\text{hind}}(\pi)$ . We are interested in characterizing the scaling of  $\text{REG}_{\mathbf{b}, T}(\pi)$  as  $T$  grows.

## 2.2 The Certainty Equivalent Heuristic

The CE heuristic is a special threshold-based policy that leverages the fluid relaxation (2) to facilitate its dynamic decision-making. Following Li and Ye (2022); Bray (2024); Jiang et al. (2022a), we consider a dual-based CE. The dual of (2) has a compact form

$$\min_{\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m} f_{\mathbf{d}}(\boldsymbol{\lambda}) \triangleq \mathbf{d}^\top \boldsymbol{\lambda} + \mathbb{E}_{(\mathbf{a}, r) \sim F} [(r - \mathbf{a}^\top \boldsymbol{\lambda})^+], \tag{3}$$

where  $f_{\mathbf{d}}(\cdot)$  is convex for any value that  $\mathbf{d}$  takes. Strong duality holds, namely we have  $V_{\mathbf{d}}^{\text{fluid}} = \min_{\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m} f_{\mathbf{d}}(\boldsymbol{\lambda})$ . We refer the reader to Li and Ye (2022); Balseiro, Besbes and Pizarro (2023) for the derivation of (3) and further discussions.

The CE heuristic solves a perturbed version of (3) at each period  $t = 1, \dots, T$  to facilitate decision making. More precisely, let  $\mathbf{b}_t^{\text{CE}}$  denote the sequence of remaining inventory of resources under the CE policy, with  $\mathbf{b}_0^{\text{CE}} = \mathbf{b}$ , and  $\mathbf{d}_t^{\text{CE}} \triangleq \frac{\mathbf{b}_{t-1}^{\text{CE}}}{T-t}$  be the corresponding normalized remaining inventory of resources. The CE heuristic solves for

$$\tilde{\boldsymbol{\lambda}}_t \in \arg \min_{\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m} \left( \mathbf{d}_t^{\text{CE}} \right)^\top \boldsymbol{\lambda} + \mathbb{E}[(r - \mathbf{a}^\top \boldsymbol{\lambda})^+] \tag{4}$$

---

<sup>1</sup>The notion ‘‘regret’’ is typically used in a learning scenario. We slightly deviate from this convention as in our setting, the underlying distribution  $F$  is known. Instead, our DM regrets for not knowing the future demand sequence in advance.

at each period  $t = 1, \dots, T$ . Then the decision follows by setting threshold at  $\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t$ :

$$x_t^{\text{CE}} = 1 \text{ if and only if } r_t \geq \mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t \text{ and } \mathbf{a}_t \leq \mathbf{b}_{t-1}^{\text{CE}}, \quad t = 1, \dots, T,$$

where  $\mathbf{b}_t^{\text{CE}} = \mathbf{b}_{t-1}^{\text{CE}} - x_t^{\text{CE}} \mathbf{a}_t$ , namely, the induced sequence of remaining inventory of resources. The inequality in  $\mathbf{a}_t \leq \mathbf{b}_{t-1}^{\text{CE}}$  is element-wise. A formal description is provided in Algorithm 1.

---

**Algorithm 1:** Certainty Equivalent (CE) Heuristic

---

**Require:** Problem instance  $(F, \mathbf{b}, T)$ .

1: **for**  $t = 1, \dots, T$  **do**

2: Observe instance  $(\mathbf{a}_t, r_t)$ .

3: Solve (4) and obtain  $\tilde{\boldsymbol{\lambda}}_t \in \arg \min_{\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m} f_{\mathbf{d}_t^{\text{CE}}}(\boldsymbol{\lambda})$ .

4: Set

$$x_t^{\text{CE}} = 1 \text{ if and only if } r_t \geq \mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t \text{ and } \mathbf{a}_t \leq \mathbf{b}_{t-1}^{\text{CE}},$$

and  $x_t^{\text{CE}} = 0$  otherwise, where  $\mathbf{b}_0^{\text{CE}} = \mathbf{b}$ ,  $\mathbf{b}_t^{\text{CE}} = \mathbf{b}_{t-1}^{\text{CE}} - x_t^{\text{CE}} \mathbf{a}_t$  is the induced sequence of remaining resources.

5: **end for**

---

We take the dual-based definition of CE in our setting because dual fluid relaxation (3), in a cleaner form, both benefits the theoretical analysis and yields a more practical algorithm. In fact, the primal problem (2) is a possibly infinite-dimensional optimization problem since we allow  $F$  to be a general continuous distribution, while the dual fluid problem (3) is a convex program with a simple feasible region (the  $m$ -dimensional positive orthant).

### 2.3 Assumptions

Our main results are presented under two sets of distributional assumptions on  $F$ . **Either** set of assumptions suffices to guarantee the desired performance of CE (cf. Theorem 3.1). We first introduce some additional notation.

**Additional notation.** Let  $F^{\mathbf{a}}(\cdot)$  denote the marginal CDF of  $\mathbf{a}$ . Further let  $F_{\mathbf{a}'}^r(\cdot)$  denote the conditional CDF of the reward  $r$  given  $\mathbf{a} = \mathbf{a}'$ . Let  $\text{supp}(F)$ ,  $\text{supp}(F^{\mathbf{a}})$  and  $\text{supp}(F_{\mathbf{a}'}^r)$  denote the support of the corresponding distributions, respectively.

**Assumption 2.1.** The joint distribution  $F$  satisfies the following conditions:

(i) (boundedness) There exist constants  $0 < \underline{A} \leq \bar{A}$  and  $\bar{r} > 0$ , s.t.  $\text{supp}(F) \in [\underline{A}, \bar{A}]^m \times [0, \bar{r}]$ .

(ii) For any  $\mathbf{a} \in \text{supp}(F^{\mathbf{a}})$ ,  $\text{supp}(F_{\mathbf{a}}^r) = [0, r_{\mathbf{a}}]$  for some  $r_{\mathbf{a}} > 0$ .

(iii) ((reverse) Hölder condition) There exist non-negative constant  $\beta$  and positive constants  $\nu, c_\nu, c_\beta$ , such that for any  $0 \leq z_1 < z_2 \leq r_{\mathbf{a}}$ ,  $c_\beta(z_2 - z_1)^{1+\beta} \leq F_{\mathbf{a}}^r(z_2) - F_{\mathbf{a}}^r(z_1) \leq c_\nu(z_2 - z_1)^\nu$ .

**Assumption 2.2.** The joint distribution  $F$  satisfies the following conditions:

- (i) (boundedness) There exist constants  $0 < \underline{A} \leq \bar{A}$  and  $\bar{r} > 0$ , s.t.  $\text{supp}(F) \in [\underline{A}, \bar{A}]^m \times [0, \bar{r}]$ .
- (ii) ( $F$  regularity)  $\text{supp}(F^{\mathbf{a}})$  is convex and compact, and the probability density associated to  $F^{\mathbf{a}}$  is bounded from below by a constant  $l_f > 0$ .  $\text{supp}(F_{\mathbf{a}}^r) = [l(\mathbf{a}), r(\mathbf{a})]$ , where  $l(\mathbf{a})$  and  $r(\mathbf{a})$  are Lipschitz continuous as functions of  $\mathbf{a}$  with Lipschitz constant  $c_L$ .
- (iii) There exist  $\mathbf{a}_o \in \text{supp}(F^{\mathbf{a}})$  and  $r_0 > 0$  s.t.  $l(\mathbf{a}) = 0$  for all  $\mathbf{a} \in \mathcal{B}(\mathbf{a}_o, r_0) \cap \text{supp}(F^{\mathbf{a}})$ .
- (iv) ((reverse) Hölder condition) There exist non-negative constant  $\beta$  and positive constants  $\nu, c_\nu, c_\beta$ , such that for any  $l(\mathbf{a}) \leq z_1 < z_2 \leq r(\mathbf{a})$ ,  $c_\beta(z_2 - z_1)^{1+\beta} \leq F_{\mathbf{a}}^r(z_2) - F_{\mathbf{a}}^r(z_1) \leq c_\nu(z_2 - z_1)^\nu$ .

Assumption 2.1 (i) enforces boundedness on  $\mathbf{a}$  and  $r$ . In (ii), we require the conditional reward distribution given any  $\mathbf{a}$  to be supported on an interval starting from zero. This is a necessary restriction for CE to achieve  $o(\sqrt{T})$  regret guarantee, when  $\mathbf{a}$  is allowed to be arbitrarily distributed (cf. Example 4.1 for bad examples violating (ii)). (iii) further excludes point masses on the conditional reward distributions ( $\nu$  bounded away from zero), and restricts the minimal rate of probability accumulation ( $\beta$  bounded away from  $\infty$ ). In particular,  $\beta = 0$  and  $\nu = 1$  corresponds to the case where the conditional distribution of the reward has lower and upper bounded density on the support. We remark on the asymmetric role of parameters  $\beta$  and  $\nu$ :  $\beta$  critically affects the best achievable regret scaling, while  $\nu$  only appears in the constant term and does not affect the regret scaling (cf. Theorem 3.1, Proposition 3.2). We refer the readers to Appendix E for more discussion on parameter  $\beta$ , in connection with the second-order growth conditions often made in the literature.

Assumption 2.2 relaxes the global requirement of Assumption 2.1 (ii) to only a local condition (iii), to hold only in the neighborhood of an arbitrary point in the support. To permit this relaxation, Assumption 2.2 (ii) enforces natural regularity of  $F$ . We refer the readers to Appendix E (cf. Example E.1) for further discussion on the condition (iii).

In the sequel, we illustrate Assumption 2.1 and Assumption 2.2 with several examples.

**Example 2.1** (Multisecretary problem). The multisecretary problem has  $m = 1$  and  $F^{\mathbf{a}}(\cdot) = \delta_1(\cdot)$ . Suppose the reward distribution is (i) without point mass, and (ii) supported on an interval starting

from zero, namely, without gaps, then Assumption 2.1 holds. A notable example is a reward distribution specified by *p.d.f.*  $h_r(x) = (1 + \beta)|1 - 2x|^\beta$ , for  $0 \leq x \leq 1$  with  $\beta \geq 0$ .

**Example 2.2** (Hyper-cube models). Suppose  $\text{supp}(F) = [1, 2]^m \times [0, 1]$ . The joint density function of  $F$  is bounded from above and below by a pair of positive constants on the support. Then both Assumptions 2.1 and 2.2 are satisfied with  $\beta = 0$  and  $\nu = 1$ .

**Example 2.3** (Generalized linear models). Let  $r = g(\mathbf{a}^\top \mathbf{z}) + \epsilon$  for a non-negative, Lipschitz continuous function  $g$ , a fixed vector  $\mathbf{z} \in \mathbb{R}_{\geq 0}^m$  and a noise random variable  $\epsilon$ , with  $F^{\mathbf{a}}$  satisfying Assumption 2.2 (i) and (ii).  $\epsilon$  is supported on an interval  $[-\mathcal{L}, \mathcal{L}]$  with *p.d.f.* bounded from above and below by a pair of positive constants. Suppose there exists  $\mathbf{a}'$  such that  $g((\mathbf{a}')^\top \mathbf{z}) < L - \eta$  for a positive constant  $\eta$ . Then Assumption 2.2 effectively holds with  $\beta = 0$  and  $\nu = 1$ , modulo the possible negative rewards which can be dealt with in a straightforward manner.

**Remark 2.1.** The simple structure of multisecretary allows for a further relaxation of Assumption 2.1 (ii), such that  $\text{supp}(F_{\mathbf{a}}^r)$  is an interval not necessarily starting from zero, under which our main results (cf. Theorem 3.1) remain valid.

**Remark 2.2.** Negative rewards, as appeared in Example 2.3, do not incur regret: the optimal decision is always to reject them. We defer a formal analysis of Example 2.3 to Appendix A.

Later in Section 3, we state the regret guarantee of CE for each of the above examples under no other assumptions (cf. Corollary 3.1). To obtain similar regret guarantees, prior work typically imposes additional fluid regularity conditions (cf. Section 3 for details). In later sections we provide a systematic review—and a comparison with our own assumptions—of these fluid regularity conditions, with a detailed investigation of non-degeneracy conditions in Section 4, and discussion on the second-order growth conditions in Appendix E.

## 3 Main Results

### 3.1 Achievable regret

**Theorem 3.1** (Achievable regret of CE). *Under either Assumption 2.1 or Assumption 2.2, the CE heuristic achieves a hindsight regret*

$$\text{REG}_{\mathbf{b}, T}(\pi^{\text{CE}}) \leq \begin{cases} \mathcal{C}(\log T)^2 & \beta = 0, \\ \tilde{\mathcal{C}} T^{\frac{1}{2} - \frac{1}{2(1+\beta)}} (\log T)^{\frac{2+\beta}{2+2\beta}} & \beta > 0, \end{cases}$$

for arbitrary  $\mathbf{b}$  and  $T > 3$ , where  $\mathcal{C}$  and  $\tilde{\mathcal{C}}$  are constants independent of  $T$  and  $\mathbf{b}$  and depend only on model primitives regarding  $F$  through the two assumptions, respectively.

**Remark 3.1.** Precise forms of  $\mathcal{C}$  and  $\tilde{\mathcal{C}}$  are provided in Appendix B.

Theorem 3.1 establishes theoretical guarantees for CE for a wide range of OLP instances, requiring only that  $F$  belongs to specific distribution classes that are both natural and easy to verify. This significantly relaxes the conditions typically needed to obtain  $o(\sqrt{T})$  regret, in particular, the non-degeneracy conditions and/or (uniform) second-order growth conditions. In Section 4 and Appendix E, we systematically examine the relationship between these fluid regularity conditions and our conditions (Assumptions 2.1 and 2.2), demonstrating that the former are not only technically unnecessary for algorithmic analysis but are often overly restrictive. In that sense, Theorem 3.1 extends the state-of-the-art understanding of CE’s range of effectiveness. We provide a proof sketch of Theorem 3.1 in Section 5. The detailed proof can be found in Appendix B. With Theorem 3.1, the following regret scaling of CE on concrete examples is an immediate corollary.

**Corollary 3.1.** *The regret guarantee in Theorem 3.1 holds for multisequential instances of Example 2.1 with each choice of  $\beta$ . For instances of Example 2.2 and Example 2.3, CE achieves  $\mathcal{O}((\log T)^2)$  regret.*

## 3.2 Fundamental regret lower bound

The achievable regret of CE stated in Theorem 3.1 is near optimal, as we formalize through a fundamental regret lower bound.

**Proposition 3.2** (Fundamental Regret Lower Bound). *There exists OLP instance  $(F, \mathbf{b}, T)$  satisfying Assumption 2.1 and Assumption 2.2, such that*

$$\inf_{\pi} \text{REG}_{\mathbf{b}, T}(\pi) \geq \frac{c}{1 + \beta} T^{\frac{1}{2} - \frac{1}{2(1+\beta)}} \mathbb{1}(\beta > 0) + c \log T \mathbb{1}(\beta = 0),$$

where  $c$  is a constant independent of  $T, \mathbf{b}$  and  $\beta$ .

**Remark 3.2.** The fundamental regret lower bound is inspired by earlier works on the multisequential problem (Bray (2024) for  $\beta = 0$  and Besbes et al. (2024) for  $\beta > 0$ ). While Besbes et al. (2024) initially considers reward distributions with a gap in their support, we observe that these gaps can be removed so that the resulting instances satisfy our Assumptions, yet their analysis still works (formally, by setting  $g = 0$  in the proof of Theorem 1 in Besbes et al. (2024)) and the regret lower bound still holds. The proof is deferred to Appendix C.

Combining Theorem 3.1 with Proposition 3.2, we find that under our assumptions, the CE heuristic can achieve a regret that matches the fundamental lower bound up to a polylogarithmic factor in  $T$ , thus demonstrating its near-optimality.

## 4 Does Degeneracy Cause CE to Fail?

Fluid degeneracy is generally believed to cause the failure of CE in stochastic control. In this section, we challenge—and refine—this perspective within the OLP framework. Building on Theorem 3.1, we demonstrate a large class of instances that, despite violating standard non-degeneracy conditions in the literature, still achieve low regret. By closely examining the concept of degeneracy, we attribute this deviation from conventional understanding to a key distinction between how degeneracy manifests in discrete versus non-discrete contexts. This analysis enables us to systematically review existing notions of non-degeneracy and shed light on the critical geometric structures of the fluid problem that fundamentally drive the regret accumulation of CE.

### 4.1 Existing non-degeneracy conditions

Non-degeneracy conditions are imposed on the associated fluid problem of an OLP instance. More precisely, we specify these conditions with respect to a fluid instance  $(F, \mathbf{d})$ , where we recall that  $\mathbf{d} = \frac{\mathbf{b}}{T}$  is the normalized (initial) resource capacity.

**Assumption 4.1** (Primal stability condition). There exists an optimal  $\mathbf{x}_{\mathbf{d}}^*$  to primal problem (2), and a neighborhood  $\mathcal{N}(\mathbf{d})$  of  $\mathbf{d}$ , such that for any  $\mathbf{d}' \in \mathcal{N}(\mathbf{d})$ , there exists an optimal  $\mathbf{x}_{\mathbf{d}'}$  to the perturbed problem (2) with RHS constraint  $\mathbf{d}'$ , for which the set of binding resource constraints remains unchanged comparing to that of  $\mathbf{x}_{\mathbf{d}}^*$ .

**Assumption 4.2** (Strict complementary slackness condition).  $\boldsymbol{\lambda}^*$  is the unique solution to the dual problem (3)  $\min_{\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m} f_{\mathbf{d}}(\boldsymbol{\lambda})$ . Furthermore, there exists an optimal  $\mathbf{x}^*$  such that strict complementary slackness is satisfied. In particular, when  $F$  has no point mass, the condition is typically stated without involving  $\mathbf{x}^*$ : for any  $i = 1, \dots, m$ ,  $\lambda_i^* = 0$  if and only if  $d_i - \mathbb{E}_{(\mathbf{a}, r) \sim F} [a_i \mathbb{1}(r > \mathbf{a}^\top \boldsymbol{\lambda}^*)] > 0$ .

Assumptions 4.1 and 4.2 are two commonly imposed “non-degeneracy” conditions in prior work (cf. Assumption 2 and SC 8 of Balseiro, Besbes and Pizarro (2023), Assumption 2(c) of Li and Ye (2022), Assumption 5 of Bray (2024)), where it is known that Assumption 4.2, together with certain smoothness conditions on  $F$  implies Assumption 4.1 (cf. Lemma 3 of Balseiro, Besbes and Pizarro

(2023)). We note that the definition of Assumption 4.1 allows for potentially multiple primal optimal solutions<sup>2</sup>. Assumption 4.2 usually appears in its simplified form that only contains  $\lambda^*$ .

Assumptions 4.1 and 4.2 are critical to the low-regret analysis in the aforementioned prior work. More precisely, these non-degeneracy conditions ensure that the binding resource constraints in the fluid problems solved in each step of Algorithm 1 remain unchanged during the algorithm’s execution, thus permitting the martingale-based argument used in these prior work that ultimately leads to  $\mathcal{O}(\log T)$  regret. The only known performance guarantee of CE that does not rely on these non-degeneracy conditions is provided by Jiang et al. (2022a), where they instead have to impose a strong, uniform version of second-order growth conditions, for which we defer more discussion to Appendix E.

In addition to Assumptions 4.1 and 4.2, the following condition is often assumed implicitly.

**Assumption 4.3** (Dual uniqueness condition).  $\lambda^*$  is the unique solution to the dual problem (3).

**Remark 4.1.** Some authors define degeneracy as the violation of dual uniqueness Assumption 4.3. However, we emphasize that Assumption 4.3 and the common non-degeneracy conditions (Assumptions 4.1 and 4.2) are not equivalent in general; the latter are (in cases, strictly) stronger (cf. Lemma 4.4). In this work, we follow the convention in the line of literature (Li and Ye, 2022; Bray, 2024; Balseiro, Besbes and Pizarro, 2023; Jiang et al., 2022a), where “degeneracy” typically refers to the violation of standard non-degeneracy conditions (Assumptions 4.1 and 4.2).

## 4.2 The (near)optimality of CE beyond non-degeneracy

All aforementioned non-degeneracy conditions impose restrictions on both the underlying distribution  $F$  and the normalized resource capacity,  $\mathbf{d}$ , whereas Assumptions 2.1 and 2.2 specify distribution classes which purely rely on the properties of  $F$ . The following lemma explicitly characterizes the distinction between Assumptions 2.1 and 2.2 and the non-degeneracy conditions in the literature.

**Lemma 4.1.** *For any distribution  $F$  satisfying Assumption 2.1 or Assumption 2.2, there exists  $\mathbf{d} \in \mathbb{R}_{\geq 0}^m$  such that  $(F, \mathbf{d})$  violates the non-degeneracy conditions (Assumptions 4.1 and 4.2).*

Together with Theorem 3.1, we immediately conclude that non-degeneracy is irrelevant to the performance of CE.

**Corollary 4.2.** *Assumptions 4.1 and 4.2 are **not** necessary for CE to achieve  $o(\sqrt{T})$  regret.*

---

<sup>2</sup>Here we say two solutions  $\mathbf{x}$  and  $\mathbf{x}'$  are different in a probability sense. Namely,  $\mathbb{P}(\mathbf{x}(\mathbf{a}, r) \neq \mathbf{x}'(\mathbf{a}, r)) > 0$

We note that the type of degeneracy characterized in Lemma 4.1 is hardly a theoretical artifact; it often arises in practical applications. In scenarios where the inventory of resources is endogenously determined, the vector  $\mathbf{d}$  frequently becomes asymptotically close to  $\mathbb{E}_{(\mathbf{a}, r) \sim F} [\mathbf{a} \mathbb{1}(r > \mathbf{a}^\top \boldsymbol{\lambda}^*)]$  to avoid waste, typically within  $\mathcal{O}\left(\frac{1}{\sqrt{T}}\right)$ , as dictated by the square-root inventory law (Jiang et al., 2022a; Bumpensanti and Wang, 2020). Now, any redundant resource with zero dual price in the fluid limit would result in degeneracy. We defer a detailed proof to Appendix D.

Corollary 4.2 contradicts the general view and prior evidence that degeneracy causes CE to fail (cf. Figure 4 in Bumpensanti and Wang (2020)), and motivates a deeper investigation into the role that degeneracy plays in the regret accumulation of CE. In the next two sections, we present two complementary perspectives, one rooted in the discrete contexts and the other in the non-discrete contexts. Together, they establish a more comprehensive understanding of whether and how degeneracy impacts the performance of CE, resolving the paradox.

### 4.3 Fluid degeneracy: the discrete perspective

Consider a discrete  $F$  supported on  $n$  request types  $(\mathbf{a}^1, r^1), \dots, (\mathbf{a}^n, r^n)$ , with corresponding probabilities  $p_1, \dots, p_n$ . This is a setting that receives dominant attention in the NRM literature. Here, the primal fluid problem (2) becomes an LP (often referred to as the Deterministic Linear Program (DLP)). Concretely,

$$\begin{aligned} \max_{\mathbf{x}} \quad & \sum_{j=1}^n p_j r^j x_j \\ \text{s.t.} \quad & \sum_{j=1}^n \mathbf{a}^j p_j x_j \leq \mathbf{d}, \quad 0 \leq \mathbf{x} \leq 1. \end{aligned} \tag{5}$$

In the NRM literature, “non-degeneracy” traditionally refers to the (LP) non-degeneracy of the DLP (Jasin and Kumar, 2012). More precisely,

**Assumption 4.4** (DLP non-degeneracy). The DLP (5) has a non-degenerate optimal solution  $\mathbf{x}^*$ ,

$$|j \in [n] : x_j^* = 0 \text{ or } x_j^* = 1| + |i \in [m] : \sum_{j=1}^n p_j a_i^j x_j^* = d_i| = n. \tag{6}$$

Bumpensanti and Wang (2020) demonstrates both numerically and theoretically that violating Assumption 4.4 indeed results in performance deterioration of CE (cf. Figure 4, Propositions 2&3 in Bumpensanti and Wang (2020)), establishing the necessity of DLP non-degeneracy for guaranteeing CE’s performance in the discrete setting. Now one may naturally wonder about the connection




between Assumption 4.4 and the aforementioned conditions, Assumptions 4.1, 4.2 and 4.3. We establish such a connection in the next lemma.

**Lemma 4.3.** *Suppose  $F$  is a discrete distribution. Premised that the optimal solution  $\mathbf{x}^*$  to DLP (5) is unique, Assumptions 4.1, 4.2, 4.3 and 4.4 are equivalent.*

**Remark 4.2.** The premise of primal  $\mathbf{x}^*$  uniqueness is not the weakest under which the equivalence in Lemma 4.3 holds, but exploring it further is beyond the current scope.

We defer the proof to Appendix D and instead provide an intuitive explanation here. Lemma 4.3 demonstrates that in the discrete setting, there is essentially only one type of degeneracy: either all non-degeneracy conditions hold simultaneously and CE performs well, or the problem exhibits degeneracy, resulting in CE’s failure. Geometrically, this dichotomy arises from the rigid, piecewise-linear structure of the fluid problem in discrete settings, as is illustrated in Figure 1.



**Figure 1:** Function  $V_d^{\text{fluid}}$  and dual function  $f_d$  in degenerate and non-degenerate settings.

We plot  $V_d^{\text{fluid}} := \min_{\lambda \in \mathbb{R}_{\geq 0}^m} f_d(\lambda)$  as a function of  $\mathbf{d}$  in Figure 1a, and  $f_d(\lambda)$  as a function of  $\lambda$  with  $\mathbf{d} = d^1$  (degenerate case), and  $\mathbf{d} = d^2$  (non-degenerate case) in Figure 1b and Figure 1c, respectively. We also plot how perturbing  $\mathbf{d}$  around  $d^1$  or  $d^2$  affects functions  $f_d(\lambda)$ . As shown in Figure 1, fluid degeneracy manifests in several equivalent forms. In Figure 1a, the degenerate point  $d^1$  corresponds to a kink in the function  $V_d^{\text{fluid}}$ . By standard LP theory (cf. Bertsimas and Tsitsiklis (1997)), the supergradient of  $V_d^{\text{fluid}}$  at  $d^1$  are the optimal dual solutions to  $\min_{\lambda \in \mathbb{R}_{\geq 0}^m} f_{d^1}(\lambda)$ . The existence of multiple supergradients at  $d^1$  thereby corresponds to multiple dual solutions, as illustrated in Figure 1b. Furthermore, we observe in Figure 1b that perturbing  $\mathbf{d}$  around  $d^1$  results in drastic changes of the dual solution, demonstrating instability. In contrast, the non-degenerate case avoids all these issues. In Figure 1a, the function  $V_d^{\text{fluid}}$  is smooth at  $d^2$ , with a unique gradient.


In Figure 1c,  $f_{d^2}(\lambda)$  has a unique minimizer, and perturbing  $\mathbf{d}$  around  $d^2$  does not change the minimizer, demonstrating strong stability.

With Lemma 4.3, the necessity of DLP non-degeneracy (Assumption 4.4) for CE to achieve low regret simultaneously extends to all notions of non-degeneracy. Thus it is with minimal loss of accuracy to assert that “degeneracy causes CE to fail.”

#### 4.4 Fluid degeneracy: the non-discrete perspective

The clear and unified picture of degeneracy in discrete settings does not extend to non-discrete settings. The following result demonstrates how the equivalence between different notions of degeneracy breaks down.

**Lemma 4.4.** *Suppose non-discrete distribution  $F$  is such that function  $V_{\mathbf{d}}^{\text{fluid}}$  is smooth in  $\mathbf{d}$ , then Assumption 4.3 always holds regardless of  $\mathbf{d}$ . In contrast, whether Assumptions 4.1 and 4.2 hold or not depend on  $\mathbf{d}$ .*



**Figure 2:** Function  $V^{\text{fluid}}$  and dual function  $f_d$  for two different parameters in a non-discrete setting.

Figure 2 visually illustrates the intuition behind Lemma 4.4, the proof of which we omit. In Figure 2a, we plot the function  $V_{\mathbf{d}}^{\text{fluid}}$ . Unlike the discrete case shown in Figure 1a,  $V_{\mathbf{d}}^{\text{fluid}}$  here is smooth, with no kinks, and its gradient exists and is unique for any  $\mathbf{d}$ . As established in standard convex optimization theory (cf. Boyd and Vandenberghe (2004)), the gradient of  $V_{\mathbf{d}}^{\text{fluid}}$  corresponds to the dual optimal solution. Consequently, Assumption 4.3 is satisfied for all  $\mathbf{d}$ , as illustrated in Figures 2b and 2c.

On the other hand, in Figure 2b we plot the dual function, projected onto a dimension in the dual space, and illustrate the degenerate scenario where both Assumptions 4.1 and 4.2 fail. At  $d^1$ ,  $\lambda = 0$

is the unique dual price. However,  $\lambda = 0$  happens to be the global minimizer of the dual function  $f_{d^1}(\lambda)$ , indicating that the corresponding resource happens to be binding, thus violating the strict complementary slackness condition. Small perturbations around  $d^1$  lead the dual price to oscillate between zero and a positive value, causing the resource’s status to alternate between binding and non-binding, violating the primal stability condition. Hence, the non-degenerate conditions fail at  $d^1$ . Consequently, Assumptions 4.1 and 4.2 fundamentally rely on  $\mathbf{d}$  to avoid such degenerate cases. Overall, Lemma 4.4 essentially implies that Assumption 4.3 and Assumptions 4.1 and 4.2 can not be equivalent in non-discrete settings with smooth  $V_{\mathbf{d}}^{\text{fluid}}$ .

A key feature of the non-discrete setting is that, the premise in Lemma 4.4, namely that  $V_{\mathbf{d}}^{\text{fluid}}$  is smooth, is fairly common and satisfied by a wide class of distributions  $F$ , in particular, those defined by Assumptions 2.1 or 2.2.


**Lemma 4.5.** *For distribution  $F$  satisfying either Assumption 2.1 or Assumption 2.2, the corresponding dual objective  $f_{\mathbf{c}}(\boldsymbol{\lambda})$  is smooth for any fixed  $\mathbf{c}$ . In particular,  $\nabla f_{\mathbf{c}}(\boldsymbol{\lambda})$  exists. Furthermore,  $V_{\mathbf{c}}^{\text{fluid}}$  is smooth, with unique gradient  $\nabla V^{\text{fluid}}$  at any  $\mathbf{c}$ .*

The proof is in Appendix D. Combining Lemma 4.5 and Lemma 4.4, Assumption 4.3 always holds under Assumption 2.1 or Assumption 2.2, yet the non-degeneracy conditions (Assumptions 4.1 and 4.2) may fail (cf. Lemma 4.1). Our main contribution essentially lies in proving that in such cases, CE achieves uniformly near-optimal regret even if these non-degeneracy conditions fail (cf. Theorem 3.1). Along the line of this discussion, we may provide some geometric intuition of our performance guarantee. Observe in Figure 2b that the smoothness of the function  $f_{\mathbf{d}}(\boldsymbol{\lambda})$  ensures the dual price does not oscillate drastically when  $\mathbf{d}$  is perturbed around the degenerate point  $d^1$ . This inherent stability, even as the resource constraint alternates between binding and non-binding, allows CE to maintain good performance. This stands in sharp contrast to the discrete setting where across all distribution  $F$ , function  $V_{\mathbf{d}}^{\text{fluid}}$  is non-smooth, and degeneracy ( $\mathbf{d}$  taking certain corner values) always comes with drastic instability phenomenon, leading to performance deterioration of CE.

Meanwhile, not all non-discrete distributions yield smooth dual objectives as in Lemma 4.5. We provide two such examples that have appeared in prior work. Besbes et al. (2024) considered multisecretary instances where the reward distributions have gaps in their supports (cf. Example 3 in Besbes et al. (2024)). Jiang et al. (2022a) studied a semi-discrete NRM instances with finitely many request types, where conditional on any fixed type, the reward distribution is supported on an arbitrary interval with lower bounded density (cf. Assumption 1 Jiang et al. (2022a)). Both

examples have non-smooth  $V_d^{\text{fluid}}$ . At those  $\mathbf{d}$  that corresponds to a kink on  $V_d^{\text{fluid}}$ , Assumption 4.3 no longer holds. CE provably fails in these cases:

**Proposition 4.6.** *Consider a multisecretary instance with  $F_{\mathbf{a}}^r = F^r \triangleq \text{Unif}[0, 1] \cup [2, 3]$  and  $\mathbf{b} = \frac{1}{2}T$ . Then  $\text{REG}_{\mathbf{b}, T}(\text{CE}) \geq c_0\sqrt{T}$  for some absolute constant  $c_0 > 0$ .*



**Figure 3:** Function  $V_d^{\text{fluid}}$  and dual function  $f_d$  for multisecretary instances with gap.

Proposition 4.6 highlights the significance of dual uniqueness (Assumption 4.3) on the performance of CE. The failure of CE in Proposition 4.6 shares similarities with how CE suffers from degeneracy in discrete settings, as illustrated in Figure 3. For detailed proof, readers are referred to Proposition 1 of Besbes et al. (2024) or Proposition 2 in Bumpensanti and Wang (2020). Novel algorithmic designs beyond CE are required for such instances, such as the CwG principle (Besbes et al., 2024) or the boundary-attracted algorithm (Jiang et al., 2022a), to achieve improved ( $o(\sqrt{T})$ ) regret. In general, OLP instances violating Assumption 4.3 can be significantly more complex than the examples mentioned, to the extent that even describing such distributions may be challenging. Developing a unified algorithm and analysis to systematically handle these instances is beyond the scope of the current work. However, we hope the methodology introduced in this paper provides a foundation for addressing such challenges in future research.

The above discussion allows us to justify one specific requirement in our Assumption 2.1, that the conditional reward distributions are all supported on intervals starting from zero. In fact, without such a restriction, it is possible to construct simple OLP instances that violates Assumption 4.3 for which CE fails.

**Example 4.1.**  $\mathbb{P}(\mathbf{a} = 1) = \mathbb{P}(\mathbf{a} = 4) = \frac{1}{2}$ .  $r|\mathbf{a} \sim \text{Unif}[1, 2]$  for both values of  $\mathbf{a}$ . The initial inventory for the resource is  $\mathbf{b} = \mathbf{d}T = \frac{1}{2}T$ .

The distribution in Example 4.1 satisfies all the conditions in Assumption 2.1 with  $\beta = 0$ , except for the requirement (ii). A straightforward calculation yields multiple dual optimal solutions  $\boldsymbol{\lambda}^* \in [\frac{1}{2}, 1]$  that violates Assumption 4.3, and leads to  $\Omega(\sqrt{T})$  regret of CE. Note that the additional regularity imposed on  $F$  in Assumption 2.2 (ii) allows us to relax the condition that all conditional reward distribution should start from zero. However, we still require this condition to hold locally, to avoid tricky corner cases. We defer more discussion to Appendix E (cf. Example E.1).

The previous discussion provides a fairly comprehensive picture regarding how fluid degeneracy affects the performance of CE. Combining both the discrete and non-discrete perspectives, it seems more accurate to describe the structural challenge faced by CE as the “curse of dual non-uniqueness” rather than the broader and less precise “curse of degeneracy.”

## 5 Proof Sketch

We rely on the concentration analysis of the hindsight dual optimal solution in each time period  $t$  to conduct our performance analysis. In period  $t$ , the primal hindsight relaxation takes the form of a multi-knapsack LP similar to Problem (1). We consider its dual, which can be thought of as the empirical version of Problem (4). In particular,

$$\boldsymbol{\lambda}_t^* = \arg \min_{\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m} \mathbf{b}_{t-1}^\top \boldsymbol{\lambda} + \sum_{j=t+1}^T (r_j - \mathbf{a}_j^\top \boldsymbol{\lambda})^+, \quad (7)$$

$$\bar{\boldsymbol{\lambda}}_t^* = \arg \min_{\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m} (\mathbf{b}_{t-1} - \mathbf{a}_t)^\top \boldsymbol{\lambda} + \sum_{j=t+1}^T (r_j - \mathbf{a}_j^\top \boldsymbol{\lambda})^+, \quad (8)$$

where we note that  $\bar{\boldsymbol{\lambda}}_t^*$  is constructed only when feasibility is satisfied at period  $t$ , namely,  $\mathbf{b}_{t-1} \geq \mathbf{a}_t$  element-wise. Note that  $\boldsymbol{\lambda}_t^*$  and  $\bar{\boldsymbol{\lambda}}_t^*$  depend both on  $\mathbf{b}_{t-1}$  (and  $\mathbf{a}_t$ ) and  $\mathcal{I}_t$ . They are not non-anticipatory, and the introduction of them serves only for the purpose of analyzing the performance of algorithms. For notational simplicity, we write  $\boldsymbol{\lambda}_t^*$  and  $\bar{\boldsymbol{\lambda}}_t^*$  with the understanding that their dependence on  $\mathbf{b}_{t-1}$  ( $\mathbf{a}_t$ ) and  $\mathcal{I}_t$  is implicit but clear from the context.

**Lemma 5.1** (Regret Decomposition of CE). *The expected regret of the CE heuristic is bounded by*

$$\begin{aligned} \text{REG}_T(\pi) \leq & C_0 \log T + \sum_{t=1}^T \mathbb{E} \left[ \left( \mathbf{a}_t^\top \bar{\boldsymbol{\lambda}}_t^* - r_t \right) \mathbb{I}_{\{\mathbf{a}_t^\top \bar{\boldsymbol{\lambda}}_t^* \leq r_t \leq \mathbf{a}_t^\top \boldsymbol{\lambda}_t^*\}} \right] \\ & + \sum_{t=1}^T \mathbb{E} \left[ \left( r_t - \mathbf{a}_t^\top \boldsymbol{\lambda}_t^* \right) \mathbb{I}_{\{\mathbf{a}_t^\top \boldsymbol{\lambda}_t^* \leq r_t \leq \mathbf{a}_t^\top \bar{\boldsymbol{\lambda}}_t^*\}} \right], \quad (9) \end{aligned}$$

where  $\mathcal{C}_0 = 2 \left(1 + \frac{m\bar{A}}{\bar{A}}\right) m\bar{r}$ , and the expectation is taken over  $\mathcal{I}_0$ .

Lemma 5.1 follows from a careful decomposition analysis similar to Jiang et al. (2022a) within the *compensated coupling* framework of Vera and Banerjee (2021). It establishes a connection between the performance of CE and the statistical properties of  $\boldsymbol{\lambda}_t^*$  and  $\bar{\boldsymbol{\lambda}}_t^*$  around  $\tilde{\boldsymbol{\lambda}}_t$ , where we recall that  $\tilde{\boldsymbol{\lambda}}_t$  (cf. Problem 4) is the fluid dual optimal solution at time  $t$  that guides the CE decision-making. Similar observations were made earlier in the literature (c.f Li and Ye (2022); Jiang et al. (2022a) for discussions on the so-called *dual convergence*). We utilize techniques from empirical processes to establish desired dual convergence results, thus leading to the ultimate proof of Theorem 3.1.

**Lemma 5.2.** *Under either **Assumption 2.1** or **Assumption 2.2**, we have*

$$\begin{aligned} \mathbb{E}_{\mathbf{a} \sim F_{\mathbf{a}}} \left[ \left( F_{\mathbf{a}}(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right) \right] &\leq \mathcal{C}_4 \left( \frac{\log(T-t)}{T-t} \right)^{\frac{2+\beta}{2+2\beta}}, \\ \mathbb{E}_{\mathbf{a} \sim F_{\mathbf{a}}} \left[ \left( F_{\mathbf{a}}(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}(\mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^*) \right) \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^* \right) \right] &\leq \mathcal{C}_4 \left( \frac{\log(T-t)}{T-t} \right)^{\frac{2+\beta}{2+2\beta}}, \end{aligned}$$

both with probability at least  $1 - \frac{9C \log(T-t)}{(T-t)^2}$ , for  $T-t \geq e^{\frac{\tilde{C}_1}{32}+1}$ , where  $\mathcal{C}_4$  and  $\tilde{C}_1$  are constants (whose forms can be found in Remark B.1) that depend on model primitives regarding  $F$  but independent of  $T-t$  and  $\mathbf{b}$ , and  $C$  is a universal constant (cf. Lemma F.7).

The expectation on the LHS in Lemma 5.2 is taken with respect to  $\mathbf{a}$ . Recall that  $\boldsymbol{\lambda}_t^*$  and  $\bar{\boldsymbol{\lambda}}_t^*$  are both random variables, therefore the LHS in Lemma 5.2 are also random variables. Lemma 5.2 establishes a high-probability concentration bound of  $\boldsymbol{\lambda}_t^*, \bar{\boldsymbol{\lambda}}_t^*$  in a specific form. It is not hard to observe that the LHS of Lemma 5.2 serve as upper bounds on the terms in the regret decomposition of Lemma 5.1, explaining why we set up bounds in such a particular form.

Observe that in the prior works, strong non-degeneracy conditions are typically imposed to regulate the concentration of  $\boldsymbol{\lambda}_t^*$  and  $\bar{\boldsymbol{\lambda}}_t^*$ . Different from the standard approach, we avoid imposing regularity conditions directly on  $\tilde{\boldsymbol{\lambda}}_t$ . Rather, conditions on the problem primitives, i.e. Assumptions **Assumption 2.1** and **Assumption 2.2** suffice to guarantee nice concentration of  $\boldsymbol{\lambda}_t^*$  and  $\bar{\boldsymbol{\lambda}}_t^*$  for us. To achieve Lemma 5.2, we utilize the properties that  $\tilde{\boldsymbol{\lambda}}_t$  and  $\boldsymbol{\lambda}_t^*, \bar{\boldsymbol{\lambda}}_t^*$  are the minimizers of the corresponding dual fluid problem and the dual LP relaxations, respectively, where the latter can be viewed as the SAA of the former. With such a view, we transform bounding the concentration of  $\boldsymbol{\lambda}_t^*$  and  $\bar{\boldsymbol{\lambda}}_t^*$  around  $\tilde{\boldsymbol{\lambda}}_t$  into bounding the worst-case concentration of certain SAA of some target functionals. We achieve this worst-case concentration through the peeling device, borrowed from

the empirical processes theory, that uses union bounds coupled with a careful choice of space partition and with carefully calculated entropy number of the space.

Lemma 5.2 may be of independent theoretical interest and the techniques developed in this work are potentially applicable to the algorithmic analysis of a broad class of online stochastic optimization problems. We defer the proof of Lemma 5.1 to Appendix F.2, the proof of Lemma 5.2 to Appendix F.3, and the proof of Theorem 3.1 to Appendix B.

## 6 Concluding Remarks

In this work, we provided near-optimal regret guarantee of the classical CE algorithm for the general OLP problem, under mild assumptions on the underlying request distributions. Our result extends the state-of-the-art understanding of CE’s range of effectiveness, revealing that the commonly imposed non-degeneracy conditions are overly restrictive and not necessary for CE to achieve low regret. We developed new algorithmic analytical techniques based on empirical processes theory, potentially applicable to a broader range of dynamic optimization problems with non-discrete distributions. Our work leave open many interesting questions.

One direction is the design and analysis of a unified algorithm that achieves near optimal performance for all OLP instances. Example 4.1 demonstrates that our performance guarantee of CE does not extend beyond the distribution classes that we identify, and novel algorithmic innovations are needed. Algorithms with uniformly near-optimal performance have previously been designed in the discrete setting (cf. Vera and Banerjee (2021)) and in the multisecretary problem (cf. Besbes et al. (2024)). The simulation-based RAMS algorithm proposed in Besbes et al. (2024) is a potential candidate that attains uniform near-optimal performance across discrete and non-discrete settings, though the existing regret analysis of this algorithm still relies on some other reference algorithms (cf. Theorem 3 in Besbes et al. (2024)).

Another direction is to relax several assumptions made in the current work. For instance, one may consider scenarios where arriving requests are not *i.i.d.*, but with certain inter-dependent structure. Also, one may attempt to establish similar results in more general settings of dynamic resource allocation beyond OLP, such as the unified framework of  $DRC^2$  proposed in Balseiro, Besbes and Pizarro (2023).

## References

Agrawal, S., Wang, Z. and Ye, Y. (2014), ‘A dynamic near-optimal algorithm for online linear programming’, *Operations Research* **62**(4), 876–890.

- Alaei, S. (2014), ‘Bayesian combinatorial auctions: Expanding single buyer mechanisms to many buyers’, *SIAM Journal on Computing* **43**(2), 930–972.
- Arlotto, A. and Gurvich, I. (2019), ‘Uniformly bounded regret in the multisecretary problem’, *Stochastic Systems* **9**(3), 231–260.
- Balseiro, S. R., Besbes, O. and Pizarro, D. (2023), ‘Survey of dynamic resource-constrained reward collection problems: Unified model and analysis’, *Operations Research* .
- Balseiro, S. R., Besbes, O. and Weintraub, G. Y. (2015), ‘Repeated auctions with budgets in ad exchanges: Approximations and design’, *Management Science* **61**(4), 864–884.
- Balseiro, S. R., Ma, W. and Zhang, W. (2023), ‘Dynamic pricing for reusable resources: The power of two prices’, *arXiv preprint arXiv:2308.13822* .
- Bertsekas, D. (2012), *Dynamic programming and optimal control: Volume I*, Vol. 4, Athena scientific.
- Bertsimas, D. and Tsitsiklis, J. N. (1997), *Introduction to linear optimization*, Vol. 6, Athena Scientific Belmont, MA.
- Besbes, O., Kanoria, Y. and Kumar, A. (2024), ‘Dynamic resource allocation: Algorithmic design principles and spectrum of achievable performances’, *Operations Research* .
- Boyd, S. and Vandenberghe, L. (2004), *Convex optimization*, Cambridge university press.
- Bray, R. L. (2024), ‘Logarithmic regret in multisecretary and online linear programs with continuous valuations’, *Operations Research* .
- Bumpensanti, P. and Wang, H. (2020), ‘A re-solving heuristic with uniformly bounded loss for network revenue management’, *Management Science* **66**(7), 2993–3009.
- Chawla, S., Devanur, N. and Lykouris, T. (2023), ‘Static pricing for multi-unit prophet inequalities’, *Operations Research* .
- Chen, C., Qian, P. and Zhang, J. (2024), ‘Incentivizing participation in decentralized dynamic matching markets’, *Available at SSRN* .
- Devanur, N. R. and Hayes, T. P. (2009), The adwords problem: online keyword matching with budgeted bidders under random permutations, in ‘Proceedings of the 10th ACM conference on Electronic commerce’, pp. 71–78.
- Feldman, J., Henzinger, M., Korula, N., Mirrokni, V. S. and Stein, C. (2010), Online stochastic packing applied to display ad allocation, in ‘European Symposium on Algorithms’, Springer, pp. 182–194.
- Gallego, G. and Van Ryzin, G. (1994), ‘Optimal dynamic pricing of inventories with stochastic demand over finite horizons’, *Management science* **40**(8), 999–1020.
- Geer, S. A. (2000), *Empirical Processes in M-estimation*, Vol. 6, Cambridge university press.
- Gupta, V. (2024), ‘Greedy algorithm for multiway matching with bounded regret’, *Operations Research* **72**(3), 1139–1155.
- Hajiaghayi, M. T., Kleinberg, R. and Sandholm, T. (2007), Automated online mechanism design and prophet inequalities, in ‘AAAI’, Vol. 7, pp. 58–65.



- Huang, L. and Neely, M. J. (2009), Delay reduction via lagrange multipliers in stochastic network optimization, *in* ‘2009 7th International Symposium on Modeling and Optimization in Mobile, Ad Hoc, and Wireless Networks’, IEEE, pp. 1–10.
- Jasin, S. and Kumar, S. (2012), ‘A re-solving heuristic with bounded revenue loss for network revenue management with customer choice’, *Mathematics of Operations Research* **37**(2), 313–345.
- Jasin, S. and Sinha, A. (2015), ‘An lp-based correlated rounding scheme for multi-item ecommerce order fulfillment’, *Operations Research* **63**(6), 1336–1351.
- Jiang, J., Ma, W. and Zhang, J. (2022a), ‘Degeneracy is ok: Logarithmic regret for network revenue management with indiscrete distributions’, *arXiv preprint arXiv:2210.07996* .
- Jiang, J., Ma, W. and Zhang, J. (2022b), Tight guarantees for multi-unit prophet inequalities and online stochastic knapsack, *in* ‘Proceedings of the 2022 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)’, SIAM, pp. 1221–1246.
- Kerimov, S., Ashlagi, I. and Gurvich, I. (2024), ‘Dynamic matching: Characterizing and achieving constant regret’, *Management Science* **70**(5), 2799–2822.
- Kleinberg, R. D. (2005), A multiple-choice secretary algorithm with applications to online auctions., *in* ‘SODA’, Vol. 5, Citeseer, pp. 630–631.
- Li, X. and Ye, Y. (2022), ‘Online linear programming: Dual convergence, new algorithms, and regret bounds’, *Operations Research* **70**(5), 2948–2966.
- Lueker, G. S. (1998), ‘Average-case analysis of off-line and on-line knapsack problems’, *Journal of Algorithms* **29**(2), 277–305.
- Meanti, M., Rinnooy Kan, A., Stougie, L. and Vercellis, C. (1990), ‘A probabilistic analysis of the multiknapsack value function’, *Mathematical Programming* **46**, 237–247.
- Molinaro, M. and Ravi, R. (2014), ‘The geometry of online packing linear programs’, *Mathematics of Operations Research* **39**(1), 46–59.
- Reiman, M. I. and Wang, Q. (2008), ‘An asymptotically optimal policy for a quantity-based network revenue management problem’, *Mathematics of Operations Research* **33**(2), 257–282.
- Sierksma, G. (2001), *Linear and integer programming: theory and practice*, CRC Press.
- Talluri, K. T. and Van Ryzin, G. J. (2006), *The theory and practice of revenue management*, Vol. 68, Springer Science & Business Media.
- Vera, A. and Banerjee, S. (2021), ‘The bayesian prophet: A low-regret framework for online decision making’, *Management Science* **67**(3), 1368–1391.
- Wei, Y., Xu, J. and Yu, S. H. (2023), Constant regret primal-dual policy for multi-way dynamic matching, *in* ‘Abstract Proceedings of the 2023 ACM SIGMETRICS International Conference on Measurement and Modeling of Computer Systems’, pp. 79–80.
- Xie, X., Gurvich, I. and Küçükyavuz, S. (2024), ‘Dynamic allocation of reusable resources: Logarithmic regret in overloaded networks’, *Operations Research* .

## A Analysis of Example 2.3

We analyze Example 2.3 in this section. Recall that in this example, reward and resource consumption follow a generalized linear model:  $r = g(\mathbf{a}^\top \mathbf{z}) + \epsilon$ .

**Lemma A.1.** *Assumption 2.2 (i), (ii) and (iv) hold for Example 2.3 with  $\beta = 0$  and  $\eta = 1$ . Furthermore, a relaxed version of (iii) holds: there exist  $\mathbf{a}_o \in \text{supp}(F^{\mathbf{a}})$  and  $r_o > 0$  s.t.  $l(\mathbf{a}) \leq 0$  for all  $\mathbf{a} \in \mathcal{B}(\mathbf{a}_o, r_o) \cap \text{supp}(F^{\mathbf{a}})$ .*

*Proof.* That (i) holds follows from the conditions on  $F^{\mathbf{a}}$  in Example 2.3, and that both function  $g$  and random variable  $\epsilon$  is bounded. That (ii) holds follows from the conditions on  $F^{\mathbf{a}}$  in Example 2.3, and the Lipschitz continuity of  $g$ . (iv) holds because  $F_{\mathbf{a}}^r(z_2) - F_{\mathbf{a}}^r(z_1) = \mathbb{P}(z_1 < r \leq z_2 | \mathbf{a}) = \mathbb{P}(z_1 - g(\mathbf{a}^\top \mathbf{z}) < \epsilon \leq z_2 - g(\mathbf{a}^\top \mathbf{z}) | \mathbf{a})$ , which is bounded by  $\underline{f}_\epsilon(z_2 - z_1)$  and  $\bar{f}_\epsilon(z_2 - z_1)$ , where  $0 < \underline{f}_\epsilon \leq \bar{f}_\epsilon$  are the upper and lower bounds on the density of  $\epsilon$ , as is assumed in Example 2.3. Finally, we argue that the relaxed version of (iii) holds. Recall that  $g((\mathbf{a}')^\top \mathbf{z}) \leq \mathcal{L} - \eta$ . Thus the support of  $r | \mathbf{a}'$  is an interval  $[-\eta, M]$ , where  $M < \infty$  since  $g$  is Lipschitz,  $\mathbf{a}$  is bounded and  $\eta$  is bounded. Namely,  $l(\mathbf{a}') = -\eta < 0$ . (iii) thus holds with a ball centered at  $\mathbf{a}'$ , where we use the Lipschitzness of  $l(\mathbf{a})$ . Q.E.D.

We next argue that the possible negative rewards can be ignored without affecting the regret. Indeed, the hindsight optimal decision when facing a request with negative reward is to reject it. Hence, any OLP instance with the underlying distribution satisfying Lemma A.1 can be WLOG restricted to the part with non-negative rewards. For that restricted probability distribution, Lemma A.1 implies that Assumption 2.2 holds.

## B Proof of Theorem 3.1

This section is devoted to the proof of Theorem 3.1.

With the help of Lemma 5.2, the proof of Theorem 3.1 is straightforward.

*Proof of Theorem 3.1.* In fact, by Lemma 5.1,

$$\begin{aligned} \text{REG}_{\mathbf{b}, T}(\pi^{\text{CE}}) &\leq \mathcal{C}_0 \log T + \sum_{t=1}^T \mathbb{E} \left[ \left( \mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t^* - r_t \right) \mathbb{I}_{\{\mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t \leq r_t \leq \mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t^*\}} \right] \\ &\quad + \sum_{t=1}^T \mathbb{E} \left[ \left( r_t - \mathbf{a}_t^\top \boldsymbol{\lambda}_t^* \right) \mathbb{I}_{\{\mathbf{a}_t^\top \boldsymbol{\lambda}_t^* \leq r_t \leq \mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t\}} \right] \\ &\leq \mathcal{C}_0 \log T + \sum_{t=1}^T \mathbb{E} \left[ \left( \mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t^* - \mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t \right) \mathbb{I}_{\{\mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t \leq r_t \leq \mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t^*\}} \right] \\ &\quad + \sum_{t=1}^T \mathbb{E} \left[ \left( \mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}_t^\top \boldsymbol{\lambda}_t^* \right) \mathbb{I}_{\{\mathbf{a}_t^\top \boldsymbol{\lambda}_t^* \leq r_t \leq \mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t\}} \right]. \end{aligned}$$

Observe that the above RHS takes the form of the LHS in Lemma 5.2, namely,

$$\begin{aligned} &\mathcal{C}_0 \log T + \sum_{t=1}^T \mathbb{E} \left[ \left( \mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t^* - \mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t \right) \mathbb{I}_{\{\mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t \leq r_t \leq \mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t^*\}} \right] + \sum_{t=1}^T \mathbb{E} \left[ \left( \mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}_t^\top \boldsymbol{\lambda}_t^* \right) \mathbb{I}_{\{\mathbf{a}_t^\top \boldsymbol{\lambda}_t^* \leq r_t \leq \mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t\}} \right] \\ &= \mathcal{C}_0 \log T + \sum_{t=1}^T \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right) \right] \end{aligned}$$

$$+ \sum_{t=1}^T \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^*) \right) \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^* \right) \right]. \quad (10)$$

Note that

$$\begin{aligned} & \sum_{t=1}^T \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^*) \right) \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^* \right) \right] \\ & \leq \sum_{t=1}^{(T-\mathcal{C}_5)_+} \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^*) \right) \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^* \right) \mathbb{I}_{\mathcal{A}_t} \right] \\ & \quad + \sum_{t=1}^{(T-\mathcal{C}_5)_+} \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^*) \right) \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^* \right) \mathbb{I}_{\mathcal{A}_t^c} \right] \\ & \quad + \sum_{t=(T-\mathcal{C}_5)_++1}^T \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^*) \right) \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^* \right) \mathbb{I}_{\{T > \mathcal{C}_5\}} \right] \\ & \leq \tilde{\mathcal{C}}_4 \sum_{t=1}^{(T-\mathcal{C}_5)_+} \left( \frac{\log(T-t)}{T-t} \right)^{\frac{2+\beta}{2+2\beta}} + \sum_{t=1}^{(T-\mathcal{C}_5)_+} \mathbb{P}(\mathcal{A}_t^c) + \mathcal{C}_5 \frac{\bar{A}}{\underline{A}} m\bar{r} \\ & \leq \tilde{\mathcal{C}}_4 \sum_{t=1}^{(T-\mathcal{C}_5)_+} \left( \frac{\log(T-t)}{T-t} \right)^{\frac{2+\beta}{2+2\beta}} + \sum_{t=1}^{(T-\mathcal{C}_5)_+} \frac{9\mathcal{C} \log(T-t) \frac{\bar{A}}{\underline{A}} m\bar{r}}{(T-t)^2} + \mathcal{C}_5 \frac{\bar{A}}{\underline{A}} m\bar{r}, \end{aligned} \quad (11)$$

where  $\mathcal{C}_5 = e^{\frac{\tilde{\mathcal{C}}_1}{32} + 1}$ , and  $\mathcal{A}_t$  denotes the event that the first inequality in Lemma 5.2 occurs. In (11), the first term is by the high probability concentration bound of Lemma 5.2, the second term is by the small probability that  $\mathcal{A}_t^c$  occurs, and the last term is by  $\|\boldsymbol{\lambda}_t^*\|, \|\tilde{\boldsymbol{\lambda}}_t\|$  are both upper bounded. Similarly,

$$\begin{aligned} & \sum_{t=1}^T \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^*) \right) \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^* \right) \right] \\ & \leq \tilde{\mathcal{C}}_4 \sum_{t=1}^{(T-\mathcal{C}_5)_+} \left( \frac{\log(T-t)}{T-t} \right)^{\frac{2+\beta}{2+2\beta}} + \sum_{t=1}^{(T-\mathcal{C}_5)_+} \frac{9\mathcal{C} \log(T-t) \frac{\bar{A}}{\underline{A}} m\bar{r}}{(T-t)^2} + \mathcal{C}_5 \frac{\bar{A}}{\underline{A}} m\bar{r}. \end{aligned} \quad (12)$$

Observe that

$$\int_1^T \left( \frac{\log t}{t} \right)^\alpha dt \leq (\log T)^\alpha \int_1^T t^{-\alpha} dt \leq \begin{cases} (\log T)^2 & \alpha = 1, \\ \frac{1}{1-\alpha} (\log T)^\alpha T^{1-\alpha} & 0 < \alpha < 1. \end{cases}$$

Thus, by the basic inequality  $\log x \leq x$ , combining (10), (11), and (12) gives us

$$\text{REG}_{\mathbf{b}, T}(\pi^{\text{CE}}) \leq \begin{cases} \mathcal{C} (\log T)^2 & \beta = 0, \\ \tilde{\mathcal{C}} T^{\frac{1}{2} - \frac{1}{2(1+\beta)}} (\log T)^{\frac{2+\beta}{2+2\beta}} & \beta > 0, \end{cases}$$

where  $\mathcal{C} = \mathcal{C}_0 + 4\tilde{\mathcal{C}}_4 + 36C \frac{\bar{A}}{\underline{A}} m\bar{r} + 2\mathcal{C}_5 \frac{\bar{A}}{\underline{A}} m\bar{r}$ , and  $\tilde{\mathcal{C}} = \mathcal{C}_0 + \frac{8+8\beta}{\beta} \tilde{\mathcal{C}}_4 + 36C \frac{\bar{A}}{\underline{A}} m\bar{r} + 2\mathcal{C}_5 \frac{\bar{A}}{\underline{A}} m\bar{r}$ . This finishes the proof of Theorem 3.1, where the explicit form of constants  $\mathcal{C}$  and  $\tilde{\mathcal{C}}$  are provided in the next remark. Q.E.D.

**Remark B.1.** From the proof of the Theorem and the lemmas, we can specify the concrete constants

appearing in the regret bound. The constants  $\mathcal{C}$  and  $\tilde{\mathcal{C}}$  in Theorem 3.1 have the specific form of

$$\begin{aligned}\mathcal{C} &= \mathcal{C}_0 + 4\mathcal{C}_4 + 36C \frac{\bar{A}}{\underline{A}} m\bar{r} + 2\mathcal{C}_5 \frac{\bar{A}}{\underline{A}} m\bar{r}, \\ \tilde{\mathcal{C}} &= \mathcal{C}_0 + \frac{8 + 8\beta}{\beta} \mathcal{C}_4 + 36C \frac{\bar{A}}{\underline{A}} m\bar{r} + 2\mathcal{C}_5 \frac{\bar{A}}{\underline{A}} m\bar{r}.\end{aligned}$$

Recall from Lemma F.7,  $C$  is an absolute constant. From the proof of Lemma 5.1 Lemma F.2, we have

$$\begin{aligned}\mathcal{C}_0 &= 2 \left( 1 + \frac{m\bar{A}}{\underline{A}} \right) m\bar{r}, \quad \mathcal{C}_5 = e^{\frac{\tilde{\mathcal{C}}_1}{32} + 1}, \quad \tilde{\mathcal{C}}_3 = 3\tilde{\mathcal{C}}_1 + \frac{4m\bar{A}\bar{r}}{\underline{A}} + \frac{2m^2\bar{A}\bar{r}}{\underline{A}}, \\ \tilde{\mathcal{C}}_1 &= 2 \left( \frac{4m\bar{r}\bar{A}}{\underline{A}} + 1 \right) \left( 10C^2 \left( 1 + 2 \left( \frac{10m}{\min\{\nu/2, 1\}} \right)^{1/2} \right)^2 \left( 4c_\nu^{\frac{1}{2}} \frac{\bar{r}}{\underline{A}} (2m\bar{A})^{\frac{\nu}{2}} + 9 \right)^2 + 2 \right).\end{aligned}$$

$\mathcal{C}_4$  is specified from the proof of Lemma 5.2. Particularly, under Assumption 2.1,

$$\mathcal{C}_4 = \tilde{\mathcal{C}}_3 \left( 1 + \max \left( \left( 2m \frac{\bar{r}\bar{A}}{\underline{A}} \right)^{\frac{1+\beta}{2+\beta}}, c_\beta^{-\frac{1}{2+\beta}} \right) \right)^{\frac{2+\beta}{1+\beta}}.$$

Under Assumption 2.2,  $\mathcal{C}_4 = \max(c_1, c_2, c_3, c_4)$  where

$$\begin{aligned}c_1 &= \tilde{\mathcal{C}}_3 + \tilde{\mathcal{C}}_3^2 \bar{A} \left( \min \left( c_\beta, 4^{-1-\beta} c_\beta c_\nu^{\frac{1+\beta}{\nu}} \left( 2\sqrt{m} \frac{\bar{r}\bar{A}}{\underline{A}} \right)^{-1-\beta} \right) C_f^{\frac{2+\beta}{2}} \left( \frac{r_1 \wedge w}{6} \sin(\theta) \right)^{\frac{(2+\beta)(m+2)}{2}} \right)^{-1} \max(1, \bar{A}), \\ c_2 &= \tilde{\mathcal{C}}_3 + \tilde{\mathcal{C}}_3 \left( \frac{2\bar{A}\sqrt{m}\bar{r}}{\underline{A}} \max \left( \tilde{\mathcal{C}}_3 4^{1+\beta} c_\beta^{-1} c_\nu^{-\frac{1+\beta}{\nu}}, \tilde{\mathcal{C}}_3^2 \frac{2\bar{A}\sqrt{m}\bar{r}}{\underline{A}} 4^{2+2\beta} c_\beta^{-2} c_\nu^{-\frac{2(1+\beta)}{\nu}} \right) \right)^{1/2}, \\ c_3 &= \tilde{\mathcal{C}}_3 \left( 1 + \max \left( \left( 2m \frac{\bar{r}\bar{A}}{\underline{A}} \right)^{\frac{1+\beta}{2+\beta}}, c_\beta^{-\frac{1}{2+\beta}} \right) \right)^{\frac{2+\beta}{1+\beta}}, \\ c_4 &= \tilde{\mathcal{C}}_3 + \tilde{\mathcal{C}}_3^2 \bar{A} \left( c_\beta C_f^{\frac{2+\beta}{2}} \left( \frac{r_0 \wedge w}{6} \sin(\theta) \right)^{\frac{(2+\beta)(m+2)}{2}} \right)^{-1} \max(1, \bar{A}),\end{aligned}$$

and  $\theta, w$  are the parameters specified by the uniform cone condition (cf. Lemma F.15), and

$$C_f = l_f \frac{\pi^{m/2}}{\Gamma(\frac{m}{2} + 1)}, \quad r_1 = \frac{1}{8} \left( \left( c_L + \sqrt{m} \frac{\bar{r}}{\underline{A}} \right)^{-1} \wedge 1 \right) c_\nu^{\frac{1}{\nu}},$$

with  $\Gamma(\cdot)$  denoting the Gamma function.

## C Proof of Proposition 3.2

*Proof of Proposition 3.2.* The multisecretary instance presented in Example 2.1 achieves the fundamental lower bound. Recall that the instances have reward *p.d.f.*:

$$h_r(x) = \begin{cases} (1 + \beta)(1 - 2x)^\beta, & 0 \leq x \leq \frac{1}{2}, \\ (1 + \beta)(2x - 1)^\beta, & \frac{1}{2} \leq x \leq 1. \end{cases}$$

For a fixed time horizon  $T > 0$ , we set the initial resource capacity at  $\lfloor \frac{T}{2} \rfloor$ . We observe that this is an instance considered in Besbes et al. (2024), only that  $g = 0$ , namely there is no gap in our case. The proof follows directly from Theorem 1 in Besbes et al. (2024). Q.E.D.

## D Proof in Section 4

### D.1 Proof of Lemma 4.1

The proof is divided into two parts. We first provide a proof of Lemma 4.5, then use this lemma to show the desired results.

*Proof of Lemma 4.5.* Direct computation shows that

$$f_{\mathbf{c}}(\boldsymbol{\lambda}) = \mathbf{c}^\top \boldsymbol{\lambda} - \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \mathbf{a}^\top \boldsymbol{\lambda} + \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \int_0^{\mathbf{a}^\top \boldsymbol{\lambda}} F_{\mathbf{a}}^r(v) dv. \quad (13)$$

Both Assumptions 2.1 and Assumption 2.2 imply that  $F_{\mathbf{a}}^r(v)$  is continuous with respect to  $v$ . Hence, by the Leibniz integral rule,  $\nabla f_{\mathbf{c}}(\boldsymbol{\lambda})$  exists. The statement about  $V_{\mathbf{c}}^{\text{fluid}}$  is obvious by the expression (13) and the fact that  $F_{\mathbf{a}}^r(v)$  is continuous. Q.E.D.

With Lemma 4.5, we proceed with the proof of Lemma 4.1.

*Proof of Lemma 4.1.* By Lemma 4.5,  $\nabla f_{\mathbf{c}}$  always exists for any  $\mathbf{c}$ . We now show that for any nonzero  $\boldsymbol{\lambda}^\circ \in \mathbb{R}_{\geq 0}^m$  satisfying  $\lambda_i^\circ = 0$  for some  $1 \leq i \leq m$ , there exists  $\mathbf{d}$  such that  $\boldsymbol{\lambda}^\circ$  is the optimal solution to the dual problem  $\min_{\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m} f_{\mathbf{d}}(\boldsymbol{\lambda})$ . Without loss of generality, let  $i = 1$ , and set  $\lambda_i^\circ > 0$  for  $i = 2, \dots, m$ . Such a  $\mathbf{d}$  can be constructed via  $\nabla f_{\mathbf{d}}(\boldsymbol{\lambda}^\circ) = \mathbf{0}$ , or more precisely  $\mathbf{d} = \mathbb{E}_{(\mathbf{a}, r) \sim F} [\mathbf{a} \mathbb{1}(r > \mathbf{a}^\top \boldsymbol{\lambda}^\circ)]$ . The resulting instance violates Assumption 4.2 since  $\lambda_1^\circ = 0$  and  $d_1 = \mathbb{E}_{(\mathbf{a}, r) \sim F} [a_1 \mathbb{1}(r > \mathbf{a}^\top \boldsymbol{\lambda}^\circ)]$ . Consider two perturbation  $\mathbf{d}' = (d_1 + \epsilon, d_2, \dots, d_m)$  and  $\mathbf{d}'' = (d_1 - \epsilon, d_2, \dots, d_m)$  of  $\mathbf{d} = (d_1, d_2, \dots, d_m)$ , where  $\epsilon > 0$ , and the optimal solutions for  $\min_{\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m} f_{\mathbf{d}'}(\boldsymbol{\lambda})$  and  $\min_{\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m} f_{\mathbf{d}''}(\boldsymbol{\lambda})$  are  $\boldsymbol{\lambda}'$  and  $\boldsymbol{\lambda}''$ , respectively. It is clear that  $\lambda_1'$  is zero while  $d_1 + \epsilon > \mathbb{E}_{(\mathbf{a}, r) \sim F} [a_1 \mathbb{1}(r > \mathbf{a}^\top \boldsymbol{\lambda}')] = d_1$ , and  $\lambda_1'' > 0$  while  $d_1 - \epsilon = \mathbb{E}_{(\mathbf{a}, r) \sim F} [a_1 \mathbb{1}(r > \mathbf{a}^\top \boldsymbol{\lambda}'')] = d_1$ . Hence, it violates Assumption 4.1. Notably, Example 2 in Jiang et al. (2022a) is a specific instance within this degenerate class. Q.E.D.

### D.2 DLP and the proof of Lemma 4.3

We write down the dual fluid problem in its LP form:

$$\begin{aligned} \min_{\boldsymbol{\lambda}} \quad & \sum_{i=1}^m d_i \lambda_i + \sum_{j=1}^n \eta_j \\ \text{s.t.} \quad & \sum_{i=1}^m a_i^j p^j \lambda_i + \eta_j \geq p^j r^j, \quad j \in [1, n], \\ & \boldsymbol{\lambda}, \boldsymbol{\eta} \geq 0. \end{aligned} \quad (14)$$

*Proof of Lemma 4.3.* (i) Assumption 4.3  $\iff$  Assumption 4.2. Since Assumption 4.2 requires dual uniqueness, it suffices to show dual uniqueness implies Assumption 4.2, namely, strict complementary slackness. By standard LP theory (cf. Exercise 4.20 in Bertsimas and Tsitsiklis (1997)), there exists a pair of primal and dual optimal solution that the strict complementary slackness condition is satisfied. Premised on  $\mathbf{x}^*$  is the unique solution to DLP (5), both primal and dual have unique optimal solution and therefore they must satisfy strict complementary slackness.

(ii) *Assumption 4.3*  $\iff$  *Assumption 4.4*. We first show *Assumption 4.3*  $\Leftarrow$  *Assumption 4.4*. Premised on primal uniqueness and under *Assumption 4.4*, we have  $\mathbf{x}^*$  is a unique, non-degenerate optimal solution to the DLP (5). Standard LP theory thus implies the dual problem (14) also has a non-degenerate and unique optimal solution  $(\boldsymbol{\lambda}^*, \boldsymbol{\eta}^*)$ , or equivalently, a unique solution  $\boldsymbol{\lambda}^*$  to problem (3) and *Assumption 4.3* is true. We now in turn prove *Assumption 4.3*  $\implies$  *Assumption 4.4*. We prove by contradiction. Assume  $\mathbf{x}^*$  is degenerate. Then by standard LP theory (cf. Theorem 4.5 in Sierksma (2001)), since both DLP (5) and the dual problem (14) are in standard (inequality) form, the dual must have multiple optimal solutions. Since  $\boldsymbol{\eta}^*$  is uniquely determined by  $\eta_j = p^j \left( r^j - \sum_{i=1}^m a_i^j \lambda_i \right)^+$  for  $j = 1, \dots, n$ , the non-uniqueness of the dual solution thus must imply the non-uniqueness of optimal  $\boldsymbol{\lambda}^*$  to the problem (3), completing the proof.

(iii) *Assumption 4.4*  $\implies$  *Assumption 4.1*. In this case,  $\mathbf{x}^*$  is the unique and non-degenerate optimal solution to DLP (5). Since the DLP has a bounded feasible region,  $\mathbf{x}^*$  must be a unique, non-degenerate optimal basic feasible solution (BFS). The primal stability in this case is immediate.

(iv) *Assumption 4.1*  $\implies$  *Assumption 4.3*. We argue by contraction. If the dual has multiple optimal solutions, then by standard LP theory (cf. Theorem 4.5 in Sierksma (2001)), the primal DLP must have a degenerate solution. Since we further assume  $\mathbf{x}^*$  is unique, it must be a BFS since the DLP has a bounded feasible region. Thus,  $\mathbf{x}^*$  is a unique, degenerate optimal BFS. There must be more than  $n$  binding constraints at  $\mathbf{x}^*$ , and  $n$  of them are linearly independent. For any neighborhood of  $\mathbf{d}$ , there always exists  $\mathbf{d}'$  in the neighborhood such that the binding constraints at  $\mathbf{x}^*$  cannot be binding simultaneously. Therefore, any optimal solution to the DLP with RHS  $\mathbf{d}'$  can not share the same set of binding resources as  $\mathbf{x}^*$ , violating *Assumption 4.1*, thus completing the proof.

Combining the above completes the proof. Q.E.D.

## E Second-Order Growth Conditions

In addition to non-degeneracy, another class of fluid regularity conditions frequently imposed in prior literature is *second-order growth* (on dual objectives). A fixed OLP instance specified by  $(F, \mathbf{b}, T)$  corresponds to a fluid instance  $(F, \mathbf{d})$ . The second-order growth conditions characterize the curvature of the function  $f_{\mathbf{d}}(\boldsymbol{\lambda})$ , as is determined by  $F$  and  $\mathbf{d}$ . Noticeably they are specific to the non-discrete settings, since in the discrete case,  $f_{\mathbf{d}}(\boldsymbol{\lambda})$  is piecewise linear, and the notion of curvature is trivial. Second-order growth conditions in the literature can be broadly classified into two categories: local and global conditions. Building on Theorem 3.1, we demonstrate that local conditions capture key structural features of the problem that fundamentally determine the achievable regret scaling of CE (and, in fact, any algorithm). By contrast, global conditions, often introduced for technical analytical purposes, are typically overly restrictive and not reflective of the actual determinants of algorithmic performance.

### E.1 Local conditions

Various forms of (local) second-order growth conditions exist in the literature. Here, we present a typical one. For ease of connecting to our own assumptions, we state it in a more general form that incorporates high-order growth. Fix a fluid instance  $(F, \mathbf{d})$  such that  $\mathbf{d} \neq \mathbf{0}$ .

**Assumption E.1** (Local higher-order growth condition). There exist a constant  $\gamma \geq 0$  and a neighborhood  $\mathcal{N}$  of the dual optimal solution  $\boldsymbol{\lambda}^*$  to  $\min_{\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m} f_{\mathbf{d}}(\boldsymbol{\lambda})$ , and positive constant  $\underline{\alpha}$ , such that for any  $\boldsymbol{\lambda} \in \mathcal{N}$ , it holds that

$$f_{\mathbf{d}}(\boldsymbol{\lambda}) - f_{\mathbf{d}}(\boldsymbol{\lambda}^*) - \nabla f_{\mathbf{d}}(\boldsymbol{\lambda}^*)^\top (\boldsymbol{\lambda} - \boldsymbol{\lambda}^*) \geq \underline{\alpha} |\boldsymbol{\lambda} - \boldsymbol{\lambda}^*|^{2+\gamma}.$$

Typically,  $\gamma = 0$  corresponds to the standard second-order growth condition.

Assumption E.1, with  $\gamma = 0$ , appears in Balseiro, Besbes and Pizarro (2023) (cf. SC 7). We note that other notions of (local) second-order growth conditions exist. For instance, Li and Ye (2022) imposes a local Lipschitz continuity condition on the conditional reward CDF (Assumption 2(b) in Li and Ye (2022)), Bray (2024) imposes a positive definiteness condition on the Hessian matrix of  $f_{\mathbf{d}}(\boldsymbol{\lambda})$  at  $\boldsymbol{\lambda}^*$  (cf. Assumption 6), while Balseiro, Besbes and Pizarro (2023) also introduces a lower downward quadratic condition on  $V_{\mathbf{d}}^{\text{fluid}}$  (cf. Assumption 2.1). For a comprehensive overview of these assumptions and their interrelations, we refer readers to Appendix A of Jiang et al. (2022a). The existing regret guarantee of an OLP instance  $(F, \mathbf{b}, T)$  typically requires the corresponding fluid instance  $(F, \mathbf{d})$  to satisfy both the second-order growth condition, **and** the non-degenerate condition (cf. Assumption 4.1 or 4.2).

We state here an observation that Assumption E.1 implies Assumption 4.3. In particular,

**Proposition E.1.** *Under the second-order growth condition of Assumption E.1 the dual solution is unique. Namely, Assumption 4.3 holds.*

*Proof.* Suppose there are two dual solutions  $\boldsymbol{\lambda}_1$  and  $\boldsymbol{\lambda}_2$  with  $\boldsymbol{\lambda}_1 \neq \boldsymbol{\lambda}_2$ . For any  $\boldsymbol{\lambda}_\theta = \theta\boldsymbol{\lambda}_1 + (1-\theta)\boldsymbol{\lambda}_2$ , by the convexity of  $f_{\mathbf{d}}(\boldsymbol{\lambda})$ , we have

$$f_{\mathbf{d}}(\boldsymbol{\lambda}_\theta) \leq \theta f_{\mathbf{d}}(\boldsymbol{\lambda}_1) + (1-\theta)f_{\mathbf{d}}(\boldsymbol{\lambda}_2),$$

which implies  $\boldsymbol{\lambda}_\theta$  is also a dual solution for any  $\theta \in [0, 1]$ . Hence, we can take  $\boldsymbol{\lambda}_\theta$  in the neighborhood of  $\boldsymbol{\lambda}_2$  stated in Assumption E.1 and  $\boldsymbol{\lambda}_\theta \neq \boldsymbol{\lambda}_2$ . By Assumption E.1, we have

$$f_{\mathbf{d}}(\boldsymbol{\lambda}_2) - f_{\mathbf{d}}(\boldsymbol{\lambda}_{\frac{\theta}{2}}) - \nabla f_{\mathbf{d}}(\boldsymbol{\lambda}_{\frac{\theta}{2}})^\top \left( \boldsymbol{\lambda}_2 - \boldsymbol{\lambda}_{\frac{\theta}{2}} \right) \geq \underline{\alpha} \left| \boldsymbol{\lambda}_2 - \boldsymbol{\lambda}_{\frac{\theta}{2}} \right|^{2+\gamma},$$

which implies

$$-\frac{1}{2} \nabla f_{\mathbf{d}}(\boldsymbol{\lambda}_{\frac{\theta}{2}})^\top (\boldsymbol{\lambda}_2 - \boldsymbol{\lambda}_\theta) \geq \underline{\alpha} \left| \frac{1}{2} (\boldsymbol{\lambda}_2 - \boldsymbol{\lambda}_\theta) \right|^{2+\gamma}.$$

Similarly, it can be seen that

$$f_{\mathbf{d}}(\boldsymbol{\lambda}_\theta) - f_{\mathbf{d}}(\boldsymbol{\lambda}_{\frac{\theta}{2}}) - \nabla f_{\mathbf{d}}(\boldsymbol{\lambda}_{\frac{\theta}{2}})^\top \left( \boldsymbol{\lambda}_\theta - \boldsymbol{\lambda}_{\frac{\theta}{2}} \right) \geq \underline{\alpha} \left| \boldsymbol{\lambda}_\theta - \boldsymbol{\lambda}_{\frac{\theta}{2}} \right|^{2+\gamma},$$

which implies

$$\frac{1}{2} \nabla f_{\mathbf{d}}(\boldsymbol{\lambda}_{\frac{\theta}{2}})^\top (\boldsymbol{\lambda}_2 - \boldsymbol{\lambda}_\theta) \geq \underline{\alpha} \left| \frac{1}{2} (\boldsymbol{\lambda}_\theta - \boldsymbol{\lambda}_2) \right|^{2+\gamma}.$$

Hence, we must have  $\boldsymbol{\lambda}_\theta = \boldsymbol{\lambda}_2$ , which contradicts the condition  $\boldsymbol{\lambda}_\theta \neq \boldsymbol{\lambda}_2$ , and finish the proof. Q.E.D.

Proposition E.1 shows that Assumption 4.3 is in fact implicitly assumed under the sufficient conditions typically made in the literature. In addition to forcing  $\boldsymbol{\lambda}^*$  to be a unique solution to  $\min_{\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m} f_{\mathbf{d}}(\boldsymbol{\lambda})$ , Assumption E.1 imposes curvature requirements of  $f_{\mathbf{d}}(\boldsymbol{\lambda})$  at  $\boldsymbol{\lambda}^*$ , captured by parameter  $\gamma$ . This parameter  $\gamma$  appears in the fundamental regret lower bound and determines the best achievable regret scaling (cf. Proposition 3.2 and Theorem 1 in Besbes et al. (2024)). In that sense, Assumption E.1 is an essential condition. In our Assumptions 2.1 and 2.2, we impose a reverse Hölder condition on the distributions with parameter  $\beta$ , which effectively captures the  $\gamma$  as appeared in Assumption E.1.

Recall that our Assumptions 2.1 and 2.2 rely purely on the properties of the underlying  $F$  and are independent of  $\mathbf{d}$ . This offers several advantages over directly imposing Assumption E.1. Beyond being more intuitive and easy-to-check, one notable benefit of our assumptions is their ability to handle cases involving scarce resources. In fact, the subtle scaling regime of a sequence

of OLP instances  $(F, \mathbf{b}_T, T)$  such that  $\mathbf{b}_T \propto T^\eta$  for  $\eta \in (0, 1)$  cannot be simply described by its “fluid” instance, in which  $\mathbf{d}$  approaches  $\mathbf{0}$ . In such cases, Assumption E.1 is no longer suitable. In contrast, our assumptions uniformly cover these regimes, for which the regret guarantee in Theorem 3.1 remains valid.

Through the lens of Assumption E.1, we can justify the requirement in our Assumption 2.2, in particular, that there exists  $\mathbf{a}_o$  such that the conditional reward distributions in a neighborhood of  $\mathbf{a}_o$  are all supported on intervals starting from zero. In fact, without such a restriction, it is possible to construct simple OLP instances with  $\beta = 0$ , yet the second-order growth condition (Assumption E.1 with  $\gamma = 0$ ) fails to hold.

**Example E.1.**  $(\mathbf{a}, r) \sim \text{Unif}[1, 2]^2$ . The initial inventory for the resource is  $\mathbf{b} = \mathbf{d}T = 1.5T$ .

The distribution in Example E.1 satisfies all the conditions in Assumption 2.2 with  $\beta = 0$ , except for the requirement (iii). A straightforward calculation reveals that  $\boldsymbol{\lambda}^* = \mathbf{0}$ , and Assumption E.1 is satisfied only with  $\gamma = 1$ , suggesting that instead of polylogarithmic in  $T$  regret, only polynomial in  $T$  regret is possible in this specific case. We note that the choice of  $\mathbf{b}$  is a boundary case. Generally, a precise characterization of the achievable regret scaling for OLP instances requires accurately tracking all such boundary cases, which can be complicated depending on the specific problem structure. In this work, we aim to strike a balance between generality and clarity. Hence we present our main results under the current set of assumptions that exclude tricky boundary cases. A further refined regret analysis is left for future research.

## E.2 Global conditions

Some prior work imposes stronger, global-version of Assumption E.1, which essentially requires the dual objective  $f_d(\boldsymbol{\lambda})$  to be quadratically lower bounded not only at the dual optimal point  $\boldsymbol{\lambda}^*$ , but at all points in a given set. We here provide a typical formulation of such global condition.

**Assumption E.2** (Uniform second-order growth condition). There exists a compact convex set  $\Omega \subseteq \mathbb{R}_{\geq 0}^m$  such that for any  $t \in [T]$ , any  $\mathbf{b}'$ , and any problem instance  $\mathcal{I}_{T-t+1}$ , the relaxed offline optimum  $V_{\mathbf{b}', t}^{\text{hind}}(\mathcal{I}_{T-t+1})$  (cf. problem (1) starting from time  $t$ ) possesses one optimal dual solution  $\tilde{\boldsymbol{\lambda}} \in \Omega$ . Moreover, there exist two positive constants  $\underline{\alpha}, \bar{\alpha}$  such that for any  $\boldsymbol{\lambda}', \boldsymbol{\lambda}'' \in \Omega$ , it holds that

$$\begin{aligned} \underline{\alpha} \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left( \mathbf{a}^\top \boldsymbol{\lambda}' - \mathbf{a}^\top \boldsymbol{\lambda}'' \right)^2 &\leq \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}') - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}'') \right) \left( \mathbf{a}^\top \boldsymbol{\lambda}' - \mathbf{a}^\top \boldsymbol{\lambda}'' \right) \right] \\ &\leq \bar{\alpha} \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left( \mathbf{a}^\top \boldsymbol{\lambda}' - \mathbf{a}^\top \boldsymbol{\lambda}'' \right)^2. \end{aligned}$$

Assumption E.2 appears as Assumption 2 in Jiang et al. (2022a). Other examples of global second-order conditions include Assumption 3(b) in Li and Ye (2022). We note that Jiang et al. (2022a) establishes an  $\mathcal{O}(\log T)$  regret guarantee of CE under Assumption E.2, also without imposing non-degeneracy conditions (i.e. Assumption 4.1 or 4.2). We here provide a comparison between their results and ours.

First, the  $\mathcal{O}(\log T)$  regret guarantee in Jiang et al. (2022a) is established in their “semi-discrete” setting, where the number of different types of resource consumption vectors is finite. We note that this discreteness is essential to their analysis, as the inverse of the probability mass of each resource consumption type enters their regret bound. In contrast, our Assumption 2.1 allows the distribution of  $\mathbf{a}$  to be arbitrary, capturing discrete, continuous and mixed-type distributions.

Second, Assumption E.2 is hard to check. In fact, to check whether Assumption E.2 holds, we need to first determine  $\Omega$ , then verify the second-order growth property for every pair of vectors in  $\Omega$ . We note that the uniformity of Assumption E.2 also makes it more restrictive than the standard local second-order conditions such as Assumption E.1. In contrast, our Assumptions 2.1 and 2.2 are distributional assumptions that are straightforward to verify.



## F Technical Proof of Lemmas

This section contains proofs of important Lemmas. In Section F.1, we introduce additional notation. In Section F.2, we provide the proof of Lemma F.1, our regret decomposition result. In Section F.3, we prove the concentration Lemma 5.2. We prove a number of auxiliary results in Section F.4. In particular, in Section F.4.1 we state and prove several results in LP theory. In Section F.4.2, we state and prove concentration results using the tools such as the peeling argument from the empirical processes theory.

### F.1 Additional Notation

We introduce additional notation in order to state our results. For any function  $g$  defined on set  $\mathcal{X}$ , let  $\|g\|_{L_\infty(\mathcal{X})} \triangleq \sup_{x \in \mathcal{X}} |g(x)|$  denote its supremum norm. In cases where  $\mathcal{X}$  is clear from the context, we use the shorthand notation  $\|g\|_\infty$  instead. For any measure  $\mathcal{Q}$  supported on  $\mathcal{X}$ , let  $\|g\|_{\mathcal{Q}} \triangleq (\int |g|^2 d\mathcal{Q})^{1/2}$  denote its  $L_2(\mathcal{Q})$  norm. Following the convention in the theory of empirical processes, we define the *entropy with bracketing* for  $L_2(\mathcal{Q})$  for a function class  $\mathcal{G}$  as follows (cf. Definition 2.2 of Geer (2000)). Let  $N_B(\delta, \mathcal{G}, \mathcal{Q})$  be the smallest value of  $N$  such that there exists pairs of functions  $\{(g_j^L, g_j^U)\}_{j=1}^N$  for which  $\|g_j^U - g_j^L\|_{\mathcal{Q}} \leq \delta$  for all  $j = 1, \dots, N$ , and such that for any  $g \in \mathcal{G}$  there exists a  $j = j(g)$  such that  $g_j^L \leq g \leq g_j^U$ . Then  $H_B(\delta, \mathcal{G}, \mathcal{Q}) = \log N_B(\delta, \mathcal{G}, \mathcal{Q})$  is called the  $\delta$ -entropy with bracketing for  $\mathcal{G}$  (for  $L_2(\mathcal{Q})$ -metric). Let

$$\begin{aligned} h_{t,\mathbf{b}}(\boldsymbol{\lambda}, \mathbf{a}, r) &= \frac{1}{T-t} \mathbf{b}^\top \boldsymbol{\lambda} + \left(r - \mathbf{a}^\top \boldsymbol{\lambda}\right)^+, \\ \phi_{t,\mathbf{b}}(\boldsymbol{\lambda}, \mathbf{a}, r) &= \frac{\partial h_{t,\mathbf{b}}(\boldsymbol{\lambda}, \mathbf{a}, r)}{\partial \boldsymbol{\lambda}} = \frac{1}{T-t} \mathbf{b} - \mathbf{a} \mathbb{I}_{\{r > \mathbf{a}^\top \boldsymbol{\lambda}\}}, \end{aligned}$$

where  $\mathbb{I}_A$  is the indicator that event  $A$  happens and  $h$  is differentiable because we assume  $r$  is continuously distributed. We also introduce functions  $f_{t,\mathbf{b}}(\cdot)$  and  $g_{t,\mathbf{b}}(\cdot)$  that are defined accordingly:

$$\begin{aligned} f_{t,\mathbf{b}}(\boldsymbol{\lambda}) &= \mathbb{E}_{(\mathbf{a},r) \sim F} [h_{t,\mathbf{b}}(\boldsymbol{\lambda}, \mathbf{a}, r)] = \frac{1}{T-t} \mathbf{b}^\top \boldsymbol{\lambda} + \mathbb{E}_{(\mathbf{a},r) \sim F} \left[ \left(r - \mathbf{a}^\top \boldsymbol{\lambda}\right)^+ \right], \\ g_{t,\mathbf{b}}(\boldsymbol{\lambda}) &= \frac{1}{T-t} \sum_{j=t+1}^T h_{t,\mathbf{b}}(\boldsymbol{\lambda}, \mathbf{a}_j, r_j) = \frac{1}{T-t} \mathbf{b}^\top \boldsymbol{\lambda} + \frac{1}{T-t} \sum_{j=t+1}^T \left(r_j - \mathbf{a}_j^\top \boldsymbol{\lambda}\right)^+, \end{aligned}$$

and by our definition,  $\tilde{\boldsymbol{\lambda}}_t, \boldsymbol{\lambda}_t^*, \bar{\boldsymbol{\lambda}}_t^*$  are optimal to problems  $\min_{\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m} f_{t,\mathbf{b}_{t-1}}(\boldsymbol{\lambda}), \min_{\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m} g_{t,\mathbf{b}_{t-1}}(\boldsymbol{\lambda})$  and  $\min_{\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m} g_{t,\mathbf{b}_{t-1}-\mathbf{a}_t}(\boldsymbol{\lambda})$ , respectively. (We generalize the definition of function  $f_{\mathbf{c}}(\cdot)$  in Section 2.2 to  $f_{t,\mathbf{b}}(\cdot)$  to be consistent with function  $g_{t,\mathbf{b}}(\cdot)$ .) Denote by  $\Omega = [0, \frac{\bar{r}}{\underline{A}}]^m$ .

### F.2 Proof of Lemma 5.1

We state a general regret decomposition lemma that holds true for any threshold policy (including CE). Let a threshold policy be specified by a sequence of threshold functions  $\{\theta_t(\cdot, \cdot)\}_{t \in [T]}$ . In particular, the DM makes an ACCEPT decision if and only if  $r_t \geq \theta_t(\mathbf{a}_t, \mathbf{b}_{t-1})$ . Note that CE is a threshold policy.

**Lemma F.1** (Regret Decomposition for Threshold-Based Policies). *Suppose  $\pi \in \Pi$  is threshold-based, associated with a sequence of threshold functions  $\{\theta_t(\cdot, \cdot)\}_{t \in [T]}$ . Then the regret of policy  $\pi$  is upper bounded by*

$$\text{REG}_T(\pi) \leq C_0 \log T + \sum_{t=1}^T \mathbb{E} \left[ \left( \mathbf{a}_t^\top \bar{\boldsymbol{\lambda}}_t^* - r_t \right) \mathbb{I}_{\{\theta_t(\mathbf{a}_t, \mathbf{b}_{t-1}) \leq r_t \leq \mathbf{a}_t^\top \bar{\boldsymbol{\lambda}}_t^*\}} \right]$$

$$+ \sum_{t=1}^T \mathbb{E} \left[ \left( r_t - \mathbf{a}_t^\top \boldsymbol{\lambda}_t^* \right) \mathbb{I}_{\{\mathbf{a}_t^\top \boldsymbol{\lambda}_t^* \leq r_t \leq \theta_t(\mathbf{a}_t, \mathbf{b}_{t-1})\}} \right], \quad (15)$$

where  $\mathcal{C}_0 = 2 \left( 1 + \frac{m\bar{A}}{A} \right) m\bar{r}$ , and the expectation is taken over  $\mathcal{I}_0$ .

*Proof of Lemma F.1.* Let  $V_t^{\text{off}} = V_{t, \mathbf{b}_t}^{\text{off}}(\mathcal{I}_t)$  denote the optimal value of the multi-knapsack LP

$$\begin{aligned} V_t^{\text{off}} &= \max \sum_{j=t+1}^T r_j x_j \\ \text{s.t.} \quad & \sum_{j=t+1}^T a_{ij} x_j \leq b_{it}, \quad i = 1, \dots, m, \\ & x_j \in [0, 1], \quad j = t+1, \dots, T. \end{aligned}$$

By LP duality theory (Lemma F.3), we also have

$$V_t^{\text{off}} = \min_{\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m} \mathbf{b}_t^\top \boldsymbol{\lambda} + \sum_{j=t+1}^T (r_j - \mathbf{a}_j^\top \boldsymbol{\lambda})^+.$$

We consider the following decomposition of regret

$$\begin{aligned} \text{REG}_T(\pi) &= \mathbb{E} \left[ V^{\text{off}} \right] - \mathbb{E} \left[ V^\pi \right] \\ &= \mathbb{E} \left[ V_0^{\text{off}} \right] - \mathbb{E} \left[ \sum_{t=1}^T r_t x_t^\pi \right] \\ &= \mathbb{E} \left[ \sum_{t=1}^T \left( V_{t-1}^{\text{off}} - V_t^{\text{off}} \right) \right] - \mathbb{E} \left[ \sum_{t=1}^T r_t x_t^\pi \right] \quad (\text{since } V_T^{\text{off}} = 0) \\ &= \sum_{t=1}^T \mathbb{E} \left[ V_{t-1}^{\text{off}} - V_t^{\text{off}} - r_t x_t^\pi \right]. \end{aligned} \quad (16)$$

By Lemma F.6, we have the equality

$$V_{t-1, \mathbf{b}_{t-1}}^{\text{off}}(\mathcal{I}_{t-1}) = r_t x_t^* + V_{t, \mathbf{b}_{t-1} - \mathbf{a}_t x_t^*}^{\text{off}}(\mathcal{I}_t),$$

where  $\{x_j^*\}_{j=t}^T$  is any primal optimal solution. The equality implies that  $V_{t-1}^{\text{off}} - V_t^{\text{off}} - r_t x_t^\pi = 0$  if  $x_t^\pi = x_t^*$ . Indeed, if  $x_t^\pi = x_t^* = 1$ , then  $\mathbf{b}_t = \mathbf{b}_{t-1} - \mathbf{a}_t x_t^\pi = \mathbf{b}_{t-1} - \mathbf{a}_t$ , and

$$V_{t-1}^{\text{off}} - V_t^{\text{off}} - r_t x_t^\pi = V_{t-1, \mathbf{b}_{t-1}}^{\text{off}} - V_{t, \mathbf{b}_t}^{\text{off}} - r_t = V_{t, \mathbf{b}_{t-1} - \mathbf{a}_t}^{\text{off}} - V_{t, \mathbf{b}_t}^{\text{off}} = 0.$$

The case  $x_t^\pi = x_t^* = 0$  is similar. In other words, if the decision under policy  $\pi$  coincides with the optimal offline solution, then we have zero regret. Thus it suffices to consider cases where  $x_t^* \neq x_t^\pi$ . To proceed, we suppose  $\mathbf{x}_t^*$  is basic feasible optimal solution for each  $\mathcal{I}_{t-1}$ , and consider the following three cases.

- **Case 1.**  $x_t^* \in (0, 1)$ . In this case  $x_t^*$  is fractional. By Lemma F.5,  $\mathbf{x}_t^*$  has at most  $m$  fractional variables. Since  $\{(\mathbf{a}_j, r_j)\}_{j=t}^T$  are *i.i.d.*, the chance that  $x_t^*$  is fractional is at most  $\frac{m}{T-t+1}$ . In such cases that  $x_t^*$  is indeed fractional, we bound the per period regret as follows

$$V_{t-1}^{\text{off}} - V_t^{\text{off}} - r_t x_t^\pi \leq V_{t-1}^{\text{off}} - V_t^{\text{off}} \leq V_{t-1, \mathbf{b}_{t-1}}^{\text{off}} - V_{t, \mathbf{b}_{t-1} - \mathbf{a}_t}^{\text{off}} \leq V_{t, \mathbf{b}_{t-1}}^{\text{off}} - V_{t, \mathbf{b}_{t-1} - \mathbf{a}_t}^{\text{off}} + \bar{r},$$

where the first inequality follows from the non-negativity of  $x_t^\pi$ , the second follows from the monotonicity of  $V_{t,\mathbf{c}}^{\text{off}}$  in  $\mathbf{c}$ , and the last inequality follows from Lemma F.6 and the monotonicity of  $V_{t,\mathbf{c}}^{\text{off}}$  in  $\mathbf{c}$ . By the boundedness of  $\mathbf{a}_t$ , we further have

$$\begin{aligned} V_{t,\mathbf{b}_{t-1}}^{\text{off}} - V_{t,\mathbf{b}_{t-1}-\mathbf{a}_t}^{\text{off}} + \bar{r} &= (T-t) (g_{t,\mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*) - g_{t,\mathbf{b}_{t-1}-\mathbf{a}_t}(\bar{\boldsymbol{\lambda}}_t^*)) + \bar{r} \\ &\leq (T-t) (g_{t,\mathbf{b}_{t-1}}(\bar{\boldsymbol{\lambda}}_t^*) - g_{t,\mathbf{b}_{t-1}-\mathbf{a}_t}(\bar{\boldsymbol{\lambda}}_t^*)) + \bar{r} \\ &= \mathbf{a}_t^\top \bar{\boldsymbol{\lambda}}_t^* + \bar{r} \leq \frac{m\bar{A}\bar{r}}{\underline{A}} + \bar{r}, \end{aligned}$$

where the last inequality follows from Lemma F.12. We then combine the above arguments and conclude that

$$\mathbb{E} \left[ \left( V_{t-1}^{\text{off}} - V_t^{\text{off}} - r_t x_t^\pi \right) \mathbb{I}_{\{x_t^* \text{ is fractional}\}} \right] \leq \frac{\left(1 + \frac{m\bar{A}}{\underline{A}}\right) m\bar{r}}{T-t+1}. \quad (17)$$

• **Case 2.**  $x_t^\pi = 1, x_t^* = 0$ . By definition of  $\pi$ ,  $x_t^\pi = 1$  implies  $r_t \geq \theta_t(\mathbf{a}_t, \mathbf{b}_{t-1})$ . By Lemma F.6,  $x_t^* = 0$  implies that  $V_{t-1,\mathbf{b}_{t-1}}^{\text{off}} = V_{t,\mathbf{b}_{t-1}}^{\text{off}} \geq V_{t,\mathbf{b}_{t-1}-\mathbf{a}_t}^{\text{off}} + r_t$ , more specifically

$$\mathbf{b}_{t-1}^\top \boldsymbol{\lambda}_t^* + \sum_{j=t+1}^T (r_j - \mathbf{a}_j^\top \boldsymbol{\lambda}_t^*)^+ \geq (\mathbf{b}_{t-1}^\top - \mathbf{a}_t) \bar{\boldsymbol{\lambda}}_t^* + \sum_{j=t+1}^T (r_j - \mathbf{a}_j^\top \bar{\boldsymbol{\lambda}}_t^*)^+ + r_t,$$

where we recall that  $\boldsymbol{\lambda}_t^*$  and  $\bar{\boldsymbol{\lambda}}_t^*$  are optimal dual solutions to Problem (7) and Problem (8), respectively. Since  $\boldsymbol{\lambda}_t^*$  minimizes the above left hand side, the above further implies

$$\begin{aligned} \mathbf{b}_{t-1}^\top \bar{\boldsymbol{\lambda}}_t^* + \sum_{j=t+1}^T (r_j - \mathbf{a}_j^\top \bar{\boldsymbol{\lambda}}_t^*)^+ &\geq (\mathbf{b}_{t-1}^\top - \mathbf{a}_t) \bar{\boldsymbol{\lambda}}_t^* + \sum_{j=t+1}^T (r_j - \mathbf{a}_j^\top \bar{\boldsymbol{\lambda}}_t^*)^+ + r_t, \\ \Leftrightarrow \mathbf{a}_t^\top \bar{\boldsymbol{\lambda}}_t^* &\geq r_t. \end{aligned}$$

Hence  $x_t^\pi = 1, x_t^* = 0$  implies  $\theta_t(\mathbf{a}_t, \mathbf{b}_{t-1}) \leq r_t \leq \mathbf{a}_t^\top \bar{\boldsymbol{\lambda}}_t^*$ .

Since  $x_t^\pi = 1$ , we have  $\mathbf{b}_t = \mathbf{b}_{t-1} - \mathbf{a}_t x_t^\pi = \mathbf{b}_{t-1} - \mathbf{a}_t$ . Therefore we may bound the per period regret as follows

$$\begin{aligned} V_{t-1}^{\text{off}} - V_t^{\text{off}} - r_t x_t^\pi &= V_{t-1,\mathbf{b}_{t-1}}^{\text{off}} - V_{t,\mathbf{b}_{t-1}-\mathbf{a}_t}^{\text{off}} - r_t \\ &= V_{t,\mathbf{b}_{t-1}}^{\text{off}} - V_{t,\mathbf{b}_{t-1}-\mathbf{a}_t}^{\text{off}} - r_t, \\ &= \mathbf{b}_{t-1}^\top \boldsymbol{\lambda}_t^* + \sum_{j=t+1}^T (r_j - \mathbf{a}_j^\top \boldsymbol{\lambda}_t^*)^+ - \left( (\mathbf{b}_{t-1} - \mathbf{a}_t)^\top \bar{\boldsymbol{\lambda}}_t^* + \sum_{j=t+1}^T (r_j - \mathbf{a}_j^\top \bar{\boldsymbol{\lambda}}_t^*)^+ \right) - r_t, \\ &\leq \mathbf{b}_{t-1}^\top \bar{\boldsymbol{\lambda}}_t^* + \sum_{j=t+1}^T (r_j - \mathbf{a}_j^\top \bar{\boldsymbol{\lambda}}_t^*)^+ - \left( (\mathbf{b}_{t-1} - \mathbf{a}_t)^\top \bar{\boldsymbol{\lambda}}_t^* + \sum_{j=t+1}^T (r_j - \mathbf{a}_j^\top \bar{\boldsymbol{\lambda}}_t^*)^+ \right) - r_t, \\ &= \mathbf{a}_t^\top \bar{\boldsymbol{\lambda}}_t^* - r_t, \end{aligned}$$

where the inequality follows from the fact that  $\boldsymbol{\lambda}_t^*$  is the minimizer of Problem (7) and that  $\bar{\boldsymbol{\lambda}}_t^*$  is feasible (non-negative) and hence must achieve a larger objective value when plugging into the objective function of Problem (7). Combining the above, we have

$$\mathbb{E} \left[ \left( V_{t-1}^{\text{off}} - V_t^{\text{off}} - r_t x_t^\pi \right) \mathbb{I}_{\{x_t^*=0, x_t^\pi=1\}} \right] \leq \mathbb{E} \left[ \left( \mathbf{a}_t^\top \bar{\boldsymbol{\lambda}}_t^* - r_t \right) \mathbb{I}_{\{\theta_t(\mathbf{a}_t, \mathbf{b}_{t-1}) \leq r_t \leq \mathbf{a}_t^\top \bar{\boldsymbol{\lambda}}_t^*\}} \right], \quad (18)$$

where the expectation is taken over  $\mathcal{I}_{t-1}$ .

• **Case 3.**  $x_t^\pi = 0, x_t^* = 1$ . This case is very similar to Case 2. We omit the detailed arguments. The conclusion can be summarized as

$$\mathbb{E} \left[ \left( V_{t-1}^{\text{off}} - V_t^{\text{off}} - r_t x_t^\pi \right) \mathbb{I}_{\{x_t^*=1, x_t^\pi=0\}} \right] \leq \mathbb{E} \left[ \left( r_t - \mathbf{a}_t^\top \boldsymbol{\lambda}_t^* \right) \mathbb{I}_{\{\mathbf{a}_t^\top \boldsymbol{\lambda}_t^* \leq r_t \leq \theta_t(\mathbf{a}_t, \mathbf{b}_{t-1})\}} \right]. \quad (19)$$

Now plugging (17), (18) and (19) into (16), we conclude that

$$\begin{aligned} \text{REG}_T(\pi) &= \sum_{t=1}^T \mathbb{E} \left[ V_{t-1}^{\text{off}} - V_t^{\text{off}} - r_t x_t^\pi \right], \\ &= \sum_{t=1}^T \mathbb{E} \left[ \left( V_{t-1}^{\text{off}} - V_t^{\text{off}} - r_t x_t^\pi \right) \mathbb{I}_{\{x_t^* \neq x_t^\pi\}} \right], \\ &= \sum_{t=1}^T \mathbb{E} \left[ \left( V_{t-1}^{\text{off}} - V_t^{\text{off}} - r_t x_t^\pi \right) \mathbb{I}_{\{x_t^* \text{ is fractional}\}} \right] \quad (\text{Case 1}) \\ &\quad + \sum_{t=1}^T \mathbb{E} \left[ \left( V_{t-1}^{\text{off}} - V_t^{\text{off}} - r_t x_t^\pi \right) \mathbb{I}_{\{x_t^*=0, x_t^\pi=1\}} \right] \quad (\text{Case 2}) \\ &\quad + \sum_{t=1}^T \mathbb{E} \left[ \left( V_{t-1}^{\text{off}} - V_t^{\text{off}} - r_t x_t^\pi \right) \mathbb{I}_{\{x_t^*=1, x_t^\pi=0\}} \right] \quad (\text{Case 3}) \\ &\leq \sum_{t=1}^T \frac{\left(1 + \frac{m\bar{A}}{A}\right) m\bar{r}}{T-t+1} + \sum_{t=1}^T \mathbb{E} \left[ \left( \mathbf{a}_t^\top \bar{\boldsymbol{\lambda}}_t^* - r_t \right) \mathbb{I}_{\{\theta_t(\mathbf{a}_t, \mathbf{b}_{t-1}) \leq r_t \leq \mathbf{a}_t^\top \bar{\boldsymbol{\lambda}}_t^*\}} \right] \\ &\quad + \sum_{t=1}^T \mathbb{E} \left[ \left( r_t - \mathbf{a}_t^\top \boldsymbol{\lambda}_t^* \right) \mathbb{I}_{\{\mathbf{a}_t^\top \boldsymbol{\lambda}_t^* \leq r_t \leq \theta_t(\mathbf{a}_t, \mathbf{b}_{t-1})\}} \right] \\ &\leq \mathcal{C} \log T + \sum_{t=1}^T \mathbb{E} \left[ \left( \mathbf{a}_t^\top \bar{\boldsymbol{\lambda}}_t^* - r_t \right) \mathbb{I}_{\{\theta_t(\mathbf{a}_t, \mathbf{b}_{t-1}) \leq r_t \leq \mathbf{a}_t^\top \bar{\boldsymbol{\lambda}}_t^*\}} \right] \\ &\quad + \sum_{t=1}^T \mathbb{E} \left[ \left( r_t - \mathbf{a}_t^\top \boldsymbol{\lambda}_t^* \right) \mathbb{I}_{\{\mathbf{a}_t^\top \boldsymbol{\lambda}_t^* \leq r_t \leq \theta_t(\mathbf{a}_t, \mathbf{b}_{t-1})\}} \right], \end{aligned}$$

where  $\mathcal{C} \triangleq 2 \left(1 + \frac{m\bar{A}}{A}\right) m\bar{r}$ . Q.E.D.

The proof of Lemma 5.1 follows immediately by plugging in  $\theta_t(\mathbf{a}_t, \mathbf{b}_{t-1}) = \mathbf{a}_t^\top \tilde{\boldsymbol{\lambda}}_t$ .

### F.3 Proof of Lemma 5.2

In this section, we provide the proof of Lemma 5.2. To this end, we need an intermediate result, which is stated in the next lemma.

**Lemma F.2.** *Denote by*

$$\begin{aligned} M_1 &= \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right)^2 \max \left( \bar{F}_a^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*), \bar{F}_a^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) \right) \right]}, \\ \bar{M}_1 &= \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right)^2 \max \left( \bar{F}_a^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*), \bar{F}_a^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) \right) \right]}, \end{aligned}$$

where  $\bar{F}_{\mathbf{a}}^r(\cdot) = 1 - F_{\mathbf{a}}^r(\cdot)$ . Then

$$\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right) \right] \leq \tilde{C}_3 \sqrt{\frac{\log(T-t)}{T-t}} \left( M_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right), \quad (20)$$

and

$$\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^*) \right) \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^* \right) \right] \leq \tilde{C}_3 \sqrt{\frac{\log(T-t)}{T-t}} \left( \bar{M}_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right), \quad (21)$$

both with probability at least  $1 - \frac{9C \log(T-t)}{(T-t)^2}$  for any  $T-t \geq e^{\frac{\tilde{C}_1}{32}+1}$ , where  $\tilde{C}_3 = 3\tilde{C}_1 + \frac{4m\bar{A}\bar{r}}{\underline{A}} + \frac{2m^2\bar{A}\bar{r}}{\underline{A}}$ , and  $\tilde{C}_1$  is as in Lemma F.9.

The proof of Lemma F.2 shall be deferred to the end of this subsection. With Lemma F.2, we now finish our proof of Lemma 5.2.

*Proof of Lemma 5.2.* We first prove the lemma under Assumption 2.1. WLOG we only provide the proof regarding  $\boldsymbol{\lambda}_t^*$ , as the proof for  $\bar{\boldsymbol{\lambda}}_t^*$  is nearly identical.

Consider the partition  $[\underline{A}, \bar{A}]^m = \mathcal{E}_1 \cup \mathcal{E}_2$ , where  $\mathcal{E}_1 = \{\mathbf{a} \in [\underline{A}, \bar{A}]^m : F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) = 1 \text{ or } F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) = 1\}$ , and  $\mathcal{E}_2 = \{\mathbf{a} \in [\underline{A}, \bar{A}]^m : F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*), F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) < 1\}$ .

**Case 1.  $\mathcal{E}_1$  happens.** In this case,  $F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) = 1$  or  $F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) = 1$ . WLOG we assume  $F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) = 1$ , then

$$\begin{aligned} & (\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*)^2 \max \left( \bar{F}_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*), \bar{F}_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) \right) \\ &= (\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*)^2 \bar{F}_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) \\ &\leq \max \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t, \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right) \left| \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right| \bar{F}_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) \\ &\leq m \frac{\bar{r}\bar{A}}{\underline{A}} \left| \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right| \bar{F}_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) \quad (\text{by Corollary F.13}) \\ &= m \frac{\bar{r}\bar{A}}{\underline{A}} \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right), \end{aligned}$$

where the last equality follows from  $F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) = 1$  and that  $\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*$  and  $F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*)$  share the same sign. If  $F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) = 1$ , the argument is nearly identical. We further bound the above RHS as follows

$$\begin{aligned} & m \frac{\bar{r}\bar{A}}{\underline{A}} \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right) \\ &\leq \left( 2m \frac{\bar{r}\bar{A}}{\underline{A}} \right)^{\frac{2+2\beta}{2+\beta}} \left( (F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*)) (\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right)^{\frac{2}{2+\beta}}, \quad (22) \end{aligned}$$

where we use the fact that  $\frac{2}{2+\beta} < 1$ ,  $|F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*)| \leq 1$  and  $|\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*| < 2m \frac{\bar{r}\bar{A}}{\underline{A}}$  by Corollary F.13.

**Case 2.  $\mathcal{E}_2$  happens.** In this case, both  $F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t)$  and  $F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*)$  are strictly bounded away from 1. WLOG, assume  $F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) \geq F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*)$ , while the other case  $F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) < F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*)$  is

almost identical. Hence, by the Hölder condition in Assumption 2.1, we have

$$c_\beta(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*)^{1+\beta} \leq F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*),$$

which leads to

$$\begin{aligned} & (\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*)^2 \max \left( \bar{F}_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*), \bar{F}_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) \right) \\ & \leq (\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*)^2 \\ & = c_\beta^{-\frac{2}{2+\beta}} (\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*)^{\frac{2}{2+\beta}} c_\beta^{\frac{2}{2+\beta}} (\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*)^{\frac{2+2\beta}{2+\beta}}, \\ & \leq c_\beta^{-\frac{2}{2+\beta}} (\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*)^{\frac{2}{2+\beta}} \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right)^{\frac{2}{2+\beta}}. \end{aligned} \quad (23)$$

Combining (22) with (23), we conclude that

$$\begin{aligned} M_1 &= \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ (\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*)^2 \max \left( \bar{F}_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*), \bar{F}_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) \right) \right]} \\ &= \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ (\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*)^2 \max \left( \bar{F}_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*), \bar{F}_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) \right) (I_{\mathcal{E}_1} + I_{\mathcal{E}_2}) \right]} \\ &\leq \max \left( \left( 2m \frac{\bar{r}\bar{A}}{\underline{A}} \right)^{\frac{1+\beta}{2+\beta}}, c_\beta^{-\frac{1}{2+\beta}} \right) \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ (\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*)^{\frac{2}{2+\beta}} \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right)^{\frac{2}{2+\beta}} \right]} \\ &\leq \max \left( \left( 2m \frac{\bar{r}\bar{A}}{\underline{A}} \right)^{\frac{1+\beta}{2+\beta}}, c_\beta^{-\frac{1}{2+\beta}} \right) \left( \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ (\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*) \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right) \right] \right)^{\frac{1}{2+\beta}} \\ &\quad \text{(by Jensen's inequality since } x^{\frac{2}{2+\beta}} \text{ is concave).} \end{aligned} \quad (24)$$

Plugging (24) into Lemma F.2,

$$\begin{aligned} & \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right) \right] \\ & \leq \tilde{C}_3 \sqrt{\frac{\log(T-t)}{T-t}} \left( M_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right) \\ & \leq \tilde{C}_3 \frac{\log(T-t)}{T-t} + \tilde{C}_3 \max \left( \left( 2m \frac{\bar{r}\bar{A}}{\underline{A}} \right)^{\frac{1+\beta}{2+\beta}}, c_\beta^{-\frac{1}{2+\beta}} \right) \sqrt{\frac{\log(T-t)}{T-t}} \\ & \quad \times \left( \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ (\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*) \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right) \right] \right)^{\frac{1}{2+\beta}}, \end{aligned}$$

which, through straightforward algebra, implies that

$$\begin{aligned} & \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right) \right] \\ & \leq \tilde{C}_3 \left( 1 + \max \left( \left( 2m \frac{\bar{r}\bar{A}}{\underline{A}} \right)^{\frac{1+\beta}{2+\beta}}, c_\beta^{-\frac{1}{2+\beta}} \right) \right)^{\frac{2+\beta}{1+\beta}} \left( \frac{\log(T-t)}{T-t} \right)^{\frac{2+\beta}{2+2\beta}}, \triangleq \tilde{C}_4 \left( \frac{\log(T-t)}{T-t} \right)^{\frac{2+\beta}{2+2\beta}}, \end{aligned}$$

with probability at least  $1 - \frac{9C \log(T-t)}{(T-t)^2}$  for any  $T - t \geq e^{\frac{\tilde{C}_1}{32} + 1}$ . Note that we have chosen

$$\tilde{C}_4 \triangleq \tilde{C}_3 \left( 1 + \max \left( \left( 2m \frac{\bar{r}\bar{A}}{\underline{A}} \right)^{\frac{1+\beta}{2+\beta}}, c_\beta^{-\frac{1}{2+\beta}} \right) \right)^{\frac{2+\beta}{1+\beta}},$$

thus concluding the proof of the lemma under Assumption 2.1.

Next, we provide the proof under Assumption 2.2. Observe that under Assumption 2.2, for any  $\mathbf{a} \in \text{supp}(F^{\mathbf{a}})$ ,  $c_\beta(r_{\mathbf{a}} - l_{\mathbf{a}})^{1+\beta} \leq F_{\mathbf{a}}^r(r_{\mathbf{a}}) - F_{\mathbf{a}}^r(l_{\mathbf{a}}) = 1 \leq c_\nu(r_{\mathbf{a}} - l_{\mathbf{a}})^\nu$ . Thus  $c_\nu^{\frac{1}{\nu}} \leq r_{\mathbf{a}} - l_{\mathbf{a}} \leq c_\beta^{-\frac{1}{1+\beta}}$  for any  $\mathbf{a}$ .

We divide the proof of Lemma 5.2 under Assumption 2.2 into several cases.

**Case 1.** There exists  $\mathbf{a} \in \text{supp}(F^{\mathbf{a}})$  such that either one of the following is satisfied

$$\frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} - \frac{1}{8}c_\nu^{\frac{1}{\nu}} \leq \mathbf{a}^\top \boldsymbol{\lambda}_t^* \leq \frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} + \frac{1}{8}c_\nu^{\frac{1}{\nu}}, \quad (25)$$

$$\frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} - \frac{1}{8}c_\nu^{\frac{1}{\nu}} \leq \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t \leq \frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} + \frac{1}{8}c_\nu^{\frac{1}{\nu}}, \quad (26)$$

where without loss of generality, we assume that (25) holds, and note that the other case, i.e., (26) holds, is nearly identical. Denote by  $r_1 \triangleq \frac{1}{8} \left( (c_L + \sqrt{m} \frac{\bar{r}}{\underline{A}})^{-1} \wedge 1 \right) c_\nu^{\frac{1}{\nu}}$ . Consider a ball  $\mathcal{B}(\mathbf{a}, r_1)$  centered at  $\mathbf{a}$  with radius  $r_1$ , where we recall from Assumption 2.2 that  $c_L$  is the Lipschitz constant of  $l_{\mathbf{a}}$  and  $r_{\mathbf{a}}$ . By Lemma F.16, for all  $\mathbf{a}' \in \mathcal{B}(\mathbf{a}, r_1)$ ,

$$\frac{l_{\mathbf{a}'} + r_{\mathbf{a}'}}{2} - \frac{1}{4}c_\nu^{\frac{1}{\nu}} \leq \mathbf{a}'^\top \boldsymbol{\lambda}_t^* \leq \frac{l_{\mathbf{a}'} + r_{\mathbf{a}'}}{2} + \frac{1}{4}c_\nu^{\frac{1}{\nu}}. \quad (27)$$

Because  $r_{\mathbf{a}} - l_{\mathbf{a}} \geq c_\nu^{\frac{1}{\nu}}$ , for all  $\mathbf{a}' \in \mathcal{B}(\mathbf{a}, r_1)$ ,  $\mathbf{a}'^\top \boldsymbol{\lambda}_t^* \in [l_{\mathbf{a}'}, r_{\mathbf{a}'}]$ . If  $\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t \in [l_{\mathbf{a}'}, r_{\mathbf{a}'}]$ , then since both  $\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t$  and  $\mathbf{a}'^\top \boldsymbol{\lambda}_t^*$  belong to the interval  $[l_{\mathbf{a}'}, r_{\mathbf{a}'}]$ , we have

$$\left| F_{\mathbf{a}'}^r(\mathbf{a}'^\top \boldsymbol{\lambda}_t^*) - F_{\mathbf{a}'}^r(\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t) \right| \geq c_\beta \left( \mathbf{a}'^\top \boldsymbol{\lambda}_t^* - \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t \right)^{1+\beta}. \quad (28)$$

If  $\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t \notin [l_{\mathbf{a}'}, r_{\mathbf{a}'}]$ , then by (27),  $\mathbf{a}'^\top \boldsymbol{\lambda}_t^*$  is at least  $\frac{1}{4}c_\nu^{\frac{1}{\nu}}$  far away from  $l_{\mathbf{a}'}$  and  $r_{\mathbf{a}'}$ , which implies

$$\begin{aligned} \left| F_{\mathbf{a}'}^r(\mathbf{a}'^\top \boldsymbol{\lambda}_t^*) - F_{\mathbf{a}'}^r(\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t) \right| &\geq \min \left( \left| F_{\mathbf{a}'}^r(\mathbf{a}'^\top \boldsymbol{\lambda}_t^*) - F_{\mathbf{a}'}^r(l_{\mathbf{a}'}) \right|, \left| F_{\mathbf{a}'}^r(\mathbf{a}'^\top \boldsymbol{\lambda}_t^*) - F_{\mathbf{a}'}^r(r_{\mathbf{a}'}) \right| \right) \\ &\geq c_\beta \left( \frac{1}{4}c_\nu^{\frac{1}{\nu}} \right)^{1+\beta} = 4^{-1-\beta} c_\beta c_\nu^{\frac{1+\beta}{\nu}}. \end{aligned} \quad (29)$$

Applying Lemma F.15, we have

$$\mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}, \mathbf{a}' \in \mathcal{B}(\mathbf{a}, r)} (\mathbf{a}'^\top \boldsymbol{\lambda}_1 - \mathbf{a}'^\top \boldsymbol{\lambda}_2)^2 \geq C_f \left( \frac{r_1 \wedge w}{6} \sin(\theta) \right)^{m+2} \|\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2\|^2, \quad (30)$$

where  $C_f = l_f \frac{\pi^{m/2}}{\Gamma(\frac{m}{2} + 1)}$ . We thus have

$$\begin{aligned} &\mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}'}^r(\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}'}^r(\mathbf{a}'^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}'^\top \boldsymbol{\lambda}_t^* \right) \right] \\ &\geq \mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}, \mathbf{a}' \in \mathcal{B}(\mathbf{a}, r_1)} \left[ \left( F_{\mathbf{a}'}^r(\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}'}^r(\mathbf{a}'^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}'^\top \boldsymbol{\lambda}_t^* \right) \right] \end{aligned}$$

$$\begin{aligned}
&\geq \mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}, \mathbf{a}' \in \mathcal{B}(\mathbf{a}, r_1)} \left[ \min \left( c_\beta \left| \mathbf{a}'^\top \boldsymbol{\lambda}_t^* - \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t \right|^{1+\beta}, 4^{-1-\beta} c_\beta c_\nu^{\frac{1+\beta}{\nu}} \right) \left| \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}'^\top \boldsymbol{\lambda}_t^* \right| \right] \quad (\text{by (28) and (29)}) \\
&= \mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}, \mathbf{a}' \in \mathcal{B}(\mathbf{a}, r_1)} \left[ \left| \mathbf{a}'^\top \boldsymbol{\lambda}_t^* - \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t \right|^{2+\beta} \min \left( c_\beta, 4^{-1-\beta} c_\beta c_\nu^{\frac{1+\beta}{\nu}} \left| \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}'^\top \boldsymbol{\lambda}_t^* \right|^{-1-\beta} \right) \right] \\
&\geq \min \left( c_\beta, 4^{-1-\beta} c_\beta c_\nu^{\frac{1+\beta}{\nu}} \left( 2\sqrt{m} \frac{\bar{r}\bar{A}}{\underline{A}} \right)^{-1-\beta} \right) \mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}, \mathbf{a}' \in \mathcal{B}(\mathbf{a}, r_1)} \left[ \left| \mathbf{a}'^\top \boldsymbol{\lambda}_t^* - \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t \right|^{2+\beta} \right] \quad (\text{by Lemma F.12}) \\
&\geq c_1 \left( \mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}, \mathbf{a}' \in \mathcal{B}(\mathbf{a}, r_1)} \left[ \left( \mathbf{a}'^\top \boldsymbol{\lambda}_t^* - \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t \right)^2 \right] \right)^{\frac{2+\beta}{2}} \quad (\text{by Jensen's inequality}) \\
&\geq c_1 C_f^{\frac{2+\beta}{2}} \left( \frac{r_1 \wedge w}{6} \sin(\theta) \right)^{\frac{(2+\beta)(m+2)}{2}} \|\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t\|^{2+\beta} \quad (\text{by (30)}), \tag{31}
\end{aligned}$$

where  $c_1 = \min \left( c_\beta, 4^{-1-\beta} c_\beta c_\nu^{\frac{1+\beta}{\nu}} \left( 2\sqrt{m} \frac{\bar{r}\bar{A}}{\underline{A}} \right)^{-1-\beta} \right)$ . We set  $c_2 = c_1 C_f^{\frac{2+\beta}{2}} \left( \frac{r_1 \wedge w}{6} \sin(\theta) \right)^{\frac{(2+\beta)(m+2)}{2}}$ . Note that

$$\begin{aligned}
M_1 &= \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right)^2 \max \left( \bar{F}_\mathbf{a}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*), \bar{F}_\mathbf{a}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) \right) \right]} \\
&\leq \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right)^2 \right]} \leq \bar{A} \|\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t\|,
\end{aligned}$$

which, together with (31) and Lemma F.2, implies

$$\begin{aligned}
c_2 \|\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t\|^{2+\beta} &\leq \mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}'}^r(\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}'}^r(\mathbf{a}'^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}'^\top \boldsymbol{\lambda}_t^* \right) \right] \\
&\leq \tilde{C}_3 \sqrt{\frac{\log(T-t)}{T-t}} \left( M_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right) \leq \tilde{C}_3 \frac{\log(T-t)}{T-t} + \tilde{C}_3 \bar{A} \sqrt{\frac{\log(T-t)}{T-t}} \|\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t\|. \tag{32}
\end{aligned}$$

Solving (32) yields

$$\|\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t\| \leq \max \left( \frac{\tilde{C}_3}{c_2} \left( \frac{\log(T-t)}{T-t} \right)^{\frac{1}{1+\beta}}, \frac{\tilde{C}_3 \bar{A}}{c_2} \left( \frac{\log(T-t)}{T-t} \right)^{\frac{1}{2(1+\beta)}} \right) \leq c_3 \left( \frac{\log(T-t)}{T-t} \right)^{\frac{1}{2(1+\beta)}}, \tag{33}$$

where  $c_3 = \max \left( \frac{\tilde{C}_3}{c_2}, \frac{\tilde{C}_3 \bar{A}}{c_2} \right)$ , and we use the fact that  $\frac{\log(T-t)}{T-t} \leq 1$  and  $\frac{1}{2+\beta} \geq \frac{1}{2(1+\beta)}$ . By (32) and (33), we obtain

$$\begin{aligned}
&\mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}'}^r(\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}'}^r(\mathbf{a}'^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}'^\top \boldsymbol{\lambda}_t^* \right) \right] \\
&\leq \tilde{C}_3 \frac{\log(T-t)}{T-t} + \tilde{C}_3 \bar{A} \sqrt{\frac{\log(T-t)}{T-t}} \|\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t\| \leq c_4 \left( \frac{\log(T-t)}{T-t} \right)^{\frac{2+\beta}{2(1+\beta)}}, \tag{34}
\end{aligned}$$

with probability at least  $1 - \frac{9C \log(T-t)}{(T-t)^2}$  and for  $T-t \geq e^{\frac{\tilde{C}_1}{32}+1}$ , with  $c_4 = \tilde{C}_3 + \tilde{C}_3 \bar{A} c_3$ .

**Case 2.** For any  $\mathbf{a} \in \text{supp}(F^{\mathbf{a}})$ , the following holds

$$\mathbf{a}^\top \boldsymbol{\lambda}_t^* \notin \left[ \frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} - \frac{1}{8} c_\nu^{\frac{1}{\nu}}, \frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} + \frac{1}{8} c_\nu^{\frac{1}{\nu}} \right],$$



$$\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t \notin \left[ \frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} - \frac{1}{8}c_{\nu}^{\frac{1}{\nu}}, \frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} + \frac{1}{8}c_{\nu}^{\frac{1}{\nu}} \right].$$

We first show that there does not exist  $\mathbf{a}', \mathbf{a}'' \in \text{supp}(F^{\mathbf{a}})$  such that

$$\mathbf{a}'^\top \boldsymbol{\lambda}_t^* \leq \frac{l_{\mathbf{a}'} + r_{\mathbf{a}'}}{2} - \frac{1}{8}c_{\nu}^{\frac{1}{\nu}}, \quad \mathbf{a}''^\top \boldsymbol{\lambda}_t^* \geq \frac{l_{\mathbf{a}''} + r_{\mathbf{a}''}}{2} + \frac{1}{8}c_{\nu}^{\frac{1}{\nu}}$$

hold. For otherwise, consider  $\mathbf{a}(p) \triangleq p\mathbf{a}' + (1-p)\mathbf{a}''$  for  $p \in [0, 1]$ , and function

$$g_{\text{gap}}(p) \triangleq \frac{l_{\mathbf{a}(p)} + r_{\mathbf{a}(p)}}{2} - \mathbf{a}(p)^\top \boldsymbol{\lambda}_t^*.$$

Then  $g_{\text{gap}}(0) \leq -\frac{1}{8}c_{\nu}^{\frac{1}{\nu}}$  and  $g_{\text{gap}}(1) \geq \frac{1}{8}c_{\nu}^{\frac{1}{\nu}}$ . By the continuity of  $l_{\mathbf{a}}$  and  $r_{\mathbf{a}}$  in  $\mathbf{a}$ ,  $g_{\text{gap}}(p)$  is a continuous function in  $p$ . By the intermediate value theorem, there must exist  $p' \in (0, 1)$  such that  $g_{\text{gap}}(p') = 0$ . Such  $p'$  corresponds to  $p'\mathbf{a}' + (1-p')\mathbf{a}''$ , which belongs to  $\text{supp}(F^{\mathbf{a}})$  by its own convexity. However, the existence of  $p'\mathbf{a}' + (1-p')\mathbf{a}''$  violates the condition of Case 2. Therefore by contradiction, it must be that either

$$\mathbf{a}^\top \boldsymbol{\lambda}_t^* < \frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} - \frac{1}{8}c_{\nu}^{\frac{1}{\nu}},$$

for all  $\mathbf{a} \in \text{supp}(F^{\mathbf{a}})$ , or

$$\mathbf{a}^\top \boldsymbol{\lambda}_t^* > \frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} + \frac{1}{8}c_{\nu}^{\frac{1}{\nu}},$$

for all  $\mathbf{a} \in \text{supp}(F^{\mathbf{a}})$ . We note that the same should be true with respect to  $\tilde{\boldsymbol{\lambda}}_t$ . We thus consider three sub-cases.

**Sub-case 1.** For all  $\mathbf{a}$ ,  $\mathbf{a}^\top \boldsymbol{\lambda}_t^* < \frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} - \frac{1}{8}c_{\nu}^{\frac{1}{\nu}}$ , yet  $\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t > \frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} + \frac{1}{8}c_{\nu}^{\frac{1}{\nu}}$ , or the other way round. In this case, for all  $\mathbf{a}' \in \text{supp}(F^{\mathbf{a}})$ ,

$$\left| F_{\mathbf{a}'}^r(\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}'}^r(\mathbf{a}'^\top \boldsymbol{\lambda}_t^*) \right| \geq F_{\mathbf{a}'}^r \left( \frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} + \frac{1}{8}c_{\nu}^{\frac{1}{\nu}} \right) - F_{\mathbf{a}'}^r \left( \frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} - \frac{1}{8}c_{\nu}^{\frac{1}{\nu}} \right) \geq c_{\beta} \left( \frac{1}{4}c_{\nu}^{\frac{1}{\nu}} \right)^{1+\beta},$$

where the second inequality follows from the Hölder condition in Assumption 2.2 and the fact that both  $\frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} - \frac{1}{8}c_{\nu}^{\frac{1}{\nu}}$  and  $\frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} + \frac{1}{8}c_{\nu}^{\frac{1}{\nu}}$  belong to the support, i.e.  $[l_{\mathbf{a}}, r_{\mathbf{a}}]$ . Thus we have

$$\mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}'}^r(\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}'}^r(\mathbf{a}'^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}'^\top \boldsymbol{\lambda}_t^* \right) \right] \geq 4^{-1-\beta} c_{\beta} c_{\nu}^{\frac{1+\beta}{\nu}} \mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}} \left| \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}'^\top \boldsymbol{\lambda}_t^* \right|. \quad (35)$$

Note that

$$\begin{aligned} M_1 &= \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right)^2 \max \left( \bar{F}_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*), \bar{F}_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) \right) \right]} \\ &\leq \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right)^2 \right]} \leq \sqrt{\frac{2\bar{A}\sqrt{m}\bar{r}}{\underline{A}} \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left| \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right|}, \quad (\text{by Lemma F.12}) \end{aligned}$$

which, together with (35) and Lemma F.2, implies

$$4^{-1-\beta} c_{\beta} c_{\nu}^{\frac{1+\beta}{\nu}} \mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}} \left| \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}'^\top \boldsymbol{\lambda}_t^* \right|$$

$$\begin{aligned}
&\leq \mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}'}^r(\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}'}^r(\mathbf{a}'^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}'^\top \boldsymbol{\lambda}_t^* \right) \right] \\
&\leq \tilde{C}_3 \sqrt{\frac{\log(T-t)}{T-t}} \left( M_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right) \\
&\leq \tilde{C}_3 \frac{\log(T-t)}{T-t} + \tilde{C}_3 \sqrt{\frac{\log(T-t)}{T-t}} \sqrt{\frac{2\bar{A}\sqrt{m\bar{r}}}{\underline{A}} \mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}} \left| \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}'^\top \boldsymbol{\lambda}_t^* \right|}. \tag{36}
\end{aligned}$$

Solving (36), we obtain

$$\mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}} \left| \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}'^\top \boldsymbol{\lambda}_t^* \right| \leq \max \left( \tilde{C}_3 4^{1+\beta} c_\beta^{-1} c_\nu^{-\frac{1+\beta}{\nu}}, \tilde{C}_3^2 \frac{2\bar{A}\sqrt{m\bar{r}}}{\underline{A}} 4^{2+2\beta} c_\beta^{-2} c_\nu^{-\frac{2(1+\beta)}{\nu}} \right) \frac{\log(T-t)}{T-t}, \tag{37}$$

Plugging back into eq. (36) yields

$$\mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}'}^r(\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}'}^r(\mathbf{a}'^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}'^\top \boldsymbol{\lambda}_t^* \right) \right] \leq c_5 \frac{\log(T-t)}{T-t}, \tag{38}$$

where  $c_5 = \tilde{C}_3 + \tilde{C}_3 \left( \frac{2\bar{A}\sqrt{m\bar{r}}}{\underline{A}} \max \left( \tilde{C}_3 4^{1+\beta} c_\beta^{-1} c_\nu^{-\frac{1+\beta}{\nu}}, \tilde{C}_3^2 \frac{2\bar{A}\sqrt{m\bar{r}}}{\underline{A}} 4^{2+2\beta} c_\beta^{-2} c_\nu^{-\frac{2(1+\beta)}{\nu}} \right) \right)^{1/2}$ . Recall that

$\frac{2+\beta}{2(1+\beta)} < 1$  and  $\frac{\log(T-t)}{T-t} \leq 1$ , we have  $\frac{\log(T-t)}{T-t} \leq \left( \frac{\log(T-t)}{T-t} \right)^{\frac{2+\beta}{2(1+\beta)}}$ . Further we note that  $c_5 \leq \tilde{C}_4$ .

We thus complete the proof of Lemma 5.2 under Assumption 2.2 in the Sub-case 1 of Case 2.

**Sub-case 2.** For all  $\mathbf{a}, \mathbf{a}'^\top \boldsymbol{\lambda}_t^* > \frac{l_{\mathbf{a}+r_{\mathbf{a}}}}{2} + \frac{1}{8}c_\nu^{\frac{1}{\nu}}$ , and  $\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t > \frac{l_{\mathbf{a}+r_{\mathbf{a}}}}{2} + \frac{1}{8}c_\nu^{\frac{1}{\nu}}$ . The proof in this case is identical to the proof under Assumption 2.1, because both  $\mathbf{a}'^\top \boldsymbol{\lambda}_t^*$  and  $\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t$  never fall below the lower bound of the support of  $F_{\mathbf{a}'}^r$  for all  $\mathbf{a} \in \text{supp}(F^{\mathbf{a}})$ , and thus Lemma 5.2 is true in this sub-case with the same constant as chosen in the previous proof under Assumption 2.1.

**Sub-case 3.** For all  $\mathbf{a}, \mathbf{a}'^\top \boldsymbol{\lambda}_t^* < \frac{l_{\mathbf{a}+r_{\mathbf{a}}}}{2} - \frac{1}{8}c_\nu^{\frac{1}{\nu}}$ , and  $\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t < \frac{l_{\mathbf{a}+r_{\mathbf{a}}}}{2} - \frac{1}{8}c_\nu^{\frac{1}{\nu}}$ . In this case, we focus on  $\mathcal{B}(\mathbf{a}_o, r_0) \cap \text{supp}(F^{\mathbf{a}})$ , where we recall that by Assumption 2.2, for  $\mathbf{a} \in \mathcal{B}(\mathbf{a}_o, r_0) \cap \text{supp}(F^{\mathbf{a}})$ ,  $l_{\mathbf{a}} = 0$ . Hence for all such  $\mathbf{a}$ , we have by the Hölder condition of Assumption 2.2 that

$$\left| F_{\mathbf{a}}^r(\mathbf{a}'^\top \boldsymbol{\lambda}_t^*) - F_{\mathbf{a}}^r(\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t) \right| \geq c_\beta \left| \mathbf{a}'^\top \boldsymbol{\lambda}_t^* - \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t \right|^{1+\beta}.$$

Therefore, we have

$$\begin{aligned}
&\mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}'}^r(\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}'}^r(\mathbf{a}'^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}'^\top \boldsymbol{\lambda}_t^* \right) \right] \\
&\geq \mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}, \mathbf{a}' \in \mathcal{B}(\mathbf{a}_o, r_0)} \left[ \left( F_{\mathbf{a}'}^r(\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}'}^r(\mathbf{a}'^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}'^\top \boldsymbol{\lambda}_t^* \right) \right] \\
&\geq \mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}, \mathbf{a}' \in \mathcal{B}(\mathbf{a}_o, r_0)} \left[ c_\beta \left| \mathbf{a}'^\top \boldsymbol{\lambda}_t^* - \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t \right|^{2+\beta} \right] \\
&\geq c_\beta \left( \mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}, \mathbf{a}' \in \mathcal{B}(\mathbf{a}_o, r_0)} \left[ \left( \mathbf{a}'^\top \boldsymbol{\lambda}_t^* - \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t \right)^2 \right] \right)^{\frac{2+\beta}{2}} \quad (\text{by Jensen's inequality}) \\
&\geq c_\beta C_f^{\frac{2+\beta}{2}} \left( \frac{r_0 \wedge w}{6} \sin(\theta) \right)^{\frac{(2+\beta)(m+2)}{2}} \|\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t\|^{2+\beta} \quad (\text{by Lemma F.15}).
\end{aligned}$$

The rest of the proof is almost identical to Case 1, and we omit the details. This gives us

$$\mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}'}^r(\mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}'}^r(\mathbf{a}'^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}'^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}'^\top \boldsymbol{\lambda}_t^* \right) \right] \leq c_6 \left( \frac{\log(T-t)}{T-t} \right)^{\frac{2+\beta}{2(1+\beta)}}, \quad (39)$$

where  $c_6 = \tilde{C}_3 + \tilde{C}_3^2 \bar{A} \left( c_\beta C_f^{\frac{2+\beta}{2}} \left( \frac{r_0 \Lambda w}{6} \sin(\theta) \right)^{\frac{(2+\beta)(m+2)}{2}} \right)^{-1} \max(1, \bar{A})$ .

Combining the above completes the proof of Lemma 5.2, where we summarize the concrete constants appearing in the bound in Remark B.1. Q.E.D.

We now prove Lemma F.2.

*Proof of Lemma F.2.* By Lemma F.10, for any  $\boldsymbol{\lambda}_1, \boldsymbol{\lambda}_2 \in \mathbb{R}_{\geq 0}^m$ ,

$$h_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_1, \mathbf{a}, r) - h_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_2, \mathbf{a}, r) = \phi_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_2, \mathbf{a}, r)^\top (\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2) + \int_{\mathbf{a}^\top \boldsymbol{\lambda}_1}^{\mathbf{a}^\top \boldsymbol{\lambda}_2} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \boldsymbol{\lambda}_2\}} \right) dv,$$

applying to functions  $f_{t, \mathbf{b}_{t-1}}$  and  $g_{t, \mathbf{b}_{t-1}}$  evaluated at  $\tilde{\boldsymbol{\lambda}}_t, \boldsymbol{\lambda}_t^*$ ,

$$\begin{aligned} & f_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*) - f_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t) \\ &= \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t, \mathbf{a}, r)^\top (\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t) + \mathbb{E}_{(\mathbf{a}, r) \sim F} \int_{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t}^{\mathbf{a}^\top \boldsymbol{\lambda}_t^*} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t\}} \right) dv, \end{aligned} \quad (40)$$

$$\begin{aligned} & f_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t) - f_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*) \\ &= \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*, \mathbf{a}, r)^\top (\tilde{\boldsymbol{\lambda}}_t - \boldsymbol{\lambda}_t^*) + \mathbb{E}_{(\mathbf{a}, r) \sim F} \int_{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t}^{\mathbf{a}^\top \boldsymbol{\lambda}_t^*} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \boldsymbol{\lambda}_t^*\}} \right) dv, \end{aligned} \quad (41)$$

and

$$\begin{aligned} & g_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*) - g_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t) \\ &= \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t, \mathbf{a}_j, r_j)^\top (\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t) + \frac{1}{T-t} \sum_{j=t+1}^T \int_{\mathbf{a}_j^\top \tilde{\boldsymbol{\lambda}}_t}^{\mathbf{a}_j^\top \boldsymbol{\lambda}_t^*} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}_j^\top \tilde{\boldsymbol{\lambda}}_t\}} \right) dv, \end{aligned} \quad (42)$$

$$\begin{aligned} & g_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t) - g_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*) \\ &= \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*, \mathbf{a}_j, r_j)^\top (\tilde{\boldsymbol{\lambda}}_t - \boldsymbol{\lambda}_t^*) + \frac{1}{T-t} \sum_{j=t+1}^T \int_{\mathbf{a}_j^\top \tilde{\boldsymbol{\lambda}}_t}^{\mathbf{a}_j^\top \boldsymbol{\lambda}_t^*} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}_j^\top \boldsymbol{\lambda}_t^*\}} \right) dv. \end{aligned} \quad (43)$$

We first prove the first bound. To this end, we observe that

$$\begin{aligned} & \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right) \right] \\ &= \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \int_{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t}^{\mathbf{a}^\top \boldsymbol{\lambda}_t^*} \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(v) \right) dv + \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \int_{\mathbf{a}^\top \boldsymbol{\lambda}_t^*}^{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t} \left( F_{\mathbf{a}}^r(v) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right) dv. \end{aligned}$$

We bound the two integrals respectively. Because  $\tilde{\boldsymbol{\lambda}}_t$  minimizes  $f_{t, \mathbf{b}_{t-1}}(\cdot)$  and  $\boldsymbol{\lambda}_t^*$  minimizes  $g_{t, \mathbf{b}_{t-1}}(\cdot)$ , by (40) and (42), we have

$$\begin{aligned} 0 &\leq f_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*) - f_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t) \\ &\leq f_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*) - f_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t) - (g_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*) - g_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t)) \end{aligned}$$

$$\begin{aligned}
&= \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t, \mathbf{a}, r)^\top (\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t) - \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t, \mathbf{a}_j, r_j)^\top (\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t) \\
&\quad + \mathbb{E}_{(\mathbf{a}, r) \sim F} \int_{\mathbf{a}^\top \boldsymbol{\lambda}_t^*}^{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t\}} \right) dv - \frac{1}{T-t} \sum_{j=t+1}^T \int_{\mathbf{a}_j^\top \boldsymbol{\lambda}_t^*}^{\mathbf{a}_j^\top \tilde{\boldsymbol{\lambda}}_t} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}_j^\top \tilde{\boldsymbol{\lambda}}_t\}} \right) dv.
\end{aligned} \tag{44}$$

Applying Lemma F.8 and Lemma F.9 to (44), together with Lemma F.11, we have as long as  $T-t \geq e^{\frac{\tilde{C}_1}{32}+1}$ , with probability at least  $1 - 6C \log(T-t)(T-t)^{-2}$  with  $C$  a universal constant,

$$\begin{aligned}
0 &\leq f_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*) - f_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t) \\
&\leq \tilde{C}_1 \sqrt{\frac{\log(T-t)}{T-t}} \left( \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} (\mathbf{a}^\top (\tilde{\boldsymbol{\lambda}}_t - \boldsymbol{\lambda}_t^*))^2 (1 - F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t))} + 2 \sqrt{\frac{\log(T-t)}{T-t}} \right. \\
&\quad \left. + \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} (\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*)^2 \left| F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right|} \right) \\
&\leq 2\tilde{C}_1 \sqrt{\frac{\log(T-t)}{T-t}} \left( M_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right),
\end{aligned} \tag{45}$$

where the first inequality is also because  $\tilde{C}_1 \geq \tilde{C}$ , where  $\tilde{C}$  and  $\tilde{C}_1$  are as in Lemma F.8 and Lemma F.9, respectively. In (45),

$$M_1 = \max \left\{ \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} (\mathbf{a}^\top (\tilde{\boldsymbol{\lambda}}_t - \boldsymbol{\lambda}_t^*))^2 (1 - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*))}, \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} (\mathbf{a}^\top (\tilde{\boldsymbol{\lambda}}_t - \boldsymbol{\lambda}_t^*))^2 (1 - F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t))} \right\}.$$

By Lemma F.14 with  $\boldsymbol{\lambda} = \boldsymbol{\lambda}_t^*$ , we have

$$\mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t, \mathbf{a}, r)^\top (\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t) \geq 0. \tag{46}$$

Plugging (46) into (40), we obtain

$$\begin{aligned}
f_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*) - f_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t) &\geq \mathbb{E}_{(\mathbf{a}, r) \sim F} \int_{\mathbf{a}^\top \boldsymbol{\lambda}_t^*}^{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t\}} \right) dv \\
&= \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \int_{\mathbf{a}^\top \boldsymbol{\lambda}_t^*}^{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t} \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(v) \right) dv,
\end{aligned}$$

which, together with (45), gives us that with probability at least  $1 - \frac{6C \log(T-t)}{(T-t)^2}$ ,

$$\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \int_{\mathbf{a}^\top \boldsymbol{\lambda}_t^*}^{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t} \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(v) \right) dv \leq 2\tilde{C}_1 \sqrt{\frac{\log(T-t)}{T-t}} \left( M_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right). \tag{47}$$

We next obtain a bound on  $\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \int_{\mathbf{a}^\top \boldsymbol{\lambda}_t^*}^{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t} (F_{\mathbf{a}}^r(v) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*)) dv$ . Note that for any  $\boldsymbol{\lambda}_1, \boldsymbol{\lambda}_2 \geq \mathbf{0}$ , we have

$$\int_{\mathbf{a}^\top \boldsymbol{\lambda}_1}^{\mathbf{a}^\top \boldsymbol{\lambda}_2} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \boldsymbol{\lambda}_2\}} \right) dv \geq 0,$$

which, by (41), yields

$$\begin{aligned}
& \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*, \mathbf{a}, r)^\top (\tilde{\boldsymbol{\lambda}}_t - \boldsymbol{\lambda}_t^*) \\
& \leq \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*, \mathbf{a}, r)^\top (\tilde{\boldsymbol{\lambda}}_t - \boldsymbol{\lambda}_t^*) + \mathbb{E}_{(\mathbf{a}, r) \sim F} \int_{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t}^{\mathbf{a}^\top \boldsymbol{\lambda}_t^*} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \boldsymbol{\lambda}_t^*\}} \right) dv \\
& = f_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t) - f_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*) \leq 0.
\end{aligned} \tag{48}$$

Lemma F.8 implies that with probability at least  $1 - \frac{3C \log(T-t)}{(T-t)^2}$ ,

$$\begin{aligned}
& \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*, \mathbf{a}, r)^\top (\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t) - \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*, \mathbf{a}_j, r_j)^\top (\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t) \\
& \quad + \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*, \mathbf{a}_j, r_j)^\top (\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t) \\
& \leq \tilde{C}_1 \sqrt{\frac{\log(T-t)}{T-t}} \left( M_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right) + \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*, \mathbf{a}_j, r_j)^\top (\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t) \\
& \leq \tilde{C}_1 \sqrt{\frac{\log(T-t)}{T-t}} \left( M_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right) + \frac{2m^2 \bar{A} \bar{r}}{(T-t) \underline{A}},
\end{aligned} \tag{49}$$

where the last inequality is by the second inequality in Lemma F.14 with  $\boldsymbol{\lambda} = \tilde{\boldsymbol{\lambda}}_t$ .

Therefore, with probability at least  $1 - \frac{3C \log(T-t)}{(T-t)^2}$ ,

$$0 \leq \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*, \mathbf{a}, r)^\top (\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t) \leq C_3 \sqrt{\frac{\log(T-t)}{T-t}} \left( M_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right), \tag{50}$$

where  $C_3 = \tilde{C}_1 + \frac{2m^2 \bar{A} \bar{r}}{\underline{A}}$ . By (48) and (50), we have with probability at least  $1 - \frac{3C \log(T-t)}{(T-t)^2}$ ,

$$\begin{aligned}
\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \int_{\mathbf{a}^\top \boldsymbol{\lambda}_t^*}^{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t} \left( F_{\mathbf{a}}^r(v) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right) dv &= \mathbb{E}_{(\mathbf{a}, r) \sim F} \int_{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t}^{\mathbf{a}^\top \boldsymbol{\lambda}_t^*} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \boldsymbol{\lambda}_t^*\}} \right) dv \\
&\leq \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*, \mathbf{a}, r)^\top (\boldsymbol{\lambda}_t^* - \tilde{\boldsymbol{\lambda}}_t) \\
&\leq C_3 \sqrt{\frac{\log(T-t)}{T-t}} \left( M_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right).
\end{aligned} \tag{51}$$

Combining (47) and (51), we obtain with probability at least  $1 - \frac{9C \log(T-t)}{(T-t)^2}$ ,

$$\begin{aligned}
& \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right) \left( \mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^* \right) \right] \\
& = \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \int_{\mathbf{a}^\top \boldsymbol{\lambda}_t^*}^{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t} \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(v) \right) dv + \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \int_{\mathbf{a}^\top \boldsymbol{\lambda}_t^*}^{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t} \left( F_{\mathbf{a}}^r(v) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*) \right) dv \\
& \leq (2\tilde{C}_1 + C_3) \sqrt{\frac{\log(T-t)}{T-t}} \left( M_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right).
\end{aligned} \tag{52}$$

Note that

$$M_1 = \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ (\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}_t^*)^2 \max \left( \bar{F}_a^r(\mathbf{a}^\top \boldsymbol{\lambda}_t^*), \bar{F}_a^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) \right) \right]}.$$

We have proven (20) regarding  $\boldsymbol{\lambda}_t^*$ .

We now turn to (21) that involves  $\bar{\boldsymbol{\lambda}}_t^*$ . The proof is very similar. Similar to (44), we have

$$\begin{aligned}
0 &\leq f_{t, \mathbf{b}_{t-1}}(\bar{\boldsymbol{\lambda}}_t^*) - f_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t) \\
&\leq f_{t, \mathbf{b}_{t-1}}(\bar{\boldsymbol{\lambda}}_t^*) - f_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t) - (g_{t, \mathbf{b}_{t-1} - \mathbf{a}_t}(\bar{\boldsymbol{\lambda}}_t^*) - g_{t, \mathbf{b}_{t-1} - \mathbf{a}_t}(\tilde{\boldsymbol{\lambda}}_t)) \\
&= \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t, \mathbf{a}, r)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) - \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1} - \mathbf{a}_t}(\tilde{\boldsymbol{\lambda}}_t, \mathbf{a}_j, r_j)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) \\
&+ \mathbb{E}_{(\mathbf{a}, r) \sim F} \int_{\mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^*}^{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^*\}} \right) dv - \frac{1}{T-t} \sum_{j=t+1}^T \int_{\mathbf{a}_j^\top \bar{\boldsymbol{\lambda}}_t^*}^{\mathbf{a}_j^\top \tilde{\boldsymbol{\lambda}}_t} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}_j^\top \bar{\boldsymbol{\lambda}}_t^*\}} \right) dv \\
&= \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t, \mathbf{a}, r)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) - \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t, \mathbf{a}_j, r_j)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) \\
&+ \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t, \mathbf{a}_j, r_j)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) - \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1} - \mathbf{a}_t}(\tilde{\boldsymbol{\lambda}}_t, \mathbf{a}_j, r_j)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) \\
&+ \mathbb{E}_{(\mathbf{a}, r) \sim F} \int_{\mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^*}^{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^*\}} \right) dv - \frac{1}{T-t} \sum_{j=t+1}^T \int_{\mathbf{a}_j^\top \bar{\boldsymbol{\lambda}}_t^*}^{\mathbf{a}_j^\top \tilde{\boldsymbol{\lambda}}_t} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}_j^\top \bar{\boldsymbol{\lambda}}_t^*\}} \right) dv \\
&= \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t, \mathbf{a}, r)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) - \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t, \mathbf{a}_j, r_j)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) \\
&+ \mathbb{E}_{(\mathbf{a}, r) \sim F} \int_{\mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^*}^{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^*\}} \right) dv - \frac{1}{T-t} \sum_{j=t+1}^T \int_{\mathbf{a}_j^\top \bar{\boldsymbol{\lambda}}_t^*}^{\mathbf{a}_j^\top \tilde{\boldsymbol{\lambda}}_t} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}_j^\top \bar{\boldsymbol{\lambda}}_t^*\}} \right) dv \\
&\quad + \frac{\mathbf{a}_t^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t)}{T-t} \\
&\leq 2\tilde{C}_1 \left( \bar{M}_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right) + \frac{2m\bar{A}\bar{r}}{\underline{A}(T-t)} \\
&\leq C_4 \sqrt{\frac{\log(T-t)}{T-t}} \left( \bar{M}_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right), \tag{53}
\end{aligned}$$

where the third inequality follows from Lemma F.8 and Lemma F.9 similar to (45), and that  $\frac{\mathbf{a}_t^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t)}{T-t} \leq \frac{2m\bar{A}\bar{r}}{\underline{A}(T-t)}$  since  $\|\mathbf{a}_t\|_\infty \leq \bar{A}$  and  $\|\bar{\boldsymbol{\lambda}}_t^*\|_\infty, \|\tilde{\boldsymbol{\lambda}}_t\|_\infty \leq \frac{\bar{r}}{\underline{A}}$  by Corollary F.13. Here  $C_4 = 2\tilde{C}_1 + \frac{2m\bar{A}\bar{r}}{\underline{A}}$ , and

$$\bar{M}_1 = \max \left\{ \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} (\mathbf{a}^\top (\tilde{\boldsymbol{\lambda}}_t - \bar{\boldsymbol{\lambda}}_t^*))^2 (1 - F_a^r(\mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^*))}, \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} (\mathbf{a}^\top (\tilde{\boldsymbol{\lambda}}_t - \bar{\boldsymbol{\lambda}}_t^*))^2 (1 - F_a^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t))} \right\}.$$

Now following nearly identical arguments as in (46), we derive a similar bound as (47), that

$$\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \int_{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t}^{\mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t} \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(v) \right) dv \leq C_4 \sqrt{\frac{\log(T-t)}{T-t}} \left( \bar{M}_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right) \quad (54)$$

with probability at least  $1 - \frac{6C \log(T-t)}{(T-t)^2}$ .

To obtain a bound on  $\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \int_{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t}^{\mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t} (F_{\mathbf{a}}^r(v) - F_{\mathbf{a}}^r(\mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t)) dv$ , we follow the argument similar to (48), which yields

$$\begin{aligned} & \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\bar{\boldsymbol{\lambda}}_t^*, \mathbf{a}, r)^\top (\tilde{\boldsymbol{\lambda}}_t - \bar{\boldsymbol{\lambda}}_t^*) \\ & \leq \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\bar{\boldsymbol{\lambda}}_t^*, \mathbf{a}, r)^\top (\tilde{\boldsymbol{\lambda}}_t - \bar{\boldsymbol{\lambda}}_t^*) + \mathbb{E}_{(\mathbf{a}, r) \sim F} \int_{\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t}^{\mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \bar{\boldsymbol{\lambda}}_t^*\}} \right) dv \\ & = f_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t) - f_{t, \mathbf{b}_{t-1}}(\bar{\boldsymbol{\lambda}}_t^*) \leq 0, \end{aligned} \quad (55)$$

where we used the fact that  $\tilde{\boldsymbol{\lambda}}_t$  is the minimizer of  $f_{t, \mathbf{b}_{t-1}}(\cdot)$ . Lemma F.8 implies that with probability at least  $1 - \frac{3C \log(T-t)}{(T-t)^2}$ ,

$$\begin{aligned} & \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\bar{\boldsymbol{\lambda}}_t^*, \mathbf{a}, r)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) - \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1} - \mathbf{a}_t}(\bar{\boldsymbol{\lambda}}_t^*, \mathbf{a}_j, r_j)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) \\ & + \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1} - \mathbf{a}_t}(\bar{\boldsymbol{\lambda}}_t^*, \mathbf{a}_j, r_j)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) \\ & = \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\bar{\boldsymbol{\lambda}}_t^*, \mathbf{a}, r)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) - \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1}}(\bar{\boldsymbol{\lambda}}_t^*, \mathbf{a}_j, r_j)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) \\ & + \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1}}(\bar{\boldsymbol{\lambda}}_t^*, \mathbf{a}_j, r_j)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) - \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1} - \mathbf{a}_t}(\bar{\boldsymbol{\lambda}}_t^*, \mathbf{a}_j, r_j)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) \\ & + \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1} - \mathbf{a}_t}(\bar{\boldsymbol{\lambda}}_t^*, \mathbf{a}_j, r_j)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) \\ & \leq \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\bar{\boldsymbol{\lambda}}_t^*, \mathbf{a}, r)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) - \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1}}(\bar{\boldsymbol{\lambda}}_t^*, \mathbf{a}_j, r_j)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) \\ & + \frac{\mathbf{a}_t^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t)}{T-t} + \frac{2m^2 \bar{A} \bar{r}}{\underline{A}(T-t)} \\ & \leq \left( \tilde{C}_1 + \frac{2m \bar{A} \bar{r}}{\underline{A}} + \frac{2m^2 \bar{A} \bar{r}}{\underline{A}} \right) \sqrt{\frac{\log(T-t)}{T-t}} \left( \bar{M}_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right), \end{aligned} \quad (56)$$

where the first inequality is by the third inequality in Lemma F.14 with  $\boldsymbol{\lambda} = \tilde{\boldsymbol{\lambda}}_t$ , and the last inequality is because  $\frac{\mathbf{a}_t^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t)}{T-t} \leq \frac{2m \bar{A} \bar{r}}{\underline{A}(T-t)}$ .

Combining (55) with (56), we have with probability at least  $1 - \frac{3 \log(T-t)}{(T-t)^2}$ ,

$$0 \leq \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\bar{\boldsymbol{\lambda}}_t^*, \mathbf{a}, r)^\top (\bar{\boldsymbol{\lambda}}_t^* - \tilde{\boldsymbol{\lambda}}_t) \leq C_5 \sqrt{\frac{\log(T-t)}{T-t}} \left( \bar{M}_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right), \quad (57)$$

where  $C_5 = \tilde{C}_1 + \frac{2m\bar{A}\bar{r}}{A} + \frac{2m^2\bar{A}\bar{r}}{A}$ . By (55) and (57), we have with probability at least  $1 - \frac{3C\log(T-t)}{(T-t)^2}$ ,

$$\begin{aligned} \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \int_{\mathbf{a}^\top \tilde{\lambda}_t^*}^{\mathbf{a}^\top \tilde{\lambda}_t} \left( F_{\mathbf{a}}^r(v) - F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\lambda}_t^*) \right) dv &= \mathbb{E}_{(\mathbf{a}, r) \sim F} \int_{\mathbf{a}^\top \tilde{\lambda}_t}^{\mathbf{a}^\top \tilde{\lambda}_t^*} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \tilde{\lambda}_t^*\}} \right) dv \\ &\leq \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\tilde{\lambda}_t^*, \mathbf{a}, r)^\top (\tilde{\lambda}_t^* - \tilde{\lambda}_t) \\ &\leq C_5 \sqrt{\frac{\log(T-t)}{T-t}} \left( \bar{M}_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right). \end{aligned} \quad (58)$$

Combining (54) and (58), we obtain with probability at least  $1 - \frac{9C\log(T-t)}{(T-t)^2}$ ,

$$\begin{aligned} &\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\lambda}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\lambda}_t^*) \right) \left( \mathbf{a}^\top \tilde{\lambda}_t - \mathbf{a}^\top \tilde{\lambda}_t^* \right) \right] \\ &= \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \int_{\mathbf{a}^\top \tilde{\lambda}_t^*}^{\mathbf{a}^\top \tilde{\lambda}_t} \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\lambda}_t) - F_{\mathbf{a}}^r(v) \right) dv + \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \int_{\mathbf{a}^\top \tilde{\lambda}_t}^{\mathbf{a}^\top \tilde{\lambda}_t^*} \left( F_{\mathbf{a}}^r(v) - F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\lambda}_t^*) \right) dv \\ &\leq (C_4 + C_5) \sqrt{\frac{\log(T-t)}{T-t}} \left( \bar{M}_1 + \sqrt{\frac{\log(T-t)}{T-t}} \right), \end{aligned} \quad (59)$$

where we note

$$\bar{M}_1 = \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left( \mathbf{a}^\top \tilde{\lambda}_t - \mathbf{a}^\top \tilde{\lambda}_t^* \right)^2 \max \left( \bar{F}_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\lambda}_t^*), \bar{F}_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\lambda}_t) \right) \right]},$$

$C_4 = 2\tilde{C}_1 + \frac{2m\bar{A}\bar{r}}{A}$ , and  $C_5 = \tilde{C}_1 + \frac{2m\bar{A}\bar{r}}{A} + \frac{2m^2\bar{A}\bar{r}}{A}$ . We have proven (21) regarding  $\tilde{\lambda}_t^*$ .

Now combining (20) and (21), we complete the proof of Lemma F.2. Q.E.D.

## F.4 Auxiliary Lemmas and Their Proof

This section contains all auxiliary lemmas that will be used in constructing our ultimate proof of the CE regret analysis. In Section F.4.1 we state and prove several results in LP theory. In Section F.4.2, we state and prove concentration results using the tools such as the peeling argument from the empirical processes theory.

### F.4.1 Properties of the Multi-Knapsack LP

Recall that the multi-knapsack LP takes the following form

$$\begin{aligned} \max \quad & \sum_{j=t+1}^T r_j x_j \\ \text{s.t.} \quad & \sum_{j=t+1}^T a_{ij} x_j \leq c_i, \quad i = 1, \dots, m, \\ & x_j \in [0, 1], \quad j = t+1, \dots, T, \end{aligned} \quad (\mathcal{P})$$

and

$$\min \quad \sum_{i=1}^m c_i \lambda_i + \sum_{j=t+1}^T \xi_j \quad (\mathcal{D-1})$$



$$\begin{aligned} \text{s.t. } \quad & \sum_{i=1}^m a_{ij} \lambda_i + \xi_j \geq r_j, \quad j = t+1, \dots, T, \\ & \lambda_i, \xi_j \geq 0, \quad j = t+1, \dots, T; \quad i = 1, \dots, m, \end{aligned}$$

We call Problem ( $\mathcal{P}$ ) the primal multi-knapsack LP and Problem ( $\mathcal{D}$ -1) the dual multi-knapsack LP that is specified by coefficients  $t, \mathbf{c}$  and  $\mathcal{I}_t = \{(\mathbf{a}_j, r_j)\}_{j=t+1}^T$ . Denote by  $V_{t, \mathbf{c}}^{\text{off}}(\mathcal{I}_t)$  their optimal value, which by strong duality coincides. Li and Ye (2022) observes that the dual problem has the following alternative, simple form.

**Lemma F.3.** *The dual problem ( $\mathcal{D}$ -1) has the alternative, simple form*

$$\min_{\lambda \in \mathbb{R}_{\geq 0}^m} \mathbf{c}^\top \lambda + \sum_{j=t+1}^T (r_j - \mathbf{a}_j^\top \lambda)^+. \quad (\mathcal{D}\text{-2})$$

*Proof.* Observe that to reach optimality, we should set  $\xi_j = 0$  if  $\sum_{i=1}^m a_{ij} \lambda_i \geq r_j$  and  $\xi_j = r_j - \sum_{i=1}^m a_{ij} \lambda_i$  otherwise, which is the same as  $\xi_j = (r_j - \sum_{i=1}^m a_{ij} \lambda_i)^+$ . Plugging into the objective function in Problem ( $\mathcal{D}$ -1) and eliminating the variables  $\xi_j$  completes the proof. Q.E.D.

**Lemma F.4 (Complementary Slackness).** *Suppose  $\mathbf{x}^*$  and  $\lambda^*$  are optimal solutions to Problem ( $\mathcal{P}$ ) and Problem ( $\mathcal{D}$ -2), respectively. Then they satisfy the complementary slackness property*

$$\begin{aligned} r_j > \mathbf{a}_j^\top \lambda & \Rightarrow x_j^* = 1, \\ r_j < \mathbf{a}_j^\top \lambda & \Rightarrow x_j^* = 0, \text{ for all } j = t+1, \dots, T. \end{aligned}$$

*Proof.* Suppose  $(\lambda^*, \xi^*)$  is an optimal solution to Problem ( $\mathcal{D}$ -1). By LP complementary slackness, we have for  $i = 1, \dots, m$  and  $j = t+1, \dots, T$ ,

$$\begin{aligned} (1 - x_j^*) \xi_j^* &= 0, \\ \left( r_j - \sum_{i=1}^m a_{ij} \lambda_i^* - \xi_j^* \right) x_j^* &= 0, \\ \left( c_i - \sum_{j=t+1}^T a_{ij} x_j^* \right) \lambda_i^* &= 0. \end{aligned}$$

If for some  $j$  we have  $r_j > \mathbf{a}_j^\top \lambda$ , then by the dual constraint of Problem ( $\mathcal{D}$ -1), we have  $\xi_j^* \geq r_j - \mathbf{a}_j^\top \lambda > 0$ . Then by the first complementary slackness equality, we must have  $1 - x_j^* = 0$ , completing the proof of the first statement.

Now if on the other hand, if we have  $r_j < \mathbf{a}_j^\top \lambda$ , then since  $\xi_j^* \geq 0$ , we have  $r_j - \sum_{i=1}^m a_{ij} \lambda_i^* - \xi_j^* \leq r_j - \sum_{i=1}^m a_{ij} \lambda_i^* < 0$ . By the second complementary slackness equality, we must have  $x_j^* = 0$ , again completing the proof of the second statement. Q.E.D.

**Lemma F.5.** *Suppose  $\mathbf{x}^*$  is an optimal basic feasible solution to Problem ( $\mathcal{P}$ ), then  $\mathbf{x}^*$  has at most  $m$  basic variables that are fractional.*

*Proof.* Refer to Meanti et al. (1990) Lemma 2.1. Q.E.D.

**Lemma F.6.** *Suppose  $\{x_j^*\}_{j=t}^T$  is an optimal solution to the primal multi-knapsack LP specified by  $t-1, \mathbf{b}$  and  $\mathcal{I}_{t-1}$ , then*

$$V_{t-1, \mathbf{b}}^{\text{off}}(\mathcal{I}_{t-1}) = r_t x_t^* + V_{t, \mathbf{b} - \mathbf{a}_t x_t^*}^{\text{off}}(\mathcal{I}_t) = \max_{x \in [0, 1]} (r_t x + V_{t, \mathbf{b} - \mathbf{a}_t x}^{\text{off}}(\mathcal{I}_t)).$$

*Proof.* By definition,  $V_{t-1, \mathbf{b}}^{\text{off}}(\mathcal{I}_{t-1}) = \sum_{j=t}^T r_j x_j^* = r_t x_t^* + \sum_{j=t+1}^T r_j x_j^*$ . On the one hand,  $\{x_j^*\}_{j=t}^T$  is feasible, concretely,  $\sum_{j=t}^T a_{ij} x_j^* \leq c_i$ ,  $i = 1, \dots, m$ , and  $x_j^* \in [0, 1]$ ,  $j = t, \dots, T$ . From the above we derive  $\sum_{j=t+1}^T a_{ij} x_j^* \leq c_i - a_{it} x_t^*$ ,  $i = 1, \dots, m$  and  $x_j^* \in [0, 1]$ ,  $j = t+1, \dots, T$ . In other words,  $\{x_j^*\}_{j=t+1}^T$  is feasible to the LP specified by  $t-1, \mathbf{b} - \mathbf{a}_t x_t^*$  and  $\mathcal{I}_t$ . Thus  $\sum_{j=t+1}^T r_j x_j^* \leq V_{t, \mathbf{b} - \mathbf{a}_t x_t^*}^{\text{off}}$  and we conclude that

$$V_{t-1, \mathbf{b}}^{\text{off}}(\mathcal{I}_{t-1}) \leq r_t x_t^* + V_{t, \mathbf{b} - \mathbf{a}_t x_t^*}^{\text{off}}(\mathcal{I}_t) \leq \max_{x \in [0, 1]} \left( r_t x + V_{t, \mathbf{b} - \mathbf{a}_t x}^{\text{off}}(\mathcal{I}_t) \right).$$

On the other hand, consider an arbitrary optimal solution  $\{\tilde{x}_j^*\}_{j=t+1}^T$  to the multi-knapsack LP specified by  $t, \mathbf{b} - \mathbf{a}_t \tilde{x}_t^*$  and  $\mathcal{I}_t$ , where  $\tilde{x}_t^* = \arg \max_{x \in [0, 1]} \left( r_t x + V_{t, \mathbf{b} - \mathbf{a}_t x}^{\text{off}}(\mathcal{I}_t) \right)$ . We have  $V_{t, \mathbf{b} - \mathbf{a}_t \tilde{x}_t^*}^{\text{off}}(\mathcal{I}_t) = \sum_{j=t+1}^T r_j \tilde{x}_j^*$ . Notice that  $(\tilde{x}_t^*, \tilde{x}_{t+1}^*, \dots, \tilde{x}_T^*)$  is a feasible solution to the original LP specified by  $t-1, \mathbf{b}$  and  $\mathcal{I}_{t-1}$ . We thus have

$$V_{t-1, \mathbf{b}}^{\text{off}}(\mathcal{I}_{t-1}) \geq r_t \tilde{x}_t^* + V_{t, \mathbf{b} - \mathbf{a}_t \tilde{x}_t^*}^{\text{off}}(\mathcal{I}_t) = \max_{x \in [0, 1]} \left( r_t x + V_{t, \mathbf{b} - \mathbf{a}_t x}^{\text{off}}(\mathcal{I}_t) \right).$$

Combining the inequalities completes the proof. Q.E.D.

#### F.4.2 Concentration and Empirical Processes

We borrow techniques from the empirical processes theory to establish concentration results that will in turn be used in proving our main theorems.

**Lemma F.7.** *Suppose that  $X$  is a random vector following distribution  $P$  supported on  $\mathcal{X}$ .  $\mathcal{G}$  is a function class such that  $\sup_{g \in \mathcal{G}} \|g\|_{L_\infty(\mathcal{X})} \leq K$ ,  $\sup_{g \in \mathcal{G}} \|g\|_P^2 \leq R^2$ . Take  $a, C_0, C_1$  satisfying*

$$a \leq C_1 \sqrt{n} R^2 / (2K), \quad (60)$$

$$a \leq 8\sqrt{2n}R, \quad (61)$$

$$a \geq C_0 \max \left\{ \int_0^{\sqrt{2}R} H_B(u/\sqrt{2}, \mathcal{G}, P)^{1/2} du, \sqrt{2}R \right\}, \quad (62)$$

$$C_0^2 \geq C^2(C_1 + 1), \quad (63)$$

where  $C$  is a universal constant. Then

$$\mathbb{P} \left( \sup_{g \in \mathcal{G}} \sqrt{n} \left| \frac{1}{n} \sum_{j=1}^n g(X_j) - \mathbb{E}_{X \sim P} g(X) \right| \geq a \right) \leq C \exp \left( -\frac{a^2}{2C^2(C_1 + 1)R^2} \right).$$

*Proof of Lemma F.7.* It suffices to show that the conditions of Theorem 5.11 of Geer (2000) are satisfied. Following the notation in Geer (2000), we use

$$\rho_K(g)^2 = 2K^2 \int \left( e^{|g|/K} - 1 - |g|/K \right) dP,$$

and let  $\mathcal{H}_{B,K}(\delta, \mathcal{G}, P)$  be the generalized entropy with bracketing (see Definition 5.1 of Geer (2000)). Lemmas 5.8 and 5.10 of Geer (2000) imply that if  $\sup_{g \in \mathcal{G}} \|g\|_{L_\infty} \leq K_1/4 < K_1/2$ , then  $\rho_{K_1}(g)^2 \leq 2R^2 = R_1^2$  with  $R_1 = \sqrt{2}R$ , and  $\mathcal{H}_{B,K_1}(\delta, \mathcal{G}, P) \leq H_B(\delta/\sqrt{2}, \mathcal{G}, P)$  for all  $\delta > 0$ . Let  $K_1 = 4K$ . Therefore, conditions of Theorem 5.11 of Geer (2000) becomes

$$a \leq C_1 \sqrt{n} R_1^2 / K_1 = C_1 \sqrt{n} R^2 / (2K),$$

$$\begin{aligned}
a &\leq 8\sqrt{n}R_1 = 8\sqrt{2n}R, \\
a &\geq C_0 \max \left\{ \int_{a/(2^6\sqrt{n})}^{R_1} \mathcal{H}_{B,K_1}(u, \mathcal{G}, P)^{1/2} du, R_1 \right\}, \\
C_0^2 &\geq C^2(C_1 + 1),
\end{aligned}$$

where the third inequality is implied by

$$a \geq C_0 \max \left\{ \int_0^{\sqrt{2}R} H_B(u/\sqrt{2}, \mathcal{G}, P)^{1/2} du, \sqrt{2}R \right\} \geq C_0 \max \left\{ \int_{a/(2^6\sqrt{n})}^{\sqrt{2}R} H_B(u/\sqrt{2}, \mathcal{G}, P)^{1/2} du, \sqrt{2}R \right\}.$$

The results of Theorem 5.11 of Geer (2000) becomes

$$\mathbb{P} \left( \sup_{g \in \mathcal{G}} \sqrt{n} \left| \frac{1}{n} \sum_{j=1}^n g(X_j) - \mathbb{E}_{X \sim P} g(X) \right| \geq a \right) \leq C \exp \left( -\frac{a^2}{C^2(C_1 + 1)R_1^2} \right),$$

which equals to  $C \exp \left( -\frac{a^2}{2C^2(C_1+1)R^2} \right)$ . This finishes the proof. Q.E.D.

We now apply Lemma F.7 to obtain two concentration results that are key components of our proof of Lemma 5.2.

**Lemma F.8.** *With probability at least  $1 - \frac{3C \log(T-t)}{(T-t)^2}$ ,*

$$\begin{aligned}
&\left| \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}}(\boldsymbol{\lambda}_1, \mathbf{a}, r)^\top (\boldsymbol{\lambda}_2 - \boldsymbol{\lambda}_1) - \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}}(\boldsymbol{\lambda}_1, \mathbf{a}_j, r_j)^\top (\boldsymbol{\lambda}_2 - \boldsymbol{\lambda}_1) \right| \\
&\leq \tilde{C} \sqrt{\frac{\log(T-t)}{T-t}} \left( \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \{(\mathbf{a}^\top (\boldsymbol{\lambda}_2 - \boldsymbol{\lambda}_1))^2 (1 - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_1))\}} + \sqrt{\frac{\log(T-t)}{T-t}} \right)
\end{aligned}$$

holds for any  $\mathbf{b} \in \mathbb{R}_{\geq 0}^m$ ,  $\boldsymbol{\lambda}_1, \boldsymbol{\lambda}_2 \in \Omega$ , as long as  $T-t \geq e^{\frac{\tilde{C}^2}{32}+1}$ , where

$$\tilde{C} = 2 \left( \frac{4m\bar{r}\bar{A}}{\underline{A}} + 1 \right) \left( 10C^2 \left( 1 + 2 \left( \frac{8m}{\min\{\nu/2, 1\}} \right)^{1/2} \right)^2 \left( c_{\nu}^{\frac{1}{2}} \frac{\bar{r}}{\underline{A}} (2m\bar{A})^{\frac{\nu}{2}} + 3 \right)^2 + 2 \right),$$

and  $C$  is a universal constant.

*Proof of Lemma F.8.* Recall that

$$\phi_{t, \mathbf{b}}(\boldsymbol{\lambda}, \mathbf{a}, r) = \frac{\partial h_{t, \mathbf{b}}(\boldsymbol{\lambda}, \mathbf{a})}{\partial \boldsymbol{\lambda}} = \frac{1}{T-t} \mathbf{b} - \mathbf{a} \mathbb{I}_{\{r > \mathbf{a}^\top \boldsymbol{\lambda}\}}.$$

Therefore, we have

$$\begin{aligned}
&\mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}}(\boldsymbol{\lambda}_1, \mathbf{a}, r)^\top (\boldsymbol{\lambda}_2 - \boldsymbol{\lambda}_1) - \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}}(\boldsymbol{\lambda}_1, \mathbf{a}_j, r_j)^\top (\boldsymbol{\lambda}_2 - \boldsymbol{\lambda}_1) \\
&= \mathbb{E}_{(\mathbf{a}, r) \sim F} \mathbf{a}^\top (\boldsymbol{\lambda}_2 - \boldsymbol{\lambda}_1) \mathbb{I}_{\{r > \mathbf{a}^\top \boldsymbol{\lambda}_1\}} - \frac{1}{T-t} \sum_{j=t+1}^T \mathbf{a}_j^\top (\boldsymbol{\lambda}_2 - \boldsymbol{\lambda}_1) \mathbb{I}_{\{r_j > \mathbf{a}_j^\top \boldsymbol{\lambda}_1\}}.
\end{aligned}$$

Define the function class  $\mathcal{G}$  parameterized by  $\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3, \mathbf{p}_4$ :

$$\mathcal{G} = \left\{ g : g(\mathbf{a}, r) = \frac{\underline{A}}{4m\bar{r}\bar{A}} \left( \mathbf{a}^\top \mathbf{p}_4 \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{p}_2\}} - \mathbf{a}^\top \mathbf{p}_1 \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{p}_3\}} \right), \mathbf{p}_j \in \Omega, j = 1, 2, 3, 4 \right\}.$$

For all  $g \in \mathcal{G}$  and  $(\mathbf{a}, r) \in \mathcal{S}$ , we have

$$\begin{aligned} |g(\mathbf{a}, r)| &= \left| \frac{\underline{A}}{4m\bar{r}\bar{A}} \left( \mathbf{a}^\top \mathbf{p}_4 \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{p}_2\}} - \mathbf{a}^\top \mathbf{p}_1 \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{p}_3\}} \right) \right| \\ &\leq \frac{\underline{A}}{4m\bar{r}\bar{A}} \left( \left| \mathbf{a}^\top \mathbf{p}_4 \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{p}_2\}} \right| + \left| \mathbf{a}^\top \mathbf{p}_1 \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{p}_3\}} \right| \right) \\ &\leq \frac{\underline{A}}{4m\bar{r}\bar{A}} \left( \left| \mathbf{a}^\top \mathbf{p}_4 \right| + \left| \mathbf{a}^\top \mathbf{p}_1 \right| \right) \\ &\leq \frac{\underline{A}}{4m\bar{r}\bar{A}} \times 2m\bar{A} \frac{\bar{r}}{\underline{A}} = \frac{1}{2} < 1, \end{aligned}$$

where in the final inequality we use  $\|\boldsymbol{\lambda}\|_\infty \leq \frac{\bar{r}}{\underline{A}}$  for any  $\boldsymbol{\lambda} \in \Omega$  and  $\|\mathbf{a}\|_\infty \leq \bar{A}, r \leq \bar{r}$  for any  $(\mathbf{a}, r) \in \mathcal{S}$ . In other words,  $\|g\|_\infty = \|g\|_{L_\infty(\mathcal{S})} \leq \frac{1}{2} < 1$  for any  $\mathbf{p}_j \in \Omega, j = 1, 2, 3, 4$ . Our plan is to approximate  $\mathcal{G}$  by its  $\delta$ -net. For any  $\delta > 0$ , consider the grid points set  $\Omega_0 \subseteq [0, \frac{\bar{r}}{\underline{A}}]^{4m}$ , where each element of each point  $\mathbf{q} = (\mathbf{q}^{(1)}, \mathbf{q}^{(2)}, \mathbf{q}^{(3)}, \mathbf{q}^{(4)}) \in \Omega_0$  has the form  $k\delta$ , with  $k = 0, 1, \dots, \lfloor \frac{\bar{r}}{\delta \underline{A}} \rfloor + 1$ , and  $\mathbf{q}^{(1)}, \mathbf{q}^{(2)}, \mathbf{q}^{(3)}, \mathbf{q}^{(4)} \in \Omega$ . Thus, we have  $\text{card}(\Omega_0) \leq (2 + \frac{\bar{r}}{\delta \underline{A}})^{4m}$ . Let  $N_0 = (2 + \frac{\bar{r}}{\delta \underline{A}})^{4m}$ . Let

$$\mathcal{G}_0 = \left\{ g : g(\mathbf{a}, r) = \frac{\underline{A}}{4m\bar{r}\bar{A}} \left( \mathbf{a}^\top \mathbf{q}^{(4)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}^{(2)}\}} - \mathbf{a}^\top \mathbf{q}^{(1)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}^{(3)}\}} \right), \mathbf{q}^{(j)} \in \Omega_0, j = 1, 2, 3, 4 \right\}.$$

For each  $g \in \mathcal{G}$ , we are going to find functions  $g_L, g_U \in \mathcal{G}_0$  such that  $g_L \leq g \leq g_U$  with  $\|g_L - g_U\|_F$  bounded by a function of  $\delta$  to be determined later, and we recall that  $\|\cdot\|_F$  is the  $L_2(F)$  norm with respect to distribution  $F$  (of  $(\mathbf{a}, r)$ ).

Since  $g \in \mathcal{G}$ , it has the form

$$g(\mathbf{a}, r) = \frac{\underline{A}}{4m\bar{r}\bar{A}} \left( \mathbf{a}^\top \mathbf{p}_4 \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{p}_2\}} - \mathbf{a}^\top \mathbf{p}_1 \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{p}_3\}} \right).$$

We choose  $\mathbf{q}_U = (\mathbf{q}_U^{(1)}, \mathbf{q}_U^{(2)}, \mathbf{q}_U^{(3)}, \mathbf{q}_U^{(4)})$  such that  $\mathbf{q}_U^{(j)} \leq \mathbf{p}_j < \mathbf{q}_U^{(j)} + \delta \mathbf{1}$ , for  $j = 1, 2$ , and  $\max\{\mathbf{q}_U^{(i)} - \delta \mathbf{1}, \mathbf{0}\} < \mathbf{p}_i \leq \mathbf{q}_U^{(i)}$ , for  $i = 3, 4$ , where the maximum is taken elementwisely. We also choose  $\mathbf{q}_L = (\mathbf{q}_L^{(1)}, \mathbf{q}_L^{(2)}, \mathbf{q}_L^{(3)}, \mathbf{q}_L^{(4)})$  such that  $\mathbf{q}_L^{(i)} \leq \mathbf{p}_i < \mathbf{q}_L^{(i)} + \delta \mathbf{1}$ , and  $\max\{\mathbf{q}_L^{(j)} - \delta \mathbf{1}, \mathbf{0}\} < \mathbf{p}_j \leq \mathbf{q}_L^{(j)}$ , for  $j = 1, 2, i = 3, 4$ .

Let

$$\begin{aligned} g_U(\mathbf{a}, r) &= \frac{\underline{A}}{4m\bar{r}\bar{A}} \left( \mathbf{a}^\top \mathbf{q}_U^{(4)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_U^{(2)}\}} - \mathbf{a}^\top \mathbf{q}_U^{(1)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_U^{(3)}\}} \right), \\ g_L(\mathbf{a}, r) &= \frac{\underline{A}}{4m\bar{r}\bar{A}} \left( \mathbf{a}^\top \mathbf{q}_L^{(4)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_L^{(2)}\}} - \mathbf{a}^\top \mathbf{q}_L^{(1)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_L^{(3)}\}} \right). \end{aligned}$$

By monotonicity and non-negativity of  $\mathbf{a}$ ,  $g_L(\mathbf{a}, r) \leq g(\mathbf{a}, r) \leq g_U(\mathbf{a}, r)$  for all  $(\mathbf{a}, r) \in \mathcal{S}$  and any  $\mathbf{p}_j \in \Omega, j = 1, 2, 3, 4$ . We next bound  $\|g_U - g_L\|_F$ . By definition

$$\|g_U - g_L\|_F^2 = \left( \frac{\underline{A}}{4m\bar{r}\bar{A}} \right)^2 \mathbb{E}_{(\mathbf{a}, r) \sim F} \left\{ (J_1(\mathbf{a}, r) - J_2(\mathbf{a}, r))^2 \right\}$$

$$\leq 2 \left( \frac{\underline{A}}{4m\bar{r}\bar{A}} \right)^2 \left( \mathbb{E}_{(\mathbf{a},r) \sim F} J_1(\mathbf{a}, r)^2 + \mathbb{E}_{(\mathbf{a},r) \sim F} J_2(\mathbf{a}, r)^2 \right),$$

where

$$\begin{aligned} J_1(\mathbf{a}, r) &= \mathbf{a}^\top \mathbf{q}_U^{(4)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_U^{(2)}\}} - \mathbf{a}^\top \mathbf{q}_L^{(4)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_L^{(2)}\}}, \\ J_2(\mathbf{a}, r) &= \mathbf{a}^\top \mathbf{q}_U^{(1)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_U^{(3)}\}} - \mathbf{a}^\top \mathbf{q}_L^{(1)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_L^{(3)}\}}. \end{aligned}$$

For  $J_1(\mathbf{a}, r)$  we have

$$\begin{aligned} & \mathbb{E}_{(\mathbf{a},r) \sim F} J_1(\mathbf{a}, r)^2 \\ &= \mathbb{E}_{(\mathbf{a},r) \sim F} \left( \mathbf{a}^\top \mathbf{q}_U^{(4)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_U^{(2)}\}} - \mathbf{a}^\top \mathbf{q}_U^{(4)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_L^{(2)}\}} + \mathbf{a}^\top \mathbf{q}_U^{(4)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_L^{(2)}\}} - \mathbf{a}^\top \mathbf{q}_L^{(4)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_L^{(2)}\}} \right)^2 \\ &\leq 2 \mathbb{E}_{(\mathbf{a},r) \sim F} \left( \left( \mathbf{a}^\top \mathbf{q}_U^{(4)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_U^{(2)}\}} - \mathbf{a}^\top \mathbf{q}_U^{(4)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_L^{(2)}\}} \right)^2 + \left( \mathbf{a}^\top \mathbf{q}_U^{(4)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_L^{(2)}\}} - \mathbf{a}^\top \mathbf{q}_L^{(4)} \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_L^{(2)}\}} \right)^2 \right) \\ &= 2 \mathbb{E}_{(\mathbf{a},r) \sim F} (J_{11}(\mathbf{a}, r)^2 + J_{12}(\mathbf{a}, r)^2). \end{aligned}$$

By our choice of  $\mathbf{q}_L$  and  $\mathbf{q}_U$ ,  $\|\mathbf{q}_L - \mathbf{q}_U\|_\infty \leq \|\mathbf{q}_L - (\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3, \mathbf{p}_4)\|_\infty + \|\mathbf{q}_U - (\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3, \mathbf{p}_4)\|_\infty \leq 2\delta$ . For  $J_{11}(\mathbf{a}, r)$ , we have

$$\begin{aligned} \mathbb{E}_{(\mathbf{a},r) \sim F} J_{11}(\mathbf{a}, r)^2 &\leq \left( m\bar{A} \frac{\bar{r}}{\underline{A}} \right)^2 \mathbb{E}_{(\mathbf{a},r) \sim F} \left( \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_U^{(2)}\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_L^{(2)}\}} \right)^2 \\ &= \left( m\bar{A} \frac{\bar{r}}{\underline{A}} \right)^2 \mathbb{E}_{(\mathbf{a},r) \sim F} \left| \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_U^{(2)}\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_L^{(2)}\}} \right| \\ &= \left( m\bar{A} \frac{\bar{r}}{\underline{A}} \right)^2 \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left| F_{\mathbf{a}}^r(\mathbf{a}^\top \mathbf{q}_L^{(2)}) - F_{\mathbf{a}}^r(\mathbf{a}^\top \mathbf{q}_U^{(2)}) \right| \\ &\leq c_\nu \left( m\bar{A} \frac{\bar{r}}{\underline{A}} \right)^2 \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} |\mathbf{a}^\top \mathbf{q}_L^{(2)} - \mathbf{a}^\top \mathbf{q}_U^{(2)}|^\nu \quad (\text{under either Assumptions}) \\ &\leq c_\nu \left( m\bar{A} \frac{\bar{r}}{\underline{A}} \right)^2 (m\bar{A})^\nu 2^\nu \delta^\nu, \end{aligned}$$

where in the last equality we use  $\mathbf{a} \in [\underline{A}, \bar{A}]^m$  and  $\|\mathbf{q}_U - \mathbf{q}_L\|_\infty \leq 2\delta$ . Now consider the second term  $J_{12}(\mathbf{a}, r)$ , we have

$$\mathbb{E}_{(\mathbf{a},r) \sim F} J_{12}(\mathbf{a}, r)^2 \leq \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left( \mathbf{a}^\top \mathbf{q}_U^{(4)} - \mathbf{a}^\top \mathbf{q}_L^{(4)} \right)^2 \leq 4m^2 \bar{A}^2 \delta^2.$$

Therefore, the term  $\mathbb{E}_{(\mathbf{a},r) \sim F} J_1(a, r)^2$  can be bounded by

$$2 \left( c_\nu \left( m\bar{A} \frac{\bar{r}}{\underline{A}} \right)^2 (m\bar{A})^\nu 2^\nu \delta^\nu + 4m^2 \bar{A}^2 \delta^2 \right).$$

The term  $\mathbb{E}_{(\mathbf{a},r) \sim F} J_2(a, r)^2$  can be bounded nearly identically. Combining the above, we have

$$\|g_U - g_L\|_F^2 \leq 2 \left( \frac{\underline{A}}{4m\bar{r}\bar{A}} \right)^2 \left( \mathbb{E}_{(\mathbf{a},r) \sim F} J_1(\mathbf{a}, r)^2 + \mathbb{E}_{(\mathbf{a},r) \sim F} J_2(\mathbf{a}, r)^2 \right)$$

$$\begin{aligned}
&\leq 4 \left( \frac{\underline{A}}{4m\bar{r}\bar{A}} \right)^2 \left( 2c_\nu \left( m\bar{A} \frac{\bar{r}}{\underline{A}} \right)^2 (m\bar{A})^\nu 2^\nu \delta^\nu + 8m^2 \bar{A}^2 \delta^2 \right) \\
&\leq c_\nu (2m\bar{A})^\nu \delta^\nu + \frac{2\underline{A}^2}{\bar{r}^2} \delta^2 \leq \left( c_\nu^{\frac{1}{2}} (2m\bar{A})^{\frac{\nu}{2}} + \frac{2\underline{A}}{\bar{r}} \right)^2 \delta^{\min\{\nu, 2\}}.
\end{aligned}$$

As a result,  $\|g_U - g_L\|_F \leq \left( c_\nu^{\frac{1}{2}} (2m\bar{A})^{\frac{\nu}{2}} + \frac{2\underline{A}}{\bar{r}} \right) \delta^{\min\{\nu/2, 1\}}$ . Consider the set of function pairs

$$\{(g_L(\mathbf{q}_1), g_U(\mathbf{q}_2)), \mathbf{q}_1, \mathbf{q}_2 \in \Omega_0\}.$$

The cardinality of this set is  $\text{card}(\Omega_0)^2 \leq N_0^2$ . By the previous argument, this set of function pairs is enough to  $\delta'$ -cover  $\mathcal{G}$  where  $\delta' = \left( c_\nu^{\frac{1}{2}} (2m\bar{A})^{\frac{\nu}{2}} + \frac{2\underline{A}}{\bar{r}} \right) \delta^{\min\{\nu/2, 1\}}$  in the following precise sense. For any  $g \in \mathcal{G}$ , we can find a pair of  $\mathbf{q}_L$  and  $\mathbf{q}_U$  in  $\Omega_0$  (given precisely earlier), such that the associated  $g_L$  and  $g_U$  satisfy  $g_L \leq g \leq g_U$ , and furthermore  $\|g_U - g_L\|_F \leq \left( c_\nu^{\frac{1}{2}} (2m\bar{A})^{\frac{\nu}{2}} + \frac{2\underline{A}}{\bar{r}} \right) \delta^{\min\{\nu/2, 1\}}$ . Denote by  $C_7 \triangleq c_\nu^{\frac{1}{2}} (2m\bar{A})^{\frac{\nu}{2}} + \frac{2\underline{A}}{\bar{r}}$ . By definition of the  $\delta$ -entropy with bracketing, we must have

$$H_B(C_7 \delta^{\min\{\nu/2, 1\}}, \mathcal{G}, F) \leq \log(N_0^2) = 8m \log \left( 2 + \frac{\bar{r}}{\delta \underline{A}} \right), \quad (64)$$

which implies

$$\begin{aligned}
H_B(u, \mathcal{G}, F) &\leq 8m \log \left( 2 + \frac{\bar{r} C_7^{\frac{1}{\min\{\nu/2, 1\}}}}{u^{\frac{1}{\min\{\nu/2, 1\}}} \underline{A}} \right) \leq 8m \log \left( \left( 2 + \frac{\bar{r} C_7}{u \underline{A}} \right)^{\frac{1}{\min\{\nu/2, 1\}}} \right) \\
&= \frac{8m}{\min\{\nu/2, 1\}} \log \left( 2 + \frac{\bar{r} C_7}{\underline{A} u} \right).
\end{aligned}$$

Thus we have, when  $\delta \leq \frac{\bar{r} C_7}{\underline{A}}$ ,

$$\int_0^\delta H_B(u, \mathcal{G}, F)^{1/2} du \leq \left( \frac{8m}{\min\{\nu/2, 1\}} \right)^{1/2} \int_0^\delta \sqrt{\log \left( 2 + \frac{\bar{r} C_7}{\underline{A} u} \right)} du \leq C_3 \delta \sqrt{\log \left( 1 + \frac{C_4}{\delta} \right)},$$

where  $C_3 = 2 \left( \frac{8m}{\min\{\nu/2, 1\}} \right)^{1/2}$  and  $C_4 = \frac{\bar{r} C_7}{\underline{A}} = c_\nu^{\frac{1}{2}} \frac{\bar{r}}{\underline{A}} (2m\bar{A})^{\frac{\nu}{2}} + 2$ . Here the integral is bounded by

$$\begin{aligned}
\int_0^\delta \sqrt{\log \left( 2 + \frac{\bar{r} C_7}{\underline{A} u} \right)} du &= \sum_{s=1}^{\infty} \int_{2^{-s}\delta}^{2^{-(s-1)}\delta} \sqrt{\log \left( 2 + \frac{\bar{r} C_7}{\underline{A} u} \right)} du \\
&\leq \sum_{s=1}^{\infty} \int_{2^{-s}\delta}^{2^{-(s-1)}\delta} \sqrt{\log \left( 2 + \frac{\bar{r} C_7}{\underline{A} 2^{-s}\delta} \right)} du \\
&= \sum_{s=1}^{\infty} 2^{-(s-1)} \delta \sqrt{\log \left( 2 + \frac{\bar{r} C_7}{\underline{A} 2^{-s}\delta} \right)} \\
&\leq \delta \sum_{s=1}^{\infty} 2^{-(s-1)} \sqrt{s \log 2 + \log \left( 1 + \frac{\bar{r} C_7}{\underline{A} \delta} \right)}
\end{aligned}$$

$$\leq 2\delta \sqrt{\log \left( 1 + \frac{\bar{r}C_7}{A\delta} \right)} \sum_{s=1}^{\infty} s2^{-(s-1)},$$

where the infinite summation converges to 1, and the last inequality holds when  $\delta \leq \frac{\bar{r}C_7}{A}$ .

Now we are ready to apply Lemma F.7. Recall that  $\sup_{g \in \mathcal{G}} \|g\|_{\infty} \leq \frac{1}{2}$ , which implies  $\sup_{g \in \mathcal{G}} \|g\|_F \leq \frac{1}{2} < 1$ . Let  $\delta_{T-t} \triangleq \sqrt{\frac{\log(T-t)}{T-t}}$ . Consider the following partition of  $\mathcal{G}$ :

$$\begin{aligned} \mathcal{G}_s &= \left\{ g \in \mathcal{G} : 2^s \delta_{T-t} < \|g\|_F \leq 2^{s+1} \delta_{T-t} \right\}, \\ \mathcal{G}_c &= \left\{ g \in \mathcal{G} : \|g\|_F \leq \delta_{T-t} \right\}, \end{aligned}$$

with  $s = 1, \dots, S$  and  $S = \lfloor \log(\delta_{T-t}^{-1}) / \log 2 \rfloor + 1$ . Clearly  $\mathcal{G} = (\cup_{s=1}^S \mathcal{G}_s) \cup \mathcal{G}_c$ . We apply the peeling device.

Consider  $\mathcal{G}_c$  first. Take  $n = T - t, K = 1, R = \delta_{T-t} = \sqrt{\frac{\log(T-t)}{T-t}}$ . Denote constant  $v = 10C^2(1 + C_3)^2(1 + C_4)^2 + 2$  where  $C_3, C_4$  are specified above and  $C$  is the universal constant as in Lemma F.7. We further take  $a = v \frac{\log(T-t)}{\sqrt{T-t}}, C_0 = \frac{v}{2(1+C_3)(1+C_4)}$  and  $C_1 = 2v$ . We first check the conditions and then apply Lemma F.7. Within the function class  $\mathcal{G}_c$ , we have  $\sup_{g \in \mathcal{G}_c} \|g\|_{\infty} \leq \sup_{g \in \mathcal{G}} \|g\|_{\infty} \leq K = 1$  and  $\sup_{g \in \mathcal{G}_c} \|g\|_F \leq \delta_{T-t} = R$ . Next we check the conditions (60) - (63). Condition (60) is  $a \leq C_1 \sqrt{n} R^2 / (2K)$ . We have that

$$C_1 \frac{\sqrt{T-t} R^2}{2K} = v \frac{\log(T-t)}{\sqrt{T-t}} = a,$$

hence condition (60) holds true. For condition (61), we need  $a \leq 8\sqrt{2(T-t)}R$ . We have

$$8\sqrt{2(T-t)}R = 8\sqrt{2\log(T-t)} \geq v \frac{\log(T-t)}{\sqrt{T-t}} = a,$$

for all  $T - t \geq e^{\frac{v^2}{32} + 1}$ . Hence condition (61) holds true. Condition (62) is

$$a \geq C_0 \max \left\{ \int_0^{\sqrt{2}R} H_B(u/\sqrt{2}, \mathcal{G}_c, F)^{1/2} du, \sqrt{2}R \right\}.$$

By our previous calculation,

$$\begin{aligned} C_0 \max \left\{ \int_0^{\sqrt{2}R} H_B(u/\sqrt{2}, \mathcal{G}_c, F)^{1/2} du, \sqrt{2}R \right\} &\leq C_0 \max \left\{ \int_0^{\sqrt{2}R} H_B(u/\sqrt{2}, \mathcal{G}, F)^{1/2} du, \sqrt{2}R \right\} \\ &= \sqrt{2}C_0 \max \left\{ \int_0^R H_B(\delta, \mathcal{G}, F)^{1/2} d\delta, R \right\} \\ &\leq \sqrt{2}C_0 \max \left\{ C_3 R \sqrt{\log \left( 1 + \frac{C_4}{R} \right)}, R \right\}, \quad (65) \end{aligned}$$

when  $\delta_{T_t} \leq \frac{\bar{r}C_7}{A}$ , which automatically holds since  $\frac{\bar{r}C_7}{A} \geq 2 \geq K \geq R$ . For the first term, we have

$$\begin{aligned}
\sqrt{2}C_0C_3R\sqrt{\log\left(1+\frac{C_4}{R}\right)} &= \sqrt{2}C_0C_3\sqrt{\frac{\log(T-t)}{T-t}}\sqrt{\log\left(1+C_4\sqrt{\frac{T-t}{\log(T-t)}}\right)}, \\
&\leq \sqrt{2}C_0C_3\sqrt{\frac{\log(T-t)}{T-t}}\sqrt{\log((1+C_4)(T-t))}, \\
&\leq \sqrt{2}C_0C_3(1+C_4)\frac{\log(T-t)}{\sqrt{T-t}} \\
&= \frac{\sqrt{2}v}{2(1+C_3)(1+C_4)}C_3(1+C_4)\frac{\log(T-t)}{\sqrt{T-t}} \\
&\leq v\frac{\log(T-t)}{\sqrt{T-t}} = a.
\end{aligned}$$

For the second term, we have

$$\sqrt{2}C_0R = \sqrt{2}C_0\sqrt{\frac{\log(T-t)}{T-t}} \leq v\sqrt{\frac{\log(T-t)}{T-t}} \leq v\frac{\log(T-t)}{\sqrt{T-t}} = a,$$

for  $T-t > 3$ . Thus condition (62) holds true. Finally, for condition (63) we need to verify that  $C_0^2 \geq C^2(C_1+1)$ . We have

$$\begin{aligned}
C^2(C_1+1) &= C^2(2v+1) = C^2(20C^2(1+C_3)^2(1+C_4)^2+5) \\
&\leq 25C^4(1+C_3)^2(1+C_4)^2 \\
&\leq \left(\frac{10C^2(1+C_3)^2(1+C_4)^2+2}{2(1+C_3)(1+C_4)}\right)^2 = C_0^2. \tag{66}
\end{aligned}$$

With all four conditions checked, Lemma F.7 implies

$$\begin{aligned}
&\mathbb{P}\left(\sup_{g \in \mathcal{G}_c} \sqrt{T-t} \left| \frac{1}{T-t} \sum_{j=t+1}^T g(\mathbf{a}_j, r_j) - \mathbb{E}_{(\mathbf{a}, r) \sim F} g(\mathbf{a}, r) \right| \geq a\right) \\
&\leq C \exp\left(-\frac{a^2}{2C^2(C_1+1)R^2}\right), \\
&= C \exp\left(-\frac{(10C^2(1+C_3)^2(1+C_4)^2+2)^2}{2C^2(20C^2(1+C_3)^2(1+C_4)^2+5)} \log(T-t)\right) \\
&\leq C \exp(-2\log(T-t)) = \frac{C}{(T-t)^2}.
\end{aligned}$$

Combining the above, we may conclude that

$$\mathbb{P}\left(\sup_{g \in \mathcal{G}_c} \left| \frac{1}{T-t} \sum_{j=t+1}^T g(\mathbf{a}_j, r_j) - \mathbb{E}_{(\mathbf{a}, r) \sim F} g(\mathbf{a}, r) \right| \geq v\frac{\log(T-t)}{T-t}\right) \leq \frac{C}{(T-t)^2}, \tag{67}$$

as long as  $T-t \geq \max\left\{e^{\frac{v^2}{32}+1}, 3\right\} = e^{\frac{v^2}{32}+1}$ , where  $v$  is an absolute constant depending only on



problem primitives and independent of  $T - t$ :

$$v = 10C^2(1 + C_3)^2(1 + C_4)^2 + 2 = 10C^2 \left( 1 + 2 \left( \frac{8m}{\min\{\nu/2, 1\}} \right)^{1/2} \right)^2 \left( c_{\bar{\nu}}^{\frac{1}{2}} \frac{\bar{r}}{\underline{A}} (2m\bar{A})^{\frac{\nu}{2}} + 3 \right)^2 + 2, \quad (68)$$

and  $e^{\frac{v^2}{32}+1} \geq e^{\frac{4}{32}+1} > 3$ .

Next, we consider  $\cup_{s=1}^S \mathcal{G}_s$ . For any  $c > 0$ , the union bound implies that

$$\begin{aligned} & \mathbb{P} \left( \sup_{g \in \cup_{s=1}^S \mathcal{G}_s} \frac{\left| \frac{1}{T-t} \sum_{j=t+1}^T g(\mathbf{a}_j, r_j) - \mathbb{E}_{(\mathbf{a}, r) \sim F} g(\mathbf{a}, r) \right|}{\|g\|_F} \geq c \right) \\ & \leq \sum_{s=1}^S \mathbb{P} \left( \sup_{g \in \mathcal{G}_s} \frac{\left| \frac{1}{T-t} \sum_{j=t+1}^T g(\mathbf{a}_j, r_j) - \mathbb{E}_{(\mathbf{a}, r) \sim F} g(\mathbf{a}, r) \right|}{\|g\|_F} \geq c \right) \\ & \leq \sum_{s=1}^S \mathbb{P} \left( \sup_{g \in \mathcal{G}_s} \left| \frac{1}{T-t} \sum_{j=t+1}^T g(\mathbf{a}_j, r_j) - \mathbb{E}_{(\mathbf{a}, r) \sim F} g(\mathbf{a}, r) \right| \geq 2^s \delta_{T-t} c \right), \end{aligned} \quad (69)$$

where the second inequality is because for any  $g \in \mathcal{G}_s$ ,  $\|g\|_F \geq 2^s \delta_{T-t}$ ,  $s = 1, \dots, S$ . In the following, we bound the tail probability

$$\mathbb{P} \left( \sup_{g \in \mathcal{G}_s} \left| \frac{1}{T-t} \sum_{j=t+1}^T g(\mathbf{a}_j, r_j) - \mathbb{E}_{(\mathbf{a}, r) \sim F} g(\mathbf{a}, r) \right| \geq 2^s \delta_{T-t} c \right)$$

for each  $\mathcal{G}_s$ ,  $s = 1, \dots, S$ .

Before applying Lemma F.7, we first check the conditions (60) - (63) hold. Take  $n = T - t$ ,  $K_s = 1$ ,  $R_s = 2^{s+1} \delta_{T-t} = 2^{s+1} \sqrt{\frac{\log(T-t)}{T-t}}$ . Recall we have set constant  $v$  as in (68), and we take  $a = v 2^{s+1} \frac{\log(T-t)}{\sqrt{T-t}}$ ,  $C_0 = \frac{v}{2(1+C_3)(1+C_4)}$  and  $C_1 = 2v$ . Within the function class  $\mathcal{G}_s$ , we have  $\sup_{g \in \mathcal{G}_s} \|g\|_{\infty} \leq \sup_{g \in \mathcal{G}} \|g\|_{\infty} \leq K_s = 1$  and  $\sup_{g \in \mathcal{G}_s} \|g\|_F \leq R_s$ . Next we check conditions (60)-(63). Condition (60) is  $a \leq C_1 \sqrt{n} R_s^2 / (2K_s)$ . We have that

$$C_1 \frac{\sqrt{T-t} R_s^2}{2K_s} = v 2^{2(s+1)} \frac{\log(T-t)}{\sqrt{T-t}} \geq v 2^{s+1} \frac{\log(T-t)}{\sqrt{T-t}} = a,$$

hence condition (60) holds true. For condition (61), we need  $a \leq 8\sqrt{2(T-t)} R_s$ . We have

$$8\sqrt{2(T-t)} R_s = 2^{s+4} \sqrt{2 \log(T-t)} \geq v 2^{s+1} \frac{\log(T-t)}{\sqrt{T-t}} = a,$$

for all  $T - t \geq e^{\frac{v^2}{32}+1}$ . Hence condition (61) holds true. Condition (62) is

$$a \geq C_0 \max \left\{ \int_0^{\sqrt{2} R_s} H_B(u/\sqrt{2}, \mathcal{G}_s, F)^{1/2} du, \sqrt{2} R_s \right\}.$$

By our previous calculation as in (65),

$$C_0 \max \left\{ \int_0^{\sqrt{2}R_s} H_B(u/\sqrt{2}, \mathcal{G}_s, F)^{1/2} du, \sqrt{2}R_s \right\} \leq \sqrt{2}C_0 \max \left\{ C_3 R_s \sqrt{\log \left( 1 + \frac{C_4}{R_s} \right)}, R_s \right\},$$

when  $\delta_{T-t} \leq \frac{\bar{r}C_7}{\underline{A}}$ , which automatically holds since  $\frac{\bar{r}C_7}{\underline{A}} \geq 2 \geq K_s \geq R_s$ . For the first term, we have

$$\begin{aligned} \sqrt{2}C_0 C_3 R_s \sqrt{\log \left( 1 + \frac{C_4}{R_s} \right)} &= \sqrt{2}C_0 C_3 2^{s+1} \sqrt{\frac{\log(T-t)}{T-t}} \sqrt{\log \left( 1 + C_4 2^{-s-1} \sqrt{\frac{T-t}{\log(T-t)}} \right)}, \\ &\leq \sqrt{2}C_0 C_3 2^{s+1} \sqrt{\frac{\log(T-t)}{T-t}} \sqrt{\log((1+C_4)(T-t))}, \\ &\leq \sqrt{2}C_0 C_3 (1+C_4) 2^{s+1} \frac{\log(T-t)}{\sqrt{T-t}} \\ &= \frac{\sqrt{2}v}{2(1+C_3)(1+C_4)} C_3 (1+C_4) 2^{s+1} \frac{\log(T-t)}{\sqrt{T-t}} \\ &\leq v 2^{s+1} \frac{\log(T-t)}{\sqrt{T-t}} = a. \end{aligned}$$

For the second term, we have

$$\sqrt{2}C_0 R_s = \sqrt{2}C_0 2^{s+1} \sqrt{\frac{\log(T-t)}{T-t}} \leq v 2^{s+1} \sqrt{\frac{\log(T-t)}{T-t}} \leq v 2^{s+1} \frac{\log(T-t)}{\sqrt{T-t}} = a,$$

for  $T-t > 3$ . Thus condition (62) holds true. Finally, for condition (63) we need to verify that  $C_0^2 \geq C^2(C_1+1)$ , which can be done identically as in (66).

With all four conditions checked, Lemma F.7 implies

$$\begin{aligned} &\mathbb{P} \left( \sup_{g \in \mathcal{G}_s} \sqrt{T-t} \left| \frac{1}{T-t} \sum_{j=t+1}^T g(\mathbf{a}_j, r_j) - \mathbb{E}_{(\mathbf{a}, r) \sim F} g(\mathbf{a}, r) \right| \geq a \right) \\ &\leq C \exp \left( -\frac{a^2}{2C^2(C_1+1)R_s^2} \right), \\ &= C \exp \left( -\frac{(10C^2(1+C_3)^2(1+C_4)^2+2)^2}{2C^2(20C^2(1+C_3)^2(1+C_4)^2+5)} \log(T-t) \right) \\ &\leq C \exp(-2\log(T-t)) = \frac{C}{(T-t)^2}. \end{aligned}$$

Recall  $a = v 2^{s+1} \frac{\log(T-t)}{\sqrt{T-t}} = v \sqrt{\log(T-t)} 2^{s+1} \delta_{T-t}$ . The tail bound can be equivalently written as

$$\mathbb{P} \left( \sup_{g \in \mathcal{G}_s} \left| \frac{1}{T-t} \sum_{j=t+1}^T g(\mathbf{a}_j, r_j) - \mathbb{E}_{(\mathbf{a}, r) \sim F} g(\mathbf{a}, r) \right| \geq v \sqrt{\frac{\log(T-t)}{T-t}} 2^{s+1} \delta_{T-t} \right) \leq \frac{C}{(T-t)^2},$$

as long as  $T-t \geq \max \left\{ e^{\frac{v^2}{32}+1}, 3 \right\} = e^{\frac{v^2}{32}+1}$ , where  $v$  is as in (68). Now we plug the above back

into (69) with  $c = 2v\sqrt{\frac{\log(T-t)}{T-t}}$ , and conclude that for  $T-t \geq e^{\frac{v^2}{32}+1}$ , it holds true that

$$\begin{aligned} & \mathbb{P} \left( \sup_{g \in \cup_{s=1}^S \mathcal{G}_s} \frac{\left| \frac{1}{T-t} \sum_{j=t+1}^T g(\mathbf{a}_j, r_j) - \mathbb{E}_{(\mathbf{a}, r) \sim F} g(\mathbf{a}, r) \right|}{\|g\|_F} \geq 2v\sqrt{\frac{\log(T-t)}{T-t}} \right) \\ & \leq \frac{CS}{(T-t)^2} \leq \frac{2C \log(T-t)}{(T-t)^2}, \end{aligned} \quad (70)$$

where the second inequality is because  $S = \lfloor \log(\delta_{T-t}^{-1}) / \log 2 \rfloor + 1 \leq 2 \log(T-t)$  for  $T-t \geq e^{\frac{v^2}{32}+1}$ .

Combining tail bounds (67) and (70) with another union bound, we conclude that

$$\begin{aligned} & \mathbb{P} \left( \left| \frac{1}{T-t} \sum_{j=t+1}^T g(\mathbf{a}_j, r_j) - \mathbb{E}_{(\mathbf{a}, r) \sim F} g(\mathbf{a}, r) \right| \geq 2v\sqrt{\frac{\log(T-t)}{T-t}} \|g\|_F + v\frac{\log(T-t)}{T-t}, \forall g \in \mathcal{G} \right) \\ & = \mathbb{P} \left( \left| \frac{1}{T-t} \sum_{j=t+1}^T g(\mathbf{a}_j, r_j) - \mathbb{E}_{(\mathbf{a}, r) \sim F} g(\mathbf{a}, r) \right| \geq 2v\sqrt{\frac{\log(T-t)}{T-t}} \|g\|_F + v\frac{\log(T-t)}{T-t}, \forall g \in (\cup_{s=1}^S \mathcal{G}_s) \cup \mathcal{G}_c \right) \\ & \leq \mathbb{P} \left( \left| \frac{1}{T-t} \sum_{j=t+1}^T g(\mathbf{a}_j, r_j) - \mathbb{E}_{(\mathbf{a}, r) \sim F} g(\mathbf{a}, r) \right| \geq 2v\sqrt{\frac{\log(T-t)}{T-t}} \|g\|_F + v\frac{\log(T-t)}{T-t}, \forall g \in \mathcal{G}_c \right) \\ & \quad + P \left( \left| \frac{1}{T-t} \sum_{j=t+1}^T g(\mathbf{a}_j, r_j) - \mathbb{E}_{(\mathbf{a}, r) \sim F} g(\mathbf{a}, r) \right| \geq 2v\sqrt{\frac{\log(T-t)}{T-t}} \|g\|_F + v\frac{\log(T-t)}{T-t}, \forall g \in \cup_{s=1}^S \mathcal{G}_s \right) \\ & \leq \mathbb{P} \left( \sup_{g \in \mathcal{G}_c} \left| \frac{1}{T-t} \sum_{j=t+1}^T g(\mathbf{a}_j, r_j) - \mathbb{E}_{(\mathbf{a}, r) \sim F} g(\mathbf{a}, r) \right| \geq v\sqrt{\frac{\log(T-t)}{T-t}} \right) \\ & \quad + P \left( \sup_{g \in \cup_{s=1}^S \mathcal{G}_s} \frac{\left| \frac{1}{T-t} \sum_{j=t+1}^T g(\mathbf{a}_j, r_j) - \mathbb{E}_{(\mathbf{a}, r) \sim F} g(\mathbf{a}, r) \right|}{\|g\|_F} \geq 2v\sqrt{\frac{\log(T-t)}{T-t}} \right) \\ & \leq \frac{C}{(T-t)^2} + \frac{2C \log(T-t)}{(T-t)^2} = \frac{3C \log(T-t)}{(T-t)^2}, \end{aligned}$$

for  $T-t \geq e^{\frac{v^2}{32}+1}$ . Consequently, we conclude that

$$\left| \frac{1}{T-t} \sum_{j=t+1}^T g(\mathbf{a}_j, r_j) - \mathbb{E}_{(\mathbf{a}, r) \sim F} g(\mathbf{a}, r) \right| \leq 2v\sqrt{\frac{\log(T-t)}{T-t}} \left( \|g\|_F + \sqrt{\frac{\log(T-t)}{T-t}} \right), \forall g \in \mathcal{G} \quad (71)$$

with probability at least  $1 - \frac{3C \log(T-t)}{(T-t)^2}$  for  $T-t \geq e^{\frac{v^2}{32}+1}$ . Note that

$$g(\mathbf{a}, r) = \frac{A}{4m\bar{r}A} \left( \mathbf{a}^\top (\boldsymbol{\lambda}_2 - \boldsymbol{\lambda}_1) \mathbb{I}_{\{r > \mathbf{a}^\top \boldsymbol{\lambda}_1\}} \right) \in \mathcal{G}$$

with  $\mathbf{p}_4 = \boldsymbol{\lambda}_2$  and  $\mathbf{p}_j = \boldsymbol{\lambda}_1$  for  $j = 1, 2, 3$ . Also,

$$\|\mathbf{a}^\top (\boldsymbol{\lambda}_2 - \boldsymbol{\lambda}_1) \mathbb{I}_{\{r > \mathbf{a}^\top \boldsymbol{\lambda}_1\}}\|_F = \left( \mathbb{E}_{(\mathbf{a}, r) \sim F} \left( \mathbf{a}^\top (\boldsymbol{\lambda}_2 - \boldsymbol{\lambda}_1) \mathbb{I}_{\{r > \mathbf{a}^\top \boldsymbol{\lambda}_1\}} \right)^2 \right)^{\frac{1}{2}}$$

$$= \sqrt{\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \{(\mathbf{a}^\top (\boldsymbol{\lambda}_2 - \boldsymbol{\lambda}_1))^2 (1 - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_1))\}}.$$

The desired result follows from (71) with

$$\tilde{C} = 2 \left( \frac{4m\bar{r}\bar{A}}{\underline{A}} + 1 \right) \left( 10C^2 \left( 1 + 2 \left( \frac{8m}{\min\{\nu/2, 1\}} \right)^{1/2} \right)^2 \left( c_{\nu}^{\frac{1}{2}} \frac{\bar{r}}{\underline{A}} (2m\bar{A})^{\frac{\nu}{2}} + 3 \right)^2 + 2 \right).$$

Q.E.D.

Similarly, we state and prove the second concentration lemma.

**Lemma F.9.** *Suppose  $T - t \geq e^{\frac{\tilde{C}_1^2}{32} + 1}$ . We have with probability at least  $1 - \frac{3C \log(T-t)}{(T-t)^2}$ , for all  $\boldsymbol{\lambda}_1, \boldsymbol{\lambda}_2 \in \Omega$ ,*

$$\left| \mathbb{E}_{(\mathbf{a}, r) \sim F} \int_{\mathbf{a}^\top \boldsymbol{\lambda}_1}^{\mathbf{a}^\top \boldsymbol{\lambda}_2} (\mathbb{I}_{\{r > u\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \boldsymbol{\lambda}_2\}}) du - \frac{1}{T-t} \sum_{j=t+1}^T \int_{\mathbf{a}_j^\top \boldsymbol{\lambda}_1}^{\mathbf{a}_j^\top \boldsymbol{\lambda}_2} (\mathbb{I}_{\{r_j > u\}} - \mathbb{I}_{\{r_j > \mathbf{a}^\top \boldsymbol{\lambda}_2\}}) du \right| \leq \tilde{C}_1 \sqrt{\frac{\log(T-t)}{T-t}} \left( \left( \mathbb{E}_{(\mathbf{a}, r) \sim F} \left( \int_{\mathbf{a}^\top \boldsymbol{\lambda}_1}^{\mathbf{a}^\top \boldsymbol{\lambda}_2} (\mathbb{I}_{\{r > u\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \boldsymbol{\lambda}_2\}}) du \right)^2 \right)^{1/2} + \sqrt{\frac{\log(T-t)}{T-t}} \right), \quad (72)$$

and

$$\left| \mathbb{E}_{(\mathbf{a}, r) \sim F} \int_{\mathbf{a}^\top \boldsymbol{\lambda}_2}^{\mathbf{a}^\top \boldsymbol{\lambda}_1} (\mathbb{I}_{\{r > u\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \boldsymbol{\lambda}_1\}}) du - \frac{1}{T-t} \sum_{j=t+1}^T \int_{\mathbf{a}_j^\top \boldsymbol{\lambda}_2}^{\mathbf{a}_j^\top \boldsymbol{\lambda}_1} (\mathbb{I}_{\{r_j > u\}} - \mathbb{I}_{\{r_j > \mathbf{a}^\top \boldsymbol{\lambda}_1\}}) du \right| \leq \tilde{C}_1 \sqrt{\frac{\log(T-t)}{T-t}} \left( \left( \mathbb{E}_{(\mathbf{a}, r) \sim F} \left( \int_{\mathbf{a}^\top \boldsymbol{\lambda}_2}^{\mathbf{a}^\top \boldsymbol{\lambda}_1} (\mathbb{I}_{\{r > u\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \boldsymbol{\lambda}_1\}}) du \right)^2 \right)^{1/2} + \sqrt{\frac{\log(T-t)}{T-t}} \right), \quad (73)$$

where

$$\tilde{C}_1 = 2 \left( \frac{4m\bar{r}\bar{A}}{\underline{A}} + 1 \right) \left( 10C^2 \left( 1 + 2 \left( \frac{10m}{\min\{\nu/2, 1\}} \right)^{1/2} \right)^2 \left( 4c_{\nu}^{\frac{1}{2}} \frac{\bar{r}}{\underline{A}} (2m\bar{A})^{\frac{\nu}{2}} + 9 \right)^2 + 2 \right),$$

and  $C$  is a universal constant.

*Proof of Lemma F.9.* The proof of Lemma F.9 is similar to the proof of Lemma F.8. To start with, define the function class parameterized by  $\mathbf{p} = \mathbf{p}_j \in \Omega$ ,  $j = 1, \dots, 5$ :

$$\mathcal{G} = \left\{ g : g(\mathbf{a}, r) = \frac{\underline{A}}{4m\bar{A}\bar{r}} \left( \int_{\mathbf{a}^\top \mathbf{p}_1}^{\mathbf{a}^\top \mathbf{p}_2} \mathbb{I}_{\{r > u\}} du - (\mathbf{a}^\top \mathbf{p}_3 - \mathbf{a}^\top \mathbf{p}_4) \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{p}_5\}} \right), \mathbf{p}_j \in \Omega, j = 1, \dots, 5 \right\}.$$

One can verify that  $\|g\|_{\infty} = \|g\|_{L_{\infty}(\mathcal{S})} \leq 1$ . Again we approximate  $\mathcal{G}$  by its  $\delta$ -net. More precisely, for any  $\delta > 0$ , we consider the grid points set  $\Omega_0 \subset [0, \frac{\bar{r}}{\underline{A}}]^{5m}$ , where each element of each point  $\mathbf{q} = (\mathbf{q}^{(j)})_{j=1, \dots, 5} \in \Omega_0$  has the form  $k\delta$ , with  $k = 0, 1, \dots, \lfloor \frac{\underline{A}}{\delta\bar{r}} \rfloor + 1$ , and  $\mathbf{q}^{(1)}, \mathbf{q}^{(2)}, \dots, \mathbf{q}^{(5)} \in \Omega$ . Thus,

we have  $\text{card}(\Omega_0) \leq (2 + \frac{\bar{r}}{\delta \bar{A}})^{5m}$ . Let  $N_0 = (2 + \frac{\bar{r}}{\delta \bar{A}})^{5m}$ . Let

$$\mathcal{G} = \left\{ g : g(\mathbf{a}, r) = \frac{\underline{A}}{4m\bar{A}\bar{r}} \left( \int_{\mathbf{a}^\top \mathbf{q}^{(1)}}^{\mathbf{a}^\top \mathbf{q}^{(2)}} \mathbb{I}_{\{r > u\}} du - \left( \mathbf{a}^\top \mathbf{q}^{(3)} - \mathbf{a}^\top \mathbf{q}^{(4)} \right) \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}^{(5)}\}} \right), \mathbf{q}^{(j)} \in \Omega_0, j = 1, \dots, 5 \right\}.$$

For a fixed  $g \in \mathcal{G}$  defined by  $\mathbf{p}$ , let's denote by  $\mathbf{q}_U$  and  $\mathbf{q}_L$  its upper and lower  $\delta$ -approximation in  $\mathcal{G}_0$ , namely,  $\max\{\mathbf{q}_U^{(j)} - \delta \mathbf{1}, \mathbf{0}\} \leq \mathbf{p}_j < \mathbf{q}_U^{(j)}$  and  $\mathbf{q}_L^{(j)} < \mathbf{p}_j \leq \mathbf{q}_L^{(j)} + \delta \mathbf{1}$  for  $j = 1, \dots, 5$  where the maximum is taken elementwisely. Then by definition  $\|\mathbf{q}_U - \mathbf{q}_L\|_\infty \leq 2\delta$ . Let

$$g_U(\mathbf{a}, r) = \frac{\underline{A}}{4m\bar{A}\bar{r}} \left( \int_{\mathbf{a}^\top \mathbf{q}_L^{(1)}}^{\mathbf{a}^\top \mathbf{q}_U^{(2)}} \mathbb{I}_{\{r > u\}} du - \left( \mathbf{a}^\top \mathbf{q}_L^{(3)} - \mathbf{a}^\top \mathbf{q}_U^{(4)} \right) \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_U^{(5)}\}} \right),$$

$$g_L(\mathbf{a}, r) = \frac{\underline{A}}{4m\bar{A}\bar{r}} \left( \int_{\mathbf{a}^\top \mathbf{q}_U^{(1)}}^{\mathbf{a}^\top \mathbf{q}_L^{(2)}} \mathbb{I}_{\{r > u\}} du - \left( \mathbf{a}^\top \mathbf{q}_U^{(3)} - \mathbf{a}^\top \mathbf{q}_L^{(4)} \right) \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_L^{(5)}\}} \right).$$

Then one can check that  $g_L(\mathbf{a}, r) \leq g(\mathbf{a}, r) \leq g_U(\mathbf{a}, r)$  for all  $(\mathbf{a}, r) \in \mathcal{S}$ , since  $\mathbf{a}$  is non-negative, and  $\mathbf{q}_L$  and  $\mathbf{q}_U$  satisfy the element-wise bound. We next bound  $\|g_U - g_L\|_F$  by

$$\begin{aligned} \|g_U - g_L\|_F^2 &= \mathbb{E}_{(\mathbf{a}, r) \sim F} (g_U(\mathbf{a}, r) - g_L(\mathbf{a}, r))^2 \\ &\leq \left( \frac{\underline{A}}{4m\bar{A}\bar{r}} \right)^2 \mathbb{E}_{(\mathbf{a}, r) \sim F} \left( \int_{\mathbf{a}^\top \mathbf{q}_L^{(1)}}^{\mathbf{a}^\top \mathbf{q}_U^{(2)}} \mathbb{I}_{\{r > u\}} du - \int_{\mathbf{a}^\top \mathbf{q}_U^{(1)}}^{\mathbf{a}^\top \mathbf{q}_L^{(2)}} \mathbb{I}_{\{r > u\}} du \right. \\ &\quad \left. + \left( \mathbf{a}^\top \mathbf{q}_U^{(3)} - \mathbf{a}^\top \mathbf{q}_L^{(4)} \right) \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_L^{(5)}\}} - \left( \mathbf{a}^\top \mathbf{q}_L^{(3)} - \mathbf{a}^\top \mathbf{q}_U^{(4)} \right) \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_U^{(5)}\}} \right)^2 \\ &\leq \left( \frac{\underline{A}}{4m\bar{A}\bar{r}} \right)^2 \mathbb{E}_{(\mathbf{a}, r) \sim F} \left( \sum_{j=1}^4 \left( \mathbf{a}^\top \mathbf{q}_U^{(j)} - \mathbf{a}^\top \mathbf{q}_L^{(j)} \right) \right. \\ &\quad \left. + \max \left( \mathbf{a}^\top \mathbf{q}_U^{(3)}, \mathbf{a}^\top \mathbf{q}_L^{(4)} \right) \left| \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_L^{(5)}\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_U^{(5)}\}} \right| \right)^2 \\ &\leq \left( \frac{\underline{A}}{4m\bar{A}\bar{r}} \right)^2 \mathbb{E}_{(\mathbf{a}, r) \sim F} \left( 8m\bar{A}\delta + \frac{m\bar{A}\bar{r}}{\underline{A}} \left| \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_L^{(5)}\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_U^{(5)}\}} \right| \right)^2 \\ &\leq 2 \left( \frac{\underline{A}}{4m\bar{A}\bar{r}} \right)^2 \mathbb{E}_{(\mathbf{a}, r) \sim F} \left( 64m^2 \bar{A}^2 \delta^2 + \left( \frac{m\bar{A}\bar{r}}{\underline{A}} \right)^2 \left| \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_L^{(5)}\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \mathbf{q}_U^{(5)}\}} \right| \right) \\ &\leq 2 \left( \frac{\underline{A}}{4m\bar{A}\bar{r}} \right)^2 \left( 64m^2 \bar{A}^2 \delta^2 + \left( \frac{m\bar{A}\bar{r}}{\underline{A}} \right)^2 \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left| F_{\mathbf{a}}^r(\mathbf{a}^\top \mathbf{q}_L^{(5)}) - F_{\mathbf{a}}^r(\mathbf{a}^\top \mathbf{q}_U^{(5)}) \right| \right) \\ &\leq 2 \left( \frac{\underline{A}}{4m\bar{A}\bar{r}} \right)^2 \left( 64m^2 \bar{A}^2 \delta^2 + \left( \frac{m\bar{A}\bar{r}}{\underline{A}} \right)^2 c_\nu \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left| \mathbf{a}^\top \mathbf{q}_L^{(5)} - \mathbf{a}^\top \mathbf{q}_U^{(5)} \right|^\nu \right) \\ &\hspace{15em} \text{(under either assumptions)} \\ &\leq 8 \left( \left( \frac{\underline{A}\delta}{\bar{r}} \right)^2 + c_\nu (2m\bar{A}\delta)^\nu \right) \leq 8 \left( \frac{2\underline{A}}{\bar{r}} + c_\nu^{\frac{1}{2}} (2m\bar{A})^{\frac{\nu}{2}} \right)^2 \delta^{\min\{2, \nu\}}. \end{aligned}$$

As a result we have  $\|g_U - g_L\|_F \leq 4 \left( \frac{2\underline{A}}{\bar{r}} + c_\nu^{\frac{1}{2}} (2m\bar{A})^{\frac{\nu}{2}} \right) \delta^{\min\{1, \frac{\nu}{2}\}}$ . Now similar to the proof of

Lemma F.8, we have

$$\int_0^\delta H_B(u, \mathcal{G}, F)^{1/2} du \leq C'_3 \delta \sqrt{\log \left( 1 + \frac{C'_4}{\delta} \right)},$$

where  $C'_3 = 2 \left( \frac{10m}{\min\{\nu/2, 1\}} \right)^{1/2}$  and  $C'_4 = 4 \left( 2 + c_{\nu}^{\frac{1}{2}} \frac{\bar{r}}{\underline{A}} (2m\bar{A})^{\frac{\nu}{2}} \right)$ .

We notice that both functions

$$\int_{\mathbf{a}^\top \boldsymbol{\lambda}_1}^{\mathbf{a}^\top \boldsymbol{\lambda}_2} (\mathbb{I}_{\{r>u\}} - \mathbb{I}_{\{r>\mathbf{a}^\top \boldsymbol{\lambda}_2\}}) du, \quad \int_{\mathbf{a}^\top \boldsymbol{\lambda}_2}^{\mathbf{a}^\top \boldsymbol{\lambda}_1} (\mathbb{I}_{\{r>u\}} - \mathbb{I}_{\{r>\mathbf{a}^\top \boldsymbol{\lambda}_2\}}) du$$

are in  $\mathcal{G}$ . The rest of the proof is identical to that of Lemma F.8, and we omit the details. Q.E.D.

### F.4.3 Other Auxiliary Lemmas

This section contains the proof of all other results appeared earlier in the paper.

**Lemma F.10.** *For any  $\boldsymbol{\lambda}_1, \boldsymbol{\lambda}_2 \in \mathbb{R}^m$  and any  $\mathbf{a} \in \mathbb{R}^m, r \in \mathbb{R}, \mathbf{b} \in \mathbb{R}^m$ , it holds true that*

$$h_{t, \mathbf{b}}(\boldsymbol{\lambda}_1, \mathbf{a}, r) - h_{t, \mathbf{b}}(\boldsymbol{\lambda}_2, \mathbf{a}, r) = \phi_{t, \mathbf{b}}(\boldsymbol{\lambda}_2, \mathbf{a}, r)^\top (\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2) + \int_{\mathbf{a}^\top \boldsymbol{\lambda}_1}^{\mathbf{a}^\top \boldsymbol{\lambda}_2} (\mathbb{I}_{\{r>v\}} - \mathbb{I}_{\{r>\mathbf{a}^\top \boldsymbol{\lambda}_2\}}) dv,$$

*Proof.* Refer to the proof of Lemma 1 in Li and Ye (2022). Q.E.D.

**Lemma F.11.** *For any  $\boldsymbol{\lambda}_1, \boldsymbol{\lambda}_2 \in \mathbb{R}_{\geq 0}^m$ ,*

$$\mathbb{E}_{(\mathbf{a}, r) \sim F} \left( \int_{\mathbf{a}^\top \boldsymbol{\lambda}_1}^{\mathbf{a}^\top \boldsymbol{\lambda}_2} (\mathbb{I}_{\{r>v\}} - \mathbb{I}_{\{r>\mathbf{a}^\top \boldsymbol{\lambda}_2\}}) dv \right)^2 \leq \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left( \left( \mathbf{a}^\top (\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2) \right)^2 \cdot \left| F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_1) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_2) \right| \right).$$

*Proof.* The Cauchy-Schwarz inequality implies that

$$\begin{aligned} & \mathbb{E}_{(\mathbf{a}, r) \sim F} \left( \int_{\mathbf{a}^\top \boldsymbol{\lambda}_1}^{\mathbf{a}^\top \boldsymbol{\lambda}_2} (\mathbb{I}_{\{r>v\}} - \mathbb{I}_{\{r>\mathbf{a}^\top \boldsymbol{\lambda}_2\}}) dv \right)^2 \\ & \leq \mathbb{E}_{(\mathbf{a}, r) \sim F} \left( \left| \int_{\mathbf{a}^\top \boldsymbol{\lambda}_1}^{\mathbf{a}^\top \boldsymbol{\lambda}_2} 1 dv \right| \cdot \left| \int_{\mathbf{a}^\top \boldsymbol{\lambda}_1}^{\mathbf{a}^\top \boldsymbol{\lambda}_2} (\mathbb{I}_{\{r>v\}} - \mathbb{I}_{\{r>\mathbf{a}^\top \boldsymbol{\lambda}_2\}}) dv \right| \right) \\ & = \mathbb{E}_{(\mathbf{a}, r) \sim F} \left( \left| \mathbf{a}^\top (\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2) \right| \cdot \left| \int_{\mathbf{a}^\top \boldsymbol{\lambda}_1}^{\mathbf{a}^\top \boldsymbol{\lambda}_2} (\mathbb{I}_{\{r>v\}} - \mathbb{I}_{\{r>\mathbf{a}^\top \boldsymbol{\lambda}_2\}}) dv \right| \right) \\ & \leq \mathbb{E}_{(\mathbf{a}, r) \sim F} \left( \left| \mathbf{a}^\top (\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2) \right| \cdot \left| \int_{\mathbf{a}^\top \boldsymbol{\lambda}_1}^{\mathbf{a}^\top \boldsymbol{\lambda}_2} (\mathbb{I}_{\{r>\mathbf{a}^\top \boldsymbol{\lambda}_1\}} - \mathbb{I}_{\{r>\mathbf{a}^\top \boldsymbol{\lambda}_2\}}) dv \right| \right) \\ & = \mathbb{E}_{(\mathbf{a}, r) \sim F} \left( \left( \mathbf{a}^\top (\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2) \right)^2 \cdot \left| \mathbb{I}_{\{r>\mathbf{a}^\top \boldsymbol{\lambda}_1\}} - \mathbb{I}_{\{r>\mathbf{a}^\top \boldsymbol{\lambda}_2\}} \right| \right) \\ & = \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left( \left( \mathbf{a}^\top (\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2) \right)^2 \cdot \left| F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_1) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}_2) \right| \right). \end{aligned}$$

Q.E.D.

**Lemma F.12.** *It holds true that  $\tilde{\boldsymbol{\lambda}}_t, \boldsymbol{\lambda}_t^*, \bar{\boldsymbol{\lambda}}_t^* \in \Omega$  for any  $\mathbf{b}_{t-1}$ , any  $\mathcal{I}_{t-1}$  and  $t = 1, \dots, T$ .*

*Proof.* We only prove the Lemma for  $\tilde{\lambda}_t$ . The argument for  $\lambda_t^*$  and  $\bar{\lambda}_t^*$  is similar. For any  $\lambda = (\lambda_1, \dots, \lambda_m)^\top \notin \Omega$  but is feasible ( $\lambda \in \mathbb{R}_{\geq 0}^m$ ), there exists  $i$  such that  $\lambda_i > \frac{\bar{r}}{\underline{A}}$ . Let  $\lambda' \triangleq (\lambda_1, \dots, \lambda_{i-1}, \frac{\bar{r}}{\underline{A}}, \lambda_{i+1}, \dots, \lambda_m)^\top$ . Since  $\mathbf{a}^\top \lambda > \mathbf{a}^\top \lambda' \geq a_i \lambda_i > \bar{r}$ , it holds path-wisely that  $(r - \mathbf{a}^\top \lambda)^+ = (r - \mathbf{a}^\top \lambda')^+ = 0$ . However,  $\mathbf{b}_{t-1}^\top \lambda > \mathbf{b}_{t-1}^\top \lambda'$  as all elements are non-negative, which leads to  $f_{t, \mathbf{b}_{t-1}}(\lambda) > f_{t, \mathbf{b}_{t-1}}(\lambda')$ . In other words,  $\lambda$  cannot be the optimal solution. Thus  $\tilde{\lambda}_t$  must be in  $\Omega$  and we have proved the Lemma. Q.E.D.

**Corollary F.13.** *The optimal solutions  $\tilde{\lambda}_t, \lambda_t^*, \bar{\lambda}_t^*$  have uniformly bounded norm for any  $\mathbf{b}_{t-1}$ , any  $\mathcal{I}_{t-1}$  and  $t = 1, \dots, T$ ,*

$$\begin{aligned} 0 &\leq \|\tilde{\lambda}_t\|_\infty, \|\lambda_t^*\|_\infty, \|\bar{\lambda}_t^*\|_\infty \leq \frac{\bar{r}}{\underline{A}}, \\ 0 &\leq \|\tilde{\lambda}_t\|, \|\lambda_t^*\|, \|\bar{\lambda}_t^*\| \leq \sqrt{m} \frac{\bar{r}}{\underline{A}}. \end{aligned}$$

**Lemma F.14.** *For any  $\lambda \in \Omega$ , any  $\mathbf{b}_{t-1} \in \mathbb{R}_{\geq 0}^m$  and any  $t = 1, \dots, T$ ,*

$$\begin{aligned} \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\tilde{\lambda}_t, \mathbf{a}, r)^\top (\lambda - \tilde{\lambda}_t) &\geq 0, \\ \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1}}(\lambda_j^*, \mathbf{a}_j, r_j)^\top (\lambda - \lambda_j^*) &\geq -\frac{2m^2 \bar{A} \bar{r}}{\underline{A}}, \quad a.s. \\ \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1} - \mathbf{a}_t}(\bar{\lambda}_j^*, \mathbf{a}_j, r_j)^\top (\lambda - \bar{\lambda}_j^*) &\geq -\frac{2m^2 \bar{A} \bar{r}}{\underline{A}}, \quad a.s. \quad (s.t. \text{ feasibility constraint}) \end{aligned}$$

*Proof.* By Lemma F.10, we have

$$f_{t, \mathbf{b}}(\lambda) - f_{t, \mathbf{b}}(\tilde{\lambda}_t) = \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}}(\tilde{\lambda}_t, \mathbf{a}, r)^\top (\lambda - \tilde{\lambda}_t) + \mathbb{E}_{(\mathbf{a}, r) \sim F} \int_{\mathbf{a}^\top \lambda}^{\mathbf{a}^\top \tilde{\lambda}_t} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \tilde{\lambda}_t\}} \right) dv.$$

We argue by contradiction. Suppose there exists  $\lambda \in \mathbb{R}_{\geq 0}^m$  such that  $\mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\tilde{\lambda}_t, \mathbf{a}, r)^\top (\lambda - \tilde{\lambda}_t) = -\eta < 0$ . Consider the following linear parameterization  $\lambda(s) = \tilde{\lambda}_t + s(\lambda - \tilde{\lambda}_t)$  for  $s \in [0, 1]$  and correspondingly

$$\begin{aligned} f_{t, \mathbf{b}_{t-1}}(\lambda(s)) - f_{t, \mathbf{b}_{t-1}}(\tilde{\lambda}_t) &= \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\tilde{\lambda}_t, \mathbf{a}, r)^\top (\lambda(s) - \tilde{\lambda}_t) + \mathbb{E}_{(\mathbf{a}, r) \sim F} \int_{\mathbf{a}^\top \lambda(s)}^{\mathbf{a}^\top \tilde{\lambda}_t} \left( \mathbb{I}_{\{r > v\}} - \mathbb{I}_{\{r > \mathbf{a}^\top \tilde{\lambda}_t\}} \right) dv \\ &= \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\tilde{\lambda}_t, \mathbf{a}, r)^\top (\lambda(s) - \tilde{\lambda}_t) + \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \int_{\mathbf{a}^\top \lambda(s)}^{\mathbf{a}^\top \tilde{\lambda}_t} \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\lambda}_t) - F_{\mathbf{a}}^r(v) \right) dv \\ &\leq \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\tilde{\lambda}_t, \mathbf{a}, r)^\top (\lambda(s) - \tilde{\lambda}_t) + \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \int_{\mathbf{a}^\top \lambda(s)}^{\mathbf{a}^\top \tilde{\lambda}_t} \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\lambda}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \lambda(s)) \right) dv \\ &= \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\tilde{\lambda}_t, \mathbf{a}, r)^\top (\lambda(s) - \tilde{\lambda}_t) + \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left\{ \left( \mathbf{a}^\top \tilde{\lambda}_t - \mathbf{a}^\top \lambda(s) \right) \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\lambda}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \lambda(s)) \right) \right\} \\ &= s \mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\tilde{\lambda}_t, \mathbf{a}, r)^\top (\lambda - \tilde{\lambda}_t) + s \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left\{ \left( \mathbf{a}^\top \tilde{\lambda}_t - \mathbf{a}^\top \lambda \right) \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\lambda}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \lambda(s)) \right) \right\} \\ &= -s\eta + s \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left\{ \left( \mathbf{a}^\top \tilde{\lambda}_t - \mathbf{a}^\top \lambda \right) \left( F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\lambda}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \lambda(s)) \right) \right\} \\ &\leq -s\eta + s \frac{2m \bar{A} \bar{r}}{\underline{A}} \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left| F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\lambda}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \lambda(s)) \right| \right], \end{aligned}$$

where in the last inequality we use the fact that both  $\tilde{\boldsymbol{\lambda}}_t$  and  $\boldsymbol{\lambda}$  belong to  $\Omega$  and that  $|\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t - \mathbf{a}^\top \boldsymbol{\lambda}|$  is upper bounded by absolute constants (cf. Lemma F.12 and Corollary F.13). By the continuity of  $F_{\mathbf{a}}^r$ , we have

$$\lim_{s \rightarrow 0} \mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left| F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}(s)) \right| \right] = 0.$$

Consequently, there must exist  $s > 0$ , such that  $\mathbb{E}_{\mathbf{a} \sim F^{\mathbf{a}}} \left[ \left| F_{\mathbf{a}}^r(\mathbf{a}^\top \tilde{\boldsymbol{\lambda}}_t) - F_{\mathbf{a}}^r(\mathbf{a}^\top \boldsymbol{\lambda}(s)) \right| \right] < \frac{\eta \underline{A}}{4m\bar{A}\bar{r}}$ . Plugging back and we conclude that  $f_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}(s)) - f_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t) \leq -\frac{s\eta}{2} < 0$ , contracting the fact that  $\tilde{\boldsymbol{\lambda}}_t$  is the minimizer of  $f_{t, \mathbf{b}_{t-1}}(\cdot)$  among all  $\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m$ . Therefore it must be that  $\mathbb{E}_{(\mathbf{a}, r) \sim F} \phi_{t, \mathbf{b}_{t-1}}(\tilde{\boldsymbol{\lambda}}_t, \mathbf{a}, r)^\top (\boldsymbol{\lambda} - \tilde{\boldsymbol{\lambda}}_t) \geq 0$  and we conclude the proof of the first case. The proof of the second and third case of the Lemma is in spirit very similar. We again argue by contradiction. Suppose there exists  $\boldsymbol{\lambda} \in \mathbb{R}_{\geq 0}^m$  such that  $\sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*, \mathbf{a}_j, r_j)^\top (\boldsymbol{\lambda} - \boldsymbol{\lambda}_t^*) = -\frac{2m^2 \bar{A} \bar{r}}{\underline{A}} - \eta < -\frac{2m^2 \bar{A} \bar{r}}{\underline{A}}$ . Consider the linear parameterization  $\boldsymbol{\lambda}(s) = \boldsymbol{\lambda}_t^* + s(\boldsymbol{\lambda} - \boldsymbol{\lambda}_t^*)$  for  $s \in [0, 1]$  and correspondingly

$$\begin{aligned} & g_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}(s)) - g_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*) \\ &= \frac{1}{T-t} \sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*, \mathbf{a}_j, r_j)^\top (\boldsymbol{\lambda}(s) - \boldsymbol{\lambda}_t^*) + \frac{1}{T-t} \sum_{j=t+1}^T \int_{\mathbf{a}_j^\top \boldsymbol{\lambda}(s)}^{\mathbf{a}_j^\top \boldsymbol{\lambda}_t^*} \left( \mathbb{I}_{\{r_j > v\}} - \mathbb{I}_{\{r_j > \mathbf{a}_j^\top \boldsymbol{\lambda}_t^*\}} \right) dv, \\ &= \frac{1}{T-t} \left( -s\eta - s \frac{2m^2 \bar{A} \bar{r}}{\underline{A}} + \sum_{j=t+1}^T \int_{\mathbf{a}_j^\top \boldsymbol{\lambda}(s)}^{\mathbf{a}_j^\top \boldsymbol{\lambda}_t^*} \left( \mathbb{I}_{\{r_j > v\}} - \mathbb{I}_{\{r_j > \mathbf{a}_j^\top \boldsymbol{\lambda}_t^*\}} \right) dv \right), \\ &\leq \frac{1}{T-t} \left( -s\eta - s \frac{2m^2 \bar{A} \bar{r}}{\underline{A}} + \sum_{j=t+1}^T \int_{\mathbf{a}_j^\top \boldsymbol{\lambda}(s)}^{\mathbf{a}_j^\top \boldsymbol{\lambda}_t^*} \left( \mathbb{I}_{\{r_j > \mathbf{a}_j^\top \boldsymbol{\lambda}(s)\}} - \mathbb{I}_{\{r_j > \mathbf{a}_j^\top \boldsymbol{\lambda}_t^*\}} \right) dv \right), \\ &= \frac{1}{T-t} \left( -s\eta - s \frac{2m^2 \bar{A} \bar{r}}{\underline{A}} + \sum_{j=t+1}^T (\mathbf{a}_j^\top \boldsymbol{\lambda}_t^* - \mathbf{a}_j^\top \boldsymbol{\lambda}(s)) \left( \mathbb{I}_{\{r_j > \mathbf{a}_j^\top \boldsymbol{\lambda}(s)\}} - \mathbb{I}_{\{r_j > \mathbf{a}_j^\top \boldsymbol{\lambda}_t^*\}} \right) \right), \\ &= \frac{1}{T-t} \left( -s\eta - s \frac{2m^2 \bar{A} \bar{r}}{\underline{A}} + s \sum_{j=t+1}^T (\mathbf{a}_j^\top \boldsymbol{\lambda}_t^* - \mathbf{a}_j^\top \boldsymbol{\lambda}) \left( \mathbb{I}_{\{r_j > \mathbf{a}_j^\top \boldsymbol{\lambda}(s)\}} - \mathbb{I}_{\{r_j > \mathbf{a}_j^\top \boldsymbol{\lambda}_t^*\}} \right) \right), \\ &\leq \frac{1}{T-t} \left( -s\eta - s \frac{2m^2 \bar{A} \bar{r}}{\underline{A}} + s \frac{2m \bar{A} \bar{r}}{\underline{A}} \sum_{j=t+1}^T \left| \mathbb{I}_{\{r_j > \mathbf{a}_j^\top \boldsymbol{\lambda}(s)\}} - \mathbb{I}_{\{r_j > \mathbf{a}_j^\top \boldsymbol{\lambda}_t^*\}} \right| \right). \end{aligned}$$

Consider the index set  $W \triangleq \{j \in [t+1, T] : r_j = \mathbf{a}_j^\top \boldsymbol{\lambda}_t^*\}$ . Since  $F_{\mathbf{a}}^r$  is a continuous distribution for any  $\mathbf{a}$ , where  $\mathbf{a} \in \mathbb{R}_{\geq 0}^m$  is a  $d$ -dimensional vector, furthermore  $\{(\mathbf{a}_j, r_j)\}_{j=t+1}^T$  are independent of each other, it follows that  $\mathbb{P}(|W| > m) = 0$  (Otherwise we have  $m+1$  *i.i.d.* drawn vectors  $(\mathbf{a}_j, r_j)$  belonging to the same  $m$ -dimensional hyperplane  $y = \mathbf{a}^\top \boldsymbol{\lambda}_t^*$ ). For fixed set of  $\{(\mathbf{a}_j, r_j)\}_{j=t+1}^T$  and  $\boldsymbol{\lambda}_t^*$ , there exists a small enough  $s > 0$ , such that

$$\mathbb{I}_{\{r_j > \mathbf{a}_j^\top \boldsymbol{\lambda}(s)\}} - \mathbb{I}_{\{r_j > \mathbf{a}_j^\top \boldsymbol{\lambda}_t^*\}} \neq 0 \quad \Leftrightarrow \quad i \in W.$$

Hence almost surely we have

$$\sum_{j=t+1}^T \left| \mathbb{I}_{\{r_j > \mathbf{a}_j^\top \boldsymbol{\lambda}(s)\}} - \mathbb{I}_{\{r_j > \mathbf{a}_j^\top \boldsymbol{\lambda}_t^*\}} \right| \leq |W| \leq m.$$



Plugging back and we conclude that  $g_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}(s)) - g_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*) \leq -\frac{s\eta}{T-t} < 0$ , contradicting the fact that  $\boldsymbol{\lambda}_t^*$  is the minimizer of  $g_{t, \mathbf{b}_{t-1}}$ . Therefore, it must be that  $\sum_{j=t+1}^T \phi_{t, \mathbf{b}_{t-1}}(\boldsymbol{\lambda}_t^*, \mathbf{a}_j, r_j)^\top (\tilde{\boldsymbol{\lambda}}_t - \boldsymbol{\lambda}_t^*) \geq -\frac{2m^2 \bar{A} \bar{r}}{\underline{A}}$ , a.s. and we conclude the second case of the proof. The third case is nearly identical to the second half, and we omit its proof. Q.E.D.

**Lemma F.15.** *Suppose Assumption 2.2 holds. Then there exists  $w > 0$  and  $\theta \in (0, 2\pi)$ , such that for any  $\mathbf{a} \in \text{supp}(F^{\mathbf{a}})$  and any  $\boldsymbol{\lambda}_1, \boldsymbol{\lambda}_2$ ,*

$$\mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}, \mathbf{a}' \in \mathcal{B}(\mathbf{a}, r)} (\mathbf{a}'^\top \boldsymbol{\lambda}_1 - \mathbf{a}'^\top \boldsymbol{\lambda}_2)^2 \geq l_f \frac{\pi^{m/2}}{\Gamma(\frac{m}{2} + 1)} \left( \frac{r \wedge w}{6} \sin(\theta) \right)^{m+2} \|\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2\|^2,$$

where  $\mathcal{B}(\mathbf{a}, r)$  is the ball centered at  $\mathbf{a}$  with radius  $r > 0$ .

*Proof.* By Assumption 2.2, since  $\text{supp}(F^{\mathbf{a}})$  is convex,  $\text{supp}(F^{\mathbf{a}})$  is a Lipschitz domain, thus satisfies the *uniform cone condition*. That is, there exist an opening angle  $\theta \in (0, \pi)$ , and a radius  $w > 0$ , such that for every point  $\mathbf{a} \in \text{supp}(F^{\mathbf{a}})$ , there exists a cone  $C_{\mathbf{a}}$  with direction represented by a unit vector  $\mathbf{v}_{\mathbf{a}} \in \mathbb{R}^m$  and vertex at  $\mathbf{a}$ , defined by

$$C_{\mathbf{a}} = \{ \mathbf{a} + s\mathbf{y} : s \in [0, w], \mathbf{y} \in \mathbb{S}^{m-1}, \langle \mathbf{y}, \mathbf{v}_{\mathbf{a}} \rangle \geq \cos(\theta) \},$$

such that  $C_{\mathbf{a}} \subset \text{supp}(F^{\mathbf{a}})$ , where  $\mathbb{S}^{m-1}$  is the unit sphere in  $\mathbb{R}^m$ .

Furthermore, the probability density associated to  $F^{\mathbf{a}}$  is bounded from below by a constant  $l_f > 0$ .

Observe that ball  $\mathcal{B}(\mathbf{a} + \frac{r \wedge w}{2} \mathbf{v}_{\mathbf{a}}, \frac{r \wedge w}{2} \sin(\theta))$  is contained in  $\mathcal{B}(\mathbf{a}, r) \cap C_{\mathbf{a}}$ . In what follows, we shall refer to this ball by  $\mathcal{B}$  for notational simplicity, and denote by  $\mathbf{q}^* \triangleq \mathbf{a} + \frac{r \wedge w}{2} \mathbf{v}_{\mathbf{a}}$  and  $r^* \triangleq \frac{r \wedge w}{2} \sin(\theta)$ . Without loss of generality, suppose  $\mathbf{q}^{*\top} (\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2) \geq 0$ , otherwise we can consider  $\mathbf{q}^{*\top} (\boldsymbol{\lambda}_2 - \boldsymbol{\lambda}_1)$  instead. Let  $\mathbf{q} = \mathbf{q}^* + \frac{2}{3} r^* \frac{(\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2)}{\|\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2\|}$ , and thus

$$\mathbf{q}^\top (\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2) = \mathbf{q}^{*\top} (\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2) + \frac{2}{3} r^* \frac{(\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2)^\top}{\|\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2\|} (\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2) \geq \frac{2}{3} r^* \|\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2\|.$$

Clearly,  $\mathcal{B}(\mathbf{q}, \frac{r^*}{3}) \subset \mathcal{B}$ . For any  $\mathbf{q}' \in \mathcal{B}(\mathbf{q}, \frac{r^*}{3})$ ,

$$\begin{aligned} \mathbf{q}'^\top (\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2) &= \mathbf{q}^\top (\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2) - (\mathbf{q} - \mathbf{q}')^\top (\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2) \\ &\geq \frac{2}{3} r^* \|\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2\| - \|\mathbf{q} - \mathbf{q}'\| \cdot \|\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2\| \\ &\geq \frac{2}{3} r^* \|\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2\| - \frac{r^*}{3} \|\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2\| = \frac{r^*}{3} \|\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2\|. \end{aligned}$$

Therefore, we have

$$\begin{aligned} \mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}, \mathbf{a}' \in \mathcal{B}(\mathbf{a}', r^*)} (\mathbf{a}'^\top \boldsymbol{\lambda}_1 - \mathbf{a}'^\top \boldsymbol{\lambda}_2)^2 &\geq \mathbb{E}_{\mathbf{a}' \sim F^{\mathbf{a}}, \mathbf{a}' \in \mathcal{B}(\mathbf{q}, \frac{r^*}{3})} (\mathbf{a}'^\top \boldsymbol{\lambda}_1 - \mathbf{a}'^\top \boldsymbol{\lambda}_2)^2, \\ &\geq \mathbb{P}\left(\mathbf{a}' \in \mathcal{B}(\mathbf{q}, \frac{r^*}{3})\right) \frac{r^{*2}}{9} \|\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2\|^2, \\ &\geq l_f \frac{\pi^{m/2}}{\Gamma(\frac{m}{2} + 1)} \frac{r^{*m}}{3^m} \frac{r^{*2}}{9} \|\boldsymbol{\lambda}_1 - \boldsymbol{\lambda}_2\|^2, \end{aligned}$$

where the last inequality follows from the fact that  $F^{\mathbf{a}}$  has lower bounded density by Assumption 2.2, the volume of the ball  $\mathcal{B}(\mathbf{q}, \frac{r^*}{3})$  is  $\frac{\pi^{m/2}}{\Gamma(\frac{m}{2} + 1)} \frac{r^{*m}}{3^m}$ , and that  $\mathcal{B} \subseteq \mathcal{B}(\mathbf{a}, r) \cap C_{\mathbf{a}} \subseteq \text{supp}(F^{\mathbf{a}})$ . The lemma thus follows from the fact that  $\mathcal{B} \subseteq \mathcal{B}(\mathbf{a}, r) \cap C_{\mathbf{a}} \subseteq \mathcal{B}(\mathbf{a}, r)$ . Q.E.D.

**Lemma F.16.** Suppose  $\mathbf{a} \in \text{supp}(F^{\mathbf{a}})$  and  $\boldsymbol{\lambda}_t^*$  satisfy

$$\frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} - \frac{1}{8}c_{\nu}^{\frac{1}{2}} \leq \mathbf{a}^{\top} \boldsymbol{\lambda}_t^* \leq \frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} + \frac{1}{8}c_{\nu}^{\frac{1}{2}}.$$

Then for all  $\mathbf{a}' \in \mathcal{B}(\mathbf{a}, r_1)$ , where  $\mathcal{B}(\mathbf{a}, r_1)$  is a ball centered at  $\mathbf{a}$  with radius  $r_1 = \frac{1}{8} \left( \left( c_L + \sqrt{m} \frac{\bar{r}}{\underline{A}} \right)^{-1} \wedge 1 \right) c_{\nu}^{\frac{1}{2}}$ , it holds that

$$\frac{l_{\mathbf{a}'} + r_{\mathbf{a}'}}{2} - \frac{1}{4}c_{\nu}^{\frac{1}{2}} \leq \mathbf{a}'^{\top} \boldsymbol{\lambda}_t^* \leq \frac{l_{\mathbf{a}'} + r_{\mathbf{a}'}}{2} + \frac{1}{4}c_{\nu}^{\frac{1}{2}}.$$

*Proof.* We only prove one direction of this inequality as the other direction is nearly identical. For any  $\mathbf{a}' \in \mathcal{B}(\mathbf{a}, r_1)$ , we have that

$$\begin{aligned} \mathbf{a}'^{\top} \boldsymbol{\lambda}_t^* &= \mathbf{a}^{\top} \boldsymbol{\lambda}_t^* + (\mathbf{a} - \mathbf{a}')^{\top} \boldsymbol{\lambda}_t^* \\ &\leq \mathbf{a}^{\top} \boldsymbol{\lambda}_t^* + \|\mathbf{a} - \mathbf{a}'\| \|\boldsymbol{\lambda}_t^*\| \quad (\text{by Cauchy-Schwarz}), \\ &\leq \frac{l_{\mathbf{a}} + r_{\mathbf{a}}}{2} + \frac{1}{8}c_{\nu}^{\frac{1}{2}} + \mathbf{a}^{\top} \boldsymbol{\lambda}_t^* + \|\mathbf{a} - \mathbf{a}'\| \|\boldsymbol{\lambda}_t^*\| \\ &\leq \frac{l_{\mathbf{a}'} + r_{\mathbf{a}'}}{2} + c_L \|\mathbf{a}' - \mathbf{a}\| + \frac{1}{8}c_{\nu}^{\frac{1}{2}} + \|\mathbf{a} - \mathbf{a}'\| \|\boldsymbol{\lambda}_t^*\| \quad (\text{by Lipschitzness of } l_{\mathbf{a}} \text{ and } r_{\mathbf{a}}) \\ &\leq \frac{l_{\mathbf{a}'} + r_{\mathbf{a}'}}{2} + \frac{1}{8}c_{\nu}^{\frac{1}{2}} + \frac{1}{8} \left( c_L + \sqrt{m} \frac{\bar{r}}{\underline{A}} \right)^{-1} c_{\nu}^{\frac{1}{2}} \times \left( c_L + \sqrt{m} \frac{\bar{r}}{\underline{A}} \right) \\ &\quad (\text{by definition of } r_1 \text{ and Lemma F.12}) \\ &= \frac{l_{\mathbf{a}'} + r_{\mathbf{a}'}}{2} + \frac{1}{4}c_{\nu}^{\frac{1}{2}}, \end{aligned}$$

completing the proof. Q.E.D.