ON ACYLINDRICAL TREE ACTIONS AND OUTER AUTOMORPHISM GROUP OF BAUMSLAG-SOLITAR GROUPS

BRATATI SOM AND DAXUN WANG

ABSTRACT. This paper explores acylindrical actions on trees, building on previous works related to the mapping class group and projection complexes. We demonstrate that the quotient action of an acylindrical action of a group on a tree by an equivariant family of subgroups remains acylindrical. We establish criteria for ensuring that this quotient action is non-elementary acylindrical, thus preserving the group's acylindrical hyperbolicity. Additionally, we show that the fundamental group of a graph of groups admits the largest acylindrical action on its Bass-Serre tree under certain conditions. As an application, we analyze the outer automorphism group of non-solvable Baumslag-Solitar groups, we prove its acylindrical hyperbolicity, highlighting the differences between various tree actions and identifying the largest acylindrical action.

1. Introduction

The study of acylindrical action on a hyperbolic space was initiated by Bowditch in [Bow08]. Roughly speaking, it is the action whereby the number of elements that simultaneously coarsely stabilize sufficiently distant points is uniformly bounded. The study of acylindrical action on hyperbolic spaces was developed by Osin [Osi16] building on the work of Bowditch [Bow08] and Sela [Sel97]. This has proven to be an effective tool to study many important families of groups. In fact, many groups (such as mapping class groups and the outer automorphism group of free groups, as well as most 3-manifold groups, etc.) admit a non-elementary acylindrical action on some hyperbolic spaces.

In this paper, we are interested in the quotient action of an acylindrical hyperbolic action. Our motivation for studying this quotient action comes from the algebraic counterpart to Thurston's Dehn filling theorem, in relation to relatively hyperbolic groups [Osi07], [GM08] and, more generally, to acylindrically hyperbolic groups [DGO17].

Specifically, similar to how new manifolds with hyperbolic structures are created from a 3-manifold with an existing hyperbolic structure via Dehn filling, in the context of group theory, taking quotients of an acylindrically hyperbolic group by certain "special" normal subgroups produces new acylindrically hyperbolic groups. In [DHS21], Dahmani, Hagen, and Sisto showed that quotients of the mapping class group by normal subgroups generated by suitable powers of Dehn twists preserve the acylindricity of the group. In proving this theorem, the authors relied on the fact established by Dahmani [Dah18] that the curve graph has the structure of a composite projection graph. Later, Clay and Mangahas in [CM22] investigated a scenario where groups act on projection complexes with a sufficient number of WPD hyperbolic elements. They demonstrated the acylindricity of the quotient group by normal closure of subgroup generated by appropriate "equivariant L-spinning family." As an application of this construction, Clay and Mangahas present new examples of acylindrically hyperbolic groups obtained from quotients of mapping class groups of orientable surfaces. More generally, for actions on hyperbolic spaces, Dahmani, Guirardel and Osin [DGO17]

introduced the notion of very rotating families which provides a natural framework for developing a geometric version of small cancellation theory. They prove similar quotient results using normal subgroups generated by the rotating families. Even though simplicial trees can be seen as projection complexes naturally (see Section 3.1, [CMM21]) as well as 0-hyperbolic spaces, none of the above mentioned theories apply to trees directly. In this paper, we not only develope a similar theory for actions on trees, but our conditions are more general as we drop both rotating and spinning condition and work with only an equivariant family of subgroups. We recall the definition of an equivariant family of subgroups below.

Let G be a group acting on a tree T. For each vertex v of T, let R_v be a subgroup of the stabilizer of v in G. We say the family of subgroups $\{R_v\}$ is an equivariant family of subgroups of G if $gR_vg^{-1}=R_{qv}$, for every g in G and every vertex v of T.

We say an action of a group on a tree is an acylindrical tree action if it is acylindrical. We first show that the quotient action of an acylindrical tree action by an equivariant family of subgroups is always acylindrical (see Theorem 4.4). Further, we provide sufficient conditions for an equivariant family of subgroups that ensure the quotient action is non-elementary acylindrical and as a consequence the quotient group remains acylindrically hyperbolic.

Theorem A. Let G be a group acting non-elementary acylindrically on a simplicial tree T, and let N be a normal subgroup of G. If N is generated by an equivariant family of subgroups $\{R_v\}$, then the quotient space T/N is a tree. Moreover, let (Γ, \mathcal{G}) be the quotient graph of groups structure of G. The quotient action $G/N \curvearrowright T/N$ is a non-elementary acylindrical tree action if $\{R_v\}$ satisfies at least one of the following conditions:

- (1) the first Betti number $b_1(\Gamma) \geq 2$ where $\Gamma = T/G$.
- (2) there are two vertices \hat{u} and \hat{v} in Γ such that $\langle G_{\hat{e}}, R_{\hat{u}} \rangle \neq G_{\hat{u}}$ and $\langle G_{\hat{f}}, R_{\hat{v}} \rangle \neq G_{\hat{v}}$, where \hat{e} and \hat{f} are the edges on the geodesic path $[\hat{u}, \hat{v}]$ adjacent to \hat{u} and \hat{v} respectively. Further, there exist two elements $g, h \in G_{\hat{v}} \setminus \langle G_{\hat{f}}, R_{\hat{v}} \rangle$ such that $gh^{-1} \notin \langle G_{\hat{f}}, R_{\hat{v}} \rangle$.
- (3) Γ contains a immersed circuit. Further, there exists a vertex \hat{v} and an edge \hat{e} adjacent to \hat{v} on the circuit such that $\langle G_{\hat{e}}, R_{\hat{v}} \rangle \neq G_{\hat{v}}$.

We note that, by Bass-Serre theory, any group G acting on a tree without edge inversions induces a graph of groups (Γ, \mathcal{G}) of G, which consists of a graph Γ together with a group for each vertex and edge of Γ , and monomorphisms from each edge group to the adjacent vertex groups. Conversely, any graph of groups (Γ, \mathcal{G}) has a canonical associated group $\pi_1(\Gamma, \mathcal{G})$, called the fundamental group of graph of groups, and a tree X, called the Bass-Serre tree, such that $\pi_1(\Gamma, \mathcal{G})$ acts on the Bass-Serre tree X. The fundamental theorem of Bass-Serre theory says that these processes are in fact mutually inverse.

Two types of graphs of groups are the simplest interesting examples. One is called an *edge of groups*, where the underlying graph consists of a single edge. The other one is called a *loop of groups*, where the underlying graph consists of a single loop. By Bass-Serre theory, the fundamental group of an edge of groups is a free amalgamated product, while the fundamental group of a loop of groups is a HNN-extension. As a consequence of Theorem A, we obtain the following two corollaries about quotient actions for these two special cases.

Corollary B. Let $G = A *_C B$ have a non-elementary acylindrical action on a tree T, and let N be a normal subgroup of G generated by an equivariant family of subgroups $\{R_v\}$. The quotient action $G/N \curvearrowright T/N$ is a non-elementary acylindrical tree action if $\{R_v\}$ satisfies the following:

(1)
$$\langle C, R_A \rangle \neq A$$
 and $\langle C, R_B \rangle \neq B$; and

(2) there exists two different elements g and h in $B \setminus \langle C, R_B \rangle$ such that $gh^{-1} \notin \langle C, R_B \rangle$.

Corollary C. Let $G = A*_{H \sim_{\phi} K}$ have a non-elementary acylindrical action on a tree T. and let N be a normal subgroup of G generated by an equivariant family of subgroups $\{R_v\}$. The quotient action $G/N \curvearrowright T/N$ is a non-elementary acylindrical tree action if $\{R_v\}$ satisfies either $\langle H, R_A \rangle \neq A$ or $\langle K, R_A \rangle \neq A$.

Note that any given group with an acylindrical action may admit many acylindrical actions on different hyperbolic spaces. It is natural to ask which acylindrical action is the "best." In [ABO19], Abbott, Balasubramanya, and Osin introduced a partial order on the set of acylindrical actions which allows us to optimize all acylindrical actions. In particular, a larger action in this poset reveals, to some extent, more information about the group. Therefore, it is natural to consider whether a largest action exists in this poset. It is proven that every hierarchically hyperbolic group [ABD21] and many graph products [Val21] admit a largest acylindrical hyperbolic action. In the second part of this paper, we study existence of such largest acylindrical actions that arise from graphs of groups. In [MO15], Minasyan and Osin provided sufficient conditions for the tree action $\pi_1(\Gamma, \mathcal{G}) \curvearrowright X$ to be acylindrical. In this paper, we give sufficient conditions for the fundamental group of graph of groups $\pi_1(\Gamma, \mathcal{G})$ to admit a largest acylindrical action.

Theorem D. Let (Γ, \mathcal{G}) be a finite graph of groups, and let T be the Bass-Serre tree of (Γ, \mathcal{G}) . Denote the fundamental group $\pi_1(\Gamma, \mathcal{G})$ by G. If the action $G \curvearrowright T$ is acylindrical, and if each vertex group acts elliptically whenever G acts acylindrically on any hyperbolic space S, then G admits the largest acylindrical action on T.

We remark that the above theorem shares a similar idea as Proposition 4.13 of [ABO19]. However, when restricting to tree actions which are specifically acylindrical, we obtained a proof of existence of a largest such action by using Bass-Serre theory.

Next, as an application of the previously mentioned results along with some additional findings, we concentrate on a specific group that exhibits interesting graph of groups structures: the outer automorphism group of non-solvable Baumslag-Solitar groups. These groups have been studied particularly in [CL83], [Cla09], [Col78] and [GHMR00]. Recall that Baumslag-Solitar groups have the following standard presentation

$$BS(p,q) = \langle x, t \mid tx^p t^{-1} = x^q \rangle$$

where p,q are both non-zero integers. For non-solvable Baumslag-Solitar groups neither |p| nor |q| equals 1. In general, acylindricity of the automorphism group $\operatorname{Aut}(G)$ has been studied by others with some conditions on G. Genevois [Gen19] proved that if G is a one-ended hyperbolic group then $\operatorname{Aut}(G)$ is acylindrically hyperbolic. Genevois and Horbez [GH21] showed that if G is an infinitely-ended finitely generated group then $\operatorname{Aut}(G)$ is acylindrically hyperbolic. For the outer automorphism group $\operatorname{Out}(G)$, it is known that $\operatorname{Out}(F_n)$ for $n \geq 2$ is acylindrically hyperbolic [BF10]. But none of these cover $\operatorname{Out}(\operatorname{BS}(p,q))$ as $\operatorname{BS}(p,q)$ is neither hyperbolic nor infinite-ended. In [Cla09], Clay computed the presentation of $\operatorname{Out}(\operatorname{BS}(p,q))$ when p properly divides q. Further, Clay showed that $\operatorname{Out}(\operatorname{BS}(p,q))$ admits two natural graph of groups structures $(\Gamma_1, \mathcal{G}_1)$ and $(\Gamma_2, \mathcal{G}_2)$, where the underlying graph Γ_1 is an infinite ray and the underlying graph Γ_2 is a single edge. By Bass-Serre theory, these two graph of groups structures correspond to two actions of $\operatorname{Out}(\operatorname{BS}(p,q))$ on different Bass-Serre trees X_1 and X_2 respectively. We examine both actions in detail and compare them. The following theorem summarizes the distinctions between these two tree actions.

Theorem E. Let G = Out(BS(p,q)) where q = pn for some p, |n| > 1. Let X_1 be the Bass-Serre tree of the infinite ray of groups $(\Gamma_1, \mathcal{G}_1)$ of G and let X_2 be the Bass-Serre tree of the edge of groups (Γ, \mathcal{G}) of G. Then we have the following results:

- (1) $G \curvearrowright X_1$ is not an acylindrical action while $G \curvearrowright X_2$ is non-elementary acylindrical, thus $G \cap X_1$ is not equivalent to $G \cap X_2$;
- (2) $G \curvearrowright X_1$ and $G \curvearrowright X_2$ are both non-elementary WPD actions;
- (3) $G \curvearrowright X_1$ and $G \curvearrowright X_2$ have the same set of hyperbolic elements;
- (4) $G \curvearrowright X_2$ is the largest acylindrical hyperbolic action.

Outline of the paper. In the first part of the paper, we establish acylindricity results for general graphs of groups. In Section 2, we recall the basic notions about Bass-Serre theory. In Section 3, we recall the definitions and results about acylindrical actions. In Section 4, we examine quotient actions of groups acting on trees, provide a proof of Theorem A, and derive Corollaries B and C. This leads to the conclusion that the quotient group by normal subgroups generated by an equivariant family of subgroups is acylindrically hyperbolic. Section 5 discusses the largest acylindrical action and aims to prove another main result of the paper, Theorem D.

Next we specifically focuses on Out(BS(p,q)) in Section 6. We briefly describe the two graph of groups structures for Out(BS(p,q)) from Clay's work in [Cla09], along with their corresponding actions on trees, and compare these actions. We demonstrate that Out(BS(p,q))has a non-elementary acylindrical action on a tree, confirming that it is an acylindrically hyperbolic group. As an application of the results from Sections 4 and 5, we show that Out(BS(p,q)) has a largest acylindrical action, and that acylindricity is preserved in quotient actions by certain normal subgroups. We illustrate the quotients of Out(BS(p,q))through Example 6.9 and Example 6.10. In addition, we prove that the two actions are both non-elementary WPD, providing an alternative proof of the acylindrical hyperbolicity of Out(BS(p,q)) when p divides q properly. Finally, at the end of Section 6, we prove our last main result, Theorem E.

Acknowledgments. We would like to thank our advisor Johanna Mangahas for her exceptional guidance and encouragement. We would also like to thank Carolyn Abbott, Sahana Balasubramanya, Indira Chatterji, Ashot Minasyan and Wenyuan Yang for helpful comments and conversations. The first author acknowledges travel support from the Simons Foundation (965204, JM).

2. Bass-Serre Theory

In this section we recall some basic notions of the Bass-Serre theory. There are two equivalent versions of this theory; details can be found in [Ser80] and [Bas93]. We will primarily work with [Ser80]. However we will occasionally use tools from [Bas93] to obtain more clear and easier proofs.

2.1. Graph of groups.

Definition 2.1. [Ser 80, Definition 2.1.1] A graph Γ consists of a set $V(\Gamma)$, a set $E(\Gamma)$ and two maps

$$E(\Gamma) \to V(\Gamma) \times V(\Gamma), \quad e \mapsto (o(e), t(e))$$

and

$$E(\Gamma) \to E(\Gamma), \quad e \mapsto \overline{e}$$

which satisfy the following condition: for each $e \in E(\Gamma)$ we have $\overline{\overline{e}} = e$, $\overline{e} \neq e$ and $o(e) = t(\overline{e})$. An element $v \in V(\Gamma)$ is called a vertex of Γ ; An element $e \in E(\Gamma)$ is called an oriented edge of Γ , and \overline{e} is called the *inverse* edge of e. The vertex $o(e) = t(\overline{e})$ is called the *origin* of e, and the vertex $t(e) = o(\overline{e})$ is called the terminus of e. These two vertices are called the extremities of e.




FIGURE 1. Two examples of graphs. The left graph consists of one edge with different extremities, whereas the right graph consists of one edge with the same extremities.

Definition 2.2. A graph of groups (Γ, \mathcal{G}) consists of a connected graph Γ , and a family of groups \mathcal{G} , together with:

- (1) a vertex group $G_v \in \mathcal{G}$ for each $v \in V(\Gamma)$;
- (2) an edge group $G_e \in \mathcal{G}$ for each $e \in E(\Gamma)$, such that $G_e = G_{\overline{e}}$; and
- (3) a monomorphism $\alpha_e : G_e \hookrightarrow G_{t(e)}$ for each $e \in E(\Gamma)$.

Further, we say (Γ, \mathcal{G}) is a *finite* graph of groups if the graph Γ is both locally finite and has finite diameter.

There are two particular cases of graph of groups, which we will describe in Example 2.3 and Example 2.4. A more general graph of groups can be seen as a combination of these following two types.

Example 2.3. An *edge of groups* is pictured as follows.

$$G_{o(e)} \xrightarrow{G_e} G_{t(e)}$$

The monomorphisms are $\alpha_e: G_e \hookrightarrow G_{t(e)}$ and $\alpha_{\overline{e}}: G_e \hookrightarrow G_{o(e)}$.

Example 2.4. A loop of groups is pictured as follows.



The monomorphisms are $\alpha_f: G_f \hookrightarrow G_v$ and $\alpha_{\overline{f}}: G_f \hookrightarrow G_v$.

Definition 2.5. Let (Γ, \mathcal{G}) be a graph of groups. Consider the group $F(\Gamma, \mathcal{G})$ generated by the vertex groups G_v for $v \in V(\Gamma)$ and the elements e of $E(\Gamma)$, subject to the following relations:

$$\overline{e} = e^{-1}$$
 and $e\alpha_e(g)e^{-1} = \alpha_{\overline{e}}(g)$ for any $e \in E(\Gamma), g \in G_e$

i.e., the group $F(\Gamma, \mathcal{G})$ is the quotient of the free product of all G_v and edges $e \in E(\Gamma)$ by the normal subgroup generated by elements $e\overline{e}$ and $e\alpha_e(g)e^{-1}\alpha_{\overline{e}}(g)^{-1}$.

Definition 2.6. [Ser80, Definition 5.1.9] Let c be a path in Γ . We let $e_1, ..., e_n$ denote the edges of c where n is the length of c, and put $v_i = o(e_{i+1}) = t(e_i)$. A word of type c is an element in $F(\Gamma, \mathcal{G})$ of the form

$$g_0e_1g_1e_2...e_ng_n$$

where $g_i \in G_{v_i}$.

2.2. Fundamental group of graph of groups.

Definition 2.7. Two equivalent definitions of the fundamental group of a graph of groups:

- (a) Let (Γ, \mathcal{G}) be a graph of groups, and let v_0 be a vertex of Γ . The fundamental group of (Γ, \mathcal{G}) at v_0 , denoted by $\pi_1(\Gamma, v_0)$, is the subgroup of $F(\mathcal{G}, \Gamma)$ consists of words whose type is a circuit based at v_0 (i.e. the path $c = (e_1, ..., e_n)$ satisfies $o(e_1) = t(e_n) = v_0$).
- (b) Let T be a maximal tree of Γ . By $E(\Gamma \setminus T)$, we denote the set of edges in $E(\Gamma)$ but not in E(T). The fundamental group of (Γ, \mathcal{G}) at T, denoted by $\pi_1(\Gamma, T)$, is the group generated by the vertex groups G_v for $v \in V(\Gamma)$ and edges e of $E(\Gamma \setminus T)$, subject to the following relations:

$$\overline{e} = e^{-1}$$
 and $e\alpha_e(g)e^{-1} = \alpha_{\overline{e}}(g)$ for any $e \in E(\Gamma \setminus T), g \in G_e$

In other words, it is the quotient of $F(\Gamma, \mathcal{G})$ by the normal subgroup generated by the set of edges in T. Each geometric edge in Γ but not in T is called a *stable letter* of $\pi_1(\Gamma, T)$. We note that the number of stable letters is independent of the choice of T. This number is called the *first Betti number* of Γ , and we denote it by $b_1(\Gamma)$ (i.e., $b_1(\Gamma) = \frac{1}{2}|E(\Gamma \setminus T)|$).

We now describe the fundamental groups of the two particular cases of graph of groups that mentioned in Example 2.3 and Example 2.4.

Example 2.8. In this example, we give a presentation of the fundamental group of an edge of groups by using definition (b) above. As pictured in Example 2.3, the maximal subtree T of an edge e is the edge itself. So the presentation of the fundamental group is given by

$$\pi_1(\Gamma, T) = \langle G_{o(e)}, G_{t(e)} \mid \alpha_e(g) = \alpha_{\overline{e}}(g) \text{ for any } g \in G_e \rangle$$

In fact, this fundamental group is the free amalgamated product of the two vertex groups over the edge group, i.e. $\pi_1(\Gamma, T) = G_{o(e)} *_{G_e} G_{t(e)}$

Example 2.9. Using definition (b) above, we now give a presentation of the fundamental group of a loop of groups. As pictured in Example 2.4, the maximal subtree T of a loop f is the vertex v. So the presentation of the fundamental group is given by

$$\pi_1(\Gamma, T) = \langle G_v, f \mid \overline{f} = f^{-1}, f\alpha_e(g)f^{-1} = \alpha_{\overline{e}}(g) \text{ for any } g \in G_f \rangle$$

In fact, this fundamental group is a HNN extension $G_v *_{\alpha_e(G_f) \sim \alpha_{\overline{e}}(G_f)}$ where f is a stable letter.

The following proposition shows that the two definitions above do not depend on which base vertex v_0 or maximal subtree T we chose. Moreover, there is a natural projection map $F(\Gamma, \mathcal{G}) \to \pi_1(\Gamma, T)$ given by mapping edges in the maximal tree T to the identity.

Proposition 2.10. [Ser80, Proposition 5.1.20] Let (Γ, \mathcal{G}) be a graph of groups, let $v_0 \in V(\Gamma)$ and let T be a maximal tree of Γ . The natural projection map $F(\Gamma, \mathcal{G}) \to \pi_1(\Gamma, T)$ induces an isomorphism of $\pi_1(\Gamma, v_0)$ onto $\pi_1(\Gamma, T)$.

Therefore, for the rest of this paper, we will sometimes use $\pi_1(\Gamma, \mathcal{G})$ to denote the fundamental group of (Γ, \mathcal{G}) .

- 2.3. Bass-Serre tree. Let (Γ, \mathcal{G}) be a graph of groups, and let T be a maximal tree of Γ . Set $G = \pi_1(\Gamma, T)$. We can construct a graph X that satisfies the following properties:
 - (a) there is an action of G on X by left multiplication;
 - (b) the quotient X/G induced by the action of G on X is isomorphic to Γ ;
 - (c) if \tilde{x} is a vertex or an edge of X, and $\tilde{y} = g\tilde{x}$ for some $g \in G$. Then the stabilizer of \tilde{y} in G is $gG_{q(\tilde{x})}g^{-1}$, where $q: X \to \Gamma$ is the quotient map.
 - (d) there exist section maps $V(\Gamma) \to V(X)$ and $E(\Gamma) \to E(X)$ of the quotient map q.

In fact, these conditions force the graph X to be as follows: the vertex set V(X) is the disjoint union of left cosets of vertex groups, i.e.

$$V(X) = \bigsqcup_{v \in V(\Gamma)} G/G_v$$

Likewise, the edge set E(X) is the disjoint union of left cosets of edge groups, i.e.

$$E(X) = \bigsqcup_{e \in E(\Gamma)} G/G_e$$

The graph X is defined by attaching the edges to the vertices in the following way:

$$o(gG_e) = gG_{o(e)}$$
 and $t(gG_e) = geG_{t(e)}$

We remark that e is the identity if e is in the maximal tree T. From the construction above, we can see that the quotient X/G has a graph of groups structure that is isomorphic to (Γ, \mathcal{G}) . We conclude this subsection with the following two theorems, known as the fundamental theorems of Bass-Serre theory.

Theorem 2.11. [Ser80, Theorem 5.3.12] Let (Γ, \mathcal{G}) be a graph of groups, and let T be a maximal tree of Γ . Set $G = \pi_1(\Gamma, T)$. Then the graph X constructed above is a tree, and we call it the Bass-Serre tree associated to (Γ, \mathcal{G}) . In particular, X is the universal cover of the quotient graph Γ .

Theorem 2.12. [Ser80, Theorem 5.3.13] Let G be a group which acts on a tree X without edge inversions (i.e. $ge \neq \bar{e}$ for any $g \in G$ and $e \in E(X)$). Then G can be identified with the fundamental group of a certain graph of groups (Γ, \mathcal{G}) , where Γ is the quotient graph X/G. Moreover, there is an G-equivariant isomorphism of the Bass-Serre tree associated to (Γ, \mathcal{G}) to X.

2.4. Action on the Bass-Serre tree.

Definition 2.13. [For02, Definition 2.4] Let G be a group acting on a tree X. An element $\gamma \in G$ is *elliptic* if it has a fixed point, and *hyperbolic* otherwise. We define the *length function* of X by

$$\ell_X(\gamma) = \min_{\substack{x \in V(X) \\ 7}} d(x, \gamma x)$$

Thus $\ell_X(\gamma) = 0$ if and only if γ is elliptic. If γ is hyperbolic, then the set

$$\{x \in X \mid d(x, \gamma x) = \ell_X(\gamma)\}\$$

is a γ -invariant linear subtree, called the *axis* of γ . We denote it by $\operatorname{Axis}(\gamma)$. The action of γ on $\operatorname{Axis}(\gamma)$ is by a translation of amplitude $\ell_X(\gamma)$.

Definition 2.14. [MO15, Definition 2.10] Let G be a group acting on a tree X. We say G is acting *elliptically*, or that G is *elliptic*, if the action of G on X has a fixed point.

3. Acylindrical tree actions

In this section, we recall definitions and results about acylindrical actions that we will require throughout the paper. For more details we refer to [Osi18].

Definition 3.1. Let G be a group acting on a hyperbolic space S by isometry. The action is called *acylindrical* if for every $\varepsilon > 0$ there exist R, N > 0 such that for every two points x, y in S with $d(x, y) \ge R$, the set

$$\{g \in G \mid d(x, gx) \le \varepsilon, \ d(y, gy) \le \varepsilon\}$$

contains at most N elements.

When we consider acylindrical action on trees in particular, we have the following definition due to Weidmann [Wei12].

Definition 3.2. [Wei12, Introduction] Let G be a group acting on a simplicial tree T and let $k \geq 0$ and C > 0 be integers. We say that the action is (k, C)-acylindrical if the pointwise stabilizer of any edge path in T of length at least k contains at most C elements.

The following theorem is due to Minasyan and Osin [MO15], which we include without proof. This theorem shows that the above two definitions are, in fact, equivalent.

Theorem 3.3. [MO15, Lemma 4.2] Let G be a group acting on a simplicial tree T by isometries. This action is acylindrical if and only if there exist constants $k \geq 0$ and $C \geq 1$ such that the action of G on T is (k, C)-acylindrical.

Definition 3.4. [Osi18, Definition 2.4] Let G be a group acting on isometrically on a hyperbolic space S. An element $g \in G$ is called *elliptic* if all orbits of g are bounded; and hyperbolic (or loxodromic) if the map $\mathbb{Z} \to S$ defined by $n \mapsto g^n s$ is a quasi-isometry for some (equivalently, any) $s \in S$. Two hyperbolic elements $g, h \in G$ are called independent if the sets $\{g^{\pm \infty}\}$ and $\{h^{\pm \infty}\}$ are disjoint.

Remark 3.5. Let G be a group acting on a tree T. It is known [Ser80, Proposition I.2.10 and Corollary] that the action of G acting on T has bounded orbits if and only if there exists some point $x \in T$ fixed by the action $G \curvearrowright T$. Therefore we can remove any ambiguity as to the definition of an elliptic action in the context of simplicial trees.

The following theorem classifies acylindrical actions on hyperbolic spaces.

Theorem 3.6. [Osi16, Theorem 1.1] Let G be a group acting acylindrically on a hyperbolic space. Then G satisfies exactly one of the following three conditions.

- (1) G has bounded orbits.
- (2) G is virtually cyclic and contains a hyperbolic element.
- (3) G contains infinitely many independent hyperbolic elements.

As an application of the above theorem, every element of a group acting acylindrically on a hyperbolic space is either elliptic or hyperbolic. In the theorem above, (1) and (2) are called *elementary* actions, whereas (3) is called a *non-elementary* action.

Definition 3.7. We say a group G is acylindrically hyperbolic if it admits a non-elementary acylindrical action on a hyperbolic metric space S by isometries.

4. Preserving acylindricity under quotient

In this section, we provide sufficient conditions on normal subgroups of G such that acylindricity of the action of G on a tree T is preserved under quotients by such normal subgroups. We begin by recalling the definition of an equivariant family of subgroups of G.

Definition 4.1. Let G be a group acting on a tree T. For each vertex $v \in V(T)$, let R_v be a subgroup of the stabilizer of v in G. We say the family of subgroups $\{R_v\}$ is an equivariant family of subgroups of G if it satisfies the following condition:

- If g lies in G and v is a vertex of T then $gR_vg^{-1}=R_{gv}$.

Remark 4.2. The equivariance condition implies that for each vertex v the subgroup R_v is normal in the stabilizer of v, and that the subgroup $\langle R_v \rangle$ of G generated by the family is normal in G. Further, if we choose orbit representatives v_i for the action of G on the vertices of T, then $\langle R_v \rangle$ is the normal closure of the set $\{R_{v_i}\}$. Letting (Γ, \mathcal{G}) be the graph of groups associated to the action of G on T, we can always find such a set of orbit representatives by choosing a lift $\tilde{\Gamma}$ of Γ in T, namely $\langle R_v \rangle = \langle \langle R_{v_i} | v_i \in V(\tilde{\Gamma}) \rangle \rangle$. Moreover, moving forward we will choose a particular lift $\tilde{\Gamma}$ of Γ such that the vertex stabilizers in $\tilde{\Gamma}$ remain the same as the vertex groups in Γ , i.e., $G_{\hat{v}} = G_v$ for lifts of vertices \hat{v} in Γ to v in $\tilde{\Gamma}$. With this choice of lift, we identify R_v 's as a subgroup of the vertex groups $G_{\hat{v}}$ in Γ for each $v \in V(\tilde{\Gamma})$. This implies that $\langle R_v \rangle = \langle \langle R_{\hat{v}} | \hat{v} \in V(\Gamma) \rangle \rangle$.

Let us consider an action on a tree. Our next result classifies normal subgroup quotients which preserve a tree structure.

Proposition 4.3. Let G be a group acting on a simplicial tree T, and let N be a normal subgroup of G. The quotient graph T/N is a tree if and only if N is generated by an equivariant family of subgroups $\{R_v\}$ i.e., $N = \langle R_v \rangle$.

Proof. Let Γ_N denote the quotient graph T/N. Consider the action of $N \curvearrowright T$ that we obtained by restricting the action of $G \curvearrowright T$ to N. Assume that Γ_N is a tree. We note that Γ_N can be lifted to a subtree $\tilde{\Gamma}_N$ in T such that $N \cdot \tilde{\Gamma}_N = T$, i.e., $\tilde{\Gamma}_N$ is a fundamental domain of T under the action $N \curvearrowright T$. By Bass-Serre theory, N admits a graph of groups structure $(\Gamma_N, \mathcal{G}_N)$ such that for each vertex \bar{v} in Γ_N , the vertex group is $G_{\bar{v}} = \operatorname{Stab}_N(v)$ where v is the lift of \bar{v} in $\tilde{\Gamma}_N$, and for each edge \bar{e} in Γ_N , the edge group is $G_{\bar{v}} = \operatorname{Stab}_N(v)$ where e is the lift of \bar{e} in $\tilde{\Gamma}_N$. Further, we have $N = \pi_1(\Gamma_N, \mathcal{G}_N)$. Since Γ_N is a tree, then $N = \langle \sqcup_{\bar{v} \in V(\Gamma_N)} G_{\bar{v}} \rangle = \langle \sqcup_{v \in V(\tilde{\Gamma}_N)} \operatorname{Stab}_N(v) \rangle$. Now let $R_v = \operatorname{Stab}_N(v) = \operatorname{Stab}_G(v) \cap N \leq \operatorname{Stab}_G(v)$ for each vertex v in $\tilde{\Gamma}_N$. Since $N \cdot \tilde{\Gamma}_N = T$, we can repeat the choice equivariantly throughout vertices in T, i.e., for any $w \in V(T)$, we have $w = h \cdot v$ for some $h \in N$ and $v \in V(\tilde{\Gamma}_N)$, we then choose $R_w = hR_vh^{-1}$. It follows that $\{R_v\}$ is an equivariant family of subgroups of G. Under this choice of $\{R_v\}$, it is clear that $N = \langle R_v \rangle$. Conversely, suppose N is a normal subgroup generated by $\{R_v\}$, an equivariant family of subgroups of G. Since each R_v is a subgroup of some vertex stabilizer, it does not contain any stable letters. In addition, quotient T by each

 R_v folds edges emanating from v. Thus, it follows from Bass-Serre theory and the definition of equivariant family that in the quotient graph, valence of each vertex might be reduced but no loop is created. Therefore the quotient graph T/N remains a tree.

Theorem 4.4. Let G be a group with an acylindrical action on a simplicial tree T, and let $\{R_v\}$ be an equivariant family of subgroups of G. Then the quotient action $G/\langle R_v\rangle$ on $T/\langle R_v\rangle$ is an acylindrical action.

Proof. For simplicity, we denote $N = \langle R_v \rangle$, denote the quotient group G/N by \overline{G} , and denote the quotient graph T/N by Γ_N . Since Γ_N is a tree, then Γ_N can be lifted to a subtree $\tilde{\Gamma}_N$ (an isometrically embedded copy of Γ_N) in T such that $N \cdot \tilde{\Gamma}_N = T$, i.e., $\tilde{\Gamma}_N$ is the fundamental domain of T under the action $N \curvearrowright T$. Similar to the proof of Proposition 4.3, we have N admits a graph of groups structure $(\Gamma_N, \mathcal{G}_N)$ for each vertex \bar{v} in Γ_N , the vertex group is $G_{\bar{v}} = \operatorname{Stab}_{N}(v)$ where v is the lift of \bar{v} in Γ_{N} , and for each edge \bar{e} in Γ_{N} , the edge group is $G_{\overline{e}} = \operatorname{Stab}_{N}(e)$ where e is the lift of \overline{e} in Γ_{N} . We now assume that $G \curvearrowright T$ is acylindrical. By Theorem 3.3, there exists constants $k \geq 0$ and $C \geq 1$ such that the action $G \curvearrowright T$ is (k,C)-acylindrical. Let $[\bar{u},\bar{v}]$ be a geodesic path in Γ_N with $d_{\Gamma_N}(\bar{u},\bar{v}) \geq k$. Further, let u be the lift of \bar{u} in $\tilde{\Gamma}_N$ and let v be the lift of \bar{v} in $\tilde{\Gamma}_N$. By the construction of $\tilde{\Gamma}_N$, we have $d_T(u,v) \geq k$. Choose coset representatives $\Sigma_N \subset G$, we can represent elements in \overline{G} by elements in Σ_N . Thus for each $g \in \mathrm{PStab}_{\overline{G}}([\bar{u},\bar{v}]) = \mathrm{PStab}_{\overline{G}}(\{\bar{u},\bar{v}\})$, we have $g \in G_{\bar{u}}$ and $g \in G_{\bar{v}}$. By the construction of $(\Gamma_N, \mathcal{G}_N)$, we have $g \in \operatorname{Stab}_N(u)$ and $g \in \operatorname{Stab}_N(v)$. This shows that $g \in \mathrm{PStab}_G([u,v])$. However, we have $|\mathrm{PStab}_G([u,v])| \leq C$. This shows that $|\operatorname{PStab}_{\overline{G}}([\bar{u},\bar{v}])| \leq C$. Thus the quotient action $G \curvearrowright \Gamma_N$ is also (k,C)-acylindrical. By Theorem 3.3 again, we have the quotient action $\overline{G} \curvearrowright \Gamma_N$ is an acylindrical action.

<u>Notations.</u> Let G be a group acting on a tree T, and let (Γ, \mathcal{G}) be a finite graph of groups associated to the action $G \curvearrowright T$. Further, let N be the normal subgroup of G generated by an equivariant family of subgroups $\{R_v\}$. We denote the quotient group G/N by \overline{G} , the quotient tree T/N by \overline{T} . Choose a coset representatives $\Sigma_N \subset G$, we then identify elements in \overline{G} by elements in Σ_N . We use v to denote vertex in T, \hat{v} to denote vertex in Γ and \bar{v} to denote vertex in \overline{T} . For any two vertices \hat{u} and \hat{v} in Γ , we denote $[\hat{v}, \hat{v}]$ the geometric geodesic path (disregrading the orientation) joining \hat{u} and \hat{v} .

We need a few lemmas before proving the main theorem, Theorem 4.7, of this section.

Lemma 4.5. Let v be a vertex in T. The vertex \bar{v} in \overline{T} has valence at least two if and only if at least one the following holds:

- (1) there is an edge \hat{e} in Γ emanating from \hat{v} such that $G_{\hat{v}} \neq \langle G_{\hat{e}}, R_{\hat{v}} \rangle$;
- (2) \hat{v} has valence at least two.

Proof. Assume that condition (1) holds. As in Remark 4.2, we can assume $\operatorname{Stab}(v) = G_{\hat{v}}$. Let e be the lift (in $\tilde{\Gamma}$) of \hat{e} that emanating from v. Since $G_{\hat{v}} \neq \langle G_{\hat{e}}, R_{\hat{v}} \rangle$, then there exists a non-trivial element $g \in G_{\hat{v}} \setminus \langle G_{\hat{e}}, R_{\hat{v}} \rangle$. Further, since $g \notin G_{\hat{e}}$, we have ge and e are two different edges emanating from v. Since $g \notin R_{\hat{v}}$. then $gG_{\hat{e}} \neq hG_{\hat{e}}$ for any $h \in R_{\hat{v}}$. This shows that ge and e lie in two different $\langle R_v \rangle$ -orbits. Thus \bar{v} has valence at least two. Now assume that condition (2) holds, then there are at least two different G-orbits of edges emanating from v. It follows that there are at least two different G-orbits of edges emanating from v. Therefore \bar{v} has valence at least two.

For the only if direction, we assume that \bar{v} has valence at least two. Then there are two different edges \bar{e} and \bar{f} emanating from \bar{v} in \bar{T} . Let e (resp. f) be an edge emanating from v in T such that $q(e) = \bar{e}$ (resp. $q(f) = \bar{f}$), where $q: T \to \bar{T}$ is the natural quotient map. There are two cases, depending on whether e and f lie in the same G-orbit or not. If e and f do not lie in the same G-orbit, then \hat{v} has valence at least two. If e and f lie in the same G-orbit \hat{e} , without loss of generality, we may assume that $\mathrm{Stab}(e) = G_{\hat{e}}$. Then there exists an element $g \in G_{\hat{v}} \setminus G_{\hat{e}}$ such that f = ge, i.e., f is represented by $gG_{\hat{e}}$. Further, since $\bar{e} \neq \bar{f}$, then $gG_{\hat{e}} \neq hG_{\hat{e}}$ for any $h \in R_{\hat{v}}$. This implies that $g \in G_{\hat{v}} \setminus \langle G_{\hat{e}}, R_{\hat{v}} \rangle$. Thus $G_{\hat{v}} \neq \langle G_{\hat{e}}, R_{\hat{v}} \rangle$. This completes the proof.

Proposition 4.6. \overline{T} is an infinite tree if at least one of the following holds:

- (1) there are two vertices \hat{u} and \hat{v} in Γ such that $\langle G_{\hat{e}}, R_{\hat{u}} \rangle \neq G_{\hat{u}}$ and $\langle G_{\hat{f}}, R_{\hat{v}} \rangle \neq G_{\hat{v}}$, where \hat{e} and \hat{f} are the edges on the geodesic path $[\hat{u}, \hat{v}]$ that adjacent to \hat{u} and \hat{v} respectively.
- (2) Γ contains a circuit.

Proof. By Proposition 4.3, \overline{T} is always a tree. Then it suffices to show that \overline{T} is infinite if at least one of the above conditions holds. Assume that condition (1) holds. Let $[\hat{u}, \hat{v}]$ be the geodesic segment in Γ joining \hat{u} and \hat{v} , we can lift $[\hat{u}, \hat{v}]$ to a geodesic segment [u, v] inside $\tilde{\Gamma}$, the partcular lift of Γ as in Remark 4.2. Note that [u, v] generates a subtree T' of T, i.e., $G \cdot [u, v] = T' \subseteq T$. Since $G_{\hat{e}} \neq G_{\hat{u}}$ and $G_{\hat{f}} \neq G_{\hat{v}}$, then T' must be an infinite tree. By Lemma 4.5, we have both \bar{u} and \bar{v} have valence at least two and the image of T' under the quotient map $q: T \to \overline{T}$ contains a bi-infinite line. More precisely, there exist $g \in G_{\hat{u}} \setminus \langle G_{\hat{e}}, R_{\hat{u}} \rangle$ and $h \in G_{\hat{v}} \setminus \langle G_{\hat{f}}, R_{\hat{v}} \rangle$ such that $g, h \in \overline{G}$, gh is hyperbolic in both G and \overline{G} , and both T and \overline{T} contain Axis(gh). Therefore \overline{T} is infinite. Now assume that condition (2) holds. This implies that G contains a stable letter t. We note that the equivariant family of subgroups $\{R_v\}$ does not contain any stable letters of G, then each axis of the stable letters survives under the quotient map $q: T \to \overline{T}$. Thus \overline{T} contains Axis(t). This implies that \overline{T} is infinite.

We now provide a proof of our first main result, which is restated as follows.

Theorem 4.7. Let G be a group acting non-elementary acylindrically on a simplicial tree T, and let N be a normal subgroup of G. If N is generated by an equivariant family of subgroups $\{R_v\}$, then the quotient space T/N is a tree. Moreover, the quotient action $G/N \curvearrowright T/N$ is a non-elementary acylindrical tree action if $\{R_v\}$ satisfies at least one of the following conditions:

- (1) the first Betti number $b_1(\Gamma) \geq 2$ where $\Gamma = T/G$.
- (2) there are two vertices \hat{u} and \hat{v} in Γ such that $\langle G_{\hat{e}}, R_{\hat{u}} \rangle \neq G_{\hat{u}}$ and $\langle G_{\hat{f}}, R_{\hat{v}} \rangle \neq G_{\hat{v}}$, where \hat{e} and \hat{f} are the edges on the geodesic path $[\hat{u}, \hat{v}]$ that are adjacent to \hat{u} and \hat{v} respectively. Further, there exist two elements $g, h \in G_{\hat{v}} \setminus \langle G_{\hat{f}}, R_{\hat{v}} \rangle$ such that $gh^{-1} \notin \langle G_{\hat{f}}, R_{\hat{v}} \rangle$.
- (3) Γ contains an immersed circuit. Further, there exists a vertex \hat{v} and an edge \hat{e} adjacent to \hat{v} on the circuit such that $\langle G_{\hat{e}}, R_{\hat{v}} \rangle \neq G_{\hat{v}}$.

Proof. We have the quotient action to be acylindrical due to Theorem 4.4. Assume that condition (1) holds. Then G contains at least two different stable letters t_1 and t_2 . Since the equivariant family of subgroups $\{R_v\}$ does not contain any stable letters of G, then we

have $t_1, t_2 \in \overline{G}$ and \overline{T} contains $\operatorname{Axis}(t_1)$ and $\operatorname{Axis}(t_2)$. This shows that t_1 and t_2 are two independent hyperbolic elements of the action $\overline{G} \curvearrowright \overline{T}$.

Assume that condition (2) holds. First, we can lift the geodesic segment $[\hat{u}, \hat{v}]$ to the geodesic segment [u, v] in $\tilde{\Gamma}$ as in Remark 4.2. Let e, f be the lifts of \hat{e} and \hat{f} on [u, v] that are adjacent to u and v respectively. By assumption, we have f, gf, hf are three different edges emanating from v. Moreover, they lie in three different $\langle R_v \rangle$ -orbits, i.e., \bar{v} has valence at least three and q(f), q(gf), q(hf) represents three different edges in \bar{T} emanating from \bar{v} . Let $g' \in G_{\hat{u}} \setminus \langle G_{\hat{e}}, R_{\hat{u}} \rangle$, similar to the proof of Proposition 4.6, we have g'g and g'h are two independent hyperbolic elements of both G and \bar{G} , and both T and \bar{T} contain Axis(g'g) and Axis(g'h).

Assume that condition (3) holds. Since Γ contains an immersed circuit, we have G contains a stable letter t. Denote the circuit by \hat{c} and choose a base vertex \hat{u} on \hat{c} . Without loss of generality, we may assume that the orientation of \hat{e} is the same as \hat{c} , for otherwise we can choose \hat{c}^{-1} instead of \hat{c} . Then we can lift \hat{c} to a geodesic segment [u, tu] in T where $q(u) = \hat{u}$, and this geodesic segment generates the axis of t in T. More precisely, we have $t^m \cdot [u, tu] = \operatorname{Axis}(t)$ for all $m \in \mathbb{Z}$. Similar to condition (1), we have \overline{T} contains $\operatorname{Axis}(t)$. Let v, e be the lift of \hat{v}, \hat{e} in [u, tu] respectively. Since $\langle G_{\hat{e}}, R_{\hat{v}} \rangle \neq G_{\hat{v}}$, similar to the proof of Lemma 4.5, there exists $g \in G_{\hat{v}} \setminus \langle G_{\hat{e}}, R_{\hat{v}} \rangle$ such that ge and e are two different edges in T emanating from v. We note that gt is a hyperbolic element of the action $G \curvearrowright T$ with translation length $\ell(gt) = \ell(t)$. In fact, the geodesic segment [gu, gtu] generates the axis of gt in T. In addition, since ge and e are two different edges in T, and ge lies on [gu, gtu]. Then gt and t are two independent hyperbolic elements of the action $G \curvearrowright T$. Recall that $g \in G_{\hat{v}} \setminus \langle G_{\hat{e}}, R_{\hat{v}} \rangle$, then $gt, t \in \overline{G}$ and \overline{T} contains both $\operatorname{Axis}(gt)$ and $\operatorname{Axis}(t)$. This shows that gt and t are two independent hyperbolic elements of the action $\overline{G} \curvearrowright \overline{T}$.

We can obtain the following two particular corollaries as immediate applications of the above theorem.

Corollary 4.8. Let $G = A *_C B$ have a non-elementary acylindrical action on a tree T, and let N be a normal subgroup of G generated by an equivariant family of subgroups $\{R_v\}$. The quotient action $G/N \curvearrowright T/N$ is a non-elementary acylindrical tree action if $\{R_v\}$ satisfies the following:

- (1) $\langle C, R_A \rangle \neq A$ and $\langle C, R_B \rangle \neq B$; and
- (2) there exists two different elements g and h in $B \setminus \langle C, R_B \rangle$ such that $gh^{-1} \notin \langle C, R_B \rangle$.

Proof. The proof follows directly from condition (2) of Theorem 4.4.

Corollary 4.9. Let $G = A*_{H \sim_{\phi} K}$ have a non-elementary acylindrical action on a tree T. and let N be a normal subgroup of G generated by an equivariant family of subgroups $\{R_v\}$. The quotient action $G/N \curvearrowright T/N$ is a non-elementary acylindrical tree action if $\{R_v\}$ satisfies either $\langle H, R_A \rangle \neq A$ or $\langle K, R_A \rangle \neq A$.

Proof. The proof follows directly from condition (3) of Theorem 4.4. \Box

We will discuss a particular application of Corollary 4.8 to the outer automorphism group of Baumslag-Solitar group in subsection 6.3.

5. Largest acylindrical action of graph of groups

In this section, our goal is to prove Theorem D. This theorem provides sufficient conditions for the fundamental group of graph of groups admitting a largest acylindrical action. This is a particular case of Proposition 4.3 in [ABO19] which gives sufficient condition for existence of a largest action of a group acting cocompactly on a connected graph. Restricting to action on trees we give a simpler proof using Bass-Serre theory. We begin by briefly recall some definitions from [ABO19] defined by Abbott, Balasubramanya and Osin.

Definition 5.1. Let X and Y be two generating sets of group G. We say that X is dominated by Y, denoted by $X \subseteq Y$, if

$$\sup_{y \in Y} |y|_X < \infty$$

where $|\cdot|_X = d_X(1,\cdot)$ denotes the word length with respect to X. It is clear that \leq defines a preorder on the generating sets of the group G and induces an equivalence relation:

$$X \sim Y \iff X \prec Y \text{ and } Y \prec X$$

Let [X] denote the equivalence class of the generating set X and $\mathcal{G}(G)$ be the set of all equivalence classes of generating sets of G. The preorder \leq induces a partial order \leq on $\mathcal{G}(G)$ by the rule

$$[X] \preccurlyeq [Y] \Longleftrightarrow X \preceq Y.$$

Given a group G with a generating set X, we denote by $\Gamma(G,X)$ the Cayley graph of G with respect to the generating set X.

Definition 5.2. Let $\mathcal{AH}(G)$ be the set of equivalence classes $[X] \in \mathcal{G}(G)$ such that $\Gamma(G, X)$ is hyperbolic and the action $G \curvearrowright \Gamma(G, X)$ is acylindrical. We say an equivalence class of generating sets is *largest* if it is the largest element in $\mathcal{AH}(G)$.

Definition 5.3. [ABO19, Section 3] We say two actions of a group G on metric spaces S and R are equivalent, denoted by $G \curvearrowright S \sim G \curvearrowright R$, if there exists a coarsely G-equivariant quasi-isometry $S \to R$.

The action of a group G on a metric space S is said to be *cobounded* if there exists a bounded subset $B \subseteq S$ such that $S = \bigcup_{g \in G} gB$. Given a cobounded acylindrical action of G on a hyperbolic space S, the following lemma provides a (possibly infinite) generating set X of G such that $\Gamma(G, X)$ is equivariantly quasi-isometric to S.

Lemma 5.4. [ABO19, Lemma 3.11] Let G be a group acting coboundedly on a geodesic metric space S. Let $B \subseteq S$ be a bounded subset such that $\bigcup_{g \in G} gB = S$. Let D = diam(B) and let b be any point of B. Then the group G is generated by the set

$$X = \{ g \in G \mid d_S(b, gb) \le 2D + 1 \}$$

and the natural action $G \curvearrowright \Gamma(G,X)$ is equivalent to $G \curvearrowright S$.

As an application of the above lemma, we say that a particular cobounded acylindrical action $G \curvearrowright S$ on a hyperbolic space is largest, when the equivalence class of the generating set associated to this action through the above correspondence, [X], that is the largest element in $\mathcal{AH}(G)$.

We now focus on cobounded acylindrical tree actions. Let (Γ, \mathcal{G}) be a finite graph of groups, and let T be the Bass-Serre tree of (Γ, \mathcal{G}) . Denote the fundamental group of graph of groups $\pi_1(\Gamma, \mathcal{G})$ by G. We first note that Γ can be lifted to a subtree $\tilde{\Gamma}$ in T such that $G \cdot \tilde{\Gamma} = T$, i.e., $\tilde{\Gamma}$

is the fundamental domain of T under the action $G \curvearrowright T$. Since Γ is a finite graph, we have $\tilde{\Gamma}$ is of finite diameter and $G \curvearrowright T$ is a cobounded action. Let $D = \operatorname{diam}(\tilde{\Gamma})$ and let \tilde{v}_0 be a base vertex of $\tilde{\Gamma}$. Then Lemma 5.4 provides a generating set $X = \{g \in G | d_T(\tilde{v}_0, g\tilde{v}_0) \leq 2D + 1\}$ of G. Further, we have $G \curvearrowright T \sim G \curvearrowright \Gamma(G, X)$. Note that G, as the fundamental group of (Γ, \mathcal{G}) , has the usual generating set $Y = (\sqcup_{v \in V(\Gamma)} G_v) \sqcup \{e \in E(\Gamma) | e \text{ is a stable letter}\}$. The following lemma shows that these two generating sets X and Y are equivalent.

Lemma 5.5. Let X and Y be the generating sets of G defined as above, then $X \sim Y$.

Proof. From Bass-Serre theory, we note that $Y \subset X$. This implies that $\sup_{Y \in Y} |y|_X = 1$. Therefore $X \preceq Y$. Conversely, without loss of generality, we may assume that \tilde{v}_0 is labeled by the group G_{v_0} . As discussed in Remark 1.18 of [Bas93], each vertex of T at distance n from G_{v_0} can be represented by a unique reduced path in Γ of the form $g_1e_1g_2e_2...g_ne_n$ where $o(e_1) = v_0$ and g_i is contained in the transversal set \sum_{e_i} for $G_{o(e_i)}/\alpha_{\overline{e_i}}(G_{e_i})$. Thus, for each $g \in X$ we have $g\tilde{v}_0 = gG_{v_0} = g_1e_1g_2e_2...g_ne_nG_{v_0}$ where $n = d_T(\tilde{v}_0, g\tilde{v}_0) \leq 2D + 1$. This implies that $g = g_1e_1g_2e_2...g_ne_nh$ for some $h \in G_{v_0}$, which shows that g has word length at most 4D + 3 with respect to the generating set Y. Therefore we have $Y \preceq X$.

Before giving the proof of our main theorem, we record the following proposition which gives sufficient conditions for an action to be the largest. Our goal is to apply this result to acylindrical tree actions.

Proposition 5.6. [ABD21, Proposition 5.2] Let G be a group, $\{H_1, \dots, H_n\}$ a finite collection of subgroups of G, and F be a finite subset of G such that $F \cup (\bigcup_{i=1}^n H_i)$ generates G. Assume that:

- (1) $\Gamma(G, F \cup (\bigcup_{i=1}^{n} H_i))$ is hyperbolic and the action of G on it is acylindrical.
- (2) Each H_i is elliptic in every acylindrical action of G on a hyperbolic space.

Then $[F \cup (\bigcup_{i=1}^{n} H_i)]$ is the largest element in $\mathcal{AH}(G)$.

We are now ready to prove our main theorem in this section.

Theorem 5.7. Let (Γ, \mathcal{G}) be a finite graph of groups, and let T be the Bass-Serre tree of (Γ, \mathcal{G}) . Denote the fundamental group $\pi_1(\Gamma, \mathcal{G})$ by G. If the action $G \curvearrowright T$ is acylindrical, and if each vertex group acts elliptically whenever G acts acylindrically on any hyperbolic space S, then G admits the largest acylindrical action on T.

Proof. Let F be the set of stable letters of Γ , and let $\{H_1, ..., H_n\}$ be the collection of vertex groups in (Γ, \mathcal{G}) . Since Γ is a finite graph, we have F is finite and $\{H_1, ..., H_n\}$ is a finite collection. Further, by Bass-Serre theory, we have $F \cup (\bigcup_{i=1}^n H_i)$ generates G. By Lemma 5.5, we have the action $G \curvearrowright T$ is equivalent to $G \curvearrowright \Gamma(G, F \cup (\bigcup_{i=1}^n H_i))$. By Lemma 3.9 of [ABO19], this is equivalent to $T \sim \Gamma(G, F \cup (\bigcup_{i=1}^n H_i))$. Thus there exists a coarsely G-equivariant quasi-isometry $T \to \Gamma(G, F \cup (\bigcup_{i=1}^n H_i))$. Note that hyperbolicity is preserved under quasi-isometries, and it is a fact that acylindricity of an action is preserved under equivariant quasi-isometries. Therefore we have $\Gamma(G, F \cup (\bigcup_{i=1}^n H_i))$ is hyperbolic and the action $G \curvearrowright \Gamma(G, F \cup (\bigcup_{i=1}^n H_i))$ is acylindrical. Thus the theorem follows directly from Proposition 5.6 above.

Corollary 5.8. Let (Γ, \mathcal{G}) be a finite graph of groups, and let T be the Bass-Serre tree of (Γ, \mathcal{G}) . Denote the fundamental group $\pi_1(\Gamma, \mathcal{G})$ by G. If the action $G \curvearrowright T$ is acylindrical,

and if each vertex group consists of finite order elements only, then G admits the largest acylindrical action on T.

Proof. Since each vertex group consists of finite order elements only, then it can not contain any hyperbolic elements whenever G acts acylindrically on any hyperbolic space S. By Theorem 3.6, this is equivalent to the statement that each vertex group acts elliptically whenever G acts acylindrically on any hyperbolic space S. Thus the corollary follows directly from Theorem 5.7 above.

We note that the outer automorphism group of some Baumslag-Solitar groups is a family of examples that satisfies Corollary 5.8 and thus Theorem 5.7. We will descirbe these groups more explicitly in subsection 6.4.

6. Outer automorphism group of BS(p,q)

Recall the Baumslag-Solitar group $\mathrm{BS}(p,q)$ is defined to be the group with the following presentation

$$BS(p,q) = \langle x, t \mid tx^p t^{-1} = x^q \rangle$$

where $p, q \in \mathbb{Z} \setminus \{0\}$. By interchanging $t \leftrightarrow t^{-1}$, one may always assume $1 \leq p \leq |q|$. Throughout this section, we only consider the non-solvable Baumslag-Solitar groups i.e. we assume that $1 < |p| \leq |q|$. We note that presentations for these outer automorphism groups are known (see Remark 6.1 and subsection 6.2 for details).

Remark 6.1. We have the following presentations for Out(BS(p, q)) when p does not divide q properly (details can be found in Section 3 of [Cla09] or [Col78] and [GHMR00]).

- Out(BS(p,q)) = $\mathbb{Z}_{2|p-q|} \rtimes \mathbb{Z}_2$ if p does not divide q.
- Out $(BS(p,q)) = \mathbb{Z} \rtimes (\mathbb{Z}_2 \times \mathbb{Z}_2)$ if p = q.
- Out(BS(p,q)) = $\mathbb{Z}_{2p} \rtimes \mathbb{Z}_2$ if p = -q.

We note that $\operatorname{Out}(\operatorname{BS}(p,q))$ is either finite or virtually $\mathbb Z$ in any of the above cases, so it suffices to investigate the case when p properly divides q. In this case, Clay showed that $\operatorname{Out}(\operatorname{BS}(p,q))$ admits a ray of groups structure (Γ_1,\mathcal{G}_1) pictured as in Figure 3. Further, he showed that (Γ_1,\mathcal{G}_1) can be collapsed to an edge of groups (Γ_2,\mathcal{G}_2) pictured as in Figure 4, and he used (Γ_2,\mathcal{G}_2) to compute the presentation of $\operatorname{Out}(\operatorname{BS}(p,q))$ via Bass-Serre theory. We will describe this presentation in subsection 6.2. In this section, our first goal is to compare the two tree actions that arise from these two graph of groups structures in the sense of Section 5. Next, we apply Corollary 4.8 to $\operatorname{Out}(\operatorname{BS}(p,q))$ and we show that the quotient action of a non-elementary acylindrical tree action of $\operatorname{Out}(\operatorname{BS}(p,q))$ by some equivariant family of subgroups is a non-elementary acylindrical tree action.

Assumption. For the remainder of this section, we will assume that q = pn where p, |n| > 1.

6.1. **Deformation space of trees.** We begin by recalling the definition of deformation space of trees, which is the tool Clay used to compute the presentation of Out(BS(p,q)).

Definition 6.2. [Cla09, Definition 1.1] Let (Γ, \mathcal{G}) be a graph of groups. An edge $e \in E(\Gamma)$ is *collapsible* if $o(e) \neq t(e)$ and $G_{t(e)} = \alpha_e(G_e)$. If one collapses $\{e, \overline{e}\}$ to the vertex o(e), the resulting graph of groups is said to be obtained from (Γ, \mathcal{G}) by a *collapse move*. The reverse of this move is called an *expansion move*. A finite sequence of these moves is called an *elementary deformation*.

Remark 6.3. We note that if an edge e (not a loop) has $G_{t(e)} = \alpha_e(G_e)$, then e contributes to a free amalgamated product $G_{o(e)} *_{G_e} G_e$ in $\pi_1(\Gamma, \mathcal{G})$. Therefore the above two moves correspond to the graph of groups isomorphism $G_{o(e)} *_{G_e} G_e \cong G_{o(e)}$. Thus we can always collapse this edge in Γ without changing the fundamental group. See Figure 2.




FIGURE 2. Example of collapse and expansion moves.

Definition 6.4. Given G-tree T, the unnormalized deformation space \mathcal{X} containing T is the set of all metric G-trees that have the same elliptic subgroups as T. We can projectivize \mathcal{X} by taking the quotient of \mathcal{X} under the action of \mathbb{R}^+ given by rescaling the metric on a given metric G-tree. The quotient \mathcal{D} is called a deformation space of G-trees.

Culler and Vogtmann's Outer space is a celebrated example of a deformation space for a finitely generated free group where the only elliptic subgroup is the trivial group. For now, the deformation space \mathcal{D} is just a set, with no additional structure. However, we can view it as an infinite dimensional simplicial complex. In particular, each projectivized G-tree $T \in \mathcal{D}$ corresponds to a n-dimensional simplex where n is the number of edges in the quotient graph T/G, and faces of this simplex correspond to projectivized G-trees T' that can be obtained from T by a sequence of collapse moves. We note that, by Forester's deformation theorem [For02, Theorem 1.1], any two projectivized G-trees $T, T' \in \mathcal{D}$ (disregarding the metric) are related by an elementary deformation. Thus the deformation space \mathcal{D} is in fact a connected simplicial complex. There is a natural action of $\operatorname{Out}(G)$ on \mathcal{D} given by precomposing of the action of G on G by an outer automorphism of G.

6.2. Two graph of groups structures of Out(BS(p,q)). In this subsection, we describe the two graph of groups structures of Out(BS(p,q)), and we briefly explain how Clay computed the presentation of Out(BS(p,q)) (see Section 4.3 in [Cla09] for details).

Let $G_{p,q}$ denote the Baumslag-Solitar group $\mathrm{BS}(p,q)$, and let $\mathcal{D}_{p,q}$ be the canonical deformation space of $G_{p,q}$. First, Clay defined an $G_{p,q}$ -invariant deformation retract $X_{p,q}$ of $\mathcal{D}_{p,q}$, where each vertex of $X_{p,q}$ corresponds to a particular type of reduced $G_{p,q}$ -tree. He showed that [Cla09, Theorem 3.9] this subcomplex $X_{p,q} \subseteq \mathcal{D}$ is in fact a tree. Furthermore, he showed that [Cla09, Proposition 4.1] the quotient graph $X_{p,q}/G_{p,q}$ is an infinite ray as pictured in Figure 3. Thus, by Bass-Serre theory, the quotient graph $X_{p,q}/G_{p,q}$ admits a graph of groups structure $(\Gamma_1, \mathcal{G}_1)$ where $\Gamma_1 = X_{p,q}/G_{p,q}$. We now discuss this graph of groups structure in details.




FIGURE 3. The graph of groups structure $(\Gamma_1, \mathcal{G}_1)$ of Out(BS(p,q)).

Clay showed that the vertex group G_{v_0} is isomorphic to the dihedral group $\mathbb{Z}_{p|n-1|} \rtimes \mathbb{Z}_2$, generated by the following automorphisms:

$$\psi: \quad x \mapsto x \qquad \qquad \iota: \quad x \mapsto x^{-1}$$
$$t \mapsto xt \qquad \qquad t \mapsto t$$

and for $k \geq 1$, the vertex group G_{v_k} is isomorphic to the dihedral group $\mathbb{Z}_{n^k|n-1|} \rtimes \mathbb{Z}_2$ generated by the following automorphisms:

$$\phi_k: \quad x \mapsto x \qquad \qquad \iota: \quad x \mapsto x^{-1}$$
$$t \mapsto (t^{-k}x^p t^k)t \qquad \qquad t \mapsto t$$

Notice that we have the following relations: $\psi^{p(n-1)} = c_x^{-p}$ (where c_x is the inner automorphism $g \mapsto xgx^{-1}$), $\iota\psi = \psi^{-1}\iota$, $\phi_{k+1}^n = \phi_k$ for $k \ge 1$, $\phi_1^n = \psi^p$ and $\iota\phi_k = \phi_k^{-1}\iota$.

In summary, we have the vertex groups of (Γ_1, \mathcal{G}) are

$$G_{v_0} = \langle \psi, \iota \mid \psi^{p(n-1)} = \iota^2 = 1, \ \iota \psi = \psi^{-1} \iota \rangle$$

and

$$G_{v_k} = \langle \phi_k, \iota \mid \iota^2 = 1, \ \phi_k^n = \phi_{k-1}, \ \iota \phi_k = \phi_k^{-1} \iota \rangle$$

The edge groups of $(\Gamma_1, \mathcal{G}_1)$ are

$$\alpha_{e_0}(G_{e_0}) = \langle \phi_1^n, \iota \rangle \leq G_{v_1}, \ \alpha_{\overline{e_0}}(G_{e_0}) = \langle \psi^p, \iota \rangle \leq G_{v_0}$$

and

$$\alpha_{e_k}(G_{e_k}) = \langle \phi_k, \iota \rangle \leq G_{v_{k+1}}, \ \alpha_{\overline{e_k}}(G_{e_k}) = G_{v_k} \text{ for } k \geq 1$$

As graph of groups, the edge e_0 contributes to a free amalgamated product as follows.

$$G_{v_0} *_{G_{e_0}} G_{v_1} = (\mathbb{Z}_{|p(n-1)|} \rtimes \mathbb{Z}_2) *_{\mathbb{Z}_{|n-1|} \rtimes \mathbb{Z}_2} (\mathbb{Z}_{|n(n-1)|} \rtimes \mathbb{Z}_2)$$

and each e_k for $k \geq 1$ contributes to a free amalgamated product as follows.

$$G_{v_k} *_{G_{e_k}} G_{v_{k+1}} = (\mathbb{Z}_{|n^k(n-1)|} \rtimes \mathbb{Z}_2) *_{\mathbb{Z}_{|n^k(n-1)|} \rtimes \mathbb{Z}_2} (\mathbb{Z}_{|n^{k+1}(n-1)|} \rtimes \mathbb{Z}_2)$$

We note that each e_k for $k \geq 1$ is in fact a collapsible edge. So the infinite ray Γ_1 can collapse to a segment pictured as in Figure 4, where one vertex corresponding to v_0 and the other vertex corresponding to the end represented by (v_1, v_2, \dots) .

$$v_0 \qquad \qquad (v_1, v_2, \dots)$$

$$e_0$$

FIGURE 4. The graph of groups structure $(\Gamma_2, \mathcal{G}_2)$ of Out(BS(p,q)).

Since each G_{v_k} is embedded in $G_{v_{k+1}}$ as a subgroup for $k \geq 1$, the stabilizer of this other end is the direct limit:

$$\lim_{n \to \infty} \mathbb{Z}_{n^k|n-1|} \rtimes \mathbb{Z}_2 = \mathbb{Z}\left[\frac{1}{|n|}\right] / |n(n-1)| \rtimes \mathbb{Z}_2$$

Thus the group $Out(G_{p,q})$ can be presented as a free amalgamated product as follows.

$$\operatorname{Out}(G_{p,q}) = (\mathbb{Z}_{|p(n-1)|} \rtimes \mathbb{Z}_2) *_{\mathbb{Z}_{|n-1|} \rtimes \mathbb{Z}_2} (\mathbb{Z}[\frac{1}{|n|}]/|n(n-1)\mathbb{Z}| \rtimes \mathbb{Z}_2)$$
(1)

6.3. Quotient action of $\operatorname{Out}(\operatorname{BS}(p,q))$. In this subsection, we apply Corollary 4.8 to $\operatorname{Out}(G_{p,q})$. Denote the Bass-Serre tree associated to the graph of groups $(\Gamma_2, \mathcal{G}_2)$ by $T_{p,q}$. We begin by showing that the tree action $\operatorname{Out}(G_{p,q}) \curvearrowright T_{p,q}$ is non-elementary acylindrical. In fact, this follows from a more general result as follows.

Proposition 6.5. Let $G = A *_D B$ be a free amalgamented product where D is finite. If $[A:D] \ge 2$ and [B:D] > 2, then the action of G on its Bass-Serre tree T is non-elementary acylindrical.

Proof. It follows from the definition that the action $G \curvearrowright T$ is (k, C)-acylindrical for k = 1 and C = |D|. Thus the action is acylindrical follows directly from Theorem 3.3. It remains to show that the action is non-elementary. Since $[A:D] \ge 2$ and [B:D] > 2, there exist a non-trivial element $a \in A \setminus D$ and two different non-trivial elements $b_1, b_2 \in B \setminus D$ such that ab_1 and ab_2 contribute to two hyperbolic elements. Since $b_1 \ne b_2$, we have b_1D, b_2D are two different edges in T. This shows that $Axis(ab_1)$ and $Axis(ab_2)$ represent two different branches of T respectively. Thus ab_1 and ab_2 are two independent hyperbolic elements. Therefore the action $G \curvearrowright T$ is non-elementary.

Corollary 6.6. $Out(G_{p,q}) \curvearrowright T_{p,q}$ is a non-elementary acylindrical action.

Proof. The proof follows directly from the presentation 1 and Proposition 6.5.

Corollary 6.7. Out(BS(p,q)) is an acylindrically hyperbolic group if and only if p divides q properly.

Proof. The proof follows directly from Remark 6.1 and Corollary 6.6. \Box

Remark 6.8. Note that when p divides q properly, it follows from the group presentation of $Out(G_{p,q})$ that the group is in fact hyperbolic relative to the infinitely generated vertex group in the amalgamented product as the edge group is finite.

We now are ready to apply Corollary 4.8 to the action $\operatorname{Out}(G_{p,q}) \curvearrowright T_{p,q}$. Recall that the presentation 1 of $\operatorname{Out}(G_{p,q})$ is a free amalgamated product, we then denote $\operatorname{Out}(G_{p,q}) = A *_C B$ where A, B and C are subgroups in presentation 1 accordingly. To have an equivariant family of subgroups, by Remark 4.2, it suffices to choose proper normal subgroups R_A in A and R_B in B. We provide two examples as follows. Consider the action $\operatorname{Out}(G_{4,12}) \curvearrowright T_{4,12}$. In Example 6.9, we provide an equivariant family of subgroups $\{R_v\}$ of $\operatorname{Out}(G_{4,12})$ that satisfies conditions of Corollary 4.8, and thus shows that the quotient action $\operatorname{Out}(G_{4,12})/\langle R_v \rangle \curvearrowright T_{2,4}/\langle R_v \rangle$ is non-elementary acylindrical and the quotient group is $\operatorname{Out}(G_{4,12})/\langle R_v \rangle$ is acylindrically hyperbolic. In contrast, we provide an equivariant family of subgroups of $\operatorname{Out}(G_{4,12})$ in Example 6.10 that violates conditions of Corollary 4.8. The quotient action is still acylindrical, however, it is an elementary acylindrical action.

Example 6.9 (Infinite quotient). First, we have the following presentation of $Out(G_{4,12})$.

$$\operatorname{Out}(G_{4,12}) = A *_{C} B = (\mathbb{Z}_{8} \rtimes \mathbb{Z}_{2}) *_{\mathbb{Z}_{2} \rtimes \mathbb{Z}_{2}} (\mathbb{Z}[\frac{1}{3}]/6\mathbb{Z} \rtimes \mathbb{Z}_{2})$$

where A is generated by $\{\psi, \iota\}$, B is generated by $\{\iota, \phi_1, ..., \phi_k, ...\}$. We also have the following relation $\psi^4 = \phi_1^3$. It follows that C is generated by $\{\psi^4, \iota\}$ (resp. $\{\phi_1^3, \iota\}$) as subgroup of A (resp. B). Further, we have $\psi^8 = \iota^2 = 1$ and $\phi_k^{3^k} = \phi_1$ for $k \geq 1$. This shows that all translates of A in $T_{4,12}$ has valence 4 whereas all translates of B in $T_{4,12}$ has infinite




FIGURE 5. A local picture of the tree $T_{4,12}$. Every translate of A has valence 4 and every translate of B has infinite valence. The two red lines represent $Axis(\psi\phi_i)$ and $Axis(\psi\phi_i)$ respectively.

valence. See Figure 5 for the picture of $T_{4,12}$. Let us now consider R_A is the subgroup of A generated by $\{\psi^2, \iota\}$ and R_B is the subgroup of B generated by $\{\phi_1, \iota\}$. Since both A and B are abelian groups, we have R_A and R_B are both normal subgroups. One can verify that, under these choices, the equivariant family of subgroups $\{R_v\}$ satisfies all conditions of Corollary 4.8. In fact, the quotient tree $T_{4,12}/\langle R_v \rangle$ is pictured as in Figure 6. In addition, each pair of elements $\psi\phi_i, \psi\phi_j$ for $i, j \geq 2$ and $i \neq j$ contribute to a pair of independent hyperbolic elements for the quotient action. Thus we have $\mathrm{Out}(G_{4,12})/\langle R_v \rangle \curvearrowright T_{4,12}/\langle R_v \rangle$ is non-elementary acylindrical, and thus the quotient group $\mathrm{Out}(G_{4,12})/\langle R_v \rangle$ is acylindrically hyperbolic.




FIGURE 6. The infinite quotient tree $T_{4,12}/\langle R_v \rangle$ in Example 6.9. The two red lines represent $Axis(\psi \phi_i)$ and $Axis(\psi \phi_i)$ respectively.

Example 6.10 (Finite quotient). Let us now consider R_A is the subgroup of A generated by $\{\psi^3, \iota\}$ and R_B is the subgroup of B still generated by $\{\phi_1, \iota\}$. Then we have $\langle C, R_A \rangle = A$

violates condition (1) of Corollary 4.8. In this case, $T_{4,12}/\langle R_v \rangle$ is a finite diameter tree pictured as in Figure 6. The quotient action $\operatorname{Out}(G_{4,12})/\langle R_v \rangle \curvearrowright T_{4,12}/\langle R_v \rangle$ is still acylindrical, however, there are no hyperbolic elements. Thus the quotient action is an elementary action.




FIGURE 7. The finite quotient tree $T_{4,12}/\langle R_v \rangle$ in Example 6.10.

6.4. Comparing tree actions of $\operatorname{Out}(\operatorname{BS}(p,q))$. In this subsection, we compare the two tree actions $\operatorname{Out}(G_{p,q}) \curvearrowright T_{p,q}$ and $\operatorname{Out}(G_{p,q}) \curvearrowright X_{p,q}$ in the sense of Section 5, and we provide a proof of our last main result, Theorem E.

We begin by applying Theorem 5.7 to the action $\operatorname{Out}(G_{p,q}) \curvearrowright T_{p,q}$ and thus showing that this action is the largest acylindrical action.

Proposition 6.11. $Out(G_{p,q})$ admits the largest acylindrical action on the Bass-Serre tree $T_{p,q}$.

Proof. Note that Proposition 6.6 shows that the action of $Out(G_{p,q})$ on $T_{p,q}$ is non-elementary acylindrical. Since the vertex groups contain finite order elements only, then the hypotheses of Corollary 5.8 are satisfied and hence the proof follows.

Now we turn our attention to the other tree action $\operatorname{Out}(G_{p,q}) \curvearrowright X_{p,q}$. We first give a short description of the tree $X_{p,q}$ (full details can be found in Section 3.2 of [Cla09]). By Bass-Serre theory, we can lift Γ_1 to an infinite ray $\tilde{\Gamma}_1$ in $X_{p,q}$ such that $\operatorname{Out}(G_{p,q}) \cdot \tilde{\Gamma}_1 = X_{p,q}$. We denote the lift of v_k in $\tilde{\Gamma}_1$ by \tilde{v}_k . Thus each vertex in $X_{p,q}$ is actually a translation of some \tilde{v}_k . We say a vertex has level k if it is a translation of \tilde{v}_k . By definition, the action $\operatorname{Out}(G_{p,q}) \curvearrowright X_{p,q}$ preserves the level of each vertex. Recall that the edge group G_{e_k} has index |n| in $G_{v_{k+1}}$ and index 1 in G_{v_k} , so each level k vertex in $X_{p,q}$ has valence |n|+1. Since G_{e_0} has index p in G_{v_0} , then each level 0 vertex in $X_{p,q}$ has valence p. For example, to highlight the level of each vertex, we have a piece of the tree $X_{2,4}$ pictured as in Figure 8. The entire tree $X_{2,4}$ can be obtained by gluing 2 pieces along a level 0 vertex, and passing to the new tree constructed by the gluing process.

We are ready now to show that the action $\operatorname{Out}(G_{p,q}) \curvearrowright X_{p,q}$ is not an acylindrical action.

Proposition 6.12. The action $Out(G_{p,q}) \curvearrowright X_{p,q}$ is not acylindrical.

Proof. By definition 3.1, we need to show that there exists an $\varepsilon > 0$ such that for any R, N > 0, there exists x, y in $X_{p,q}$ with $d(x, y) \geq R$ and

$$|\{g \in \operatorname{Out}(G_{p,q}) \mid d(x,gx) \le \varepsilon, \ d(y,gy) \le \varepsilon\}| > N$$

Consider $\varepsilon = \frac{1}{2}$. Let R, N > 0 be fixed. We can choose k > 0 such that $|n^k(n-1)| > N$. Let \tilde{v}_l be a vertex in $\tilde{\Gamma}_1$ such that l > k and $d(\tilde{v}_l, \tilde{v}_k) \geq R$. Then, by choosing $x = \tilde{v}_l$ and $y = \tilde{v}_k$, we have




FIGURE 8. A piece of the Bass-Serre tree $X_{2,4}$.

$$|\{g \in \operatorname{Out}(G_{p,q}) \mid d(\tilde{v}_l, g\tilde{v}_l) \leq \varepsilon, \ d(\tilde{v}_k, g\tilde{v}_k) \leq \varepsilon\}| = |G_{v_k}| > |n^k(n-1)| > N$$

Hence the action is not acylindrical.

We next turn toward a brief discussion of weakly properly discontinuously action (WPD action). The definition of this type of action is essentially due to Bestvina and Fujiwara [BF02]. However, to obtain clearer and simpler proofs, we will follow an equivalent definition in [MO15].

Definition 6.13. [MO15, Definition 3.5] An element h of a group G acting isometrically on a metric space S satisfies the weak proper discontinuity condition (or h is a WPD element) if for some $s \in S$ (or, equivalently, for all $s \in S$) and every $\varepsilon \geq 0$, there is an $M \geq 0$ such that

$$|\{g \in G | d(s, gs) \le \varepsilon, \ d(h^M s, gh^M s) \le \varepsilon\}| < \infty$$

Definition 6.14. An action of a group G on a hyperbolic space S isometrically is called non-elementary WPD if there are at least two independent hyperbolic elements that satisfy the WPD condition.

The following theorem by Osin gives us different ways to prove a group to be acylindrically hyperbolic.

Theorem 6.15. [Osi16, Theorem 1.2] For any group G, the following are equivalent.

- (1) There exists a generating set X of G such that corresponding Cayley graph $\Gamma(G, X)$ is hyperbolic, $|\Gamma(G, X)| > 2$, and the natural action of G on $\Gamma(G, X)$ is acylindrical.
- (2) G admits a non-elementary acylindrical action on a hyperbolic space.
- (3) G is not virtually cyclic and admits an action on a hyperbolic space such that at least one element of G is loxodromic and satisfies the WPD condition.
- (4) G contains a proper infinite hyperbolically embedded subgroup.

It is clear that a non-elementary acylindrical action is a non-elementary WPD action. However the converse is not true in general. We have shown in Proposition 6.12 that the action $\operatorname{Out}(G_{p,q}) \curvearrowright X_{p,q}$ is not acylindrical. Next, we show that this action is in fact a non-elementary WPD action.

Lemma 6.16. For any level 0-vertex w in $X_{p,q}$, the set $\{g \in Out(G_{p,q}) \mid g\tilde{v}_0 = w\}$ is finite.

Proof. For simplicity, we denote $S_w = \{g \in \operatorname{Out}(G_{p,q}) \mid g\tilde{v}_0 = w\}$. Let $m = d(w, \tilde{v}_0)$. As discussed in the proof of Theorem 1.17 of [Bas93], w can be uniquely represented by $g_1f_1g_2f_2...g_mf_mG_{v_0}$ where $(f_1, f_2, ..., f_m)$ is a reduced edge path in Γ_1 with $o(f_1) = t(f_m) = v_0$, and g_i is contained in the transversal set Σ_{f_i} for $G_{o(f_i)}/\alpha_{\overline{f}_i}(G_{f_i})$. If $g \in S_w$, we have $gG_{v_0} = g_1f_1g_2f_2...g_mf_mG_{v_0}$ which implies that $g \in g_1f_1g_2f_2...g_mf_mG_{v_0}$. Since every vertex group G_{v_k} and every edge group G_{e_k} in Γ_1 is finite, then there are only finitely many choices for Σ_{f_i} and each $|\Sigma_{f_i}|$ is finite. Thus g has only finitely many choices, which implies that S_w is finite.

Proposition 6.17. The action $Out(G_{p,q}) \curvearrowright X_{p,q}$ is a non-elementary WPD action.

Proof. For any $\varepsilon \geq 0$. Let $s = \tilde{v}_0 \in X_{p,q}$. We first note that, similar to the proof of Proposition 6.5, every pair of elements $\psi \phi_i$ and $\psi \phi_j$ for $i, j \geq 1$ and $i \neq j$ contributes to a pair of independent hyperbolic elements for the action $\operatorname{Out}(G_{p,q}) \curvearrowright X_{p,q}$. We then claim that each $\psi \phi_i$ satisfies WPD condition. By Definition 6.13, it suffices to show that

$$|\{g \in \operatorname{Out}(G_{p,q}) \mid d(\tilde{v}_0, g\tilde{v}_0) \le \varepsilon\}| < \infty$$

Note that there are only finitely many translates of \tilde{v}_0 inside a ε -neighborhood of \tilde{v}_0 . Thus, by Lemma 6.16, the set $\{g \in \text{Out}(G_{p,q}) \mid d(\tilde{v}_0, g\tilde{v}_0) \leq \varepsilon\}$ is finite. This completes the proof. \square

We remark that, by Theorem 6.15, the proposition above provides an alternative proof of acylindrical hyperbolicity of Out(BS(p,q)). We are ready now to prove our last main result, Theorem E, which can be restated as follows.

Theorem 6.18. Let $G = Out(G_{p,q})$ where q = pn for some p, |n| > 1. Then we have the following results:

- (1) $G \curvearrowright X_{p,q}$ is not an acylindrical action while $G \curvearrowright T_{p,q}$ is non-elementary acylindrical, thus $G \curvearrowright X_{p,q}$ is not equivalent to $G \curvearrowright T_{p,q}$;
- (2) $G \curvearrowright X_{p,q}$ and $G \curvearrowright T_{p,q}$ are both non-elementary WPD actions;
- (3) $G \curvearrowright X_{p,q}$ and $G \curvearrowright T_{p,q}$ have the same set of hyperbolic elements;
- (4) $G \curvearrowright T_{p,q}$ is the largest acylindrical hyperbolic action.

Proof. (1), (2) and (4) follows directly from Corollary 6.6, Proposition 6.12, Proposition 6.11 and Proposition 6.17. For (3), we note that $T_{p,q}$ can be obtained from $X_{p,q}$ by a sequence of G-equivariant collapse moves. This implies that an element $g \in G$ is elliptic in $G \curvearrowright X_{p,q}$ if and only if g is elliptic in $G \curvearrowright T_{p,q}$. Thus the two actions $G \curvearrowright X_{p,q}$ and $G \curvearrowright T_{p,q}$ have the same set of elliptic elements. It follows that they have the same set of hyperbolic elements.

References

- [ABD21] Carolyn Abbott, Jason Behrstock, and Matthew Gentry Durham. Largest acylindrical actions and stability in hierarchically hyperbolic groups. *Trans. Amer. Math. Soc. Ser. B*, 8:66–104, 2021. With an appendix by Daniel Berlyne and Jacob Russell.
- [ABO19] Carolyn Abbott, Sahana H. Balasubramanya, and Denis Osin. Hyperbolic structures on groups. Algebr. Geom. Topol., 19(4):1747–1835, 2019.
- [Bas93] Hyman Bass. Covering theory for graphs of groups. J. Pure Appl. Algebra, 89(1-2):3-47, 1993.
- [BF02] Mladen Bestvina and Koji Fujiwara. Bounded cohomology of subgroups of mapping class groups. Geom. Topol., 6:69–89, 2002.

- [BF10] Mladen Bestvina and Mark Feighn. A hyperbolic $Out(F_n)$ -complex. Groups Geom. Dyn., 4(1):31-58, 2010.
- [Bow08] Brian H. Bowditch. Tight geodesics in the curve complex. *Invent. Math.*, 171(2):281–300, 2008.
- [CL83] Donald J. Collins and Frank Levin. Automorphisms and Hopficity of certain Baumslag-Solitar groups. Arch. Math. (Basel), 40(5):385–400, 1983.
- [Cla09] Matt Clay. Deformation spaces of G-trees and automorphisms of Baumslag-Solitar groups. Groups Geom. Dyn., 3(1):39–69, 2009.
- [CM22] Matt Clay and Johanna Mangahas. Hyperbolic quotients of projection complexes. *Groups Geom. Dyn.*, 16(1):225–246, 2022.
- [CMM21] Matt Clay, Johanna Mangahas, and Dan Margalit. Right-angled Artin groups as normal subgroups of mapping class groups. *Compos. Math.*, 157(8):1807–1852, 2021.
- [Col78] Donald J. Collins. The automorphism towers of some one-relator groups. *Proc. London Math. Soc.* (3), 36(3):480–493, 1978.
- [Dah18] Fran, cois Dahmani. The normal closure of big Dehn twists and plate spinning with rotating families. Geom. Topol., 22(7):4113–4144, 2018.
- [DGO17] F. Dahmani, V. Guirardel, and D. Osin. Hyperbolically embedded subgroups and rotating families in groups acting on hyperbolic spaces. *Mem. Amer. Math. Soc.*, 245(1156):v+152, 2017.
- [DHS21] François Dahmani, Mark Hagen, and Alessandro Sisto. Dehn filling Dehn twists. *Proc. Roy. Soc. Edinburgh Sect. A*, 151(1):28–51, 2021.
- [For02] Max Forester. Deformation and rigidity of simplicial group actions on trees. *Geom. Topol.*, 6:219–267, 2002.
- [Gen19] Anthony Genevois. Negative curvature in automorphism groups of one-ended hyperbolic groups. J. Comb. Algebra, 3(3):305–329, 2019.
- [GH21] Anthony Genevois and Camille Horbez. Acylindrical hyperbolicity of automorphism groups of infinitely ended groups. *J. Topol.*, 14(3):963–991, 2021.
- [GHMR00] N. D. Gilbert, J. Howie, V. Metaftsis, and E. Raptis. Tree actions of automorphism groups. *J. Group Theory*, 3(2):213–223, 2000.
- [GM08] Daniel Groves and Jason Fox Manning. Dehn filling in relatively hyperbolic groups. Israel J. Math., 168:317-429, 2008.
- [MO15] Ashot Minasyan and Denis Osin. Acylindrical hyperbolicity of groups acting on trees. Math. Ann., 362(3-4):1055-1105, 2015.
- [Osi07] Denis V. Osin. Peripheral fillings of relatively hyperbolic groups. *Invent. Math.*, 167(2):295–326, 2007.
- [Osi16] D. Osin. Acylindrically hyperbolic groups. Trans. Amer. Math. Soc., 368(2):851–888, 2016.
- [Osi18] Denis V. Osin. Groups acting acylindrically on hyperbolic spaces. In Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. II. Invited lectures, pages 919–939. World Sci. Publ., Hackensack, NJ, 2018.
- [Sel97] Z. Sela. Acylindrical accessibility for groups. Invent. Math., 129(3):527–565, 1997.
- [Ser80] Jean-Pierre Serre. Trees. Springer-Verlag, Berlin-New York, 1980. Translated from the French by John Stillwell.
- [Val21] Motiejus Valiunas. Acylindrical hyperbolicity of groups acting on quasi-median graphs and equations in graph products. *Groups Geom. Dyn.*, 15(1):143–195, 2021.
- [Wei12] Richard Weidmann. On accessibility of finitely generated groups. Q. J. Math., 63(1):211–225, 2012.
 - B. Som: Department of Mathematics, University at Buffalo *Email address*: bratatis@buffalo.edu
 - D. WANG: YAU MATHEMATICAL SCIENCES CENTER, TSINGHUA UNIVERSITY, BEIJING, CHINA *Email address*: wangdaxun@mail.tsinghua.edu.cn