

SOME REMARKS ON PLANE CURVES RELATED TO FREENESS

ALEXANDRU DIMCA

ABSTRACT. For a free curve C of degree d with exponents (d_1, d_2) there is a simple formula relating d, d_1, d_2 and the total Tjurina number of C . Our first result discusses how this result changes when the curve C is no longer free. For a free line arrangement with exponents (d_1, d_2) , the Poincaré polynomial coincides with the Betti polynomial $B(t)$ and with the product $P(t) = (1 + d_1 t)(1 + d_2 t)$. Our second result shows that for any curve C , the difference $P(t) - B(t)$ is a polynomial $at + bt^2$, with a and b non-negative integers. Moreover $a = 0$ or $b = 0$ if and only if C is a free line arrangement. We also give a combinatorial restriction which a potential minimal counter-example to Terao's Conjecture must satisfy. Moreover, we make more precise a result by H. Schenck concerning the relation between the exponents and the maximal multiplicity of points in any line arrangement.

1. INTRODUCTION

We denote by $S = \mathbb{C}[x, y, z]$ the polynomial ring in three variables x, y, z with complex coefficients, and by $C : f = 0$ a reduced curve of degree $d \geq 3$ in the complex projective plane \mathbb{P}^2 . Let J_f be the Jacobian ideal of f , i.e., the homogeneous ideal in S spanned by the partial derivatives f_x, f_y, f_z of f , and by $M(f) = S/J_f$ the corresponding graded quotient ring, called the Jacobian (or Milnor) algebra of f . Consider the graded S -module of Jacobian syzygies of f or, equivalently, the module of derivations killing f , namely

$$(1.1) \quad D_0(f) = \{\theta \in \text{Der}(S) : \theta(f) = 0\}.$$

We say that $C : f = 0$ is an *m-syzygy curve* if the module $D_0(f)$ is minimally generated by m homogeneous syzygies, say $\rho_1, \rho_2, \dots, \rho_m$, of degrees $d_j = \deg \rho_j$ ordered such that

$$d_1 \leq d_2 \leq \dots \leq d_m.$$

We call these degrees (d_1, \dots, d_m) the *exponents* of the curve C . The smallest degree d_1 is sometimes denoted by $\text{mdr}(f)$ and is called the minimal degree of a Jacobian relation for f .

The S -module $D_0(f)$ is free if and only if $m = 2$, and in this case the curve C is said to be free. The exponents (d_1, d_2) of a free curve C of degree d are known to satisfy two key relations, namely

$$(1.2) \quad d_1 + d_2 = d - 1$$

2010 *Mathematics Subject Classification.* Primary 14H50; Secondary 13D02.

Key words and phrases. plane curve, free curve, Milnor algebra, minimal resolution, Tjurina number, Betti numbers, line arrangement.

and

$$(1.3) \quad d_1 d_2 = (d - 1)^2 - \tau(C),$$

where $\tau(C)$ denotes the total Tjurina number of C , see for instance [4, 9]. It is known that the condition (1.2) characterizes the free curves, that is a curve C which is not free satisfies

$$(1.4) \quad d_1 + d_2 \geq d,$$

with equality exactly for the plus-one generated curves, see [8, Theorem 2.3] for details. The first main result of this note is to completely characterize the curves satisfying the second condition (1.3), see Theorem 2.2. Beyond the free curves, the equality (1.3) is satisfied only for 3-syzygy curves with $d_3 = d - 1$ and for all the other curves one has

$$(1.5) \quad d_1 d_2 > (d - 1)^2 - \tau(C).$$

When $C = \mathcal{A}$ is a line arrangement in \mathbb{P}^2 , one can define its Poincaré polynomial $\pi(\mathcal{A}, t)$ and it is well known that

$$\pi(\mathcal{A}, t) = (1 + d_1 t)(1 + d_2 t)$$

if \mathcal{A} is free with exponents (d_1, d_2) , see [15, 3]. Moreover, the Poincaré polynomial $\pi(\mathcal{A}, t)$ coincides with the Betti polynomial $B(M(\mathcal{A}))(t)$ of the complement $M(\mathcal{A}) = \mathbb{P}^2 \setminus \mathcal{A}$, see (3.1) for the general definition of the Betti polynomial and see [15, 3] for the equality $\pi(\mathcal{A}, t) = B(M(\mathcal{A}))(t)$. Hence the above relation may be written as

$$(1.6) \quad B(M(\mathcal{A}))(t) = (1 + d_1 t)(1 + d_2 t).$$

The second main result of this note is Theorem 3.1, which describes the relation between $B(M(C))(t)$ and the product $(1 + d_1 t)(1 + d_2 t)$ for any curve C in \mathbb{P}^2 . It shows that the difference

$$(1 + d_1 t)(1 + d_2 t) - B(M(C))(t)$$

is a polynomial $a(C)t + b(C)t^2$, with $a(C)$ and $b(C)$ non-negative integers. Moreover $a(C) = 0$ or $b(C) = 0$ implies that C is a free line arrangement.

As applications of Theorems 2.2 and 3.1 we show that any irreducible component of a free curve with $d_1 = 1$ is rational, see Remark 3.3 and Corollary 3.4.

The first result in Section 5 gives a combinatorial restriction on the number of multiple points on any line in the arrangement \mathcal{A} , which a potential minimal counter-example \mathcal{A} to Terao's Conjecture must satisfy, see Theorem 4.3. The second result in Section 5 makes more precise a claim in [17, Lemma 5.2] concerning the relation between the exponents and the maximal multiplicity of multiple points in a line arrangement, see Theorem 4.5. We thank Piotr Pokora for drawing our attention on this question.

2. ON PLANE CURVES SATISFYING $\tau(C) = (d-1)^2 - d_1d_2$

We recall first the construction of the Bourbaki ideal $B(C, \rho_1)$ associated to a degree d reduced curve $C : f = 0$ and to a minimal degree non-zero syzygy $\rho_1 \in AR(f)$, see [7]. For any choice of the syzygy $\rho_1 = (a_1, b_1, c_1)$ with minimal degree d_1 , we have a morphism of graded S -modules

$$(2.1) \quad S(-d_1) \xrightarrow{u} AR(f), \quad u(h) = h \cdot \rho_1.$$

For any homogeneous syzygy $\rho = (a, b, c) \in AR(f)_m$, consider the determinant $\Delta(\rho) = \det M(\rho)$ of the 3×3 matrix $M(\rho)$ which has as first row x, y, z , as second row a_1, b_1, c_1 and as third row a, b, c . Then it turns out that $\Delta(\rho)$ is divisible by f , see [4], and we define thus a new morphism of graded S -modules

$$(2.2) \quad AR(f) \xrightarrow{v} S(d_1 - d + 1), \quad v(r) = \Delta(\rho)/f,$$

and a homogeneous ideal $B(C, \rho_1) \subset S$ such that $\text{im } v = B(C, \rho_1)(d_1 - d + 1)$. It is known that the ideal $B(C, \rho_1)$, when C is not a free curve, defines a 0-dimensional subscheme in \mathbb{P}^2 , which is locally a complete intersection, see [7, Theorem 5.1].

Secondly, for the reader's convenience, we recall below the main claim of [8, Theorem 3.5], in a better formulation. Recall that $t(C) = d_1 + d_2 - d + 1$ is the type of the curve C , see [2, Definition 1.2].

Theorem 2.1. *Let $C : f = 0$ be a reduced curve of degree $d \geq 3$ with exponents $d_1 \leq \dots \leq d_m$ with $m \geq 3$, and let ρ_1 be a non-zero syzygy of minimal degree d_1 . Let d' be the smallest integer such that*

$$d_3 \leq d' \leq \min(d_m, d - 1)$$

and the linear system $B(C, \rho_1)_{d_1+d'-d+1}$ has a 0-dimensional base locus. Then

$$(2.3) \quad \tau(C) \geq (d-1)^2 - d_1d_2 + (d-1-d')t(C).$$

and equality holds if and only if C is a 3-syzygy curve and then $d' = d_3$.

Proof. As explained in the proof of [8, Theorem 3.5], the integer d' above does indeed exist and one has

$$\tau(C) \geq (d-1)(d-d_1-1) + d_1^2 - [d_1 - (d-1-d_2)][d_1 - (d-1-d')].$$

A direct simple transformation of the right hand member of the inequality in Theorem 2.1 yields the new inequality

$$(2.4) \quad \tau(C) \geq (d-1)^2 - d_1d_2 + (d-1-d')(d_1 + d_2 - d + 1).$$

This yields our new formulation of [8, Theorem 3.5]. □

By definition of d' we have

$$d-1-d' \geq 0$$

and moreover

$$t(C) = d_1 + d_2 - d + 1 > 0$$

when C is not free, by (1.4). This implies the following result.

Theorem 2.2. *Let C be a reduced plane curve of degree d in \mathbb{P}^2 with exponents $d_1 \leq d_2 \leq \dots \leq d_m$, where $m \geq 3$. Then $\tau(C) = (d-1)^2 - d_1d_2$ if and only if $m = 3$ and $d_3 = d-1$. In all the other cases one has*

$$\tau(C) > (d-1)^2 - d_1d_2.$$

Proof. If $m > 3$, then Theorem 2.1 tells us that the inequality (2.4) is strict, and hence $\tau(C) > (d-1)^2 - d_1d_2$. When $m = 3$, it follows by definition that $d' = d_3$. Then Theorem 2.1 tells us that the inequality (2.4) is an equality and hence we have

$$\tau(C) = (d-1)^2 - d_1d_2 + (d-1-d_3)(d_1+d_2-d+1),$$

which is equivalent to the formula given in [8, Proposition 2.1 (4)] for $\tau(C)$. Therefore we get that $\tau(C) = (d-1)^2 - d_1d_2$ in this case if and only if $d_3 = d-1$. \square

Corollary 2.3. *Let \mathcal{A} be an arrangement of d lines in the projective plane \mathbb{P}^2 with exponents $d_1 \leq d_2 \leq \dots \leq d_m$, where $m \geq 2$. Then the following inequality holds*

$$\tau(\mathcal{A}) \geq (d-1)^2 - d_1d_2 + t(\mathcal{A}),$$

where $t(\mathcal{A})$ is the type of the arrangement \mathcal{A} . In particular, \mathcal{A} is free if and only if

$$\tau(\mathcal{A}) = (d-1)^2 - d_1d_2.$$

Proof. Recall that for line arrangements the maximal degree d_m in the exponents of C is bounded by $d-2$, see [17, Corollary 3.5]. Then the first claim follows from the inequality (2.3). The second claim follows by (1.3) and Theorem 2.2. \square

Corollary 2.4. *Let C be a reduced plane curve of degree d in \mathbb{P}^2 with exponents $d_1 \leq d_2 \leq \dots \leq d_m$, where $m \geq 2$. Then*

$$(d-1)^2 - d_1d_2 \leq \tau(C) \leq (d-1)^2 - d_1(d-1-d_1).$$

Proof. Theorem 2.2 gives the first inequality, while the second inequality comes from the upper-bound on $\tau(C)$ given in [9]. \square

Note that C is free if and only if $d_2 = d-1-d_1$ and then both inequalities in Corollary 2.4 become equalities.

Remark 2.5. Since $d_2 \leq d-1$, see for instance [8, Theorem 2.4], it follows that

$$\tau(C) \geq (d-1)^2 - d_1d_2 \geq (d-1)^2 - d_1(d-1) = (d-1)(d-1-d_1).$$

Hence our Corollary 2.4 can be regarded as an improvement of the lower bound for $\tau(C)$ given in [9]. Moreover, the equality $\tau(C) = (d-1)(d-1-d_1)$ holds if and only if C is a 3-syzygy curve with $d_2 = d_3 = d-1$, see [8, Theorem 3.5]

Example 2.6. Examples of 3-syzygy curves such that $d_2 = d_3 = d-1$ are given by the Thom-Sebastiani curves described in [8, Example 4.5] and by the union of a smooth degree $(d-1)$ curve with one of its generic secants, see [8, Example 4.3 (i)].

Examples of 3-syzygy curves such that $d_2 = d-2$ and $d_3 = d-1$ are given by the union of a smooth Fermat degree $(d-1)$ curve with a inflectional tangent meeting the curve in a single point, see [8, Example 4.3 (ii)].

An example consisting of the Klein quartic and 4 bitangents that is a plus-one generated curve with $d_3 = d - 1$ can be found in [12, Proposition 4.11].

3. EULER POLYNOMIALS OF COMPLEMENTS OF PLANE CURVES

For any topological space M , having the homotopy type of a finite CW -complex of dimension 2, we define the Betti polynomial $B(M)$ of M by the formula

$$(3.1) \quad B(M)(t) = b_0(M) + b_1(M)t + b_2(M)t^2,$$

where $b_j(M)$ denotes the j -th Betti number of M . In particular, we can define the polynomial $B(M(C))$, where $M(C)$ is the complement $\mathbb{P}^2 \setminus C$ of a reduced plane curve C in \mathbb{P}^2 . In this case, we have the following result.

Theorem 3.1. *Let C be a reduced plane curve of degree d in \mathbb{P}^2 with exponents $d_1 \leq d_2 \leq \dots \leq d_m$, where $m \geq 2$. Let*

$$\alpha(C) = \tau(C) - ((d-1)^2 - d_1d_2).$$

If e denotes the number of irreducible components of C , then one has

$$(1 + d_1t)(1 + d_2t) - B(M(C))(t) = a(C)t + b(C)t^2,$$

where $a(C) = d_1 + d_2 - e + 1$ and $b(C) = \mu(C) - \tau(C) + d - e + \alpha(C)$, with $\mu(C)$ denoting the total Milnor number of C . In particular, one has the following.

- (1) $a(C) = t(C) + (d - e) \geq 0$ and $b(C) \geq 0$;
- (2) *any of the two equalities $a(C) = 0$ and $b(C) = 0$ holds if and only if C is a free line arrangement, and then the other equality also holds.*

Proof. Recall that $b_1(M(C)) = e - 1$ for any curve C . On the other hand, the Euler numbers of the spaces under consideration satisfy

$$E(M(C)) = E(\mathbb{P}^2) - E(C) = 3 - E(C) \text{ and } E(C) = E(C_d) + \mu(C),$$

where C_d is a smooth curve of degree d . Hence

$$E(C_d) = 2 - 2g = 2 - (d-1)(d-2).$$

Putting these formula together yields

$$b_2(M(C)) = (d-1)^2 - \mu(C) - (d-e),$$

and this clearly prove our first claim. For the second claim we notice that in general we have

$$d_1 + d_2 \geq d - 1 \geq e - 1$$

which implies that $a(C) \geq 0$. The equality $a(C) = 0$ implies $t(C) = d_1 + d_2 - d + 1 = 0$ and $e = d$, namely that C is a free line arrangement.

Moreover, $\mu(C) \geq \tau(C)$, with equality if and only if all the singularities of C are quasi homogeneous. Corollary 2.4 shows that $\alpha(C) \geq 0$ and since $d \geq e$ obviously, we get $b(C) \geq 0$. Finally, $b(C) = 0$ implies that $d = e$, and hence C is a line arrangement, and also yields $\alpha(C) = 0$. Corollary 2.3 implies that C is free and this completes the proof of the second claim. \square

Corollary 3.2. *Let C be a reduced plane curve of degree d , having e irreducible components. If C is free with exponents (d_1, d_2) , then*

$$(1 + d_1t)(1 + d_2t) = B(M(C))(t) + (d - e)t(1 + t) + (\mu(C) - \tau(C))t^2.$$

In particular, if in addition all the singularities of C are quasi homogeneous, one has

$$(1 + d_1t)(1 + d_2t) = B(M(C))(t) + (d - e)t(1 + t)$$

and the Euler number $E(M(C))$ satisfies

$$E(M(C)) = B(M(C))(-1) = (d_1 - 1)(d_2 - 1).$$

Proof. For a free curve C we have $d_1 + d_2 = d - 1$ and $\alpha(C) = 0$, recall (1.2) and (1.3). \square

We notice that the equality (1.6) follows also from Corollary 3.2, since $e = d$ in this case, see also [16].

Remark 3.3. The reduced plane curves with $d_1 = 1$ were studied in [10], where the interested reader may find out many explicit examples. It turns out that the curves with $d_1 = 1$ are either free or nearly free, that is 3-syzygy curves such that the exponents are $(1, d - 1, d - 1)$. This follows from [10, Proposition 1.3 (ii)] and the characterization of nearly free curves given in [4]. Moreover, such a curve admits a 1-dimensional connected group of symmetries H , see [10]. Since H is either (\mathbb{C}^*, \cdot) or $(\mathbb{C}, +)$, it follows that the closures of the 1-dimensional orbits of H are rational curves. This implies that any irreducible component of a plane curve C with $d_1 = 1$ is rational. When C is free, we can give the following alternative proof for this result.

Corollary 3.4. *Let C be a free curve with exponents (d_1, d_2) such that $d_1 = 1$. Then any irreducible component of C is a rational curve.*

Proof. Using Corollary 3.2 we get

$$E(C) = B(M(C))(-1) = (d_1 - 1)(d_2 - 1) - (\mu(C) - \tau(C)) \leq 0,$$

since $\mu(C) \geq \tau(C)$. Then using a conjecture of W. Veys in [18], proved by A. de Jong and J. Steenbrink in [13] and by R. Gurjar and A. Parameswaran in [11], it follows that any irreducible component of C is rational. \square

4. TWO RESULTS RELATED TO TERAQ'S CONJECTURE

First we recall two main results from [6] in the setting of line arrangements in \mathbb{P}^2 . Similar results appear also in [1, Theorem 1.11] and in [14, Theorems 3.5 and 3.6]. Since all the singularities of line arrangements are quasi-homogeneous, it follows that all the invariants ϵ in the quoted results are 0. First we restate [6, Theorem 1.3].

Theorem 4.1. *Let \mathcal{A}' be a line arrangement in \mathbb{P}^2 , L a line in \mathbb{P}^2 , which is not in \mathcal{A}' . We assume that the union $\mathcal{A} = \mathcal{A}' \cup L$ is a free curve with exponents (d_1, d_2) . Then the exponents (d'_1, d'_2) (resp. (d'_1, d'_2, d'_3)) of the free (resp. plus-one generated) arrangement \mathcal{A}' and the number $r = |\mathcal{A}' \cap L|$ of intersection points satisfy one of the following conditions, and all these three cases are possible.*

- (1) $d_1 < d_2$, $d'_1 = d_1$ and $d'_2 = d_2 - 1$. In this case \mathcal{A}' is free and $r = d_1 + 1$.

- (2) $d'_1 = d_1 - 1$ and $d'_2 = d_2$. In this case \mathcal{A}' is free and $r = d_2 + 1$.
(3) $d'_1 = d_1$ and $d'_2 = d_2$. In this case \mathcal{A}' is plus-one generated and

$$r = |\mathcal{A}'| - d'_3 \leq |\mathcal{A}| - 1 - d_2 = d_1.$$

In particular, \mathcal{A}' is free if and only if $r \geq d_1 + 1$.

Now we restate [6, Theorem 1.4].

Theorem 4.2. *Let \mathcal{A}' be a line arrangement in \mathbb{P}^2 , L a line in \mathbb{P}^2 , which is not in \mathcal{A}' . We consider the union $\mathcal{A} = \mathcal{A}' \cup L$ and assume that \mathcal{A}' is a free arrangement with exponents (d'_1, d'_2) . Then the exponents (d_1, d_2) (resp. (d_1, d_2, d_3)) of the free (resp. plus-one generated) \mathcal{A} and the number $r = |\mathcal{A}' \cap L|$ of intersection points satisfy one of the following conditions, and all these three cases are possible.*

- (1) $d_1 = d'_1$ and $d_2 = d'_2 + 1$. In this case \mathcal{A} is free and $r = d'_1 + 1$.
(2) $d'_1 < d'_2$, $d_1 = d'_1 + 1$ and $d_2 = d'_2$. In this case \mathcal{A} is free and $r = d'_2 + 1$.
(3) $d_1 = d'_1 + 1$ and $d_2 = d'_2 + 1$. In this case \mathcal{A} is plus-one generated and

$$r = d_3 + 1 \geq d_2 + 1 = d'_2 + 2.$$

In particular, \mathcal{A} is free if and only if $r \leq d'_2 + 1$.

We recall that (a partial case of) Terao's conjecture says that if \mathcal{A} and \mathcal{B} are two line arrangements in \mathbb{P}^2 having the same combinatorics, and if \mathcal{A} is free, then \mathcal{B} is also free. By a *minimal counter example to Terao's Conjecture* we mean a pair \mathcal{A}, \mathcal{B} as above such that Terao's conjecture holds for all pairs $\mathcal{A}', \mathcal{B}'$ with $|\mathcal{A}'| = |\mathcal{B}'| < |\mathcal{A}| = |\mathcal{B}|$, but not for the pair \mathcal{A}, \mathcal{B} . Our next result gives some information on such a minimal counter example to Terao's Conjecture, supposing it exists.

Theorem 4.3. *Let \mathcal{A}, \mathcal{B} be a minimal counter example to Terao's Conjecture, such that \mathcal{A} is free with exponents $d_1 \leq d_2$. Then the following property holds for the free arrangement \mathcal{A} : for any line $L \in \mathcal{A}$, the number r_L of multiple points of \mathcal{A} situated on L satisfies the inequality*

$$r_L \leq d_1.$$

Proof. We prove that if there is a line $L \in \mathcal{A}$ such that

$$r_L \geq d_1 + 1$$

then \mathcal{B} is also free, and hence \mathcal{A}, \mathcal{B} is not a counter example to Terao's Conjecture. If such a line L exists, then we denote by L' the line in \mathcal{B} corresponding to the line $L \in \mathcal{A}$ with $r_L \geq d_1 + 1$ under the isomorphism of intersection lattices $L(\mathcal{A}) \simeq L(\mathcal{B})$. Let $\mathcal{A}' = \mathcal{A} \setminus \{L\}$ and $\mathcal{B}' = \mathcal{B} \setminus \{L'\}$ be the two deleted arrangements obtained from \mathcal{A} and \mathcal{B} by deleting the line L and respectively L' . Clearly

$$r_{L'} = |\mathcal{B}' \cap L'| = r_L.$$

We apply Theorem 4.1 and conclude that \mathcal{A}' is a free line arrangement. Indeed, this happens in the cases (1) and (2) of Theorem 4.1. In the remaining case (3), we have $r_L = |\mathcal{A}' \cap L| \leq d_1$, hence this case cannot occur in our situation.

On the other hand, it is clear that the intersection lattice isomorphism $L(\mathcal{A}) \simeq L(\mathcal{B})$ yields a new intersection lattice isomorphism $L(\mathcal{A}') \simeq L(\mathcal{B}')$. Since \mathcal{A}, \mathcal{B} was

supposed to be a *minimal* counter example to Terao's Conjecture, it follows that Terao's Conjecture holds for the pair \mathcal{A}' , \mathcal{B}' and hence \mathcal{B}' is also a free line arrangement with the same exponents as \mathcal{A}' , call them $d'_1 \leq d'_2$. We apply now Theorem 4.2 to the arrangements \mathcal{B}' and \mathcal{B} , and note that in case (1) of Theorem 4.1 we have

$$r_{L'} = r_L = d'_1 + 1 = d_1 + 1 \leq d_2 + 1$$

and hence \mathcal{B} is free by Theorem 4.2. Similarly, in case (2) of Theorem 4.1 we have

$$r_{L'} = r_L = d_1 + 1 = d_2 + 1 = d'_2 + 1$$

and hence again \mathcal{B} is free. □

Remark 4.4. The monomial line arrangement

$$\mathcal{A} = \mathcal{A}(m, m, 3) : (x^m - y^m)(y^m - z^m)(x^m - z^m) = 0$$

is free with exponents $d_1 = m + 1$ and $d_2 = 2m - 2$ for $m \geq 3$, see for instance [3, Example 8.6 (i)]. For any line $L \in \mathcal{A}$ one has $r_L = m + 1$ so the condition $r_L \leq d_1$ from Theorem 4.3 holds. On the other hand, the full monomial line arrangement

$$\mathcal{A} = \mathcal{A}(m, 1, 3) : xyz(x^m - y^m)(y^m - z^m)(x^m - z^m) = 0$$

is free with exponents $d_1 = m + 1$ and $d_2 = 2m + 1$ for $m \geq 2$, see for instance [3, Example 8.6 (ii)]. For any line $L \in \mathcal{A}$ one has $r_L = m + 2$ so the condition $r_L \leq d_1$ from Theorem 4.3 does not hold. Similarly, this condition fails for the Hessian line arrangement which is free with exponents $d_1 = 4$ and $d_2 = 7$ and for any line L in it one has $r_L = 5$, see for instance [3, Example 8.6 (i)]. We believe that the condition $r_L \leq d_1$ from Theorem 4.3 limits a lot the possibilities for \mathcal{A} , but we have no result in this direction.

Let \mathcal{A} be a line arrangement with exponents $d_1 \leq \dots \leq d_m$ and let $m(\mathcal{A})$ be the maximal multiplicity of a multiple point in \mathcal{A} . Then Lemma 5.2 in [17] states (without a proof and in a rather cryptic way) that for the line arrangement \mathcal{A} one has

$$(4.1) \quad d_m \geq m(\mathcal{A}) - 1,$$

see Theorem 5.4 claim 5. in [17] for a clearer statement, where $M = m(\mathcal{A}) - 1$. In fact this claim can be improved as follows.

Theorem 4.5. *With the above notation, one has*

$$d_2 \geq t(\mathcal{A}) + m(\mathcal{A}) - 1 \geq m(\mathcal{A}) - 1,$$

where $t(\mathcal{A}) = d_1 + d_2 - d + 1 \geq 0$ is the type of the arrangement \mathcal{A} . In particular, the equality $d_2 = m(\mathcal{A}) - 1$ implies that \mathcal{A} is free, and for a non free arrangement \mathcal{A} one has

$$d_2 \geq m(\mathcal{A}).$$

Proof. The inequality

$$d_2 \geq (d_1 + d_2 - d + 1) + m(\mathcal{A}) - 1$$

is clearly equivalent to the inequality

$$d_1 \leq d - m(\mathcal{A}),$$

which is proven in [5, Theorem 1.2].

□

Remark 4.6. Consider the line arrangement

$$\mathcal{A} : f = (x^5 - y^5)(x + 2y + z)(x + 3y - 5z) = 0$$

consisting of 5 lines through a point plus two generic lines. A direct computation shows that \mathcal{A} has exponents $(2, 5, 5)$, hence \mathcal{A} is a nearly free arrangement satisfying $d_2 \geq m(\mathcal{A}) = 5$. The corresponding Poincaré polynomial is

$$\pi(\mathcal{A}, t) = (1 + 3t)^2.$$

This is a special case of Example 5.3 in [17], which seem to suggest that in that paper Lemma 5.2 *refers only to free arrangements*. Indeed, the author says that we can use this result in order to show that \mathcal{A} is not free, perhaps thinking that the Poincaré polynomial yields the exponents as in the free case.

Remark 4.7. There is the following relation between Theorems 4.3 and 4.5 above. Consider the arrangement \mathcal{A} in Theorem 4.3 and let $p \in \mathcal{A}$ be a point of maximal multiplicity $m(\mathcal{A})$. If L is a line in \mathcal{A} not passing through p , it is clear that

$$r_L \geq m(\mathcal{A})$$

since any line through p intersects L in a distinct point. If this inequality is an equality for any line L not passing through p , then p is a modular point, and hence \mathcal{A} is a supersolvable line arrangement. Since for supersolvable line arrangements Terao's conjecture holds, it follows that this is not the case, and hence there is a line L not through p with $r_L > m(\mathcal{A})$. This yields the stronger inequality

$$d_1 > m(\mathcal{A}),$$

for the arrangement \mathcal{A} in Theorem 4.3, which also follows from [5, Corollary 1.4]. This result says that for free arrangements \mathcal{A} with $m(\mathcal{A}) \geq d_1$ Terao's Conjecture holds.

REFERENCES

- [1] T. Abe, Plus-one generated and next to free arrangements of hyperplanes, *Int. Math. Res. Not.* 2021(2019), Issue 12, 9233–9261. 4
- [2] T. Abe, A. Dimca, P. Pokora, A new hierarchy for complex plane curves, arXiv:2410.11479. 2
- [3] A. Dimca, *Hyperplane Arrangements: An Introduction*, Universitext, Springer, 2017 1, 4.4
- [4] A. Dimca, Freeness versus maximal global Tjurina number for plane curves, *Math. Proc. Cambridge Phil. Soc.* 163 (2017), 161–172. 1, 2, 3.3
- [5] A. Dimca, Curve arrangements, pencils, and Jacobian syzygies, *Michigan Math. J.* 66 (2017), 347–365. 4, 4.7

- [6] A. Dimca, On plus-one generated curves arising from free curves. *Bull. Math. Sci.*, Vol. 14, No. 3 (2024) 2450007 (21 pages). 4, 4
- [7] A. Dimca, G. Sticlaru, On the jumping lines of bundles of logarithmic vector fields along plane curves, *Publ. Mat.* 64 (2020), 513–542. 2, 2
- [8] A. Dimca, G. Sticlaru, Plane curves with three syzygies, minimal Tjurina curves, and nearly cuspidal curves, *Geometriae Dedicata* 207 (2020), 29–49. 1, 2, 2, 2, 2, 2.5, 2.6
- [9] A.A. du Plessis, C.T.C. Wall, Application of the theory of the discriminant to highly singular plane curves, *Math. Proc. Cambridge Phil. Soc.*, 126(1999), 259-266. 1, 2, 2.5
- [10] A.A. du Plessis, C.T.C. Wall, Curves in $\mathbb{P}^2(\mathbb{C})$ with 1-dimensional symmetry, *Revista Mat. Complutense* 12 (1999), 117–131. 3.3
- [11] R. Gurjar, A. Parameswaran, Open surfaces with non-positive Euler characteristic, *Compositio Math.* 99(1995), 213-229. 3
- [12] M. Janasz, P. Pokora, M. Zieliński, On arrangements of smooth plane quartics and their bitangents, arXiv:2308.16514 2.6
- [13] A. de Jong, J. Steenbrink, Proof of a conjecture of W. Veys, *Indag. Math.* 6(1995), 99-104. 3
- [14] A. Măcinic and P. Pokora, Addition–deletion results for plus-one generated curves. *J Algebr. Comb.*, <https://doi.org/10.1007/s10801-024-01350-x>, (2024). 4
- [15] P. Orlik, H. Terao, *Arrangements of Hyperplanes*, Springer-Verlag, Berlin Heidelberg New York, 1992. 1
- [16] P. Pokora, On Poincaré polynomials for plane curves with quasi-homogeneous singularities, arXiv:2412.08436. 3
- [17] H. Schenck, Elementary modifications and line configurations in P^2 , *Comm. Math. Helv.* 78 (2003), 447–462. 1, 2, 4, 4, 4.6
- [18] W. Veys, Numerical data of resolutions of singularities and Igusa’s local zeta function, Ph. D. thesis, Univ. Leuven,1991. 3

UNIVERSITÉ CÔTE D’AZUR, CNRS, LJAD, FRANCE AND SIMION STOILow INSTITUTE OF MATHEMATICS, P.O. BOX 1-764, RO-014700 BUCHAREST, ROMANIA

Email address: Alexandru.DIMCA@univ-cotedazur.fr