

# ZYGMUND THEOREM FOR HARMONIC QUASIREGULAR MAPPINGS

DAVID KALAJ

ABSTRACT. Let  $K \geq 1$ . We prove Zygmund theorem for  $K$ -quasiregular harmonic mappings in the unit disk  $\mathbb{D}$  in the complex plane by providing a constant  $C(K)$  in the inequality

$$\|f\|_1 \leq C(K)(1 + \|\operatorname{Re}(f)\|_1 \log^+ \|\operatorname{Re} f\|_1).$$

Moreover for a quasiregular harmonic mapping  $f = (f_1, \dots, f_n)$  defined in the unit ball  $\mathbb{B} \subset \mathbb{R}^n$ , we prove the asymptotically sharp inequality

$$\|f\|_1 - |f(0)| \leq (n-1)K^2(\|f_1 \log f_1\|_1 - f_1(0) \log f_1(0)),$$

when  $K \rightarrow 1$ , provided that  $f_1$  is positive.

## CONTENTS

1. Introduction	1
1.1. Hardy class	1
1.2. Motivation	2
1.3. Quasiregular and harmonic mappings	2
1.4. Main results	3
2. Proof of main results	4
Funding	8
Data availability	8
Ethics declarations	8
References	8

## 1. INTRODUCTION

In this paper  $\mathbb{D}$  is the unit disk in the complex plane  $\mathbb{C}$  and  $\mathbb{B}$  is the unit ball in the Euclidean space  $\mathbb{R}^n$ . By  $\mathbb{S}$  we denote the unit sphere.

1.1. **Hardy class.** Let  $p > 0$  and

$$\mathbf{h}^p(\mathbb{B}) = \{f : \mathbb{B} \rightarrow \mathbb{R}^n, \text{ so that } \|f\|_p = \sup_{0 < r < 1} M_p(r, f) < \infty\},$$

where

$$M_p(r, f) = \left( \int_{\mathbb{S}} |f(r\zeta)|^p d\sigma(\zeta) \right)^{1/p}.$$

---

2020 *Mathematics Subject Classification.* Primary 30H10 .

*Key words and phrases.* Harmonic mappings, quasiregular mappings, Riesz inequality, Zygmund inequality.

Here  $\sigma$  is the standard normalized area measure of the unit sphere  $\mathbb{S}$ . Now define for  $x > 0$ ,  $\log^+(x) := \max\{\log x, 0\}$ . Then define the space  $\mathbf{h} \log^+ \mathbf{h}$  of functions defined in the unit ball so that

$$\sup_{0 < r < 1} \int_{\mathbb{S}} |f(r\zeta)| \log^+ |f(r\zeta)| d\sigma(\zeta) < \infty.$$

**1.2. Motivation.** We formulate the following Zygmund theorem [15, p. 254], which is the motivation for our study. Assume that  $f = u + iv$  is a holomorphic function defined in the unit disk  $\mathbb{D}$ . Assume further that  $|u| \log^+ |u| \in \mathbf{h}^1(\mathbb{D})$ . We write it shortly by  $u \in \mathbf{h} \log^+ \mathbf{h}$ . Then  $f \in \mathbf{h}^1$  and we have the inequality

$$(1.1) \quad \|f\|_1 \leq A(\| |u| \log^+ |u| \|_1 + 1),$$

where  $A$  is an absolute constant. It is well-known that the condition  $u \in \mathbf{h}^1(\mathbb{D})$  does not imply that  $f \in \mathbf{h}^1(\mathbb{D})$ , so (1.1) is in some sense the best possible inequality [2, p. 60]. On the other hand for  $p > 1$ ,  $u \in \mathbf{h}^p(\mathbb{D})$  imply that  $f \in \mathbf{h}^p(\mathbb{D})$  and moreover there is a constant  $c_p$  so that  $\|f\|_p \leq c_p \|u\|_p$  whenever  $\text{Im} f(0) = 0$ . This is a well-known M. Riesz theorem. For some recent extensions of M. Riesz theorem we refer to the papers [6, 10, 11, 12, 1, 7, 8, 5].

The sharpness of constant  $A$  in Zygmund theorem as well as its connection with M. Riesz theorem is discussed by Pichorides in [14].

This paper aims to extend Zygmund theorem to the class of quasiregular harmonic mappings in the unit disk and in the unit ball.

**1.3. Quasiregular and harmonic mappings.** A continuous and nonconstant mapping  $f : G \rightarrow \mathbb{R}^n$ ,  $n \geq 2$ , in the local Sobolev space  $W_{loc}^{1,n}(G, \mathbb{R}^n)$  is  $K$ -quasiregular,  $K \geq 1$ , if

$$|Df(x)| \leq K \ell(f'(x))$$

for almost every  $x \in G$ , where  $G$  is an open subset of  $\mathbb{R}^n$ . Here  $Df(x)$  is the formal differential matrix and

$$|Df(x)| = \sup_{|h|=1} |f'(x)h|, \quad \ell(Df(x)) = \inf_{|h|=1} |f'(x)h|.$$

A smooth mapping  $w : G \rightarrow \mathbb{R}^n$  is called harmonic if it satisfies the Laplace equation  $\Delta w = 0$ . The solution of the equation  $\Delta w = g$  (in the sense of distributions see [4]) in the ball  $B_R = R \cdot \mathbb{B}$ , satisfying the boundary condition  $w|_{S_R} = f \in L^1(S_R)$ , where  $S_R = R \cdot \mathbb{S}$  is given by

$$(1.2) \quad w(x) = \int_{S_R} P(x, \eta) f(\eta) d\sigma(\eta) - \int_{B_R} G(x, y) g(y) dV(y), \quad |x| < 1.$$

Here

$$(1.3) \quad P(x, \eta) = \frac{R^2 - |x|^2}{R|x - \eta|^n}$$

is the Poisson kernel and  $d\sigma$  is the surface  $n - 1$  dimensional measure of the Euclidean sphere which satisfies the condition:  $\int_{\mathbb{S}} P(x, \eta) d\sigma(\eta) \equiv 1$ . The first integral in (1.2) is called the Poisson integral and is usually denoted by  $u(x) = P[f](x)$ . Then  $u$  is a harmonic mapping. The function

$$(1.4) \quad G(x, y) = \begin{cases} \frac{1}{2\pi} \log \frac{R|x-y|}{|R^2-xy|}, & \text{for } n = 2; \\ c_n \left( \frac{1}{|x-y|^{n-2}} - \frac{1}{(R^2+|x|^2|y|^2/R^2-2(x,y))^{(n-2)/2}} \right), & \text{for } n \geq 3, \end{cases}$$

where

$$(1.5) \quad c_n = \frac{1}{(n-2)\omega_{n-1}}$$

and  $\omega_{n-1}$  is the measure of  $\mathbb{S}$ , is the Green function of the ball  $B_R$ .

If  $n = 2$ , then we use simultaneously the notation of  $k$ -quasiregular mappings for  $k = (K - 1)/(K + 1)$ . If  $f : \mathbb{D} \rightarrow \mathbb{C}$  is harmonic mapping, then there exist two holomorphic functions  $g$  and  $h$  ( $h(0) = 0$ ), in the unit disk, so that  $f(z) = g(z) + \overline{h(z)}$ . Then  $f$  is sense-preserving and  $k$ -quasiregular, if and only if  $|h'(z)| \leq k|g'(z)|$ ,  $z \in \mathbb{D}$ .

**1.4. Main results.** The following three theorems are main results.

**Theorem 1.1.** *Let  $f = g + \bar{h}$  be a  $K$ -quasiregular mapping defined in the unit disk  $\mathbb{D}$ . Assume further that  $u = \operatorname{Re} f \in \mathbf{h} \log^+ \mathbf{h}$ . Then  $f \in \mathbf{h}^1$  and we have the inequality*

$$(1.6) \quad \int_0^{2\pi} |f(re^{it})| \frac{dt}{2\pi} \leq C(K) \left( 1 + \int_0^{2\pi} |u(re^{it})| \log^+ |u(re^{it})| \frac{|dt|}{2\pi} \right)$$

for every  $r \in (0, 1)$  and in particular

$$(1.7) \quad \|f\|_1 \leq C(K) (\|u \log^+ u\|_1 + 1).$$

For a quasiregular mapping with a positive real part, we have the following asymptotically sharp result.

**Theorem 1.2.** *Assume that  $f = u + iv$  is a harmonic  $K$ -quasiregular function defined in the unit disk  $\mathbb{D}$  so that  $v(0) = 0$  and assume further that  $u(z) > 0$  for  $z \in \mathbb{D}$ . If  $u \in \mathbf{h} \log^+ \mathbf{h}$ , then  $f \in \mathbf{h}^1$  and we have the inequality*

$$(1.8) \quad \int_0^{2\pi} |f(re^{it})| \frac{dt}{2\pi} \leq K^2 \left( e^{-1+K^{-2}} + \int_0^{2\pi} u(re^{it}) \log u(re^{it}) \frac{|dt|}{2\pi} \right)$$

for every  $r \in (0, 1)$  and in particular

$$(1.9) \quad \|f\|_1 \leq K^2 \left( e^{-1+K^{-2}} + \int_{\mathbb{T}} u \log u \frac{|dz|}{2\pi} \right)$$

and

$$\|v\|_1 \leq \|f\|_1 \leq K^2 (\|u \log^+ u\|_1 + 1).$$

The constants are asymptotically sharp when  $K \rightarrow 1$ .

A straightforward consequence of Theorem 1.2 is the following corollary which is interesting in its own right.

**Corollary 1.3.** *If  $f = u + iv$  is an analytic function of the unit disk into the vertical strip  $(0, 1) \times \mathbb{R}$  so that  $v(0) = 0$ , then*

$$\|f\|_{H^1} < 1.$$

The constant 1 is sharp as can be seen by the following example

$$f(z) = \frac{n}{(n+1)} + \frac{z}{n+1},$$

when  $n \rightarrow \infty$ .

For higher-dimensional settings we prove

**Theorem 1.4.** *Assume that  $f = (f_1, \dots, f_n) : \mathbb{B} \rightarrow \mathbb{R}^n$  is a harmonic  $K$ -quasiregular function defined in the unit ball  $\mathbb{B}$  so that  $u(x) = f_1(x) > 0$  for  $z \in \mathbb{B}$ . If  $u \in \mathbf{h} \log^+ \mathbf{h}$ , then  $f \in \mathbf{h}^1$  and we have the inequality*

$$(1.10) \quad \|f\|_1 - |f(0)| \leq K^2(n-1) \left( \int_{\mathbb{S}} (u(\zeta) \log u(\zeta) - u(0) \log u(0)) d\sigma(\zeta) \right)$$

and if  $|f(0)| = u(0)$ , then

$$(1.11) \quad \|f\|_1 \leq K^2(n-1)(e^{-1+K^{-2}/(n-1)} + \|u \log^+ u\|_1).$$

The constant  $K^2(n-1)$  in (1.10) is asymptotically sharp as  $K \rightarrow 1$ .

## 2. PROOF OF MAIN RESULTS

*Proof of Theorem 1.1.* We lose no generality in assuming that  $r = 1$ . Assume first that  $f(0) = 0$  and thus  $h(0) = g(0) = 0$ . Note first that if  $H$  is an analytic function defined in the unit disk with  $H(0) = 0$ , then we have the double inequality

$$(2.1) \quad \|H\|_p \leq c_1(p) \|G[H]\|_p \leq c_2(p) \|H\|_p,$$

where

$$(2.2) \quad G[H](z) = \left( \int_0^1 |H'(\rho z)|^2 (1-\rho) d\rho \right)^{1/2}.$$

This holds due to Calderon type theorem of Pavlović [13, Theorem 10.6]. We apply this inequality for  $p = 1$  and  $H \in \{g, h, g+h\}$ . Let  $u = \operatorname{Re}(g+h)$ , then  $\tilde{u} = \operatorname{Im}(g+h) \in H^1$  and from the classical Zygmund theorem [2, p. 57], we have

$$\|\tilde{u}\|_{H^1} \leq A \| |u| \log^+ |u| \|_{L^1} + B,$$

where

$$A = 1, \quad B = 6\pi e.$$

Since  $|u| \leq 1 + |u| \log^+ |u|$ , we get the following

$$\|g+h\|_{H^1} \leq \|u\|_1 + \|\tilde{u}\|_1 \leq 2 \| |u| \log^+ |u| \|_{L^1} + 6\pi e + 1.$$

Since  $f$  is quasiregular we have

$$(2.3) \quad |g'(z)| \leq \frac{|g'(z) + h'(z)|}{1-k}, \quad |h'(z)| \leq \frac{k|g'(z) + h'(z)|}{1-k},$$

where  $k = (K-1)/(K+1)$ . Denote the constants in (2.2) by  $c_1 = c_1(1)$  and  $c_2 = c_2(1)$ . Then by (2.1) and (2.3), we have

$$\|g\|_{H^1} \leq c_1 \|G(g)\|_1 \leq \frac{c_1}{1-k} \|G(g+h)\|_1 \leq \frac{c_1 \cdot c_2}{1-k} \|g+h\|_{H^1}$$

and

$$\|h\|_{H^1} \leq c_1 \|G(h)\|_1 \leq \frac{c_1 k}{1-k} \|G(g+h)\|_1 \leq \frac{c_1 \cdot c_2 k}{1-k} \|g+h\|_{H^1}.$$

Thus

$$\|f\|_1 \leq CK (\|g\|_{H^1} + \|h\|_{H^1}) \leq CK (2 \| |u| \log^+ |u| \|_{L^1} + 6\pi e + 1),$$

where  $C = c_1 \cdot c_2$ .

The case  $f(0) \neq 0$  follows from the following inequality  $|f(0)| \leq \|f\|_1$ . The last inequality follows from the fact that  $|f|$  is a subharmonic function.

Thus for  $C(K) = 2(6\pi e + 1)c_1c_2K$  we proved (1.6) and (1.7). □

To prove Theorem 1.2, we need the following lemma

**Lemma 2.1.** *If  $f = u + iv = g + \bar{h}$  is quasiregular, then*

$$\Delta|f| \leq K^2 \Delta(u \log u)$$

*provided that  $u$  is positive.*

*Proof of Lemma 2.1.* Since  $u = \operatorname{Re}(g + h)$ , it follows that  $|\nabla u| = |g' + h'|$ . Further

$$|f| = ((g + \bar{h})(\bar{g} + h))^{1/2}.$$

Thus

$$|f|_z = \frac{1}{2}((g + \bar{h})(\bar{g} + h))^{1/2-1}(g'(\bar{g} + h) + (g + \bar{h})h')$$

and

$$\begin{aligned} |f|_{z\bar{z}} &= -\frac{1}{4}((g + \bar{h})(\bar{g} + h))^{1/2-2}|g'(\bar{g} + h) + (g + \bar{h})h'|^2 \\ &\quad + \frac{1}{2}((g + \bar{h})(\bar{g} + h))^{1/2-1}(|g'|^2 + |h'|^2). \end{aligned}$$

Then, after some simple manipulations we get

$$\Delta|f| = 4|f|_{z\bar{z}} = \frac{|g' - \frac{f}{f}h'|^2}{|f|}.$$

Further

$$\Delta(u \log u) = \frac{|\nabla u|^2}{u} = \frac{|g' + h'|^2}{u}.$$

Then

$$\frac{\Delta|f|}{\Delta(u \log u)} = \frac{|g' - \frac{f}{f}h'|^2}{|g' + h'|^2} \frac{u}{|f|} \leq \frac{(|g'| + |h'|)^2}{(|g'| - |h'|)^2} \frac{u}{|f|} \leq K^2.$$

□

*Proof of Theorem 1.2.* By Proposition 1.2, we have

$$|f(0)| = \int_{\mathbb{T}} |f(rz)| \frac{|dz|}{2\pi} - \frac{1}{2\pi} \int_{|z| < r} \Delta|f(z)| \log \frac{r}{|z|} dx dy.$$

Let  $\mathbb{D}_r = \{z : |z| < r\}$ . Now, because  $|f(0)| = u(0)$ , and using again Proposition 1.2,

$$\begin{aligned} \int_{\mathbb{T}} |f(rz)| \frac{|dz|}{2\pi} &= u(0) + \frac{1}{2\pi} \int_{\mathbb{D}_r} \Delta|f(z)| \log \frac{r}{|z|} dx dy \\ &\leq u(0) + K^2 \frac{1}{2\pi} \int_{\mathbb{D}_r} \Delta(u(z) \log u(z)) \log \frac{r}{|z|} dx dy \\ &= u(0) + K^2 \left( -u(0) \log u(0) + \frac{1}{2\pi} \int_{\mathbb{T}} u(rz) \log u(rz) |dz| \right) \\ &= \Phi(u(0)) + K^2 \frac{1}{2\pi} \int_{\mathbb{T}} u(rz) \log u(rz) |dz|, \end{aligned}$$

where

$$\Phi(\xi) := \xi - K^2 \xi \log \xi.$$

Then  $\Phi$  attains its maximum for  $\xi = u(0) = e^{-1+1/K^2}$ , which is equal to  $K^2 e^{1-1/K^2}$ . This implies (1.8).  $\square$

*Proof of Theorem 1.4.* For any fixed  $y \neq 0$  we assume that  $\{U_1, \dots, U_{n-1}, N\}$  is a system of mutually orthogonal vectors of the unit norm, where  $N = \frac{y}{|y|}$  and the vectors  $\{U_1, \dots, U_{n-1}\}$  are arbitrarily chosen. For a differentiable mapping  $h$  defined in domain  $\mathbb{B}$  define  $h_N(x) = Dh(x)N$  and set  $h_{U_i}(x) = Dh(x)U_i$  for  $i = 1, \dots, n-1$ . Since the Hilbert-Schmidt norm of  $Dh$  is independent of basis, we have

$$(2.4) \quad \|Dh\|^2 = |h_N|^2 + |h_{U_1}|^2 + |h_{U_2}|^2 + \dots + |h_{U_{n-1}}|^2.$$

Let  $S(x) = \frac{f(x)}{|f(x)|}$  where  $f(x) \neq 0$  because  $f_1(x) > 0$ . Fix  $x$  and let  $y = f(x)$  and  $N = y/|y|$ . Then

$$(2.5) \quad \begin{aligned} \|DS(x)\|^2 &= |S_N(x)|^2 + |S_{U_1}(x)|^2 + |S_{U_2}(x)|^2 + \dots + |S_{U_{n-1}}(x)|^2 \\ &= |S_{U_1}(x)|^2 + |S_{U_2}(x)|^2 + \dots + |S_{U_{n-1}}(x)|^2, \end{aligned}$$

because  $S_N(x) = \langle DS(x), S(x) \rangle = \nabla|S(x)| = \nabla(1) = 0$ .

Since

$$S_{U_j} = |f|^{-1} f_{U_j} - |f|^{-3} \langle f, f_{U_j} \rangle f,$$

it follows that

$$|S_{U_j}| = \frac{|f_{U_j}|^2}{|f|^2} - \frac{\langle f, f_{U_j} \rangle^2}{|f|^4} \leq \frac{|f_{U_j}|^2}{|f|^2}.$$

Further by straightforward computation (see e.g. [9, Lemma 1.4]),

$$\Delta|f(x)| = |f(x)| \|DS(x)\|^2$$

and

$$\Delta(f_1 \log f_1) = \frac{|\nabla f_1|^2}{f_1}.$$

Since  $F$  is  $K$ -quasiregular, we have

$$|f_{U_j}| \leq K |\nabla f_1|,$$

for every  $j$ . Thus

$$(2.6) \quad \frac{\Delta|f|}{\Delta(f_1 \log f_1)} \leq K^2(n-1).$$

Then by (1.2) we have

$$\|f\|_1 = |f(0)| + c_n \int_{\mathbb{B}} \Delta|f(x)| (|x|^{2-n} - 1) dV(x),$$

and for  $u = f_1$

$$\int_{\mathbb{S}} u \log u d\sigma(\zeta) = u(0) \log u(0) + c_n \int_{\mathbb{B}} \Delta(u \log u) (|x|^{2-n} - 1) dV(x).$$

Further, we have

$$\|f\|_1 - |f(0)| \leq K^2(n-1) \left( \int_{\mathbb{S}} (u(\zeta) \log u(\zeta) - u(0) \log u(0)) d\sigma(\zeta) \right).$$

Then as in proof of Theorem 1.2, if  $|f(0)| = u(0)$ , we have

$$\|f\|_1 \leq K^2(n-1)(e^{-1+K^{-2}/(n-1)} + \|u \log^+ u\|_1).$$

To prove sharpness, demonstrate it firstly for  $n = 3$  and let

$$f(x) = m^2 e_1 + mx.$$

Then in spherical coordinates,

$$f(x) = (m^2 + m \cos t, m \sin s \sin t, m \cos s \sin t), \quad t \in [0, \pi], \quad s \in [0, 2\pi],$$

and then

$$\|f\|_1 - |f(0)| = -m^2 + \frac{1}{4\pi} \int_0^{2\pi} \int_0^\pi m \sin t \sqrt{1 + m^2 + 2m \cos t} dt ds = \frac{1}{3}.$$

Moreover for  $u = m^2 + m \cos t$

$$2 \left( \int_{\mathbb{S}} (u(\zeta) \log u(\zeta) - u(0) \log u(0)) d\sigma(\zeta) \right) = \phi(m)$$

where

$$\phi(m) = \frac{m}{2} \left( (1+m)^2 \log(m+1) - (2m + (m-1)^2 \log(m-1) + 4m \log m) \right).$$

Moreover  $\lim_{m \rightarrow \infty} \phi(m) = \frac{1}{3}$ .

For general  $n$  we can take  $f(x) = ax + (1, 0, \dots, 0)$ ,  $0 < a < 1$ , and define

$$X(f) = \|f\|_1 - |f(0)|,$$

and

$$Y(f) = \left( \int_{\mathbb{S}} (u(\zeta) \log u(\zeta) - u(0) \log u(0)) d\sigma(\zeta) \right).$$

Then for  $C_n = \frac{\Gamma[\frac{n}{2}]}{\sqrt{\pi} \Gamma[\frac{1}{2}(-1+n)]}$  we have

$$X(f) = C_n \int_0^\pi \sin^{n-2} t \left( \sqrt{1 + a^2 + 2a \cos t} - 1 \right) dt$$

and

$$Y(f) = C_n \int_0^\pi \sin^{n-2} t (1 + a \cos t) \log(1 + a \cos t) dt.$$

Since

$$\left( \sqrt{1 + a^2 + 2a \cos t} - 1 \right) = a \cos(t) + \left( \frac{1}{2} - \frac{\cos^2 t}{2} \right) a^2 + o(a^2)$$

and

$$(1 + a \cos t) \log(1 + a \cos t) = a \cos t + \frac{a^2}{2} \cos^2 t + o(a^2),$$

we get

$$X(f) = C_n \frac{a^2 \sqrt{\pi} \Gamma[\frac{1+n}{2}]}{2\Gamma[1 + \frac{n}{2}]} + o(a^2) = \frac{a^2(n-1)}{2n} + o(a^2)$$

$$Y(f) = C_n \frac{a^2 \sqrt{\pi} \Gamma[\frac{1}{2}(-1+n)]}{4\Gamma[1 + \frac{n}{2}]} + o(a^2) = \frac{a^2}{2n} + o(a^2).$$

Then we obtain

$$\lim_{a \rightarrow 0} \frac{X(f)}{Y(f)} = n - 1.$$

Hence the constant  $n - 1$  is the best possible for the case  $K = 1$ .  $\square$

**Funding.** This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

**Data availability.** Data sharing is not applicable to this article since no data sets were generated or analyzed

#### ETHICS DECLARATIONS

The author declares that he has not conflict of interest.

#### REFERENCES

- [1] S. L. Chen and M. Huang, Riesz type theorems for  $\kappa$ -pluriharmonic mappings, invariant harmonic quasiregular mappings and harmonic quasiregular mappings, arXiv:2310.15452.
- [2] P. Duren, *Theory of  $H^p$  spaces*. Pure and Applied Mathematics (Academic Press) 38. New York and London: Academic Press. XII, 258 p. (1970).
- [3] D. Gilbarg and N. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Vol. 224, 2nd Edition, Springer 1983.
- [4] L. Hörmander, *Notions of convexity*, Progress in Mathematics, Birkhäuser Boston Inc., Boston, MA, 1994.
- [5] B. Hollenbeck, N. J. Kalton, I. E. Verbitsky, *Best constants for some operators associated with the Fourier and Hilbert transforms*. *Studia Math.* 157 (2003), no. 3, 237–278.
- [6] D. Kalaj, On Riesz type inequalities for harmonic mappings on the unit disk, *Trans. Am. Math. Soc.* **372** (2019), 4031–4051.
- [7] D. Kalaj, Riesz and Kolmogorov inequality for harmonic quasiregular mappings, arXiv:2310.12643.
- [8] D. Kalaj, On M. Riesz conjugate function theorem for harmonic functions, arXiv:2310.00464
- [9] D. Kalaj, On the univalent solution of PDE  $\Delta u = f$  between spherical annuli, *J. Math. Anal. Appl.*, **327**, No. 1, (2007) 1–11.
- [10] J. Liu and J. F. Zhu, Riesz conjugate functions theorem for harmonic, quasiconformal mappings, *Adv. Math.* **434** (2023), 109321.
- [11] P. Melentijević, Hollenbeck-Verbitsky conjecture on best constant inequalities for analytic and co-analytic projections, *Math. Ann.* (2023). <https://doi.org/10.1007/s00208-023-02639-1>.
- [12] P. Melentijević, M. Marković, Best Constants in Inequalities Involving Analytic and Co-Analytic Projections and Riesz's Theorem in Various Function Spaces, *Potential Anal.* (2022). <https://doi.org/10.1007/s11118-022-10021-0>.
- [13] M. Pavlović, *Function classes on the unit disc. An introduction*. 2nd revised and extended edition, De Gruyter Studies in Mathematics 52. Berlin: De Gruyter (ISBN 978-3-11-062844-9/hbk; 978-3-11-063085-5/ebook). xv, 553 p. (2019).
- [14] S. K. Pichorides, On the best values of the constants in the theorems of M. Riesz, Zygmund and Kolmogorov, *Studia Math.* **44** (1972), 165–179.
- [15] A. Zygmund, *Trigonometric series*. Volumes I and II combined. With a foreword by Robert Fefferman. 3rd ed. Cambridge Mathematical Library. Cambridge: Cambridge University Press (ISBN 0-521-89053-5/pbk). xiii, 364 p. (2002).

UNIVERSITY OF MONTENEGRO, FACULTY OF NATURAL SCIENCES AND MATHEMATICS, CETINJSKI PUT B.B.  
81000 PODGORICA, MONTENEGRO  
*Email address:* davidk@ucg.ac.me