Distributed Framework Construction for Affine Formation Control

Huiming Li¹, Hao Chen^{1,2}, Xiangke Wang¹ Senior Member, IEEE, Zhongkui Li³, Senior Member, IEEE, Lincheng Shen¹

Abstract—In affine formation control problems, the construction of the framework with universal rigidity and affine localizability is a critical prerequisite, but it has not yet been well addressed, especially when additional agents join the formation or link/agent failures emerge. Motivated by this observation, we investigate the problem of constructing affine formation frameworks in three scenarios, including vertex addition, edge deletion and vertex deletion. Our approach starts from the original affine formation and uses geometric methods to locally adjust the structure of the weighted graph to describe the topology, so that the modified framework maintains the universal rigidity and affine localizability. Notably, the developed strategies only utilize local measurements and exhibit distributed characteristics, laying the foundation for applications in multi-agent systems. To demonstrate the compatibility with affine formation control proposals, we present a case study on affine formation tracking in a multi-UAV formation, demonstrating the effectiveness of our algorithms in constructing eligible frameworks in aforementioned scenarios. Moreover, a comparative simulations is also conducted to highlight the low time complexity of our distributed algorithm relative to the centralized optimization-based method.

Index Terms—Affine formations, framework construction, rigidity maintenance, robot swarm

I. INTRODUCTION

FFINE formation control problem has recently attracted an increasing attention of scholars, which allows configuration transformations including translation, rotation, scaling, shearing and their combinations [1]–[3]. Various studies have been conducted to investigate the collaborative control scheme for affine formations, such as considering different system dynamics, e.g., linear dynamics [2], [4], unicycle models [5] and Euler-Lagrange models [6], or considering different operating modes, e.g., switching topologies [7], event-triggered

*This work was supported in part by the National Natural Science Foundation of China under Grant 62303483, U23B2032, U2241214, and 62376280, in part by the Research Project of the National University of Defense Technology under Grant ZK21-05. The corresponding author is Xiangke Wang.

¹College of Intelligence Science and Technology, National University of Defense Technology, Changsha 410073, China.

² Laboratory of Science and Technology on Integrated Logistics Support, National University of Defense Technology, Changsha 410073, China.

³ The State Key Laboratory for Turbulence and Complex Systems, Department of Mechanics and Engineering Science, College of Engineering, Peking University, Beijing 100871, China.

Email: huiminglhm@163.com, chenhao09@nudt.edu.cn, xkwang@nudt.edu.cn, zhongkli@pku.edu.cn, lcshen@nudt.edu.cn.

conditions [8], etc. Undoubtedly, the implementation of the above distributed cooperative control schemes heavily relies on the *framework* of affine formations, which is described by assigning coordinates in Euclidean space to the topology represented by a weighted graph [1]. A determined framework fixes the direction of information flow within a multiagent system, which directly influences the convergence of distributed collaborative control schemes. Moreover, in affine formations, the weighted graph used to describe the topology quantifies the strength of interaction among agents, which also affects the convergence speed of the control proposals. Most related studies used pre-designed frameworks, which generally suffers from inevitable changes during the mission execution process [9]. To address the potential various scenarios, the construction of affine formation frameworks is of great value and practical significance.

From a mathematical perspective, the problem of designing frameworks for multi-agent systems is usually transformed into the problem of constructing appropriate graph-related matrices (e.g., the bearing-based Laplacian matrix [10] and the stress matrix [11]), which not only reflect the underlying communication among agents but also affect the control gain of the close-loop system. Based on the theoretical analysis in [1], [2], there are two critical properties, *universal rigidity* and *affine localizability*, that need to be carefully considered to construct affine formation frameworks due to their direct impact on stabilizability. Up to now, some efforts have been put in to construct affine formation frameworks.

Starting with the rigidity required by an affine formation framework, a classic method, the Henneberg Construction (HC), is widely applied in many papers. HC was proposed in [12] to grow the minimally rigid graphs based on the graph structures and the properties of related matrices. Subsequently, lots of studies followed this idea to analyze how the graph rigidity changes after HCs, and how to obtain graphs with the desired characteristics through HCs. Connelly proved the universal rigidity is maintained after HCs [13], and a series of algorithms are proposed for building and reconstructing rigid graphs, not only the classical distance rigidity [14], but also the bearing rigidity [10], [15], [16]. In [17] and [18], the issue of growing rigid tensegrity frameworks were addressed with the aid of HCs. Nevertheless, the affine localizability is rarely considered simultaneously in the related work, which requires the proposed framework construction algorithms to consider more constraints.

To meet specific constraints and address the framework

construction issue, a normal idea is to establish an optimization problem to obtain a legal topology in which the minimum energy is spent among all the possible networks. In some existing works, the authors studied framework construction problem by using optimization methods to improve resilience [19], [20] and achieve stable coordination [21], [22]. In [11], Xiao et al. formulated the affine formation framework construction problem into a centralized mixed integer semidefinite programming (MISDP) problem to obtain the proper stress matrix for affine formation maneuvers while maximizing the convergence speed and minimizing the communication cost. The ready-to-use solver was applied to construct suitable frameworks, which had a significant impact on the computational time of the NP-hard mixed integer programming problem. However, all the researches mentioned above rely on global information (such as global positions required by [11]) of multi-agent systems and solve the optimization problem in a centralized way. On the one hand, limited by practical situations, global measurements and central agents are not guaranteed in some real robot systems, for example, only bearing measurements are available in [23] and the distributed architecture is necessary for large-scale swarms in [24]. On the other hand, the centralized optimization algorithm requires powerful computing resources, which adds difficulty to the update of the framework in real-time conditions. In [11], the computational cost of growing affine formation frameworks increased rapidly with the increase of swarm size. Accordingly, beyond existing research, it is a meaningful and challenging issue to construct affine formation frameworks using local measurements in a distributed way.

Motivated by these observations, we investigate the distributed strategies to construct affine formation frameworks to provide fundamental support for formation control strategies. Different from centralized approaches, we design the framework construction strategies based on HCs by using local measurements in three specific scenarios. Theoretical analysis and simulations are provided to demonstrate that our algorithms preserve not only universal rigidity but also affine localizability to stabilize affine formations. The main contributions of this paper can be summarized as follows.

- (1) We propose distributed strategies to construct affine formation frameworks to address various application scenarios, including vertex addition, edge deletion and vertex deletion. Our approaches do not rely on global measurements, making it well-suited for extensive application in multi-agent systems due to their distributed nature.
- (2) We provide comprehensive theoretical analysis on the mathematical conditions of our proposed framework construction strategies to guarantee the characteristics of affine formations, namely universal rigidity and affine localizability. The presented simulations demonstrate the low time complexity and the widespread practicality in multi-agent systems.

The remainder of this paper is organized as follows. We provide foundational definitions and lemmas in Section II. In Section III, we address the key problem and propose distributed algorithms on affine formation constructions. Then, we provide some discussions about our strategies in Section IV. The effectiveness is validated by simulations in Section V. The conclusions are summarized in Section VI.

Notation : Throughout the paper, the partial order $A \succ 0$ $(A \succeq 0)$ means that the matrix A is positive definite (positive semi-definite).

II. PROBLEM FORMULATION

In this section, we introduce some basic definitions and supportive lemmas, and then establish the framework construction problems to be studied.

A. Affine Formation

We model the interaction topology of a *n*-agent system as a simple undirected graph $\mathcal{G} = (\mathcal{V}, \mathcal{E})$, where $\mathcal{V} = \{v_1 \cdots v_n\}$ is the set of vertices, $\mathcal{E} \subseteq \mathcal{V} \times \mathcal{V}$ is the set of edges, and $|\mathcal{E}| = m$. An edge e_{ij} exists if and only if there is a bidirectional interaction between vertex v_i and v_j . Hence, the communication neighbor set of vertex v_i is denoted by $\mathcal{N}_i := \{v_j \in \mathcal{V} \mid e_{ij} \in \mathcal{E}\}.$ Let $p_i \in \mathbb{R}^d$ be the position of vertex v_i , and $p := [p_i^T \cdots p_n^T]^T$ describes the *configuration* of the multi-agent system. We define a framework as a graph associated with p, i.e., (\mathcal{G}, p) . The set of points $\{p_i\}_{i=1}^n$ are called affinely dependent if there exist scalars $\{a_i\}_{i=1}^{n}$ that are not all zero such that $\sum_{i=1}^{n} a_i p_i = 0$ and $\sum_{i=1}^{n} a_i = 0$, and affinely independent otherwise. A configuration p (or a framework $(\mathcal{G}, \boldsymbol{p})$ in \mathbb{R}^d is said to be in general position if no subset of the points $\{p_i\}_{i=1}^n$ of cardinality d+1 is affinely dependent. For example, a set of points in the plane are in general position if no three of them lie on a straight line. Considering the limited perception distance of agents, we define the perceived neighbors of agent i as $\mathcal{N}_i^p := \{v_j \in$ $\mathcal{V} \mid \|\boldsymbol{p}_i - \boldsymbol{p}_j\| \leq d_{per}$, where d_{per} is the perception distance. We assume that all perceptible agents can communicate, i.e., $\mathcal{N}_i \subseteq \mathcal{N}_i^p$.

We assign a *stress*, i.e., a set of scalars $\{\varpi_{ij}\}_{(i,j)\in\mathcal{E}}$, for all edges in the undirected graph. That is, a weight ϖ_{ij} is assigned for the edge e_{ij} , and $\varpi_{ij} = \varpi_{ji}$. An *equilibrium stress* is established if and only if $\sum_{v_j\in\mathcal{N}_i} \varpi_{ij} (p_j - p_i) = \mathbf{0}$ holds for any vertex v_i in the framework (\mathcal{G}, p) . Its matrix form can be written as $(\mathbf{\Omega} \otimes \mathbf{I}_d) \mathbf{p} = \mathbf{0}$, where the sign \otimes represents the Kronecker product and $\mathbf{\Omega} \in \mathbb{R}^{n \times n}$ is termed the *stress matrix*, defined as

$$[\mathbf{\Omega}]_{ij} = \begin{cases} -\varpi_{ij} & \text{if } i \neq j \text{ and } v_j \in \mathcal{N}_i \\ 0 & \text{if } i \neq j \text{ and } v_j \notin \mathcal{N}_i \\ \sum_{k \in \mathcal{N}_i} \varpi_{ik} & \text{if } i = j \end{cases}$$

The matrix I_d denotes the identity matrix in \mathbb{R}^d . The following lemma reveals the connection between the stress matrix and the rigidity.

Lemma 1 ([25]): Let (\mathcal{G}, p) be an d-dimensional framework on n vertices in \mathbb{R}^d , for some $d \leq n-2$. Then (\mathcal{G}, p) is universally rigid if the following two conditions hold.

- (1) $(\mathcal{G}, \boldsymbol{p})$ has a positive semi-definite (PSD) stress matrix Ω with rank n d 1.
- (2) For each vertex v_i , the set $\{p_i\} \cup \{p_j : v_j \in \mathcal{N}_j\}$ is in general position in \mathbb{R}^d .

Recently, the stress matrix has been widely applied to stabilize affine formations of multiple vehicles [1]–[3], [5]–[8], [26], [27]. Different from rigid formations, affine formations allow the rotation, translation, scaling, shearing and combinations of them, which have a great advantage in enhancing formation maneuverability. In the affine formation control scheme, the leader-follower structure is adopted naturally. Let n_l vertices $\mathcal{V}_l = \{v_1, \cdots, v_{n_l}\}$ be the leaders, and the remaining $n_f = n - n_l$ vertices in the framework be the followers, denoted by $\mathcal{V}_f = \mathcal{V} \setminus \mathcal{V}_l$. Define a constant configuration in general position as nominal configuration $\boldsymbol{r} = \left[\boldsymbol{r}_{l}^{T}, \boldsymbol{r}_{f}^{T}\right]^{T}$, where \boldsymbol{r}_{l} represents the configuration of n_{l} leaders and \vec{r}_f describes the configuration of the followers. Accordingly, the *nominal framework* is denoted by $(\mathcal{G}, \mathbf{r})$, and the time-varying target configuration is expressed as $p^*(t) = [I_n \otimes A(t)] r + \mathbf{1}_n \otimes b(t)$, where the invertible matrix $A(t) \in \mathbb{R}^{d \times d}$ and translation vector $b(t) \in \mathbb{R}^{d}$ are continuous on t. In this paper, we assume that the nominal framework $(\mathcal{G}, \mathbf{r})$ satisfy $n \geq d+2$, where n_l agents that affinely span (refer to [2]) \mathbb{R}^d are selected as leaders. Since the affine span of \mathbb{R}^d requires at least d+1 points, $n_l = d+1$ is assumed to be held, following the analysis in [2], [5].

Lemma 2 ([2]): Assume that the nominal formation $(\mathcal{G}, \mathbf{r})$ has a PSD stress matrix satisfying rank $(\mathbf{\Omega}) = n - d - 1$. The following conditions are equivalent to each other

(i) $\{\boldsymbol{r}_i\}_{i=1}^n$ affinely span \mathbb{R}^d ; (ii) null($\boldsymbol{\Omega}$) = col $\left(\begin{bmatrix} \boldsymbol{r}_1 & \cdots & \boldsymbol{r}_n \\ 1 & \cdots & 1 \end{bmatrix}^T \right)$.

Let the stress matrix Ω associated with $(\mathcal{G}, \mathbf{r})$ be $\Omega = \begin{bmatrix} \Omega_{ll} & \Omega_{lf} \\ \Omega_{fl} & \Omega_{ff} \end{bmatrix}$, where $\Omega_{ll} \in \mathbb{R}^{n_l \times n_l}$, $\Omega_{lf} \in \mathbb{R}^{n_l \times n_f}$, $\Omega_{fl} \in \mathbb{R}^{n_f \times n_l}$, and $\Omega_{ff} \in \mathbb{R}^{n_f \times n_f}$. The definition of affine localizability is proposed in [2].

Definition 1 (Affine localizability [2]): The nominal formation $(\mathcal{G}, \mathbf{r})$ is affinely localizable by the leaders if for any $\mathbf{p} = \begin{bmatrix} \mathbf{p}_l^T, \mathbf{p}_f^T \end{bmatrix} \in \mathcal{A}(\mathbf{r}), \ \mathbf{p}_f$ can be uniquely determined by \mathbf{p}_l , where $\mathcal{A}(\mathbf{r}) = \{\mathbf{p}: \mathbf{p} = (\mathbf{I}_n \otimes \mathbf{A})\mathbf{r} + \mathbf{1}_n \otimes \mathbf{b}, \mathbf{A} \in \mathbb{R}^{d \times d}, \mathbf{b} \in \mathbb{R}^d \}.$

Lemma 3 ([2]): If the nominal formation $(\mathcal{G}, \mathbf{r})$ is universally rigid and $\{\mathbf{r}_i\}_{i=1}^n$ affinely span \mathbb{R}^d , $(\mathcal{G}, \mathbf{r})$ is affinely localizable if and only if $\mathbf{\Omega}_{ff}$ is nonsingular.

In the affine formation control scheme, a general dynamic of the closed-loop system is described by $\dot{\delta}_f = -\Omega_{ff} \delta_f$, where δ_f represents the followers' formation tracking error. Obviously, the properties of Ω_{ff} greatly affect the convergence and stability of the system. Combined with Lemma 1 and 3, it is easily deduced that Ω_{ff} is positive definite, if $(\mathcal{G}, \mathbf{r})$ is universally rigid and affinely localizable. Accordingly, to achieve desired affine formations, the following conditions are necessary for the nominal formation $(\mathcal{G}, \mathbf{r})$.

- (i) [Universal Rigidity] The stress matrix Ω of (G, r) is PSD and rank(Ω) = n d 1;
- (ii) [Affine Localizability] The nominal formation $(\mathcal{G}, \mathbf{r})$ is affinely localized by the selected leaders.

Specifically, the first condition indicates that the framework $(\mathcal{G}, \mathbf{r})$ is universally rigid [1], [13]. Combined with Lemma 2,

Fig. 1. An unmanned aerial vehicle formation maneuvers through complicated environments. Three scenarios are presented, including vertex addition, edge deletion and vertex deletion.

the null space of Ω is obtained. The second condition implies Ω_{ff} is positive definite to stabilize the formation tracking error. Accordingly, the universal rigidity and affine localizability are two key properties to ensure the feasibility of the framework under various affine formation control schemes, which are also the focus of our subsequent theoretical analysis.

B. Problem Statement

In a robot swarm, due to the time-varying target tasks and complex environments, the configuration and topology are usually not constant. It is crucial to flexibly generate and reconstruct framework according to different situations. Take the scenario in Fig. 1 as an example, when a group of unmanned aerial vehicles (UAVs) pass through dense buildings or woods, situations including UAV joining, crashing, and communication disruptions may occur. From the perspective of graph theory, these situations can be described by the vertex addition, edge deletion and vertex deletion (corresponding to the UAV joining, communication disruption and crashing), respectively. The original topology of the swarm is changed, and even the inherent properties (such as connectivity, rigidity, etc.) are disrupted. Considering the supportive role of topology in cooperative control schemes, it is particularly crucial to timely construct the topology in various situations, which gives rise to the demand for framework construction strategies. Furthermore, considering the lack of understanding of the global states and measurements for perception-limited robots, the study of distributed algorithms is motivated in this paper.

In this paper, we investigate how to construct an affine formation framework to quickly adapt to different operating conditions. From the perspective of graph theory, the connections between vertices and the weights of edges are reshaped to guarantee the universal rigidity and affine localizability. Given an original affine formation framework (\mathcal{G}, p) with a stress matrix Ω , design distributed algorithms to guarantee the universal rigidity and affine localizability of the obtained framework after adding vertices to (\mathcal{G}, p) , or removing vertices and edges from (\mathcal{G}, p) .

Assumption 1: Suppose that the original framework $(\mathcal{G}, \mathbf{p})$ in general position is affinely localizable and universally rigid, where \mathcal{G} contains a (d+1)-lateration graph as a spanning subgraph.

A graph of n vertices is called a (d + 1)-lateration graph if there is a permutation π of the vertices, $\pi(1)$, $\pi(2)$, \cdots , $\pi(n)$,



such that (i) the first (d+1) vertices, $\pi(1), \dots, \pi(d+1)$, form a clique, and (ii) each remaining vertex $\pi(j)$, for $j = (d+2), \dots, n$, is adjacent to (d+1) vertices in the set $\{\pi(1), \pi(2) \dots \pi(j-1)\}$. Based on Assumption 1, it is deduced that (\mathcal{G}, p) admits a positive PSD stress matrix with rank (n-d-1) [28]. The problems studied in this paper are established as follows.

Problem 1 (Vertex Addition): Given an affine formation framework $(\mathcal{G}, \mathbf{p})$ and a new vertex v_u , design a strategy to link v_u with $(\mathcal{G}, \mathbf{p})$ and then rearrange the stress so that the obtained framework $(\mathcal{G}^+, \mathbf{p}^+)$ is universally rigid and affinely localizable, where $\mathcal{G}^+ = (\mathcal{V} \cup \{v_u\}, \mathcal{E}^+)$ and $\mathbf{p}^+ = [\mathbf{p}^T, \mathbf{p}_u^T]^T$.

Problem 2 (Edge Deletion): Given an affine formation framework $(\mathcal{G}, \mathbf{p})$ with an edge e_{jk} (i.e., $j, k \in \mathcal{V}, e_{jk} \in \mathcal{E}$), design a strategy to delete e_{jk} from \mathcal{E} while maintaining the universal rigidity and affine localizability of the obtained framework $(\mathcal{G}_{ed}, \mathbf{p}_{ed})$, where $e_{jk} \notin \mathcal{E}_{ed}$.

Problem 3 (Vertex Deletion): Given an affine formation framework $(\mathcal{G}, \mathbf{p})$ with a specific vertex v_u , design a strategy to delete v_u from the vertex set \mathcal{V} while maintaining the universal rigidity and affine localizability of the obtained framework $(\mathcal{G}_{vd}, \mathbf{p}_{vd})$, where $v_u \notin \mathcal{V}_{vd}$.

III. CONSTRUCTION OF AFFINE FORMATION FRAMEWORKS IN 2D

In this section, we aim to develop affine formation framework construction strategies to solve the problems proposed in Section II. We consider d = 2 in this section, and the methods are further extended to three-dimensional spaces in Section IV.

A. Vertex Addition

Given an affine formation $(\mathcal{G}, \mathbf{p})$ and a new vertex v_u , our objective here is to merge v_u to the original framework $(\mathcal{G}, \mathbf{p})$ with appropriate edges and weights, resulting in an new framework $(\mathcal{G}^+, \mathbf{p}^+)$ with universal rigidity and affine localizability. Naturally, the leaders in $(\mathcal{G}, \mathbf{p})$ can be inherited by $(\mathcal{G}^+, \mathbf{p}^+)$, i.e, $\mathcal{V}_l^+ = \mathcal{V}_l$.

Assumption 2: Given an affine formation (\mathcal{G}, p) in general position, suppose that the perception distance of the new vertex v_u is large enough to ensure $|\mathcal{N}_u^p| \geq 3$, and the set $\{p_u\} \cup \{p_j : j \in \mathcal{N}_u^p\}$ is in general position.

We can choose three vertices in \mathcal{N}_{u}^{p} , denoted by v_{i} , v_{j} and v_{k} . According to Assumption 2, we have that no three of v_{i} , v_{j} , v_{k} and v_{u} are collinear in \mathbb{R}^{2} . To generate an eligible affine formation framework, we add edges between v_{u} and all of v_{i} , v_{j} and v_{k} , the resulting framework $(\mathcal{G}^{+}, \mathbf{p}^{+})$ is shown as in Fig. 2, where $\mathcal{G}^{+} = (\mathcal{V} \cup \{v_{u}\}, \mathcal{E} \cup \{e_{iu}, e_{ju}, e_{ku}\})$ and $\mathbf{p}^{+} = [\mathbf{p}^{T}, \mathbf{p}_{u}^{T}]^{T}$. To maintain the universal rigidity and affine localibility of $(\mathcal{G}^{+}, \mathbf{p}^{+})$, the stress in \mathcal{G}^{+} need to be rearranged, as discussed below.

Reorder the elements in \mathcal{V} to place the selected three vertices v_i , v_j , v_k at the end, i.e., $\mathcal{V} \triangleq \{\cdots, v_{i-1}, v_{i+1}, \cdots, v_{j-1}, v_{j+1}, \cdots, v_{k-1}, v_{k+1}, \cdots, v_i, v_j, v_k\}$ if i < j < k. The configuration p is also reordered based



Fig. 2. An example of vertex addition. (a)The original framework (\mathcal{G}, p) in general position. (b) The obtained graph \mathcal{G}^+ , where the vertex v_u (labeled by a orange circle) and three edges are added.

on the index of \mathcal{V} . Thus, for the graph \mathcal{G} , the corresponding stress matrix Ω can be described as below.

$$\mathbf{\Omega} = \begin{bmatrix} \mathbf{\Omega}^{P_1} & \mathbf{\Omega}^{P_1} & \mathbf{\Omega}^{P_2} \\ \mathbf{\Omega}^{P_2} & \mathbf{\Omega}^{P_1} & \mathbf{\Omega}^{P_2} \\ (\mathbf{\Omega}^{P_2})^T & \mathbf{\Omega}^{P_1} & \mathbf{\Omega}^{P_2} & \mathbf{\Omega}^{P_2} \\ \mathbf{\Omega}^{P_1} & \mathbf{\Omega}^{P_2} & \mathbf{\Omega}^{P_2} \\ \mathbf{\Omega}^{P_1} & \mathbf{\Omega}^{P_2} & \mathbf{\Omega}^{P_2} \\ \mathbf{\Omega}^{P_2} & \mathbf{\Omega}^{P_2} & \mathbf{\Omega}^{P_2} \\ \mathbf{\Omega}^{P_1} & \mathbf{\Omega}^{P_2} & \mathbf{\Omega}^{P_2} \\ \mathbf{\Omega}^{P_2} & \mathbf{\Omega}^{P_2} & \mathbf{\Omega}^{P_2} \\ \mathbf{\Omega$$

where Ω is indexed based on the elements in \mathcal{V} . Inherited from Lemma 2, we have

$$\begin{bmatrix} p_1 & \cdots & p_i & p_j & p_k \\ 1 & \cdots & 1 & 1 & 1 \end{bmatrix} \mathbf{\Omega} = \mathbf{0}.$$
 (2)

After adding a new vertex v_u and three edges, e_{iu} , e_{ju} , e_{ku} (as shown in Fig. 2), the following equation (3) is established based on the definition of a equilibrium stress.

$$\varpi_{iu} \left(\boldsymbol{p}_i - \boldsymbol{p}_u \right) + \varpi_{ju} \left(\boldsymbol{p}_j - \boldsymbol{p}_u \right) + \varpi_{ku} \left(\boldsymbol{p}_k - \boldsymbol{p}_u \right) = \boldsymbol{0}, \quad (3)$$

where ϖ_{iu} , ϖ_{ju} and ϖ_{ku} are the weights of edges e_{iu} , e_{ju} , e_{ku} . The resulting stress matrix Ω^+ is expressed as follows.

$$\mathbf{\Omega}^{+} = \begin{bmatrix} -\mathbf{\Omega}_{--}^{P1} & -\mathbf{\Omega}_{ii}^{+} & -\mathbf{\Omega}_{ij}^{P2} & \mathbf{0} \\ -\mathbf{\Omega}_{ii}^{-} & \mathbf{\Omega}_{ij}^{+} & \mathbf{\Omega}_{ij}^{+} & \mathbf{\Omega}_{ik}^{-} & -\mathbf{\omega}_{iu} \\ (\mathbf{\Omega}_{-}^{P2})^{T} & \mathbf{\Omega}_{ji}^{+} & \mathbf{\Omega}_{jj}^{+} & \mathbf{\Omega}_{jk}^{+} & -\mathbf{\omega}_{ju} \\ & + \mathbf{\Omega}_{ki}^{+} & \mathbf{\Omega}_{kj}^{+} & \mathbf{\Omega}_{kk}^{+} & -\mathbf{\omega}_{ku} \\ \mathbf{0} & -\mathbf{\omega}_{iu} & -\mathbf{\omega}_{ju} & -\mathbf{\omega}_{ku} & \mathbf{\omega}_{uu} \end{bmatrix}, \quad (4)$$

where $\varpi_{uu} = \varpi_{iu} + \varpi_{ju} + \varpi_{ku}$. Similarly, we have

$$\begin{bmatrix} \boldsymbol{p}_1 & \cdots & \boldsymbol{p}_i & \boldsymbol{p}_j & \boldsymbol{p}_k & \boldsymbol{p}_u \\ 1 & \cdots & 1 & 1 & 1 & 1 \end{bmatrix} \boldsymbol{\Omega}^+ = \boldsymbol{0}.$$
 (5)

Combining eqs. (2)-(5), we get

$$\underbrace{\begin{bmatrix} \boldsymbol{p}_{i} & \boldsymbol{p}_{j} & \boldsymbol{p}_{k} & \boldsymbol{p}_{u} \\ 1 & 1 & 1 & 1 \end{bmatrix}}_{\triangleq \boldsymbol{P}_{u}} \\ \begin{bmatrix} \Omega_{ii}^{+} - \Omega_{ii} & \Omega_{ij}^{+} - \Omega_{ij} & \Omega_{ik}^{+} - \Omega_{ik} & -\varpi_{iu} \\ \Omega_{ij}^{+} - \Omega_{ij} & \Omega_{jj}^{+} - \Omega_{jj} & \Omega_{jk}^{+} - \Omega_{jk} & -\varpi_{ju} \\ \Omega_{ik}^{+} - \Omega_{ik} & \Omega_{jk}^{+} - \Omega_{jk} & \Omega_{kk}^{+} - \Omega_{kk} & -\varpi_{ku} \\ -\varpi_{iu} & -\varpi_{ju} & -\varpi_{ku} & \varpi_{uu} \end{bmatrix}}_{\triangleq \boldsymbol{\Omega}_{u}} = \boldsymbol{0}.$$

$$(6)$$

Since v_u is not collinear with any two of v_i , v_j , v_k , the vectors $(\mathbf{p}_i - \mathbf{p}_u)$, $(\mathbf{p}_j - \mathbf{p}_u)$ and $(\mathbf{p}_k - \mathbf{p}_u)$ are linearly independent, which implies that rank $(\mathbf{P}_u) = 3$. Accordingly, there exists a nonzero vector $\boldsymbol{\phi} = [\phi_1, \phi_2, \phi_3, \phi_4]^T$ satisfying $\mathbf{P}_u \boldsymbol{\phi} = \mathbf{0}$, which means

$$\phi_1 + \phi_2 + \phi_3 + \phi_4 = 0, \ \phi_1 p_i + \phi_2 p_j + \phi_3 p_k + \phi_4 p_u = 0.$$
 (7)

Accordingly, the matrix Ω_u can be designed as below.

$$\boldsymbol{\Omega}_{u} = s\boldsymbol{\phi}\boldsymbol{\phi}^{T} = s \begin{vmatrix} \phi_{1}^{2} & \phi_{1}\phi_{2} & \phi_{1}\phi_{3} & \phi_{1}\phi_{4} \\ \phi_{1}\phi_{2} & \phi_{2}^{2} & \phi_{2}\phi_{3} & \phi_{2}\phi_{4} \\ \phi_{1}\phi_{3} & \phi_{2}\phi_{3} & \phi_{3}^{2} & \phi_{3}\phi_{4} \\ \phi_{1}\phi_{4} & \phi_{2}\phi_{4} & \phi_{3}\phi_{4} & \phi_{4}^{2} \end{vmatrix} , \quad (8)$$

where s is a scaling parameter. Thus, we have $\varpi_{iu} =$ $-s\phi_1\phi_4, \ \varpi_{ju} = -s\phi_2\phi_4, \ \varpi_{ku} = -s\phi_3\phi_4 \text{ and } \varpi_{uu} = s\phi_4^2.$ Then, the corresponding stress matrix of the newly obtained framework (\mathcal{G}^+, p^+) is established as follows.

$$\Omega^{+} = \underbrace{\left[\begin{array}{c} \Omega \\ \hline \mathbf{0}_{1\times n} \\ \hline \mathbf{0}_{n\times 1} \\ \hline \mathbf{0}_$$

where Ω_u is described as eq. (8) with s > 0 and Ω is the original stress matrix related to $(\mathcal{G}, \boldsymbol{p})$.

Next, a comprehensive theoretical analysis is provided to demonstrate the effectiveness of our strategy in constructing affine frameworks.

Theorem 1: Under Assumption 1-2, consider a new vertex v_u and an affine formation framework $(\mathcal{G}, \boldsymbol{p})$ in \mathbb{R}^2 . With a positive scaling parameter s, after adding three edges connecting the vertex v_u and the existing vertices v_i , v_j and v_k to $(\mathcal{G}, \boldsymbol{p})$, the obtained framework $(\mathcal{G}^+, \boldsymbol{p}^+)$ is universally rigid and affinely localizable, with the corresponding stress matrix determined by (8) and (9).

Proof: Please refer to Appendix I for details.

According to Theorem 1, the distributed nature of our proposed strategy is fully presented. Specifically, the global positions are not necessary based on the definition of a equilibrium stress (3). The weight of an edge is determined by the relative position of the corresponding two endpoints when new edges are added to the framework, e.g., Ω_u in (8). Accordingly, the local measurements are enough to update the stress matrix. Moreover, only several vertices are involved in the generation and reconstruction of affine formation frameworks, which supports the distributed execution of our methods. The construction of affine formation framework is solved by local measurements and communication, leading to the great application prospects in multi-robot systems. In fact, our algorithms can be integrated with most, if not all, distributed affine formation control schemes. A simulation is carried out in Section V-B to verify the compatibility, where an example of formation tracking control law [5] is applied.

Furthermore, we present an algorithm for adding m vertices to $(\mathcal{G}, \boldsymbol{p})$, as shown in Algorithm 1. With Algorithm 1, we can easily grow the original affine formation framework to a larger scale, with a hierarchical structure. Intuitively, based on Theorem 1, we only use the local position measurements of a limited number of vertices, and incremental operations are executed to update the stress matrix. Different from [11], global position measurements and centralized optimization are not necessary in our proposed approach.

B. Edge Deletion

Note that removing an edge from the original framework $(\mathcal{G}, \boldsymbol{p})$ is equivalent to adjusting the related weight to zero,

AI	gorithm 1: Vertex Addition Algorithm in \mathbb{R}^2
I	nput: An affine formation framework $(\mathcal{G}, \boldsymbol{p})$ with a stress
	matrix $\mathbf{\Omega}$, q new vertices v_{a1}, \cdots, v_{aq} with
	$oldsymbol{p}_{a1},\cdots,oldsymbol{p}_{aq}.$
C	Dutput: An augmented framework $(\mathcal{G}_{add}, \boldsymbol{p}_{add})$ with a new
	stress matrix $\mathbf{\Omega}_{add}$.
1 S	et the positive scaling parameter s;
2 fo	or $\ell = 1, \ \cdots, \ q$ do
3	$v_u \leftarrow v_{a\ell}, p_u \leftarrow p_{a\ell};$
4	Choose three perceived vertices v_i , v_j and v_k , where
	$v_i, v_j, v_k \in \mathcal{V};$
5	$\mathcal{G}_{add} = (\mathcal{V}_{add}, \mathcal{E}_{add})$ where $\mathcal{V}_{add} \leftarrow \mathcal{V} \cup \{v_u\}$,
	$\mathcal{E}_{add} \leftarrow \mathcal{E} \cup \{e_{iu}, \ e_{ju}, \ e_{ku}\};$
6	$\boldsymbol{\Omega}_{u} \leftarrow s \boldsymbol{\phi} \boldsymbol{\phi}^{T}$ based on Eq. (8). $\boldsymbol{p}_{add} \leftarrow \left[\boldsymbol{p}^{T}, \boldsymbol{p}_{u}^{T} \right]^{T}$;
7	$\mathbf{\Omega}_{add} \leftarrow \mathbf{\Omega}_a + \mathbf{\Omega}_b$ defined in Eq. (9);

 $m{s} \hspace{0.1 in} \left| \hspace{0.1 in} \mathcal{G} \leftarrow \mathcal{G}_{add}, \hspace{0.1 in} m{p} \leftarrow m{p}_{add}, \hspace{0.1 in} m{\Omega} \leftarrow m{\Omega}_{add};
ight.$

9 Return
$$(\mathcal{G}_{add}, \boldsymbol{p}_{add})$$
 and $\boldsymbol{\Omega}_{add}$



Fig. 3. Possible regions for placing four vertices in general position.

without damaging the universal rigidity and affine localizability. Using the parameter s in eq. (8), a natural idea is generated to eliminate a certain edge, as detailed below.

Take the removal of edge e_{jk} between vertex v_j and v_k in $(\mathcal{G}, \boldsymbol{p})$ as an example. If we select two suitable vertices v_i and v_q in $\mathcal{N}_i^p \cap \mathcal{N}_k^p$, the vertices v_i , v_j and v_k can form a triangle, and the regions of the fourth vertex v_q lies in can be labeled by a, \dots, g , as shown in Fig. 3. For four non-collinear vertices, we get a non-zero vector ϕ satisfying $oldsymbol{p}_i$ $oldsymbol{p}_j$ $oldsymbol{p}_k$ $oldsymbol{p}_q$ ϕ = 0. In the original framework $1 \quad 1$ 1 1 $(\mathcal{G}, \boldsymbol{p})$, the stress matrix $\boldsymbol{\Omega}$ is indexed based on the elements in \mathcal{V} , which are reordered such that v_i , v_j , v_k , v_q are listed at the end. The entries involved in these four vertices in the stress matrix Ω can be described as below.

$$\mathbf{\Omega}_{v} = \begin{bmatrix} \Omega_{ii} & \Omega_{ij} & \Omega_{ik} & \Omega_{iq} \\ \Omega_{ij} & \Omega_{jj} & \Omega_{jk} & \Omega_{jq} \\ \Omega_{ik} & \Omega_{jk} & \Omega_{kk} & \Omega_{kq} \\ \Omega_{iq} & \Omega_{jq} & \Omega_{kq} & \Omega_{qq} \end{bmatrix}.$$
(10)

After eliminating e_{jk} , the framework is described by $(\mathcal{G}^-, \mathbf{p})$, and the corresponding block matrix is established as follows.

$$\boldsymbol{\Omega}_{v}^{-} = \begin{bmatrix} \Omega_{ii}^{-} & \Omega_{ij}^{-} & \Omega_{ik}^{-} & \Omega_{iq}^{-} \\ \Omega_{ij}^{-} & \Omega_{jj}^{-} & 0 & \Omega_{jq}^{-} \\ \Omega_{ik}^{-} & 0 & \Omega_{kk}^{-} & \Omega_{kq}^{-} \\ \Omega_{iq}^{-} & \Omega_{jq}^{-} & \Omega_{kq}^{-} & \Omega_{qq}^{-} \end{bmatrix}.$$
(11)

Inspired by Eqs. (6) and (8), we have

$$\begin{bmatrix} \boldsymbol{p}_i & \boldsymbol{p}_j & \boldsymbol{p}_k & \boldsymbol{p}_q \\ 1 & 1 & 1 & 1 \end{bmatrix} (\boldsymbol{\Omega}_v^- - \boldsymbol{\Omega}_v) = \boldsymbol{0}.$$
(12)

TABLE I The sign of the weight of the edges connected to v_a

Regions	a	b	c	ð	e	f	g
k_1	+	+	_	+	+	_	_
k_2	_	+	+	_	+	+	_
$arpi_{iq}$	+	+	_	_	_	+	+
$arpi_{jq}$	_	+	+	+	—	_	+
$arpi_{kq}$	_	_	_	+	+	+	+

Let
$$\Omega_u \triangleq \Omega_v^- - \Omega_v = s_{ed} \phi \phi^T$$
, where $\phi = [\phi_1, \cdots, \phi_4]^T$

$$\mathbf{\Omega}_v^- = \mathbf{\Omega}_v + \mathbf{\Omega}_u$$

$$= \mathbf{\Omega}_{v} + \frac{1}{\varpi_{qq}} \begin{bmatrix} \varpi_{iq}^{2} & \varpi_{iq}\varpi_{jq} & \varpi_{iq}\varpi_{kq} & -\varpi_{iq}\varpi_{qq} \\ \varpi_{iq}\varpi_{jq} & \varpi_{jq}^{2} & \varpi_{jq}\varpi_{kq} & -\varpi_{jq}\varpi_{qq} \\ \varpi_{iq}\varpi_{kq} & \varpi_{jq}\varpi_{kq} & \varpi_{kq}^{2} & -\varpi_{kq}\varpi_{qq} \\ -\varpi_{iq}\varpi_{qq} & -\varpi_{jq}\varpi_{qq} & -\varpi_{kq}\varpi_{qq} & \varpi_{qq}^{2} \end{bmatrix}$$
(13)

where $\varpi_{iq} = -s_{ed}\phi_1\phi_4$, $\varpi_{jq} = -s_{ed}\phi_2\phi_4$, $\varpi_{kq} =$ $-s_{ed}\phi_3\phi_4$ and $\varpi_{qq} = s_{ed}\phi_4^2$. To eliminate e_{jk} , the following equation is established

$$\Omega_{jk} + \frac{\varpi_{jq}\varpi_{kq}}{\varpi_{qq}} = 0 \quad \Rightarrow s_{ed} = -\frac{\Omega_{jk}}{\phi_2\phi_3}.$$
 (14)

To maintain the universal rigidity and affine localizability of the obtained framework $(\mathcal{G}^-, \mathbf{p}), s_{ed} > 0$ is required based on Theorem 1, implying that

$$\Omega_{jk}\varpi_{jq}\varpi_{kq} < 0. \tag{15}$$

With a specific Ω_{ik} , the sign of s_{ed} is determined by the signs of ϖ_{jq} and ϖ_{kq} , which depend on the relative positions among v_i , v_j , v_k and v_q . In detail, since v_q is not collinear with any two of v_i , v_j and v_k , the vectors $p_q - p_i$ and $p_q - p_j$ can serve as a set of bases in \mathbb{R}^2 . That is,

$$\boldsymbol{p}_{q} - \boldsymbol{p}_{k} = k_{1} \left(\boldsymbol{p}_{q} - \boldsymbol{p}_{i} \right) + k_{2} \left(\boldsymbol{p}_{q} - \boldsymbol{p}_{j} \right), \qquad (16)$$

where k_1 and k_2 are two real parameters that are not simultaneously zero. Combining with eq. (3), we have

$$\left(\varpi_{iq}+\varpi_{kq}k_{1}\right)\left(\boldsymbol{p}_{q}-\boldsymbol{p}_{i}\right)+\left(\varpi_{jq}+\varpi_{kq}k_{2}\right)\left(\boldsymbol{p}_{q}-\boldsymbol{p}_{j}\right)=\boldsymbol{0}.$$

Accordingly, the following equations are deduced.

$$\varpi_{iq} = -k_1 \varpi_{kq}, \quad \varpi_{jq} = -k_2 \varpi_{kq}, \quad (17)$$

which mean $\varpi_{iq} \varpi_{jq} = k_1 k_2 \varpi_{kq}^2$, $\varpi_{iq} \varpi_{kq} = -k_1 \varpi_{kq}^2$ and $\varpi_{jq} \varpi_{kq} = -k_2 \varpi_{kq}^2$. When v_q lies in different regions, the signs of k_1 , k_2 and the weights are listed in Table. I, which serves as a guideline to determine whether the edge e_{ik} can be deleted without damaging universal rigidity and affine localizability. For example, if the weight of e_{jk} is positive, i.e., $\Omega_{jk} < 0$, a vertex v_q in \mathcal{G} lying in Region \mathfrak{a} , \mathfrak{d} or \mathfrak{g} should be chosen to ensure a positive s_{ed} based on (15) and Table. I. However, it is also possible that there are no vertices lying in these regions. Accordingly, based on whether e_{ik} can be eliminated through the rearrangement of edge weights in \mathcal{G} , there are three possible cases presented as follows.

Case (1): There are two vertices $v_i, v_q \in \mathcal{N}_i^p \cap \mathcal{N}_k^p$ to ensure that s_{ed} in (14) is positive.

In this case, we can delete the edge e_{ik} directly to obtain a new framework $(\mathcal{G}_{ed}, \boldsymbol{p}_{ed})$, where $\mathcal{G}_{ed} = (\mathcal{V}, \mathcal{E} \setminus \{e_{jk}\})$, $p_{ed} = p$. The stress matrix Ω_{ed} for $(\mathcal{G}_{ed}, p_{ed})$ is shown as below,

$$\boldsymbol{\Omega}_{ed} = \boldsymbol{\Omega} + \left[-\frac{\boldsymbol{0}_{(n-4)\times(n-4)}}{\boldsymbol{0}_{4\times(n-4)}} \right] \left[-\frac{\boldsymbol{0}_{(n-4)\times4}}{\boldsymbol{\Omega}_u} \right], \quad (18)$$

where $\Omega_u = -\frac{\Omega_{jk}}{\phi_2 \phi_3} \phi \phi^T$. The key point is to find the eligible vertices in $\mathcal{N}_j^p \cap \mathcal{N}_k^p$. In practice, the number of perceived neighbors is limited so that the local traversal is usually acceptable. For higher efficiency, we can also set a time upper limit on the search for suitable neighbors. If we cannot find proper neighbors within the specified time, a relay strategy is activated. That is, we can eliminate a specific edge by adding new vertices, and the following two cases are considered.

Case (2): There are no vertices in $\mathcal{N}_i^p \cap \mathcal{N}_k^p$ to guarantee positive s_{ed} , but $\mathcal{N}_i^p \cap \mathcal{N}_k^p \neq \emptyset$.

In this case, there are no suitable vertices to ensure that the positive scaling parameter s_{ed} is available. To complete our edge deletion algorithm, we can eliminate specific edges by adding a new relay vertex. The following assumption describes the criteria for selecting the relay vertices.

Assumption 3: Suppose that the vertices $v_i \in \mathcal{N}_i^p \cap \mathcal{N}_k^p$ and a relay vertex v_r meet the following conditions.

- $v_i, v_j, v_k \in \mathcal{N}^p_r$, and the set $\{ \boldsymbol{p}_i, \ \boldsymbol{p}_j, \ \boldsymbol{p}_k, \ \boldsymbol{p}_r \}$ is in general position;
- If the weight $\varpi_{jk} = -\Omega_{jk} > 0$, v_r lies in Region \mathfrak{a} , \mathfrak{d} or \mathfrak{g} of a triangle formed by v_i , v_j and v_k . If the weight $\varpi_{jk} = -\Omega_{jk} < 0, v_r$ lies in Region b, c, c or f, as shown in Fig. 3.

Under Assumption 3, we can reconstruct the a framework without edge e_{ik} as follows. Remove e_{ik} in the original affine formation framework $(\mathcal{G}, \boldsymbol{p})$, and then add v_r along with three edges e_{ir} , e_{jr} and e_{kr} based on Theorem 1. Accordingly, a new framework $(\mathcal{G}_{ed}, \boldsymbol{p}_{ed})$ is built, where $\mathcal{G}_{ed} =$ $(\mathcal{V} \cup \{v_r\}, \mathcal{E} \cup \{e_{ir}, e_{jr}, e_{kr}\} \setminus \{e_{jk}\}), \ \boldsymbol{p}_{ed} = [\boldsymbol{p}^T, \boldsymbol{p}_r^T]^T.$ The appropriate stress exist and present in (19) to guarantee that $(\mathcal{G}_{ed}, \boldsymbol{p}_{ed})$ is universally rigid and affinely localizable.

$$\boldsymbol{\Omega}_{ed} = \begin{bmatrix} \mathbf{\Omega} & \mathbf{0}_{n \times 1} \\ \mathbf{\overline{0}}_{1 \times n} & \mathbf{\overline{0}}_{n} \end{bmatrix} + \begin{bmatrix} \mathbf{0}_{(n-3) \times (n-3)} & \mathbf{0}_{n-3} \\ \mathbf{\overline{0}}_{4 \times (n-3)} & \mathbf{\overline{\Omega}}_{u} \end{bmatrix}$$
(19)

where $s_{ed} = -\frac{\Omega_{jk}}{\bar{\phi}_2 \bar{\phi}_3}$ and $\Omega_u = s_{ed} \bar{\phi} \bar{\phi}^T$. The vector $\bar{\phi} =$ $\begin{bmatrix} \bar{\phi}_1, \ \bar{\phi}_2, \ \bar{\phi}_3, \ \bar{\phi}_4 \end{bmatrix}^T \text{ satisfies } \begin{bmatrix} \mathbf{p}_i & \mathbf{p}_j & \mathbf{p}_k & \mathbf{p}_r \\ 1 & 1 & 1 & 1 \end{bmatrix} \bar{\phi} = \mathbf{0}.$ Case (3): $\mathcal{N}_j^p \cap \mathcal{N}_k^p = \emptyset.$

In this case, there is no vertex in $\mathcal{N}_j^p \cap \mathcal{N}_k^p$ such that two relay vertices, v_{r_1} and v_{r_2} , are needed with the following assumption.

Assumption 4: Suppose that the vertices v_{r_1} and v_{r_2} meet the following conditions.

- Any two vertices of v_j , v_k , v_{r_1} and v_{r_2} can measure their relative positions to each other. The set $\{p_j, p_k, p_{r_1}, p_{r_2}\}$ is in general position;
- If the weight $\varpi_{jk} = -\Omega_{jk} > 0$, v_{r_2} lies in Region \mathfrak{a} , \mathfrak{d} or \mathfrak{g} of a triangle formed by v_{r_1} , v_j and v_k . If the weight



Fig. 4. An example of edge deletion. (a) An affine formation framework (\mathcal{G}, p) with $p = [0, 1, 1, 0, 0, -1, -1, 0, 1, -1]^T$, where e_{23} is going to be deleted. (b) An affine formation framework (\mathcal{G}, p) with $p = [0, 1, 1, 0, 0, -1, -1, 0]^T$, where e_{23} is going to be deleted. (c) The obtained framework $(\mathcal{G}_{ed}, p_{ed})$ without the edge e_{23} .

 $\varpi_{jk} = -\Omega_{jk} < 0, v_{r_2} \text{ lies in Region } \mathfrak{b}, \mathfrak{c}, \mathfrak{e} \text{ or } \mathfrak{f}, \text{ as shown in Fig. 3.}$

Choose v_j and v_k to be the neighbors of v_{r_1} , and add the vertex v_{r_1} to the original framework $(\mathcal{G}, \boldsymbol{p})$ by using Algorithm 1. Then, we merge v_{r_2} to the framework with three edges, $e_{r_1r_2}$, e_{jr_2} and e_{kr_2} , based on Theorem 1 again. We can set the scaling parameter as $s_{ed} = -\frac{\Omega_{jk}}{\bar{\phi}_2\bar{\phi}_3}$, where $\begin{bmatrix} \boldsymbol{p}_{r_1} & \boldsymbol{p}_j & \boldsymbol{p}_k & \boldsymbol{p}_{r_2} \\ 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} \bar{\phi}_1, \ \bar{\phi}_2, \ \bar{\phi}_3, \ \bar{\phi}_4 \end{bmatrix}^T = \mathbf{0}$. Consequently, the framework $(\mathcal{G}_{ed}, \boldsymbol{p}_{ed})$ is constructed, where $\mathcal{G}_{ed} =$

Juently, the framework $(\mathcal{G}_{ed}, \boldsymbol{p}_{ed})$ is constructed, where $\mathcal{G}_{ed} = (\mathcal{V} \cup \{v_{r_1}, v_{r_2}\}, \mathcal{E} \cup \{e_{\sim r_1}, e_{jr_2}, e_{kr_2}, e_{r_1r_2}\} \setminus \{e_{jk}\}), \boldsymbol{p}_{ed} = [\boldsymbol{p}^T, \boldsymbol{p}_{r_1}^T, \boldsymbol{p}_{r_2}^T]^T$. In **Case (3)**, the vertex v_{r_1} plays the role similar to v_i in **Case (2)**. Thus, the similar steps and analysis is omitted.

To explain our proposed edge deletion strategy more clearly, an example is presented as below.

Example 1 (Edge Deletion): To remove e_{23} in Fig. 4(a) and 4(b), two strategies mentioned above are executed, respectively. Suppose that the distance between any two vertices is less than the perception distance d_{per} . In Fig. 4(a), j = 2, k=3 so that $\mathcal{N}_{j}^{p}\cap\mathcal{N}_{k}^{p}=\{1,4,5\}$. To delete the edge e_{23} , let i = 1 and q = 5. We have $\phi = \frac{1}{\sqrt{10}} [1, -2, -1, 2]^T$ so that $s_{ed} = -\frac{\Omega_{23}}{\phi_2 \phi_3} = -\frac{-0.8}{0.2} = 4 > 0$. Thus, Corollary 1 is applied and the obtained affine formation framework is shown in Fig. 4(c). Nevertheless, in Fig. 4(b), $\mathcal{N}_i^p \cap \mathcal{N}_k^p = \{1, 4\} \neq \emptyset$. If i = 1 and q = 4, we can calculate $s_{ed} = -4 < 0$, which implies that there is no suitable vertices to ensure a positive scaling parameter, indicating that a relay vertex is necessary. Thus, following **Case** (2), choose a relay vertex v_5 with $\boldsymbol{p}_5 = [1, -1]^T$, lying in Region \mathfrak{d} . Three edges, $e_{15}, e_{25}, e_{35}, e$ are added to the original framework with $s_{ed} = 2$, resulting $\varpi_{23} = 0$ after the operation. The obtained affine formation framework is also shown in Fig. 4(c).

Corollary 1: Given a framework $(\mathcal{G}, \mathbf{p})$ containing an edge e_{jk} with the weight $-\Omega_{jk}$, if there are proper vertices v_i and v_q in $\mathcal{N}_j^p \cap \mathcal{N}_k^p$ to ensure a positive scaling parameter s_{ed} shown in eq. (14), the obtained framework $(\mathcal{G}_{ed}, \mathbf{p}_{ed})$ with the stress matrix (18) is universally rigid and affinely localizable.

The proof of Corollary 1 directly follows a similar sketch with Theorem 1, and is omitted here. With the help of Corollary 1, we can rearrange the stress to drive the weight of e_{jk} to zero, which is equivalent to removing the edge from the original framework. The edge deletion strategies proposed



Fig. 5. An example of the removal of a inner node. (a) The inner node v_5 (red, with three children v_7 , v_8 , v_{10}) is removed from the framework, and v_7 is selected to inherit its role. (b) The parent sets of v_7 , v_8 , v_{10} are rebuilt, and the hierarchical structure is reconstructed.

in different cases share the same fundamental idea presented in Corollary 1, with the differences lying in the selection of neighbor vertices and the design of scaling parameter s_{ed} .

Remark 1: The edge deletion strategy using relay vertices in **Case (2) and (3)** is directly inspired by Theorem 1. To ensure a positive scaling parameter, the position of relay vertex needs to be carefully selected, taking Table I as a reference. In this paper, the relay vertex is assumed to be on standby and respond immediately. In a real robot system, more technical details need to be designed to guarantee the smooth operation of relay robots.

C. Vertex Deletion

In this subsection, we consider an inverse operation of "vertex addition" presented in Section III-A, the removal of a vertex from an affine formation framework (\mathcal{G}, p) .

Based on Algorithm 1, the hierarchical structure of an affine formation framework (\mathcal{G}, p) is naturally built. Denote the hierarchy of vertices in an original affine formation framework $(\mathcal{G}_0, \boldsymbol{p}_0)$ as 0. Define the hierarchy $h(v_i)$ of a vertex v_i as the length of its longest path from v_i to the vertices in $(\mathcal{G}_0, \boldsymbol{p}_0)$. Given a series of free vertices to be integrated into $(\mathcal{G}_0, \boldsymbol{p}_0)$, Algorithm 1 can be applied to generate $(\mathcal{G}, \boldsymbol{p})$. To add a free v_u to the existing graph, three existing vertices are selected to be parents, denoted by v_i, v_j, v_k . The hierarchy of vertex v_u is defined as $h(v_u) = \max(h(v_i), h(v_j), h(v_k)) + 1$. According to whether a vertex has a child or not, the vertices in the graph can be classified into two categories: (a) outer node, a vertex with no child (e.g., v_9 and v_{10} in Fig. 5(a)); (b) inner node, a vertex with at least one child (e.g., v_5 and v_7 in Fig. 5(a)). We consider the vertex deletion problem in both cases in this section.

Case 1 (Deletion of an Outer Node): We first consider the case with an outer node. Consider an affine framework (\mathcal{G}, p) with $n \ (n \ge 5)$ vertices, containing an outer node v_u . After removing vertex v_u and all edges connected to v_u , we can obtain a framework $(\mathcal{G}_{vd}, p_{vd})$.

For the affine formation framework $(\mathcal{G}, \boldsymbol{p})$, the stress matrix

can be described as follows.

$$\boldsymbol{\Omega} = \begin{bmatrix} -\boldsymbol{\Omega}_{vd}^{B1} & \boldsymbol{\Omega}_{ii} & \boldsymbol{\Omega}_{vd}^{B2} & \boldsymbol{\Omega}_{ik} & \boldsymbol{\Omega}_{iu} \\ -\boldsymbol{\Omega}_{vd}^{B2} & \boldsymbol{\Omega}_{ii} & \boldsymbol{\Omega}_{ij} & \boldsymbol{\Omega}_{ik} & \boldsymbol{\Omega}_{iu} \\ \begin{pmatrix} \boldsymbol{\Omega}_{vd}^{B2} \end{pmatrix}^{T} & \boldsymbol{\Omega}_{ij} & \boldsymbol{\Omega}_{jj} & \boldsymbol{\Omega}_{jk} & \boldsymbol{\Omega}_{ju} \\ -\boldsymbol{\Omega}_{iu}^{-1} & \boldsymbol{\Omega}_{ij} & \boldsymbol{\Omega}_{ji} & \boldsymbol{\Omega}_{jk} & \boldsymbol{\Omega}_{ju} \\ -\boldsymbol{\Omega}_{iu}^{-1} & \boldsymbol{\Omega}_{iu}^{-1} & \boldsymbol{\Omega}_{ju}^{-1} & \boldsymbol{\Omega}_{iu}^{-1} & \boldsymbol{\Omega}_{iu}^{-1} \\ \end{bmatrix}, \quad (20)$$

Notably, $\Omega_{uu} > 0$. Based on the definition of an equilibrium stress, we have the following equation for v_u ,

$$\Omega_{iu} \left(\boldsymbol{p}_{i} - \boldsymbol{p}_{u}\right) + \Omega_{ju} \left(\boldsymbol{p}_{j} - \boldsymbol{p}_{u}\right) + \Omega_{ku} \left(\boldsymbol{p}_{k} - \boldsymbol{p}_{u}\right) = \mathbf{0}$$

$$\Rightarrow \underbrace{\begin{bmatrix} \boldsymbol{p}_{i} & \boldsymbol{p}_{j} & \boldsymbol{p}_{k} & \boldsymbol{p}_{u} \\ 1 & 1 & 1 & 1 \end{bmatrix}}_{\triangleq \boldsymbol{P}_{u}} \underbrace{\begin{bmatrix} \Omega_{iu} & \Omega_{ju} & \Omega_{ku} & \Omega_{uu} \end{bmatrix}^{T}}_{\triangleq \boldsymbol{\Omega}_{\phi}} = \mathbf{0}.$$
(21)

Since $(\mathcal{G}, \mathbf{p})$ is in general position so that v_u is not collinear with any two of v_i , v_j , v_k , it is deduced that rank $(\mathbf{P}_u) = 3$ and nullity $(\mathbf{P}_u) = 1$. Define an auxiliary matrix $\mathbf{\Omega}_{vd}^u$ as below,

$$\mathbf{\Omega}_{vd}^{u} = \frac{1}{\Omega_{uu}} \mathbf{\Omega}_{\phi} \mathbf{\Omega}_{\phi}^{T} \succeq 0.$$
(22)

Then, we can establish the stress matrix Ω_{vd} for the framework $(\mathcal{G}_{vd}, \boldsymbol{p}_{vd})$ as below.

$$\begin{bmatrix} \boldsymbol{\Omega}_{vd} & \boldsymbol{0} \\ \boldsymbol{0} & \boldsymbol{0} \end{bmatrix} = \boldsymbol{\Omega} - \begin{bmatrix} \boldsymbol{0} & \boldsymbol{0} \\ \boldsymbol{0} & \boldsymbol{\Omega}_{vd}^{u} \end{bmatrix}, \quad (23)$$

Thus,

Then, we establish the following theorem to prove $(\mathcal{G}_{vd}, \boldsymbol{p}_{vd})$ is universally rigid and affinely localizable.

Theorem 2: Consider an affine framework (\mathcal{G}, p) with $n \ (n \geq 5)$ vertices, containing an outer node v_u . After removing vertex v_u and all edges connected to v_u , we can obtain a universally rigid and affinely localizable framework $(\mathcal{G}_{vd}, p_{vd})$ with a stress matrix Ω_{vd} shown in (24), where $\mathcal{G}_{vd} = (\mathcal{V} \setminus \{v_u\}, \mathcal{E} \setminus \{e_{iu}, e_{ju}, e_{ku}\}).$

Proof: It is obvious that $(\mathcal{G}_{vd}, p_{vd})$ is in general position. Then, we need to clarify that $\Omega_{vd} \succeq 0$, rank $(\Omega_{vd}) = n-4$ and the block matrix describing the stress among followers $\Omega_{ff}^{vd} \succ 0$. The following proof has a similar sketch with Theorem 1 in Appendix I. Based on the characteristics of Ω_{vd} , we conclude that $(\mathcal{G}_{vd}, p_{vd})$ is universally rigid and affinely localizable.

By directly applying Theorem 2, we can remove outer nodes without damaging the universal rigidity and affine localizability of the remaining framework.

Case 2 (Deletion of an Inner Node): When an inner node leaves the framework (e.g., v_5 in Fig. 5(a)), the universal rigidity and affine localizability of the framework are destroyed and need to be repaired. Hence, a strategy inspired by inheritance is derived.

Assumption 5: For an inner node v_u , denote the parents as $\mathcal{P}_u = \{v_{p_i}^u, v_{p_j}^u, v_{p_k}^u\}$ and the children as $\mathcal{C}_u = \{v_{c_1}^u, v_{c_2}^u \cdots v_{c_m}^u\}$, where $h(v_{c_1}^u) \leq \cdots \leq h(v_{c_m}^u)$. Suppose that the perception distance is large enough to ensure $v_{\mathcal{P}} \in$ $\mathcal{N}_{\mathcal{C}}^p$, where $v_{\mathcal{P}} \in \mathcal{P}_u$ and $v_{\mathcal{C}} \in \mathcal{C}_u$. The set $\{p_{\mathcal{P}}\} \cup \{p_{\mathcal{C}}\}$ is in general position, where $p_{\mathcal{P}}$ is the position of $v_{\mathcal{P}}$, and $p_{\mathcal{C}}$ is defined analogously.

Assumption 5 lays the foundation for establishing edges between the parents and children of v_u . Based on Theorem 1, a matrix Ω_b in (9) is constructed to update the stress matrix when adding a free vertex. Accordingly, denote the auxiliary matrix to add v_u , $v_{c_1}^u \cdots v_{c_m}^u$ as Ω_b^u , $\Omega_b^{uc_1} \cdots \Omega_b^{uc_m} \in \mathbb{R}^{n \times n}$, and their orders are unified by inserting zeros. The steps of removing an inner node v_u is presented as follows.

- S1. Remove all edges connecting v_u , and update the stress matrix as $\Omega_{cache} = \Omega \Omega_b^u \Omega_b^{uc_1} \dots \Omega_b^{uc_m}$. Obviously, the elements in the *u*-th row and *u*-th column of Ω_{cache} become zero. Delete the *u*-th row and the *u*-th column, and a simplified matrix is shown as $\tilde{\Omega}_{cache} \in \mathbb{R}^{(n-1)\times(n-1)}$.
- S2. The vertex v_{c_1} , a child of v_u that has the highest hierarchy, is chosen to inherit its role in the framework. Choose one vertex from $\mathcal{P}_u \setminus \mathcal{P}_{c_1}$, labeled by $v_{p_1}^{c_1}$. Rebuild the parent set of v_{c_1} , $\overline{\mathcal{P}}_{c_1}$, with $v_{p_1}^{c_1}$ and the remaining vertices in \mathcal{P}_{c_1} . Using the positions of v_{c_1} and its parents, construct the auxiliary matrix $\mathbf{\Omega}_b^{c_1} \in \mathbb{R}^{(n-1)\times(n-1)}$;
- S3. Rebuild the parent sets of other vertices in $C_u \setminus \{v_{c_1}\}$. Taking the vertex v_{c_2} as an example, if v_{c_1} is not a parent of v_{c_2} (i.e., $\mathcal{P}_{c_2} = \{v_u, v_{p_j}^{c_2}, v_{p_k}^{c_2}\}$), the new parent set is $\overline{\mathcal{P}}_{c_2} = \{v_{c_1}, v_{p_j}^{c_2}, v_{p_k}^{c_2}\}$, where v_u is replaced by v_{c_1} . If $v_{c_1} \in \mathcal{P}_{c_2}$, we can choose one vertex in $\mathcal{P}_u \setminus \mathcal{P}_{c_2}$ to replace v_u , e.g., $\overline{\mathcal{P}}_{c_2} = \{v_{p_i}^u, v_{c_1}, v_{p_k}^{c_2}\}$. The remaining children of v_u follow the strategy. An example is presented in Fig. 5(b).
- S4. Construct auxiliary matrices for vertices in $C_u \setminus \{v_{c_1}\}$ using their own and parental positions, i.e., $\Omega_b^{c_1c_2}, \dots, \Omega_b^{c_1c_m} \in \mathbb{R}^{(n-1)\times(n-1)}$ (the order is unified by inserting zero). Accordingly, a new affine formation framework $(\mathcal{G}_{vd}, \mathbf{p}_{vd})$ is reconstructed without v_u , and the corresponding stress matrix is

$$\Omega_{vd} = \tilde{\Omega}_{cache} + \Omega_b^{c_1 c_2} + \dots + \Omega_b^{c_1 c_m}.$$
 (25)

The universal rigidity and affine localizability of $(\mathcal{G}_{vd}, \boldsymbol{p}_{vd})$ can be proved by analyzing the characteristic of Ω_{vd} shown in Eq. (25). Combining the proof sketch of Theorem 1 and 2, a similar process can be implemented, which is omitted here. To explain our method more clearly, an example is presented.

Example 2 (Vertex Deletion): Using Algorithm 1, an affine formation framework shown in Fig. 6(a) is generated, where p $\left[8, 0, 0, 8, -8, 0, 0, -8, 9, -10, 0, -12, 11, 1, 14, -14, -7, -5\right]^T$ v_7 , v_8 , v_9 are outer nodes and v_5 , v_6 are inner nodes. To remove an outer node (e.g., v_9), Theorem 2 is directly applied. After calculating, the eigenvalues of the stress matrix are 0, 0, 0, 0.107, 0.398, 0.91, 1.82, 4.77, which demonstrates the universal rigidity and affine localizability of the obtained framework without v_9 , as shown in Fig. 6(b). To verify the strategy of deleting an inner node, take the removal of v_5 as an example. After removing all edges connected to v_5 , the graph is shown in Fig. 6(c). Obviously, the rigidity of framework is destroyed. To repair the universal rigidity and affine localizability, the second and third step, S2 and S3, are



Fig. 6. An example of the vertex deletion algorithm. (a) The original affine formation framework, where $v_1, v_2, v_3 \in \mathcal{V}_l$. (b) The removal of an outer node v_9 . (c) The removal of an inner node v_5 . S1-S2: delete all edges connected to v_5 , and v_6 is chosen to inherit of the role of v_5 . (c) The removal of an inner node v_5 . S3-S4: repair the universal rigidity and affine localizability.

executed. We choose v_6 to inherit the role of v_5 , so that the hierarchical relationship between vertices is reconstructed, as presented in Tab. 2. The final framework is shown in Fig. 6(d), and the eigenvalues of the corresponding stress matrix obtained from (25) are 0, 0, 0, 0.204, 0.36, 1.19, 1.38, 4.87. In a word, the effectiveness of our proposed approached in reconstructing a universally rigid and affinely localizable framework has been validated.

The proposed vertex deletion strategy is also distributed, as it only utilizes local measurements to change the local structure of the affine framework. Taking the deletion of a inner node shown in Fig. 6(c) and 6(d) as an example, the connections among the inner node's parents and children are reconstructed, and the remaining part in the framework is preserved. The affine formation framework generated by Algorithm 1 has a clear hierarchical structure with specific parents and children, making it convenient for further research. For any other affine formation framework, as long as a similar hierarchical structure can be established, our proposed vertex deletion strategy can be extended.

IV. DISCUSSION

A. Extension to three-dimensional space

The strategies proposed in Section III concentrate on the affine formation framework construction in \mathbb{R}^2 . Actually, our method can be extended to three-dimensional scenarios following the same idea. Taking the vertex addition strategy as an example, a new vertex v_u is intended to be added to $(\mathcal{G}, \mathbf{p})$ in three-dimensional space. Inspired by Theorem 1, four edges are introduced to the original framework $(\mathcal{G}, \mathbf{p})$, and the corresponding stress are determined by the position of the new vertex v_u with respect to the four affinely independent vertices in \mathcal{N}_u^p , denoted by v_i , v_j , v_k and v_g . Similar to Section III-A, suppose that the set { \mathbf{p}_i , \mathbf{p}_j , \mathbf{p}_k , \mathbf{p}_g , \mathbf{p}_u } is in general

position.

To grow the affine framework with a new vertex v_u , four edges e_{iu} , e_{ju} , e_{ku} , e_{gu} are linked to $(\mathcal{G}, \boldsymbol{p})$ to obtain a extended framework $(\mathcal{G}^+, \boldsymbol{p}^+)$, where $\boldsymbol{p}^+ = [\boldsymbol{p}^T, \boldsymbol{p}_u^T]^T$ and $\mathcal{G}^+ = (\mathcal{V} \cup \{v_u\}, \mathcal{E} \cup \{e_{iu}, e_{ju}, e_{ku}, e_{gu}\})$. For the new vertex v_u , we have the following equation due to the definition of the equilibrium stress,

$$\begin{aligned} \varpi_{iu} \left(\boldsymbol{p}_i - \boldsymbol{p}_u \right) + \varpi_{ju} \left(\boldsymbol{p}_j - \boldsymbol{p}_u \right) \\ + \varpi_{ku} \left(\boldsymbol{p}_k - \boldsymbol{p}_u \right) + \varpi_{gu} \left(\boldsymbol{p}_g - \boldsymbol{p}_u \right) = \boldsymbol{0}, \end{aligned}$$
 (26)

where ϖ_{iu} , ϖ_{ju} , ϖ_{ku} and ϖ_{gu} are the weights of edges e_{iu} , e_{ju} , e_{ku} , e_{gu} . Define P_u as $P_u \triangleq \begin{bmatrix} p_i & p_j & p_k & p_g & p_u \\ 1 & 1 & 1 & 1 \end{bmatrix}$. Since $\operatorname{rank}(P_u) = 4$, there is a nonzero vector $\boldsymbol{\phi} = [\phi_1, \phi_2, \phi_3, \phi_4, \phi_5]^T$ satisfying $P_u \boldsymbol{\phi} = \mathbf{0}$. Similarly, define $\Omega_u = s \boldsymbol{\phi} \boldsymbol{\phi}^T$, where s > 0 is the scaling parameter. Accordingly, the weights can be represented by $\varpi_{iu} = -s \phi_1 \phi_5$, $\varpi_{ju} = -s \phi_2 \phi_5$, $\varpi_{ku} = -s \phi_3 \phi_5$ and $\varpi_{gu} = -s \phi_4 \phi_5$. The augmented stress matrix Ω^+ can be rewritten as

$$\mathbf{\Omega}^{+} = \left[\begin{array}{c} \mathbf{\Omega} & \mathbf{0}_{n \times 1} \\ \mathbf{\overline{0}}_{1 \times n} & \mathbf{0} \end{array} \right] + \left[\begin{array}{c} \mathbf{0}_{(n-4) \times (n-4)} & \mathbf{0}_{(n-4) \times 5} \\ \mathbf{\overline{0}}_{5 \times (n-4)} & \mathbf{0}_{u} \end{array} \right].$$
(27)

Correspondingly, we have the following result on vertex addition in \mathbb{R}^3 .

Corollary 2: Under Assumption 1, for a given affine formation framework $(\mathcal{G}, \mathbf{p})$ in \mathbb{R}^3 , adding a new vertex v_u and four edges between v_u with four vertices v_i, v_j, v_k and v_g , the universal rigidity and affine localizability are inherited by the extended affine formation framework $(\mathcal{G}^+, \mathbf{p}^+)$ with a stress matrix $\mathbf{\Omega}^+$ described in eq. (27).

The similar scheme employed in the proof of Theorem 1 can be applied to explain Corollary 2, which is omitted here for simplicity. Inspired by Corollary 2, the strategies for edge and vertex deletion in \mathbb{R}^3 can be extended along the lines of their counterparts in two-dimensional space.

B. Complexity Analysis

It is evident that our proposed framework reconstruction strategies are incremental, inherently possessing a relatively low time complexity. When the topology of the formation changes, we need a little computational resources to reconstruct the new framework, because only low dimensional matrix calculations are required in our algorithm. Taking Algorithm 1 as an example, providing that there are q new vertices joining the formation, the time complexity can be calculated as O(qn). Note that we can simultaneously apply Algorithm 1 to all new vertices if the vertex addition process happens in parallel. Then, the time complexity will be further reduced to O(n). Therefore, in practice, our algorithm is portable enough to be applied to robots with limited computing power to deal with a variety of collaborative tasks in time.

V. SIMULATIONS

In this section, two simulations and a comparison are carried out to validate our proposed algorithms.

Vertex	Before Operation			After Operation			
	$\operatorname{Hierarchy}(h\left(\cdot ight))$	Parents	Children	Hierarchy $(h(\cdot))$	Parents	Children	
v_1, v_2, v_3, v_4	0	None	_	0	None	_	
v_5	1	v_1, v_3, v_4	v_6, v_7, v_8	None	None	None	
v_6	2	v_1, v_4, v_5	v_8, v_9	1	v_1, v_3, v_4	v_7, v_8, v_9	
v_7	2	v_1, v_2, v_5	None	2	v_1, v_2, v_6	None	
v_8	3	v_1, v_5, v_6	None	2	v_1, v_3, v_6	None	
v_9	3	v_3, v_4, v_6	None	2	v_3, v_4, v_6	None	

TABLE II

"None" represents no data; "-" represents omitting data.

TABLE III AVERAGE RUNTIME WITH NETWORK SIZE WHEN DIFFERENT FRAMEWORK GROWING STRATEGY IS IMPOSED.

Average Runtime (s) Method Number of Agents	Xiao's [11]	Ours
5 Agents	0.8886	0.0010822
6 Agents	27.9123	0.0013588
7 Agents	755.8002	0.001694
8 Agents	NA	0.00217
20 Agents	NA	0.01111
50 Agents	NA	0.06963
100 Agents	NA	0.27694
200 Agents	NA	1.30814

A. Simulation 1: Growing Framework for Affine Formations

By using Algorithm 1, we present simulations to grow an eligible affine formation framework with different number of agents, as shown in Fig. 7. An original framework with four vertices is given in Fig. 7(a), then a series of vertices with random positions are considered to be added to the framework. As described in Fig. 7(b)-7(d), frameworks with different swarm size are generated while maintaining universal rigidity and affine localizability, which implying the effectiveness of the proposed distributed vertex addition strategy. We further demonstrate the rigidity of constructing affine frameworks by comparative simulations with [11], where a MISDP problem is established and an optimization-based topology design scheme is designed. We use two methods to grow affine formation frameworks with the same number of agents. The programs run on a laptop with AMD Ryzen 7 and 16 GB memory. Each simulation is repeated ten times and the average runtime is recorded in Table. III. It is obvious that Xiao's method takes much more time than ours, due to the fact that the number of variables that need to be optimized in the MISDP algorithm increases rapidly with the number of vertices, resulting in a sharp increase in runtime. As a comparison, our method only focuses on several neighbors near the newly added vertex, bringing about a significant reduction in computational cost, which reveals the potential of our method in onboard applications in robot swarms.



Fig. 7. An example to grow affine formation frameworks using Algorithm 1. (a) The original framework with 4 agents. (b) The obtained framework after adding 50 agents. (c) The framework after adding after adding 100 agents. (d) The framework after adding after adding 200 agents.

B. Simulation 2: Edge Deletion & Vertex Addition and Deletion

In this subsection, a scenario is designed to validate the effectiveness of our proposed agent-level framework construction algorithms, namely the proposed vertex addition, edge deletion and vertex deletion strategies in section III. We couple the affine formation framework construction strategies with the formation tracking control law proposed in [5], with the former providing the nominal framework for the latter. In the scenario, we consider a group of fixed-wing unmanned aerial vehicles (UAVs) modeled by unicycles moving in two-dimensional space while tracking moving leaders and achieving affine transformations. The convergence of affine formation tracking errors can serve as a powerful evidence to prove the effectiveness of our framework construction algorithms. Moreover, the adopted control scheme [5] is just an example and can be replaced by other similar affine formation control laws.

We consider a group of six agents with an original framework shown in Fig. 8(a), where the first three red vertices are regarded as leaders. Followers are driven to track leaders while achieving affine transformations. During the maneuver, agents adjust the topological connection between each other, as shown



Fig. 8. An example of edge deletion, vertex addition and deletion. (a) The original framework. (b) The obtained framework after deleting e_{23} and e_{46} . (c) The framework after adding two vertices, denoted by v_7 and v_8 . (d) The topology after three vertices deletion.



Fig. 9. The trajectory of fixed-wing UAVs.

in Fig. 8(b). Then, two additional agents join the formation (Fig. 8(c)) following Algorithm 1. After a period of formation flight, three agents leave the formation to perform other tasks, and the framework is reshaped as shown in Fig. 8(d). Fig. 9 depicts the trajectory of UAVs and Fig. 10 shows the evolution of formation tracking errors $||p_i - p_i^*||$, where p_i represents the position of the *i*-th UAV and p_i^* represents the desired one. It can be seen that the target formations are performed to pass through various environments, implying that the obtained framework is universally rigid and affine localizable.

Based on the simulation results, it is evident that our proposed distributed framework construction algorithms are highly effective to generate and reconstruct affine formation frameworks, demonstrating the potential for extensive applications in robot swarms.

VI. CONCLUSION

This paper have addressed the issue of constructing affine formation frameworks in a distributed manner in three scenarios, i.e., vertex addition, edge deletion and vertex deletion.



Fig. 10. The evolution of formation tracking errors for fixed-wing UAVs in Scenario 1.

We have designed strategies to construct frameworks with universal rigidity and affine localizability from the geometric perspective based on the structure of weighted graphs used to describe topology of affine formations. Naturally, comprehensive theoretical analysis has been provided to demonstrate the effectiveness, and simulations have been presented to verify the compatibility with distributed affine formation control proposals. Moreover, the comparative simulation has demonstrated the rapidity in constructing large-scale affine formations. Our approach relaxes the requirements for global information and computing resources, making it more easy to implement for robot swarms to perform various tasks. One future research direction is to extend distributed framework construction algorithms to directed graphs.

APPENDIX I THE PROOF OF THEOREM 1

To complete the proof of Theorem 1, the following lemmas are introduced for the analysis presented in the sequel.

Lemma 4: [17] Given PSD matrices $X \in \mathbb{R}^{n \times n}$ and $Y \in \mathbb{R}^{n \times n}$, let Z = X + Y. Then for any nonzero vector $\varepsilon \in \mathbb{R}^n$, $\varepsilon \in \text{null}(Z)$ if and only if $\varepsilon \in \text{null}(X)$ and $\varepsilon \in \text{null}(Y)$.

Lemma 5 (Rank-Nullity Theorem [29]): If there is a matrix A with x rows and y columns over a field, then rank(A) + nullity(A) = y, where nullity(A) means the nullity of the matrix A, that is, the dimension of the kernel of A.

Lemma 6 ([30]): For any symmetric matrix M of the form

$$M = \left[\begin{array}{cc} A & B \\ B^T & C \end{array} \right], \tag{28}$$

if C is invertible then the following properties hold:

(1) $M \succ 0 \iff C \succ 0$ and $A - BC^{-1}B^T \succ 0$.

(2) If $C \succ 0$, then $M \succeq 0 \iff A - BC^{-1}B^T \succeq 0$.

Then, let us complete the proof of Theorem 1.

Proof: Fist of all, with a positive s, we can prove that $\varpi_{uu} = s\phi_4^2$ in Eq. (8) is positive. Specifically, it is equivalent to prove $\phi_4 \neq 0$ because of s > 0. Suppose $\phi_4 = 0$, we get the following equations based on eq.(7),

$$\phi_1 + \phi_2 + \phi_3 = 0, \ \phi_1 (\mathbf{p}_i - \mathbf{p}_k) + \phi_2 (\mathbf{p}_j - \mathbf{p}_k) = \mathbf{0}.$$
 (29)

Since $(\mathcal{G}, \mathbf{p})$ is in general position, three vertices v_i , v_j , v_k are non-collinear. The vectors $(\mathbf{p}_i - \mathbf{p}_k)$ and $(\mathbf{p}_j - \mathbf{p}_k)$ are linearly independent, implying $\phi_1 = \phi_2 = 0$. Then, $\phi_3 = 0$ holds. Consequently, $\phi = \mathbf{0}$ is obtained, which contradicts

with the fact that ϕ is a nonzero vector. Therefore, we have $\phi_4 \neq 0$ so that ϖ_{uu} is positive when s > 0.

Then, in the newly obtained framework (\mathcal{G}^+, p^+) , we have $\mathcal{V}_l^+ = \mathcal{V}_l, \mathcal{V}_f^+ := \mathcal{V}_f \cup \{v_u\}$ and $\Omega^+ := \begin{bmatrix} \Omega_{ll}^+ & \Omega_{lf}^+ \\ \Omega_{fl}^+ & \Omega_{ff}^+ \end{bmatrix}$. Combining Lemmas 1 and 3, to demonstrate (\mathcal{G}^+, p^+) is universally rigid and affinely localizable, it is equivalent to prove the following conditions: (1) Ω^+ is PSD; (2) rank $(\Omega^+) = n-2$; (3) the block matrix Ω_{ff}^+ is positive definite. These are proven one by one as follows.

- (1) Based on Assumption 1 and eq. (8), we have Ω and Ω_u are both PSD, it is obvious that Ω_a and Ω_b in (9) are both PSD. Then according to Lemma 4, the matrix Ω^+ is PSD naturally.
- (2) Then, we analyze the rank of Ω^+ . According to Lemma 5, Let us start with the kernel of Ω^+ . Based on Lemma 2, we obtain the null space of Ω_a in eq. (9) is

$$\mathbb{S}_{a} = \operatorname{span}\left(\begin{bmatrix} \boldsymbol{p}_{1,1} \\ \vdots \\ \boldsymbol{p}_{i,1} \\ \boldsymbol{p}_{j,1} \\ \boldsymbol{p}_{k,1} \\ \boldsymbol{q}_{u,1} \end{bmatrix}, \begin{bmatrix} \boldsymbol{p}_{1,2} \\ \vdots \\ \boldsymbol{p}_{i,2} \\ \boldsymbol{p}_{j,2} \\ \boldsymbol{p}_{k,2} \\ \boldsymbol{q}_{u,2} \end{bmatrix}, \begin{bmatrix} 1 \\ \vdots \\ 1 \\ 1 \\ \vdots \\ 1 \\ \vdots \\ c_{u} \end{bmatrix} \right)$$
(30)
$$\triangleq \operatorname{span}\left(\boldsymbol{n}_{1}^{a}, \ \boldsymbol{n}_{2}^{a}, \ \boldsymbol{n}_{3}^{a} \right),$$

where $p_{i,1}$ is the first component of p_i $(i \in \{1, \dots, n\})$, and $p_{i,2}$ is defined analogously. $q_{u,\sim}$ and c_u are any arbitrary scalars. Similarly, the null space of Ω_b is

$$S_{b} = \operatorname{span} \left(\begin{bmatrix} q_{1,1} \\ \vdots \\ q_{(i-1)\cdot 1} \\ p_{i,1} \\ p_{j,1} \\ p_{k,1} \\ p_{u,1} \end{bmatrix}, \begin{bmatrix} q_{1,2} \\ \vdots \\ q_{(i-1)\cdot 2} \\ p_{i,2} \\ p_{j,2} \\ p_{k,2} \\ p_{u,2} \end{bmatrix}, \begin{bmatrix} c_{1} \\ \vdots \\ c_{i} \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \right)$$
$$\triangleq \operatorname{span} \left(\boldsymbol{n}_{1}^{b}, \boldsymbol{n}_{2}^{b}, \boldsymbol{n}_{3}^{b} \right), \qquad (21)$$

According to Lemma 4 and eq. (9), the null space of Ω^+ can be presented as null $(\Omega^+) = \mathbb{S}_a \cap \mathbb{S}_b$, whose nominal form is shown as

$$\alpha_1 n_1^a + \alpha_2 n_2^a + \alpha_3 n_3^a = \beta_1 n_1^b + \beta_2 n_2^b + \beta_3 n_3^b,$$
 (32)

where α_i and β_i (i = 1, 2, 3) are scalars that are not all zero. Because \mathbb{S}_a and \mathbb{S}_b share three identical elements, the following equation is deduced,

$$(\alpha_1 - \beta_1) \begin{bmatrix} \boldsymbol{p}_{i,1} \\ \boldsymbol{p}_{j,1} \\ \boldsymbol{p}_{k,1} \end{bmatrix} + (\alpha_2 - \beta_2) \begin{bmatrix} \boldsymbol{p}_{i,2} \\ \boldsymbol{p}_{j,2} \\ \boldsymbol{p}_{k,2} \end{bmatrix} + (\alpha_3 - \beta_3) \begin{bmatrix} 1 \\ 1 \\ 1 \\ (33) \end{bmatrix} = \mathbf{0},$$

which means

$$\underbrace{\begin{bmatrix} \boldsymbol{p}_i & \boldsymbol{p}_j & \boldsymbol{p}_k \\ 1 & 1 & 1 \end{bmatrix}}_{\triangleq \boldsymbol{P}_k} \begin{bmatrix} \alpha_1 - \beta_1 \\ \alpha_2 - \beta_2 \\ \alpha_3 - \beta_3 \end{bmatrix} = \boldsymbol{0}.$$
(34)

Because vertices v_i , v_j and v_k are not collinear, rank $(\mathbf{P}_k) = 3$ holds. Accordingly, we have $\alpha_i =$ $\beta_i, \forall i \in \{1, 2, 3\}$. Assume there is a vector $v \in \text{null}(\Omega^+)$, then we will show that v can be represented by $p_{i,1}^+, p_{i,2}^+$ $(i \in \{1, \dots, n+1\})$ and $\mathbf{1}_{n+1}$. According to Lemma 4, $v \in \text{null}(\Omega^+)$ implies $v \in \mathbb{S}_a$ and $v \in \mathbb{S}_b$. That is,

$$oldsymbol{v} = lpha_1 oldsymbol{n}_1^a + lpha_2 oldsymbol{n}_2^a + lpha_3 oldsymbol{n}_3^a = lpha_1 oldsymbol{n}_1^b + lpha_2 oldsymbol{n}_2^b + lpha_3 oldsymbol{n}_3^b$$

Thus, let the last entry in n_i^a equal the last entry in n_i^b , and the first *n* entries in n_i^b equal the first *n* entries in n_i^a . It follows that

$$\boldsymbol{v} = \alpha_{1} \begin{bmatrix} \boldsymbol{p}_{1,1} \\ \vdots \\ \boldsymbol{p}_{i,1} \\ \boldsymbol{p}_{j,1} \\ \boldsymbol{p}_{k,1} \\ \boldsymbol{p}_{u,1} \end{bmatrix} + \alpha_{2} \begin{bmatrix} \boldsymbol{p}_{1,2} \\ \vdots \\ \boldsymbol{p}_{i,2} \\ \boldsymbol{p}_{j,2} \\ \boldsymbol{p}_{k,2} \\ \boldsymbol{p}_{u,2} \end{bmatrix} + \alpha_{3} \begin{bmatrix} 1 \\ \vdots \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$
(35)

which means

null
$$(\mathbf{\Omega}^+) = \operatorname{col}\left(\underbrace{\left[\begin{array}{cccc} \mathbf{p}_1 & \cdots & \mathbf{p}_i & \mathbf{p}_j & \mathbf{p}_k & \mathbf{p}_u \\ 1 & \cdots & 1 & 1 & 1 & 1 \end{array}\right]^T}_{\triangleq \overline{P}(\mathbf{p}^+)}\right).$$

Since rank $(\bar{P}(p^+)) = 3$, we have nullity $(\Omega^+) = 3$ so that rank $(\Omega^+) = n + 1 - 3 = n - 2$. According to Lemma 1, the framework (\mathcal{G}^+, p^+) is universally rigid.

- (3) Now consider the block matrix Ω_{ff}^+ . The added vertex v_u plays the role of a follower in (\mathcal{G}^+, p^+) . Depending on whether the selected the vertices v_i , v_j and v_k represent leaders or not, there are four situations in total as below.
 - All the vertices $v_i, v_j, v_k \in \mathcal{V}_l$. Based on Assumption 1, we get the block stress matrix Ω_{ff} associated with Ω is positive definite. If $v_i, v_j, v_k \in \mathcal{V}_l$, then Ω_{ff}^+ can be described by $\Omega_{ff}^+ = \left[-\frac{\Omega_{ff}}{\mathbf{0}_{1 \times n_f}} - \frac{\mathbf{0}_{n_f \times 1}}{\varpi_{uu}} - \right]$. Since $\varpi_{uu} > 0$ when $s > 0, \Omega_{ff}^+ \succ 0$ holds naturally.
 - Two of the vertices v_i , v_j , v_k belong to \mathcal{V}_l , and one belongs to \mathcal{V}_f .

Without losing generality, assume $v_i \in \mathcal{V}_f$ and $v_j, v_k \in \mathcal{V}_l$ so that Ω_{ff} can be divided into $\Omega_{ff} = \begin{bmatrix} \Omega_{ff}^{P1} & \Omega_{ff}^{P2} \\ \Omega_{ff}^{P1} & \Omega_{ff}^{P2} \end{bmatrix} \succ 0$. Based on Lemma 6, we

have $\Omega_{ii} > 0$ and $\Omega_{ff}^{P1} - \frac{1}{\Omega_{ii}} \Omega_{ff}^{P2} \left(\Omega_{ff}^{P2} \right)^T \succ 0$. After adding a new vertex v_u , based on eq. (9), the block stress matrix Ω_{ff}^+ associated with the augmented framework $(\mathcal{G}^+, \mathbf{p}^+)$ is described as follows.

$$\boldsymbol{\Omega}_{ff}^{+} = \begin{bmatrix} \boldsymbol{\Omega}_{ff}^{P1} & \boldsymbol{\Omega}_{ff}^{P2} & \boldsymbol{0}_{(n_{f}-1)\times 1} \\ \boldsymbol{\Omega}_{ff}^{P2} & \boldsymbol{\Omega}_{ii} + \frac{\boldsymbol{\varpi}_{iu}}{\boldsymbol{\varpi}_{uu}} & -\boldsymbol{\varpi}_{iu} \\ \boldsymbol{0}_{1\times(n_{f}-1)} & -\boldsymbol{\varpi}_{iu} & \boldsymbol{\varpi}_{uu} \end{bmatrix}.$$
(36)

Based on Lemma 6, we have

$$\begin{bmatrix} \Omega_{ii} + \frac{\varpi_{iu}^2}{\varpi_{uu}} & -\varpi_{iu} \\ -\varpi_{iu} & \varpi_{uu} \end{bmatrix} \succ 0.$$

Hence, the following equation is deduced.

$$\boldsymbol{\Omega}_{ff}^{P1} - \begin{bmatrix} \boldsymbol{\Omega}_{ff}^{P2} & \boldsymbol{0}_{(n_f-1)\times 1} \end{bmatrix} \cdot \begin{bmatrix} \Omega_{ii} + \frac{\varpi_{iu}^2}{\varpi_{uu}} & -\varpi_{iu} \\ -\varpi_{iu} & \varpi_{uu} \end{bmatrix}^{-1} \begin{bmatrix} \left(\boldsymbol{\Omega}_{ff}^{P2} \right)^T \\ \boldsymbol{0}_{1\times (n_f-1)} \end{bmatrix}$$
(37)
$$=\boldsymbol{\Omega}_{ff}^{P1} - \frac{1}{\Omega_{ii}} \boldsymbol{\Omega}_{ff}^{P2} \left(\boldsymbol{\Omega}_{ff}^{P2} \right)^T \succ 0.$$

Thus, Ω_{ff}^+ in (36) is positive definite. • Two of the vertices v_i , v_j , v_k belong to \mathcal{V}_f , and one belongs to \mathcal{V}_l .

In this situation, we apply Lemma 6 repeatedly in a similar way to prove that the symmetric matrix Ω_{ff}^+ is positive definite. Without losing its generality, assume $v_i, v_j \in \mathcal{V}_f$ and $v_k \in \mathcal{V}_l$, and Ω_{ff} can be presented as $\Omega_{ff} = \begin{bmatrix} -\frac{\Omega_{ff}^{P1}}{(\Omega_{ff}^{P2})^{-1} & \Omega_{ij}^{P2} & \Omega_{ij} \\ (\Omega_{ff}^{P2})^{-1} & \Omega_{ij} & \Omega_{jj} \end{bmatrix}$. Obviously, due to $\Omega_{ff} \succ 0$, we have $\begin{bmatrix} \Omega_{ii} & \Omega_{ij} \\ \Omega_{ij} & \Omega_{jj} \end{bmatrix} \succ 0$ and

$$\boldsymbol{\Omega}_{ff}^{P1} - \boldsymbol{\Omega}_{ff}^{P2} \begin{bmatrix} \Omega_{ii} & \Omega_{ij} \\ \Omega_{ij} & \Omega_{jj} \end{bmatrix} \quad \left(\boldsymbol{\Omega}_{ff}^{P2}\right)^T \succ 0. \quad (38)$$

Based on eq. (9), the block stress matrix Ω_{ff}^+ is established as follows.

$$\boldsymbol{\Omega}_{ff}^{+} = \begin{bmatrix} \boldsymbol{\Omega}_{ff}^{P1} & \boldsymbol{\Omega}_{ff}^{P2} & \boldsymbol{0}_{(n_{f}-2)\times 1} \\ \boldsymbol{\Omega}_{ff}^{P2} & \boldsymbol{\Pi}_{ff}^{T} & \boldsymbol{\Pi}_{ff}^{P3+} \\ \boldsymbol{0}_{1\times(n_{f}-2)} & \boldsymbol{\Omega}_{ff}^{P3+} \end{bmatrix},$$
(39)

where

$$\mathbf{\Omega}_{ff}^{P3+} \triangleq \begin{bmatrix} \Omega_{ii} + \frac{\varpi_{iu}^2}{\varpi_{uu}} & \Omega_{ij} + \frac{\varpi_{iu}\varpi_{ju}}{\varpi_{uu}} & -\varpi_{iu} \\ \Omega_{ij} + \frac{\varpi_{iu}\varpi_{ju}}{\varpi_{uu}} & \Omega_{jj} + \frac{\varpi_{ju}^2}{\varpi_{uu}} & -\varpi_{ju} \\ -\varpi_{iu} & -\varpi_{ju} & \varpi_{uu} \end{bmatrix}$$

Since $\varpi_{uu} > 0$ and

$$\begin{bmatrix} \Omega_{ii} + \frac{\varpi_{iu}^2}{\varpi_{uu}} & \Omega_{ij} + \frac{\varpi_{iu}\varpi_{ju}}{\varpi_{uu}} \\ \Omega_{ij} + \frac{\varpi_{iu}\varpi_{ju}}{\varpi_{uu}} & \Omega_{jj} + \frac{\varpi_{ju}^2}{\varpi_{uu}} \end{bmatrix} - \frac{1}{\varpi_{uu}} \\ \begin{bmatrix} -\varpi_{iu} \\ -\varpi_{ju} \end{bmatrix} \begin{bmatrix} -\varpi_{iu} & -\varpi_{ju} \end{bmatrix} = \begin{bmatrix} \Omega_{ii} & \Omega_{ij} \\ \Omega_{ij} & \Omega_{jj} \end{bmatrix} \succ 0,$$

we have $\Omega_{ff}^{P3+} \succ 0$. It follows from eq. (39) and eq. (38) that

$$\boldsymbol{\Omega}_{ff}^{P1} - \begin{bmatrix} \boldsymbol{\Omega}_{ff}^{P2} & \boldsymbol{0} \end{bmatrix} \begin{pmatrix} \boldsymbol{\Omega}_{ff}^{P3+} \end{pmatrix}^{-1} \begin{bmatrix} \begin{pmatrix} \boldsymbol{\Omega}_{ff}^{P2} \end{pmatrix}^T \\ \boldsymbol{0}_{1\times(n_f-2)} \end{bmatrix} \succ 0.$$
(40)

By applying Lemma 6 again, we have $\Omega_{ff}^+ \succ 0$. • All the vertices $v_i, v_j, v_k \in \mathcal{V}_f$.

The analysis shares the same idea as in the above two cases. The block stress matrix Ω_{ff} is positive definite and can be divided into

$$\boldsymbol{\Omega}_{ff} = \begin{bmatrix} -\boldsymbol{\Omega}_{ff}^{P1} & \boldsymbol{\Omega}_{ii}^{P2} & \boldsymbol{\Omega}_{ij}^{P2} \\ \left(\boldsymbol{\Omega}_{ff}^{P2} \right)^T & \boldsymbol{\Omega}_{ij} & \boldsymbol{\Omega}_{ij} & \boldsymbol{\Omega}_{jk} \\ & \boldsymbol{\Omega}_{ij} & \boldsymbol{\Omega}_{jj} & \boldsymbol{\Omega}_{jk} \\ & \boldsymbol{\Omega}_{ik} & \boldsymbol{\Omega}_{jk} & \boldsymbol{\Omega}_{kk} \\ & \boldsymbol{\Pi}_{ff} & \boldsymbol{\Pi}_{ff} \end{bmatrix} \succ \boldsymbol{0}.$$

After adding a new vertex v_u as a follower, we have

$$\boldsymbol{\Omega}_{ff}^{+} = \begin{bmatrix} \boldsymbol{\Omega}_{ff}^{P1} & \boldsymbol{\Omega}_{ff}^{P2} & \boldsymbol{0}_{(n_{f}-3)\times1} \\ \boldsymbol{\Omega}_{ff}^{P2} & \boldsymbol{\Omega}_{ff}^{P2} & \boldsymbol{\Omega}_{ff}^{P3+} \\ \boldsymbol{0}_{1\times(n_{f}-3)} & \boldsymbol{\Omega}_{ff}^{P3+} \end{bmatrix}$$

where $\mathbf{\Omega}_{ff}^{P3+} = \begin{bmatrix} \mathbf{\Omega}_{ff}^{P3} & \mathbf{0} \\ \mathbf{0} & 0 \end{bmatrix} + \mathbf{\Omega}_{u}$. By applying Lemma 6 again, we can prove the matrix Ω_{ff}^+ is

also positive definite.

According to Lemma 2, the stress matrix Ω^+ associated with the augmented framework (\mathcal{G}^+, p^+) follows that

$$\left(\boldsymbol{\Omega}^{+}\otimes\boldsymbol{I}_{d}\right)\boldsymbol{p}^{+}=\left[\begin{array}{cc}\bar{\boldsymbol{\Omega}}_{ll}^{+}&\bar{\boldsymbol{\Omega}}_{lf}^{+}\\\bar{\boldsymbol{\Omega}}_{fl}^{+}&\bar{\boldsymbol{\Omega}}_{ff}^{+}\end{array}\right]\left[\begin{array}{cc}\boldsymbol{p}_{l}^{+}\\\boldsymbol{p}_{f}^{+}\end{array}\right]=\boldsymbol{0}.$$
 (41)

Accordingly, we have $\bar{\Omega}_{fl}^+ p_l^+ + \bar{\Omega}_{ff}^+ p_f^+ = 0$. We clarify that the augmented framework (\mathcal{G}^+, p^+) is affinely localized by the selected leaders since $\Omega_{ff}^+ \succ 0$, which leads

to
$$\boldsymbol{p}_f^+ = -\left(\bar{\boldsymbol{\Omega}}_{ff}^+\right)^{-1} \bar{\boldsymbol{\Omega}}_{fl}^+ \boldsymbol{p}_l^+.$$

With all the discussions above, the augmented framework $(\mathcal{G}^+, \boldsymbol{p}^+)$ obtained from adding a new v_u and three weighted edges meets the requirements for universal rigidity and affine localizability.

REFERENCES

- [1] Z. Lin, L. Wang, Z. Chen, M. Fu, and Z. Han, "Necessary and sufficient graphical conditions for affine formation control," IEEE Transactions on Automatic Control, vol. 61, no. 10, pp. 2877-2891, 2016.
- [2] S. Zhao, "Affine formation maneuver control of multiagent systems," IEEE Transactions on Automatic Control, vol. 63, no. 12, pp. 4140-4155, 2018.
- [3] H. Wang and J. Dong, "Neural network-based hierarchical fault-tolerant affine formation control for heterogeneous nonlinear multi-agent systems," IEEE Transactions on Intelligent Transportation Systems, pp. 1-9. 2023.
- [4] Y. Lin, Z. Lin, Z. Sun, and B. D. O. Anderson, "A unified approach for finite-time global stabilization of affine, rigid, and translational formation," IEEE Transactions on Automatic Control, vol. 67, no. 4, pp. 1869-1881, 2022.
- [5] H. Li, H. Chen, and X. Wang, "Affine formation tracking control of unmanned aerial vehicles," Frontiers of Information Technology & Electronic Engineering, vol. 23, no. 7, pp. 909-919, 2022.
- [6] Y. Xu, D. Luo, D. Li, Y. You, and H. Duan, "Target-enclosing affine formation control of two-layer networked spacecraft with collision avoidance," Chinese Journal of Aeronautics, vol. 32, no. 12, pp. 2679-2693, 2019.
- [7] J. Yang, F. Xiao, and T. Chen, "Formation tracking of nonholonomic systems on the special euclidean group under fixed and switching topologies: An affine formation strategy," SIAM Journal on Control and Optimization, vol. 59, no. 4, pp. 2850-2874, 2021.
- [8] C. Zhu, B. Huang, Y. Lu, X. Li, and Y. Su, "Distributed affine formation maneuver control of autonomous surface vehicles with event-triggered data transmission mechanism," IEEE Transactions on Control Systems Technology, vol. 31, no. 3, pp. 1006-1017, 2023.
- H. Chen, X. Wang, Z. Liu, Z. Li, and L. Shen, "Survivable networks for [9] consensus," IEEE Transactions on Control of Network Systems, vol. 9, no. 2, pp. 588-600, 2022.

- [10] Y. Zhang, S. Oğuz, S. Wang, E. Garone, X. Wang, M. Dorigo, and M. K. Heinrich, "Self-reconfigurable hierarchical frameworks for formation control of robot swarms," *IEEE Transactions on Cybernetics*, pp. 1–14, 2023.
- [11] F. Xiao, Q. Yang, X. Zhao, and H. Fang, "A framework for optimized topology design and leader selection in affine formation control," *IEEE Robotics and Automation Letters*, vol. 7, no. 4, pp. 8627–8634, 2022.
- [12] T.-S. Tay and W. Whiteley, "Generating isostatic frameworks," *Structural Topology*, pp. 21–69, 1985.
- [13] R. Connelly, "Generic global rigidity," Discrete & Computational Geometry, vol. 33, pp. 549–563, 2005.
- [14] A. Gasparri, R. K. Williams, A. Priolo, and G. S. Sukhatme, "Decentralized and parallel constructions for optimally rigid graphs in R^2 ," *IEEE Transactions on Mobile Computing*, vol. 14, no. 11, pp. 2216–2228, 2015.
- [15] T. Eren, "Formation shape control based on bearing rigidity," *International Journal of Control*, vol. 85, no. 9, pp. 1361–1379, 2012.
- [16] M. H. Trinh, S. Zhao, Z. Sun, D. Zelazo, B. D. O. Anderson, and H.-S. Ahn, "Bearing-based formation control of a group of agents with leader-first follower structure," *IEEE Transactions on Automatic Control*, vol. 64, no. 2, pp. 598–613, 2019.
- [17] Q. Yang, M. Cao, and B. D. O. Anderson, "Growing super stable tensegrity frameworks," *IEEE Transactions on Cybernetics*, vol. 49, no. 7, pp. 2524–2535, 2019.
- [18] Q. Yang, M. Cao, H. Fang, and J. Chen, "Constructing universally rigid tensegrity frameworks with application in multiagent formation control," *IEEE Transactions on Automatic Control*, vol. 64, no. 1, pp. 381–388, 2019.
- [19] R. K. Ramachandran, P. Pierpaoli, M. Egerstedt, and G. S. Sukhatme, "Resilient monitoring in heterogeneous multi-robot systems through network reconfiguration," *IEEE Transactions on Robotics*, vol. 38, no. 1, pp. 126–138, 2022.
- [20] Q. Feng, X. Hai, B. Sun, Y. Ren, Z. Wang, D. Yang, Y. Hu, and R. Feng, "Resilience optimization for multi-uav formation reconfiguration via enhanced pigeon-inspired optimization," *Chinese Journal of Aeronautics*, vol. 35, no. 1, pp. 110–123, 2022.
- [21] D. Groß and O. Stursberg, "Optimized distributed control and network topology design for interconnected systems," in 2011 50th IEEE Conference on Decision and Control and European Control Conference, 2011, pp. 8112–8117.
- [22] P. Mukherjee, M. Santilli, A. Gasparri, and R. K. Williams, "Optimal topology selection for stable coordination of asymmetrically interacting multi-robot systems," in 2020 IEEE International Conference on Robotics and Automation (ICRA), 2020, pp. 6668–6674.
- [23] J. Li, Z. Ning, S. He, C.-H. Lee, and S. Zhao, "Three-dimensional bearing-only target following via observability-enhanced helical guidance," *IEEE Transactions on Robotics*, vol. 39, no. 2, pp. 1509–1526, 2023.
- [24] Z. Liu, X. Wang, L. Shen, S. Zhao, Y. Cong, J. Li, D. Yin, S. Jia, and X. Xiang, "Mission-oriented miniature fixed-wing uav swarms: A multilayered and distributed architecture," *IEEE Transactions on Systems, Man, and Cybernetics: Systems*, vol. 52, no. 3, pp. 1588–1602, 2022.
- [25] A. Alfakih and V.-H. Nguyen, "On affine motions and universal rigidity of tensegrity frameworks," *Linear Algebra and its Applications*, vol. 439, no. 1, pp. 3134–3147, 2013.
- [26] D. Li, G. Ma, Y. Xu, W. He, and S. S. Ge, "Layered affine formation control of networked uncertain systems: A fully distributed approach over directed graphs," *IEEE Transactions on Cybernetics*, vol. 51, no. 12, pp. 6119–6130, 2021.
- [27] Y. Zheng, L. Zhang, B. Huang, and Y. Su, "Distributed event-triggered affine formation control for multiple underactuated marine surface vehicles," *Ocean Engineering*, vol. 265, p. 112607, 2022.
- [28] A. Alfakih, N. Taheri, and Y. Ye, "On stress matrices of (d+1)-lateration frameworks in general position," *Mathematical Programming*, vol. 137, no. 1, pp. 1–17, 2013.
- [29] S. Lang, Introduction to Linear Algebra. New York, USA: Springer New York, NY, 1986.
- [30] F. Zhang, *The Schur Complement and Its Applications*. New York, USA: Springer New York, NY, 2006.