Tim Ophelders \square

Department of Information and Computing Sciences, Utrecht University, the Netherlands Department of Mathematics and Computer Science, TU Eindhoven, the Netherlands

Anna Schenfisch \square

Department of Mathematics and Computer Science, TU Eindhoven, the Netherlands

— Abstract

Simplicial complexes can be used to represent high-dimensional shapes, and are suitable for computational purposes. They generalize point clouds in that they capture connectivity by means of edges, triangles, and higher-dimensional analogues, collectively called simplices. Simplicial complexes arising from real-world settings may not be directly observable. Hence, for an unknown simplicial complex in Euclidean space, we want to efficiently reconstruct it by querying local structure.

In particular, we are interested in queries for the *indegree* of a simplex σ in some direction: the number of cofacets of σ contained in some halfspace "below" σ . Fasy et al. proposed a method that, given the vertex set of a simplicial complex, uses indegree queries to reconstruct the set of edges. In particular, they use a sweep algorithm through the vertex set, identifying edges adjacent to and above each vertex in the sweeping order. The algorithm relies on a natural but crucial property of the sweeping order: at a given vertex v, all edges adjacent to v contained in the halfspace below v have another endpoint that appeared earlier in the order.

The edge reconstruction algorithm does not immediately extend to higher-dimensional simplex reconstruction. In particular, it is not possible to sweep through a set of *i*-simplices in a fixed direction and maintain that all (i + 1)-cofacets of a given simplex σ that come below σ are known. We circumvent this by defining a *sweeping order* on a set of *i*-simplices, that additionally pairs each *i*-simplex σ with a direction perpendicular to σ . Analogous to Fasy et al., our order has the crucial property that, at any *i*-simplex σ paired with direction *s*, each (i + 1)-dimensional coface of σ that lies in the halfspace below σ with respect to the direction *s* has an *i*-dimensional face that appeared earlier in the order. We show how to compute such an order and use it to extend the edge reconstruction algorithm of Fasy et al. to simplicial complex reconstruction. Our algorithm can reconstruct arbitrary embedded simplicial complexes.

2012 ACM Subject Classification Theory of computation \rightarrow Computational Geometry

Keywords and phrases Simplicial complex, reconstruction, topological descriptors, sweep algorithm

Funding *Tim Ophelders*: Partially supported by the Dutch Research Council (NWO) under project no. VI.Veni.212.260.

Anna Schenfisch: Supported by the Dutch Research Council (NWO) under project no. P21-13.

1 Introduction

Since their introduction in [24] to compute or count intersections of geometric objects, sweep algorithms have become ubiquitous in computational geometry, see, e.g., [7, 27, 14, 13, 20, 8, 28, 25]. For problems in \mathbb{R}^d , such algorithms typically use a (d-1)-dimensional hyperplane as their *sweeping object*, swept in some constant direction through the space. Sweep algorithms maintain solutions to subproblems associated with the region that was swept thus far. Throughout a sweep, there is a discrete set of locations, called *sweep events*, where the solution to the subproblem may change combinatorially. For efficient sweep algorithms, this solution can be derived efficiently from solutions to the previous subproblems.

We are particularly motivated by the sweep algorithm of [11], which sweeps up through a

given set of vertices to reconstruct the edge set. At each vertex in the sweep, they use a query called *indegree* to count edges adjacent to a vertex in a particular halfspace. Using indegree and a radial order of candidate edges around v, they identify all edges adjacent to v in the upper halfspace. Central to this process is the ability to maintain that all edges with a vertex below the sweep height have already been identified. Indegree has a higher-dimensional generalization, namely, the count of cofacets of a given simplex in a particular halfspace.

While we generally consider indegree as a given function in this paper, it is possible to compute indegree from other geometric or topological information. In particular, indegree is computed in [11, 10, 3] using verbose persistence diagrams, a tool that records homological changes throughout a directional filtration. Indeed, simplicial complex reconstruction has close ties to the study of faithful discretizations of *directional transforms*, which are sets of (topological) descriptors corresponding to height/lower-star filtrations in various directions that completely characterize the shape.

There were two main challenges preventing [11] from extending their results to higherdimensional simplex reconstruction. First, at the time of [11], there was no method to compute higher-dimensional indegree using topological descriptors. However, [10] developed exactly this tool, using an inclusion-exclusion type argument. But the second challenge remained; a straightforward higher-dimensional generalization of [11] was not possible. As the authors note, a height-based sweep of the vertex set is no longer feasible, since "radially ordering higher dimensional simplices [around a common vertex] is not well-defined, and this issue prevents the methods [...] from being immediately transferable [to general simplicial complexes]." If the sweeping plane moves in a direction perpendicular to a given *i*-simplex σ (so the sweeping plane eventually contains σ), it *is* possible to radially order the cofacets of σ around σ . However, in general, no single direction is perpendicular to all simplices of some fixed positive dimension. We could try moving the sweeping plane non-linearly to contain each *i*-simplex, but in what order? An arbitrary order does not guarantee the ability to maintain that all (*i* + 1)-simplices below the sweeping plane have already been identified.

In this paper, we show how to compute a *sweeping order*, or a way to move the sweeping plane so that all the properties necessary for the algorithm in [11] are maintained. This sweeping order is the missing ingredient needed for us to then describe the generalization of edge reconstruction to higher-dimensional simplex reconstruction.

Outline In Section 2, we provide a brief overview of foundational topics and important notation. We define our main tool in Section 3—a *sweeping order* of a set of *i*-simplices—and show how such an order can be computed. The edge reconstruction method of [11] uses a search through radially ordered *candidate edges* adjacent to a given vertex. We generalize the notion of candidates to cofacets of a higher-dimensional simplex in Section 4, and show how to order them radially around such a simplex in Section 5. We determine which candidates are actually in the to-be-reconstructed K using a radial search that uses *indegree queries*, which count the cofacets of a simplex in a particular halfspace. Finally, in Section 6, we extend the edge reconstruction methods of [11] to reconstruct the (i + 1)-simplices of K, given all lower-dimensional simplices. Our entire complex K can then be reconstructed iteratively from just its vertex set and indegree queries.

2 Preliminaries

In this section, we define our most extensively used terms, and refer the reader to [9, 16] for further information. We start simplices and simplicial complexes.



Figure 1 Examples of complexes in \mathbb{R}^2 that are (a) both locally injective and embedded, (b) locally injective but not embedded, and (c) neither locally injective nor embedded.

▶ **Definition 1** (Simplex). An abstract *i*-simplex σ is a set of i + 1 elements, called vertices. The dimension of σ is then *i*, denoted dim(σ). If σ and τ are abstract simplices and $\sigma \subseteq \tau$, we call σ a face of τ and τ a coface of σ . If additionally dim(σ) = dim(τ) - 1, we call σ a facet of τ and τ a cofacet of σ . An *i*-simplex in \mathbb{R}^d is an abstract *i*-simplex where each vertex maps to a distinct point in \mathbb{R}^d . We geometrically interpret a simplex as the convex hull of these points.

▶ **Definition 2** (Simplicial Complex). An abstract simplicial complex K is a set of abstract simplices, such that if $\sigma \in K$ and $\rho \subseteq \sigma$, then $\rho \in K$. A simplicial complex in \mathbb{R}^d consists of an abstract simplicial complex, where each vertex is mapped injectively to a point in \mathbb{R}^d . Geometrically, we think of the simplicial complex as the union of convex hulls of its simplices.

Our work always interprets simplicial complexes geometrically, so going forward, we may be somewhat less precise with terminology. Definition 3 distinguishes various types of simplicial complexes based on how their simplices intersect, see also Figure 1.

▶ **Definition 3** (Embedded and locally injective). We call a pair of simplices σ and σ' an injective pair if the intersection of their convex hulls is either empty or the convex hull of a common face (i.e., a simplex ρ that is a face of σ and σ'). We call a simplicial complex in \mathbb{R}^d embedded if all pairs of simplices are injective pairs. We call a simplicial complex in \mathbb{R}^d locally injective if all pairs of simplices that have a common face are injective pairs.

Let K be a simplicial complex in \mathbb{R}^d . Let dim(K) denote the maximum dimension over all simplices in K. We denote the *i*-skeleton of K (the subset of simplices with dimension at most *i*) by K_i and the number of *i*-simplices by n_i . For an *i*-simplex σ , let $cof(\sigma) \subseteq K_{i+1}$ denote the set of all cofacets of σ . For a simplex $\sigma \in K$, we denote its affine hull by aff (σ) .

The set of all unit vectors in \mathbb{R}^d is parameterized by the unit (d-1)-sphere, denoted \mathbb{S}^{d-1} , and a unit vector is called a *direction*. Denote by $\perp_{\sigma} \subseteq \mathbb{S}^{d-1}$ the set of directions perpendicular to σ and note that, if dim(aff(σ)) = $i' \leq d-1$, then \perp_{σ} is a (d-i'-1)-sphere. With respect to some $s \in \perp_{\sigma}$, all points in σ have the same height, which we refer to as the *s*-coordinate of σ . We may abuse dot product notation and write $s \cdot \sigma$ to denote this *s*-coordinate. For $s \in \perp_{\sigma}$, the set of *down-cofacets* of σ , denoted $\operatorname{cof}_s^<(\sigma) \subseteq \operatorname{cof}(\sigma)$, consists of the cofacets $\sigma \cup \{v\}$ for which $s \cdot v < s \cdot \sigma$, i.e. the vertex v lies in the open halfspace below σ with respect to direction s. Similarly, the set of *up-cofacets* of σ , denoted $\operatorname{cof}_s^\geq(\sigma)$, consists of those cofacets $\sigma \cup \{v\}$ for which $s \cdot v \geq s \cdot \sigma$. Regardless of our choice of $s \in \perp_{\sigma}$, we have $\operatorname{cof}_s^<(\sigma) \cup \operatorname{cof}_s^\geq(\sigma) = \operatorname{cof}(\sigma)$.

Conceptually, our algorithms involve ordering points by rotating a hyperplane around some central space. In the following definition, we ensure that we make a consistent choice of normal direction to associate with each point we encounter.

▶ Definition 4 (γ -Normal of p). Let $\gamma : [0, 2\pi) \to \mathbb{S}^1$ angularly parameterize the unit circle $\mathbb{S}^1 \subset \mathbb{R}^2$. Then for some point $p \in \mathbb{R}^2 \setminus \{(0, 0)\}$, consider the unique angle α such that the ray



Figure 2 $\gamma(\alpha)$ is the γ -normal of (a) p, (b) p relative to c, (c) p as well as $q \in \mathbb{R}^3$.

from (0,0) through $\gamma(\alpha - \pi/2 \mod 2\pi)$ passes through p. We refer to $\gamma(\alpha)$ as the γ -normal of p. We define the γ -normal of the origin to be $\gamma(0)$. For some (center) point $c \in \mathbb{R}^2$, we define the γ -normal of p relative to c to be the γ -normal of p - c. See Figure 2 (a-b).

In \mathbb{R}^d with $d \geq 2$, if $\gamma : [0, 2\pi) \to \mathbb{S}^{d-1}$ angularly parameterizes a unit-circle centered at the origin (i.e., a rotated copy of \mathbb{S}^1), we can for any point $q \in \mathbb{R}^d$ consider its orthogonal projection p onto the 2-dimensional linear subspace $\operatorname{span}(\gamma)$ that contains γ , and define the γ -normal of q as the γ -normal of p based on this subspace. See Figure 2 (c). Similarly, for points $q, c \in \mathbb{R}^d$, we can define the γ -normal of q relative to c as the γ -normal of q - c.

Now consider some simplex $\sigma \subset \mathbb{R}^d$ with $\dim(aff(\sigma)) < d-1$, let γ be some angularly parameterized circle of directions all perpendicular to σ , and let q be some point in \mathbb{R}^d . Then, the γ -normal of q relative to any point c in the $aff(\sigma)$ is the same, so we unambiguously define the γ -normal of q relative to σ to be the γ -normal of q relative to any such point c.

Finally, consider a simplex $\sigma \subset \mathbb{R}^d$ with $\dim(aff(\sigma)) = d - 1$. Let s be one of the two (antipodal) directions perpendicular to σ , and let $\gamma: [0, 2\pi) \to \mathbb{S}^{d-1}$ be an angularly parameterized circle of directions with $\gamma(0) = s$. For any point $p \in \mathbb{R}^d$, define the γ -normal of p relative to σ to be $\gamma(0)$ if the s-coordinate of p is at least that of σ , and $\gamma(\pi)$ otherwise.

If $v_1 \neq v_2$ are vertices such that $\operatorname{aff}(\sigma \cup \{v_1\}) = \operatorname{aff}(\sigma \cup \{v_2\})$, but v_1 and v_2 are on "opposite sides" of σ , and if the γ -normal of v_1 relative to σ is $\gamma(\alpha)$, then the normal for v_2 is $-\gamma(\alpha)$.

3 Sweeping Orders

A main goal of this paper is to show how to compute an order on simplices (along with a corresponding list of directions perpendicular to the simplices) in order to emulate properties characteristic of sweepline algorithms with discrete points or vertices as events.



Figure 3 Here, every edge *e* has other edges in the halfspaces on either side of *e*. This general higher-dimensional phenomenon contrasts the special zero-dimensional case; in every direction, we can find at least one vertex with no other vertices below it. However, for each dimension of simplex, there *do* exist simplices with no *cofacets* in a halfspace below it, which we will see is true in general.

As illustrated in Figure 3, we cannot guarantee that all *i*-simplices below a particular *i*-simplex in our sweep appeared earlier in the order, so we instead focus on cofacets of simplices in the sweep. More specifically, we introduce a sequence such that, for a given *i*-simplex σ and a corresponding direction $s \in \perp_{\sigma}$, all cofacets of σ contained in $\operatorname{cof}_{s}^{<}(\sigma)$ are cofacets of some prior *i*-simplex in the sequence. (Indeed, this is a property of the usual sweep through a geometric graph, where vertices are sweep events). This is formalized below.

▶ Definition 5 (Sweeping Order). A sweeping order of K_i is any sequence $SO_i := ((\sigma_j, s_j))_{j=1}^{n_i}$ that satisfies the following three properties.

- **1.** Each s_j is a direction perpendicular to σ_j .
- **2.** Each *i*-simplex of K appears exactly once in SO_i .
- **3.** For any *i*-simplex σ_j , any cofacet in $cof_{s_i}^{\leq}(\sigma_j)$ is also a cofacet of some σ_h , for h < j.

Note that if Property 3 holds, then there exists some initial (σ_1, s_1) , so that, with respect to s_1 , the simplex σ_1 has *no* cofacets in the halfspace below it.

3.1 Computing a Sweeping Order

In this section, we show how to compute a sweeping order SO_i of K_i . For K_0 , we simply pick an arbitrary direction s, order vertices by their s-coordinate (breaking ties arbitrarily), and output (v, s) for each vertex v in that order. See Algorithm 1.

To compute a sweeping order for K_i with i > 0, we assume that we already know a sweeping order SO_{i-1} of K_{i-1} (for example because we have computed SO_{i-1} recursively). Then SO_i is the sequence of pairs output by $ORDER(K_i, SO_{i-1})$ in Algorithm 1. We use the convention that ρ , σ , and τ represent a simplex of dimension i - 1, i, and i + 1, respectively. We consider the (i - 1)-simplices ρ in the order prescribed by SO_{i-1} . For ease of exposition, assume that dim(aff(ρ)) $\leq d - 2$. We radially order the cofacets σ of ρ that have not yet been output. Specifically, if SO_{i-1} pairs ρ with direction s, then the radial order of its cofacets is based on a parameterized circle $\gamma_{\rho} \colon [0, 2\pi) \to \mathbb{S}^{d-1}$ of directions, rotating around ρ , starting at s. Each direction $\gamma_{\rho}(\alpha)$ corresponds to the unique halfspace whose boundary contains ρ , and whose exterior normal points in direction $\gamma_{\rho}(\alpha)$. For each cofacet $\sigma = \rho \cup \{v\}$ that has not yet been output, we intuitively consider the angle α_v for which σ enters this halfspace. Specifically, we consider α_v , the angle such that $\gamma_{\rho}(\alpha_v)$ is the γ_{ρ} -normal of v relative to ρ . We then output these cofacets $\rho \cup \{v\}$ in increasing order based on α_v , breaking ties arbitrarily, and pair them with the corresponding direction $\gamma_{\rho}(\alpha_v)$. Figure 4 illustrates the order in which various *i*-simplices are output.

If we encounter a simplex ρ whose affine hull has dimension d-1, there are only two (antipodal) directions perpendicular to it; in this case, we simply choose an \mathbb{S}^1 containing these two directions, so that all its cofaces have one of two possible angles. For clearer exposition, we encompass these two cases in the following definition.

▶ **Definition 6** (Maximally Perpendicular Circle). Let K be a simplicial complex in \mathbb{R}^d and let $\sigma \subseteq K$ be an *i*-simplex with dim(aff(σ)) < d. We say that a circle of directions, γ_{σ} , is maximally perpendicular (to σ), if

1. When dim $(aff(\sigma)) < d-1$, we have $\gamma_{\sigma} \subseteq \perp_{\sigma}$, or;

2. When dim $(aff(\sigma)) = d - 1$, we have $\perp_{\sigma} \subseteq \gamma_{\sigma}$.

That is, generally, γ_{σ} "rotates around" σ , except in the case that σ only has two directions perpendicular to it, in which case, these directions are contained in γ_{σ} . In either case, γ_{σ} is "as perpendicular" to σ as a circle of directions possibly can be.



Figure 4 A simulation of $ORDER(K_0)$ with direction s, and $ORDER(K_i, SO_{i-1})$ for $i \in \{1, 2\}$. Some circles γ_{ρ} relevant to the order are shown. Each γ_{v_i} lies in a plane parallel to the page. The circle γ_{e_6} lies in a plane perpendicular to edge e_6 . A dotted line connecting a simplex σ to a facet ρ indicates that ρ is the facet that outputs σ . A solid line indicates the γ_{ρ} -normal with which some σ is output. Indices correspond to the order in simplices are output.

Algorithm 1 Computing sweeping orders: SO_0 , and SO_i given SO_{i-1} and K_i .

¹ **procedure** ORDER(complex K_0) $s \leftarrow arbitrary direction$ 2 for vertex $v \in V(K_0)$ sorted increasingly by s-coordinate, breaking ties arbitrarily do 3 **output** (v, s)4 ⁵ **procedure** ORDER(complex K_i , sweeping order SO_{i-1} of K_{i-1}) \triangleright s is a direction perpendicular to the (i-1)-simplex ρ for (ρ, s) in SO_{i-1} do 6 $\gamma_{\rho} \leftarrow$ arbitrary circle $\gamma_{\rho} \colon [0, 2\pi) \to \mathbb{S}^{d-1}$ of directions maximally perpendicular to 7 ρ , where the angle between s and $\gamma_{\rho}(\alpha)$ is α , so $\gamma_{\rho}(0) = s$ and $\gamma_{\rho}(\pi) = -s$ $U_{\rho} \leftarrow \{ \text{vertex } v \text{ of } K_0 \mid \rho \cup \{v\} \text{ is an } i \text{-simplex of } K_i \text{ that was not yet output} \}$ 8 for $v \in U_{\rho}$ do 9 $\alpha_v \leftarrow \alpha$ such that $\gamma_{\rho}(\alpha)$ is the γ_{ρ} -normal to v relative to ρ . 10 for $v \in U_{\rho}$ sorted increasingly by α_v , breaking ties arbitrarily do 11 output $(\rho \cup \{v\}, \gamma_{\rho}(\alpha_v))$ 12

▶ Remark 7. In \mathbb{R}^0 or \mathbb{R}^1 , we cannot find any \mathbb{S}^1 of directions in the ambient space. In this very particular case, we invite the reader to imagine \mathbb{R}^0 or \mathbb{R}^1 along with any simplicial complex it contains as being included into \mathbb{R}^2 , e.g., along the *x*-axis. Then we are able to proceed just as we do in the general case. Thus, we assume that d > 1 for ease of exposition.

Since our method requires the use of maximally perpendicular circles of directions, and a sweeping order pairs simplices with directions perpendicular to them, we require every simplex to have at least some \mathbb{S}^0 of perpendicular directions. We formalize this in Assumption 1, which we henceforth assume is satisfied by K.

▶ Assumption 1 (General Position for Enough Perpendiculars). Let K be a simplicial complex in \mathbb{R}^d . For every simplex $\sigma \in K_i$, we require that dim $(aff(\sigma)) \leq d-1$.

Note that this is a rather lenient condition. Assumption 1 automatically holds when i < d, so when K_i is at most (d - 1)-dimensional. Additionally, Assumption 1 allows for simplices of higher dimension as long as their affine hulls are sufficiently low-dimensional.

Proceeding with this assumption, $ORDER(K_i, SO_{i-1})$, Algorithm 1, takes as input K_i and SO_{i-1} , a sweeping order for K_{i-1} , and outputs a sequence of *i*-simplices and directions. The main result of this section is Theorem 10, which says $ORDER(K_i, SO_{i-1})$ is a sweeping order for K_i . We first show that it satisfies Property 1 and 3 of Definition 5.

▶ Lemma 8 (Directions are Perpendicular to their Paired Simplices). Let $0 \le i \le \dim(K) - 1$. If i = 0, let $SO_i = ORDER(K_i)$. If i > 0, let $SO_i = ORDER(K_i, SO_{i-1})$ for some sweeping order SO_{i-1} . For all elements $(\sigma, s_{\sigma}) \in SO_i$, the direction s is perpendicular to σ . That is, the output of Algorithm 1 satisfies Property 1 of Definition 5.

Proof. Consider an arbitrary $(\sigma, s_{\sigma}) \in SO_i$. If i = 0, then σ is a vertex, and any direction is perpendicular to σ , including s. So consider the case i > 0. Then $\sigma = \rho \cup \{v\}$ for some (i-1)-simplex ρ and vertex v, where (ρ, s_{ρ}) is an index for the loop in Line 6, and v is an element of U_p (Line 9). On Line 7, we find the angle α_v such that $\gamma_{\rho}(\alpha_v)$ is the γ_{ρ} -normal of v relative to ρ . By Assumption 1, we have dim(aff(ρ)) $\leq d-1$.

If dim(aff(ρ)) < d-1, then γ_{ρ} is perpendicular to ρ , so the γ_{ρ} -normal $\gamma_{\rho}(\alpha_{v})$ is well-defined and hence perpendicular to σ . If dim(aff(ρ)) = d-1, then we must have aff(σ) = aff(ρ), otherwise dim(aff(σ)) would be greater than d-1, violating Assumption 1. Then the γ_{ρ} normal of v relative to ρ is $\gamma_{\rho}(0) = s_{\rho}$, which is perpendicular ρ and hence also to σ .

▶ Lemma 9 (Halfspace Property). Suppose that $ORDER(K_i, SO_{i-1}) = ((\sigma_j, s_j))_{j=1}^{n_i}$. Then, for any $1 \le j \le n_i$, for each simplex τ of $cof_{s_j}^{\le}(\sigma_j)$, τ is a cofacet of some σ_h with h < j. That is, Algorithm 1 satisfies Property 3 of Definition 5.

Proof. When i = 0, on Line 4, we output vertices ordered by their height with respect to some direction s, breaking ties arbitrarily. Then any edge of a vertex v_j whose other endpoint v_h lies in the open halfspace below v_j with respect to s is trivially also an edge of v_h , and v_h comes before v_j in the ordering induced by s.

Next, consider the case i > 0 and suppose that $ORDER(K_i, SO_{i-1})$, outputs (σ_j, s_j) in the iteration (ρ, s) . Let v_j be the vertex such that $\sigma_j = \rho \cup \{v_j\}$. We show for any simplex $\tau = \sigma_j \cup \{v_h\} \in cof_{s_i}^{<}(\sigma_j)$, that the simplex $\sigma_h = \rho \cup \{v_h\}$ satisfies the claim.



Figure 5 For some $\sigma_j = \rho \cup \{v_j\}$, we consider a cofacet $\tau = \rho \cup \{v_j, v_h\}$ (unshaded) for which the vertex v_h lies below σ_j with respect to the direction $\gamma_\rho(\alpha_{v_j})$. The simplex $\sigma_h = \rho \cup \{v_h\}$ is a cofacet of ρ that also has τ as a cofacet. The simplex ρ has a perpendicular circle of directions γ_ρ .

Since σ_h is a cofacet of ρ , and the iteration of (ρ, s) outputs all cofacets of ρ that have not yet been output, σ_h is output either in iteration of (ρ, s) , or in a previous iteration. If σ_h was output in a previous iteration, then σ_h comes before σ_j in the order, so h < j. Consider the remaining case that σ_h is output during iteration (ρ, s) . Since (ρ, s) is an element of a sweeping order for K_{i-1} , it satisfies Property 3 of Definition 5, so both σ_j and σ_h must be elements of $\operatorname{cof}_s^{\geq}(\rho)$, otherwise they would have been output in a previous iteration. Then we have $s \cdot v_j \geq s \cdot \rho$ and $s \cdot v_h \geq s \cdot \rho$, where $s = \gamma_{\rho}(0)$.

Let $\gamma_{\rho}(\alpha_{v_j})$ and $\gamma_{\rho}(\alpha_{v_h})$ denote γ_{ρ} -normals of v_j and v_h relative to ρ , respectively. Because $\sigma_j \cup \{v_h\} \in \operatorname{cof}_{s_j}^{<}(\sigma_j)$, and since $s_j = \gamma_{\rho}(\alpha_{v_j})$, we have $\gamma_{\rho}(\alpha_{v_j}) \cdot v_h < \gamma_{\rho}(\alpha_{v_j}) \cdot \sigma_j$. Then v_h lies (as illustrated by the shaded sector in Figure 5) in the open halfspace that contains σ_j in its boundary with exterior normal direction $\gamma_{\rho}(\alpha_{v_j})$, but not in the open halfspace that contains ρ in its boundary with exterior normal direction $\gamma_{\rho}(0)$. This means that $0 \leq \alpha_h < \alpha_j$, so σ_h is output before σ_j on Line 11, and the claim is satisfied.

Algorithm 1 moreover satisfies Property 2 of Definition 5 trivially, so Theorem 10 follows.

Theorem 10. $ORDER(K_i, SO_{i-1})$, Algorithm 1, outputs a sweeping order for K_i .

4 Candidate Simplices

Our reconstruction method reconstructs K_{i+1} from K_i by determining all (i + 1)-simplices of K. If not all facets of an (i + 1)-simplex τ (not necessarily in K_{i+1}), lie in K_i , then we know that τ is not part of K and we do not need to consider τ in our search. Moreover, if we happen to know K satisfies extra properties, e.g., that it is embedded, and adding τ to K_i would violate that property, there is no reason to consider τ in our search. The following definition specifies which potential simplices we should consider in our search. See Figure 6.

▶ Definition 11 (Candidate Vertices and Candidate Cofacets). Let K be a simplicial complex in \mathbb{R}^d with property X (locally injective, embedded, no condition, etc.), and let $\sigma \subseteq K$ be an *i*-simplex for some $i < \dim(K)$. Suppose that there exists a vertex $v \in K_0$ such that the simplices in K_i defined on $\sigma \cup \{v\}$ form the boundary of an (i+1)-simplex τ such that $K_i \cup \tau$ is a subcomplex of some simplicial complex with property X. We call v a candidate vertex of σ . Furthermore, we call the (i+1)-simplex defined by $\sigma \cup \{v\}$ a candidate cofacet of σ .

Intuitively, for purposes of reconstruction, a candidate cofacet of σ is an (i + 1)-simplex that may or may not be contained in K. Because our general position assumption is stated



Figure 6 Candidate edges (a–b) and triangles (c–d) of an (unknown) simplicial complex depending on whether it is known to be locally injective or embedded. Only (a) does not satisfy Assumption 2.



Figure 7 Suppose that (a) displays the candidate edges for v_1, v_2, v_3 , and v_4 , which violates Assumption 2. Note that this can lead to non-reconstructible simplicial complexes, as (b) and (c) have the same indegree information for all directions.

in terms of candidate simplices, specifying property X may reduce the number of candidate vertices, and allow us to reconstruct more general simplicial complexes. We assume that the set of candidates of dimension i+1 can be determined once we have correctly reconstructed K_i . We henceforth assume that these candidates are known to us. We also make the following general position assumption on K_i and its candidate simplices.

▶ Assumption 2 (General Position for (i + 1)-Reconstruction). Let K be a simplicial complex in \mathbb{R}^d . Suppose, for every i-simplex σ , the set of all candidate cofacets of σ along with K_i is locally injective in \mathbb{R}^d . Then we say that K is in general position for (i + 1)-reconstruction.

▶ Observation 12. Assumption 2 is automatically satisfied for candidates of embedded simplicial complexes. To see this, consider two candidate cofacets τ_1 and τ_2 of an i-simplex σ . Because K_i is embedded, the boundaries of τ_1 and τ_2 intersect only in σ . If τ_1 and τ_2 is not a locally injective pair, the convex hull of one contains the boundary of the other. Without loss of generality, assume that the boundary of τ_1 is contained in the convex hull of τ_2 . But then τ_2 would not be a candidate cofacet for σ , as $K_i \cup \tau_2$ is not embedded.

See Figure 7 for an instance where violating Assumption 2 corresponds to non-reconstructible simplicial complexes. Furthermore, Assumption 2 implies that, for a candidate cofacet τ , we have dim $(\tau) = \dim(\operatorname{aff}(\tau)) = i + 1$. Assumption 2 implies the following property.

▶ Lemma 13. Let σ be an *i*-simplex of a simplicial complex K in \mathbb{R}^d where i < d. Let C_{i+1} denote the candidate cofacets of σ . If $K_i \cup C_{i+1}$ is locally injective, then only two candidate cofacets of σ can share the same affine hull, and such candidates are separated by aff(σ).

Proof. Let τ be a candidate cofacet of σ . Because $K_i \cup C_{i+1}$ is locally injective, the dimension of $\operatorname{aff}(\tau)$ is one more than the dimension of $\operatorname{aff}(\sigma)$. The space $\operatorname{aff}(\sigma)$ is a separating hyperplane in $\operatorname{aff}(\tau)$, so there is a well-defined notion of being on a particular side of $\operatorname{aff}(\sigma)$ in $\operatorname{aff}(\tau)$. Let τ' be some other candidate cofacet of σ and suppose, towards a contradiction, that $\operatorname{aff}(\tau) = \operatorname{aff}(\tau')$, and both τ and τ' lie on the same side of $\operatorname{aff}(\sigma)$. Then σ is the largest common face of τ and τ' , but their intersection contains more than σ , so they are not an injective pair, contradicting our assumption of local injectivity for $K_i \cup C_{i+1}$. Finally, since there are only two sides of $\operatorname{aff}(\sigma)$ in an (i + 1)-dimensional hull, candidate cofacets of σ that share a common affine hull can only come in pairs on opposite sides.

5 Radially Ordering Candidate Cofacets

In this paper, we use circles of maximally perpendicular directions in two separate but related contexts. We have already seen one instance; in Section 3, we found a sweeping order for K_i by rotating around circles maximally perpendicular to each (i - 1)-simplex. In this section, we discuss the existence of the second type of maximally perpendicular circle, around which we rotate to order candidate cofacets of some central *i*-simplex in the reconstruction process.



Figure 8 Here, σ is a vertex in \mathbb{R}^3 . Then our pointset *P* consists of a single vertex *p*, chosen to avoid the affine hulls of pairs of (candidate) cofacets of σ (in this simple example, there is one such plane to avoid). Notice that the resulting circle γ_{σ} does not contain a direction simultaneously orthogonal to τ and τ' ; this would only occur if τ and τ' were colinear or if *p* were chosen to lie in the purple plane, i.e., if all four vertices were coplanar. The arrows on γ_{σ} foreshadow our eventual aim; it can be used to define a total radial order on the candidate cofacets of σ .

▶ Definition 14 (Candidate-Ordering Circle). Let K be a simplicial complex in \mathbb{R}^d and consider an i-simplex $\sigma \subseteq K$ for which there exists some parameterized circle of directions, $\gamma_{\sigma} : [0, 2\pi) \to \mathbb{R}^d$, that is maximally perpendicular to σ . If every candidate vertex of σ has a unique γ_{σ} -normal relative to σ , then we say γ_{σ} is candidate-ordering.

Since orders from different angular parameterizations are simply cyclic permutations of each other, we generally do not specify the parameterization of a candidate-ordering circle. Towards candidate-ordering circles, the following lemma shows that, for an simplex σ with dimension i < d - 1, we can build a circle of directions γ_{σ} so that any direction in it is perpendicular to at most two candidate cofacets of σ simultaneously, and such candidate pairs share affine hulls.

▶ Lemma 15. Let K be a simplicial complex in \mathbb{R}^d for $d \ge 2$, and let $\sigma \subseteq K$ be an *i*-simplex with i < d - 1. Let C_{i+1} denote the candidate cofacets of σ . If $K_i \cup C_{i+1}$ is locally injective, then a circle of directions γ_{σ} maximally perpendicular to σ exists so that whenever a direction of γ_{σ} is normal to two candidate cofacets τ and τ' of σ , the affine hulls of τ and τ' are equal.

Proof. We show the existence of γ_{σ} constructively. First, suppose i = d - 2. Then $\gamma_{\sigma} = \perp_{\sigma}$ is a maximally perpendicular circle, and the affine hulls of candidate cofacets of σ are (d-1)-planes normal to some direction in γ_{σ} . By Lemma 13, these planes contain at most two candidate cofacets and the claim is satisfied.

Next, consider i < d-2. Let τ and τ' be candidate cofacets of σ . Then $\operatorname{aff}(\tau \cup \tau')$ is (i+2)-dimensional (if $\operatorname{aff}(\tau) \neq \operatorname{aff}(\tau')$), or else is (i+1)-dimensional (if $\operatorname{aff}(\tau) = \operatorname{aff}(\tau')$). In particular, the arrangement of the planes $\operatorname{aff}(\tau \cup \tau')$ for every pair of candidate cofacets τ and τ' is less than d-dimensional. Then we can choose a set of d-i-2 points not contained in this arrangement, which we denote P, in such a way so that $\operatorname{aff}(\sigma \cup \{P\})$ is (d-2)-dimensional. Let γ_{σ} be the unique \mathbb{S}^1 of directions (maximally) perpendicular to $\operatorname{aff}(\sigma \cup \{P\})$.

Now, suppose there is some $s \in \gamma_{\sigma}$ perpendicular to two candidate cofacets, $\sigma \cup \{v\}$ and $\sigma \cup \{v'\}$. That is, for all $p \in \sigma \cup \{P\}$, we have $s \cdot v = s \cdot v' = s \cdot p$. Then $\sigma \cup \{P\} \cup \{v, v'\}$ is a set of (d-1)+2 = d+1 points that all lie on the same (d-1)-plane (normal to s). But by construction, dim(aff($\sigma \cup \{P\}$)) = d-2, so it must be that aff(τ) = aff(τ').

See Figure 8 for a low-dimensional illustration of the construction described above.

We combine Lemmas 13 and 15 to show that a candidate-ordering circle always exists.

▶ Lemma 16. Let K be a simplicial complex, let σ be an i-simplex for $i \leq d-1$, and suppose that K satisfies Assumption 2. Then there exists a candidate-ordering γ_{σ} .

Proof. We proceed with a constructive argument. If i = d - 1, then any candidate cofacets of σ are contained in the same *d*-plane, and so, by Lemma 13, σ can have at most two

candidate cofacets, contained on opposite sides of σ . Then, for any circle γ_{σ} of directions maximally perpendicular to σ , the γ_{σ} -normal relative to σ of one candidate cofacet is $\gamma_{\sigma}(0)$, and that of the other is $\gamma_{\sigma}(\pi)$; these are distinct, so the claim holds.

Next, suppose i < d-1, and suppose that $\gamma_{\sigma} \subseteq \perp_{\sigma}$ is the circle maximally perpendicular to σ , constructed as in the proof of Lemma 15. Choose an angular parametrization of γ_{σ} by $[0, 2\pi)$. By Lemma 15, whenever a direction $\gamma_{\sigma}(\alpha)$ is orthogonal to multiple candidate cofacets, they share affine hulls. By Lemma 13, such candidate cofacets lie on opposite sides of σ , and can only come in pairs. Thus, such candidate cofacet pairs have opposite (and distinct) γ_{σ} -normals relative to σ . Thus, γ_{σ} is a candidate-ordering circle for σ .

In Section 6, we consider sweeping orders where, for each pair (σ, s) , the direction s is part of some candidate-ordering circle around σ . We call this special type of sweeping order a *candidate-ordering-compatible* sweeping order. To show this type of order always exists, we begin by establishing two helpful lemmas. The first serves as our eventual base case.

▶ Lemma 17 (Candidate-Ordering for All Vertices). Let K be a simplicial complex in \mathbb{R}^d . Then there is a circle of directions that is candidate-ordering for every vertex of K_0 simultaneously.

Proof. First, consider the case d = 2. By Lemma 16, each vertex $v \in K_0$ has some candidateordering circle. Then the unique circle of directions is candidate-ordering, and the claim is satisfied trivially. Suppose then that d > 2. Each triple of affinely independent vertices defines a plane; let $S \subset \mathbb{S}^{d-1}$ be the set of directions normal to such planes, formed by considering all triples in K_0 . Since K_0 is finite, so is S, and we can find some circle γ such that $\gamma \cap S = \emptyset$. We claim γ is candidate-ordering for all vertices.

Consider some $v \in K_0$, and suppose, towards a contradiction, that γ is *not* candidateordering for v. That is, when rotating around v by the circle γ , there are two candidate vertices of v, which we call v_1 and v_2 , that appear at the same "angle" around γ . In particular, this means that, for some $s \in \gamma$, we have $s \cdot v = s \cdot v_1 = s \cdot v_2$, but aff (v, v_1, v_2) is two-dimensional. But then this s is normal to aff (v, v_1, v_2) , i.e., $s \in S$, which contradicts our assumption that γ is disjoint from S.

Next, we observe that the candidate vertices of a simplex are also candidate vertices for a facet of that simplex.

▶ Lemma 18. Let K be a simplicial complex in \mathbb{R}^d , and, for 0 < i < d - 1 consider an (i-1)-simplex $\rho \in K$. Let σ be a cofacet of ρ . Then, if $v \in K_0$ is a candidate vertex of σ , it is also a candidate vertex of ρ .

Proof. Recall from Definition 11 that the simplices of K_i defined on $\sigma \cup \{v\}$ form the boundary of an (i + 1)-simplex τ such that $K_i \cup \tau$ is subcomplex of some simplicial complex K' that satisfies Assumption 2, and possibly additional constraints. Note that the simplices defined on $\sigma \cup \{v\}$ are a superset of the simplices defined on $\rho \cup \{v\}$; in particular, simplices on $\rho \cup \{v\}$ in K_{i-1} form the boundary of an *i*-simplex σ' . Then $K_{i-1} \cup \{\sigma'\}$ is also a subcomplex of K'. Therefore, v is also a candidate vertex for ρ .

Finally, we are ready to prove Lemma 19.

▶ Lemma 19. Let K be a simplicial complex in \mathbb{R}^d , and $i \leq d$. Suppose that K is in general position for (i'+1)-reconstruction, for all $i' \leq i \leq d-1$. Then a sweeping order $((\sigma_j, s_j))_{j=1}^{n_i}$ exists such that, for all $1 \leq j \leq n_i$, there is a candidate-ordering circle around σ_j that contains s_j .

Proof. We claim that we can construct such a sweeping order by iteratively calling Algorithm 1 and specifically using candidate-ordering circles on Line 7 for each call. We proceed by induction on *i*. Consider the base case i = 0. By Lemma 17 there exists some circle of directions that is candidate-ordering for *all* vertices. Choosing some arbitrary direction *s* from this circle on Line 2 in Algorithm 1 results in the sweeping sequence $((v_j, s))_{j=1}^{n_0}$, which satisfies the claim.

Next, for some $\ell - 1 \ge 0$, suppose that $((\rho_j, s_j))_{j=1}^{n_{\ell}-1}$ is a sweeping order for $K_{\ell-1}$, where every s_j is part of a candidate-ordering circle γ_j around ρ_j , and where we specifically rotate around γ_j in Line 7 to order cofacets of ρ_j . Suppose that we compute a sweeping sequence for K_{ℓ} using $((\rho_j, s_j))_{j=1}^{n_{\ell}-1}$ as input to Algorithm 1. Let (σ, s) be an arbitrary term of the output, and suppose that it is output in the iteration of Line 6 that corresponds to (ρ_k, s_k) . Then ρ_k is the first $(\ell - 1)$ -simplex in the sweeping order for $K_{\ell-1}$ that contains σ as a cofacet. Furthermore, γ_k used on Line 7 is candidate-ordering for ρ_k and $s \in \gamma_k$.

Because σ was chosen arbitrarily, it suffices to show that there exists a (maximally perpendicular) candidate-ordering circle around σ that contains s. Assume without loss of generality that γ is a maximally perpendicular circle that contains s and minimizes the number of candidate pairs that have the same γ -normal relative to σ .

If $\ell = d - 1$, any maximally perpendicular circle contains the two directions normal to σ , so the two possible candidate cofacets of σ would be assigned unique γ -normals, i.e., γ is candidate-ordering. Similarly, if $\ell = d - 2$, γ is the unique maximally perpendicular circle to σ , and so by Lemma 19, it is candidate-ordering.

Suppose then, that $\ell < d-2$; note that this implies that \perp_{σ} is at least a two-sphere, and so, contains an uncountable number of maximally perpendicular (sub-)circles. Suppose, towards a contradiction, that γ is not candidate-ordering for σ . Then there exist two candidate vertices for σ , v_1 and v_2 , that have the same γ -normal $\gamma(t)$ relative to σ . First, consider the case that $\gamma(t)$ is not s or -s. Because $\ell < d-2$, there exists some direction $s' \notin \{s, -s\}$, that is perpendicular to $\operatorname{aff}(\sigma)$, but not perpendicular to $\operatorname{aff}(\sigma \cup \{v_1, v_2\})$. Note that there is a unique circle perpendicular to σ that contains s, -s, and s', which we denote by γ' . The circles γ and γ' both lie in the space of circles through s and -s that are maximally perpendicular to σ , and this space is parameterized by the equator of \perp_{σ} with s and -sas poles. Consider rotating γ infinitesimally towards γ' along a geodesic in this space. We claim that such a rotation results in another maximally perpendicular circle, γ'' , such that no additional pairs of candidates get the same γ'' -normal, but v_1 and v_2 no longer have the same γ'' -normal. If not, all circles along the geodesic of rotation would assign the same normal to v_1 and v_2 , which we know is not the case by construction of γ' . Thus, γ did not minimize the number of candidate pairs with the same γ -normal. Hence, whenever v_1 and v_2 have the same γ -normal, their γ -normal is s or -s.

In that case, the s-coordinates of σ , v_1 , and v_2 are equal. If $\operatorname{aff}(\sigma \cup \{v_1\}) = \operatorname{aff}(\sigma \cup \{v_2\})$, then by Lemma 13, v_1 and v_2 lie on "opposite sides" of σ and would have received distinct γ -normals; namely, s and -s. Therefore, $\operatorname{aff}(\sigma \cup \{v_1\}) \neq \operatorname{aff}(\sigma \cup \{v_2\})$. By Lemma 18, we know that v_1 and v_2 are also candidate vertices for ρ_k . Since ρ_k is a facet of σ , the s-coordinates of σ and ρ_k are equal, and hence the s-coordinates of ρ_k , v_1 , and v_2 are equal. However, note that $\operatorname{aff}(\rho_k \cup \{v_1\}) \neq \operatorname{aff}(\rho_k \cup \{v_2\})$, so the two candidates would be assigned the same γ_k -angle relative to ρ_k . This contradicts our assumption that γ_k was candidate-ordering for ρ_k . The claim now follows by induction.

As an immediate consequence of the proof of this Lemma 19, we see that computing this particular type of sweeping order via Algorithm 1 is possible.

► Corollary 20. If we input a candidate-ordering-compatible sweeping order to Algorithm 1, the output is also candidate-ordering-compatible.

6 Application to Simplicial Complex Reconstruction

In this section, we give details of an application where the properties of our sweeping order are central, namely, extending the edge reconstruction algorithm of [11] to simplicial complex reconstruction. Just as in [11], we sweep through *i*-simplices, finding all cofacets above a given *i*-simplex at each step, and maintaining that all cofacets below have already been found. We iterate the process of reconstructing K_{i+1} from K_i until we have reconstructed K.

6.1 Finding Cofacets above a Single *i*-Simplex

First, we discuss our central subroutine; identifying all cofacets of and above a single *i*-simplex σ , supposing that all cofacets of and below σ have already been found, and where "above" and "below" are with respect to some direction perpendicular to σ . Justifying that we can complete this step of reconstruction with a near copy of the algorithm given in [11] is a primary goal of this section. Our main tool is the following function, which counts cofacets of a simplex in the halfspace below the simplex with respect to some direction.

▶ Definition 21 (Indegree). Given σ , some simplex of a simplicial complex K, and $s \in \perp_{\sigma}$, Indeg (σ, s) returns the number of cofacets of σ that have a vertex strictly below σ with respect to direction s. That is, Indeg $(\sigma, s) = |cof_s^{<}(\sigma)|$.

For now, we assume $\text{Indeg}(\sigma, s)$ as a well-defined subroutine; we provide further discussion about its actual existing implementations and corresponding limitations in Appendix A.

Our method to find cofacets of and above a particular *i*-simplex is essentially identical to the method to find edges adjacent to and above a particular vertex in [11, Algorithm 2], so we begin with a short summary. The reader may be aided by Figure 9. Informally, the algorithm considers a central vertex v, whose adjacent edges in some lower halfspace are known, but for which its adjacent edges in the upper halfspace are unknown, where "above" and "below" are with respect to a fixed direction, s. The algorithm maintains a partition of the upper halfspace into *edge arc objects*, which correspond to a wedge-shaped region and contain a list of the radially sorted candidate vertices contained in the region, as well as the number of actual cofacets (edges). We initialize the first edge arc to be the entire upper halfspace; computing INDEGREE(v, -s) gives us the correct number of actual



Figure 9 The first few steps of UpEdges, or, if σ is a more general simplex, the first few steps of UpCofacets. We initialize the first arc as the entire upper halfplane (left). We find the indegree of σ in direction -s is two, but we don't yet know which pair of v_1, v_2 , or v_3 are part of these two cofacets of σ . Next, we use a direction s' that divides the set of candidate vertices above σ in half (right). The indegree of σ in direction s' is two, but since we know $\sigma \cup \{v_4\}$ is a cofacet of σ , we know there is exactly one cofacet in the blue arc. Since there is only one candidate vertex in this arc, we know that $\sigma \cup \{v_3\}$ is a cofacet of σ . Next, we would process the green arc in a similar manner.

cofacets of v (i.e., $|cof_s^{\geq}(v)|$). We then begin a binary search through the radially sorted candidate vertices in the upper halfspace. Choosing a direction s' for which half the candidate vertices are above v and half are below, we split the initial edge arc object in half, using given angle information to avoid choosing some s' that causes a candidate vertex and v to have the same s'-coordinate. The indegree query INDEGREE (σ, s') may now also count edges in $cof_s^{<}(\sigma)$, but since these edges are known, we subtract them from the indegree count to obtain the correct count of actual cofacets of v in this new edge arc. We continue splitting edge arcs in half, until the number of candidate vertices matches the number of cofacets of v. In this way, we identify all edges of $cof_s^{\geq}(\sigma)$.

The following lemma asserts the correctness of [11, Algorithm 2].

▶ Lemma 22 (Adapted [11, Theorem 8]). Let K be a simplicial complex, v be a vertex of K, let V be the set of candidate vertices above v with respect to some direction s, totally radially ordered by a candidate-ordering circle containing s, and let θ denote the minimum nonzero angle between any three vertices in K₀. Then UPEDGES(v, V, cof[<]_s(\sigma), θ , -) = cof[>]_s(v).

We emphasize that having a known lower halfspace is a crucial condition. Rotating the query halfspace may change the count, and we must be able to distinguish changes to the count arising from the candidate cofacets above σ from that from candidate cofacets below.

Note that the last input to UpEdges is left blank in Lemma 22. In the original setting, this input is a verbose persistence diagram used to compute indegree. However, we are supposing indegree is a well-defined subroutine and are not yet concerned with its actual calculation. This difference in perspective also results in slightly different general position assumptions taken between the current paper and [11]. Note that our definition of indegree (Definition 21) gives us a count of cofacets in a closed halfspace; our more general perspective allows us to avoid restrictions on other vertices having the same s-coordinate as v, and allows us output cofacets of v in the *closed* halfspace above v. See Appendix A.1 for a more thorough discussion on how actual implementations of indegree impact general position requirements.

Finally, the algorithm UpEdges in [11] discusses a *clockwise* ordering of candidate vertices. This is to clarify sidedness, but is otherwise taken without loss of generality; we therefore proceed without referencing clockwise or counterclockwise.

We are now ready to generalize UpEdges and define our routine for computing the cofacets above an *i*-simplex. Given an *i*-simplex with i < d - 1, and a candidate-ordering circle, the corresponding radial ordering of candidate cofacets behaves nearly identically to the



Figure 10 Projecting σ and its candidate cofacets to the plane containing a candidate-ordering circle γ_{σ} results in a star graph with the image of σ at the center, and the order of candidate vertices in the image remains unchanged. Note that $\sigma \cup \{v_2\}$ is not an actual cofacet of σ ; then, despite appearing as an edge in the projection, it would never contribute to an indegree count, as indegree only counts actual cofacets. It is also possible that an (i + 1)-simplex τ is attached to σ , but is not a cofacet of σ (not pictured). Again, despite corresponding to edge(s) in the projection, τ does not contribute to the indegree of σ , nor even as a candidate cofacet, and is essentially ignored.

radial ordering of edges adjacent to some central vertex. Figure 10 highlights this connection. For i = d - 1, we can no longer rotate around the simplex, but a simple indegree check confirms the existence or absence of an upper cofacet.

Algorithm 2 For an *i*-simplex $\sigma \in K$ and direction *s*, UPCOFACETS $(\sigma, s, \gamma_{\sigma}, V, \text{cof}_{s}^{<}(\sigma), \theta)$ computes $\text{cof}_{s}^{\geq}(\sigma)$, where γ_{σ} is a candidate-ordering circle containing *s*, *V* are the candidate vertices of and above σ (ordered by γ_{σ}), and θ is the minimum nonzero angle between any three vertices of K_{0} .

1 procedure UPCOFACETS $(\sigma, s, \gamma_{\sigma}, V, \operatorname{cof}_{s}^{<}(\sigma), \theta)$ 2 if i < d-1 then 3 return UPEDGES $(\sigma, V, \operatorname{cof}_{s}^{<}(\sigma), \theta, -)$, using indegree is as in Definition 21, and the radial order induced by γ_{σ} 4 if i = d-1 then 5 return \emptyset if INDEGREE $(\sigma, -s) = 0$ or $\sigma \cup \{V\}$ if INDEGREE $(\sigma, -s) = 1$

▶ Lemma 23. For σ , an i-simplex of a simplicial complex K, γ_{σ} , a candidate-ordering circle and, $cof_s^{<}(\sigma)$ for some $s \in \gamma$, and V, the set of candidate vertices above σ radially ordered by γ , and θ , the minimum nonzero angle between any three vertices of K_0 , we have UPCOFACETS $(\sigma, s, \gamma, V, cof_s^{<}(\sigma), \theta) = cof_s^{\geq}(\sigma)$. That is, Algorithm 2 is correct.

Proof. For i < d-1, we simply run UpEdges on Line 3, and the proof is a straightforward adaptation of [11, Theorem 8]. For i = d - 1, by Lemma 13, at most one candidate vertex v may lie above σ , i.e., V is either empty, or a single vertex. If there is no actual cofacet above σ , then INDEGREE $(\sigma, -s) = 0$, and we correctly return \emptyset on Line 5. If there actually is a cofacet above σ , then INDEGREE $(\sigma, -s) = 1$, and we correctly return $\sigma \cup \{V\} = \sigma \cup \{v\}$.

6.2 Reconstructing K

By processing *i*-simplices in a sweeping order, Algorithm 3 reconstructs K_{i+1} .

Algorithm 3 RECONSTRUCTNEXT(K_i , SO_i), for a candidate-ordering-compatible sweeping sequence, SO_i , computes K_{i+1} .

¹ procedure RECONSTRUCTNEXT(K_i, SO_i)	
2	$\theta \leftarrow \text{minimum nonzero angle between any three vertices in } K_0$
3	Found $\leftarrow \{\}$
4	for (σ, s) in SO_i do
5	$\gamma_{\sigma} \leftarrow \text{candidate-ordering circle that contains } s$
6	$Up Verts \leftarrow array of candidate vertices v for \sigma with v \cdot s \ge \sigma \cdot s, ordered by \gamma_{\sigma}$
7	$C \leftarrow \text{array of cofacets of } \sigma \text{ contained in } Found \text{ with a vertex below } \sigma, \text{ sorted}$
	around σ in the radial order induced by γ_{σ}
8	$Found = Found \cup \texttt{UpCofacets}(\sigma, s, \gamma_{\sigma}, \textit{UpVerts}, C, \theta)$
${}_{\mathfrak{s}} \ _ \ \mathbf{return} \ K_i \cup Found$	

▶ Lemma 24. Algorithm 3 is correct. That is, $RECONSTRUCTNEXT(K_i, SO_i) = K_{i+1}$.

Proof. We claim that the loop in Lines 4-8 satisfies the following invariant: entering iteration j, the list *Found* contains all cofacets of any previously processed *i*-simplices. This is initially trivially true, since we enter the loop after initializing *Found* to empty on Line 3.

Next, suppose that this loop invariant is true when entering iteration j for some $1 \le j < n_i$, and denote the pair indexing iteration j by (σ_j, s_j) . That is, for each h < j, we have

 $\operatorname{cof}(\sigma_h) \subseteq \operatorname{Found}$. Since the *i*-simplices are processed according to the sweeping order, the direction s_j is perpendicular to σ_j by Property 1 of Definition 5, so $\operatorname{cof}_{s_j}^<(\sigma_j)$ is well-defined. Furthermore, by Property 3 of Definition 5, all cofacets of σ_j contained in $\operatorname{cof}_{s_j}^<(\sigma_j)$ are cofacets of *i*-simplices that appeared earlier in the sweeping order. By assumption, such (i+1)-simplices are already recorded in Found, meaning that $\operatorname{cof}_{s_j}^<(\sigma_j) \subseteq \operatorname{Found}$. In particular, on line Line 7, we have $C = \operatorname{cof}_{s_j}^<(\sigma_j)$. Then by Lemma 23, calling UpCofacets $(\sigma_j, s_j, \gamma_{\sigma_j}, C, \theta)$ on Line 8, correctly returns $\operatorname{cof}_{s_j}^\geq(\sigma_j)$. Adding this to Found means that, entering iteration j + 1, all cofaces of previously processed *i*-simplices are in Found, maintaining the invariant.

By Property 2 of Definition 5, all *i*-simplices appear exactly once in SO_i , so the loop iterates over all *i*-simplices in K. Since each (i + 1)-simplex is necessarily a cofacet of some *i*-simplex, when the loop terminates, all (i + 1)-simplices have been found.

Finally, we present Algorithm 4, which reconstructs K given the vertex set K_0 .

```
Algorithm 4 RECONSTRUCTALL(K_0) computes K, satisfying Assumption 2 for all i < \dim(K).
```

1 procedure RECONSTRUCTALL(K_0) $SO_0 \leftarrow \text{ORDER}(K_0)$, computed to be candidate-ordering-compatible $i \leftarrow 0$ 4 while K_i has *i*-simplices do $\left[\begin{array}{c} i \leftarrow i+1 \\ K_i \leftarrow \text{RECONSTRUCTNEXT}(K_{i-1}, SO_{i-1}) \\ SO_i \leftarrow \text{ORDER}(K_i, SO_{i-1}), \text{ choosing a candidate-ordering } \gamma_{\rho} \text{ on Line } 7$ $\left[\begin{array}{c} \text{return } K_i \end{array}\right]$

▶ Lemma 25. Algorithm 4 is correct. That is, $RECONSTRUCTALL(K_0) = K$.

Proof. We claim the loop in Lines 4–7 satisfies the following invariant: entering iteration i, we know K_{i-1} as well as a sweeping order for K_{i-1} . Since the vertex set is taken as input, we know K_0 initially. In Line 2, we call $ORDER(K_0)$, which, by Theorem 10, computes a sweeping order for K_0 . Thus, the loop invariant holds before entering the loop. Now suppose this invariant is true when entering iteration j for some $j \ge 1$. That is, we know K_{j-1} and a sweeping order for K_{j-1} , which we use as inputs in the call to ReconstructNext. on Line 6. By Lemma 24, this produces K_j . Finally on Line 7, we use Order (Algorithm 1) to compute a sweeping order for K_j , which, again, is correct by Theorem 10. Thus, the loop invariant holds. Since K is a finite simplicial complex, index i will eventually reach dim(K). At this point, the i-skeleton K_i equals the full simplicial complex K. When we set $i = \dim(K) + 1$ in Line 5, there are no i-simplices to find or order, so the output on Lines 6 and 7 is empty. Thus, we exit the loop and correctly return $K_i = K$.

We now consider the special case where the to-be-reconstructed complex K is known to be embedded. In that case, there are no (d+1)-dimensional simplices, and we can terminate the loop of Line 4 after constructing K_d (and before constructing SO_d). Whenever the algorithm computes SO_i , we have i < d, so Assumption 1 is automatically satisfied. Assumption 2 is satisfied for candidates of embedded complexes by Observation 12. Theorem 26 follows.

▶ **Theorem 26.** Let K be an embedded simplicial complex. There is an algorithm that under the promise that K is embedded, can reconstruct K using indegree.

7 Discussion

We note a possible improvement to our reconstruction procedure; in Algorithm 2, we search through all candidate vertices of σ in an upper half space. However, in practice, some of these candidate vertices may be part of cofacets of σ that were already computed at an earlier step (a statement that applies to higher-dimensional simplices, but not a standard single-direction sweep through vertices). It would be more efficient to only consider candidate vertices in the upper halfspace that do not already correspond to known cofacets of σ . We did not include this in Algorithm 2, so as to keep the connection to the central algorithm of [11] more explicit, but this adaptation would be fairly straightforward.

We defined sweeping orders with the particular motivation of simplicial complex reconstruction, but we note that Algorithm 1 could be adapted to order objects with a more general cell-structure. In particular, we expect that Algorithm 1 can be adapted to order faces of hyperplane arrangements. With such an order, we are curious what other traditional sweepline algorithms could be adapted to higher-dimensional generalizations.

— References

- 1 Erik J Amézquita, Michelle Y Quigley, Tim Ophelders, Jacob B Landis, Daniel Koenig, Elizabeth Munch, and Daniel H Chitwood. Measuring hidden phenotype: Quantifying the shape of barley seeds using the Euler characteristic transform. *in silico Plants*, 4(1):diab033, 2022.
- 2 Robin Lynne Belton, Brittany Terese Fasy, Rostik Mertz, Samuel Micka, David L. Millman, Daniel Salinas, Anna Schenfisch, Jordan Schupbach, and Lucia Williams. Learning simplicial complexes from persistence diagrams. In *Canadian Conference on Computational Geometry*, August 2018.
- 3 Robin Lynne Belton, Brittany Terese Fasy, Rostik Mertz, Samuel Micka, David L. Millman, Daniel Salinas, Anna Schenfisch, Jordan Schupbach, and Lucia Williams. Reconstructing embedded graphs from persistence diagrams. *Computational Geometry: Theory and Applications*, 2020. doi:10.1016/j.comgeo.2020.101658.
- 4 Leo M. Betthauser. Topological Reconstruction of Grayscale Images. PhD thesis, University of Florida, 2018.
- 5 Lorin Crawford, Anthea Monod, Andrew X Chen, Sayan Mukherjee, and Raúl Rabadán. Predicting clinical outcomes in glioblastoma: An application of topological and functional data analysis. *Journal of the American Statistical Association*, pages 1–12, 2019.
- **6** Justin Curry, Sayan Mukherjee, and Katharine Turner. How many directions determine a shape and other sufficiency results for two topological transforms. *Transactions of the American Mathematical Society, Series B*, 9(32):1006–1043, 2022.
- 7 Vid Domiter and Borut Žalik. Sweep-line algorithm for constrained Delaunay triangulation. International Journal of Geographical Information Science, 22(4):449–462, 2008.
- 8 Herbert Edelsbrunner and Leonidas J Guibas. Topologically sweeping an arrangement. In Proceedings of the eighteenth annual ACM symposium on Theory of computing, pages 389–403, 1986.
- **9** Herbert Edelsbrunner and John L Harer. *Computational topology: an introduction*. American Mathematical Society, 2022.
- 10 Brittany Terese Fasy, Samuel Micka, David L. Millman, Anna Schenfisch, and Lucia Williams. A faithful discretization of the verbose persistent homology transform, 2024. URL: https: //arxiv.org/abs/1912.12759, arXiv:1912.12759.
- 11 Brittany Terese Fasy, Samuel Micka, David L Millman, Anna Schenfisch, and Lucy Williams. Efficient graph reconstruction and representation using augmented persistence diagrams. In *Canadian Conference on Computational Geometry*, 2022.

- 12 Brittany Terese Fasy, David L Millman, and Anna Schenfisch. How small can faithful sets be? ordering topological descriptors. In *Canadian Conference on Computational Geometry*, 2024.
- 13 Cháulio Ferreira, Marcus VA Andrade, Salles VG Magalhães, W Randolph Franklin, and Guilherme C Pena. A parallel sweep line algorithm for visibility computation. In *GeoInfo*, pages 85–96. Citeseer, 2013.
- 14 Steven Fortune. A sweepline algorithm for Voronoi diagrams. In Proceedings of the second annual symposium on Computational geometry, pages 313–322, 1986.
- **15** Robert Ghrist, Rachel Levanger, and Huy Mai. Persistent homology and Euler integral transforms. *Journal of Applied and Computational Topology*, 2(1-2):55–60, 2018.
- 16 Allen Hatcher. Algebraic Topology. Cambridge Univ. Press, Cambridge, 2002.
- 17 Christoph Hofer, Roland Kwitt, Marc Niethammer, Yvonne Höller, Eugen Trinka, and Andreas Uhl. Constructing shape spaces from a topological perspective. In *International Conference* on *Information Processing in Medical Imaging*, pages 106–118. Springer, 2017.
- 18 Mattie Ji. On the injectivity of Euler integral transforms with hyperplanes and quadric hypersurfaces. *Mathematics*, 12(15):2339, 2024.
- 19 Qitong Jiang, Sebastian Kurtek, and Tom Needham. The weighted Euler curve transform for shape and image analysis. In *Proceedings of the IEEE/CVF Conference on Computer Vision* and Pattern Recognition Workshops, pages 844–845, 2020.
- 20 Niko Lukač, Borut Žalik, and Krista Rizman Žalik. Sweep-hyperplane clustering algorithm using dynamic model. *Informatica*, 25(4):563–580, 2014.
- 21 Clément Maria, Steve Oudot, and Elchanan Solomon. Intrinsic Topological Transforms via the Distance Kernel Embedding. In Sergio Cabello and Danny Z. Chen, editors, 36th International Symposium on Computational Geometry (SoCG 2020), volume 164 of Leibniz International Proceedings in Informatics (LIPIcs), pages 56:1–56:15, Dagstuhl, Germany, 2020. Schloss Dagstuhl-Leibniz-Zentrum für Informatik. doi:10.4230/LIPIcs.SoCG.2020.56.
- 22 Facundo Mémoli and Ling Zhou. Ephemeral Persistence Features and the Stability of Filtered Chain Complexes. In 39th International Symposium on Computational Geometry (SoCG 2023), volume 258 of Leibniz International Proceedings in Informatics (LIPIcs), pages 51:1–51:18, Dagstuhl, Germany, 2023.
- 23 Samuel Adam Micka. Searching and Reconstruction: Algorithms with Topological Descriptors. PhD thesis, Montana State University, 2020.
- 24 Michael Ian Shamos and Dan Hoey. Geometric intersection problems. In 17th Annual Symposium on Foundations of Computer Science (sfcs 1976), pages 208–215. IEEE, 1976.
- 25 Frank Y Shih and Vijayalakshmi Gaddipati. Geometric modeling and representation based on sweep mathematical morphology. *Information Sciences*, 171(1-3):213–231, 2005.
- 26 Katharine Turner, Sayan Mukherjee, and Doug M. Boyer. Persistent homology transform for modeling shapes and surfaces. *Information and Inference: A Journal of the IMA*, 3(4):310–344, 2014.
- 27 Borut Žalik. An efficient sweep-line Delaunay triangulation algorithm. Computer-Aided Design, 37(10):1027–1038, 2005.
- 28 Borut Žalik, Damjan Strnad, David Podgorelec, Ivana Kolingerová, Andrej Nerat, Niko Lukač, Štefan Kohek, and Luka Lukač. Geometric shape characterisation based on a multi-sweeping paradigm. Symmetry, 15(6):1212, 2023.
- **29** Ling Zhou. Beyond Persistent Homology: More Discriminative Persistent Invariants. PhD thesis, The Ohio State University, 2023.

A Connections to Directional Transforms

The persistent homology transform (PHT) and Euler characteristic transform (ECT), first defined in [26], map a geometric simplicial complex K to the set of persistence diagrams or Euler characteristic functions, respectively, corresponding to lower-star/sublevel set filtrations of K in each direction. In [26, 15], we see that such sets (parameterized by the sphere of directions) uniquely correspond to K; that is, they are *faithful*. In [18], we see conditions for the faithfulness of the *quadratic ECT*, which replaces the hyperplane sweeps used in sublevel set filtrations with quadratic hypersurfaces. Directional transforms can be useful in shape classification applications, see, e.g., [5, 26, 19, 4, 17, 21, 1]. Such applications necessarily use a *discretization* of the directional transform, i.e., a finte set of descriptors corresponding to a finite sample of the parameterizing directions. Finding finite sets of directional topological descriptors from which a simplicial complex can still be reconstructed has been explored in several contexts [2, 3, 4, 6, 23, 11, 10]. In particular, this is the perspective taken by [11]; since indegree is computable using information contained in verbose persistence diagrams, the sweep algorithm for edge reconstruction produces a set of directions for which the corresponding discretization of the verbose PHT is faithful.

Verbose topological descriptors are, roughly speaking, a record of how topological invariants change during a filtration, where we note the effects of adding a single simplex at a time, even if multiple simplices are added at a single parameter value. This varies from the more traditional *concise* topological descriptors, which only consider the coarse information of how the topological invariants change for each subcomplex of the filtration. Note that for an *index filtration* (where each simplex appears at a distinct parameter value), verbose and concise topological descriptors are equivalent. Also note that verbose and concise are sometimes called augmented and non-augmented, respectively.

If we choose to record homology, Betti number, or Euler characteristic, the resulting topological descriptor is the (verbose/concise) persistence diagram, Betti function, or Euler characteristic function, respectively. While precise definitions of these functions are not necessary for our discussion, we refer the reader to [12, Section 3 and Appendix A] for a more careful treatment.

A.1 Indegree from Verbose Persistence Diagrams and Betti Functions

We note that the definition of indegree given in [11, Def. 3] is restricted to vertices and edges (a special case of our definition), and the definition of k-indegree given in [10, Def. 21] allows us to choose the dimension of cofaces to count (a generalization of our definition).

In the case of verbose persistence diagrams and (with trivial adaptations) verbose Betti functions, we can immediately determine the number of (i + 1)-simplices that appear at the height of σ . From a single diagram corresponding to the direction in question, we simply count the number of *i*-dimensional points whose birth height is the same as the height of σ plus the number of (i + 1)-dimensional deaths whose birth height is the height of σ ; this total is the number of (i + 1)-simplices that appear at the height of σ (see, e.g., [10, Lemma 11]). However, this count may be more than the actual indegree of σ , since (i + 1)-simplices that are not cofacets of σ may have their highest vertex at the height of σ .

In [10, Alg. 1], such extra simplices that are "attached" to a proper face of σ are eliminated from the count by recursively finding the number of (i + 1)-simplices adjacent to faces of σ , beginning with σ itself, then with (i - 1)-dimensional faces of σ , then with (i - 2)-dimensional faces, and so on. Through an inclusion-exclusion computation, these totals are added and subtracted to obtain the true indegree of σ . This operation is limited to computing indegree

in directions s that isolate σ in a hyperplane normal to s, and also requires various tilts to isolate subsets of vertices. Thus, for this implementation of indegree, we require the following general position assumption.

▶ Assumption 3 (General Position for Indegree from Verbose Descriptors). Let K be a simplicial complex in \mathbb{R}^d . To compute indegree using verbose persistence diagrams (or verbose Betti functions) using the methods of [10], we require every collection of i vertices for $1 \le i < d$ to be affinely independent.

The specifics of actually computing indegree using verbose persistence diagrams or verbose Betti functions mean that, in practice, computing indegree in a single direction requires verbose descriptors for filtrations from multiple directions.

▶ Lemma 27 (Complexity of Computing Indegree). Let $\sigma \in K$ be an *i*-simplex, and suppose that $s \in \mathbb{S}^{d-1}$ is perpendicular to σ , and no other vertices of K have the same s-coordinate as σ . Then the indegree of σ with respect to s can be computed using $2^{i+1} - 1$ verbose persistence diagrams or verbose Betti functions.

The proof combines ideas from Algorithm 1 and Theorem 39 of [10].

A.2 Challenges Computing Indegree with other Descriptor Types

As detailed in [10], if we choose to use verbose Euler characteristic functions rather than verbose persistence diagrams or verbose Betti functions, we can no longer compute indegree. Instead, we can only compute *even-/odd-degree*, or the number of even- (odd-) dimensional cofaces of and "below" a given simplex. While mild adaptations to the reconstruction algorithm of [10] allows for this alternate type of query in the reconstruction process, our radial search algorithm (Algorithm 2) unavoidably relies on knowing the dimension of a coface. Namely, if the number of, e.g., even-dimensional cofaces of a given *i*-simplex σ in some cofacet arc matches the number we might expect if all candidates were true cofacets of σ , we still cannot guarantee that all candidates are true cofacets of σ , since lower (or higher) dimensional cofaces may also contribute to this count.

We are not aware of any methods to compute indegree using concise descriptors, other than reconstructing the entirety of K using other methods, and using this information to report indegree (which rather defeats the purpose). Development of such a method seems unlikely. In various senses, concise descriptors have been shown to be weaker than their verbose counterparts [12, 29, 22]. In particular, concise descriptors are sensitive to the "flatness" of a simplicial complex, or how close to affinely-dependent any subset of vertices is. Existing bounds on the size of faithful discretizations of consise directional transforms bound this flatness [6], and without such a bound, there are strict requirements on what directions are necessary [12, Corollary 1].

A.3 Connecting Reconstruction Algorithms to Faithful Discretizations

If a set of topological descriptors can be used to reconstruct a simplicial complex, it is a faithful set. Therefore, if we consider the set of topological descriptors that would be used to perform all our indegree queries (along with a set to reconstruct K_0), we would arrive at a faithful discretization of the associated directional transform, i.e., either the verbose persistent homology transform or the verbose Betti function transform.

In Section 6, we take a similar perspective as [11, 3]; namely, given no initial information about a simplicial complex K other than general position and possibly K_0 , we then aim to

reconstruct K using indegree queries. This perspective is fundamentally different from [10], which, using *total* information about K, defines a set of directions from which K could be reconstructed (i.e., corresponding to a faithful discretization of the associated topological transform). Using initial knowledge of K to define the direction set allows [10] to lessen the impact of unnecessary "exploratory" queries. In particular, when reconstructing K_{i+1} given K_i , the reconstruction algorithm of [10] certifies the presence of each (i + 1)-simplex, and since the total number of (i + 1)-simplices can be read off any single diagram, there is no need to test remaining candidates that are not true (i + 1)-simplices.

Since we assume no initial knowledge of K, unlike [10], we don't know where to look in order to certify only the true (i+1)-simplices; we inevitably encounter and test candidate (i+1)-simplices that are not in K_{i+1} . While this results in "extra" queries from the perspective of purely building a finite faithful set of directions, it is necessary for our framework.