A Satisfiability algorithm based on Simple Spinors of the Clifford algebra of $\mathbb{R}^{n,n}$

Marco Budinich

University of Trieste and INFN, Trieste, Italy mbh@ts.infn.it

January 7, 2025

Abstract

We refine the formulation of the Boolean satisfiability problem with n Boolean variables in Clifford algebra $\mathcal{C}\ell(\mathbb{R}^{n,n})$ [3] and exploit this continuous setting to design a new unsatisfiability test. This algorithm is not combinatorial and proves unsatisfiability in polynomial time.

Keywords: Clifford algebra; simple spinors; satisfiability; orthogonal group.

1 Introduction

Clifford algebra is a remarkably powerful tool initially developed to deal with automorphisms of quadratic spaces [14] that got its major achievements with spinors of mathematical physics. Since then it has been successfully applied to very many different fields, including combinatorial problems [5, 4].

On the other side the Boolean SATisfiability problem is the progenitor of many combinatorial problems and, suprisingly, fits smoothly in the Boolean algebra embedded in $\mathcal{C}\ell(\mathbb{R}^{n,n})$, the Clifford algebra of $\mathbb{R}^{n,n}$, the realm of simple spinors [7]. This is not the first encounter of mathematical physics with SAT: also statistical mechanics applied to SAT brought interesting results [12].

Our first contribution is a solid foundation of Boolean algebra within idempotents of $\mathcal{C}\ell(\mathbb{R}^{n,n})$ and we use this framework to examine different formulations of SAT from a unified standpoint. Subsequently we focus on the continuous formulation of SAT in the group O(n), with its equivalent spinorial and vectorial representations. The second, and main, contribution is to exploit this formulation to design an algorithm for an unsatisfiability test. The core of the algorithm relies on linear combinations of simple spinors induced by the clauses. More in detail in sections 2 and 3 we succintly recall the Boolean Satisfiability problem and the Clifford algebra $\mathcal{C}\ell(\mathbb{R}^{n,n})$. Section 4 provides a sturdy formulation of Boolean algebra with idempotents of $\mathcal{C}\ell(\mathbb{R}^{n,n})$ tailored to our needs while in section 5 we apply these results to give a neat encoding of SAT in $\mathcal{C}\ell(\mathbb{R}^{n,n})$ together with a first unsatisfiability condition. To prepare the SAT formulation in a more specific setting in section 6 we introduce, with a simple formalism, the isomorphism between the set of all totally null subspaces of maximal dimension n of $\mathbb{R}^{n,n}$ and the continuous group O(n). In section 7 we use this to transform a SAT problem into the problem of forming a cover for O(n) that gives a continuous formulation of SAT [3] with another unsatisfiability condition. In final section 8, subdivided in three parts, we build on previous results to produce an unsatisfiability test that exploits the linear space properties of the simple spinors of $\mathcal{C}\ell(\mathbb{R}^{n,n})$, in one to one correspondence with O(n). This test is not combinatorial and a simple analysis indicates that it is polynomial.

For the convenience of the reader we tried to make this paper as elementary and self-contained as possible.

2 The Boolean Satisfiability problem

The Boolean Satisfiability Problem [12, Section 7.2.2.2] seeks an assignment of *n* Boolean variables $\rho_i \in \{T, F\}$ (true, false), that makes T, *satisfies*, a given Boolean formula S put in Conjunctive Normal Form (CNF) e.g.

$$\mathcal{S} \equiv (\rho_1 \vee \overline{\rho}_2) \land (\rho_2 \vee \rho_3) \land (\overline{\rho}_1 \vee \overline{\rho}_3) \land (\overline{\rho}_1 \vee \overline{\rho}_2 \vee \rho_3) \land (\rho_1 \vee \rho_2 \vee \overline{\rho}_3) \quad (1)$$

as a logical AND (\wedge) of *m* clauses C_j , the expressions in parenthesis, each clause being composed by the logical OR (\vee) of *k* or less *literals* possibly preceded by logical NOT ($\neg \rho_i, \overline{\rho_i}$ for short). In (1) n = 3, m = 5 and k = 3. To underline the difference with algebraic equality = in what follows we use \equiv to represent logical equivalence, namely that for all possible values taken by the Boolean variables the two expressions are equal. A solution is either an assignment of ρ_i that gives $S \equiv T$ or a proof that such an assignment does not exist and $S \equiv F$ in all cases.

SAT was the first combinatorial problem proven to be NP-complete [8]; in particular while the case of k = 3, 3SAT, can be solved only in a time that grows exponentially with n, 2SAT and 1SAT problems can be solved in polynomial time, that is *fast*.

Using the distributive properties of the logical operators \lor , \land any given kSAT S expands in a logical OR of up to k^m terms each term being a 1SAT problem namely a logical AND of m Boolean variables. Since $\rho_i \land \overline{\rho_i} \equiv F$ the presence of a variable together with its logical complement is a necessary and sufficient condition for making a 1SAT formula F, namely *unsatisfiable*, and thus these terms can be omitted and so k^m is just an upper bound to

the number of terms. Conversely a satisfiable 1SAT formula has only one assignment of its Boolean variables that makes it T and that can be read scanning the formula; thus in the sequel we will freely use 1SAT formulas for assignments.

The final expanded expression can be further simplified and reordered exploiting the commutativity of the logical operators \lor , \land and the properties $\rho_i \land \rho_i \equiv \rho_i \lor \rho_i \equiv \rho_i$. All "surviving" terms of this expansion, the Disjunctive Normal Form (DNF), are 1SAT terms, each of them representing an assignment that satisfies the problem. On the contrary if the DNF is empty, as happens for (1), this is a proof that there are no assignments that make the formula T: the problem is unsatisfiable.

Expansion to DNF is a dreadful algorithm for solving SAT: first of all the method is an overkill since it produces all possible solutions whereas one would be enough; in second place this brute force approach gives a running time proportional to the number of expansion terms $\mathcal{O}\left(\left(k^{\frac{m}{n}}\right)^n\right)$ whereas a modern SAT solver [13] run in $\mathcal{O}(1.307^n)$. Nevertheless DNF will play a central role in the formulation of SAT in Clifford algebra.

3 $\mathbb{R}^{n,n}$ and its Clifford algebra

We review some properties of $\mathbb{R}^{n,n}$ and of its Clifford algebra $\mathcal{C}\ell(\mathbb{R}^{n,n})$ that are at the heart of the following results.

 $\mathcal{C}\ell(\mathbb{R}^{n,n})$ is isomorphic to the algebra of real matrices $\mathbb{R}(2^n)$ [14] and this algebra is more easily manipulated exploiting the properties of its Extended Fock basis (EFB, see [2] and references therein) with which any algebra element is a linear superposition of simple spinors. The 2n generators of the algebra e_i form an orthonormal basis of the linear space $\mathbb{R}^{n,n}$

$$e_i e_j + e_j e_i := \{e_i, e_j\} = 2 \begin{cases} \delta_{ij} & \text{for } i \le n \\ -\delta_{ij} & \text{for } i > n \end{cases}$$
 $i, j = 1, 2, \dots, 2n$ (2)

and we define the Witt, or null, basis of $\mathbb{R}^{n,n}$:

$$\begin{cases} p_i = \frac{1}{2}(e_i + e_{i+n}) \\ q_i = \frac{1}{2}(e_i - e_{i+n}) \end{cases} \quad i = 1, 2, \dots, n$$
(3)

that, with $e_i e_j = -e_j e_i$ for $i \neq j$, gives

$$\{p_i, p_j\} = \{q_i, q_j\} = 0 \qquad \{p_i, q_j\} = \delta_{ij} \tag{4}$$

showing that all p_i, q_i are mutually orthogonal, also to themselves, that implies $p_i^2 = q_i^2 = 0$ and are thus null vectors. Defining

$$\begin{cases} P = \operatorname{Span}(p_1, p_2, \dots, p_n) \\ Q = \operatorname{Span}(q_1, q_2, \dots, q_n) \end{cases}$$
(5)

P and Q are two totally null subspaces of maximum dimension n and form a Witt decomposition [14] of $\mathbb{R}^{n,n}$ since $P \cap Q = \{0\}$ and $P \oplus Q = \mathbb{R}^{n,n}$.

The 2^{2n} simple spinors forming EFB are given by all possible sequences

$$\psi = \psi_1 \psi_2 \cdots \psi_i \cdots \psi_n \qquad \psi_i \in \{q_i p_i, p_i q_i, p_i, q_i\} \qquad i = 1, 2, \dots, n \qquad (6)$$

where each ψ_i takes one of its 4 possible values [2] and each ψ_i is uniquely identified by two "bits" $h_i, g_i = \pm 1$: $h_i = 1$ if the leftmost vector of ψ_i is q_i , -1 otherwise; $g_i = 1$ if ψ_i is even, -1 if odd. The *h* and *g* signatures of ψ are respectively the vectors (h_1, h_2, \ldots, h_n) and (g_1, g_2, \ldots, g_n) .

Since $e_i e_{i+n} = q_i p_i - p_i q_i := [q_i, p_i]$ in EFB the identity 1 and the volume element ω (scalar and pseudoscalar) assume similar expressions [2]:

$$1 := \{q_1, p_1\} \{q_2, p_2\} \cdots \{q_n, p_n\} \omega := e_1 e_2 \cdots e_{2n} = (-1)^{\frac{n(n-1)}{2}} [q_1, p_1] [q_2, p_2] \cdots [q_n, p_n]$$
(7)

and since $\mathcal{C}\ell(\mathbb{R}^{n,n})$ is a simple algebra, the algebra identity is also the sum of its 2^n primitive (indecomposable) idempotents \mathbb{p}_i we gather in set \mathbb{P}

$$\mathbb{1} = \sum_{i=1}^{2^n} \mathbb{p}_i \qquad \mathbb{p}_i \in \mathbb{P} \quad . \tag{8}$$

Comparing the two expressions of 1 we observe that the full expansion of the anticommutators of (7) contains 2^n terms each term being one of the primitive idempotents *and* a simple spinor. \mathbb{P} is thus a proper subset of EFB (6) and its elements are

$$\mathbb{p} = \psi_1 \psi_2 \cdots \psi_i \cdots \psi_n \qquad \psi_i \in \{q_i p_i, p_i q_i\} \qquad i = 1, 2, \dots, n \quad . \tag{9}$$

We recall the standard properties of primitive idempotents

$$\mathbf{p}_i^2 = \mathbf{p}_i \quad (\mathbb{1} - \mathbf{p}_i)^2 = \mathbb{1} - \mathbf{p}_i \quad \mathbf{p}_i(\mathbb{1} - \mathbf{p}_i) = 0 \quad \mathbf{p}_i \mathbf{p}_j = \delta_{ij} \mathbf{p}_i \tag{10}$$

and define the set

$$\mathcal{I} := \left\{ \sum_{i=1}^{2^n} \delta_i \mathbb{p}_i : \delta_i \in \{0, 1\}, \mathbb{p}_i \in \mathbb{P} \right\}$$
(11)

in one to one correspondence with the power set of \mathbb{P} . \mathcal{I} is closed under Clifford product but not under addition and is thus not even a subspace. With (10) we easily prove

Proposition 1. For any $s \in \mathcal{I}$ then $s^2 = s$.

 \mathcal{I} is thus the set of the idempotents, in general not primitive; a simple consequence is that for any $s \in \mathcal{I}$ also $(\mathbb{1} - s) \in \mathcal{I}$.

Any EFB element of (6) is a simple spinor, uniquely identified in EFB by its h signature, while the minimal left ideal, or spinor space S, to which it belongs is identified by its $h \circ g = (h_1g_1, h_2g_2, \ldots, h_ng_n)$ signature [2]. The algebra, as a linear space, is the direct sum of these 2^n spinor spaces that, in isomorphic matrix algebra $\mathbb{R}(2^n)$, are usually associated to linear spaces of matrix columns.

For each of these 2^n spinor spaces \mathbb{S} its 2^n simple spinors (6) [2] form a Fock basis \mathcal{F} and any spinor $\psi \in \mathbb{S}$ is a linear combination of the simple spinors $\psi_{\lambda} \in \mathcal{F}$ [6, 2], namely

$$\psi = \sum_{\lambda} \alpha_{\lambda} \psi_{\lambda} \qquad \alpha_{\lambda} \in \mathbb{R}, \quad \psi_{\lambda} \in \mathcal{F} \quad .$$
(12)

We illustrate this with the simplest example in $\mathbb{R}^{1,1}$, the familiar Minkowski plane of physics, here $\mathcal{C}\ell(\mathbb{R}^{1,1}) \cong \mathbb{R}(2)$ and the EFB (6) is formed by just 4 elements: $\{qp_{++}, pq_{--}, p_{-+}, q_{+-}\}$ with the subscripts indicating respectively h and $h \circ g$ signatures that give the binary form of the integer matrix indexes; its EFB matrix is

$$+ - + \begin{pmatrix} qp & q \\ p & pq \end{pmatrix}$$

and, as anticipated, we can write the generic element $\mu \in \mathcal{C}\ell(\mathbb{R}^{1,1})$ in EFB

$$\mu = \xi_{++}qp_{++} + \xi_{--}pq_{--} + \xi_{-+}p_{-+} + \xi_{+-}q_{+-} \qquad \xi \in \mathbb{R}$$

The two columns are two minimal left ideals namely two (equivalent) spinor spaces S_+ and S_- . The two elements of each column are the simple spinors of Fock basis \mathcal{F} while qp and pq are the primitive idempotents and qp+pq=1.

In turn simple spinors of a Fock basis \mathcal{F} are in one to one correspondence with $\mathbb{R}^{n,n}$ null subspaces of maximal dimension n. For any $\psi \in \mathcal{F}$ we define its associated maximal null subspace $M(\psi)$ as

$$M(\psi) = \text{Span}(x_1, x_2, \dots, x_n) \quad x_i = \begin{cases} p_i & \text{iff } \psi_i = p_i, p_i q_i \\ q_i & \text{iff } \psi_i = q_i, q_i p_i \end{cases} \quad i = 1, 2, \dots, n$$
(13)

and x_i is determined by the *h* signature of ψ in EFB [6, 2]. For example in $\mathcal{C}\ell(\mathbb{R}^{3,3})$ given the simple spinor $\psi = p_1q_1q_2p_2q_3p_3$

$$\psi = p_1 q_1 q_2 p_2 q_3 p_3 \quad \Longrightarrow \quad M(\psi) = \operatorname{Span}\left(p_1, q_2, q_3\right)$$

and with (4) we see that for any $v \in M(\psi)$ then $v\psi = 0$.

We gather these 2^n maximal null subspaces of $\mathbb{R}^{n,n}$ in set \mathcal{M}_n each of its elements being the span of the *n* null vectors obtained choosing one null vector from each couple (p_i, q_i) (3). The set \mathcal{M}_n is the same for all the 2^n different possible spinor spaces being identified by the h signatures of ψ in EFB and so it can be defined also starting from primitive idempotents (9) and in summary we can give three equivalent definitions for \mathcal{M}_n

$$\mathcal{M}_{n} = \begin{cases} \{M(\psi) : \psi \in \mathcal{F}\} \\ \{\operatorname{Span}(x_{1}, x_{2}, \dots, x_{n}) : x_{i} \in \{p_{i}, q_{i}\}\} \\ \{M(p) : p \in \mathbb{P}\} \end{cases}$$
(14)

4 The Boolean algebra of $\mathcal{C}\ell(\mathbb{R}^{n,n})$

We exploit the known fact that in any associative, unital, algebra every family of commuting, orthogonal, idempotents generates a Boolean algebra to prove that the 2^{2^n} idempotents of \mathcal{I} (11) form a Boolean algebra.

A finite Boolean algebra is a set equipped with the inner operations of logical AND, OR and NOT that satisfy well known properties but we will use an axiomatic definition [11, 9] that needs only a binary and a unary inner operations satisfying three simple axioms to prove:

Proposition 2. The set \mathcal{I} equipped with the two inner operations

is a finite Boolean algebra.

Proof. We already observed that \mathcal{I} is closed under operations (15) that moreover satisfy Boolean algebra axiomatic definition [11]: the binary operation is associative since Clifford product is and commutative because all \mathcal{I} elements commute. The third (Huntington's) axiom requires that for any $s_1, s_2 \in \mathcal{I}$

$$(\mathbb{1} - (\mathbb{1} - s_1)s_2)(\mathbb{1} - (\mathbb{1} - s_1)(\mathbb{1} - s_2)) = s_1$$

that is easily verified. \Box

We remark that \mathcal{I} is not a subalgebra of $\mathcal{C}\ell(\mathbb{R}^{n,n})$ since it is not closed under addition. Any finite Boolean algebra is isomorphic to the power set of its Boolean atoms [9]. In this case \mathcal{I} elements are in one to one correspondence with the power set of \mathbb{P} and we thus identify the Boolean atoms with the 2^n primitive idempotents (9).

With simple manipulations we get all Boolean expressions in $\mathcal{C}\ell(\mathbb{R}^{n,n})$: in the unary operation of (15) we recognize the logical NOT and associating the logical AND to Clifford product from $s(\mathbb{1} - s) = 0 \in \mathcal{I}$ we deduce that 0 stands for F and consequently that $\mathbb{1}$ stands for T. For the logical OR we use De Morgan's relations

$$\rho_1 \vee \rho_2 \equiv \overline{\rho_1 \wedge \overline{\rho_2}} \quad \to \quad \mathbb{1} - (\mathbb{1} - s_1)(\mathbb{1} - s_2) = s_1 + s_2 - s_1 s_2$$

and we can easily verify that \mathcal{I} is closed also under this binary operation.

We formulate Boolean expressions in $\mathcal{C}\ell(\mathbb{R}^{n,n})$ associating Boolean variables ρ_i to idempotents and we gather associations in this table where p_i and q_i are vectors of the Witt basis (3)

$$\begin{array}{rcl}
\mathbf{F} & \to & 0\\
\mathbf{T} & \to & 1\\
\rho_i & \to & q_i p_i\\
\overline{\rho}_i & \to & 1 - q_i p_i = p_i q_i\\
\rho_i \wedge \rho_j & \to & q_i p_i q_j p_j\\
\rho_i \vee \rho_j & \to & q_i p_i + q_j p_j - q_i p_i q_j p_j \end{array} \tag{16}$$

For example given some simple Boolean expressions with (4) we easily verify

and from now on we will use ρ_i and $\overline{\rho}_i$ also in $\mathcal{C}\ell(\mathbb{R}^{n,n})$ meaning respectively $q_i p_i$ and $p_i q_i$ and Clifford product will stand for logical AND \wedge and in full generality we can prove [4]

Proposition 3. Any Boolean expression S with n Boolean variables is represented in $\mathcal{C}\ell(\mathbb{R}^{n,n})$ by $S \in \mathcal{I}$ obtained with substitutions (16) moreover \overline{S} is represented by 1 - S both being idempotents of $\mathcal{C}\ell(\mathbb{R}^{n,n})$. Given another Boolean expression Q the logical equivalence $S \equiv Q$ holds if and only if S = Q for their respective idempotents in $\mathcal{C}\ell(\mathbb{R}^{n,n})$.

In summary with substitutions (16) we can safely encode any Boolean expression, and thus SAT problems, in Clifford algebra.

5 SAT in Clifford algebra $\mathcal{C}\ell(\mathbb{R}^{n,n})$

The straightest way of encoding a SAT problem in CNF (1) in Clifford algebra is exploiting De Morgan relations to rewrite its clauses as

$$\mathcal{C}_j \equiv (\rho_{j_1} \lor \rho_{j_2} \lor \cdots \lor \rho_{j_k}) \equiv \overline{\rho_{j_1}} \overline{\rho_{j_2}} \cdots \overline{\rho}_{j_k}$$

and thus the expression of a clause in Clifford algebra is

$$\mathcal{C}_j \to \mathbb{1} - \overline{\rho}_{j_1} \overline{\rho}_{j_2} \cdots \overline{\rho}_{j_k} := \mathbb{1} - z_j \tag{17}$$

and the expression of a SAT problem in CNF with m clauses is

$$S = \prod_{j=1}^{m} (\mathbb{1} - z_j) \tag{18}$$

and from Proposition 3 easily descends

Proposition 4. Given a SAT problem S then $S \equiv F$ if and only if, for the corresponding algebraic expression in $\mathcal{C}\ell(\mathbb{R}^{n,n})$ (18) S = 0

that transforms a Boolean problem in an algebraic one. To master the implications of (18) we need the full expression of a 1SAT formula e.g. $\rho_1 \overline{\rho}_2$ namely $q_1 p_1 p_2 q_2$: with (7), (8) and (9)

$$q_1 p_1 \ p_2 q_2 = q_1 p_1 \ p_2 q_2 \ \mathbb{1} = q_1 p_1 \ p_2 q_2 \prod_{j=3}^n \{q_j, p_j\}$$
(19)

since $q_1p_1 \{q_1, p_1\} = q_1p_1$ and $p_2q_2 \{q_2, p_2\} = p_2q_2$ and the full expansion of this expression is the sum of 2^{n-2} primitive idempotents $\mathbb{p}(9)$ and thus $q_1p_1 p_2q_2$ is an idempotent of \mathcal{I} . From the Boolean standpoint this can be interpreted as the property that given the 1SAT formula $\rho_1\overline{\rho}_2$ the other, unspecified, n-2 Boolean variables ρ_3, \ldots, ρ_n can take all possible 2^{n-2} values or, more technically, that $\rho_1\overline{\rho}_2$ has a *full* DNF made of 2^{n-2} Boolean atoms.

More in general any 1SAT formula with m Boolean variables is a sum of 2^{n-m} primitive idempotents, namely Boolean atoms. With (16) we can rewrite (9) as

$$\mathbf{p} = \psi_1 \psi_2 \cdots \psi_i \cdots \psi_n \qquad \psi_i \in \{\rho_i, \overline{\rho}_i\} \qquad i = 1, 2, \dots, n$$

showing that the 2^n primitive idempotents p are just the possible 2^n 1SAT formulas with n Boolean variables, the Boolean atoms, for example:

$$\rho_1 \overline{\rho}_2 \rho_3 \cdots \rho_n \to q_1 p_1 \ p_2 q_2 \ q_3 p_3 \cdots q_n p_n \in \mathbb{P} \quad . \tag{20}$$

By Proposition 3 $S \in \mathcal{I}$ (11) and is thus the sum of primitive idempotents (9) that now we know represent Boolean atoms and ultimately (18) gives the full DNF expansion of the SAT problem S each term being one assignment that makes the problem T while if the expansion is empty the problem is unsatisfiable and thus expansion of the CNF S of (18) reproduces faithfully the Boolean expansion to DNF outlined in section 2.

From the computational side Proposition 4 is not a big deal since the expansion of (18) corresponds to the DNF expansion that in section 2 we named a "dreadful" algorithm. But porting SAT to Clifford algebra offers other advantages since we can exploit algebra properties. For example the unsatisfiability condition S = 0 makes S a scalar whereas if satisfiable S is not a scalar. Exploiting scalar properties in Clifford algebra we proved [4]

Theorem 1. A given nonempty SAT problem in $\mathcal{C}\ell(\mathbb{R}^{n,n})$ (18) is unsatisfiable (S = 0) if and only if, for all generators (2) of $\mathcal{C}\ell(\mathbb{R}^{n,n})$

$$e_i S e_i^{-1} = S \qquad \forall \ 1 \le i \le 2n \quad . \tag{21}$$

This result gives an unsatisfiability test based on the symmetry properties of its CNF expression S (18). We remark that as far as computational performances are concerned an efficient unsatisfiability test would bring along also an efficient solution algorithm. Suppose the test (21) fails and thus that S is satisfiable, to get an actual solution we choose a Boolean variable, e.g. ρ_i , and replace it with T and apply again the test to the derived problem S_i . If the test on S_i fails as well this means that $\rho_i \equiv T$ otherwise, necessarily, $\rho_i \equiv F$ and repeating this procedure n times for all Boolean variables we obtain an assignment that satisfies S. The algorithmic properties of unsatisfiability test (21) have been preliminarly explored in [4].

 $\mathcal{C}\ell(\mathbb{R}^{n,n})$ epitomizes the geometry of linear space $\mathbb{R}^{n,n}$ and thus SAT encoding (18) brings along also a geometric interpretation that is at the root of further encodings of SAT in Clifford algebra.

S (18) is ultimately a sum of primitive idempotents (9) that are in one to one correspondence with the null maximal subspaces of \mathcal{M}_n (14).

It follows that S, and more in general any \mathcal{I} element, induces a subset of \mathcal{M}_n , the empty subset if S = 0. More precisely the elements of this subset are all and only those maximal totally null subspaces (14) corresponding to the Boolean atoms making S. For any $s \in \mathcal{I}$ (11) let I_s such that

$$s = \sum_{i \in I_s} \mathbb{p}_i \qquad I_s \subseteq \{1, 2, \dots 2^n\}$$

$$(22)$$

and so all $s \in \mathcal{I}$ induce a subset of \mathcal{M}_n

$$\mathcal{T}'_{s} := \{ M(\mathbb{p}_{i}) : i \in I_{s} \} \subseteq \mathcal{M}_{n} \implies \mathcal{T}'_{\mathbb{1}-s} = \mathcal{M}_{n} - \mathcal{T}'_{s} \quad .$$
(23)

Applying this definition to clauses idempotents (17) Proposition 4 becomes:

Proposition 5. Given a SAT problem S in $C\ell(\mathbb{R}^{n,n})$ (18) then the problem is unsatisfiable (S = 0) if and only if

$$\bigcup_{j=1}^{m} \mathcal{T}'_{z_j} = \mathcal{M}_n \quad . \tag{24}$$

Proof. For any $s_1, s_2 \in \mathcal{I}$ from \mathcal{I} definition (11) we easily get

$$\mathcal{T}'_{s_1s_2} = \mathcal{T}'_{s_1} \cap \mathcal{T}'_{s_2}$$

and in this setting Proposition 4 S states that S = 0 if and only if

$$\mathcal{T}'_{S} = \bigcap_{j=1}^{m} \mathcal{T}'_{\mathbb{1}-z_{j}} = \emptyset \quad . \tag{25}$$

The thesis follows by (23) and by elementary set properties. \Box

The SAT problem has now the form of a problem of subsets of \mathcal{M}_n that provides also an interpretation of (18). Since z_j is the unique assignment

of the k literals of \mathcal{C}_j that give $\mathcal{C}_j \equiv \mathbf{F}$, and thus S = 0, then if the union of all these cases (24) covers \mathcal{M}_n the problem is unsatisfiable. This in turn implies that any $M(\mathbb{p}) \notin \bigcup_{j=1}^m \mathcal{T}'_{z_j}$ is a solution of S.

Given z_j with k literals we define $M(z_j)$ as the k dimensional null subspace (13) induced by the literals of z_j , and adopting the lighter notation \mathcal{T}'_j for \mathcal{T}'_{z_i} , it is easy to see that

$$\mathcal{T}'_j := \{ M(\mathfrak{p}_i) : M(z_j) \subseteq M(\mathfrak{p}_i) \} \subseteq \mathcal{M}_n \quad .$$
(26)

To proceed further we review the isomorphism between the set of all totally null subspaces of maximal dimension of $\mathbb{R}^{n,n}$ and the group O(n).

6 The orthogonal group O(n) and the set \mathcal{N}_n

Let \mathcal{N}_n be the set of all totally null subspaces of maximal dimension n of $\mathbb{R}^{n,n}$, a quadric Grassmannian for Ian Porteous [14, Chapter 14]. \mathcal{N}_n is isomorphic to subgroup O(n) of O(n, n) and O(n) acts transitively on \mathcal{N}_n .

We review these relations: seeing the linear space $\mathbb{R}^{n,n}$ as $\mathbb{R}^n \times \mathbb{R}^n$ we can write its generic element as (x, y) and $(x, y)^2 = x^2 - y^2$. Any *n* dimensional subspace of $\mathbb{R}^{n,n}$ may be represented as the image of an injective map $\mathbb{R}^n \to \mathbb{R}^n \times \mathbb{R}^n$; $x \to (s(x), t(x))$ for $s, t \in \operatorname{GL}(n)$. This subspace is made by all pairs (s(x), t(x)) and we denote it with $(s, t) \in \operatorname{GL}(n) \times \operatorname{GL}(n)$. By same mechanism, with s = 1 and $t \in O(n)$, for any $x \in \mathbb{R}^n (x, t(x)) \in \mathbb{R}^{n,n}$ is a null vector since $(x, t(x))^2 = x^2 - t(x)^2 = 0$ and it belongs to the *n* dimensional null subspace (1, t). Given that from now on $s \equiv 1$ it is natural to identify \mathbb{R}^n as $\mathbb{R}^n \times \{0\}$, the spacelike subspace of $\mathbb{R}^{n,n}$, and we will do so unless differently specified.

Isometries (orthogonal transformations) $t \in O(n)$ establish the quoted isomorphism since any subspace $(\mathbb{1}, t) \subset \mathbb{R}^{n,n}$ is in \mathcal{N}_n and conversely any element of \mathcal{N}_n can be written as $(\mathbb{1}, t)$ [14, Corollary 14.13] and thus

$$\mathcal{N}_n = \{(\mathbb{1}, t) : t \in \mathcal{O}(n)\}\tag{27}$$

and the isomorphism between \mathcal{N}_n and O(n) is realized by map

$$\mathcal{N}_n \to \mathcal{O}(n); (\mathbb{1}, t) \to t$$
 . (28)

For example, assuming that the map $(\mathbb{1},\mathbb{1}): \mathbb{R}^n \to \mathbb{R}^n \times \mathbb{R}^n$ is such that $e_i \to (e_i, e_{i+n})$, then two generic null vectors of P and Q (5) are respectively (x, x) and (y, -y) and in this notation P and Q are thus

$$\begin{cases}
P = (1, 1) \\
Q = (1, -1)
\end{cases}$$
(29)

The action of O(n) is transitive on \mathcal{N}_n since for any $t, u \in O(n)$, $(\mathbb{1}, ut) \in \mathcal{N}_n$ and the action of O(n) is trivially transitive on O(n). We examine isomorphism (28) when restricted to subset $\mathcal{M}_n \subset \mathcal{N}_n$ (14) taking $P = (\mathbb{1}, \mathbb{1})$ as our "reference" element of \mathcal{M}_n . Let $\lambda_i \in O(n)$ be the hyperplane reflection inverting timelike vector e_{i+n} , namely

$$\lambda_i(e_j) = \begin{cases} -e_j & \text{for } j = i+n \\ e_j & \text{otherwise} \end{cases} \quad i = 1, 2, \dots, n \quad j = n+1, n+2, \dots, 2n$$

its action on the Witt basis (3) exchanges the null vectors p_i and q_i . Starting from P (5) we can get any other \mathcal{M}_n element inverting a subset of the n timelike vectors e_{i+n} . Each isometry λ_i is represented, in the vectorial representation of O(n), by a diagonal matrix $\lambda \in \mathbb{R}(n)$ with ± 1 on the diagonal and these matrices form the group

$$O(1) \times O(1) \cdots \times O(1) = \stackrel{n}{\times} O(1) := O^{n}(1)$$

immediate to get since $O(1) = \{\pm 1\}$. $O^n(1)$ is a discrete, abelian, subgroup of involutions of O(n), namely linear maps t such that $t^2 = 1$. It is thus clear that since P = (1, 1) for any $M(p) \in \mathcal{M}_n$ there exists a unique $\lambda \in O^n(1)$ such that

$$M(\mathbf{p}) = (\mathbb{1}, \lambda)$$

and thus we proved constructively

Proposition 6. Isomorphism (28) restricted to $\mathcal{M}_n \subset \mathcal{N}_n$ has for image subgroup $O^n(1)$ of O(n)

$$\mathcal{M}_n = \{(\mathbb{1}, \lambda) : \lambda \in O^n(1)\} \qquad \Longrightarrow \qquad \mathcal{M}_n \to O^n(1); (\mathbb{1}, \lambda) \to \lambda \quad (30)$$

Given reference P let $\psi_1 = p_1 q_1 \ p_2 q_2 \ p_3 q_3 \cdots p_n q_n$ be the *reference* simple spinor, such that $M(\psi_1) = P = (1, 1)$, the vacuum spinor of physics, we resume concisely the action of $\lambda \in O^n(1)$ on spinors and vectors with (see e.g. [6, 2] for more extensive treatments)

$$M(\lambda(\psi_1)) := M(\psi_\lambda) = (1, \lambda) \tag{31}$$

and by the action of λ we get respectively from $\psi_{\mathbb{1}}$ all spinors of the Fock basis \mathcal{F} and from null subspace P all \mathcal{M}_n elements.

Isomorphism (30) adds a fourth definition of \mathcal{M}_n (14) with which we can port SAT within group $O^n(1)$ and, later, O(n). We redefine \mathcal{T}'_s (23) as a subset of $O^n(1)$

$$\mathcal{T}'_s := \{\lambda \in \mathcal{O}^n(1) : (\mathbb{1}, \lambda) = M(\mathbb{p}_i), i \in I_s\} \subset \mathcal{O}^n(1)$$
(32)

and with this definition we can transform Proposition 5 to

Proposition 7. Given a SAT problem S in $C\ell(\mathbb{R}^{n,n})$ (18) then the problem is unsatisfiable (S = 0) if and only if

$$\cup_{j=1}^{m} \mathcal{T}'_j = O^n(1) \tag{33}$$

that gives the first formulation of SAT problems in group language. From the computational point of view of there are no improvements since $O^n(1)$ is a discrete group and checking if subsets \mathcal{T}'_j form a cover essentially requires testing all 2^n group elements, just the same as testing all 2^n Boolean atoms to see if any solves SAT.

We resume all this in a commutative diagram in which numbers refer to formulas

$$\begin{array}{c}
\mathbb{p} \in \mathbb{P} \\
\stackrel{(14)}{\swarrow} & \stackrel{(31)}{\searrow} \\
M(\mathbb{p}) \in \mathcal{M}_n & \stackrel{(30)}{\longleftrightarrow} \lambda \in \mathcal{O}^n(1)
\end{array}$$

that authorizes us, from now on, to freely switch between:

- the 2^n primitive idempotents p, or Boolean assignments (atoms) of n Boolean variables,
- the 2^n simple spinors of the Fock basis $\psi_{\lambda} \in \mathcal{F}$,
- the 2^n totally null subspaces of maximal dimension $M(\psi_{\lambda}) \in \mathcal{M}_n$,
- the 2^n elements of the discrete, abelian group $\lambda \in O^n(1)$.

7 SAT in orthogonal group O(n)

Before going on we remind some basics properties of spinor space S, a minimal left ideal of $\mathcal{C}\ell(\mathbb{R}^{n,n})$, and the particular case of *simple spinors* that we will indicate with $\mathbb{S}_s \subset \mathbb{S}$. Simple spinors [7] are an elusive subject that rarely surfaces in recent literature, a noteworthy exception being [6]. In a nutshell: we saw that to $\psi_{\lambda} \in \mathcal{F}$ are associated $M(\psi_{\lambda}) = (\mathbb{1}, \lambda) \in \mathcal{M}_n$ and more in general simple spinors are those spinors that are associated to a null subspace of maximal dimension n, namely

$$\psi_t \in \mathbb{S}$$
 such that $M(\psi_t) = (\mathbb{1}, t) \in \mathcal{N}_n$ $t \in \mathcal{O}(n)$ (34)

this relation between simple spinors and O(n) being bijective [6].

In the last step we show that, when problem S is unsatisfiable, subsets induced by clauses not only form a cover of $O^n(1)$ (33) but also of its parent group O(n) that opens new computational perspectives and we summarize here the needed results of [3] to which the reader is addressed for a more exhaustive treatment. We start extending the definition of isometries induced by a clause (26) with (32) to

$$\mathcal{T}_j := \{ t \in \mathcal{O}(n) : M(z_j) \subseteq (\mathbb{1}, t) \} \subset \mathcal{O}(n)$$
(35)

this being an obvious generalization of (26), moreover $O^n(1) \subset O(n)$ implies $\mathcal{T}'_j \subseteq \mathcal{T}_j$ and we can give [3] different equivalent definitions for \mathcal{T}_j e.g.

$$\mathcal{T}_{j} = \begin{cases} \{t \in \mathcal{O}(n) : M(z_{j}) \subseteq (\mathbb{1}, t)\} \subset \mathcal{O}(n) \\ \{\psi_{t} \in \operatorname{Span}(\psi_{\lambda}) : \lambda \in \mathcal{T}'_{j} \subset \mathcal{F} \text{ and } \psi_{t} \in \mathbb{S}_{s} \text{ with } M(\psi_{t}) = (\mathbb{1}, t)\} \subset \mathbb{S}_{s} \\ \{t \in \mathcal{O}(n) : (\mathbb{1}, t) = M(\psi_{t}) \text{ for } \psi_{t} \text{ as above} \end{cases}$$

and moreover for any \mathcal{T}_j then

$$\mathcal{T}_j \cap \mathrm{O}^n(1) = \mathcal{T}'_j$$
 .

Given \mathcal{T}_j definition as the span of the subset of the Fock basis given by \mathcal{T}'_j and being spinor space S a linear space, we can define the set

$$\mathcal{T}_{j} + \mathcal{T}_{k} := \left\{ \begin{aligned} & \alpha, \beta \in \mathbb{R} \\ \psi = \alpha \psi_{j} + \beta \psi_{k} : & \psi_{j} \in \mathcal{T}_{j}, \quad \psi_{k} \in \mathcal{T}_{k}, \\ & \text{such that} \quad \psi \in \mathbb{S}_{s} \end{aligned} \right\}$$
(36)

with which [3]


Theorem 2. A given SAT problem S in $C\ell(\mathbb{R}^{n,n})$ (18) with n Boolean variables is unsatisfiable if and only if the isometries induced by its m clauses (35), (36) form a cover for O(n):

$$\sum_{j=1}^{m} \mathcal{T}_j = O(n) \quad . \tag{37}$$

Here $\sum_{j=1}^{m} \mathcal{T}_j$ does not imply an addition between sets \mathcal{T} 's but only stands for the set of spinors linear combinations of spinors taken from different \mathcal{T} 's.

We remark that this theorem is not a straightforward generalization of Proposition 7 and that definition (36) is pivotal: replacing \sum with \cup in (37) the result does not hold: for example the SAT problem with 2 Boolean variables given by the 4 clauses: $\rho_1\rho_2$, $\rho_1\overline{\rho}_2$, $\overline{\rho}_1\rho_2$, $\overline{\rho}_1\overline{\rho}_2$ is clearly unsatisfiable but the corresponding $2^2 = 4$ diagonal matrices of $\mathbb{R}(2)$ satisfy Proposition 7 but do not form a cover of O(2).

With these (and following) results we can port the commutative diagram of Section 6 to the continuous case and to simple spinors



that authorizes us, from now on, to *freely switch* between:

- simple spinors $\psi_t \in \mathbb{S}_s$,
- isometries $t \in O(n)$,
- totally null subspaces of maximal dimension $(1, t) \in \mathcal{N}_n$.

8 An unsatisfiability test made with simple spinors: the idea

To apply the new formulation to the computational side of SAT we focus our attention to the continuous setting of Theorem 2 where we need to prove that with $\sum_{j=1}^{m} \mathcal{T}_j$ we can cover O(n) since we already remarked [3] that an unsatisfiability test exploiting Proposition 7 offers no real advantages.

A known characteristic of SAT problems is that while checking if a single assignment is in $\bigcup_{j=1}^{m} \mathcal{T}'_{j}$ is polynomial (easy), to give a proof that all 2^{n} assignments are in this set, providing an unsatisfiability certificate, can require $\mathcal{O}(2^{n})$ tests (hard).

In the continuous setting of Theorem 2 we can also presume that checking if a single $t \in O(n)$ is in $\sum_{j=1}^{m} \mathcal{T}_{j}$ is easy. But in O(n) things are quite different and we show that, for some t, just two of these tests can provide a certificate of unsatisfiability.

With isomorphisms (34) to any $t \in O(n)$ corresponds the simple spinor $\psi_t \in \mathbb{S}_s$ such that $M(\psi_t) = (\mathbb{1}, t)$ and this spinor can be expanded in Fock basis \mathcal{F} (12) and we define its *support*

$$\sup \psi_t := \{\psi_\lambda \in \mathcal{F} : \alpha_\lambda \neq 0 \text{ in } (12)\} \subseteq \mathcal{F}$$
(38)

so that $\psi_t \in \text{Span}(\sup \psi_t)$ and clearly $0 < |\sup \psi_t| \le 2^n$. So any $t \in O(n)$ induces $\sup \psi_t$, namely a set of $\lambda \in O^n(1)$ that in turn can be seen as a set of Boolean assignments (30). Applying this to our case, given any $t \in \mathcal{T}_j$, since $\mathcal{T}_j \cap O^n(1) = \mathcal{T}'_j$ [3, Lemma 1] then

$$\sup \psi_t \subseteq \mathcal{T}'_j \implies \mathcal{T}_j \subseteq \operatorname{Span}\left(\mathcal{T}'_j\right)$$

provided we identify $\psi_{\lambda} \in \mathcal{F}$ with corresponding $\lambda \in O^{n}(1)$ (more precisely $M(\psi_{\lambda}) = (\mathbb{1}, \lambda)$) and where in second relation we put \subseteq since not all linear combinations of $\psi_{\lambda} \in \mathcal{T}'_{j}$ are simple spinors. In other words $\sup \psi_{t}$ are the Boolean assignments induced by t that make the problem unsatisfiable. In following Corollary 15 we show that this generalizes to any $t \in \sum_{j} \mathcal{T}_{j}$ of Theorem 2 and that, if $t \in \sum_{j} \mathcal{T}_{j}$, all Boolean assignments of $\sup \psi_{t}$ make the problem at hand unsatisfiable.

In the next step we show that there exist simple spinors such that $|\sup \psi_t| = 2^{n-1}$ and thus, if one of them is in $\sum_{j=1}^m \mathcal{T}_j$, this excludes 2^{n-1} SAT assignments in one shot. With another $t' \in O(n)$ we can exclude the complementary 2^{n-1} terms (corresponding respectively to cases of det $t, t' = \pm 1$) and so we can conclude that if $t, t' \in \sum_{j=1}^m \mathcal{T}_j$ then $\bigcup_{j=1}^m \mathcal{T}'_j$ covers the entire Fock basis \mathcal{F} and thus the problem at hand is unsatisfiable by Proposition 7.

The advantage of the continuous formulation is now manifest: in the discrete formulation a single $\lambda \in \bigcup_{j=1}^{m} \mathcal{T}'_{j}$ excludes just *one* assignment whereas in the continuous case $t \in \sum_{j=1}^{m} \mathcal{T}_{j}$ excludes up to 2^{n-1} .

Our SAT problem is defined in the Clifford algebra of $\mathbb{R}^{n,n}$ and precisely in the spinorial representation of O(n) and since $\mathcal{C}\ell(\mathbb{R}^{n,n}) \cong \mathbb{R}(2^n)$, SAT is a problem in the algebra of real matrices of dimension $2^n \times 2^n$ and, from the computational side, the situation looks problematic. The turning point is that the spinorial representation of O(n) is equivalent to its vectorial representation corresponding to the much more manageable and familiar algebra of real $n \times n$ matrices $\mathbb{R}(n)$.

In a nutshell to any $t \in O(n)$ corresponds an orthogonal matrix $T \in \mathbb{R}(n)$ of the vectorial representation and the action of T on $u \in \mathbb{R}^n$ is given by Tu. In the spinorial representation of t in $\mathcal{C}\ell(\mathbb{R}^{n,n}) \cong \mathbb{R}(2^n)$ the action of t on uis given by

$$(-1)^{k} v_{1} v_{2} \cdots v_{k} \ u \ (v_{1} v_{2} \cdots v_{k})^{-1} \tag{39}$$

for some $v_1, v_2, \ldots, v_k \in \{0\} \times \mathbb{R}^n$, linearly independent and with $k \leq n$, this being nothing else than the Cartan theorem for Euclidean spaces in disguise [14, Theorem 5.15]: vectors v_1, v_2, \ldots, v_k give the directions of the $k \leq n$ hyperplane reflections in which any isometry of O(n) can be decomposed.

Through equivalence between the representations of O(n) and exploiting commutative diagram of Section 7, our results can be equally formulated either in $\mathbb{R}(2^n)$ or in $\mathbb{R}(n)$ (and in particular Theorem 2) but we do not insist on this.

Summarizing there are two main ingredients in this recipe for SAT: the first is that a single $t \in \sum_{j=1}^{m} \mathcal{T}_j$ can rule out 2^{n-1} assignments while the second is the equivalence between the spinorial and vectorial representations of O(n) respectively in matrix algebras $\mathbb{R}(2^n)$ and $\mathbb{R}(n)$. Moreover in this formulation there is no combinatorics since \mathcal{F} is a proper basis of the linear space of spinors \mathbb{S} and expansion (12) is unique and if an element of the Fock basis $\psi_{\lambda} \notin \sum_{j=1}^{m} \mathcal{T}_{j}$ necessarily it is a solution of the SAT problem.

8.1 The theory

We now proceed proving formally various results we will need in last part and we do it within the frame of $\mathcal{C}\ell(\mathbb{R}^{n,n}) \cong \mathbb{R}(2^n)$ that offers a more structured theoretical setting since it contains vectors, bivectors, spinors and the neatly defined Boolean algebra we exploited to formulate SAT problems.

We warn the reader that subsequent steps assume good familiarity with Clifford algebras and simple spinors but she/he can find all details in quoted references, excluding few new results and adaptations to SAT. However, through equivalence between the spinorial and vectorial representations of O(n) and exploiting commutative diagram of Section 7, *all* following results could also be equally formulated in the familiar matrix algebra $\mathbb{R}(n)$, without resorting to spinors.

We start by some very general simple spinors properties [1]:

Proposition 8. All simple spinors ψ of $\mathcal{C}\ell(\mathbb{R}^{n,n})$ can be written as

$$\psi = v_1 v_2 \cdots v_k \psi_{\mathbb{1}} \tag{40}$$

for some $v_1, v_2, \ldots v_k \in \{0\} \times \mathbb{R}^n$, linearly independent and with $k \leq n$.

This is again a consequence of the Cartan theorem for Euclidean spaces [14, Theorem 5.15] : vectors $v_1, v_2, \ldots v_k$ give the $\{0\} \times \mathbb{R}^n$ directions of the $k \leq n$ reflections in which any isometry of O(n) can be decomposed (39). We just hint that expanding each couple $v_i v_{i+1} = v_i \cdot v_{i+1} + v_i \wedge v_{i+1}$ we can expand ψ and moreover expanding each $v_i \wedge v_{i+1}$ in bivector basis $e_k e_l$

$$v_i \wedge v_{i+1} = \left(\sum_k \alpha_{i,k} e_k\right) \wedge \left(\sum_l \alpha_{i+1,l} e_l\right) = \sum_{k,l;l>k} (\alpha_{i,k} \alpha_{i+1,l} - \alpha_{i,l} \alpha_{i+1,k}) e_k e_l$$

we finally arrive at ψ expansion (12) since all $e_{i_1}e_{i_2}\cdots e_{i_r}\psi_{\mathbb{1}} \in \mathcal{F}$. This shows how the expansion of a simple spinor in the Fock basis is deeply intertwined with bivectors of $\mathcal{C}\ell(\mathbb{R}^{n,n})$ and the corresponding Lie algebra. In the vectorial representation of O(n) this explains how any $t \in SO(n)$ can be decomposed in $g \leq {n \choose 2} SO(2)$ rotations acting in subspaces $\text{Span}(e_i, e_j)$ giving the Givens expansion of t [10] but we do not insist on this.

Given any two null subspaces of $\mathbb{R}^{n,n}$ $(\mathbb{1}, t_1), (\mathbb{1}, t_2) \in \mathcal{N}_n$ they have necessarily an intersection of dimension r with $0 \leq r \leq n$, their *incidence*, that is given by all vectors $u \in \mathbb{R}^n$ such that $(u, t_1u) = (u, t_2u)$, namely $t_1u = t_2u$. We give a pivotal property of simple spinors [6, Proposition 5] here slightly adapted to our needs:

Proposition 9. Given any two linearly independent simple spinors $\psi, \phi \in \mathbb{S}_s$ then their linear combinations $\alpha \psi + \beta \phi$ ($\alpha, \beta \in \mathbb{R}$) are simple if and only if the incidence of their associated n dimensional null subspaces is n - 2 namely

$$\dim \left(M(\psi) \cap M(\phi) \right) = n - 2$$

and then $M(\psi) \cap M(\alpha \psi + \beta \phi) = M(\phi) \cap M(\alpha \psi + \beta \phi) = M(\psi) \cap M(\phi)$.

We put this result in a more usable form:

Proposition 10. Given any simple spinors ψ of $\mathcal{C}\ell(\mathbb{R}^{n,n})$ and given any two, linearly independent, $v_1, v_2 \in \{0\} \times \mathbb{R}^n$ then

$$v_1v_2\psi = (v_1 \cdot v_2 + v_1 \wedge v_2)\psi$$

is a simple spinor and if $M(\psi) = (1,t)$ then $M(v_1v_2\psi) = (1,tt_{\theta})$ where $t_{\theta} \in SO(2)$ acts in Span (v_1,v_2) ; moreover their common null subspace is

$$M(\psi) \cap M(v_1v_2\psi) = \{(u,tu) \in \mathbb{R}^{n,n} : u \in Span(v_1,v_2)^{\perp} \subset \{0\} \times \mathbb{R}^n\}$$
.

Proof. By standard properties of Clifford product $v_1 \wedge v_2$ identifies the two dimensional subspace $\text{Span}(v_1, v_2)$ and any $u \in \text{Span}(v_1, v_2)^{\perp}$ commutes with v_1v_2 . By hypothesis $M(\psi) = (\mathbb{1}, t) = \{(x, tx) \in \mathbb{R}^{n,n} : (x, tx)\psi = 0\}$ and moreover $M(v_1v_2\psi) = v_1v_2M(\psi)(v_1v_2)^{-1}$ [6] this being the spinorial representation of $t_{\theta} \in \text{SO}(2)$ acting in $\text{Span}(v_1, v_2)$ (in matrix formalism $T_{\theta}x = (\mathbb{1} - 2v_1v_1^T)(\mathbb{1} - 2v_2v_2^T)x)$. It follows that $(x, tx)\psi = 0 =$ $v_1v_2(x, tx)(v_1v_2)^{-1}v_1v_2\psi$ so that for any $u \in \text{Span}(v_1, v_2)^{\perp}$ then $(u, tu) \in$ $M(\psi) \cap M(v_1v_2\psi)$; the incidence of the two null subspaces is n - 2 and, by Proposition 9, $v_1v_2\psi$ is a simple spinor. □

All simple spinors are Weyl [6, Proposition 3], namely eigenvectors of the volume element of $\mathcal{C}\ell(\mathbb{R}^{n,n})$ of eigenvalue ± 1 : their *helicity* and with (34) given $\psi \in \mathbb{S}_s$ and $t \in \mathcal{O}(n)$ such that $M(\psi) = (\mathbb{1}, t)$ the helicity of ψ is equal to det $t = \pm 1$.

Lemma 1. Given any simple $\psi \in S_s$ and its expansion in the Fock basis (12) of $|\sup \psi| = r$ and given any generator e_i of $\mathbb{R}^{n,n}$, then $e_i\psi$: is a simple spinor of opposite helicity, $|\sup e_i\psi| = r$ and $\sup \psi \cap \sup e_i\psi = \emptyset$.

Proof. Given $\psi \in \mathbb{S}_s$ of given helicity and its Fock basis expansion (12), being Fock basis elements simple spinors themselves, it is easy to see that all $\psi_{\lambda} \in \sup \psi$ must have the same helicity of ψ . For any generator $e_i, e_i \psi$ is a simple spinor of opposite helicity [6] and thus all its Fock basis elements also have opposite helicity thus proving that $\sup \psi \cap \sup e_i \psi = \emptyset$. To prove that the size of the support of the two spinors is identical we remark that for any ψ_{λ} of (12) there exist one and only one $e_i \psi_{\lambda} \in \mathcal{F}$ (that in the language of [2] has opposite h and g-signatures with respect to ψ_{λ} and thus same $h \circ g$ -signature being in the same spinor space). \Box

Proposition 11. For any n > 1 in $\mathcal{C}\ell(\mathbb{R}^{n,n})$ there exist infinite simple spinors $\psi \in S_s$ (34) such that $|\sup \psi| = 2^{n-1}$.

Proof. We proceed by induction on n starting from n = 2, in this case a Fock basis of spinor space is given by e.g. [2]

$$\mathcal{F} = \{q_1 q_2, q_1 p_2 q_2, p_1 q_1 q_2, p_1 q_1 p_2 q_2\}$$

and thus $S = \text{Span}(q_1 q_2, q_1 p_2 q_2, p_1 q_1 q_2, p_1 q_1 p_2 q_2)$. By Proposition 9 any spinor of the linear subspace $\text{Span}(q_1 q_2, p_1 q_1 p_2 q_2)$ is simple: e.g. $\psi = \cos \frac{\theta}{2} q_1 q_2 + \sin \frac{\theta}{2} p_1 q_1 p_2 q_2$ is simple with $M(\psi) = (\mathbb{1}, \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix})$ moreover $e_2 \psi = (p_2 + q_2) \psi = \sin \frac{\theta}{2} p_1 q_1 q_2 - \cos \frac{\theta}{2} q_1 p_2 q_2$ is simple with $M(e_2\psi) = (\mathbb{1}, \begin{pmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{pmatrix})$ and so the proposition is true for n = 2. We remark that the helicities of ψ and $e_2\psi$ are opposite and that the determinant of $M(\psi)$ and $M(e_2\psi)$ are respectively 1 and -1. For the induction step let the proposition be true for n-1 and let ϕ be a simple spinor of $\mathcal{C}\ell(\mathbb{R}^{n-1,n-1})$ of $|\sup \phi| = 2^{n-2}$ and we move to $\mathcal{C}\ell(\mathbb{R}^{n,n})$: here $\varphi = \phi p_n q_n$ is a simple spinor of support 2^{n-2} with $M(\varphi) = M(\phi) \oplus \mathbb{R}\{p_n\}$ how it is simple to check. Let $v = \alpha e_n + \beta e_{n-1}$ with $\alpha, \beta \in \mathbb{R}$ and $\alpha\beta \neq 0$ and with $e_{n-1}\phi$ with support of size 2^{n-2} in $\mathcal{C}\ell(\mathbb{R}^{n-1,n-1})$ and opposite helicity to ϕ . Spinor

$$\psi = e_n v \varphi = e_n v \phi p_n q_n = (e_n \cdot v + e_n \wedge v) \phi p_n q_n = (\alpha + \beta e_n e_{n-1}) \phi p_n q_n = \\ = \alpha \phi p_n q_n - \beta e_{n-1} \phi q_n$$

where in last equality we exploited the fact that e_n has no effect on spinor ϕ of $\mathcal{C}\ell(\mathbb{R}^{n-1,n-1})$ and, without loss of generality, we have assumed that e_n commutes with ϕ . The two final terms have equal helicities and by induction hypothesis have support in Fock basis of $\mathcal{C}\ell(\mathbb{R}^{n,n})$ of size 2^{n-1} . The spinor

$$e_n\psi = e_n(\alpha\phi p_nq_n - \beta e_{n-1}\phi q_n) = \alpha\phi q_n + \beta e_{n-1}\phi p_nq_n$$

is of opposite helicity and covers the missing half of the Fock basis of $\mathcal{C}\ell(\mathbb{R}^{n,n})$ thus completing the proof. \Box

Thus given any $\varphi \in S_s$ such that $|\sup \varphi| = 2^{n-1}$ by Lemma 1 we have $\sup \varphi \oplus \sup e_i \varphi = \mathcal{F}$. We already defined before (31) the *reference* spinor $\psi_1 = p_1 q_1 p_2 q_2 p_3 q_3 \cdots p_n q_n$; clearly $\psi_1 \in \mathcal{F}$ and $M(\psi_1) = \text{Span}(p_1, p_2, \dots, p_n) = P = (1, 1)$ and with ψ_1 we build explicitly one of these spinors:

Corollary 12. For any n > 1 given $\mathcal{C}\ell(\mathbb{R}^{n,n})$ the simple spinor

$$\varphi := \prod_{i=1}^{n-1} (\mathbb{1} + e_i e_{i+1}) \psi_{\mathbb{1}}$$

$$\tag{41}$$

has $|\sup \varphi| = 2^{n-1}$, moreover $\sup \varphi \oplus \sup e_i \varphi = \mathcal{F}$.

Proof. Instead of proceeding by induction like in Proposition 11 we give a simpler proof observing that the product contains n-1 terms in parenthesis each of them being the sum of identity and bivector $e_i e_{i+1}$ so that the full expansion of φ contains all the 2^{n-1} subsets of the product obtained choosing one of the two terms in each parentheses and observing that it is impossible to get complete cancellations between bivector terms. \Box

We remark that φ definition (41) implicitly defines also the succession

$$\varphi_{k} = \prod_{i=1}^{k-1} (\mathbb{1} + e_{i}e_{i+1})\psi_{\mathbb{1}} \qquad \varphi_{k} = (\mathbb{1} + e_{k-1}e_{k})\varphi_{k-1} \qquad (42)$$

where we assume $\varphi_1 := \psi_1$ and $\varphi_n = \varphi$. Given φ the corresponding $M(\varphi) = (1, t) \in \mathcal{N}_n$ has $t = \prod_{i=1}^{n-1} t_{i,i+1}$ where $t_{i,i+1} \in \mathrm{SO}(2)$ acts in $\mathrm{Span}(e_i, e_{i+1})$ with an angle of $\pi/2$ and clearly det t = 1.

Aiming at our goal we now apply Proposition 10 to the case of $\psi = \psi_j + \psi_k \in \mathcal{T}_j + \mathcal{T}_k$ (36) and we remark that any of these spinors can be also in $\mathcal{T}_j \cup \mathcal{T}_k$ or not, this second case resulting the most interesting one since it pinpoints the difference between Proposition 7 and Theorem 2: for example it is easy to prove that for (41) $\varphi \notin \bigcup_{j=1}^m \mathcal{T}_j$. In following propositions we will keep trace of the two possibilities. We start with a technical result:

Lemma 2. Given any $t \in \mathcal{T}_j$ (35) and any $v \in \{0\} \times \mathbb{R}^n$ then $t_v := -vtv^{-1} \in \mathcal{T}_j$ if and only if $v \in Span(\sup z_j)^{\perp}$.

Proof. For any $t \in \mathcal{T}_j \ M(z_j) \subseteq (1,t)$ and its action on u is that of (39) while the action of t_v is given by $(-1)^{k+1}vv_1v_2\cdots v_k \ u \ (vv_1v_2\cdots v_k)^{-1}$. Let $v \in \operatorname{Span}(\sup z_j)^{\perp}$ then v commutes with all $e_i \in \sup z_j$ and thus $M(z_j) \subseteq (1,t_v)$, namely $t_v \in \mathcal{T}_j$. Conversely given $t_v \in \mathcal{T}_j$ necessarily $M(z_j) \subseteq (1,t_v)$ and since we know that also $M(z_j) \subseteq (1,t)$ it follows that v commutes with all $e_i \in \sup z_j$ and thus, necessarily, $v \in \operatorname{Span}(\sup z_j)^{\perp}$. \Box

Proposition 13. Given two clauses z_j, z_k with induced sets $\mathcal{T}_j, \mathcal{T}_k$ then $\psi = \alpha \psi_j + \beta \psi_k \in \mathcal{T}_j + \mathcal{T}_k$ $(\alpha, \beta \in \mathbb{R})$ if and only if the incidence of $M(\psi_j), M(\psi_k)$ is n-2 and thus if $\psi = v_1 v_2 \psi_j$ like in Proposition 10 and thus $v_1 \wedge v_2 \psi_j \in \mathcal{T}_k$ and in particular only in the three following cases depending on the number of common opposite generators (Boolean variables) of z_j and z_k :

- no common, opposite generators (but possibly equal ones): if $v_1, v_2 \in Span(\sup z_j \cap \sup z_k)^{\perp}$ and in this case in general $\psi \in \mathcal{T}_j \cup \mathcal{T}_k$. If moreover $v_1 \in Span(\sup z_j \sup z_j \cap \sup z_k)$ and $v_2 \in Span(\sup z_k \sup z_j \cap \sup z_k)$ then $\psi \notin \mathcal{T}_j \cup \mathcal{T}_k$,
- 1 common, opposite generator e_i : if $v_1 = e_i$ and $v_2 \in Span(\sup z_j \cap \sup z_k \{e_i\})^{\perp}$,
- 2 common, opposite generators e_i, e_l : if $v_1 = e_i$ and $v_2 = e_l$.

Proof. We start observing that $\mathcal{T}_j \cap \mathcal{T}_k = \emptyset$ if and only if z_j and z_k have one or more common opposite generators since for any $\psi \in \mathcal{T}_j \cap \mathcal{T}_k$ then both $M(z_j), M(z_k) \subseteq M(\psi)$ and this may happen only if clauses are compatible, namely only if they have no common opposite generators (they still can have common equal generators). For the purposes of this Proposition cases with more than two opposite generators are excluded because incidence n-2would be impossible.

If there is one common, opposite coordinate e_i necessarily it is not in the common part and by Proposition 10 there must exist another v_2 , linearly independent from $v_1 = e_i$, such that $v_1 \wedge v_2 \psi_j \in \mathcal{T}_k$. Any common coordinate of $\sup z_j \cap \sup z_k - \{e_i\}$ is necessarily equal and thus in common part $M(\psi_j) \cap M(\psi_k)$ and by Proposition 10 follows $v_2 \in \text{Span}(\sup z_j \cap \sup z_k - \{e_i\})^{\perp}$; in this case, because of opposite e_i , for $\alpha\beta \neq 0$ $\psi \notin \mathcal{T}_j \cup \mathcal{T}_k$.

For two common opposite coordinates, namely e.g. e_i, e_l then, being incidence of $M(\psi_j)$ and $M(\psi_k)$ n-2 and being necessarily e_i, e_l not in the incidence part, by Proposition 10, necessarily $v_1 \wedge v_2 \psi_j = e_i e_l \psi_j \in \mathcal{T}_k$ and also in this case for $\alpha \beta \neq 0 \ \psi \notin \mathcal{T}_j \cup \mathcal{T}_k$.

We conclude with the case of no common opposite generators, namely $\mathcal{T}_j \cap \mathcal{T}_k \neq \emptyset$: given $\psi = \alpha \psi_j + \beta \psi_k \in \mathcal{T}_j + \mathcal{T}_k$ by Proposition 10 $\psi = v_1 v_2 \psi_j$ and the common part is given by all $(u, tu) \in \mathbb{R}^{n,n}$ with $u \in \text{Span}(v_1, v_2)^{\perp} \subset \{0\} \times \mathbb{R}^n$ it follows that any $v_1, v_2 \in \text{Span}(\sup z_j \cap \sup z_k)^{\perp}$ can do since $\sup z_j \cap \sup z_k$ can contain only common equal generators, excluded by Proposition 10.

We remark that choosing $v_1, v_2 \in (\sup z_j)^{\perp}$ by Lemma 2 $\psi \in \mathcal{T}_j$ (and similarly for \mathcal{T}_k) and so in general in these cases $\psi \in \mathcal{T}_j \cup \mathcal{T}_k$. On the other hand $\psi \notin \mathcal{T}_j \cup \mathcal{T}_k$ (36) if and only if $M(z_j), M(z_k) \not\subset M(\psi)$ that by Lemma 2 requires e.g. $v_1 \in \text{Span}(\sup z_j - \sup z_j \cap \sup z_k)$ and $v_2 \in$ Span $(\sup z_k - \sup z_j \cap \sup z_k)$.

To prove the converse we remark that in all cases $v_1 \wedge v_2 \psi_i \in \mathcal{T}_k$. \Box

Corollary 14. Given two clauses z_j, z_k with their induced sets $\mathcal{T}_j, \mathcal{T}_k$ then $\mathcal{T}_j + \mathcal{T}_k \neq \emptyset$ (36) if and only if the clauses have 0,1 or 2 common opposite generators.

Some other technical Lemmas:

Lemma 3. Given a SAT problem let J be any non empty subset of the m clauses, then

$$\left(\sum_{j\in J}\mathcal{T}_j\right)\cap O^n(1)=\cup_{j\in J}\mathcal{T}'_j$$

Proof. We already know that $\mathcal{T}_j \cap \mathcal{O}^n(1) = \mathcal{T}'_j$ [3, Lemma 1] so we just need to prove that $(\mathcal{T}_j + \mathcal{T}_k) \cap \mathcal{O}^n(1) = \mathcal{T}'_j \cup \mathcal{T}'_k$ but this follows trivially from the definition of $\mathcal{T}_j + \mathcal{T}_k$ (36) and from the fact that \mathcal{F} is a proper basis of \mathbb{S} and thus all elements of $\mathcal{T}_j + \mathcal{T}_k$ are necessarily in Span $(\mathcal{T}'_j \cup \mathcal{T}'_k)$ and the relation is correctly = (and not only \subseteq) since all $\psi_\lambda \in \mathcal{T}'_j$ are in \mathcal{T}_j . \Box

Corollary 15. Given $\psi \in S_s$ such that $\psi \in \sum_{j \in J} T_j$, where J is any non empty subset of the m clauses, then

$$\sup \psi \subseteq \cup_{j \in J} \mathcal{T}'_j \ .$$

Thus ψ excludes all Boolean assignments of $\sup \psi$ since any $\psi_{\lambda} \in \sup \psi$ is necessarily in at least one \mathcal{T}'_{j} and thus is an assignment that renders F the problem at hand and thus also φ (41) excludes 2^{n-1} assignments.

With (36) we can reformulate Theorem 2 in S_s exploiting the one to one correspondence between simple spinors and O(n) but we put the same result in a form more amenable to an actual algorithm testing unsatisfiability:

Theorem 3. A given SAT problem is unsatisfiable if and only if given a single $\varphi \in S_s$ with $|\sup \varphi| = 2^{n-1}$ like (41) and any single e_i then the isometries induced by its m clauses (35) are such that:

$$\varphi, e_i \varphi \in \sum_{j=1}^m \mathcal{T}_j \quad . \tag{43}$$

Proof. Supposing that (43) holds this implies that the Fock basis expansion of φ , with $|\sup \varphi| = 2^{n-1}$, is in $\sum_{j=1}^{m} \mathcal{T}_j$ and by Corollary 15 $\sum_{j=1}^{m} \mathcal{T}_j$ covers the first half of the Fock basis. Repeating the same procedure with $e_i\varphi$ by Lemma 1 we cover the second half of the Fock basis and thus $\sum_{j=1}^{m} \mathcal{T}_j$ contains the full Fock basis and the problem is unsatisfiable by Proposition 7; the converse follows immediately from same Proposition. \Box

An unsatisfiability test exploiting Theorem 3 in essence requires:

- build $\varphi \in S_s$ of maximal support (we did it in Corollary 12),
- verify whether $\varphi \stackrel{?}{\in} \sum_{j=1}^{m} \mathcal{T}_{j}$ (we will do it in Proposition 16),
- perform the same test on $e_i\varphi$.

and if both tests succeed we can conclude that the SAT problem at hand is unsatisfiable. The remarkable fact is that we will not need to check the 2^{n-1} Fock basis components of φ (41) but we can exploit the recursive construction of the succession φ_k (42) breaking the initial problem into *n* subproblems each of them corresponding to the application of Propositions 10 and 13 to φ_k , in turn corresponding to the Givens expansion of φ .

The crucial pending question is the computational complexity of checking whether φ is in $\sum_{j=1}^{m} \mathcal{T}_{j}$ that we address in next section and we start recalling some standard properties of \mathbb{R}^{n} and of its isometry group O(n).

It is well known that any $t \in O(n)$ has only 3 eigenvalues: ± 1 and couples of complex conjugates and that all eigenvectors corresponding to different eigenvalues are reciprocally orthogonal. Moreover any $t \in O(n)$ having only ± 1 eigenvalues is an *involution*: $t^2 = 1$ and in $\mathbb{R}(n)$ $T = T^T$, a prominent, but not exhaustive, example being the 2^n involutions $\lambda \in O^n(1)$.

It follows that any $t \in O(n)$ defines univocally three reciprocally orthogonal subspaces of \mathbb{R}^n corresponding respectively to eigenvectors of ± 1 eigenvalues and to their orthogonal complement and at least one of these subspaces have non null dimensions. These subspaces are in close connection with the Wall parametrization of $t \in O(n)$ [15, Chapter 11] but we do not insist on this here.

We define inv (t) to be the *involutory subspace* of t, namely the subspace of \mathbb{R}^n spanned by the eigenvectors corresponding to ± 1 eigenvalues,

namely inv $(t) = \ker(t + 1) \oplus \ker(t - 1)$; obviously inv $(t)^{\perp}$ is the subspace corresponding to complex eigenvalues and

$$\operatorname{inv}(t) \oplus \operatorname{inv}(t)^{\perp} = \ker(t+1) \oplus \ker(t-1) \oplus \operatorname{inv}(t)^{\perp} = \mathbb{R}^{n}$$

and moreover the restrictions of t to any couple of these subspaces commute.

Applying this definition to our case for any $t \in \mathcal{T}_j$ then $\text{Span}(\sup z_j) \subseteq \text{inv}(t)$ where the inclusion can be strict since inv(t) can contain other directions in subspace $\text{Span}(\sup z_j)^{\perp}$. Since for our problem all $\sup z_j$ are subsets of generators we will focus here on cases in which inv(t) is made by subset of generators but in general it can contain any $\{0\} \times \mathbb{R}^n$ direction.

Given $t \in \mathcal{T}_j$ for any $e_i \in \sup z_j$ it follows $te_i = \pm e_i$ and thus for the corresponding simple spinor ψ_t that either p_i or q_i are in $M(\psi_t)$ (29) and thus all $\psi_{\lambda} \in \sup \psi_t$ share corresponding fixed coordinates $p_i q_i$ or q_i the proof being that p_i or q_i is in $M(\psi_t)$ and in all $M(\psi_{\lambda})$ of $\sup \psi_t$.

We thus define also for simple spinor ψ , inv (ψ) as the subset of fixed coordinates $p_i q_i$ or q_i in $\psi_{\lambda} \in \sup \psi$ corresponding to p_i or q_i in $M(\psi)$ and in all $M(\psi_{\lambda})$ of its support.

8.2 An actual algorithm

We put at work the theory of previous section to outline an algorithm that tests unsatisfiability and we start characterizing φ (41) and its associated succession (42).

Lemma 4. Given φ_k (42) then $p_{k+1}, p_{k+2}, \ldots, p_n \in M(\varphi_k)$ and thus $inv(\varphi_k) = p_{k+1}q_{k+1} p_{k+2}q_{k+2} \cdots p_n q_n$. The corresponding $t_k \in O(n)$, such that $M(\varphi_k) = (\mathbb{1}, t_k)$, has $inv(t_k) = Span(e_{k+1}, e_{k+2}, \ldots, e_n)$ its non involutory part being $Span(e_1, e_2, \ldots, e_k)$.

Proof. That $p_{k+1}, p_{k+2}, \ldots, p_n \in M(\varphi_k)$ is clear from φ_k definition (42) since in the product there are no terms with $\{e_{k+1}, e_{k+2}, \ldots, e_n\}$ that can alter $\psi_{1} = p_1q_1 \ p_2q_2 \cdots p_nq_n$ so we need just to prove that for k > 1 all $v \in \text{Span}(e_1, e_2, \ldots, e_k)$ are non involutory, namely that $t_k v \neq \pm v$ that we prove by induction on k: for k = 2 it is well known that for any $t_2 \in \text{SO}(2)$ inv $(t_2) = \{0\}$; supposing this true for k-1 we move to $\varphi_k = (1+e_{k-1}e_k)\varphi_{k-1}$ and thus $t_k = t_{\frac{\pi}{2}}t_{k-1}$ with $t_{\frac{\pi}{2}} \in \text{SO}(2)$ acting in $\text{Span}(e_{k-1}, e_k)$: should there exist v such that $t_k v = \pm v$ we would have $t_{k-1}v = \pm t_{-\frac{\pi}{2}}v$ and we can write $v = \alpha e_k + \beta e_{k-1} + v_2$ with $v_2 \in \text{Span}(e_1, e_2, \ldots, e_{k-2})$ and thus we would have $t_{-\frac{\pi}{2}}(\alpha e_k + \beta e_{k-1} + v_2) = \alpha' e_k + \beta' e_{k-1} + v_2$ where v_2 is unmodified since $t_{-\frac{\pi}{2}} \arctan (e_{k-1}, e_k)$ and it should be equal to $t_{k-1}(\alpha e_k + \beta e_{k-1} + v_2) =$ $\alpha e_k + t_{k-1}(\beta e_{k-1} + v_2)$ since t_{k-1} acts in $\text{Span}(e_1, e_2, \ldots, e_{k-1})$ and thus to have equality we should have $\alpha = \pm \alpha'$ that, since inv $(t_{-\frac{\pi}{2}}) = \{0\}$, can be realized only if $\alpha = \beta = 0$ that would imply that $t_{k-1}v_2 = \pm v_2$ against induction hypothesis. \Box **Lemma 5.** Given φ_k (42) for k > 1 then $\varphi_k \in \mathcal{T}_j$ if and only if $\sup z_j \subseteq \{e_{k+1}, e_{k+2}, \ldots, e_n\}$ with all generators of z_j in positive form.

Proof. $\varphi_k \in \mathcal{T}_j$ if and only if $M(z_j) \subseteq M(\varphi_k)$ and being the involutory part of φ_k such that $\text{Span}(p_{k+1}, p_{k+2}, \ldots, p_n) \subseteq M(\varphi_k)$ it follows $M(z_j) \subseteq \text{Span}(p_{k+1}, p_{k+2}, \ldots, p_n)$ and thus the thesis; the converse is immediate. \Box

We remark that the case $\varphi_1 = \psi_1$ is different since in this case the condition is $\sup z_j \subseteq \{e_1, e_2, \ldots, e_n\}$.

Lemma 6. Given φ_k (42) then $\varphi_k \in \sum_{j \in J} \mathcal{T}_j$, $J \subseteq \{1, 2, \ldots, m\}$ being a subset of clauses, if and only if for all $l \in [1, k]$, $\varphi_l \in \sum_{j \in J} \mathcal{T}_j$, moreover $\sup \varphi_1 \subset \sup \varphi_2 \subset \cdots \subset \sup \varphi_k$ of sizes $1, 2, \ldots, 2^{k-1}$.

Proof. Let $\varphi_k \in \sum_{j \in J} \mathcal{T}_j$, we just saw that $p_k q_k$ is a fixed coordinate of φ_{k-1} and thus also in all $\psi_{\lambda} \in \sup \varphi_{k-1}$ and thus in $\sup e_{k-1}e_k\varphi_{k-1}$ the *k*-th coordinate is q_k : this proves that $\sup \varphi_{k-1} \subset \sup \varphi_k$ and that $|\sup \varphi_k| = 2|\sup \varphi_{k-1}|$ and thus for any $l \in [1, k] \sup \varphi_l \subset \sup \varphi_k$ that proves the result; the converse is immediate. \Box

We are interested to study cases in which φ_k belongs to a sum of \mathcal{T}_j and we start with the simplest example, namely the sum of two clauses $\mathcal{T}_j + \mathcal{T}_l$. We purposely exclude the trivial case in which both φ_k and φ_{k-1} belongs to a single \mathcal{T}_j , by Lemma 5 the case in which $\sup z_j \subseteq \{e_{k+1}, e_{k+2}, \ldots, e_n\}$ all in positive form.

Lemma 7. Given $\varphi_k = (\mathbb{1} + e_{k-1}e_k)\varphi_{k-1}$ and \mathcal{T}_j and \mathcal{T}_l such that $\varphi_{k-1} \begin{cases} \in \mathcal{T}_j \\ \notin \mathcal{T}_l \end{cases}$ and $e_{k-1}e_k\varphi_{k-1} \begin{cases} \notin \mathcal{T}_j \\ \in \mathcal{T}_l \end{cases}$ then $\varphi_k \in \mathcal{T}_j + \mathcal{T}_l$ with the involutory part containing $z_j \cup z_l - \{e_k\}$ namely a set that contains all the generators of z_j and z_l but e_k , all in positive form. We can thus associate to φ_k a composed clause $z_j \cup z_l - \{e_k\}$ that contains a number of generators strictly lower than $|\sup z_j| + |\sup z_l|.$

Proof. By Lemma 5 sup $z_j \subseteq \{e_k, e_{k+1}, \ldots, e_n\}$ and since $e_{k-1}e_k\varphi_{k-1} \notin \mathcal{T}_j$ by Lemma 2 $e_k \in \sup z_j$ in positive form and thus, since $e_{k-1}e_k\varphi_{k-1} \in \mathcal{T}_l$ then also $e_k \in \sup z_l$ but in negative form. This does not violate Lemma 5 that does not apply to $e_{k-1}e_k\varphi_{k-1}$. It follows that z_j and z_l are clauses with one common opposite generator e_k , while they can have common equal, positive generators in $\{e_{k+1}, e_{k+2}, \ldots, e_n\}$, and Propositions 10 and 13 apply. For any $\psi = \alpha\psi_j + \beta\psi_l \in \mathcal{T}_j + \mathcal{T}_l (\alpha, \beta \in \mathbb{R}), \psi = v_1v_2\psi_j$ with $v_1 \wedge v_2\psi_j \in \mathcal{T}_l$ and moreover for $v_1 = e_k, v_2 = e_k - e_{k-1}$ we get exactly $v_1v_2\varphi_{k-1} = (\mathbb{1} + e_{k-1}e_k)\varphi_{k-1}$. By Proposition 10 Span $(v_1, v_2) =$ Span (e_{k-1}, e_k) identifies the non incident part of the two null subspaces induced by the simple spinors being summed while all other directions must be in common. It follows that all generators $z_j \cup z_k - \{e_k\}$ must be present in the involutory part of the sum, their number being $|\sup z_j| + |\sup z_l| - 2$ or less if there are also common equal generators. \Box

We can thus easily generalize the set \mathcal{T}_j (35) associated to clause z_j to a set \mathcal{T}_x associated to an arbitrary, given, involutory part z_x ,

$$\mathcal{T}_x := \{ \psi \in \mathbb{S}_s : M(z_x) \subseteq M(\psi) \} \subset \mathbb{S}_s$$

the involutory part typically being, like in previous Lemma, associated to a sum of simple spinors that builds φ_k

$$z_x := z_j \cup z_l - \{e_k\} \tag{44}$$

that we define a *composed* clause and gives the involutory part of $\alpha \psi_j + \beta \psi_l$. Proceeding to successive φ_k we will use \mathcal{T}_x while keeping separate track of the origin of the terms of the sum: $\varphi_k \in \mathcal{T}_j + \mathcal{T}_l$. We remark that given $\psi = \psi_j + \psi_l \in \mathcal{T}_j + \mathcal{T}_l$ when we want to iterate the procedure and add another simple spinor e.g. $\phi \in \mathcal{T}_r$ to get $\phi + \psi$, Proposition 10 applies again and thus necessarily $\phi + \psi = (\mathbb{1} + u_1 u_2)\psi = (\mathbb{1} + u_1 u_2)(\mathbb{1} + v_1 v_2)\psi_j$ and we can iterate previous Lemma to get that the involutions associated to $\phi + \psi$ will be $((z_j \cup z_l - \{e_k\}) \cup z_r - \{e_{k+1}\})$ where we introduced this notation with brackets to indicate that this operation is *not* associative and that in general $\phi + (\psi_j + \psi_l) \neq (\phi + \psi_j) + \psi_l$ since in general $(\phi + \psi_j)$ is not simple: for example let $\psi = \psi_1 + e_1 e_2 \psi_1 = (\mathbb{1} + e_1 e_2) \psi_1$ that is simple and we may add $\phi = e_3 e_4 \psi$ to get $\phi + \psi = (\mathbb{1} + e_3 e_4) \psi = e_3 e_4 \psi + (\psi_1 + e_1 e_2 \psi_1)$ where the parenthesis is strictly necessary since $e_3 e_4 \psi + \psi_1$ is not a simple spinor since incidence of the two terms is n - 4.

We remark that Lemma 7 is of general validity since it deals with *two* simple spinors at the time and thus that we can apply it several times to form sum of more than two addends, each being associated to a plain or composed clause. It is simple to verify that composed clauses have all properties of real clauses like e.g. Lemma 2, both being essentially identified by a given involutory part.

Proposition 16. Given $\varphi \in S_s$ (41) and m sets \mathcal{T}_j induced by clauses of a given SAT problem, checking wether

$$\varphi \stackrel{?}{\in} \sum_{j=1}^{m} \mathcal{T}_j \tag{45}$$

can be done in polynomial time.

Proof. By Lemma 6 $\varphi \in \sum_{j=1}^{m} \mathcal{T}_{j}$ if and only if all terms of the succession φ_{k} (42) are in this set and thus our procedure to verify (45) starts from $\varphi_{1} = \psi_{1}$ and look after clause z_{j} such that $\varphi_{1} \in \mathcal{T}_{j}$ and continues increasing k and searching a subset of clauses containing φ_{k} .

We start with $\varphi_1 = \psi_1$ and in $\mathcal{O}(n)$ we can find all clauses z_j with all generators in positive form such that $\varphi_1 \in \mathcal{T}_j$ (obviously there may be many; should there be none by Lemma 6 $\varphi \notin \sum_{j=1}^m \mathcal{T}_j$ and 1 is a solution for the given SAT problem). For the next step $\varphi_2 = (1 + e_1 e_2)\varphi_1$: either $e_1 e_2 \varphi_1$ is also in \mathcal{T}_j , that thus contains also φ_2 , or we have to find another clause z_l such that $e_1 e_2 \varphi_1 \in \mathcal{T}_l$. Again: if $\varphi_1 \in \mathcal{T}_l$ then both φ_1 and φ_2 are in \mathcal{T}_l , if $\varphi_1 \notin \mathcal{T}_l$ then $\varphi_2 \in \mathcal{T}_j + \mathcal{T}_l$. We remark that φ_2 case is the only case (with respect to Lemma 7) in which z_j and z_l may have also 2 common opposite generators, see Proposition 13, but also in this case it is easy to define the associated composed clause $z_x := z_j \cup z_l - \{e_1, e_2\}$. If such a clause z_l do not exist then $e_1 e_2 \varphi_1$ is a solution and we can stop here. At worst the time to find $\varphi_2 \in \mathcal{T}_x$ is $\mathcal{O}(n^2)$ since we look for a couple of clauses z_j and z_l with one or two common opposite generators.

Before going to cases φ_k with k > 2 we remark that we will need composed clauses derived from repeated applications of Lemma 7 and we can start calculating, preliminarly and once for all, the composed clauses we can form with our sets \mathcal{T}_i by repeated application of composition rule (44).

To do this we define an undirected graph of adjacency matrix G with a node for each clause and an edge between clauses that have a common opposite generator. Let there exists an edge between clauses z_j and z_l with common opposite generator e_i : this edge induces a composed clause given by $z_j \cup z_l - \{e_i\}$. If moreover z_l has an edge with z_r , with different common opposite generator e_j , this second edge induces a further composed clause given by $((z_j \cup z_l - \{e_i\}) \cup z_r - \{e_j\})$ since $e_j \in \sup z_l$ is also $e_j \in \sup(z_j \cup z_l - \{e_i\})$.

This process can be iterated and it is simple to see that all composed clauses can be obtained in this fashion. The composed clauses obtained in r iterations are given by matrix powers G^r that is an $\mathcal{O}(n^3)$ process. We remark that G^r contains also illegally composed clauses in which the same common opposite generator is used more than once. The maximum power of the adjacency matrix is n since there are at maximum n different common opposite generators.

For any r, G^r contains $\mathcal{O}(n^2)$ composed clauses and so all possible composed clauses are $\mathcal{O}(n^3)$ and to calculate them we repeated n times an $\mathcal{O}(n^3)$ matrix multiplication. In summary to build the $\mathcal{O}(n^3)$ composed clauses we use $\mathcal{O}(n^4)$ time but this process is done just once.

We are now ready to go to the generic case of φ_k and we assume $\varphi_{k-1} \in \mathcal{T}_x$, z_x being a plain or composed clause. There are only three mutually exclusive possibilities that we examine in succession:

- 1. $e_{k-1}e_k\varphi_{k-1} \in \mathcal{T}_x$, by Lemma 2 this happens if and only if $e_k \notin \sup z_x$ that can be easily verified in $\mathcal{O}(n)$ simply searching $\sup z_x$;
- 2. there exists one plain or a composed clause z_y , such that $e_{k-1}e_k\varphi_{k-1} \in \mathcal{T}_y$. Since we are not in case 1, $e_{k-1}e_k\varphi_{k-1} \notin \mathcal{T}_x$ and there are only

two possibilities: if also $\varphi_{k-1} \in \mathcal{T}_y$ then both φ_{k-1} and φ_k are in \mathcal{T}_y and this can be verified in $\mathcal{O}(n)$ checking involutory parts of z_y and of φ_{k-1} . If $\varphi_{k-1} \notin \mathcal{T}_y$ then we are in the conditions of Lemma 7 and to prove that $\varphi_k \in \mathcal{T}_x + \mathcal{T}_y$ we have to find z_y , plain or composed, such that $e_{k-1}e_k\varphi_{k-1} \in \mathcal{T}_y$ (we assume to be in cases with k > 2where Lemma 7 applies strictly). Any clause z_y , plain or composed, with the following characteristics: no generators in $\{e_1, e_2, \ldots, e_{k-1}\}$, generator e_k in negative form and any generator in $\{e_{k+1}, e_{k+2}, \ldots, e_n\}$ in positive form, constitutes a valid candidate. If such a z_y exists it is necessarily in the set $\{G, G^2, \ldots, G^n\}$ and can be found in $\mathcal{O}(n^3)$;

3. if neither of previous cases occurred than we have a certificate that $e_{k-1}e_k\varphi_{k-1} \notin \sum_{j=1}^m \mathcal{T}_j$ and thus φ_k and by Lemma 6 also φ and we can stop searching.

So at every level checking for φ_k is an $\mathcal{O}(n^3)$ task and the process repeated for n levels is at most an $\mathcal{O}(n^4)$ task. \Box

In summary we proposed a proof-of-concept algorithm working for both 2 and 3SAT problems. It is intriguing that it shows loose similarities with *resolution*, when it composes two clause with one common opposite literal, and with *transitive closure*, when it builds the graph of composed clauses.

Dedication

I dedicate this paper to the memory of my father Paolo Budinich who passed away in November 2013 not before transferring me his overwhelming enthusiasm for simple spinors, at the heart of this work. 17 years ago we moved together the first steps along this winding track [5].

References

- [1] Marco Budinich. On spinors and null vectors. Journal of Physics A: Mathematical and Theoretical, 47(11):115201, March 2014.
- Marco Budinich. On spinors transformations. Journal of Mathematical Physics, 57(7):071703-1-11, July 2016. arXiv:1603.02181 [math-ph] 7 Mar 2016.
- [3] Marco Budinich. The Boolean SATisfiability Problem and the orthogonal group O(n), April 2019. arXiv:1904.02629 [math.CO].
- Marco Budinich. The Boolean SATisfiability Problem in Clifford algebra. *Theoretical Computer Science*, 784:1–10, September 2019. arXiv:1704.02942v3 [math-ph] 17 May 2018.
- [5] Marco Budinich and Paolo Budinich. A spinorial formulation of the maximum clique problem of a graph. *Journal of Mathematical Physics*, 47(4):043502, 2006.
- [6] Paolo Budinich and Andrzej Mariusz Trautman. Fock Space Description of Simple Spinors. Journal of Mathematical Physics, 30(9):2125– 2131, September 1989.
- [7] Élie Cartan. The Theory of Spinors. Hermann, Paris, 1966. first edition in French: La theorie des spineurs I and II, Actuatités Sci. et Industr., No. 643 and 701, Hermann, Paris, 1938.
- [8] Stephen A. Cook. The Complexity of Theorem-proving Procedures. In Proceedings of the Third Annual ACM Symposium on Theory of Computing, STOC '71, pages 151–158, New York, NY, USA, 1971. ACM.
- Steven Givant and Paul Halmos. Introduction to Boolean Algebras. Undergraduate Texts in Mathematics. Springer-Verlag, Springer-Verlag New York, 2009.
- [10] David K. Hoffman, Richard C. Raffenetti, and Klaus Ruedenberg. Generalization of euler angles to n-dimensional orthogonal matrices. *Jour*nal of Mathematical Physics, 13(4):528–533, 1972.
- [11] Edward V. Huntington. New sets of independent postulates for the algebra of logic, with special reference to Whitehead and Russell's Principia Mathematica. *Transactions of the American Mathematical Soci*ety, 35:274–304, 1933.
- [12] Donald Ervin Knuth. The Art of Computer Programming. Combinatorial Algorithms, volume IV. Addison-Wiley, Reading, MA, release in fascicles edition, 2015.

- [13] Ramamohan Paturi, Pavel Pudlák, Michael E. Saks, and Francis Zane. An Improved Exponential-time Algorithm for k-SAT. *Journal of the* ACM, 52(3):337–364, May 2005.
- [14] Ian Robertson Porteous. Clifford Algebras and the Classical Groups. Cambridge Studies in Advanced Mathematics: 50. Cambridge University Press, 1995.
- [15] Donald E. Taylor. The geometry of the classical groups, volume 9 of Sigma series in pure mathematics. Heldermann Verlag, Berlin, 1992.